

The complexity of the Quantified CSP having the polynomially generated powers property.

Dmitriy Zhuk

October 19, 2021

Abstract

It is known that if an algebra of polymorphisms of the constraint language has the Polynomially Generated Powers (PGP) Property then the Quantified CSP can be reduced to the CSP over the same constraint language with constants. The only limitation of this reduction is that it is applicable only for the constraint languages with constants. We drastically simplified the reduction and generalized it for constraint languages without constants. As a result, we completely classified the complexity of the QCSP for constraint languages having the PGP property.

1 Introduction

Let Γ be a finite set of relations on a set A . *The Constraint Satisfaction Problem over the constraint language Γ* , denoted by $\text{CSP}(\Gamma)$, is the following decision problem: given a formula of the form

$$R_1(v_{1,1}, \dots, v_{1,n_1}) \wedge \dots \wedge R_s(v_{s,1}, \dots, v_{s,n_s}),$$

where $R_1, \dots, R_s \in \Gamma$ and each $v_{i,j} \in \{x_1, \dots, x_n\}$; decide whether the formula is satisfiable. A natural generalization of the CSP is the *Quantified Constraint Satisfaction Problem (QCSP)*, where we allow to use both existential and universal quantifiers. Formally, for a constraint language Γ , $\text{QCSP}(\Gamma)$ is the problem to evaluate a sentence of the form

$$\forall x_1 \exists y_1 \dots \forall x_n \exists y_n R_1(v_{1,1}, \dots, v_{1,n_1}) \wedge \dots \wedge R_s(v_{s,1}, \dots, v_{s,n_s}),$$

where $R_1, \dots, R_s \in \Gamma$, and $v_{i,j} \in \{x_1, \dots, x_n, y_1, \dots, y_n\}$ for every i, j (see [1, 7, 9, 10]).

Unlike CSP, in general the Quantified CSP can be PSpace-hard and the complexity of $\text{QCSP}(\Gamma)$ is not known. In this paper we consider one of the main ideas that allows to reduce the QCSP to the usual CSP and therefore reduce the complexity. To explain the idea let us consider an easier problem. By QCSP^{Π_2} we denote the Π_2 -restriction of the QCSP, that is, given a sentence

$$\forall x_1 \dots \forall x_n \exists y_1 \dots \exists y_m \Phi \tag{1}$$

where Φ is a conjunction of relations from Γ ; decide whether it holds. To solve this problem it suffices to check that Φ has a solution for all evaluations of x_1, \dots, x_n , hence it suffices to look at (the conjunction of) $|A|^n$ instances of $\text{CSP}(\Gamma^*)$, where $\Gamma^* = \Gamma \cup \{(x = a) \mid a \in A\}$. It is not hard to show (see [8]) that, if A^n can be generated under $\text{Pol}(\Gamma)$ from some subset $\Sigma \subseteq A^n$, then it is sufficient to check only evaluations from Σ , and to solve $|\Sigma|$ instances of $\text{CSP}(\Gamma^*)$. Hence, if Σ can be chosen small enough and can be efficiently computed then this simple idea gives a polynomial reduction to the CSP.

Example. Suppose $A = \{0, 1\}$ and Γ consists of linear equations in \mathbb{Z}_2 . Let us check that the instance (1) holds for $(x_1, \dots, x_n) = (0, \dots, 0)$ and $(x_1, \dots, x_n) = (0, \dots, 0, 1, 0, \dots, 0)$ for any position of 1. To do this we just solve $n + 1$ CSP instances of $\text{CSP}(\Gamma^*)$ and each of these instances is a system of linear equations and can be solved in polynomial time. If at least one of the instances does not have a solution, then the instance (1) does not hold. Assume that all of them are satisfiable, then consider the relation $\Delta(x_1, \dots, x_n) = \exists y_1 \dots \exists y_m \Phi$. Since Γ is preserved by $x + y + z$, Δ is also preserved by $x + y + z$. Applying this operation to the tuples $(0, 0, \dots, 0), (1, 0, \dots, 0), (0, 1, 0, \dots, 0), \dots, (0, 0, \dots, 0, 1) \in \Delta$ coordinatewise we derive that $\Delta = \{0, 1\}^n$, that is Δ contains all tuples and (1) holds. Thus, we build a reduction of $\text{QCSP}^{\text{II}_2}(\Gamma)$ to $\text{CSP}(\Gamma)$ and proved that $\text{QCSP}^{\text{II}_2}(\Gamma^*)$ is solvable in polynomial time.

This idea can be generalized as follows. We say that a set of operations F (or an algebra $(A; F)$) has the *polynomially generated powers (PGP)* property if there exists a polynomial $p(n)$ such that A^n can be generated from $p(n)$ tuples using operations of F . Another behaviour that might arise is that there is an exponential function f so that the smallest generating sets for A^n require size at least $f(n)$. In this case we say that it has the *exponentially generated powers (EGP)* property.

Theorem 1. [12] *Every finite algebra $(A; F)$ either has the PGP property, or has the EGP property.*

Moreover, it was shown that a polynomial size generating set in the PGP case can be chosen very simple and efficiently computable. We say that a tuple (a_1, \dots, a_n) has a *switch* in the i -th position if $a_{i-1} \neq a_i$. For example, the tuple $(1, 1, 0, 2, 0, 0, 0)$ has 3 switches. We say that an algebra $(A; F)$ is *k-switchable* if A^n is generated from all tuples with at most k switches. We say that an algebra $(A; F)$ is *switchable* if it is k -switchable for some k . Since we have only polynomially many tuples with at most k switches, switchability implies the PGP property. It turned out that the backward implication also holds.

Theorem 2. [12] *A finite algebra $(A; F)$ has the PGP property if and only if it is switchable.*

Thus, as a generating set of polynomial size we can take the set of all tuples with at most k switches, which gives a polynomial reduction of $\text{QCSP}^{\text{II}_2}(\Gamma)$ to $\text{CSP}(\Gamma^*)$ if $\text{Pol}(\Gamma)$ has the PGP property.

It turns out that a similar reduction can be made for the general $\text{QCSP}(\Gamma)$.

Theorem 3. [8, 5] *Suppose Γ contains all constants relations and $\text{Pol}(\Gamma)$ has the PGP property. Then $\text{QCSP}(\Gamma)$ is polynomially reducible to $\text{CSP}(\Gamma)$.*

The proof of this theorem was rather complicated. First, in [8] Hubie Chen proved that the existence of a switchable algebra (with a slightly different type of switchability) allows to do this reduction for any constraint language with constants. Combining the equivalence of the two types of switchability [4] with Theorem 2 we derive Theorem 3. Another disadvantage of the Hubie Chen reduction is that it works only for constraint languages with constants.

Nevertheless, Theorem 3 and the classification of the complexity of the CSP [2, 3, 13, 11] give the following classification of the complexity for the PGP case with constants.

Corollary 4. [8, 5] *Suppose Γ contains all constants relations and $\text{Pol}(\Gamma)$ has the PGP property. If Γ admits a WNU polymorphism, then $\text{QCSP}(\Gamma)$ is solvable in polynomial time; $\text{QCSP}(\Gamma)$ is NP-complete otherwise.*

For many years there was a Conjecture suggested by Hubie Chen [9, 6] claiming that $\text{QCSP}(\Gamma)$ is PSpace-hard whenever $\text{Pol}(\Gamma)$ has the EGP property. If this conjecture was true, then, by Theorem 1, it would complete the classification of the complexity at least for constraint languages with constants. Recently, this conjecture was disproved [14] but the PGP case remains a very important case for the study of the complexity of the Quantified CSP.

2 Main Results

In this paper we present a simpler proof of Theorem 3. Moreover, our reduction works not only for constraint languages with constants but for any constraint languages Γ such that $\text{Pol}(\Gamma)$ has the PGP property. As a result we obtain a stronger claim and a complete classification of the complexity of the Quantified CSP for any constraint language Γ whose polymorphisms have the PGP property.

Theorem 5. *Suppose $\text{Pol}(\Gamma)$ has the PGP property. Then $\text{QCSP}(\Gamma)$ is polynomially reducible to $\text{CSP}(\Gamma^*)$.*

By $\text{QCSP}_{n(\forall)\leq k}^{\Pi_2}(\Gamma)$ we denote the modification of $\text{QCSP}(\Gamma)$ where we allow only Π_2 -sentences with at most k universal quantifiers, that is, $\forall x_1 \dots \forall x_n \exists y_1 \dots \exists y_m \Phi$ for $n \leq k$.

Theorem 6. *Suppose $\text{Pol}(\Gamma)$ has the PGP property. Then $\text{QCSP}(\Gamma)$ is polynomially equivalent to $\text{QCSP}_{n(\forall)\leq |A|}^{\Pi_2}(\Gamma)$.*

Let M be the matrix whose rows are all the tuples of A^k listed in the lexicographic order. Thus, M is a matrix with k columns and $|A|^k$ rows. Let $\gamma_1^k, \dots, \gamma_k^k \in A^{|A|^k}$ be the k columns of M . For a relation R of arity s on a domain A by R^k we denote the relation on the domain A^k defined by

$$((a_1^1, \dots, a_1^k), (a_2^1, \dots, a_2^k), \dots, (a_s^1, \dots, a_s^k)) \in R^k \Leftrightarrow \forall i: (a_1^i, \dots, a_s^i) \in R.$$

Put $\Gamma^k = \{R^k \mid R \in \Gamma\}$. Then we have the following reduction to the CSP over the domain $A^{|A|^{|A|}}$.

Theorem 7. *Suppose $\text{Pol}(\Gamma)$ has the PGP property. Then $\text{QCSP}(\Gamma)$ is polynomially equivalent to $\text{CSP}(\Gamma^{|A|^{|A|}} \cup \{\gamma_1^{|A|}, \dots, \gamma_{|A|}^{|A|}\})$.*

This implies the following complexity classification.

Theorem 8. *Suppose $\text{Pol}(\Gamma)$ has the PGP property. Then $\text{QCSP}(\Gamma)$ is solvable in polynomial time if $\Gamma^{|A|^{|A|}} \cup \{\gamma_1^{|A|}, \dots, \gamma_{|A|}^{|A|}\}$ admits a WNU polymorphism; $\text{QCSP}(\Gamma)$ is NP-complete otherwise.*

The paper is organized as follows. In Section 3 we formulate the main reduction and prove the main results of the paper using this reduction. In Section 4 we prove four transformations of formulas that allow to remove universal quantifiers, reduce the number of universal quantifiers, and transform it into a Π_2 -form. In Section 5 we prove the main reduction of the paper.

3 Main Reduction

We will need the following transformation of a quantified sentence. Suppose $0 = n_0 < n_1 < n_2 < \dots < n_k < n_{k+1} = n + 1$ and $\Psi = \exists y_1 \forall x_1 \exists y_2 \forall x_2 \dots \exists y_n \forall x_n \Phi$. Define a mapping $\omega_{n_1, \dots, n_k}: \{x_1, \dots, x_n, y_1, \dots, y_n\} \rightarrow \{x_1, \dots, x_n, y_1, \dots, y_n, z_0, z_1, \dots, z_k\}$ by

$$\omega_{n_1, \dots, n_k}(u) = \begin{cases} y_i, & \text{if } u = y_i \text{ for some } i \\ x_{n_j}, & \text{if } u = x_{n_j} \text{ for some } j \\ z_j, & \text{if } u = x_i \text{ and } n_j < i < n_{j+1} \end{cases}$$

By $\omega_{n_1, \dots, n_k}(\Psi)$ we denote the sentence such that

1. every variable u is replaced by the variable $\omega_{n_1, \dots, n_k}(u)$;
2. the quantification $\forall x_i$ is removed whenever $\omega_{n_1, \dots, n_k}(x_i) \neq x_i$;
3. the outer quantification $\forall z_0 \forall z_1 \dots \forall z_k$ is added.

The key result of this paper is as follows.

Theorem 9. *Suppose $\text{Pol}(\Gamma)$ is r -switchable, $\Psi = \exists y_1 \forall x_1 \exists y_2 \forall x_2 \dots \exists y_n \forall x_n \Phi$, where Φ is a conjunctive formula over Γ . Then Ψ holds if and only if for every $1 \leq n_1 < n_2 < \dots < n_k \leq n$, $k \leq r$, the sentence $\omega_{n_1, \dots, n_k}(\Psi)$ holds.*

Thus, we have a reduction of the Quantified CSP to polynomially many instances with at most $2r + 1$ universal quantifiers. To derive the main results we will need the following technical lemmas proved in Section 4.

Lemma 10. *$\text{QCSP}_{n(\forall) \leq k}(\Gamma)$ is polynomially reducible to $\text{CSP}(\Gamma^*)$.*

Lemma 11. *$\text{QCSP}_{n(\forall) \leq k}(\Gamma)$ is polynomially reducible to $\text{QCSP}_{n(\forall) \leq |A|^k}^{\Pi_2}(\Gamma)$.*

Lemma 12. *$\text{QCSP}_{n(\forall) \leq k}^{\Pi_2}(\Gamma)$ is polynomially reducible to $\text{QCSP}_{n(\forall) \leq |A|}^{\Pi_2}(\Gamma)$.*

Now we can prove the reduction to the CSP with constants.

Theorem 5. *Suppose $\text{Pol}(\Gamma)$ has the PGP property. Then $\text{QCSP}(\Gamma)$ is polynomially reducible to $\text{CSP}(\Gamma^*)$.*

Proof. By Theorem 2, $\text{Pol}(\Gamma)$ is r -switchable for some r . By Theorem 9 an instance I of $\text{QCSP}(\Gamma)$ is equivalent to a conjunction of polynomially many instances I_1, \dots, I_t of $\text{QCSP}(\Gamma)$ such that each instance has at most $2r + 1$ universally quantified variables. By Lemma 10, for each I_j we can build an equivalent instance I'_j of $\text{CSP}(\Gamma^*)$. Then $I'_1 \wedge \dots \wedge I'_t$ is an instance of $\text{CSP}(\Gamma^*)$ equivalent to the original instance I . \square

Theorem 6. *Suppose $\text{Pol}(\Gamma)$ has the PGP property. Then $\text{QCSP}(\Gamma)$ is polynomially equivalent to $\text{QCSP}_{n(\forall) \leq |A|}^{\Pi_2}(\Gamma)$.*

Proof. By Theorem 2, $\text{Pol}(\Gamma)$ is r -switchable for some r . Again, by Theorem 9 an instance I of $\text{QCSP}(\Gamma)$ is equivalent to a conjunction of polynomially many instances I_1, \dots, I_t of $\text{QCSP}(\Gamma)$ such that each instance has at most $2r + 1$ universally quantified variables. By Lemma 11 for each I_j we can build an equivalent instance I'_j of $\text{QCSP}_{n(\forall) \leq |A|^{2r+1}}^{\Pi_2}(\Gamma)$. We additionally require that existential variables in I'_1, \dots, I'_t to be different and universal variables to be the same. Then the conjunction $I'_1 \wedge \dots \wedge I'_t$ can be easily transformed into an equivalent instance J of $\text{QCSP}_{n(\forall) \leq |A|^{2k+1}}^{\Pi_2}(\Gamma)$ by moving all universal quantifiers left. It remains to apply Lemma 12 to obtain an equivalent instance of $\text{QCSP}_{n(\forall) \leq |A|}^{\Pi_2}(\Gamma)$. \square

Recall that $\gamma_1^k, \dots, \gamma_k^k \subseteq A^{|A|^k}$ are the k columns of the matrix whose rows are all the tuples of A^k listed in the lexicographic order. For $\alpha \in A^{|A|^k}$ by $\alpha^{c_1, \dots, c_{|A|}}$ we denote the i -th element of α where the i -th row of the matrix is $(c_1, \dots, c_{|A|})$.

Theorem 7. *Suppose $\text{Pol}(\Gamma)$ has the PGP property. Then $\text{QCSP}(\Gamma)$ is polynomially equivalent to $\text{CSP}(\Gamma^{|A|^{|A|}} \cup \{\gamma_1^{|A|}, \dots, \gamma_{|A|}^{|A|}\})$.*

Proof. \Rightarrow . Suppose \mathcal{I} is an instance of $\text{QCSP}(\Gamma)$. By Theorem 6, we can build an equivalent instance of $\text{QCSP}_{n(\forall)\leq|A|}^{\Pi_2}(\Gamma)$, that is an instance $\Psi = \forall x_1 \dots \forall x_{|A|} \exists y_1 \dots \exists y_n \Phi$, where Φ is a conjunction of relations from Γ . Let us build an equivalent instance \mathcal{J} of $\text{CSP}(\Gamma^{|A|^{|A|}} \cup \{\gamma_1^{|A|}, \dots, \gamma_{|A|}^{|A|}\})$. To obtain \mathcal{J} from Φ we

1. replace each relation $R \in \Gamma$ by $R^{|A|^{|A|}} \in \Gamma^{|A|^{|A|}}$;
2. add constraint $\gamma_i(x_i)$ for each $i \in \{1, 2, \dots, |A|\}$.

Let us show that Ψ and \mathcal{J} are equivalent. If Ψ holds, then Φ has a solution $(y_1, \dots, y_n) = (\beta_1^{c_1, \dots, c_{|A|}}, \dots, \beta_n^{c_1, \dots, c_{|A|}})$ for any evaluation of $(x_1, \dots, x_{|A|}) = (c_1, \dots, c_{|A|})$. If we interpret each β_i as a tuple from $A^{|A|^{|A|}}$ then $(x_1, \dots, x_{|A|}, y_1, \dots, y_n) = (\gamma_1, \dots, \gamma_{|A|}, \beta_1, \dots, \beta_n)$ is a solution of \mathcal{J} . Similarly, if \mathcal{J} has a solution $(x_1, \dots, x_{|A|}, y_1, \dots, y_n) = (\gamma_1, \dots, \gamma_{|A|}, \beta_1, \dots, \beta_n)$ then $(y_1, \dots, y_n) = (\beta_1^{c_1, \dots, c_{|A|}}, \dots, \beta_n^{c_1, \dots, c_{|A|}})$ is a solution of Φ corresponding to $(x_1, \dots, x_{|A|}) = (c_1, \dots, c_{|A|})$.

\Leftarrow . Suppose \mathcal{J} is an instance of $\text{CSP}(\Gamma^{|A|^{|A|}} \cup \{\gamma_1^{|A|}, \dots, \gamma_{|A|}^{|A|}\})$. To build the quantifier-free part Φ from \mathcal{J} we

1. introduce $|A|$ new variables $x_1, \dots, x_{|A|}$;
2. if \mathcal{J} contains $\gamma_i(z)$ and $\gamma_j(z)$ for some variables z and $i \neq j$, then we reduce \mathcal{J} to any no-instance of $\text{QCSP}(\Gamma)$;
3. replace each relation $R^{|A|^{|A|}} \in \Gamma^{|A|^{|A|}}$ by $R \in \Gamma$;
4. remove each constraint $\gamma_i(z)$ and replace the variable z by x_i .

It is not hard to check that \mathcal{J} is equivalent to $\Psi = \forall x_1 \dots \forall x_{|A|} \exists y_1 \dots \exists y_n \Phi$. \square

As a corollary we obtain the following result.

Theorem 8. *Suppose $\text{Pol}(\Gamma)$ has the PGP property. Then $\text{QCSP}(\Gamma)$ is solvable in polynomial time if $\Gamma^{|A|^{|A|}} \cup \{\gamma_1^{|A|}, \dots, \gamma_{|A|}^{|A|}\}$ admits a WNU polymorphism; $\text{QCSP}(\Gamma)$ is NP-complete otherwise.*

4 Transformations

In this section we define four transformations of instances, which are probably well-known.

4.1 Universal quantifiers removal.

Suppose $A = \{a_1, \dots, a_s\}$, $\Psi = \exists y_1 \dots \exists y_j \forall x \Phi$ is a QCSP sentence (Φ may contain quantifiers). Then Ψ is equivalent to

$$\exists y_1 \dots \exists y_j \exists x^1 \dots \exists x^s \bigwedge_{i=1}^s (\Phi_i \wedge x^i = a_i),$$

where Φ_i is obtained from Φ by replacement of x by x^i . If the number of universal quantifiers is bounded, we can remove all the universally quantified variables in polynomial time, and the result can be written in the prenex form, which is a CSP sentence. Thus, we have the following lemma.

Lemma 10. *$\text{QCSP}_{n(\forall)\leq k}(\Gamma)$ is polynomially reducible to $\text{CSP}(\Gamma^*)$.*

4.2 Moving universal quantifiers left.

Suppose $|A| = s$, $\Psi = \exists y_1 \dots \exists y_j \forall x \Phi$ is a QCSP sentence (Φ may contain quantifiers). Then Ψ is equivalent to

$$\forall x^1 \dots \forall x^s \exists y_1 \dots \exists y_j \bigwedge_{i=1}^s (\Phi_i),$$

where Φ_i is obtained from Φ by replacement of x by x^i . In this way we can move all the universal quantifiers left starting with the right one, and the result can be written in the prenex form to get a QCSP sentence. Thus, we have the following lemma.

Lemma 11. $\text{QCSP}_{n(\forall) \leq k}(\Gamma)$ is polynomially reducible to $\text{QCSP}_{n(\forall) \leq |A|^k}^{\Pi_2}(\Gamma)$.

4.3 Reducing the number of quantifiers

Suppose $A = \{a_1, \dots, a_s\}$. Suppose we have an instance $\Psi = \forall x_1 \dots \forall x_k \exists y_1 \dots \exists y_n \Phi$ of $\text{QCSP}_{n(\forall) \leq k}^{\Pi_2}$. We can make it easier by reducing the number of universal quantifiers using the following trick. Since k is a constant, there are only $|A|^k$ possible evaluations of (x_1, \dots, x_k) we need to check. We do not want to use constants, that is why we do the following.

For each mapping φ from $[k] := \{1, 2, \dots, k\}$ to $[s] := \{1, 2, \dots, s\}$ we define the sentence Ψ_φ where each variable x_i is replaced by $z_{\varphi(i)}$, and the universal quantification is $\forall z_1 \dots \forall z_s$ instead of $\forall x_1 \dots \forall x_k$. Then Ψ is equivalent to $\bigwedge_{\varphi: [k] \rightarrow [s]} \Psi_\varphi$, which can be transformed to a Π_2 -sentence whose universally quantified variables are $\forall z_1 \dots \forall z_s$. Thus, we proved the following lemma.

Lemma 12. $\text{QCSP}_{n(\forall) \leq k}^{\Pi_2}(\Gamma)$ is polynomially reducible to $\text{QCSP}_{n(\forall) \leq |A|^k}^{\Pi_2}(\Gamma)$.

4.4 Transformation to a Π_2 -formula.

Suppose $\Psi = \exists y_1 \forall x_1 \exists y_2 \forall x_2 \dots \exists y_n \forall x_n \Phi$. We define a transformation ζ that transforms Ψ into an equivalent Π_2 -formula. Note that $\zeta(\Psi)$ is of exponential size on the size of Ψ .

For each tuple $(a_1, \dots, a_n) \in A^n$ the formula Φ_{a_1, \dots, a_n} is obtained from Φ by

1. replacement of each variable y_i by $y_i^{a_1, \dots, a_{i-1}}$
2. replacement of each variable x_i by $x_i^{a_1, \dots, a_i}$

By $\zeta(\Psi)$ we denote the sentence such that

1. its quantifier-free part is $\bigwedge_{(a_1, \dots, a_n) \in A^n} \Phi_{a_1, \dots, a_n}$
2. first we universally quantify the x -variables, then we existentially quantify the y -variables.

Let us show that Ψ and $\zeta(\Psi)$ are equivalent.

Lemma 13. $\Psi \rightarrow \zeta(\Psi)$.

Proof. Suppose $(f_1(), x_1, f_2(x_1), x_2, f_3(x_1, x_2), x_3, \dots)$ is a solution of Ψ . Let us show that a solution of $\zeta(\Psi)$ can be defined by

$$y_i^{a_1, \dots, a_{i-1}} = f_i(x_1^{a_1}, x_2^{a_1, a_2}, x_3^{a_1, a_2, a_3}, \dots, x_{i-1}^{a_1, \dots, a_{i-1}}).$$

In fact, each conjunctive formula Φ_{a_1, \dots, a_n} holds as it is just Φ with all the variables renamed. \square

Lemma 14. $\zeta(\Psi) \rightarrow \Psi$.

Proof. Set $x_i^{a_1, \dots, a_i} = a_i$ for every i and $a_1, \dots, a_i \in A$. Since $\zeta(\Psi)$ holds, for each assignment of the x -variables there exists a proper assignment of the y -variables. We define a solution to Ψ by $(f_1(), x_1, f_2(x_1), x_2, f_3(x_1, x_2), x_3, \dots)$, where $f_i(a_1, \dots, a_{i-1}) = y_i^{a_1, \dots, a_{i-1}}$. To check that the conjunctive formula Φ holds for $(x_1, \dots, x_n) = (a_1, \dots, a_n)$ we consider Φ_{a_1, \dots, a_n} . \square

5 Proof of Theorem 9

Theorem 9. *Suppose $\text{Pol}(\Gamma)$ is r -switchable, $\Psi = \exists y_1 \forall x_1 \exists y_2 \forall x_2 \dots \exists y_n \forall x_n \Phi$, where Φ is a conjunctive formula over Γ . Then Ψ holds if and only if for every $1 \leq n_1 < n_2 < \dots < n_k \leq n$, $k \leq r$, the sentence $\omega_{n_1, \dots, n_k}(\Psi)$ holds.*

Proof. To obtain $\omega_{n_1, \dots, n_k}(\Psi)$ from Ψ we just identify universally quantified variables and move universal quantifiers left, hence $\omega_{n_1, \dots, n_k}(\Psi)$ holds whenever Ψ holds.

Assume that $\omega_{n_1, \dots, n_k}(\Psi)$ holds for every $1 \leq n_1 < n_2 < \dots < n_k \leq n$. Let us show that Ψ holds. By Lemmas 13 and 14 Ψ and $\zeta(\Psi)$ are equivalent, hence it is sufficient to show that $\zeta(\Psi)$ holds. Let $R \subseteq A^t$ be the relation defined by $\zeta(\Psi)$ where all universal quantifiers are removed. We assume that the order of coordinates in R corresponds to the order of variables x_1, x_2, \dots, x_n .

We need to show that R is full. Since $\text{Pol}(R)$ is r -switchable, it is sufficient to show that all tuples with at most r switches are in R . Consider a tuple α whose switches are in the positions corresponding to the variables $x_{n_1}, x_{n_2}, \dots, x_{n_k}$, where $k \leq r$. Consider a strategy for the Existential Player in $\omega_{n_1, \dots, n_k}(\Psi)$, that is $y_i = f_i(z_0, z_1, \dots, z_k, x_{n_1}, \dots, x_{n_{\ell(i)}})$, where $\ell(i)$ is the maximal number such that $n_{\ell(i)} < i$. Then an assignment to the y -variables (in the definition of R) corresponding to α can be defined by $y_i^{a_1, \dots, a_{i-1}} = f_i(b_0, b_1, \dots, b_k, c_1, \dots, c_{\ell(i)})$, where

- b_0 is the value of the coordinates in α corresponding to the x -variables before x_{n_1} ;
- each b_j is the value of the coordinates in α corresponding to the x -variables between x_{n_j} and $x_{n_{j+1}}$;
- b_k is the value of the coordinates in α corresponding to the x -variables after x_{n_k} ;
- each c_j is the value of the coordinates in α corresponding to the variable $x_{n_j}^{a_1, \dots, a_{n_j}}$.

Since the functions f_1, \dots, f_n give a solution to $\omega_{n_1, \dots, n_k}(\Psi)$, we defined a proper assignment to $\xi(\Psi)$. Hence $\alpha \in R$ and R is full. \square

References

- [1] Ferdinand Börner, Andrei A. Bulatov, Hubie Chen, Peter Jeavons, and Andrei A. Krokhin. The complexity of constraint satisfaction games and QCSP. *Inf. Comput.*, 207(9):923–944, 2009.
- [2] Andrei A. Bulatov. A dichotomy theorem for nonuniform CSPs. In *2017 IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 319–330, 2017.
- [3] Andrei A. Bulatov. A dichotomy theorem for nonuniform CSPs. *CoRR*, abs/1703.03021, 2017.

- [4] C. Carvalho, F. Madelaine, and B. Martin. From complexity to algebra and back: Digraph classes, collapsibility, and the PGP. In *30th Annual ACM/IEEE Symposium on Logic in Computer Science*, pages 462–474, July 2015.
- [5] Catarina Carvalho, Florent R. Madelaine, and Barnaby Martin. From complexity to algebra and back: digraph classes, collapsibility and the PGP. In *30th Annual IEEE Symposium on Logic in Computer Science (LICS)*, 2015.
- [6] Catarina Carvalho, Barnaby Martin, and Dmitriy Zhuk. The complexity of quantified constraints using the algebraic formulation. In *42nd International Symposium on Mathematical Foundations of Computer Science, MFCS 2017, August 21-25, 2017 - Aalborg, Denmark*, pages 27:1–27:14, 2017.
- [7] Hubie Chen. The complexity of quantified constraint satisfaction: Collapsibility, sink algebras, and the three-element case. *SIAM J. Comput.*, 37(5):1674–1701, 2008.
- [8] Hubie Chen. Quantified constraint satisfaction and the polynomially generated powers property. *Algebra universalis*, 65(3):213–241, 2011. An extended abstract appeared in ICALP B 2008.
- [9] Hubie Chen. Meditations on quantified constraint satisfaction. In *Logic and Program Semantics - Essays Dedicated to Dexter Kozen on the Occasion of His 60th Birthday*, pages 35–49, 2012.
- [10] Barnaby Martin. Quantified Constraints in Twenty Seventeen. In Andrei Krokhin and Stanislav Zivny, editors, *The Constraint Satisfaction Problem: Complexity and Approximability*, volume 7 of *Dagstuhl Follow-Ups*, pages 327–346. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, Dagstuhl, Germany, 2017.
- [11] D. Zhuk. A proof of CSP dichotomy conjecture. In *2017 IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 331–342, Oct 2017.
- [12] Dmitriy Zhuk. The size of generating sets of powers. *Journal of Combinatorial Theory, Series A*, 167:91–103, 2019.
- [13] Dmitriy Zhuk. A proof of the CSP dichotomy conjecture. *Journal of the ACM (JACM)*, 67(5):1–78, 2020.
- [14] Dmitriy Zhuk and Barnaby Martin. QCSP monsters and the demise of the Chen conjecture. In *Proceedings of the 52nd Annual ACM SIGACT Symposium on Theory of Computing*, pages 91–104, 2020.