SECOND OBSTRUCTION TO PSEUDOISOTOPY I

KIYOSHI IGUSA

ABSTRACT. A pseudoisotopy of M is a diffeomorphism of $M \times I$ which is the identity on $M \times 0$. We give an explicit construction of pseudoisotopies of 4-manifolds which realize certain elements of the "second obstruction to pseudoisotopy". The next paper will give precise statements of our results for 5-manifolds which had been announced long ago.

CONTENTS

Introduction	1
1. Definition of second obstruction to pseudoisotopy	3
1.1. Basic setup	3
1.2. The graphic	5
1.3. Second obstruction	5
1.4. Computation of second obstruction	7
1.5. Dependence on markings	10
1.6. Involution	11
2. Constructing pseudoisotopies using two disks	12
3. Realizing the framing obstruction	15
4. Construction of the two disks	16
Acknowledgements	17
References	17

INTRODUCTION

Recall that, for any smooth compact manifold M, a *pseudoisotopy* of M is a diffeomorphism of $M \times I$ which is the identity on $M \times 0 \cup \partial M \times I$. The restriction of this diffeomorphism to $M \times 1$ gives a diffeomorphism of M which is the identity on ∂M . This diffeomorphism is said to be *pseudoisotopic* to the identity diffeomorphism of Mrel ∂M . Let $\mathcal{C}(M)$ be the group of pseudoisotopies of M. Thus

 $\mathcal{C}(M) = Diff(M \times I \, rel \, M \times 0 \cup \partial M \times I).$

²⁰²⁰ Mathematics Subject Classification. 19J10: 58K60.

Key words and phrases. Morse theory, 4-manifolds, diffeomorphisms.

Supported by the Simons Foundation.

We give $\mathcal{C}(M)$ a suitable topology so that, up to canonical homotopy, a path in $\mathcal{C}(M)$ is given by a diffeomorphism of $M \times I \times I$ which preserves the last coordinate. (Section 1.) Since $\mathcal{C}(M)$ is a topological group, $\pi_0 \mathcal{C}(M)$ will be a group.

For M simply connected with dim $M \ge 5$, Jean Cerf [1] showed that $\mathcal{C}(M)$ is connected. For dim $M \ge 6$, Hatcher and Wagoner [2] showed that there is an exact sequence

(0.1)
$$Wh_1^+(\pi_1 M; \mathbb{Z}_2 \oplus \pi_2 M) \to \pi_0 \mathcal{C}(M) \to Wh_2(\pi) \to 0.$$

When dim $M \leq 5$, we apply a "suspension functor" $\sigma : \mathcal{C}(M) \to \mathcal{C}(M \times I)$ several times to get to the *stable pseudoisotopy space* $\mathcal{P}(M) = \operatorname{colim} \mathcal{C}(M \times I^N)$ [6] which should be used instead of $\mathcal{C}(M)$ in the above exact sequence.

In [5], [4] and [3] the sequence (0.1) was extended to an exact sequence

(0.2)
$$K_3\mathbb{Z}[\pi_1 M] \xrightarrow{\chi} Wh_1^+(\pi_1 M; \mathbb{Z}_2 \oplus \pi_2 M) \xrightarrow{\theta} \pi_0\mathcal{P}(M) \to Wh_2(\pi) \to 0$$

The question arises: Which elements of $\pi_0 \mathcal{P}(M)$ are realized when dim M < 6, i.e., which elements of $\pi_0 \mathcal{P}(M)$ lie in the image of the stabilization map $\pi_0 \mathcal{C}(M) \to \pi_0 \mathcal{P}(M)$? This question was answered in dimension 5 by the author in a preliminary version of [7] and partially answered in the topological category in dimension 4 by Kwasik in [8] using the techniques of [2] and one idea from [7]. In both [7] and [8] we need an assumption which trivializes the image of χ in (0.2).

This series of papers discusses this question for dim M = 4, 5. In the current paper we show that, for certain 4-dimensional manifolds M, some elements of the second obstruction group

$$Wh_1^+(\pi_1 M; \mathbb{Z}_2 \oplus \pi_2 M) = Wh_1^+(\pi_1 M; \mathbb{Z}_2) \oplus Wh_1^+(\pi_1 M; \pi_2 M)$$

can be realized by pseudoisotopies of M. Given a group G and G-module A, $Wh_1^+(G; A)$ is a quotient of $A[G]/A[e] = A \otimes \mathbb{Z}[G]/\mathbb{Z}[e]$ (Definition 1.7). In particular, $Wh_1^+(\pi_1 M; \pi_2 M)$ is generated by elements $\alpha[\sigma]$ where $\alpha \in \pi_2 M$ and σ is a nontrivial element of $\pi_1 M$.

Theorem A (Theorem 2.3). Let M be the connected sum of $\mathbb{C}P^2$ with a nonsimply connected 4-manifold X. Then the elements of $Wh_1^+(\pi_1M;\pi_2M)$ of the form $\alpha[\sigma]$ where α is represented by the 2-sphere $\mathbb{C}P^1 \subset \mathbb{C}P^2$ lift to $\pi_0\mathcal{C}(M)$ and map to nontrivial elements of the kernel of $\pi_0\mathcal{P}(M) \twoheadrightarrow Wh_2(\pi)$.

The proof relies on the fact that $\mathbb{C}P^1 \subset \mathbb{C}P^2$ is an embedded 2-sphere which gives an element of $\pi_2 M$ which is not hit by the Postnikov invariant $k_1 \in H^3(\pi_1 M; \pi_2 M)$. We know by the formula for χ given in [5] that this element of $\pi_2 M$, together with any nontrivial element of $\pi_1 M$, gives an element of the second obstruction group $Wh_1^+(\pi_1 M; \pi_2 M)$ whose image in $\pi_0 \mathcal{P}(M)$ is nontrivial. The assertion in this paper is that this second obstruction element of $\pi_0 \mathcal{P}(M)$ lifts to $\pi_0 \mathcal{C}(M)$ where dim M = 4.

The proof relies on the space $\mathcal{D}(M)$ of "lens-space models" for pseudoisotopies of M(Definition 1.4). This is the space considered by Hatcher and Wagoner to determine the second obstruction to pseudoisotopy. We also consider the space $\mathcal{D}_0(M)$ of "marked lens-space models" for M since $\pi_0 \mathcal{D}_0(M)$ forms a group (Lemma 1.6). We recall that $Wh_1^+(\pi_1 M; \mathbb{Z}_2)$, is $\mathbb{Z}_2[\pi_1 M]/\mathbb{Z}_2[e]$ modulo the conjugation action of $\pi_1 M$. For example, when $\pi_1 M$ is finite,

$$Wh_1^+(\pi_1M;\mathbb{Z}_2) = \bigoplus_{c-1} \mathbb{Z}_2$$

is the direct sum of c-1 copies of \mathbb{Z}_2 where c is the number of conjugacy classes of elements of $\pi_1 M$. In dimension 4, we do not know whether all elements of $Wh_1^+(\pi_1 M; \mathbb{Z}_2)$ can be realized by pseudoisotopies of M and any realization might have infinite order. We show in this paper that there are homomorphisms $\tilde{\theta}_0 : \pi_0 \mathcal{D}_0(M) \to \pi_0 \mathcal{C}(M)$ and $\overline{\lambda}_0 : \pi_0 \mathcal{D}_0(M) \to Wh_1^+(\pi_1 M; \mathbb{Z}_2 \oplus \pi_2 M)$ making the following diagram commute. Furthermore, when $M = X \# \mathbb{C}P^2$, $\overline{\lambda}_0$ maps onto $Wh_1^+(\pi_1 M; \mathbb{Z}_2)$.

$$\begin{array}{c|c} \pi_0 \mathcal{D}_0(M) & \xrightarrow{\widetilde{\theta}_0} & \pi_0 \mathcal{C}(M) \\ & & & \downarrow \\ & & & \downarrow \\ Wh_1^+(\pi_1 M; \mathbb{Z}_2 \oplus \pi_2 M) \xrightarrow{\theta} & \pi_0 \mathcal{P}(M) \end{array}$$

This proves the second main theorem:

Theorem B (Theorem 3.1). Let M be the connected sum of $\mathbb{C}P^2$ with a nonsimply connected 4-manifold X. Then every elements in the image of $Wh_1^+(\pi_1 M; \mathbb{Z}_2) \to \pi_0 \mathcal{P}(M)$ lift to $\pi_0 \mathcal{C}(M)$.

The same proof works in a slightly more general case, when M has an embedded 2-sphere with odd self-intersection number. (Theorem 3.2.) A topological version of this result for smooth 4-manifolds was obtained by Kwasik [8, Proposition 4.1].

1. Definition of second obstruction to pseudoisotopy

We will review the basic theory, due to Cerf [1], relating diffeomorphisms of $M \times I$ and Morse theory. Then we focus on the second obstruction, due to Hatcher and Wagoner [2]. We assume in this section that M is a compact connected smooth manifold of dimension $n \ge 4$.

1.1. **Basic setup.** Let $\mathcal{F}(M)$ be the space of all smooth functions $f: M \times I \to I$ so that f is equal to projection to I in some neighborhood of

$$\partial (M \times I) = \partial M \times I \cup M \times 0 \cup M \times 1.$$

Then $\mathcal{F}(M)$ is clearly contractible: $\mathcal{F}(M) \simeq *$. Let $\mathcal{E}(M)$ be the space of all $f \in \mathcal{F}(M)$ with no singularities. Then, Cerf observed that

$$\mathcal{E}(M) \simeq \mathcal{C}(M)$$

where the homotopy equivalence $\mathcal{C}(M) \to \mathcal{E}(M)$ is given by sending $g: M \times I \to M \times I$ to $\pi_I \circ g: M \times I \to I$ where $\pi_I: M \times I \to I$ is the projection map. Given $f \in \mathcal{E}(M)$, the corresponding element of $\mathcal{C}(M)$ is constructed uniquely up to contractible choice by integrating the gradient of f. As a consequence

$$\pi_0 \mathcal{C}(M) \cong \pi_1(\mathcal{F}(M), \mathcal{E}(M)).$$

For clarity we repeat: This bijection sends the isotopy class of $g \in \mathcal{C}(M)$ to the homotopy class of the 1-parameter family f_t if and only if $f_1 = \pi_I \circ g$ (and $f_0 = \pi_I$).

Proposition 1.1. Let $(f_t^a), (f_t^b)$ be elements of $\pi_1(\mathcal{F}(M), \mathcal{E}(M))$ corresponding to $g^a, g^b \in \pi_0 \mathcal{C}(M)$. Thus $f_1^a = \pi_I \circ g^a, f_1^b = \pi_I \circ g^b$. Then the product in $\pi_0 \mathcal{C}(M)$, given by composition: $g^a \circ g^b$ is f_t^{ab} given by putting f_t^b to the left of f_t^a :

$$f_t^{ab} = \begin{cases} f_{2t}^b & \text{if } t \le \frac{1}{2} \\ f_{2t-1}^a \circ g^b & \text{if } t > \frac{1}{2}. \end{cases}$$

Proof. First note that the family of function f_t^{ab} lies in $\pi_1(\mathcal{F}(M), \mathcal{E}(M))$ since, at t = 0, $f_0^b = \pi_I$ and, at $t = \frac{1}{2}$, $f_1^b = \pi_I \circ g^b = f_0^a \circ g^b$. It is also easy to see that f^{ab} corresponds to $g^a g^b$ since $f_1^{ab} = f_1^a \circ g^b = \pi_I \circ g^a \circ g^b$.

Corollary 1.2. The inverse of $(f_t) \in \pi_1(\mathcal{F}(M), \mathcal{E}(M))$ is the 1-parameter family $f'_t = f_{1-t} \circ g^{-1} : M \times I \to I$

where $g \in \mathcal{C}(M)$ is a diffeomorphism of $M \times I$ whose composition with projection π_I : $M \times I \to I$ is $f_1 \in \mathcal{E}(M)$ (so that $f'_0 \circ g = f_1 = \pi_I \circ g$, making $f'_0 = \pi_I$.)

Proof. Let $f_t^a = f_t'$ and $f_t^b = f_t$. By Proposition 1.1, the product $f^{ab} = f^a f^b$ at t = 1 is $f_1^{ab} = f_1^a \circ g^b = f_0^b \circ (g^b)^{-1} \circ g^b = f_0^b = \pi_I$.

This corresponds to the identity in $\pi_0 \mathcal{C}(M)$. So, $f^a = (f'_t)$ is the inverse of $f^b = (f_t)$. \Box

Here is an alternate version of composition in $\pi_0 \mathcal{C}(M)$ in terms of functions.

Proposition 1.3. The group structure (composition) on $\pi_0 \mathcal{C}(M)$ corresponds to stacking of functions $f_a, f_b : M \times I \to I$ in $\mathcal{E}(M)$ given by "putting f_b on top of f_a ":

$$(f_a \bullet f_b)(x,t) = \begin{cases} \frac{1}{2}f_a(x,2t) & \text{if } t \le \frac{1}{2}\\ \frac{1}{2} + \frac{1}{2}f_b(x,2t-1) & \text{if } t > \frac{1}{2} \end{cases}$$

Proof. Let $g_a, g_b \in \mathcal{C}(M)$ be given by $g_a(x,t) = (h_a(x,t), f_a(x,t))$ and $g_b = (h_b, f_b)$. Then g_a, g_b are isotopic to $g'_a, g'_b \in \mathcal{C}(M)$ given by "pushing g_a down and g_b up":

$$g'_a(x,t) = \begin{cases} (h_a(x,2t), \frac{1}{2}f_a(x,2t)) & \text{if } t \le \frac{1}{2} \\ (r(x),t) & \text{if } t > \frac{1}{2} \end{cases}$$

where $r \in Diff(M)$ is the restriction of g_a to $M \times 1$, i.e., $g_a(x, 1) = (r(x), 1)$ and

$$g'_b(x,t) = \begin{cases} (x,t) & \text{if } t \le \frac{1}{2} \\ (h_b(x,2t-1),\frac{1}{2} + \frac{1}{2}f_b(x,2t-1)) & \text{if } t > \frac{1}{2} \end{cases}$$

Then $g_a \circ g_b$ is isotopic to $g'_a \circ g'_b$ given by

$$(g'_a \circ g'_b)(x,t) = \begin{cases} (h_a(x,2t), \frac{1}{2}f_a(x,2t)) & \text{if } t \le \frac{1}{2} \\ (rh_b(x,2t-1), \frac{1}{2} + \frac{1}{2}f_b(x,2t-1)) & \text{if } t > \frac{1}{2} \end{cases}$$

Taking the second coordinate gives $f_a \bullet f_b$.

The group structure on $\pi_1(\mathcal{F}(M), \mathcal{E}(M))$ is also given by stacking functions.

1.2. The graphic. The fiberwise singular set $\Sigma(f)$ of f is the set

 $\Sigma(f) := \{ (x, s, t) \in M \times I^2 : (x, s) \in M \times I \text{ is a singularity of } f_t \}.$

Since each function f_t will generically have a finite singular set, and f_0 , f_1 are nonsingular, $\Sigma(f)$ will, generically, be a closed 1-dimensional manifold. If $\Sigma(f)$ is empty or if f is homotopic relative to $\partial(M \times I^2)$ to a family of function with empty singular set, then f represents the trivial element of $\pi_1(\mathcal{F}(M), \mathcal{E}(M))$. Thus, $\pi_1(\mathcal{F}(M), \mathcal{E}(M))$ is the obstruction to the elimination of the singular set $\Sigma(f)$ by deformation of f.

We visualize $\Sigma(f)$ via its graphic which is defined to be the set of all $(s,t) \in I^2$ so that s is a critical value of f_t . Two examples are given in Figure 1. The graphic on the right side of Figure 1 is called an "eye" or "lens".

Definition 1.4. A lens-shaped graphic or eye is, by definition, the graphic of a family of functions $f_t : M \times I \to I$ so that f_t is Morse with two critical points p_t, q_t with $f_t(p_t) < f_t(q_t)$ for $t_0 < t < t_1, f_{t_0}, f_{t_1}$ have only one critical point called a *birth-death point* and f_t is nonsingular for $t \notin [t_0, t_1]$. A family of functions (f_t) having a lens-shaped graphic will be called a *lens-shaped model* for M.



FIGURE 1. Two possible graphics of 1-parameter families of functions on $M \times I$. The one on the right is called an "eye" or "lens".

1.3. Second obstruction. There are two steps in the process of simplification of the singular set $\Sigma(f)$ of f. Hatcher and Wagoner [2] showed that, for dim $M \ge 6$, there is a surjective homomorphism

$$p: \pi_0 \mathcal{C}(M) \to Wh_2(\pi_1 M)$$

where, for any group π , $Wh_2(\pi)$ is a quotient of $K_2(\mathbb{Z}[\pi])$. They called this invariant the "first obstruction to pseudo-isotopy".

Hatcher and Wagoner also showed that the kernel of the map p is generated by "lensshaped models" (Definition 1.4) in the middle indices k, k + 1 where $k = \lfloor \frac{n}{2} \rfloor$ is the greatest integer $\leq \frac{n}{2}$ where $n = \dim M$. The "second obstruction to pseudo-isotopy" is an invariant defined on this set of 1-parameter families of function.

To define these invariants, we will add extra structure, called "markings", to these families of functions, define invariants on the marked lens-shaped models, then determine how these invariants change when we change the markings.

Definition 1.5. Let $\mathcal{D}(M)$ denote the space of lens-shaped models (f_t) for M with critical points p_t, q_t in indices k, k + 1 where $k = \lfloor \frac{n}{2} \rfloor$. Let $\mathcal{D}_0(M)$ denote the space of "marked" lens-shaped models for M. These are defined to be lens-shaped models (f_t) together with:

- (1) A homotopy class of paths γ from a fixed base point $* = (x_0, 0, 0) \in M \times I^2$ to the birth point b_0 of f_t (at $t = t_0$ in Fig. 1).
- (2) An orientation of the negative eigenspace of $D^2 f_t$ at the birth point b_0 .

The group $\pi_1(M) \times \mathbb{Z}_2$ acts on the set of marking, $\pi_1(M)$ acting on the paths by composition and \mathbb{Z}_2 acting by reversing the orientation. We denote by $\tilde{\theta} : \pi_0 \mathcal{D}(M) \to \pi_0 \mathcal{C}(M)$ the set map which sends a lens-shaped model (f_t) to the corresponding element of $\pi_0 \mathcal{C}(M) \cong$ $\pi_1(\mathcal{F}(M), \mathcal{E}(M))$. Composing with the surjective mapping $\pi_0 \mathcal{D}_0(M) \twoheadrightarrow \pi_0 \mathcal{D}(M)$ we get a mapping

$$\hat{\theta}_0: \pi_0 \mathcal{D}_0(M) \to \pi_0 \mathcal{C}(M)$$

which we will show is a homomorphism of groups (Lemma 1.6).

The group structure on $\pi_0 \mathcal{D}_0(M)$ is given as follows where, for convenience of notation, we double the size of the parameter interval to [0, 2]. Let f^a, f^b be two marked lensshaped models for M. These are 1-parameter families of functions $f_t^a, f_t^b : M \times I \to I$. The new function $f^a * f^b$ will be given by rescaling a family of function $f_t^{ab} : M \times I \to I$, $t \in [0, 2]$ by $(f^a * f^b)_t = f_{2t}^{ab}$. The functions f_t^{ab} are constructed in two stages.

- (1) Starting with $f^c = f^a \cdot f^b$ given by putting f^b to the left of f^a as in Proposition 1.1 but with the parameter domain stretched out. Thus $f^c_t = f^b_t$ for $t \leq 1$ and $f^c_t = f^a_{t-1} \circ g^b$ for t > 1 where $g^b \in \mathcal{C}(M)$ so that $f^b_1 = \pi_I \circ g^b$. See Fig. 2.
- (2) "Merging" the birth point b_2 of $f^a \circ g^b$ (related to the birth point b'_0 of f^a by $b_2 = (g^b)^{-1}(b'_0)$) with the death point b_1 of f^b along the dashed path w_t in Fig. 2 using the markings. The result is f^{ab}_t for $t \in [0, 2]$.



FIGURE 2. The family of functions f_t^c on $M \times I$ for $t \in [0, 2]$ is given to be f_t^b on the left side where $t \in [0, 1]$ and $f_{t-1}^a \circ g^b$ on the right side where $t \in [1, 2]$. The death point b_1 at t_1 is cancelled with the birth point b_2 at t_2 along the path w_t to produce the marked lens shaped model f_t^{ab} .

The family of functions f_t^c , $t \in [0, 2]$, is indicated in Figure 2. The "merging" process is given as follows. We have functions f_t^c for $t \in [t_1, t_2]$ which are nonsingular for $t_1 < t < t_2$ and so that each of $f_{t_1}^c$, $f_{t_2}^c$ has exactly one critical point b_1, b_2 , resp., which is a birth-death point of index $k = \lfloor \frac{n}{2} \rfloor$. Furthermore, the markings on f^b , f^a (when we compose with $(g^b)^{-1}$ in the case of f^a) give paths from b_1, b_2 to a common basepoint. In the case of b_1 this path goes first along the path p_t of index k critical points of f^b_t to the birth point b_0 of f^b and then along the path to the basepoint given by the marking. We combine these to give a path $w_t \in M \times I$ for $t \in [t_1, t_2]$ so that $w_t = b_i$ for $t = t_i$ and f^c_t is nonsingular at w_t for all $t \in (t_1, t_2)$.

At each point w_t for $t \in [t_1, t_2]$ we let v_t be the tangent vector which points in the direction of the gradient of f_t^c for $t \in (t_1, t_2)$ and which points in the positive cubic direction of f_t^c in the kernel of $D^2 f_t^c$ for $t = t_1, t_2$. Using the second part of the markings for f^a, f^b (composed with $D(g^b)^{-1}$ in the case of f^a), we have an oriented tangent k plane at b_1, b_2 which is orthogonal to v_t . For $n \geq 3$ the space of oriented k-planes in \mathbb{R}^n is simply connected. So, this can be extended uniquely up to homtopy to a family of oriented tangent k planes along w_t orthogonal to v_t at each w_t .

We can choose local coordinates $(x, y, z) \in \mathbb{R}^k \times \mathbb{R}^{n-k} \times \mathbb{R}$ for $M \times I$ centered at w_t so that

$$f_t^c(x, y, z) = h_t(z) + ||y||^2 - ||x||^2$$

for all $t \in [t_1, t_2]$ where $h_t(z)$ is a smooth family of functions in one variable z so that $h_t(0) = f_t^c(w_t)$ for each $t \in [t_1, t_2]$. Working only with h_t we can modify the family of function f_t^c in a small neighborhood of the path w_t so that the new function has a birth-death point at w_t for all $t \in [t_1, t_2]$. We can then resolve these degenerate critical points into two Morse critical points in "cancelling position". This gives the family of functions f_t^{ab} which is well-defined up to homotopy. The product is $(f^a * f^b)_t = f_{2t}^{ab}$. The marking for $f^a * f^b$ is given by the marking for f_t^b .

Lemma 1.6. For $n \geq 3$, $\pi_0 \mathcal{D}_0(M)$ forms a groups as outlined above and the mapping $\tilde{\theta}_0 : \pi_0 \mathcal{D}_0(M) \to \pi_0 \mathcal{C}(M)$, given in Definition 1.5 is a group homomorphism.

Proof. The multiplication rule is $(f^a)(f^b) = f^a * f^b$ as described above. It is straightforward to show that this is associative. It is well-defined since the operation is given by the path in the space of oriented k-planes in the kernel of df_t which is given by the marking. Since this multiplication rule is a deformation of the product given in Proposition 1.1, the mapping $\tilde{\theta}_0 : \pi_0 \mathcal{D}_0(M) \to \pi_0 \mathcal{C}(M)$ is a homomorphism of semigroups.

The identity in $\pi_0 \mathcal{D}_0(M)$ is given by any lens-shaped model (f_t) which is symmetrical in the sense that $f_t = f_{1-t}$ for all t. The left inverse of (f_t^b) is f_t^a given by

$$f_t^a = f_{1-t}^b \circ g^{-1} : M \times I \to I$$

where $g \in \mathcal{C}(M)$ is given up to isotopy by $f_1^b = \pi_I \circ g : M \times I \to I$. The marking on f^a is given by $g^b(\gamma)\beta$ where γ is the path from the basepoint to the birth point of f^b which maps by g^b to the death point of f^a and β is the path along the Morse point set p_t back to the birth point of f^a . This represents the inverse of f^b in $\pi_1(\mathcal{F}(M), \mathcal{E}(M))$ by Proposition 1.2. However, f_t^{ab} is symmetrical since $f_t^c = f_b$ and $f_{2-t}^c = f_{1-t}^a \circ g^b = f_t^b$ and the markings match. So, $\pi_0 \mathcal{D}_0(M)$ is a group and $\tilde{\theta}_0$ is a homomorphism.

1.4. Computation of second obstruction. To compute $\pi_0 \mathcal{D}_0(M)$, we need the following. For any group π and left π -module A, we consider $A[\pi] := A \otimes \mathbb{Z}[\pi]$, the direct

sum of copies of A, one for each element of π . This is a π -module with the diagonal action of π where π acts on $\mathbb{Z}[\pi]$ by conjugation:

(1.1)
$$\sigma\left(\sum a_i \otimes g_i\right) = \sum \sigma a_i \otimes \sigma g_i \sigma^{-1}.$$

Each conjugacy class in π gives a submodule of $A[\pi]$, namely the sum of all $A \otimes \tau$ indexed by the conjugates τ of some fixed $\sigma \in \pi$. Let $A[e] = A \otimes e$ be the submodule indexed by the identity $e \in \pi$. We consider the quotient module $A[\pi]/A[e]$.

Definition 1.7. We define $Wh_1^+(\pi; A)$ to be the additive group of coinvariants of the action of π on $A[\pi]/A[e]$:

$$Wh_1^+(\pi; A) := (A[\pi]/A[e])_{\pi}.$$

When the action of π on A is trivial, this is the direct sum of copies of A, one for every nonidentity conjugacy class in π . More generally

$$Wh_1^+(\pi; A) \cong \bigoplus A_{C(\sigma_\alpha)}$$

This is the direct sum over all nonidentity conjugacy classes in π of the coinvariants of the action of $C(\sigma_{\alpha})$ on A where $C(\sigma_{\alpha})$ is the centralizer of a representative σ_{α} of the conjugacy class.

The main example is $\pi = \pi_1 M$ and $A = \mathbb{Z}_2 \oplus \pi_2 M$. For example, if π is finite with trivial action on $\pi_2 M$ then

$$Wh_1^+(\pi_1M; \mathbb{Z}_2 \oplus \pi_2M) = \bigoplus_{c-1} (\mathbb{Z}_2 \oplus \pi_2M)$$

where c is the number of conjugacy classes of elements of $\pi_1 M$.

Then the theorem of Hatcher and Wagoner can be restated as follows.

Theorem 1.8. [2] For dim $M = n \ge 6$ we have an isomorphism of groups:

(1.2) $\lambda_0: \pi_0 \mathcal{D}_0(M) \cong (\mathbb{Z}_2 \oplus \pi_2 M) \otimes \mathbb{Z}[\pi_1 M] / \mathbb{Z}[e] = (\mathbb{Z}_2 \oplus \pi_2 M)[\pi_1 M] / (\mathbb{Z}_2 \oplus \pi_2 M)[e].$

Furthermore, this isomorphism is equivariant with respect to the action of $\pi_1 M \times \mathbb{Z}_2$ given by changing the markings on $\mathcal{D}_0(M)$ and with the diagonal action of $\pi_1 M$ given in (1.1) and the trivial action of \mathbb{Z}_2 . Thus $\pi_0 \mathcal{D}_0(M)_{\pi_1 M} = Wh_1^+(\pi_1 M; \mathbb{Z}_2 \oplus \pi_2 M)$. Furthermore, we have an exact sequence

$$Wh_1^+(\pi_1M; \mathbb{Z}_2 \oplus \pi_2M) \xrightarrow{\overline{\theta}_0} \pi_0\mathcal{C}(M) \to Wh_2(\pi_1M) \to 0$$

where $\overline{\theta}_0$ is the homomorphism induced by $\widetilde{\theta}_0 : \pi_0 \mathcal{D}_0(M) \to \pi_0 \mathcal{C}(M)$.

The idea is that, by Morse theory, a 1-parameter family of functions gives a family of chain complexes. In this case the chain complexes are acyclic. Hatcher and Wagoner show that this family can be deformed until all critical points are in two indices (the "Two-index Theorem"). This gives a family of invertible matrices. There is a K-theoretic obstruction to simplifying this family into the constant family of an $m \times m$ identity matrix. The resulting graphic is a disjoint union of m "eyes" which can be deformed into one "eye" by choosing markings and using the product structure of $\pi_0 \mathcal{D}_0(M)$.

We now recall the definition of the invariant λ_0 in (1.2). Consider a marked lensshaped model (f_t) for M^n with birth-death points at parameter values t_0, t_1 . For each $t_0 < t < t_1$, f_t has critical points p_t, q_t in indices k, k + 1 where $k = \lfloor \frac{n}{2} \rfloor$ with critical values $f_t(p_t) < \frac{1}{2} < f_t(q_t)$. The level surface $V_t^n = f_t^{-1}(\frac{1}{2})$ has an embedded k-sphere S_q^k which is the stable manifold of q_t and an embedded n - k sphere S_p^{n-k} which is the unstable manifold of p_t . For t slightly greater than t_0 and slightly less than t_1 , these spheres meet transversely at one point. However, for values of t between t_0 and t_1 , these spheres may meet in more than one point.

Definition 1.9. [2] The framed bordism invariant of the marked lens-shaped model f_t is defined as follows. The orientation of the negative eigenspace of $D^2 f_t$ at the birth point b_0 of f_{t_0} gives an orientation for S_q^k and a normal framing of S_p^{n-k} . Thus, their intersection is a framed 1-dimensional submanifold $J^1 \subset S_q^k \times (t_0, t_1)$ which has one open component going from t_0 to t_1 . Let J_0 be the union of the closed components of J. Each point in J represents a trajectory of the negative gradient $-\nabla f_t$ from q_t down to p_t . Together with the paths p_t, q_t to the birth point b_0 then the marking which is a path from the basepoint of $M \times I^2$ to b_0 , we obtain a mapping

$$J^1 \to \Omega M$$

Restricting this to J_0 gives an element in the framed bordism group

$$[J_0] \in \Omega_1^{fr}(\Omega M) = (\mathbb{Z}_2 \oplus \pi_2 M)[\pi_1 M].$$

The open component of J maps to the identity component of ΩM . This makes any framed bordism class in $(\mathbb{Z}_2 \oplus \pi_2 M)[e]$ not well-defined since it could be absorbed into the open component of J. Modulo this we obtain the *framed bordism invariant*

$$\lambda_0(f_t) \in (\mathbb{Z}_2 \oplus \pi_2 M)[\pi_1 M]/(\mathbb{Z}_2 \oplus \pi_2 M)[e].$$

If we take the product of two families f_t , f'_t in $\mathcal{D}_0(M)$, we obtain the disjoint union of the closed components J_0, J'_0 of the framed manifolds J, J' for f_t, f'_t . Therefore, λ_0 gives a homomorphism of groups:

$$\lambda_0: \pi_0 \mathcal{D}_0(M) \to (\mathbb{Z}_2 \oplus \pi_2 M)[\pi_1 M]/(\mathbb{Z}_2 \oplus \pi_2 M)[e].$$

Lemma 1.10. [2] For dim $M \ge 6$, λ_0 is an isomorphism.

Proof. In this range of dimensions, $n - k \ge k \ge 3$. So, $S_q^k \times (t_0, t_1)$ has dimension ≥ 4 and there is no obstruction to deforming framed 1-manifolds in $S_q^k \times (t_0, t_1)$. If the framed bordism invariant of f_t is trivial, i.e., if it lies in $(\mathbb{Z}_2 \oplus \pi_2 M)[e]$, each component of J can be absorbed into the open component by connected sum. Since dim $V \ge 6$, the open component can be straightened making f_t into the identity element of $\pi_0 \mathcal{D}_0(M)$. Thus λ_0 is a monomorphism.

To show that λ_0 is surjective, take any generator $(a, b)[\sigma] \in (\mathbb{Z}_2 \oplus \pi_2 M)[\pi_1 M]$. This is represented by a framed circle J_0 in $S_q^k \times (t_0, t_1)$ together with a map $J_0 \to \Omega M$. Since dim $M \ge 6$ this can be realized as a one parameter family of Morse functions. We create two critical points p_t, q_t of index k, k + 1 in cancelling position. Then create two extra geometric incidences which add and subtract σ from the incidence number. Then cancel them again after moving them around using $(a, b) \in \mathbb{Z}_2 \oplus \pi_2 M$. The result has only one geometric incidence between the two critical points and thus they can be cancelled. This creates the desired lens-space module showing that λ_0 is surjective.

We want to extend the framed bordism invariant for $\pi_0 \mathcal{D}_0(M)$ to the case n = 4 (and n = 5). In both n = 4, 5, we have k = 2. So, $S_q^k \times (t_0, t_1)$ has dimension 3 and the framed 1-dimensional manifold J_0 in Definition 1.9 has integer framing invariant (for the same reason that $\pi_3 S^2 = \mathbb{Z}$). In dimension n = 4 we cannot prove the analogue of Lemma 1.10. (The case n = 5 will be discussed further in [7].) However, the framed bordism invariant of Definition 1.9 is still defined and can be refined as follows.

Proposition 1.11. For n = 4 or 5, the framed bordism invariant factors through a well-defined integer framed bordism invariant which is a homomorphism

$$\lambda_0: \pi_0 \mathcal{D}_0(M) \to (\mathbb{Z} \oplus \pi_2 M)[\pi_1 M]/(\mathbb{Z} \oplus \pi_2 M)[e].$$

1.5. Dependence on markings. We show that the integer framed bordism invariant $\widetilde{\lambda}_0$ is $\pi_1 M \times \mathbb{Z}_2$ equivariant.

Lemma 1.12. For n = 4, 5, the integer framed bordism invariant λ_0 from Proposition 1.11 is equivariant with respect to the action of $\pi_1 M \times \mathbb{Z}_2$ given on $\mathcal{D}_0(M)$ by changing the markings and on $(\mathbb{Z} \oplus \pi_2 M)[\pi_1 M]$ by the diagonal action of $\pi_1 M$ as in (1.1) and the sign action of \mathbb{Z}_2 on \mathbb{Z} .

To clarify the statement, the group of coinvariants of the action of $\pi_1 M \times \mathbb{Z}_2$ on $(\mathbb{Z} \oplus \pi_2 M)[\pi_1 M]/(\mathbb{Z} \oplus \pi_2 M)[e]$ is equal to $Wh_1^+(\pi_1 M; \mathbb{Z}_2 \oplus \pi_2 M)$. In other words, the sign of the integer framing number of J_0 is not intrinsic. It just comes from the marking. This implies the following.

Proposition 1.13. For n = 4, 5 the integer framed bordism invariant induces a welldefined mapping

$$\overline{\lambda}: \pi_0 \mathcal{D}(M) \to Wh_1^+(\pi_1 M; \mathbb{Z}_2 \oplus \pi_2 M).$$

To understand the proof of Lemma 1.12, it helps to consider a concrete equivalent problem. What happens to the generator of $\pi_3 S^2$ when we reverse the orientation for both source and target? When we reverse the orientation of S^2 , nothing happens! This is because the Hopf map $S^3 \to \mathbb{C}P^1 = S^2$ commutes with complex conjugation which is homotopic to the identity on S^3 but reverses the orientation of $\mathbb{C}P^1 = S^2$. Reversing the orientation of S^3 is negation on $\pi_3 S^2$ by definition of its group structure. Therefore, reversing the orientation of both S^2 and S^3 acts on $\pi_3 S^2 = \mathbb{Z}$ by changing sign.

Proof of Lemma 1.12. Each closed component of the framed 1-manifold J_0 gives an oriented cycle in the loop space of $M \times I^2$ at the birth point of f_t . This contributes $(n, a)[\sigma]$ to the integer framing invariant where $n \in \mathbb{Z}$, $a \in \pi_2 M$ and $\sigma \in \pi_1 M$. When we change the orientation of the descending plane of the birth point, we change the orientation of the descending sphere $S_q^2 \times (t_0, t_1)$ and the normal orientation of the ascending sphere $S_p^{n-2} \times (t_0, t_1)$. The result is that tangential orientation of the intersection $J \subset S_q^2 \times (t_0, t_1)$ is unchanged. However, the self-intersection number of J in $S_q^2 \times (t_0, t_1)$ changes sign. This will change the invariant from $(n, a)[\sigma]$ to $(-n, a)[\sigma]$. This is similar to what happens with $\pi_3 S^2$.

The other part of the marking is the path from the basepoint of $M \times I^2$ to the birth point of f_t . is changed by, say $\gamma \in \pi_1 M$, this invariant will change to $(n, \gamma a)[\gamma \sigma \gamma^{-1}]$. The framing number n does not change since it does not use the path to the basepoint. \Box

10

1.6. **Involution.** We also have a very useful involution ε which flips the function f "upside-down". It is given by

$$\varepsilon f(x,t) = 1 - f(x,1-t).$$

This is projection to the second coordinate of the function εg given for each g(x,t) = (h(x,t), f(x,t)) in $\mathcal{C}(M)$ by

$$\varepsilon g(x,t) = (r^{-1}h(x,1-t), 1 - f(x,1-t))$$

where r is the automorphism of M given by g(x, 1) = (r(x), 1). Since ε switches the stacking direction for functions (Proposition 1.3) we have

$$\varepsilon(f \bullet f') = \varepsilon(f') \bullet \varepsilon(f).$$

The involution ε turns the graphic of a function upside-down. Also, it changes the index k of a Morse critical point (x,t) to index n + 1 - k (and the critical point moves to (x, 1 - t)). Thus, in the case when dim M = n is even, this involution sends $\mathcal{D}(M)$ to $\mathcal{D}(M)$.

Proposition 1.14. Suppose that $n \ge 4$ is even and $(f_t) \in \mathcal{D}(M)$ with $\overline{\lambda}(f_t) = (n, \alpha)[\sigma] \in Wh_1^+(\pi_1M; \mathbb{Z}_2 \oplus \pi_2M)$ where $n \in \mathbb{Z}_2$, $\alpha \in \pi_2M$ and $\sigma \in \pi_1M$. Then

$$\overline{\lambda}(\varepsilon(f_t)) = (n + w_2(\alpha), -\alpha)[\sigma^{-1}].$$

where $w_2 \in H^2(M; \mathbb{Z}_2)$ is the second Stiefel-Whitney class of M.

Proof. First, choose a marking for (f_t) . By linearity of the invariant, we can restrict attention to the contribution of one closed component, say C, of J, the intersection of the descending sphere $S_q^k \times (t_0, t_1)$ with the ascending sphere $S_p^k \times (t_0, t_1)$ in $V^{2k} \times (t_0, t_1)$.

To compute the invariant for $\varepsilon(f_t)$, we take the same family of functions f_t with the same critical points p_t, q_t but with the negative gradient $-\nabla f_t$ replaced with the positive gradient. The loop representing σ in $\overline{\lambda}(f_t)$ goes from the birth point b_0 up to q_t , then down through C to p_t and back to b_0 . The loop for $\overline{\lambda}(\varepsilon(f_t))$ goes the other way (from p_t up to q_t through C). So, σ is replaced with σ^{-1} .

The tangential orientation of J is unchanged since, by definition, it points from the birth point to the death point on the open component. So, the component C is tangentially oriented in the same direction for both (f_t) and $\varepsilon(f_t)$. But, the σ direction is changed. So, the $\pi_2 M$ element changes sign from α to $-\alpha$. This element $\alpha \in \pi_2 M$ is represented by an embedded 2-sphere with equator C, upper hemisphere D^2_+ contained in the descending manifold $D_q^{k+1} \times (t_0, t_1)$ of q_t and lower hemisphere D^2_- contained in the ascending manifold $D_p^{k+1} \times (t_0, t_1)$ of p_t . For $\varepsilon(f_t)$ the roles of D^2_+ , D^2_- are switched.

The framing number n of C in $\overline{\lambda}(f_t)$ is given by comparing the normal framing of $S_p^k \times (t_0, t_1)$ along C, which extends to a normal framing of D_-^2 in $D_p^{k+1} \times (t_0, t_1)$, with the tangential framing of D_+^2 in $D_q^{k+1} \times (t_0, t_1)$. The framing number n' of C in $\overline{\lambda}(\varepsilon(f_t))$ is given by comparing the normal framing of D_+^2 in $D_q^{k+1} \times (t_0, t_1)$ with the tangential framing of D_-^2 in $D_p^{k+1} \times (t_0, t_1)$. So, the sum n+n' compares the normal bundle of D_+^2 in $M \times I^2$ with the normal bundle of D_-^2 . In other words, $n+n' = w_2(D_+^2 \cup D_-^2) = w_2(\alpha)$.

Proposition 1.14 will guide us in the remainder of this paper. We will assume that M is a 4-manifold with an embedded 2-sphere with odd w_2 . For example, we could take connected sum with $\mathbb{C}P^2$. Using an embedded 2-sphere we will construct a lens-shaped model for M with invariant $\alpha[\sigma]$ where $\alpha \in \pi_2 M$ is the class of the embedded 2-sphere and σ is any element of $\pi_1 M$. The explicit construction will allow us to compute the framing number. It will be the degree of the normal bundle of the embedded 2-sphere in M, i.e., its self-intersection number. We do the same using σ^{-1} and apply the involution ε . By Proposition 1.14, the sum will give all nontrivial elements of $Wh_1^+(\pi_1 M; \mathbb{Z}_2)$.

2. Constructing pseudoisotopies using two disks

To prove Theorems A and B we need embedded disks: $D_q^3, D_p^2 \subset M^4$ as shown in Figure 3. We use the fact that the self-intersection number of the 2-sphere $\mathbb{C}P^1$ in $\mathbb{C}P^2$, i.e., the degree of its normal bundle, is 1. Conversely, if M contains an embedded 2-sphere with self-intersection number 1, it must have the form $M = X \# \mathbb{C}P^2$.

Lemma 2.1. Let $S_0^2 \subset M$ be an embedded 2-sphere with self-intersection number w. Given any $\sigma \in \pi_1 M$, there are embedded disks $D_q^3 \subset M^4$ and $D_p^2 \subset M^4$ with boundaries $S_q^2 = \partial D_q^3$, $S_p^1 = \partial D_p^2$ with the following properties (Figure 3).

- $(1) S_q^2 \cap D_p^2 = \{a\} \subset int D_p^2$ $(2) D_q^3 \cap S_p^1 = \{b\} \subset int D_q^3$ $(3) D_q^3 \cap D_p^2 \text{ is the disjoint union of}$ (a) a line segment from a to b (denoted J)(b) a circle S^1_* .
- (4) Any path from b to S^1_* in D^3_q followed by any path from S^1_* to b in D^2_p is homotopic to (a conjugate of) the chosen element $\sigma \in \pi_1 M$.
- (5) S^1_* bounds two 2-disks $B^2_q \subset D^3_q$ and $B^2_p \subset D^2_p$ whose union is homotopic to S^2_0 . (6) The normal framing of S^1_* in D^3_q given by any trivialization of the normal bundle of D_n^2 in M has winding number w.

Remark 2.2. We choose a path from b to the base point of M so that the loop at b represents the element $\sigma \in \pi_1 M$. We also fix an orientation of the normal bundle of D_p^2 in M (giving a normal orientation of S_p^1 in M) and an orientation of D_q^3 so that these orientations agree at $D_q^3 \cap S_p^1 = \{b\}$. These choices will give a marking on the lens-shaped model we are constructing (Definition 1.5.)

We shall now construct a 1-parameter family of functions (f_t) on $M^4 \times I$ assuming we are given embedded disks $D_q^3, D_p^2 \subset M$ as described above and shown in Figure 3. Start with $f_0 = \pi_I$, the projection to I = [0, 1]. To show that the result is a nontrivial element of $\pi_0 \mathcal{C}(M)$ we need to assume that w = 1.

We introduce a pair of cancelling Morse critical points p_t, q_t of indices 2,3 for $t > t_0$ where f_{t_0} has a birth-death point of index 2. Choose a path from this birth-death point to the basepoint of $M \times I^2$.





FIGURE 3. Two embedded disks $D_p^2, D_q^3 \subset M$ with $D_p^2 \cap D_q^3 = S_*^1 \coprod J$. In this figure, D_p^2 is flat and D_q^3 is drawn as a boundary connected sum of two 3-disks along a path given by σ . $B_p^2 \subset D_p^2$ is a 2-disk with $\partial B_p^2 = S_*^1$. \tilde{b} is the normal vector field of S_p^1 pointing inward into D_p^2 .



For t slightly greater than t_0 , the regular levels of the Morse function f_t start at the top: $M^4 = M \times 1$ as shown in Figure 3, with an embedded 2-disk $D_p^2 \subset$ M^4 (shaded on the left), performs surgery on $\partial D_p^2 =$ S_p^1 at q_t to produce the intermediate level surface V^4 shown in Figure 4 with an embedded sphere S_L^2 which is the stable sphere of q_t . $D_p^2 \subset M$ descends to $S_p^2 \subset V$, the unstable sphere of p_t .

Thus S_L^2 is the stable sphere of q_t , the set of all points in V^4 which are carried by trajectories of ∇f_t to q_t . Points on the surgery circle $S_p^1 = \partial D_p^2$ give normal vector fields for S_L^2 in V^4 and, in the direction corresponding to the point $b = \partial D_p^2 \cap D_q^3$, a small 2-sphere in D_q^3 centered b in M, under the flow generated by $-\nabla f_t$ goes to a sphere $\tilde{S}_L^2 \subset V^4$ parallel to S_L^2 .

Let $S_p^2 \subset V^4$ be the unstable sphere of p_t . Since p_t, q_t are a cancelling pair of critical points, the two spheres S_L^2, S_p^2 intersect transversely at one point. We can take S_p^2 to be the image of D_p^2 in V^4 .

For $t_1 \leq t \leq t_2$ we do an isotopy of the descending sphere of the critical point q_t from S_L^2 to S_q^2 along the cylinder $\widetilde{D}_q^3 = S^2 \times \widetilde{J}$. This intersects S_p^2 along the circle S_*^1 .





FIGURE 4. The level surface V^4 obtained by doing surgery on $S_p^1 \subset M$. The disk $D_p^2 \subset M$ becomes the sphere $S_p^2 \subset V$ which meets $S_L^2 \subset V$, the sphere dual to S_p^1 , transversely at \tilde{b} . In V, the disk $D_q^3 \subset M$ becomes $\tilde{D}_q^3 = S^2 \times \tilde{J}$ with boundary $\partial \tilde{D}_q^3 = S_L^2 \coprod S_q^2$. In this figure, \tilde{D}_q^3 is flat.

By construction, any normal framing of \widetilde{D}_q^3 in $V \times I$ gives a trivial framing for S_*^1 in $S_p^2 \times I$ and, by Condition (6), for n odd, any normal framing of $S_p^2 \times I$ gives a nontrivial framing of S_*^1 in \widetilde{D}_q^3 , the trace of the deformation of S_L^2 to S_q^2 . By (5), the trajectories of $-\nabla f_t$ through S_*^1 gives a 2-sphere homotopic to S_0^2 . Also, each of these trajectories, together with the paths p_t , q_t to the birth-death point of f_{t_0} followed by the chosen path to the basepoint of $M \times I^2$ gives the chosen element $\sigma \in \pi_1 M$.

When n = 1 and $M = X \# \mathbb{C}P^2$, M contains a subspace $Y = X - D^4$ with $\pi_1 Y = \pi_1 M$ and $\mathbb{C}P^1$ is disjoint from Y. So, the projection of the first Postnikov invariant $k_1 \in H^3(\pi_1 M; \pi_2 M)$ to $H^3(\pi_1 M; \mathbb{Z}) = H^3(\pi_1 M; \pi_2 \mathbb{C}P^2)$ is trivial. Thus, in the exact sequence:

$$K_3(\mathbb{Z}[\pi_1M]) \xrightarrow{\chi} Wh_1^+(\pi_1; \mathbb{Z}_2 \oplus \pi_2M) \to \pi_1\mathcal{P}(M) \to Wh_2(\pi_1M) \to 0$$

the boundary map χ does not hit the element $\tau[\sigma] \in Wh_1^+(\pi_1 M; \pi_2 M)$ where $\tau \in \pi_2 M$ is the class of $\mathbb{C}P^1$ since, by [5], this factor of χ is given by:

$$K_3(\mathbb{Z}[\pi_1 M]) \to H_3(\pi_1 M; \pi_2 M) \xrightarrow{\cap k_1} H_0(\pi_1 M; \pi_2 M) \to Wh_1^+(\pi_1 M; \pi_2 M)$$

which does not hit the summand $Wh_1^+(\pi_1 M; \pi_2 M)$ generated by $\tau[\pi_1 M]$.

At t_2 we can cancel the critical points at a birth-death point to give a marked lens-shaped model with invariant $\tau[\sigma] \in Wh_1^+(\pi_1 M; \pi_2 M)$ where $\tau \in \pi_2 M$ is the homotopy class of $\mathbb{C}P^1$.



More generally, this same construction proves the following.

Theorem 2.3 (Theorem A). Given a compact 4-manifold M and $\tau \in \pi_2 M$ represented by an embedded 2-sphere and $\sigma \in \pi_1 M$, the obstruction element $\tau[\sigma] \in Wh_1^+(\pi_1 M; \pi_2 M)$ lifts to $\mathcal{C}(M)$. Furthermore, when the embedded 2-sphere has self-intersection number ± 1 , this element of $\pi_0 \mathcal{C}(M)$ is stably nontrivial (survives to a nontrivial element of $\pi_0 \mathcal{P}(M)$).

Remark 2.4. The fact that the obstruction element $\tau[\sigma]$ lifts to an element of $\pi_0 \mathcal{C}(M)$ does not imply that its image in $\pi_0 \mathcal{C}(M)$ is nonzero except in the case when the self-intersection number of τ is $n = \pm 1$ since the Postnikov invariant argument given above only works in that case.

3. Realizing the framing obstruction

We use the construction from the previous section to construct elements realizing every element of the framing obstruction group

$$Wh_{1}^{+}(\pi_{1}M;\mathbb{Z}_{2}).$$

However, the comment in Remark 2.4 still applies. Lifting these obstruction elements to $\pi_0 \mathcal{C}(M)$ does not imply that their images are nonzero, except in the following case.

Suppose that $\pi_1 M$ is free abelian of finite nonzero rank. Then, as observed by Hatcher, $Wh_1^+(\pi_1 M; \mathbb{Z}_2)$ will be an infinitely generated group. But $K_3(\mathbb{Z}[\pi_1 M])$ is finitely generated by the fundamental theorem of K-theory. From the exact sequence

$$K_3(\mathbb{Z}[\pi_1 M]) \xrightarrow{\chi} Wh_1^+(\pi_1 M; \mathbb{Z}_2) \oplus Wh_1^+(\pi_1 M; \pi_2 M) \to \pi_0 \mathcal{P}(M)$$

we can then conclude that there are infinitely many nontrivial elements of $\pi_0 \mathcal{C}(M)$ coming from $Wh_1^+(\pi_1 M; \mathbb{Z}_2)$. However, these elements might not be 2-torsion when M is 4-dimensional.

For nontrivial $\sigma \in \pi_1 M$ and $\tau \in \pi_2 M$ represented by an embedded 2-sphere in Mwith self-intersection number ± 1 we realize the obstruction $\tau[\sigma]$ as in Theorem 2.3 and do it again for $\tau[\sigma^{-1}]$ where we choose markings for our lens-shaped models (lifting them to elements of $\mathcal{D}_0(M)$) to make these obstructions well-defined (Definition 1.5). However, the second time, we apply the involution ε which turns the Morse functions upside down to make the intersection circle S^1_* trivially framed in \widetilde{D}^2_q and nontrivially framed in $S^2_p \times I$. As in the proof of Proposition 1.14, the product of these two elements of $\mathcal{D}_0(M)$ will map to zero in $\pi_2 M[\pi_1 M]/\pi_2 M[e]$ since $\tau[\sigma] - \tau[\sigma] = 0$. The integer framed bordism invariant (Proposition 1.11) will be $\pm[\sigma]$, a generator of $\mathbb{Z}[\pi_1 M]/\mathbb{Z}[e]$. This gives:

Theorem 3.1 (Theorem B). Let $M = X \# \mathbb{C}P^2$ be the connected sum of a nonsimply connected 4-manifold X with $\mathbb{C}P^2$. Then the integer framed bordism invariant $\widetilde{\lambda}_0$: $\pi_0 \mathcal{D}_0(M) \to \mathbb{Z}[\pi_1 M]/\mathbb{Z}[e]$ is surjective. In particular, every element in the image of the second obstruction map $\theta : Wh_1^+(\pi_1 M; \mathbb{Z}_2) \to \pi_0 \mathcal{P}(M)$ lifts to $\pi_0 \mathcal{C}(M)$.

More generally, an analogous construction gives the following.

Theorem 3.2. Suppose M^4 contains an embedded 2-sphere with odd self-intersection number. Then every element of the image of θ : $Wh_1^+(\pi_1 M; \mathbb{Z}_2) \to \pi_0 \mathcal{P}(M)$ lifts to $\pi_0 \mathcal{C}(M)$.

4. Construction of the two disks

We prove Lemma 2.1 by constructing the two disks D_p^2 , D_q^3 first in the special case when the embedded 2-sphere is $\mathbb{C}P^1$, then more generally for any embedded 2-sphere by modifying the construction in the special case.

The disk D_q^3 will be the boundary connected sum of two 3-disks D_0^3 and D_1^3 where D_p^2, D_0^3, D_1^3 will all be contained in a punctured $\mathbb{C}P^2$ contained in M.

Let D_p^2 be the set of all point in $\mathbb{C}P^2$ with coordinates [1, z, 0] where $|z| \leq 3$ and let B_p^2 be the subset of those points [1, z, 0] with $|z| \leq 1$. These are 2-disks with $\partial B_p^2 = S_*^1$ being the set of all [1, x, 0] where |x| = 1.

Let $S^1_* \times D^2$ denote the set of all points in $\mathbb{C}P^2$ of the form [1, x, z] where |x| = 1and $|z| \leq 1$. This is the restriction to S^1_* of the normal disk bundle of D^2_p in $\mathbb{C}P^2$. Thus $S^1_* \times D^2$ meets D^2_p transversely along S^1_* as shown in Figure 5.



FIGURE 5. The 2-disk $D_p^2 \subset \mathbb{C}P^1 \subset \mathbb{C}P^2$ with subdisk B_p^2 . The boundary $\partial B_p^2 = S_*^1$ is in red. In blue are $S_*^1 \times D^2$, meeting D_p^2 transversely along S_*^1 and D_0^3 meeting ∂D_p^2 transversely at one point with $J = D_0^3 \cap D_p^2$ in red.

Let D_0^3 be the 3-disk in $\mathbb{C}P^2$ which meets $\partial D_p^2 = S_p^1$ transversely at one point and so that $D_0^3 \cap D_p^2$ is the line segment J as shown in Figure 5.

To construct D_1^3 , we take the twisted circle \widetilde{S}^1_* in $\partial(S^1_* \times D^2)$ given by

$$S^1_* = \{ [1, x, x] \in \mathbb{C}P^2 \mid |x| = 1 \}.$$

This bounds a 2-disk in $\mathbb{C}P^2$ in the complement of $\mathbb{C}P^1$ by taking the set of all $[t, x, x] \in \mathbb{C}P^2$ with |x| = 1 and $t \in [0, 1]$. This is a 2-disk since $[t, x, x] = [t\overline{x}, 1, 1]$. The union of B_p^2 with this 2-disk is a 2-sphere homotopic to $\mathbb{C}P^1$. Use this 2-disk to attach a handle to $S_*^1 \times D^2$ to obtain a 3-disk D_1^3 which meets D_p^2 transversely along S_*^1 .

Finally, let D_q^3 be given by taking the boundary connect sum of the two 3-disks D_1^3 , D_0^3 along a path in M given by the chosen element $\sigma \in \pi_1 M$. The result is as shown in Figure 3 proving Lemma 2.1 in the special case then $S_0^2 = \mathbb{C}P^1$.

Proof of Lemma 2.1 in general. Let $S_0^2 \subset M$ be an embedded 2-sphere with self-intersection number n. We follow the same outline as above using Figure 5.

Let D_p^2 in Figure 5 be the southern hemisphere of S_0^2 with subdisk $B_p^2 \subset D_p^2 \subset S_0^2$ and $S_*^1 = \partial B_p^2$. For the circle bundle $\partial S_*^1 \times D^2 \to S_*^1$ take the section \tilde{S}_*^1 with winding number n. This is the section which extends to a section B_q^2 of the normal circle bundle over $S_0^2 \setminus B_p^2$. We can use this section to attach a 3-dimensional 2-handle to $S_*^1 \times D^2$. This gives a 3-disk D_1^3 which intersects D_p^2 transversely along S_*^1 so that $B_q^2 \cup B_p^2 \subset D_1^3 \cup D_p^2$ is homotopic to S_0^2 . Let D_q^3 be the boundary connected sum of D_1^3 with a 3-disk D_0^3 given as before using the path σ to do the connected sum. This produces the disks D_p^2 and D_q^3 with all the required properties.

Acknowledgements

The author would like to thank Sander Kupers for encouraging him to write up these results and for commenting on earlier versions of this manuscript. He also thanks Bjørn Jahren and Tom Goodwillie for useful correspondences. The author is indebted to Allen Hatcher for explaining pseudoisotopy theory to him and for posing the question of what happens in low dimensions. The author is supported by the Simons Foundation.

References

- Cerf, Jean. La stratification naturelle des espaces de fonctions différentiables réelles et le théoreme de la pseudo-isotopie. Publications Mathématiques de l'IHÉS 39 (1970): 5–173.
- [2] Hatcher, Allen, and John Wagoner. Pseudo-isotopies of compact manifolds. Société mathématique de France, 1973.
- [3] Igusa, Kiyoshi. The $Wh_3(\pi)$ Obstruction for Pseudoisotopy. PhD thesis. 1978.
- [4] _____. On the algebraic K-theory of A_{∞} -ring spaces. Lecture notes in Math 967: Algebraic K-theory. Springer, Berlin, Heidelberg, 1982. 146–194.
- [5] _____. What happens to the formula of Hatcher and Wagoner when the first Postnikov invariant of M is nontrivial? Lecture notes in Math 1046: Algebraic K-Theory, Number Theory, Geometry and Analysis, pp. 104–172. 1982.
- [6] _____. The stability theorem for smooth pseudoisotopies. K-theory 2, no. 1–2 (1988).
- [7] _____. Second obstruction to pseudoisotopy II: dimension 5. under revision.
- [8] Kwasik, Slawomir. Low-dimensional concordances, Whitney towers and isotopies. Math Proc of Cambridge Phil Soc, vol. 102, no. 1, pp. 103–119 (1987).

DEPARTMENT OF MATHEMATICS, BRANDEIS UNIVERSITY, WALTHAM, MA 02454 Email address: igusa@brandeis.edu