

# FAMILY BAUER–FURUTA INVARIANT, EXOTIC SURFACES AND SMALE CONJECTURE

JIANFENG LIN, ANUBHAV MUKHERJEE

**ABSTRACT.** We establish the existence of a pair of exotic surfaces in a punctured  $K3$  which remains exotic after one external stabilization and have diffeomorphic complements. A key ingredient in the proof is a vanishing theorem of the family Bauer–Furuta invariant for diffeomorphisms on a large family of spin 4-manifolds, which is proved using the tom Dieck splitting theorem in equivariant stable homotopy theory. In particular, we prove that the  $S^1$ -equivariant family Bauer–Furuta invariant of any orientation-preserving diffeomorphism on  $S^4$  is trivial and that the  $\text{Pin}(2)$ -equivariant family Bauer–Furuta invariant for a diffeomorphism on  $S^2 \times S^2$  is trivial if the diffeomorphism acts trivially on the homology. Therefore, these invariants do not detect exotic self-diffeomorphisms on  $S^4$  or  $S^2 \times S^2$ . Furthermore, our theorem also applies to certain exotic loops of diffeomorphisms on  $S^4$  (as recently discovered by Watanabe) and show that these loops have trivial family Bauer–Furuta invariants. En route, we observe a curious element in the  $\text{Pin}(2)$ -equivariant stable homotopy group of spheres which could potentially be used to detect an exotic diffeomorphism on  $S^4$ .

## 1. INTRODUCTION

The Bauer–Furuta invariant of smooth 4-manifolds is a stable homotopy refinement of the classical Seiberg–Witten invariant [11]. It contains subtle but important information about the underlying 4-manifolds which is invisible from other invariants. In particular, the Bauer–Furuta invariant can detect new exotic smooth structures (for example, on  $K3\#K3$ ) and its family version [47, 53] is a powerful tool in detecting exotic diffeomorphisms [31, 35]. For this reason, this invariant has been actively studied in recent years [8, 9, 13, 26, 29, 31, 41, 46, 48]. While the classical Seiberg–Witten invariant is an integer, the Bauer–Furuta invariant is an element in the (equivariant) stable homotopy group of spheres. This allows one to apply powerful techniques in equivariant topology (e.g. equivariant K-theory and equivariant homology) to extract information on the Bauer–Furuta invariant and to study 4-manifolds. This idea has resulted in many remarkable theorems including Furuta’s  $\frac{10}{8}$ -theorem [18] (see also [19, 25]) and Manolescu’s disproof of the triangulation conjecture [36] (see also [34]). In this paper, we follow this line of thought and use the equivariant stable homotopy theory to prove a general vanishing result for the family Bauer–Furuta invariant of diffeomorphism on a large family of spin 4-manifolds (Theorem 1.9). This vanishing theorem has several applications, which we discuss first.

**1.1. Exotic surfaces.** The first application concerns exotic surfaces in 4-manifolds. Recall that a pair of compact surfaces  $\Sigma, \Sigma'$  in a 4-manifold  $X$  with  $\partial\Sigma = \partial\Sigma' \subset \partial X$  are called *exotic surfaces* if there exists a topological isotopy of  $X$  relative to the boundary that takes  $\Sigma$  to  $\Sigma'$  but there is no such smooth isotopy. Understanding exotic surfaces in 4-manifolds is of great interest since they are closely related to exotic smooth structures on 4-manifolds. In recent years, various methods have been used to detect examples of exotic surfaces (e.g., the Donaldson’s invariants [15], the

Seiberg–Witten invariants [16], Heegaard–Floer homology [27], symplectic & contact geometry [23] and Khovanov homology [24]). In this paper, we add the family Bauer–Furuta invariant to this list. As we will see, this method reveals a new phenomenon : some exotic surfaces can survive a stabilization.

To further explain this, we mention an important principle in 4-dimensional topology, discovered by Wall in the 1960s [51]. This principle says that exotic phenomena are all eliminated by sufficiently many stabilizations, i.e., taking connected sum with  $S^2 \times S^2$ 's. Since then, it has been a fundamental problem to search for exotic phenomena that survives one stabilization. In the case of exotic surfaces, work of Perron [38] and Quinn [39] showed that exotic surfaces always become smoothly isotopic after taking sufficiently many stabilizations (away from the surfaces). *It is a long standing open-question, whether a single stabilization is enough for this purpose?* (See [2, 5] for studies of this problem.) Using the 4-dimensional light bulb theorem (proved by Gabai [20] and Scheniderman–Teichner [42]), Auckly–Kim–Melvin–Ruberman–Schwartz [6] proves a surprising result in the positive direction: They showed that for exotic pairs of connected surfaces in non-characteristic homology classes with simply-connected complements, a single stabilization is enough. In this paper, we prove a result in the negative direction by producing the first pair of orientable exotic surfaces that remain non-isotopic after a single stabilization.

**Theorem 1.1.** *There exists a link  $L$  on the boundary  $S^3$  of punctured  $K3$  that bounds a pair of slice disks  $D_L$  and  $D'_L$  which are topologically isotopic relative to the boundary but not smoothly so. Moreover, they remain exotic after a single stabilization, i.e. connected sum with  $S^2 \times S^2$ , and their complements are diffeomorphic.*

*Remark 1.2.* As mentioned in Theorem 1.1, another interesting feature of our example is that the complements of  $D_L$  and  $D'_L$  are diffeomorphic. Examples with this feature was produced by Schwartz [44] (for closed manifolds, using non-triviality of an h-cobordism) and Akbulut [3] (for embedded disks in the 4-ball, using corks). Note that recent work of Juhász–Miller–Zemke [27] and Hayden [23] give some fascinating examples of knots in  $S^3$  that bounds exotic surfaces in 4-ball. In those examples it is not clear if the complements of the surfaces are diffeomorphic or not.

We sketch the proof of Theorem 1.1 as follows (see Section 3 for details). Let  $D_L = \bigsqcup_{22} D^2$  be the disjoint union of the cores of 2-handles in the punctured  $K3$  and let  $D'_L$  be the image of  $D_L$  under the Dehn twist on the punctured  $K3$  along its boundary. By a result in [30] and [37], this Dehn twist is topologically isotopic to the identity map relative to the boundary. Hence  $D_L$  and  $D'_L$  are topologically isotopic relative to the boundary. However,  $D_L$  and  $D'_L$  are not smoothly isotopic relative to the boundary even after a stabilization. Suppose they were. Then we show that there is a smooth isotopy between  $\delta \# \text{id}_{S^2 \times S^2}$  and  $\text{id}_{K3 \# K3} \# f$ , where  $\delta$  denote the Dehn twist on  $K3 \# K3$  and  $f$  is a diffeomorphism  $S^2 \times S^2$ . By the connected sum formula of the Bauer–Furuta invariant and our vanishing theorem (Theorem 1.9) in the case of  $S^2 \times S^2$ , the  $\text{Pin}(2)$ -equivariant family Bauer–Furuta invariant of  $\text{id}_{K3 \# K3} \# f$  is trivial. This contradicts with a result of the first author [35], which states that the corresponding invariant of  $\delta \# \text{id}_{S^2 \times S^2}$  is nontrivial.

It is worth noting that the argument in the current paper is essentially different from the argument in [35] by the first author. In [35], there is a specific diffeomorphism  $\delta \# \text{id}_{S^2 \times S^2}$ . By partially computing it's family Bauer–Furuta invariant, one can show that the invariant is non vanishing. However, in the current paper, we just have a map of the form  $\text{id}_{K3 \# K3} \# f$  with  $f$

being a hypothetical diffeomorphism on  $S^2 \times S^2$ . Since we don't have much control over  $f$ , we have to prove a general vanishing theorem for the family Bauer–Furuta invariant that works for arbitrary  $f$  that acts trivially on homology. This is where extra arguments in equivariant stable homotopy theory have to get involved.

We also emphasize that in the proof of Theorem 1.1, the  $\text{Pin}(2)$ -equivariance is essentially used multiple times for different purposes. First, the  $\text{Pin}(2)$ -symmetry is included in the definition of the equivariant Bauer–Furuta invariant, so that this invariant can be nonzero even after a stabilization. This special feature, which is different from many other gauge theoretic invariants (e.g., the Seiberg–Witten invariant, Donaldson's polynomial invariant, the  $S^1$ -equivariant and the non-equivariant Bauer–Furuta invariant), makes the  $\text{Pin}(2)$ -equivariant Bauer–Furuta invariant particularly suitable for detecting exotic phenomena under stabilizations. Second, because of this  $\text{Pin}(2)$ -symmetry, the map between spheres which underlies the Bauer–Furuta invariant must be a  $\text{Pin}(2)$ -equivariant map. This actually imposes a very strong constraint on the possible values of the Bauer–Furuta invariant. This constraint is crucial in the proof of our vanishing theorem.

*Remark 1.3.* From the construction of  $D_L$  and  $D'_L$ , we see that  $L$  can be taken to be the attaching link in any Kirby diagram of  $K3$  (without 1-handles and 3-handles). Different choices of  $L$  will give different exotic surfaces. The following proposition gives an upper bound of the stabilizations needed to make these surfaces smoothly isotopic. It is proved by combining the argument in [6] with the recent work of Schwartz [43], which generalizes the light bulb theorem to embedded disks.

*Proposition 1.4.* (1) Let  $L_0 \subset L$  be a sublink and let  $D_{L_0} \subset D_L$  and  $D'_{L_0} \subset D'_L$  be the subsurfaces that contain all the components with boundary in  $L_0$ . Suppose  $L_0$  is an unlink. Then  $D_{L_0}$  is smoothly isotopic to  $D'_{L_0}$  relative to the boundary (without any stabilizations).

(2) Suppose  $L$  contains an  $m$ -component unlink ( $m \geq 0$ ). Then after  $22 - m$  stabilizations,  $D_L$  and  $D'_L$  are smoothly isotopic relative to the boundary. In particular, for any  $L$ ,  $D_L$  and  $D'_L$  are smoothly isotopic after 22 stabilizations.

In Section 3, we give an example of  $L$  with all components being the unknot. By Proposition 1.4, each component of  $D_L$  is smoothly isotopic to the corresponding component of  $D'_L$ . Therefore, in this example, the exotica that persists a stabilization is the knotting between different components. Proposition 1.4 also motivates the following question,

*Question 1.* Given  $L$ , what is the minimal number of stabilizations that can make  $D_L$  and  $D'_L$  smoothly isotopic?

**1.2. Smale Conjecture.** Our second motivation comes from a famous conjecture by Smale. In dimension 4, this conjecture states that the natural inclusion  $O(5) \hookrightarrow \text{Diff}(S^4)$  is a homotopy equivalence [Kirby's Problem List [28], Problem 4.34, 4.126]. Recently, Watanabe [52] disproved Smale Conjecture by showing that the induced maps on higher homotopy groups are not surjective. Here we state a special case of his result which is most relevant to our discussion:

**Theorem 1.5** (Watanabe, [52]). *There is a non-torsion element  $\gamma_1 \in \pi_1(\text{Diff}(S^4))$  such that for any nonzero integer  $d$ , one has*

$$d\gamma_1 \notin \text{image}(\pi_1(O(5)) \rightarrow \pi_1(\text{Diff}(S^4))).$$

*Furthermore,  $\gamma_1$  is an exotic loop of diffeomorphisms. Namely,  $\gamma_1$  is null-homotopic in the homeomorphism group of  $S^4$ .*

We note that the following important case is still open:

**Conjecture 1** (Smale). *Any orientation preserving self-diffeomorphism of  $S^4$  is smoothly isotopic to Identity.*

Watanabe’s proof is not gauge theoretic. Instead, he used Kontsevich’s characteristic classes to detect  $\gamma_1$ . Note that gauge theory has also been very successful in detecting exotic diffeomorphisms on 4-manifolds [2, 5, 10, 31, 35, 40]. In an upcoming paper by Auckly–Ruberman [7], they succeeded in using gauge theory to detect exotic loops (and elements in higher homotopy groups) in the space of diffeomorphisms of a 4-manifold. Given these developments, one may ask the following two general questions:

**Question 2.** *Can  $\gamma_1$  and its multiples be detected by gauge theoretic invariants?*

**Question 3.** *Can we use gauge theoretic invariants to attack Conjecture 1? (One can see Akbulut’s recent work [4] towards this conjecture.)*

Note that a negative answer to Question 2 would also be very interesting because it implies that even on  $S^4$ , there exist exotic phenomena that are invisible from gauge theory.

Since the family Bauer–Furuta invariant is a homotopy refinement of the family Seiberg–Witten invariant [9, 11], it is natural to specialize these questions to the the family Bauer–Furuta invariants. Our results (Theorem 1.6 and Theorem 1.7) provide partial negative answers to these questions.

**Theorem 1.6.** *Let  $f$  be an orientation-preserving diffeomorphism of  $S^4$ . Let  $\tilde{f}$  be a lift of  $f$  to an automorphism of the spin structure of  $S^4$ . Then we have*

$$\text{FBF}^{S^1}(S^4, f, \tilde{f}) = 0 \quad \text{and} \quad \text{FBF}^{\{e\}}(S^4, f, \tilde{f}) = 0.$$

**Theorem 1.7.** *Let  $\gamma$  be an exotic loop in  $\text{Diff}(S^4)$  then one has*

$$\text{FBF}^{S^1}(S^4, \gamma) = 0 \quad \text{and} \quad \text{FBF}^{\{e\}}(S^4, \gamma) = 0.$$

*If we further assume that  $\gamma$  is divisible by 2 in  $\pi_1(\text{Diff}(S^4))$  (e.g.,  $\gamma = 2k\gamma_1$ ), then*

$$\text{FBF}^{\text{Pin}(2)}(S^4, \gamma) = 0.$$

Theorem 1.6 and Theorem 1.7 together imply that the  $S^1$ -equivariant version and the non-equivariant version of the family Bauer–Furuta invariant do not detect exotic (loops of) diffeomorphisms on  $S^4$ . In particular, Theorem 1.6 rules out the possibility that the non-equivariant Bauer–Furuta invariant of a diffeomorphism (which is an element in  $\pi_1^s(S^0)$ ) equals the Hopf map. And Theorem 1.7 rules out the possibility that the non-equivariant Bauer–Furuta invariant of an exotic loop of diffeomorphisms (which is an element in  $\pi_2^s(S^0)$ ) equals the square of the Hopf map.

Note that Theorem 1.6 has a generalization (Theorem 1.9) that applies to a large family of spin 4-manifolds. However, we do not have a vanishing result for the  $\text{Pin}(2)$ -equivariant family Bauer–Furuta invariants that applies to all diffeomorphisms on  $S^4$ . Actually, we discovered a curious 2-torsion element in the  $\text{Pin}(2)$ -equivariant stable homotopy group of spheres. This element satisfies a basic requirement, called “*BF-type*”, of being realizable as  $\text{FBF}^{\text{Pin}(2)}(S^4, f, \tilde{f})$  for some exotic diffeomorphism  $f$ . Roughly speaking, this condition means that the equivariant homotopy class is trivial when restricted to the  $S^1$ -fixed points. We refer to Definition 5.1 for a precise definition.

**Theorem 1.8.** *There is an unique nonzero element  $\alpha_0 \in \pi_{1, \mathcal{M}}^{\text{Pin}(2)}(S^0)$  that is of BF-type.*

Theorem 1.8 suggests that the  $\text{Pin}(2)$ -equivariant family Bauer–Furuta invariant could potentially be useful to detect a counter example of Conjecture 1. It would be extremely interesting to calculate the  $\text{Pin}(2)$ -equivariant family Bauer–Furuta invariant of existing potential counter examples [14, 21]. Furthermore, Theorem 1.6 and Theorem 1.8 together provide strong evidence indicating that the  $j$ -symmetry is an indispensable ingredient if Seiberg–Witten theory can be used to attack Conjecture 1. For this reason, it is natural to try to detect this element  $\alpha_0$  by studying the induced map on the  $G$ -equivariant Borel (co)homology for various subgroups  $G$  of  $\text{Pin}(2)$ , because they correspond to various versions of monopole Floer homology. However, we will give an explicit space level description of  $\alpha_0$  in Section 5 and prove that this method does not detect  $\alpha_0$ .

**1.3. The vanishing theorem.** Now we state the vanishing theorem for the family Bauer–Furuta invariant of a diffeomorphism. As we mentioned, this theorem generalizes Theorem 1.6 and it plays an essential role in the proof of Theorem 1.1.

**Theorem 1.9.** *Let  $X$  be an oriented spin 4-manifold with  $b_1(X) = 0$ . We fix a homology orientation and a spin structure on  $X$ . Let  $f : X \rightarrow X$  be a diffeomorphism which preserves the spin structure and acts trivially on  $H^*(X; \mathbb{R})$ . Then for any lift  $\tilde{f}$  of  $f$  to an automorphism on the spin structure, we have the following results:*

- (1) *Suppose  $2\chi(X) + 3\sigma(X) = 4$ . Then the invariants*

$$\text{FBF}^{S^1}(X, f, \tilde{f}) \quad \text{and} \quad \text{FBF}^{\{e\}}(X, f, \tilde{f})$$

*are both vanishing.*

- (2) *Suppose  $2\chi(X) + 3\sigma(X) > 4$ . Then the invariants*

$$\text{FBF}^{\text{Pin}(2)}(X, f, \tilde{f}), \quad \text{FBF}^{S^1}(X, f, \tilde{f}) \quad \text{and} \quad \text{FBF}^{\{e\}}(X, f, \tilde{f})$$

*are all vanishing.*

We note that for any spin 4-manifold, we have  $4 \mid 2\chi(X) + 3\sigma(X)$ . We also note that the quantity  $2\chi(X) + 3\sigma(X)$ , which equals  $c_1(TX)^2[X]$  when  $X$  is equipped with an almost complex structure, plays an important role in the geography problem of irreducible and symplectic 4-manifolds (see, for example, [17]). Actually, all known examples of simply-connected, irreducible 4-manifolds satisfies the inequality

$$(1) \quad 2\chi(X) + 3\sigma(X) \geq 0.$$

Moreover, by a result of Taubes [49], the inequality (1) holds for any irreducible symplectic 4-manifolds. In particular, this implies that Theorem 1.9 applies to any simply-connected, spin, symplectic 4-manifold that is not homeomorphic to an elliptic surface  $E(2k)$ . And it also applies to many reducible 4-manifolds (e.g.  $K3\#(S^2 \times S^2)$ ). We refers to Remark 5.4 for examples where  $2\chi(X) + 3\sigma(X) = -4, 0$  or  $4$  but  $\text{FBF}^{\text{Pin}(2)}(X, f, \tilde{f}) \neq 0$ .

Theorem 1.9 is proved using tools from equivariant stable homotopy theory (in particular, the tom Dieck splitting theorem and the Adams' isomorphism). The geometric input from gauge theory is the following observation: When restricted to the  $S^1$ -fixed points, the family Bauer–Furuta invariant can be defined via the linear operator  $d^+ \oplus d^*$ , which only records the induced map on  $H^2(X; \mathbb{R})$  and hence can not distinguish  $f$  with the identity map on  $X$ . As a result,

the family Bauer–Furuta invariant becomes trivial once we restrict it to the  $S^1$ -fixed points. In a certain range of dimensions, this is enough to show that the family Bauer–Furuta invariant itself is trivial even before passing to the  $S^1$ -fixed points. So we can prove Theorem 1.9.

As a final remark, we mention that Theorem 1.9 doesn't say anything about those  $\text{spin}^c$  structures not coming from a spin structure. Actually, the family Bauer–Furuta invariants and the family Seiberg–Witten invariants for these  $\text{spin}^c$  structures do detect exotic diffeomorphisms (see [10, 40]). Note that when defining the family Bauer–Furuta invariant in the absence of a spin structure, there are some extra complications in choosing the framing of the moduli space (see [31, Section 3]).

**Organization of the paper:** In Section 2, we recall the definition of the  $\text{Pin}(2)$ -equivariant family Bauer–Furuta invariant. Then in Section 3, we prove Theorem 1.1 (existence of exotic surfaces under a stabilization) by assuming that Theorem 1.9 (the vanishing theorem for the family Bauer–Furuta invariant) holds for  $S^2 \times S^2$ . We also prove Proposition 1.4 in this section. The rest of the paper is devoted to the proof of the vanishing results Theorem 1.7, Theorem 1.8 and Theorem 1.9. In Section 4, we first review some standard constructions and results in equivariant stable homotopy theory. Then we prove several algebraic vanishing results for the  $\text{Pin}(2)$ -equivariant stable homotopy groups (Proposition 4.8 ~ Proposition 4.11). These vanishing results will be used in Section 5 to finish the proof of Theorem 1.7, Theorem 1.8 and Theorem 1.9. Section 4 and 5 mainly consist of algebraic topology.

**Acknowledgments:** The authors are extremely thankful to John Etnyre and Danny Ruberman for having conversations & email exchanges and providing some key insights related to this work. The first author wishes to thank Mike Hill and XiaoLin Danny Shi for pointing out the references [12, 33]. The second author is especially grateful to Nobuo Iida, Hokuto Konno and Masaki Taniguchi for their priceless friendship and being patience while teaching him Bauer–Furuta theory and having many discussions related to this work. We are thankful to Kyle Hayden, Hyunki Min, Lisa Piccirillo and Ian Zemke for showing great interest in this project. We are also thankful to Selman Akbulut, Sander Kupers and Markus Szymik for giving helpful comments on an earlier version of the paper. The second author is partially supported by the NSF grant DMS-1906414.

## 2. THE FAMILY BAUER–FURUTA INVARIANT.

Our proof of Theorem 1.1 heavily uses the family Bauer–Furuta invariant, which was defined by Bauer–Furuta [11] (for a single 4-manifold), Szymik [47] and Xu [53] (for families). So for completeness, we first recall its definition following the expositions in [31, 35]. We start by introducing some notations. Consider the 1-dimensional Lie group

$$\text{Pin}(2) = \{e^{i\theta}\} \cup \{j \cdot e^{i\theta}\} \subset \mathbb{H}.$$

We are mainly interested in the following three representations of  $\text{Pin}(2)$ :

- $\mathbb{H}$ : the 4-dimensional representation, where  $\text{Pin}(2)$  acts as the left multiplication.
- $\widetilde{\mathbb{R}}$ : the 1-dimensional representation, where the unique component  $S^1 \subset \text{Pin}(2)$  acts trivially and the other component acts as multiplication by  $-1$ . Note that this is also the adjoint representation of  $\text{Pin}(2)$  on its Lie algebra.
- $\mathbb{R}$ : the 1-dimensional trivial representation.

For a finite dimensional  $\text{Pin}(2)$ -representation  $V$ , we use  $S^V$  to denote the one point compactification of  $V$  and use  $S(V)$  to denote the unit sphere in  $V$ . The space  $S^V$  is called the “representation sphere” and it admits a  $\text{Pin}(2)$ -action that fixes the point  $\infty$ , which we choose as the base point.

**2.1. The invariant of a single diffeomorphism.** Now we start the construction of the family Bauer–Furuta invariant of a self-diffeomorphism. Let  $X$  be an oriented 4-manifold with  $b_1(X) = 0$ . We equip  $X$  with a cohomological orientation (i.e. an orientation of  $H_+^2(X; \mathbb{R})$ ) and a spin structure. We use  $P$  to denote the corresponding  $\text{Spin}(4)$ -bundle. Let  $f$  be a self-diffeomorphism on  $X$  such that  $f^*$  acts trivially on  $H^*(X; \mathbb{R})$ . Then we form the mapping torus

$$Tf = ([0, 1] \times X)/(0, x) \sim (1, f(x))$$

and treat it as a fiber bundle over  $S^1$  with fiber  $X$ . Next, let  $\tilde{f} : P \rightarrow P$  be a bundle automorphism of  $P$  which lifts  $f$ . (There are two choices of  $\tilde{f}$  and we pick one.) Then we form the  $\text{Spin}(4)$ -bundle

$$([0, 1] \times P)/(0, x) \sim (1, \tilde{f}(x)),$$

which gives a family spin structure on  $Tf$ . This family spin structure allows us to define the family Seiberg–Witten equations on  $Tf$ . By doing the *finite dimensional approximation* on these equations, we obtain a  $\text{Pin}(2)$ -equivariant map

$$(2) \quad \widetilde{sw} : U \rightarrow V$$

, where  $V$  is finite dimensional representation space of  $\text{Pin}(2)$  and  $U$  is a fiber bundle over  $S^1$  whose fibers are finite dimensional representations spaces of  $\text{Pin}(2)$ . Moreover, because of the compactness property of the Seiberg–Witten moduli space, by choosing  $U, V$  large enough, one can ensure that  $\widetilde{sw}$  satisfies the following nice property: There exists a large  $R$  and a small  $\epsilon$ , such that

$$\widetilde{sw}(S_R(U)) \cap B_\epsilon(V) = \emptyset.$$

Here  $S_R(U) \subset U$  denotes the sphere bundle with radius  $R$  and  $B_\epsilon(V) \subset V$  denotes the disk with radius  $\epsilon$ . As a consequence, we get an induced map

$$\widetilde{sw}^+ : U^+ \cong B_R(U)/S_R(U) \rightarrow V/(V \setminus \text{int}(B_\epsilon(V))) \cong V^+$$

between one-point compactifications. Furthermore, as explained in [31, 35], by using the homology orientation on  $X$  and the  $\text{Pin}(2)$ -actions on  $U, V$ , one can choose canonical (up to homotopy) identifications

$$(3) \quad U \cong S^1 \times (\widetilde{\mathbb{R}}^{n'} \oplus \mathbb{H}^{m' - \frac{\sigma(X)}{16}}), \quad V \cong \widetilde{\mathbb{R}}^{n'+b_2^+(X)} \oplus \mathbb{H}^{m'}$$

for some  $m', n' \gg 0$ . Using these trivialisations, we can rewrite  $\widetilde{sw}^+$  as a map

$$(4) \quad \widetilde{sw}^+ : S_+^1 \wedge S^{(m' - \frac{\sigma(X)}{16})\mathbb{H} + n'\widetilde{\mathbb{R}}} \rightarrow S^{m'\mathbb{H} + (n'+b_2^+(X))\widetilde{\mathbb{R}}}.$$

Here  $S_+^1$  denotes union of  $S^1$  with a disjoint base point. To obtain a map between two spheres, we embed  $S^1$  as the unit sphere  $S(2\mathbb{R})$  in  $\mathbb{R}^2$  and let  $c_0 : S^{2\mathbb{R}} \rightarrow S^{\mathbb{R}} \wedge S(2\mathbb{R})_+$  be the Pontragin–Thom collapsing map which collapses all the points outside a tubular neighborhood of  $S(2\mathbb{R})$ .

Consider the composition

$$(5) \quad S^{2\mathbb{R}} \wedge S^{(m' - \frac{\sigma(X)}{16})\mathbb{H} + n'\tilde{\mathbb{R}}} \xrightarrow{c_0 \wedge \text{id}_{S^{(m' - \frac{\sigma(X)}{16})\mathbb{H} + n'\tilde{\mathbb{R}}}}} S^{\mathbb{R}} \wedge (S(2\mathbb{R})_+) \wedge S^{(m' - \frac{\sigma(X)}{16})\mathbb{H} + n'\tilde{\mathbb{R}}} \\ \xrightarrow{\text{id}_{S^{\mathbb{R}}} \wedge \tilde{s}w^+} S^{\mathbb{R}} \wedge S^{m'\mathbb{H} + (n' + b_2^+(X))\tilde{\mathbb{R}}}.$$

It is a  $\text{Pin}(2)$ -equivariant map between two spheres and thus it defines an element in the  $\text{Pin}(2)$ -equivariant stable homotopy of spheres. This element is exactly the  $\text{Pin}(2)$ -equivariant family Bauer–Furuta invariant of  $f$ , denoted by  $\text{FBF}^{\text{Pin}(2)}(X, f, \tilde{f})$ . More precisely, we have

$$\text{FBF}^{\text{Pin}(2)}(X, f, \tilde{f}) \in \begin{cases} \{S^{\mathbb{R} + (-\frac{\sigma(X)}{16})\mathbb{H}}, S^{b^+(X)\tilde{\mathbb{R}}}\}_{\mathcal{U}}^{\text{Pin}(2)} & \text{if } \sigma(X) \leq 0 \\ \{S^{\mathbb{R}}, S^{b^+(X)\tilde{\mathbb{R}} + \frac{\sigma(X)}{16}\mathbb{H}}\}_{\mathcal{U}}^{\text{Pin}(2)} & \text{if } \sigma(X) > 0 \end{cases}.$$

(See Definition 4.1 for a precise description of these groups.)

The  $S^1$ -equivariant Bauer–Furuta invariant  $\text{FBF}^{S^1}(X, f, \tilde{f})$  and the non-equivariant Bauer–Furuta invariant  $\text{FBF}^{\{e\}}(X, f, \tilde{f})$  are defined to be the restriction of  $\text{FBF}^{\text{Pin}(2)}(X, f, \tilde{f})$  to the subgroups  $S^1, \{e\} \subset \text{Pin}(2)$  respectively. Therefore, we have

$$\text{FBF}^{\text{Pin}(2)}(X, f, \tilde{f}) = 0 \implies \text{FBF}^{S^1}(X, f, \tilde{f}) = 0 \implies \text{FBF}^{\{e\}}(X, f, \tilde{f}) = 0.$$

All these invariants are unchanged under smooth isotopy. In particular, when  $f$  is smoothly isotopic to  $\text{id}_X$  and  $\tilde{f}$  is the lift that corresponding to the identity map on the spin bundle, all of them are zero. To get an invariant of  $f$ , we can consider the unordered pair

$$\{\text{FBF}^{\text{Pin}(2)}(X, f, \tilde{f}_1), \text{FBF}^{\text{Pin}(2)}(X, f, \tilde{f}_2)\}$$

for two possible lifts  $\tilde{f}_1, \tilde{f}_2$ . If  $f$  is smoothly isotopic to  $\text{id}_X$ , then at least one element in this pair must be vanishing.

*Remark 2.1.* We briefly explain the motivation of the composition (5). Since  $S^1_+$  is stably homotopy equivalent to  $S^1 \vee S^0$ , the map  $\tilde{s}w^+$  admits a splitting into  $\tilde{s}w^+_0 \vee \tilde{s}w^+_1$  in the stable homotopy category. The component  $\tilde{s}w^+_0$  represent the Bauer–Furuta invariant of the fiber  $X$  and the component  $\tilde{s}w^+_1$  captures information about  $f$ . Since we are studying exotic diffeomorphisms on a fixed smooth manifold  $X$ , our main interest is  $\tilde{s}w^+_1$ . The effect of composing with a suspension of  $c_0$  in (5) is exactly keeping  $\tilde{s}w^+_1$  and ignoring  $\tilde{s}w^+_0$ . This also explains why the family Bauer–Furuta invariant can be trivial even when the Bauer–Furuta invariant of the fiber is nontrivial.

**2.2. The invariant of a loop of diffeomorphisms.** Let  $X$  be as before. Now we construct the family Bauer–Furuta invariant of a loop of diffeomorphisms on  $X$ . Let  $\gamma : S^1 \rightarrow \text{Diff}(X)$  be a loop based at  $\text{id}_X$ . We use  $\gamma$  as a clutching function and form a smooth fiber bundle

$$X \hookrightarrow T\gamma \rightarrow S^2.$$

**Definition 2.2.**  $\gamma$  is *spin-liftable* if the second Stiefel–Whitney class  $w_2(T\gamma)$  vanishes.

Note that this condition is always satisfied if  $\gamma$  is null-homotopic in the homeomorphism group of  $X$ . Because in this case,  $T\gamma$  is homeomorphic to  $X \times S^2$ . By Wu’s formula, the Seifert–Whitney class is independent of the smooth structure, we have  $w_2(T\gamma) = w_2(X \times S^2) = 0$ .

From now on, we assume  $\gamma$  is spin-liftable. Then  $T\gamma$  carries a family spin structure. Since the inclusion  $X \hookrightarrow T\gamma$  induces an isomorphism on  $H^1(-; \mathbb{Z}/2)$ , such a family spin structure is



unique if we require that its restriction the fiber is the given spin structure on  $X$ . (This is different from the previous case of a single diffeomorphism.)

Now we repeat the same construction as before and get a map

$$(6) \quad \widetilde{sw}^+ : S_+^2 \wedge S^{(m' - \frac{\sigma(X)}{16})\mathbb{H} + n'\widetilde{\mathbb{R}}} \rightarrow S^{m'\mathbb{H} + (n' + b_2^+(X))\widetilde{\mathbb{R}}}.$$

for some  $m', n' \gg 0$ . We embed  $S^2$  as the unit sphere in  $S^{3\mathbb{R}}$  and form the composition between (suspensions of)  $\widetilde{sw}^+$  and the Pontryagin-Thom collapsing map  $S^{3\mathbb{R}} \rightarrow S^{\mathbb{R}} \wedge S(3\mathbb{R})_+$ . This gives us a map

$$S^{3\mathbb{R} + (m' - \frac{\sigma(X)}{16})\mathbb{H} + n'\widetilde{\mathbb{R}}} \rightarrow S^{\mathbb{R} + m'\mathbb{H} + (n' + b_2^+(X))\widetilde{\mathbb{R}}}.$$

The  $\text{Pin}(2)$ -equivariant family Bauer–Furuta invariant  $\text{FBF}^{\text{Pin}(2)}(X, \gamma)$  is defined as the equivariant stable homotopy class of this map. More precisely, we have

$$\text{FBF}^{\text{Pin}(2)}(X, \gamma) \in \begin{cases} \{S^{2\mathbb{R} + (-\frac{\sigma(X)}{16})\mathbb{H}}, S^{b^+(X)\widetilde{\mathbb{R}}}\}_{\mathcal{U}}^{\text{Pin}(2)} & \text{if } \sigma(X) \leq 0 \\ \{S^{2\mathbb{R}}, S^{b^+(X)\widetilde{\mathbb{R}} + \frac{\sigma(X)}{16}\mathbb{H}}\}_{\mathcal{U}}^{\text{Pin}(2)} & \text{if } \sigma(X) > 0 \end{cases}.$$

The invariants  $\text{FBF}^{S^1}(X, \gamma)$  and  $\text{FBF}^{\{e\}}(X, \gamma)$  are defined by restricting  $\text{FBF}^{\text{Pin}(2)}(X, \gamma)$  to the subgroups  $\{e\}, S^1$  respectively. Again, these invariants are unchanged under any smooth isotopies.

*Remark 2.3.* As noted in [31], there are extra complications if one tries to generalize this construction and define invariants of elements in  $\pi_k(\text{Diff}(X))$  with  $k \geq 2$ . Because the symplectic group  $\text{Sp}(n)$  is 2-connected, every quaternion bundle over  $S^{k+1}$  has a trivialization which is unique up to homotopy when  $k \leq 1$ . This allows us to define the canonical trivialisations of the bundles involved in construction (see (3)). However, this argument breaks down when  $k \geq 2$ .

### 3. EXOTIC SURFACES AND STABILIZATION

**3.1. Exotic surfaces that survive a stabilization.** We will now prove Theorem 1.1, which shows the existence of exotic surfaces that remains exotic after a single stabilization. The proof here relies on Theorem 1.9 in the case of  $S^2 \times S^2$ . Theorem 1.9 will be proved in Section 5 in full generality.

We start with the following technical lemma.

**Lemma 3.1.** *Let  $D = \bigsqcup D^2 \hookrightarrow X$  be a smooth, proper embedding of a finite disjoint union of disks into a 4-manifold  $X$  with boundary. Consider a self-diffeomorphism  $f : X \rightarrow X$  that fixes all points in a neighborhood of  $\partial X$  and all points on  $D$ . Then  $f$  is smoothly isotopic (relative to  $\partial X$ ) to some  $f'$  which fixes all points in a neighborhood of  $(\partial X) \cup D$ .*

*Proof.* Let  $N$  be the normal bundle of  $D$  in  $X$ . By choosing a trivialization of this bundle, we get an identification  $N = D \times \mathbb{R}^2$ . Since  $f$  fixes  $D$  pointwisely, its differential induces a bundle automorphism  $l' : N \rightarrow N$ . Let  $\rho : N \rightarrow \nu(D)$  be a diffeomorphism to a tubular neighborhood of  $D$ . We use  $l$  to denote the composition

$$\rho \circ l' \circ \rho^{-1} : \nu(D) \rightarrow \nu(D).$$

For  $\epsilon > 0$ , we use  $\nu_\epsilon(D)$  to denote  $\rho(D \times D_\epsilon)$ , where  $D_\epsilon \subset \mathbb{R}^2$  is the ball centered at the origin with radius  $\epsilon$ . We pick  $\epsilon$  small enough such that  $f(\nu_\epsilon(D)) \subset \nu(D)$ . We first construct a smooth isotopy  $h_1 : [0, 1] \times \nu_\epsilon(D) \rightarrow \nu(D)$  between the maps

$$f|_{\nu_\epsilon(D)} : \nu_\epsilon(D) \rightarrow \nu(D) \quad \text{and} \quad l|_{\nu_\epsilon(D)} : \nu_\epsilon(D) \rightarrow \nu(D).$$

(Isotopy here means that for any  $t$ , the map  $h_1(t, -)$  is a diffeomorphism onto its image.) This is done via the standard rescaling trick: For any  $t > 0$ , we use  $\phi_t : \nu(D) \rightarrow \nu(D)$  to denote the self-diffeomorphism defined by

$$\phi_t(\rho(x, \vec{v})) = \rho(x, t\vec{v}) \quad \forall x \in D, \vec{v} \in \mathbb{R}^2.$$

Then the isotopy  $h_1$  is given by

$$h_1(t, y) := \begin{cases} \phi_t^{-1} \circ f \circ \phi_t(t, y) & \text{if } t \in (0, 1] \\ l(y) & \text{if } t = 0 \end{cases}.$$

Next, we construct an smooth isotopy  $h_2 : [0, 1] \times \nu_\epsilon(D) \rightarrow \nu(D)$  from the map  $l|_{\nu_\epsilon(D)}$  to the inclusion map  $i : \nu_\epsilon(D) \rightarrow \nu(D)$ . This is done via a simple obstruction argument: The map  $l'$  is linear on each fiber so it has the form

$$l'(x, \vec{v}) = (x, \tau(x) \cdot \vec{v})$$

for some  $\tau : D \rightarrow \text{GL}^+(\mathbb{R}^2)$  that equals 1 near  $\partial D$ . Since  $\text{GL}^+(\mathbb{R}^2)$  is homotopy equivalent to  $S^1$  and  $\pi_2(S^1) = 0$ , there exists a smooth homotopy (relative to  $\partial D$ )

$$\tilde{\tau} : [0, 1] \times D \rightarrow \text{GL}^+(\mathbb{R}^2)$$

from  $\tau$  to the constant map 1. we define  $\tilde{l}' : [0, 1] \times N \rightarrow N$  by

$$\tilde{l}'(t, x, \vec{v}) = (x, \tilde{\tau}(t, x) \cdot \vec{v}) \quad \text{for } t \in [0, 1], x \in D, \vec{v} \in \mathbb{R}^2.$$

Then we define  $h_2(t, y) := \rho(\tilde{l}'(t, \rho^{-1}(y)))$ . By compositing  $h_1$  with  $h_2$ , we obtain a smooth isotopy between the maps

$$f|_{\nu_\epsilon(D)} : \nu_\epsilon(D) \rightarrow \nu(D) \quad \text{and} \quad i : \nu_\epsilon(D) \rightarrow \nu(D).$$

Furthermore, this isotopy is trivial near  $\partial X$ . Therefore, we can apply the isotopy extension theorem to produce an isotopy of  $f$  to some  $f'$  that fixed all points in a neighborhood of  $(\partial X) \cup D$ .  $\square$

Next, we briefly recall the construction of the Dehn twist on the punctured  $K3$ . We pick an essential loop

$$(7) \quad \gamma : [0, 1] \rightarrow SO(4)$$

such that  $\gamma(t) = 1$  for  $t$  near 0, 1. Then we define the self-diffeomorphism

$$\tilde{\gamma} : [0, 1] \times S^3 \rightarrow [0, 1] \times S^3$$

by  $\tilde{\gamma}(t, x) = (t, \gamma(t)x)$ . Let  $K3^\circ$  be the punctured  $K3$ . We fix a diffeomorphism from  $[0, 1] \times S^3$  to a collar neighborhood of  $K3^\circ$  and define the Dehn twist as follows

$$(8) \quad \delta_{K3^\circ} := \tilde{\gamma} \cup \text{id}_{K3^\circ \setminus ([0, 1] \times S^3)} : K3^\circ \rightarrow K3^\circ.$$

Now we are ready to show the existence of exotic surfaces that survive even after a single stabilization.

*Proof of Theorem 1.1.* Recall that one can construct  $K3^\circ$  by starting with  $S^3 \times I$  and then adding a bunch of 2-handles along  $S^3 \times \{1\}$  and cap it off with a 4-ball. Let  $L$  be the set of attaching spheres on  $S^3$ . Then  $L$  is a link with 22 components. Let  $D_L$  be the union

$$(L \times I) \cup_{L \times \{1\}} \{\text{cores of the two handles}\}.$$

Then  $D_L$  is a slice disk bounded by  $L \subset \partial(K3^\circ)$ .

Consider the Dehn-twist  $\delta_{K3^\circ}$  defined in (8). In [31], Kronheimer–Mrowka showed that  $\delta_{K3^\circ}$  is not smoothly isotopic to  $\text{id}_{K3^\circ}$  relative to the boundary. Let  $D'_L = \delta_{K3^\circ}(D_L)$ , then  $D'_L$  and  $D_L$  are topologically isotopic relative to the boundary, since the result of Krannich–Kupers [30, Remark 6] and Orson–Powell [37] implies  $\delta_{K3^\circ}$  is indeed topologically isotopic to  $\text{id}_{K3^\circ}$  relative to the boundary. Also it is obvious now that their complements are diffeomorphic.

The claim is that  $D_L$  and  $D'_L$  are not smoothly isotopic even after a single stabilization. Suppose this were not true. Denote  $\delta \# \text{id}_{S^2 \times S^2}$  by  $\delta_{K3^\circ \# (S^2 \times S^2)}$ . By the isotopy extension theorem, there exists a self-diffeomorphism  $g_1 : K3^\circ \# (S^2 \times S^2) \rightarrow K3^\circ \# (S^2 \times S^2)$  that is smoothly isotopic to the identity relative to the boundary and sends  $D_L$  to  $D'_L$ . Then  $g_1 \circ \delta_{K3^\circ \# (S^2 \times S^2)}$  fixes all points on  $D_L$  and is smoothly isotopic to  $\delta_{K3^\circ \# (S^2 \times S^2)}$  relative to the boundary. By Lemma 3.1, we can further isotope  $g_1 \circ \delta_{K3^\circ \# (S^2 \times S^2)}$  relative to the boundary to a diffeomorphism

$$g_2 : K3^\circ \# (S^2 \times S^2) \rightarrow K3^\circ \# (S^2 \times S^2)$$

that is identity on the neighbourhood of  $D_L \cup S^3 \times \{0\}$ . Notice that the complement of this set in  $K3^\circ \# (S^2 \times S^2)$  is a punctured  $S^2 \times S^2$ . Thus  $g_2$  can be decompose as  $\text{id}_{K3^\circ} \# f$  for some  $f \in \text{Diff}(S^2 \times S^2)$  which acts trivially on the cohomology. Furthermore, let  $B$  be the small disk in  $S^2 \times S^2$  that we apply the connected sum construction. Then  $f$  fixes  $B$  pointwisely.

So we have shown that  $\delta_{K3^\circ \# S^2 \times S^2}$  is smoothly isotopic to  $\text{id}_{K3^\circ} \# f$ . This further implies that the Dehn twist

$$\delta_{K3 \# K3 \# (S^2 \times S^2)} : K3 \# K3 \# (S^2 \times S^2) \rightarrow K3 \# K3 \# (S^2 \times S^2)$$

along the neck between the two  $K3$ 's is smoothly isotopic to  $(\text{id}_{K3 \# K3}) \# f$ .

Let  $\text{id}_{S_{K3 \# K3}}$  be the identity map on the spin bundle over  $K3 \# K3$ . There are two lifts of  $f$  to the spin bundle over  $S^2 \times S^2$ . We set  $\tilde{f}$  to be the one that is identity on all the fibers over  $B$ . Then we can form the connected sum  $\text{id}_{S_{K3 \# K3}} \# \tilde{f}$ , which is a lift of  $\text{id}_{K3 \# K3} \# f$  to an automorphism on the spin bundle.

It follows the connected sum formula [35, Proposition 2.26] that

$$\begin{aligned} & \text{FBF}^{\text{Pin}(2)}(K3 \# K3 \# S^2 \times S^2, \text{id}_{K3 \# K3} \# f, \text{id}_{S_{K3 \# K3}} \# \tilde{f}) \\ &= \text{BF}^{\text{Pin}(2)}(K3 \# K3) \wedge \text{FBF}^{\text{Pin}(2)}(S^2 \times S^2, f, \tilde{f}). \end{aligned}$$

But Theorem 1.9 implies that  $\text{FBF}^{\text{Pin}(2)}(S^2 \times S^2, f, \tilde{f}) = 0$ , and hence

$$\text{FBF}^{\text{Pin}(2)}(K3 \# K3 \# S^2 \times S^2, \text{id}_{K3 \# K3} \# f, \text{id}_{S_{K3 \# K3}} \# \tilde{f}) = 0.$$

But in [35, Proof of Theorem 1.2], the first author proves that

$$\text{FBF}^{\text{Pin}(2)}(K3 \# K3 \# S^2 \times S^2, \delta_{K3 \# K3 \# (S^2 \times S^2)}, \tilde{\delta}) \neq 0$$

for any lift  $\tilde{\delta}$  of  $\delta_{K3\#K3\#(S^2 \times S^2)}$  to the spin structure of  $K3\#K3\#S^2 \times S^2$ .<sup>1</sup> Since  $\delta_{K3\#K3\#(S^2 \times S^2)}$  is smoothly isotopic to  $\text{id}_{K3\#K3\#f}$ , this contracts the isotopy invariance of the family Bauer–Furuta invariant. Thus  $D_L$  and  $D'_L$  are not smoothly isotopic relative to the boundary in  $K3^\circ\#S^2 \times S^2$ .  $\square$

**3.2. An upper bound on the number of stabilizations.** Now we move on to the proof of Proposition 1.4, which gives an upper bound on the stabilizations needed to make  $D_L$  and  $D'_L$  smoothly isotopic. We start with part (1).

*Proof of Proposition 1.4 (1).* Since  $L_0$  is an unlink, there is a diffeomorphism  $f : S^3 \rightarrow S^3$  that is smoothly isotopic to  $\text{id}_{S^3}$  and sends  $L_0$  to the standard unlink

$$L'_0 = \{(\cos \theta, \sin \theta, i) \mid \theta \in [0, 2\pi], i = 1, \dots, m\} \subset \mathbb{R}^3 \cup \{\infty\} = S^3.$$

By taking the rotations along the  $z$ -axis, we can construct an essential loop  $\gamma' : [0, 1] \rightarrow SO(4)$  such that  $\gamma'(t)$  fixes  $L'_0$  as a subset for any  $t$ . Then we consider the self-diffeomorphism

$$\tilde{\gamma}' : [0, 1] \times S^3 \rightarrow [0, 1] \times S^3$$

defined by

$$\tilde{\gamma}'(t, x) := (t, f^{-1} \circ \gamma'(t) \circ f(x)).$$

By extending  $\tilde{\gamma}'$  with the identity map on  $K3^\circ \setminus ([0, 1] \times S^3)$ , we obtain a self-diffeomorphism  $\delta'_{K3^\circ}$  on  $K3^\circ$ . Since  $\gamma'$  is homotopic to the loop  $\gamma$  in (7), and  $f$  is smoothly isotopic to  $\text{id}_{S^3}$ , we see that  $\delta'_{K3^\circ}$  is smoothly isotopic to  $\delta_{K3^\circ}$  relative to the boundary. Therefore,  $\delta'_{K3^\circ}(D_{L_0})$  and  $D'_{L_0}$  are smoothly isotopic relative to the boundary. Furthermore, since  $f^{-1} \circ \gamma'(t) \circ f$  fixes  $L_0$  as a subset, the embedded surfaces  $\delta'_{K3^\circ}(D_{L_0})$  and  $D_{L_0}$  differ from each other by a Dehn twist on the domain  $\bigsqcup_m D^2$ . Since this Dehn twist is smoothly isotopic to the identity map relative to the boundary, the surfaces  $D_{L_0}$  and  $\delta'_{K3^\circ}(D_{L_0})$  are also smoothly isotopic relative to the boundary.  $\square$

Figure 1 is a Kirby diagram of  $K3$  [22, Page 305]. The attaching link has 21 unknotted components and a single knotted component (which is a trefoil). By doing a handle slide of this knotted component along one of the unknotted components, we can change the middle crossing and make it into the unknot. This gives an example of  $L$  whose all components are unknots. By Proposition 1.4 (1), all components of  $D_L$  are smoothly isotopic to the corresponding components of  $D'_L$  relative to the boundary.

Now we move on to the proof of Proposition 1.4 (2). We start by the following lemma.

**Lemma 3.2.** *Given a sublink  $L'$  of  $L$ , let  $\nu(D_{L'})$  be an open tubular neighborhood of  $D_{L'}$  in  $K3^\circ$ . Then we have the following results.*

- (i)  $K3^\circ \setminus \nu(D_{L'})$  and  $K3^\circ \setminus \nu(D'_{L'})$  are simply connected.
- (ii) For any  $n \geq 0$ , the inclusion  $(K3^\circ \setminus \nu(D_{L'}))\#n(S^2 \times S^2) \hookrightarrow K3^\circ\#n(S^2 \times S^2)$  induces an injection on the second homotopy group  $\pi_2(-)$ .

<sup>1</sup>There is a subtle difference between the setting of the current paper and [35]. Here we are using an incomplete universe  $\mathbb{H}^\infty \oplus \tilde{\mathbb{R}}^\infty \oplus \mathbb{R}^\infty$  to set up an equivariant stable homotopy category, while in [35], a complete universe is used. (See Section 4 for further explanations.) This difference does not cause a problem because by definition,  $\text{FBF}^{\text{Pin}(2)}(K3\#K3\#S^2 \times S^2, \delta_{K3\#K3\#(S^2 \times S^2)}, \tilde{\delta}) \neq 0$  in a complete universe implies the same result in any universe.




FIGURE 1. A Kirby Diagram of  $K3$ .

*Proof.* (i) We prove the case of  $D_{L'}$  and the other case is similar. For each component  $K$  of  $L'$ , let  $m_k \subset \partial(K3^\circ \setminus \nu(D_{L'}))$  be the meridian of the embedded disk  $D_K$ . Since  $K3^\circ$  is simply-connected, we see that  $\pi_1(K3^\circ \setminus \nu(D_{L'}))$  is normally generated by  $\{m_K\}$ . So it suffices to show that  $m_K$  is null-homotopic in  $K3^\circ \setminus \nu(D_{L'})$ . To prove this, we consider the cocore of the 2-handle attached to  $K$ . This cocore becomes a cylinder in  $K3^\circ \setminus \nu(D_{L'})$  and it provides a homotopy between  $m_K$  and a knot on the boundary of the 4-handle  $D^4$  (which doesn't intersect  $\nu(D_{L'})$ ). Since  $D^4$  is contractible, this knot (and hence  $m_K$ ) is null-homotopic.

(ii) By the Mayer-Vietoris sequence, the map

$$H_2((K3^\circ \setminus \nu(D_{L'})) \# n(S^2 \times S^2); \mathbb{Z}) \rightarrow H_2(K3^\circ \# n(S^2 \times S^2); \mathbb{Z})$$

induced by the inclusion is injective. By the Hurewicz theorem, this map is also the induced map on  $\pi_2(-)$ .  $\square$

The next proposition is a variation of the main theorem in [6].

**Proposition 3.3.** *Let  $X$  be a simply-connected, smooth 4-manifold with boundary. Let  $D_K$  and  $D'_K$  be two smoothly embedded disks bounded by the same knot  $K \subset \partial X$ . Suppose the following conditions are all satisfied:*

- (i)  $X \setminus D_K$  and  $X \setminus D'_K$  are simply-connected.
- (ii)  $D_K$  and  $D'_K$  are homotopic relative to the boundary.
- (iii)  $[D_K] \in H_2(X, \partial X; \mathbb{Z}/2)$  is ordinary (i.e. it is not the Poincaré dual of  $w_2(X)$ ).

*Then in  $X \# (S^2 \times S^2)$ ,  $D_K$  and  $D'_K$  are smoothly isotopic relative to the boundary.*

*Proof.* The proof is adapted from the corresponding proof in [6]. For completeness, we sketch the argument, with an emphasize on the adaptations needed here. First, since  $X \setminus D_K$  is simply connected, we can find an immersed sphere  $\Sigma$  that intersects  $D_K$  transversely at a single point. Using condition (iii), one can modify  $\Sigma$  so that it has an even self-intersection number. Since  $D'_K$  is homotopic to  $D_K$  relative to the boundary, the algebraic intersection number  $\Sigma \cdot D'_K$  equals 1. After suitable finger moves and the (immersed) Whitney moves, we can isotope  $D'_K$  and  $\Sigma$  so that  $\Sigma$  intersects both  $D_K$  and  $D'_K$  transversely at a single point. To arrange for  $\Sigma$  to be embedded, we take a connected sum with  $S^2 \times S^2$  away from  $D_K \cup D'_K \cup \Sigma$ . Take coordinate 2-spheres  $S = S^2 \times \{\text{pt}\}$  and  $T = \{\text{pt}\} \times S^2$ . By replacing  $\Sigma$  with its connected sum with  $S$  along a tube, we may assume that  $\Sigma$  has a geometric dual  $T$  (i.e.,  $T$  intersects transverse  $\Sigma$  at a single point). Then we can eliminate the double points of  $\Sigma$  by the Norman trick, tubing to parallel copies of

$T$ . This gives us an embedded  $\Sigma$  with even self-intersection number. After tubing additional copies of  $T$ , we can arrange the self-intersection number of  $\Sigma$  to be zero. Therefore,  $D_K$  and  $D'_K$  have a common geometric dual  $\Sigma$  in  $X \# (S^2 \times S^2)$ . No essential changes are necessary until step. Then we apply the light bulb theorem for embedded disks [43, Corollary 3.3], which states that  $D_K$  and  $D'_K$  are smoothly isotopic relative to the boundary if the following three conditions are verified:

- (1)  $D_K$  and  $D'_K$  are homotopic in  $X \# (S^2 \times S^2)$  relative to the boundary.
- (2) The Dax invariant  $\text{Dax}(D_K, D'_K)$  vanishes.
- (3) The map  $\pi_1((X \# (S^2 \times S^2)) \setminus \Sigma) \rightarrow \pi_1(X \# (S^2 \times S^2))$  induced by the inclusion is an isomorphism. (I.e.  $(X \# (S^2 \times S^2)) \setminus \Sigma$  is simply connected.)

(1) follows from our assumption (ii). (2) is trivial since any surfaces in a simply connected 4-manifold have trivial Dax invariant. (3) is satisfied because  $\Sigma$  has an geometric dual  $T$  (so the meridian of  $\Sigma$  bounds a disk in  $X \setminus \Sigma$ ). This finishes the proof.  $\square$

*Proof of Proposition 1.4 (2).* Let  $L_0 \subset L$  be the  $m$ -component unlink. Denote the components of  $L \setminus L_0$  by  $K_1, K_2, \dots, K_{22-m}$ . For  $1 \leq n \leq 22 - m$ , we set

$$L_n = L_0 \bigsqcup K_1 \bigsqcup \dots \bigsqcup K_n \quad \text{for } 1 \leq n \leq 22 - m.$$

We will prove the following claim by induction.

**Claim 1.** *For any all  $0 \leq n \leq 22 - m$ ,  $D_{L_n}$  and  $D'_{L_n}$  are smoothly isotopic relative to the boundary in  $K3^\circ \# n(S^2 \times S^2)$ .*

The case  $n = 0$  is implied by Proposition 1.4 (1). Now suppose the case  $n = i$  has been proved. Denote  $K3^\circ \# i(S^2 \times S^2)$  by  $X_i$ . Then by our induction hypothesis, there exists a diffeomorphism  $f : X_i \rightarrow X_i$  which is smoothly isotopic to  $\text{id}_{X_i}$  relative to the boundary and satisfies  $f(D'_{L_i}) = D_{L_i}$ . Note that  $D_{K_{i+1}}$  and  $f(D'_{K_{i+1}})$  are disks embedded in  $X_i \setminus \nu(D_{L_i})$  bounded by the same knot  $K_{i+1}$ . To prove the case  $n = i + 1$ , it suffices to show that  $D_{K_{i+1}}$  and  $f(D'_{K_{i+1}})$  are smoothly isotopic relative to the boundary in  $(X_i \setminus \nu(D_{L_i})) \# (S^2 \times S^2)$ . (The connected sum is taken near  $\partial X_i$  so that the isotopy between  $f$  and  $\text{id}_{X_i}$  can be trivially extended.) By Proposition 3.3, we just need to check the following three conditions:

- (1)  $X_i \setminus \nu(D_{L_i})$ ,  $X_i \setminus \nu(D_{L_i} \bigsqcup D_{K_{i+1}})$  and  $X_i \setminus \nu(D_{L_i} \bigsqcup f(D'_{K_{i+1}}))$  are all simply connected.
- (2) The homology class  $[D_{K_{i+1}}] \in H_2(X_i \setminus \nu(D_{L_i}); \mathbb{Z}/2)$  is ordinary.
- (3)  $D_{K_{i+1}}$  and  $f(D'_{K_{i+1}})$  are homotopic in  $X_i \setminus \nu(D_{L_i})$  relative to the boundary.

Note that

$$\begin{aligned} X_i \setminus \nu(D_{L_i} \bigsqcup D_{K_{i+1}}) &\cong (K3^\circ \setminus \nu(D_{L_{i+1}})) \# i(S^2 \times S^2), \\ X_i \setminus \nu(D_{L_i} \bigsqcup f(D'_{K_{i+1}})) &\cong (K3^\circ \setminus \nu(D'_{L_{i+1}})) \# i(S^2 \times S^2). \end{aligned}$$

Therefore, (1) is directly implied by Lemma 3.2 (i).

To prove (2), we consider the cocore of the two-handle  $h_2^{i+1}$  attached to  $K_{i+1}$ . The boundary of this cocore (i.e., the belt sphere) is a knot on the boundary of the 4-handle  $D^4$ . Hence it bounds a properly embedded surface  $F$  in  $D^4$ . We consider the surface

$$F' = \{\text{cocore of } h_2^{i+1}\} \cup F \hookrightarrow K3^\circ \setminus \nu(D_{L_i}) \subset K3^\circ.$$

Since  $K3^\circ$  has an even intersection form, the self-intersection number of  $F'$  is even. Note that

$$D_{K_{i+1}} \cdot F' = 1 \not\equiv F' \cdot F' \pmod{2}.$$

So  $[D_{K_{i+1}}]$  is ordinary.

To prove (3), we note that  $D_{K_{i+1}}$  and  $D'_{K_{i+1}}$  are topologically isotopic relative to the boundary in  $K3^\circ$ . Hence they are also topologically isotopic relative to the boundary in  $X_i$ . Since  $D'_{K_{i+1}}$  is smoothly isotopic to  $f(D'_{K_{i+1}})$  relative to the boundary, we see that  $D_{K_{i+1}}$  and  $f(D'_{K_{i+1}})$  are topologically isotopic relative to the boundary in  $X_i$ . In particular, they are homotopic to each other relative to the boundary in  $X_i$ . Consider the map  $\tau : S^2 \rightarrow X_i$  which sends the upper hemisphere and the lower hemisphere to  $D_{K_{i+1}}$  and  $f(D'_{K_{i+1}})$  respectively. Since  $D_{K_{i+1}}$  and  $f(D'_{K_{i+1}})$  are homotopic in  $X_i$  relative to the boundary,  $\tau(S^2)$  is null-homotopic in  $X_i$ . By Lemma 3.2 (2),  $\tau(S^2)$  is also null-homotopic in  $X_i \setminus \nu(D_{L_i})$ . Therefore,  $D_{K_{i+1}}$  and  $f(D'_{K_{i+1}})$  are homotopic in  $X_i \setminus \nu(D_{L_i})$  relative to the boundary. This proves (3).  $\square$

#### 4. EQUIVARIANT STABLE HOMOTOPY THEORY

In this section, we briefly recall the definitions of equivariant stable homotopy groups and state some important properties about them. In particular, we state a version of the tom Dieck splitting theorem for an incomplete universe, which turns the computation of the equivariant stable group of a  $G$ -space into the computation of certain non-equivariant stable homotopy groups. Then we use this theorem to prove some algebraic propositions about the  $\text{Pin}(2)$ -equivariant stable homotopy groups of spheres. These results will be essentially used in Section 5 to prove the vanishing theorem for the family Bauer–Furuta invariant. Our exposition here follows the classical method of Adams [1] (which assumes  $G$  is finite). We refer to [45, 32] for the general case of compact Lie groups and more complete discussions using the language of equivariant spectra.

**4.1. The tom Dieck splitting theorem.** Let  $G$  be a compact Lie group. We consider an infinite dimensional  $G$ -representation  $\mathcal{U}$  that satisfies the following two properties:

- There is a canonical embedding from the infinite dimensional trivial representation  $\bigoplus_{\infty} \mathbb{R}$  to  $\mathcal{U}$ .
- For any finite dimensional subrepresentation  $V$  of  $\mathcal{U}$ , there exists an embedding from  $\bigoplus_{\infty} V$  to  $\mathcal{U}$ .

We call such  $\mathcal{U}$  a “ $G$ -universe”. If all finite dimensional representation of  $G$  can be embedded into  $\mathcal{U}$ , then we call  $\mathcal{U}$  a complete universe. Otherwise, we say  $\mathcal{U}$  is incomplete. For our topological applications, it is necessary to consider incomplete universes.

We equip  $\mathcal{U}$  with a  $G$ -invariant inner product. For any pointed  $G$ -spaces  $X, Y$  and any finite dimensional subrepresentations  $V \subset W \subset \mathcal{U}$ , there is a map

$$[S^V \wedge X, S^V \wedge Y]^G \xrightarrow{\text{id}_{S^{W-V}} \wedge -} [S^W \wedge X, S^W \wedge Y]^G,$$

where  $\text{id}_{S^{W-V}}$  denotes the identity map on the representation sphere of the orthogonal complement of  $V$  in  $W$ .

**Definition 4.1.** For any pointed spaces  $X, Y$ , the equivariant stable homotopy group is defined as

$$\{X, Y\}_{\mathcal{U}}^G := \varinjlim_{V \subset \mathcal{U}} [S^V \wedge X, S^V \wedge Y]^G,$$

where the limit is taken over all finite dimensional  $G$ -invariant subspace  $V \subset \mathcal{U}$ . (Just like the non-equivariant setting,  $\{X, Y\}_{\mathcal{U}}^G$  has a abelian group structure.) For  $k \in \mathbb{N}$ , we use  $\pi_{k, \mathcal{U}}^G(X)$  to denote  $\{S^k, X\}_{\mathcal{U}}^G$ .

When  $G$  is the trivial group, there is essentially a unique choice of  $\mathcal{U}$ . In this case, we denote  $\{X, Y\}_{\mathcal{U}}^G$  by  $\{X, Y\}$  and denote  $\pi_{k, \mathcal{U}}^G(X)$  by  $\pi_k^s(X)$ .

Given a closed subgroup  $H$  of  $G$ , we use  $\text{res}_H^G(\mathcal{U})$  to denote the same space  $\mathcal{U}$ , regarded it as an  $H$ -representation space. Then there is a well-defined restriction map

$$\text{res}_H^G : \{X, Y\}_{\mathcal{U}}^G \rightarrow \{X, Y\}_{\text{res}_H^G(\mathcal{U})}^H$$

defined by only remembering the  $H$ -action.

Also, let  $N_G H$  be the normalizer of  $H$  in  $G$  and let  $W_G H = N_G H / H$  be the Weyl group. Then the  $H$ -fixed point of  $\mathcal{U}$ , which we denote by  $\mathcal{U}^H$ , is an  $W_G H$ -universe. There is a well-defined map

$$(9) \quad (-)_{Y, X, G}^H : \{Y, X\}_{\mathcal{U}}^G \rightarrow \{Y^H, X^H\}_{\mathcal{U}^H}^{W_G H},$$

which sends a maps to its restriction to the  $H$ -fixed points. In the special case  $Y = S^k$ , we have the map

$$(-)_{S^k, X, G}^H : \pi_{k, \mathcal{U}}^G(X) \rightarrow \pi_{k, \mathcal{U}^H}^{W_G H}(X^H).$$

When  $X, Y, G$  are obvious from the context, we will suppress them from our notations and simply write  $(-)^H$ .

Next, for any pointed  $G$ -space  $X$ , we introduce a map

$$(10) \quad \zeta_H^G : \pi_{k, \mathcal{U}^H}^{W_G H}(EW_G H_+ \wedge X^H) \rightarrow \pi_{k, \mathcal{U}}^G(X)$$

which is crucial in the tom Dieck splitting theorem. The precise definition of  $\zeta_H^G$  can be found in [45, Page 60]. For our purpose, it suffices to consider the case that  $H$  is a normal subgroup of  $G$ . In this case, we have  $N_G H = G$ ,  $W_G H = G/H$  and the map  $\zeta_H^G$  has a simple description: We have taken an element  $[f] \in \pi_{k, \mathcal{U}^H}^{G/H}((EG/H)_+ \wedge X^H)$ , represented by a  $G/H$ -equivariant map

$$g : S^V \wedge S^k \rightarrow S^V \wedge (EG/H)_+ \wedge X^H$$

for some  $G/H$ -representation  $V \subset \mathcal{U}^H \subset \mathcal{U}$ . We regard  $f$  as a  $G$ -equivariant map and consider the composition

$$(11) \quad \tilde{g} : S^V \wedge S^k \xrightarrow{f} S^V \wedge (EG/H)_+ \wedge X^H \xrightarrow{1 \wedge p_0 \wedge 1} S^V \wedge S^0 \wedge X^H = S^V \wedge X^H \xrightarrow{1 \wedge i} S^V \wedge X.$$

Here  $p_0 : (EG/H)_+ \rightarrow S^0$  is the map that preserves the base point and collapses the whole  $EG/H$  to the other point in  $S^0$ , and the map  $i : X^H \rightarrow X$  is the inclusion map. Note that both  $p_0$  and  $i$  are  $G$ -equivariant because  $H$  is normal. We define  $\zeta_H^G[g] := [\tilde{g}]$ .

Now we state the tom Dieck splitting theorem for an incomplete universe. This theorem was originally proved in [50] for a complete universe. The version given below is a special case of [33, Theorem 3.3]. See [12, 32] for various generalizations to equivariant spectra.



**Theorem 4.2.** *Let  $X$  be a pointed  $G$ -space. Then the map*

$$(12) \quad \sum_{[H]} \zeta_H^G : \bigoplus_{[H]} \pi_{k, \mathcal{U}^H}^{W_G H} (EW_G H_+ \wedge X^H) \rightarrow \pi_{k, \mathcal{U}}^G(X)$$

*is an isomorphism for any  $k \geq 0$ . Here the direct sum is taken over all conjugacy classes of subgroups  $H$  that arises as the isotropy group of some point in  $\mathcal{U}$ .*

Note that the  $W_G H$ -action on  $(EW_G H)_+ \wedge X^H$  is free away from the base point. By the Adams' isomorphism theorem [32, Corollary 7.2], we have an isomorphism

$$(13) \quad \pi_k^s((S^{\text{Ad}(W_G H)} \wedge (EW_G H)_+ \wedge X^H)/W_G H) \xrightarrow{\cong} \pi_{k, \mathcal{U}^H}^{W_G H}((EW_H)_+ \wedge X^H).$$

Here  $\text{Ad}(W_G H)$  denotes the adjoint representation of  $W_G H$  on its Lie algebra. Note this isomorphism relies on the fact that  $W_G H$  can be embedded into  $\mathcal{U}^H$  because  $H$  is an isotropy group. (See [33, Theorem 12.2].) Using (13), we can rewrite the isomorphism (12) as the isomorphism

$$(14) \quad \bigoplus_{[H]} \pi_k^s((S^{\text{Ad}(W_G H)} \wedge (EW_G H)_+ \wedge X^H)/W_G H) \xrightarrow{\cong} \pi_{k, \mathcal{U}}^G(X).$$

**4.2. Some additional lemmas.** We introduce a few additional lemmas that will be useful in our later computations.

**Lemma 4.3.** *Let  $X$  be a pointed space and  $Y$  be a pointed  $G$ -space. Then  $\{G_+ \wedge X, Y\}_{\mathcal{U}}^G \cong \{X, Y\}$ .*

*Proof.* This comes from the standard adjunction  $(G_+ \wedge -) \rightleftharpoons \text{res}_{\{e\}}^G(-)$ . (See [32, Theorem 4.7] and [1, Theorem 5.1]).  $\square$

**Lemma 4.4.** *Let  $X$  be a  $d$ -dimensional  $G$ -CW complex and let  $Y$  a pointed  $G$ -space. Suppose the  $G$ -action on  $X$  is free away from the base point and that the non-equivariant stable homotopy group  $\pi_m^s(Y)$  is trivial for any  $0 \leq m \leq d$ . Then we have  $\{X, Y\}_{\mathcal{U}}^G \cong 0$ .*

*Proof.* This follows from a standard induction argument. Recall that a  $G$ -CW complex is obtained by attaching the  $G$ -cells  $(G/H) \times D^k$  for various closed subgroups  $H$  inductively over  $k$ . Since the  $G$ -action on  $X$  is free away from the base point, all cells (except the base point) are of the form  $G \times D^k$ . Let  $X_k$  be the  $k$ -skeleton of  $X$ . Then

$$X_k/X_{k-1} = G_+ \wedge \left( \bigvee_{A(k)} S^{k\mathbb{R}} \right)$$

where  $A(k)$  denotes the set of  $k$ -cells in  $X$ . For any  $k \leq d$ , by Lemma 4.3, we have

$$\{G_+ \wedge \left( \bigvee_{A(k)} S^{k\mathbb{R}} \right), Y\}_{\mathcal{U}}^G \cong \left\{ \bigvee_{A(k)} S^{k\mathbb{R}}, Y \right\} \cong 0.$$

Therefore, we can inductively prove that  $\{X_k, Y\}_{\mathcal{U}}^G \cong 0$  using the long exact sequence

$$\cdots \rightarrow \{X_k/X_{k-1}, Y\}_{\mathcal{U}}^G \rightarrow \{X_k, Y\}_{\mathcal{U}}^G \rightarrow \{X_{k-1}, Y\}_{\mathcal{U}}^G \rightarrow \cdots$$

induced by the cofiber sequence  $X_{k-1} \rightarrow X_k \rightarrow X_k/X_{k-1}$ .  $\square$

**Lemma 4.5.** *Let  $X$  be a space with a free  $G$ -action and let  $V$  be a  $n$ -dimensional  $G$ -representation. Consider the vector bundle  $\tilde{V} := (V \times X)/G$  over the space  $X/G$ . Suppose  $\tilde{V}$  is a trivial bundle. Then there is a base-point preserving,  $G$ -equivariant homeomorphism from  $S^V \wedge X_+$  to  $S^n \wedge X_+$ .*

*Proof.* Since  $\tilde{V}$  is a trivial bundle, there exists a map  $\rho_0 : (V \times X)/G \rightarrow \mathbb{R}^n$  which restricts to a linear isomorphism on any fiber. We compose this map with the quotient map  $V \times X \rightarrow (V \times X)/G$  to get an  $G$ -equivariant map  $\rho_1 : V \times X \rightarrow \mathbb{R}^n$ . Then we consider the  $G$ -equivariant homeomorphism

$$\rho : V \times X \rightarrow \mathbb{R}^n \times X$$

defined by  $\rho(\vec{v}, x) := (\rho_1(\vec{v}, x), x)$ . After adding the base point at infinity,  $\rho$  induces the homeomorphism from  $S^V \wedge X_+$  to  $S^n \wedge X_+$ .  $\square$

**4.3. Pin(2)-equivariant stable homotopy groups.** From now on, we set  $\mathcal{U}$  to be the incomplete Pin(2)-universe  $\mathbb{H}^\infty \oplus \tilde{\mathbb{R}}^\infty \oplus \mathbb{R}^\infty$ . From this universe, we can define four other universes  $\text{res}_{S^1}^{\text{Pin}(2)}(\mathcal{U})$ ,  $\text{res}_{\{e\}}^{\text{Pin}(2)}(\mathcal{U})$ ,  $\mathcal{U}^{S^1}$  and  $\mathcal{U}^{\text{Pin}(2)}$ . We denote them by  $\mathcal{U}_1, \mathcal{U}_2, \mathcal{U}_3$  and  $\mathcal{U}_4$  respectively. The corresponding Lie groups are  $S^1, \{e\}, C_2$  and  $\{e\}$  respectively. Here  $C_2 = \text{Pin}(2)/S^1$  denotes the cyclic group of order 2.

We will use  $S(\mathbb{H}^\infty)$  as a specific model for  $E\text{Pin}(2)$ . Then  $B\text{Pin}(2)$  is the quotient of  $BS^1 = \mathbb{C}\mathbb{P}^\infty$  under the involution

$$[(z_0, z_1, z_2, z_3, \dots)] \rightarrow [(-\bar{z}_1, \bar{z}_0, -\bar{z}_3, \bar{z}_2, \dots)].$$

We start with the following lemma:

**Lemma 4.6.** *Let  $X$  be a Pin(2)-CW complex. Then for any  $k \geq 0$  the kernel of the map*

$$(15) \quad (-)_{S^k, X, \text{Pin}(2)}^{S^1} : \pi_{k, \mathcal{U}}^{\text{Pin}(2)}(X) \rightarrow \pi_{k, \mathcal{U}_3}^{C_2}(X)$$

is exactly the image of the injective map  $\zeta_{\{e\}}^{\text{Pin}(2)}$ . In particular, the kernel of the map (15) is isomorphic to the non-equivariant stable homotopy group  $\pi_k^s((S^{\tilde{\mathbb{R}}} \wedge E\text{Pin}(2)_+ \wedge X)/\text{Pin}(2))$ .

*Proof.* We consider the following diagram

$$\begin{array}{ccc} \pi_{k, \mathcal{U}}^{\text{Pin}(2)}(E\text{Pin}(2)_+ \wedge X) \oplus \pi_{k, \mathcal{U}_3}^{C_2}((EC_2)_+ \wedge X^{S^1}) \oplus \pi_{k, \mathcal{U}_4}(X^{\text{Pin}(2)}) & \xrightarrow{\zeta_{\{e\}}^{\text{Pin}(2)} \oplus \zeta_{S^1}^{\text{Pin}(2)} \oplus \zeta_{\text{Pin}(2)}^{\text{Pin}(2)}} & \pi_{k, \mathcal{U}}^{\text{Pin}(2)}(X) \\ \downarrow \pi & & \downarrow (-)_{S^k, X, \text{Pin}(2)}^{S^1} \\ \pi_{k, \mathcal{U}_3}^{C_2}((EC_2)_+ \wedge X^{S^1}) \oplus \pi_{k, \mathcal{U}_4}(X^{\text{Pin}(2)}) & \xrightarrow{\zeta_{C_2}^{C_2} \oplus \zeta_{\{e\}}^{C_2}} & \pi_{k, \mathcal{U}_3}^{C_2}(X^{S^1}) \end{array},$$

where  $\pi$  is the projection to the second and third summand. By checking the explicit description of  $\zeta_H^G$  in (11), it is straightforward to see that this diagram commutes. Therefore, the kernel of  $(-)_{S^k, X, \text{Pin}(2)}^{S^1}$  is exactly the image of  $\zeta_{\{e\}}^G$ . We then apply (13) to finish the proof.  $\square$

Using the same argument, one can prove the following lemma.

**Lemma 4.7.** *Let  $X$  be an  $S^1$ -CW complex. Then the kernel of the map*

$$(-)_{S^k, X, S^1}^{S^1} : \pi_{k, \mathcal{U}_3}^{S^1}(X) \rightarrow \pi_k^s(X^{S^1})$$

equals the image of the injective map  $\zeta_{\{e\}}^{S^1} : \pi_{k, \mathcal{U}_3}^{S^1}(ES_+^1 \wedge X) \rightarrow \pi_{k, \mathcal{U}_3}^{S^1}(X)$ . In particular, the kernel is isomorphic to  $\pi_{k-1}^s((ES_+^1 \wedge X)/S^1)$ .

The following propositions (Proposition 4.8~ Proposition 4.10) are the key results in this section. They are the homotopy theoretic version of Theorem 1.9.

**Proposition 4.8.** *Suppose  $m, n \geq 0$  and are not both zero. Then the map*

$$(-)_{S^1, S^{m\mathbb{H}+n\tilde{\mathbb{R}}}, \text{Pin}(2)}^{S^1} : \pi_{1, \mathcal{U}}^{\text{Pin}(2)}(S^{m\mathbb{H}+n\tilde{\mathbb{R}}}) \rightarrow \pi_{1, \mathcal{U}_3}^{C_2}(S^{n\tilde{\mathbb{R}}})$$

*is injective.*

**Proposition 4.9.** *Suppose  $n > 4m \geq 0$ . Then the map*

$$(-)_{S^{\mathbb{R}+m\mathbb{H}}, S^{n\tilde{\mathbb{R}}}, \text{Pin}(2)}^{S^1} : \{S^{\mathbb{R}+m\mathbb{H}}, S^{n\tilde{\mathbb{R}}}\}_{\mathcal{U}}^{\text{Pin}(2)} \rightarrow \pi_{1, \mathcal{U}_3}^{C_2}(S^{n\tilde{\mathbb{R}}})$$

*is injective.*

**Proposition 4.10.** *Suppose  $n = 4m \geq 0$ . Then the map*

$$(-)_{S^{\mathbb{R}+m\mathbb{H}}, S^{n\tilde{\mathbb{R}}}, \text{Pin}(2)}^{S^1} : \{S^{\mathbb{R}+m\mathbb{H}}, S^{n\tilde{\mathbb{R}}}\}_{\mathcal{U}}^{\text{Pin}(2)} \rightarrow \pi_{1, \mathcal{U}_3}^{C_2}(S^{n\tilde{\mathbb{R}}})$$

*has a kernel isomorphic to  $\mathbb{Z}/2$ . Furthermore, let  $\alpha$  be the nonzero element in this kernel. Then we have  $\text{res}_{S^1}^{\text{Pin}(2)}(\alpha) = 0$ .*

**Proposition 4.11.** *The map*

$$(-)_{S^{2\mathbb{R}}, S^0, \text{Pin}(2)}^{S^1} : \pi_{2, \mathcal{U}}^{\text{Pin}(2)}(S^0) \rightarrow \pi_{2, \mathcal{U}_3}^{C_2}(S^0)$$

*has a kernel isomorphic to  $\mathbb{Z}/2$ . Furthermore, let  $\alpha'$  be the nonzero element in this kernel. Then we have  $\text{res}_{S^1}^{\text{Pin}(2)}(\alpha') = 0$ .*

*Proof of Proposition 4.8.* By Lemma 4.6, we have

$$(16) \quad \text{Ker}(-)_{S^1, S^{m\mathbb{H}+n\tilde{\mathbb{R}}}, \text{Pin}(2)}^{S^1} \cong \pi_1^s((S^{m\mathbb{H}+(n+1)\tilde{\mathbb{R}}} \wedge E\text{Pin}(2)_+)/\text{Pin}(2)).$$

Let  $\xi_{\tilde{\mathbb{R}}}$  (resp.  $\xi_{\mathbb{H}}$ ) be the bundle associated to the universal bundle  $\text{Pin}(2) \hookrightarrow E\text{Pin}(2) \rightarrow B\text{Pin}(2)$  and the representation  $\mathbb{R}$  (resp.  $\mathbb{H}$ ). Then we have

$$(S^{m\mathbb{H}+(n+1)\tilde{\mathbb{R}}} \wedge E\text{Pin}(2)_+)/\text{Pin}(2) = \text{Th}(B\text{Pin}(2), m\xi_{\mathbb{H}} \oplus (n+1)\xi_{\tilde{\mathbb{R}}}).$$

Here  $\text{Th}(-)$  denotes the Thom space.

Suppose  $m, n \geq 0$  are not both zero. Then  $m\xi_{\mathbb{H}} \oplus (n+1)\xi_{\tilde{\mathbb{R}}}$  is a vector bundle of dimension  $\geq 2$ . Therefore, other than the base point, the space  $\text{Th}(B\text{Pin}(2), m\xi_{\mathbb{H}} \oplus (n+1)\xi_{\tilde{\mathbb{R}}})$  has no cells in dimension 0 and 1. By the skeletal approximation theorem, we have

$$\text{Ker}(-)_{S^1, S^{m\mathbb{H}+n\tilde{\mathbb{R}}}, \text{Pin}(2)}^{S^1} \cong \pi_1^s(\text{Th}(B\text{Pin}(2), m\xi_{\mathbb{H}} \oplus (n+1)\xi_{\tilde{\mathbb{R}}})) \cong 0.$$

□

*Proof of Proposition 4.9.* The case  $m = 0$  has been covered by Proposition 4.8 so we assume  $m \geq 1$ , which implies that  $n \geq 5$ . Take

$$\beta \in \text{Ker}((-)_{S^{\mathbb{R}+m\mathbb{H}}, S^{n\tilde{\mathbb{R}}}, \text{Pin}(2)}^{S^1}).$$

Take the inclusion  $i_0 : S^{\mathbb{R}} \rightarrow S^{\mathbb{R}+m\mathbb{H}}$ . We also use  $i_0$  to denote the corresponding element in  $\{S^{\mathbb{R}}, S^{\mathbb{R}+m\mathbb{H}}\}_{\mathcal{U}}^{\text{Pin}(2)}$ . Then we have

$$(17) \quad (-)_{S^{\mathbb{R}}, S^{n\mathbb{R}}, \text{Pin}(2)}^{S^1}(\beta \cdot i_0) = (-)_{S^{\mathbb{R}+m\mathbb{H}}, S^{n\mathbb{R}}, \text{Pin}(2)}^{S^1}(\beta) \cdot (-)_{S^{\mathbb{R}}, S^{\mathbb{R}+m\mathbb{H}}, \text{Pin}(2)}^{S^1}(i_0) = 0.$$

But  $\text{Ker}((-)_{S^{\mathbb{R}}, S^{n\mathbb{R}}, \text{Pin}(2)}^{S^1})$  is trivial by (1). Hence we see that  $\beta \cdot i_0 = 0$ .

Next, we consider the long exact sequence

$$\dots \rightarrow \{S^{2\mathbb{R}} \wedge (S(m\mathbb{H})_+), S^{m\mathbb{R}}\}_{\mathcal{U}}^{\text{Pin}(2)} \rightarrow \{S^{\mathbb{R}+m\mathbb{H}}, S^{n\mathbb{R}}\}_{\mathcal{U}}^{\text{Pin}(2)} \xrightarrow{-i_0} \{S^{\mathbb{R}}, S^{m\mathbb{R}}\}_{\mathcal{U}}^{\text{Pin}(2)} \rightarrow \dots$$

induced by the cofiber sequence  $S^0 \rightarrow S^{m\mathbb{H}} \rightarrow S^{\mathbb{R}} \wedge S(m\mathbb{H})_+$ . Note that  $\text{Pin}(2)$  acts freely on  $S^{2\mathbb{R}} \wedge (S(m\mathbb{H})_+)$  away from the base point, and the quotient is a CW complex of dimension  $4m$ . Since  $n > 4m$ , by Lemma 4.4, we see that  $\{S^{2\mathbb{R}} \wedge (S(m\mathbb{H})_+), S^{n\mathbb{R}}\}_{\mathcal{U}}^{\text{Pin}(2)} \cong 0$ . Hence the map  $- \cdot i_0$  is injective. Since  $\beta \cdot i_0 = 0$ , we get  $\beta = 0$ .  $\square$

We need some preparations in order to prove Proposition 4.10.

**Lemma 4.12.** (1) For  $m > 0$ , we have  $\{S^{2\mathbb{R}} \wedge S(m\mathbb{H})_+, S^{4m\mathbb{R}}\}_{\mathcal{U}}^{\text{Pin}(2)} \cong \mathbb{Z}/2$ . (2) Let  $\gamma$  be the unique nonzero element in  $\{S^{2\mathbb{R}} \wedge S(m\mathbb{H})_+, S^{4m\mathbb{R}}\}$ . Then  $\text{res}_{S^1}^{\text{Pin}(2)}(\gamma) = 0$ .

*Proof.* (1) The  $\text{Pin}(2)$ -action on  $S^{2\mathbb{R}} \wedge S((m-1)\mathbb{H})_+$  is free away from the base point and the quotient is a CW-complex of dimension  $4m-4$ . By Lemma 4.4, we have

$$\{S^{2\mathbb{R}} \wedge S((m-1)\mathbb{H})_+, S^{4m\mathbb{R}}\}_{\mathcal{U}}^{\text{Pin}(2)} \cong \{S^{3\mathbb{R}} \wedge S((m-1)\mathbb{H})_+, S^{4m\mathbb{R}}\}_{\mathcal{U}}^{\text{Pin}(2)} \cong 0.$$

Consider the cofiber sequence

$$S((m-1)\mathbb{H})_+ \hookrightarrow S(m\mathbb{H})_+ \rightarrow S(m\mathbb{H})/S((m-1)\mathbb{H}).$$

Since  $S(m\mathbb{H})/S((m-1)\mathbb{H}) = S^{(m-1)\mathbb{H}} \wedge (S(\mathbb{H})_+)$ , we have the induced long exact sequence of homotopy groups

$$\{S^{2\mathbb{R}} \wedge S(m\mathbb{H})_+, S^{4m\mathbb{R}}\}_{\mathcal{U}}^{\text{Pin}(2)} \cong \{S^{2\mathbb{R}+(m-1)\mathbb{H}} \wedge (S(\mathbb{H})_+), S^{4m\mathbb{R}}\}_{\mathcal{U}}^{\text{Pin}(2)}.$$

To compute this group, we consider the vector bundles  $\xi'_{\mathbb{H}}$  (resp.  $\xi'_{4\mathbb{R}}$ ) associated to the principal bundle

$$\text{Pin}(2) \hookrightarrow S(\mathbb{H}) \rightarrow \mathbb{R}\mathbb{P}^2$$

for the representations  $\mathbb{H}$  (resp.  $4\mathbb{R}$ ). Since the structure group of both bundles can be lifted to  $SU(2)$  and any  $SU(2)$  bundles is trivial on a 2-dimensional CW complex. We see that  $\xi'_{\mathbb{H}}$  and  $\xi'_{4\mathbb{R}}$  are both trivial. By Lemma 4.5, we have the following  $\text{Pin}(2)$ -equivariant homeomorphisms

$$S^{4\mathbb{R}+(m-1)\mathbb{H}} \wedge (S(\mathbb{H})_+) \cong S^{4m\mathbb{R}} \wedge (S(\mathbb{H})_+) \cong S^{4m\mathbb{R}} \wedge (S(\mathbb{H})_+).$$

They give the following isomorphism

$$\begin{aligned} \{S^{2\mathbb{R}+(m-1)\mathbb{H}} \wedge (S(\mathbb{H})_+), S^{4m\mathbb{R}}\}_{\mathcal{U}}^{\text{Pin}(2)} &\cong \{S^{4\mathbb{R}+(m-1)\mathbb{H}} \wedge (S(\mathbb{H})_+), S^{4m\mathbb{R}+2\mathbb{R}}\}_{\mathcal{U}}^{\text{Pin}(2)} \\ &\cong \{S^{4m\mathbb{R}} \wedge (S(\mathbb{H})_+), S^{4m\mathbb{R}+2\mathbb{R}}\}_{\mathcal{U}}^{\text{Pin}(2)} \\ &\cong \{S(\mathbb{H})_+, S^{2\mathbb{R}}\}_{\mathcal{U}}^{\text{Pin}(2)} \\ &\cong \{\mathbb{R}\mathbb{P}_+^2, S^2\} \\ &\cong \mathbb{Z}/2. \end{aligned}$$

(2) The element  $\text{res}_{S^1}^{\text{Pin}(2)}(\gamma)$  belongs to the group

$$\{S^{2\mathbb{R}} \wedge S(m\mathbb{H})_+, S^{4m\tilde{\mathbb{R}}}\}_{\mathcal{U}_3}^{S^1} \cong \{S(\mathbb{H})_+, S^{(4m-2)\tilde{\mathbb{R}}}\}_{\mathcal{U}_3}^{S^1}.$$

Since  $S^1$  acts freely on  $S(m\mathbb{H})$  and trivially on  $S^{(4m-2)\tilde{\mathbb{R}}}$ , we have

$$\{S(m\mathbb{H})_+, S^{(4m-2)\tilde{\mathbb{R}}}\}_{\mathcal{U}_3}^{S^1} \cong \{(S(m\mathbb{H})_+)/S^1, S^{(4m-2)\tilde{\mathbb{R}}}\} \cong \{\mathbb{C}\mathbb{P}_+^{2m-1}, S^{4m-2}\} \cong \mathbb{Z}.$$

Since  $\gamma$  is 2-torsion,  $\text{res}_{S^1}^{\text{Pin}(2)}(\gamma)$  must be zero.  $\square$

**Lemma 4.13.** *For  $m > 0$ , the map*

$$(18) \quad q^* : \{S^{2\mathbb{R}} \wedge (S(m\mathbb{H})_+), S^{4m\tilde{\mathbb{R}}}\}_{\mathcal{U}}^{\text{Pin}(2)} \rightarrow \{S^{\mathbb{R}+m\mathbb{H}}, S^{4m\tilde{\mathbb{R}}}\}_{\mathcal{U}}^{\text{Pin}(2)}$$

*induced by the quotient map  $q : S^{m\mathbb{H}} \rightarrow S^{m\mathbb{H}}/S^0 = S^{\mathbb{R}} \wedge (S(m\mathbb{H})_+)$  is injective.*

*Proof.* We consider the long exact sequence

$$\dots \rightarrow \{S^{2\mathbb{R}}, S^{4m\tilde{\mathbb{R}}}\}_{\mathcal{U}}^{\text{Pin}(2)} \xrightarrow{p_1^*} \{S^{2\mathbb{R}} \wedge (S(m\mathbb{H})_+), S^{4m\tilde{\mathbb{R}}}\}_{\mathcal{U}}^{\text{Pin}(2)} \xrightarrow{q^*} \{S^{\mathbb{R}+m\mathbb{H}}, S^{4m\tilde{\mathbb{R}}}\}_{\mathcal{U}}^{\text{Pin}(2)} \rightarrow \dots$$

induced by the cofiber sequence  $S(m\mathbb{H})_+ \xrightarrow{p_1} S^0 \xrightarrow{i_1} S^{m\mathbb{H}}$ , where  $i_1$  is the inclusion map and  $p_1 : S(m\mathbb{H})_+ \rightarrow S^0$  is the map that collapses the whole  $S(m\mathbb{H})$  to the non-base point in  $S^0$ . It suffices to show that the map  $p_1^*$  is trivial. By (14), we have

$$\{S^{2\mathbb{R}}, S^{4m\tilde{\mathbb{R}}}\}_{\mathcal{U}}^{\text{Pin}(2)} \cong \{S^2, (S^{(4m+1)\tilde{\mathbb{R}}} \wedge E\text{Pin}(2)_+)/\text{Pin}(2)\} \oplus \{S^2, (S^{4m\tilde{\mathbb{R}}} \wedge (EC_2)_+)/C_2\} \oplus \{S^2, S^0\}.$$

Since  $m > 0$ , the first two summands vanishes by the skeletal approximation theorem. Hence we have

$$\{S^{2\mathbb{R}}, S^{4m\tilde{\mathbb{R}}}\}_{\mathcal{U}}^{\text{Pin}(2)} \cong \{S^2, S^0\} \cong \mathbb{Z}/2,$$

and it generated by  $\xi_{\text{Pin}(2)}^{\text{Pin}(2)}(\eta^2)$ . (Here  $\eta \in \pi_1(S^0)$  is the Hopf map.) We need to show that  $p_1^*(\xi_{\text{Pin}(2)}^{\text{Pin}(2)}(\eta^2)) = 0$ . By the definition of  $\xi_{\text{Pin}(2)}^{\text{Pin}(2)}$  in (11),  $\xi_{\text{Pin}(2)}^{\text{Pin}(2)}(\eta^2)$  is equal to the following composition of stable maps

$$S^2 \xrightarrow{\eta^2} S^0 \xrightarrow{i_1} S^{4m\tilde{\mathbb{R}}},$$

where  $i_1 : S^0 \rightarrow S^{4m\tilde{\mathbb{R}}}$  is the inclusion. Therefore, in order to show that  $p_1^*$  is zero, it suffices to show that the composition

$$(19) \quad S(m\mathbb{H})_+ \xrightarrow{p_1} S^0 \xrightarrow{i_1} S^{4m\tilde{\mathbb{R}}}$$

is stably null-homotopic. By the exact sequence

$$(20) \quad \dots \rightarrow \{S^{m\mathbb{H}}, S^{4m\tilde{\mathbb{R}}}\}_{\mathcal{U}}^{\text{Pin}(2)} \rightarrow \{S^0, S^{4m\tilde{\mathbb{R}}}\}_{\mathcal{U}}^{\text{Pin}(2)} \xrightarrow{p^*} \{S(m\mathbb{H})_+, S^{4m\tilde{\mathbb{R}}}\}_{\mathcal{U}}^{\text{Pin}(2)} \rightarrow \dots,$$

the composition (19) is trivial if  $i_1 : S^0 \rightarrow S^{4m\tilde{\mathbb{R}}}$  can be extended to a map  $\tilde{i} : S^{m\mathbb{H}} \rightarrow S^{4m\tilde{\mathbb{R}}}$ . Note that there is a  $\text{Pin}(2)$ -equivariant map  $\varphi : S^{\mathbb{H}} \rightarrow S^{3\tilde{\mathbb{R}}}$  defined by

$$\varphi(z + wj) := (|z|^2 - |w|^2, \text{Re}(z\bar{w}), \text{Im}(z\bar{w})).$$

By smashing  $m$  copies of  $\varphi$  and the inclusion map  $S^0 \hookrightarrow S^{m\tilde{\mathbb{R}}}$ , we get the desired extension  $\tilde{i}$ .  $\square$

*Proof of Proposition 4.10.* We first consider the case  $m = n = 0$ . By (16), we get

$$\mathrm{Ker}(-)_{S^{\mathbb{R}}, S^0, \mathrm{Pin}(2)}^{S^1} \cong \pi_1^s(\mathrm{Th}(B\mathrm{Pin}(2), \xi_{\mathbb{R}})).$$

Note that the 3-skeleton of  $B\mathrm{Pin}(2)$  is  $\mathbb{R}\mathbb{P}^2$  and the restriction of  $\xi_{\mathbb{R}}$  to the 3-skeleton is exactly the tautological line bundle over it. Therefore, the 4-skeleton of  $\mathrm{Th}(B\mathrm{Pin}(2), \xi_{\mathbb{R}})$  is  $\mathbb{R}\mathbb{P}^3$ . This implies that

$$\mathrm{Ker}(-)_{S^1, S^0, \mathrm{Pin}(2)}^{S^1} \cong \pi_1^s(\mathbb{R}\mathbb{P}^3) \cong \mathbb{Z}/2.$$

Let  $\alpha$  be the nonzero element in this kernel. By definition,  $\mathrm{res}_{S^1}^{\mathrm{Pin}(2)}(\alpha)$  belongs to the kernel of the map

$$(-)_{S^1, S^{\mathbb{R}}, S^1}^{S^1} : \pi_{1, \mathcal{U}_2}^{S^1}(S^0) \rightarrow \pi_1^s(S^0).$$

By Lemma 4.7, we have

$$\mathrm{Ker}(-)_{S^1, S^{\mathbb{R}}, S^1}^{S^1} \cong \pi_0^s(\mathbb{C}\mathbb{P}_+^\infty) \cong \mathbb{Z}.$$

Since  $\alpha$  is 2-torsion, we must have  $\mathrm{res}_{S^1}^{\mathrm{Pin}(2)}(\alpha) = 0$ .

Now we consider the case  $n = 4m > 0$ . We take a element

$$\beta \in \mathrm{Ker}(-)_{S^{\mathbb{R}+m\mathbb{H}}, S^{n\mathbb{R}}, \mathrm{Pin}(2)}^{S^1}.$$

Similar to the proof of Proposition 4.9, we can show that  $\beta$  belongs to the image of the map

$$\{S^{2\mathbb{R}}, S^{4m\mathbb{R}}\}_{\mathcal{U}}^{\mathrm{Pin}(2)} \xrightarrow{p_1^*} \{S^{2\mathbb{R}} \wedge (S(m\mathbb{H})_+), S^{4m\mathbb{R}}\}_{\mathcal{U}}^{\mathrm{Pin}(2)}.$$

By Lemma 4.12,  $\{S^{2\mathbb{R}}, S^{4m\mathbb{R}}\}_{\mathcal{U}}^{\mathrm{Pin}(2)} = \{0, \gamma\}$ . So  $\beta$  is either 0 or  $p_1^*(\gamma)$ . By Lemma 4.13,  $p_1^*(\gamma) \neq 0$ . Also note that  $(-)_{S^1}^{S^1}(\gamma) = 0$  because the  $S^1$ -action on its domain  $S^{2\mathbb{R}} \wedge S(m\mathbb{H})_+$  is free away from the base point. This implies that  $p_1^*(\gamma)$  actually belongs to  $\mathrm{Ker}((-)_{S^{\mathbb{R}+m\mathbb{H}}, S^{n\mathbb{R}}, \mathrm{Pin}(2)}^{S^1})$ . To this end, we see that

$$\mathrm{Ker}((-)_{S^{\mathbb{R}+m\mathbb{H}}, S^{n\mathbb{R}}, \mathrm{Pin}(2)}^{S^1}) = \{0, p_1^*(\gamma)\} \cong \mathbb{Z}/2.$$

Since  $\mathrm{res}_{S^1}^{\mathrm{Pin}(2)}(\gamma) = 0$  by Lemma 4.12, we have  $\mathrm{res}_{S^1}^{\mathrm{Pin}(2)}(p_1^*(\gamma)) = 0$ .  $\square$

*Proof of Proposition 4.11.* By Lemma 4.6, we have

$$\mathrm{Ker}(-)_{S^{2\mathbb{R}}, S^0, \mathrm{Pin}(2)}^{S^1} \cong \pi_2^s(\mathrm{Th}(B\mathrm{Pin}(2), \xi_{\mathbb{R}})) \cong \pi_2^s(\mathbb{R}\mathbb{P}^3).$$

Stably,  $\mathbb{R}\mathbb{P}^3$  splits into  $S^3 \vee \mathbb{R}\mathbb{P}^2$ . Note that  $\mathbb{R}\mathbb{P}^2$  fits in to the cofiber sequence

$$S^1 \xrightarrow{\cdot 2} S^1 \rightarrow \mathbb{R}\mathbb{P}^2$$

Using the induced long exact sequence of stable homotopy groups, we get an isomorphism

$$\mathrm{Ker}(-)_{S^{2\mathbb{R}}, S^0, \mathrm{Pin}(2)}^{S^1} \cong \pi_2^s(\mathbb{R}\mathbb{P}^3) \xrightarrow[\eta \cdot -]{\cong} \pi_1^s(\mathbb{R}\mathbb{P}^3) \cong \mathbb{Z}/2,$$

where  $\eta \in \pi_1^s(S^0)$  denotes the Hopf map. Let  $\alpha'$  be the nonzero element in this kernel. Then since the smash product with  $\eta$  commutes with the Adams isomorphism (13), we have  $\alpha' = \eta \cdot \alpha$ , where  $\alpha \in \mathrm{Ker}(-)_{S^{\mathbb{R}}, S^0, \mathrm{Pin}(2)}^{S^1}$  is the element established in Proposition 4.10. By that proposition, we get

$$\mathrm{res}_{S^1}^{\mathrm{Pin}(2)}(\alpha') = \eta \cdot \mathrm{res}_{S^1}^{\mathrm{Pin}(2)}(\alpha) = 0.$$

$\square$

## 5. PROPERTIES OF THE FAMILY BAUER–FURUTA INVARIANT

In this section, we use the homotopy theoretic results that we proved in Section 4 to study the family Bauer–Furuta invariant. We will start with proving the vanishing properties of the family Bauer–Furuta invariant. Then we give a space level construction of the stable homotopy class  $\alpha_0$  given in Theorem 1.8, which provides a potential tool to detect exotic diffeomorphisms on  $S^4$ .

**5.1. Proof of the vanishing results.** To prove Theorem 1.9, we give the following definition, which is motivated by the construction of the family Bauer–Furuta map defined in Section 2.

**Definition 5.1.** Let  $\alpha$  be an element in one of the following stable homotopy groups:

- $\{S^{k\mathbb{R}} \wedge S^{m\mathbb{H}}, S^{n\tilde{\mathbb{R}}}\}_{\mathcal{U}}^{\text{Pin}(2)}$  for some integers  $k, m, n \geq 0$ , or
- $\{S^{k\mathbb{R}}, S^{n\tilde{\mathbb{R}}+(-m)\mathbb{H}}\}_{\mathcal{U}}^{\text{Pin}(2)}$  for some integers  $k, n \geq 0$  and  $m < 0$ .

We say  $\alpha$  is an element of *BF-type* if it satisfies the following condition

$$(-)^{S^1}(\alpha) = 0 \in \{S^{k\mathbb{R}}, S^{n\tilde{\mathbb{R}}}\}_{\mathcal{U}_3}^{C_2}.$$

Namely, the restriction of  $\alpha$  to the  $S^1$ -fixed points is trivial as a  $C_2$ -equivariant stable homotopy class.

**Lemma 5.2.** *Let  $X$  be a spin 4-manifold with  $b_1(X) = 0$ . Then we have the following results:*

- Let  $f$  be a orientation preserving self-diffeomorphism on  $X$  that preserves the spin structure and acts trivially on  $H^*(X; \mathbb{R})$ . Let  $\tilde{f}$  be an automorphism of the spin structure on  $X$  that lifts  $f$ . Then  $\text{FBF}^{\text{Pin}(2)}(X, f, \tilde{f})$  is an element of *BF-type*.
- Let  $\gamma : S^1 \rightarrow \text{Diff}(X)$  be a loop based at  $\text{id}_X$ . Suppose  $\gamma$  is spin-liftable (see Definition 2.2). Then  $\text{FBF}^{\text{Pin}(2)}(X, \gamma)$  is an element of *BF-type*.

*Proof.* We focus on the proof of (i) since (ii) is similar. Consider the approximated Seiberg–Witten map  $\tilde{sw} : U \rightarrow V$  (see equation (2)). Recall that both the domain and the target of this map consists of pairs  $(\alpha, \phi)$ , where  $\alpha$  is a differential form on  $X$  and  $\phi$  is a spinor on  $X$ . The symmetry group  $S^1$  is the constant gauge group. Hence it acts trivially on  $\alpha$  and acts as the scalar multiplication on  $\phi$ . From this description, we see that  $(\alpha, \phi)$  is an  $S^1$ -fixed point if and only if  $\phi = 0$ . As a result, when restricted to the  $S^1$ -fixed points,  $\tilde{sw}$  is a finite dimensional approximation of the family of linear operators

$$\{(d^*, d^+)_s : \Omega^1(X_s; i\mathbb{R}) \rightarrow (\Omega^0(X_s; i\mathbb{R})/\mathbb{R}) \oplus \Omega_+^2(X_s; i\mathbb{R})\}_{s \in S(2\mathbb{R})}.$$

The trivialization (3) induce isomorphisms

$$U \cap \Omega^1(X_s; i\mathbb{R}) \cong \mathbb{R}^{n'},$$

$$V \cap (\Omega^0(X_s; i\mathbb{R})/\mathbb{R}) \oplus \Omega^2(X_s; i\mathbb{R}) \cong \mathbb{R}^{n'+b_2^+(X_s)}.$$

Furthermore, by checking the construction of these trivializations (3) in [31, 35], we see that under the above isomorphisms, the operator  $(d^*, d^+)_s$  is exactly the inclusion  $\mathbb{R}^{n'} \hookrightarrow \mathbb{R}^{n'+b_2^+(X)}$  into the first  $n'$  components. From this discussion, we see that when we restrict the induced map

$$\tilde{sw}^+ : S_+^1 \wedge S^{(m' - \frac{\sigma(X)}{16})\mathbb{H} + n'\tilde{\mathbb{R}}} \rightarrow S^{m'\mathbb{H} + (n'+b_2^+(X))\tilde{\mathbb{R}}}$$

to the  $S^1$ -fixed points, it is homotopic to the following map

$$(21) \quad (S(2\mathbb{R})_+)^{\wedge} \wedge S^{n'\tilde{\mathbb{R}}} \xrightarrow{p_2 \wedge i_2} S^0 \wedge S^{(n+n')\tilde{\mathbb{R}}}$$

where  $p_2 : S(2\mathbb{R})_+ \rightarrow S^0$  preserves the base point and collapse  $S(2\mathbb{R})$  to the other point in  $S^0$ , and  $i_2 : S^{n'}\tilde{\mathbb{R}} \rightarrow S^{(n+n')\tilde{\mathbb{R}}}$  is the inclusion into the first  $n'$  components.

Recall that  $\text{FBF}^{\text{Pin}(2)}(X, f, \tilde{f})$  is defined via the composition (5). From our discussion above, the element  $(-)^{S^1}(\text{FBF}^{\text{Pin}(2)}(X, f, \tilde{f}))$  can be represented by the following composition

$$S^{2\mathbb{R}} \wedge S^{n'\tilde{\mathbb{R}}} \xrightarrow{c_0 \wedge \text{id}_{S^{n'\tilde{\mathbb{R}}}}} S^{\mathbb{R}} \wedge (S(2\mathbb{R})_+) \wedge S^{n'\tilde{\mathbb{R}}} \xrightarrow{\text{id}_{S^{\mathbb{R}}} \wedge p_2 \wedge i_2} S^{\mathbb{R}} \wedge S^0 \wedge S^{(n+n')\tilde{\mathbb{R}}}.$$

Note that  $c_0 \circ (\text{id}_{S^{\mathbb{R}}} \wedge p_2) : S^{2\mathbb{R}} \rightarrow S^{\mathbb{R}}$  is null-homotopic. Thus  $(-)^{S^1}(\text{FBF}^{\text{Pin}(2)}(X, f, \tilde{f})) = 0$ .  $\square$

The following proposition is just a restatement of Proposition 4.8  $\sim$  Proposition 4.11.

- Proposition 5.3.** (i) Suppose  $m, n \geq 0$  and are not both zero. Then  $\pi_{1, \mathcal{U}}^{\text{Pin}(2)}(S^{m\mathbb{H}+n\tilde{\mathbb{R}}})$  doesn't contain any nonzero element of BF-type.
- (ii) Suppose  $n > 4m \geq 0$ . Then  $\{S^{\mathbb{R}+m\mathbb{H}}, S^{n\tilde{\mathbb{R}}}\}_{\mathcal{U}}^{\text{Pin}(2)}$  doesn't contain any nonzero element of BF-type.
- (iii) Suppose  $n = 4m \geq 0$ . Then  $\{S^{\mathbb{R}+m\mathbb{H}}, S^{n\tilde{\mathbb{R}}}\}_{\mathcal{U}}^{\text{Pin}(2)}$  contains a unique nonzero element of BF-type, denoted by  $\alpha_0$ . Furthermore, we have  $\text{res}_{S^1}^{\text{Pin}(2)}(\alpha_0) = 0$ .
- (iv)  $\pi_2^{\text{Pin}(2)}(S^0)$  contains a unique nonzero element of BF-type, denoted by  $\alpha_1$ . Furthermore, we have  $\text{res}_{S^1}^{\text{Pin}(2)}(\alpha_1) = 0$ .

*Proof of Theorem 1.7.* Let  $\gamma$  an exotic loop in  $\text{Diff}(S^4)$ . Then  $\gamma$  is spin-liftable. By Lemma 5.2,  $\text{FBF}^{\text{Pin}(2)}(S^4, \gamma)$  is an element in  $\pi_2^{\text{Pin}(2)}(S^0)$  of BF-type. By Proposition 5.3 (iv),  $\text{FBF}^{\text{Pin}(2)}(S^4, \gamma)$  is either 0 or  $\alpha_1$ . So we have

$$\text{FBF}^{S^1}(S^4, \gamma) = \text{res}_{S^1}^{\text{Pin}(2)}(\text{FBF}^{\text{Pin}(2)}(S^4, \gamma)) = 0.$$

This further implies that  $\text{FBF}^{\{e\}}(S^4, \gamma) = 0$ .

Now suppose  $\gamma$  is homotopic to  $2\gamma'$  for some loop  $\gamma'$  in  $\text{Diff}(S^4)$ . We may assume that  $\gamma'$  is also spin-liftable by compositing it with an essential loop in  $\text{SO}(5) \subset \text{Diff}(S^4)$ . By Proposition 5.3 (iv) again,  $\text{FBF}^{\text{Pin}(2)}(S^4, \gamma')$  is either 0 or  $\alpha_1$ . Since  $\alpha_1$  is 2-torsion, we have

$$\text{FBF}^{\text{Pin}(2)}(S^4, \gamma) = 2 \text{FBF}^{\text{Pin}(2)}(S^4, \gamma') = 0.$$

$\square$

*Proof of Theorem 1.8.* This is a special case Proposition 5.3 (iii) for  $m = n = 0$ .  $\square$

*Proof of Theorem 1.9:* Assume the intersection form of  $X$  is isomorphic to  $2mE8 \oplus n\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  for some  $m \in \mathbb{Z}$  and  $n \in \mathbb{N}$ . Then  $2\chi(X) + 3\sigma(X) = 4 + 4 \cdot (n - 4m)$ .

Suppose  $2\chi(X) + 3\sigma(X) = 4$ . Then  $m = 4n$ . By Lemma 5.2,  $\text{FBF}^{\text{Pin}(2)}(X, f, \tilde{f})$  is an element in  $\{S^{\mathbb{R}+m\mathbb{H}}, S^{n\tilde{\mathbb{R}}}\}_{\mathcal{U}}^{\text{Pin}(2)}$  of BF-type. Therefore, by Proposition 5.3 (iii), we have

$$\text{FBF}^{S^1}(X, f, \tilde{f}) = \text{res}_{S^1}^{\text{Pin}(2)}(\text{FBF}^{\text{Pin}(2)}(X, f, \tilde{f})) = 0.$$

This proves (1).

Suppose  $2\chi(X) + 3\sigma(X) > 4$ . Then  $n > 4m$ . Again by Lemma 5.2,  $\text{FBF}^{\text{Pin}(2)}(X, f, \tilde{f})$  is an element of BF-type in  $\pi_{1, \mathcal{U}}^{\text{Pin}(2)}(S^{-m\mathbb{H}+n\tilde{\mathbb{R}}})$  if  $m < 0$  and element of BF-type in  $\{S^{\mathbb{R}+m\mathbb{H}}, S^{n\tilde{\mathbb{R}}}\}_{\mathcal{U}}^{\text{Pin}(2)}$



if  $m \geq 0$ . Then we apply Proposition 5.3 (i) and (ii) to conclude that  $\text{FBF}^{\text{Pin}(2)}(X, f, \tilde{f}) = 0$ . This proves (2).  $\square$

*Remark 5.4.* We mention various examples of nonvanishing family Bauer–Furuta invariants. These examples show that various conditions in Theorem 1.9 are indeed necessary.

- Consider the Dehn twist  $\delta_{K3\#K3}$  along the separating neck in  $K3\#K3$ . Then Kronheimer–Mrowka [31] prove that

$$\text{FBF}^{\{e\}}(K3\#K3, \delta_{K3\#K3}, \tilde{\delta}) = \eta^3 \neq 0$$

for any spin bundle automorphism  $\tilde{\delta}$  which lifts  $\delta_{K3\#K3}$ . Here  $\eta \in \pi_1^s(S^0)$  denotes the Hopf element. Note that  $2\chi(K3\#K3) + 3\sigma(K3\#K3) = -4$ .

- Consider the covering transformation on the spin bundle over  $K3$ . This is a lift of  $\text{id}_{K3}$ , denoted by  $\tilde{\text{id}}_{K3}$ . Then Kronheimer–Mrowka [31] proves that

$$\text{FBF}^{\{e\}}(K3, \text{id}_{K3}, \tilde{\text{id}}_{K3}) = \eta^2 \neq 0.$$

Note that  $2\chi(K3) + 3\sigma(K3) = 0$ .

- Consider the covering transformation on the spin bundle over  $K3\#(S^2 \times S^2)$ , denoted by  $\tilde{\text{id}}_{K3\#(S^2 \times S^2)}$ . Then by [35, Proof of Theorem 1.2], we have

$$\text{FBF}^{\text{Pin}(2)}(K3\#(S^2 \times S^2), \text{id}_{K3\#(S^2 \times S^2)}, \tilde{\text{id}}_{K3\#(S^2 \times S^2)}) \neq 0.$$

Note that  $2\chi(K3\#(S^2 \times S^2)) + 3\sigma(K3\#(S^2 \times S^2)) = 4$ .

**5.2. A space level description of  $\alpha_0$ .** The element  $\alpha_0 \in \pi_{1, \mathcal{U}}^{\text{Pin}(2)}(S^0)$  given in Theorem 1.8 is of special interest because it could arise as the  $\text{Pin}(2)$ -equivariant family Bauer–Furuta invariant of an exotic diffeomorphism on  $S^4$ . In this section, we explicitly construct a map

$$l : S^{\mathbb{R}+\mathbb{H}} \rightarrow S^{\mathbb{H}}$$

which represents  $\alpha_0$ .

We treat  $\text{Pin}(2)$  as a submanifold of  $\mathbb{H}$ , with the standard embedding

$$\text{Pin}(2) = \{e^{i\theta}\} \cup \{je^{i\theta}\} \hookrightarrow \mathbb{H}.$$

Denote the tangent bundle and the normal bundle of  $\text{Pin}(2)$  by  $T_*\text{Pin}(2)$  and  $N_*\text{Pin}(2)$  respectively. Both bundles are  $\text{Pin}(2)$ -bundles. Namely, there are  $\text{Pin}(2)$ -actions on both bundles that cover the left-multiplication action of  $\text{Pin}(2)$  on itself. Furthermore, we have trivializations

$$(22) \quad T_*\text{Pin}(2) \cong \mathbb{R} \times \text{Pin}(2),$$

$$(23) \quad T_*\text{Pin}(2) \oplus N_*\text{Pin}(2) \cong \mathbb{H} \times \text{Pin}(2).$$

Here (22) is induced by the Lie group structure on  $\text{Pin}(2)$  and (23) is restricted from the trivialization of  $T_*\mathbb{H}$ . These two trivializations induce equivariant homeomorphisms

$$\varphi_1 : S^{\mathbb{R}} \wedge \text{Th}(\text{Pin}(2), N_*\text{Pin}(2)) \xrightarrow{\cong} \text{Th}(\text{Pin}(2), T_*\text{Pin}(2) \oplus N_*\text{Pin}(2)),$$

$$\varphi_2 : \text{Th}(\text{Pin}(2), T_*\text{Pin}(2) \oplus N_*\text{Pin}(2)) \xrightarrow{\cong} S^{\mathbb{H}} \wedge (\text{Pin}(2)_+).$$

Let  $\nu(\text{Pin}(2))$  be a  $\text{Pin}(2)$ -equivariant, closed tubular neighborhood of  $\text{Pin}(2)$  in  $\mathbb{H}$ . Then there exists an equivariant homeomorphism

$$\varphi_3 : \nu(\text{Pin}(2))/\partial\nu(\text{Pin}(2)) \xrightarrow{\cong} \text{Th}(\text{Pin}(2), N_*\text{Pin}(2)).$$

Then we consider the following composition

$$\begin{aligned}
(24) \quad l : S^{\mathbb{R}} \wedge S^{\mathbb{H}} &\xrightarrow{\text{id}_{S^{\mathbb{R}}} \wedge c_1} S^{\mathbb{R}} \wedge (\nu(\text{Pin}(2))/\partial\nu(\text{Pin}(2))) \\
&\xrightarrow{\text{id}_{S^{\mathbb{R}}} \wedge \varphi_3} S^{\mathbb{R}} \wedge \text{Th}(\text{Pin}(2), N_*\text{Pin}(2)) \\
&\xrightarrow{\varphi_1} \text{Th}(\text{Pin}(2), T_*\text{Pin}(2) \oplus N_*\text{Pin}(2)) \\
&\xrightarrow{\varphi_2} S^{\mathbb{H}} \wedge (\text{Pin}(2)_+) \\
&\xrightarrow{\text{id}_{S^{\mathbb{H}}} \wedge p_3} S^{\mathbb{H}}
\end{aligned}$$

Here  $c_1 : S^{\mathbb{H}} \rightarrow \nu(\text{Pin}(2))/\partial\nu(\text{Pin}(2))$  is the Pontryagin–Thom collapsing map. The map  $p_3 : \text{Pin}(2)_+ \rightarrow S^0$  preserves the base point and collapses the whole  $\text{Pin}(2)$  into the other point in  $S^0$ .

**Proposition 5.5.**  $\alpha_0$  is represented by the map  $l$ .

*Proof.* By the proof of Proposition 4.9,  $\alpha_0 = \zeta_{\{e\}}^{\text{Pin}(2)}(\alpha_1)$ , where

$$\zeta_{\{e\}}^{\text{Pin}(2)} : \pi_{1, \mathcal{U}}^{\text{Pin}(2)}(E\text{Pin}(2)_+) \rightarrow \pi_{1, \mathcal{U}}^{\text{Pin}(2)}(S^0)$$

is the map defined in (10) and  $\alpha_1$  is any generator of  $\pi_{1, \mathcal{U}}^{\text{Pin}(2)}(E\text{Pin}(2)_+)$ . We first give a space level description of  $\alpha_1$ . Consider the following commutative diagram

$$\begin{array}{ccc}
(25) \quad \pi_{1, \mathcal{U}}^{\text{Pin}(2)}(\text{Pin}(2)_+) & \xrightarrow{i_1} & \pi_{1, \mathcal{U}}^{\text{Pin}(2)}(E\text{Pin}(2)_+) \\
j \uparrow \cong & & j' \uparrow \cong \\
\pi_1^s((S^{\mathbb{R}} \wedge \text{Pin}(2)_+)/\text{Pin}(2)) & \xrightarrow{i'_1} & \pi_1^s((S^{\mathbb{R}} \wedge E\text{Pin}(2)_+)/\text{Pin}(2))
\end{array}$$

Here  $i_*$  and  $i'_*$  are both induced by the inclusion  $i : \text{Pin}(2)_+ \hookrightarrow E\text{Pin}(2)_+$ . And  $j, j'$  are the Adams' isomorphisms (13). By our discussion in the proof of Proposition 4.9, the group  $\pi_1^s((S^{\mathbb{R}} \wedge E\text{Pin}(2)_+)/\text{Pin}(2))$  is isomorphic to  $\mathbb{Z}/2$  and it is generated by the inclusion of the 1-skeleton:

$$S^{\mathbb{R}} \cong (S^{\mathbb{R}} \wedge \text{Pin}(2)_+)/\text{Pin}(2) \hookrightarrow (S^{\mathbb{R}} \wedge E\text{Pin}(2)_+)/\text{Pin}(2).$$

Therefore, the map  $i'_*$  is just the surjective map  $\mathbb{Z} \rightarrow \mathbb{Z}/2$ .

By the diagram (25), we see that  $\alpha_1 = i_* \circ j(1)$ . By the explicit description of the Adams' isomorphism [32, Page 84], we see that  $j(1) \in \pi_{1, \mathcal{U}}^{\text{Pin}(2)}(\text{Pin}(2)_+)$  is represented by the composition

$$(26) \quad l_1 := \varphi_2 \circ \varphi_1 \circ (\text{id}_{S^{\mathbb{R}}} \wedge \varphi_3) \circ (\text{id}_{S^{\mathbb{R}}} \wedge c_1) : S^{\mathbb{R}+\mathbb{H}} \rightarrow S^{\mathbb{H}} \wedge (\text{Pin}(2)_+).$$

(i.e., the composition of the first four maps in (24)). Hence  $\alpha_1$  is presented by the composition

$$l_2 : S^{\mathbb{R}+\mathbb{H}} \xrightarrow{l_1} S^{\mathbb{H}} \wedge (\text{Pin}(2)_+) \xrightarrow{\text{id}_{S^{\mathbb{H}}} \wedge i} S^{\mathbb{H}} \wedge (E\text{Pin}(2)_+).$$

By the construction of  $\zeta_{\{e\}}^{\text{Pin}(2)}$  in (11),  $\alpha_0 = \zeta_{\{e\}}^{\text{Pin}(2)}(\alpha_1)$  is represented by the composition

$$l_3 : S^{\mathbb{R}+\mathbb{H}} \xrightarrow{l_2} S^{\mathbb{H}} \wedge (E\text{Pin}(2)_+) \xrightarrow{\text{id}_{S^{\mathbb{H}}} \wedge p_0} S^{\mathbb{H}}.$$

Since  $p_0 \circ i : \text{Pin}(2)_+ \rightarrow S^0$  is exactly the map  $p_4$ , we see that the maps  $l_3$  and  $l$  are the same.  $\square$

*Remark 5.6.* It is worth noting that  $l$  actually represents an element in  $\{S^1, S^0\}_{\mathcal{U}' }^{\text{Pin}(2)}$ , where  $\mathcal{U}' = \mathbb{H}^\infty \oplus \widetilde{\mathbb{R}}^\infty$  is a universe that only contains the nontrivial representations  $\widetilde{\mathbb{R}}$  and  $\mathbb{H}$ . Compared to the universe  $\mathcal{U} = \mathbb{H}^\infty \oplus \widetilde{\mathbb{R}}^\infty \oplus \mathbb{R}^\infty$  (which we have been working with so far), this smaller universe actually fits better with our setting in gauge theory. This is because the trivializations (3) only involve  $\widetilde{\mathbb{R}}$  and  $\mathbb{H}$ . However, since  $\mathcal{U}'$  doesn't contain the trivial representation  $\mathbb{R}$ , the corresponding stable homotopy category behaves badly. (For example, for general  $X, Y$ , the set  $\{X, Y\}_{\mathcal{U}' }^{\text{Pin}(2)}$  does not even carry a group structure.) This is the main reason why we choose to work with  $\mathcal{U}$  instead of  $\mathcal{U}'$ . We also note that when restricting to the subgroup  $S^1$ , the universes  $\mathcal{U}$  and  $\mathcal{U}'$  are isomorphic.

We end the paper with the following proposition. It states that  $\alpha_0$  cannot be detected by Borel (co)homology, which corresponds to various versions of monopole Floer (co)homology.

**Proposition 5.7.** *Let  $A$  be an abelian group and let  $G$  be a closed subgroup of  $\text{Pin}(2)$ . Then the induced maps*

$$\begin{aligned} \alpha_{0,*} : \tilde{H}_*^G(S^{\mathbb{R}+\mathbb{H}}; A) &\rightarrow \tilde{H}_*^G(S^{\mathbb{H}}; A), \\ \alpha_0^* : \tilde{H}_G^*(S^{\mathbb{H}}; A) &\rightarrow \tilde{H}_G^*(S^{\mathbb{R}+\mathbb{H}}; A). \end{aligned}$$

*between the  $G$ -equivariant reduced Borel (co)homology groups (with coefficient  $A$ ) are both trivial.*

*Proof.*  $\alpha_0$  is represented by the map  $l$ . Since  $l$  factors through the map  $l_1$  in (26). It suffices to show that the induced maps

$$\begin{aligned} l_{1,*} : \tilde{H}_*^G(S^{\mathbb{R}+\mathbb{H}}; A) &\rightarrow \tilde{H}_*^G(S^{\mathbb{H}} \wedge (\text{Pin}(2)_+); A), \\ l_1^* : \tilde{H}_G^*((\text{Pin}(2)_+) \wedge S^{\mathbb{H}}; A) &\rightarrow \tilde{H}_G^*(S^{\mathbb{R}+\mathbb{H}}; A). \end{aligned}$$

are trivial. To prove this, we consider the fiber bundle

$$SU(2)/G \hookrightarrow BG \rightarrow BSU(2),$$

which makes  $H^*(BG; A)$  a module over

$$H^*(BSU(2); \mathbb{Z}) = H^*(\mathbb{H}\mathbb{P}^\infty; \mathbb{Z}) = \mathbb{Z}[v].$$

An elementary argument using the Serre spectral sequence shows that the map

$$v \cdot - : H^m(BG; A) \rightarrow H^{m+4}(BG; A)$$

is an isomorphism for any  $m \geq 0$ . By the definition of Borel cohomology,  $\tilde{H}_G^*(S^{\mathbb{R}+\mathbb{H}}; A)$  is module over  $H^*(BG; \mathbb{Z})$ . Hence  $\tilde{H}_G^*(S^{\mathbb{R}+\mathbb{H}}; A)$  is also over  $\mathbb{Z}[v]$ . By the Thom isomorphism theorem, we have an isomorphism

$$H^*(BG; A) \cong \tilde{H}_G^{*+5}(S^{\mathbb{R}+\mathbb{H}}; A)$$

between two  $\mathbb{Z}[v]$ -modules. Therefore,  $\tilde{H}_G^*(S^{\mathbb{R}+\mathbb{H}}; A)$  is supported in degrees greater or equal 5 and the map

$$v \cdot - : \tilde{H}_G^m(S^{\mathbb{R}+\mathbb{H}}; A) \rightarrow \tilde{H}_G^{m+4}(S^{\mathbb{R}+\mathbb{H}}; A)$$

is an isomorphism for any  $m \geq 5$ . Moreover, since the  $G$ -action on  $S^{\mathbb{H}} \wedge (\text{Pin}(2)_+)$  is free away from the base point, the group

$$\tilde{H}_G^*(S^{\mathbb{H}} \wedge (\text{Pin}(2)_+); A) = \tilde{H}^*((S^{\mathbb{H}} \wedge (\text{Pin}(2)_+)/G; A)$$

is supported in degrees less or equal to 5 (less or equal to 4 is  $G$  is not a finite group).

To this end, we see that  $l_1^*$  can only be nontrivial in degree 5. However, suppose  $l_1^*(x) \neq 0$  for some  $x \in \tilde{H}_G^5(S^{\mathbb{H}} \wedge (\text{Pin}(2)_+); A)$ . Then

$$0 = l_1^*(0) = l_1^*(v \cdot x) = v \cdot l_1^*(x) \neq 0.$$

This contradiction shows that  $l_1^* = 0$  in all degrees. Hence we have  $\alpha_0^* = 0$ . The proof that  $\alpha_{0,*} = 0$  is essentially the same.  $\square$

## REFERENCES

- [1] J. F. Adams. Prerequisites (on equivariant stable homotopy) for Carlsson’s lecture. In *Algebraic topology, Aarhus 1982 (Aarhus, 1982)*, volume 1051 of *Lecture Notes in Math.*, pages 483–532. Springer, Berlin, 1984.
- [2] Selman Akbulut. Isotoping 2-spheres in 4-manifolds. In *Proceedings of the Gökova Geometry-Topology Conference 2014*, pages 264–266. Gökova Geometry/Topology Conference (GGT), Gökova, 2015.
- [3] Selman Akbulut. Corks and exotic ribbons in  $B^4$ . *arXiv preprint arXiv:2103.13967*, 2021.
- [4] Selman Akbulut. On the Smale Conjecture for  $\text{Diff}(S^4)$ , 2021.
- [5] Dave Auckly, Hee Jung Kim, Paul Melvin, and Daniel Ruberman. Stable isotopy in four dimensions. *J. Lond. Math. Soc. (2)*, 91(2):439–463, 2015.
- [6] Dave Auckly, Hee Jung Kim, Paul Melvin, Daniel Ruberman, and Hannah Schwartz. Isotopy of surfaces in 4-manifolds after a single stabilization. *Adv. Math.*, 341:609–615, 2019.
- [7] Dave Auckly and Daniel Ruberman. Exotic phenomena in dimension four: diffeomorphism groups and embedding spaces. *preprint*.
- [8] David Baraglia and Hokuto Konno. A note on the Nielsen realization problem for K3 surfaces. *arXiv preprint arXiv:1908.03970*, 2019.
- [9] David Baraglia and Hokuto Konno. On the Bauer-Furuta and Seiberg-Witten invariants of families of 4-manifolds. *arXiv preprint arXiv:1903.01649*, 2019.
- [10] David Baraglia and Hokuto Konno. A gluing formula for families Seiberg-Witten invariants. *Geom. Topol.*, 24(3):1381–1456, 2020.
- [11] Stefan Bauer and Mikio Furuta. A stable cohomotopy refinement of Seiberg-Witten invariants. I. *Invent. Math.*, 155(1):1–19, 2004.
- [12] Andrew J Blumberg and Michael A Hill. Equivariant stable categories for incomplete systems of transfers. *Journal of the London Mathematical Society*, 2021.
- [13] Jim Bryan. Seiberg-witten theory and  $z/2^p$  actions on spin 4-manifolds. *arXiv preprint dg-ga/9704010*, 1997.
- [14] Ryan Budney and David Gabai. Knotted 3-balls in  $S^4$ . *arXiv preprint arXiv:1912.09029*, 2019.
- [15] Sergey Finashin, Matthias Kreck, and O. Viro. Exotic knotings of surfaces in the 4-sphere. *Bulletin of The American Mathematical Society - BULL AMER MATH SOC*, 17, 04 1987.
- [16] Ronald Fintushel and Ronald J. Stern. Surfaces in 4-manifolds. *Math. Res. Lett.*, 4(6):907–914, 1997.
- [17] Ronald Fintushel and Ronald J. Stern. Six lectures on four 4-manifolds. In *Low dimensional topology*, volume 15 of *IAS/Park City Math. Ser.*, pages 265–315. Amer. Math. Soc., Providence, RI, 2009.
- [18] M. Furuta. Monopole equation and the  $\frac{11}{8}$ -conjecture. *Math. Res. Lett.*, 8(3):279–291, 2001.
- [19] Mikio Furuta, Yukio Kametani, Hirofumi Matsue, and Norihiko Minami. Homotopy theoretical considerations of the bauer-furuta stable homotopy Seiberg-Witten invariants. *Geometry & Topology Monographs*, 10:155–166, 2007.
- [20] David Gabai. The 4-dimensional light bulb theorem. *J. Amer. Math. Soc.*, 33(3):609–652, 2020.
- [21] David T. Gay. Diffeomorphisms of the 4-sphere, Cerf theory and Montesinos twins. 2021.
- [22] Robert E. Gompf and András I. Stipsicz. *4-manifolds and Kirby calculus*, volume 20 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1999.
- [23] Kyle Hayden. Exotically knotted disks and complex curves. 2021.
- [24] Kyle Hayden and Isaac Sundberg. Khovanov homology and exotic surfaces in the 4-ball. 2021.
- [25] Michael J Hopkins, Jianfeng Lin, XiaoLin Danny Shi, and Zhouli Xu. Intersection forms of spin 4-manifolds and the pin (2)-equivariant mahowald invariant. *arXiv preprint arXiv:1812.04052*, 2018.
- [26] Nobuo Iida. A Bauer-Furuta type refinement of Kronheimer-Mrowka’s invariant for 4-manifolds with contact boundary. *arXiv preprint arXiv:1906.07938*, 2019.

- [27] András Juhász, Maggie Miller, and Ian Zemke. Transverse invariants and exotic surfaces in the 4-ball. *Geometry & Topology*, 2020.
- [28] Rob Kirby. Problems in low-dimensional topology. <https://math.berkeley.edu/kirby/>, 1995.
- [29] Hokuto Konno, Jin Miyazawa, and Masaki Taniguchi. Involutions, knots, and Floer K-theory. *arXiv preprint arXiv:2110.09258*, 2021.
- [30] Manuel Krannich and Alexander Kupers. On Torelli groups and Dehn twists of smooth 4-manifolds. *arXiv preprint arXiv:2105.08904*, 2021.
- [31] P. B. Kronheimer and T. S. Mrowka. The Dehn twist on a sum of two  $K3$  surfaces. *Math. Res. Lett.*, 27(6):1767–1783, 2020.
- [32] L. G. Lewis, Jr., J. P. May, M. Steinberger, and J. E. McClure. *Equivariant stable homotopy theory*, volume 1213 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1986. With contributions by J. E. McClure.
- [33] L. Gaunce Lewis, Jr. Splitting theorems for certain equivariant spectra. *Mem. Amer. Math. Soc.*, 144(686):x+89, 2000.
- [34] Francesco Lin. *A Morse-Bott approach to monopole Floer homology and the triangulation conjecture*, volume 255. American Mathematical Society, 2018.
- [35] Jianfeng Lin. Isotopy of the Dehn twist on  $K3\#K3$  after a single stabilization. 2020.
- [36] Ciprian Manolescu. Pin(2)-equivariant Seiberg-Witten Floer homology and the triangulation conjecture. *J. Amer. Math. Soc.*, 29(1):147–176, 2016.
- [37] Patrick Orson and Mark Powell. Mapping class groups of simply-connected 4-manifolds with boundary. *In preparation*.
- [38] B. Perron. Pseudo-isotopies et isotopies en dimension quatre dans la catégorie topologique. *Topology*, 25(4):381–397, 1986.
- [39] Frank Quinn. Isotopy of 4-manifolds. *J. Differential Geom.*, 24(3):343–372, 1986.
- [40] Daniel Ruberman. An obstruction to smooth isotopy in dimension 4. *Math. Res. Lett.*, 5(6):743–758, 1998.
- [41] Birgit Schmidt. Spin 4-manifolds and Pin(2)-equivariant homotopy theory. *Ph. D. thesis, Universität Bielefeld*, 2003.
- [42] Rob Schneiderman and Peter Teichner. Homotopy versus isotopy: spheres with duals in 4-manifolds. *arXiv: Geometric Topology*, 2019.
- [43] Hannah Schwartz. A 4-dimensional light bulb theorem for disks. 2021.
- [44] Hannah R. Schwartz. Equivalent non-isotopic spheres in 4-manifolds. *J. Topol.*, 12(4):1396–1412, 2019.
- [45] Stefan Schwede. Lectures on equivariant stable homotopy theory. *Available on the author’s website*, 38, 2014.
- [46] Markus Szymik. Stable diffeomorphism groups of 4-manifolds. *Math. Res. Lett.*, 15(5):1003–1016, 2008.
- [47] Markus Szymik. Characteristic cohomotopy classes for families of 4-manifolds. *Forum Math.*, 22(3):509–523, 2010.
- [48] Markus Szymik. Bauer-Furuta invariants and Galois symmetries. *Q. J. Math.*, 63(4):1033–1054, 2012.
- [49] Clifford Henry Taubes. *Seiberg Witten and Gromov invariants for symplectic 4-manifolds*, volume 2 of *First International Press Lecture Series*. International Press, Somerville, MA, 2000. Edited by Richard Wentworth.
- [50] Tammo tom Dieck. Orbittypen und äquivariante Homologie. II. *Arch. Math. (Basel)*, 26(6):650–662, 1975.
- [51] C. T. C. Wall. On simply-connected 4-manifolds. *J. London Math. Soc.*, 39:141–149, 1964.
- [52] Tadayuki Watanabe. Some exotic nontrivial elements of the rational homotopy groups of  $\text{Diff}(S^4)$ . 2019.
- [53] Ming Xu. *The Bauer-Furuta invariant and a cohomotopy refined Ruberman invariant*. ProQuest LLC, Ann Arbor, MI, 2004. Thesis (Ph.D.)—State University of New York at Stony Brook.

YAU MATHEMATICAL SCIENCES CENTER, TSINGHUA UNIVERSITY, BEIJING, CHINA.

*Email address:* linjian5477@gmail.com

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GEORGIA, USA.

*Email address:* anubhavmaths@gatech.edu