A note on multicolor Ramsey number of small odd cycles versus a large clique

Zixiang Xu^{a,*}and Gennian Ge^{a,†}

^a School of Mathematical Sciences, Capital Normal University, Beijing 100048, China.

Abstract

Let $R_k(H; K_m)$ be the smallest number N such that every coloring of the edges of K_N with k + 1 colors has either a monochromatic H in color i for some $1 \leq i \leq k$, or a monochromatic K_m in color k + 1. In this short note, we study the lower bound for $R_k(H; K_m)$ when H is C_5 or C_7 , respectively. We show that

$$R_k(C_5; K_m) = \Omega(m^{\frac{3k}{8}+1} / (\log m)^{\frac{3k}{8}+1}),$$

and

$$R_k(C_7; K_m) = \Omega(m^{\frac{2k}{9}+1} / (\log m)^{\frac{2k}{9}+1}),$$

for fixed positive integer k and $m \to \infty$. These slightly improve the previously known lower bound $R_k(C_{2\ell+1}; K_m) = \Omega(m^{\frac{k}{2\ell-1}+1}/(\log m)^{k+\frac{2k}{2\ell-1}})$ obtained by Alon and Rödl. The proof is based on random block constructions and random blowups argument.

Key words and phrases: Multicolor Ramsey problems, odd cycles *AMS subject classifications*: 05C55.

1 Introduction

For a given graph H and a positive integer k, the multicolor Ramsey number $R_k(H; K_m)$ is the smallest number N such that every edge coloring of K_N with k + 1 colors has either

^{*}e-mail: zxxu8023@qq.com.

[†]e-mail: gnge@zju.edu.cn. Research supported by the National Key Research and Development Program of China under Grant No. 2020YFA0712100, National Natural Science Foundation of China under Grant No. 11971325, and Beijing Scholars Program.

a monochromatic H in color i for some $1 \leq i \leq k$, or a monochromatic K_m in color k + 1. Obtaining good upper and lower bounds on $R_k(H; K_m)$ is a central question in Ramsey theory, and despite more than 80 years of effort, still not much is known. For example, the most well-studied case is that of k = 1 and $H = K_m$, and the theoretical bounds $2^{\frac{m}{2}} \leq R_1(K_m; K_m) \leq 2^{2m}$ were obtained by Erdős and Szekeres [9, 11] in 1940s. However, only lower term improvements [5, 15, 17] have been made to these bounds. For larger values of $k \geq 2$, even less has been known. When $H = K_m$, a recent breakthrough of Conlon and Ferber [6] yielded an exponential improvement on the lower bounds for multicolor diagonal Ramsey numbers $R_k(K_m; K_m)$ using mixed random algebraic approaches. More recently, the lower bounds for $R_k(K_m; K_m)$ were improved by Wigderson [20] and Sawin [16] via different methods.

In this paper, we mainly study the lower bound for $R_k(H; K_m)$ when H is an odd cycle $C_{2\ell+1}$. For k = 1, Erdős, Faudree, Rousseau and Schelp [10] proved that $R_1(C_\ell; K_m) = O(m^{1+\frac{1}{\ell}})$, where $t = \lceil \frac{\ell}{2} \rceil - 1$. These were later improved by Sudakov [18], who proved that $R_1(C_{2\ell+1}; K_m) = O(\frac{m^{1+1/\ell}}{\log^{1/\ell}m})$. When $\ell = 1$, the matched lower bound for $R_1(C_3, K_m) = \Theta(\frac{m^2}{\log m})$ was shown in [1, 13]. Moreover, by analyzing the C_ℓ -free process, Bohman and Keevash [3] proved that $R_1(C_\ell; K_m) = \Omega(\frac{m^{(\ell-1)/(\ell-2)}}{\log m})$. For $\ell \in \{5, 6, 7, 10\}$, the best known lower bounds for $R_1(C_\ell; K_m)$ were improved by Mubayi and Verstraëte [8] via different approaches. When $k \ge 2$, Alon and Rödl [2] proved that $R_k(C_3; K_m) = \Theta(m^{k+1} \text{poly} \log m)$ and $R_k(C_{2\ell+1}; K_m) = \Omega(m^{1+\frac{k}{2\ell-1}}/(\log m)^{k+\frac{2k}{2\ell-1}})$. For more literature on the Ramsey problems, we refer the readers to the dynamic survey [14] and the references therein. Inspired by the random homomorphisms method in the recent work [20] on multicolor diagonal Ramsey number, we prove the lower bound for $R_k(H; K_m)$ when H is C_5 or C_7 , respectively. The appropriate base graphs come from the random block construction in [8].

Theorem 1.1. For fixed positive integer k and $m \to \infty$, we have

$$R_k(C_5; K_m) = \Omega\left(\left(\frac{m}{\log m}\right)^{\frac{3k}{8}+1}\right),$$

and

$$R_k(C_7; K_m) = \Omega\left(\left(\frac{m}{\log m}\right)^{\frac{2k}{9}+1}\right)$$

These slightly improve the previously known lower bounds $R_k(C_5; K_m) = \Omega(m^{\frac{k}{3}+1} \operatorname{poly} \log m)$ and $R_k(C_7; K_m) = \Omega(m^{\frac{k}{5}+1} \operatorname{poly} \log m)$ respectively.

2 Proof of Theorem 1.1

Our first step is to construct a base graph which is C_5 -free and has no large independent set. The construction is based on a recent work of Mubayi and Verstraëte [8]. By the existence of generalized hexagons of order (q, q^3) (see [7, 12, 19]), there exists a graph G with bipartition $G = A \cup B$ of sizes $|A| = (q+1)(q^8 + q^4 + 1)$ and $|B| = (q^3 + 1)(q^8 + q^4 + 1)$ such that every vertex in A has degree $q^3 + 1$ and every vertex in B has degree q + 1. Moreover, the girth of G is at least 12, that is, there is no cycle of length less than 12 in G. For a vertex $a \in A$, we can randomly partition the neighbors $N_G(a)$ into two parts S_a and T_a uniformly, and then we form a complete bipartite graph between S_a and T_a . Furthermore, we repeat these operations for all vertices in A and finally obtain a new graph F on vertex set V(F) = B. Note that there is no multiple edge in the new graph F since G is C_4 -free. Moreover, F is C_5 -free, otherwise by definition of graph F, there will be a copy of C_{10} in G, a contraction. We also claim that there is no independent set of size $t = (1 + o(1))q^8$ with positive probability. To see this, let I be a subset of F with |I| = t. Set $t_a = |I \cap N_G(a)|$ for $a \in A$, it is easy to see the probability that $I \cap N_G(a)$ is an independent set in F is 2^{1-t_a} . Since the partitions are independently at random for different $a \in A$, the probability that I forms an independent set in F is $\prod_{a \in A} 2^{1-t_a}$.

and the expected number of independent sets of size t in F is $\binom{|F|}{t} \prod_{a \in A} 2^{1-t_a} < 1$.

Now we fix a graph F which is C_5 -free and has no independent set of size $t = (1 + o(1))q^8$. We say F' is an r-blowup of the graph F means the graph is obtained by replacing each vertex of F by an independent set of size r and each edge of F by a complete bipartite graph. Set $r = c_k q^{3(k-1)}$ and $m = d_k q^8 \log q$ for some suitable constants $c_k, d_k > 0$. We claim that the number of independent sets of size m in F' is at most $\binom{|F|}{t} \binom{tr}{m}$. To see this, since there is no independent set of size t in F, there are at most $\binom{|F|}{t}$ blocks which contain the independent sets in F'. Moreover, each vertex of the independent set in F' of size m is one of the tr vertices in these blocks since each block contains exactly r vertices.

We then show that there exists a sequence of k + 1 graphs $F_1, F_2, \ldots, F_k, F_{k+1}$ on vertex set V(F'), such that F_i is isomorphic to F' with $1 \leq i \leq k$ and F_{k+1} is the graph whose edges are all pairs of vertices of V(F') that do not lie in any F_i with $1 \leq i \leq k$. Moreover, F_{k+1} is K_m -free. To show this, we pick k random copies F_1, F_2, \ldots, F_k of F' on V(F') independently, then the graph F_{k+1} is determined by the above rules. For a fixed subset M in V(F') with |M| = m, the probability that M forms an independent set in any F_i is $\frac{\binom{|F|}{m}}{\binom{|F|}{m}}$, thus the probability that M forms a clique in F_{k+1} is at most $\left(\frac{\binom{|F|}{m}}{\binom{|F|}{m}}\right)^k$, and the expected number of K_m in F_{k+1} is at

most $\binom{|F'|}{m} \left(\frac{\binom{|F|}{t}\binom{tr}{m}}{\binom{|F'|}{m}}\right)^k < 1$ by choices of the parameters r and m. Hence, the required graph F_{k+1} exists, with positive probability.

Finally, we fix a required sequence of graphs $F_1, F_2, \ldots, F_{k+1}$ as above. Let N := r|F| and color each edge of K_N by minimum index $i \in \{1, 2, \ldots, k\}$ such that the edge belongs to F_i . Moreover, if no such *i* exists, we color the edge by color k + 1. Through the above analysis, we can see there is no C_5 in color *i* with $1 \leq i \leq k$ and no K_m in color k + 1. This gives $R_k(C_5; K_m) = \Omega\left(\left(\frac{m}{\log m}\right)^{\frac{3k}{8}+1}\right).$

We can also obtain the lower bound for $R_k(C_7; K_m)$ through the similar analysis. The required base graphs of girth at least 16 can be constructed from Ree-Tits octagons (see [7, 12, 19]), we omit the details.

3 Remarks

In [20], Wigderson pointed out that the random induced subgraphs and random blowups are closely related and are both part of a more general framework of random homomorphisms. We do agree that the framework of random homomorphisms is great, and in some sense we can regard the random induced subgraphs and the random blowups as the same thing. However, in this paper, we prefer to use the random blowups method, since our C_5 -free base graph from the random block construction has too many independent sets of small order. In diagonal Ramsey case [6, 20], the base graph is constructed algebraically, which provides a good estimation of the number of small independent sets. The random blowups method has certain limitations, for example, it cannot work when the graph H is bipartite, unless one chooses other ways to define the "blowups".

It seems to be very hard to obtain the tight asymptotic bound for $R_k(C_{2\ell+1}; K_m)$, even for the simplest case $R_1(C_5; K_m)$. The best known upper bound $R_1(C_5; K_m) = O(\frac{m^{3/2}}{\sqrt{\log m}})$ was proven in [4]. For multicolor case, we conjecture that $R_k(C_5; K_m) = O(m^{\frac{k}{2}+1} \text{poly} \log m)$ holds.

Acknowledgements

The authors thank Xin Wei for carefully reading an early draft of this paper.

References

- M. Ajtai, J. Komlós, and E. Szemerédi. A note on Ramsey numbers. J. Combin. Theory Ser. A, 29(3):354–360, 1980.
- [2] N. Alon and V. Rödl. Sharp bounds for some multicolor Ramsey numbers. Combinatorica, 25(2):125–141, 2005.
- [3] T. Bohman and P. Keevash. The early evolution of the *H*-free process. Invent. Math., 181(2):291–336, 2010.
- [4] Y. Caro, Y. Li, C. C. Rousseau, and Y. Zhang. Asymptotic bounds for some bipartite graph: complete graph Ramsey numbers. *Discrete Math.*, 220(1-3):51–56, 2000.

- [5] D. Conlon. A new upper bound for diagonal Ramsey numbers. Ann. of Math. (2), 170(2):941–960, 2009.
- [6] D. Conlon and A. Ferber. Lower bounds for multicolor Ramsey numbers. Adv. Math., 378:Paper No. 107528, 5, 2021.
- [7] B. De Bruyn. An introduction to incidence geometry. Frontiers in Mathematics. Birkhäuser/Springer, Cham, 2016.
- [8] J. V. Dhruv Mubayi. A note on pseudorandom ramsey graphs. *arXiv preprint*, arXiv: 1909.01461, 2019.
- [9] P. Erdős. Some remarks on the theory of graphs. Bull. Amer. Math. Soc., 53:292–294, 1947.
- [10] P. Erdős, R. J. Faudree, C. C. Rousseau, and R. H. Schelp. On cycle-complete graph Ramsey numbers. J. Graph Theory, 2(1):53–64, 1978.
- [11] P. Erdős and G. Szekeres. A combinatorial problem in geometry. Compositio Math., 2:463–470, 1935.
- [12] E. Govaert and H. Van Maldeghem. Some combinatorial and geometric characterizations of the finite dual classical generalized hexagons. J. Geom., 68(1-2):87–95, 2000.
- [13] J. H. Kim. The Ramsey number R(3,t) has order of magnitude $t^2/\log t$. Random Structures Algorithms, 7(3):173–207, 1995.
- [14] S. P. Radziszowski. Small Ramsey numbers. *Electron. J. Combin.*, 1:Dynamic Survey 1, 30, 1994.
- [15] A. Sah. Diagonal ramsey via effective quasirandomness. arXiv preprint, arXiv: 2005.09251, 2020.
- [16] W. Sawin. An improved lower bound for multicolor ramsey numbers and the halfmultiplicity ramsey number problem. arXiv preprint, arXiv: 2105.08850, 2021.
- [17] J. Spencer. Ramsey's theorem—a new lower bound. J. Combinatorial Theory Ser. A, 18:108–115, 1975.
- [18] B. Sudakov. A note on odd cycle-complete graph Ramsey numbers. *Electron. J. Combin.*, 9(1):Note 1, 4, 2002.

- [19] H. Van Maldeghem. Generalized polygons. Modern Birkhäuser Classics. Birkhäuser/Springer Basel AG, Basel, 1998. [2011 reprint of the 1998 original] [MR1725957].
- [20] Y. Wigderson. An improved lower bound on multicolor Ramsey numbers. Proc. Amer. Math. Soc., 149(6):2371–2374, 2021.