## On the divisibility of sums of even powers of *q*-binomial coefficients

Ji-Cai Liu and Xue-Ting Jiang

Department of Mathematics, Wenzhou University, Wenzhou 325035, PR China jcliu2016@gmail.com, xtjiang2021@foxmail.com

**Abstract.** We prove the divisibility conjecture on sums of even powers of *q*-binomial coefficients, which was recently proposed by Guo, Schlosser and Zudilin. Our proof relies on two *q*-harmonic series congruences due to Shi and Pan.

Keywords: q-binomial coefficients; q-congruences; cyclotomic polynomials

MR Subject Classifications: 11A07, 11B65, 05A10

## 1 Introduction

In 2012, Guo and Zeng [9, (1.9)] established the following divisibility result to prove two amazing supercongruences involving the Apéry numbers:

$$\sum_{k=0}^{n-1} \binom{n+k}{k}^2 \binom{n-1}{k}^2 \equiv 0 \pmod{n}.$$

$$(1.1)$$

Guo and Zeng [9, Conjecture 5.13] also conjectured a q-analogue of (1.1) as follows:

$$\sum_{k=0}^{n-1} q^{(n-k)^2} {n+k \brack k}^2 {n-1 \brack k}^2 \equiv q^{(n-1)^2} [n] \pmod{[p]_{q^{n/p}}^2}, \tag{1.2}$$

where n is any power of a prime p.

To understand (1.2), we shall recall some notation. The Gaussian *q*-binomial coefficients are defined as

$$\begin{bmatrix} n\\k \end{bmatrix} = \begin{bmatrix} n\\k \end{bmatrix}_q = \begin{cases} \frac{(q;q)_n}{(q;q)_k(q;q)_{n-k}} & \text{if } 0 \leqslant k \leqslant n, \\ 0 & \text{otherwise,} \end{cases}$$

where the q-shifted factorials are given by  $(a;q)_n = (1-a)(1-aq)\cdots(1-aq^{n-1})$  for  $n \ge 1$  and  $(a;q)_0 = 1$ . The q-integers are defined by  $[n] = [n]_q = (1-q^n)/(1-q)$ . The nth cyclotomic polynomial is given by

$$\Phi_n(q) = \prod_{\substack{1 \le k \le n \\ (n,k)=1}} (q - \zeta^k),$$

where  $\zeta$  denotes a primitive *n*th root of unity.

For polynomials  $A_1(q), A_2(q), P(q) \in \mathbb{Z}[q]$ , the q-congruence

$$A_1(q)/A_2(q) \equiv 0 \pmod{P(q)}$$

is understood as  $A_1(q)$  is divisible by P(q) and  $A_2(q)$  is coprime with P(q). In general, for rational functions  $A(q), B(q) \in \mathbb{Q}(q)$  and polynomial  $P(q) \in \mathbb{Z}[q]$ ,

$$A(q) \equiv B(q) \pmod{P(q)} \iff A(q) - B(q) \equiv 0 \pmod{P(q)}.$$

In the past few years, congruences for q-binomial coefficients as well as basic hypergeometric series attracted many experts' attention (see, for example, [1, 2, 4-7, 10-14, 17-19]).

In 2020, Gu and Guo [3, Theorem 1.2] confirmed the conjectural q-congruence (1.2) by establishing the following more general result:

$$\sum_{k=0}^{n-1} q^{(n-k)^2} {n+k \brack k}^2 {n-1 \brack k}^2 \equiv q[n] \pmod{\Phi_n(q)^2},$$
(1.3)

where n is any positive integer. They also conjectured that (1.3) holds modulo  $[n]\Phi_n(q)^2$  (see [3, Conjecture 1.3]), which is still open up to now.

Recently, Guo, Schlosser and Zudilin [8, Theorem 8] generalize (1.3) in a different form:

$$\sum_{k=0}^{n-1} q^{r(n-k)^2 + (r-1)k} {n+k \brack k}^{2r} {n-1 \brack k}^{2r} \equiv q[n] \pmod{\Phi_n(q)^2}, \tag{1.4}$$

where r is any positive integer. They pointed out that via the same method as in the proof of [9, Theorem 5.3], one can easily show that

$$\sum_{k=0}^{n-1} q^{r(n-k)^2 + (r-1)k} {n+k \brack k}^{2r} {n-1 \brack k}^{2r} \equiv 0 \pmod{[n]}.$$
 (1.5)

It follows that (1.4) is also true modulo  $[n]\Phi_n(q)$ . Guo, Schlosser and Zudilin [8, Conjecture 1] also conjectured that both (1.4) and the conjecture [3, Conjecture 1.3]) due to Gu and Guo possess the following unified generalization:

Conjecture 1.1 (Guo–Schlosser–Zudilin) Let n be a positive integer and r an arbitrary integer. Then

$$\sum_{k=0}^{n-1} q^{r(n-k)^2 + (r-1)k} {n+k \brack k}^{2r} {n-1 \brack k}^{2r}$$
$$\equiv q^{(r-1)n+1}[n] - \frac{r(2r-1)(n-1)^2 q(1-q)^2}{4} [n]^3 \pmod{[n]\Phi_n(q)^3}.$$
(1.6)

At the end of their paper, Guo, Schlosser and Zudilin [8] mentioned that Conjecture 1.1 is even still open modulo  $[n]\Phi_n(q)^2$ . The motivation of this paper is to give a positive answer to Conjecture 1.1.

**Theorem 1.2** The q-congruence (1.6) is true.

The important ingredients in the proof of Theorem 1.2 are the following two q-harmonic series congruences:

$$\sum_{j=1}^{n-1} \frac{1}{1-q^j} \equiv \frac{n-1}{2} + \frac{(n^2-1)(1-q^n)}{24} \pmod{\Phi_n(q)^2},\tag{1.7}$$

and

$$\sum_{j=1}^{n-1} \frac{1}{(1-q^j)^2} \equiv -\frac{(n-1)(n-5)}{12} \pmod{\Phi_n(q)}.$$
 (1.8)

As Pan [16, Proposition 2.1] mentioned, the q-congruence (1.7) can be easily proved by the same method as in the proof of [15, Theorem 1]. In fact, one can also prove (1.8) in a similar way as in the proof of [15, (5)]. In the next section, we shall present the detailed proof of Theorem 1.2.

## 2 Proof of Theorem 1.2

Noting that

$$(1-q^{n-j})(1-q^{n+j}) = (1-q^n)^2 - (1-q^j)^2 q^{n-j},$$

we obtain

$$\begin{bmatrix} n+k\\k \end{bmatrix} \begin{bmatrix} n-1\\k \end{bmatrix} = \frac{1}{(q;q)_k^2} \prod_{j=1}^k (1-q^{n-j})(1-q^{n+j})$$
$$= \frac{1}{(q;q)_k^2} \prod_{j=1}^k \left( (1-q^n)^2 - (1-q^j)^2 q^{n-j} \right)$$
$$\equiv (-1)^k q^{nk-\frac{k(k+1)}{2}} \left( 1 - \frac{(1-q^n)^2}{q^n} \sum_{j=1}^k \frac{q^j}{(1-q^j)^2} \right) \pmod{\Phi_n(q)^4},$$
(2.1)

where we have used the fact that  $1 - q^n \equiv 0 \pmod{\Phi_n(q)}$ .

By (2.1) and the binomial theorem, we have

$$\binom{n+k}{k}^{2r} \binom{n-1}{k}^{2r} \equiv q^{2rnk-rk(k+1)} \left(1 - \frac{2r(1-q^n)^2}{q^n} \sum_{j=1}^k \frac{q^j}{(1-q^j)^2}\right) \pmod{\Phi_n(q)^4}.$$

It follows that

$$\sum_{k=0}^{n-1} q^{r(n-k)^2 + (r-1)k} {n+k \choose k}^{2r} {n-1 \choose k}^{2r}$$
$$\equiv q^{n^2 r} \sum_{k=0}^{n-1} q^{-k} - 2rq^{n(nr-1)}(1-q^n)^2 \sum_{k=0}^{n-1} q^{-k} \sum_{j=1}^k \frac{q^j}{(1-q^j)^2} \pmod{\Phi_n(q)^4}.$$
(2.2)

It is clear that

$$q^{n^2r} \sum_{k=0}^{n-1} q^{-k} = q^{n(nr-1)+1}[n].$$
(2.3)

Furthermore, we have

$$\sum_{k=0}^{n-1} q^{-k} \sum_{j=1}^{k} \frac{q^j}{(1-q^j)^2} = \sum_{j=1}^{n-1} \frac{q^j}{(1-q^j)^2} \sum_{k=j}^{n-1} q^{-k}$$
$$= \frac{1}{q^{n-1}(q-1)} \sum_{j=1}^{n-1} \frac{q^n - q^j}{(1-q^j)^2}$$
$$= \frac{1}{q^{n-1}(q-1)} \left( \sum_{j=1}^{n-1} \frac{1}{1-q^j} - (1-q^n) \sum_{j=1}^{n-1} \frac{1}{(1-q^j)^2} \right).$$
(2.4)

Substituting (1.7) and (1.8) into the right-hand side of (2.4), we obtain

$$\sum_{k=0}^{n-1} q^{-k} \sum_{j=1}^{k} \frac{q^j}{(1-q^j)^2}$$
$$\equiv \frac{1}{q^{n-1}(q-1)} \left(\frac{n-1}{2} + \frac{(n-1)(n-3)(1-q^n)}{8}\right) \pmod{\Phi_n(q)^2}. \tag{2.5}$$

Combining (2.2), (2.3) and (2.5) gives

$$\sum_{k=0}^{n-1} q^{r(n-k)^2 + (r-1)k} {n+k \brack k}^{2r} {n-1 \brack k}^{2r} \equiv q^{n(nr-1)+1}[n] + 2rq^{n(nr-2)+1}(1-q)[n]^2 \\ \times \left(\frac{n-1}{2} + \frac{(n-1)(n-3)(1-q)[n]}{8}\right) \pmod{\Phi_n(q)^4}.$$
(2.6)

Note that

$$q^{sn} = 1 - (1 - q^{sn})$$
  
= 1 - (1 - q^n)(1 + q^n + q^{2n} + \dots + q^{(s-1)n})  
= 1 - s(1 - q^n) \pmod{\Phi\_n(q)^2}.

It follows that

$$q^{sn} = 1 - (1 - q^n)(1 + q^n + q^{2n} + \dots + q^{(s-1)n})$$
  

$$\equiv 1 - (1 - q^n)\left(s - \frac{s(s-1)(1 - q^n)}{2}\right) \pmod{\Phi_n(q)^3}$$
  

$$= 1 - s(1 - q)[n] + \frac{s(s-1)(1 - q)^2}{2}[n]^2.$$
(2.7)

Applying (2.7) to the right-hand side of (2.6) yields

$$\sum_{k=0}^{n-1} q^{r(n-k)^2 + (r-1)k} {n+k \brack k}^{2r} {n-1 \brack k}^{2r} \equiv q[n] - q(1-q)(r-1)[n]^2 - \frac{q(1-q)^2 (2n^2r^2 - n^2r - 4nr^2 + 2nr + 5r - 4)}{4} [n]^3 \pmod{\Phi_n(q)^4}.$$
(2.8)

On the other hand, by (2.7) we have

$$q^{(r-1)n+1}[n] - \frac{r(2r-1)(n-1)^2q(1-q)^2}{4}[n]^3 \equiv q[n] - q(1-q)(r-1)[n]^2 - \frac{q(1-q)^2(2n^2r^2 - n^2r - 4nr^2 + 2nr + 5r - 4)}{4}[n]^3 \pmod{\Phi_n(q)^4}.$$
 (2.9)

Finally, comparing the right-hand sides of (2.8) and (2.9) gives

$$\sum_{k=0}^{n-1} q^{r(n-k)^2 + (r-1)k} {n+k \brack k}^{2r} {n-1 \brack k}^{2r}$$
$$\equiv q^{(r-1)n+1}[n] - \frac{r(2r-1)(n-1)^2 q(1-q)^2}{4} [n]^3 \pmod{\Phi_n(q)^4}.$$
(2.10)

Then the proof of (1.6) follows from (1.5) and (2.10).

Acknowledgments. The first author was supported by the National Natural Science Foundation of China (grant 12171370).

## References

- M. El Bachraoui, On supercongruences for truncated sums of squares of basic hypergeometric series, Ramanujan J. 54 (2021), 415–426.
- [2] O. Gorodetsky, q-Congruences, with applications to supercongruences and the cyclic sieving phenomenon, Int. J. Number Theory 15 (2019), 1919–1968.
- [3] C.-Y. Gu and V.J.W. Guo, q-Analogues of two supercongruences of Z.-W. Sun, Czechoslovak Math. J. 70 (2020), 757–765.
- [4] V.J.W. Guo, A q-analogue of the (A.2) supercongruence of Van Hamme for primes  $p \equiv 1 \pmod{4}$ , Rev. R. Acad. Cienc. Exactas Fís. Nat., Ser. A Mat. 114 (2020), Art. 123.
- [5] V.J.W. Guo, Proof of a generalization of the (C.2) supercongruence of Van Hamme, Rev. R. Acad. Cienc. Exactas Fís. Nat., Ser. A Mat. 115 (2021), Art. 45.
- [6] V.J.W. Guo and J.-C. Liu, q-Analogues of two Ramanujan-type formulas for 1/π, J. Difference Equ. Appl. 24 (2018), 1368–1373.
- [7] V.J.W. Guo and M.J. Schlosser, Some q-supercongruences modulo the square and cube of a cyclotomic polynomial, Rev. R. Acad. Cienc. Exactas Fís. Nat., Ser. A Mat. 115 (2021), Art. 132.
- [8] V.J.W. Guo, M.J. Schlosser and W. Zudili, New quadratic identities for basic hypergeometric series and q-congruences, preprint (2021). Available at http://math.ecnu.edu.cn/~jwguo/maths/quad.pdf
- [9] V.J.W. Guo and J. Zeng, New congruences for sums involving Apéry numbers or central Delannoy numbers, Int. J. Number Theory 8 (2012), 2003–2016.
- [10] V.J.W. Guo and W. Zudilin, A q-microscope for supercongruences, Adv. Math. 346 (2019), 329–358.
- [11] L. Li and S.-D. Wang, Proof of a q-supercongruence conjectured by Guo and Schlosser, Rev. R. Acad. Cienc. Exactas Fís. Nat., Ser. A Mat. 114 (2020), Art. 190.
- [12] J.-C. Liu, On a congruence involving q-Catalan numbers, C. R. Math. Acad. Sci. Paris 358 (2020), 211–215.
- [13] J.-C. Liu and Z.-Y. Huang, A truncated identity of Euler and related q-congruences, Bull. Aust. Math. Soc. 102 (2020), 353–359.
- [14] J.-C. Liu and F. Petrov, Congruences on sums of q-binomial coefficients, Adv. in Appl. Math. 116 (2020), Art. 102003.
- [15] L.-L. Shi and H. Pan, A q-analogue of Wolstenholme's harmonic series congruence, Amer. Math. Monthly 114 (2007), 529–531.
- [16] H. Pan, A generalization of Wolstenholme's harmonic series congruence, Rocky Mountain J. Math. 38 (2008), 1263–1269.
- [17] C. Wang and H.-X. Ni, Some q-congruences arising from certain identities, Period. Math. Hungar., online, doi: 10.1007/s10998-021-00416-8.
- [18] X. Wang and M. Yu, Some new q-congruences on double sums, Rev. R. Acad. Cienc. Exactas Fís. Nat., Ser. A Mat. 115 (2021), Art. 9.
- [19] C. Wei, Some q-supercongruences modulo the fourth power of a cyclotomic polynomial, J. Combin. Theory Ser. A 182 (2021), Art. 105469.