# Paths With Three Blocks In Digraphs

Maidoun Mortada<sup>1</sup>, Amine El Sahili<sup>2</sup>,

Zahraa Mohsen <sup>1,3</sup>

#### Abstract

A path with three blocks P(k, l, r) is an oriented path formed by kforward arcs followed by l-backward arcs then r-forward arcs. We prove that any (2k + 1)-chromatic digraph contains a path P(1, k, 1). However the existence of P(1, l, 1) with  $l \ge k$  is established in any (k+4)-chromatic digraph. In general, we establish a quadratic bound for paths with three blocks.

## 1 Introduction

The digraphs considered here have no loops or multiple edges. An oriented graph is a digraph in which, for every two vertices x and y, at most one of (x, y) or (y, x) is an edge. The chromatic number of a digraph is the chromatic number of its underlying graph. A graph G is said to be k-critical if  $\chi(G) = k$  and  $\chi(G - v) = k - 1$  for any vertex v in V(G).

A block of a path in a digraph is a maximal directed subpath. We recall that the length of a path is the number of its edges.

Given an *n*-chromatic digraph D, which oriented paths of length n - 1 can be found in D?

Havet and Thomassé proved that any *n*-tournament contains any oriented path of length n - 1 except in three cases: the directed 3-cycle, the regular tournament on five vertices, and the Paley tournament on seven vertices; in these cases D contains no antidirected path of length n - 1 [HT00].

In general, when D is an *n*-chromatic digraph, the situation is quite different. Addario.et.al. proved that any *n*-chromatic digraph D, contains any path with two blocks of length (n - 1)[ABHT07]. Moreover, Mortada.et.al proved that any (n + 1)-chromatic digraph contains a three blocks path of length n - 1, in which two consecutive blocks are of length 1 each[MESEJ21]. Regardless Burr's result[Bur80], no more results have been established in this direction. Burr's result states that any  $(n - 1)^2$ -chromatic digraph contains any oriented tree of order n. Consequently, we easily deduce that any  $(n - 1)^2$ -chromatic digraph

<sup>&</sup>lt;sup>1</sup>Lebanese University, KALMA Laboratory, Baalbeck.

<sup>&</sup>lt;sup>2</sup>Lebanese University, KALMA Laboratory, Beirut.

<sup>&</sup>lt;sup>3</sup>Paris Diderot University, IMJ Laboratory, Paris.

contains any oriented path of order n.

In this paper we are interested in studying three blocks paths. In the second section, we improve the bound to  $\frac{3}{4}n^2$  for paths with three blocks. In the third section, we prove that any 2k + 1-chromatic digraph contains a path P(1, k, 1), for k positive integer. Moreover, we show that this becomes k + 4 if we searched for the existence of P(1, l, 1), with  $l \ge k$ .

## 2 Paths with Three Blocks in Digraphs with Large Chromatic Number

We follow in this section the same reasoning as El Sahili's result.[ES04]

Let f(n) be the smallest positive integer such that every f(n)- chromatic digraph D contains a P(k, l, r) where k, l, r are positive integers such that k + l + r = n - 1. It follows from the result of Burr [Bur80] that  $f(n) \leq (n - 1)^2$ . Given a path P, we denote by  $\overline{P}$  the path obtained from P by reversing the directions of all arcs of P.

**Lemma 2.1.** Let G be a graph containing no  $K_{2n+1}$ ,  $n \ge 2$ . Suppose that we can orient G in such a way that each vertex has in-degree at most n, then  $\chi(G) \le 2n$ .

**Definition 2.1.** Define the sequence g(m,i) for  $m \ge 4$  and  $0 \le i \le \frac{m}{2} - 1$  by:

g(m,0) = m and g(4,1) = 4 $g(m,i) = g(m-1,i-1) + 2(m-3) \text{ for } m \ge 5 \text{ and } i \ge 1.$ 

**Lemma 2.2.** Any g(m, i)-chromatic digraph D contains any path P(k, m-1-k-i, i) for k positive integer less than m-1.

*Proof.* We will proceed by induction on i.

The case is solved by Addario.et.al [ABHT07] for i = 0. The case is deduced from El Sahili [ES03], and see also [MESEJ21] for i = 1, m = 4. For  $i \ge 1$ ,  $m \ge 5$ , suppose to the contrary that D is a g(m, i)-chromatic with no P(k, m - 1 - k - i, i). Let H be the subdigraph of D induced by the vertices of out-degree in D at least m - 2. Set H' the subdigraph induced by V(D - H). We claim that H contains no path of the form P(k, m - 1 - k - i, i - 1), since else let Q =P(k, m - 1 - k - i, i - 1) in H, and let v be the end vertex of Q which is also the end vertex of the block of length i - 1. Since  $d^+(v) \ge m - 2$ , there exists a vertex  $u \in N^+(v) - Q$ , hence  $Q \cup (v, u)$  is a path P(k, m - 1 - k - i, i) in H and so in D, a contradiction. Hence by the induction hypothesis  $\chi(H) < g(m - 1, i - 1)$ . Now using Lemma 2.1, we can clearly observe that  $\chi(H') \le 2(m - 3)$ , since else H' contains a tournament of order 2m-5 and so contains P(k, m-1-k-i, i) by [Tho86], contradiction. Therefore  $\chi(D) \leq \chi(H) + \chi(H') < g(m-1, i-1) + 2(m-3) = g(m, i)$ , a contradiction.

In the same way we can prove the following lemma:

**Lemma 2.3.** Any g(m, i)-chromatic digraph D contains any path  $\overline{P}(k, m-1-k-i, i)$  for k positive integer less that m-1.

**Lemma 2.4.** For every  $m \ge 4$ , the sequence g(m, i) is increasing with respect to *i*.

 $\begin{array}{l} \textit{Proof.} \mbox{ We argue by induction on } i. \\ \textit{For } m = 4, \ g(4,1) = 4 \geq g(4,0). \\ \textit{For } m \geq 5, \ g(m,1) = g(m-1,0) + 2(m-3) = m-1 + 2(m-3) \geq m = g(m,0). \\ \textit{Suppose that } g(m,i) \geq g(m,i-1). \\ \textit{Then } g(m,i+1) = g(m-1,i) + 2(m-3) \geq g(m-1,i-1) + 2(m-3) = g(m,i). \end{array}$ 

**Theorem 2.1.**  $f(m) \le \frac{3}{4}m^2, m \ge 4.$ 

*Proof.* Since every g(m, i)-chromatic digraph contains any P(k, m - 1 - k - i, i)and any  $\overline{P}(k, m - 1 - k - i, i)$ , then  $f(m) \leq g\left(m, \frac{m}{2} - 1\right)$  for m even and  $f(m) \leq g\left(m, \frac{m-3}{2}\right)$  for m odd. For m even,

$$g\left(m, \frac{m}{2} - 1\right)$$

$$= g\left(m - 1, \left(\frac{m}{2} - 1\right) - 1\right) + 2(m - 3)$$

$$= g\left(m - 2, \left(\frac{m}{2} - 1\right) - 2\right) + 2((m - 1) - 3) + 2(m - 3)$$

$$= g\left(m - \left(\frac{m}{2} - 2\right), 1\right) + 2(m - 3) + \dots + 2\left(m - \left(\frac{m}{2} - 3\right) - 3\right)$$

$$= g\left(m - \left(\frac{m}{2} - 1\right), 0\right) + 2(m - 3) + \dots + 2\left(m - \left(\frac{m}{2} - 2\right) - 3\right)$$

$$= g\left(\frac{m}{2} + 1, 0\right) + 2\sum_{k=0}^{m/2-2} ((m - k) - 3)$$

$$= \frac{m}{2} + 1 + \frac{3m^2}{4} - \frac{7m}{2} + 4$$

$$= \frac{3}{4}m^2 - 3m + 5.$$

For m odd,

$$\begin{split} g\left(m,\frac{m-3}{2}\right) \\ &= g\left(m-1,\left(\frac{m-3}{2}\right)-1\right)+2(m-3) \\ &= g\left(m-2,\left(\frac{m-3}{2}\right)-2\right)+2((m-1)-3)+2(m-3) \\ &= g\left(m-\left(\frac{m-3}{2}-1\right),1\right)+2(m-3)+\ldots+2\left(m-\left(\frac{m-5}{2}-1\right)-3\right) \\ &= g\left(m-\left(\frac{m-3}{2}-1\right)-1,0\right)+2(m-3)+\ldots+2\left(m-\left(\frac{m-5}{2}\right)-3\right) \\ &= g\left(\frac{m+3}{2},0\right)+2\sum_{k=0}^{m/2-5/2}((m-k)-3) \\ &= \frac{m+3}{2}+\frac{3m^2}{4}-4m+\frac{21}{4} \\ &= \frac{3}{4}m^2-\frac{7m}{2}+\frac{27}{4}. \\ & \text{Hence, } f(m) \leq \frac{3}{4}m^2 \text{ for all } m. \end{split}$$

## 3 Paths with Three Blocks, P(1,k,1)

In this section, we are going to improve the bound we found in section 2 for specific forms of P(k, l, r).

**Theorem 3.1.** Any (k + 4)-chromatic digraph D contains a P(1, l, 1) for some  $l \ge k$ .

*Proof.* Let D be a (k + 4)-chromatic digraph. Suppose that D contains no P(1, l, 1) for all  $l \ge k$ . Suppose without loss of generality that D is k + 4-critical, and so  $d(v) \ge k + 3$  for all  $v \in V(D)$ . Note that D contains a P(k, 1). Let  $P = P(s, 1) = x_1 x_2 \cdots x_{s+1} y$  be such that  $s \ge k$  and s is maximal. Due to the maximality of s, all the in-neighbors of  $x_1$ , if exist, belong to P. Besides, all the out-neighbors of  $x_1$  belong to P, since else let  $z \in N^+(x_1) - P$ , so  $(x_1, z) \cup P$  is a P(1, s, 1) for  $s \ge k$ , contradiction.

Let *i* be the minimal integer such that  $x_i \in N(x_1)$  and  $i \geq 3$ . Note that  $|[x_i, x_{s+1}]| \geq |N(x_1) - \{x_2, y\}| \geq k + 1$ . If  $(x_1, x_i) \in E(D)$  then the path  $(x_1, x_2) \cup (x_1, x_i) \cup [x_i, x_{s+1}]) \cup (y, x_{s+1})$  is P(1, l, 1) for  $l \geq k$ , contradiction. Else if  $(x_i, x_1) \in E(D)$  then the path  $(x_i, x_1) \cup [x_i, x_{s+1}] \cup (y, x_{s+1})$  is P(1, l, 1) for  $l \geq k$ , contradiction.

**Conjecture 3.1.** Any (k + 3)-chromatic digraph D contains a P(1, l, 1) for some  $l \ge k$ .

The problem of existence of P(1, k, 1) in a digraph D is quite different. Define f(k) to be the smallest integer such that any f(k)-chromatic digraph contains a P(1, k, 1). Based on the existence of paths of the form P(k, 1), we may show that  $f(k) \leq 3k + 4$ . Indeed, let D be a (3k + 4)-chromatic digraph with no P(1, k, 1). Let U be the set of all origins of any P(k, 1) in D. It is clear that  $U \neq \emptyset$ . If all the vertices in U has an out-degree in D less than k + 1 then, by lemma 2.1,  $\chi(U) \leq 2k + 2$ . Then  $\chi(D - U) \geq k + 2$  and so D - U contains a P(k, 1), a contradiction. Hence there exists  $u \in U$  such that  $d^+(u) > k + 1$ . let  $P_u(k, 1)$  be a path of the form P(k, 1) in D with origin u. Let  $z \in N^+(u) - P_u(k, 1)$  and so the path  $P_u(k, 1) \cup (u, z)$  is P(1, k, 1), contradiction.

In the sequel, we improve the bound above of f(k) to 2k + 1.

#### **Theorem 3.2.** $f(k) \le 2k + 1$ .

Proof. Let D be a digraph with  $\chi(D) = 2k + 1$  with  $k \ge 2$  positive integer. Suppose without loss of generality that D is critical. We argue by the way of contradiction assuming that D has no P(1, k, 1). Let  $D_1$  be the subdigraph of D induced by the vertices of D with  $d_D^-(v) \ge k+1$  and  $D_2$  the subdigraph of Dinduced by  $V(D) - V(D_1)$ . Note that all the vertices in  $D_2$  have an out-degree in D is greater than or equal k. We either have  $\chi(D_2) \ge k$  or  $\chi(D_1) \ge k+2$ .

Suppose that  $\chi(D_2) \geq k$ . Let P be a directed path of maximal length in  $D_2$ . It's clear that  $l(P) \geq k - 1$ . We claim that  $l(P) \geq k$ . Indeed, suppose to the contrary that l(P) = k - 1. Set  $P = x_1 \cdots x_k$ . Note that  $N_{D_2}^+(x_k) \subset P$  then  $x_k$  has at least two out-neighbors in  $D_1$ , say  $a_1$  and  $a_2$ . Since  $d^+(x_1) \geq k$ , let v an out-neighbor of  $x_1$  outside P. If  $v \notin \{a_1, a_2\}$  then  $N^-(a_i) = P \cup \{v\}$  for i = 1, 2, since else let  $z \in N^-(a_i) - (P \cup \{v\})$  then  $(z, a_i) \cup (x_k, a_i) \cup P \cup (x_1, v)$  is a P(1, k, 1), contradiction. But now  $(v, a_2) \cup (x_k, a_2) \cup P \cup (x_1, a_1)$  is a P(1, k, 1), contradiction. Thus  $v \in \{a_1, a_2\}$ . Without loss of generality consider that  $v = a_2$ . Note that  $N^-(a_1) = P \cup \{v\}$  and  $N^+(x_1) \subseteq P \cup \{v\}$ . Since  $(a_2, a_1) \in D$  and  $d^-(a_2) \geq k + 1$ , then  $a_2$  has an in-neighbor outside  $P \cup \{a_1\}$ , say x. Set  $(x, a_2) \cup (x_k, a_2) \cup P \cup (x_1, a_1)$  is a P(1, k, 1), contradiction.

Set then  $P = x_1 \cdots x_{r+k}$  with  $r \ge 1$  and  $P_1 = x_r \cdots x_{r+k}$ .

We now claim that  $N^+(x_{r+k}) \cap D_1 = \emptyset$ . Suppose to the contrary that  $N^+(x_{r+k}) \cap D_1 \neq \emptyset$ . If  $|N^+(x_{r+k}) \cap D_1| \geq 2$ . Then a contradiction can be reached as before. Then  $|N^+(x_{r+k}) \cap D_1| = 1$ . Let  $a_1 \in N^+(x_{r+k}) \cap D_1$ . We claim that  $x_r \in N^-(a_1)$ . Suppose not, then there exists  $w \in N^-(a_1) - P_1$ . If  $(x_{r+k-1}, a_1) \in E(D)$ , then  $N^+(x_r) = [x_{r+1}, x_{r+k}]$  and so  $(x_r, x_{r+2}) \in E(D)$ . Consequently,  $(x_r, x_{r+1}) \cup (x_r, x_{r+2}) \cup [x_{r+2}, x_{r+k}] \cup (x_{r+k}, a_1) \cup (w, a_1)$  is a

P(1, k, 1), contradiction. Hence  $(x_{r+k-1}, a_1) \notin E(D)$  and so there exists  $w' \in N^-(a_1) - P_1 - \{w\}$ . In this case  $N^+(x_{r+1}) = [x_{r+2}, x_{r+k}] \cup \{a_1\}$ . And so  $N^+(x_r) = [x_{r+1}, x_{r+k}]$  in particular  $(x_r, x_{r+2}) \in E(D)$ , contradiction as before. Hence  $x_r \in N^-(a_1)$ . Consequently,  $N^+(x_{r+1}) = [x_{r+2}, x_{r+k}] \cup \{a_1\}$ . In particular  $(x_{r+1}, x_{r+k}) \in E(D)$ , but  $N^+(x_{r+k}) \ge k$  then there exists  $x_j \in N^+(x_{r+k})$  for  $1 \le j < r$  so  $r \ge 2$ . As before  $x_{r-1} \in N^-(a_1)$ . Then  $x_{r+k}$  has at least two out-neighbors in  $[x_1, x_r[$ , let  $x_i$  be one of them with i maximal, then  $(x_{i-1}, x_i) \cup [x_{r+1}, x_{r+k}] \cup (x_{r+k}, x_i) \cup (x_{r+1}, a_1)$  is a P(1, k, 1), contradiction.

Next, if  $(x_{r+k}, x_r) \in E(D)$  then  $(x_{r+k}, x_1) \in E(D)$ . Suppose to the contrary that  $(x_{r+k}, x_1) \notin D$ . Since  $(x_{r+k}, x_r) \in E(D)$  then  $N^+(x_{r+1}) \subseteq [x_{r+2}, x_{r+k}] \cup \{x_{r-1}\}$ , since else we get a P(1, k, 1), a contradiction. But  $d^+(x_{r+1}) \ge k$ , then  $N^+(x_{r+1}) = [x_{r+2}, x_{r+k}] \cup \{x_{r-1}\}$ . In particular,  $(x_{r+1}, x_{r+k}) \in E(D)$ . Since  $d^+(x_{r+k}) \ge k$ , then there exists 1 < i < r-1 such that  $x_i \in N^+(x_{r+1})$ . Hence  $(x_{i-1}, x_i) \cup [x_{r+1}, x_{r+k}] \cup (x_{r+k}, x_i) \cup (x_{r+1}, x_{r-1})$  is P(1, k, 1), a contradiction.

Now if  $|N^+(x_{r+k}) \cap [x_2, x_r]| \geq 3$ , then we get P(1, k, 1), contradiction. If  $|N^+(x_{r+k}) \cap [x_2, x_r]| = 2$  then so  $(x_{r+k}, x_1) \in E(D)$ , since else we get that  $(x_{r+k}, x_{r+1}) \in E(D)$ , but since  $d^+(x_{r+1}) \geq k$  then  $x_{r+1}$  has at least two out neighbors outside  $P_1$  which gives a contradiction as before. If  $|N^+(x_{r+k}) \cap$  $[x_2, x_r] \leq 1$ , then  $(x_{r+k}, x_1) \in E(D)$ . Hence  $x_i$  can play the role of  $x_{r+k}$  for every  $x_i \in V(P)$ . We claim that  $N^-(x_i) \subset P$  for every  $x_i \in V(P)$ . Otherwise, without loss of generality suppose that there exists  $z \in N^{-}(x_{r+k}) - P$ .  $z \in D_1$ , since else  $(z, x_{r+k}) \cup (x_{r+k}, x_1) \cup [x_1, x_{r+k-1}]$  is a directed path in  $D_2$  longer than P, a contradiction. Since  $N^+(x_r) \subseteq P$  then  $N^+(x_r) = [x_{r+1}, x_{r+k}]$ , otherwise we get a P(1, k, 1), in particular,  $(x_r, x_{r+2})$  and  $(x_r, x_{r+k}) \in E(D)$ . But  $d^+(x_{r+k}) \geq k$ , then there exists  $x_j \in N^+(x_{r+k})$  such that 1 < j < r. Hence  $(x_r, x_{r+1}) \cup (x_r, x_{r+2}) \cup [x_{r+2}, x_{r+k}] \cup (x_{r+k}, x_j) \cup (x_{j-1}, x_j)$  is a P(1, k, 1), a contradiction. Hence  $N^{-}(x_i) \subset P$  for every  $x_i \in V(P)$ . Since also  $N^{+}(x_i) \subset P$ and D is critical then  $V(D) = V(D_2) = V(P)$ . Since  $d^+(x_i) \ge k$  and  $d^-(x_i) \le k$ for every  $x_i \in V(P)$ , then  $d^+(x_i) = d^-(x_i) = k$  for every  $x_i \in V(P)$  and so D is a 2k-regular digraph. Since  $\chi(D) = 2k + 1$ , then D is a 2k + 1-tournament and so it contains a P(1, k, 1), a contradiction.

Consequently,  $\chi(D_2) < k$  and so  $\chi(D_1) \ge k+2$ , then let  $P = z_1 \cdots z_{k+r}$  be a directed path in  $D_1$  with maximal length in  $(r \ge 2)$  and let  $P_2 = z_1 \cdots z_{k+1}$ . We prove that  $N^-(z_1) \cap D_2 = \emptyset$ . Suppose to the contrary that  $N^-(z_1) \cap D_2 \neq \emptyset$ . Let  $a_1 \in D_2 \cap N^-(z_1)$ . Since  $d^-(z_k) \ge k+1$ , then there exists  $v \in N^-(z_k) - [z_1, z_{k+1}] - \{a_1\}$ . Note that  $d^+(a_1) \ge k$ . If there exists  $w \in N^+(a_1) - [z_1, z_k] - \{v\}$ , we get a P(1, k, 1), a contradiction. Hence  $N^+(a_1) \subseteq ([z_1, z_k] \cup \{v\})$ . If  $(a_1, z_2) \in E(D)$  then  $N^-(z_{k+1}) = [z_1, z_k] \cup \{a_1\}$  since else we get a P(1, k, 1), a contradiction. But  $N^+(a_1) \subseteq ([z_1, z_k] \cup \{v\})$ , a contradiction. And so  $(a_1, z_3) \in E(D)$ . Since  $d^-(z_{k+2}) \ge k+1$  and  $z_{k+2} \notin N^+(a_1)$ , then there exists  $x \in N^-(z_{k+2}) - ([z_3, z_k] \cup \{a_1, z_1\})$ . Hence  $(x, z_{k+2}) \cup [z_3, z_{k+2}] \cup (a_1, z_3) \cup (a_1, z_1)$ is a P(1, k, 1), contradiction.

Since  $d^{-}(z_1) \geq k+1$  then there exists  $z_{t_1}, z_{t_2} \in N^{-}(z_1)$  such that  $k < t_1 <$ 

 $t_2 < k + r$ . Also since  $d^-(z_k) \ge k + 1$  then  $|N^-(z_k) - [z_1, z_{k-1}]| \ge 2$ . Set  $I = N^-(z_k) - [z_1, z_{k-1}]$ .  $I \subseteq \{z_{t_1}, z_{t_1+1}\} \cap \{z_{t_2}, z_{t_2+1}\}$  since else we get a P(1, k, 1), a contradiction. But  $|I| \ge 2$ , and so  $|\{z_{t_1}, z_{t_1+1}\} \cap \{z_{t_2}, z_{t_2+1}\}| \ge 2$  then  $\{t_1, t_1 + 1\} = \{t_2, t_2 + 1\}$ , a contradiction.

Acknowledgment. The authors would like to acknowledge the National Council for Scientific Research of Lebanon (CNRS-L) and the Agence Universitaire de la Francophonie in cooperation with Lebanese University for granting a doctoral fellowship to Zahraa Mohsen.

### References

- [ABHT07] L. Addario-Berry, F. Havet, and S. Thomassé, Paths with two blocks in nchromatic digraphs, J. Combin. Theory Ser. B 97 (2007), no. 4, 620–626. MR2325801
  - [Bur80] S. A. Burr, Subtrees of directed graphs and hypergraphs, Congr. Numer. 28 (1980), 227–239. MR608430
  - [ES03] A. El Sahili, Functions and line digraphs, J. Graph Theory 44 (2003), no. 4, 296–303. MR2019237
  - [ES04] \_\_\_\_\_, Paths with two blocks in k-chromatic digraphs, Discrete Math. 287 (2004), no. 1-3, 151–153. MR2094709
  - [HT00] F. Havet and S. Thomassé, Oriented Hamiltonian paths in tournaments: a proof of Rosenfeld's conjecture, J. Combin. Theory Ser. B 78 (2000), no. 2, 243–273. MR1750898
- [MESEJ21] M. Mortada, A. El Sahili, and M. El Joubbeh, About paths with three blocks, Australas. J. Combin. 80 (2021), 99–105. MR4249458
  - [Tho86] A. Thomason, Paths and cycles in tournaments, Trans. Amer. Math. Soc. 296 (1986), no. 1, 167–180. MR837805