ALMOST COHERENT MODULES AND ALMOST COHERENT SHEAVES

BOGDAN ZAVYALOV

ABSTRACT. We review the theory of almost coherent modules that was introduced in "Almost Ring Theory" [GR03] by Gabber and Ramero. Then we globalize it by developing a new theory of almost coherent sheaves on schemes and on a class of "nice" formal schemes. We show that these sheaves satisfy many properties similar to usual coherent sheaves, i.e. the Almost Proper Mapping Theorem, the Formal GAGA, etc. We also construct an almost version of the Grothendieck twisted image functor $f^!$ and verify its properties. Lastly, we study sheaves of p-adic nearby cycles on admissible formal models of rigid spaces and show that these sheaves provide examples of almost coherent sheaves. This gives a new proof of the finiteness result for étale cohomology of proper rigid spaces obtained before in the work of Peter Scholze "p-adic Hodge Theory For Rigid-Analytic Varities" [Sch13].

Contents

1. Introduction	3
1.1. Motivation	3
1.2. Foundations of Almost Mathematics (Sections 2-5)	4
1.3. p -adic Nearby Cycles Sheaves (Section 6)	8
1.4. Acknowledgements	9
1.5. Notation	10
2. Almost Commutative Algebra	11
2.1. The Category of Almost Modules	11
2.2. Basic Functors on the Categories of Almost Modules	15
2.3. Derived Category of Almost Modules	18
2.4. Basic Functors on the Derived Categories of Almost Modules	19
2.5. Almost Finitely Generated and Almost Finitely Presented Modules	24
2.6. Almost Coherent Modules and Almost Coherent Rings	34
2.7. Base Change for Almost Modules	39
2.8. Almost Finitely Generated Modules over Adhesive Rings	44
3. Almost Mathematics on Ringed Sites	46
3.1. The Category of \mathcal{O}_X^a -modules	46
3.2. Basic Functors on the Category Of \mathcal{O}_X^a -Modules	50
3.3. The Projection Formula	57
3.4. Derived Category of \mathcal{O}_X^a -Modules	60
3.5. Basic Functors on the Derived Categories of \mathcal{O}_X^a -modules	61
4. Almost Coherent Sheaves on Schemes and Formal Schemes	66
4.1. Schemes. The Category of Almost Coherent \mathcal{O}_X^a -modules	66
4.2. Schemes. Basic Functors on Almost Coherent \mathcal{O}_X^a -modules	72
4.3. Schemes. Derived Category of Almost Coherent \mathcal{O}_X^a -modules	75
4.4. Formal Schemes. The Category of Almost Coherent $\mathcal{O}^a_{\mathfrak{X}}$ -modules	80
4.5. Formal Schemes. Basic Functors on Almost Coherent $\mathcal{O}^a_{\mathfrak{X}}$ -modules	87

4.6. Formal Schemes. Derived Category of Almost Coherent $\mathcal{O}^a_{\mathfrak{r}}$ -modules	90
4.7. Formal Schemes. Basic Functors on the Derived Categories of $\mathcal{O}_{\mathfrak{r}}^{a}$ -modules	93
5. Cohomological Properties of Almost Coherent Sheaves	97
5.1. Almost Proper Mapping Theorem	97
5.2. Characterization of Quasi-Coherent, Almost Coherent Complexes	102
5.3. The GAGA Theorem	103
5.4. The Formal Function Theorem	109
5.5. Almost Version of Grothendieck Duality	110
6. Almost Coherence of " <i>p</i> -adic Nearby Cycles"	113
6.1. Introduction	113
6.2. The Complex $\mathbf{R}\nu_*(\mathbf{L}\otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$ is Almost Quasi-Coherent and Commutes with Ét	ale
Base Change	115
6.3. The Complex $\mathbf{R}\nu_*(\mathbf{L}\otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$ is Almost Coherent for Smooth X	121
6.4. Proof of Theorem 6.1.1 for $\mathbf{R}\nu_*(\mathbf{L}\otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$	127
6.5. Proof of Theorem 6.1.1 for $\mathbf{R}\nu_*(\mathbf{L}\otimes_{\widehat{\mathbf{Z}}_n}\widehat{\mathcal{O}}_X^+)$	136
Appendix	142
Appendix A. Complexes over Adhesive Rings	142
Appendix B. Achinger's Result in the Non-Noetherian Case	145
Appendix C. Locally Constant and Lisse Sheaves on the Pro-Étale Topology	149
References	153

1. INTRODUCTION

1.1. Motivation. The purpose of this paper is twofold. The first goal is to develop a sufficiently rich theory of almost coherent sheaves on schemes and a class of formal schemes. The second goal is to provide the reader with one interesting source of examples of almost coherent sheaves. Namely, we show that the complex of *p*-adic nearby cycles $\mathbf{R}\nu_*(\mathcal{O}_X^+/p)$ has almost coherent cohomology sheaves for any admissible formal \mathcal{O}_C -scheme \mathfrak{X} with generic fiber $X = \mathfrak{X}_C$.

Before we discuss the content of each chapter in detail, we explain the motivation behind the work done in this manuscript.

Scholze showed in [Sch13] that there is an almost isomorphism

$$\mathrm{H}^{i}(X, \mathbf{F}_{p}) \otimes \mathcal{O}_{C}/p \simeq^{a} \mathrm{H}^{i}(X, \mathcal{O}_{X}^{+}/p)$$

for any proper rigid-analytic variety X over a p-adic algebraically closed field C. This almost isomorphism allows us to reduce studying certain properties of $\mathrm{H}^{i}(X, \mathbf{F}_{p})$ for a p-adic proper rigidanalytic space X to studying almost properties of the cohomology groups $\mathrm{H}^{i}(X, \mathcal{O}_{X}^{+}/p)$, or the full complex $\mathbf{R}\Gamma(X, \mathcal{O}_{X}^{+}/p)$. For instance, Scholze shows that $\mathrm{H}^{i}(X, \mathbf{F}_{p})$ are finite groups by deducing it from almost coherence of $\mathrm{H}^{i}(X, \mathcal{O}_{X}^{+}/p)$ over \mathcal{O}_{C}/p .

Scholze's argument does not involve any choice of an admissible formal model for X and is performed entirely on the generic fiber. However, a different natural approach to studying $\mathbf{R}\Gamma(X, \mathcal{O}_X^+/p)$ is to rewrite this complex as

$$\mathbf{R}\Gamma(X, \mathcal{O}_X^+/p) \simeq \mathbf{R}\Gamma(\mathfrak{X}, \mathbf{R}\nu_*\mathcal{O}_X^+/p)$$

for a choice of an admissible formal \mathcal{O}_C -model \mathfrak{X} and the natural morphism of ringed sites

$$\nu \colon (X_{\text{pro\acute{e}t}}, \mathfrak{O}_X^+/p) \to (\mathfrak{X}_{0,\text{Zar}}, \mathfrak{O}_{\mathfrak{X}_0})$$

with \mathfrak{X}_0 the mod-*p* fiber of \mathfrak{X} . Then we can separately study the complex $\mathbf{R}\nu_*(\mathfrak{O}_X^+/p)$ and the functor $\mathbf{R}\Gamma(\mathfrak{X}, -)$. In order to make this strategy work, we develop the notion of almost coherent sheaves on \mathfrak{X} and \mathfrak{X}_0 and show its various properties similar to the properties of coherent sheaves. This occupies Chapters 2-5. Chapter 6 is devoted to showing that the complex $\mathbf{R}\nu_*(\mathfrak{O}_X^+/p)$ has almost coherent cohomology groups. Combining it with the Almost Proper Mapping Theorem 1.2.6, we reprove [Sch13, Lemma 5.8 and Theorem 5.1]

Theorem 1.1.1. (Corollary 6.1.3) Let X be a proper rigid-analytic variety over an algebraically closed field, complete, rank-1 valuation field C of mixed characteristic (0, p). Let **L** be an \mathbf{F}_p -local system on X. Then $\mathrm{H}^i(X, \mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$ are almost finitely generated over \mathcal{O}_C/p , and $\mathrm{H}^i(X, \mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p) \simeq^a 0$ for $i > 2 \dim X$.

Theorem 1.1.2. (Corollary 6.1.4) Let X, L, and C be as above. Then $\mathrm{H}^{i}(X, \mathbf{L})$ are finite, and $\mathrm{H}^{i}(X, \mathbf{L}) \simeq 0$ for $i > 2 \dim X$.

Similar ideas are used in our upcoming paper [Zav21] to establish Poincaré duality for \mathbf{F}_p local systems on smooth and proper rigid spaces over non-archimedean fields of mixed characteristic (0, p). In particular, that proof extensively uses the almost version of Grothendieck Duality and almost coherentness of $\mathbf{R}\nu_*(\mathcal{O}_X^+/p)$ established in this paper.

That being said, we now discuss content and the main results of each Section in more detail.

1.2. Foundations of Almost Mathematics (Sections 2-5). Section 2.1 is devoted to defining the category of almost modules and studying its main properties. This section is very motivated by [GR03]. However, it seems that some results that we need later in the paper are not present in [GR03], so we give an (almost) self-contained introduction to almost commutative algebra. We define the notions of the category of almost modules (see the discussion after Corollary 2.1.4), their tensor products (see Proposition 2.2.1(1)), almost Hom functor alHom_{R^a}(-, -) (see Proposition 2.2.1(3)), almost finitely generated (see Definitions 2.5.1), almost finitely presented (see Definition 2.5.2), and almost coherent modules (see Definition 2.6.1). We show that almost coherent modules satisfy most natural properties similar to the properties of classical coherent modules. We summarize some of them in the theorem below:

Theorem 1.2.1. (Lemma 2.6.7, Propositions 2.6.14, 2.6.15, 2.6.16, and Lemma 2.7.6) Let R be a ring with an ideal \mathfrak{m} such that $\widetilde{\mathfrak{m}} := \mathfrak{m} \otimes_R \mathfrak{m}$ is R-flat and $\mathfrak{m}^2 = \mathfrak{m}$.

- (1) Almost coherent R^a -modules form a Weak Serre subcategory of \mathbf{Mod}_R .
- (2) If R is an almost coherent ring (i.e. free rank-1 R-module is almost coherent), and M^a, N^a two objects in $\mathbf{D}^-_{acoh}(R)^a$. Then $M^a \otimes_{R^a}^L N^a \in \mathbf{D}^-_{acoh}(R)^a$.
- (3) If R is an almost coherent ring, and $M^a \in \mathbf{D}^-_{acoh}(R)^a$, $N^a \in \mathbf{D}^+_{acoh}(R)^a$. Then

$$\mathbf{R}$$
al $\operatorname{Hom}_{R^a}(M^a, N^a) \in \mathbf{D}^+_{acoh}(R)^a$

- (4) If R is an almost coherent ring, $M^a \in \mathbf{D}^-_{acoh}(R)^a$, $N^a \in \mathbf{D}^+(R)^a$, and P^a an almost flat R^a -module. Then the natural map $\mathbf{R}\operatorname{Hom}_{R^a}(M^a, N^a) \otimes_{R^a} P^a \to \mathbf{R}\operatorname{Hom}_{R^a}(M^a, N^a \otimes_{R^a} P^a)$ is an almost isomorphism.
- (5) Let $R \to S$ be a faithfully flat map, and let M^a be an R^a -module. Suppose that $M^a \otimes_{R^a} S^a$ is almost finitely generated (resp. almost finitely presented, resp. almost coherent) S^a -module. Then so is M^a .

We discuss the extension of almost mathematics to ringed sites in Section 3. The main goal is to generalize all constructions from almost mathematics to a general ringed site. We define two notions of almost \mathcal{O}_X -modules on a ringed site (X, \mathcal{O}_X) (see Definition 3.1.9) and of \mathcal{O}_X^a -modules (see Definition 3.1.10) and show that these notions are equivalent:

Theorem 1.2.2. (Theorem 3.1.19) Let R be as in Theorem 1.2.1 and (X, \mathcal{O}_X) a ringed R-site. Then the functor

$$(-)^a \colon \mathbf{Mod}^a_{\mathcal{O}_X} \to \mathbf{Mod}_{\mathcal{O}^a_X}$$

is an equivalence of categories.

We also define the functors $-\otimes -$, $\operatorname{Hom}_{\mathcal{O}_X^a}(-,-)$, $\operatorname{alHom}_{\mathcal{O}_X^a}(-,-)$, $\underline{\mathcal{Hom}}_{\mathcal{O}_X^a}(-,-)$, $\underline{al\mathcal{Hom}}_{\mathcal{O}_X^a}(-,-)$, f_* , and f^* on the category of \mathcal{O}_X^a -modules. We refer to Section 3.2 for an extensive discussion of these functors. Then we study the derived category of \mathcal{O}_X^a -modules and derived analogues of the functors mentioned above. This is done in Sections 3.4 and 3.5.

We develop the theory of almost finitely presented and almost (quasi-)coherent sheaves on schemes and a class of formal schemes in Section 4.1. The main goal is to show that these sheaves behave similarly to the classical coherent sheaves in many aspects.

We roughly define almost finitely presented \mathcal{O}_X^a -modules as modules such that, for any finitely generated sub-ideal $\mathfrak{m}_0 \subset \mathfrak{m}$, can be locally approximated by finitely presented \mathcal{O}_X -modules up to modules annihilated by \mathfrak{m}_0 (see Definition 4.1.3 for a precise definition). Sections 4.1-4.3 are mostly concerned with local properties of these sheaves. We summarize some of the main results below:

Theorem 1.2.3. (Corollary 4.1.12, Theorem 4.3.6, Lemmas 4.3.8, 4.3.7, 4.3.9, and 4.3.10) Let R be a ring with an ideal \mathfrak{m} such that $\widetilde{\mathfrak{m}} := \mathfrak{m} \otimes_R \mathfrak{m}$ is R-flat and $\mathfrak{m}^2 = \mathfrak{m}$.

- (1) For any *R*-scheme *X*, almost coherent \mathcal{O}_X^a -modules form a Weak Serre subcategory of $\mathbf{Mod}_{\mathcal{O}_Y}^a$.
- (2) The functor

$$(-): \mathbf{D}_*(R)^a \to \mathbf{D}_{aqc,*}(\operatorname{Spec} R)^a$$

is a *t*-exact equivalence of triangulated categories for $* \in \{", a \text{coh}\}$. Its quasi-inverse is given by $\mathbf{R}\Gamma(\operatorname{Spec} R, -)$. In particular, an almost quasi-coherent $\mathcal{O}^a_{\operatorname{Spec} R}$ -module \mathcal{F}^a is almost coherent if and only if $\mathcal{F}^a(\operatorname{Spec} R)$ is an almost coherent R^a -module.

- (3) The natural morphism $\widetilde{M^a \otimes_{R^a}^L N^a} \to \widetilde{M^a} \otimes_{\mathcal{O}^a_{\operatorname{Spec} R}}^L \widetilde{N^a}$ is an isomorphism for any $M^a, N^a \in \mathbf{D}(R)^a$.
- (4) Let that $f: \operatorname{Spec} B \to \operatorname{Spec} A$ is an *R*-morphism of affine schemes. Then $\operatorname{\mathbf{L}} f^*(\widetilde{M^a})$ is functorially isomorphic to $\widetilde{M^a \otimes_{A^a}^L B^a}$ for any $M^a \in \mathbf{D}(A)^a$.
- (5) Let $f: X \to Y$ be a quasi-compact and quasi-separated morphism of R-schemes. Suppose that Y is quasi-compact. Then $\mathbf{R}f_*$ carries $\mathbf{D}^*_{aqc}(X)^a$ to $\mathbf{D}^*_{aqc}(Y)^a$ for any $* \in \{", -, +, b\}$.
- (6) Suppose that R is almost coherent. Then the natural maps

$$\begin{aligned} \mathbf{R} & \operatorname{alHom}_{R^{a}}(M^{a}, N^{a}) \to \mathbf{R} \underline{al\mathcal{H}om}_{\mathcal{O}_{\operatorname{Spec} R}^{a}}(\widetilde{M^{a}}, \widetilde{N^{a}}), \\ & \widetilde{\mathbf{R}} \operatorname{Hom}_{R^{a}}(M^{a}, N^{a}) \to \mathbf{R} \underline{\mathcal{H}om}_{\mathcal{O}_{\operatorname{Spec} R}^{a}}(\widetilde{M^{a}}, \widetilde{N^{a}}) \end{aligned}$$

are almost isomorphisms for $M^a \in \mathbf{D}^-_{acoh}(R)^a$, $N^a \in \mathbf{D}^+(R)^a$.

We also establish one non-trivial global result on almost finitely presented \mathcal{O}_X^a -modules. Namely, we show that even though the definition of almost finitely presented \mathcal{O}_X^a -modules is local, we can find good approximations by finitely presented \mathcal{O}_X -modules globally under some mild assumption on X. This result is systematically used in Chapter 5 to get global properties of almost coherent \mathcal{O}_X^a -modules.

Theorem 1.2.4. (Lemma 4.1.8) Let X be a quasi-compact and quasi-separated R-scheme, and let \mathcal{F} be an almost quasi-coherent \mathcal{O}_X -module. Then \mathcal{F} is almost finitely presented if and only if for any finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$ there is a morphism $f : \mathcal{G} \to \mathcal{F}$ such that \mathcal{G} is a quasi-coherent finitely presented \mathcal{O}_X -module, $\mathfrak{m}_0(\ker f) = 0$ and $\mathfrak{m}_0(\operatorname{Coker} f) = 0$.

We now discuss the content of Sections 4.4-4.7. The main goal there is to prove analogues of the results in Theorem 1.2.3 for a class of formal schemes. In order to achieve this we restrict our attention to the class of topologically finitely presented schemes over a topologically universally adhesive ring R (see Setup 4.4.1). This, in particular, includes admissible formal schemes over a mixed characteristic, *p*-adically complete rank-1 valuation ring \mathcal{O}_C with algebraically closed fraction field C.

One of the main difficulties in developing a good theory of almost coherent $\mathcal{O}^a_{\mathfrak{X}}$ -modules on a formal scheme \mathfrak{X} is that there is no good abelian theory of "quasi-coherent" on \mathfrak{X} . This was an important auxiliary tool used in developing the theory of almost coherent sheaves on schemes that does not have an immediate counterpart in the world of formal schemes.

We overcome this issue in two different ways: we use the notion of adically quasi-coherent $\mathcal{O}_{\mathfrak{X}}$ -modules introduced in [FK18] (see Definition 4.4.2) and the notion of derived quasi-coherent

 $\mathcal{O}_{\mathfrak{X}}$ -modules introduced in [Lur20] (see Definition 4.6.1). The first notion has the advantage that every adically quasi-coherent $\mathcal{O}_{\mathfrak{X}}$ -module is an actual $\mathcal{O}_{\mathfrak{X}}$ -module, but these modules do not form a Weak Serre subcategory inside $\mathbf{Mod}_{\mathcal{O}_{\mathfrak{X}}}$, so they are not always very useful in practice. The latter definition has the advantage that derived quasi-coherent $\mathcal{O}_{\mathfrak{X}}$ -modules form a triangulated subcategory inside $\mathbf{D}(\mathfrak{X})$, it is quite convenient for certain purposes. However, derived quasicoherent $\mathcal{O}_{\mathfrak{X}}$ -modules are merely objects of $\mathbf{D}(\mathfrak{X})$ and not actual $\mathcal{O}_{\mathfrak{X}}$ -modules in the classical sense. Therefore, we usually use adically quasi-coherent $\mathcal{O}_{\mathfrak{X}}$ -modules when needed except for Section 4.6, where the notion of derived quasi-coherent $\mathcal{O}_{\mathfrak{X}}$ -modules seems to be more useful for our purposes. In particular, it allows us to define the functor

$$(-)^{\Delta} \colon \mathbf{D}_{acoh}(A)^{a} \to \mathbf{D}_{acoh}(\operatorname{Spf} A)^{a}$$

for any topologically finitely presented *R*-algebra *A* in a way that "extends" the classical functor $(-)^{\Delta}$: $\mathbf{Mod}_{A}^{\mathrm{acoh}} \to \mathbf{Mod}_{\mathcal{O}_{\mathfrak{X}}}$ (see Definition 4.6.7 and Lemma 4.6.13).

Theorem 1.2.5. (Lemma 4.4.23, Corollary 4.6.16, Lemmas 4.7.4, 4.7.3, 4.7.4) Let R be a ring with a finitely generated ideal I such that R is I-adically complete, I-adically topologically universally adhesive, I-torsion free with an ideal \mathfrak{m} such that $I \subset \mathfrak{m}, \mathfrak{m}^2 = \mathfrak{m}$ and $\widetilde{\mathfrak{m}} := \mathfrak{m} \otimes_R \mathfrak{m}$ is R-flat.

- (1) For any topologically finitely presented formal *R*-scheme \mathfrak{X} , almost coherent $\mathcal{O}_{\mathfrak{X}}^{a}$ -modules form a Weak Serre subcategory of $\mathbf{Mod}_{\mathcal{O}_{\mathfrak{X}}}^{a}$.
- (2) The functor

$$\mathbf{R}\Gamma(\operatorname{Spf} R, -) \colon \mathbf{D}_{acoh}(\operatorname{Spf} R)^a \to \mathbf{D}_{acoh}(R)^a$$

is a *t*-exact equivalence of triangulated categories.

- (3) The natural morphism $(M^a \otimes_{R^a}^L N^a)^{\Delta} \to (M^a)^{\Delta} \otimes_{\mathcal{O}^a_{\text{Spf }R}}^L (N^a)^{\Delta}$ is an isomorphism for any for any $M^a, N^a \in \mathbf{D}_{acoh}(R)^a$.
- (4) Let that $\mathfrak{f} \colon \operatorname{Spf} B \to \operatorname{Spf} A$ be a morphism of topologically finitely presented affine formal R-schemes. Then $\mathbf{L}\mathfrak{f}^*\left((M^a)^{\Delta}\right)$ is functorially isomorphic to $(M^a \otimes_{A^a}^L B^a)^{\Delta}$ for any $M^a \in \mathbf{D}_{acoh}(A)^a$.
- (5) The natural map

$$(\mathbf{R}alHom_{R^{a}}(M^{a}, N^{a}))^{\Delta} \to \mathbf{R}\underline{alHom}_{\mathcal{O}_{\mathrm{Spf}\,R}^{a}}\left((M^{a})^{\Delta}, (N^{a})^{\Delta}\right),$$
$$(\mathbf{R}Hom_{R^{a}}(M^{a}, N^{a}))^{\Delta} \to \mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_{\mathrm{Spf}\,R}^{a}}\left((M^{a})^{\Delta}, (N^{a})^{\Delta}\right)$$

are almost isomorphisms for $M^a \in \mathbf{D}^-_{acoh}(R)^a$, $N^a \in \mathbf{D}^+_{acoh}(R)^a$.

We discuss global properties of almost coherent sheaves in Chapter 5. Namely, we generalize certain cohomological properties of classical coherent sheaves to the case of almost coherent sheaves. We start with the almost version of the Proper Mapping Theorem:

Theorem 1.2.6. (Theorem 5.1.3) Let R be a universally coherent¹ ring with an ideal \mathfrak{m} such that $\widetilde{\mathfrak{m}} := \mathfrak{m} \otimes_R \mathfrak{m}$ is R-flat and $\mathfrak{m}^2 = \mathfrak{m}$. And let $f: X \to Y$ be a proper morphism of finitely presented R-schemes with quasi-compact Y. Then $\mathbf{R}f_*$ carries $\mathbf{D}^*_{acoh}(X)^a$ to $\mathbf{D}^*_{acoh}(Y)^a$ for $* \in \{ ", -, +, b \}$.

The essential idea of the proof is to reduce Theorem 1.2.6 to the classical Proper Mapping Theorem over an universally coherent base [FK18, Theorem I.8.1.3]. The key input to make this reduction work is Theorem 1.2.4.

¹Any finitely presented R-algebra A is coherent

7

We also prove a version of the Almost Proper Mapping Theorem for a morphism of formal schemes. We do not know if an analogue of Theorem 1.2.4 holds for coherent modules on a formal scheme. So we proceed in a different way and reduce the formal version of the Almost Proper Mapping Theorem to the schematic version stated above. For a technical reason, we need an extra assumption on the ideal of almost mathematics. Namely, we need to assume that $\mathfrak{m} = \bigcup_{n=1}^{\infty} (\varpi^{1/n})$ for a non-zero divisor $\varpi \in R$. We expect this assumption to be an artifact of our proof, but we were not able to eliminate it. We note that this condition is satisfied in our main case of interest: $R = \mathcal{O}_C$ for a complete, rank-1 valuation ring of mixed characteristic (0, p) with algebraically closed fraction field C. In this case $\mathfrak{m} = \bigcup_{n=1}^{\infty} (p^{1/n})$.

Theorem 1.2.7. (Theorem 5.1.6) Let R be a ring with a finitely generated ideal I such that R is I-adically complete, I-adically topologically universally adhesive an ideal \mathfrak{m} such that $I \subset \mathfrak{m} = \bigcup_{n=1}^{\infty} (\varpi^{1/n})$ for a non-zero divisor $\varpi \in R$, $\mathfrak{m}^2 = \mathfrak{m}$ and $\widetilde{\mathfrak{m}} := \mathfrak{m} \otimes_R \mathfrak{m}$ is R-flat. And let $\mathfrak{f} : \mathfrak{X} \to \mathfrak{Y}$ be a proper morphism of finitely presented formal R-schemes with quasi-compact Y. Then $\mathbf{R}\mathfrak{f}_*$ carries $\mathbf{D}^*_{acoh}(\mathfrak{X})^a$ to $\mathbf{D}^*_{acoh}(\mathfrak{Y})^a$ for $* \in \{ ", -, +, b \}$.

Then we provide a characterization of quasi-coherent, almost coherent complexes on finitely presented, separated schemes over a universally coherent base ring R. This is an almost analogue of [Sta21, Tag 0CSI]. We follow the same proof strategy but adjust it in certain places to make it work in the almost setting. This result is important for us as it will later play a crucial role in the proof of the Formal GAGA Theorem for almost coherent sheaves.

Theorem 1.2.8. (Theorem 5.2.3) Let R be a universally coherent² ring with an ideal \mathfrak{m} such that $\widetilde{\mathfrak{m}} := \mathfrak{m} \otimes_R \mathfrak{m}$ is R-flat and $\mathfrak{m}^2 = \mathfrak{m}$. And let X be a separated, finitely presented R-scheme. Let $\mathcal{F} \in \mathbf{D}^-_{ac}(X)$ be an object such that

$$\mathbf{R}\operatorname{Hom}_X(\mathcal{P},\mathcal{F}) \in \mathbf{D}^-_{acoh}(R)$$

for any $\mathcal{P} \in \operatorname{Perf}(X)$. Then $\mathcal{F} \in \mathbf{D}^{-}_{ac.acoh}(X)$.

Theorem 1.2.9. (Corollary 5.3.3) Let R be as in Theorem 1.2.7, and X a finitely presented R-scheme. Then the functor

$$\mathbf{L}c^*: \mathbf{D}^*_{acoh}(X)^a \to \mathbf{D}^*_{acoh}(\mathfrak{X})^c$$

induces an equivalence of categories for $* \in \{", ", +, -, b\}$.

We note that the standard proof of the classical formal GAGA theorem via projective methods has no chance to work in the almost coherent situation (due to a lack of "finiteness" for almost coherent sheaves). Instead, we "explicitly" construct a pseudo-inverse to $\mathbf{L}c^*$ in the derived world by adapting an argument from the paper of J. Hall [Hal18].

The last thing we discuss in Section 5 is the almost version of the Grothendieck Duality. This is an important technical tool in our proof of Poincaré Duality in [Zav21]. So we develop some foundations of the $f^!$ functor in the almost world in this manuscript. We summarize the main properties of this functors below:

Theorem 1.2.10. (Theorem 5.5.8) Let R be as in Theorem 1.2.6, and FPS_R be the category of finitely presented, separated R-schemes. Then there is a well-defined functor (-)! from FPS_R into the 2-category of categories such that

(1) $(X)! = \mathbf{D}^+_{aqc}(X)^a$,

(2) for a smooth morphism $f: X \to Y$ of pure relative dimension $d, f' \simeq \mathbf{L} f^*(-) \otimes_{\mathcal{O}_X^d}^L \Omega^d_{X/Y}[d]$.

²Any finitely presented R-algebra A is coherent

(3) for a proper morphism $f: X \to Y$, $f^!$ is right adjoint to $\mathbf{R}f_*: \mathbf{D}^+_{acoh}(X)^a \to \mathbf{D}^+_{acoh}(Y)^a$.

1.3. *p*-adic Nearby Cycles Sheaves (Section 6). The main goal of Section 6 is to give the main non-trivial example of almost coherent sheaves. These are the so-called *p*-adic nearby cycles sheaves.

We fix a complete, rank-1 valuation ring C of mixed characteristic (0, p) with algebraically closed fraction field C, and a rigid space X over C with an admissible formal \mathcal{O}_C -model \mathfrak{X} .

The rigid space X comes with a morphism of ringed sites

$$\nu \colon (X_{\text{pro\acute{e}t}}, \widehat{\mathbb{O}}_X^+) \to (\mathfrak{X}_{\text{Zar}}, \mathbb{O}_{\mathfrak{X}})$$

and

$$\nu \colon (X_{\text{pro\acute{e}t}}, \mathfrak{O}_X^+/p) \to (\mathfrak{X}_{0,\text{Zar}}, \mathfrak{O}_{\mathfrak{X}_0})$$

where \mathfrak{X}_0 is the mod-*p* fiber of \mathfrak{X} . The sheaves $\mathrm{R}^i \nu_* (\mathfrak{O}_X^+/p)$ are called the *p*-adic nearby cycles sheaves. These sheaves turn out to be almost coherent $\mathcal{O}_{\mathfrak{X}_0}$ -modules. Moreover, the same holds for $\mathrm{R}^i \nu_* (\mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p)$ for any étale \mathbf{F}_p local system \mathbf{L} . This is the crucial step in proving Theorem 1.1.1.

Theorem 1.3.1. (Theorem 6.1.1) Let \mathcal{O}_C , X, and \mathfrak{X} be as above. Suppose that dim X = d. Then

$$\mathbf{R}\nu_*(\mathbf{L}\otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p) \in \mathbf{D}_{acoh}^{[0,2d]}(X_0) \text{ and } \mathbf{R}\nu_*(\mathbf{L}\otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p)^a \in \mathbf{D}_{acoh}^{[0,d]}(X_0)^a$$

for an \mathbf{F}_p -local system **L**. More precisely, for an affine admissible $\mathfrak{X} = \operatorname{Spf} A$ with the adic generic fiber X, the natural maps

$$\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\mathbf{F}_{p}} \mathfrak{O}_{X}^{+}/p) \to \mathrm{R}^{i}\nu_{*}(\mathbf{L} \otimes_{\mathbf{F}_{p}} \mathfrak{O}_{X}^{+}/p)$$

are almost isomorphisms.

Moreover, the formation of $\mathrm{R}^{i}\nu_{*}(\mathbf{L}\otimes_{\mathbf{F}_{p}} \mathcal{O}_{X}^{+}/p)$ almost commute with étale base change, i.e., for any étale morphism $\mathfrak{f}: \mathfrak{X} \to \mathfrak{Y}$, the natural morphism

$$\mathfrak{f}^*\left(\mathrm{R}^i\nu_{\mathfrak{X}_C,*}(\mathbf{L}\otimes_{\mathbf{F}_p}\mathfrak{O}^+_{\mathfrak{X}_C}/p)\right)\to\mathrm{R}^i\nu_{\mathfrak{Y}_C,*}(\mathfrak{f}^*_C\mathbf{L}\otimes_{\mathbf{F}_p}\mathfrak{O}^+_{\mathfrak{Y}_C}/p)$$

is an almost isomorphism.

We also show the integral version of Theorem 1.3.1:

Theorem 1.3.2. (Theorem 6.1.1) Let \mathcal{O}_C , X, and \mathfrak{X} be as above. Suppose that dim X = d. Then

$$\mathbf{R}\nu_*(\mathbf{L}\otimes_{\mathbf{Z}_p}\widehat{\mathcal{O}}_X^+)\in\mathbf{D}_{acoh}^{[0,2d]}(\mathfrak{X})\text{ and }\mathbf{R}\nu_*(\mathbf{L}\otimes_{\mathbf{Z}_p}\widehat{\mathcal{O}}_X^+)^a\in\mathbf{D}_{acoh}^{[0,d]}(\mathfrak{X})^a$$

for a lisse $\widehat{\mathbf{Z}}_p$ sheaf **L** (see Definition C.7). More precisely, for an affine admissible $\mathfrak{X} = \operatorname{Spf} A$ with the adic generic fiber X, the natural maps

$$\mathrm{H}^{i}(X,\mathbf{L}\otimes_{\widehat{\mathbf{Z}}_{p}}\widehat{\mathrm{O}}_{X}^{+})^{\Delta}\to\mathrm{R}^{i}\nu_{*}(\mathbf{L}\otimes_{\widehat{\mathbf{Z}}_{p}}\widehat{\mathrm{O}}_{X}^{+})$$

are almost isomorphisms.

We now explain the main steps of our proof of Theorems 1.3.1 and 1.3.2:

(1) We first show that the sheaves $\mathbb{R}^i \nu_*(\mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$ are almost quasi-coherent and, more precisely, the natural morphisms

$$\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathrm{O}}_{X}^{+})^{\Delta} \to \mathrm{R}^{i} \nu_{*}(\mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathrm{O}}_{X}^{+})$$

are almost isomorphisms for affine $\mathfrak{X} = \operatorname{Spf} R$. The main key input is the construction of the "universal covering space" of $\mathfrak{X} = \operatorname{Spf} A$. This universal covering space turns out to

be an affinoid perfectoid, so we can almost compute étale cohomology $\mathrm{H}^{i}(\mathfrak{X}_{C}, \mathbf{L} \otimes_{\mathbf{F}_{p}} \mathfrak{O}_{X}^{+}/p)$ via the generic fiber of this covering. The construction of the "universal covering space" is explained just after Lemma 6.2.5.

- (2) The same ideas can be used to show that the formation of $\mathrm{R}^{i}\nu_{*}(\mathbf{L}\otimes_{\widehat{\mathbf{Z}}_{p}}\widehat{\mathcal{O}}_{X}^{+})$ commutes with étale base change.
- (3) We show that the $\mathcal{O}_{\mathfrak{X}_0}$ -modules $\mathrm{R}^i \nu_* \left(\mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+ / p \right)$ are almost coherent for smooth X. This is done in three steps: we firstly find an admissible blow-up $\mathfrak{X}' \to \mathfrak{X}$ such that \mathfrak{X}' has an open affine covering $\mathfrak{X}' = \bigcup_{i \in I} \mathfrak{U}_i$ such that each $\mathfrak{U}_i = \mathrm{Spf} A_i$ admits a finite rig-étale morphism to $\widehat{\mathbf{A}}^d_{\mathcal{O}_C}$, then we show that the cohomology groups $\mathrm{H}^i(\mathfrak{U}_{i,C}, \mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+ / p)$ are almost coherent over A_i/p , and finally we conclude the almost coherence of $\mathrm{R}^i \nu_* \left(\mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+ / p \right)$.

The first step is the combination of [BLR95, Proposition 3.7] and Theorem B.4. The first result allows to choose an admissible blow-up $\mathfrak{X}' \to \mathfrak{X}$ with an open affine covering $\mathfrak{X}' = \bigcup_{i \in I} \mathfrak{U}_i$ such that each \mathfrak{U}_i admits a rig-étale morphism $\mathfrak{U}_i \to \widehat{\mathbf{A}}^d_{\mathcal{O}_C}$. Then Theorem B.4 guarantees that actually we can change these morphisms so that $\mathfrak{U}_i \to \widehat{\mathbf{A}}^d_{\mathcal{O}_C}$ are *finite* and rig-étale. This is the non-noetherian generalization of Achinger's result [Ach17, Proposition 6.6.1] proven over a discretely valued ring.

The second step follows the strategy presented in [Sch13]. We construct an explicit affinoid perfectoid cover of \mathfrak{U}_i that is a $\mathbf{Z}_p(1)^d$ -torsor. So we reduce studying of $\mathrm{H}^i(\mathfrak{U}_{i,C}, \mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p)$ to studying cohomology groups of $\mathbf{Z}_p(1)^d$ that can be explicitly understood via the Koszul complex.

The last step is the consequence of the Almost Proper Mapping Theorem 1.2.6 and the already obtained results.

- (4) The next step is to show that $\mathrm{R}^{i}\nu_{*}\left(\mathbf{L}\otimes_{\mathbf{F}_{p}}\mathcal{O}_{X}^{+}/p\right)$ is almost coherent for a general X. This is done by choosing a proper hypercovering by smooth spaces X_{\bullet} and then use a version of cohomological v-descent to conclude almost coherence of the p-adic nearby cycles sheaves. As an important technical tool, we use the theory of diamonds developed in [Sch17].
- (5) Next we show that $\mathbf{R}\nu_* \left(\mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p\right)$ is concentrated in degrees [0, 2d] and almost concentrated in degrees [0, d]. The former claim is quite formal and follows from [Hub96, Proposition 2.8.1]. The latter claim is more subtle. The key input in its proof is the version of the purity theorem [BS19, Theorem 10.11] that implies that any *finite* (but not necessarily étale) adic space over an affinoid perfectoid has a diamond that is isomorphic to a diamond of an affinoid perfectoid. This allows us to reduce the question of cohomological bounds of $\mathbf{R}\nu_* \left(\mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p\right)^a$ to the question of cohomological dimension of the pro-finite group $\mathbf{Z}_p(1)^d$. This can be understood quite explicitly via Koszul complexes. This finishes the proof of Theorem 1.3.1.
- (6) Finally, we show Theorem 1.3.2 by reducing it to Theorem 1.3.1. This reduction goes similarly to the reduction of Theorem 1.2.7 to Theorem 1.2.6.

1.4. Acknowledgements. We are very grateful to B. Bhatt, B. Conrad, S. Petrov, and D. B. Lim for many fruitful discussions. We express additional gratitude to B. Bhatt for bringing [BS19, Theorem 10.11] and [Guo19] to our attention. We are thankful to B. Conrad for reading the first draft of this paper and making useful suggestions on how to improve the exposition of this paper. Part of this work was carried out at the mathematics department of the University of Michigan. We thank them for their hospitality.

1.5. Notation. We follow [Sta21, Tag 02MN] for the definition of a (Weak) Serre subcategory of an abelian category \mathcal{A} .

Namely, we say that a non-empty full subcategory \mathcal{C} of an abelian category \mathcal{A} is a *Serre subcategory* if, for any exact sequence $A \to B \to C$ with $A, C \in \mathcal{C}$, we have $B \in \mathcal{C}$. We say that \mathcal{C} is a *Weak Serre subcategory* if, for any exact sequence

$$A_0 \to A_1 \to A_2 \to A_3 \to A_4$$

with $A_0, A_1, A_3, A_4 \in \mathbb{C}$, we have $A_2 \in \mathbb{C}$. Look at [Sta21, Tag 02MP] and [Sta21, Tag 0754] for an alternative way to describe (Weak) Serre subcategories.

If C is a Serre subcategory of an abelian category \mathcal{A} we define the *quotient category* as a pair $(\mathcal{A}/\mathbb{C}, F)$ of an abelian category \mathcal{A}/\mathbb{C} and an exact functor

$$F: \mathcal{A} \to \mathcal{A}/\mathcal{C}$$

such that, for any exact functor $G: \mathcal{A} \to \mathcal{B}$ to an abelian category \mathcal{B} with $\mathcal{C} \subset \ker G$, there is a factorization $G = H \circ F$ for a unique exact functor $H: \mathcal{A}/\mathcal{C} \to \mathcal{B}$. The quotient category always exists by [Sta21, Tag 02MS].

If \mathcal{B} is a full triangulated subcategory of a triangulated category \mathcal{D} we define the Verdier quotient as a pair $(\mathcal{D}/\mathcal{B}, F)$ of a triangulated category \mathcal{D}/\mathcal{B} and an exact functor

$$F: \mathcal{D} \to \mathcal{D}/\mathcal{B}$$

such that, for any exact functor $G: \mathcal{D} \to \mathcal{D}'$ to a pre-triangulated category \mathcal{D}' with $\mathcal{B} \subset \ker G$, there is a factorization $G = H \circ F$ for a unique exact functor $H: \mathcal{D}/\mathcal{B} \to \mathcal{D}'$. The Verdier quotient always exists by [Sta21, Tag 05RJ].

We say that a diagram of categories

$$\begin{array}{ccc} \mathcal{A} & \stackrel{J}{\longrightarrow} & \mathcal{B} \\ \downarrow^{h} & \stackrel{\alpha}{\longleftarrow} & \downarrow^{g} \\ \mathcal{C} & \stackrel{k}{\longrightarrow} & \mathcal{D} \end{array}$$

is 2-commutative if $\alpha: k \circ h \Rightarrow g \circ f$ is a natural isomorphism of functors.

For an abelian group M and commuting endomorphisms f_1, \ldots, f_n , we define the Koszul complex

$$K(M; f_1, \dots, f_n) \coloneqq M \to M \otimes_{\mathbf{Z}} \mathbf{Z}^n \to M \otimes_{\mathbf{Z}} \wedge^2(\mathbf{Z}^n) \to \dots \to M \otimes_{\mathbf{Z}} \wedge^n(\mathbf{Z}^n)$$

viewed as a chain complex in cohomological degrees $0, \ldots, n$. The differential

$$d^{k} \colon M \otimes_{\mathbf{Z}} \wedge^{k} (\mathbf{Z}^{n}) \simeq \bigoplus_{1 \le i_{1} < \dots < i_{k} \le n} M \to M \otimes_{\mathbf{Z}} \wedge^{i+1} (\mathbf{Z}^{n}) \simeq \bigoplus_{1 \le j_{1} < \dots < j_{k+1} \le n} M$$

from M in spot $i_1 < \cdots < i_k$ to M in spot $j_1 < \cdots < j_{k+1}$ is nonzero only if $\{i_1, \ldots, i_k\} \subset \{j_1, \ldots, j_{k+1}\}$, in which case it is given by $(-1)^{m-1}f_{j_m}$, where $m \in \{1, \ldots, k+1\}$ is the unique integer such that $j_m \notin \{i_1, \ldots, i_k\}$.

If M is an R-module and f_i are elements of R the complex $K(M; f_1, \ldots, f_n)$ is a complex of R-modules and can be identified with

$$M \to M \otimes_R R^n \to M \otimes_R \wedge^2 (R^n) \to \dots \to M \otimes_R \wedge^n (R^n)$$

2. Almost Commutative Algebra

This chapter is devoted to the study of almost coherent modules. We recall some basic definitions of almost mathematics in Section 2.1. Then we discuss the main properties of almost finitely generated and almost finitely presented modules in Section 2.5. These two sections closely follow the discussion of almost mathematics in [GR03]. Section 2.6 is dedicated to almost coherent modules and almost coherent rings. We show that almost coherent modules from a Weak Serre subcategory of R-modules, and they coincide with almost finitely presented ones in the case of almost coherent rings. We discuss base change results in Section 2.7. Finally, we develop some topological aspects of almost finitely generated modules over "topologically universally adhesive rings" in Section 2.8. We also point out Appendix A is close in spirit to this Chapter, but it deals with more delicate (and technical) aspects of this theory that are used only in the proof of Theorem 5.1.6. So we decided to put that discussion into an appendix.

2.1. The Category of Almost Modules. We begin this section by recalling some basic definitions of almost mathematics from [GR03]. We fix some "base" ring R with an ideal \mathfrak{m} such that $\mathfrak{m}^2 = \mathfrak{m}$ and $\tilde{\mathfrak{m}} = \mathfrak{m} \otimes_R \mathfrak{m}$ is flat. We always do almost mathematics with respect to \mathfrak{m} .

Lemma 2.1.1. Let M be an R-module. Then the following are equivalent:

- (1) The module $\mathfrak{m}M$ is the zero module.
- (2) The module $\mathfrak{m} \otimes_R M$ is the zero module.
- (3) The module $\widetilde{\mathfrak{m}} \otimes_R M$ is the zero module.
- (4) The module M is annihilated by ε for every $\varepsilon \in \mathfrak{m}$.

Proof. Note that the multiplication map $\mathfrak{m} \otimes_R \mathfrak{m} \to \mathfrak{m}$ is surjective as $\mathfrak{m}^2 = \mathfrak{m}$. This implies that we have surjections

$$\widetilde{\mathfrak{m}} \otimes_R M \twoheadrightarrow \mathfrak{m} \otimes_R M \twoheadrightarrow \mathfrak{m} M.$$

This shows that (3) implies (2), and (2) implies (1). It is clear that (2) implies (3), and (1) is equivalent to (4). So the only thing we are left to show is that (1) implies (2).

Suppose that $\mathfrak{m}M \simeq 0$. Pick an arbitrary element $a \otimes m \in \mathfrak{m} \otimes_R M$ with $a \in \mathfrak{m}, m \in M$. Since $\mathfrak{m}^2 = \mathfrak{m}$ there is a finite number of elements $y_1, \ldots, y_k, x_1, \ldots, x_k \in \mathfrak{m}$ such that

$$a = \sum_{i=1}^{k} x_i y_i.$$

Then we have an equality

$$a \otimes m = \sum_{i=1}^{k} x_i y_i \otimes m = \sum_{i=1}^{k} x_i \otimes y_i m = 0.$$

Definition 2.1.2. An *R*-module M is almost zero, if any of the equivalent conditions of Lemma 2.1.1 is satisfied for M.

Lemma 2.1.3. Under the assumption as above, the "multiplication" morphism $\widetilde{\mathfrak{m}} \otimes_R \widetilde{\mathfrak{m}} \to \widetilde{\mathfrak{m}}$ is an isomorphism.

Proof. We consider a short exact sequence

$$0 \to \mathfrak{m} \to R \to R/\mathfrak{m} \to 0.$$

Note that $(R/\mathfrak{m}) \otimes_R \mathfrak{m} = \mathfrak{m}/\mathfrak{m}^2 = 0$, so we get a short exact sequence

$$0 \to \operatorname{Tor}_1^R(R/\mathfrak{m},\mathfrak{m}) \to \widetilde{\mathfrak{m}} \to \mathfrak{m} \to 0.$$

Since $\operatorname{Tor}_1^R(R/\mathfrak{m},\mathfrak{m})$ is almost zero, Lemma 2.1.1 says that after applying the functor $-\otimes_R \widetilde{\mathfrak{m}}$ we get an isomorphism

$$\mathfrak{m} \otimes_R \mathfrak{m} \simeq \mathfrak{m} \otimes_R \mathfrak{m}.$$

Since $\widetilde{\mathfrak{m}}$ is *R*-flat, we also see that $\mathfrak{m} \otimes_R \widetilde{\mathfrak{m}}$ injects into $\widetilde{\mathfrak{m}}$. Moreover, it maps isomorphically onto its image $\mathfrak{m}\widetilde{\mathfrak{m}} = \widetilde{\mathfrak{m}}$ as $\mathfrak{m}^2 = \mathfrak{m}$. Altogether it shows that

$$\widetilde{\mathfrak{m}}\otimes_R \widetilde{\mathfrak{m}}\simeq \widetilde{\mathfrak{m}}$$

It is straightforward to see that the constructed isomorphism is the "multiplication" map. \Box

We denote by Σ_R the category of almost zero *R*-modules considered as a full subcategory of \mathbf{Mod}_R .

Corollary 2.1.4. The category Σ_R is a Serre subcategory of Mod_R^3 .

Proof. This follows directly from criterion (3) from Lemma 2.1.1, flatness of $\widetilde{\mathfrak{m}}$ and [Sta21, Tag 02MP].

This corollary allows us to define the quotient category $\mathbf{Mod}_R^a := \mathbf{Mod}_R / \Sigma_R$ that we call as the category of almost *R*-modules⁴. Note that the localization functor

$$(-)^a: \mathbf{Mod}_R \to \mathbf{Mod}_R^a$$

is an exact and essentially surjective functor. We refer to elements of \mathbf{Mod}_R^a as almost *R*-modules or R^a -modules. We will usually denote them by M^a in order to distinguish almost *R*-modules from *R*-modules.

Definition 2.1.5. A morphism $f: M \to N$ is called an almost isomorphism (resp. almost injection, resp. almost surjection) if the corresponding morphism $f^a: M^a \to N^a$ is an isomorphism (resp. injection, resp. surjection) in \mathbf{Mod}_R^a .

Remark 2.1.6. For any *R*-module *M*, the natural morphism $\pi : \widetilde{\mathfrak{m}} \otimes_R M \to M$ is an almost isomorphism. Indeed, it suffices to show that

$$\widetilde{\mathfrak{m}} \otimes_R \ker \pi \simeq 0$$
 and $\widetilde{\mathfrak{m}} \otimes_R \operatorname{Coker} \pi \simeq 0$.

Using R-flatness of $\widetilde{\mathfrak{m}}$, we can reduce the question to showing that the the map

$$\widetilde{\mathfrak{m}} \otimes_R \pi \colon \widetilde{\mathfrak{m}} \otimes_R \widetilde{\mathfrak{m}} \otimes_R M \to \widetilde{\mathfrak{m}} \otimes_R M$$

is an isomorphism. This follows from Lemma 2.1.3.

Definition 2.1.7. Two *R*-modules *M* and *N* are called *almost isomorphic* if M^a is isomorphic to N^a in \mathbf{Mod}_R^a .

Lemma 2.1.8. Let $f: M \to N$ be an morphism of *R*-modules, then

- (1) The morphism f is an almost injection (resp. almost surjection, resp. almost isomorphism) if and only if ker(f) (resp. Coker(f), resp. ker(f) and Coker(f)) is an almost zero module.
- (2) We have a functorial bijection $\operatorname{Hom}_R(\widetilde{\mathfrak{m}} \otimes_R M, N) = \operatorname{Hom}_{\operatorname{\mathbf{Mod}}_R^a}(M^a, N^a).$

 $^{^{3}}$ We refer to [Sta21, Tag 02MN] for the discussion of (Weak) Serre categories.

 $^{^{4}}$ We refer to [Sta21, Tag 02MS] for the discussion of quotient categories.

(3) Modules M and N are almost isomorphic (not necessary via a morphism f) if and only if $\widetilde{\mathfrak{m}} \otimes_R M \simeq \widetilde{\mathfrak{m}} \otimes_R N$.

Proof. (1) just follows from definition of the quotient category. (2) is discussed in detail in [GR03, page 12, (2.2.4)].

Now we show that (3) follows from (1) and (2). Remark 2.1.6 implies that M and N are almost isomorphic if $\widetilde{\mathfrak{m}} \otimes_R M \simeq \widetilde{\mathfrak{m}} \otimes_R N$.

Now suppose that there is an almost isomorphism $\varphi : M^a \to N^a$. It has a representative $f: \widetilde{\mathfrak{m}} \otimes_R M \to N$ by (2). Now (1) and *R*-flatness of $\widetilde{\mathfrak{m}}$ imply that $\widetilde{\mathfrak{m}} \otimes_R f: \widetilde{\mathfrak{m}} \otimes_R \widetilde{\mathfrak{m}} \otimes_R M \to \widetilde{\mathfrak{m}} \otimes_R N$ is an isomorphism. Now $\widetilde{\mathfrak{m}} \otimes_R \widetilde{\mathfrak{m}} \simeq \widetilde{\mathfrak{m}}$ by Lemma 2.1.3, so $\widetilde{\mathfrak{m}} \otimes_R f$ gives an isomorphism

$$\widetilde{\mathfrak{m}} \otimes_R f : \widetilde{\mathfrak{m}} \otimes_R M \to \widetilde{\mathfrak{m}} \otimes_R N.$$

We now define the *functor of almost sections*

$$(-)_* \colon \mathbf{Mod}_R^a \to \mathbf{Mod}_R$$

as

$$(M^a)_* \coloneqq \operatorname{Hom}_{\operatorname{\mathbf{Mod}}_R^a}(R^a, M^a) = \operatorname{Hom}_R(\widetilde{\mathfrak{m}}, M)$$

for any R^a -module M^a with an R-module representative M. The construction is clearly functorial in M^a , so it does define the functor $(-)_* \colon \mathbf{Mod}_R^a \to \mathbf{Mod}_R$.

The functor of almost sections is going to be the right adjoint to the almostification functor $(-)^a$. Before we discuss why this is the case, we need to define the unit and counit transformations.

We start with the unit of the adjunction. For any R-module M, there is a functorial morphism

$$\eta_{M,*} \colon M \to \operatorname{Hom}_R(\widetilde{\mathfrak{m}}, M) = M_*^{\mathfrak{c}}$$

that can easily be seen to be an almost isomorphism.

This allows us to define a functorial morphism

$$\varepsilon_{N^a,*} \colon (N^a_*)^a \to N^a$$

for any R^a -module N^a . Namely, the map $\eta_{N,*} \colon N \to N^a_*$ is an almost isomorphism, so we can invert it in the almost category and define

$$\varepsilon_{N^a,*} \coloneqq (\eta^a_{N,*})^{-1} \colon (N^a_*)^a \to N^a$$

Now we define another functor

$$(-)_! \colon \mathbf{Mod}_R^a \to \mathbf{Mod}_R$$

that is going to be a left adjoint to the almostification functor $(-)^a$. Namely, we put

$$(M^a)_! \coloneqq (M^a)_* \otimes_R \widetilde{\mathfrak{m}} \xleftarrow{\sim} M \otimes_R \widetilde{\mathfrak{m}}$$

for any \mathbb{R}^a -module M^a with an \mathbb{R} -module representative M. This construction is clearly functorial in M^a , so it does define a functor. Similarly to the discussion above, for any \mathbb{R} -module M, we define the transformation

$$\varepsilon_{M,!} \colon (M^a)_! = \widetilde{\mathfrak{m}} \otimes_R M \to M$$

as the map induces by the the natural morphism $\widetilde{\mathfrak{m}} \to R$. Clearly, $\varepsilon_{M,!}$ is an almost isomorphism for any M. So, this actually allows us to define the morphism

$$\eta_{N^a,!} \colon N^a \to (\widetilde{\mathfrak{m}} \otimes_R N)^a \simeq (N^a_!)^a$$

as $\eta_{N^a,!} = (\varepsilon^a_{N,!})^{-1}$.

We summarize the main properties of these functors in the lemma below:

Lemma 2.1.9. Let R and \mathfrak{m} be as above. Then

- (1) The functor $(-)_*$ is the right adjoint to $(-)^a$. In particular, it is left exact.
- (2) The unit of the adjunction is equal to $\eta_{M,*}$, the counit of the adjunction is equal to $\varepsilon_{N^a,*}$. In particular, both of the are isomorphisms.
- (3) The functor $(-)_{!}$ is the left adjoint to the localization functor $(-)^{a}$.
- (4) The functor $(-)_! : \mathbf{Mod}_R^a \to \mathbf{Mod}_R$ is exact.
- (5) The unit of the adjunction is equal to $\eta_{N^a,!}$, the counit of the adjunction is equal to $\varepsilon_{M,!}$. In particular, both of the are almost isomorphisms.

Proof. This is explained [GR03, Proposition 2.2.13 and Proposition 2.2.21].

The last thing we need to address in this section is how almost mathematics interacts with base change. We want to be able to speak about preservation of various properties of modules under a base change along a map $R \to S$. The issue here is to define the corresponding ideal \mathfrak{m}_S as in the definition of almost mathematics. It turns out that the most naive ideal $\mathfrak{m}_S := \mathfrak{m}S$ works well, but the reason is that the assumptions on the ideal defining almost mathematics are rather weak. More specifically, we could have required the flatness of the ideal \mathfrak{m} (instead of $\tilde{\mathfrak{m}}$), and then the ideal $\mathfrak{m}S$ would not serve well for defining almost mathematics on S. The next lemma shows that everything works well in the current setup.

Lemma 2.1.10. Let $f : R \to S$ be a ring homomorphism, and let \mathfrak{m}_S be the ideal $\mathfrak{m}_S \subset S$. Then we have an equality $\mathfrak{m}_S^2 = \mathfrak{m}_S$ and the S-module $\widetilde{\mathfrak{m}_S} := \mathfrak{m}_S \otimes_S \mathfrak{m}_S$ is S-flat.

Proof. The equality $\mathfrak{m}_S^2 = \mathfrak{m}_S$ follows from the analogous assumption on \mathfrak{m} and the construction of \mathfrak{m}_S . As for the flatness issue, we claim that $\mathfrak{m}_S \otimes_S \mathfrak{m}_S \simeq (\mathfrak{m} \otimes_R S) \otimes_S (\mathfrak{m} \otimes_R S)$. That would certainly imply that desired flatness statement. In order to prove this claim, we look at a short exact sequence

$$0 \to \mathfrak{m} \to R \to R/\mathfrak{m} \to 0$$

We apply $-\otimes_R S$ to get a short exact sequence

$$0 \to \operatorname{Tor}_1^R(R/\mathfrak{m}, S) \to \mathfrak{m} \otimes_R S \to \mathfrak{m} S \to 0.$$

We observe that $\operatorname{Tor}_{1}^{R}(R/\mathfrak{m}, S)$ is almost zero, so both $\operatorname{Tor}_{1}^{R}(R/\mathfrak{m}, S) \otimes_{S} \mathfrak{m}S$ and $\operatorname{Tor}_{1}^{R}(R/\mathfrak{m}, S) \otimes_{S} \mathfrak{m}S$ ($\mathfrak{m} \otimes_{R} S$) are zero modules by Lemma 2.1.1. So we use functors $- \otimes_{S} (\mathfrak{m} \otimes_{R} S)$ and $- \otimes_{S} \mathfrak{m}S$ to obtain isomorphisms

$$(\mathfrak{m} \otimes_R S) \otimes_S (\mathfrak{m} \otimes_R S) \simeq \mathfrak{m} S \otimes_R (\mathfrak{m} \otimes_R S) \simeq (\mathfrak{m} S) \otimes_S (\mathfrak{m} S).$$

Thus we get the desired equality.

Lemma 2.1.11. Let $f: R \to S$ be a ring homomorphism, and $F: \mathbf{Mod}_R \to \mathbf{Mod}_S$ an *R*-linear functor (resp. $F: \mathbf{Mod}_R^{op} \to \mathbf{Mod}_S$ an *R*-linear functor). Then *F* sends almost zero *R*-modules to almost zero *S*-modules.

Proof. Suppose that M is an almost zero R-module, so $\varepsilon M = 0$ for any $\varepsilon \in \mathfrak{m}$. Then $\varepsilon F(M) = 0$ because F is R-linear, so F(M) is almost zero by Lemma 2.1.1.

Corollary 2.1.12. Let $f: R \to S$ be a ring homomorphism, and $F: \operatorname{Mod}_R \to \operatorname{Mod}_S$ a left or right exact *R*-linear functor (resp. $F: \operatorname{Mod}_R^{op} \to \operatorname{Mod}_S$ a left or right exact *R*-linear functor). Then *F* preserves almost isomorphisms.

Proof. We only show the case of a left exact functor $F: \mathbf{Mod}_R \to \mathbf{Mod}_S$, all other cases are analogous to the this one.

Choose any almost isomorphism $f: M' \to M''$, we want to show that F(f) is an almost isomorphism. Consider the following exact sequences:

$$0 \to K \to M' \to M \to 0,$$
$$0 \to M \to M'' \to Q \to 0.$$

We know that K and Q are almost zero by our assumption on f. Now, the above short exact sequences induce the following exact sequences:

$$0 \to F(K) \to F(M') \to F(M) \to \mathbb{R}^1 F(K),$$
$$0 \to F(M) \to F(M'') \to F(Q).$$

Lemma 2.1.11 guarantees that F(K), $\mathbb{R}^1 F(K)$, and F(Q) are almost zero S-modules. Therefore, the morphisms $F(M') \to F(M)$ and $F(M) \to F(M'')$ are both almost isomorphisms. In particular, the composition $F(M') \to F(M'')$ is an almost isomorphism as well.

2.2. Basic Functors on the Categories of Almost Modules. The category of almost modules admits certain natural functors induced from the category of *R*-modules. It has two versions of the Hom-functor and the tensor product functor. We summarize properties of these functors in the following proposition:

Proposition 2.2.1. Let R, \mathfrak{m} be as above. Then

(1) We define tensor product functor $-\otimes_{R^a} -: \mathbf{Mod}_R^a \times \mathbf{Mod}_R^a \to \mathbf{Mod}_R^a$ as

 $(M^a, N^a) \mapsto (M^a_! \otimes_R N^a_!)^a$.

Then there is a natural transformation of functors

$$\begin{array}{c} \mathbf{Mod}_R \times \mathbf{Mod}_R & \xrightarrow{-\otimes_R -} & \mathbf{Mod}_R \\ & \downarrow^{(-)^a \times (-)^a} & \downarrow^{(-)^a} \\ \mathbf{Mod}_R^a \times \mathbf{Mod}_R^a & \xrightarrow{-\otimes_R a -} & \mathbf{Mod}_R^a \end{array}$$

that makes the diagram 2-commutative. In particular, there is a functorial isomorphism $(M \otimes_R N)^a \simeq M^a \otimes_{R^a} N^a$ for any $M, N \in \mathbf{Mod}_R$.

(2) There is a functorial isomorphism

$$\operatorname{Hom}_{R^a}(M^a, N^a) \simeq \operatorname{Hom}_R(\widetilde{\mathfrak{m}} \otimes M, N)$$
.

for any $M, N \in \mathbf{Mod}_R$. In particular, there is a canonical structure of an *R*-module on the group $\operatorname{Hom}_{R^a}(M^a, N^a)$; thus defines the functor

$$\operatorname{Hom}_{R^a}(-,-)\colon \operatorname{\mathbf{Mod}}_R^{a,op} \times \operatorname{\mathbf{Mod}}_R^a \to \operatorname{\mathbf{Mod}}_R$$

(3) We define the functor $\operatorname{alHom}_{R^a}(-,-): \operatorname{Mod}_R^{a,op} \times \operatorname{Mod}_R^a \to \operatorname{Mod}_R^a$ of almost homomorphisms as

$$(M^a, N^a) \mapsto \operatorname{Hom}_{R^a}(M, N)^a$$
.

Then there is a natural transformation of functors

$$\begin{array}{ccc} \mathbf{Mod}_R^{op} \times \mathbf{Mod}_R & \xrightarrow{\mathrm{Hom}_R(-,-)} & \mathbf{Mod}_R \\ & & & \downarrow (-)^a \times (-)^a & & \downarrow (-)^a \\ & & \mathbf{Mod}_R^{a,op} \times \mathbf{Mod}_R^a \xrightarrow{\mathrm{alHom}_Ra\,(-,-)} & \mathbf{Mod}_R^a \end{array}$$

that makes the diagram 2-commutative. In particular, $\operatorname{alHom}_{R^a}(M^a, N^a) \cong^a \operatorname{Hom}_R(M, N)^a$ for any $M, N \in \operatorname{\mathbf{Mod}}_R$.

Proof. (1). We define

 $\rho_{M,N} \colon (M^a_! \otimes_R N^a_!)^a \to (M \otimes_R N)^a$

to be the morphism induced by

 $M^a_{\mathsf{I}} \simeq \widetilde{\mathfrak{m}} \otimes_R M \to M \text{ and } N^a_{\mathsf{I}} \simeq \widetilde{\mathfrak{m}} \otimes_R N \to N.$

It is clear that $\rho_{M,N}$ is functorial in both variables, so it defines a natural transformation of functors ρ . We also need to check that $\rho_{M,N}$ is an isomorphism for any M and N. This follows from the following two observations: $\rho_{M,N}$ is an isomorphism if and only if $\rho_{M,N} \otimes_R \widetilde{\mathfrak{m}}$ is an isomorphism; and $\rho_{M,N} \otimes_R \widetilde{\mathfrak{m}}$ is easily seen to be an isomorphism as $\widetilde{\mathfrak{m}} \otimes_R \widetilde{\mathfrak{m}} \to \widetilde{\mathfrak{m}}$ is an isomorphism.

(2) is just a reformulation of Lemma 2.1.8(2).

In order to show (3), we need to define a functorial morphism

$$\rho_{M,N} \colon \operatorname{Hom}_R(M,N)^a \to \operatorname{alHom}_{R^a}(M^a,N^a).$$

We start by using the functorial identification

alHom_{$$R^a$$} $(M^a, N^a) \cong^a \operatorname{Hom}_R(\widetilde{\mathfrak{m}} \otimes M, N)^a$

from (2). Namely, we define $\rho_{M,N}$ as the morphism $\operatorname{Hom}_R(M, N)^a \to \operatorname{Hom}_R(\widetilde{\mathfrak{m}} \otimes M, N)^a$ induced by the map $\widetilde{\mathfrak{m}} \otimes M \to M$. This is clearly functorial in both variables, so it defines the natural transformation ρ .

We also need to check that $\rho_{M,N}$ is an isomorphism for any M and N. This boils down to the fact that $\operatorname{Hom}_R(-, N)$ sends almost isomorphisms to almost isomorphisms. This, in turn, follows from Corollary 2.1.12.

Remark 2.2.2. It is straightforward to check that if N has a structure of an S^a -module for some R-algebra S, then the R^a -modules $M^a \otimes_{R^a} N^a$, alHom_{$R^a}(<math>M^a, N^a$) have functorial-in- M^a structures of S^a -modules. This implies that the functors $- \otimes_{R^a} N^a$, alHom_{R^a}($-, N^a$) naturally land in **Mod**^a_S, i.e. define functors</sub>

$$-\otimes_{R^a} N^a \colon \mathbf{Mod}_R^a \to \mathbf{Mod}_S^a$$
, and $\mathrm{alHom}_{R^a}(-, N^a) \colon \mathbf{Mod}_R^{a,op} \to \mathbf{Mod}_S^a$

Similarly, $\operatorname{Hom}_{R^a}(-, N^a)$ defines a functor $\operatorname{Mod}_R^a \to \operatorname{Mod}_S$.

The functor of almost homomorphisms is quite important as it turns out to be the *inner Hom* functor, i.e. it is right adjoint to the tensor product.

Lemma 2.2.3. Let $f: R \to S$ be a ring homomorphism, and let M^a be an R^a -module and N^a, K^a be S^a -modules. Then there is a functorial S-linear isomorphism

$$\operatorname{Hom}_{S^a}(M^a \otimes_{R^a} N^a, K^a) \simeq \operatorname{Hom}_{R^a}(M^a, \operatorname{alHom}_{S^a}(N^a, K^a)) .$$

Proof. This is a consequence of the usual \otimes -Hom-adjunction, Proposition 2.2.1, and the fact that $\widetilde{\mathfrak{m}}^{\otimes 2} \simeq \widetilde{\mathfrak{m}}$. Indeed, we have the following sequence of functorial isomorphisms

$$\operatorname{Hom}_{S^{a}}(M^{a} \otimes_{R^{a}} N^{a}, K^{a}) \simeq \operatorname{Hom}_{S}(\widetilde{\mathfrak{m}} \otimes_{R} M \otimes_{R} N, K)$$
$$\simeq \operatorname{Hom}_{S}((\widetilde{\mathfrak{m}} \otimes_{R} M) \otimes_{R} (\widetilde{\mathfrak{m}} \otimes_{R} N), K)$$
$$\simeq \operatorname{Hom}_{R}(\widetilde{\mathfrak{m}} \otimes_{R} M, \operatorname{Hom}_{S}(\widetilde{\mathfrak{m}} \otimes_{R} N, K))$$
$$\simeq \operatorname{Hom}_{R^{a}}(M, \operatorname{alHom}_{S^{a}}(N^{a}, K^{a})) .$$

The first isomomorphism follows from Proposition 2.2.1(1), (2), the second isomorphism follows from the observation $\widetilde{\mathfrak{m}}^{\otimes 2} \simeq \widetilde{\mathfrak{m}}$, the third isomorphism is just the classical \otimes -Hom-adjunction, and the last isomorphism is a consequence of Proposition 2.2.1(2), (3).

- **Corollary 2.2.4.** (1) Let N be an R^a -module, then the functor $-\otimes_{R^a} N^a$ is left adjoint to the functor alHom_{R^a} $(N^a, -)$.
 - (2) Let $R \to S$ be a ring homomorphism. Then the functor $-\otimes_{R^a} S^a \colon \mathbf{Mod}_R^a \to \mathbf{Mod}_S^a$ is left adjoint to the forgetful functor.

Proof. Part (1) follows from Lemma 2.2.3 by taking S to be equal to R. Part (2) follows from Lemma 2.2.3 by taking N^a to be equal to S^a .

We finish the section by introducing the certain types of R^a -modules that will be used throughout the paper.

Definition 2.2.5. • We say that an R^a -module M^a is almost flat if the functor $M^a \otimes_{R^a} -: \operatorname{Mod}_R^a \to \operatorname{Mod}_R^a$ is exact.

- We say that an R^a -module I^a is *injective* if the functor $\operatorname{Hom}_{R^a}(-, I^a)$: $\operatorname{Mod}_{R}^{a,op} \to \operatorname{Mod}_{R}^{a}$ is exact.
- We say that an R^a -module P^a is almost projective if the functor alHom_{R^a} $(P^a, -)$: $\mathbf{Mod}_R^a \to \mathbf{Mod}_R^a$ is exact.

Lemma 2.2.6. The functor $(-)^a \colon \mathbf{Mod}_R \to \mathbf{Mod}_R^a$ sends flat (resp. injective, resp. projective) R-modules to almost flat (resp. injective, resp. almost projective) R^a -modules.

Proof. The case of flat modules is clear from Lemma 2.2.1(1). The case of injective modules follows from the fact that $(-)^a$ admits an exact left adjoint functor $(-)_!$. The case of projective modules is clear from the definition.

Corollary 2.2.7. Any bounded above complex $C^{\bullet,a} \in \mathbf{Comp}^-(\mathbb{R}^a)$ admits a resolution $P^{\bullet,a} \to C^{\bullet}$ by a bounded above complex of almost projective modules.

Proof. We consider the complex $C_!^{\bullet,a} \in \mathbf{Comp}^-(R)$, this complex admits a resolution by complex of free modules $p: P^{\bullet} \to C_!^{\bullet,a}$. Now we apply $(-)^a$ to this morphism to get the map

$$P^{\bullet,a} \xrightarrow{p^a} (C^{\bullet,a}_!)^a \xleftarrow{\varepsilon} C^{\bullet,a}$$
.

The map ε is an *isomorphism* in $\operatorname{Comp}(R^a)$ by Lemma 2.1.9, and p^a is a quasi-isomorphism. Thus $\varepsilon^{-1} \circ p^a \colon P^{\bullet,a} \to C^{\bullet,a}$ is a quasi-isomorphism in $\operatorname{Comp}(R^a)$. Now note that each term of $P^{\bullet,a}$ is almost projective by Lemma 2.2.6.

2.3. Derived Category of Almost Modules. We define the derived category of almost modules in two different ways and show that these definitions coincide. Later we define certain derived functors on the derived category of almost modules. We pay some extra attention to show that the functors in this section are well-defined on unbounded derived categories.

We start the section by introducing two different notions of the derived category of almost modules and then show that they are actually the same.

Definition 2.3.1. We define the *derived category of almost* R-modules as $\mathbf{D}(R^a) := \mathbf{D}(\mathbf{Mod}_R^a)$.

We define the bounded version of derived category of almost R-modules $\mathbf{D}^*(\mathbb{R}^a)$ for $* \in \{+, -, b\}$ as the full subcategory consisting of bounded below (resp. bounded above, resp. bounded) complexes.

Definition 2.3.2. We define the almost derived category of *R*-modules as the Verdier quotient $\mathbf{D}(R)^a \coloneqq \mathbf{D}(\mathbf{Mod}_R)/\mathbf{D}_{\Sigma_R}(\mathbf{Mod}_R).$

We recall that Σ_R is the Serre subcategory of \mathbf{Mod}_R that consists of almost zero modules, and $\mathbf{D}_{\Sigma_R}(\mathbf{Mod}_R)$ is the full triangulated category of elements in $\mathbf{D}(\mathbf{Mod}_R)$ with almost zero cohomology modules.

We note that the functor $(-)^a \colon \mathbf{Mod}_R \to \mathbf{Mod}_R^a$ is exact and additive. Thus it can be derived to the functor $(-)^a \colon \mathbf{D}(R) \to \mathbf{D}(R^a)$. Similarly, the functor $(-)_! \colon \mathbf{Mod}_R^a \to \mathbf{Mod}_R$ is additive and exact, thus it can be derived to the functor $(-)_! \colon \mathbf{D}(R^a) \to \mathbf{D}(R)$. The standard argument shows that $(-)_!$ is a left adjoint functor to the functor $(-)^a$ as this already happens on the level of abelian categories. Now we also want to derive the functor $(-)_* \colon \mathbf{Mod}_R^a \to \mathbf{Mod}_R$. In order to do this on the level of unbounded derived categories, we need to show that $\mathbf{D}(R^a)$ has "enough K-injective objects".

Definition 2.3.3. We say that a complex of R^a -module $I^{\bullet,a}$ is *K*-injective if $\operatorname{Hom}_{K(R^a)}(C^{\bullet,a}, I^{\bullet,a}) = 0$ for any acyclic complex $C^{\bullet,a}$ of R^a -modules.

Remark 2.3.4. We remind the reader that $K(R^a)$ stands for the homotopy category of R^a -modules.

The first thing we need to show is that $\mathbf{Comp}(\mathbb{R}^a)$ has "enough" K-injective objects. This will allow us derive many functors.

Lemma 2.3.5. The functor $(-)^a$: **Comp** $(R) \to$ **Comp** (R^a) sends *K*-injective *R*-complexes to *K*-injective R^a -complexes.

Proof. We note that $(-)^a$ admits an exact left adjoint $(-)_!$ thus [Sta21, Tag 08BJ] ensures that $(-)^a$ preserves K-injective complexes.

Corollary 2.3.6. Every object $M^{\bullet,a} \in \mathbf{Comp}(\mathbb{R}^a)$ is quasi-isomorphic to a K-injective complex.

Proof. We know that the complex $M^{\bullet} \in \mathbf{Comp}(R)$ is quasi-isomorphic to a K-injective complex I^{\bullet} by [Sta21, Tag 090Y] (or [Sta21, Tag 079P]). Now we use Lemma 2.3.5 to say that $I^{\bullet,a}$ is a K-injective complex that is quasi-isomorphic to $M^{\bullet,a}$.

Now as the first application of Corollary 2.3.6 we define the functor $(-)_*: \mathbf{D}(\mathbb{R}^a) \to \mathbf{D}(\mathbb{R})$ as the derived functor of $(-)_*: \mathbf{Mod}_{\mathbb{R}}^a \to \mathbf{Mod}_{\mathbb{R}}$. This functor exists by [Sta21, Tag 070K]. **Lemma 2.3.7.** (1) The functors $\mathbf{D}(R) \xrightarrow[(-)^a]{(-)^a} \mathbf{D}(R^a)$ are adjoint. Moreover, the unit (resp.

counit) morphism

$$(M^a)_! \to M \text{ (resp. } N \to (N_!)^a)$$

is an almost isomorphism (resp. isomorphism) for any $M \in \mathbf{D}(R), N \in \mathbf{D}(R^a)$. In particular, the functor $(-)^a$ is essentially surjective.

(2) The functors $\mathbf{D}(R) \xrightarrow[(-)_*]{(-)_*} \mathbf{D}(R^a)$ are adjoint. Moreover, the unit (resp. counit) morphism

$$M \to (M^a)_* \text{ (resp. } (N_*)^a \to N)$$

is an almost isomorphism (resp. isomorphism) for any $M \in \mathbf{D}(R), N \in \mathbf{D}(R^a)$.

Proof. We start the proof by showing (1). Firstly we note that the functors $(-)_!$ and $(-)^a$ are adjoint by the discussion above. Now we show the cone of the counit map is always in $\mathbf{D}_{\Sigma_R}(R)$. As both functors $(-)^a$ and $(-)_!$ are exact on the level of abelian categories, it suffices to show the claim for $M \in \mathbf{Mod}_R^a$. But then the statement follows from Lemma 2.1.9(5). The same argument shows that the unit map $N \to (N_!)^a$ is an isomorphism for any $N \in \mathbf{D}(R^a)$.

Now we go to (2). We define the functor $(-)_*: \mathbf{D}(R^a) \to \mathbf{D}(R)$ as the right derived functor of the left exact additive functor $(-)_*: \mathbf{Mod}_R^a \to \mathbf{Mod}_R$. This functor exists by [Sta21, Tag 070K] and Corollary 2.3.6. The functor $(-)_*$ is right adjoint to $(-)^a$ by [Sta21, Tag 0DVC].

We check that the natural map $M \to (M^a)_*$ is an almost isomorphism for any $M \in \mathbf{D}(R)$. We choose some K-injective resolution $M \xrightarrow{\sim} I^{\bullet}$. Then Lemma 2.3.5 guarantees that $M^a \to I^{\bullet,a}$ is a K-injective resolution of the complex M^a . The map $M \to (M^a)_*$ has a representative

$$I^{\bullet} \to (I^{\bullet,a})_*$$

This map is an almost isomorphism of complexes by Lemma 2.1.9(2). Thus the map $M \to (M^a)_*$ is an almost isomorphism. A similar argument shows that the counit map $(N_*)^a \to N$ is an (almost) isomorphism for any $N \in \mathbf{D}(\mathbb{R}^a)$.

Theorem 2.3.8. The functor $(-)^a : \mathbf{D}(R) \to \mathbf{D}(R^a)$ induces an equivalence of triangulated categories $(-)^a : \mathbf{D}(R)^a \to \mathbf{D}(R^a)$.

Proof. We recall that the Verdier quotient is constructed as the localization of $\mathbf{D}(R)$ along the morphisms f such that $\operatorname{cone}(f) \in \mathbf{D}_{\Sigma_R}(R)$. For instance, this is the definition of Verdier quotient at [Sta21, Tag 05RI]. Now we see that a morphism $f^a \colon C^a \to C'^a$ in invertible in $\mathbf{D}(R^a)$ if and only if $\operatorname{cone}(f) \in \mathbf{D}_{\Sigma_R}(R)$ by the definition of Σ_R and exactness of $(-)^a$. Moreover, $(-)^a$ admits a right adjoint such that $(-)^a \circ (-)_* \to \operatorname{Id}$ is an isomorphism of functors. Thus we can apply [GZ67, Proposition 1.3] to say that the induced functor $(-)^a \colon \mathbf{D}(R)^a \to \mathbf{D}(R^a)$ must be an equivalence. \Box

Remark 2.3.9. Theorem 2.3.8 shows that the two notions of the derived category of almost modules are the same. In what follows, we do not distinguish $\mathbf{D}(R^a)$ and $\mathbf{D}(R)^a$ anymore.

2.4. Basic Functors on the Derived Categories of Almost Modules. Now we can "derive" certain functors constructed in previous section. We start with defining the derived versions of different Hom functors, after that we move to the case of the derived tensor product functor.

Definition 2.4.1. We define the *derived Hom* functor

 $\mathbf{R}\operatorname{Hom}_{R^a}(-,-): \mathbf{D}(R)^{a,op} \times \mathbf{D}(R)^a \to \mathbf{D}(R)$

as it is done in [Sta21, Tag 0A5W] using the fact that $\mathbf{Comp}(\mathbb{R}^a)$ has enough K-injective complexes. We define the *Ext modules* as *R*-modules defined as

 $\operatorname{Ext}_{R^a}^i(M^a, N^a) \coloneqq \operatorname{H}^i(\mathbf{R}\operatorname{Hom}_{R^a}(M^a, N^a))$

for $M^a, N^a \in \mathbf{Mod}_R^a$.

Explicitly, for any $M^a, N^a \in \mathbf{D}(R)^a$, the construction of the complex $\mathbf{R}\operatorname{Hom}_{R^a}(M^a, N^a)$ goes as follows. We pick any a $C^{\bullet,a} \to M^a$ and a K-injective resolution $N^a \to I^{\bullet,a}$. Then we set $\mathbf{R}\operatorname{Hom}_{R^a}(M^a, N^a) = \operatorname{Hom}_{R^a}^{\bullet}(C^{\bullet,a}, I^{\bullet,a})$. This construction is independent of the choices and functorial in both variables. We are not going to review this theory here, but rather refer to [Sta21, Tag 0A5W] for the details.

Remark 2.4.2. We see that [Sta21, Tag 0A64] implies that there is a functorial isomorphism

 $\mathrm{H}^{i}(\mathbf{R}\mathrm{Hom}_{R^{a}}(M^{a},N^{a}))\simeq\mathrm{Hom}_{\mathbf{D}(R)^{a}}(M^{a},N^{a}[i])$.

Lemma 2.4.3. (1) There are functorial isomorphisms

 $\operatorname{Hom}_{\mathbf{D}(R)^a}(M^a, N^a) \simeq \operatorname{Hom}_{\mathbf{D}(R)}(M^a_!, N)$ and $\operatorname{\mathbf{R}Hom}_{R^a}(M^a, N^a) \simeq \operatorname{\mathbf{R}Hom}_R(M^a_!, N)$ for any $M, N \in \mathbf{D}(R)$.

(2) For any chosen $M^a \in \mathbf{Mod}_R^a$, the functor $\mathbf{R}\mathrm{Hom}_{R^a}(M^a, -) \colon \mathbf{D}(R)^a \to \mathbf{D}(R)$ is isomorphic to the (right) derived functor of $\mathrm{Hom}_{R^a}(M^a, -)$.

Proof. The first claim easily follows from the fact $(-)^a$ is a right adjoint to the exact functor $(-)_!$. We leave the details to the reader.

The second claim follows from [Sta21, Tag 070K] and Corollary 2.3.6.

Definition 2.4.4. We define the *derived functor of almost homomorphisms*

$$\mathbf{R}$$
al $\operatorname{Hom}_{R^a}(-,-): \mathbf{D}(R)^{a,op} \times \mathbf{D}(R)^a \to \mathbf{D}(R)^a$

as

$$\mathbf{R}$$
al $\operatorname{Hom}_{R^a}(M^a, N^a) \coloneqq \mathbf{R}\operatorname{Hom}_{R^a}(M^a, N^a)^a = \mathbf{R}\operatorname{Hom}_R(M^a_!, N)^a$.

We define the *almost Ext modules* as R^a -modules defined by

 $\operatorname{alExt}_{R^a}^i(M^a, N^a) \coloneqq \operatorname{H}^i(\operatorname{RalHom}_{R^a}(M^a, N^a))$

for $M^a, N^a \in \mathbf{Mod}_R^a$.

Definition 2.4.5. We define the *the complex of almost homomorphisms* $\operatorname{alHom}_{R^a}^{\bullet}(K^{\bullet,a}, L^{\bullet,a})$ for $K^{\bullet,a}, L^{\bullet,a} \in \operatorname{Comp}(R^a)$ as follows:

$$\mathrm{alHom}_{R^a}^n(K^{\bullet,a}, L^{\bullet,a}) \coloneqq \prod_{n=p+q} \mathrm{alHom}_{R^a}(K^{-q,a}, L^{p,a})$$

with the differential

$$d(f) = d_{L^{\bullet,a}} \circ f - (-1)^n f \circ d_{K^{\bullet,a}}$$

Lemma 2.4.6. Let $P^{\bullet,a}$ be a bounded above complex of R^a -modules with almost projective cohomology modules. Suppose that $M^{\bullet,a} \to N^{\bullet,a}$ is an almost quasi-isomorphism of bounded below complex of R^a -modules. Then the natural morphism

$$\operatorname{alHom}_{R^a}^{\bullet}(P^{\bullet,a}, M^{\bullet,a}) \to \operatorname{alHom}_{R^a}^{\bullet}(P^{\bullet,a}, N^{\bullet,a})$$

is an almost quasi-isomorphism.

20

Proof. We note that as in the case of the usual Hom-complexes, there are convergent⁵ spectral sequences

Moreover, there is a natural morphism of spectral sequences $E_1^{i,j} \to E_1^{\prime i,j}$. Thus it suffices to show that the associated map on the first page is an almost isomorphism on each entry. Now we use the fact that $\operatorname{alHom}_{R^a}(P^{-i,a}, -)$ is exact to rewrite the first page of this spectral sequence as

$$\mathbf{E}_{1}^{i,j} = \operatorname{alHom}_{R^{a}} \left(P^{-i,a}, \mathbf{H}^{j}(M^{\bullet,a}) \right)$$

and the same for $E_{1}^{i,j}$. So the question boils down to show that the natural morphisms

alHom_{$$R^a$$} $\left(P^{-i,a}, \mathrm{H}^{j}(M^{\bullet,a})\right) \to \mathrm{alHom}_{R^a}\left(P^{-i,a}, \mathrm{H}^{j}(N^{\bullet,a})\right)$

are almost isomorphisms. But this is clear as $M^{\bullet,a} \to N^{\bullet,a}$ is an almost quasi-isomorphism. \Box

Lemma 2.4.7. Let $P_1^{\bullet,a} \to P_2^{\bullet,a}$ be an almost quasi-isomorphism of bounded above complexes with almost projective cohomology modules. Suppose that $M^{\bullet,a}$ is a bounded below complex of R^a -modules. Then the natural morphism

$$\mathrm{alHom}_{R^a}^{\bullet}(P_2^{\bullet,a}, M^{\bullet,a}) \to \mathrm{alHom}_{R^a}^{\bullet}(P_1^{\bullet,a}, M^{\bullet,a})$$

is an almost quasi-isomorphism.

Proof. We choose some injective resolution $M^{\bullet,a} \to I^{\bullet,a}$ of the bounded below complex $M^{\bullet,a}$. Then we have a commutative diagram

The bottom horizontal arrow is an almost quasi-isomorphism by the standard categorical argument with injective resolutions. The vertical maps are almost quasi-isomorphism by Lemma 2.4.6. \Box

Proposition 2.4.8. (1) There is a natural transformation of functors

$$\mathbf{D}(R)^{op} \times \mathbf{D}(R) \xrightarrow{\mathbf{R} \operatorname{Hom}_{R}(-,-)} \mathbf{D}(R)$$

$$\downarrow^{(-)^{a} \times (-)^{a}} \qquad \downarrow^{(-)^{a}} \qquad \downarrow^{(-)^{a}}$$

$$\mathbf{D}(R)^{a,op} \times \mathbf{D}(R)^{a} \xrightarrow{\mathbf{R} \operatorname{alHom}_{R^{a}}(-,-)} \mathbf{D}(R)^{a}$$

that makes the diagram 2-commutative. In particular,

$$\mathbf{R}$$
al $\operatorname{Hom}_{R^a}(M^a, N^a) \cong^a \mathbf{R}\operatorname{Hom}_R(M, N)^a$

for any $M, N \in \mathbf{D}(R)$.

- (2) For any chosen $M^a \in \mathbf{Mod}_R^a$, the functor $\mathbf{RalHom}_{R^a}(M^a, -) \colon \mathbf{D}(R)^a \to \mathbf{D}(R)^a$ is isomorphic to the (right) derived functor of $\mathrm{alHom}_{R^a}(M^a, -)$.
- (3) For any chosen $N^a \in \mathbf{Mod}_R^a$, the functor $\mathbf{RalHom}_{R^a}(-, N^a) \colon \mathbf{D}^-(R)^{a,op} \to \mathbf{D}(R)^a$ is isomorphic to the (right) derived functor of $\mathrm{alHom}_{R^a}(-, N^a)$.

⁵Here we use that $P^{\bullet,a}$ is bounded above, $M^{\bullet,a}$ and $N^{\bullet,a}$ are bounded below

Proof. In order to show Part (1), we construct functorial morphisms

$$\rho_{M,N} \colon \mathbf{R}\mathrm{Hom}_R(M,N)^a \to \mathbf{R}\mathrm{a}\mathrm{Hom}_{R^a}(M^a,N^a)$$
.

for any $M, N \in \mathbf{D}(R)$. We recall that there is a functorial identification

$$\mathbf{R}$$
al $\operatorname{Hom}_{R^a}(M^a, N^a) \cong^a \mathbf{R}\operatorname{Hom}_R(M^a_!, N)^a \cong^a \mathbf{R}\operatorname{Hom}_R(\widetilde{\mathfrak{m}} \otimes_R M, N)^a)$

So we define

$$\rho_{M,N} \colon \mathbf{R}\mathrm{Hom}_R(M,N)^a \to \mathbf{R}\mathrm{Hom}_R(\widetilde{\mathfrak{m}} \otimes_R M,N)^a$$

as the morphism induced by the canonical map $\widetilde{\mathfrak{m}} \otimes_R M \to M$. This is clearly functorial, so it defines the natural transformation of functors. The only thing we are left to show is that $\rho_{M,N}$ is an almost isomorphism for any $M, N \in \mathbf{D}(R)$.

Let us recall that the way we compute $\mathbf{R}\operatorname{Hom}_R(M, N)$. It is isomorphic to $\operatorname{Hom}_R^{\bullet}(C^{\bullet}, I^{\bullet})$ for any choice of a K-injective resolution of $N \xrightarrow{\sim} I^{\bullet}$ and any resolution $M \xrightarrow{\sim} C^{\bullet}$. Since $\widetilde{\mathfrak{m}} \otimes_R C^{\bullet}$ is a resolution of $\widetilde{\mathfrak{m}} \otimes_R M$ by R-flatness of $\widetilde{\mathfrak{m}}$, we reduce the question to show that the natural map

$$\operatorname{Hom}_R^{\bullet}(C^{\bullet}, I^{\bullet}) \to \operatorname{Hom}_R^{\bullet}(\widetilde{\mathfrak{m}} \otimes_R C^{\bullet}, I^{\bullet})$$

is an almost quasi-isomorphism of complexes. We actually show more, we show that it is an almost isomorphism of complexes. Indeed, the degree n part of this map is the map

$$\prod_{p+q=n} \operatorname{Hom}_R(C^{-q}, I^p) \to \prod_{p+q=n} \operatorname{Hom}_R(\widetilde{\mathfrak{m}} \otimes_R C^{-q}, I^p)$$

Since the (infinite) product is an exact functor in \mathbf{Mod}_R^a , and any (infinite) product of almost zero modules is almost zero, it is actually sufficient to show that each particular map $\operatorname{Hom}_R(C^{-q}, I^p) \to \operatorname{Hom}_R(\widetilde{\mathfrak{m}} \otimes_R C^{-q}, I^p)$ is an almost isomorphism. This follows from Proposition 2.2.1(3).

Part (2) is similar to that of Proposition 2.4.3.

Part (3) is also similar to Part (2) of Proposition 2.4.3, but there are some subtleties due to the fact that \mathbf{Mod}_R^a does not have enough projective objects. We fix this issue by using instead [Sta21, Tag 06XN] of [Sta21, Tag 070K]. We apply it to the subset \mathcal{P} being the set of bounded above complexes with almost projective terms. This result is indeed applicable in our situation due to Corollary 2.2.7 and Lemma 2.4.7.

Now we deal with the case of the derived tensor product functor.

Definition 2.4.9. We say that a complex of R^a -module $K^{\bullet,a}$ is almost K-flat if the naive tensor product complex $C^{\bullet,a} \otimes_{R^a}^{\bullet} K^{\bullet,a}$ is acyclic for any acyclic complex $C^{\bullet,a}$ of R^a -modules

Lemma 2.4.10. The functor $(-)^a$: $\mathbf{Comp}(R) \to \mathbf{Comp}(R^a)$ sends K-flat R-complexes to almost K-flat R^a -complexes.

Proof. Suppose that $C^{\bullet,a}$ is an acyclic complex of \mathbb{R}^a -modules and K^{\bullet} is a K-flat compelx. Then we see that

$$C^{\bullet,a} \otimes_{R^a}^{\bullet} K^{\bullet,a} \cong^a (C^{\bullet} \otimes_R^{\bullet} K^{\bullet})^a \cong^a (\widetilde{\mathfrak{m}} \otimes_R C^{\bullet} \otimes_R^{\bullet} K^{\bullet})^a \cong^a ((\widetilde{\mathfrak{m}} \otimes_R C^{\bullet}) \otimes_R^{\bullet} K^{\bullet})^a .$$

The latter complex is acyclic as $\widetilde{\mathfrak{m}} \otimes C^{\bullet}$ is acyclic and K^{\bullet} is K-flat.

Corollary 2.4.11. Every object $M^{\bullet,a} \in \mathbf{Comp}(\mathbb{R}^a)$ is quasi-isomorphic to an almost K-flat complex.

Proof. We know that the complex $M^{\bullet} \in \mathbf{Comp}(\mathbb{R}^{a})$ is quasi-isomorphic to a K-flat complex K^{\bullet} by [Sta21, Tag 06Y4]. Now we use Lemma 2.4.10 to say that $K^{\bullet,a}$ is almost K-flat complex that is quasi-isomorphic to $M^{\bullet,a}$.

Definition 2.4.12. We define the *derived tensor product functor*

$$-\otimes_{R^a}^L -: \mathbf{D}(R)^a \times \mathbf{D}(R)^a \to \mathbf{D}(R)^a$$

by the rule $(M^a, N^a) \mapsto (M_! \otimes_R^L N_!)^a$ for any $M^a, N^a \in \mathbf{D}(R)^a$.

Proposition 2.4.13. (1) There is a natural transformation of functors

that makes the diagram 2-commutative. In particular, there is a functorial isomorphism $(M \otimes_R^L N)^a \simeq M^a \otimes_{R^a}^L N^a$ for any $M, N \in \mathbf{D}(R)$.

(2) For any chosen $M^a \in \mathbf{Mod}_R^a$, the functor $M^a \otimes_{R^a}^L -: \mathbf{D}(R)^a \to \mathbf{D}(R)^a$ is isomorphic to the (left) derived functor of $M^a \otimes_{R^a} -$.

Proof. The proof of Part (1) is similar to that of Lemma 2.2.1(1). We leave details to the reader.

The proof of Part (2) is similar to that of Proposition 2.4.8(2). The claim follows by applying [Sta21, Tag 06XN] with \mathcal{P} being the subset of almost K-flat complexes. This result is indeed applicable in our situation due to Corollary 2.4.11 and the almost version of [Sta21, Tag 064L]. \Box

Lemma 2.4.14. Let $M^a, N^a, K^a \in \mathbf{D}(R)^a$, then we have a functorial isomorphism

$$\mathbf{R}\operatorname{Hom}_{R^{a}}(M^{a}\otimes_{R^{a}}^{L}N^{a},K^{a})\simeq\mathbf{R}\operatorname{Hom}_{R^{a}}(M^{a},\mathbf{R}\operatorname{a}\operatorname{Hom}_{R^{a}}(N^{a},K^{a}))$$

In particular, the functors **R**alHom_{R^a} $(N^a, -)$: **D** $(R)^a \xrightarrow{}$ **D** $(R)^a$: $-\otimes_{R^a}^L N^a$ are adjoint.

Proof. The claim follows from the following sequence of canonical identifications:

$$\begin{aligned} \mathbf{R} \operatorname{Hom}_{R^{a}}(M^{a} \otimes_{R^{a}}^{L} N^{a}, K^{a}) &\simeq \mathbf{R} \operatorname{Hom}_{R}((\widetilde{\mathfrak{m}} \otimes_{R} M) \otimes_{R}^{L} (\widetilde{\mathfrak{m}} \otimes_{R} N), K) & \text{Lemma 2.4.3(1)} \\ &\simeq \mathbf{R} \operatorname{Hom}_{R}(\widetilde{\mathfrak{m}} \otimes_{R} M, \mathbf{R} \operatorname{Hom}_{R}(\widetilde{\mathfrak{m}} \otimes_{R} N, K)) & [\text{Sta21, Tag 0A5W}] \\ &\simeq \mathbf{R} \operatorname{Hom}_{R^{a}}(M^{a}, \mathbf{R} \operatorname{Hom}_{R}(\widetilde{\mathfrak{m}} \otimes_{R} N, K)^{a}) & \text{Lemma 2.4.3(1)} \\ &\simeq \mathbf{R} \operatorname{Hom}_{R^{a}}(M^{a}, \mathbf{R} \operatorname{alHom}_{R^{a}}(N^{a}, K^{a})) & \text{Definition 2.4.4} \end{aligned}$$

Definition 2.4.15. Let $f: R \to S$ be a ring homomorphism. We define the base change functor $- \otimes_{R^a}^L S^a: \mathbf{D}(R)^a \to \mathbf{D}(S)^a$

by the rule $M^a \mapsto (M_! \otimes_R^L S)^a$ for any $M^a \in \mathbf{D}(R)^a$.

Proposition 2.4.16. (1) There is a natural transformation of functors

$$\mathbf{D}(R) \xrightarrow{-\otimes_R^L S} \mathbf{D}(S)$$

$$(-)^a \downarrow \xrightarrow{\rho} \qquad \downarrow (-)^a$$

$$\mathbf{D}(R)^a \xrightarrow{\otimes_{R^a} S^a} \mathbf{D}(S)^a$$

that makes the diagram 2-commutative. In particular, there is a functorial isomorphism $(M \otimes_R^L S)^a \simeq M^a \otimes_{R^a}^L S^a$ for any $M \in \mathbf{D}(R)$.

(2) The functor $-\otimes_{R^a}^L S^a \colon \mathbf{D}(R)^a \to \mathbf{D}(S)^a$ is isomorphic to the (left) derived functor of $-\otimes_{R^a}^L S^a$.

Proof. The proof is identical to Proposition 2.4.13.

Lemma 2.4.17. Let $R \to S$ be a ring homomorphism, and let $M^a \in \mathbf{D}(R)^a$, $N^a \in \mathbf{D}(S)^a$. Then we have a functorial isomorphism

$$\mathbf{R}\operatorname{Hom}_{S^a}(M^a \otimes_{R^a}^L S^a, N^a) \simeq \mathbf{R}\operatorname{Hom}_{R^a}(M^a, N^a)$$
.

In particular, the functors Forget: $\mathbf{D}(S)^a \xrightarrow{\longleftarrow} \mathbf{D}(R)^a \colon -\otimes_{R^a}^L S^a$ are adjoint.

Proof. The proof is similar to that of Lemma 2.4.14.

2.5. Almost Finitely Generated and Almost Finitely Presented Modules. We discuss the notions of almost finitely generated and almost finitely presented modules in section. The discussion follows [GR03] closely. The main difference is that we avoid any use of "uniform structures" in our treatment, we think that it simplifies the exposition. We recall that we fixed some "base" ring R with an ideal \mathfrak{m} such that $\mathfrak{m}^2 = \mathfrak{m}$ and $\tilde{\mathfrak{m}} = \mathfrak{m} \otimes_R \mathfrak{m}$ is flat, and we always do almost mathematics with respect to this ideal.

Definition 2.5.1. An *R*-module *M* is called *almost finitely generated*, if for any $\varepsilon \in \mathfrak{m}$ there is an integer n_{ε} and an *R*-homomorphism

$$R^{n_{\varepsilon}} \xrightarrow{J} M$$

such that $\operatorname{Coker}(f)$ is killed by ε .

Definition 2.5.2. An *R*-module *M* is called *almost finitely presented*, if for any $\varepsilon, \delta \in \mathfrak{m}$ there are integers $n_{\varepsilon,\delta}$, $m_{\varepsilon,\delta}$ and a complex

$$R^{m_{\varepsilon,\delta}} \xrightarrow{g} R^{n_{\varepsilon,\delta}} \xrightarrow{J} M$$

such that $\operatorname{Coker}(f)$ is killed by ε and $\delta(\ker f) \subset \operatorname{Im} g$.

Remark 2.5.3. Clearly, any almost finitely presented *R*-module is almost finitely generated.

Remark 2.5.4. A typical example of an almost finitely presented module that is not finitely generated is $M = \bigoplus_{n\geq 1} \mathcal{O}_C / p^{1/n} \mathcal{O}_C$ for an algebraically closed non-archimedean field C of mixed characteristic (0, p).

The next few lemmas discuss the most basic properties of almost finitely generated and almost finitely presented modules. For example, it is not entirely obvious that these notions transfer across almost isomorphisms. We show that this is actually the case, so these notions descend to \mathbf{Mod}_R^a . We also show that almost finitely generated and almost finitely presented modules have many good properties that we have for the usual finitely generated and finitely presented modules. Although all proofs below are elementary, they require some accuracy to rigorously prove them (and some proofs are actually tricky).

Our first main goal is to get some other useful criteria for a module to be almost finitely generated (resp. almost finitely presented) and finally show that this notion does not depend on a class of almost isomorphism.

Lemma 2.5.5. Let M be an R-module, then M is almost finitely generated if and only if for any finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$ there a morphism $R^n \xrightarrow{f} M$ such that $\mathfrak{m}_0(\operatorname{Coker} f) = 0$.

Proof. The "if" part is clear, so we only need to deal with the "only if" part. We choose a set of generators $(\varepsilon_0, \ldots, \varepsilon_n)$ for an ideal \mathfrak{m}_0 . Then we have *R*-morphisms

$$f_i \colon \mathbb{R}^{n_{\varepsilon_i}} \to M$$

such that $\varepsilon_i(\operatorname{Coker} f_i) = 0$ for all *i*. Then the sum of these morphisms

$$f \coloneqq \bigoplus_{i=1}^{n} f_i \colon R^{\sum n_{\varepsilon_i}} \to M$$

defines a map such that $\mathfrak{m}_0(\operatorname{Coker} f) = 0$. Since \mathfrak{m}_0 was an arbitrary morphism, this finishes the proof.

Lemma 2.5.6. Let M be an almost finitely presented R-module, and let $\varphi : \mathbb{R}^n \to M$ be an R-homomorphism such that $\mathfrak{m}_1(\operatorname{Coker} \varphi) = 0$ for some ideal $\mathfrak{m}_1 \subset \mathfrak{m}$. Then for every finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}_1\mathfrak{m}$ there is morphism $\psi : \mathbb{R}^m \to M$ such that

$$R^m \xrightarrow{\psi} R^n \xrightarrow{\varphi} M$$

is a three-term complex and $\mathfrak{m}_0(\operatorname{Ker} \varphi) \subset \operatorname{Im}(\psi)$.

Proof. Since M is almost finitely presented, for any two elements $\varepsilon_1, \varepsilon_2 \in \mathfrak{m}$, we can find a complex

$$R^{m_2} \xrightarrow{g} R^{m_1} \xrightarrow{f} M$$

such that $\varepsilon_1(\operatorname{Coker} f) = 0$ and $\varepsilon_2(\ker f) \subset \operatorname{Im} g$. Now we choose some element $\delta \in \mathfrak{m}_1$, and we shall define morphisms

$$\alpha: \mathbb{R}^{m_1} \to \mathbb{R}^n \text{ and } \beta: \mathbb{R}^n \to \mathbb{R}^m$$

such that $\varphi \circ \alpha = \delta f$ and $f \circ \beta = \varepsilon_1 \varphi$. Here is the corresponding picture:

$$\begin{array}{cccc} R^{m_2} & \xrightarrow{g} & R^{m_1} & \xrightarrow{f} & M \\ & & & & & & & \\ & & & & & \\ & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & \\ & & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & &$$

We define α and β in the following way: we fix a basis e_1, \ldots, e_{m_1} of \mathbb{R}^{m_1} and a basis e'_1, \ldots, e'_n of \mathbb{R}^n , then we define

 $\alpha(e_i) = y_i \in \mathbb{R}^n$ for some y_i such that $\varphi(y_i) = \delta f(e_i)$,

$$\beta(e'_i) = x_i \in \mathbb{R}^{m_1}$$
 for some x_i such that $f(x_i) = \varepsilon_1 \varphi(e'_i)$

and then extend these maps by linearity. It is clear that $\varphi \circ \alpha = \delta f$ and $f \circ \beta = \varepsilon_1 \varphi$ as it holds on basis elements.

Now we can define a morphism $\psi: \mathbb{R}^n \oplus \mathbb{R}^{m_2} \to \mathbb{R}^n$ by the rule

$$\psi(x,y) = \alpha \circ \beta(x) - (\varepsilon_1 \delta)x + \alpha \circ g(y).$$

We now show that

$$\varphi \circ \psi = 0$$
 and $\varepsilon_1 \varepsilon_2 \delta \operatorname{Ker} \varphi \subset \operatorname{Im} \psi$

We start by showing that $\varphi \circ \psi = 0$: it suffices to prove that

$$(\alpha \circ g)(y) \in \operatorname{Ker} \varphi \text{ for } y \in \mathbb{R}^{m_2}, \text{ and } (\alpha \circ \beta)(x) - (\varepsilon_1 \delta)x \in \operatorname{Ker} \varphi \text{ for } x \in \mathbb{R}^n$$

We note that we have an equality

$$(\varphi \circ \alpha \circ g)(y) = \delta(f \circ g)(y) = \delta 0 = 0,$$

so $(\alpha \circ g)(y) \in \operatorname{Ker}(\varphi)$. We also have an equality

$$\begin{aligned} (\varphi \circ (\alpha \circ \beta - \varepsilon_1 \delta))(x) &= (\varphi \circ \alpha \circ \beta)(x) - \varepsilon_1 \delta \varphi(x) \\ &= \delta(f \circ \beta)(x) - \varepsilon_1 \delta \varphi(x) \\ &= \delta \varepsilon_1 \varphi(x) - \varepsilon_1 \delta \varphi(x) \\ &= 0. \end{aligned}$$

this shows that $(\alpha \circ \beta)(x) - (\varepsilon_1 \delta)x \in \operatorname{Ker}(\varphi)$ as well.

We show that $(\varepsilon_1 \varepsilon_2 \delta) \operatorname{Ker} \varphi \subset \operatorname{Im}(\psi)$: we observe that for any $x \in \operatorname{Ker} \varphi$ we have $\beta(x) \subset \operatorname{Ker} f$ as $f \circ \beta = \varepsilon_1 \varphi$. This implies that $\varepsilon_2 \beta(x) \in \operatorname{Im} g$ since $\varepsilon_2 \operatorname{Ker} f \subset \operatorname{Im} g$. Thus there is $y \in \mathbb{R}^{m_2}$ such that $g(y) = \varepsilon_2 \beta(x)$, so $(\alpha \circ g)(y) = \varepsilon_2 \alpha \circ \beta(x)$. This shows that

$$\psi(-\varepsilon_2 x, y) = -\varepsilon_2(\alpha \circ \beta)(x) + \varepsilon_1 \varepsilon_2 \delta x + (\alpha \circ g)(y) = \\ -\varepsilon_2(\alpha \circ \beta)(x) + \varepsilon_1 \varepsilon_2 \delta x + \varepsilon_2(\alpha \circ \beta)(x) = \varepsilon_1 \varepsilon_2 \delta x$$

We conclude that $\varepsilon_1 \varepsilon_2 \delta x \in \operatorname{Im}(\psi)$ for any $x \in \operatorname{Ker}(\varphi)$.

Finally, we recall that \mathfrak{m}_0 is a finitely generated ideal, and that $\mathfrak{m}_0 \subset \mathfrak{m}_1 \mathfrak{m} = \mathfrak{m}_1 \mathfrak{m}^2 \subset \mathfrak{m}_1$. This means that we can find a finite set I, and a finite set of elements $\varepsilon_{i,1}, \varepsilon_{i,2} \in \mathfrak{m}, \delta_i \in \mathfrak{m}_1$ such that \mathfrak{m}_0 is contained is the ideal $J := (\varepsilon_{i,1}\varepsilon_{i,2}\delta_i)_{i\in I}$ (the ideal generated by all the products $\varepsilon_{i,1}\varepsilon_{i,2}\delta_i$). The previous discussion implies that for each $i \in I$, we have a map $\psi_i : \mathbb{R}^{k_i} \to \mathbb{R}^n$ such that $\varphi \circ \psi_i = 0$ and $(\varepsilon_{i,1}\varepsilon_{i,2}\delta_i)(\operatorname{Ker} \varphi) \subset \operatorname{Im} \psi_i$. By passing to the homomorphism

$$\psi \coloneqq \bigoplus_{i \in I} \psi_i \colon R^{\sum k_i} \to R^n$$

we get a map ψ such that $\varphi \circ \psi = 0$ and $\mathfrak{m}_0(\operatorname{Ker} \varphi) \subset \operatorname{Im}(\psi)$. Therefore ψ does the job.

Lemma 2.5.7. Let M be an R-module. Then the following conditions are equivalent:

- (1) The R-module M is almost finitely presented.
- (2) For any finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$ there exist a finitely presented *R*-module *N* and a homomorphism $f: N \to M$ such that $\mathfrak{m}_0(\ker f) = 0$ and $\mathfrak{m}_0(\operatorname{Coker} f) = 0$.
- (3) For any finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$ there exist integers n, m and a three-term complex

$$R^m \xrightarrow{g} R^n \xrightarrow{f} M$$

such that $\mathfrak{m}_0(\operatorname{Coker} f) = 0$ and $\mathfrak{m}_0(\operatorname{Ker} f) \subset \operatorname{Im} g$.

Proof. It is clear that the condition (3) implies both conditions (1) and (2).

We show that (1) implies (3). Since M is an almost finitely generated R-module, Lemma 2.5.5 guarantees that for any finitely generated ideal $\mathfrak{m}' \subset \mathfrak{m}$ there is a morphism $R^n \xrightarrow{f} M$ such that $\mathfrak{m}'(\operatorname{Coker} f) = 0$.

We know that $\mathfrak{m}_0 \subset \mathfrak{m} = \mathfrak{m}^2$, this easily implies that there is a finitely generated ideal $\mathfrak{m}_1 \subset \mathfrak{m}$ such that $\mathfrak{m}_0 \subset \mathfrak{m}_1 \mathfrak{m} \subset \mathfrak{m}_1$. So, using $\mathfrak{m}' = \mathfrak{m}_1$, we can find a homomorphism $R^n \xrightarrow{\varphi} M$ such that $\mathfrak{m}_1(\operatorname{Coker} \varphi) = 0$. Lemma 2.5.6 claims that we can also find a homomorphism $\psi \colon R^m \to R^n$ such that

$$R^m \xrightarrow{\psi} R^n \xrightarrow{\varphi} M$$

is a three-term complex and $\mathfrak{m}_0(\ker \varphi) \subset \operatorname{Im} \psi$. Since $\mathfrak{m}_0 \subset \mathfrak{m}_1$ and $\mathfrak{m}_1(\operatorname{Coker} \varphi) = 0$, we get that $\mathfrak{m}_0(\operatorname{Coker} \varphi) = 0$ as well. This finishes the proof since \mathfrak{m}_0 was an arbitrary finitely generated sub-ideal of \mathfrak{m} .

Now we show that (2) implies (3). We pick an arbitrary finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$, and we try to find a three-term complex

$$R^m \xrightarrow{g} R^n \xrightarrow{f} M$$

such that $\mathfrak{m}_0(\operatorname{Coker} f) = 0$ and $\mathfrak{m}_0(\ker f) \subset \operatorname{Im}(g)$. In order to achieve this we use the assumption (2) to find a morphism $h: N \to M$ such that N is a finitely presented R-module, $\mathfrak{m}_0(\operatorname{Coker} h) = 0$, and $\mathfrak{m}_0(\ker h) = 0$. Since N is finitely presented we can find a short exact sequence

$$R^m \xrightarrow{g} R^n \xrightarrow{f'} N \to 0$$

It is straightforward to see that a three-term complex

$$R^m \xrightarrow{g} R^n \xrightarrow{f := h \circ f'} M$$

satisfies the condition that $\mathfrak{m}_0(\operatorname{Coker} f) = 0$ and $\mathfrak{m}_0(\ker f) \subset \operatorname{Im}(g)$.

Lemma 2.5.8. Let M be an R-module, and suppose that for any finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$ there exists a morphism $f: N \to M$ such that $\mathfrak{m}_0(\ker f) = 0$, $\mathfrak{m}_0(\operatorname{Coker} f) = 0$ and N is almost finitely generated (resp. almost finitely presented). Then M is also almost finitely generated (resp. almost finitely presented).

Proof. We give a proof only in the almost finitely presented case; the other case is easier. We pick an arbitrary finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$ and another finitely generated ideal $\mathfrak{m}_1 \subset \mathfrak{m}$ such that $\mathfrak{m}_0 \subset \mathfrak{m}_1^2$. Then we use the assumption to get a morphism

$$f: N \to M$$

such that $\mathfrak{m}_1(\operatorname{Ker} f) = 0, \mathfrak{m}_1(\operatorname{Coker} f) = 0$ and N is an almost finitely presented R-module. Lemma 2.5.7 guarantees that there is a three-term complex

$$R^m \xrightarrow{h} R^n \xrightarrow{g} N$$

such that $\mathfrak{m}_1(\operatorname{Coker} g) = 0$ and $\mathfrak{m}_1(\operatorname{Ker} g) \subset \operatorname{Im} h$. Then we can consider a three-term complex

$$R^m \xrightarrow{h} R^n \xrightarrow{f':=f \circ g} M,$$

it is easily seen that $\mathfrak{m}_1^2(\operatorname{Coker} f') = 0$ and $\mathfrak{m}_1^2(\ker f') \subset \operatorname{Im}(h)$. Since $\mathfrak{m}_0 \subset \mathfrak{m}_1^2$ we conclude that $\mathfrak{m}_0(\operatorname{Coker} f') = 0$ and $\mathfrak{m}_0(\ker f') \subset \operatorname{Im}(h)$. This shows that M is almost finitely presented. \Box

Corollary 2.5.9. Let M and N be two almost isomorphic R-modules (see Definition 2.1.7). Then M is almost finitely generated (resp. almost finitely presented) if and only if so is N.

Proof. Lemma 2.1.8 implies that $\widetilde{\mathfrak{m}} \otimes_R M \simeq \widetilde{\mathfrak{m}} \otimes_R N$. So in order to prove the claim, it suffices to show that an *R*-module is almost finitely generated (resp. almost finitely presented) if $\widetilde{\mathfrak{m}} \otimes_R M$ is. Now Lemma 2.5.8 shows that if $\widetilde{\mathfrak{m}} \otimes_R M$ is almost finitely generated (resp. almost finitely presented), then so is M.

Now we suppose that M is almost finitely generated and show that $\mathfrak{m} \otimes_R M$ is as well. It actually suffices to show that $\mathfrak{m} \otimes_R M$ is almost finitely generated because

$$\widetilde{\mathfrak{m}} \otimes_R M \simeq \mathfrak{m} \otimes_R \mathfrak{m} \otimes_R M,$$

so we show that $\mathfrak{m} \otimes_R M$ is almost finitely generated if so is M.

We choose a finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$ and another finitely generated ideal $\mathfrak{m}_1 \subset \mathfrak{m}$ such that $\mathfrak{m}_0 \subset \mathfrak{m}_1^3$. Then we apply Lemma 2.5.5 to find an *R*-homomorphism $f : M' \to M$ such that M' is finitely generated, $\mathfrak{m}_1(\operatorname{Coker} f) = 0$ and $\mathfrak{m}_1(\operatorname{Ker} f) = 0$. We define a morphism $g : \mathfrak{m}_1 \otimes_R M' \to \mathfrak{m} \otimes_R M$ as the *R*-homomorphism induced by f and the natural inclusion $\mathfrak{m}_1 \to \mathfrak{m}$. So we have a commutative diagram:

$$\mathfrak{m}_1 \otimes_R M' \xrightarrow{g} \mathfrak{m} \otimes_R M \\ \downarrow^h \qquad \qquad \downarrow^p \\ M' \xrightarrow{f} M.$$

Note that all kernels and cokernels of f, h and p are killed by \mathfrak{m}_1 . This is straightforward to see that it implies that kernel and cokernel of g are killed by \mathfrak{m}_1^3 . In particular, they are killed by \mathfrak{m}_0 by the choice of \mathfrak{m}_1 . Since \mathfrak{m}_1 and M' are finitely generated, their tensor product is finitely generated as well. Therefore, we conclude that $\mathfrak{m} \otimes_R M$ is almost finitely generated by Lemma 2.5.5.

Now we deal with the almost finitely presented case. As in the finitely generated case, we only need to show that $\mathfrak{m} \otimes_R M$ is almost finitely presented if so is M. We start by choosing a finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$ and another finitely generated ideal $\mathfrak{m}_1 \subset \mathfrak{m}$ such that $\mathfrak{m}_0 \subset \mathfrak{m}_1^3$. We use Lemma 2.5.7 to get a morphism $f: M' \to M$ such that M' is finitely presented, $\mathfrak{m}_1(\operatorname{Coker} f) = 0$ and $\mathfrak{m}_1(\operatorname{Ker} f) = 0$. However, there is a problem that even though M' is finitely presented we cannot control finite presentation of $\mathfrak{m}_1 \otimes_R M'$ because the ideal \mathfrak{m}_1 is not necessary finitely presented. Instead we do a better trick (following the ideas from [GR03, page 19]).

We choose some generators $(\varepsilon_1, \varepsilon_2, \ldots, \varepsilon_n)$ of the ideal \mathfrak{m}_1 and we write a truncated in degree 2 Koszul complex $K_{\bullet}(\varepsilon_1, \ldots, \varepsilon_n)$. Explicitly this is just a four-term complex

$$\wedge^2 R^n \xrightarrow{\partial_2} R^n \xrightarrow{\partial} R \to 0,$$

where ∂_2 is defined as the unique *R*-homomorphism such that

$$\partial_2(e_i \wedge e_j) = \varepsilon_j e_i - \varepsilon_i e_j$$

for the standard basis (e_1, \ldots, e_n) of a free module \mathbb{R}^n , and ∂ is defined as the unique \mathbb{R} -linear homomorphism such that

$$\partial(e_i) = \varepsilon_i$$

We note that $\operatorname{Im}(\partial) = \mathfrak{m}_1$. And we define $\mathfrak{m}'_1 := \operatorname{Coker}(\partial_2)$, so the differential ∂ induces a morphism $\partial : \mathfrak{m}'_1 \to \mathfrak{m}_1$. We also have an *R*-linear homomorphism $\overline{e_i} : R \to \mathfrak{m}'_1$ defined as

 $\overline{e_i}(x) = [xe_i] \in \mathfrak{m}'_1$, where e_i is an element of a standard basis and $x \in R$.

Then we claim that the composition

$$\mathfrak{m}_1' \xrightarrow{\partial} \mathfrak{m}_1 \hookrightarrow R \xrightarrow{\overline{e_i}} \mathfrak{m}_1'$$

is equal the multiplication by ε_i . Indeed, it suffices to show it on generators $[e_i]$. Then we have

$$\overline{e_i}(\partial(e_j)) = \overline{e_i}(\varepsilon_j) = [\varepsilon_j e_i] = [\varepsilon_i e_j],$$

where the last equality comes from the definition of \mathfrak{m}'_1 . This implies that we have a diagram

$$\mathfrak{m}'_1 \otimes_R M' \xrightarrow{\partial \otimes_R M'} M' \longrightarrow \mathfrak{m}'_1 \otimes_R M'.$$

for each i. Therefore, we see that $\mathfrak{m}_1(\ker h) = 0$ for $h \coloneqq \partial \otimes_R M'$. We also have a composition

$$R \xrightarrow{\overline{e_i}} \mathfrak{m}'_1 \xrightarrow{\partial} \mathfrak{m}_1 \hookrightarrow R$$

that is easily seen to be equal to the multiplication by ε_i . The similar argument guarantees that $\mathfrak{m}_1(\operatorname{Coker} h) = 0$. Again, we consider the commutative square:

$$\mathfrak{m}'_1 \otimes_R M' \xrightarrow{g} \mathfrak{m} \otimes_R M$$
$$\downarrow^h \qquad \qquad \downarrow^p$$
$$M' \xrightarrow{f} M$$

where all kernels and cokernels of f, h and p are annihilated by \mathfrak{m}_1 . But now the advantage is that \mathfrak{m}'_1 is finitely presented by the construction. So the tensor product $\mathfrak{m}'_1 \otimes_R M'$ is also finitely presented. Then the same argument as in the finitely generated case implies that $\mathfrak{m}_0(\ker g) = 0$ and $\mathfrak{m}_0(\operatorname{Coker} g) = 0$. Since \mathfrak{m}_0 was an arbitrary finitely generated ideal in \mathfrak{m} , we conclude that $\mathfrak{m} \otimes_R M$ is almost finitely presented. \Box

Definition 2.5.10. We say that an R^a -module $M^a \in \mathbf{Mod}_R^a$ is almost finitely generated (resp. almost finitely presented) if its representative $M \in \mathbf{Mod}_R$ is almost finitely generated (resp. almost finitely presented). This definition does not depend on a choice of representative by Lemma 2.5.9

We now want to establish certain good properties of almost finitely presented modules in short exact sequences. This will be crucial later to develop a good theory of almost coherent modules.

Lemma 2.5.11. Let $0 \to M' \xrightarrow{\varphi} M \xrightarrow{\psi} M'' \to 0$ be an exact sequence of *R*-modules, then

- (1) If M is almost finitely generated, then so is M''.
- (2) If M' and M'' are almost finitely generated (resp. finitely presented), then so is M.
- (3) If M is almost finitely generated and M'' is almost finitely presented, then M' is almost finitely generated.
- (4) If M is almost finitely presented and M' is almost finitely generated, then M'' is almost finitely presented.

Proof. (1): We suppose that M is almost finitely generated. This means that for any $\varepsilon \in \mathfrak{m}$ there is an integer n_{ε} and a morphism $\mathbb{R}^{n_{\varepsilon}} \xrightarrow{f} M$ such that $\varepsilon(\operatorname{Coker} f) = 0$. We define $f'': \mathbb{R}^{n_{\varepsilon}} \to M''$ as the composition $\psi \circ f$. It is easy to see that $\varepsilon(\operatorname{Coker} f'') = 0$, so M' is indeed almost finitely generated.

(2): Firstly, we deal with almost finitely generated case. So, we suppose that M' and M'' are almost finitely generated *R*-modules, and we show that so is M.

We choose any finitely generated ideal \mathfrak{m}_0 and a finitely generated ideal \mathfrak{m}_1 such that $\mathfrak{m}_0 \subset \mathfrak{m}_1^2$. Now we apply Lemma 2.5.5 to find morphisms

$$f' \colon \mathbb{R}^{n_1} \to M' \text{ and } \overline{f''} \colon \mathbb{R}^{n_2} \to M''$$

such that $\mathfrak{m}_1(\operatorname{Coker} f') \simeq \mathfrak{m}_1(\operatorname{Coker} \overline{f''}) \simeq 0$. We lift $\overline{f''}$ in an arbitrary way to a morphism $f'': \mathbb{R}^{n_2} \to M$, so we have a commutative diagram:

An easy application of the Snake Lemma implies that $\mathfrak{m}_1^2(\operatorname{Coker}(f' \oplus f'')) = 0$. In particular, we have $\mathfrak{m}_0(\operatorname{Coker}(f' \oplus f'') = 0$. Thus we conclude that M is almost finitely generated as \mathfrak{m}_0 was an arbitrary finitely generated subideal of \mathfrak{m} .

Now we deal with the finitely presented case. The idea will be similar. We choose a finitely generated ideal \mathfrak{m}_0 and another finitely generated ideal $\mathfrak{m}_1 \subset \mathfrak{m}$ such that $\mathfrak{m}_0 \subset \mathfrak{m}_1^3$. Then we use Lemma 2.5.7 to find complexes

$$R^{m_1} \xrightarrow{g'} R^{n_1} \xrightarrow{f'} M'$$
, and $R^{m_2} \xrightarrow{\overline{g''}} R^{n_2} \xrightarrow{\overline{f''}} M''$

such that $\mathfrak{m}_1(\operatorname{Coker}(f')) \simeq \mathfrak{m}_1(\operatorname{Coker}(\overline{f''})) \simeq 0$, $\mathfrak{m}_1(\ker f') \subset \operatorname{Im}(g')$ and $\mathfrak{m}_1(\ker \overline{f''}) \subset \operatorname{Im}(\overline{g''})$. We lift $\overline{f''}$ in an arbitrary way to a morphism $f'': \mathbb{R}^{n_2} \to M$, so we have a commutative diagram:

We use the Snake Lemma to get an exact sequence:

$$0 \to \ker f' \to \ker f' \oplus f'' \to \ker \overline{f''} \to \operatorname{Coker} f$$

with Coker f' being annihilated by \mathfrak{m}_1 . Now we note that the composition $\overline{g''} \circ \overline{f''} = 0$, so $\overline{g''}$ induces a morphism $\overline{g''}: \mathbb{R}^{m_2} \to \ker f''$. We choose some generators of $\mathfrak{m}_1 = (\varepsilon_1, \ldots, \varepsilon_k)$. Then each of $\varepsilon_j g''(e_i)$ can be lifted to an element $x_{i,j} \in \ker f' \oplus f''$. We define the morphism $g'': \mathbb{R}^{m_2k} \to \ker f' \oplus f''$ as $g''(e_{i,j}) = x_{i,j}$. Then we have a three-term complex

$$R^{n_2} \bigoplus R^{m_2 k} \xrightarrow{g' \oplus g''} R^{n_1} \bigoplus R^{n_2} \xrightarrow{f' \oplus f''} M.$$

It is easy to see that $\mathfrak{m}_1^2(\operatorname{Coker} f' \oplus f'') = 0$ and $\mathfrak{m}_1^3(\ker f' \oplus f'') \subset \operatorname{Im} g' \oplus g''$. In particular, we get that $\mathfrak{m}_0(\operatorname{Coker}(f' \oplus f'')) = 0$ and $\mathfrak{m}_0(\operatorname{Ker}(f' \oplus f'')) \subset \operatorname{Im}(g' \oplus g'')$. Since \mathfrak{m}_0 was an arbitrary finitely generated ideal in \mathfrak{m} , we conclude that M is almost finitely presented.

(3): Now we suppose that M is almost finitely generated and M'' is almost finitely presented, and we want to show that M' is almost finitely generated. We pick some finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$, and another finitely generated ideal $\mathfrak{m}_1 \subset \mathfrak{m}$ such that $\mathfrak{m}_0 \subset \mathfrak{m}_1^3$. We also choose some three arbitrary elements $\varepsilon_1, \varepsilon_2, \varepsilon_3 \in \mathfrak{m}_1$.

Recall that M'' is almost finitely presented, so we can find a complex

$$R^m \xrightarrow{g} R^n \xrightarrow{f} M''$$

such that $\varepsilon_1(\operatorname{Coker} f) = 0$ and $\varepsilon_2(\operatorname{Ker} f) \subset \operatorname{Im} g$. By the universal property of free modules we can extend it to a commutative diagram

Also, we know that M is almost finitely generated, so we can find a homomorphism $\mathbb{R}^k \xrightarrow{r} M$ such that $\varepsilon_3(\operatorname{Coker}(r)) = 0$. We consider elements $x'_1 := r(e_1), \ldots, x'_k := r(e_k)$ with e_i being elements of

the standard basis of R^k . Since Coker f is killed by ε_1 we can find some elements $y_1, y_2, \ldots, y_k \in R^n$ such that $f(y_i) = \varepsilon_1 \psi(x'_i)$. And we define elements

$$x_i := \varepsilon_1 x_i' - \alpha(y_i) \in M.$$

We note that actually $x_i \in M'$ for each i = 1, ..., k (we slightly abuse notations and identify M' with its image in M). We finally define a morphism

$$R^k \oplus R^m \xrightarrow{h} M'$$

as
$$h(a_1e_1 \oplus a_2e_2 \oplus \cdots \oplus a_ke_k \oplus y) = a_1x_1 + a_2x_2 + \cdots + a_kx_k + \beta(y)$$

where a_i are any elements of R, e_i are elements of the standard basis of R^k , and y is an arbitrary element of R^m .

We claim that $(\varepsilon_3\varepsilon_2\varepsilon_1)(\operatorname{Coker} h) = 0$. Let us choose an arbitrary element $x \in M'$ and show that $(\varepsilon_3\varepsilon_2\varepsilon_1)x \in \operatorname{Im} h$. We know that there exist some elements a_1, \ldots, a_k such that $\varepsilon_3 x = \sum_{i=1}^k a_i x'_i$. So we see that

$$\varepsilon_1 \varepsilon_3 x - \sum_{i=1}^k a_i x_i = \sum_{i=1}^k a_i (\varepsilon_1 x_i' - x_i) = \alpha \left(\sum_{i=1}^k a_i y_i \right)$$
(2.1)

by definition of x_i . We also see that

$$f\left(\sum_{i=1}^{k} a_i y_i\right) = (\psi \circ \alpha) \left(\sum_{i=1}^{k} a_i y_i\right) = \psi \left(\varepsilon_1 \varepsilon_3 x - \sum_{i=1}^{k} a_i x_i\right) = 0,$$

where the last equality holds because $\varepsilon_1 \varepsilon_3 x - \sum_{i=1}^k a_i x_i$ lies in M' as x and x_i already lie in M'. This implies that $\varepsilon_2(\sum_{i=1}^k a_i y_i) \in \text{Im}(g)$ by the choice of g and f. Let us say that $\varepsilon_2(\sum_{i=1}^k a_i y_i) = g(b)$.

Finally, we use injectivity of φ and equation 2.1 to get that

$$\varepsilon_{3}\varepsilon_{2}\varepsilon_{1}x = \varepsilon_{2}\left(\sum_{i=1}^{k}a_{i}x_{i}\right) + \alpha(\varepsilon_{2}\sum_{i=1}^{k}a_{i}y_{i})$$
$$= \varepsilon_{2}\left(\sum_{i=1}^{k}a_{i}x_{i}\right) + \alpha(g(b))$$
$$= \varepsilon_{2}\left(\sum_{i=1}^{k}a_{i}x_{i}\right) + \beta(b) \in \operatorname{Im} h.$$

The standard argument as in the proofs of Lemmas 2.5.5 and 2.5.7 shows that there exists a morphism $F : \mathbb{R}^N \to M'$ such that $\mathfrak{m}_1^3(\operatorname{Coker} F) = 0$. In particular, we see that $\mathfrak{m}_0(\operatorname{Coker} F) = 0$, and this implies that M' is almost finitely generated.

(4): We suppose that M is almost finitely presented and M' is almost finitely generated, and we want to show that M'' is also almost finitely presented. We pick an arbitrary finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$, and another finitely generated ideal $\mathfrak{m}_1 \subset \mathfrak{m}$ such that $\mathfrak{m}_0 \subset \mathfrak{m}_1^3$. Lemmas 2.5.5 and 2.5.7 guarantee that there exists a complex

$$R^m \xrightarrow{g} R^n \xrightarrow{f} M$$

and a morphism

 $R^k \xrightarrow{r} M'$

such that $\mathfrak{m}_1(\operatorname{Coker} f) \simeq \mathfrak{m}_1(\operatorname{Coker} r) \simeq 0$ and $\mathfrak{m}_1(\ker f) \subset \operatorname{Im} g$. We define the morphism

 $F\colon \mathbb{R}^n\to M''$

as the composition $\psi \circ f$. It is straightforward to see $\mathfrak{m}_1(\operatorname{Coker} F) = 0$. Now we pick any $\varepsilon \in \mathfrak{m}_1$ and choose any $\alpha \colon \mathbb{R}^k \to \mathbb{R}^n$ such that $f \circ \alpha = \varepsilon \varphi \circ r$, this is possibly as $\varepsilon(\operatorname{Coker} f) = 0$. Then we define $G_{\varepsilon} \colon \mathbb{R}^k \oplus \mathbb{R}^m \to \mathbb{R}^n$ as the direct sum $\alpha \oplus g$. A diagram chase proves that $F \circ G_{\varepsilon} = 0$ and $\varepsilon \mathfrak{m}_1^2 \operatorname{Ker} F \subset \operatorname{Im} G_{\varepsilon}$. The standard argument as in the proofs of Lemmas 2.5.5 and 2.5.7 shows that there exists a morphism $G \colon \mathbb{R}^N \to \mathbb{R}^n$ such that $\mathfrak{m}_1^3(\operatorname{Ker} F) \subset \operatorname{Im} G$. In particular, $\mathfrak{m}_0 \operatorname{Ker} F \subset \operatorname{Im} G$.

Corollary 2.5.12. Let $0 \to M'^a \xrightarrow{\varphi} M^a \xrightarrow{\psi} M''^a \to 0$ be an exact sequence of R^a -modules. Then all the conclusions of Lemma 2.5.11 still hold.

Proof. We use Lemma 2.1.9(4), (5) to see that the sequence

$$0 \to (M'^a)_! \xrightarrow{\varphi_!} (M^a)_! \xrightarrow{\psi_!} (M''^a)_! \to 0$$

is exact and almost isomorphic to the original one. Moreover, Corollary 2.5.9 says that each of those modules $N_!^a$ is almost finitely generated (resp. almost finitely presented) if and only if so is the corresponding N^a . Thus the problem is reduced to Lemma 2.5.11.

Lemma 2.5.13. Let M^a, N^a be two almost finitely generated (resp. almost finitely presented) R^a -modules, then so is $M^a \otimes_{R^a} N^a$. Similarly, $M \otimes_R N$ is almost finitely generated (resp. almost finitely presented) for any almost finitely generated (resp. almost finitely presented) R-modules M and N.

Proof. We show the claim only in the case of almost finitely presented modules, the case of almost finitely generated modules is significantly easier. Moreover, we use Proposition 2.2.1(1) to reduce the question to show that the tensor product of two almost finitely presented *R*-modules is almost finitely presented.

Step 1. The case of finitely presented modules: If both M and N are finitely presented, then this is a standard fact proven in [Bou98, II, §3.6, Proposition 6].

Step 2. The case of M being finitely presented: Now we deal with the case of a finitely presented R-module M and merely almost finitely presented N. We fix a finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$ and a finitely generated ideal \mathfrak{m}_1 such that $\mathfrak{m}_0 \subset \mathfrak{m}_1^2$. Now we use Lemma 2.5.7(2) to find a finitely presented module N' and a morphism $f: N' \to N$ such that $\ker(f)$ and $\operatorname{Coker}(f)$ are annihilated by \mathfrak{m}_0 . We denote the image of f by N'' and consider the short exact sequences

$$\begin{array}{l} 0 \rightarrow K \rightarrow N' \rightarrow N'' \rightarrow 0 \ , \\ 0 \rightarrow N'' \rightarrow N \rightarrow Q \rightarrow 0 \end{array}$$

with K and Q being annihilated by \mathfrak{m}_0 . After applying the functor $M \otimes_R -$, we get the following exact sequences:

$$M \otimes_R K \to M \otimes_R N' \to M \otimes_R N'' \to 0 ,$$

$$\operatorname{Tor}_1^R(M, Q) \to M \otimes_R N'' \to M \otimes_R N \to M \otimes_R Q \to 0$$

We note that $M \otimes_R K$, $\operatorname{Tor}_1^R(M, Q)$, and $M \otimes_R Q$ are annihilated by \mathfrak{m}_0 . Now it is straightforward to conclude that the map

$$M \otimes_R f \colon M \otimes N' \to M \otimes N$$

has kernel and cokernel annihilated by $\mathfrak{m}_1 \subset \mathfrak{m}_0^2$. Moreover, $M \otimes N'$ is a finitely presented module by Step 1. Since \mathfrak{m}_1 was an arbitrary finitely generated subideal of \mathfrak{m} , we conclude that $M \otimes N$ is almost finitely presented by Lemma 2.5.7(2).

Step 3. The general case: Repeat the argument of Step 2 once again using Step 2 in place of Step 1 at the end, and Lemma 2.5.8 in place of Lemma 2.5.7(2). \Box

Lemma 2.5.14. Let M be an almost finitely presented R-module, let N be any R-module, and let P be an almost flat R-module. Then the natural map $\operatorname{Hom}_R(M, N) \otimes_R P \to \operatorname{Hom}_R(M, N \otimes_R P)$ is an almost isomorphism.

Similarly, $\operatorname{Hom}_{R^a}(M^a, N^a) \otimes_{R^a} P^a \to \operatorname{Hom}_{R^a}(M^a, N^a \otimes_{R^a} P^a)$ is an almost isomorphism for any almost finitely presented R^a -module M^a , any R^a -module N^a , and an almost flat R^a -module P^a .

Proof. Proposition 2.2.1(1) and (3) ensure that it suffices to prove the claim for the case of honest R-modules M, N, and P.

Step 1. The case of a finitely presented module M: We choose a presentation of M:

$$R^n \to R^m \to M \to 0$$

Then we use that P is almost flat to get a morphism of almost exact sequences:

Clearly, the second and third vertical arrows are (almost) isomorphisms, so the first vertical arrow is an almost isomorphism as well.

Step 2. The General Case: The case of almost finitely presented module M follows from the finitely presented case by approximating it by finitely presented ones. This is similar to the strategy used in Lemma 2.5.13, we leave the details to the reader.

The last thing that we will need is the interaction between properties of an R-module M and its "reduction" M/I for some finitely generated ideal $I \subset \mathfrak{m}$. For example, we know that for an ideal $I \subset \operatorname{rad}(R)$ and a finite module M, Nakayama's lemma states that M/I = 0 if and only if M = 0. Another thing is that an I-adically complete module M is R-finite if and only if M/I is R/I-finite. It turns out that both facts have their "almost" analogues.

Lemma 2.5.15. Let $I \subset \mathfrak{m} \cap \operatorname{rad}(R)$ be a finitely generated ideal. If M is an almost finitely generated R-module such that $M/IM \simeq 0$. Then $M \simeq 0$. If $M/IM \cong^a 0$, then $M \cong^a 0$.

Proof. We use a definition of an almost finitely generated module to find a finite submodule N that contains IM. If M/IM is isomorphic to the zero module, then the containment $IM \subset N \subset M$ implies that N = M. Thus M is actually finitely generated, now we use the usual Nakayama's Lemma to finish the proof.

If M/IM is merely almost isomorphic to the zero module, then we see that the inclusion $IM \subset M$ is an almost isomorphism. In particular, $\mathfrak{m}M$ is contained in IM. Using that $\mathfrak{m}^2 = \mathfrak{m}$, we obtain an *equality*

$$\mathfrak{m}M = \mathfrak{m}^2 M = \mathfrak{m}(IM) = I(\mathfrak{m}M)$$

Thus we can apply the argument from above to conclude that $\mathfrak{m}M = 0$. This finishes the proof as $\mathfrak{m}M \cong^a M$.

Lemma 2.5.16. Let R be I-adically complete for some finitely generated $I \subset \mathfrak{m}$. Then an I-adically complete R-module M is almost finitely generated if and only if M/IM is almost finitely generated.

Proof. [GR03, Lemma 5.3.18]

2.6. Almost Coherent Modules and Almost Coherent Rings. This section is devoted to the study of "almost coherent" modules. They are supposed to be "almost" analogues of usual coherent modules. We show that they always form a Weak Serre subcategory in \mathbf{Mod}_R . Then we study the special case of almost coherent modules over an almost coherent ring, and show that in this case almost coherent modules are the same as almost finitely presented modules. We recall that we fixed some "base" ring R with an ideal \mathfrak{m} such that $\mathfrak{m}^2 = \mathfrak{m}$ and $\tilde{\mathfrak{m}} = \mathfrak{m} \otimes_R \mathfrak{m}$ is flat, and always we do almost mathematics with respect to this ideal.

Definition 2.6.1. We say that an (almost) *R*-module *M* is almost coherent if it is almost finitely generated and every almost finitely generated almost submodule $N^a \subset M^a$ is almost finitely presented.

Remark 2.6.2. An almost submodule $N^a \subset M^a$ does not necessarily give rise to a submodule $N' \subset M$ for some $(N')^a \simeq N$. The most we can say is that there is an injection $f_! \colon (N^a)_! \to (M^a)_!$ whose almostification is equal to the the morphism f (this follows from Lemma 2.1.8(2)).

Lemma 2.6.3. Let M^a be an almost *R*-module with a representative $M \in \mathbf{Mod}_R$. Then the following are equivalent

- (1) The almost module M^a is almost coherent.
- (2) The actual *R*-module $(M^a)_*$ is almost finitely generated, and any almost finitely generated *R*-submodule of $(M^a)_*$ is almost finitely presented.
- (3) The actual *R*-module $(M^a)_!$ is almost finitely generated, and any almost finitely generated *R*-submodule of $(M^a)_!$ is almost finitely presented.

Proof. First of all, we note that Corollary 2.5.9 guarantees that M is almost finitely generated if and only if so is $(M^a)_*$. Secondly, Lemma 2.1.9 implies that the functor $(-)_*$ is left exact. Therefore, any almost submodule $N^a \subset M^a$ gives rise to an actual submodule $(N^a)_* \subset (M^a)_*$ that is almost isomorphic to N. In reverse, any submodule $N \subset (M^a)_*$ gives rise to an almost submodule of M^a . Hence, we see that all almost finitely generated almost submodules of M^a are almost finitely presented if and only if all actual almost finitely generated submodules of M_* are almost finitely presented (here we again use Corollary 2.5.9). This shows the equivalence of (1) and (2). The same argument shows that (1) is equivalent to (3).

Note that it is not that clear whether a coherent R-module is almost coherent. The issue is that in the definition of almost coherent modules we need to be able to handle all almost finitely generated almost submodules and not only finitely generated. The lemma below is a useful tool to deal with such problems, in particular, it turns out (Corollary 2.6.6) that all coherent modules are indeed almost coherent, but we do not know a direct way to see it.

Lemma 2.6.4. Let M be an R-module. Then M is an almost coherent module if one of the following holds:

(1) For any finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$ there exists a coherent *R*-module *N* and morphism $f: N \to M$ such that $\mathfrak{m}_0(\ker f) = 0$ and $\mathfrak{m}_0(\operatorname{Coker} f) = 0$.

(2) For any finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$ there exists an *almost* coherent *R*-module *N* and morphism $f: N \to M$ such that $\mathfrak{m}_0(\ker f) = 0$ and $\mathfrak{m}_0(\operatorname{Coker} f) = 0$.

Proof. We start the proof by noting that M comes with the natural almost isomorphism $M \to M_*^a$, so both of the assumptions on M pass through this almost isomorphism. Thus, Lemma 2.6.3 implies that it suffices to show that $M_* := M_*^a$ is almost coherent.

Lemma 2.5.7 guarantees that M_* is almost finitely generated. Thus we only need to check the second condition from Definition 2.6.1. So we pick an arbitrary almost finitely generated Rsubmodule $M_1 \subset M_*$, we want to show that it is almost finitely presented. We choose an arbitrary finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$ and another finitely generated ideal $\mathfrak{m}_1 \subset \mathfrak{m}$ such that $\mathfrak{m}_0 \subset \mathfrak{m}_1^2$.

We use Lemma 2.5.8 to find a morphism $\varphi : \mathbb{R}^n \to M_1$ such that $\mathfrak{m}_1(\operatorname{Coker} \varphi) = 0$. Let e_1, \ldots, e_n be the standard basis in \mathbb{R}^n , and define $x_i := \varphi(e_i)$ to be their images. We also choose some set of generators $(\varepsilon_1, \ldots, \varepsilon_m)$ for the ideal \mathfrak{m}_1 .

Now we recall that by our assumption there is a morphism $f: N \to M_*$ with a(n) (almost) coherent *R*-module *N* such that $\mathfrak{m}_1(\operatorname{Coker} f) = 0$ and $\mathfrak{m}_1(\ker f) = 0$. This implies that $\varepsilon_i x_j$ is in the image of *f* for any $i = 1, \ldots, m, j = 1, \cdots n$. Let us choose some $y_{i,j} \in N$ such that $f(y_{i,j}) = \varepsilon_i x_j$, and we define an *R*-module *N'* as the submodule of *N* generated by all $y_{i,j}$, this is a finite *R*-module by the construction. Since *N* is a (almost) coherent module, we conclude that N' is actually (almost) finitely presented.

We observe that $f' := f|_{N'}$ naturally lands in M_1 , and we have $\mathfrak{m}_1(\ker f') = 0$ and $\mathfrak{m}_1^2(\operatorname{Coker} f') = 0$. Since $\mathfrak{m}_0 \subset \mathfrak{m}_1^2$ this shows that the morphism

$$N' \xrightarrow{f'} M_1$$

has a kernel and cokernel killed by \mathfrak{m}_0 . Lemma 2.5.8 shows that M_1 is almost finitely presented. \Box

Remark 2.6.5. Does the converse of this Lemma hold?

Corollary 2.6.6. Any coherent R-module M is almost coherent.

The next thing that we want to show is that almost coherent modules from a Weak Serre subcategory of \mathbf{Mod}_R . This is an almost analogue of the corresponding statement in the classical case.

Lemma 2.6.7. Let R and \mathfrak{m} as above. Then

- (1) An almost finitely generated almost submodule of an almost coherent module is almost coherent.
- (2) Let $\varphi : N^a \to M^a$ be an almost homomorphism from an almost finitely generated R^a -module to an almost coherent R^a -module, then ker φ is almost finitely generated R^a -module.
- (3) Let $\varphi : N^a \to M^a$ be an injective almost homomorphism of almost coherent R^a -modules, then Coker φ is almost coherent R^a -module.
- (4) Let $\varphi : N^a \to M^a$ be an almost homomorphism of almost coherent R^a -modules, then ker φ and Coker φ are almost coherent R^a -modules.
- (5) Given a short exact sequence of R^a -modules $0 \to M'^a \to M^a \to M''^a \to 0$ if two out of three are almost coherent so is the third.

Proof. (1): This is evident from the definition of an almost coherent almost module.

(2): Let us define $N''^a := \operatorname{Im} \varphi$ and $N'^a := \ker \varphi$, then Corollary 2.5.12(1) implies that N''^a is an almost finitely generated almost submodule of M^a . It is actually almost finitely presented since

 M^a is almost coherent, we use Corollary 2.5.12(3) to get that N' is almost finitely generated as well.

(3): We denote Coker φ by $M^{\prime\prime a}$, then we have a short exact sequence

$$0 \to N^a \to M^a \to M''^a \to 0.$$

Corollary 2.5.12(1) implies that M''^a is almost finitely generated. Let us choose any almost finitely generated almost submodule $M''_1 \subset M''^a$ and denote its pre-image in M^a by M_1^a . Then we have a short exact sequence

$$0 \to N^a \to M_1^a \to M_1^{\prime\prime a} \to 0.$$

Corollary 2.5.12(2) guarantees that M_1^a is an almost finitely generated almost submodule of M^a . Since M^a is almost coherent, we see that M_1^a is an almost finitely presented R^a -module. Therefore, Corollary 2.5.12(4) implies that $M_1''^a$ is also almost finitely presented. Hence, the R^a -module M''^a is almost coherent.

(4): We know that $N'^a := \ker \varphi$ is almost finitely generated by (2). Since N^a is almost coherent, we conclude that N'^a is almost coherent by (1). We define $N''^a := \operatorname{Im} \varphi$ and $M''^a := \operatorname{Coker} \varphi$, then we note that we have two short exact sequences

$$\begin{split} 0 &\to N'^a \to N^a \to N''^a \to 0, \\ 0 &\to N''^a \to M^a \to M''^a \to 0. \end{split}$$

We observe that (3) shows that N''^a is almost coherent, then we use (3) once more to conclude that M''^a is also almost coherent.

(5): The only thing that we are left to show is that if M'^a and M''^a are almost coherent so is M^a . It is almost finitely generated by Corollary 2.5.12(2). Now to check the second condition from Definition 2.6.1, we choose an almost finitely generated almost submodule $M_1^a \subset M^a$. Let us denote by $M_1''^a$ its image in M''^a , and by $M_1'^a$ the kernel of this map. So we have a short exact sequence

$$0 \to M_1^{\prime a} \to M_1^a \to M_1^{\prime \prime a} \to 0.$$

Corollary 2.5.12 guarantees that $M_1''^a$ is an almost finitely generated almost submodule of the almost coherent R^a -module M''^a . Hence, (1) implies that $M_1''^a$ is almost coherent, in particular, it is almost finitely presented. Moreover, we can now use (2) to get that $M_1'^a$ is an almost finitely generated almost submodule of M'^a . Since M'^a is almost coherent, we conclude that $M_1'^a$ is actually almost finitely presented. Finally, Corollary 2.5.12(2) shows that M_1^a is almost finitely presented as well. This finishes the proof of almost coherence of the R^a -module M.

Corollary 2.6.8. Let M^a be an almost finitely presented R^a -modules and let N^a be an almost coherent R^a -module. Then $M^a \otimes_{R^a} N^a$ and alHom_{R^a} (M^a, N^a) are almost coherent.

Proof. We use Proposition 2.2.1(1),(3) to reduce the question to show that $M \otimes_R N$ and $\operatorname{Hom}_R(M, N)$ are almost coherent *R*-modules for any almost finitely presented *R*-module *M* and almost coherent *R*-module *N*.

Step 1. The case of finitely presented module M: In this case we pick a presentation of M as the quotient

$$R^n \to R^m \to M \to 0$$
.

Then we have short exact sequences

$$N^n \to N^m \to M \otimes_R N \to 0$$

36
and

$$0 \to \operatorname{Hom}_R(M, N) \to N^m \to N^n$$

We note that Lemma 2.6.7(5) implies that N^m and N^n are almost coherent. Thus Lemma 2.6.7(5) guarantees that both $M \otimes_R N$ and $\operatorname{Hom}_R(M, N)$ are almost coherent as well.

Step 2. The General Case: The argument is similar to the one used in Step 2 of the proof of Lemma 2.5.13. We approximate M with finitely presented R-modules. This gives us an approximations of $M^a \otimes_{R^a} N^a$ and alHom_{R^a} (M^a, N^a) by almost coherent modules. Now Lemma 2.6.4 guarantees that these modules are actually almost coherent. We leave details to the interested reader.

We define ACoh_R (resp. ACoh_R^a) to be the strictly full⁶ subcategoty of Mod_R (resp. Mod_R^a) consisting of almost coherent *R*-modules (resp. R^a -modules).

Corollary 2.6.9. The category ACoh_R (resp. ACoh_R^a) is a Weak Serre subcategory of Mod_R^a (resp. Mod_R^a).

Corollary 2.6.9 and the discussion in [Sta21, Tag 06UP] ensure that $\mathbf{D}_{acoh}(R)$ and $\mathbf{D}_{acoh}(R)^{a7}$ are strictly full saturated⁸ triangulated subcategories of $\mathbf{D}(R)$ and $\mathbf{D}(R)^a$ respectively. We define $\mathbf{D}^+_{acoh}(R) \coloneqq \mathbf{D}_{acoh}(R) \cap \mathbf{D}^+(R)$ and similarly for all other bounded versions.

The last part of this subsection is dedicated to the study of almost coherent rings and almost coherent modules over almost coherent rings. Recall that coherent modules over a coherent ring coincide with finitely presented ones. Similarly, we will show that almost coherent modules over an almost coherent ring turn out to be the same as almost finitely presented ones.

Definition 2.6.10. We say that a ring R is almost coherent if the rank-1 free module R is almost coherent as an R-module.

Lemma 2.6.11. A coherent ring R is almost coherent.

Proof. Apply Corrollary 2.6.6 to a rank-1 free module R.

Lemma 2.6.12. If R is an almost coherent ring, then any almost finitely presented R-module M is almost coherent.

Proof. Step 1: If M is finitely presented over R, then we can write it as a cokernel of a map between free finite rank modules. A free finite rank module over an almost coherent ring is almost coherent by Lemma 2.6.7(5). A cokernel of a map of almost coherent modules is almost coherent by Lemma 2.6.7(4). Therefore, any finitely presented M is almost coherent.

Step 2: Suppose that M is merely almost finitely presented. Lemma 2.5.7 guarantees that, for any finitely generated $\mathfrak{m}_0 \subset \mathfrak{m}$, we can find a finitely presented module N and a map $f: N \to M$ such that ker f and Coker f are annihilated by \mathfrak{m}_0 . N is almost coherent by Step 1. Therefore, Lemma 2.6.4(2) implies that M is almost coherent as well.

Corollary 2.6.13. Let R be an almost coherent ring. Then an R-module M is almost coherent if and only if it is almost finitely presented.

⁶i.e. full subcategory that is closed under isomorphisms.

⁷These are full subcategories of $\mathbf{D}(R)$ and $\mathbf{D}(R)^a$ of complexes with almost coherent cohomology modules, respectively.

⁸A strictly full subcategory \mathcal{D}' of a triangulated category \mathcal{D} is saturated if $X \oplus Y \in \mathcal{D}'$ implies $X, Y \in \mathcal{D}'$.

Proof. The "only if" part is clear from the definition, the "if" part follows from Lemma 2.6.12. \Box

Proposition 2.6.14. Let R be an almost coherent ring, and let M^a , N^a be two objects in $\mathbf{D}^-_{acoh}(R)^a$. Then $M^a \otimes_{R^a}^L N^a \in \mathbf{D}^-_{acoh}(R)^a$.

Proof. Step 1. Reduce to the case $M, N \in \mathbf{Mod}_R$: We firstly notice that the question is equivalent to showing that $M \otimes_R^L N$ is almost coherent for any $M, N \in \mathbf{D}^-_{acoh}(R)$. Indeed, we notice that $(-)_!: \mathbf{D}(R)^a \to \mathbf{D}(R)$ sends $\mathbf{D}^-_{acoh}(R)^a$ to $\mathbf{D}^-_{acoh}(R)$, and then use Proposition 2.4.13 to make the reduction.

Now we use that almost coherent modules are closed under extensions (Lemma 2.6.7(5)) and the convergent spectral sequence

$$\mathbf{E}_{2}^{p,q} = \mathbf{H}^{q} \left(\mathbf{H}^{p} \left(M \right) \otimes_{R}^{L} N \right) \Rightarrow \mathbf{H}^{p+q} \left(M \otimes_{R}^{L} N \right)$$

to conclude that it is sufficient to show that $\mathrm{H}^{q}(\mathrm{H}^{p}(M) \otimes_{R}^{L} N) \in \mathbf{ACoh}_{R}$ for any p, q, or equivalently $\mathrm{H}^{p}(M) \otimes_{R}^{L} N \in \mathbf{D}_{acoh}(R)$ for any integer p. This reduces to the case of $M \in \mathbf{Mod}_{R}$. Now we repeat the argument once again to reduce to the case $M, N \in \mathbf{Mod}_{R}$ (so really in \mathbf{ACoh}_{R}).

Step 2. We show that $M \otimes_R^L N \in \mathbf{D}_{acoh}^-(R)$ for $M, N \in \mathbf{Mod}_R$: The claim is equivalent to show that $\operatorname{Tor}_i^R(M, N) \in \mathbf{ACoh}_R$ for any $i \geq 0$. The case of i = 0 is done in Corollary 2.6.8. Now we use the dimension shift trick to give an argument for i > 0. As always, we distinguish the case of i = 1 and i > 1.

We firstly deal with the case i = 1. We fix any finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$. Now we use the definition of almost finitely generated modules to find map $f: \mathbb{R}^n \to M$ such that $Q \coloneqq \operatorname{Coker}(f)$ is annihilated by \mathfrak{m}_0 . We note that \mathbb{R}^n is almost coherent since \mathbb{R} was assumed to be an almost coherent ring. Thus Lemma 2.6.7(4) implies that $K \coloneqq \ker(f)$ is almost coherent as well. We denote $M' \coloneqq \operatorname{Im}(f)$ and consider the short exact sequences

$$0 \to K \to R^n \to M' \to 0 ,$$

$$0 \to M' \to M \to Q \to 0 .$$
(2.2)

The first one implies that we have an exact sequence

$$0 \to \operatorname{Tor}_1^R(M', N) \to K \otimes_R N \to N^n$$

Now we note that both $K \otimes_R N$ and N^n are almost coherent by Corollary 2.6.8 and Lemma 2.6.7(4), respectively. Therefore, Lemma 2.6.7(4) ensures that $\operatorname{Tor}_1^R(M', N)$ is also almost coherent as the kernel of a morphism of almost coherent modules.

Moreover, we see from (2.2) that the map $\operatorname{Tor}_1^R(M', N) \to \operatorname{Tor}_1^R(M, N)$ has kernel and cokernel killed by \mathfrak{m}_0 . So Lemma 2.6.4 implies that $\operatorname{Tor}_1^R(M, N)$ is almost coherent as well since \mathfrak{m}_0 was an arbitrary finitely generated subideal of \mathfrak{m} .

The case of i > 1 is similar. We suppose that the claim is proven for all almost coherent modules M' and the integer i - 1 > 0. We show that the statement also holds for the integer i. We repeat the same argument as in the case of i = 1 to find short exact sequences

$$0 \to K \to R^n \to M' \to 0 ,$$

$$0 \to M' \to M \to Q \to 0 .$$

with K being an almost coherent module, and Q annihilated by \mathfrak{m}_0 . Then the corresponding long exact sequences of Tor-groups imply that $\operatorname{Tor}_i^R(M', N) = \operatorname{Tor}_{i-1}^R(K, N)$ is almost coherent by the induction hypothesis, and the map

$$\operatorname{Tor}_{i}^{R}(M', N) \to \operatorname{Tor}_{i}^{R}(M, N)$$

has kernel and cokernel annihilated by \mathfrak{m}_0 . Thus Lemma 2.6.4 again implies that $\operatorname{Tor}_i^R(M, N)$ is almost coherent since \mathfrak{m}_0 was an arbitrary finitely generated subideal of \mathfrak{m} .

Proposition 2.6.15. Let R be an almost coherent ring, and let $M^a \in \mathbf{D}^-_{acoh}(R)^a, N^a \in \mathbf{D}^+_{acoh}(R)^a$. Then \mathbf{R} alHom_{R^a} $(M^a, N^a) \in \mathbf{D}^+_{acoh}(R)^a$.

Proof. The proof is similar to that of Proposition 2.6.14. We use Proposition 2.4.8 and the same argument with spectral sequences to reduce the claim to showing that $\operatorname{Ext}_{R}^{i}(M, N) \in \mathbf{D}_{acoh}^{-}(R)$ for any $M, N \in \mathbf{ACoh}_{R}$ and $i \geq 0$. The case of i = 0 is done in Corollary 2.6.8. Then the same trick with dimension shifting does the job for all other i.

Proposition 2.6.16. Let R be an almost coherent ring, let $M \in \mathbf{D}^-_{acoh}(R)$, $N \in \mathbf{D}^+(R)$, and let P be an almost flat R-module. Then the natural map $\mathbf{R}\operatorname{Hom}_R(M, N) \otimes_R P \to \mathbf{R}\operatorname{Hom}_R(M, N \otimes_R P)$ is an almost isomorphism.

Similarly, $\mathbf{R}\operatorname{Hom}_{R^a}(M^a, N^a) \otimes_{R^a}^L P^a \to \mathbf{R}\operatorname{Hom}_{R^a}(M^a, N^a \otimes_{R^a}^L P^a)$ is an almost isomorphism for any $M^a \in \mathbf{D}^-_{acoh}(R)^a$, $N^a \in \mathbf{D}^+(R)^a$, and let P^a a flat R^a -module.

Proof. The proof is similar to that of the lemmas above with the caveat that we need to work in the almost category to leverage almost flatness of the module P. Namely, we firstly do the spectral sequence argument in \mathbf{Mod}_{R}^{a} to reduce the question to showing that

$$\operatorname{Ext}_{R^a}^p(M^a, N^a) \otimes_{R^a} P^a \to \operatorname{Ext}_{R^a}^p(M^a, N^a \otimes_{R^a} P^a)$$

is an almost isomorphism for any $M \in \mathbf{ACoh}_R^a$, $N \in \mathbf{Mod}_R^a$, almost flat *R*-module *P*, and any integer *p*. The case p = 0 was done in Lemma 2.5.14. Then we argue by dimension shifting similar to what was done in Proposition 2.6.14.

Corollary 2.6.17. Let R be an almost coherent ring, let $M^a \in \mathbf{D}^-_{acoh}(R^a)$, $N \in \mathbf{D}^+(R^a)$, and let P^a be an almost flat R^a -module. Then the natural map

$$\mathbf{R}$$
al $\operatorname{Hom}_{R^a}(M^a, N^a) \otimes_{R^a} P^a \to \mathbf{R}$ al $\operatorname{Hom}_{R^a}(M^a, N^a \otimes_{R^a} P^a)$

is an isomorphism in $\mathbf{D}(\mathbb{R}^a)$.

2.7. Base Change for Almost Modules. The last topic that we want to discuss about almost modules over general rings is their behavior with respect to base change. Recall that given a ring homomorphism $\varphi: R \to S$ we always do almost mathematics on S-modules with respect to the ideal $\mathfrak{m}_S := \mathfrak{m}S$; look at Lemma 2.1.10 to see why $\widetilde{\mathfrak{m}_S}$ is flat.

Lemma 2.7.1. Let $\varphi : R \to S$ be a ring homomorphism, and let M^a be an almost finitely generated (resp. almost finitely presented) R^a -module. Then the module $M^a_S := M^a \otimes_{R^a} S^a$ is almost finitely generated (resp. almost finitely presented).

Proof. The claim follows from Lemma 2.5.7(2) and the fact that for any finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}_S$ there is a finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$ such that $\mathfrak{m}_0' \subset \mathfrak{m}_0 S$. We only give a complete proof in the case of finitely presented modules as the other case is an easier version of the same.

Firstly, we note that it suffices to show that $M \otimes_R S$ is almost finitely presented. Now the observation above implies that it suffices to check the condition of Lemma 2.5.7(2) only for ideals of the form $\mathfrak{m}_0 S$ for a finitely generated subideal $\mathfrak{m}_0 \subset \mathfrak{m}$. Then we choose some finitely generated ideal $\mathfrak{m}_1 \subset \mathfrak{m}$ such that $\mathfrak{m}_0 \subset \mathfrak{m}_1^2$ and we use Lemma 2.5.7(2) to find a finitely presented module N and a map $f: N \to M$ such that $\mathfrak{m}_1(\operatorname{Ker} f) = \mathfrak{m}_1(\operatorname{Coker} f) = 0$. Consider an exact sequence

$$0 \to K \to N \xrightarrow{J} M \to Q \to 0$$

and denote the image f by M'. Then we have the following exact sequences:

$$K \otimes_R S \to N \otimes_R S \to M' \otimes_R S \to 0$$

$$\operatorname{For}_1^R(Q, S) \to M' \otimes_R S \to M \otimes_R S \to Q \otimes_R$$

S

Since $K \otimes_R S$, $\operatorname{Tor}_1^R(Q, S)$ and $Q \otimes_R S$ are killed by $\mathfrak{m}_1 S$, we conclude that $\operatorname{Coker}(f \otimes_R S)$ and $\ker(f \otimes_R S)$ are annihilated by $\mathfrak{m}_1^2 S$. In particular, they are killed by $\mathfrak{m}_0 S$. Since $N \otimes_R S$ is finitely presented over S, Lemma 2.5.7 finishes the proof.

Corollary 2.7.2. Let $R \to S$ be a ring homomorphism with an almost coherent S, and let M^a be an object of $\mathbf{D}^-_{acoh}(R)^a$. Then $M^a \otimes_{R^a}^L S^a \in \mathbf{D}^-_{acoh}(S)^a$.

Proof. The proof is similar to that of Proposition 2.6.14. We use Proposition 2.4.16 and the same argument with spectral sequences to reduce the claim to showing that $\mathrm{H}^{i}(M \otimes_{R}^{L} S) \in \mathbf{D}_{acoh}^{-}(S)$ for any $M \in \mathbf{ACoh}_{R}$ and $i \geq 0$. The case of i = 0 follows from Lemma 2.7.1 as S is almost coherent. Then the same trick with dimension shifting does the job for all other i.

Lemma 2.7.3. Let S be a R-algebra that is finite (resp. finitely presented) as an R-module, and let M^a be an S^a -module. Then M^a is almost finitely generated (resp. almost finitely presented) over R^a if and only if it is almost finitely generated (resp. almost finitely presented) over S^a .

Proof. As always, we firstly reduce the question to the case of an honest S-module M. Now we use the observation that it suffices to check the condition of Lemma 2.5.7(2) only for the ideals of the form $\mathfrak{m}_0 S$ for some finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m} \subset R$. Then the only non-trivial direction is to show that M is almost finitely presented over S if it is almost finitely presented over R. This is proven in a more general situation in Lemma 2.7.4

Lemma 2.7.4. Let S be a possibly non-commutative R-algebra that is finite as a left (resp. right) R-module, and let M be a left (resp. right) S-module that is almost finitely presented over R. Then M is almost finitely presented over S (i.e. for every finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$, there exists a finitely presented left (resp. right) S-module N and a map $N \to M$ such that ker f and Coker f are annihilated by \mathfrak{m}_0).

Remark 2.7.5. This lemma will actually be used for a non-commutative ring S in the proof of Theorem 5.2.1 that, in turn, will be used in the proof of formal GAGA for almost coherent sheaves Theorem 5.3.2. Namely, we will apply to result to $S = \text{End}_{\mathbf{P}^N}(\mathcal{O} \oplus \mathcal{O}(1) \oplus \ldots \mathcal{O}(N))$.

Besides this application, Lemma 2.7.4 will be mostly used for almost coherent commutative rings R and S, where the proof can be significantly simplified.

Proof. We give a proof for left S-modules, the proof for right S-modules is the same. We start the proof by choosing some generators x_1, \ldots, x_n of S as an R-module. Then we pick a finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$ and another finitely generated ideal \mathfrak{m}_1 such that $\mathfrak{m}_0 \subset \mathfrak{m}_1^2$. And we also choose some generators $(\varepsilon_1, \ldots, \varepsilon_k) = \mathfrak{m}_1$ and find a three-term complex

$$R^t \xrightarrow{g} R^m \xrightarrow{f} M$$

such that $\mathfrak{m}_1(\operatorname{Coker} f) = 0$ and $\mathfrak{m}_1(\ker f) \subset \operatorname{Im} g$. We consider the images $y_i \coloneqq f(e_i) \in M$ of the standard basis elements in \mathbb{R}^m . Then we can find some $\beta_{i,j,s,r} \in \mathbb{R}$ such that

$$\varepsilon_s x_i y_j = \sum_{r=1}^m \beta_{i,j,s,r} \cdot y_r$$
 with $\beta_{i,j,s,r} \in R$

for any s = 1, ..., k; i = 1, ..., n; j = 1, ..., m. Moreover, we have t "relations"

$$\sum_{j=1}^{m} \alpha_{i,j} y_j = 0 \text{ with } \alpha_{i,j} \in R$$

such that for any relation $\sum_{i=1}^{m} b_i y_i = 0$ with $b_i \in R$ and any $\varepsilon \in \mathfrak{m}_1$, we have that the vector $\{\varepsilon b_i\}_{i=1}^{m} \in R^m$ lives in the *R*-subspace generated by vectors $\{\alpha_{i,j}\}_{i=1}^{m}$ for $j = 1, \ldots, t$. Or, in other words, if $\sum_{j=1}^{m} \alpha_{i,j} y_j = 0$ then $\varepsilon(\sum_{j=1}^{m} \alpha_{i,j} e_j) \in \operatorname{Im}(g)$ for any $\varepsilon \in \mathfrak{m}_1$.

Now we are finally ready to define a three-term complex

$$S^{nmk+t} \xrightarrow{\psi} S^m \xrightarrow{\varphi} M$$

We define the map φ as the unique S-linear homomorphism such that $\varphi(e_i) = y_i$ for the standard basis in S^m . We define ψ as the unique S-linear homomorphism such that

$$\psi(f_{i,j,s}) = \varepsilon_s x_i e_j - \sum_{r=1}^m \beta_{i,j,s,r} \cdot e_r \text{ and } \psi(f'_l) = \sum_{j=1}^m \alpha_{l,j} e_j$$

for the standard basis

$$\left\{f_{i,j,s}, f_l'\right\}_{i \le n, j \le m, s \le k, l \le t} \in S^{nmk+t}$$

Then we clearly have that $\varphi \circ \psi = 0$ and that $\mathfrak{m}_1(\operatorname{Coker} \varphi) = 0$. We claim that $\mathfrak{m}_1^2(\ker \varphi) \subset \operatorname{Im} \psi$.

Let $\varphi(\sum_{i=1}^{m} c_i e_i) = 0$ for some elements $c_i \in S$. We can write each

$$c_i = \sum_{j=1}^n r_{i,j} x_j \text{ with } r_{i,j} \in R$$
(2.3)

because x_1, \ldots, x_n are *R*-module generators of *S*. Thus, the condition that $\varphi(\sum_{i=1}^m c_i e_i) = 0$ is equivalent to $\sum_{i,j} r_{i,j} x_j y_i = 0$. Now recall that for any $s = 1, \ldots, k$ we have

$$\varepsilon_s x_j y_i = \sum_{r=1}^m \beta_{j,i,s,r} \cdot y_r.$$

Therefore, multiplying equation 2.3 by ε_s , we get an equality

$$0 = \varepsilon_s \left(\sum_{i,j} r_{i,j} x_j y_i \right) = \sum_{i,j} r_{i,j} \left(\sum_{r=1}^m \beta_{j,i,s,r} \cdot y_r \right) = \sum_{r=1}^m \left(\sum_{i,j} r_{i,j} \beta_{j,i,s,r} \right) y_r$$

This means that for any s' = 1, ..., k the vector $\{\varepsilon_{s'}(\sum_{i,j} r_{i,j}\beta_{j,i,s,r})\}_{r=1}^m \in \mathbb{R}^m$ lives in an R-subspace generated by vectors $\{\alpha_{i,j}\}_{i=1}^m$. In particular, for any r and s', $\varepsilon_{s'}(\sum_{i,j} r_{i,j}\beta_{j,i,s,r}e_r)$ is equal to ψ (some sum of f'_l) by definition of ψ .

After unwinding all the definitions we get the following:

$$\begin{split} \varepsilon_{s'}\varepsilon_s\left(\sum_{i=1}^m c_i e_i\right) &= \varepsilon_{s'}\varepsilon_s\left(\sum_{i,j} r_{i,j} x_j e_i\right) \\ &= \varepsilon_{s'}\left(\sum_{i,j} r_{i,j}\left(\varepsilon_s x_j e_i - \sum_r \beta_{j,i,s,r} e_r + \sum_r \beta_{j,i,s,r} e_r\right)\right) \\ &= \varepsilon_{s'}\left(\sum_{i,j} r_{i,j}\left(\varepsilon_s x_j e_i - \sum_r \beta_{j,i,s,r} e_r\right)\right) + \varepsilon_{s'}\left(\sum_r \left(\sum_{i,j} r_{i,j} \beta_{j,i,s,r}\right) e_r\right) \\ &= \psi\left(\varepsilon_{s'}\sum_{i,j} r_{i,j} f_{j,i,s}\right) + \psi \text{ (some sum of } f_l') \end{split}$$

So we see that $\mathfrak{m}_1^2 \ker(\varphi) \subset \operatorname{Im} \psi$. In particular, we have $\mathfrak{m}_0 \ker(\varphi) \subset \operatorname{Im} \psi$. Now we replace the map $\varphi \colon S^n \to M$ with the induced map

 $\overline{\varphi} \colon \operatorname{Coker}(\psi) \to M$

to get a map from a finitely presented left S-module such that $\ker(\overline{\varphi})$ and $\operatorname{Coker}(\overline{\varphi})$ are annihilated by \mathfrak{m}_0 .

Lemma 2.7.6. Let $\varphi : R \to S$ be a faithfully flat as above, and let M^a be an R^a -module. Suppose that $M^a \otimes_{R^a} S^a$ is almost finitely generated (resp. almost finitely presented) S^a -module. Then so is M^a .

Proof. As always, we firstly reduce the questions to the case of an honest S-module M, i.e. we show that an R-module M is almost finitely generated (resp. almost finitely presented) if so is the S-module $M \otimes_R S$.

We start with the almost finitely generated case. So we suppose that $M \otimes_R S$ is almost finitely generated, thus given any $\varepsilon \in \mathfrak{m}$ we can choose a morphism $g: S^n \to M \otimes_R S$ such that $\varepsilon(\operatorname{Coker} g) = 0$. Let us consider the standard basis e_1, \ldots, e_n of S^n , and we write

$$g(e_i) = \sum_j m_{i,j} \otimes s_{i,j}$$
 with $m_{i,j} \in M, s_{i,j} \in S$.

We define an *R*-module *F* to be a finite free *R*-module with a basis $e_{i,j}$. Then we define morphism

$$h: F \to M$$

as a unique *R*-linear homomorphism with $h(e_{i,j}) = m_{i,j}$. It is easy to see that $\varepsilon(\operatorname{Coker}(h \otimes_R S)) = 0$. Since *f* is faithfully flat, this implies that $\varepsilon(\operatorname{Coker} h) = 0$. We conclude that *M* is almost finitely generated as ε was an arbitrary element of \mathfrak{m} .

Now we deal with the almost finitely presented case. We pick some finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$, and another finitely generated ideal $\mathfrak{m}_1 \subset \mathfrak{m}$ such that $\mathfrak{m}_0 \subset \mathfrak{m}_1\mathfrak{m}$. We try to find a three-term complex

$$R^m \xrightarrow{g} R^n \xrightarrow{f} M$$

such that $\mathfrak{m}_0(\operatorname{Coker} f) = 0$ and $\mathfrak{m}_0(\ker f) \subset \operatorname{Im} g$.

The settled almost finitely generated case implies that M is at least almost finitely generated. In particular, we have some morphism

$$R^n \xrightarrow{f} M$$

such that $\mathfrak{m}_1(\operatorname{Coker} f) = 0$, thus $\mathfrak{m}_1(\operatorname{Coker}(f \otimes_R S)) = 0$ as well. Therefore, we can apply Lemma 2.5.6 to find a homomorphism $g' \colon S^m \to S^n$ such that $\mathfrak{m}_0(\ker(f \otimes_R S)) \subset \operatorname{Im}(g')$ and $(f \otimes_R S) \circ g' = 0$. This implies that g' actually lands inside $\ker(f \otimes_R S) = \ker(f) \otimes_R S$ by *R*-flatness of *S*.

Now we do the same trick as above: we write

$$g(e_i) = \sum_j m_{i,j} \otimes s_{i,j}$$
 with $m_{i,j} \in \ker(f), s_{i,j} \in S$.

We define an *R*-module *F* to be a finite free *R*-module with a basis $e_{i,j}$. Then we define a morphism

$$g \colon F \to \ker(f)$$

as the unique *R*-linear morphism such that $g(e_{i,j}) = m_{i,j}$. Then we see that $\mathfrak{m}_0(\ker(f \otimes_R S)) \subset \operatorname{Im}(g \otimes_R S)$. Since *S* is faithfully flat we conclude that $\mathfrak{m}_0(\ker f) \subset \operatorname{Im}(g)$ as well. This shows that a three-term complex

$$F \xrightarrow{g} R^n \xrightarrow{f} M$$

does the job. Therefore, M is an almost finitely presented R-module.

Corollary 2.7.7. Let $\varphi \colon R \to S$ be a faithfully flat as above, and let M^a be an R^a -module. Suppose that $M^a \otimes_{R^a} S^a$ is almost coherent S^a -module. Then so is M^a .

Proof. This follows directly from Lemma 2.7.6

We also show that the Hom functor almost commutes with flat base change under some finiteness assumptions.

Lemma 2.7.8. Let $R \to S$ be a flat map, M an almost finitely presented R-module, and N an R-module. Then the natural map

$$\operatorname{Hom}_R(M, N) \otimes_R S \to \operatorname{Hom}_S(M \otimes_R S, N \otimes_R S)$$

is an almost isomorphism.

Proof. This follows from the classical \otimes -Hom adjunction and Lemma 2.5.14.

Lemma 2.7.9. Let R be an almost coherent ring, $R \to S$ be a flat map, and $M \in \mathbf{D}^-_{acoh}(R)$, $N \in \mathbf{D}^+(R)$. Then the natural map

$$\mathbf{R}\operatorname{Hom}_R(M,N)\otimes^L_R S \to \mathbf{R}\operatorname{Hom}_S(M\otimes^L_R S,N\otimes^L_R S)$$

is an almost isomorphism.

Proof. We recall that we always have a canonical isomorphism $\mathbf{R}\operatorname{Hom}_R(K, L) \simeq \mathbf{R}\operatorname{Hom}_S(K \otimes_R^L S, L)$ for any $K \in \mathbf{D}^-(R)$ and any $L \in \mathbf{D}^+(S)$. This implies that it suffices to show that the natural map

$$\mathbf{R}\operatorname{Hom}_R(M,N)\otimes_R^L S \to \mathbf{R}\operatorname{Hom}_R(M,N\otimes_R^L S)$$

is an almost isomorphism. This follows from Proposition 2.6.16.

2.8. Almost Finitely Generated Modules over Adhesive Rings. This section discusses some basic aspects of almost finitely generated modules over adhesive rings. The motivation for this discussion will be the notion of almost coherent sheaves on formal schemes that we develop in Section 4.4. The results of this Section would be crucial in verifying certain good properties of adically quasi-coherent, almost coherent sheaves on "good" formal schemes. One of the main ingredients that we would need is the "Weak" version of the Artin-Rees Lemma (Lemma 2.8.6) and Lemma 2.8.7. Recall that these properties are already known for finite modules over the so-called "adhesive" rings. This is explained in a beautiful paper [FGK11]. The main goal of this section is to extend these result to the case of almost finitely generated modules.

That being said, let us introduce the Setup for this section. We start with the definition of an adhesive ring:

Definition 2.8.1. [FGK11, Definition 7.1.1] An adically topologized ring R endowed with the adic topology defined by a finitely generated ideal $I \subset R$ is said to be *(I-adically) adhesive* if it is Noetherian outside I^9 and satisfies the following condition: for any finitely generated R-module M, its I^{∞} -torsion part $M[I^{\infty}]$ is finitely generated.

Remark 2.8.2. Following the convention of [FGK11] we do not require a ring R with adic topology to be either *I*-adically complete or separated.

Set-up 2.8.3. We fix an *I*-adically adhesive ring *R* with an ideal \mathfrak{m} such that $I \subset \mathfrak{m}, \mathfrak{m}^2 = \mathfrak{m}$ and $\widetilde{\mathfrak{m}} := \mathfrak{m} \otimes_R \mathfrak{m}$ is flat. We always do almost mathematics with respect to the ideal \mathfrak{m} .

The main example of an adhesive ring is a (topologically) finitely presented algebra over a complete microbial valuation ring. This follows from [FGK11, Proposition 7.2.2] and [FGK11, Theorem 7.3.2]. For example, any topologically finitely presented algebra over a complete rank-1 valuation ring is adhesive.

Lemma 2.8.4. Let R be as in the Setup 2.8.3, and let M be an I-torsionfree almost finitely generated module. Then M is almost finitely presented. Similarly, any saturated submodule¹⁰ of an almost finitely generated R-module is almost finitely generated.

Proof. As M is almost finitely generated, we can find a finitely generated submodule $N \subset M$ that contains $\mathfrak{m}_0 M$ for a choice of a finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$. Since N is a submodule of M, it is itself *I*-torsionfree. Then [FGK11, Proposition 7.1.2] shows that N is finitely presented. Then Lemma 2.5.7(2) implies that M is almost finitely presented.

Now let M be an almost finitely generated R-module, and let $M' \subset M$ be a saturated submodule. Then M/M' is almost finitely generated by Lemma 2.5.11(1) and it is *I*-torsionfree. Therefore, it is almost finitely presented by the argument above. Then Lemma 2.5.11(3) guarantees that M' is almost finitely generated.

Lemma 2.8.5. Let R be as in the Setup 2.8.3, and let M be an almost finitely generated R-module. Then the I^{∞} -torsion module $M[I^{\infty}]$ is bounded (i.e. there is an integer n such that $M[I^n] = M[I^{\infty}]$).

Proof. Since M is almost finitely generated and the ideal $I \subset \mathfrak{m}$ is finitely generated, we conclude that there exists a finitely generated submodule $N \subset M$ such that $IM \subset N$. Then $I(M[I^{\infty}]) \subset$ $N[I^{\infty}]$, and $N[I^{\infty}]$ is finitely generated by adhesiveness of the ring R. In particular, there is an integer n such that $N[I^{\infty}]$ is annihilated by I^n . This implies that any element of $M[I^{\infty}]$ is annihilated by n+1.

⁹By definition, this means that the scheme Spec $A \setminus V(I)$ is noetherian.

¹⁰A submodule $N \subset M$ is saturated if $M/N[I^{\infty}] = 0$.

Lemma 2.8.6. Let R be as in the Setup 2.8.3, and let M be an almost finitely generated R-module. Suppose that $N \subset M$ is a submodule of M. For any integer n, there is an integer m such that $N \cap I^m M \subset I^n N$. In particular, the induced topology on the module N coincides with the I-adic one.

Proof. If M is finitely generated, then this is [FGK11, Theorem 4.2.2]. In general we use the definition of almost finitely generated module to find a submodule $M' \subset M$ such that M' is finitely generated and $IM \subset M'$. We define $N' := N \cap M'$ as the intersection of those modules. Then the established "weak" form of the Artin-Rees Lemma for finitely generated R-modules provides us with an integer m such that $N' \cap I^m M' \subset I^n N'$. In particular, we have

$$I^{m+1}M \cap N' \subset I^m M' \cap N' \subset I^n N' \subset I^n N.$$

Then we conclude that

$$I^{m+2}M \cap N \subset I^{m+1}M \cap M' \cap N \subset I^{m+1}M \cap N' \subset I^nN.$$

Since n was arbitrary, we conclude the claim.

Lemma 2.8.7. Let R be as in the Setup 2.8.3, and let M be an almost finitely generated R-module. Then the natural morphism $M \otimes_R \widehat{R} \to \widehat{M}$ is an isomorphism. In particular, any almost finitely generated module over a complete adhesive ring is complete.

Proof. We know that the claim holds for finitely generated modules by [FGK11, Proposition 4.3.4]. Now we deal with the almost finitely generated case. We choose a finitely generated submodule $N \subset M$ such that $IM \subset N$. Lemma 2.8.6 implies that the induced topology on N coincides with the *I*-adic topology on N. Thus the short exact sequence

$$0 \to N \to M \to M/N \to 0$$

remains exact after completion. Since $R \to \hat{R}$ is flat by [FGK11, Proposition 4.3.4], we conclude that we have a morphism of short exact sequences

Note that φ_N is an isomorphism as N is finitely generated, and $\varphi_{M/N}$ is isomorphism since it is an *I*-torsion module so $M/N \simeq (M/N) \otimes_R \widehat{R} \simeq \widehat{M/N}$. The five-lemma implies that φ_M is an isomorphism as well.

Corollary 2.8.8. Let R be as in the Setup 2.8.3, and let $M \in \mathbf{D}_{acoh}(R)$. Then M is *I*-adically derived complete¹¹.

Proof. First of all, we note that [Sta21, Tag 091P] implies that M is derived complete if and only if so are $H^i(M)$ for any integer i. So it suffices to show that any almost coherent R-module is derived complete. Lemma 2.8.7 gives that any such module is classically complete, and [Sta21, Tag 091T] ensures that any clasically complete module is derived complete.

 $^{^{11}}$ Look at [Sta21, Tag 091N] for the definition of derived completeness

3. Almost Mathematics on Ringed Sites

The main goal of this Chapter is to "globalize" results from Chapter 2. The two main cases of interest are almost coherent sheaves on schemes and "good" formal schemes. In order to treat those case somewhat uniformly we define some notions in the most general set-up of locally ringed spaces and check their basic properties. This is the content of Section 3.1. Sections 4.1 are 4.4 are devoted to the setting up foundations of almost coherent sheaves on schemes and formal schemes, respectively. In particular, we show that the notion of almost finitely generated (resp. presented, resp. coherent) module globalizes well on schemes and some "good" formal schemes. We prove the Proper Mapping Theorems in Section 5.1 both in the algebraic and formal Setups. Finally, we show the formal GAGA Theorem for adically quasi-coherent, almost coherent sheaves in Section 5.3. This is perhaps the most surprising result in this chapters as almost coherent sheaves are usually not finite type sheaves, so the "classical" proofs of Formal GAGA Theorem cannot really work in that situation.

3.1. The Category of \mathcal{O}_X^a -modules. We start this section by fixing a ring R with an ideal \mathfrak{m} such that $\mathfrak{m} = \mathfrak{m}^2$ and $\tilde{\mathfrak{m}} = \mathfrak{m} \otimes_R \mathfrak{m}$ is R-flat. We always do almost mathematics with respect to this ideal. The main goal of this section is to globalize the notion of almost mathematics on ringed R-sites.

The main object of our study in this Section will be a ringed site (X, \mathcal{O}_X) with \mathcal{O}_X being a sheaf of *R*-algebras. We call such sites as *ringed R-sites*. On each open *U*, it makes sense to speak about almost mathematics on $\mathcal{O}_X(U)$ -modules with respect to the ideal $\mathfrak{m}\mathcal{O}_X(U)^{12}$. Usually definitions of many notions in almost mathematics involve tensoring against the module $\tilde{\mathfrak{m}}$. We globalize this procedure in the following definition:

Definition 3.1.1. Let (X, \mathcal{O}_X) be a ringed *R*-site, and let \mathcal{F} be any \mathcal{O}_X -module. Then we define the sheaf $\widetilde{\mathfrak{m}} \otimes \mathcal{F}$ as the sheafification of the the presheaf that is defined as

$$U \mapsto \widetilde{\mathfrak{m}} \otimes_R \mathfrak{F}(U)$$

Remark 3.1.2. We note that this definition coincides with the tensor product $\underline{\widetilde{\mathfrak{m}}} \otimes_R \mathcal{F}$, where $\underline{\widetilde{\mathfrak{m}}}$ is the constant sheaf associated with the *R*-module \mathfrak{m} . Using flatness of the *R*-module $\widetilde{\mathfrak{m}}$, it is easy to see that the functor $-\otimes \widetilde{\mathfrak{m}}$ is exact and descends to a functor on the derived categories:

$$-\otimes \widetilde{\mathfrak{m}} \colon \mathbf{D}(X) \to \mathbf{D}(X)$$

where we denote by $\mathbf{D}(X)$ the derived category of \mathcal{O}_X -modules. Another way to think about it is to introduce the sheaf $\underline{\widetilde{\mathfrak{m}}}_X := \underline{\widetilde{\mathfrak{m}}} \otimes_R \mathcal{O}_X$. Then one easily see that there is a functorial isomorphism $\overline{\widetilde{\mathfrak{m}}} \otimes \mathcal{F} \simeq \underline{\widetilde{\mathfrak{m}}}_X \otimes_{\mathcal{O}_X} \mathcal{F}$ for any \mathcal{O}_X -module \mathcal{F} .

Definition 3.1.3. We say that an \mathcal{O}_X -module \mathcal{F} is *almost zero* if $\widetilde{\mathfrak{m}} \otimes \mathcal{F}$ is zero. We denote the category of almost zero \mathcal{O}_X -modules by Σ_X .

Remark 3.1.4. Since $\widetilde{\mathfrak{m}}$ is an *R*-flat module, we easily see that the category of almost zero \mathcal{O}_X -modules form a Serre subcategory of $\mathbf{Mod}_{\mathcal{O}_X} = \mathbf{Mod}_X$.

Lemma 3.1.5. Let (X, \mathcal{O}_X) be a ringed *R*-site, and let \mathcal{F} be an \mathcal{O}_X -module. Suppose that \mathcal{U} is a base of topology on *X*. Then the following conditions are equivalent:

- (1) $\mathcal{F} \otimes \widetilde{\mathfrak{m}}$ is the zero sheaf.
- (2) For any $\varepsilon \in \mathfrak{m}$, $\varepsilon \mathcal{F} = 0$.

 $^{^{12}}$ look at Lemma 2.1.10 for the reason why this makes sense.

- (3) For any $U \in \mathcal{U}$, the module $\widetilde{\mathfrak{m}} \otimes \mathfrak{F}(U)$ is zero.
- (4) For any $U \in \mathcal{U}$, the module $\mathfrak{m} \otimes \mathfrak{F}(U)$ is zero.
- (5) For any $U \in \mathcal{U}$, the module $\mathfrak{m}(\mathcal{F}(U))$ is zero.

Proof. We firstly show that (1) implies (2). We pick an element $\varepsilon \in \mathfrak{m} = \mathfrak{m}^2$ and write it as $\varepsilon = \sum x_i \cdot y_i$ for some $x_i, y_i \in \mathfrak{m}$. So the multiplication by ε map can be decomposed as

$$\mathcal{F} \xrightarrow{s \mapsto s \otimes \sum x_i \otimes y_i} \mathcal{F} \otimes \widetilde{\mathfrak{m}} \xrightarrow{m} \mathcal{F}$$

where the last map is induced by the multiplication by $\widetilde{\mathfrak{m}} \to R$. Then if $\mathcal{F} \otimes \widetilde{\mathfrak{m}} = 0$, then the multiplication by ε map is zero for any $\varepsilon \in \mathfrak{m}$. Now (2) easily implies (5). Lemma 2.1.1 ensures that (3), (4), and (5) are equivalent. Finally, (3) clearly implies (1).

Lemma 3.1.6. Let (X, \mathcal{O}_X) be a ringed *R*-site, and let \mathcal{F} be an almost zero \mathcal{O}_X -module. Then $\mathrm{H}^i(U, \mathcal{F}) \cong^a 0$ for any open $U \in X^{13}$ and any $i \geq 0$.

Proof. If \mathcal{F} is almost zero, then $\varepsilon \mathcal{F} = 0$ for any $\varepsilon \in \mathfrak{m}$ by Lemma 3.1.5. Since the functors $\mathrm{H}^{i}(X, -)$ are *R*-linear, we conclude that $\varepsilon \mathrm{H}^{i}(U, \mathcal{F}) = 0$ for any open *U* and any $\varepsilon \in \mathfrak{m}, i \geq 0$. Thus Lemma 2.1.1 ensures that $\mathrm{H}^{i}(U, \mathcal{F}) \cong^{a} 0$.

Definition 3.1.7. We say that a homomorphism $\varphi \colon \mathcal{F} \to \mathcal{G}$ of \mathcal{O}_X -modules is an *almost isomorphism* if ker(φ) and Coker(φ) are almost zero.

Lemma 3.1.8. A homomorphism $\varphi \colon \mathcal{F} \to \mathcal{G}$ of \mathcal{O}_X -modules is an almost isomorphism if and only if $\varphi(U) \colon \mathcal{F}(U) \to \mathcal{G}(U)$ is an almost isomorphism of $\mathcal{O}_X(U)$ -modules for any open $U \in X$.

Proof. The \Leftarrow implication is clear from the definitions. We give a proof of the \Rightarrow implication.

Suppose that φ is an almost isomorphism. We define the auxiliary \mathcal{O}_X -modules: $\mathcal{K} := \ker(\varphi), \mathcal{F}' := \operatorname{Im}(\varphi), \mathcal{Q} := \operatorname{Coker}(\varphi)$. Lemma 3.1.6 implies that the maps

$$\mathfrak{F}(U) \to \mathfrak{F}'(U) \text{ and } \mathfrak{F}'(U) \to \mathfrak{G}(U)$$

are almost isomorphisms. In particular, the composition $\mathfrak{F}(U) \to \mathfrak{G}(U)$ must also be an almost isomorphism.

Now we discuss the notion of almost \mathcal{O}_X -modules on a ringed *R*-site (X, \mathcal{O}_X) . This notion can be defined in two different ways: either as the quotient of the category of \mathcal{O}_X -modules by the Serre subcategory of almost zero modules or as modules over the almost structure sheaf \mathcal{O}_X^a . We need to explain these two notions in more detail now.

Definition 3.1.9. We define the category of almost O_X -modules as the quotient category

$$\mathbf{Mod}^a_{\mathcal{O}_X} \coloneqq \mathbf{Mod}_{\mathcal{O}_X} / \Sigma_X$$
.

Now we want to define the category $\mathbf{Mod}_{\mathcal{O}_X^a}$ of \mathcal{O}_X^a -modules that we will show to be equivalent to $\mathbf{Mod}_{\mathcal{O}_X}^a$. We recall that the almostification functor $(-)^a$ is exact on the level of modules and commutes with arbitrary products. This allows us to define the almost structure sheaf:

Definition 3.1.10. The almost structure sheaf \mathcal{O}_X^a is the sheaf¹⁴ of R^a -modules \mathcal{O}_X^a : $(\mathrm{Ob}(X))^{op} \to \mathrm{Mod}_R^a$ defined as $U \mapsto \mathcal{O}_X(U)^a$.

¹³An open $U \in X$ is by definition an object $U \in Ob(X)$ of the category underlying the site X.

¹⁴It is a sheaf exactly because $(-)^a$ is exact and commutes with arbitrary products.

Definition 3.1.11. We define the category of \mathcal{O}_X^a -modules $\operatorname{Mod}_{\mathcal{O}_X^a}$ as the category of the modules over $\mathcal{O}_X^a \in \operatorname{Shv}(X, R^a)$ in the categorical sense. More precisely, the objects are sheaves of R^a modules \mathcal{F} with a map $\mathcal{F} \otimes_{R^a} \mathcal{O}_X^a \to \mathcal{F}$ over R^a satisfying the usual axioms for a module. Morphisms are defined in the evident way.

We now define the functor

 $(-)^a \colon \mathbf{Mod}_{\mathcal{O}_X} \to \mathbf{Mod}_{\mathcal{O}_X^a}$

that sends a sheaf to its "almostification", i.e. it applies the functor $(-)^a \colon \mathbf{Mod}_R \to \mathbf{Mod}_R^a$ section-wise. Since the almostification functor $(-)^a$ is exact and commutes with arbitrary product, it is evident that \mathcal{F}^a is actually a sheaf for any \mathcal{O}_X -module F. Moreover, it is clear that $\mathcal{F}^a \simeq 0$ for any almost zero \mathcal{O}_X -module \mathcal{F} . Thus, it induces the functor

$$(-)^a \colon \mathbf{Mod}^a_{\mathcal{O}_X} \to \mathbf{Mod}_{\mathcal{O}^a_X}$$
.

The claim is that this functor induces the equivalence of categories. The first step towards the proof is to construct the right adjoint to $(-)^a \colon \mathbf{Mod}_{\mathcal{O}_X} \to \mathbf{Mod}_{\mathcal{O}_X^a}$. Our construction of the right adjoint functor will use the existence of the left adjoint functor. So we slightly postpone the proof of the mentioned above equivalence and discuss adjoints to $(-)^a$.

We start with the definition of the left adjoint functor. The idea is to apply the functor $(-)_!$: $\mathbf{Mod}_R^a \to \mathbf{Mod}_R^a$ section-wise, though this strategy does not quite work as $(-)_!$ does not commute with infinite products.

Definition 3.1.12. • We define the functor $(-)_1^p \colon \operatorname{Mod}_{\mathcal{O}_X}^{o_a} \to \operatorname{Mod}_{\mathcal{O}_X}^{p-15}$ as $\mathcal{F} \mapsto (U \mapsto \mathcal{F}(U)_1)$

• We define the functor $(-)_!$: $\mathbf{Mod}_{\mathcal{O}_X^a} \to \mathbf{Mod}_{\mathcal{O}_X}$ as the composition $(-)_! \coloneqq (-)^{\#} \circ (-)_!^p$, where $(-)^{\#}$ is the sheafification functor.

Lemma 3.1.13. Let (X, \mathcal{O}_X) be a ringed *R*-site. Then

(1) The functor

$$(-)_! \colon \mathbf{Mod}_{\mathcal{O}_X^a} \to \mathbf{Mod}_{\mathcal{O}_X}$$

is the left adjoint to the localization functor $(-)^a \colon \mathbf{Mod}_{\mathcal{O}_X} \to \mathbf{Mod}_{\mathcal{O}_X^a}$. In particular, we have a functorial isomorphism

$$\operatorname{Hom}_{\mathcal{O}_X^a}(\mathcal{F}, \mathcal{G}^a) \simeq \operatorname{Hom}_{\mathcal{O}_X}(\mathcal{F}_!, \mathcal{G})$$

for any $\mathcal{F} \in \mathbf{Mod}_{\mathcal{O}_X^a}, \mathcal{G} \in \mathbf{Mod}_{\mathcal{O}_X}$.

- (2) The functor $(-)_{!}: \operatorname{Mod}_{\mathcal{O}_{X}} \to \operatorname{Mod}_{\mathcal{O}_{X}}$ is exact.
- (3) The counit morphism $(\mathcal{F}^a)_! \to \mathcal{F}$ is an almost isomorphism for any $\mathcal{F} \in \mathbf{Mod}_{\mathcal{O}_X}$. The unit morphism $\mathcal{G} \to (\mathcal{G}_!)^a$ is an isomorphism for any $\mathcal{G} \in \mathbf{Mod}_{\mathcal{O}_X^a}$. In particular, the functor $(-)^a$ is essentially surjective.

Proof. (1) follows from Lemma 2.1.9(3) and the adjunction between sheafication and the forgetful functor. More precisely, we have the following functorial isomorphisms

$$\operatorname{Hom}_{\mathcal{O}_X^a}(\mathcal{F}, \mathcal{G}^a) \simeq \operatorname{Hom}_{\operatorname{\mathbf{Mod}}_{\mathcal{O}_X}^p}(\mathcal{F}_!^p, \mathcal{G}) \simeq \operatorname{Hom}_{\mathcal{O}_X}(\mathcal{F}_!, \mathcal{G})$$

We show (2). It is easy to see that $(-)_{!}$ is left exact from Lemma 2.1.9(4) and the exactness of the sheafification functor. It is also right exact since it is a left adjoint functor to $(-)^{a}$.

¹⁵ $\mathbf{Mod}_{\mathcal{O}_X}^p$ stands for the category of modules over \mathcal{O}_X in the category of presheaves

Now we show (3). Lemma 2.1.9(5) ensures that the kernel and cokernel of the counit map of presheaves $(\mathcal{F}^a)_!^p \to \mathcal{F}$ are annihilated by any $\varepsilon \in \mathfrak{m}$. Then the same holds after sheafification, proving the $(\mathcal{F}^a)_!^p \to \mathcal{F}$ is an almost isomorphism by Lemma 3.1.5.

We consider the unit map $\mathcal{G} \to (\mathcal{G}_!)^a$, we note that using the adjuction $((-)_!, (-)^a)$ section-wise, we can refine this map

$$\mathfrak{G} \to (\mathfrak{G}_!^p)^a \to (\mathfrak{G}_!)^a$$
.

It suffices to show that both maps are isomorphisms, the first map is an isomorphism by Lemma 2.1.9(5). In particular, this implies that $(\mathcal{G}_{!}^{p})^{a}$ is already a sheaf of almost \mathbb{R}^{a} -modules, but then we see that the natural map $(\mathcal{G}_{!}^{p})^{a} \to (\mathcal{G}_{!})^{a}$ must also be an isomorphism as it coincides with the sheafification in the category of presheaves of \mathbb{R}^{a} -modules.

Remark 3.1.14. In what follows, we denote the objects of $\operatorname{Mod}_{\mathcal{O}_X^a}$ by \mathcal{F}^a to distinguish \mathcal{O}_X and \mathcal{O}_X^a -modules. This notation does not cause any confusion as $(-)^a$ is indeed essentially surjective.

Now we construct the right adjoint functor to $(-)^a$. The naive idea of applying $(-)_*$ section-wise works well in this case. The only thing we emphasize here is that essential surjectivity of $(-)^a$ is used in our definition of $(-)_*$.

Definition 3.1.15. The functor of almost sections $(-)_* \colon \operatorname{Mod}_{\mathcal{O}_X} \to \operatorname{Mod}_{\mathcal{O}_X}$ is defined as

$$\mathcal{F}^a \mapsto (U \mapsto \operatorname{Hom}_R(\widetilde{\mathfrak{m}}, \mathcal{F}(U)))$$

with the structure of \mathcal{O}_X -module coming from the structure of \mathcal{O}_X -module on \mathcal{F} .

Remark 3.1.16. The functor $(-)_*$ is well-defined, i.e. is independent of a choice of \mathcal{F} and defines a *sheaf* of \mathcal{O}_X -modules. The first claim follows from Lemma 2.1.8(2) and Lemma 3.1.8, the second claim follows from the fact that $\operatorname{Hom}_R(\widetilde{\mathfrak{m}}, -)$ commutes with arbitrary products.

Lemma 3.1.17. Let (X, \mathcal{O}_X) be a ringed *R*-site. Then

- (1) The functor $(-)_*: \operatorname{Mod}_{\mathcal{O}_X^a} \to \operatorname{Mod}_{\mathcal{O}_X}$ is the right adjoint to the exact localization functor $(-)^a: \operatorname{Mod}_{\mathcal{O}_X} \to \operatorname{Mod}_{\mathcal{O}_X^a}$. In particular, it is left exact.
- (2) The unit morphism $\mathcal{F} \to (\mathcal{F}^a)_*$ is an almost isomorphism for any $\mathcal{F} \in \mathbf{Mod}_{\mathcal{O}_X}$. The counit morphism $(\mathcal{G}^a_*)^a \to \mathcal{G}^a$ is an isomorphism for any $\mathcal{G}^a \in \mathbf{Mod}_{\mathcal{O}^a_Y}$.

Proof. It is sufficient to check both claims section-wise. This, in turn, follow from Lemma 2.1.9(1) and Lemma 2.1.9(2) respectively.

Corollary 3.1.18. Let (X, \mathcal{O}_X) be a ringed *R*-site. Then the functor

$$(-)^a \colon \mathbf{Mod}_{\mathcal{O}_X} \to \mathbf{Mod}_{\mathcal{O}_X^a}$$

is exact.

Proof. The functor $(-)^a$ is exact as it has both left and right adjoints.

Theorem 3.1.19. Let (X, \mathcal{O}_X) be a ringed *R*-site. Then the functor

$$(-)^a \colon \mathbf{Mod}^a_{\mathcal{O}_X} \to \mathbf{Mod}_{\mathcal{O}^a_X}$$

is an equivalence of categories.

Proof. Lemma 3.1.17 implies that the functor $(-)^a \colon \mathbf{Mod}_{\mathcal{O}_X} \to \mathbf{Mod}_{\mathcal{O}_X^a}$ has right adjoint functor $(-)_*$ such that the counit morphism $(-)^a \circ (-)_* \to \mathrm{Id}$ is an isomorphism of functors. Moreover, exactness of $(-)^a$ implies that a morphism $\varphi \colon \mathcal{F} \to \mathcal{G}$ is an almost isomorphism if and only if $\varphi^a \colon \mathcal{F}^a \to \mathcal{G}^a$ is an isomorphism. Thus [GZ67, Proposition 1.3] guarantees that the induced functor $(-)^a \colon \mathbf{Mod}_{\mathcal{O}_X} \to \mathbf{Mod}_{\mathcal{O}_X}^a \to \mathrm{Mod}_{\mathcal{O}_X}^a$ is an equivalence.

Remark 3.1.20. In what follows, we do not distinguish $\operatorname{Mod}_{\mathcal{O}_X^a}$ and $\operatorname{Mod}_{\mathcal{O}_X}^a$. Moreover, we denote both categories by Mod_X^a to simplify the notations.

3.2. Basic Functors on the Category Of \mathcal{O}_X^a -Modules. We discuss how to define certain basic functors on \mathbf{Mod}_X^a . Our main functors of interest are Hom, alHom, \otimes , f^* , and f_* . We define their almost analogues and their relation with the original functors. As a by-product we give a slightly more intrinsic definition of $(-)_*: \mathbf{Mod}_X^a \to \mathbf{Mod}_X$ along the lines of the definition of the \mathbf{Mod}_R^a -version of this functor.

For the rest of the section we fix a ringed *R*-site (X, \mathcal{O}_X) .

Definition 3.2.1. • The global Hom functor

 $\operatorname{Hom}_{\mathcal{O}_{X}^{a}}(-,-)\colon \operatorname{\mathbf{Mod}}_{X}^{a,op}\times\operatorname{\mathbf{Mod}}_{X}^{a}\to\operatorname{\mathbf{Mod}}_{R}$

is defined as $(\mathcal{F}^a, \mathcal{G}^a) \mapsto \operatorname{Hom}_{\mathcal{O}_{\mathbf{v}}^a}(\mathcal{F}^a, \mathcal{G}^a).$

• The local Hom functor

$$\underline{\mathcal{H}om}_{\mathcal{O}_X^a}(-,-)\colon \mathbf{Mod}_X^{a,op}\times \mathbf{Mod}_X^a\to \mathbf{Mod}_X$$

is defined as $(\mathcal{F}^a, \mathcal{G}^a) \mapsto \left(U \mapsto \operatorname{Hom}_{\mathcal{O}_U^a}(\mathcal{F}^a|_U, \mathcal{G}^a|_U) \right)$. The standard argument shows that this functor is well-defined, i.e. $\underline{\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}, \mathcal{G})$ is indeed a sheaf of \mathcal{O}_X -modules.

Lemma 3.2.2. Let U be an open in X, and let $\mathcal{F}^a, \mathcal{G}^a$ be \mathcal{O}^a_X -modules. Then the natural map

$$\Gamma\left(U, \underline{\mathcal{H}om}_{\mathcal{O}_X^a}\left(\mathcal{F}^a, \mathcal{G}^a\right)\right) \to \operatorname{Hom}_{\mathcal{O}_U^a}\left(\mathcal{F}^a|_U, \mathcal{G}^a|_U\right)$$

is an isomorphism of $\mathcal{O}_X(U)$ -modules.

Proof. This is evident from the definition.

Lemma 3.2.3. Let (X, \mathcal{O}_X) be a ringed *R*-site. Then there is a functorial isomorphism of \mathcal{O}_X -modules

$$\underline{\mathcal{H}om}_{\mathcal{O}_{\mathbf{Y}}^{a}}(\mathcal{F}^{a},\mathcal{G}^{a})\xrightarrow{\sim}\underline{\mathcal{H}om}_{\mathcal{O}_{\mathbf{Y}}}((\mathcal{F}^{a})_{!},\mathcal{G})$$

for $\mathfrak{F}^a \in \mathbf{Mod}_X^a$ and $\mathfrak{G} \in \mathbf{Mod}_X$.

Proof. Lemma 3.2.2 and Lemma 3.1.13 ensure that the desired isomorphism exists section-wise. It glues to a global isomorphism of sheaves since these section-wise isomorphisms are functorial in U.

Now we move on to show a promised more intrinsic definition of the functor $(-)_*$. As a warm-up we need the following result:

Lemma 3.2.4. Suppose that the ringed *R*-site (X, \mathcal{O}_X) has a final object. By slightly abusing the notation, we also denote the final object by *X*. Then the evaluation map

$$\operatorname{ev}_X \colon \operatorname{Hom}_{\mathcal{O}_X^a} (\mathcal{O}_X^a, \mathcal{G}^a) \to \operatorname{Hom}_{\mathcal{O}_X(X)^a} (\mathcal{O}_X^a(X), \mathcal{G}^a(X))$$

$$\varphi \mapsto \varphi(X)$$

is an isomorphism of $\mathcal{O}_X(X)$ -modules for any $\mathcal{G}^a \in \mathbf{Mod}_X^a$.

Proof. As $(-)^a$ is essentially surjective by Lemma 3.1.13(3), there actually exists some \mathcal{O}_X -module \mathcal{G} with almostification being equal to \mathcal{G}^a . Now we recall that the data of an \mathcal{O}^a_X -linear homomorphism $\varphi \colon \mathcal{O}^a_X \to \mathcal{G}^a$ is equivalent to the data of homomorphisms $\varphi_U \in \operatorname{Hom}_{\mathcal{O}_X(U)^a}(\mathcal{O}^a_X(U), \mathcal{G}^a(U))$ for each open U in X such the diagram

$$\begin{array}{ccc} \mathfrak{O}_{X}(U)^{a} & \xrightarrow{\varphi_{U}} & \mathfrak{G}(U)^{a} \\ & & & \downarrow^{r_{\mathfrak{O}_{X}^{a}}|_{V}^{U}} & \downarrow^{r_{\mathfrak{G}^{a}}|_{V}^{U}} \\ \mathfrak{O}_{X}(V)^{a} & \xrightarrow{\varphi_{V}} & \mathfrak{G}(V)^{a} \end{array}$$

commutes for any $V \subset U$. Now we note that φ_U uniquely determines φ_V in such a diagram. Indeed, this follows from the equality

$$\operatorname{Hom}_{\mathcal{O}_{X}(V)^{a}}(\mathcal{O}_{X}(V)^{a}, \mathfrak{G}(V)^{a}) = \operatorname{Hom}_{\mathcal{O}_{X}(V)}(\widetilde{\mathfrak{m}} \otimes \mathcal{O}_{X}(V), \mathfrak{G}(V))$$
$$= \operatorname{Hom}_{\mathcal{O}_{X}(V)}(\widetilde{\mathfrak{m}} \otimes \mathcal{O}_{X}(U) \otimes_{\mathcal{O}_{X}(U)} \mathcal{O}_{X}(V), \mathfrak{G}(V))$$
$$= \operatorname{Hom}_{\mathcal{O}_{X}(U)}(\widetilde{\mathfrak{m}} \otimes \mathcal{O}_{X}(U), \mathfrak{G}(V))$$
$$= \operatorname{Hom}_{\mathcal{O}_{X}(U)^{a}}(\mathcal{O}_{X}(U)^{a}, \mathfrak{G}(V)^{a}).$$

Now we use the assumption that X is the final object to conclude that any homomorphism $\varphi \colon \mathcal{O}_X^a \to \mathcal{G}^a$ is uniquely defined by $\varphi(X)$. \Box

Corollary 3.2.5. Let (X, \mathcal{O}_X) be an *R*-ringed site, and let $U \in X$ be an open. Then the evaluation map

$$\operatorname{ev}_{U} \colon \operatorname{Hom}_{\mathcal{O}_{U}^{a}}(\mathcal{O}_{U}^{a}, \mathcal{G}|_{U}^{a}) \to \operatorname{Hom}_{\mathcal{O}_{U}(U)^{a}}(\mathcal{O}_{U}^{a}(U), \mathcal{G}^{a}(U))$$
$$\varphi \mapsto \varphi(U)$$

is an isomorphism of $\mathcal{O}_X(U)$ -modules for any $\mathcal{G}^a \in \mathbf{Mod}_X^a$.

Proof. For the purpose of the proof, we can change the site X by the slicing site X/U of objects over U. Then U automatically becomes the final object in X/U, so we can just apply Lemma 3.2.4 to finish the proof.

Now we are ready to prove a new description of the sheaf version of the functor $(-)_*$.

Lemma 3.2.6. Let (X, \mathcal{O}_X) be a ringed *R*-site. Then there is a functorial isomorphism of \mathcal{O}_X -modules

$$\underline{\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{O}_X^a, \mathcal{F}^a) \to \mathcal{F}^a_*$$

for $\mathcal{F}^a \in \mathbf{Mod}_X^a$.

Proof. Lemma 3.2.2 and Corollary 3.2.5 imply that there is an isomorphism of $\mathcal{O}_X(U)$ -modules

$$\Gamma\left(U,\underline{\mathcal{H}om}_{\mathcal{O}_{X}^{a}}\left(\mathcal{O}_{X}^{a},\mathcal{F}^{a}\right)\right)\xrightarrow{\sim}\operatorname{Hom}_{\mathcal{O}_{U}(U)^{a}}\left(\mathcal{O}_{U}^{a}\left(U\right),\mathcal{F}^{a}\left(U\right)\right)$$

that is functorial in both U and \mathcal{F}^a . Now we use the functorial isomorphism of $\mathcal{O}_X(U)$

$$\operatorname{Hom}_{\mathcal{O}_{U}(U)^{a}}\left(\mathcal{O}_{U}(U)^{a}, \mathcal{F}^{a}(U)\right) \simeq \operatorname{Hom}_{R^{a}}\left(R^{a}, \mathcal{F}^{a}(U)\right) = (\mathcal{F}^{a})_{*}(U)$$

to construct a functorial isomorphism

$$\Gamma\left(U, \underline{\mathcal{H}om}_{\mathcal{O}_X^a}\left(\mathcal{O}_X^a, \mathcal{F}^a\right)\right) \xrightarrow{\sim} (\mathcal{F}^a)_*(U) \ .$$

Functoriality in U ensures that it glues to the global isomorphism of \mathcal{O}_X -modules

$$\underline{\mathcal{H}om}_{\mathcal{O}_X^a}\left(\mathcal{O}_X^a, \mathcal{F}^a\right) \xrightarrow{\sim} \mathcal{F}^a_* \ .$$

Now we discuss the functor of almost homomorphisms.

Definition 3.2.7. • The global alHom functor

alHom
$$_{\mathcal{O}_X^a}(-,-)\colon \mathbf{Mod}_X^{a,op} \times \mathbf{Mod}_X^a \to \mathbf{Mod}_R^a$$

is defined as

$$(\mathfrak{F}^a,\mathfrak{G}^a)\mapsto \operatorname{Hom}_{\mathfrak{O}_X^a}(\mathfrak{F}^a,\mathfrak{G}^a)^a\simeq \operatorname{Hom}_{\mathfrak{O}_X}((\mathfrak{F}^a)_!,\mathfrak{G})^a$$

• The local alHom functor

$$\underline{al\mathcal{H}om}_{\mathcal{O}_X^a}(-,-)\colon \mathbf{Mod}_X^{a,op}\times \mathbf{Mod}_X^a\to \mathbf{Mod}_X^a$$

is defined as

$$(\mathfrak{F}^a, \mathfrak{G}^a) \mapsto \left(U \mapsto \mathrm{alHom}_{\mathcal{O}_U^a} (\mathfrak{F}^a|_U, \mathfrak{G}^a|_U)^a \right) \;.$$

Remark 3.2.8. At this point we have not checked that $\underline{alHom}_{\mathcal{O}_X^a}(\mathcal{F}^a, \mathcal{G}^a)$ is actually a sheaf. However, this follows from the following lemma.

Lemma 3.2.9. The natural map

$$\underline{\mathcal{H}om}_{\mathcal{O}_{X}}(\widetilde{\mathfrak{m}}\otimes\mathfrak{F},\mathfrak{G})^{a}\rightarrow\underline{al\mathcal{H}om}_{\mathcal{O}_{Y}^{a}}(\mathfrak{F}^{a},\mathfrak{G}^{a})$$

is an almost isomorphism of \mathcal{O}_X^a -modules for any $\mathcal{F}^a, \mathcal{G}^a \in \mathbf{Mod}^a_{\mathcal{O}_X}$. In particular, <u>*alHom*</u> $_{\mathcal{O}_X^a}(\mathcal{F}^a, \mathcal{G}^a)$ is a sheaf of \mathcal{O}_X^a -modules.

Proof. This follows from the sequence of functorial in U isomorphisms:

$$\frac{\mathcal{H}om_{\mathcal{O}_X}(\widetilde{\mathfrak{m}}\otimes\mathcal{F},\mathcal{G})(U)^a}{\simeq^a \operatorname{alHom}_{\mathcal{O}_U^a}(\mathcal{F}^a|_U,\mathcal{G}^a|_U)} \simeq^a \operatorname{alHom}_{\mathcal{O}_U^a}(\mathcal{F}^a|_U,\mathcal{G}^a|_U) \simeq^a \underline{al\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}^a,\mathcal{G}^a)(U)$$

In order to make Definition 3.2.7, we need to show that these functors can actually be computed by using any representative for \mathcal{F}^a and \mathcal{G}^a .

Proposition 3.2.10. Let (X, \mathcal{O}_X) be a ringed *R*-site. Then:

(1) There is a natural transformation of functors



that makes the diagram 2-commutative. In particular, $\operatorname{alHom}_{\mathcal{O}_X^a}(\mathcal{F}^a, \mathcal{G}^a) \simeq \operatorname{Hom}_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G})^a$ for any $\mathcal{F}, \mathcal{G} \in \operatorname{\mathbf{Mod}}_{\mathcal{O}_X}$.

(2) Then there is a natural transformation of functors

$$\begin{array}{c} \mathbf{Mod}_X^{op} \times \mathbf{Mod}_X \xrightarrow{\underline{\mathcal{H}om}_{\mathbb{O}_X}(-,-)} \mathbf{Mod}_X \\ & \downarrow^{(-)^a \times (-)^a} & \downarrow^{(-)^a} \\ \mathbf{Mod}_X^{a,op} \times \mathbf{Mod}_X^a \xrightarrow{\underline{\mathcal{H}om}_{\mathbb{O}_X}^a(-,-)} \mathbf{Mod}_X^a \end{array}$$

that makes the diagram 2-commutative. In particular, $\underline{al\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}^a, \mathcal{G}^a) \simeq \underline{\mathcal{H}om}_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G})^a$ for any $\mathcal{F}, \mathcal{G} \in \mathbf{Mod}_{\mathcal{O}_X}$.

Proof. The proof is similar to the proof of Proposition 2.2.1(3). The only new thing is that we need to prove an analogue of Corollary 2.1.12, i.e. that the functors $\operatorname{alHom}_{\mathcal{O}_X}(-, \mathcal{G})$, $\operatorname{alHom}_{\mathcal{O}_X}(-, \mathcal{G})$ preserve almost isomorphisms. It essentially boils down to showing that $\operatorname{Ext}^i_{\mathcal{O}_X}(\mathcal{K}, \mathcal{G}) \cong^a 0$ and $\operatorname{Ext}^i_{\mathcal{O}_X}(\mathcal{K}, \mathcal{G}) \cong^a 0$ for any $\mathcal{K} \in \Sigma_X, \mathcal{G} \in \operatorname{Mod}_X$, and an integer $i \geq 0$.

Now Lemma 3.1.5 implies that $\varepsilon \mathcal{K} = 0$ for any $\varepsilon \in \mathfrak{m}$. Thus we see that $\operatorname{Ext}^{i}_{\mathcal{O}_{X}}(\mathcal{K}, \mathcal{G})$ and $\underline{\operatorname{Ext}^{i}}_{\mathcal{O}_{X}}(\mathcal{K}, \mathcal{G})$ are also annihilated by any $\varepsilon \in \mathfrak{m}$ since the functors $\operatorname{Ext}^{i}_{\mathcal{O}_{X}}(-,\mathcal{G})$, $\underline{\operatorname{Ext}^{i}}_{\mathcal{O}_{X}}(-,\mathcal{G})$ are R-linear. Thus $\operatorname{Ext}^{i}_{\mathcal{O}_{X}}(\mathcal{K},\mathcal{G})$ and $\underline{\operatorname{Ext}^{i}}_{\mathcal{O}_{X}}(\mathcal{K},\mathcal{G})$ are almost zero by Lemma 2.1.1 and Lemma 3.1.5 respectively.

Definition 3.2.11. The tensor product functor $-\otimes_{\mathcal{O}_X^a} -: \mathbf{Mod}_X^a \times \mathbf{Mod}_X^a \to \mathbf{Mod}_X^a$ is defined as

$$(\mathfrak{F}^a,\mathfrak{G}^a)\mapsto\mathfrak{F}^a_!\otimes_{\mathfrak{O}_X}\mathfrak{G}^a_!$$

Proposition 3.2.12. There is a natural transformation of functors

$$\begin{array}{ccc} \mathbf{Mod}_X \times \mathbf{Mod}_X & \xrightarrow{-\otimes_{\mathfrak{O}_X} -} & \mathbf{Mod}_X \\ & & \downarrow^{(-)^a \times (-)^a} & \downarrow^{(-)^a} \\ & \mathbf{Mod}_X^a \times \mathbf{Mod}_X^a & \xrightarrow{-\otimes_{\mathfrak{O}_X} -} & \mathbf{Mod}_X^a \end{array}$$

that makes the diagram 2-commutative. In particular, there is a functorial isomorphism

$$(\mathfrak{F} \otimes_{\mathfrak{O}_X} \mathfrak{G})^a \simeq \mathfrak{F}^a \otimes_{\mathfrak{O}_X^a} \mathfrak{G}^a$$

for any $\mathcal{F}, \mathcal{G} \in \mathbf{Mod}_X$.

Proof. The proof is absolutely analogous to that of Propisition 2.2.1(1).

The tensor product is adjoint to $\underline{\mathcal{H}om}$ as it happens in the case of R^a -modules. We give a proof of the local version of this statement.

Lemma 3.2.13. Let (X, \mathcal{O}_X) be a ringed *R*-site, and let $\mathcal{F}^a, \mathcal{G}^a, \mathcal{H}^a$ be \mathcal{O}_X^a -modules. Then there is a functorial isomorphism

$$\underline{\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}^a \otimes_{\mathcal{O}_X^a} \mathcal{G}^a, \mathcal{H}^a) \simeq \underline{\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}^a, \underline{al\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{G}^a, \mathcal{H}^a)) \ .$$

After passing to the global sections, this gives the isomorphism

$$\operatorname{Hom}_{\mathcal{O}_X^a}(\mathcal{F}^a \otimes_{\mathcal{O}_X^a} \mathcal{G}^a, \mathcal{H}^a) \simeq \operatorname{Hom}_{\mathcal{O}_X^a}(\mathcal{F}^a, \underline{al\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{G}^a, \mathcal{H}^a)) \ .$$

And after passing to the almostifications, it gives an isomorphism

 $\underline{al\mathcal{H}om}_{\mathcal{O}_{Y}^{a}}(\mathcal{F}^{a}\otimes_{\mathcal{O}_{Y}^{a}}\mathcal{G}^{a},\mathcal{H}^{a})\simeq\underline{al\mathcal{H}om}_{\mathcal{O}_{Y}^{a}}(\mathcal{F}^{a},\underline{al\mathcal{H}om}_{\mathcal{O}_{Y}^{a}}(\mathcal{G}^{a},\mathcal{H}^{a})).$

Proof. We compute $\Gamma(U, \underline{\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}^a \otimes_{\mathcal{O}_X^a} \mathcal{G}^a, \mathcal{H}^a))$ by using Lemma 3.2.2 and the standard \otimes - $\underline{\mathcal{H}om}$ adjunction. Namely,

$$\Gamma\left(U, \underline{\mathcal{H}om}_{\mathbb{O}_{X}^{a}}\left(\mathfrak{F}^{a}\otimes_{\mathbb{O}_{X}^{a}}\mathfrak{G}^{a}, \mathfrak{H}^{a}\right)\right) \simeq \operatorname{Hom}_{\mathbb{O}_{U}^{a}}\left(\mathfrak{F}^{a}|_{U}\otimes_{\mathbb{O}_{U}^{a}}\mathfrak{G}^{a}|_{U}, \mathfrak{H}^{a}|_{U}\right) \qquad \text{Lemma 3.2.2}$$

$$\simeq \operatorname{Hom}_{\mathbb{O}_{U}^{a}}\left(\left(\mathfrak{F}|_{U}\otimes_{\mathbb{O}_{U}}\mathfrak{G}|_{U}\right)^{a}, \mathfrak{H}^{a}|_{U}\right) \qquad \text{Proposition 3.2.12}$$

$$\simeq \operatorname{Hom}_{\mathbb{O}_{U}}\left(\widetilde{\mathfrak{m}}\otimes\left(\mathfrak{F}|_{U}\otimes_{\mathbb{O}_{U}}\mathfrak{G}|_{U}\right), \mathfrak{H}|_{U}\right) \qquad \text{Lemma 3.1.13}$$

$$\simeq \operatorname{Hom}_{\mathbb{O}_{U}}\left(\left(\widetilde{\mathfrak{m}}\otimes\mathfrak{F}|_{U}\right)\otimes_{\mathbb{O}_{U}}\left(\widetilde{\mathfrak{m}}\otimes\mathfrak{G}|_{U}\right), \mathfrak{H}|_{U}\right) \qquad \widetilde{\mathfrak{m}}^{\otimes 2}\simeq\widetilde{\mathfrak{m}}$$

$$\simeq \operatorname{Hom}_{\mathbb{O}_{U}}\left(\widetilde{\mathfrak{m}}\otimes\mathfrak{F}|_{U}, \underline{\mathcal{H}om}_{\mathbb{O}_{U}}\left(\widetilde{\mathfrak{m}}\otimes\mathfrak{G}|_{U}, \mathfrak{H}|_{U}\right)\right) \qquad \otimes -\mathcal{H}om \text{ adjunction}$$

$$\simeq \operatorname{Hom}_{\mathbb{O}_{U}^{a}}\left(\mathfrak{F}^{a}|_{U}, \underline{al\mathcal{H}om}_{\mathbb{O}_{U}}\left(\widetilde{\mathfrak{m}}\otimes\mathfrak{G}|_{U}, \mathfrak{H}|_{U}\right)\right) \qquad \text{Lemma 3.1.13}$$

$$\simeq \Gamma\left(U, \underline{\mathcal{H}om}_{\mathbb{O}_{X}^{a}}\left(\mathfrak{F}^{a}, \underline{al\mathcal{H}om}_{\mathbb{O}_{X}^{a}}\left(\mathfrak{G}^{a}, \mathfrak{H}^{a}\right)\right)\right) \qquad \text{Lemma 3.2.2}$$

Since these identifications are functorial in U, we can glue them to a global isomorphism

$$\underline{\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}^a \otimes_{\mathcal{O}_X^a} \mathcal{G}^a, \mathcal{H}^a) \simeq \underline{\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}^a, \underline{al\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{G}^a, \mathcal{H}^a)) \ .$$

This finishes the proof.

Corollary 3.2.14. Let (X, \mathcal{O}_X) be a ringed *R*-site, and let \mathcal{F}^a be an \mathcal{O}^a_X -module. Then the functor $-\otimes_{\mathcal{O}^a_X} \mathcal{F}^a$ is left adjoint to $\underline{al\mathcal{H}om}_{\mathcal{O}^a_X}(\mathcal{F}^a, -)$.

For what follows, we fix a map $f: (X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ of ringed *R*-sites. We are going to define the almost version of the pullback and pushforward functors.

Definition 3.2.15. The *pullback functor* $f_a^* \colon \mathbf{Mod}_X^a \to \mathbf{Mod}_Y^a$ is defined as

$$\mathfrak{F}^a \mapsto (f^*(\mathfrak{F}^a_!))^a$$

In what follows, we will often abuse notation and simply write f^* instead of f_a^* . This is "allowed" by Proposition 3.2.19.

As always, we want to show that this functor can be actually computed by applying f^* to any representative of \mathcal{F}^a . The main ingredient is to show that f^* sends almost isomorphisms to almost isomorphisms. The following lemma shows slightly more, and will be quite useful later on.

Lemma 3.2.16. Let $f: (X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ be a morphism of ringed *R*-sites. Then for any \mathcal{O}_X -module \mathcal{F} , there is a natural isomorphism $\varphi_f(\mathcal{F}): f^*(\widetilde{\mathfrak{m}} \otimes \mathcal{F}) \to \widetilde{\mathfrak{m}} \otimes f^*\mathcal{F}$ functorial in \mathcal{F} .

Proof. We use Remark 3.1.2 to say that $\widetilde{\mathfrak{m}} \otimes \mathcal{F}$ is functorially isomorphic to $\widetilde{\mathfrak{m}}_Y \otimes_{\mathcal{O}_Y} \mathcal{F}$, where $\underline{\widetilde{\mathfrak{m}}}_Y := \underline{\widetilde{\mathfrak{m}}} \otimes_R \mathcal{O}_Y$. Now we note that $f^*(\underline{\widetilde{\mathfrak{m}}}_Y) \simeq \underline{\widetilde{\mathfrak{m}}}_X$ as can be easily seen (using the $\widetilde{\mathfrak{m}}$ is *R*-flat) from the very definitions. Therefore, $\varphi_f(\mathcal{F})$ comes from the fact that the pullback functor commutes with the tensor product. More precisely, we define it as the composition

$$f^*(\widetilde{\mathfrak{m}}\otimes\mathfrak{F})\xrightarrow{\sim} f^*(\underline{\widetilde{\mathfrak{m}}}_Y\otimes_{\mathbb{O}_Y}\mathfrak{F})\xrightarrow{\sim} f^*(\underline{\widetilde{\mathfrak{m}}}_Y)\otimes_{\mathbb{O}_X}f^*(\mathfrak{F})\xrightarrow{\sim} \underline{\widetilde{\mathfrak{m}}}_X\otimes_{\mathbb{O}_X}f^*(\mathfrak{F}) \ .$$

We now also show a derived version of Lemma 3.2.16 that will be used later in the text.

Lemma 3.2.17. Let $f: (X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ be a morphism of ringed *R*-sites. Then for any $\mathcal{F} \in \mathbf{D}(X)$, there is a natural isomorphism

$$\varphi_f(\mathfrak{F})\colon \mathbf{L}f^*(\widetilde{\mathfrak{m}}\otimes\mathfrak{F})\to\widetilde{\mathfrak{m}}\otimes\mathbf{L}f^*\mathfrak{F}$$

functorial in \mathcal{F} . In particular, $f^*(\widetilde{\mathfrak{m}} \otimes \mathcal{F}) \simeq \widetilde{\mathfrak{m}} \otimes f^*(\mathcal{F})$ for any $\mathcal{F} \in \mathbf{Mod}_X$.

Proof. Similarly, we use Remark 3.1.2 to say that $\widetilde{\mathfrak{m}} \otimes \mathcal{F}$ is functorially isomorphic to $\widetilde{\mathfrak{m}}_Y \otimes_{\mathfrak{O}_Y} \mathcal{F}$, where $\underline{\widetilde{\mathfrak{m}}}_Y \coloneqq \underline{\widetilde{\mathfrak{m}}} \otimes_R \mathfrak{O}_Y$. Now we note that $\mathbf{L}f^*(\underline{\widetilde{\mathfrak{m}}}_Y) \simeq f^*(\underline{\widetilde{\mathfrak{m}}}_Y) \simeq \underline{\widetilde{\mathfrak{m}}}_X$ as $\widetilde{\mathfrak{m}}$ is *R*-flat. The rest of the proof is the same using the $\mathbf{L}f^*$ functorially commutes with the derived tensor product. \Box

Corollary 3.2.18. Let $f: (X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ be a morphism of ringed *R*-sites, and let $\varphi: \mathcal{F} \to \mathcal{G}$ be an almost isomorphism of \mathcal{O}_Y -modules. Then the homomorphism $f^*(\varphi): f^*(\mathcal{F}) \to f^*(\mathcal{G})$ is an almost isomorphism.

Proof. The question boils down to show that the homomorphism

$$\widetilde{\mathfrak{m}} \otimes f^*(\mathfrak{F}) \to \widetilde{\mathfrak{m}} \otimes f^*(\mathfrak{G})$$

is an isomorphism. Lemma 3.2.16 ensures that it is sufficient to prove that the map

$$f^*(\widetilde{\mathfrak{m}}\otimes\mathfrak{F})\to f^*(\widetilde{\mathfrak{m}}\otimes\mathfrak{G})$$

is an isomorphism. But this is clear as the map $\widetilde{\mathfrak{m}} \otimes \mathfrak{F} \to \widetilde{\mathfrak{m}} \otimes \mathfrak{G}$ is already an isomorphism. \Box

Proposition 3.2.19. Let $f: (X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ be a morphism of ringed *R*-sites. Then there is a natural transformation of functors



that makes the diagram 2-commutative. In particular, there is a functorial isomorphism $(f^*\mathcal{F})^a \simeq f_a^*(\mathcal{F}^a)$ for any $\mathcal{F} \in \mathbf{Mod}_X$.

Proof. The proof is similar to Proposition 2.2.1. We define $\rho_{\mathcal{F}}: f^*(\widetilde{\mathfrak{m}} \otimes \mathcal{F})^a \to f^*(\mathcal{F})^a$ as the map induced by the natural homomorphism $\widetilde{\mathfrak{m}} \otimes \mathcal{F} \to \mathcal{F}$. It is clearly functorial in \mathcal{F} , and it is an isomorphism by Corollary 3.2.18.

Definition 3.2.20. The pushforward functor $f_*^a: \operatorname{Mod}_X^a \to \operatorname{Mod}_Y^a$ is defined as

$$\mathfrak{F}^a \mapsto (f_*(\mathfrak{F}^a_!))^a$$

In what follows, we will often abuse the notations and simply write f_* instead of f_*^a . This is "allowed" by Proposition 3.2.24.

Definition 3.2.21. The global sections functor $\Gamma^a(X, -)$: $\mathbf{Mod}_X^a \to \mathbf{Mod}_R^a$ is defined as

$$\mathfrak{F}^a \mapsto \Gamma(X, \mathfrak{F}^a)^a$$
.

In what follows, we will often abuse the notations and simply write Γ instead of Γ_a . This is also "allowed" by Proposition 3.2.24.

Remark 3.2.22. The global section functor can be realized as the pushforward along the map $(X, \mathcal{O}_X) \to (*, R)$.

Lemma 3.2.23. Let $f: (X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ be a morphism of ringed *R*-sites, and let $\varphi: \mathcal{F} \to \mathcal{G}$ be an almost isomorphism. Then the morphism $f_*(\varphi): f_*(\mathcal{F}) \to f_*(\mathcal{G})$ is an almost isomorphism.

Proof. The standard argument with considering the kernel and cokernel of φ shows that it is sufficient to prove that $f_*\mathcal{K} \cong^a 0$, $\mathbb{R}^1 f_*\mathcal{K} \cong^a 0$ for any almost zero \mathcal{O}_X -module \mathcal{K} . This follows from *R*-linearity of f_* and Lemma 3.1.5.

Proposition 3.2.24. Let $f: (X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ be a morphism of ringed *R*-spaces. Then there is a natural transformation of functors



that makes the diagram 2-commutative. In particular, there is a functorial isomorphism $(f_*\mathcal{F})^a \simeq f_*^a(\mathcal{F}^a)$ for any $\mathcal{F} \in \mathbf{Mod}_X$. The same results hold true for $\Gamma^a(X, -)$.

Proof. We define $\rho_{\mathcal{F}}: f_*(\widetilde{\mathfrak{m}} \otimes \mathfrak{F})^a \to f_*(\mathfrak{F})^a$ as the map induced by the natural homomorphism $\widetilde{\mathfrak{m}} \otimes \mathfrak{F} \to \mathfrak{F}$. It is clearly functorial in \mathfrak{F} , and it is an isomorphism by Lemma 3.2.23.

Lemma 3.2.25. Let (X, \mathcal{O}_X) be a ringed *R*-site, and let \mathcal{F}, \mathcal{G} be \mathcal{O}_X^a -modules. Then there is a natural morphism

$$\Gamma\left(U, \underline{al\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}^a, \mathcal{G}^a)\right) \to \mathrm{alHom}_{\mathcal{O}_U^a}(\mathcal{F}^a|_U, \mathcal{G}^a|_U)$$

is an isomorphism of \mathbb{R}^a -modules for any open $U \subset X$.

Proof. The claim easily follows from Lemma 3.2.2, Proposition 3.2.10(2), and Proposition 3.2.24

Lemma 3.2.26. Let $f: (X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ be a morphism of ringed *R*-site, and let $\mathcal{F}^a \in \mathbf{Mod}_Y^a$, and $\mathcal{G}^a \in \mathbf{Mod}_X^a$. Then there is a functorial isomorphism of \mathcal{O}_Y -modules

$$f_* \underline{\mathcal{H}om}_{\mathcal{O}_X^a}(f^*(\mathcal{F}^a), \mathcal{G}^a) \simeq \underline{\mathcal{H}om}_{\mathcal{O}_Y^a}(\mathcal{F}^a, f_*(\mathcal{G}^a))$$
.

After passing to the global sections, this gives the isomorphism of $\mathcal{O}_Y(Y)$ -modules

$$\operatorname{Hom}_{\mathcal{O}_{Y}^{a}}(f^{*}(\mathcal{F}^{a}),\mathcal{G}^{a}) \simeq \operatorname{Hom}_{\mathcal{O}_{Y}^{a}}(\mathcal{F}^{a},f_{*}(\mathcal{G}^{a}))$$

And after passing to the almostifications, it gives the isomorphism of \mathcal{O}_V^a -modules

$$f_*\underline{al\mathcal{H}om}_{\mathcal{O}_{\mathbf{V}}^a}(f^*(\mathcal{F}^a),\mathcal{G}^a) \cong^a \underline{al\mathcal{H}om}_{\mathcal{O}_{\mathbf{V}}^a}(\mathcal{F}^a,f_*(\mathcal{G}^a))$$

Proof. This is a combination of the classical (f^*, f_*) -adjunction, Lemma 3.1.13, Lemma 3.2.16, Proposition 3.2.19, and Proposition 3.2.24. Indeed, we choose an open $U \subset Y$ and denote its preimage by $V \coloneqq f^{-1}(U)$. We also define $\mathcal{F}^a_U \coloneqq \mathcal{F}^a|_U$ and $\mathcal{G}^a_V \coloneqq \mathcal{G}^a|_V$. The claim follows follows from the sequence of functorial isomorphisms

$$\begin{split} \Gamma\left(U,\underline{\mathcal{H}om}_{\mathbb{O}_{Y}^{a}}\left(\mathcal{F}^{a},f_{*}\left(\mathcal{G}^{a}\right)\right)\right) &\simeq \operatorname{Hom}_{\mathbb{O}_{U}^{a}}\left(\mathcal{F}_{U}^{a},f_{*}\left(\mathcal{G}_{V}^{a}\right)\right) & \text{Lemma 3.2.2} \\ &\simeq \operatorname{Hom}_{\mathbb{O}_{U}^{a}}\left(\mathcal{F}_{U}^{a},f_{*}\left(\mathcal{G}_{V}\right)^{a}\right) & \text{Proposition 3.2.24} \\ &\simeq \operatorname{Hom}_{\mathbb{O}_{U}}\left(\widetilde{\mathfrak{m}}\otimes\mathcal{F}_{U},f_{*}\left(\mathcal{G}_{V}\right)\right) & \text{Lemma 3.1.13} \\ &\simeq \operatorname{Hom}_{\mathbb{O}_{V}}\left(f^{*}\left(\widetilde{\mathfrak{m}}\otimes\mathcal{F}_{U}\right),\mathcal{G}_{V}\right) & (f^{*},f_{*})\text{-adjunction} \\ &\simeq \operatorname{Hom}_{\mathbb{O}_{V}}\left(\widetilde{\mathfrak{m}}\otimes f^{*}\left(\mathcal{F}_{U}\right),\mathcal{G}_{V}\right) & \text{Lemma 3.2.16} \\ &\simeq \operatorname{Hom}_{\mathbb{O}_{V}^{a}}\left(f^{*}\left(\mathcal{F}_{U}\right)^{a},\mathcal{G}_{V}^{a}\right) & \text{Lemma 3.1.13} \\ &\simeq \operatorname{Hom}_{\mathbb{O}_{V}^{a}}\left(f^{*}\left(\mathcal{F}_{U}^{a}\right),\mathcal{G}_{V}^{a}\right) & \text{Proposition 3.2.19} \\ &\simeq \Gamma\left(U,f_{*}\underline{\mathcal{H}om}_{\mathbb{O}_{X}^{a}}\left(f^{*}\left(\mathcal{F}^{a}\right),\mathcal{G}^{a}\right)\right) & \text{Lemma 3.2.2} \end{split}$$

Since these identifications are functorial in U, we can glue them to a global isomorphism

 $f_* \underline{\mathcal{H}om}_{\mathcal{O}^a_X}(f^*(\mathcal{F}^a), \mathcal{G}^a) \simeq \underline{\mathcal{H}om}_{\mathcal{O}^a_V}(\mathcal{F}^a, f_*(\mathcal{G}^a)) \ .$

57

Corollary 3.2.27. Let $f: (X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ be a morphism of ringed *R*-site. Then the functors $\operatorname{Mod}_X^a \xleftarrow{f^*}_{f_*} \operatorname{Mod}_Y^a$ are adjoint.

3.3. The Projection Formula. The definition of \mathcal{O}_X -modules behaves especially nicely on locally spectral spaces¹⁶. For instance, we show that we can explicitly describe sections of $\widetilde{\mathfrak{m}} \otimes \mathfrak{F}$ on a basis of opens for such spaces. Moreover, we show that the projection formula holds for spectral morphisms of locally spectral spaces.

Remark 3.3.1. We mention one problem of working with locally spectral spaces that we deliberately avoid in all of our proofs. Suppose that X is a locally ringed space and $U \subset X$ is an open spectral subspace then the natural map $U \to X$ need not be quasi-compact. In particular, an intersection of two open spectral subspaces in X need not be spectral itself.

In order to get such examples, one can consider X to a scheme that is not quasi-separated and U an open affine subscheme. Then the inclusion map $U \to X$ is usually not quasi-compact.

Lemma 3.3.2. Let (X, \mathcal{O}_X) be a locally spectral, locally ringed *R*-space. Then for any spectral¹⁷ open subset $U \subset X$ the natural morphism

$$\widetilde{\mathfrak{m}} \otimes_R \mathfrak{F}(U) \to (\widetilde{\mathfrak{m}} \otimes \mathfrak{F})(U)$$

is an isomorphism of $\mathcal{O}_X(U)$ -modules.

Proof. As spectral subspaces form a basis of topology on X, it suffices to show that the functor

$$U \to \widetilde{\mathfrak{m}} \otimes_R \mathfrak{F}(U)$$

restricted to spectral open subsets satisfy the sheaf condition. In particular, we can assume that X itself is spectral.

As any open spectral U is quasi-compact, we conclude that any open covering $U = \bigcup_{i \in I} U_i$ admits a refinement by a finite one. Thus, it is sufficient to check the sheaf condition for finite coverings of a spectral spaces by spectral open subspaces. Thus, we need to show that, for any finite covering $U = \bigcup_{i \in I} U_i$, the sequence

$$0 \to \widetilde{\mathfrak{m}} \otimes_R \mathfrak{F}(U) \to \prod_{i=1}^n (\widetilde{\mathfrak{m}} \otimes_R \mathfrak{F}(U_i)) \to \prod_{i,j=1}^n (\widetilde{\mathfrak{m}} \otimes_R \mathfrak{F}(U_i \cap U_j)).$$

is exact. But this follows from flatness of $\widetilde{\mathfrak{m}}$ and the fact that tensor product commutes with *finite* direct products.

Now we want to show a version of the projection formula for the functor $\widetilde{\mathfrak{m}} \otimes -$, it will take some time to rigorously prove it. We recall that a map of locally spectral spaces is called *spectral*, if the pre-image of any spectral open subset is spectral.

Lemma 3.3.3. Let (X, \mathcal{O}_X) be a spectral locally ringed *R*-space. Then for any injective \mathcal{O}_X -module \mathcal{I} the \mathcal{O}_X -module $\widetilde{\mathfrak{m}} \otimes \mathcal{I}$ is an $\mathrm{H}^0(X, -)$ -acyclic.

¹⁶We refer to [Sta21, Tag 08YF] and [Wed19, §3] for a comprehensive discussion of (locally) spectral spaces

¹⁷We remind the reader that actually any quasi-compact quasi-separated open subset of a locally spectral space is spectral. This can be easily seen from the definitions.

Proof. We start the proof by noting that [Sta21, Tag 01EV] guarantees that it suffices to show that the Čech cohomology groups $\check{\mathrm{H}}^{i}(U, \widetilde{\mathfrak{m}} \otimes \mathfrak{I})$ vanish for all open subsets $U \subset X$ and i > 0. Since any open subset of a locally spectral space is locally spectral, it suffices to show that $\check{\mathrm{H}}^{i}(U, \widetilde{\mathfrak{m}} \otimes \mathfrak{I}) = 0$ for i > 0.

We note that quasi-compact opens form a basis for the topology on X. Since X is quasi-compact, finite coverings by quasi-compact opens form a cofinal subsystem in the system of coverings of X. Thus it is enough to check vanishing of higher $\check{H}^{i}(\mathcal{U}, \widetilde{\mathfrak{m}} \otimes \mathcal{I})$ for any such coverings \mathcal{U} of X.

We pick such a covering $\mathcal{U}: X = \bigcup_{i=1}^{n} U_i$ and observe that all the intersections $U_{i_1,\ldots,i_m} = \bigcap_{k=1}^{m} U_{i_k}$ are again quasi-compact by spectrality of X. In particular, they are spectral. Now we invoke [Sta21, Tag 0A36] to say that it suffices to show that

$$(\widetilde{\mathfrak{m}}\otimes\mathfrak{I})(V)\xrightarrow{r_{\widetilde{\mathfrak{m}}\otimes\mathfrak{I}}|_U^V}(\widetilde{\mathfrak{m}}\otimes\mathfrak{I})(U)$$

is surjective for any inclusion of any *spectral* open subsets $U \hookrightarrow V$. Lemma 3.3.2 says that this map $r_{\widetilde{\mathfrak{m}}\otimes \mathbb{J}}|_{U}^{V}$ is identified with the map

$$\widetilde{\mathfrak{m}} \otimes_R \mathfrak{I}(V) \xrightarrow{\widetilde{\mathfrak{m}} \otimes_R r_{\mathfrak{I}}|_U^V} \widetilde{\mathfrak{m}} \otimes_R \mathfrak{I}(U).$$

But now we note that $r_{\mathfrak{I}}|_U^V$ is surjective since any injective \mathcal{O}_X -module is flasque by [Sta21, Tag 01EA], and therefore the map $\widetilde{\mathfrak{m}} \otimes_R r_{\mathfrak{I}}|_U^V$ is surjective as well.

Corollary 3.3.4. Let $f: (X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ be a spectral morphism of locally spectral, locally ringed *R*-spaces, and let \mathfrak{I} be an injective \mathcal{O}_X -module. Then $\widetilde{\mathfrak{m}} \otimes \mathfrak{I}$ is an $f_*(-)$ -acyclic

Proof. It suffices to show that for any open spectral $U \subset Y$ the higher cohomology groups

$$\mathrm{H}^{i}(X_{U}, (\widetilde{\mathfrak{m}} \otimes \mathfrak{I})|_{X_{U}})$$

vanish. This follows from Lemma 3.3.3 since X_U is spectral by the assumption on f.

Lemma 3.3.5. Let $f: (X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ be a spectral morphism of locally spectral, locally ringed R-spaces, and let \mathcal{F} be an \mathcal{O}_X -module. Then there is an isomorphism

$$\beta:\widetilde{\mathfrak{m}}\otimes f_*\mathcal{F}\to f_*(\widetilde{\mathfrak{m}}\otimes\mathcal{F})$$

functorial in \mathcal{F} .

Proof. It suffices to define a morphism on a basis of spectral open subspaces $U \subset Y$. For any such $U \subset Y$ we define

$$\beta_U : (\widetilde{\mathfrak{m}} \otimes f_* \mathfrak{F})(U) \to f_*(\widetilde{\mathfrak{m}} \otimes \mathfrak{F})(U)$$

as the composition of isomorphisms

$$(\widetilde{\mathfrak{m}} \otimes f_* \mathfrak{F})(U) \xrightarrow{\alpha_U^{-1}} \widetilde{\mathfrak{m}} \otimes_R (f_* \mathfrak{F})(U) = \widetilde{\mathfrak{m}} \otimes_R \mathfrak{F}(X_U) \xrightarrow{\alpha_{X_U}} (\widetilde{\mathfrak{m}} \otimes \mathfrak{F})(X_U) = f_*(\widetilde{\mathfrak{m}} \otimes \mathfrak{F})(U)$$

with α_U and α_{X_U} isomorphisms from Lemma 3.3.2. Since the construction of α was functorial in U we conclude that β defines a morphism of sheaves. It is an isomorphism because we constructed β_U to be isomorphism and basis of Y.

Lemma 3.3.6. Let $f: (X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ be a spectral morphism of locally spectral, locally ringed R-spaces. Then for any $\mathcal{F} \in \mathbf{D}(X)$, there is a morphism

$$\rho_f(\mathcal{F}): \widetilde{\mathfrak{m}} \otimes \mathbf{R} f_* \mathcal{F} \to \mathbf{R} f_*(\widetilde{\mathfrak{m}} \otimes \mathcal{F})$$

functorial in F. This map is an isomorphism in either of the following cases:

• The complex \mathcal{F} is bounded below, i.e. $\mathcal{F} \in \mathbf{D}^+(X)$, or

• The space X is locally of uniformly bounded Krull dimension and $\mathcal{F} \in \mathbf{D}(X)$.

Proof. We start the proof by constructing the map $\rho_f(\mathcal{F})$. Note that by the adjunction, it suffices to construct a map

$$\mathbf{L}f^*(\widetilde{\mathfrak{m}}\otimes\mathbf{R}f_*\mathcal{F})\to\widetilde{\mathfrak{m}}\otimes\mathcal{F}$$

We also denote the counit of the adjunction between $\mathbf{L}f^*$ and $\mathbf{R}f_*$ by

$$\eta_{\mathfrak{F}} \colon \mathbf{L}f^*\mathbf{R}f_*\mathfrak{F} \to \mathfrak{F}$$

Then we define the map

$$\mathbf{L}f^*(\widetilde{\mathfrak{m}}\otimes\mathbf{R}f_*\mathfrak{F})\to\widetilde{\mathfrak{m}}\otimes\mathfrak{F}$$

as the composition

$$\mathbf{L}f^{*}(\widetilde{\mathfrak{m}}\otimes\mathbf{R}f_{*}\mathcal{F})\xrightarrow{\varphi_{f}(\mathbf{R}f_{*}\mathcal{F})}\widetilde{\mathfrak{m}}\otimes\mathbf{L}f^{*}\mathbf{R}f_{*}\mathcal{F}\xrightarrow{\widetilde{\mathfrak{m}}\otimes\eta_{\mathcal{F}}}\widetilde{\mathfrak{m}}\otimes\mathcal{F}$$

where the first map is the isomorphism coming from Lemma 3.2.17 and the second map comes from the adjunction morphism $\varepsilon_{\mathcal{F}}$.

Now we show that $\rho_f(\mathcal{F})$ is an isomorphism for $\mathcal{F} \in \mathbf{D}^+(X)$. We choose an injective resolution $\mathcal{F} \to \mathcal{I}^{\bullet}$. In this case we use Corollary 3.3.4 to note that β is the natural map

$$\widetilde{\mathfrak{m}} \otimes f_*(\mathfrak{I}^{\bullet}) \to f_*(\widetilde{\mathfrak{m}} \otimes \mathfrak{I}^{\bullet})$$

that is an isomorphism by Lemma 3.3.5.

The last thing we need to show is that $\rho_f(\mathcal{F})$ is an isomorphism for any $\mathcal{F} \in \mathbf{D}(X)$ if X is locally of uniformly bounded Krull dimension. The claim is local, so we may and do assume that both X and Y are spectral spaces. As X is quasi-compact (as it is spectral now) and locally of finite Krull dimension, we conclude that X has finite Krull dimension, say $N := \dim X$. Then [Sch92, Corollary 4.6] (another reference is [Sta21, Tag 0A3G]) implies that $\mathrm{H}^i(U, \mathcal{G}) = 0$ for any open spectral $U \subset X$, $\mathcal{G} \in \mathbf{Mod}_X$, and i > N. In particular, $\mathrm{R}^i f_* \mathcal{G} = 0$ for any $\mathcal{G} \in \mathbf{Mod}_X$, and i > N. Thus we see that the assumptions of [Sta21, Tag 0D6U] are verified in this case (with $\mathcal{A} = \mathbf{Mod}_X$ and $\mathcal{A}' = \mathbf{Mod}_Y$), so the natural map

$$\mathcal{H}^{j}(\mathbf{R}f_{*}\mathcal{F}) \to \mathcal{H}^{j}(\mathbf{R}f_{*}(\tau^{\geq -n}\mathcal{F}))$$

is an isomorphism for any $\mathcal{F} \in \mathbf{D}(X)$, $j \geq N - n$. As $\widetilde{\mathfrak{m}}$ is *R*-flat, we get the commutative diagram

with the vertical arrows being isomorphisms for $j \ge N - n$, and the bottom horizontal map is an isomorphism as $\tau^{\ge -n} \mathcal{F} \in \mathbf{D}^+(X)$. Thus, by choosing an appropriate $n \ge 0$, we see that $\mathcal{H}^j(\rho_{\mathcal{F}})$ is an isomorphism for any j; so $\rho_{\mathcal{F}}$ is an isomorphism itself. \Box

3.4. Derived Category of \mathcal{O}_X^a -Modules. This section is a global analogue of Section 2.3. We give two different definitions of the derived category of almost \mathcal{O}_X -modules and show that they coincide.

Definition 3.4.1. We define the *derived category of* \mathcal{O}^a_X *-modules* as $\mathbf{D}(X^a) \coloneqq \mathbf{D}(\mathbf{Mod}^a_X)$.

We define the bounded version of derived category of almost *R*-modules $\mathbf{D}^*(X^a)$ for $* \in \{+, -, b\}$ as the full subcategory of $\mathbf{D}(X^a)$ consisting of bounded below (resp. bounded above, resp. bounded) complexes.

Definition 3.4.2. We define the almost derived category of \mathcal{O}_X -modules as the Verdier quotient¹⁸ $\mathbf{D}(X)^a \coloneqq \mathbf{D}(\mathbf{Mod}_X)/\mathbf{D}_{\Sigma_X}(\mathbf{Mod}_X).$

Remark 3.4.3. We recall that Σ_X is the Serre subcategory of \mathbf{Mod}_X that consists of almost zero \mathcal{O}_X -modules.

We note that the functor $(-)^a \colon \mathbf{Mod}_X \to \mathbf{Mod}_X^a$ is exact and additive. Thus it can be derived to the functor $(-)^a \colon \mathbf{D}(X) \to \mathbf{D}(X^a)$. Similarly, the functor $(-)_! \colon \mathbf{Mod}_X^a \to \mathbf{Mod}_X$ can be derived to the functor $(-)_! \colon \mathbf{D}(X^a) \to \mathbf{D}(X)$. The standard argument shows that $(-)_!$ is a left adjoint functor to the functor $(-)^a$ as this already happens on the level of abelian categories.

We also want to establish a derived version of the functor $(-)_*$. But since functor is only left exact, we do need to do some work to derive it. Namely, we need to ensure that \mathcal{O}_X^a -modules admit enough *K*-injective complexes.

Definition 3.4.4. We say that a complex of \mathcal{O}_X^a -module $I^{\bullet,a}$ is *K*-injective if $\operatorname{Hom}_{K(\mathcal{O}_X^a)}(C^{\bullet,a}, I^{\bullet,a}) = 0$ for any acyclic complex $C^{\bullet,a}$ of R^a -modules.

Remark 3.4.5. We remind the reader that $K(\mathcal{O}_X^a)$ stands for the homotopy category of \mathcal{O}_X^a -modules.

Lemma 3.4.6. The functor $(-)^a$: $\mathbf{Comp}(\mathcal{O}_X) \to \mathbf{Comp}(\mathcal{O}_X^a)$ sends *K*-injective \mathcal{O}_X^a -complexes to *K*-injective \mathcal{O}_X^a -complexes.

Proof. We note that $(-)^a$ admits an exact left adjoint $(-)_!$ thus [Sta21, Tag 08BJ] ensures that $(-)^a$ preserves K-injective complexes.

Corollary 3.4.7. Let (X, \mathcal{O}_X) be a ringed *R*-site. Then every object $\mathcal{F}^{\bullet,a} \in \mathbf{Comp}(\mathcal{O}_X^a)$ is quasiisomorphic to a *K*-injective complex.

Proof. The proof of Corollary 2.3.6 works verbatim with the only exception that one needs to use [Sta21, Tag 079P] instead of [Sta21, Tag 090Y].

Now, similarly to the case of R^a -modules, we define the functor $(-)_*: \mathbf{D}(X^a) \to \mathbf{D}(X)$ as the derived functor of $(-)_*: \mathbf{Mod}_X^a \to \mathbf{Mod}_X$. This functor exists by [Sta21, Tag 070K].

Lemma 3.4.8. Let (X, \mathcal{O}_X) be a ringed *R*-site. Then

(1) The functors $\mathbf{D}(X) \xleftarrow{(-)!}{(-)^a} \mathbf{D}(X^a)$ are adjoint. Moreover, the counit (resp. unit) mor-

phism

$$(\mathfrak{F}^a)_! \to \mathfrak{F} \text{ (resp. } \mathfrak{G} \to (\mathfrak{G}_!)^a)$$

is an almost isomorphism (resp. isomorphism) for any $\mathcal{F} \in \mathbf{D}(X), \mathcal{G} \in \mathbf{D}(X^a)$. In particular, the functor $(-)^a$ is essentially surjective.

¹⁸We refer to [Sta21, Tag 05RA] for an extensive discussion of Verdier quotients of triangulated categories.

(2) The functor $(-)^a : \mathbf{D}(X) \to \mathbf{D}(X^a)$ also admits a right adjoint functor $(-)_* : \mathbf{D}(X^a) \to \mathbf{D}(X)$. Moreover, the unit (resp. counit) morphism

$$\mathfrak{F} \to (\mathfrak{F}^a)_* \text{ (resp. } (\mathfrak{G}_*)^a \to \mathfrak{G})$$

is an almost isomorphism (resp. isomorphism) for any $\mathcal{F} \in \mathbf{D}(X), \mathcal{G} \in \mathbf{D}(X^a)$.

Proof. The proof is absolutely similar to Lemma 2.3.7.

Theorem 3.4.9. The functor $(-)^a : \mathbf{D}(X) \to \mathbf{D}(X^a)$ induces an equivalence of triangulated categories $(-)^a : \mathbf{D}(X)^a \to \mathbf{D}(X^a)$.

Proof. The proof is similar to that of Theorem 2.3.8.

Remark 3.4.10. Theorem 3.4.9 shows that the two notions of the derived category of almost modules are the same. In what follows, we do not distinguish $\mathbf{D}(X^a)$ and $\mathbf{D}(X)^a$ anymore.

3.5. Basic Functors on the Derived Categories of \mathcal{O}_X^a -modules. Now we can "derive" certain functors constructed in section 3.2. For the rest of the section we fix a ringed *R*-site (X, \mathcal{O}_X) . The section follows the exposition of section 2.4 very closely.

Definition 3.5.1. We define the *derived Hom* functors

$$\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_X^a}(-,-)\colon \mathbf{D}(X)^{a,op} \times \mathbf{D}(X)^a \to \mathbf{D}(X), \text{ and}$$
$$\mathbf{R}\mathrm{Hom}_{\mathcal{O}_X^a}(-,-)\colon \mathbf{D}(X)^{a,op} \times \mathbf{D}(X)^a \to \mathbf{D}(R)$$

as it is done in [Sta21, Tag 08DH] and [Sta21, Tag 0B6A], respectively.

Definition 3.5.2. We define the *global Ext-modules* as the *R*-modules $\operatorname{Ext}^{i}_{\mathcal{O}^{a}_{X}}(\mathcal{F}^{a}, \mathcal{G}^{a}) \coloneqq \operatorname{H}^{i}(\mathbf{R}\operatorname{Hom}_{\mathcal{O}^{a}_{X}}(\mathcal{F}^{a}, \mathcal{G}^{a}))$ for $\mathcal{F}^{a}, \mathcal{G}^{a} \in \operatorname{\mathbf{Mod}}^{a}_{X}$.

We define the *local Ext-sheaves* as the \mathcal{O}_X -modules $\underline{\mathcal{E}xt}^i_{\mathcal{O}^a_X}(\mathcal{F}^a, \mathcal{G}^a) := \mathcal{H}^i(\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}^a_X}(\mathcal{F}^a, \mathcal{G}^a))$ for $\mathcal{F}^a, \mathcal{G}^a \in \mathbf{Mod}^a_X$.

Remark 3.5.3. We see that [Sta21, Tag 0A64] implies that there is a functorial isomorphism

$$\mathrm{H}^{i}\left(\mathbf{R}\mathrm{Hom}_{\mathbb{O}_{X}^{a}}\left(\mathcal{F}^{a},\mathcal{G}^{a}\right)\right)\simeq\mathrm{Hom}_{\mathbf{D}(R)^{a}}\left(\mathcal{F}^{a},\mathcal{G}^{a}[i]\right)$$

for $\mathcal{F}^a, \mathcal{G}^a \in \mathbf{D}(X)^a$.

Remark 3.5.4. The standard argument shows that there is a functorial isomorphism

$$\mathbf{R}\Gamma(U, \mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_{\mathbf{V}}^{a}}(\mathcal{F}^{a}, \mathcal{G}^{a})) \simeq \mathbf{R}\mathrm{Hom}_{\mathcal{O}_{U}^{a}}(\mathcal{F}^{a}|_{U}, \mathcal{G}^{a}|_{U})$$

for any open $U \in X$, $\mathcal{F}^a, \mathcal{G}^a \in \mathbf{D}(X)^a$.

Now we give show a local version of the $((-)_{!}, (-)^{a})$ -adjunction, and relate $\mathbb{R}\underline{\mathcal{H}om}$ (resp. \mathbb{R} Hom) to the certain derived functor. This goes in complete analogy with the situation in the usual (not almost) world.

Lemma 3.5.5. Let (X, \mathcal{O}_X) be a ringed *R*-site. Then

(1) There is a functorial isomorphism

 $\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_{\mathbf{Y}}^{a}}(\mathcal{F}^{a},\mathcal{G}^{a})\simeq\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_{\mathbf{Y}}}(\mathcal{F}^{a}_{!},\mathcal{G})$

for any $\mathfrak{F}^a \in \mathbf{D}(X)^a$ and $\mathfrak{G} \in \mathbf{D}(X)$. In particular, this isomorphism induces functorial isomorphisms

 $\mathbf{R}\mathrm{Hom}_{\mathcal{O}_X^a}(\mathcal{F}^a, \mathcal{G}^a) \simeq \mathbf{R}\mathrm{Hom}_{\mathcal{O}_X}(\mathcal{F}^a_!, \mathcal{G}) \text{ and } \mathrm{Hom}_{\mathbf{D}(X)^a}(\mathcal{F}^a, \mathcal{G}^a) \simeq \mathbf{R}\mathrm{Hom}_{\mathbf{D}(X)}(\mathcal{F}^a_!, \mathcal{G}) .$

61

- (2) For any chosen $\mathcal{F}^a \in \mathbf{Mod}_X^a$, the functor $\mathbf{R}\mathrm{Hom}_{\mathcal{O}_X^a}(\mathcal{F}^a, -) \colon \mathbf{D}(X)^a \to \mathbf{D}(R)$ is isomorphic to the (right) derived functor of $\mathrm{Hom}_{\mathcal{O}_X^a}(\mathcal{F}^a, -)$.
- (3) For any chosen $\mathcal{F}^a \in \mathbf{Mod}_X^a$, the functor $\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}^a, -) \colon \mathbf{D}(X)^a \to \mathbf{D}(X)$ is isomorphic to the (right) derived functor of $\operatorname{Hom}_{\mathcal{O}_X^a}(\mathcal{F}^a, -)$.

Proof. We prove Part (1). We firstly define the map

$$\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}^a, \mathcal{G}^a) \to \mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_X}(\mathcal{F}^a_!, \mathcal{G}) \ .$$

We choose some representation $\mathcal{F}^{\bullet,a}$ of \mathcal{F}^a and a quasi-isomorphism $\mathcal{G} \xrightarrow{\sim} \mathcal{I}^{\bullet}$ of \mathcal{G} to a K-injective complex \mathcal{I}^{\bullet} . Then we know that $\mathcal{I}^{\bullet,a}$ is a K-injective resolution of \mathcal{G}^a by Lemma 3.4.6. Therefore, the construction of the derived hom says that we have isomorphisms

$$\frac{\mathbf{R}\underline{\mathcal{H}}om_{\mathcal{O}_{X}^{a}}(\mathcal{F}^{a}, \mathcal{G}^{a}) \simeq \underline{\mathcal{H}}om_{\mathcal{O}_{X}^{a}}^{\bullet}(\mathcal{F}^{\bullet,a}, \mathcal{I}^{\bullet,a})}{\mathbf{R}\underline{\mathcal{H}}om_{\mathcal{O}_{X}}(\mathcal{F}^{a}_{!}, \mathcal{G}) \simeq \underline{\mathcal{H}}om_{\mathcal{O}_{X}}^{\bullet}(\mathcal{F}^{\bullet,a}_{!}, \mathcal{I}^{\bullet})}$$

Now we recall that term-wise we have the following equalities:

$$\underline{\mathcal{H}om}^{n}_{\mathcal{O}^{a}_{X}}(\mathcal{F}^{\bullet,a},\mathcal{I}^{\bullet,a}) = \prod_{p+q=n} \underline{\mathcal{H}om}_{\mathcal{O}^{a}_{X}}(\mathcal{F}^{-q,a},\mathcal{I}^{p,a})$$

$$\underline{\mathcal{H}om}^{n}_{\mathcal{O}_{X}}(\mathcal{F}^{\bullet,a}_{!},\mathcal{I}^{\bullet}) = \prod_{p+q=n} \underline{\mathcal{H}om}_{\mathcal{O}_{X}}(\mathcal{F}^{-q,a}_{!},\mathcal{I}^{p})$$

Thus we can apply Lemma 3.2.3 term-wise to produce an isomorphism

$$\kappa_n \colon \underline{\mathcal{H}om}^n_{\mathcal{O}^a_X}(\mathcal{F}^{\bullet,a},\mathcal{I}^{\bullet,a}) \to \underline{\mathcal{H}om}^n_{\mathcal{O}_X}(\mathcal{F}^{\bullet,a}_!,\mathcal{I}^{\bullet})$$

for each n. It is then straightforward to see that κ_n commute with the differential, and thus induce the isomorphism of complexes

$$\kappa \colon \underline{\mathcal{H}om}^{\bullet}_{\mathcal{O}^{a}_{X}}(\mathcal{F}^{\bullet,a},\mathcal{I}^{\bullet,a}) \xrightarrow{\sim} \underline{\mathcal{H}om}^{\bullet}_{\mathcal{O}_{X}}(\mathcal{F}^{\bullet,a}_{!},\mathcal{I}^{\bullet}) \ .$$

In particular, it produces the desired isomorphism $\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}^a, \mathcal{G}^a) \xrightarrow{\sim} \mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}^a, \mathcal{G}^a)$. The construction is clearly functorial in both \mathcal{F}^a and \mathcal{G} .

Parts (2) and (3) are identical to Lemma 2.4.3(2).

Definition 3.5.6. We define the *derived almost Hom* functors

$$\mathbf{R}\underline{al\mathcal{H}om}_{\mathcal{O}_X^a}(-,-)\colon \mathbf{D}(X)^{a,op} \times \mathbf{D}(X)^a \to \mathbf{D}(X)^a$$
$$\mathbf{R}al\mathcal{H}om_{\mathcal{O}_X^a}(-,-)\colon \mathbf{D}(X)^{a,op} \times \mathbf{D}(X)^a \to \mathbf{D}(R)^a$$

as

$$\begin{aligned} \mathbf{R}_{\underline{alHom}_{\mathbb{O}_X^a}}(\mathbb{F}^a, \mathbb{G}^a) &\coloneqq \mathbf{R}_{\underline{\mathcal{H}om}_{\mathbb{O}_X^a}}(\mathbb{F}^a, \mathbb{G}^a)^a = \mathbf{R}_{\underline{\mathcal{H}om}_{\mathbb{O}_X}}(\mathbb{F}^a_!, \mathbb{G})^a \\ \mathbf{R}_{alHom_{\mathbb{O}_X^a}}(\mathbb{F}^a, \mathbb{G}^a) &\coloneqq \mathbf{R}_{Hom_{\mathbb{O}_X}}(\mathbb{F}^a, \mathbb{G}^a)^a = \mathbf{R}_{Hom_{\mathbb{O}_X}}(\mathbb{F}^a_!, \mathbb{G})^a \end{aligned}$$

Definition 3.5.7. We define the global almost Ext modules as the R^a -modules $\operatorname{alExt}^i_{\mathcal{O}^a_X}(\mathcal{F}^a, \mathcal{G}^a) := \operatorname{H}^i(\operatorname{RalHom}_{\mathcal{O}^a_Y}(\mathcal{F}^a, \mathcal{G}^a))$ for $\mathcal{F}^a, \mathcal{G}^a \in \operatorname{\mathbf{Mod}}^a_X$.

We define the *local almost Ext sheaves* as the \mathcal{O}_X^a -modules $\underline{al\mathcal{E}xt}_{\mathcal{O}_X^a}^i(\mathcal{F}^a, \mathcal{G}^a) := \mathcal{H}^i(\mathbf{R}\underline{al\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}^a, \mathcal{G}^a))$ for $\mathcal{F}^a, \mathcal{G}^a \in \mathbf{Mod}_X^a$.

Proposition 3.5.8. Let (X, \mathcal{O}_X) be a ringed *R*-site. Then:

(1) There is a natural transformation of functors

$$\begin{array}{c} \mathbf{D}(X)^{op} \times \mathbf{D}(X) \xrightarrow{\mathbf{R} \underline{\mathcal{H}om}_{\mathcal{O}_{X}}(-,-)} \mathbf{D}(X) \\ & \downarrow^{(-)^{a} \times (-)^{a}} \xrightarrow{\rho} \downarrow^{(-)^{a}} \\ \mathbf{D}(X)^{a,op} \times \mathbf{D}(X)^{a} \xrightarrow{\mathcal{D}(X)^{a}} \mathbf{D}(X)^{a} \end{array}$$

that makes the diagram 2-commutative. In particular, $\mathbf{R}_{\underline{alHom}_{\mathcal{O}_X}}(\mathcal{F}^a, \mathcal{G}^a) \simeq \mathbf{R}_{\underline{\mathcal{H}om}_{\mathcal{O}_X}}(\mathcal{F}, \mathcal{G})^a$ for any $\mathcal{F}, \mathcal{G} \in \mathbf{D}(\mathcal{O}_X)$.

- (2) For any chosen $\mathcal{F}^a \in \mathbf{Mod}_R^a$, the functor $\mathbf{R}_{\underline{al}\mathcal{H}om_{\mathcal{O}_X^a}}(\mathcal{F}^a, -) \colon \mathbf{D}(X)^a \to \mathbf{D}(X)^a$ is isomorphic to the (right) derived functor of $\underline{al}\mathcal{H}om_{\mathcal{O}_X^a}(\mathcal{F}^a, -)$.
- (3) The analogous results hold true for the functor $\operatorname{RalHom}_{\mathcal{O}_X^a}(-,-)$.

Proof. The proof is identical to that of Proposition 2.4.8. One only needs to use Proposition 3.2.10 in place of Proposition 2.2.1(3).

Now we deal with the case of the derived tensor product functor. We will show that our definition of the derived tensor product functor makes $\mathbf{R}_{\underline{alHom}}_{\mathcal{O}_X^a}(-,-)$ into the inner Hom functor.

Definition 3.5.9. We say that a complex of \mathcal{O}_X^a -module $\mathcal{F}^{\bullet,a}$ is *almost K-flat* if the naive tensor product complex $\mathcal{C}^{\bullet,a} \otimes_{\mathcal{O}_X^a}^{\bullet} \mathcal{F}^{\bullet,a}$ is acyclic for any acyclic complex $\mathcal{C}^{\bullet,a}$ of \mathcal{O}_X^a -modules.

Lemma 3.5.10. The functor $(-)^a \colon \mathbf{Comp}(\mathcal{O}_X) \to \mathbf{Comp}(\mathcal{O}_X^a)$ sends K-flat \mathcal{O}_X -complexes to almost K-flat \mathcal{O}_X^a -complexes.

Proof. The proof Lemma 2.4.10 applies verbatim.

Lemma 3.5.11. Let $f: (X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ be a morphism of ringed *R*-sites, and let $\mathcal{F}^{\bullet,a} \in \mathbf{Comp}(\mathcal{O}_Y^a)$ be an almost *K*-flat complex. Then $f^*(\mathcal{F}^{\bullet,a}) \in \mathbf{Comp}(\mathcal{O}_X^a)$ is almost *K*-flat.

Proof. The proof of [Sta21, Tag 06YW] works verbatim in this situation.

Corollary 3.5.12. Every object $\mathcal{F}^{\bullet,a} \in \mathbf{Comp}(\mathcal{O}_X^a)$ is quasi-isomorphic to an almost K-flat complex.

Proof. The proof of Corollary 2.4.11 applies verbatim with the only difference that one needs to use [Sta21, Tag 06YF] in place of [Sta21, Tag 06Y4]. \Box

Definition 3.5.13. We define the *derived tensor product functor*

$$-\otimes^{L}_{\mathbb{O}^{a}_{X}} -: \mathbf{D}(\mathbb{O}_{X})^{a} \times \mathbf{D}(\mathbb{O}_{X})^{a} \to \mathbf{D}(\mathbb{O}_{X})^{a}$$

by the rule $(\mathcal{F}^a, \mathcal{G}^a) \mapsto (\mathcal{G}_! \otimes^L_{\mathcal{O}_X} \mathcal{G}_!)^a$ for any $\mathcal{F}^a, \mathcal{G}^a \in \mathbf{D}(X)^a$.

Proposition 3.5.14. (1) There is a natural transformation of functors

that makes the diagram 2-commutative. In particular, there is a functorial isomorphism $(\mathfrak{F} \otimes_{\mathfrak{O}_X}^L \mathfrak{G})^a \simeq \mathfrak{F}^a \otimes_{\mathfrak{O}_X}^L \mathfrak{G}^a$ for any $\mathfrak{F}, \mathfrak{G} \in \mathbf{D}(X)$.

(2) For any chosen $\mathfrak{F}^a \in \mathbf{Mod}_X^a$, the functor $\mathfrak{F}^a \otimes_{R^a}^L -: \mathbf{D}(X)^a \to \mathbf{D}(X)^a$ is isomorphic to the (left) derived functor of $\mathfrak{F}^a \otimes_{\mathfrak{O}_X^a} -$.

Proof. Again, the proof is identical to that of Proposition 3.5.14. The only non-trivial input that we need is existence of sufficiently many K-flat complexes of \mathcal{O}_X^a -modules. But this is guaranteed by Lemma 3.5.12.

Remark 3.5.15. For any $\mathcal{F}^a, \mathcal{G}^a \in \mathbf{D}(X)^a$, there is a canonical morphism

$$\mathbf{R}\underline{al}\mathcal{H}\underline{om}_{\mathcal{O}_X^a}(\mathcal{F}^a, \mathcal{G}^a) \otimes^L_{\mathcal{O}_X} \mathcal{F}^a \to \mathcal{G}^a$$

that, after the identifications from Proposition 3.5.8 and Proposition 3.5.14, is the almostification of the canonical morphism

$$\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_X}(\mathcal{F}^a_!, \mathcal{G}^a_!) \otimes^L_{\mathcal{O}_X} \mathcal{F}^a_! \to \mathcal{G}^a_!$$

from [Sta21, Tag 0A8V].

Lemma 3.5.16. Let (X, \mathcal{O}_X) be a ringed *R*-site, and let $\mathcal{F}^a, \mathcal{G}^a, \mathcal{H}^a \in \mathbf{D}(\mathcal{O}_X)^a$. Then we have a functorial isomorphism

$$\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}^a \otimes_{\mathcal{O}_X^a}^L \mathcal{G}^a, \mathcal{H}^a) \simeq \mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}^a, \mathbf{R}\underline{al\mathcal{H}om}_{R^a}(\mathcal{G}^a, \mathcal{H}^a)) \ .$$

This induces functorial isomorphisms

$$\begin{aligned} \mathbf{R}\mathrm{Hom}_{\mathcal{O}_{X}^{a}}(\mathcal{F}^{a}\otimes_{\mathcal{O}_{X}^{a}}^{L}\mathcal{G}^{a},\mathcal{H}^{a}) &\simeq \mathbf{R}\mathrm{Hom}_{\mathcal{O}_{X}^{a}}(\mathcal{F}^{a},\mathbf{R}\underline{al\mathcal{H}om}_{R^{a}}(\mathcal{G}^{a},\mathcal{H}^{a})) \ , \\ \mathbf{R}\underline{al\mathcal{H}om}_{\mathcal{O}_{X}^{a}}(\mathcal{F}^{a}\otimes_{\mathcal{O}_{X}^{a}}^{L}\mathcal{G}^{a},\mathcal{H}^{a}) &\simeq \mathbf{R}\underline{al\mathcal{H}om}_{\mathcal{O}_{X}^{a}}(\mathcal{F}^{a},\mathbf{R}\underline{al\mathcal{H}om}_{R^{a}}(\mathcal{G}^{a},\mathcal{H}^{a})) \ , \\ \mathbf{R}\mathrm{alHom}_{\mathcal{O}_{X}^{a}}(\mathcal{F}^{a}\otimes_{\mathcal{O}_{Y}^{a}}^{L}\mathcal{G}^{a},\mathcal{H}^{a}) &\simeq \mathbf{R}\mathrm{alHom}_{\mathcal{O}_{X}^{a}}(\mathcal{F}^{a},\mathbf{R}\underline{al\mathcal{H}om}_{R^{a}}(\mathcal{G}^{a},\mathcal{H}^{a})) \ . \end{aligned}$$

Proof. The proof of the first isomorphism is very similar to that of Lemma 2.4.14. We leave the details to the interested reader. The second isomorphism comes from the fist one by applying the functor $\mathbf{R}\Gamma(X, -)$. The third and the fourth isomorphisms are obtained by applying $(-)^a$ to the first and the second isomorphisms respectively. Here we implicitly use Proposition 3.5.8.

Corollary 3.5.17. Let (X, \mathcal{O}_X) be a ringed *R*-site, and let $\mathcal{G}^a \in \mathbf{D}(X)^a$. Then the functors

$$\mathbf{R}_{\underline{al}\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{G}^a, -) \colon \mathbf{D}(X)^a \xleftarrow{} \mathbf{D}(X)^a \colon - \otimes_{\mathcal{O}_X^a}^L \mathcal{G}^a$$

are adjoint.

The next two functors we deal with are the derived pullback and derived pushforward. We start with the derived pullback.

Definition 3.5.18. Let $f: (X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ be a morphism of ringed *R*-sites. We define the *derived pullback functor*

$$\mathbf{L}f^*: \mathbf{D}(Y)^a \to \mathbf{D}(X)^a$$

as the derived functor of the right exact, additive functor $f^* \colon \mathbf{Mod}_Y^a \to \mathbf{Mod}_X^a$.

Remark 3.5.19. We need to explain why the desired derived functor exists and how it can be computed. It turns out that it can be constructed by choosing K-flat resolutions, the argument for this is identical to [Sta21, Tag 06YY]. We only emphasize that three main inputs are Lemma 3.5.11, Lemma 3.5.10 and an almost analogue of [Sta21, Tag 06YG].

Proposition 3.5.20. Let $f: (X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ be a morphism of ringed *R*-sites. Then there is a natural transformation of functors



that makes the diagram 2-commutative. In particular, there is a functorial isomorphism $(\mathbf{L}f^*\mathcal{F})^a \simeq \mathbf{L}f^*(\mathcal{F}^a)$ for any $\mathcal{F} \in \mathbf{D}(Y)$.

Proof. We construct the natural tranformation $\rho: \mathbf{L}f^* \circ (-)^a \Rightarrow (-)^a \circ \mathbf{L}f^*$ as follows. Pick any object $\mathcal{F} \in \mathbf{D}(Y)$ and its K-flat representative \mathcal{K}^{\bullet} , then \mathcal{K}^{\bullet} is adapted to compute the usual derived pullback $\mathbf{L}f^*$. Lemma 3.5.11 ensures $\mathcal{K}^{\bullet,a}$ is also adapted to compute the almost version of the derived pullback $\mathbf{L}f^*$. So we define the morphism

$$\rho_{\mathcal{F}} \colon (f^*(\widetilde{\mathfrak{m}} \otimes \mathcal{K}^{\bullet}))^a \to f^*(\mathcal{K}^{\bullet})^a$$

as the natural morphism induced by $\widetilde{\mathfrak{m}} \otimes \mathcal{K}^{\bullet} \to \mathcal{K}^{\bullet}$. This map is clearly functorial, so it defines a transformation of functors ρ . In order to show that it is an isomorphism of functors, it suffices to show that the map

$$f^*(\widetilde{\mathfrak{m}}\otimes\mathfrak{K}^{\bullet})\to f^*(\mathfrak{K}^{\bullet})$$

is an almost isomorphism of complexes for any K-flat complex K^{\bullet} . But this is clear as $\widetilde{\mathfrak{m}} \otimes \mathcal{K}^{\bullet} \to \mathcal{K}^{\bullet}$ is an almost isomorphism, and Corollary 3.2.18 ensures that f^* preserves almost isomorphisms. \Box

Definition 3.5.21. Let $f: (X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ be a morphism of ringed *R*-sites. We define the *derived pushforward functor*

$$\mathbf{R}f_*: \mathbf{D}(X)^a \to \mathbf{D}(Y)^a$$

as the derived functor of the left exact, additive functor $f_*: \operatorname{Mod}_X^a \to \operatorname{Mod}_Y^a$.

We define the *derived global sections functor* $\mathbf{R}\Gamma(U, -) \colon \mathbf{D}(X)^a \to \mathbf{D}(R)^a$ in a similar way for any open $U \subset X$.

Remark 3.5.22. This functor exists by abstract nonsense (i.e. [Sta21, Tag 070K]) as the category Mod_X^a has enough *K*-injective complexes by Lemma 3.4.7.

Proposition 3.5.23. Let $f: (X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ be a morphism of ringed *R*-sites. Then there is a natural transformation of functors



that makes the diagram 2-commutative. In particular, there is a functorial isomorphism $(\mathbf{R}f_*\mathcal{F})^a \simeq \mathbf{R}f_*(\mathcal{F}^a)$ for any $\mathcal{F} \in \mathbf{D}(X)$. The analogous results hold for the functor $\mathbf{R}\Gamma(U, -)$.

Proof. The proof is very similar to that of Proposition 3.5.20. The main essential ingredients are: $(-)^a$ sends K-injective complexes to K-injective complexes, and f_* preserves almost isomorphisms. These two results were shown in Lemma 3.4.6 and Lemma 3.2.23.

Lemma 3.5.24. Let (X, \mathcal{O}_X) be a ringed *R*-site, let \mathcal{F} be an \mathcal{O}_X^a -module, and let $U \in X$ be an open object. Then we have a canonical isomorphism

$$\mathbf{R}\Gamma(U, \mathbf{R}\underline{al}\mathcal{H}om_{\mathcal{O}_{\mathbf{V}}^{a}}(\mathcal{F}^{a}, \mathcal{G}^{a})) \simeq \mathbf{R}alHom_{\mathcal{O}_{U}^{a}}(\mathcal{F}^{a}|_{U}, \mathcal{G}^{a}|_{U})$$

Proof. This easily follows from Remark 3.5.4, Proposition 3.5.8, and Proposition 3.5.23.

Lemma 3.5.25. Let $f: (X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ be a locally ringed morphism of *R*-spaces. Then there is a functorial isomorphism

$$\mathbf{R}f_*\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_X^a}(\mathbf{L}f^*\mathcal{F}^a, \mathcal{G}^a) \simeq \mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_Y^a}(\mathcal{F}^a, \mathbf{R}f_*\mathcal{G}^a)$$

for $\mathcal{F}^a \in \mathbf{D}(Y)^a$, $\mathcal{G}^a \in \mathbf{D}(X)^a$. This isomorphism induces isomorphisms

 $\mathbf{R} f_* \mathbf{R} \underline{al \mathcal{H} om}_{\mathcal{O}_X^a} (\mathbf{L} f^* \mathcal{F}^a, \mathcal{G}^a) \simeq \mathbf{R} \underline{al \mathcal{H} om}_{\mathcal{O}_Y^a} (\mathcal{F}^a, \mathbf{R} f_* \mathcal{G}^a) ,$ $\mathbf{R} \mathrm{Hom}_{\mathcal{O}_X^a} (\mathbf{L} f^* \mathcal{F}^a, \mathcal{G}^a) \simeq \mathbf{R} \mathrm{Hom}_{\mathcal{O}_Y^a} (\mathcal{F}^a, \mathbf{R} f_* \mathcal{G}^a) ,$ $\mathbf{R} \mathrm{al Hom}_{\mathcal{O}_X^a} (\mathbf{L} f^* \mathcal{F}^a, \mathcal{G}^a) \simeq \mathbf{R} \mathrm{al Hom}_{\mathcal{O}_V^a} (\mathcal{F}^a, \mathbf{R} f_* \mathcal{G}^a) .$

Proof. It is a standard exercise to show that the first isomorphism implies all other isomorphisms by applying certain functors to it, so we deal only with the first one. The proof of the first one is also quite standard and similar to Lemma 3.2.26, but we spell it out for the reader's convenience. The desired isomorphism comes from a sequence of canonical identifications:

$$\begin{split} \mathbf{R}f_*\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_X^a}(\mathbf{L}f^*(\mathcal{F}^a),\mathcal{G}^a) &\simeq \mathbf{R}f_*\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_X^a}(\mathbf{L}f^*(\mathcal{F})^a,\mathcal{G}^a) & \text{Proposition 3.5.20} \\ &\simeq \mathbf{R}f_*\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_X}(\widetilde{\mathfrak{m}}\otimes\mathbf{L}f^*(\mathcal{F}),\mathcal{G}) & \text{Lemma 3.5.5(1)} \\ &\simeq \mathbf{R}f_*\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_X}(\mathbf{L}f^*(\widetilde{\mathfrak{m}}\otimes\mathcal{F}),\mathcal{G}) & \text{Lemma 3.2.17} \\ &\simeq \mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_Y}(\widetilde{\mathfrak{m}}\otimes\mathcal{F},\mathbf{R}f_*(\mathcal{G})) & \text{Classical} \\ &\simeq \mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_Y^a}(\mathcal{F}^a,\mathbf{R}f_*(\mathcal{G})^a) & \text{Lemma 3.5.5(1)} \\ &\simeq \mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_Y^a}(\mathcal{F}^a,\mathbf{R}f_*(\mathcal{G})^a) & \text{Proposition 3.5.23.} \end{split}$$

Corollary 3.5.26. Let $f: (X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ be a locally ringed morphism of *R*-spaces. Then the functors $\mathbf{R}f_*(-): \mathbf{D}(X)^a \longleftrightarrow \mathbf{D}(Y)^a: \mathbf{L}f^*(-)$ are adjoint.

4. Almost Coherent Sheaves on Schemes and Formal Schemes

4.1. Schemes. The Category of Almost Coherent \mathcal{O}_X^a -modules. In this Section we discuss the notion of almost quasi-coherent, almost finite type, almost finitely presented and almost coherent sheaves on an arbitrary scheme. The main content of this Section is to make sure that almost coherent sheaves form a Weak Serre subcategory in \mathcal{O}_X -modules. Another important statement is the "approximation" Lemma 4.1.8 that is the key fact to reduce many statements about almost finitely presented \mathcal{O}_X -modules to the "classical" case of finitely presented \mathcal{O}_X -modules. In particular, we follow this approach in our proof of the Almost Proper Mapping Theorem in Section 5.1.

As always, we fix a ring R with an ideal \mathfrak{m} such that $\mathfrak{m} = \mathfrak{m}^2$ and $\tilde{\mathfrak{m}} = \mathfrak{m} \otimes_R \mathfrak{m}$ is R-flat. We always do almost mathematics with respect to this ideal. In what follows X will always denote an R-scheme. Note that this implies that X is a locally spectral, ringed R-site, so the results of the previous sections apply.

We begin with some definitions:

Definition 4.1.1. We say that an \mathcal{O}_X^a -module \mathcal{F}^a is almost quasi-coherent if $\mathcal{F}_!^a \simeq \widetilde{\mathfrak{m}} \otimes \mathcal{F}$ is a quasi-coherent \mathcal{O}_X -module.

We say that an \mathcal{O}_X -module \mathcal{F} is almost quasi-coherent if \mathcal{F}^a is an almost quasi-coherent \mathcal{O}^a_X -module.

Remark 4.1.2. We denote by $\mathbf{AQcoh}_X \subset \mathbf{Mod}_X^a$ the full subcategory consisting of almost quasicoherent \mathcal{O}_X^a -modules. It is straightforward¹⁹ to see that the "forgetful" functor induces an equivalence

$$\mathbf{AQcoh}_X \simeq \mathbf{Qcoh}_X / (\Sigma_X \cap \mathbf{Qcoh}_X) ,$$

i.e. \mathbf{AQcoh}_X is equivalent to the quotient category of quasi-coherent \mathcal{O}_X -modules by the full subcategory of almost zero, quasi-coherent \mathcal{O}_X -modules.

Definition 4.1.3. We say that an \mathcal{O}_X^a -module \mathcal{F}^a is of almost finite type (resp. almost finitely presented) if \mathcal{F}^a is almost quasi-coherent, and there is a covering of X by open affines $\{U_i\}_{i \in I}$ such that $\mathcal{F}^a(U_i)$ is an almost finitely generated (resp. almost finitely presented) $\mathcal{O}_X^a(U_i)$ -module.

We say that an \mathcal{O}_X -module \mathcal{F} is of almost finite type (resp. almost finitely presented) if so is \mathcal{F}^a .

Remark 4.1.4. We denote by $\mathbf{Qcoh}_X^{\text{aft}}$ (resp. $\mathbf{Qcoh}_X^{\text{afp}}$) the full subcategory of \mathbf{Qcoh}_X consisting of almost finite type (resp. almost finitely presented) \mathcal{O}_X -modules. Similarly, we denote by $\mathbf{AQcoh}_X^{\text{aft}}$ (resp. $\mathbf{AQcoh}_X^{\text{afp}}$) the full subcategory of \mathbf{AQcoh}_X consisting of almost finite type (resp. almost finitely presented) \mathcal{O}_X^a -modules. Again, it is straightforward to see that the "forgetful" functor induces the equivalences

$$\mathbf{AQcoh}_X^{\mathrm{aft}} \simeq \mathbf{Qcoh}_X^{\mathrm{aft}} / (\Sigma_X \cap \mathbf{Qcoh}_X^{\mathrm{aft}}), \ \mathbf{AQcoh}_X^{\mathrm{afp}} \simeq \mathbf{Qcoh}_X^{\mathrm{afp}} / (\Sigma_X \cap \mathbf{Qcoh}_X^{\mathrm{afp}})$$

Remark 4.1.5. In the usual theory of \mathcal{O}_X -modules, finite type \mathcal{O}_X -modules are usually not required to be quasi-coherent. However, it is much more convenient for our purposes to put almost quasi-coherence in the definition of almost finite type modules.

The first thing that we need to check is that these notions do not depend on a choice of an affine covering.

Lemma 4.1.6. Let \mathcal{F}^a be an almost finite type (resp. almost finitely presented) \mathcal{O}^a_X -module on an *R*-scheme *X*. Then $\mathcal{F}^a(U)$ is an almost finitely generated (resp. almost finitely presented) $\mathcal{O}^a_X(U)$ -module for any open affine $U \subset X$.

Proof. First of all, Corollary 2.5.9 and Lemma 3.3.2 imply that for any open affine U, $\mathcal{F}^{a}(U)$ is almost finitely generated (resp. almost finitely presented) if and only if so is $(\widetilde{\mathfrak{m}} \otimes \mathcal{F}^{a})(U)$. Thus we can replace \mathcal{F}^{a} by $\mathcal{F}^{a}_{1} \simeq \widetilde{\mathfrak{m}} \otimes \mathcal{F}$ to assume that \mathcal{F} is an honest quasi-coherent \mathcal{O}_{X} -module.

Now Lemma 2.7.1 guarantees that the problem is local on X. So we can assume that X = U is an affine scheme and we need to show that $\mathcal{F}(X)$ is almost finitely generated (resp. almost finitely presented).

We pick some covering $X = \bigcup_{i=1}^{n} U_i$ by open affines U_i such that $\mathcal{F}(U_i)$ is almost finitely generated (resp. almost finitely presented) as an $\mathcal{O}_X(U_i)$ -module. We note that since \mathcal{F} is quasi-coherent we have an isomorphism

$$\mathfrak{F}(U_i) \simeq \mathfrak{F}(X) \otimes_{\mathfrak{O}_X(X)} \mathfrak{O}_X(U_i).$$

 $^{^{19}}$ The proof is completely similar to the proof of Theorem 3.1.19 or Theorem 3.4.9.

Now we see that a map $\mathcal{O}_X(X) \to \prod_{i=1}^n \mathcal{O}_X(U_i)$ is faithfully flat, and the module

$$\mathcal{F}(X) \otimes_{\mathcal{O}_X(X)} \left(\prod_{i=1}^n \mathcal{O}_X(U_i)\right) \simeq \left(\prod_{i=1}^n \mathcal{O}_X(U_i)\right) \otimes_{\mathcal{O}_X(X)} \mathcal{F}(X)$$

is almost finitely generated (resp. almost finitely presented) over $\prod_{i=1}^{n} \mathcal{O}_X(U_i)$. Therefore, Lemma 2.7.6 guarantees that $\mathcal{F}(X)$ is almost finitely generated (resp. almost finitely presented) as an $\mathcal{O}_X(X)$ -module.

Corollary 4.1.7. Let $X = \operatorname{Spec} A$ be an affine *R*-scheme, and let \mathcal{F}^a be an almost quasi-coherent \mathcal{O}^a_X -module. Then \mathcal{F}^a is almost finite type (resp. almost finitely presented) if and only if $\Gamma(X, \mathcal{F}^a)$ is almost finitely generated (resp. almost finitely presented) *A*-module.

We want to globalize Lemma 2.5.7 in the case of almost finitely presented \mathcal{O}_X -modules. The definition of almost finitely presented \mathcal{O}_X -modules is local, so it might be difficult to prove some global statements such as the Almost Proper Mapping Theorem without some global understanding of those sheaves. This is the reason we want to get a more global description of almost finitely presented \mathcal{O}_X -modules. That description is achieved in the following lemma:

Lemma 4.1.8. Let X be a quasi-compact and quasi-separated R-scheme, and let \mathcal{F} be an almost quasi-coherent \mathcal{O}_X -module. Then \mathcal{F} is of almost finite type (resp. almost finitely presented) if and only if for any finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$ there is a morphism $f : \mathcal{G} \to \mathcal{F}$ such that \mathcal{G} is a quasi-coherent finite type (resp. finitely presented) \mathcal{O}_X -module, $\mathfrak{m}_0(\ker f) = 0$ and $\mathfrak{m}_0(\operatorname{Coker} f) = 0$.

Proof. The "if" part is easy. We pick a finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$ and a quasi-coherent finite type (resp. finitely presented) \mathcal{O}_X -modules \mathcal{G} and a map $f : \mathcal{G} \to \mathcal{F}$ such that the kernel and cokernel are annihilated by \mathfrak{m}_0 . We note that $\mathcal{G}(U)/\mathcal{F}(U) \subset (\mathcal{G}/\mathcal{F})(U)$, so $\mathfrak{m}_0(\ker f(U)) = 0$ and $\mathfrak{m}_0(\operatorname{Coker} f(U)) = 0$ for any open affine $U \subset X$. Then we use Lemma 2.5.7(2) to see that $\mathcal{F}(U)$ is almost finitely generated (resp. almost finitely presented). Corollary 2.5.9 guarantees that $\widetilde{\mathfrak{m}} \otimes_R \mathcal{F}(U)$ is almost finitely presented as well, so \mathcal{F} is indeed of almost finite type (resp. almost finitely presented).

Now we prove the "only if" part only for almost finitely presented sheaves as the other case is significantly easier. We start the proof by noting that it is harmless to pass from \mathcal{F} to $\widetilde{\mathfrak{m}} \otimes \mathcal{F}$. Indeed, the morphism $\widetilde{\mathfrak{m}} \otimes \mathcal{F} \to \mathcal{F}$ is an almost isomorphism. So if there is a morphism $f: \mathcal{G} \to \widetilde{\mathfrak{m}} \otimes \mathcal{F}$ with \mathcal{G} being quasi-coherent finitely type (resp. finitely presented) \mathcal{O}_X -module such that $\mathfrak{m}_0(\ker f) = 0$ and $\mathfrak{m}_0(\operatorname{Coker} f) = 0$, then the composition $\mathcal{G} \to \mathcal{F}$ satisfies the same conditions for \mathfrak{m}_0^2 . The standard argument (by choosing $\mathfrak{m}_1 \subset \mathfrak{m}$ such that $\mathfrak{m}_0 \subset \mathfrak{m}_1^2$) shows that it is harmless to replace \mathcal{F} with $\widetilde{\mathfrak{m}} \otimes \mathcal{F}$. Thus we may and do assume that \mathcal{F} is quasi-coherent.

We pick an arbitrary finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$ and another finitely generated ideal $\mathfrak{m}_1 \subset \mathfrak{m}$ such that $\mathfrak{m}_0 \subset \mathfrak{m}_1^2$. We try to construct a morphism

 $\mathcal{G} \to \mathcal{F}$

such that \mathcal{G} is a finitely presented \mathcal{O}_X -module, and kernel and cokernel of this morphism are annihilated by \mathfrak{m}_0 .

The first non-trivial input is [EGA I, Corollaire (6.9.12)] that says that $\mathcal{F} \simeq \operatorname{colim}_{i \in I} \mathcal{G}_i$ is a filtered colimit of finitely presented \mathcal{O}_X -modules. We choose some finite covering of X by open affine subschemes U_j that is possible as X is quasi-compact. Then we use [Sta21, Tag 009F] to see that

$$\operatorname{colim}_{i\in I} \mathfrak{G}_i(U_i) \to \mathfrak{F}(U_i)$$

69

is an isomorphism for all j. As each of $\mathcal{F}(U_j)$ is almost finitely generated, we can find a finite Rsubmodule $M \subset \mathcal{F}(U_j)$ containing $\mathfrak{m}_1 \mathcal{F}(U_j)$ for any j. We observe that all the generators of M lie in the image of $\mathcal{G}_{i_j}(U_j)$ for some large $i_j \in I$. Thus the cokernel of the natural map $\mathcal{G}_{i_j}(U_j) \to \mathcal{F}(U_j)$ is annihilated by \mathfrak{m}_1 . This implies that the cokernel of the homomorphism $\mathcal{G}_{i_j}|_{U_j} \to \mathcal{F}|_{U_j}$ is killed by \mathfrak{m}_1 as both are quasi-coherent sheaves on an affine scheme. Finally, we use that I is a filtered system to find some $i \in I$ that dominates all i_j . We rename \mathcal{G}_i as \mathcal{G} and define $f: \mathcal{G} \to \mathcal{F}$ as the morphism coming from the filtered system. Note that \mathcal{G} is a finitely presented \mathcal{O}_X -module by definition and Coker f is annihilated by \mathfrak{m}_1 . The problem is that ker f might still be pretty huge (not killed by \mathfrak{m}_1 or any of its powers). We need to fix this problem.

We consider a quasi-coherent \mathcal{O}_X -module $\mathcal{K} := \ker f$. We want to find some quasi-coherent finite type \mathcal{O}_X -submodule $\mathcal{K}' \subset \mathcal{K}$ such that \mathcal{K}/\mathcal{K}' is annihilated by \mathfrak{m}_1^2 . We choose some finite covering of X by open affines U_i and denote a ring $\mathcal{O}_X(U_i)$ by A_i . Note that for any such U_i we have an exact sequence

$$0 \to \mathcal{K}(U_i) \to \mathcal{G}(U_i) \to \mathcal{F}(U_i)$$

with $\mathcal{G}(U_i)$ a finitely presented A_i -module and $\mathcal{F}(U_i)$ an almost finitely presented A_i -module. So choose an integer n and a surjective homomorphism $\pi \colon A_i^n \to \mathcal{G}(U_i)$. We define a homomorphism $g \colon A_i^n \to \mathcal{F}(U_i)$ as a composition $\pi \circ f$. Now comes the key part, as $\mathcal{F}(U_i)$ is almost finitely presented we use Lemma 2.5.6 to find a three-term complex

$$A_i^m \xrightarrow{g'} A_i^n \xrightarrow{g} \mathcal{F}(U_i)$$

such that $\mathfrak{m}_1^2(\ker g) \subset \operatorname{Im} g'$. Note that a natural morphism $A_i^m \to \mathcal{F}(U_i)$ factors through an inclusion $\mathcal{K}(U_i) \hookrightarrow \mathcal{G}(U_i)$ as $f \circ \pi \circ g' = g \circ g' = 0$. Thus we have a commutative diagram



We also note that $\pi(\ker g) = \mathcal{K}(U_i)$ as $g = f \circ \pi$ and π is surjective. Therefore, the condition $\mathfrak{m}_1^2(\ker g) \subset \operatorname{Im} g'$ implies that

$$\mathfrak{m}_1^2(\mathfrak{K}(U_i)) \subset \pi'(A_i^m).$$

Let us denote the submodule $\pi'(A_i^m) \subset \mathcal{K}(U_i)$ by N_i ; this is a finitely generated A_i -module since it is a quotient of finite free A_i -module. This submodule defines a quasi-coherent, finite type \mathcal{O}_{U_i} -submodule $\widetilde{N_i} \subset \mathcal{K}|_{U_i}$ such that $(\mathcal{K}|_{U_i})/\widetilde{N_i}$ is annihilated by \mathfrak{m}_1^2 .

So what we finally come up with is a set of quasi-coherent, finite type \mathcal{O}_{U_i} -submodules $\widetilde{N_i} \subset \mathcal{K}|_{U_i}$ such that $(\mathcal{K}|_{U_i})/\widetilde{N_i}$ is annihilated by \mathfrak{m}_1^2 for any *i*. We use [EGA I, Lemme (6.9.6)] to extend all \mathcal{O}_{U_i} -modules $\widetilde{N_i}$ to some quasi-coherent, finite type \mathcal{O}_X -submodule $\mathcal{K}'_i \subset \mathcal{K}$ such that $\mathcal{K}'_i|_{U_i} = \widetilde{N_i}$. We finally take the product of all \mathcal{K}'_i in \mathcal{K} , which is a quasi-coherent \mathcal{O}_X -module of finite type since this a finite product. We denote this product by \mathcal{K}' , and we see that $\widetilde{N_i} \subset \mathcal{K}'|_{U_i}$. Therefore the \mathcal{O}_X -module \mathcal{K}/\mathcal{K}' is annihilated by \mathfrak{m}_1^2 .

Finally, we define the quasi-coherent \mathcal{O}_X -module $\mathcal{G}' \coloneqq \mathcal{G}/\mathcal{K}'$; it is finitely presented since it is a quotient of finitely presented sheaf by a quasi-coherent, finite type sheaf. Since $\mathcal{K}' \subset \ker f$, the morphism f descends to a morphism

$$f' \colon \mathfrak{G}' \to \mathfrak{F}.$$

It is easily seen that $\mathfrak{m}_1(\operatorname{Coker} f') = 0$ and we also know that $\mathfrak{m}_1^2(\ker f') = \mathfrak{m}_1^2(\mathcal{K}/\mathcal{K}') = 0$ since $\mathfrak{m}_1^2(\mathcal{K}) \subset \mathcal{K}'$. Recall that we have chosen \mathfrak{m}_1 such that $\mathfrak{m}_0 \subset \mathfrak{m}_1^2$. So what we have finally constructed

is a morphism

$$f': \mathfrak{G}' \to \mathfrak{F}$$

such that \mathcal{G}' is a finitely presented \mathcal{O}_X -module, $\mathfrak{m}_0(\ker f') = 0$ and $\mathfrak{m}_0(\operatorname{Coker} f') = 0$.

Now we check that almost finite type and almost finitely presented \mathcal{O}_X^a behave nicely in short exact sequences.

Lemma 4.1.9. Let $0 \to \mathcal{F}'^a \xrightarrow{\varphi} \mathcal{F}^a \xrightarrow{\psi} \mathcal{F}''^a \to 0$ be an exact sequence of \mathcal{O}^a_X -modules. Then

- (1) If \mathcal{F}^a is almost of finite type and \mathcal{F}''^a is almost quasi-coherent, then \mathcal{F}''^a is almost finite type.
- (2) If $\mathcal{F}^{\prime a}$ and $\mathcal{F}^{\prime\prime a}$ are of almost finite type (resp. finitely presented), then so is \mathcal{F}^{a} .
- (3) If \$\mathcal{F}^a\$ is of almost finite type and \$\mathcal{F}''^a\$ is almost finitely presented, then \$\mathcal{F}'^a\$ is of almost finite type.
- (4) If \$\mathcal{F}^a\$ is almost finitely presented and \$\mathcal{F}'^a\$ is of almost finite type, then \$\mathcal{F}''^a\$ is almost finitely presented.

Proof. First of all, we apply the exact functor $(-)_!$ to all \mathcal{O}_X^a -modules in question to assume the short sequence is an exact sequence of \mathcal{O}_X -modules and all \mathcal{O}_X -modules in this sequence are quasicoherent. Note that we implicitly use here that quasi-coherent modules form a Serre subcategory of all \mathcal{O}_X -modules by [Sta21, Tag 01IE]. Then we check the statement on the level of global sections on all open affine subschemes $U \subset X$ using that quasi-coherent sheaves have vanishing higher cohomology on affine schemes. And that is done in Lemma 2.5.11.

Definition 4.1.10. We say that an \mathcal{O}_X^a -module \mathcal{F}^a is almost coherent if \mathcal{F}^a is almost finite type, and for any open set U any almost finite type \mathcal{O}_U^a -submodule $\mathcal{G}^a \subset (\mathcal{F}^a|_U)$ is an almost finitely presented \mathcal{O}_U^a -module.

We say that an \mathcal{O}_X -module \mathcal{F} is almost coherent if \mathcal{F}^a is an almost coherent \mathcal{O}^a_X -module.

Lemma 4.1.11. Let X be an R-scheme, and let \mathcal{F}^a be an \mathcal{O}^a_X -module. Then the following are equivalent:

- (1) \mathcal{F}^a is almost coherent.
- (2) \mathcal{F}^a is almost quasi-coherent, and the $\mathcal{O}^a_X(U)$ -module $\mathcal{F}^a(U)$ is almost coherent for any open affine subscheme $U \subset X$.
- (3) \mathcal{F}^a is almost quasi-coherent, and there is a covering of X by open affine subschemes $(U_i)_{i \in I}$ such that $\mathcal{F}^a(U_i)$ is almost coherent for each *i*.

In particular, if $X = \operatorname{Spec} A$ is an affine *R*-scheme and \mathcal{F}^a is an almost quasi-coherent \mathcal{O}^a_X -module. Then \mathcal{F}^a is almost coherent if and only if $\mathcal{F}^a(X)$ is almost coherent as an *A*-module.

Proof. We start the proof by noting that we can replace \mathcal{F}^a by $\mathcal{F}^a_!$ to assume that \mathcal{F} is a quasi-coherent \mathcal{O}_X -module.

Firstly, we check that (1) implies (2). Given any affine open $U \subset X$ and any almost finitely generated almost submodule $M^a \subset \mathcal{F}(U)^a$, we define an almost subsheaf $(M^a)_! \subset (\mathcal{F}|_U)^a$. We see that $(M^a)_!$ must be an almost finitely presented \mathcal{O}_U -module, so Lemma 4.1.6 guarantees that $M^a_!$ is almost finitely presented $\mathcal{O}_X(U)$ -module. Therefore, any almost finitely generated submodule $M^a \subset \mathcal{F}(U)^a$ is almost finitely presented. This shows that $\mathcal{F}(U)$ is almost coherent.

Now we show that (2) implies (1). Suppose that \mathcal{F} is almost quasi-coherent and $\mathcal{F}(U)$ is almost coherent for any open affine $U \subset X$. First of all, it shows that \mathcal{F} is of almost finite type, since this

notion is local by definition. Now suppose that we have an almost finite type almost \mathcal{O}_X -submodule $\mathcal{G} \subset (\mathcal{F}|_U)^a$ for some open U. It is represented by a homomorphism

$$\widetilde{\mathfrak{m}} \otimes \mathfrak{G} \xrightarrow{g} \mathfrak{F}$$

with \mathcal{G} being an \mathcal{O}_X -module of almost finite type, and $\widetilde{\mathfrak{m}} \otimes \ker g \simeq 0$. We want to show that \mathcal{G} is almost finitely presented as \mathcal{O}_X -module. This is a local question, so we can assume that U is affine. Then Lemma 3.3.2 implies that the natural morphism

$$g(U): \widetilde{\mathfrak{m}} \otimes_R \mathfrak{G}(U) \to \mathfrak{F}(U)$$

defines an almost submodule of $\mathcal{F}(U)$. We conclude that $\widetilde{\mathfrak{m}} \otimes_R \mathcal{G}(U)$ is almost finitely presented by the assumption on $\mathcal{F}(U)$. Since the notion of almost finitely presented \mathcal{O}_X -module is local, we see that \mathcal{G} is almost finitely presented.

Clearly, (2) implies (3). And it is easy to see that Lemma 2.7.7 guarantees that (3) implies (2). \Box

Corollary 4.1.12. Let X be an R-scheme, then:

- (1) Any almost finite type \mathcal{O}_X^a -submodule of an almost coherent \mathcal{O}_X^a -module is almost coherent.
- (2) Let $\varphi \colon \mathcal{F}^a \to \mathcal{G}^a$ be a homomorphism from an almost finite type \mathcal{O}^a_X -module to an almost coherent \mathcal{O}^a_X -module, then ker (φ) is an almost finite type \mathcal{O}^a_X -module.
- (3) Let $\varphi \colon \mathcal{F}^a \to \mathcal{G}^a$ be a homomorphism of almost coherent \mathcal{O}^a_X -modules, then ker(φ) and Coker(φ) are almost coherent \mathcal{O}^a_X -modules.
- (4) Given a short exact sequence of \mathcal{O}_X^a -modules

$$0 \to \mathcal{F}'^a \to \mathcal{F}^a \to \mathcal{F}''^a \to 0$$

if two out of three are almost coherent so is the third.

Proof. The proofs (1), (2) and (3) are quite straightforward. As usually, we apply $(-)_!$ to assume that all sheaves in the question are quasi-coherent \mathcal{O}_X -modules. Then the question is local and it is sufficient to work on global sections over all affine open subschemes $U \subset X$. So the problem is reduced to Lemma 2.6.7.

The proof of part (4) is similar, but we only need to invoke that given a short exact sequence of \mathcal{O}_X -modules

$$0 \to \mathcal{F}_{!}^{\prime a} \to \mathcal{F}_{!}^{a} \to \mathcal{F}_{!}^{\prime \prime a} \to 0$$

if two of these sheaves are quasi-coherent, so is the third one. This is proven in the affine case in [Sta21, Tag 01IE], the general case reduces to the affine one. The rest of the argument is the same. \Box

Definition 4.1.13. We define the categories \mathbf{Qcoh}_X^{acoh} and \mathbf{AQcoh}_X^{acoh} as the full subcategories of \mathbf{Qcoh}_X and \mathbf{AQcoh}_X , respectively, consisting of almost coherent modules. As always, it is straightforward to see that the "forgetful" functor induces the equivalence

$$\mathbf{AQcoh}_X^{acoh} \simeq \mathbf{Qcoh}_X^{acoh} / (\Sigma_X \cap \mathbf{Qcoh}_X^{acoh})$$

Moreover, Corollary 4.1.12 ensures that $\mathbf{Qcoh}_X^{acoh} \subset \mathbf{Qcoh}_X$ and $\mathbf{AQcoh}_X^{acoh} \subset \mathbf{AQcoh}_X$ are weak Serre subcategories.

Definition 4.1.14. We define the category Acoh_X as the full subcategories of Mod_X consisting of almost coherent \mathcal{O}_X -modules. Corollary 4.1.12 ensures that $\operatorname{Qcoh}_X^{acoh} \subset \operatorname{Qcoh}_X$ and $\operatorname{AQcoh}_X^{acoh} \subset \operatorname{AQcoh}_X$ are Weak Serre subcategories.

The last thing that we discuss here is the notion of almost coherent schemes.

Definition 4.1.15. We say that an *R*-scheme X is almost coherent if the sheaf \mathcal{O}_X is an almost coherent \mathcal{O}_X -module.

Lemma 4.1.16. Let X be a coherent R-scheme. Then X is also almost coherent.

Proof. The structure sheaf \mathcal{O}_X is quasi-coherent by definition. Lemma 4.1.11 says that it suffices to show that $\mathcal{O}_X(U)$ is an almost coherent $\mathcal{O}_X(U)$ -module for any open affine $U \subset X$. Since X is coherent, we conclude that $\mathcal{O}_X(U)$ is coherent as an $\mathcal{O}_X(U)$ -module. Then Lemma 2.6.6 implies that it is actually almost coherent.

Lemma 4.1.17. Let X be an almost coherent R-scheme. Then an \mathcal{O}_X^a -module \mathcal{F}^a is almost coherent if and only if it is of almost finite presentation.

Proof. The "only if" part is easy since any almost coherent \mathcal{O}_X^a -module is of almost finite presentation by the definition. The "if" part is a local question, so we can assume that X is affine, then the claim follows from Lemma 2.6.12.

4.2. Schemes. Basic Functors on Almost Coherent \mathcal{O}_X^a -modules. This section is devoted to study how certain functors defined in Section 3.2 interact with the notions of almost (quasi-) coherent \mathcal{O}_X^a -modules defined in the previous section.

We start with the affine situation, i.e. $X = \operatorname{Spec} A$. In this case, we note that the functor $(-): \operatorname{Mod}_A \to \operatorname{Qcoh}_X$ sends almost zero A-modules to almost zero \mathcal{O}_X -modules. Thus it induces the functor

$$(-): \mathbf{Mod}_A^a \to \mathbf{AQcoh}_X$$
.

Lemma 4.2.1. Let $X = \operatorname{Spec} A$ be an affine R-scheme. Then the functor (-): $\operatorname{Mod}_A \to \operatorname{Qcoh}_X$ induces equivalences (-): $\operatorname{Mod}_A^* \to \operatorname{Qcoh}_X^*$ for any $* \in \{\operatorname{aft}, \operatorname{afp}, \operatorname{acoh}\}$. The quasi-inverse functor is given by $\Gamma(X, -)$.

Proof. We note that the functor (-): $\mathbf{Mod}_A \to \mathbf{Qcoh}_X$ is an equivalence with the quasi-inverse $\Gamma(X, -)$. Now we note Lemma 4.1.7 and Lemma 4.1.11 guarantee that a quasi-coherent \mathcal{O}_X -module \mathcal{F} is almost finite type (resp. almost finitely presented, resp. almost coherent) if $\Gamma(X, \mathcal{F})$ is almost finitely generated (resp. almost finitely presented, resp. almost coherent) as an A-module. \Box

Lemma 4.2.2. Let $X = \operatorname{Spec} A$ be an affine R-scheme. Then the functors (-): $\operatorname{Mod}_A^a \to \operatorname{AQcoh}_X$ induces equivalences $(-)^a$: $\operatorname{Mod}_A^{a,*} \to \operatorname{AQcoh}_X^*$ for any $* \in \{$ "", aft, afp, acoh $\}$. The quasi-inverse functor is given by $\Gamma(X, -)$.

Proof. We note that (-): $\mathbf{Mod}_A \to \mathbf{Qcoh}_X$ induces an equivalence between almost zero A-modules and almost zero, quasi-coherent \mathcal{O}_X -modules. Thus the claim follows from Lemma 4.2.1, Remark 4.1.2, Remark 4.1.4, Definition 4.1.13 and the analogous presentations of $\mathbf{Mod}_A^{a,*}$ as quotients of $\mathbf{Mod}_A^{a,*}$ for any $* \in \{aft, afp, acoh\}$.

Now we show that the pullback functor preserves almost finite type and almost finitely presented \mathcal{O}_X^a -modules.

Lemma 4.2.3. Let $f: X \to Y$ be a morphism of *R*-scheme.

(1) Suppose that $X = \operatorname{Spec} B$, $Y = \operatorname{Spec} A$ are affine *R*-schemes. Then $f^*(\overline{M^a})$ is functorially isomorphic to $\widetilde{M^a \otimes_{A^a} B^a}$ for any $M^a \in \operatorname{\mathbf{Mod}}_A^a$.
- (2) The functor f^* preserves almost quasi-coherence (resp. almost finite type, resp. almost finitely presented) for O-modules.
- (3) The functor f^* preserves almost quasi-coherence (resp. almost finite type, resp. almost finitely presented) for \mathcal{O}^a -modules.

Proof. (1) follows from Proposition 3.2.19 and the analogous result for quasi-coherent \mathcal{O}_Y -modules. More precisely, Proposition 3.2.19 provides with the functorial isomorphism

$$f^*\left(\widetilde{M^a}\right) \xrightarrow{\sim} \left(f^*(\widetilde{M})\right)^a \simeq \left(\widetilde{M\otimes_A B}\right)^a \simeq M^{\widetilde{a}} \otimes_{A^a} B^a$$

(2) and (3) are local on X and Y, so we may and do assume that $X = \operatorname{Spec} B$, $Y = \operatorname{Spec} A$ are affine *R*-schemes. In this case, Lemma 4.2.2 guarantees that any almost quasi-coherent \mathcal{O}_X^a -module is isomorphic to $\widetilde{M^a}$ for some A^a -module M^a . Now (1) ensures that $f^*(\widetilde{M^a}) \simeq M^a \otimes_{A^a} B^a$ as almost quasi-coherent \mathcal{O}_X^a -modules. The other claims from (2) and (3) are proven similarly using Lemma 4.2.2 and Lemma 2.7.1.

The next thing we discuss is how the finiteness properties interact with tensor products.

Lemma 4.2.4. Let X be an R-scheme.

- (1) Suppose that $X = \operatorname{Spec} A$ is an affine *R*-scheme. Then $\widetilde{M^a} \otimes_{\mathcal{O}_X^a} \widetilde{N^a}$ is functorially isomorphic to $\widetilde{M^a} \otimes_{A^a} N^a$ for any $M^a, N^a \in \operatorname{\mathbf{Mod}}_A^a$.
- (2) Let $\mathcal{F}^a, \mathcal{G}^a$ be two almost finite type (resp. almost finitely presented) \mathcal{O}^a_X -modules. Then the \mathcal{O}^a_X -module $\mathcal{F}^a \otimes_{\mathcal{O}^a_X} \mathcal{G}^a$ is almost finite type (resp. almost finitely presented). The analogous result holds for \mathcal{O}_X -modules \mathcal{F}, \mathcal{G} .
- (3) Let \mathcal{F}^a be an almost coherent \mathcal{O}^a_X -module, and \mathcal{G}^a be an almost finitely presented \mathcal{O}^a_X -module. Then $\mathcal{F}^a \otimes_{\mathcal{O}^a_X} \mathcal{G}^a$ is an almost coherent \mathcal{O}^a_X -module. The analogous result holds for \mathcal{O}_X -modules \mathcal{F}, \mathcal{G} .

Proof. The proof is similar to the proof of Lemma 4.2.3. The only difference is that one needs to use Proposition 3.2.12 in place of Proposition 3.2.19 to prove Part (1). Part (2) follows from Lemma 2.5.13, and Part (3) follows from Corollary 2.6.8.

We show f_* preserves almost quasi-coherence of \mathcal{O}^a -modules for a quasi-compact and quasiseparated morphism f. Later on, we will be able to show that f_* also preserves almost coherence of \mathcal{O}^a -modules for certain proper morphisms.

Lemma 4.2.5. Let $f: X \to Y$ be a quasi-compact and quasi-separated morphism of *R*-schemes. Then

- (1) The \mathcal{O}_Y -module $f_*(\mathcal{F})$ is almost quasi-coherent for any almost quasi-coherent \mathcal{O}_X -module \mathcal{F} .
- (2) The \mathcal{O}_Y^a -module $f_*(\mathcal{F}^a)$ is almost quasi-coherent for any almost quasi-coherent \mathcal{O}_X^a -module \mathcal{F}^a .

Proof. Since \mathcal{F} is almost quasi-coherent, we conclude that $\widetilde{\mathfrak{m}} \otimes \mathcal{F}$ is a quasi-coherent \mathcal{O}_X -module. Thus $f_*(\widetilde{\mathfrak{m}} \otimes \mathcal{F})$ is a quasi-coherent \mathcal{O}_Y -module by [Sta21, Tag 01LC]. Recall that the projection formula (Lemma 3.3.5) ensures that

$$f_*(\widetilde{\mathfrak{m}}\otimes \mathfrak{F})\simeq \widetilde{\mathfrak{m}}\otimes f_*\mathfrak{F}$$
.

Thus, we see that $\widetilde{\mathfrak{m}} \otimes f_* \mathcal{F} \simeq f_*(\mathcal{F}^a)_!$ is a quasi-coherent \mathcal{O}_Y -module. This shows that both $f_*(\mathcal{F})$ and $f_*(\mathcal{F}^a)$ are almost quasi-coherent over \mathcal{O}_Y and \mathcal{O}_Y^a respectively. This finishes the proof of the both parts.

Finally, we deal with the $\underline{\mathcal{H}om}_{\mathcal{O}_X^a}(-,-)$ functor. This is probably the most subtle functor considered in this section. We start with the following preparatory lemma:

Lemma 4.2.6. Let X be an R-scheme.

(1) Suppose $X = \operatorname{Spec} A$ is an affine *R*-scheme. Then the canonical map

$$\operatorname{Hom}_{A}(M,N) \to \underline{\operatorname{Hom}}_{\mathcal{O}_{X}}(\widetilde{M},\widetilde{N})$$

$$(4.1)$$

is an almost isomorphism of \mathcal{O}_X -modules for any almost finitely presented A-module M and any A-module N.

(2) Suppose $X = \operatorname{Spec} A$ is an affine *R*-scheme. Then there is a functorial isomorphism

$$\operatorname{alHom}_{A^{a}}(M^{a}, N^{a}) \simeq \underline{al\mathcal{H}om}_{\mathcal{O}_{X}^{a}}(\widetilde{M^{a}}, \widetilde{N^{a}})$$

$$(4.2)$$

of \mathcal{O}_X^a -modules for any almost finitely presented A^a -module M^a , and any A^a -module N^a . We also get a functorial almost isomorphism

$$\operatorname{Hom}_{A}(M,N) \simeq^{a} \underline{\operatorname{Hom}}_{\mathcal{O}_{X}^{a}}(\widetilde{M^{a}},\widetilde{N^{a}})$$

$$(4.3)$$

of \mathcal{O}_X -modules for any almost finitely presented A-module M, and any A-module N.

- (3) Suppose \$\mathcal{F}\$ is an almost finitely presented \$\mathcal{O}_X\$-module and \$\mathcal{G}\$ an almost quasi-coherent \$\mathcal{O}_X\$-module, then \$\frac{\mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G})\$ is an almost quasi-coherent \$\mathcal{O}_X\$-module.
- (4) Suppose \mathcal{F}^a is an almost finitely presented \mathcal{O}^a_X -module and \mathcal{G}^a an almost quasi-coherent \mathcal{O}^a_X -module, then $\underline{\mathcal{H}om}_{\mathcal{O}^a_X}(\mathcal{F}^a, \mathcal{G}^a)$ (resp. $\underline{al\mathcal{H}om}_{\mathcal{O}^a_X}(\mathcal{F}^a, \mathcal{G}^a)$) is an almost quasi-coherent \mathcal{O}_X -module (resp. \mathcal{O}^a_X -module).

Proof. (1): We note that we have a canonical isomorphism $\operatorname{Hom}_A(M, N) \to \operatorname{Hom}_{\mathcal{O}_X}(M, N)$ for any A-modules M, N. This induces a morphism

$$\operatorname{Hom}_{A}(\widetilde{M}, N) \to \underline{\mathcal{H}om}_{\mathcal{O}_{X}}(\widetilde{M}, \widetilde{N})$$
.

In order to show that it is an almost isomorphism for an almost finitely presented M, it suffices to show that the natural map

$$\operatorname{Hom}_{A}(M, N) \otimes_{A} A_{f} \to \operatorname{Hom}_{A_{f}}(M \otimes_{A} A_{f}, N \otimes_{A} A_{f})$$

is an almost isomorphism for any $f \in A$. This follows from Lemma 2.7.8.

(2) follows easily from (1). Indeed, we apply the functorial isomorphism

$$\underline{\mathcal{H}om}_{\mathcal{O}_X}(\mathcal{F},\mathcal{G})^a \simeq \underline{al\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}^a,\mathcal{G}^a)$$

from Proposition 3.2.10(2) to the almost isomorphism in Part (1) to get the functorial isomorphism

$$\operatorname{Hom}_{A}^{\widetilde{}}(\widetilde{M},N)^{a} \simeq \underline{al} \mathfrak{Hom}_{\mathbb{O}_{X}^{a}}^{}(\widetilde{M^{a}},\widetilde{N^{a}}) \ .$$

Now we use Proposition 2.2.1(3) to get the functorial isomorphism

alHom_{A^a} $(M^a, N^a) \simeq \text{Hom}_A(M, N)^a$.

Applying the functor (-) to it and composing with the isomorphism above, we get the functorial isomorphism

$$\mathrm{alHom}_{A^a}^{\widetilde{}}(M^a,N^a)\simeq\underline{al\mathcal{H}om}_{\mathcal{O}^a_X}(\widetilde{M^a},\widetilde{N^a})\ .$$

The construction of the isomorphism (4.3) is similar and even easier.

(3) is a local question, so we can assume that $X = \operatorname{Spec} A$. We note that

$$\underline{\mathcal{H}om}_{\mathcal{O}_X}(\mathcal{F},\mathcal{G}) \simeq^a \underline{\mathcal{H}om}_{\mathcal{O}_X}(\widetilde{\mathfrak{m}} \otimes \mathcal{F}, \widetilde{\mathfrak{m}} \otimes \mathcal{G})$$

by Proposition 3.2.10(2). Thus, we can assume that both \mathcal{F} and \mathcal{G} are quasi-coherent. Then the claim follows from (1) and Lemma 4.2.1.

(4) is similarly just a consequence of (2) and Lemma 4.2.2.

Corollary 4.2.7. Let X be an R-scheme.

- (1) Let \$\mathcal{F}\$ be an almost finitely presented \$\mathcal{O}_X\$-module, and let \$\mathcal{G}\$ be an almost coherent \$\mathcal{O}_X\$-module.
 Then \$\frac{\mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G})\$ is an almost coherent \$\mathcal{O}_X\$-module.
- (2) Let \mathcal{F}^a be an almost finitely presented \mathcal{O}^a_X -module, and let \mathcal{G}^a be an almost coherent \mathcal{O}^a_X -module. Then $\underline{\mathcal{H}om}_{\mathcal{O}^a_X}(\mathcal{F}^a, \mathcal{G}^a)$ (resp. $\underline{al\mathcal{H}om}_{\mathcal{O}^a_X}(\mathcal{F}^a, \mathcal{G}^a)$) is an almost coherent \mathcal{O}_X -module (resp. \mathcal{O}^a_X -module).

Proof. We start the proof by observing that $\underline{\mathcal{H}om}_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}) \simeq^a \underline{\mathcal{H}om}_{\mathcal{O}_X}(\tilde{\mathfrak{m}} \otimes \mathcal{F}, \tilde{\mathfrak{m}} \otimes \mathcal{G})$ by Proposition 3.2.10(2). Thus we can assume that both \mathcal{F} and \mathcal{G} are actually quasi-coherent. In that case we use Lemma 4.2.6(1) and Lemma 4.1.11 to reduce the question to showing that $\mathcal{H}om_A(M, N)$ is an almost coherent for any almost finitely presented M and almost coherent N. However, this has already been done in Corollary 2.6.8.

Part (2) follows from Part (1) as $\underline{\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}^a, \mathcal{G}^a) \simeq \underline{\mathcal{H}om}_{\mathcal{O}_X}(\mathcal{F}^a_!, \mathcal{G})$ and $\underline{al\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}^a, \mathcal{G}^a) \simeq \underline{\mathcal{H}om}_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G})^a$.

4.3. Schemes. Derived Category of Almost Coherent \mathcal{O}_X^a -modules. The goal of this section is to define different categories that can be called "derived category of almost coherent shaves". Namely, we define the categories $\mathbf{D}_{acoh}(X)$, $\mathbf{D}_{qc,acoh}(X)$, and $\mathbf{D}_{acoh}(X)^a$. Then we show that many derived functors of interest preserve almost coherence in an appropriate sense.

Definition 4.3.1. We define $\mathbf{D}_{aqc}(X)$ (resp. $\mathbf{D}_{aqc}(X)^a$) to be the full triangulated subcategory of $\mathbf{D}(X)$ (resp. $\mathbf{D}(X)^a$) consisting of complexes with almost quasi-coherent cohomology sheaves.

Definition 4.3.2. We define $\mathbf{D}_{acoh}(X)$ (resp. $\mathbf{D}_{qc,acoh}(X)$, resp. $\mathbf{D}_{acoh}(X)^a$) to be the full triangulated subcategory of $\mathbf{D}(X)$ (resp. $\mathbf{D}(X)$, resp. $\mathbf{D}(X)^a$) consisting of complexes with almost coherent (resp. quasi-coherent and almost coherent, resp. almost coherent) cohomology sheaves.

Remark 4.3.3. The definition above makes sense as the categories Acoh_X , $\operatorname{Qcoh}_X^{\operatorname{acoh}}$, and $\operatorname{AQcoh}_X^{\operatorname{acoh}}$ are weak Serre subcategories of Mod_X , Mod_X , and Mod_X^a respectively.

Now suppose that $X = \operatorname{Spec} A$ is an affine *R*-scheme. Then we note that the functor

$$(-): \mathbf{Mod}_A \to \mathbf{Mod}_X$$

is additive and exact, thus can be easily derived to the functor

$$(-): \mathbf{D}(A) \to \mathbf{D}_{qc}(X)$$
.

Lemma 4.3.4. Let $X = \operatorname{Spec} A$ be an affine *R*-scheme. Then the functor

$$(-): \mathbf{D}(A) \to \mathbf{D}_{qc}(X)$$

is a *t*-exact equivalence of triangulated categories²⁰ with quasi-inverse given by $\mathbf{R}\Gamma(X, -)$. Moreover, these two functors induce the equivalence

$$(-): \mathbf{D}^*_{acoh}(A) \longleftrightarrow \mathbf{D}^*_{qc,acoh}(X): \mathbf{R}\Gamma(X, -)$$

for any $* \in \{", ", +, -, b\}$.

Proof. The first part is just [Sta21, Tag 06Z0]. In particular, it shows that $\mathrm{H}^{i}(\mathbf{R}\Gamma(X, \mathcal{F})) \simeq \mathrm{H}^{0}(X, \mathcal{H}^{i}(\mathcal{F}))$ for any $\mathcal{F} \in \mathbf{D}_{qc}(X)$. Now Lemma 4.1.11 implies that $\mathcal{H}^{i}(\mathcal{F})$ is almost coherent if and only if so is $\mathrm{H}^{0}(X, \mathcal{H}^{i}(\mathcal{F}))$. So the functor $\mathbf{R}\Gamma(X, -)$ sends $\mathbf{D}^{*}_{qc,acoh}(X)$ to $\mathbf{D}^{*}_{acoh}(A)$.

We also observe that the functor (-) clearly sends $\mathbf{D}_{acoh}(A)$ to $\mathbf{D}_{qc,acoh}(X)$. Thus we conclude that (-) and $\mathbf{R}\Gamma(X, -)$ induce an equivalence of $\mathbf{D}_{acoh}(A)$ and $\mathbf{D}_{qc,acoh}(X)$. The bounded versions follow from *t*-exactness of both functors.

Lemma 4.3.5. Let $X = \operatorname{Spec} A$ be an affine *R*-scheme. Then the almostification functor

$$(-)^a \colon \mathbf{D}^*_{qc}(X) \to \mathbf{D}^*_{aqc}(X)$$

induces an equivalence $\mathbf{D}_{qc}^*(X)/\mathbf{D}_{qc,\Sigma_X}^*(X) \xrightarrow{\sim} \mathbf{D}_{aqc}^*(X)^a$ for any $* \in \{ ", ", +, -, b \}$. Similarly, the induced functor

$$\mathbf{D}^*_{qc,acoh}(X)/\mathbf{D}^*_{qc,\Sigma_X}(X) \xrightarrow{\sim} \mathbf{D}^*_{acoh}(X)^a$$

is an equivalence for any $* \in \{", ", +, -, b\}$.

Proof. The functor $(-)_{!}: \mathbf{D}_{aqc}^{*}(X)^{a} \to \mathbf{D}_{qc}^{*}(X)$ gives the left adjoint to $(-)^{a}$ such that $\mathrm{Id} \to (-)_{!} \circ (-)^{a}$ is an isomorphism and the kernel of $(-)^{a}$ consists exactly of the morphisms f such that $\mathrm{cone}(f) \in \mathbf{D}_{qc,\Sigma_{X}}(X)$. Thus the dual version of [GZ67, Proposition 1.3] finishes the proof of the first claim. The proof of the second claim is similar once one notices that $\widetilde{M^{a}}$ is almost coherent for any almost coherent A^{a} -module M^{a} . The latter fact follows from Lemma 4.1.11.

Lemma 4.1.11 ensures that $\mathbf{D}(A)^a \simeq \mathbf{D}(A)/\mathbf{D}_{\Sigma_A}(A)$. As (-) clearly sends $\mathbf{D}_{\Sigma_A}(A)$ into $\mathbf{D}^*_{qc,\Sigma_X}(X)$, we conclude that it induces the functor

$$(\widetilde{(-)}: \mathbf{D}^*(A)^a \to \mathbf{D}^*_{aqc}(X)^a$$
.

Theorem 4.3.6. Let $X = \operatorname{Spec} A$ be an affine *R*-scheme. Then the functor

$$(-): \mathbf{D}(A)^a \to \mathbf{D}_{aqc}(X)^a$$

is a *t*-exact equivalence of triangulated categories with quasi-inverse given by $\mathbf{R}\Gamma(X, -)$. Moreover, these two functors induce equivalences

$$\widetilde{(-)} \colon \mathbf{D}^*_{acoh}(A)^a \longleftrightarrow \mathbf{D}^*_{acoh}(X)^a \colon \mathbf{R}\Gamma(X,-)$$

for any $* \in \{"", +, -, b\}.$

 $^{^{20}}$ with their standard *t*-structures

Proof. We note that Lemma 4.3.4 ensures that $(-): \mathbf{D}_{qc,acoh}^*(X) \to \mathbf{D}_{acoh}^*(X)^a$ is an equivalence with quasi-inverse $\mathbf{R}\Gamma(X, -)$. Moreover, $(-)^a$ induces an equivalence between $\mathbf{D}_{\Sigma_A}(A)$ and $\mathbf{D}_{qc,\Sigma_X}(X)$; we leave the verification to the interested reader. Thus, Lemma 4.3.5 ensures that (-) gives an equivalence

$$\mathbf{D}(A)^a \simeq \mathbf{D}(A)/\mathbf{D}_{\Sigma_A}(A) \xrightarrow{\sim} \mathbf{D}_{qc}(X)/\mathbf{D}_{qc,\Sigma_X}(X) \simeq \mathbf{D}_{aqc}(X)^a$$
.

Its inverse is given by the functor $\mathbf{D}_{aqc}(X)^a \to \mathbf{D}(A)^a$ induced by $\mathbf{R}\Gamma(X, -)$ on $\mathbf{D}_{qc}(X)$ that exactly coincides with $\mathbf{R}\Gamma(X, -): \mathbf{D}_{aqc}(X)^a \to \mathbf{D}(A)^a$ by Proposition 3.5.23.

The version with almost coherent cohomology sheaves is similar to the analogous statement from Lemma 4.3.4. $\hfill \Box$

Lemma 4.3.7. Let $f: X \to Y$ be a morphism of *R*-schemes.

- (1) Suppose that $X = \operatorname{Spec} B$, $Y = \operatorname{Spec} A$ are affine *R*-schemes. Then $\mathbf{L}f^*(\widetilde{M^a})$ is functorially isomorphic to $\widetilde{M^a \otimes_{A^a}} B^a$ for any $M^a \in \mathbf{D}(A)^a$.
- (2) The functor $\mathbf{L}f^*$ carries an object of $\mathbf{D}^*_{aqc}(Y)$ to an object of $\mathbf{D}^*_{aqc}(X)$ for $* \in \{",",-\}$.
- (3) The functor $\mathbf{L}f^*$ carries an object of $\mathbf{D}^*_{agc}(Y)^a$ to an object of $\mathbf{D}^*_{agc}(X)^a$ for $* \in \{ ", ", \}$.
- (4) Suppose that X is an almost coherent R-scheme. Then the functor $\mathbf{L}f^*$ carries an object of $\mathbf{D}^-_{qc,acoh}(Y)$ (resp. $\mathbf{D}^-_{acoh}(Y)$) to an object of $\mathbf{D}^-_{qc,acoh}(X)$ (resp. $\mathbf{D}^-_{acoh}(X)$).
- (5) Suppose that X is an almost coherent R-scheme. Then the functor $\mathbf{L}f^*$ carries an object of $\mathbf{D}^-_{acoh}(Y)^a$ to an object of $\mathbf{D}^-_{acoh}(X)^a$.

Proof. We start with (1). We use Proposition 3.5.20 to see that $\mathbf{L}f^*(\widetilde{M^a}) \simeq (\mathbf{L}f^*(\widetilde{M}))^a$. Thus it suffices to show that $\mathbf{L}f^*(\widetilde{M}) \simeq \widetilde{M \otimes_A^L B}$ as $(\widetilde{M \otimes_A^L B})^a \simeq \widetilde{M^a \otimes_{A^a}^L B^a}$ by Proposition 2.4.16. But the result for quasi-coherent complexes is classical.

Now we show (2). We note that Lemma 3.2.17 implies that $\mathbf{L}f^*(\widetilde{\mathfrak{m}} \otimes \mathcal{F}) \simeq \widetilde{\mathfrak{m}} \otimes \mathbf{L}f^*(\mathcal{F})$ for any $\mathcal{F} \in \mathbf{D}(Y)$. Thus we can replace \mathcal{F} with $\widetilde{\mathfrak{m}} \otimes \mathcal{F}$ to assume that it is quasi-coherent. Then it is a standard fact that $\mathbf{L}f^*$ sends $\mathbf{D}_{qc}^*(Y)$ to $\mathbf{D}_{qc}^*(X)$ for $* \in \{ ", -\}$.

(3) follows from (2) by noting that $\mathbf{L}f^*(\mathcal{F}^a) \simeq (\mathbf{L}f^*(\mathcal{F}^a))^a$ according to Proposition 3.5.20.

To prove (4), we again use the isomorphism $\mathbf{L}f^*(\widetilde{\mathfrak{m}}\otimes \mathfrak{F})\simeq \widetilde{\mathfrak{m}}\otimes \mathbf{L}f^*(\mathfrak{F})$ to assume that \mathfrak{F} is in $\mathbf{D}^-_{ac\,acob}(X)$. Then we use the spectral sequence

$$\mathbf{E}_{2}^{p,q} = \mathbf{L}^{-p} f^{*}(\mathcal{H}^{q}(\mathcal{F})) \Rightarrow \mathcal{H}^{p+q}(\mathbf{L}f^{*}\mathcal{F})$$

Since $\mathbf{Qcoh}_X^{\mathrm{acoh}}$ is a Weak Serre subcategory of \mathbf{Mod}_X , it suffices to show that each of the \mathcal{O}_X -modules $\mathrm{L}^{-p} f^*(\mathcal{H}^q(\mathcal{F}))$ is almost coherent. In particular, we can assume that \mathcal{F} is a quasi-coherent, almost coherent \mathcal{O}_Y -module.

As X is almost coherent, Lemma 4.1.17 guarantees that it is actually enough to show that each of those sheaves is almost finitely presented. Since the statement is local, we can assume that both X and Y are affine. Thus Lemma 4.3.4 guarantees that there is $M \in \mathbf{Mod}_A^{\mathrm{acoh}}$ such that $\widetilde{M} \simeq \mathcal{F}$. Thus Part (1) and Lemma 4.1.11 ensure that it is sufficient to show that $M^a \otimes_{A^a}^L B^a \simeq (M \otimes_A^L B)^a$ has almost finitely presented cohomology modules. This is exactly the content of Corollary 2.7.2.

5) follows from (4) as
$$\mathbf{L}f^*(\mathcal{F}^a) \simeq (\mathbf{L}f^*(\mathcal{F}^a_!))^a$$
.

Lemma 4.3.8. Let X be an R-scheme.

- (1) Suppose that $X = \operatorname{Spec} A$ is an affine *R*-scheme. Then $\widetilde{M^a} \otimes_{\mathcal{O}_X^a}^{L} \widetilde{N^a}$ is functorially isomorphic to $\widetilde{M^a} \otimes_{A^a}^{L} N^a$ for any $M^a, N^a \in \mathbf{D}(A)^a$.
- (2) Let $\mathcal{F}, \mathcal{G} \in \mathbf{D}^*_{aqc}(X)$, then $\mathcal{F} \otimes^L_{\mathcal{O}_X} \mathcal{G} \in \mathbf{D}_{aqc}(X)$ for $* \in \{", ", -\}$.
- (3) Let $\mathcal{F}^a, \mathcal{G}^a \in \mathbf{D}^*_{aqc}(X)^a$, then $\mathcal{F}^a \otimes^L_{\mathcal{O}^a_X} \mathcal{G}^a \in \mathbf{D}_{aqc}(X)^a$ for $* \in \{ ", ", \}$.
- (4) Suppose that X is an almost coherent R-scheme, and let $\mathcal{F}, \mathcal{G} \in \mathbf{D}^{-}_{qc,acoh}(X)$ (resp. $\mathbf{D}^{-}_{acoh}(X)$). Then $\mathcal{F} \otimes^{L}_{\mathcal{O}_{X}} \mathcal{G} \in \mathbf{D}^{-}_{qc,acoh}(X)$ (resp. $\mathbf{D}^{-}_{acoh}(X)$).
- (5) Suppose that X is an almost coherent R-scheme, and let $\mathcal{F}^a, \mathcal{G}^a \in \mathbf{D}^-_{acoh}(X)^a$. Then $\mathcal{F}^a \otimes_{\mathcal{O}^a_X}^L \mathcal{G}^a \in \mathbf{D}^-_{acoh}(X)^a$.

Proof. The proof is basically identical to that of Lemma 4.3.7 and left to the reader. We only mention that one has to use Proposition 2.6.14 in place of Corollary 2.7.2. \Box

Lemma 4.3.9. Let $f: X \to Y$ be a quasi-compact and quasi-separated morphism of *R*-schemes. Suppose that *Y* is quasi-compact. Then

- (1) The functor $\mathbf{R}f_*$ carries $\mathbf{D}^*_{aac}(X)$ to $\mathbf{D}^*_{aac}(Y)$ for any $* \in \{", ", -, +, b\}$.
- (2) The functor $\mathbf{R}f_*$ carries $\mathbf{D}^*_{aqc}(X)^a$ to $\mathbf{D}^*_{aqc}(Y)^a$ for any $* \in \{", ", -, +, b\}$.

Proof. Proposition 3.5.23 guarantees that $(\mathbf{R}f_*\mathfrak{F})^a \simeq \mathbf{R}f_*\mathfrak{F}^a$. Since $(\widetilde{\mathfrak{m}} \otimes \mathfrak{F})^a \simeq \mathfrak{F}^a$, we see that it suffices to show that the functor

 $\mathbf{R}f_*(\widetilde{\mathfrak{m}}\otimes -)$

carries $\mathbf{D}_{aqc}^*(X)$ to $\mathbf{D}_{aqc}^*(Y)$ for any $* \in \{ ", -, +, b \}$. Since $\widetilde{\mathfrak{m}} \otimes \mathcal{F}$ is in $\mathbf{D}_{qc}(X)$, we conclude that it is enough to show that $\mathbf{R}f_*(-)$ carries $\mathbf{D}_{qc}^*(X)$ to $\mathbf{D}_{qc}^*(Y)$ for any $* \in \{ ", -, +, b \}$. This is proven in [Sta21, Tag 08D5].

Before going to the case of the derived Hom-functors, we recall the construction of the functorial map

$$\psi \colon \operatorname{\mathbf{R}Hom}_{A}(M,N) \to \operatorname{\mathbf{R}}_{\operatorname{\operatorname{\mathcal H}Om}_{\mathcal{O}_{X}}}(\widetilde{M},\widetilde{N})$$

for any $M \in \mathbf{D}^{-}(A)$, $N \in \mathbf{D}^{+}(A)$ on an affine scheme $X = \operatorname{Spec} A$. Indeed, the functor (-) is left adjoint to the global section functor $\Gamma(X, -)$ on the abelian level. Thus after deriving these functors, we get that $\widetilde{-}$ is left adjoint to $\mathbf{R}\Gamma(X, -)$. Therefore, for any $\mathcal{F} \in \mathbf{D}(X)$, there is a canonical morphism $\mathbf{R}\Gamma(X, \mathcal{F}) \to \mathcal{F}$. We apply it to $\mathcal{F} = \mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_X}(\widetilde{M}, \widetilde{N})$ to get the desired morphism

$$\psi \colon \mathbf{R}\mathrm{Hom}_{A}(M,N) \to \mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_{X}}(\widetilde{M},\widetilde{N}) \;.$$

Lemma 4.3.10. Let X be an almost coherent R-scheme.

(1) Suppose $X = \operatorname{Spec} A$ is an affine *R*-scheme. The canonical map

$$\psi \colon \mathbf{R} \widetilde{\mathrm{Hom}}_A(M,N) \to \mathbf{R} \underline{\mathcal{H}om}_{\mathcal{O}_X}(\widetilde{M},\widetilde{N})$$

is an almost isomorphism for $M \in \mathbf{D}^{-}_{acoh}(A), N \in \mathbf{D}^{+}(A)$.

(2) Suppose $X = \operatorname{Spec} A$ is an affine *R*-scheme. There is a functorial isomorphism

 \mathbf{R} al $\mathrm{Hom}_{A^a}(M^a, N^a) \simeq \mathbf{R}\underline{al}\mathcal{H}om_{\mathcal{O}_X^a}(\widetilde{M^a}, \widetilde{N^a})$

for $M^a \in \mathbf{D}^-_{acoh}(A)^a$, $N^a \in \mathbf{D}^+(A)^a$. We also get a functorial almost isomorphism

$$\mathbf{R}\operatorname{Hom}_{A^a}(M^a, N^a) \simeq^a \mathbf{R} \underbrace{\mathcal{H}om}_{\mathcal{O}^a_{\mathcal{V}}}(M^a, N^a)$$

for $M \in \mathbf{D}^{-}_{acoh}(A), N \in \mathbf{D}^{+}(A)$.

- (3) Suppose $\mathcal{F} \in \mathbf{D}^{-}_{acoh}(X)$ and $\mathcal{G} \in \mathbf{D}^{+}_{aqc}(X)$. Then $\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_{X}}(\mathcal{F},\mathcal{G}) \in \mathbf{D}^{+}_{aqc}(X)$.
- (4) Suppose $\mathfrak{F}^{a} \in \mathbf{D}^{-}_{acoh}(X)^{a}$ and $\mathfrak{G}^{a} \in \mathbf{D}^{+}_{aqc}(X)^{a}$. Then $\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_{X}^{a}}(\mathfrak{F}^{a},\mathfrak{G}^{a}) \in \mathbf{D}^{+}_{aqc}(X)$ and $\mathbf{R}\underline{al\mathcal{H}om}_{\mathcal{O}_{X}^{a}}(\mathfrak{F}^{a},\mathfrak{G}^{a}) \in \mathbf{D}^{+}_{aqc}(X)^{a}$.

Proof. We start with (1). We use the convergent compatible spectral sequences

$$\begin{aligned} & \mathbf{E}_{2}^{p,q} = \mathrm{Ext}_{A}^{p}(\widetilde{\mathbf{H}^{-q}(M)}, N) \Rightarrow \mathrm{Ext}_{A}^{p+q}(M, N) \\ & \mathbf{E}_{2}^{\prime p,q} = \mathcal{E}xt_{\mathcal{O}_{X}}^{p}\left(\widetilde{\mathbf{H}^{-q}(M)}, \widetilde{N}\right) \Rightarrow \mathcal{E}xt_{\mathcal{O}_{X}}^{p+q}\left(\widetilde{M}, \widetilde{N}\right) \end{aligned}$$

to see that we may assume that $M \in \mathbf{Acoh}_A$ is just a module in degree 0. Similarly, we use the compatible spectral sequences

$$\begin{split} \mathbf{E}_{2}^{p,q} &= \mathrm{Ext}_{A}^{q}(\widetilde{M}, \mathrm{H}^{p}(N)) \Rightarrow \mathrm{Ext}_{A}^{\widetilde{p+q}}(\widetilde{M}, N) \\ \mathbf{E}_{2}^{\prime p,q} &= \mathcal{E}xt_{\mathbb{O}_{X}}^{q}(\widetilde{M}, \widetilde{\mathrm{H}^{p}(N)}) \Rightarrow \mathcal{E}xt_{\mathbb{O}_{X}}^{p+q}(\widetilde{M}, \widetilde{N}) \end{split}$$

to also assume that $N \in \mathbf{Mod}_A$. So the claim boils down to showing that the natural maps

$$\operatorname{Ext}_{A}^{\widetilde{p}}(\widetilde{M}, N) \to \mathcal{E}xt_{\mathcal{O}_{X}}^{p}(\widetilde{M}, \widetilde{N})$$

are almost isomorphisms for any $M \in \operatorname{Acoh}_A$, $N \in \operatorname{Mod}_A$, and $p \ge 0$. Lemma 3.1.5 says that it is sufficient to say that kernel and cokernel are annihilated by any finitely generated sub-ideal $\mathfrak{m}_0 \subset \mathfrak{m}$.

Recall that, for any \mathcal{O}_X -modules \mathcal{F} , \mathcal{G} , the sheaf $\mathcal{E}xt^p_{\mathcal{O}_X}(\mathcal{F},\mathcal{G})$ is canonically isomorphic to sheaffication of the presheaf

$$U \mapsto \operatorname{Ext}_{\mathcal{O}_U}^p(\mathcal{F}|_U, \mathcal{G}|_U)$$
.

Thus, in order to show that the map $\operatorname{Ext}_{A}^{\widetilde{p}}(M, N) \to \mathcal{E}xt_{\mathcal{O}_{X}}^{p}(\widetilde{M}, \widetilde{N})$ is an almost isomorphism, it suffices to show that

$$\operatorname{Ext}_{A}^{p}(M,N) \otimes_{A} A_{f} \to \operatorname{Ext}_{\mathcal{O}_{X_{f}}}^{p}(\widetilde{M_{f}},\widetilde{N_{f}})$$

is an almost isomorphism. Now we use canonical isomorphisms

$$\operatorname{Ext}_{\mathcal{O}_{X_f}}^{p}(\widetilde{M_f}, \widetilde{N_f}) \simeq \operatorname{Hom}_{\mathbf{D}(X_f)}(\widetilde{M_f}, \widetilde{N_f}[p])$$
$$\simeq \operatorname{Hom}_{\mathbf{D}(A_f)}(M_f, N_f[p])$$
$$\simeq \operatorname{Ext}_{A_f}^{p}(M_f, N_f),$$

where the second isomorphism uses that $(\tilde{-})$ induces a *t*-exact equivalence $(\tilde{-})$: $\mathbf{D}(A_f) \to \mathbf{D}_{qc}(\operatorname{Spec} A)$. Thus, the question boils down to showing that the natural map

$$\operatorname{Ext}_{A}^{p}(M,N) \otimes_{A} A_{f} \to \operatorname{Ext}_{A_{f}}^{p}(M_{f},N_{f})$$

is an almost isomorphism. This follows from Proposition 2.7.9.

(2) formally follows from (1) by using Proposition 3.5.8(1).

(3) is also a basic consequence of (2). Indeed, the claim is local, so we can assume that X = Spec A is an affine R-scheme. In that case we use Theorem 4.3.6 to say that $\mathcal{F} \simeq \widetilde{M}$, $\mathcal{G} \simeq \widetilde{N}$ for

some $M \in \mathbf{D}^-_{acoh}(A)$, $N \in \mathbf{D}^+(A)$. Then $\mathbf{R} \underline{\mathcal{H}om}_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}) \simeq \mathbf{R} \mathrm{Hom}_A(M, N)$ by (2), and the latter complex has almost quasi-coherent cohomology sheaves by design.

(4) easily follows from (3) and the isomorphisms

$$\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_X}(\mathcal{F}^a, \mathcal{G}^a) \simeq \mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_X}(\mathcal{F}^a_!, \mathcal{G})$$

$$\mathbf{R}\underline{al\mathcal{H}om}_{\mathcal{O}_X^a}(\mathcal{F}^a, \mathcal{G}^a) \simeq \mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_X}(\mathcal{F}^a_!, \mathcal{G})^a$$

that come from Lemma (1) and Definition 3.5.6.

Corollary 4.3.11. Let X be an almost coherent R-scheme.

- (1) Let $\mathcal{F} \in \mathbf{D}^{-}_{aqc,acoh}(X), \ \mathcal{G} \in \mathbf{D}^{+}_{aqc,acoh}(X)$. Then $\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_{X}}(\mathcal{F},\mathcal{G}) \in \mathbf{D}^{+}_{aqc,acoh}(X)$.
- (2) Let $\mathcal{F}^a \in \mathbf{D}^-_{acoh}(X)^a$, $\mathcal{G}^a \in \mathbf{D}^+_{acoh}(X)^a$. Then $\mathbf{R}\underline{al\mathcal{H}om}_{\mathcal{O}^a_{\mathbf{Y}}}(\mathcal{F}^a, \mathcal{G}^a) \in \mathbf{D}^+_{acoh}(X)^a$.

Proof. The question is local on X, so we can suppose that X = Spec A is affine. Then Lemma 4.3.10, Theorem 4.3.6, and Lemma 4.1.11 reduce both question to showing that $\mathbb{R}\text{Hom}_A(M, N) \in \mathbb{D}^+_{acoh}(A)$ for $M \in \mathbb{D}^-_{acoh}(A)$ and $N \in \mathbb{D}^+_{acoh}(A)$. This is the content of Proposition 2.6.15.

4.4. Formal Schemes. The Category of Almost Coherent $\mathcal{O}^a_{\mathfrak{X}}$ -modules. In this Section we discuss the notion of almost coherent sheaves on "good" formal schemes. One of the main complications is that there is no good notion of a "quasi-coherent" sheaf on a formal scheme. Namely, even though there is a notion of adically quasi-coherent sheaves on a big class of formal schemes due to [FK18, §I.3], this notion does not really behave well. For example, the category of adically quasi-coherent sheaves usually is not abelian. One of the main difficulties in working with adically quasi-coherent sheaves is the lack of the Artin-Rees lemma beyond the case of finitely generated modules. More precisely, many operations with adically quasi-coherent sheaves require taking completions, but it is difficult to control the effect of it without the use of the Artin-Rees lemma.

The way we deal with this problem is to use a version of the Artin-Rees lemma (Lemma 2.8.6) for almost finitely generated modules over "good" rings. The presence of the Artin-Rees lemma suggests that it is reasonable to expect that we might have a good notion of adically quasi-coherent, almost coherent $\mathcal{O}_{\mathfrak{X}}$ -modules on some "good" class of formal schemes.

We start by giving the Setup in which we can develop the theory of almost coherent sheaves

Set-up 4.4.1. We fix a ring R with a finitely generated ideal I such that R is I-adically complete, I-adically topologically universally adhesive²¹, and I-torsion free with an ideal \mathfrak{m} such that $I \subset \mathfrak{m}$, $\mathfrak{m}^2 = \mathfrak{m}$ and $\mathfrak{\tilde{m}} := \mathfrak{m} \otimes_R \mathfrak{m}$ is R-flat.

The basic example of such a ring is a complete microbial²² valuation ring R with algebraically closed fraction field K. We pick a pseudo-uniformizer ϖ and define $I := (\varpi)$, $\mathfrak{m} := \bigcup_{i=1}^{\infty} (\varpi^{1/n})$ for some compatible choice of roots of ϖ . We note that R is topologically universally adhesive by [FGK11, Theorem 7.3.2].

We note that the assumptions in Setup 4.4.1 imply that any finitely presented algebra over a topologically finitely presented R-algebra is coherent and I-adically adhesive. Coherence follows from [FGK11, Proposition 7.2.2] and adhesiveness basically follows from the definition. In what follows, we will use those facts without saying.

²¹This means that the algebra $R(X_1, \ldots, X_n)[T_1, \ldots, T_m]$ is *I*-adically adhesive for any *n* and *m*

²²A valuation ring R is microbial if there is a non-zero topologically nilpotent element $\varpi \in R$. Any such element is called a pseudo-uniformizer.

In what follows, \mathfrak{X} will always mean a topologically finitely presented formal *R*-scheme. We will denote by $\mathfrak{X}_k := \mathfrak{X} \times_{\operatorname{Spf} R} \operatorname{Spec} R/I^{k+1}$ the "reduction" schemes. They come together with the closed immersion $i_k : \mathfrak{X}_k \to \mathfrak{X}$. Also, given any $\mathcal{O}_{\mathfrak{X}}$ -module \mathcal{F} , we will always denote its "reduction" $i_k^* \mathcal{F}$ by \mathcal{F}_k .

Definition 4.4.2. [FK18, Definition I.3.1.3] An $\mathcal{O}_{\mathfrak{X}}$ -module \mathcal{F} on a formal scheme \mathfrak{X} of finite ideal type is called *adically quasi-coherent* if $\mathcal{F} \to \lim_n \mathcal{F}_n$ is an isomorphism and, for any open formal subscheme $\mathfrak{U} \subset \mathcal{F}$ and any ideal of definition \mathcal{I} of finite type, the sheaf $\mathcal{F}/\mathcal{I}\mathcal{F}$ is a quasi-coherent sheaf on the scheme $(\mathfrak{U}, \mathcal{O}_{\mathfrak{U}}/\mathcal{I})$.

Definition 4.4.3. We say that an $\mathcal{O}^a_{\mathfrak{X}}$ -module \mathfrak{F}^a is almost adically quasi-coherent if $\mathfrak{F}^a_! \simeq \widetilde{\mathfrak{m}} \otimes \mathfrak{F}$ is an adically quasi-coherent $\mathcal{O}_{\mathfrak{X}}$ -module. We denote this category by $\mathbf{AQcoh}_{\mathfrak{X}}$.

We say that an $\mathcal{O}_{\mathfrak{X}}$ -module \mathfrak{F} is almost adically quasi-coherent if \mathfrak{F}^a is an almost quasi-coherent $\mathcal{O}^a_{\mathfrak{X}}$ -module.

Remark 4.4.4. In general, we can not say that an adically quasi-coherent $\mathcal{O}_{\mathfrak{X}}$ -module \mathfrak{F} is almost adically quasi-coherent. The problem is that the sheaf $\widetilde{\mathfrak{m}} \otimes \mathfrak{F}$ might not be complete, i.e. the map $\widetilde{\mathfrak{m}} \otimes \mathfrak{F} \to \lim_k \widetilde{\mathfrak{m}} \otimes \mathfrak{F}_k$ is a priori only an almost isomorphism.

Lemma 4.4.5. Let \mathfrak{X} be a topologically finitely presented formal *R*-scheme for *R* as in the Setup 4.4.1, and let \mathcal{F}^a be an almost adically quasi-coherent $\mathcal{O}^a_{\mathfrak{X}}$ -module. Then \mathcal{F}^a_k is almost quasi-coherent for all *k*. Moreover, if an $\mathcal{O}^a_{\mathfrak{X}}$ -module \mathcal{G}^a is annihilated by some I^{n+1} . Then \mathcal{G}^a is almost adically quasi-coherent if and only if so is \mathcal{G}^a_n .

Proof. In order to prove the first claim, it is sufficient to show that $\widetilde{\mathfrak{m}} \otimes \mathcal{F}_k$ is quasi-coherent provided that $\widetilde{\mathfrak{m}} \otimes \mathcal{F}$ is adically quasi-coherent. We use Corollary 3.2.18 to say that $\widetilde{\mathfrak{m}} \otimes \mathcal{F}_k \simeq (\widetilde{\mathfrak{m}} \otimes \mathcal{F})_k$ and the reduction of an adically quasi-coherent module is quasi-coherent. Therefore, each \mathcal{F}_k^a is almost adically quasi-coherent.

Now if \mathfrak{G} is annihilated by I^{n+1} then $\mathfrak{G} = i_{n,*}\mathfrak{G}_n$. We use the Projection Formula (Corollary 3.3.6) to say that $\widetilde{\mathfrak{m}} \otimes \mathfrak{G} \simeq i_{n,*}(\mathfrak{G}_n \otimes \widetilde{\mathfrak{m}})$. Clearly, $i_{n,*}$ sends quasi-coherent sheaves to adically quasi-coherent sheaves. So \mathfrak{G}^a is almost adically quasi-coherent if so is \mathfrak{G}_n^a .

Definition 4.4.6. We say that an $\mathcal{O}^a_{\mathfrak{X}}$ -module \mathcal{F}^a is of almost finite type (resp. almost finitely presented) if \mathcal{F}^a is almost adically quasi-coherent, and there is a covering of \mathfrak{X} by open affines $\{\mathfrak{U}_i\}_{i\in I}$ such that $\mathcal{F}^a(\mathfrak{U}_i)$ is an almost finitely generated (resp. almost finitely presented) $\mathcal{O}^a_{\mathfrak{X}}(\mathfrak{U}_i)$ -module. We denote these categories by $\mathbf{AQcoh}^{aft}_{\mathfrak{X}}$ and $\mathbf{AQcoh}^{afp}_{\mathfrak{X}}$ respectively.

We say that an $\mathcal{O}_{\mathfrak{X}}$ -module \mathcal{F} is of almost finite type (resp. almost finitely presented) if so is \mathcal{F}^a .

Definition 4.4.7. We say that an $\mathcal{O}_{\mathfrak{X}}$ -module \mathfrak{F} is *adically quasi-coherent of almost finite type* (resp. *adically quasi-coherent almost finitely presented*) if it is adically quasi-coherent and there is a covering of \mathfrak{X} by open affines $\{\mathfrak{U}_i\}_{i\in I}$ such that $\mathfrak{F}(\mathfrak{U}_i)$ is an almost finitely generated (resp. almost finitely presented) $\mathcal{O}_{\mathfrak{X}}(\mathfrak{U}_i)$ -module. We denote these categories by $\mathbf{Qcoh}_{\mathfrak{X}}^{\mathrm{aff}}$ and $\mathbf{Qcoh}_{\mathfrak{X}}^{afp}$ respectively.

Remark 4.4.8. If \mathcal{F}^a is a finite type (resp. finitely presented) $\mathcal{O}^a_{\mathfrak{X}}$ -module, then $(\mathcal{F}^a)_!$ is adically quasi-coherent of almost finite type (resp. almost finite presentation).

Remark 4.4.9. We note that, a priori, it is not clear if \mathcal{F}^a is an almost finite type (resp. almost finitely presented) $\mathcal{O}^a_{\mathfrak{X}}$ -module for an adically quasi-coherent almost finite type (resp. almost finitely presented) $\mathcal{O}_{\mathfrak{X}}$ -module \mathcal{F} . The problem is that our definition of adically quasi-coherent almost finite type (resp. almost finitely presented) module does not require $\widetilde{\mathfrak{m}} \otimes \mathcal{F}$ to be adically quasi-coherent. However, we will show in Lemma 4.4.10 that it is automatic in this case.

Lemma 4.4.10. Let \mathfrak{X} be a topologically finitely presented formal *R*-scheme for *R* as in the Setup 4.4.1, and let \mathcal{F} be an adically quasi-coherent of almost finite type (resp. almost finitely presented) $\mathcal{O}_{\mathfrak{F}}$ -module. Then $\widetilde{\mathfrak{m}} \otimes \mathfrak{F}$ is adically quasi-coherent. In particular, \mathfrak{F} is almost of finite type (resp. almost finitely presented).

Proof. Corollary 2.5.9 and Lemma 3.3.2 imply that the only condition we really need to check is that $\widetilde{\mathfrak{m}} \otimes \mathcal{F}$ is adically quasi-coherent. Therefore, it suffices to prove the result for an adically quasi-coherent almost finite type $\mathcal{O}_{\mathfrak{X}}$ -module \mathcal{F} .

Since the question is local on \mathfrak{X} , we can assume that $\mathfrak{X} = \operatorname{Spf} A$ is affine and $M \coloneqq \mathfrak{F}(X)$ is almost finitely generated over A. Then we use [FK18, Theorem I.3.2.8] to say that \mathcal{F} is isomorphic to M^{Δ} . We claim that $\widetilde{\mathfrak{m}} \otimes \mathfrak{F}$ is isomorphic to $(\widetilde{\mathfrak{m}} \otimes_A M)^{\Delta}$ as that would imply that $\widetilde{\mathfrak{m}} \otimes \mathfrak{F}$ is adically quasi-coherent by [FK18, Proposition I.3.2.2]. In order to show that $\widetilde{\mathfrak{m}} \otimes \mathfrak{F}$ is isomorphic to $(\widetilde{\mathfrak{m}} \otimes_R M)^{\Delta}$ we need to check two things: for any open affine $\operatorname{Spf} B = \mathfrak{U} \subset \mathfrak{X}$ the B-module $(\widetilde{\mathfrak{m}} \otimes \mathfrak{F})(\mathfrak{U})$ is *I*-adically complete, and then the natural map $(\widetilde{\mathfrak{m}} \otimes_R M)\widehat{\otimes}_A B \to (\widetilde{\mathfrak{m}} \otimes \mathfrak{F})(\mathfrak{U})$ is an isomorphism.

We start with the first claim. Lemma 3.3.2 says that $(\widetilde{\mathfrak{m}} \otimes \mathfrak{F})(\mathfrak{U})$ is isomorphic to $\widetilde{\mathfrak{m}} \otimes_R \mathfrak{F}(\mathfrak{U})$. Since \mathcal{F} is adically quasi-coherent, $\mathcal{F}(\mathfrak{U}) \simeq A \widehat{\otimes}_A B$, so $(\widetilde{\mathfrak{m}} \otimes \mathcal{F})(\mathfrak{U}) \simeq \widetilde{\mathfrak{m}} \otimes_R (M \widehat{\otimes}_A B)$. Lemma 2.7.1 says that $M \otimes_A B$ is almost finitely generated over B, so it is already I-adically complete by Lemma 2.8.7. Therefore, we see that $\widetilde{\mathfrak{m}} \otimes_R \mathfrak{F}(\mathfrak{U}) \simeq \widetilde{\mathfrak{m}} \otimes_R (M \otimes_A B)$, and the latter is almost finitely generated over B by Corollary 2.5.9. Thus we use Lemma 2.8.7 once more to show its completeness.

Now we show that the natural morphism $(\widetilde{\mathfrak{m}} \otimes_R M) \widehat{\otimes}_A B \to (\widetilde{\mathfrak{m}} \otimes \mathfrak{F})(\mathfrak{U})$ is an isomorphism. Again, using the same results as above we can get rid of any completions and identify this map with the "identity" map

$$(\widetilde{\mathfrak{m}} \otimes_R M) \otimes_A B \to \widetilde{\mathfrak{m}} \otimes_R (M \otimes_A B)$$

This finishes the proof.

Lemma 4.4.11. Let \mathfrak{X} be a topologically finitely presented formal *R*-scheme for *R* as in the Setup 4.4.1, and let \mathcal{F}^a be an almost finite type (resp. almost finitely presented) $\mathcal{O}^a_{\mathfrak{X}}$ -module. Then the $\mathcal{O}^a_{\mathfrak{X}_k}$ -module \mathcal{F}^a_k is almost finite type (resp. almost finitely presented) for any integer k.

Proof. Lemma 4.4.5 implies that each \mathcal{F}_k^a is an almost quasi-coherent \mathcal{O}_{X_k} -module. So it is sufficient to find a covering of \mathfrak{X}_k by open affines $\mathfrak{U}_{i,k}$ such that $\mathcal{F}_k^a(\mathfrak{U}_{i,k})$ is almost finitely generated (resp. almost finitely presented) over $\mathcal{O}^a_{\mathfrak{X}_k}(\mathfrak{U}_{i,k})$. We note that underlying topological spaces of \mathfrak{X} and \mathfrak{X}_k are the same, so we can choose some covering of \mathfrak{X} by open affines \mathfrak{U}_i such that $\mathfrak{F}^a(\mathfrak{U}_i)$ are almost finitely generated (resp. almost finitely presented) over $\mathcal{O}_{\mathfrak{X}}^{a}(\mathfrak{U}_{i})$, and define $\mathfrak{U}_{i,k}$ as the "reductions" of \mathfrak{U}_i . Then using the vanishing result for higher cohomology groups of adically quas-coherent sheaves on affine formal schemes of finite type [FK18, Theorem I.7.1.1] and Lemma 3.3.2, we deduce that

$$\begin{aligned} \mathcal{F}_k^a(\mathfrak{U}_{i,k}) \simeq (\widetilde{\mathfrak{m}} \otimes \mathcal{F}_k^a)(\mathfrak{U}_{i,k})^a \simeq \left(\widetilde{\mathfrak{m}} \otimes \mathcal{F}(\mathfrak{U}_i) / I^{k+1}\right)^a \\ \text{ted (resp. almost finitely presented) over } \mathcal{O}_{\mathfrak{T}_k}(\mathfrak{U}_{i,k}). \end{aligned}$$

is almost finitely generated (resp. almost finitely presented) over $\mathcal{O}_{\mathfrak{X}_{k}}(\mathfrak{U}_{i,k})$.

Lemma 4.4.12. Let \mathfrak{X} be a locally topologically finitely presented formal R-scheme for R as in the Setup 4.4.1, and let \mathcal{F}^a be an almost finite type (resp. almost finitely presented) $\mathcal{O}^a_{\mathfrak{X}}$ -module. Then $\mathcal{F}^{a}(\mathfrak{U})$ is an almost finitely generated (resp. almost finitely presented) $\mathcal{O}^{a}_{\mathfrak{X}}(\mathfrak{U})$ -module for any open affine $\mathfrak{U} \subset \mathfrak{X}$.

Proof. Corollary 2.5.9 and Lemma 3.3.2 guarantee that we can replace \mathcal{F} with $\widetilde{\mathfrak{m}} \otimes \mathcal{F}$ for the purpose of the proof. Thus we may and do assume that \mathcal{F} is an adically quasi-coherent almost finitely generated (resp. almost finitely presented) $\mathcal{O}_{\mathfrak{X}}$ -module. Then using Lemma 2.7.1 and

Lemma 2.8.7 we can use the argument as in the proof of Lemma 4.4.10 show that the restriction of \mathcal{F} to any open formal subscheme is still adically quasi-coherent of almost finite type (resp. finitely

 \mathcal{F} to any open formal subscheme is still adically quasi-coherent of almost finite type (resp. finitely presented), so we may and do assume that $\mathfrak{X} = \operatorname{Spf} A$ is an affine formal *R*-scheme. Since now \mathfrak{X} is quasi-compact, we can choose a finite refinement of the covering $\mathfrak{X} = \bigcup \mathfrak{U}_i$ such that $\mathcal{F}(\mathfrak{U}_i)$ is almost finitely generated (resp. almost finitely presented) over $\mathcal{O}(\mathfrak{U}_i)$. Thus we may and do assume that a covering (\mathfrak{U}_i) is finite.

Now we have an affine topologically finitely presented formal *R*-scheme $\mathfrak{X} = \text{Spf } A$, a finite covering of \mathfrak{X} by affines $\mathfrak{U}_i = \text{Spf } A_i$, and an adically quasi-coherent $\mathcal{O}_{\mathfrak{X}}$ -module \mathcal{F} such that $\mathcal{F}(\mathfrak{U}_i)$ is almost finitely generated (resp. almost finitely presented) A_i -module. We want to show that $\mathcal{F}(\mathfrak{X})$ is an almost finitely generated (resp. almost finitely presented) A-module.

We firstly deal with the almost finitely generated case. We note that Lemma 4.1.6, Lemma 4.4.11, and [FK18, Theorem I.7.1.1] imply that $\mathcal{F}(\mathfrak{X})/I$ is almost finitely generated. We know that \mathcal{F} is adically quasi-coherent so $\mathcal{F}(\mathfrak{X})$ must be an *I*-adically complete *A*-module. Therefore, Corollary 2.5.16 guarantees that $\mathcal{F}(X)$ is an almost finitely generated *A*-module.

Now we move to the almost finitely presented case. We already now that $\mathcal{F}(\mathfrak{X})$ is almost finitely generated over A. Thus the standard argument with Lemma 2.8.7 implies that $\mathcal{F}(\mathfrak{U}_i) = \mathcal{F}(\mathfrak{X}) \otimes_A A_i$ for any i. Recall that [FK18, Proposition I.4.8.1] implies²³ that each $A \to A_i$ is flat. Since Spf A_i form a covering of Spf A, we conclude that $A \to \prod_{i=1}^{n} A_i$ is faithfully flat. Now the result follows from faithfully flat descent for almost finitely presented modules that is proven in Lemma 2.7.6. \Box

Corollary 4.4.13. Let $\mathfrak{X} = \text{Spf } A$ be a topologically finitely presented affine formal R-scheme for R as in the Setup 4.4.1, and let \mathcal{F}^a be an almost adically quasi-coherent $\mathcal{O}^a_{\mathfrak{X}}$ -module. Then \mathcal{F}^a is almost finite type (resp. almost finitely presented) if and only if $\mathcal{F}^a(\mathfrak{X})$ is almost finitely generated (resp. almost finitely presented) A^a -module.

Similarly, an adically quasi-coherent $\mathcal{O}_{\mathfrak{X}}$ -module \mathcal{F} is almost finite type (resp. almost finitely presented) if and only if $\mathcal{F}(\mathfrak{X})$ is almost finitely generated (resp. almost finitely presented) *A*-module.

Lemma 4.4.14. Let $\mathfrak{X} = \operatorname{Spf} A$ be a topologically finitely presented affine formal *R*-scheme for *R* as in the Setup 4.4.1, let $\varphi \colon N \to M$ be a homomorphism of almost finitely generated *A*-modules. Then the following sequence

$$0 \to (\ker \phi)^{\Delta} \to N^{\Delta} \xrightarrow{\varphi^{\Delta}} M^{\Delta} \to (\operatorname{Coker} \phi)^{\Delta} \to 0$$

is exact. Moreover, $\operatorname{Im}(\varphi)^{\Delta} \simeq \operatorname{Im}(\varphi^{\Delta})$.

Proof. We denote the kernel ker ϕ by K, the image $\text{Im}(\varphi)$ by M', and the cokernel Coker ϕ by Q.

We start with ker φ^{Δ} : We note that $(\ker \varphi^{\Delta})(\mathfrak{X}) = K$, this induces a natural morphism $\alpha \colon K^{\Delta} \to \ker \varphi^{\varphi}$. We show that it is an isomorphism, it suffices to check that it induces an isomorphism on values over a basis of principal open subsets. Now recall that for any A-module L, we have an equality $L^{\Delta}(\operatorname{Spf} A_{\{f\}}) = \widehat{L_f}$ where the completion is taken with respect to the *I*-adic topology. Thus in order to check that α is an isomorphism it suffices to show that $\widehat{K_f}$ is naturally identified with $(\ker \varphi)(\operatorname{Spf} A_{\{f\}}) = \ker(\widehat{N_f} \to \widehat{M_f})$. Using the Artin-Rees Lemma 2.8.6 over the adhesive ring A_f , we conclude that the induced topologies on K_f and M'_f coincide with the *I*-adic ones. This implies that

$$\widehat{K_f} = \lim K_f / I^n K_f = \lim K_f / (I^n N_f \cap K_f) \text{ and } \widehat{M'_f} = \lim M'_f / I^n M'_f = \lim M'_f / (I^n M_f \cap M'_f)$$

²³Topologically universally adhesive rings are by definition "t. u. rigid-Noetherian"

This guarantees that we have two exact sequences:

$$0 \to \widehat{K_f} \to \widehat{M_f} \to \widehat{M_f'} \to 0,$$
$$0 \to \widehat{M_f'} \to \widehat{N_f}$$

In particular, we get that the natural map $\widehat{K_f} \to \ker(\widehat{M_f} \to \widehat{N_f})$ is an isomorphism. That shows that $K^{\Delta} \simeq \ker(\varphi^{\Delta})$.

We prove the claim for $\operatorname{Im} \varphi^{\Delta}$: We note that since the category of $\mathcal{O}_{\mathfrak{X}}$ -modules is abelian, we can identify $\operatorname{Im} \varphi^{\Delta} \simeq \operatorname{Coker}(K^{\Delta} \to N^{\Delta})$. We observe that [FK18, Theorem I.7.1.1] and the established fact above that ker φ is adically quasi-coherent imply that the natural map $\mathcal{F}(\mathfrak{U})/K^{\Delta}(\mathfrak{U}) \to (\operatorname{Im} \varphi^{\Delta})(\mathfrak{U})$ is an isomorphism for any affine open formal subscheme \mathfrak{U} . In particular, we have $(\operatorname{Im} \varphi^{\Delta})(\mathfrak{X}) = M/K = M'$. Therefore, we have a natural map $(M')^{\Delta} \to \operatorname{Im} \varphi^{\Delta}$ and we show that it is an isomorphism. Again it suffices to show that this map is an isomorphism on values over a basis of principal open subsets. Then we use the identification $\mathcal{F}(\mathfrak{U})/K^{\Delta}(\mathfrak{U}) \simeq (\operatorname{Im} \varphi)(\mathfrak{U})$ and the short exact sequence

$$0 \to \widehat{K_f} \to \widehat{M_f} \to \widehat{M'_f} \to 0,$$

to finish the proof.

We show the claim for $\operatorname{Coker} \varphi^{\Delta}$: The argument is identical to the argument for $\operatorname{Im} \varphi$ once we proved that $\operatorname{Im} \varphi = \ker(\mathfrak{G} \to \operatorname{Coker} \varphi)$ is adically quasi-coherent.

Corollary 4.4.15. Let $\mathfrak{X} = \text{Spf } A$ be a topologically finitely presented affine formal *R*-scheme for *R* as in the Setup 4.4.1, let *M* an almost finitely generated *A*-module, and let *N* be any *A*-submodule of *M*. Then the following sequence

$$0 \to N^{\Delta} \xrightarrow{\varphi^{\Delta}} M^{\Delta} \to (M/N)^{\Delta} \to 0$$

is exact.

Proof. We just apply Lemma 4.4.14 to the homomorphism $M \to M/N$ of almost finitely generated A-modules.

Corollary 4.4.16. Let \mathfrak{X} be a topologically finitely presented formal *R*-scheme for *R* as in the Setup 4.4.1, and let $\varphi: \mathfrak{F} \to \mathfrak{G}$ be a morphism of adically quasi-coherent, almost finite type $\mathcal{O}_{\mathfrak{X}}$ -modules. Then ker φ , Coker φ , and Im φ are adically quasi-coherent $\mathcal{O}_{\mathfrak{X}}$ -modules.

Corollary 4.4.17. Let \mathfrak{X} be a topologically finitely presented formal *R*-scheme for *R* as in the Setup 4.4.1, and let $\varphi \colon \mathfrak{F}^a \to \mathfrak{G}^a$ be a morphism of almost almost finite type $\mathcal{O}^a_{\mathfrak{X}}$ -modules. Then ker φ , Coker φ and Im φ are almost adically quasi-coherent $\mathcal{O}^a_{\mathfrak{X}}$ -modules.

Proof. We apply the exact functor $(-)_{!}$ to the map φ and reduce the claim to Corollary 4.4.16.

Now we are ready to show that almost finite type and almost finitely presented $\mathcal{O}_{\mathfrak{X}}$ -modules share many good properties we would expect. The only subtle thing is that we do not know whether an adically quasi-coherent quotient of an adically quasi-coherent, almost finite type $\mathcal{O}_{\mathfrak{X}}$ -module is of almost finite type. The main extra complication here is that we need to be very careful with the adically quasi-coherent condition in the definition of almost finite type (resp. almost finitely presented) modules since that condition does not behave well in general.

Lemma 4.4.18. Let $0 \to \mathcal{F}' \xrightarrow{\varphi} \mathcal{F} \xrightarrow{\psi} \mathcal{F}'' \to 0$ be an exact sequence of $\mathcal{O}_{\mathfrak{X}}$ -modules, then

- (1) If \mathcal{F} is adically quasi-coherent of almost finite type, and \mathcal{F}' is adically quasi-coherent then \mathcal{F}'' is adically quasi-coherent of almost finite type.
- (2) If \$\mathcal{F}'\$ and \$\mathcal{F}''\$ are adically quasi-coherent of almost finite type (resp. almost finitely presented) then so is \$\mathcal{F}\$.
- (3) If \mathcal{F} is adically quasi-coherent of almost finite type and \mathcal{F}'' is adically quasi-coherent almost finitely presented then \mathcal{F}' is adically quasi-coherent of almost finite type.
- (4) If \mathcal{F} is adically quasi-coherent of almost finitely presented and \mathcal{F}' is adically quasi-coherent of almost finite type then \mathcal{F}'' is adically quasi-coherent, almost finitely presented.

Proof. (1): Without loss of generality, we can assume that $\mathfrak{X} = \operatorname{Spf} A$ is an affine formal scheme. Then $\mathfrak{F} \cong M^{\Delta}$ for some almost finitely generated A-module M, and $\mathfrak{F}' \cong N^{\Delta}$ for some A-submodule $N \subset M$. Then Corollary 4.4.15 ensures that $\mathfrak{F}'' \simeq (M/N)^{\Delta}$. In particular, it is adically quasicoherent. Then the claim is an easy consequence of the vanishing theorem [FK18, Theorem I.7.1.1] and Lemma 2.5.11(1).

(2): The difficult part is to show that \mathcal{F} is adically quasi-coherent. In fact once we know that \mathcal{F} is adically quasi-coherent, it is automatically of almost finite type (resp. almost finitely presented) by [FK18, Theorem I.7.1.1] and Lemma 2.5.11(2).

In order to show that \mathcal{F} is adically quasi-coherent, we may and do assume that $\mathfrak{X} = \operatorname{Spf} A$ is an affine formal *R*-scheme for some adhesive ring *A*. Then let us introduce *A*-modules M' := $\mathcal{F}'(\mathfrak{X}), M := \mathcal{F}(\mathfrak{X}), \text{ and } M'' := \mathcal{F}''(\mathfrak{X})$. We have the natural morphism $M^{\Delta} \to \mathcal{F}$ and we show that it is an isomorphism. The vanishing theorem [FK18, Theorem I.7.1.1] implies that we have a short exact sequence:

$$0 \to M' \to M \to M'' \to 0$$

Thus M is almost finitely generated (resp. almost finitely presented) by Lemma 2.5.11(2). Then Corollary 4.4.14 gives that we have a short exact sequence

$$0 \to M'^{\Delta} \to M^{\Delta} \to M''^{\Delta} \to 0$$

Using the vanishing theorem [FK18, Theorem I.7.1.1] once again we get a commutative diagram

where the rows are exact, and left and right vertical arrows are isomorphisms. That implies that the map $M^{\Delta} \to \mathcal{F}$ is an isomorphism.

- (3): This easily follows from Lemma 2.5.11(3), Lemma 4.4.16 and [FK18, Theorem I.7.1.1].
- (4): This also easily follows from Lemma 2.5.11(4), Lemma 4.4.16 and [FK18, Theorem I.7.1.1].

We also give the almost version of this Lemma:

Corollary 4.4.19. Let $0 \to \mathcal{F}'^a \xrightarrow{\varphi} \mathcal{F}^a \xrightarrow{\psi} \mathcal{F}''^a \to 0$ be an exact sequence of $\mathcal{O}^a_{\mathfrak{X}}$ -modules, then

(1) If \mathcal{F}^a is of almost finite type, and \mathcal{F}'^a is almost adically quasi-coherent then \mathcal{F}''^a is of almost finite type.

- (2) If $\mathcal{F}^{\prime a}$ and $\mathcal{F}^{\prime \prime a}$ are of almost finite type (resp. almost finitely presented) then so is \mathcal{F}^{a} .
- (3) If \$\mathcal{F}^a\$ is of almost finite type and \$\mathcal{F}''^a\$ is almost finitely presented then \$\mathcal{F}'^a\$ is of almost finite type.
- (4) If \mathcal{F}^a is of almost finitely presented and \mathcal{F}'^a is of almost finite type then \mathcal{F}''^a is almost finitely presented.

Definition 4.4.20. We say that an $\mathcal{O}^a_{\mathfrak{X}}$ -module \mathcal{F}^a is *almost coherent* if \mathcal{F}^a is almost finite type and for any open set \mathfrak{U} any finite type $\mathcal{O}^a_{\mathfrak{X}}$ -submodule $\mathcal{G}^a \subset (\mathcal{F}|_{\mathfrak{U}})^a$ is an almost finitely presented $\mathcal{O}_{\mathfrak{U}}$ -module.

We say that \mathcal{F} is *(adically quasi-coherent) almost coherent* $\mathcal{O}_{\mathfrak{X}}$ -module if \mathcal{F}^a almost coherent (and \mathcal{F} is adically quasi-coherent).

Remark 4.4.21. Lemma 4.4.10 ensures that any adically quasi-coherent almost coherent \mathcal{O}_X -module \mathcal{F} is almost coherent.

Lemma 4.4.22. Let \mathcal{F}^a be an $\mathcal{O}^a_{\mathfrak{X}}$ -module on a topologically finitely presented formal *R*-scheme \mathfrak{X} . Then the following are equivalent:

- (1) \mathcal{F}^a is almost coherent.
- (2) \mathcal{F}^a is almost quasi-coherent and the $\mathcal{O}^a_{\mathfrak{X}}(\mathfrak{U})$ -module $\mathcal{F}^a(\mathfrak{U})$ is almost coherent for any open affine formal subscheme $\mathfrak{U} \subset \mathfrak{X}$.
- (3) \mathcal{F}^a is almost quasi-coherent and there is a covering of \mathfrak{X} by open affine subschemes $(\mathfrak{U}_i)_{i\in I}$ such that $\mathcal{F}^a(\mathfrak{U}_i)$ is almost coherent for each *i*.

In particular, an $\mathcal{O}^a_{\mathfrak{F}}$ -module \mathfrak{F}^a is almost coherent if and only if it almost finitely presented.

Proof. The proof that these three notions are equivalent is identical to the proof of Lemma 4.4.22 modulo facts that we have already established in this chapter, especially Corollary 4.4.14.

As for the last claim, we recall that \mathfrak{X} is topologically finitely presented over a topologically universally adhesive ring, so $\mathcal{O}_{\mathfrak{X}}(\mathfrak{U})$ is coherent for any open affine \mathfrak{U} [FK18, Prop. 0.8.5.23, Lemma I.1.7.4, Prop. I.2.3.3]. Then Lemma 2.6.11 and Lemma 2.6.13 prove the equivalence.

Even though Lemma 4.4.22 says that the notion of almost coherence coincides with the notion of almost finite presentation, it shows that almost coherence is morally "the correct" definition. In what follows, we prefer to use the terminology of almost coherent sheaves as it is shorter and gives a better intuition from our point of view.

Lemma 4.4.23. (1) Any almost finite type $\mathcal{O}^a_{\mathfrak{X}}$ -submodule of an almost coherent $\mathcal{O}^a_{\mathfrak{X}}$ -module is almost coherent.

- (2) Let $\varphi \colon \mathcal{F}^a \to \mathcal{G}^a$ be a homomorphism from an almost finite type $\mathcal{O}^a_{\mathfrak{X}}$ -module to an almost coherent $\mathcal{O}^a_{\mathfrak{X}}$ -module. Then ker φ is an almost finite type $\mathcal{O}^a_{\mathfrak{X}}$ -module.
- (3) Let $\varphi \colon \mathcal{F}^a \to \mathcal{G}^a$ be a homomorphism of almost coherent $\mathcal{O}^a_{\mathfrak{X}}$ -modules. Then ker φ and Coker φ are almost coherent $\mathcal{O}_{\mathfrak{X}}$ -modules.
- (4) Given a short exact sequence of $\mathcal{O}^a_{\mathfrak{X}}$ -modules

$$0 \to \mathcal{F}^{\prime a} \to \mathcal{F}^{a} \to \mathcal{F}^{\prime\prime a} \to 0,$$

if two out of three are almost coherent so is the third one.

Remark 4.4.24. There is also an evident version of this corollary for adically quasi-coherent almost coherent $\mathcal{O}_{\mathfrak{X}}$ -modules.

Proof. The proof is identical to Corollary 4.1.12 once we have Lemma 4.4.16 and equivalence of almost coherent and almost finitely presented $O_{\mathfrak{X}}$ -modules from Lemma 4.4.22.

Corollary 4.4.25. Let \mathfrak{X} be topologically finitely presented formal *R*-scheme for *R* as in the Setup 4.4.1. Then the category $\mathbf{Qcoh}_{\mathfrak{X}}^{\mathrm{acoh}}$ (resp. $\mathbf{AQcoh}_{\mathfrak{X}}^{\mathrm{acoh}}$) of adically quasi-coherent almost coherent $\mathcal{O}_{\mathfrak{X}}$ -modules (resp. $\mathcal{O}_{\mathfrak{X}}^{a}$ -modules) is a Weak Serre subcategory of $\mathbf{Mod}_{\mathfrak{X}}$ (resp. $\mathbf{Mod}_{\mathfrak{X}}^{a}$).

4.5. Formal Schemes. Basic Functors on Almost Coherent $\mathcal{O}^a_{\mathfrak{X}}$ -modules. This section is devoted to study how certain functors defined in Section 3.2 interact with the notions of almost (quasi-)coherent $\mathcal{O}^a_{\mathfrak{X}}$ -modules. The exposition follows Section 4.2 very closely.

We start with the affine situation, i.e. $\mathfrak{X} = \operatorname{Spf} A$. In this case, we note that the functor $(-)^{\Delta} \colon \operatorname{Mod}_A \to \operatorname{Qcoh}_{\mathfrak{X}}$ sends almost zero A-modules to almost zero $\mathcal{O}_{\mathfrak{X}}$ -modules. Thus, it induces a functor

$$(-)^{\Delta} \colon \mathbf{Mod}^a_A \to \mathbf{AQcoh}_{\mathfrak{X}}$$
.

Lemma 4.5.1. Let $\mathfrak{X} = \operatorname{Spf} A$ be an affine formal *R*-scheme for *R* as in the Setup 4.4.1. Then the functor $(-)^{\Delta} \colon \operatorname{Mod}_A \to \operatorname{Qcoh}_{\mathfrak{X}}$ induces an equivalence $(-)^{\Delta} \colon \operatorname{Mod}_A^* \to \operatorname{Qcoh}_{\mathfrak{X}}^*$ for any $* \in \{\operatorname{aft}, \operatorname{acoh}\}$. The quasi-inverse functor is given by $\Gamma(\mathfrak{X}, -)$.

Proof. We note that the functor $(-)^{\Delta}$: $\operatorname{Mod}_A \to \operatorname{Qcoh}_{\mathfrak{X}}$ induces an equivalence between the category of *I*-adically complete *A*-modules and adically quasi-coherent $\mathcal{O}_{\mathfrak{X}}$ -modules by [FK18, Theorem I.3.2.8]. Recall that all almost finite type modules are complete by Lemma 2.8.7. Thus it suffices to show that an adically quasi-coherent $\mathcal{O}_{\mathfrak{X}}$ -module is almost finitely generated (resp. almost coherent) if and only if so is $\Gamma(X, \mathcal{F})$. Now this follows from Lemma 4.4.13 and Lemma 4.4.22.

Lemma 4.5.2. Let $\mathfrak{X} = \operatorname{Spf} A$ be an affine formal *R*-scheme for *R* as in the Setup 4.4.1. Then the functor $(-)^{\Delta} \colon \operatorname{Mod}_A \to \operatorname{Qcoh}_{\mathfrak{X}}$ induces equivalences $(-)^{\Delta} \colon \operatorname{Mod}_A^{a,*} \to \operatorname{AQcoh}_{\mathfrak{X}}^*$ for any $* \in \{\operatorname{aft}, \operatorname{acoh}\}$. The quasi-inverse functor is given by $\Gamma(X, -)$.

Proof. The proof is analogous to Lemma 4.2.2 once Lemma 4.5.1 is verified.

Now recall that for any *R*-scheme *X*, we can define the *I*-adic completion of *X* as a colimit $\operatorname{colim}(X_k, \mathcal{O}_{X_k})$ of the reductions $X_k \coloneqq X \times_R \operatorname{Spec} R/I^{k+1}$ in the category of formal schemes. We refer to [FK18, §1.4(c)] for more details. This completion comes with a map of locally ringed spaces

$$c\colon \widehat{X} \to X$$
.

One important example of a completion is $\widehat{\operatorname{Spec} A} = \operatorname{Spf} \widehat{A}$ for any *R*-algebra A^{24} . We study the properties of the completion map in the case of a finitely presented *R*-scheme or an affine scheme Spec *A* for a topologically finitely presented *R*-algebra *A*.

Lemma 4.5.3. Let X = Spec A be an affine R-scheme for R as in the Setup 4.4.1. Suppose that A is either finitely presented or topologically finitely presented over R. Then the morphism $c: \widehat{X} \to X$ is flat and there is a functorial isomorphism $M^{\Delta} \cong c^*(\widetilde{M})$ for any almost finitely generated A-module M.

Proof. The flatness assertion is just [FK18, Proposition I.1.4.7 (2)]. The natural map

$$M \to \mathrm{H}^0(\mathfrak{X}, c^*(\tilde{M}))$$

induces the map $M^{\Delta} \to c^*(\widetilde{M})$. In order to show that it is an isomorphism, it is enough to show that the map

$$\widehat{M_f} \to M_f \otimes_{A_f} \widehat{A_f}$$

²⁴We note that \widehat{A} is *I*-adically complete by [Sta21, Tag 05GG] since *I* is finitely generated.

is an isomorphism for any $f \in A$. This follows from Lemma 2.8.7, as each such A_f is *I*-adically adhesive.

Corollary 4.5.4. Let X be a locally finitely presented R-scheme for R as in the Setup 4.4.1. Then the morphism $c: \widehat{X} \to X$ is flat and c^* sends almost finite type \mathcal{O}_X^a -modules (resp. almost coherent \mathcal{O}_X^a -modules) to almost finite type $\mathcal{O}_{\mathfrak{X}}^a$ -modules (resp. almost coherent $\mathcal{O}_{\mathfrak{X}}^a$ -modules).

Similarly, c^* sends quasi-coherent almost finite type \mathcal{O}_X -modules (resp. quasi-coherent almost coherent \mathcal{O}_X -modules) to adically quasi-coherent almost finite type $\mathcal{O}_{\mathfrak{X}}$ -modules (resp. adically quasi-coherent almost coherent $\mathcal{O}_{\mathfrak{X}}$ -modules)

Proof. The statement is local, so we can assume that X = Spec A. Then the claim follows from Lemma 4.5.3.

Now we show that the pullback functor preserves almost finite type and almost coherent $\mathcal{O}^a_{\mathfrak{X}}$ -modules.

Lemma 4.5.5. Let $\mathfrak{f}: \mathfrak{X} \to \mathfrak{Y}$ be a morphism of locally finitely presented formal *R*-scheme for *R* as in the Setup 4.4.1.

- (1) Suppose that $\mathfrak{X} = \operatorname{Spf} B$, $\mathfrak{Y} = \operatorname{Spf} A$ are affine formal *R*-schemes. Then $\mathfrak{f}^*(M^{\Delta})$ is functorially isomorphic to $(M \otimes_A B)^{\Delta}$ for any $M \in \operatorname{\mathbf{Mod}}_A^{\operatorname{aft}}$.
- (2) Suppose that $\mathfrak{X} = \operatorname{Spf} B, \mathfrak{Y} = \operatorname{Spf} A$ are affine formal *R*-schemes. Then $\mathfrak{f}^*(M^{a,\Delta})$ is functorially isomorphic to $(M^a \otimes_{A^a} B^a)^{\Delta}$ for any $M^a \in \operatorname{\mathbf{Mod}}_A^{a,\operatorname{aff}}$.
- (3) The functor f^* sends $\operatorname{Qcoh}_{\mathfrak{Y}}^{\operatorname{aft}}$ (resp. $\operatorname{Qcoh}_{\mathfrak{Y}}^{\operatorname{acoh}}$) to $\operatorname{Qcoh}_{\mathfrak{X}}^{\operatorname{aft}}$ (resp. $\operatorname{Qcoh}_{\mathfrak{X}}^{\operatorname{acoh}}$).
- (4) The functor f^* sends $\mathbf{AQcoh}_{\mathfrak{Y}}^{\mathrm{aft}}$ (resp. $\mathbf{AQcoh}_{\mathfrak{Y}}^{\mathrm{acoh}}$) to $\mathbf{Qcoh}_{\mathfrak{X}}^{\mathrm{aft}}$ (resp. $\mathbf{Qcoh}_{\mathfrak{X}}^{\mathrm{acoh}}$).

Proof. We prove (1), the proofs of other parts follow from it similarly to the proof Lemma 4.2.3. We consider a commutative diagram

$$\begin{array}{ccc} \operatorname{Spf} B & \stackrel{c_B}{\longrightarrow} & \operatorname{Spec} B \\ & & & & \downarrow^f \\ \operatorname{Spf} A & \stackrel{c_A}{\longrightarrow} & \operatorname{Spec} A \end{array}$$

where the map $f: \operatorname{Spec} B \to \operatorname{Spec} A$ is the map induced by $\mathfrak{f}^{\#}: A \to B$. Then we have that $M^{\Delta} \simeq c_{A}^{*} \widetilde{M}$ by Lemma 4.5.3. Therefore,

$$\mathfrak{f}^*(M^{\Delta}) \simeq c_B^*(f^*\widetilde{M}) \simeq c_B^*(\widetilde{M \otimes_A B}) \simeq (M \otimes_A B)^{\Delta}$$

where the last isomorphism follows from Lemma 4.5.3 again.

The next thing we discuss is how the finiteness properties interact with tensor products.

Lemma 4.5.6. Let \mathfrak{X} be a topologically finitely presented formal *R*-scheme for *R* as in the Setup 4.4.1.

- (1) Suppose that $\mathfrak{X} = \operatorname{Spf} A$ is affine. Then $M^{\Delta} \otimes_{\mathfrak{O}_{\mathfrak{X}}} N^{\Delta}$ is functorially isomorphic to $(M \otimes_A N)^{\Delta}$ for any $M, N \in \operatorname{\mathbf{Mod}}_{A}^{\operatorname{aft}}$.
- (2) Suppose that $\mathfrak{X} = \operatorname{Spf} A$ is affine. Then $M^{a,\Delta} \otimes_{\mathcal{O}_{\mathfrak{X}}^a} N^{a,\Delta}$ is functorially isomorphic to $(M^a \otimes_{A^a} N^a)^{\Delta}$ for any $M^a, N^a \in \operatorname{\mathbf{Mod}}_A^{a,\operatorname{aff}}$.

- (3) Let \mathcal{F}, \mathcal{G} be two adically quasi-coherent almost finite type (resp. almost finitely presented) $\mathcal{O}_{\mathfrak{X}}$ -modules. Then the \mathcal{O}_X -module $\mathcal{F} \otimes_{\mathcal{O}_{\mathfrak{X}}} \mathcal{G}$ is adically quasi-coherent of almost finite type (resp. almost finitely presented).
- (4) Let $\mathcal{F}^a, \mathcal{G}^a$ be two almost finite type (resp. almost coherent) $\mathcal{O}^a_{\mathfrak{X}}$ -modules. Then the $\mathcal{O}^a_{\mathfrak{X}}$ -module $\mathcal{F}^a \otimes_{\mathcal{O}^a_{\mathfrak{X}}} \mathcal{G}^a$ is of almost finite type (resp. almost coherent). The analogous result holds for \mathcal{O}_X -modules \mathcal{F}, \mathcal{G} .

Proof. Again, we only show (1) as the other parts follow from this similarly to the proof of Lemma 4.2.4 with the simplification that almost coherent and almost finitely presented modules coincide by our assumption on \mathfrak{X} and R.

The proof of (1) is, in turn, similar to that of Lemma 4.5.5 (1). We consider the completion morphism $c: \operatorname{Spf} A \to \operatorname{Spec} A$. Then we have a sequence of isomorphisms

$$M^{\Delta} \otimes_{\mathcal{O}_{\mathfrak{X}}} N^{\Delta} \simeq c^{*}(\widetilde{M}) \otimes_{\mathcal{O}_{\mathfrak{X}}} c^{*}(\widetilde{N}) \simeq c^{*}(\widetilde{M} \otimes_{\mathcal{O}_{\operatorname{Spec} A}} \widetilde{N}) \simeq c^{*}(\widetilde{M} \otimes_{A} N) \simeq (M \otimes_{A} N)^{\Delta}.$$

Finally, we deal with the functor $\underline{\mathcal{H}om}_{\mathcal{O}_X^a}(-,-)$. This is probably the most subtle functor considered in this section. We start with the following preparatory lemma:

Lemma 4.5.7. Let \mathfrak{X} be a locally topologically finitely presented formal *R*-scheme for *R* as in the Setup 4.4.1.

(1) Suppose $\mathfrak{X} = \operatorname{Spf} A$ is affine. Then the canonical map

$$\operatorname{Hom}_{A}(M, N)^{\Delta} \to \underline{\mathcal{H}om}_{\mathcal{O}_{\mathfrak{X}}}(M^{\Delta}, N^{\Delta}) \tag{4.4}$$

is an almost isomorphism for any almost coherent A-modules M and N.

(2) Suppose $\mathfrak{X} = \operatorname{Spf} A$ is affine. Then there is a functorial isomorphism

$$\mathrm{alHom}_{A^{a}}(M^{a}, N^{a})^{\Delta} \simeq \underline{alHom}_{\mathcal{O}_{\mathfrak{X}}^{a}}(M^{a,\Delta}, N^{a,\Delta})$$

$$(4.5)$$

for any almost coherent A^a -module M^a and N^a . We also get a functorial almost isomorphism

$$\operatorname{Hom}_{A^{a}}(M^{a}, N^{a})^{\Delta} \simeq^{a} \underline{\operatorname{Hom}}_{\mathcal{O}_{\mathfrak{X}}^{a}}(M^{a,\Delta}, N^{a,\Delta})$$

$$(4.6)$$

for any almost coherent A^a -module M^a and N^a .

- (3) Suppose \mathcal{F} and \mathcal{G} are almost coherent $\mathcal{O}_{\mathfrak{X}}$ -modules. Then $\underline{\mathcal{H}om}_{\mathcal{O}_{\mathfrak{X}}}(\mathcal{F},\mathcal{G})$ is an almost coherent $\mathcal{O}_{\mathfrak{X}}$ -module.
- (4) Suppose \mathcal{F}^a and \mathcal{G}^a are almost coherent $\mathcal{O}^a_{\mathfrak{F}}$ -modules. Then

$$\underline{\mathcal{H}om}_{\mathcal{O}_{\mathfrak{X}}^{a}}(\mathcal{F}^{a}, \mathcal{G}^{a}) \text{ (resp. } \underline{al\mathcal{H}om}_{\mathcal{O}_{\mathfrak{X}}^{a}}(\mathcal{F}^{a}, \mathcal{G}^{a}))$$

is an almost coherent $\mathcal{O}_{\mathfrak{X}}$ -module (resp. \mathcal{O}_{X}^{a} -module).

Proof. Again, the proof is absolutely analogous to Lemma 4.2.6 and Corollary 4.2.7 once (1) is proven. So we only give a proof of (1) here.

We note that both M and N are I-adically complete by Lemma 2.8.7. Now we use [FK18] to say that the natural map $\operatorname{Hom}_A(M, N) \to \operatorname{Hom}_{\mathcal{O}_{\mathfrak{X}}}(M^{\Delta}, N^{\Delta})$ is an isomorphism. This induces a morphism

$$\operatorname{Hom}_A(M,N)^{\Delta} \to \underline{\mathcal{H}om}_{\mathcal{O}_{\mathfrak{X}}}(M^{\Delta},N^{\Delta})$$

In order to prove that it is an almost isomorphism, it suffices to show that the natural map

 $\operatorname{Hom}_{A}(M,N)\widehat{\otimes}_{A}A_{\{f\}} \to \operatorname{Hom}_{A_{\{f\}}}(M\widehat{\otimes}_{A}A_{\{f\}},N\widehat{\otimes}_{A}A_{\{f\}})$

is an almost isomorphism for any $f \in A$. Now we note that $\operatorname{Hom}_A(M, N)$ is almost coherent by Corollary 2.6.8. Thus, $\operatorname{Hom}_A(M, N) \otimes_A A_{\{f\}}$ is already complete, so the completed tensor product coincides with the usual one. Similarly, $M \widehat{\otimes}_A A_{\{f\}} \simeq M \otimes_A A_{\{f\}}$ and $N \widehat{\otimes}_A A_{\{f\}} \simeq N \otimes_A A_{\{f\}}$. Therefore, the question boils down to showing that the natural map

$$\operatorname{Hom}_{A}(M, N) \otimes_{A} A_{\{f\}} \to \operatorname{Hom}_{A_{\{f\}}}(M \otimes_{A} A_{\{f\}}, N \otimes_{A} A_{\{f\}})$$

is an almost isomorphism. This, in turn, follows from Lemma 2.7.8.

4.6. Formal Schemes. Derived Category of Almost Coherent $\mathcal{O}^a_{\mathfrak{X}}$ -modules. We discuss the notion of the derived category of almost coherent sheaves on a formal scheme \mathfrak{X} . One major issue is that there the derived category of $\mathcal{O}_{\mathfrak{X}}$ -modules with adically quasi-coherent cohomology sheaves is not well-defined, as adically quasi-coherent sheaves is not a Weak Serre subcategory of $\mathbf{Mod}_{\mathfrak{X}}$, so it is not even an abelian category. However, it would be useful for certain technical reasons to be able to work with that category.

In order to overcome this issue, we follow the strategy used in [Lur20] and define " $\mathbf{D}_{qc}(\mathfrak{X})$ " completely on the derived level without really defining a good abelian notion of (adically) quasicoherent sheaves. For the rest of the section, we fix a base ring R as in the Setup 4.4.1.

Definition 4.6.1. Let \mathfrak{X} be a locally topologically finitely presented *R*-scheme. Then we define the *derived category of adically quasi-coherent sheaves* " $\mathbf{D}_{qc}(\mathfrak{X})$ " as a full subcategory of $\mathbf{D}(\mathfrak{X})$ with elements \mathfrak{F} such that

- For every open affine $\mathfrak{U} \subset \mathfrak{X}$, $\mathbf{R}\Gamma(\mathfrak{U}, \mathfrak{F}) \in \mathbf{D}(\mathcal{O}_{\mathfrak{X}}(\mathfrak{U}))$ is derived *I*-adically complete.
- For every inclusion $\mathfrak{U} \subset \mathfrak{V}$ of affine formal subschemes of \mathfrak{X} , the natural morphism

$$\mathbf{R}\Gamma(\mathfrak{V},\mathfrak{F})\widehat{\otimes}_{\mathfrak{O}_{\mathfrak{X}}(\mathfrak{V})}^{L}\mathfrak{O}_{\mathfrak{X}(\mathfrak{U})}\to\mathbf{R}\Gamma(\mathfrak{U},\mathfrak{F})$$

is an isomorphism, where the completion is understood in the derived sense.

Remark 4.6.2. We refer to [Sta21, Tag 091N] and [Sta21, Tag 0995] for a self-contained discussion of the derived completion of modules and sheaves of modules respectively.

We want to give an interpretation of " $\mathbf{D}_{qc}(\mathfrak{X})$ " in terms of A-modules for an affine formal scheme $\mathfrak{X} = \operatorname{Spf} A$. We recall that in the case of schemes, we have a natural equivalence $\mathbf{D}_{qc}(\operatorname{Spec} A) \simeq \mathbf{D}(A)$ and the map is induced by $\mathbf{R}\Gamma(\operatorname{Spec} A, -)$. In the case of formal schemes, it is not literally true. We need to impose certain completeness conditions.

Definition 4.6.3. Let A be a ring with a finitely generated ideal I. We define the $\mathbf{D}_{comp}(A, I) \subset \mathbf{D}(A)$ as a full triangulated subcategory consisting of I-adically derived complete objects.

Suppose now that $\mathfrak{X} = \operatorname{Spf} A$ be an affine scheme, topologically finitely presented over R. We note that the natural functor $\mathbf{R}\Gamma(\mathfrak{X}, -) \colon \mathbf{D}(\mathfrak{X}) \to \mathbf{D}(A)$ induces a functor

$$\mathbf{R}\Gamma(\mathfrak{X},-)$$
: " $\mathbf{D}_{qc}(\mathfrak{X})$ " $\to \mathbf{D}_{comp}(A,I)$

The main claim is that this functor is an equivalence. This is the main content of [Lur20, Corollary 8.2.4.15]. We need to prove one technical result to ensure that our definitions are consistent with the definitions in Lurie's book.

Lemma 4.6.4. Let A be a topologically finitely presented R-algebra for R as in the Setup 4.4.1, let $f \in A$ be any element, and let $(x_1, \ldots, x_d) = I$ be a choice of generators for the ideal of definition of R. Denote by $K(A_f; x_1^n, \ldots, x_d^n)$ the Koszul complexes for the sequence (x_1^n, \ldots, x_d^n) . Then the pro-systems $\{K(A_f; x_1^n, \ldots, x_d^n)\}$ and $\{A_f/I^n\}$ are isomorphic in $Pro(\mathbf{D}(A_f))$.

Proof. The proof is the same [Sta21, Tag 0921]. The only difference that one needs to use [FGK11, Theorem 4.2.2(2)(b)] in place of the usual the Artin-Rees lemma.

Lemma 4.6.5. Let A be a topologically finitely presented R-algebra for R as in the Setup 4.4.1, let $f \in A$ be any element. Then the completed localization $A_{\{f\}}$ coincides with the I-adic derived completion of A_f .

Proof. Choose some generators $I = (x_1, \ldots, x_d)$. Then we know that the derived completion completion of A_f is given by $\mathbf{R} \lim_n K(A_f; x_1^n, \ldots, x_d^n)$ where $K(A_f; x_1^n, \ldots, x_d^n)$ is the Koszul complex for the sequence (x_1^n, \ldots, x_d^n) . Lemma 4.6.4 implies that the pro-systems $\{K(A_f; x_1^n, \ldots, x_d^n)\}$ and $\{A_f/I^n\}$ are naturally pro-isomorphic. Thus we have an isomorphism

$$\mathbf{R}\lim_{n} K(A_{f}; x_{1}^{n}, \dots, x_{d}^{n}) \cong \mathbf{R}\lim_{n} A_{f}/I^{n} \simeq A_{\{f\}} .$$

The last isomorphism uses the Mittag-Leffler criterion to ensure vanishing of lim¹.

Theorem 4.6.6. [Lur20, Corollary 8.2.4.15] Let $\mathfrak{X} = \text{Spf } A$ be an affine, finitely presented formal scheme over R as in the Setup 4.4.1. Then functor $\mathbf{R}\Gamma(\mathfrak{X}, -)$: " $\mathbf{D}_{qc}(\mathfrak{X})$ " $\to \mathbf{D}_{comp}(A, I)$ is an equivalence of categories.

Proof. The statement can be deduced from [Lur20, Corollary 8.2.4.15] by applying π_0 to it. We note that even though [Lur20, Corollary 8.2.4.15] uses ∞ -categories, the cited proof can be rephrased in our situation without using any derived geometry. However, it would require quite a big digression, so instead we explain why our definitions are compatible with definitions in [Lur20].

Lemma 4.6.5 implies that the definition of Spf A in [Lur20] is compatible with the classical one. Now [Lur20, Proposition 8.2.4.18] ensures that our definition of " $\mathbf{D}_{qc}(\mathfrak{X})$ " is equivalent to $\pi_0(\operatorname{Qcoh}(\mathfrak{X}))$ in the sense of [Lur20].

Definition 4.6.7. We denote by

$$(-)^{\Delta} \colon \mathbf{D}_{comp}(A, I) \to \mathbf{D}_{qc}(\mathfrak{X})^{"}$$

the pseudo-inverse to $\mathbf{R}\Gamma(\mathfrak{X}, -)$: " $\mathbf{D}_{qc}(\mathfrak{X})$ " $\to \mathbf{D}_{comp}(A, I)$. We note that clearly $\mathbf{R}\Gamma(\operatorname{Spf} A_{\{f\}}, M^{\Delta}) \simeq M \widehat{\otimes}_A A_{\{f\}}$ for any $M \in \mathbf{D}_{comp}(A, I)$.

Remark 4.6.8. This functor $(-)^{\Delta}$ is not compatible with the one used the previous sections. We will emphasize which version we use whenever it may cause any confusion.

Our real goal is to show that there is an equivalence between $\mathbf{D}_{acoh}(A)$ and $\mathbf{D}_{qc,acoh}(\mathfrak{X})$. Theorem 4.6.6 will be a useful tool to prove this equivalence. We now give a precise definition of $\mathbf{D}_{qc,acoh}(\mathfrak{X})$.

Definition 4.6.9. We define $\mathbf{D}_{qc,acoh}(\mathfrak{X})$ (resp. $\mathbf{D}_{acoh}(\mathfrak{X})^a$) to be the full triangulated subcategory of $\mathbf{D}(\mathfrak{X})$ (resp. $\mathbf{D}(\mathfrak{X})^a$) consisting of complexes with adically quasi-coherent, almost coherent (resp. almost coherent) cohomology sheaves (resp. almost sheaves).

Remark 4.6.10. An argument similar to one in the proof of Lemma 4.3.5 shows that $\mathbf{D}_{acoh}(\mathfrak{X})^a$ is equivalent to the Verdier quotient $\mathbf{D}_{qc,acoh}(\mathfrak{X})/\mathbf{D}_{qc,\Sigma_{\mathfrak{X}}}(\mathfrak{X})$.

In order to show an equivalence $\mathbf{D}_{qc,acoh}(\mathfrak{X}) \simeq \mathbf{D}_{acoh}(A)$, our first goal is to show that $\mathbf{D}_{qc,acoh}$ lies inside " $\mathbf{D}_{qc}(\mathfrak{X})$ ". Even though it looks very plausible, it requires a proof that is not entirely trivial.

Lemma 4.6.11. Let $\mathfrak{X} = \operatorname{Spf} A$ be an affine topologically finitely presented formal *R*-scheme for R as in the Setup 4.4.1. Then the functor $\mathbf{R}\Gamma(\mathfrak{X}, -): \mathbf{D}_{qc,acoh}(\mathfrak{X}) \to \mathbf{D}(A)$ is *t*-exact (with respect to the evident *t*-structures on both sides) and factors through $\mathbf{D}_{acoh}(A)$. More precisely, there is an isomorphism

$$\mathrm{H}^{i}\left(\mathbf{R}\Gamma\left(\mathfrak{X},\mathfrak{F}\right)\right)\simeq\mathrm{H}^{0}\left(\mathfrak{X},\mathcal{H}^{i}\left(\mathfrak{F}\right)\right)\in\mathbf{Mod}_{A}^{\mathrm{acoh}}$$

for any object $\mathcal{F} \in \mathbf{D}_{qc,acoh}(\mathfrak{X})$.

Proof. We note that the vanishing theorem [FK18, Theorem I.7.1.1] implies that we can use [Sta21, Tag 0D6U] with N = 0. Thus we see that the map $\mathrm{H}^{i}(\mathbf{R}\Gamma(\mathfrak{X}, \mathcal{F})) \to \mathrm{H}^{i}(\mathbf{R}\Gamma(\mathfrak{X}, \tau^{\geq i}\mathcal{F}))$ is an isomorphism for any integer i, and that $\mathbf{R}\Gamma(\mathfrak{X}, \mathcal{F}) \in \mathbf{D}_{acoh}(A)$ for any $\mathcal{F} \in \mathbf{D}_{qc,acoh}(\mathfrak{X})$. Combining it with the canonical isomorphism $\mathrm{H}^{i}(\mathbf{R}\Gamma(\mathfrak{X}, \tau^{\geq i}\mathcal{F})) \simeq \mathrm{H}^{0}(\mathfrak{X}, \mathcal{H}^{i}(\mathcal{F}))$ we get the desired result. \Box

Lemma 4.6.12. Let \mathfrak{X} be an locally topologically finitely presented formal *R*-scheme for *R* as in the Setup 4.4.1. Then $\mathbf{D}_{qc,acoh}(\mathfrak{X})$ is naturally a full triangulated subcategory of " $\mathbf{D}_{qc}(\mathfrak{X})$ ".

Proof. Both $\mathbf{D}_{qc,acoh}(\mathfrak{X})$ and " $\mathbf{D}_{qc}(\mathfrak{X})$ " are full triangulated subcategories of $\mathbf{D}(\mathfrak{X})$. Thus, it suffices to show that any $\mathcal{F} \in \mathbf{D}_{qc,acoh}(\mathfrak{X})$ lies in " $\mathbf{D}_{qc}(\mathfrak{X})$ ".

Lemma 4.6.11 and Corollary 2.8.8 imply that $\mathbf{R}\Gamma(\mathfrak{U}, \mathfrak{F}) \in \mathbf{D}_{comp}(A, I)$ for any open affine $\mathfrak{U} \subset \mathfrak{X}$. Now suppose $\mathfrak{U} \subset \mathfrak{V}$ is an inclusion of open affine formal subschemes in \mathfrak{X} . We consider the natural morphism

$$\mathbf{R}\Gamma(\mathfrak{V},\mathfrak{F})\widehat{\otimes}_{\mathfrak{O}_{\mathfrak{X}}(\mathfrak{V})}^{L}\mathcal{O}_{\mathfrak{X}}(\mathfrak{U})\to\mathbf{R}\Gamma(\mathfrak{U},\mathfrak{F})$$

We note that $\mathcal{O}_{\mathfrak{X}}(\mathfrak{U})$ is flat over $\mathcal{O}_{\mathfrak{X}}(\mathfrak{V})$ by [FK18, Proposition I.4.8.1]. Thus, the complex

$$\mathbf{R}\Gamma(\mathfrak{V},\mathfrak{F})\otimes^{L}_{\mathfrak{O}_{\mathfrak{X}}(\mathfrak{V})}\mathfrak{O}_{\mathfrak{X}(\mathfrak{U})}$$

lies in $\mathbf{D}_{acoh}(\mathcal{O}_{\mathfrak{X}}(\mathfrak{U}))$ by Lemma 2.7.1. Therefore, it also lies in $\mathbf{D}_{comp}(A, I)$ by Corollary 2.8.8. So we conclude that

$$\mathbf{R}\Gamma(\mathfrak{V},\mathfrak{F})\widehat{\otimes}^{L}_{\mathfrak{O}_{\mathfrak{X}}(\mathfrak{V})}\mathcal{O}_{\mathfrak{X}}(\mathfrak{U})\simeq\mathbf{R}\Gamma(\mathfrak{V},\mathfrak{F})\otimes^{L}_{\mathfrak{O}_{\mathfrak{Y}}(\mathfrak{V})}\mathcal{O}_{\mathfrak{X}}(\mathfrak{U})$$

Using $\mathcal{O}_{\mathfrak{X}}(\mathfrak{V})$ -flatness of $\mathcal{O}_{\mathfrak{X}}(\mathfrak{U})$, we conclude that the question boils down to show that

$$\mathrm{H}^{i}(\mathfrak{V},\mathcal{F})\otimes_{\mathfrak{O}_{\mathfrak{X}}(\mathfrak{V})}\mathcal{O}_{\mathfrak{X}(\mathfrak{U})}\to\mathrm{H}^{i}(\mathfrak{U},\mathcal{F})$$

is an isomorphism for all i. Now Lemma 4.6.11 implies that this, in turn, reduces to showing that the natural map

$$\Gamma(\mathfrak{V},\mathcal{H}^{i}(\mathcal{F}))\otimes_{\mathcal{O}_{\mathfrak{X}}(\mathfrak{V})}\mathcal{O}_{\mathfrak{X}(\mathfrak{U})}\to\Gamma(\mathfrak{U},\mathcal{H}^{i}(\mathcal{F}))$$

is an isomorphism. Without loss of generality, we may assume that $\mathfrak{X} = \mathfrak{V} = \operatorname{Spf} A$. Then $\mathcal{H}^{i}(\mathfrak{F})$ is an adically quasi-coherent, almost coherent $\mathcal{O}_{\mathfrak{X}}$ -module, so it is isomorphic to M^{Δ} for some $M \in \operatorname{\mathbf{Mod}}_{A}^{\operatorname{acoh}}$ by Lemma 4.5.1. The $(-)^{\Delta}$ notation is used here in the classical sense. So the desired claim follows from [FK18, Lemma 3.6.4] and the observation that $M \otimes_{\mathcal{O}_{\mathfrak{X}}(\mathfrak{V})} \mathcal{O}_{\mathfrak{X}}(\mathfrak{U})$ is already *I*-adically complete by Lemma 2.8.7.

Now we show that $(-)^{\Delta}$ functor sends $\mathbf{D}_{acoh}(A)$ to $\mathbf{D}_{qc,acoh}(\operatorname{Spf} A)$. This is also not entirely obvious as this derived version of $(-)^{\Delta}$ a priori has nothing to do with the classical version of $(-)^{\Delta}$ -functor defined on classically *I*-adically complete modules. The key is to show that these functors coincide on $\mathbf{Mod}_{A}^{\operatorname{acoh}}$.

Lemma 4.6.13. Let $\mathfrak{X} = \text{Spf } A$ be an affine topologically finitely presented formal R-scheme for R as in the Setup 4.4.1. Then the functor $(-)^{\Delta} \colon \mathbf{D}_{acoh}(A) \to \mathbf{D}_{qc}(\mathfrak{X})$ " factors through $\mathbf{D}_{qc,acoh}(\mathfrak{X})$. Moreover, for any $M \in \mathbf{D}_{acoh}(A)$, there are functorial isomorphisms

$$\mathrm{H}^{i}(M)^{\Delta} \simeq \mathcal{H}^{i}(M^{\Delta})$$

where $(-)^{\Delta}$ should be understood in the derived sense on the right side and it the classical sense on the left side.

Proof. We note that $\mathrm{H}^{i}(\mathfrak{X}, M^{\Delta}) \simeq \mathrm{H}^{i}(M)$ by its very construction. As $\mathcal{H}^{i}(M^{\Delta})$ is canonically isomorphic to the sheafification of the presheaf

$$\mathfrak{U} \mapsto \mathrm{H}^{i}(\mathfrak{U}, M^{\Delta})$$

we get that there is a canonical map $\mathrm{H}^{i}(M) \to \Gamma(\mathfrak{X}, \mathcal{H}^{i}(M^{\Delta}))$. Thus the universal property of the classical $(-)^{\Delta}$ functor, we get a functorial morphism

$$\mathrm{H}^{i}(M)^{\Delta} \to \mathcal{H}^{i}(M^{\Delta})$$

Since $H^i(M)$ is almost coherent, it suffices to show that this map is an isomorphism for any *i*. This boils down (using almost coherence of $H^i(M)$) to show that

$$\mathrm{H}^{i}(M) \otimes_{A} A_{\{f\}} \to \mathrm{H}^{i}(\mathrm{Spf}\,A_{\{f\}}, M^{\Delta})$$

for all $f \in A$. Now recall that $\mathbf{R}\Gamma(\operatorname{Spf} A_{\{f\}}, M^{\Delta}) \simeq M \widehat{\otimes}_A^L A_{\{f\}}$ for any $f \in A$. Using that $M \in \mathbf{D}_{acoh}(A)$, $A_{\{f\}}$ is flat over A, and that almost coherent complexes are derived complete by Lemma 2.8.8, we conclude that the natural map

$$\mathrm{H}^{i}(M) \otimes_{A} A_{\{f\}} \to \mathrm{H}^{i}(\mathrm{Spf}\,A_{\{f\}}, M^{\Delta})$$

is an isomorphism finishing the proof.

Corollary 4.6.14. Let $\mathfrak{X} = \operatorname{Spf} A$ be an affine topologically finitely presented formal *R*-scheme for *R* as in the Setup 4.4.1. Suppose that $M \in \mathbf{D}(A)$ has almost zero cohomology modules. Then $\mathfrak{H}^{i}(M^{\Delta})$ is an almost zero, adically quasi-coherent $\mathcal{O}_{\mathfrak{X}}$ -module for all integer *i*. In particular, $(-)^{\Delta}$ induces the functor $(-)^{\Delta} \colon \mathbf{D}_{coh}(A)^{a} \to \mathbf{D}_{qc,acoh}(\mathfrak{X})^{a}$.

Proof. This follows directly from the observation that any almost zero A-modules is almost coherent and the formula $\mathrm{H}^{i}(M)^{\Delta} \simeq \mathcal{H}^{i}(M^{\Delta})$ established in Lemma 4.6.13.

Theorem 4.6.15. Let $\mathfrak{X} = \text{Spf } A$ be an affine topologically finitely presented formal R-scheme for R as in the Setup 4.4.1. Then the functor $\mathbf{R}\Gamma(\mathfrak{X}, -): \mathbf{D}_{qc,acoh}(\mathfrak{X}) \to \mathbf{D}_{acoh}(A)$ is *t*-exact equivalence of triangulated categories with the psedo-inverse $(-)^{\Delta}$.

Proof. Lemma 4.6.11 implies that $\mathbf{R}\Gamma(\mathfrak{X}, -)$ induces the functor $\mathbf{D}_{qc,acoh}(\mathfrak{X}) \to \mathbf{D}_{acoh}(A)$ and that this functor is *t*-exact. Lemma 4.6.12 and Theorem 4.6.6 ensures that it is sufficient to show that $(-)^{\Delta}$ sends $\mathbf{D}_{acoh}(A)$ to $\mathbf{D}_{qc,acoh}(\mathfrak{X})$, this follows from Lemma 4.6.13.

Now we can pass to the almost categories using Remark 4.6.10 to get the almost version of Theorem 4.6.15.

Corollary 4.6.16. Let $\mathfrak{X} = \operatorname{Spf} A$ be an affine topologically finitely presented formal *R*-scheme for *R* as in the Setup 4.4.1. Then the functor $\mathbf{R}\Gamma(\mathfrak{X}, -): \mathbf{D}_{acoh}(\mathfrak{X})^a \to \mathbf{D}_{acoh}(A)^a$ is *t*-exact equivalence of triangulated categories with the psedo-inverse $(-)^{\Delta}$.

4.7. Formal Schemes. Basic Functors on the Derived Categories of $\mathcal{O}^a_{\mathfrak{X}}$ -modules. We discuss the derived analogue of the results in Section 4.5. We show that the derived completion, derived tensor product, derived pullback, and derived almost Hom functors preserve complexes with almost coherent cohomology sheaves under certain conditions. For the rest of the section, we fix a ring R as in the Setup 4.4.1.

We start with the completion functor. We recall that we have defined the morphism of locally ringed spaces $c: \hat{X} \to X$ for any *R*-scheme *X*. If *X* is locally finitely presented over *R* or *X* = Spec *A*

for a topologically finitely presented R-algebra A, then c is a flat morphism as was shown in Lemma 4.5.3 and Corollary 4.5.4.

Lemma 4.7.1. Let $X = \operatorname{Spec} A$ be an affine R-scheme for R as in the Setup 4.4.1. Suppose that A is either finitely presented or topologically finitely presented over R. Suppose $M \in \mathbf{D}_{acoh}(A)$. Then $M^{\Delta} \simeq \mathbf{L}c^*(\widetilde{M})$.

Proof. First of all, we show that $\mathbf{L}c^*(\widetilde{M}) \in \mathbf{D}_{qc,acoh}(\widehat{X})$. Indeed, the functor c^* is exact as c is flat. Thus, Lemma 4.5.3 guarantees that we have a sequence of isomorphisms

$$\mathcal{H}^{i}\left(\mathbf{L}c^{*}\left(\widetilde{M}\right)\right)\simeq c^{*}\left(\widetilde{\mathrm{H}^{i}\left(M\right)}\right)\simeq\left(\mathrm{H}^{i}\left(M\right)\right)^{\Delta}\ .$$

In particular, Theorem 4.6.6 ensures that the natural morphism

$$M \simeq \mathbf{R}\Gamma(X, M) \to \mathbf{R}\Gamma(\widehat{X}, \mathbf{L}c^*(M))$$

induces the morphism $M^{\Delta} \to \mathbf{L}c^{*}(\widetilde{M})$. As c^{*} is exact, Lemma 4.6.13 implies that it is sufficient to show that the natural map

$$\mathrm{H}^{i}(M)^{\Delta} \to c^{*}(\widetilde{\mathrm{H}^{i}(M)})$$

is an isomorphism for all i. This follows from Lemma 4.5.3.

Corollary 4.7.2. Let X be a locally finitely presented R-scheme for R as in the Setup 4.4.1. Then $\mathbf{L}c^*$ induces functors $\mathbf{L}c^* \colon \mathbf{D}^*_{qc,acoh}(X) \to \mathbf{D}^*_{qc,acoh}(\widehat{X})$ (resp. $\mathbf{L}c^* \colon \mathbf{D}^*_{acoh}(X)^a \to \mathbf{D}^*_{acoh}(\widehat{X})^a$) for any $* \in \{ \text{```}, -, b, + \}$.

Proof. The claim is local, so it suffices to assume that X = Spec A. Then it follows from exactness of c^* and Lemma 4.7.1.

Lemma 4.7.3. Let $\mathfrak{f}: \mathfrak{X} \to \mathfrak{Y}$ be a morphism of locally finitely presented formal *R*-scheme for *R* as in the Setup 4.4.1.

- (1) Suppose that $\mathfrak{X} = \operatorname{Spf} B$, $\mathfrak{Y} = \operatorname{Spf} A$ are affine formal *R*-schemes. Then $L\mathfrak{f}^*(M^{\Delta})$ is functorially isomorphic to $(M \otimes_A B)^{\Delta}$ for any $M \in \mathbf{D}_{acoh}(A)$.
- (2) Suppose that $\mathfrak{X} = \operatorname{Spf} B$, $\mathfrak{Y} = \operatorname{Spf} A$ are affine formal *R*-schemes. Then $\mathbf{L}\mathfrak{f}^*(M^{a,\Delta})$ is functorially isomorphic to $(M^a \otimes_{A^a} B^a)^{\Delta}$ for any $M^a \in \mathbf{D}_{acoh}(A)$.
- (3) The functor $\mathbf{L}\mathfrak{f}^*$ carries $\mathbf{D}^-_{qc,acoh}(\mathfrak{Y})$ to $\mathbf{D}^-_{qc,acoh}(\mathfrak{X})$.
- (4) The functor $\mathbf{L}\mathfrak{f}^*$ carries $\mathbf{D}^-_{acoh}(\mathfrak{Y})^a$ to $\mathbf{D}^-_{acoh}(\mathfrak{X})^a$.

Proof. The proof is similar to Lemma 4.5.5. We use Lemma 4.7.1 to reduce to the analogous algebraic fact that was already proven in Lemma 4.2.3. \Box

Lemma 4.7.4. Let \mathfrak{X} be a locally topologically finitely presented formal *R*-scheme for *R* as in the Setup 4.4.1.

- (1) Suppose that $\mathfrak{X} = \operatorname{Spf} A$ is affine. Then $M^{\Delta} \otimes_{\mathfrak{O}_{\mathfrak{X}}}^{L} N^{\Delta}$ is functorially isomorphic to $(M \otimes_{A}^{L} N)^{\Delta}$ for any $M, N \in \mathbf{D}_{acoh}(A)$.
- (2) Suppose that $\mathfrak{X} = \operatorname{Spf} A$ is affine. Then $M^{a,\Delta} \otimes_{\mathfrak{O}_{\mathfrak{X}}^a}^L N^{a,\Delta}$ is functorially isomorphic to $(M^a \otimes_{A^a}^L N^a)^{\Delta}$ for any $M^a, N^a \in \mathbf{D}_{acoh}(A)^a$.
- (3) Let $\mathfrak{F}, \mathfrak{G} \in \mathbf{D}^{-}_{qc,acoh}(\mathfrak{X})$. Then $\mathfrak{F} \otimes^{L}_{\mathfrak{O}_{\mathfrak{X}}} \mathfrak{G} \in \mathbf{D}^{-}_{qc,acoh}(\mathfrak{X})$.
- (4) Let $\mathfrak{F}^a, \, \mathfrak{G}^a \in \mathbf{D}^-_{acoh}(\mathfrak{X})^a$. Then $\mathfrak{F}^a \otimes^L_{\mathfrak{O}^a_{\mathfrak{X}}} \mathfrak{G}^a \in \mathbf{D}^-_{acoh}(\mathfrak{X})^a$.

Proof. Similarly to Lemma 4.7.3, we use Lemma 4.7.1 to reduce to the analogous algebraic fact that was already proven in Lemma 4.2.4. \Box

Now we discuss the $\mathbf{R}_{\underline{alHom}_{\mathcal{O}_{\mathfrak{X}}}}(-,-)$ functor. Our strategy of showing that $\mathbf{R}_{\underline{alHom}}(-,-)$ preserves almost coherent complexes will be slightly different from the schematic case. The main technical problem is to define the map $\mathbf{R}_{\underline{alHom}_{A^a}}(M^a, N^a)^{\Delta} \to \mathbf{R}_{\underline{alHom}_{\mathcal{O}_{\mathfrak{X}}^a}}(M^{\Delta}, N^{\Delta})$ in the affine case.

The main issue is that we do not know if $(-)^{\Delta}$ is a left adjoint to the functor of global section on the whole category $\mathbf{D}(\mathfrak{X})$; we only know that it becomes a pseudo-inverse to $\mathbf{R}\Gamma(\mathfrak{X}, -)$ after restriction to " $\mathbf{D}_{qc}(\mathfrak{X})$ ". However, the complex $\mathbf{R}\underline{\mathcal{H}om}_{\mathfrak{O}_{\mathfrak{X}}}(M^{\Delta}, N^{\Delta})$ itself does not usually lie inside " $\mathbf{D}_{qc}(\mathfrak{X})$ ". To overcome this issue, we will show that $\widetilde{\mathfrak{m}} \otimes \mathbf{R}\underline{\mathcal{H}om}_{\mathfrak{O}_{\mathfrak{X}}}(M^{\Delta}, N^{\Delta})$ does lie in " $\mathbf{D}_{qc}(\mathfrak{X})$ " if $M \in \mathbf{D}_{acoh}^{-}(A)$ and $N \in \mathbf{D}_{acoh}^{+}(A)$.

Since " $\mathbf{D}_{qc}(\mathfrak{X})$ " was defined in a bit abstract way, it is probably the easiest way to show that $\widetilde{\mathfrak{m}} \otimes \mathbf{R} \underline{\mathcal{H}}om_{\mathfrak{O}_{\mathfrak{X}}}(M^{\Delta}, N^{\Delta})$ actually lies in $\mathbf{D}_{qc,acoh}(\mathfrak{X})$. That is sufficient by Lemma 4.6.12.

Lemma 4.7.5. Let $\mathfrak{X} = \text{Spf } A$ be a topologically finitely presented formal *R*-scheme for *R* as in the Setup 4.4.1. Let $M, N \in \mathbf{Mod}_A^{\operatorname{acoh}}$ there are natural almost isomorphisms

$$\operatorname{Ext}_{A}^{p}(M,N)^{\Delta} \xrightarrow{\sim} \mathcal{E}xt_{\mathcal{O}_{\mathfrak{X}}}^{p}(M^{\Delta},N^{\Delta})$$

for all integer p.

Proof. We recall that $\mathcal{E}xt^p_{\mathcal{O}_{\mathfrak{T}}}(M^{\Delta}, N^{\Delta})$ is canonically isomorphic to sheafification of the presheaf

$$\mathfrak{U} \mapsto \operatorname{Ext}^p_{\mathcal{O}_{\mathfrak{U}}}(M^{\Delta}|_{\mathfrak{U}}, N^{\Delta}|_{\mathfrak{U}}) \;.$$

In particular, there is a canonical map $\operatorname{Ext}^p_{\mathcal{O}_{\mathfrak{X}}}(M^{\Delta}, N^{\Delta}) \to \Gamma(\mathfrak{X}, \mathcal{E}xt^p_{\mathcal{O}_{\mathfrak{X}}}(M^{\Delta}, N^{\Delta}))$. It induces the morphism

$$\operatorname{Ext}^{p}_{\mathcal{O}_{\mathfrak{X}}}(M^{\Delta}, N^{\Delta})^{\Delta} \to \mathcal{E}xt^{p}_{\mathcal{O}_{\mathfrak{X}}}(M^{\Delta}, N^{\Delta}) \ . \tag{4.7}$$

Now we note that the classical $(-)^{\Delta}$ functor and the derived version coincide on almost coherent modules by Lemma 4.6.13. Hence, the equivalence " $\mathbf{D}_{qc}(\mathfrak{X})$ " $\simeq \mathbf{D}_{comp}(A, I)$ coming from Theorem 4.6.6 ensures that $\operatorname{Ext}_{\mathcal{O}_{\mathfrak{X}}}^{p}(M^{\Delta}, N^{\Delta}) \simeq \operatorname{Ext}_{A}^{p}(M, N)$. So the map (4.7) becomes the map

$$\operatorname{Ext}_{A}^{p}(M,N)^{\Delta} \to \mathcal{E}xt_{\mathcal{O}_{\mathfrak{F}}}^{p}(M^{\Delta},N^{\Delta})$$

We note that $\operatorname{Ext}_{A}^{p}(M, N)$ is an almost coherent A-module by Proposition 2.6.15. Using that almost coherent modules are complete, we conclude that it suffices to show that

$$\operatorname{Ext}_{A}^{p}(M,N) \otimes_{A} A_{\{f\}} \to \operatorname{Ext}_{\operatorname{Spf}A_{\{f\}}}^{p}(M^{\Delta}|_{\operatorname{Spf}A_{\{f\}}}, N^{\Delta}|_{\operatorname{Spf}A_{\{f\}}})$$

is an almost isomorphism. Using Lemma 4.5.5 and the equivalence " $\mathbf{D}_{qc}(\mathfrak{X})$ " $\simeq \mathbf{D}_{comp}(A, I)$ as above, we see that the map above becomes the canonical map

$$\operatorname{Ext}_{A}^{p}(M,N) \otimes_{A} A_{\{f\}} \to \operatorname{Ext}_{A_{\{f\}}}^{p}(M \otimes_{A} A_{\{f\}}, N \otimes_{A} A_{\{f\}}) .$$

Finally, this map is an almost isomorphism by Proposition 2.7.9.

Corollary 4.7.6. Let \mathfrak{X} be a locally topologically finitely presented formal *R*-scheme for *R* as in the Setup 4.4.1. Let $\mathfrak{F} \in \mathbf{D}^-_{qc,acoh}(\mathfrak{X})$, and $\mathfrak{G} \in \mathbf{D}^+_{qc,acoh}(\mathfrak{X})$. Then $\widetilde{\mathfrak{m}} \otimes \mathbf{R} \underline{\mathcal{H}om}_{\mathfrak{O}_{\mathfrak{X}}}(\mathfrak{F}, \mathfrak{G}) \in \mathbf{D}^+_{qc,acoh}(\mathfrak{X})$.

Proof. The claim is local, so we can assume that $\mathfrak{X} = \operatorname{Spf} A$. Then Theorem 4.6.15 imples that $\mathfrak{F} \cong M^{\Delta}$ for $M \in \mathbf{D}^{-}_{acoh}(A)$ and $\mathfrak{G} \cong N^{\Delta}$ for $N \in \mathbf{D}^{+}_{acoh}(A)$. Then Lemma 4.7.5 guarantees that $\mathcal{H}^{p}\left(\mathbf{R} \underbrace{\mathcal{H}om}_{\mathfrak{O}_{\mathfrak{X}}}(\mathfrak{F},\mathfrak{G})\right) \simeq^{a} \operatorname{Ext}_{A}^{p}(M, N)^{\Delta}$. In other words,

$$\widetilde{\mathfrak{m}} \otimes \mathcal{H}^p\left(\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_{\mathfrak{T}}}(\mathcal{F},\mathcal{G})\right) \simeq^a \widetilde{\mathfrak{m}} \otimes \operatorname{Ext}_A^p(M,N)^{\Delta}$$

Now $\operatorname{Ext}_{A}^{p}(M, N)^{\Delta}$ is an adically quasi-coherent, almost coherent $\mathcal{O}_{\mathfrak{X}}$ -module by Proposition 2.6.15 and Lemma 4.5.1. So Lemma 4.4.10 guarantees that $\widetilde{\mathfrak{m}} \otimes \operatorname{Ext}_{A}^{p}(M, N)^{\Delta}$ is also adically quasi-coherent and almost coherent. Therefore, $\widetilde{\mathfrak{m}} \otimes \mathbf{R} \underline{\mathcal{H}om}_{\mathcal{O}_{\mathfrak{X}}}(\mathfrak{F}, \mathfrak{G}) \in \mathbf{D}_{qc,acoh}^{+}(\mathfrak{X})$.

Lemma 4.7.7. Let \mathfrak{X} be a locally topologically finitely presented formal *R*-scheme for *R* as in the Setup 4.4.1.

(1) Suppose $\mathfrak{X} = \operatorname{Spf} A$ is affine. Then there is a functorial isomorphism

$$\mathbf{R}$$
al $\operatorname{Hom}_{A^a}(M^a, N^a)^{\Delta} \to \mathbf{R}\underline{al} \mathcal{H}om_{\mathcal{O}^a_{\infty}}(M^{a,\Delta}, N^{a,\Delta})$

for $M \in \mathbf{D}^{-}_{acoh}(A)^a$ and $N \in \mathbf{D}^{+}_{acoh}(A)^a$.

(2) Suppose $\mathcal{F}^a \in \mathbf{D}^+_{acoh}(\mathfrak{X})^a$ and $\mathcal{G}^a \in \mathbf{D}^-_{acoh}(\mathfrak{X})$ are almost coherent $\mathcal{O}^a_{\mathfrak{X}}$ -modules. Then $\mathbf{R}_{\underline{alHom}_{\mathcal{O}^a_{\mathfrak{X}}}}(\mathcal{F}^a, \mathcal{G}^a) \in \mathbf{D}^+_{acoh}(\mathfrak{X})^a$.

Proof. We start with (1). Proposition 3.5.8 implies the map

 $(\widetilde{\mathfrak{m}}\otimes\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_{\mathfrak{X}}}(M^{\Delta},N^{\Delta}))^{a}\rightarrow\mathbf{R}\underline{al\mathcal{H}om}_{\mathcal{O}_{\mathfrak{X}}^{a}}(M^{a,\Delta},N^{a,\Delta})$

is an isomorphism in $\mathbf{D}(\mathfrak{X})^a$. Similarly, the map

 $(\widetilde{\mathfrak{m}} \otimes \mathbf{R}\mathrm{Hom}_A(M,N)^{\Delta})^a \to \mathbf{R}\mathrm{a}\mathrm{Hom}_{A^a}(M^a,N^a)^{\Delta}$

is an isomorphism by Lemma 4.7.5. This implies that it suffices to construct a functorial isomorphism

$$\widetilde{\mathfrak{m}} \otimes \mathbf{R} \operatorname{Hom}_A(M, N)^{\Delta} \to \widetilde{\mathfrak{m}} \otimes \mathbf{R} \underline{\mathcal{H}om}_{\mathcal{O}_{\mathfrak{X}}}(M^{\Delta}, N^{\Delta})$$
.

Now Corollary 4.7.6 guarantees that $\widetilde{\mathfrak{m}} \otimes \mathbf{R} \underline{\mathcal{H}om}_{\mathcal{O}_{\mathfrak{X}}}(M^{\Delta}, N^{\Delta}) \in \mathbf{D}_{qc,acoh}(\mathfrak{X})$. Proposition 2.6.15, Lemma 4.5.1, and Lemma 4.4.10 also guarantee that $\widetilde{\mathfrak{m}} \otimes \mathbf{R} \operatorname{Hom}_{A}(M, N)^{\Delta} \in \mathbf{D}_{qc,acoh}(\mathfrak{X})$. Thus, Theorem 4.6.6 ensures that in order to construct the desired isomorphism it suffices to do it after applying $\mathbf{R}\Gamma(\mathfrak{X}, -)$. Projection Formula (Theorem 3.3.6) provides us with functorial isomorphisms

$$\mathbf{R}\Gamma(\mathfrak{X}, \widetilde{\mathfrak{m}} \otimes \mathbf{R}\mathrm{Hom}_{A}(M, N)^{\Delta}) \simeq \widetilde{\mathfrak{m}} \otimes \mathbf{R}\mathrm{Hom}_{A}(M, N)$$
$$\mathbf{R}\Gamma(\mathfrak{X}, \widetilde{\mathfrak{m}} \otimes \mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_{\mathfrak{X}}}(M^{\Delta}, N^{\Delta})) \simeq \widetilde{\mathfrak{m}} \otimes \mathbf{R}\Gamma(\mathfrak{X}, \mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_{\mathfrak{X}}}(M^{\Delta}, N^{\Delta}))$$
$$\simeq \widetilde{\mathfrak{m}} \otimes \mathbf{R}\mathrm{Hom}_{\mathcal{O}_{X}}(M^{\Delta}, N^{\Delta})$$
$$\simeq \widetilde{\mathfrak{m}} \otimes \mathbf{R}\mathrm{Hom}_{A}(M, N)$$

where the last isomorphism uses equivalence from Theorem 4.6.6. Thus, we see

$$\mathbf{R}\Gamma(\mathfrak{X},\widetilde{\mathfrak{m}}\otimes\mathbf{R}\mathrm{Hom}_{A}(M,N)^{\Delta})\simeq\mathbf{R}\Gamma(\mathfrak{X},\widetilde{\mathfrak{m}}\otimes\mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_{\mathfrak{X}}}(M^{\Delta},N^{\Delta})).$$

As a consequence, we have a functorial isomorhism

$$\widetilde{\mathfrak{m}} \otimes \mathbf{R}\mathrm{Hom}_A(M,N)^{\Delta} \xrightarrow{\sim} \widetilde{\mathfrak{m}} \otimes \mathbf{R}\underline{\mathcal{H}om}_{\mathcal{O}_{\mathfrak{T}}}(M^{\Delta},N^{\Delta})$$
.

This induces the desired isomorphism

$$\mathbf{RalHom}_{A^a}(M^a, N^a)^{\Delta} \xrightarrow{\sim} \mathbf{R}\underline{al\mathcal{H}om}_{\mathcal{O}^a_{\mathfrak{X}}}(M^{a,\Delta}, N^{a,\Delta})$$

(2) is an easy consequence of (1), Proposition 2.6.15, and Corollary 4.6.14.

5. Cohomological Properties of Almost Coherent Sheaves

5.1. Almost Proper Mapping Theorem. The main goal of this section is to prove the "Almost Proper Mapping Theorem" both in setup of both "nice" schemes and "nice" formal schemes. The theorem roughly says the derived pushforward of an almost coherent \mathcal{O}_X -module along a (topolog-ically) finitely presented proper map is almost coherent.

The idea of the proof in the schematic case is rather easy: we "approximate" an almost finitely presented \mathcal{O}_X -module by finitely presented using Lemma 4.1.8 and then the usual Proper Mapping Theorem. However, there is a subtlety that the usual Proper Mapping Theorem is usually proven only for a (locally) noetherian base, and we are really interested in non-noetherian situation. So we use a more general version (in so-called "universally coherent" case) of the Proper Mapping Theorem from the book [FK18].

The proof in the formal case is more subtle. We do not know if an analogue of Lemma 4.1.8 holds in the formal setup. So, it is not clear if we can run a similar argument using the Proper Mapping Theorem for usual schemes as an input. Instead we imitate the proof of the latter in the almost setup following the ideas from [FK18].

Definition 5.1.1. We say that a scheme Y is *universally coherent* if any scheme X that is locally of finite presentation over Y is coherent (i.e. the structure sheaf \mathcal{O}_X is coherent).

Theorem 5.1.2 (Proper Mapping Theorem). [FK18, Theorem I.8.1.3] Let Y be a universally coherent quasi-compact scheme, and $f: X \to Y$ a proper morphism of finite presentation. Then the functor $\mathbf{R}f_*$ sends $\mathbf{D}^*_{coh}(X)$ to $\mathbf{D}^*_{coh}(Y)$ for any $* \in \{"", +, -, b\}$.

We want to generalize this theorem to the "almost world". So we pick a ring R and a fixed ideal $\mathfrak{m} \subset R$ such that $\mathfrak{m}^2 = \mathfrak{m}$ and $\tilde{\mathfrak{m}} = \mathfrak{m} \otimes_R \mathfrak{m}$ is R-flat. In this section, we always consider almost mathematics with respect to this ideal.

Theorem 5.1.3 (Almost Proper Mapping Theorem). Let Y be a universally coherent quasicompact R-scheme, and let $f: X \to Y$ be a proper, finitely presented morphism. Then

- The functor $\mathbf{R}f_*$ sends $\mathbf{D}^*_{qc,acoh}(X)$ to $\mathbf{D}^*_{qc,acoh}(Y)$ for any $* \in \{", ", +, -, b\}$.
- The functor $\mathbf{R}f_*$ sends $\mathbf{D}^*_{acoh}(X)^a$ to $\mathbf{D}^*_{acoh}(Y)^a$ for any $* \in \{", ", +, -, b\}$.
- The functor $\mathbf{R}f_*$ sends $\mathbf{D}^+_{acoh}(X)$ to $\mathbf{D}^+_{acoh}(Y)$.
- If Y has finite Krull dimension, then $\mathbf{R}f_*$ sends $\mathbf{D}^*_{acoh}(X)$ to $\mathbf{D}^*_{acoh}(Y)$ for any $* \in \{", ", +, -, b\}$.

Lemma 5.1.4. Let Y be a quasi-compact scheme of finite Krull dimension, and let $f : X \to Y$ be a finite type, quasi-separated morphism. Then X has finite Krull dimension, and f_* is of finite cohomological dimension on \mathbf{Mod}_X .

Proof. First of all, we show that X has finite Krull dimension. Indeed, the morphism $f: X \to Y$ is quasi-compact, therefore X is quasi-compact. Then it suffices to show that locally X has finite Krull dimension. So we can assume that $X = \operatorname{Spec} B$, $Y = \operatorname{Spec} A$, and the map is given by a finite type morphism $A \to B$. In that situation we have dim $Y = \dim A$ and dim $X = \dim B$. Thus, it is enough to show that the Krull dimension of a finite type A-algebra is finite. This readily reduces the question to the case of a polynomial algebra dim $A[X_1, \ldots, X_n]$. Now [AM69, Chapter 11, Exercise 6] implies that dim $A[X_1, \ldots, X_n] \leq \dim A + 2n$.

Now we prove that f_* has finite cohomological dimension. We note that it suffices to show that there is an integer N such that for any open affine $U \subset Y$ the cohomology groups $\mathrm{H}^i(X_U, \mathcal{F})$ vanish for $i \geq N$ and any \mathcal{O}_{X_U} -module \mathcal{F} . We recall that f is quasi-separated, so X_U is quasi-compact,

quasi-separated and dim $X_U \leq \dim X$ for any open $U \subset X$. Therefore, it is sufficient to show that on any spectral space X we have $\mathrm{H}^i(X, \mathcal{F}) = 0$ for $i > \dim X$ and $\mathcal{F} \in \mathcal{A}b(X)$. This is proven in [Sch92, Corollary 4.6] (another reference is [Sta21, Tag 0A3G]). Thus we see that $N = \dim X$ does the job.

Proof of Theorem 5.1.3. Step 0. Reduction to the case of bounded below derived categories: We note that f_* always has bounded cohomological dimension on \mathbf{Qcoh}_X . Indeed, for any $\mathcal{F} \in \mathbf{Qcoh}_X$ on a separated scheme X, we can compute $\mathrm{H}^i(X, \mathcal{F})$ by the alternating Čech complex for some finite covering of X by affines. Therefore, if X can be covered by N affines, the functor f_* restricted to \mathbf{Qcoh}_X has cohomological dimension at most N.

Now we use [Sta21, Tag 0D6U] (alternatively, one can use [Lim19, Lemma 3.4]) to reduce the question of proving the claim for any $\mathcal{F} \in \mathbf{D}_{qc,acoh}(X)$ to the question of proving the claim for all its truncations $\tau^{\geq a}\mathcal{F}$. In particular, we can assume that $\mathcal{F} \in \mathbf{D}^+_{qc,acoh}(X)$. The case $\mathcal{F}^a \in \mathbf{D}^*_{acoh}(X)^a$ can be shown similarly. Actually, Proposition 3.5.23 and the observation that $\mathcal{F}^a_! \in \mathbf{D}_{qc,acoh}(X)$ imply that the results for $\mathbf{D}^*_{qc,acoh}(X)$ and $\mathbf{D}^*_{acoh}(X)^a$ are equivalent.

The same argument also works for $\mathbf{D}^+_{acoh}(X)$ provided that X, Y and f_* has finite cohomological dimension. Lemma 5.3.4 and [Sta21, Tag 0A3G] say that it holds whenever Y has finite Krull dimension.

Step 1. Reduction to the case quasi-coherent almost coherent sheaves: Using the Projection Formula (Lemma 3.3.6) (resp. Proposition 3.5.23), we see that in order to show $\mathbf{R}f_*$ sends $\mathbf{D}^+_{acoh}(X)$ to $\mathbf{D}^+_{acoh}(Y)$ (resp. $\mathbf{D}^+_{acoh}(X)^a$ to $\mathbf{D}^+_{acoh}(Y)^a$) it is sufficient to show the analogous result for $\mathbf{D}^+_{acoh}(X)$. Moreover, we can use the spectral sequence

$$\mathbf{E}_{2}^{p,q} = \mathbf{R}^{p} f_{*} \mathcal{H}^{q}(\mathcal{F}) \Rightarrow \mathbf{R}^{p+q} f_{*}(\mathcal{F})$$

to reduce the claim to the fact that higher derived pushforwards of a quasi-coherent, almost coherent sheaf are quasi-coherent, almost coherent.

Step 2. The case of a quasi-coherent, almost coherent \mathcal{O}_X -module \mathcal{F} : We show that $\mathbb{R}^i f_* \mathcal{F}$ is a quasi-coherent, almost coherent \mathcal{O}_Y -module for any quasi-coherent, almost coherent \mathcal{O}_X -module \mathcal{F} and any *i*. First of all, we note that $\mathbb{R}^i f_* \mathcal{F}$ is quasi-coherent as higher pushforwards along quasi-compact, quasi-separated morphisms preserve quasi-coherence.

Now we show almost coherence of $\mathbb{R}^i f_* \mathcal{F}$. Note that it is sufficient to show that $\mathbb{R}^i f_* \mathcal{F}$ is almost finitely presented as Y is a coherent scheme (this follows from Lemma 4.1.16 and Lemma 4.1.17). We choose some finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$ and another finitely generated ideal $\mathfrak{m}_1 \subset \mathfrak{m}$ such that $\mathfrak{m}_0 \subset \mathfrak{m}_1^2$. Then we use Lemma 4.1.8 to find a finitely presented \mathcal{O}_X -module \mathcal{G} and a morphism

$$\varphi: \mathfrak{G} \to \mathfrak{F}$$

such that ker(φ) and Coker(φ) are annihilated by \mathfrak{m}_1 . We define \mathcal{O}_X -modules

 $\mathcal{K} := \ker \varphi, \ \mathcal{M} := \operatorname{Im} \varphi \text{ and } \mathcal{Q} := \operatorname{Coker} \varphi,$

so we have two short exact sequences

$$\begin{array}{l} 0 \to \mathcal{K} \to \mathcal{G} \to \mathcal{M} \to 0 \\ 0 \to \mathcal{M} \to \mathcal{F} \to \mathcal{Q} \to 0 \end{array}$$

with sheaves \mathcal{K} and \mathcal{Q} killed by \mathfrak{m}_1 . This easily shows that the natural homomorphisms

$$\mathrm{R}^{i}f_{*}(\varphi):\mathrm{R}^{i}f_{*}\mathcal{G}\to\mathrm{R}^{i}f_{*}\mathcal{F}$$

have kernels and cokernels annihilated by \mathfrak{m}_1^2 . Since $\mathfrak{m}_0 \subset \mathfrak{m}_1^2$ we see that $\mathfrak{m}_0(\ker \mathrm{R}^i f_*(\varphi)) = 0$ and $\mathfrak{m}_0(\operatorname{Coker} \mathrm{R}^i f_*(\varphi)) = 0$. Moreover, we know that $\mathrm{R}^i f_* \mathcal{G}$ is a finitely presented \mathcal{O}_Y -module by Theorem 5.1.2 (\mathcal{G} is a coherent \mathcal{O}_X -module since X is a coherent scheme). Therefore we use Lemma 4.1.8 to conclude that $\mathrm{R}^i f_* \mathcal{F}$ is an almost finitely presented \mathcal{O}_Y -module for any $i \geq 0$. And this implies the almost coherence of $\mathrm{R}^i f_* \mathcal{F}$ as explained above. \Box

Now we go to the formal version of this theorem. Unfortunately, we can not establish the Almost Proper Mapping Theorem in the most general desired form; we need to restrict our attention to the case of a principal ideal of definition (and a little bit more).

Set-up 5.1.5. We fix ϖ -adically complete, and ϖ -adically topologically universally adhesive ring R with some nonzerodivisor ϖ , and we assume there is a compatible system of roots $\varpi^{1/n}$ for each integer $n \ge 0$. We define the ideal of almost mathematics as $\mathfrak{m} = \bigcup_n \varpi^{1/n} R$. It is clear that $\mathfrak{m}^2 = \mathfrak{m}$ and [GR03, Proposition 2.1.7] implies that $\widetilde{\mathfrak{m}} = \mathfrak{m} \otimes_R \mathfrak{m}$ is flat.

Note that any such ring satisfies the conditions of the Setup 4.4.1 and Setup A.1. So, we have a good notion of an (adically quasi-coherent) almost coherent $\mathcal{O}_{\mathfrak{X}}$ -modules on any topologically finitely presented formal *R*-scheme, and the results of Appendix A are applicable in our setting.

Theorem 5.1.6 (Almost Proper Mapping Theorem). Let \mathfrak{Y} be a topologically finitely presented formal *R*-scheme for *R* as in the Setup 5.1.5. And let $\mathfrak{f} : \mathfrak{X} \to \mathfrak{Y}$ be a proper, topologically finitely presented morphism. Then

- The functor $\mathbf{R}\mathfrak{f}_*$ sends $\mathbf{D}^*_{ac,acoh}(\mathfrak{X})$ to $\mathbf{D}^*_{ac,acoh}(\mathfrak{Y})$ for any $* \in \{", ", +, -, b\}$.
- The functor $\mathbf{R}_{\mathfrak{f}_*}$ sends $\mathbf{D}^*_{acoh}(\mathfrak{X})^a$ to $\mathbf{D}^*_{acoh}(\mathfrak{Y})^a$ for any $* \in \{", ", +, -, b\}$.
- The functor $\mathbf{R}\mathfrak{f}_*$ sends $\mathbf{D}^+_{acoh}(\mathfrak{X})$ to $\mathbf{D}^+_{acoh}(\mathfrak{Y})$.
- If $Y_0 := \mathfrak{Y} \times_{\operatorname{Spf} R} (\operatorname{Spec} R/\varpi)$ has finite Krull dimension, then $\mathbf{R}\mathfrak{f}_*$ sends $\mathbf{D}^*_{acoh}(\mathfrak{X})$ to $\mathbf{D}^*_{acoh}(\mathfrak{Y})$ for any $* \in \{ "", +, -, b \}$.

Moreover, if $\mathfrak{Y} = \operatorname{Spf} A$ is an affine scheme and \mathcal{F} is an adically quasi-coherent, almost coherent $\mathcal{O}_{\mathfrak{X}}$ -module, then the natural map $\operatorname{H}^{n}(\mathfrak{X}, \mathcal{F})^{\Delta} \to \operatorname{R}^{n} f_{*} \mathcal{F}$ is an isomorphism of $\mathcal{O}_{\mathfrak{Y}}$ -modules for $n \geq 0$.

Lemma 5.1.7. Let \mathfrak{Y} be a quasi-compact adic formal *R*-scheme, and let $\mathfrak{f} : \mathfrak{X} \to \mathfrak{Y}$ be a topologically finite type, quasi-separated morphism. Suppose that the reduction $Y_0 = \mathfrak{Y} \times_{\mathrm{Spf} R} (\mathrm{Spec} R/\varpi)$ (or equivalently the "special fiber" $\overline{\mathfrak{Y}} = \mathfrak{Y} \times_{\mathrm{Spf} R} \mathrm{Spec} R/\mathrm{Rad}(\varpi)$) is of finite Krull dimension. Then \mathfrak{X} has finite Krull dimension, and \mathfrak{f}_* is of finite cohomological dimension on $\mathrm{Mod}_{\mathfrak{X}}$.

Proof. The proof is identical to Lemma 5.1.4 once we notice that the underlying topological spaces of \mathfrak{Y}, Y_0 and $\overline{\mathfrak{Y}}$ are canonically identified.

Also, before going to the proof of Theorem 5.1.6 we need to establish one preliminary lemma.

Lemma 5.1.8. Let $\mathfrak{f}: \mathfrak{X} \to \mathfrak{Y} = \operatorname{Spf} A$ be a morphism as in Theorem 5.1.6 with affine \mathfrak{Y} , and let $\mathcal{F} \in \operatorname{\mathbf{Mod}}_{\mathfrak{X}}$ be an adically quasi-coherent, almost coherent sheaf. Then $\mathrm{R}^q\mathfrak{f}_*\mathcal{F}$ is an adically quasi-coherent, almost coherent $\mathfrak{O}_{\mathfrak{Y}}$ -module if

- (1) the A-module $\mathrm{H}^{q}(\mathfrak{X}, \mathfrak{F})$ is almost coherent for any $q \geq 0$,
- (2) for any $g \in A$, the canonical map

 $\mathrm{H}^{q}(\mathfrak{X}, \mathfrak{F}) \otimes_{A} A_{\{q\}} \to \mathrm{H}^{q}(\mathfrak{X}_{\mathfrak{U}}, \mathfrak{F}),$

where $\mathfrak{U} = \operatorname{Spf} A_{\{q\}} \to \mathfrak{Y} = \operatorname{Spf} A$, is an isomorphism for any $q \ge 0$.

Proof. Set $M := \mathrm{H}^q(\mathfrak{X}, \mathfrak{F})$ that is almost coherent by the hypothesis 1. Therefore, Lemma 2.8.7 guarantees that M is ϖ -adically complete, and so M^{Δ} is an adically quasi-coherent, almost coherent $\mathcal{O}_{\mathfrak{X}}$ -module. Now note that $\mathrm{R}^q \mathfrak{F}$ is the sheafification of the presheaf

$$\mathfrak{U}\mapsto \mathrm{H}^q(\mathfrak{X}_\mathfrak{U},\mathfrak{F})$$

Thus there is a canonical map $M \to \mathrm{H}^0(\mathfrak{Y}, \mathrm{R}^q f_* \mathcal{F})$ that induces the morphism

 $M^{\Delta} \to \mathbf{R}^q f_* \mathcal{F}$

The second hypothesis together with Lemma 2.7.1 and Lemma 2.8.7 ensures this map is an isomorphism on stalks (as the sheafification process preserves stalks). Therefore, $M^{\Delta} \to \mathbb{R}^{q} f_{*} \mathcal{F}$ is an isomorphism of $\mathcal{O}_{\mathfrak{X}}$ -modules. In particular, $\mathbb{R}^{q}\mathfrak{f}_{*}\mathcal{F}$ is adically quasi-coherent and almost coherent.

Proof of Theorem 5.1.6. We use the same reduction as in the proof of Theorem 5.1.3 to reduce to the situation of an adically quasi-coherent, almost coherent $\mathcal{O}_{\mathfrak{X}}$ -module \mathfrak{F} . Moreover, the statement is local on \mathfrak{Y} , so we can assume that $\mathfrak{Y} = \operatorname{Spf} A$ is affine. The assumptions on the R imply that the ϖ -adic ring A fits into the Setup A.1.

Now we show that both conditions in Lemma 5.1.8 are satisfied in our situation. We choose some finite covering \mathfrak{X} by affine open formal schemes $\mathfrak{X} = \bigcup_{i=1}^{m} \mathfrak{U}_i$. As \mathfrak{X} is separated, the intersections $\mathfrak{U}_{i,j} := \mathfrak{U}_i \cap \mathfrak{U}_j$ are also affine. Thus the vanishing theorem [FK18, Theorem I.7.1.1] guarantees that we can compute cohomology of \mathcal{F} as the cohomology of the *alternating* Čech complex $K^{\bullet} := C^{\bullet}(\mathfrak{U}, \mathcal{F})$. Note that, in particular, all these cohomology groups vanish in high degree that does not depend on an adically quasi-coherent sheaf \mathcal{F} . So we shall verify both hypothesis of Lemma 5.1.8 via descending induction on the degree of a cohomology group. From now one we assume that both conditions are verified for some q + 1 and *any* adically quasi-coherent, almost coherent sheaf \mathcal{F} .

We intoduce two auxillary complexes $\varpi^{k+1}K^{\bullet} = C^{\bullet}(\mathfrak{U}, \varpi^{k+1}\mathfrak{F})$ and $K_k^{\bullet} \coloneqq C^{\bullet}(U_k, \mathfrak{F}_k)$, where \mathfrak{F}_k is the k-th reduction of the $\mathcal{O}_{\mathfrak{X}}$ -module \mathfrak{F} , and each $U_{i,k}$ is the k-th reduction of the formal scheme \mathfrak{U}_i . So we have a short exact sequence of complexes:

$$0 \to \varpi^{k+1} K^{\bullet} \to K^{\bullet} \to K^{\bullet}_k \to 0.$$

Note that Lemma 4.4.11 states that the sheaf \mathcal{F}_k is a quasi-coherent, almost coherent sheaf on X_k (as each X_k is a coherent scheme). Moreover, Corollary 4.4.18 implies that $\varpi^{k+1}\mathcal{F}$ is also an adically quasi-coherent, almost coherent $\mathcal{O}_{\mathfrak{X}}$ -module.

We show that $\mathrm{H}^{q}(K^{\bullet})$ is almost finitely generated: Combining the induction hypothesis, Lemma 2.7.3 and Theorem 5.1.3, we can assume that the cohomology groups $\mathrm{H}^{q+1}(\varpi^{k+1}K^{\bullet})$ and $\mathrm{H}^{q}(K_{k}^{\bullet})$ are almost coherent over A (the assumption on R and A imply that A is universally coherent). Also, we note that Lemma 2.8.5 and Lemma 2.8.7 show that each term of the complexes K^{\bullet} and $\varpi^{k+1}K^{\bullet}$ have bounded ϖ^{∞} -torsion and ϖ -adically complete. Thus Lemma A.4 says that there is some ksuch that

$$\mathbf{F}^{k+1}\mathbf{H}^{q}(K^{\bullet}) = \mathrm{Im}\left(\mathbf{H}^{q}(\varpi^{k+1}K^{\bullet}) \to \mathbf{H}^{q}(K^{\bullet})\right) \subset \varpi\mathbf{H}^{q}(K^{\bullet}).$$

We choose such k, and use the long exact sequence

$$\mathrm{H}^{q}(\varpi^{k+1}K^{\bullet}) \to \mathrm{H}^{q}(K^{\bullet}) \to \mathrm{H}^{q}(K^{\bullet}_{k}) \xrightarrow{\delta^{q}_{k}} \mathrm{H}^{q+1}(\varpi^{k+1}K^{\bullet})$$

to see that $\mathrm{H}^{q}(K^{\bullet})/\mathrm{F}^{k+1}\mathrm{H}^{q}(K^{\bullet}) = \ker \delta_{k}^{q}$ must be a coherent A-module by Lemma 2.6.7. In particular, it is an almost finitely generated A-module. But then the module $\mathrm{H}^{q}(K^{\bullet})/\varpi$ is almost almost finitely generated as a quotient of $\mathrm{H}^{q}(K^{\bullet})/\mathrm{F}^{k+1}\mathrm{H}^{q}(K^{\bullet})$. Finally we observe that $\mathrm{H}^{q}(K^{\bullet})$ is equal to $\operatorname{Coker}(K^{q-1} \xrightarrow{d^{q-1}} \ker d^q)$ where the first module is ϖ -adically complete and the second module is ϖ -adically separated by the proof of Lemma A.4. Therefore, Lemma A.2 guarantees that $\operatorname{H}^q(K^{\bullet})$ is almost finitely generated over A.

We show that $\mathrm{H}^{q}(K^{\bullet})$ is almost finitely presented: Now we show that $\mathrm{H}^{q}(K^{\bullet})$ is almost finitely presented provided that it is almost finitely generated. Namely, we consider the submodule T := $\mathrm{H}^{q}(K^{\bullet})[\varpi^{\infty}]$ and the quotient module $\mathrm{H}^{q}(K^{\bullet})/T$. Note that Lemma 2.8.4 states that T is an almost finitely generated module and $\mathrm{H}^{q}(K^{\bullet})/T$ is almost finitely presented. Thus it is sufficient to show that T is almost finitely presented in order to conclude that so is $\mathrm{H}^{q}(K^{\bullet})$.

We recall that Lemma 2.8.5 says that $\mathrm{H}^{q}(K^{\bullet})$ has bounded ϖ^{∞} -torsion. In particular, it implies that there is an integer n such that

$$\varpi^{n} \mathrm{H}^{q}(K^{\bullet}) \cap T = \varpi^{n}(T \cap \mathrm{H}^{q}(K^{\bullet})) = 0.$$

Using Lemma A.4 once again, we conclude that there is some large k such that $F^{k+1}H^q(K^{\bullet})\cap T = 0$. Another reformulation of this is that $\ker \delta_k^q \cap T = 0$, so T can be identified with a submodule of $H^q(K_k^{\bullet})$ for some large k. However, T is almost finitely generated and $H^q(K_k^{\bullet})$ is almost coherent by Almost Proper Mapping Theorem 5.1.3. Therefore, T is actually almost finitely presented over A by the very definition of almost coherent modules!²⁵ That finishes the proof of almost finite presentation of $H^q(K^{\bullet})$.

We verify the second condition of Lemma 5.1.8: Given any open principal open affine $\operatorname{Spf} B = \mathfrak{V} \subset \mathfrak{Y}$, we have a finite covering of $\mathfrak{X}_{\mathfrak{V}}$ by the intersections $\mathfrak{V} \cap \mathfrak{U}_i$. Denote this covering by $\mathfrak{U}_{\mathfrak{V}}$, then we introduce the alternating Čech complex $L^{\bullet} := C^{\bullet}(\mathfrak{U}_{\mathfrak{V}}, \mathcal{F})$ that computes the cohomology groups $\operatorname{H}^q(\mathfrak{V}, \mathcal{F})$. We need to check that the natural A-linear map $r: K^{\bullet} \to L^{\bullet}$ induces isomorphisms

$$\mathrm{H}^{q}(r) \colon \mathrm{H}^{q}(K^{\bullet}) \otimes_{A} B \to \mathrm{H}^{q}(L^{\bullet})$$

for all q and all adically quasi-coherent, almost coherent \mathcal{F} . Note that both groups are zero in high dimensions as we observed above. Therefore, we can prove this claim by descending induction on q. Suppose that we know the claim in degree q + 1 for all adically quasi-coherent, almost coherent \mathcal{F} , and we want to show it in degree q.

Recall that since \mathcal{F} is adically quasi-coherent and almost coherent, all the terms of the complexes K^{\bullet} and L^{\bullet} are ϖ -adically complete and have bounded ϖ^{∞} -torsion. Moreover, we have already proven that the cohomologies of these complexes are almost finitely presented over A and B, respectively. Finally, we note that the map r induces the map $r_k \colon K_k^{\bullet} \to L_k^{\bullet}$ and the induced maps

$$\mathrm{H}^{q}(r_{k}) \colon \mathrm{H}^{q}(K_{k}^{\bullet}) \otimes_{A_{k}} B_{k} \to \mathrm{H}^{q}(L_{k}^{\bullet})$$

are isomorphisms since $A_k \to B_k$ is flat and those complexes computes cohomology groups $\mathrm{H}^q(X_k, \mathcal{F}_k)$ and $\mathrm{H}^q(X_{V,k}, \mathcal{F})$. Finally, we note that the map $A \to B$ is flat by [FK18, Proposition I.4.8.1] and

$$\mathrm{H}^{q+1}(\varpi^{k+1}r)\colon\mathrm{H}^{q+1}(\varpi^{k+1}K^{\bullet})\otimes_{A}B\to\mathrm{H}^{q}(\varpi^{k+1}L^{\bullet})$$

is an isomorphism for any k by the induction hypothesis. Thus all the assumptions of Lemma A.5 are satisfied, so the map $H^q(r)$ is indeed an isomorphism for any q and any \mathcal{F} .

²⁵We crucially use here that A is a coherent ring

5.2. Characterization of Quasi-Coherent, Almost Coherent Complexes. The main goal of this Section is to show an almost analogue of [Sta21, Tag 0CSI]. This gives a useful characterization of objects in $\mathbf{D}^{b}_{qcmacoh}(X)$ on a separated, finitely presented *R*-scheme for a universally coherent *R*. This will be crucially used in our proof of the almost version of the Formal GAGA Theorem 5.3.2.

Our proof is very close to the proof of [Sta21, Tag 0CSI], but we need to make certain adjustments to make the arguments work in the almost coherent setting.

Theorem 5.2.1. Let R be an universally coherent ring with an ideal \mathfrak{m} such that $\mathfrak{m}^2 = \mathfrak{m}$ and $\widetilde{\mathfrak{m}} := \mathfrak{m} \otimes_R \mathfrak{m}$ is flat. Suppose that $\mathcal{F} \in \mathbf{D}_{qc}(\mathbf{P}_R^N)$ an element such that $\mathbf{R}\operatorname{Hom}_{\mathbf{P}^N}(\mathcal{P}, \mathcal{F}) \in \mathbf{D}_{acoh}^-(R)$ for $\mathcal{P} = \bigoplus_{i=0}^N \mathcal{O}(i)$. Then $\mathcal{F} \in \mathbf{D}_{qc,acoh}^-(\mathbf{P}_R^N)$.

Proof. We follow the ideas of [Sta21, Tag 0CSG]. Denote the dg algebra \mathbb{R} Hom_X(\mathcal{P}, \mathcal{P}) by S. A computation of cohomology groups of line bundles on \mathbb{P}_R^N implies that S is a "discrete" non-commutative algebra that is finite and flat over R. [Sta21, Tag 0BQU]²⁶ guarantees that the functor

$$-\otimes_{S}^{\mathbf{L}} \mathcal{P} \colon \mathbf{D}(S) \to \mathbf{D}_{qc}(\mathbf{P}^{N})$$

is an equivalence of categories, and the map in the other direction is given by

$$\mathbf{R}$$
Hom $(\mathcal{P}, -)$: $\mathbf{D}_{qc}(\mathbf{P}^N) \to \mathbf{D}(S)$

So if we define $M := \mathbf{R}\operatorname{Hom}(\mathcal{P}, \mathcal{F}) \in \mathbf{D}(S)$, our assumptions imply that that the image of M in $\mathbf{D}(R)$ lands inside $\mathbf{D}_{acoh}^{-}(R)$. We need to show that this assumption guarantees that $\mathcal{F} \simeq M \otimes_{S}^{\mathbf{L}} \mathcal{P}$ lives in $\mathbf{D}_{ac.acoh}^{-}(\mathbf{P}^{N})$. Moreover, using the convergence spectral sequence

$$\mathbf{E}_{2}^{p,q} = \mathcal{H}^{p}(\mathbf{H}^{q}(M) \otimes_{S}^{\mathbf{L}} \mathcal{P}) \Rightarrow \mathcal{H}^{p+q}(M \otimes_{S}^{\mathbf{L}} \mathcal{P})$$

shows that it is sufficient to assume that M is just an S-module. Then Lemma 2.7.4 implies that for any finitely generated ideal $\mathfrak{m}_0 \subset \mathfrak{m}$, there is a finitely presented right S-module N with a morphism $f: N \to M$ such that ker f and Coker f are annihilated by \mathfrak{m}_0 . The universal coherence of R and [Sta21, Tag 0CSF] imply that $N \otimes_S^L \mathcal{P} \in \mathbf{D}^-_{ac.coh}(\mathbf{P}^N)$. Now we note that the functor

$$-\otimes_{S}^{\mathbf{L}} \mathfrak{P} \colon \mathbf{D}(S) \to \mathbf{D}_{qc}(\mathbf{P}^{N})$$

is R-linear, so the standard argument shows that the cone of the morphism

$$f \otimes_{S}^{\mathbf{L}} \mathfrak{P} \colon N \otimes_{S}^{\mathbf{L}} \mathfrak{P} \to M \otimes_{S}^{\mathbf{L}} \mathfrak{P}$$

has cohomology sheaves anihilated by $\mathfrak{m}_0 \mathcal{O}_X$. Finally, Lemma 2.5.7 says that $M \otimes_S^{\mathbf{L}} \mathcal{P}$ is in $\mathbf{D}^-_{ac.acoh}(\mathbf{P}^N)$.

Lemma 5.2.2. Let R be a universally coherent ring, and let X be a scheme separated and of finite presentation over R. Let $K \in \mathbf{D}_{qc}(X)$. If $\mathbf{R}\Gamma(X, E \otimes_{\mathfrak{O}_X}^{\mathbf{L}} K)$ is in $\mathbf{D}_{acoh}^-(R)$ for every $E \in \mathbf{D}_{coh}^-(X)$, then $K \in \mathbf{D}_{ac.acoh}^-(X)$.

Proof. We follow the proof of [Sta21, Tag 0CSL]. The condition that $K \in \mathbf{D}_{qc,acoh}^{-}(X)$ is local on X as X is quasi-compact. Therefore, we can prove it locally around each point x. We use [Sta21, Tag 0CSJ] to find

- An open subset $U \subset X$ containing x.
- An open subset $V \subset \mathbf{P}_{B}^{n}$.
- A closed subset $Z \subset X \times_R \mathbf{P}_R^n$ with a point $z \in Z$ lying over x

 $^{^{26}}$ Note that they have slightly different notations for R and S

• An object $E \in \mathbf{D}_{coh}^{-}(X \times_{R} \mathbf{P}_{R}^{n})$.

with a lot of properties listed in the cited lemma. Even though the notations are pretty heavy, the only real properties of these object that we will use are that $x \in U$ and

$$\mathbf{R}q_*(\mathbf{L}p^*K \otimes^{\mathbf{L}} E)|_V = \mathbf{R}(U \to V)_*(K|_U)$$

The last formula is proven in [Sta21, Tag 0CSK] and we refer to this lemma for a discussion of the morphism $U \to V$ that turns out to be a finitely presented closed immersion.

That being said, we note that the argument above shows that it is sufficient to show that $K|_U$ is almost coherent for each such U. Moreover, the formula $\mathbf{R}q_*(\mathbf{L}p^*K \otimes^{\mathbf{L}} E)|_V = \mathbf{R}(U \to V)_*(K|_U)$, the fact that $U \to V$ is a finitely presented closed immersion and Lemma 2.7.4 imply that it is sufficient to show that $\mathbf{R}(U \to V)_*(K|_U) = \mathbf{R}q_*(\mathbf{L}p^*K \otimes^{\mathbf{L}} E)|_V$ lies in $\mathbf{D}^-_{qc,acoh}(V)$. In particular, it is enough to show that $\mathbf{R}q_*(\mathbf{L}p^*K \otimes^{\mathbf{L}} E) \in \mathbf{D}^-_{qc,acoh}(\mathbf{P}^n_R)$. The key is that we can check that condition using Theorem 5.2.1.

We define a sheaf $\mathcal{P} \coloneqq \bigoplus_{i=0}^{n} \mathcal{O}_{\mathbf{P}^{n}}(i)$ and we compute

$$\begin{aligned} \mathbf{R} \mathrm{Hom}_{\mathbf{P}^{n}}(\mathcal{P}, \mathbf{R}q_{*}(\mathbf{L}p^{*}K \otimes^{\mathbf{L}} E)) &= \mathbf{R}\Gamma(\mathbf{P}^{n}, \mathbf{R}q_{*}(\mathbf{L}p^{*}K \otimes^{\mathbf{L}} E) \otimes^{\mathbf{L}}_{\mathcal{O}_{\mathbf{P}^{n}}} \mathcal{P}^{\vee}) \\ &= \mathbf{R}\Gamma(\mathbf{P}^{n}, \mathbf{R}q_{*}(\mathbf{L}p^{*}K \otimes^{\mathbf{L}} E \otimes^{\mathbf{L}} \mathbf{L}q^{*}\mathcal{P}^{\vee})) \\ &= \mathbf{R}\Gamma(X \times_{R} \mathbf{P}^{n}_{R}, \mathbf{L}p^{*}K \otimes^{\mathbf{L}} E \otimes^{\mathbf{L}} \mathbf{L}q^{*}\mathcal{P}^{\vee}) \\ &= \mathbf{R}\Gamma(X, \mathbf{R}p_{*}(\mathbf{L}p^{*}K \otimes^{\mathbf{L}} E \otimes^{\mathbf{L}} \mathbf{L}q^{*}\mathcal{P}^{\vee})) \\ &= \mathbf{R}\Gamma(X, K \otimes^{\mathbf{L}}_{\mathcal{O}_{\mathbf{V}}} \mathbf{R}p_{*}(E \otimes^{\mathbf{L}} \mathbf{L}q^{*}\mathcal{P}^{\vee})) \end{aligned}$$

where the second and fifth equality come from the projection formula [Sta21, Tag 08EU]. Finally, we note that the Proper Mapping Theorem 5.1.2 implies that $\mathbf{R}p_*(E \otimes^{\mathbf{L}} \mathbf{L}q^*\mathcal{P}^{\vee}) \in \mathbf{D}^-_{coh}(X)$, so the assumption says that

$$\mathbf{R}\mathrm{Hom}_{\mathbf{P}^{n}}(\mathcal{P}, \mathbf{R}q_{*}(\mathbf{L}p^{*}K \otimes^{\mathbf{L}} E)) = \mathbf{R}\Gamma(X, K \otimes^{\mathbf{L}}_{\mathcal{O}_{X}} \mathbf{R}p_{*}(E \otimes^{\mathbf{L}} \mathbf{L}q^{*}\mathcal{P}^{\vee})) \in \mathbf{D}_{coh}^{-}(R)$$

Now Theorem 5.2.1 finishes the proof.

Theorem 5.2.3. Let R be a universally coherent ring, and let X be a separated, finitely presented R-scheme. Let $\mathcal{F} \in \mathbf{D}^-_{qc}(X)$ be an object such that $\mathbf{R}\operatorname{Hom}_X(\mathcal{P},\mathcal{F}) \in \mathbf{D}^-_{acoh}(R)$ for any $\mathcal{P} \in \operatorname{Perf}(X)$, then $\mathcal{F} \in \mathbf{D}^-_{qc,acoh}(X)$. Analogously, if $\mathbf{R}\operatorname{Hom}_X(\mathcal{P},\mathcal{F}) \in \mathbf{D}^b_{acoh}(R)$ for any $\mathcal{P} \in \operatorname{Perf}(X)$, then $\mathcal{F} \in \mathbf{D}^b_{ac,acoh}(X)$.

Proof. Once we have Lemma 5.2.2 and the equality $\operatorname{RHom}_X(\mathcal{P}, \mathcal{F}) = \operatorname{R}\Gamma(X, \mathcal{P}^{\vee} \otimes_{\mathcal{O}_X}^{\mathbf{L}} \mathcal{F})$, the first part of the Theorem is absolutely analogous to [Sta21, Tag 0CSH]. The second part now follows directly from [Sta21, Tag 09IS] and [BNP17, Lemma 3.0.14].

5.3. The GAGA Theorem. The main goal of this section is to prove the formal GAGA Theorem for almost coherent sheaves. It roughly says that any adically quasi-coherent, almost coherent sheaf on a completion of a proper, finitely presented scheme admits an essentially unique algebraization, and the same holds for morphisms of those sheaves.

We start by recalling the statement of the classical formal GAGA Theorem. We start with a proper A-scheme for some complete adic noetherian ring A with the ideal of definition \mathfrak{m} . Then we consider the \mathfrak{m} -adic completion \mathfrak{X} as a formal scheme over Spf A. It comes with the natural morphism $c: \mathfrak{X} \to X$ of locally ringed spaces that induces a functor

$$c^* \colon \mathbf{Coh}_X \to \mathbf{Coh}_{\mathfrak{X}}$$

The GAGA Theorem says that it is an equivalence of categories. Let us say few words about the "classical" proof of this theorem. There are essentially three independent steps in the proof: the first one is to show that the morphism c is flat; the second one is to show that the functor c^* induces an isomorphism

$$c^* \colon \mathrm{H}^{i}(X, \mathrm{F}) \to \mathrm{H}^{i}(\mathfrak{X}, c^*\mathrm{F})$$

for any any $F \in \mathbf{Coh}_X$ and any integer *i*. And the last one is to prove that any coherent sheaf $\mathcal{G} \in \mathbf{Coh}_{\mathbf{P}^N}$ admits a surjection of the form $\bigoplus_i \mathcal{O}(n_i)^{m_i} \to \mathcal{G}$. Though the first two steps generalizes to our Setup, there is no chance to have any analogue of the last statement. The reason is easy: existence of such a surjection would automatically imply that the sheaf \mathcal{G} is of finite type, however almost coherent sheaves are usually not of finite type.

This issue suggests that we should take another approach to GAGA Theorems recently developed by J. Hall in his paper [Hal18]. The main advantage of this approach is that it firstly *constructs* a candidate for the algebraization, and only then he proves that this candidate works.

We start with the discussion of the GAGA functor. In what follows, we assume that R is a ring from the Setup 5.1.5. We pick a finitely presented R-scheme X, and we consider its ϖ -adic completion \mathfrak{X} that is a topologically finitely presented formal R-scheme. The formal scheme \mathfrak{X} comes equipped with the canonical morphism of locally ringed spaces

$$c\colon (\mathfrak{X}, \mathfrak{O}_{\mathfrak{X}}) \to (X, \mathfrak{O}_X)$$

that induces the functor

$$\mathbf{L}c^*: \mathbf{D}(X) \to \mathbf{D}(\mathfrak{X})$$

We now want to check that this functor "preserves" quasi-coherent, almost coherent objects. That is necessarily even to formulate the GAGA statement.

Lemma 5.3.1. Let R be a ring as in the Setup 5.1.5, and let X be a topologically finitely presented scheme over R. Then the morphism c is flat, and the funtor $c^* \colon \mathbf{Mod}_{\mathcal{O}_X} \to \mathbf{Mod}_{\mathcal{O}_{\mathfrak{X}}}$ sends (quasi-coherent and) almost coherent sheaves to (adically quasi-coherent and) almost coherent sheaves. In particular, it induces functors

$$\mathbf{L}c^*: \mathbf{D}^*_{qc,acoh}(X) \to \mathbf{D}^*_{qc,acoh}(\mathfrak{X})$$

for any $* \in \{", ", +, -, b\}$.

Proof. The flatness assertion is just [FK18, Proposition I.1.4.7 (2)]. Flatness of c implies that it suffices to show that $c^*(G)$ is adically quasi-coherent, almost coherent $\mathcal{O}_{\mathfrak{X}}$ -module for a quasi-coherent, almost coherent \mathcal{O}_X -module G. This claim is Zariski-local on X. Thus we can assume that $X = \operatorname{Spec} A$ is affine, so $G \simeq \widetilde{M}$ for some almost finitely presented A-module M. This case is done in Lemma 4.5.3.

Theorem 5.3.2. Let R be a ring as in the Setup 5.1.5, and let X be a proper, finitely presented scheme over a ring R. Then the functor

$$\mathbf{L}c^*: \mathbf{D}^*_{qc,acoh}(X) \to \mathbf{D}^*_{qc,acoh}(\mathfrak{X})$$

induces an equivalence of categories for $* \in \{", ", +, -, b\}$.

Corollary 5.3.3. Let R and X be as in Theorem 5.3.2. Then the functor

$$\mathbf{L}c^*: \mathbf{D}^*_{acoh}(X)^a \to \mathbf{D}^*_{acoh}(\mathfrak{X})^a$$

induces an equivalence of categories for $* \in \{``",+,-,b\}.$

We follow Jack Hall's proof of the GAGA Theorem very closely with according simplifications due to the flatness of the functor c^* . As he works entirely in the setting of the pseudo-coherent objects, and almost coherent sheaves may not be pseudo-coherent, we repeat some arguments in our setting.

Before going to the proof, we need to define the functor in the other direction. Recall that we always have a functor

$$\mathbf{R}c_*: \mathbf{D}(\mathfrak{X}) \to \mathbf{D}(X)$$

This functor is t-exact as $c: \mathfrak{X} \to X$ is topologically just a closed immersion. In particular, it preserves boundedness of complexes (in any direction). However, that functor usually does not preserve (almost) coherent objects as can be seen in the example of $\mathbf{R}c_*\mathcal{O}_{\mathfrak{X}} = c_*\mathcal{O}_{\mathfrak{X}}$. A way to fix it is to use a "so-called" quasi-coherator functor

$$\mathbf{R}Q_X : \mathbf{D}(X) \to \mathbf{D}_{qc}(X)$$

that is defined as the right adjoint to the inclusion $\iota: \mathbf{D}_{qc}(X) \to \mathbf{D}(X)$. It exists by [Sta21, Tag 0CR0]. So this allows us to define a functor

$$\mathbf{R}c_{qc}\colon \mathbf{D}(\mathfrak{X})\to \mathbf{D}_{qc}(X)$$

as the composition $\mathbf{R}c_{qc} \coloneqq \mathbf{R}Q_X \circ \mathbf{R}c_*$.

Combining the adjunctions $(\mathbf{L}c^*, \mathbf{R}c_*)$ and $(\iota, \mathbf{R}Q_X)$, we conclude that we have a pair of the adjoint functors:

$$\mathbf{L}c^*: \mathbf{D}_{qc}(X) \rightleftharpoons \mathbf{D}(\mathfrak{X}) : \mathbf{R}c_{qc}$$

That gives us the unit and counit morphisms

$$\eta \colon \mathrm{Id} \to \mathbf{R}c_{qc}\mathbf{L}c^* \text{ and } \varepsilon \colon \mathbf{L}c^*\mathbf{R}c_{qc} \to \mathrm{Id}$$

For future reference, we also note that the adjuntion and the monoidal property of the functor $\mathbf{L}c^*$ define a projection morphism

$$\pi_{\mathbf{G},\mathcal{F}} \colon \mathbf{G} \otimes_{\mathcal{O}_X}^{\mathbf{L}} (\mathbf{R}c_{qc}\mathcal{F}) \to \mathbf{R}c_{qc}(\mathbf{L}c^*\mathbf{G} \otimes_{\mathcal{O}_{\mathfrak{X}}}^{\mathbf{L}} \mathcal{F})$$

for any $G \in \mathbf{D}_{qc}(X)$ and any $\mathcal{F} \in \mathbf{D}(\mathfrak{X})$. Before discussing the actual proof of Theorem 5.3.2, we need to establish some formal properties of these functors. In particular, we need to verify that the unit and counit morphisms are isomorphisms in some easy special cases.

Lemma 5.3.4. Let R be a ring as in the Setup 5.1.5, and X a finitely presented R-scheme. Then there is an integer N = N(X) such that $\mathbf{R}c_{qc}$ carries $\mathbf{D}_{qc,acoh}^{\leq n}(\mathfrak{X})$ to $\mathbf{D}_{qc}^{\leq n+N}(X)$ (resp. $\mathbf{D}_{qc,acoh}^{[a,n]}(\mathfrak{X})$ to $\mathbf{D}_{qc}^{[a,n+N]}(X)$) for any integer n. In particular, the natural map

$$\tau^{\geq a} \mathbf{R} c_{qc} \mathcal{F} \to \tau^{\geq a} (\mathbf{R} c_{qc} \tau^{\geq a - N} \mathcal{F})$$

is an isomorphism for any $\mathcal{F} \in \mathbf{D}_{qc,acoh}(\mathfrak{X})$ and any integer a.

Proof. We explain the proof that \mathbf{R}_{cqc} carries $\mathbf{D}_{qc,acoh}^{\leq n}(\mathfrak{X})$ to $\mathbf{D}_{qc}^{\leq n+N}(X)$; the case of $\mathbf{D}_{qc,acoh}^{[a,n]}(\mathfrak{X})$ is similar.

We start the proof by verifying the assumptions of [Sta21, Tag 0CSA] in our setting. Namely, we fix an object $\mathcal{F} \in \mathbf{D}_{qc,acoh}^{\leq n}(\mathfrak{X})$ and show that $\mathrm{H}^{i}(\mathbf{R}\Gamma(U, c_{*}\mathcal{F})) = 0$ for any open affine $U \subset X$ and any $i \geq n$. Indeed, we know that the functor $c_{*} \colon \mathbf{Mod}_{\mathcal{O}_{\mathfrak{X}}} \to \mathbf{Mod}_{\mathcal{O}_{\mathfrak{X}}}$ is exact as c is topologically just a closed immersion. Therefore, we see that

$$\mathrm{H}^{i}(\mathbf{R}\Gamma(U, c_{*}\mathcal{F})) = \mathrm{H}^{i}(\mathbf{R}\Gamma(\widehat{U}, \mathcal{F})) = \mathrm{H}^{i}(\widehat{U}, \mathcal{F}|_{\widehat{U}})$$

Lemma 4.6.11 implies that $\mathrm{H}^{i}(\widehat{U}, \mathcal{F}|_{\widehat{U}}) = 0$ for any $i \geq n$. Moreover, we know that $\mathbf{R}c_{*}\mathcal{F} \in \mathbf{D}^{\leq n}(X)$ as c_{*} is exact on $\mathbf{Mod}_{\mathfrak{X}}$ and $\mathcal{F} \in \mathbf{D}^{\leq n}(\mathfrak{X})$.

Now we apply [Sta21, Tag 0CSA] for $K = \mathbf{R}c_*\mathcal{F}$, $a = -\infty$ and b = n to finish the proof of the first claim in the Lemma. One can check that the proof of that Lemma works well for $a = -\infty$.

The second claim of the lemma follows from the first claim and the distinguished triangle

 $\tau^{\leq a-N-1}\mathcal{F} \to \mathcal{F} \to \tau^{\geq a-N}\mathcal{F} \to \tau^{\leq a-N-1}\mathcal{F}[1]$

Namely, we apply the exact functor $\mathbf{R}c_{qc}$ to this distinguished triangle to get that

$$\mathbf{R}c_{qc}\left(\tau^{\leq a-N-1}\mathcal{F}\right) \to \mathbf{R}c_{qc}\mathcal{F} \to \mathbf{R}c_{qc}\left(\tau^{\geq a-N}\mathcal{F}\right) \to \mathbf{R}c_{qc}\left(\tau^{\leq a-N-1}\mathcal{F}[1]\right)$$

is a distinguished triangle in $\mathbf{D}_{qc}(X)$ and that $\mathbf{R}c_{qc}(\tau^{\leq a-N-1}\mathcal{F}) \in \mathbf{D}_{qc}^{\leq a-1}(X)$. This implies that the map

$$\tau^{\geq a} \mathbf{R} c_{qc} \mathcal{F} \to \tau^{\geq a} \mathbf{R} c_{qc} \left(\tau^{\geq a-N} \mathcal{F} \right)$$

is an isomorphism.

Lemma 5.3.5. Let X be as in Theorem 5.3.2, $\mathcal{F} \in \mathbf{D}^-_{qc,acoh}(\mathfrak{X})$ and $\mathbf{G} \in \mathbf{D}^-_{qc}(X)$. Suppose that for each *i* there is n_i such that $\varpi^{n_i} \mathcal{H}^i(\mathcal{F}) = 0$ and $\varpi^{n_i} \mathcal{H}^i(\mathbf{G}) = 0$. Then the natural morphisms $\eta_{\mathbf{G}}$ and $\varepsilon \mathcal{F}$ are isomorphisms.

Proof. We prove the claim only for \mathcal{F} as the other claim is similar.

Reduction to the case $\mathcal{F} \in \mathbf{D}^{b}_{qc,acoh}(\mathfrak{X})$: First of all, we note that it suffices to show that the natural maps

$$\tau^{\geq a} \mathcal{F} \to \tau^{\geq a} \mathbf{L} c^* \mathbf{R} c_{qc} \mathcal{F}$$

is an isomorphism for any a. Moreover, we note that t-exactness of $\mathbf{L}c^*$ and Lemma 5.3.4 imply that there is an integer N such that the natural map $\tau^{\geq a}\mathbf{L}c^*\mathbf{R}c_{qc}\mathcal{F} \to \tau^{\geq a}\mathbf{L}c^*\mathbf{R}c_{qc}\tau^{\geq a-N}\mathcal{F}$ is an isomorphism for any integer a. In particular, we have a commutative diagram

$$\begin{array}{ccc} \tau^{\geq a-N} \mathcal{F} & \longrightarrow \mathbf{L}c^* \mathbf{R}c_{qc}(\tau^{\geq a-N} \mathcal{F}) \\ & \downarrow & & \downarrow \\ \tau^{\geq a} \mathcal{F} & \longrightarrow \tau^{\geq a} \mathbf{L}c^* \mathbf{R}c_{qc} \mathcal{F} \simeq \tau^{\geq a} \mathbf{L}c^* \mathbf{R}c_{qc} \tau^{\geq a-N} \mathcal{F} \end{array}$$

where the horizontal maps induce isomorphisms in degree $\geq a$. Therefore, it suffices to prove the claim for $\tau^{\geq a-N} \mathcal{F}$. So we may and do assume that \mathcal{F} is bounded.

Proof for a bounded \mathcal{F} : The case of a bounded \mathcal{F} easily reduces to the case of an adically quasicoherent, almost coherent $\mathcal{O}_{\mathfrak{X}}$ -module concentrated in degree 0. In that situation we have an adically quasi-coherent module \mathcal{F} such that $\varpi^{k+1}\mathcal{F} = 0$ for some k. That implies that $\mathcal{F} = i_{k,*}\mathcal{F}_k = \mathbf{R}i_{k,*}\mathcal{F}_k$ for the closed immersion $i_k \colon X_k \to \mathfrak{X}$. Now it is straightforward to see that the canonical map

$$\mathbf{R}_{i_{k,*}} \mathcal{F}_k \to \mathbf{L}c^* \mathbf{R}c_{qc}(\mathbf{R}_{i_{k,*}} \mathcal{F}_k)$$

is an isomorphism. The key is flatness of c and the observation that $\mathbf{R}c_*(\mathbf{R}i_{k,*}\mathcal{F}_k)$ is already quasi-coherent, so the quasi-coherator does nothing in this case.

The other thing we need to check is that the map $\pi_{G,\mathcal{F}}$ is an isomorphism for $\mathcal{G} \in \operatorname{Perf}(X)$. As this statement is proven in [Hal18] without any pseudo-coherence assumption on $\mathcal{F} \in \mathbf{D}(\mathfrak{X})$, we just cite it here.

Lemma 5.3.6. If $G \in D_{ac}(X)$ and $\mathcal{F} \in D(\mathfrak{X})$, then the natural projection morphism

 $\pi_{\mathbf{G}, \mathfrak{F}} \colon \mathbf{G} \otimes^{\mathbf{L}}_{\mathcal{O}_{X}} \mathbf{R}c_{qc} \mathfrak{F} \to \mathbf{R}c_{qc}(\mathbf{L}c^{*}\mathbf{G} \otimes^{\mathbf{L}}_{\mathfrak{O}_{\mathfrak{F}}} \mathfrak{F})$

is an isomorphism if G is perfect.

Proof. [Hal18, Lemma 4.3]

Now we come to the key input ingredient. Even though $\mathbf{R}c_{qc}$ is quite abstract and difficult to compute in practice, it turns out that the Almost Proper Mapping Theorem allows us to check that this functor sends $\mathbf{D}^{-}_{ac.acoh}(\mathfrak{X})$ to $\mathbf{D}^{-}_{ac.acoh}(X)$. That would give us a candidate for the algebraization.

Lemma 5.3.7. Let X be a proper, finitely presented scheme over R that is in the Setup 5.1.5. Then $\mathbf{R}c_{qc}$ sends $\mathbf{D}^*_{qc,acoh}(\mathfrak{X})$ to $\mathbf{D}^*_{qc,acoh}(X)$ for $* \in \{-, b\}$.

Proof. We prove only the bounded below case as the other one is analogous. We pick any $\mathcal{F} \in \mathbf{D}^-_{qc,acoh}(\mathfrak{X})$ and we use Theorem 5.2.3 to say that it is sufficient to show that $\mathbf{R}\operatorname{Hom}_X(\mathbf{P}, \mathbf{R}c_*\mathcal{F}) \in \mathbf{D}^-_{acoh}(R)$ for any perfect complex $\mathbf{P} \in \operatorname{Perf}(X)$. That turns out to be a formal consequence of the Almost Propper Mapping Theorem 5.1.6. Indeed, we have

$$\begin{aligned} \mathbf{R} \mathrm{Hom}_{X}(\mathrm{P}, \mathbf{R} c_{qc} \mathcal{F}) &= \mathbf{R} \mathrm{Hom}_{\mathfrak{X}}(\mathbf{L} c^{*} \mathrm{P}, \mathcal{F}) \\ &= \mathbf{R} \mathrm{Hom}_{\mathfrak{X}}(\mathcal{O}_{\mathfrak{X}}, (\mathbf{L} c^{*} \mathrm{P})^{\vee} \otimes_{\mathcal{O}_{\mathfrak{X}}}^{\mathbf{L}} \mathcal{F}) \\ &= \mathbf{R} \Gamma(\mathfrak{X}, (\mathbf{L} c^{*} \mathrm{P})^{\vee} \otimes_{\mathcal{O}_{\mathfrak{X}}}^{\mathbf{L}} \mathcal{F}) \in \mathbf{D}_{acoh}^{-}(R), \end{aligned}$$

where the last formula comes from the fact that derived pushforward and derived dual operations preserve perfect complexes, and for any $\mathcal{P} \in \operatorname{Perf}(\mathfrak{X})$ we have $\mathcal{P} \otimes_{\mathcal{O}_{\mathfrak{X}}}^{\mathbf{L}} \mathcal{F} \in \mathbf{D}_{ac.acoh}^{-}(\mathfrak{X})$.

Finally, we are ready to give a proof of the GAGA Theorem.

Proof of Theorem 5.3.2. Claim 0: It suffices to show the theorem for * = -, i.e. for bounded above derived categories. Indeed, flatness of c^* implies that $\mathbf{L}c^*$ preserve boundedness (resp. boundedness above, resp. boundedness below), so it suffices to show that the natural morphisms

$$\eta_{\mathbf{G}} \colon \mathbf{G} \to \mathbf{R}c_{qc}\mathbf{L}c^{*}\mathbf{G}$$
$$\varepsilon_{\mathfrak{F}} \colon \mathbf{L}c^{*}\mathbf{R}c_{ac}\mathfrak{F} \to \mathfrak{F}$$

are isomorphisms for any $G \in \mathbf{D}_{qc,acoh}(X)$ and $\mathfrak{F} \in \mathbf{D}_{qc,acoh}(\mathfrak{X})$.

We fix N as in Lemma 5.3.4. Then flatness of c^* and Lemma 5.3.4 guarantee that

$$\mathbf{R}c_{qc}\mathbf{L}c^{*}\tau^{\geq a}\mathbf{G}\in\mathbf{D}^{[a,\infty]}(X)$$
$$\mathbf{L}c^{*}\mathbf{R}c_{qc}\tau^{\geq a}\mathcal{F}\in\mathbf{D}^{[a,\infty]}(\mathfrak{X}).$$

Therefore, we see that $\eta_{\rm G}$ is an isomorphism on \mathcal{H}^i for i < a if and only if the same holds for $\eta_{\tau \geq a_{\rm G}}$. Since a was arbitrary, we conclude that it suffices to show that $\eta_{\rm G}$ is an isomorphism for ${\rm G} \in {\mathbf{D}}^-_{qc,acoh}(X)$. Similar argument shows that it suffices to show that $\varepsilon_{\mathcal{F}}$ is an isomorphism for $\mathcal{F} \in {\mathbf{D}}^-_{ac,acoh}(\mathfrak{X})$. So it suffices to prove the theorem for * = -.

Before we formulate the next claim, we need to use the so-called "approximation by perfect complexes" [Sta21, Tag 08EL] to find some $P \in Perf(X)$ such that $\tau^{\geq 0}P \simeq \mathcal{O}_X/\varpi \simeq \mathcal{O}_{X_0}$ and whose support is equal to X_0 . We note that it implies that all cohomology sheaves $\mathcal{H}^i(P)$ are killed by some power of ϖ . We also denote its (derived) pullback by $\mathcal{P} := \mathbf{L}c^*P$.

Claim 1: If $G \in \mathbf{D}^{-}_{qc,acoh}(X)$ such that $G \otimes^{\mathbf{L}}_{\mathcal{O}_{X}} P \simeq 0$, then $G \simeq 0$. Similarly, if $\mathcal{F} \in \mathbf{D}^{-}_{qc,acoh}(\mathfrak{X})$ such that $\mathcal{F} \otimes^{\mathbf{L}}_{\mathcal{O}_{X}} \mathcal{P} \simeq 0$, then $\mathcal{F} \simeq 0$.

We choose the maximal m (assuming that $G \not\simeq 0$)such that $\mathcal{H}^m(G) \neq 0$. Then we see that $\mathcal{H}^m(G \otimes_{\mathcal{O}_X}^{\mathbf{L}} P) \simeq \mathcal{H}^m(G) \otimes_{\mathcal{O}_X} \mathcal{O}_{X_0} = \mathcal{H}^m(G)/\varpi$. We have $(\mathcal{H}^m(G)/\varpi)(U) = \mathcal{H}^m(G)(U)/\varpi \simeq 0$ on any open affine U. So Nakayama's Lemma 2.5.15 implies that $\mathcal{H}^m(G)(U) \simeq 0$ for any such U. This contradicts the choice of m. The proof in the formal setup is the same once we notice that $\mathcal{H}^0(\mathcal{P}) = \mathcal{O}_{\mathfrak{X}}/\varpi$.

Claim 2: The map $\eta_G \colon G \to \mathbf{R}c_{qc}\mathbf{L}c^*G$ is an isomorphism for any $G \in \mathbf{D}^-_{ac,acoh}(X)$.

Claim 1 implies that it is sufficient to show that the map

$$\varepsilon_{\mathbf{G}} \otimes_{\mathcal{O}_{X}}^{\mathbf{L}} \mathbf{P} \colon \mathbf{G} \otimes_{\mathcal{O}_{X}}^{\mathbf{L}} \mathbf{P} \to \mathbf{R}c_{qc}\mathbf{L}c^{*}\mathbf{G} \otimes_{\mathcal{O}_{X}}^{\mathbf{L}} \mathbf{P}$$
(5.1)

is an isomorphism. Recall that the cohomology sheaves of P are killed by some power of ϖ . This property passes to $G \otimes_{\mathcal{O}_X}^{\mathbf{L}} P$, so we can use Lemma 5.3.5 to get that the map

$$\varepsilon_{\mathbf{G}\otimes_{\mathcal{O}_{X}}^{\mathbf{L}}\mathbf{P}}\colon\mathbf{G}\otimes_{\mathcal{O}_{X}}^{\mathbf{L}}\mathbf{P}\rightarrow\mathbf{R}c_{qc}\left(\mathbf{L}c^{*}\left(\mathbf{G}\otimes_{\mathcal{O}_{X}}^{\mathbf{L}}P\right)\right)$$

is an isomorphism. Now comes the key: we fit the morphism $\varepsilon_{G \otimes_{\mathcal{O}_X}^{\mathbf{L}} P}$ into the following commutative triangle:

$$\begin{array}{c} \mathbf{G} \otimes_{\mathbb{O}_{X}}^{\mathbf{L}} \mathbf{P} \xrightarrow{\varepsilon_{\mathbf{G}} \otimes_{\mathbb{O}_{X}}^{\mathbf{L}} \mathbf{P}} & \mathbf{R}c_{qc}\mathbf{L}c^{*}\mathbf{G} \otimes_{\mathbb{O}_{X}}^{\mathbf{L}} \mathbf{P} \\ \xrightarrow{\varepsilon_{\mathbf{G}} \otimes_{\mathbb{O}_{X}}^{\mathbf{L}} \mathbf{P}} & & & \downarrow^{\pi_{\mathbf{P},\mathbf{L}c^{*}\mathbf{G}}} \\ \mathbf{R}c_{qc}(\mathbf{L}c^{*}(\mathbf{G} \otimes_{\mathbb{O}_{X}}^{\mathbf{L}} P)) \xrightarrow{} & \mathbf{R}c_{qc}(\mathbf{L}c^{*}\mathbf{G} \otimes_{\mathbb{O}_{Y}}^{\mathbf{L}} \mathbf{L}c^{*}\mathbf{P}) \end{array}$$

where the bottom horizontal arrow is the isomorphism map induced by the monoidal structure on $\mathbf{L}c^*$. Moreover, we have already established that the left vertical arrow is an isomorphism, and right vertical arrow is an isomorphism due to Lemma 5.3.6. That shows that the top horizontal must be also an isomorphism.

Claim 3: The map $\varepsilon_{\mathcal{F}} \colon \mathbf{L}c^*\mathbf{R}c_{qc}\mathcal{F} \to \mathcal{F}$ is an isomorphism for any $\mathcal{F} \in \mathbf{D}^-_{ac,acoh}(\mathfrak{X})$.

We use Claim 1 again to say that it is sufficient to show that the map

$$\varepsilon_{\mathfrak{F}} \otimes^{\mathbf{L}}_{\mathcal{O}_{\mathfrak{F}}} \mathbf{L}c^* \mathbf{P} \colon \mathbf{L}c^* \mathbf{R}c_{qc} \mathfrak{F} \otimes^{\mathbf{L}}_{\mathcal{O}_{\mathfrak{F}}} \mathbf{L}c^* \mathbf{P} \to \mathfrak{F} \otimes^{\mathbf{L}}_{\mathcal{O}_{\mathfrak{F}}} \mathbf{L}c^* \mathbf{P}$$

is an isomorphism. But that map fits into the commutative diagram:

where the left vertical morphism is the canonical isomorphism induced by the monoidal structure on $\mathbf{L}c^*$, the bottom morphism is an isomorphism by Lemma 5.3.6, and the right vertical morphism is
an isomorphism by Lemma 5.3.5. This implies that the top horizontal morphism is an isomorphism and that finishes the proof. $\hfill \Box$

5.4. The Formal Function Theorem. We derive certain consequences from the GAGA Theorem established in the previous section. Namely, we prove a version of the Formal Function Theorem for almost coherent sheaves. We note that it is possible to give a proof of it independent of the GAGA Theorem.

Let us introduce the setup for this section. We pick a ring R as in the Setup 5.1.5 and a finitely presented R-algebra A. We note that A and \widehat{A} are also topologically universally adhesive by [FK18, Proposition 0.8.5.19], and they are (topologically universally) coherent by [FK18, Proposition 0.8.5.23]. We consider a proper, finitely presented A-scheme X, and an almost coherent \mathcal{O}_X -module \mathcal{F} . We denote the ϖ -adic completion of X by \mathfrak{X} , so we have a commutative diagram:

$$\begin{array}{c} \mathfrak{X} & \stackrel{c}{\longrightarrow} X \\ & \downarrow \widehat{f} & \qquad \downarrow f \\ \operatorname{Spf}(\widehat{A}) & \longrightarrow \operatorname{Spec} A \end{array}$$

Given this diagram we can consider four different cohomology groups:

$$\mathrm{H}^{i}(\mathfrak{X}, c^{*}\mathcal{F}), \ \mathrm{H}^{\widehat{i}}(X, \widetilde{\mathcal{F}}), \ \mathrm{H}^{i}(X, \mathcal{F}) \otimes_{A} \widehat{A}, \ \mathrm{and} \ \lim_{n} \mathrm{H}^{i}(X_{n}, \mathcal{F}_{n}).$$

All these groups have a natural structure of \widehat{A} -module, and it is straightforward to construct functorial in \mathcal{F} homomorphisms

Theorem 5.4.1. In the notation as above, all the maps $\alpha_{\mathcal{F}}^i$, $\beta_{\mathcal{F}}^i$, $\gamma_{\mathcal{F}}^i$, $\phi_{\mathcal{F}}^i$ are almost isomorphisms for any almost coherent \mathcal{O}_X -module \mathcal{F} . If \mathcal{F} is quasi-coherent, almost coherent, then these maps are isomorphisms.

Proof. Step 0. Reduction to the case of a quasi-coherent, almost coherent sheaf \mathcal{F} : We observe that Lemma 3.3.2, Lemma 3.2.17 and the fact that limits of two almost isomorphic direct systems are almost the same allow us to replace \mathcal{F} with $\tilde{\mathfrak{m}} \otimes \mathcal{F}$ to assume that \mathcal{F} is quasi-coherent and almost coherent.

Step 1. The map $\alpha_{\mathcal{F}}^i$ is an (almost) isomorphism: This is just a consequence of Lemma 2.8.7 as we established in Theorem 5.1.3 that $\mathrm{H}^i(X,\mathcal{F})$ is an almost coherent A-module.

Step 2. The map $\beta_{\mathfrak{F}}^i$ is an (almost) isomorphism: We note that the assumptions on A imply that the map $A \to \widehat{A}$ is flat by [FK18, Proposition 0.8.218]. Thus the flat base change for quasicoherent cohomology groups implies that $\mathrm{H}^i(X, \mathfrak{F}) \otimes_A \widehat{A} \simeq \mathrm{H}^i(X_{\widehat{A}}, \mathfrak{F}_{\widehat{A}})$. Therefore, we may and do assume that A is ϖ -adically complete. Then the map $\mathrm{H}^i(X, \mathfrak{F}) \to \mathrm{H}^i(\mathfrak{X}, c^*\mathfrak{F})$ is an isomorphism by Theorem 5.3.2.

Step 3. The map $\gamma_{\mathcal{F}}^i$ is an (almost) isomorphism: Again, we can reduce to the case when $A = \widehat{A}$ as we did in Step 2. Then we see that $c^*\mathcal{F}$ is adically quasi-coherent, almost coherent by Lemma 5.3.1.

Now we note that $\lim_{n} H^{i}(X_{n}, \mathcal{F}_{n})$ is the completion of the A-module $H^{i}(\mathfrak{X}, c^{*}\mathcal{F})$ with respect to the topology defined by

$$\mathbf{F}^{n}(\mathbf{H}^{i}(\mathfrak{X}, c^{*}\mathfrak{F})) = \mathrm{Im}(\mathbf{H}^{i}(\mathfrak{X}, \varpi^{n}c^{*}\mathfrak{F}) \to \mathbf{H}^{i}(\mathfrak{X}, c^{*}\mathfrak{F}))$$

But the proof of Theorem 5.1.6 implies²⁷ that this topology is equivalent to the ϖ -adic topology. Thus completions coincide, however $\mathrm{H}^{i}(\mathfrak{X}, c^{*}\mathfrak{F})$ is already complete in its ϖ -adic topology due to Theorem 5.1.6 and Lemma 2.8.7.

Step 4. The map $\phi_{\mathcal{F}}^i$ is an (almost) isomorphism: This is a consequence of the three Steps above and commutativity of the diagram (5.2).

We now fix a ring R as in the Setup 5.1.5 and consider a proper, finitely presented R-scheme X with the ϖ -adic completion $c: \mathfrak{X} \to X$. Then, for any $K \in \mathbf{D}_{acoh}(X)$, we have a natural morphism $\beta_K: \mathbf{R}\Gamma(X, K) \to \mathbf{R}\Gamma(\mathfrak{X}, \mathbf{L}c^*K)$. The next theorem asserts that this map is always an almost isomorphism.

Theorem 5.4.2. Let X be as above, and let $K \in \mathbf{D}_{gc,acoh}(X)$. Then the natural map

$$\beta_K \colon \mathbf{R}\Gamma(X, K) \to \mathbf{R}\Gamma(\mathfrak{X}, \mathbf{L}c^*K)$$

is an isomorphism. Moreover, the map β_K is an almost isomorphism for $K \in \mathbf{D}_{acoh}(X)$.

Proof. Step 0: Reduction to the case $K \in \mathbf{D}_{qc,acoh}(X)$: If $K \in \mathbf{D}_{acoh}(X)$ then we use Proposition 3.5.23 and Proposition 3.5.20 to replace K with $\widetilde{\mathfrak{m}} \otimes K \in \mathbf{D}_{ac,acoh}(X)$.

Step 1: Reduce to the case of the $\mathcal{F} \in \mathbf{Acoh}_X \cap \mathbf{Qcoh}_X$: The standard argument using flatness of c and [Sta21, Tag 0D6U] shows that we can reduce to the case of a bounded below complex K. Then the standard argument using the spectral sequence $\mathbf{E}_2^{p,q} = \mathbf{R}^p \Gamma(X, \mathcal{H}^q(K)) \Rightarrow \mathbf{R}^{p+q} \Gamma(X, K)$ and flatness of c reduces the question to the case $K = \mathcal{F} \in \mathbf{Acoh}_X \cap \mathbf{Qcoh}_X$.

Step 2. The case of $\mathcal{F} \in \mathbf{Acoh}_X \cap \mathbf{Qcoh}_X$: This just follows directly from Theorem 5.4.1. \Box

5.5. Almost Version of Grothendieck Duality. For this section, we fix a universally coherent ring R with an ideal \mathfrak{m} such that $\widetilde{\mathfrak{m}} := \mathfrak{m} \otimes_R \mathfrak{m}$ is R-flat and $\mathfrak{m}^2 = \mathfrak{m}$. Since R is universally coherent, there is a good theory of $f^!$ functor for separated morphisms between finitely presented R-schemes²⁸.

Proposition 5.5.1. Let $f: X \to Y$ be a separated morphism of finitely presented *R*-schemes. Then $f^!$ sends $\mathbf{D}^+_{qc,acoh}(Y)$ to $\mathbf{D}^+_{qc,acoh}(X)$.

Proof. The only thing that we need to check here is that $f^!$ preserves almost coherence of cohomology sheaves. This statement is local, so we can assume that both X and Y are affine. Then we can choose a closed embedding $X \to \mathbf{A}_Y^n \to Y$. So, it suffices to prove the claim for a finitely presented closed immersion and for the morphism $\mathbf{A}_Y^n \to Y$.

In the case $f: X \to Y$ a finitely presented closed immersion, we know that

$$f^{!}(\mathcal{F}) \simeq \underline{\mathbf{R}}\mathcal{H}om_{Y}(f_{*}\mathcal{O}_{X},\mathcal{F})$$

 $^{^{27}}$ This claim essentially boils down to Lemma A.4.

 $^{^{28}}$ All proofs from [Sta21, Tag 0DWE] just work as written in this situation, but it does not seem to be addressed in the literature in this generality

for any $\mathcal{F} \in \mathbf{D}_{qc}^+(Y)$. Since Y is a coherent scheme and f is finitely presented, we conclude that $f_*\mathcal{O}_X$ is an almost coherent \mathcal{O}_Y -module. Therefore, $f^!(\mathcal{F}) = \underline{\mathbf{R}}\mathcal{H}om_Y(f_*\mathcal{O}_X, \mathcal{F}) \in \mathbf{D}_{qc,acoh}(X)$ by Corollary 4.3.11.

Now we consider the case of a relative affine space $f: X = \mathbf{A}_Y^n \to Y$. In this case, the formula for $f^!$ is $f^!(\mathfrak{F}) \simeq \mathbf{L} f^* \mathfrak{F} \otimes_{\mathcal{O}_X}^L \Omega^n_{X/Y}[n]$. Then $\mathbf{L} f^*(\mathfrak{F}) \in \mathbf{D}^+_{qc,acoh}(X)$ by Lemma 4.3.7(4), and so $\mathbf{L} f^* \mathfrak{F} \otimes_{\mathcal{O}_X}^L \Omega^n_{X/Y}[n] \in \mathbf{D}^+_{qc,acoh}(X)$ by Lemma 4.3.8(4).

Now we use Proposition 5.5.1 to define the almost version of the upper shrick functor:

Definition 5.5.2. Let $f: X \to Y$ be a separated morphism of finitely presented *R*-schemes. We define $f_a^!: \mathbf{D}_{aqc}^+(Y)^a \to \mathbf{D}_{aqc}^+(X)^a$ as $f_a^!(\mathfrak{F}) \coloneqq (f^!(\mathfrak{F}_!))^a$.

Remark 5.5.3. In what follows, we will usually denote the functor $f_a^!$ simply by $f^!$ as it will not cause any confusion.

Lemma 5.5.4. Let $f: X \to Y$ be a separated morphism of finitely presented *R*-schemes. Then $f^!$ carries $\mathbf{D}^+_{acoh}(Y)^a$ to $\mathbf{D}^+_{acoh}(X)^a$.

Proof. This follows from Proposition 5.5.1.

Theorem 5.5.5. Let $f: X \to Y$ be as above. Suppose that f is proper. Then $f^!: \mathbf{D}^+_{aqc}(Y)^a \to \mathbf{D}^+_{aqc}(X)^a$ is right adjoint to the functor $\mathbf{R}f_*: \mathbf{D}^+_{aqc}(Y)^a \to \mathbf{D}^+_{aqc}(X)^a$.

We note that the theorem makes sense as $\mathbf{R}f_*$ carries $\mathbf{D}^+_{aqc}(X)^a$ into $\mathbf{D}^+_{aqc}(Y)$ by Lemma 4.3.9.

Proof. This follows from a sequence of canonical isomorphisms:

$$\begin{split} \mathbf{R} \mathrm{Hom}_{\mathbf{D}(Y)^{a}}(\mathbf{R} f_{*} \mathcal{F}^{a}, \mathcal{G}^{a}) &\simeq \mathbf{R} \mathrm{Hom}_{\mathbf{D}(Y)}(\widetilde{\mathfrak{m}} \otimes \mathbf{R} f_{*} \mathcal{F}, \mathcal{G}) & \text{Lemma 3.1.13} \\ &\simeq \mathbf{R} \mathrm{Hom}_{\mathbf{D}(Y)}(\mathbf{R} f_{*}(\widetilde{\mathfrak{m}} \otimes \mathcal{F}), \mathcal{G}) & \text{Lemma 3.3.6} \\ &\simeq \mathbf{R} \mathrm{Hom}_{\mathbf{D}(X)}(\widetilde{\mathfrak{m}} \otimes \mathcal{F}, f^{!}(\mathcal{G})) & \text{Grothendieck Duality} \\ &\simeq \mathbf{R} \mathrm{Hom}_{\mathbf{D}(X)^{a}}(\mathcal{F}^{a}, f^{!}(\mathcal{G})^{a}) & \text{Lemma 3.1.13}. \end{split}$$

Now suppose that $f: X \to Y$ be a proper morphism of finitely presented *R*-schemes, $\mathcal{F}^a \in \mathbf{D}^+_{aqc}(X)^a$, and $\mathcal{G}^a \in \mathbf{D}^+_{aqc}(Y)^a$. Then we want to construct a canonical morphism

$$\mathbf{R}f_*\mathbf{R}\underline{al\mathcal{H}om}_X(\mathcal{F}^a, f^!(\mathcal{G}^a)) \to \mathbf{R}\underline{al\mathcal{H}om}_Y(\mathbf{R}f_*(\mathcal{F}^a), \mathcal{G}^a).$$

Lemma 3.5.16 says that such a map is equivalent to a map

$$\mathbf{R}f_*\mathbf{R}\underline{al\mathcal{H}om}_X(\mathcal{F}^a, f^!(\mathcal{G}^a))\otimes^L_{\mathcal{O}_X}\mathbf{R}f_*(\mathcal{F}^a)\to \mathcal{G}^a.$$

We construct the latter map as the composition

$$\mathbf{R}f_*\mathbf{R}\underline{al\mathcal{H}om}_X(\mathcal{F}^a, f^!(\mathcal{G}^a)) \otimes_{\mathcal{O}_X}^L \mathbf{R}f_*(\mathcal{F}^a) \to \mathbf{R}f_*\left(\mathbf{R}\underline{al\mathcal{H}om}_X(\mathcal{F}^a, f^!(\mathcal{G}^a)) \otimes_{\mathcal{O}_X}^L \mathcal{F}^a\right) \to \mathbf{R}f_*f^!\mathcal{G}^a \to \mathcal{G}^a$$

where the first map is induced by the relative cup product ([Sta21, Tag 0B68]), the second map comes from Remark 3.5.15, and the last map is the counit of the ($\mathbf{R}f_*, f^!$)-adjunction.

Lemma 5.5.6. Let $f: X \to Y$ be a proper morphism of finitely presented *R*-schemes, $\mathcal{F}^a \in \mathbf{D}^+_{agc}(X)^a$, and $\mathcal{G}^a \in \mathbf{D}^+_{agc}(Y)^a$. Then the map

$$\mathbf{R}f_*\mathbf{R}\underline{al\mathcal{H}om}_X(\mathcal{F}^a, f^!(\mathcal{G}^a)) \to \mathbf{R}\underline{al\mathcal{H}om}_Y(\mathbf{R}f_*(\mathcal{F}^a), \mathcal{G}^a).$$

is an (almost) isomorphism in $\mathbf{D}^+_{aqc}(X)^a$.

Proof. This follows from [Sta21, Tag 0A9Q] after unravelling all the definitions.

Theorem 5.5.7. Let $f: X \to Y$ be as above. Suppose that f is smooth of pure dimension d. Then $f^{!}(-) \simeq \mathbf{L}f^{*}(-) \otimes_{\mathcal{O}_{X}}^{L} \Omega^{d}_{X/Y}[d]$

Proof. It follows from the corresponding statement in the classical Grothendieck Duality. \Box

We summarize all the results of this section in the following theorem:

Theorem 5.5.8. Let R be a universally coherent ring with an ideal \mathfrak{m} such that $\widetilde{\mathfrak{m}} := \mathfrak{m} \otimes_R \mathfrak{m}$ is R-flat and $\mathfrak{m}^2 = \mathfrak{m}$, and FPS_R be the category of finitely presented, separated R-schemes. Then there is a well-defined functor (-)! from FPS_R into the 2-category of categories such that

- (1) $(X)! = \mathbf{D}^+_{aqc}(X)^a$,
- (2) for a smooth morphism $f: X \to Y$ of pure relative dimension $d, f' \simeq \mathbf{L} f^*(-) \otimes_{\mathcal{O}_X}^L \Omega^d_{X/Y}[d]$.
- (3) for a proper morphism $f: X \to Y$, $f^!$ is right adjoint to $\mathbf{R}f_*: \mathbf{D}^+_{aqc}(X)^a \to \mathbf{D}^+_{aqc}(Y)^a$.

6. Almost Coherence of "p-adic Nearby Cycles"

6.1. Introduction. We study the "Nearby Cycles" sheaves $\mathbf{R}\nu_*\widehat{\mathcal{O}}_X^+$ and $\mathbf{R}\nu_*\mathcal{O}_X^+/p$ for a rigidanalytic space X. It turns out that these complexes have almost coherent cohomology sheaves that are almost concentrated in degrees $[0, \dim X]$. This gives us the main non-trivial example of almost coherent sheaves.

Before giving precise definitions, let us explain our motivation to study these sheaves. We recall that Scholze proved ([Sch13, Theorem 5.1]) that the cohomology groups $\mathrm{H}^{i}(X, \mathbf{F}_{p})$ are finite for any smooth, proper rigid-analytic variety X over an algebraically closed field C of mixed characteristic (0, p). The essential content of his proof is to show that cohomology groups $\mathrm{H}^{i}(X, \mathcal{O}_{X}^{+}/p)$ are almost finitely generated \mathcal{O}_{C}/p -modules for all $i \geq 0$. His proof uses properness in a very elaborate way; namely, he constructs some "good covering" of X by affinoids and then shows that there is enough cancelation in the Čech-to-Derived spectral sequence associated with that covering. We refer to [Sch13, §5] for the details of his proof.

What one can try to do instead is to separate this problem into two different problems. We can choose some admissible formal model \mathfrak{X} of the space X and consider the associated morphism of ringed topoi

$$t: (X_{\text{\acute{e}t}}, \mathbb{O}_X^+) \to (\mathfrak{X}_{\text{Zar}}, \mathbb{O}_{\mathfrak{X}})$$

that induces the morphism

$$t: (X_{\text{\'et}}, \mathfrak{O}_X^+/p) \to (\mathfrak{X}_{\operatorname{Zar}}, \mathfrak{O}_{\mathfrak{X}}/p) = (X_0, \mathfrak{O}_{X_0})$$

where $X_0 := \mathfrak{X} \times_{\operatorname{Spf} \mathfrak{O}_C} \operatorname{Spec} \mathfrak{O}_C / p$ is the reduction scheme of \mathfrak{X} . Then one can write $\mathbf{R}\Gamma(X, \mathfrak{O}_X^+ / p) \simeq \mathbf{R}\Gamma(\mathfrak{X}, \mathbf{R}t_*\mathfrak{O}_X^+ / p)$. The key is that now \mathfrak{X} is proper over $\operatorname{Spf} \mathfrak{O}_K$ by [Lüt90, Lemma 2.6]. Thus the Almost Proper Mapping Theorem 5.1.3 implies that it is sufficient to show that $\mathbf{R}t_*\mathfrak{O}_X^+ / p \in \mathbf{D}_{coh}^+(X)$ in order to get almost finite generation of $\mathbf{R}\Gamma(X, \mathfrak{O}_X^+ / p)$. The main advantage now is that we can study $\mathbf{R}t_*\mathfrak{O}_X^+ / p$ locally on the formal model \mathfrak{X} and this simplifies the argument, we do not need to construct rather elaborate coverings of X in order to get the finiteness result. Actually the only place where we use properness of X is to get properness of the formal model \mathfrak{X} , the rest of the proof is completely local on \mathfrak{X} .

The way we prove almost coherence of $\mathbf{R}t_*\mathcal{O}_X^+/p$ is very close in spirit to [Sch13, Lemma 5.6]. Although we need to be slightly more accurate as we can only work locally on \mathfrak{X} and not on X itself. The main technical tool to avoid this issue is Achinger's result [Ach17, Proposition 6.6.1] that says that in mixed characteristic any étale map from an affinoid space to a closed unit disc can be modified into a finite étale map. Unfortunately, the proof of this claim in [Ach17] is given when the ground field is discretely values, and we need to use this result over an algebraically closed field that is never discretely valued. Thus we provide the reader with a detailed proof of this claim over a general field in Appendix B.

That being said we can finally explain the notations and formulate the main theorem of this Chapter. We recall that Scholze defined a pro-étale site $X_{\text{proét}}$ ([Sch13, Definition 3.3] and [Sch16]), where coverings of X are roughly coverings by means of étale morphisms and infinite towers of finite étale morphisms. It comes with a morphism of sites

$$\lambda \colon X_{\text{pro\acute{e}t}} \to X_{\text{\acute{e}t}}$$

and we define "the completed integral structure sheaf" $\widehat{\mathcal{O}}_X^+ := \lim_n \lambda^{-1}(\mathcal{O}_X^+)/p^n$. In what follows, we slightly abuse notations and denote $\lambda^{-1}(\mathcal{O}_X^+)$ just by \mathcal{O}_X^+ . It turns out ([Sch13, Lemma 4.2]) and Lemma 4.10]) that the natural map $\widehat{\mathcal{O}}_X^+/p^n \to \mathcal{O}_X^+/p^n$ is an isomorphism for any $n \ge 0$,

and $\widehat{\mathcal{O}}_X^+(U) \simeq R^+$ for any affinoid perfectoid $U \in X_{\text{pro\acute{e}t}}$ with $\widehat{U} = \text{Spa}(R, R^+)^{29}$. Moreover, the adjunction $\mathcal{F} \to \mathbf{R}\lambda_*\lambda^{-1}\mathcal{F}$ is a quasi-isomorphism for any $\mathcal{F} \in \text{Shv}(X_{\acute{e}t})$ by [Sch13, Corollary 3.17]. In particular, $\mathbf{R}t_*\mathcal{O}_X^+/p \simeq \mathbf{R}\mu_*\mathcal{O}_X^+/p$. Altogether, we have a commutative diagram of ringed topoi

$$(X_{\text{pro\acute{e}t}}, \widehat{\mathbb{O}}_X^+) \xrightarrow{\nu} (X_{\text{\acute{e}t}}, \mathbb{O}_X^+) \xrightarrow{t} (\mathfrak{X}, \mathbb{O}_{\mathfrak{X}})$$

that also induces a commutative diagram

$$(X_{\text{pro\acute{e}t}}, \mathbb{O}_X^+/p) \xrightarrow{\nu} 30$$

$$\downarrow^{\lambda} \xrightarrow{\nu} (X_{\acute{e}t}, \mathbb{O}_X^+/p) \xrightarrow{t} (\mathfrak{X}, \mathbb{O}_{\mathfrak{X}}/p) = (X_0, \mathbb{O}_{X_0})$$

We refer to Appendix C, especially to Definition C.1, Theorem C.4, Remark C.5 and Definition C.6, for our conventions on the notions of local systems, locally constant sheaves, and lisse sheaves. Here we only recall that the definition of lisse $\hat{\mathbf{Z}}_p$ -sheaves, as this is probably the least standard one.

A pro-étale sheaf $\widehat{\mathbf{Z}}_p$ is defined as $\widehat{\mathbf{Z}}_p \coloneqq \lim_n \underline{\mathbf{Z}}/p^n \mathbf{Z}$ in $\mathbf{Shv}(X_{\text{proét}})$. A lisse $\widehat{\mathbf{Z}}_p$ -sheaf on $X_{\text{proét}}$ is a sheaf \mathbf{L} of $\widehat{\mathbf{Z}}_p$ -modules on $X_{\text{proét}}$, such that locally in $X_{\text{proét}}$, \mathbf{L} is isomorphic to $\widehat{\mathbf{Z}}_p \otimes_{\mathbf{Z}_p} M$, where M is a finitely generated \mathbf{Z}_p -module. We refer to [Sch13, Section 8] for an equivalence of this notion and the classical notion of lisse étale \mathbf{Z}_p -sheaves.

Now we are ready to formulate the main theorem of this Section.

Theorem 6.1.1. Let C be a complete, algebraically closed rank-1 valuation field of mixed characteristic (0, p). Let \mathcal{O}_C be its valuation ring, and let \mathfrak{X} an admissible formal \mathcal{O}_C -scheme with the adic generic fiber $X \coloneqq \mathfrak{X}_C$ of dimension d. We define the ideal of almost mathematics \mathfrak{m} to be the maximal ideal of the valuation ring $\mathfrak{m}_C \subset \mathcal{O}_C^{31}$. Then

$$\mathbf{R}\nu_*(\mathbf{L}\otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p) \in \mathbf{D}_{acoh}^{[0,2d]}(X_0) \text{ and } \mathbf{R}\nu_*(\mathbf{L}\otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p)^a \in \mathbf{D}_{acoh}^{[0,d]}(X_0)^a$$

for an \mathbf{F}_p -local system **L**. Similarly,

$$\mathbf{R}\nu_*(\mathbf{L}\otimes_{\widehat{\mathbf{Z}}_p}\widehat{\mathbb{O}}_X^+)\in\mathbf{D}^{[0,2d]}_{acoh}(\mathfrak{X}) \text{ and } \mathbf{R}\nu_*(\mathbf{L}\otimes_{\widehat{\mathbf{Z}}_p}\widehat{\mathbb{O}}_X^+)^a\in\mathbf{D}^{[0,d]}_{acoh}(\mathfrak{X})^a$$

for any lisse \mathbf{Z}_p -sheaf **L**. More precisely, for an affine admissible $\mathfrak{X} = \operatorname{Spf} A$ with the adic generic fiber X, the natural maps

$$\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\mathbf{F}_{p}} \mathcal{O}_{X}^{+}/p) \to \mathrm{R}^{i}\nu_{*}(\mathbf{L} \otimes_{\mathbf{F}_{p}} \mathcal{O}_{X}^{+}/p) \text{ and } \mathrm{H}^{i}(X, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathcal{O}}_{X})^{\Delta} \to \mathrm{R}^{i}\nu_{*}(\mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathcal{O}}_{X}^{+})$$

are almost isomorphisms.

Moreover, the formation of $\mathrm{R}^{i}\nu_{*}(\mathbf{L}\otimes_{\mathbf{F}_{p}} \mathfrak{O}_{X}^{+}/p)$ (resp. $\mathrm{R}^{i}\nu_{*}(\mathbf{L}\otimes_{\mathbf{Z}_{p}} \widehat{\mathcal{O}}_{X}^{+}))$ almost commute with étale base change, i.e., for any étale morphism $\mathfrak{f}: \mathfrak{Y} \to \mathfrak{X}$, the natural morphism

$$\mathfrak{f}^*\left(\mathrm{R}^i\nu_{\mathfrak{X}_C,*}(\mathbf{L}\otimes_{\mathbf{F}_p}\mathfrak{O}^+_{\mathfrak{X}_C}/p)\right)\to\mathrm{R}^i\nu_{\mathfrak{Y}_C,*}(\mathfrak{f}^*_C\mathbf{L}\otimes_{\mathbf{F}_p}\mathfrak{O}^+_{\mathfrak{Y}_C}/p)$$

²⁹Look at [Sch13, §3] for the definition of \hat{U} .

³⁰As usually, X_0 stands here for the reduction $\mathfrak{X} \times_{\operatorname{Spf} \mathcal{O}_C} \operatorname{Spec} \mathcal{O}_C / p$.

³¹We note that $\widetilde{\mathfrak{m}} \coloneqq \mathfrak{m} \otimes_{\mathfrak{O}_C} \mathfrak{m}$ is naturally isomorphic to \mathfrak{m} in this situation

$$\left(\text{resp. }\mathfrak{f}^*\left(\mathrm{R}^i\nu_{\mathfrak{X}_C,*}(\mathbf{L}\otimes_{\mathbf{Z}_p}\widehat{\mathcal{O}}^+_{\mathfrak{X}_C})\right)\to\mathrm{R}^i\nu_{\mathfrak{Y}_C,*}(\mathfrak{f}^*_C\mathbf{L}\otimes_{\mathbf{Z}_p}\widehat{\mathcal{O}}^+_{\mathfrak{Y}_C})\right)$$

is an almost isomorphism for any \mathbf{F}_p -local system \mathbf{L} (resp. lisse \mathbf{Z}_p -sheaf \mathbf{L} .).

For the rest of the paper we fix an algebraically closed, complete rank-1 valuation field C that has a mixed characteristic (0, p). We also choose some pseudo-uniformizer $\varpi \in \mathcal{O}_C$ such that $\varpi^p = p$.

Corollary 6.1.2. Let $\mathfrak{X} = \operatorname{Spf} A$ be an affine admissible formal \mathcal{O}_C -scheme with the adic generic fiber $X = \mathfrak{X}_C$. Then the cohomology groups $\operatorname{H}^i(X, \widehat{\mathcal{O}}_X^+)$ are almost finitely presented over A. In particular, they are *p*-adically complete and have bounded torsion p^{∞} -torsion.

Proof. This is a straightforward consequence of Theorem 6.1.1, Lemma 2.8.5 and Lemma 2.8.7 \Box

Corollary 6.1.3. Let X be a proper rigid-analytic variety over C of dimension d, and let **L** be an \mathbf{F}_p -local system on X. Then the cohomology groups $\mathrm{H}^i(X, \mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$ are almost finitely generated over \mathcal{O}_C/p , and $\mathrm{H}^i(X, \mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p) \cong^a 0$ for i > 2d.

Proof. We start a proof by choosing some admissible formal model \mathfrak{X} of the space X. It is necessarily proper by [Lüt90, Lemma 2.6]. Now Theorem 6.1.1 implies that

$$\mathbf{R}\nu_*\left(\mathbf{L}\otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p\right) \simeq \mathbf{R}t_*\left(\mathbf{L}\otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p\right) \in \mathbf{D}_{acoh}^{[0,d]}(X_0)^a$$

where $X_0 := \mathfrak{X} \times_{\operatorname{Spf} \mathcal{O}_C} \operatorname{Spec} \mathcal{O}_C / p$ is the reduction of \mathfrak{X} . Recall that the underlying topological spaces of X_0 and $\overline{\mathfrak{X}} := \mathfrak{X} \times_{\operatorname{Spf} \mathcal{O}_C} \operatorname{Spec} \mathcal{O}_C / \mathfrak{m}_C$ are the same. Thus [FK18, Corollary II.10.1.11] implies that X_0 has Krull dimension d. Therefore, Theorem 5.1.3, [Sta21, Tag 0A3G] and Lemma 3.3.6 imply that

$$\mathbf{R}\Gamma(X,\mathbf{L}\otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p) \simeq \mathbf{R}\Gamma\left(X_0,\mathbf{R}\nu_*\left(\mathbf{L}\otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p\right)\right) \in \mathbf{D}_{acoh}^{[0,2d]}(\mathfrak{O}_C)^a$$

Corollary 6.1.4. Let X be a proper rigid variety over C of dimension d. Let **L** be an \mathbf{F}_p -local system. Then the cohomology groups $\mathrm{H}^i(X, \mathbf{L})$ are finite, and $\mathrm{H}^i(X, \mathbf{L}) \simeq 0$ for i > 2d.

Proof. The proof of [Sch13, Theorem 5.1] carries over once we established Corollary 6.1.3.

6.2. The Complex $\mathbf{R}\nu_*(\mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$ is Almost Quasi-Coherent and Commutes with Étale Base Change. We start the proof Theorem 6.1.1 in this Section. Namely, we show that the complex $\mathbf{R}\nu_*(\mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$ is almost quasi-coherent for any \mathcal{O}_C -admissible formal scheme \mathfrak{X} with the adic generic fiber X, and any \mathbf{F}_p -local system \mathbf{L} . And we show that it commutes with étale base change. Throughout this section we do not even require the adic generic fiber to be smooth. We actually show that the canonical map

$$\mathrm{H}^{i}(X,\mathbf{L}\otimes_{\mathbf{F}_{p}}\mathbb{O}_{X}^{+}/p)\to\mathrm{R}^{i}\nu_{*}(\mathbf{L}\otimes_{\mathbf{F}_{p}}\mathbb{O}_{X}^{+}/p)$$

is an almost isomorphism in the case of an affine \mathfrak{X} . We note that a priori almost quasi-coherence of $\mathrm{R}^{i}\nu_{*}(\mathbf{L}\otimes_{\mathbf{F}_{p}} \mathcal{O}_{X}^{+}/p)$ does not say anything about its relation to $\mathrm{H}^{i}(X, \mathbf{L}\otimes_{\mathbf{F}_{p}} \mathcal{O}_{X}^{+}/p)$ as the sheafification is involved in the construction of $\mathrm{R}^{i}\nu_{*}(\mathbf{L}\otimes_{\mathbf{F}_{p}} \mathcal{O}_{X}^{+}/p)$, so it is not clear why it does not destroy any connection with $\mathrm{H}^{i}(X, \mathbf{L}\otimes_{\mathbf{F}_{p}} \mathcal{O}_{X}^{+}/p)$.

Before discussing the proof, we need to recall the notion of a perfectoid pair and "universal covers" of affine schemes. Recall that $\varpi \in \mathcal{O}_C$ is a pseudo-uniformizer with $\varpi^p = p$.

Definition 6.2.1. We say that a complete (C, \mathcal{O}_C) -Tate-Huber pair (R, R^+) is a *perfectoid pair* if R is uniform, and the Frobenius homomorphism $R^{\circ}/\varpi \xrightarrow{x \mapsto x^p} R^{\circ}/p$ is an isomorphism.

We say that a ϖ -torsionfree \mathcal{O}_C -algebra R is *integrally perfectoid* if R is ϖ -torsionfree, and the Frobenius homomorphism $R/\varpi \xrightarrow{x \mapsto x^p} R/p$ is an isomorphism.

Remark 6.2.2. This definition is consistent with [BMS18, Definition 3.5] due to [BMS18, Lemma 3.10]

We now explain the construction of the "universal covering space" of an affinoid space. Suppose we have an admissible³² non-zero \mathcal{O}_C -algebra R such that $R\left[\frac{1}{p}\right]$ has no non-trivial idempotents (so Spec $R\left[\frac{1}{p}\right]$ is connected), and we choose a geometric point \overline{x} : Spec $\Omega \to \text{Spec } R\left[\frac{1}{p}\right]$ for an algebraically closed field Ω . We consider the category of pointed, connected, finite étale Galois morphisms

$$(\operatorname{Spec} R_i, \overline{x_i}) \to (\operatorname{Spec} R\left[\frac{1}{p}\right], \overline{x}).$$

A standard argument shows that this system is filtered (we point out that it crucially uses the connectedness assumption). We define $R' \coloneqq \operatorname{colim}_{i \in I} R_i$ and we get that $\operatorname{Spec} R' = \lim_{i \in I} \operatorname{Spec} R_i$ using that I is filtered. We also define a geometric point $\operatorname{Spec} k(\overline{y}) \to \operatorname{Spec} R'$ as some geometric point above a topological point " $\lim_{i \in I} \overline{x_i}$ ". It is easy to see that $\operatorname{Spec} R'$ is connected as any idempotent must be defined on some finite level. Moreover, we claim that $\operatorname{Spec} R'$ has no non-trivial connected, finite étale covers.

Lemma 6.2.3. The scheme Spec R' does not have any non-trivial connected, finite étale covers.

Proof. It suffices to show that any finite étale cover $f : \operatorname{Spec} R'' \to \operatorname{Spec} R'$ has a section. Since finite étale morphism are finitely presented, and $\operatorname{Spec} R'' = \lim_{i \in I} \operatorname{Spec} R_i$ is a filtered limit of affine schemes, we can use the spreading out techniques from [EGA IV₃] to assume that f comes as a base change of a finite étale morphism $f_i: \operatorname{Spec} R''_i \to \operatorname{Spec} R_i$ for some $i \in I$. It suffices to show that f_i has a section after a pullback along $v_{j,i}: \operatorname{Spec} R_j \to \operatorname{Spec} R_i$ for some $j \geq i$.

We recall that $\operatorname{Spec} R''_i$ has a finite number of connected components by [Sta21, Tag 07VB]. It implies that each connected of $\operatorname{Spec} R''_i$ is open and closed. Therefore, we may and do assume that $\operatorname{Spec} R''_i$ is connected. Now we use the fact that any connected finite étale cover of a connected scheme can be dominated by a connected Galois cover. This means that there is a scheme Spec S and a morphism

$$g: \operatorname{Spec} S \to \operatorname{Spec} R_i''$$

such that the composition

$$\operatorname{Spec} S \xrightarrow{g} \operatorname{Spec} R_i'' \to \operatorname{Spec} R_i \to \operatorname{Spec} R\left[\frac{1}{\pi}\right]$$

is a connected Galois cover of Spec $R\left[\frac{1}{p}\right]$. Thus we can choose some geometric point \overline{z} in Spec S above the geometric point $\overline{x_i}$ in Spec R_i so that the pair (Spec S, \overline{z}) is isomorphism to an object (Spec $R_j, \overline{x_j}$) for some $j \in J$. Then it is clear that f_i has a section after the base change along the map Spec $R_j \to$ Spec R_i , this finishes the proof.

³²Flat and topologically finitely generated

We now define \overline{R} as the integral closure of R in R', it is easy to see that $\overline{R}[\frac{1}{p}] = R'$. We finally set $(\widehat{\overline{R}}[\frac{1}{p}], \widehat{\overline{R}})$ to be the *p*-adic completion of the pair $(\overline{R}[\frac{1}{p}], \overline{R})$. We observe that $\widehat{\overline{R}}$ is open and integrally closed in $\widehat{\overline{R}}[\frac{1}{p}]$ by [Bha, Lemma 5.1.2]. We call these construction the "universal cover" of Spf R. We note that a group

$$\Delta := \pi_1(\operatorname{Spec} R[\frac{1}{p}], \overline{x}) = \operatorname{Gal}(\overline{R}[\frac{1}{p}]/R[\frac{1}{p}]))$$

continuously acts on $(\widehat{\overline{R}}[\frac{1}{p}], \widehat{\overline{R}})$.

Lemma 6.2.4. Let $(\widehat{\overline{R}}[\frac{1}{p}], \widehat{\overline{R}})$ be as above. Then it is a perfectoid pair.

Proof. We showed above that $\widehat{\overline{R}}$ is integrally closed in $\widehat{\overline{R}}[\frac{1}{p}]$. It is clearly open in $\widehat{\overline{R}}[\frac{1}{p}]$, and it is contained in $\widehat{\overline{R}}[\frac{1}{p}]^{\circ}$. Now [Sch12, Lemma 5.3 (iv)] says that $\widehat{\overline{R}}_*$ is *p*-adically complete and so [Bha, Lemma 5.1.2] reads that $\widehat{\overline{R}}_*$ is integrally closed in $\widehat{\overline{R}}[\frac{1}{p}]$. Therefore, [Bha, Proposition 5.2.5 and Proposition 5.2.6] implies that $\widehat{\overline{R}}[\frac{1}{p}]$ is uniform. Finally, [Sch17, Lemma 15.3] shows that $\widehat{\overline{R}}[\frac{1}{p}]$ is a perfectoid algebra, and it finishes the proof.

Lemma 6.2.5. Let $(\widehat{\overline{R}}[\frac{1}{p}], \widehat{\overline{R}})$ be as above. Then the adic space $\widetilde{X} = \text{Spa}(\widehat{\overline{R}}[1/p], \widehat{\overline{R}})$ is connected and simply-connected (i.e. does not have any non-split finite étale covers).

Proof. We firstly prove that \widetilde{X} is connected. It suffices to show that $\overline{\widehat{R}}[1/p]$ does not have non-trivial idempotents. Any idempotent is clearly integral over \mathbb{Z} . So if there was an idempotent, it would lie in $\overline{\widehat{R}}$. But now we observe that \overline{R} is henselian along p as a colimit of p-adically complete rings. The ring $\overline{\widehat{R}}$ is also henselian along p as it is p-adically complete. Now [Ray70, XI, §2, Proposition 1] reads that we have bijections

$$\{0,1\} = \operatorname{Idem}(\overline{R}) = \operatorname{Idem}(\overline{R}/p) = \operatorname{Idem}(\widehat{\overline{R}}/p) = \operatorname{Idem}(\widehat{\overline{R}})$$

We conclude that \widetilde{X} is connected.

Now we show that there are no non-split finite étale covers of \widetilde{X} . Indeed, [Sch12, Proposition 7.6 and Theorem 7.9] reads that $\widetilde{X}_{\text{fét}} \simeq \widehat{\overline{R}}[1/p]_{\text{fét}}$. And [GR03, Proposition 5.4.53] implies that $\widehat{\overline{R}}[1/p]_{\text{fét}} = \overline{R}[1/p]_{\text{fét}}$. Finally, we note that $\overline{R}[1/p] = R'$ and $R'_{\text{fét}}$ does not have any non-split finite étale covers by Lemma 6.2.3.

We define R_i^+ to be the integral closure of R_i in R. Similarly, we define R^+ to be the integral closure of R in R[1/p], and we define $X \coloneqq \operatorname{Spa}(R[1/p], R^+)$. Then we see that the system " $\lim_{i \in U} \operatorname{Spa}(R_i^+[1/p], R_i^+)$ " $\to X$ defines a pro-étale covering of $\operatorname{Spa}(R[1/p], R^+)$ by an affinoid perfectoid space. Moreover, each morphism $\operatorname{Spa}(R_i^+[1/p], R_i^+)$ is actually $\operatorname{Gal}(R_i/R[1/p])$ -torsor. Therefore, the map " $\lim_{i \in I} \operatorname{Spa}(R_i^+[1/p], R_i^+)$ " $\to X$ is a Δ -torsor in $X_{\operatorname{pro\acute{e}t}}$. Moreover, Lemma 6.2.5 states that $\operatorname{Spa}(\overline{R}[1/p], \overline{R})$ is connected and simply-connected, so the \mathbf{F}_p -local system \mathbf{L} becomes isomorphism to $\mathbf{F}_p^{\operatorname{rk}\mathbf{L}}$ on $\operatorname{Spa}(\overline{R}[1/p], \overline{R})$. We say that $\operatorname{Spa}(\overline{R}[1/p], \overline{R})$ is the universal covering space of $\mathfrak{X} = \operatorname{Spf} R$. [Sch13, Lemma 3.16 and Lemma 4.10] imply that the Čech complex for the covering " $\lim_{i \in I} \operatorname{Spa}(R_i^+[1/p], R_i^+)$ " $\to X$ and the pro-étale sheaf $\mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p$ is almost isomorphic to the complex

$$\mathbf{R}\Gamma_{cont}(\Delta, \Gamma(X, \mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)) = \mathbf{R}\Gamma_{cont}(\Delta, \overline{R}/p^{\oplus \operatorname{rk} \mathbf{L}})$$

for some twisted R/p-linear action of Δ on $\overline{R}/p^{\oplus \operatorname{rk} \mathbf{L}}$. Moreover, the map

$$\mathbf{R}\Gamma_{cont}(\Delta, \overline{R}/p^{\oplus \operatorname{rk} \mathbf{L}}) \to \mathbf{R}\Gamma(X, \mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$$

is an almost isomorphism. The main profit of this construction is that it reduces questions about abstract cohomology groups to continuous cohomology of profinite groups. Therefore, the question of showing that $\mathrm{R}^{i}\nu_{*}(\mathbf{L}\otimes_{\mathbf{F}_{p}} \mathcal{O}_{X}^{+}/p)$ is almost quasi-coherent reduces to two different problems: show that this construction "commutes" with open immersions in an appropriate sense, and that continuous cohomology of profinite groups commutes with flat base change. We address both questions below.

Lemma 6.2.6. Let $f: \operatorname{Spf} S \to \operatorname{Spf} R$ be an étale morphism of admissible affine formal \mathcal{O}_C -schemes. And let R_{∞} be a *p*-torsionfree *R*-algebra that is integrally perfected over \mathcal{O}_C . Then $S_{\infty} := S \widehat{\otimes}_{\mathcal{O}_C} R_{\infty}$ is *p*-torsionfree integrally perfected \mathcal{O}_C -algebra.

Proof. Firstly, we note that $R \to S$ is a flat morphism by [FK18, Proposition I.4.8.1], so $S \otimes_R R_{\infty}$ is ϖ -torsion free. Since a *p*-adic completion of a *p*-torsionfree (equivalently ϖ -torsionfree) is *p*-torsionfree, we conclude that $S \otimes_R R_{\infty}$ is ϖ -torsionfree. We see that the only thing we are left to show is that the Frobenius morphism

$$S_{\infty}/\varpi \to S_{\infty}/p$$

is an isomorphism. This map coincides with the map

$$S/\varpi \otimes_{R/\varpi} R_{\infty}/\varpi \to S/p \otimes_{R/p} R_{\infty}/p$$

and we show that the latter is an isomorphism. We consider the commutative diagram

$$\frac{\operatorname{Spec}\left(S\otimes_{R}R_{\infty}\right)/\varpi}{F} \xrightarrow{\Phi_{S}^{*}} \frac{\Phi_{S}^{*}}{\Phi_{R}^{*}\times S} \operatorname{Spec}S_{\infty}/p}$$

$$\int_{f_{\infty}/\varpi} \xrightarrow{\int_{f_{\infty}/\varpi}} \frac{f_{\infty}/\varphi}{\int_{Spec}R_{\infty}/\varphi} \frac{\Phi_{R}^{*}}{\Phi_{R}^{*}} \xrightarrow{\operatorname{Spec}} \frac{f_{\infty}/p}{\Phi_{R}^{*}}$$

where the square is commutative. Moreover, we know that f_{∞}/p and f_{∞}/ϖ are étale morphisms and the Frobenious Φ_R^* an isomorphism. Therefore, the base change map $\operatorname{Spec} S_{\infty}/p \otimes_{R_{\infty}/p} R_{\infty}/\varpi \to$ $\operatorname{Spec} R_{\infty}/\varpi$ is étale, and the base change morphism $\Phi_R^* \times S$ is an isomorphism. Now we also observe that F is étale as a morphism between étale R_{∞}/ϖ -schemes. Therefore, Φ_S^* is étale as well, but it is also a bijective radiciel morphism (since it is an absolute Frobenius). Now we use [SGA 1, Théorème 5.1] to conclude that it must be an isomorphism. \Box

Corollary 6.2.7. Let $f: \operatorname{Spf} S \to \operatorname{Spf} R$ be an étale morphism of admissible affine formal \mathcal{O}_C -schemes. And let $(\widehat{\overline{R}}[1/p], \widehat{\overline{R}})$ be the "universal cover" constructed above. Then

$$\left(\left(S\widehat{\otimes}_R\widehat{\overline{R}}\right)[1/p],S\widehat{\otimes}_R\widehat{\overline{R}}\right)$$

is a perfectoid pair.

Proof. We know that $(\widehat{\overline{R}}[\frac{1}{p}], \widehat{\overline{R}})$ is a perfectoid pair by Lemma 6.2.4. Then [BMS18, Lemma 3.20] implies that $\widehat{\overline{R}}$ is an integral perfectoid. Then Lemma 6.2.6 states that $S \otimes_R \widehat{\overline{R}}$ is also a *p*-torsionfree

integral perfectoid. Moreover, it is integrally closed in $S \widehat{\otimes}_R \widehat{\overline{R}}[1/p]$ by [Bha, Lemma 5.1.2]. Thus [BMS18, Lemma 3.20] guarantees that $\left(\left(S \widehat{\otimes}_R \widehat{\overline{R}}\right)[1/p], S \widehat{\otimes}_R \widehat{\overline{R}}\right)$ is a perfectoid pair.

Corollary 6.2.8. Let $f: \operatorname{Spf} S \to \operatorname{Spf} R$ be an étale morphism of admissible affine \mathcal{O}_C -schemes. Let $\widetilde{X} = \operatorname{Spa}(\widehat{\overline{R}}[1/p], \widehat{\overline{R}})$ and $\widetilde{Y} = \operatorname{Spa}((S \otimes_R \widehat{\overline{R}})[1/p], S \otimes_R \widehat{\overline{R}})$. Then the natural morphism

$$\Gamma(X, \mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p) \otimes_{R/p} S/p \to \Gamma(Y, \mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p)$$

is an almost isomorphism.

Proof. This is follows from [Sch13, Lemma 4.12] and Corollary 6.2.7. Namely, Corollary 6.2.7 reads that the space \tilde{Y} is affinoid perfectoid, and then [Sch13, Lemma 4.12] says that the map

$$\Gamma(\widetilde{X}, \mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p) \otimes_{\overline{R}/p} (\overline{R} \otimes_R S)/p \to \Gamma(\widetilde{Y}, \mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p)$$

is an almost isomorphism. This map is easily identified with the map

$$\Gamma(\widetilde{X}, \mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p) \otimes_{R/p} S/p \to \Gamma(\widetilde{Y}, \mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p)$$

that finishes the proof.

The last thing we need to address before showing almost quasi-coherence of $R\nu_* \mathcal{O}_X^+/p$ is that continuous cohomology of profinite groups commutes with flat base change.

Lemma 6.2.9. Let G be a profinite group, and let M be a discrete R-module that has a continuous R-linear action of G. Suppose that $R \to S$ is a flat homomorphisms of rings. Then the canonical morphism $\operatorname{H}^{i}_{cont}(G, M) \otimes_{R} S \to \operatorname{H}^{i}_{cont}(G, M \otimes_{R} S)$ is an isomorphism for $i \geq 0$.

Proof. This is a combination of two facts: filtered colimits commute with tensor product, cohomology of finite groups commute with flat base change (in particular, invariants commute with base change). Indeed, Lemma follows from a sequence of isomorphisms

$$\begin{aligned} \mathrm{H}^{i}_{cont}(G,M)\otimes_{R}S &\cong (\mathrm{colim}_{H\triangleleft G,\mathrm{open}}\mathrm{H}^{i}(G/H,M^{H}))\otimes_{R}S \\ &\simeq \mathrm{colim}_{H\triangleleft G,\mathrm{open}}(\mathrm{H}^{i}(G/H,M^{H})\otimes_{R}S) \\ &\simeq \mathrm{colim}_{H\triangleleft G,\mathrm{open}}\mathrm{H}^{i}(G/H,M^{H}\otimes_{R}S) \\ &\simeq \mathrm{colim}_{H\triangleleft G,\mathrm{open}}\mathrm{H}^{i}(G/H,(M\otimes_{R}S)^{H}) \\ &\simeq \mathrm{H}^{i}_{cont}(G,M\otimes_{R}S) \end{aligned}$$

Lemma 6.2.10. Let $\mathfrak{f}: \operatorname{Spf} S \to \operatorname{Spf} R$ be an étale morphism of admissible formal \mathcal{O}_C -schemes with generic fiber $\mathfrak{f}_C: Y \to X$. Suppose that **L** is an étale \mathbf{F}_p local system on X. Then the natural morphism

$$\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\mathbf{F}_{p}} \mathfrak{O}_{X}^{+}/p) \otimes_{R/p} S/p \to \mathrm{H}^{i}(Y, \mathbf{L} \otimes_{\mathbf{F}_{p}} \mathfrak{O}_{Y}^{+}/p)$$

is an almost isomorphism.

Proof. We consider the "universal cover" $\widetilde{X} := \operatorname{Spa}(\widehat{\overline{R}}[\frac{1}{p}], \widehat{\overline{R}}) \to X$ with the Galois group Δ . Then the discussion after Lemma 6.2.4 implies that the map

$$\lim_{i \in I} \operatorname{Spa}(R_i^+[1/p], R_i^+) \to \operatorname{Spa}(R[1/p], R^+) = X$$

is a pro-étale cover by an affinoid perfectoid. Moreover, this map is a Δ -torsor, so we saw that canonical map

$$\mathrm{H}^{i}_{cont}(\Delta, M) \to \mathrm{H}^{i}(X, \mathbf{L} \otimes_{\mathbf{F}_{p}} \mathcal{O}^{+}_{X}/p)$$

is an almost isomorphism, where $M \coloneqq \Gamma(\widetilde{X}, \mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$. Now Lemma 6.2.7 implies that the morphism

$$\lim_{i \in I} \operatorname{Spa}((S \otimes_R R_i^+)[1/p], S \otimes_R R_i^+)) \to \operatorname{Spa}(S[1/p], S^+) = Y$$

is a pro-étale cover by an affinoid perfectoid that is a Δ -torsor. So we have a commutative diagram

$$\begin{array}{ccc} \mathrm{H}^{i}_{cont}(\Delta, M) \otimes_{R/p} S/p & \longrightarrow \mathrm{H}^{i}(X, \mathbf{L} \otimes_{\mathbf{F}_{p}} \mathcal{O}^{+}_{X}/p) \otimes_{R/p} S/p \\ & & \downarrow \\ \mathrm{H}^{i}_{cont}(\Delta, M \otimes_{R/p} S/p) & \longrightarrow \mathrm{H}^{i}(Y, \mathbf{L} \otimes_{\mathbf{F}_{p}} \mathcal{O}^{+}_{Y}/p) \end{array}$$

where the horizontal maps are almost isomorphisms by the argument above and Corollary 6.2.8, and the left vertical map is an isomorphism by Lemma 6.2.9. This implies that the right vertical map is also an almost isomorphism, finishing the proof.

Theorem 6.2.11. Let \mathfrak{X} be an admissible formal \mathcal{O}_C -scheme with the adic generic fiber $X = \mathfrak{X}_C$. Let \mathbf{L} be an \mathbf{F}_p -local system. Then the \mathcal{O}_{X_0} -modules $\mathbf{R}^i \nu_* (\mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$ are almost quasi-coherent. More precisely, if $\mathfrak{X} = \operatorname{Spf} R$ is affine, then the natural map

$$\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\mathbf{F}_{p}} \mathfrak{O}_{X}^{+}/p) \to \mathrm{R}^{i}\nu_{*}(\mathbf{L} \otimes_{\mathbf{F}_{p}} \mathfrak{O}_{X}^{+}/p)$$

is an almost isomorphism for any $i \ge 0$.

Proof. The first claim is clearly Zariski-local on \mathfrak{X} , so we can assume that $\mathfrak{X} = \operatorname{Spf} R$ is an affine admissible formal scheme with the reduction $X_0 = \operatorname{Spec} R/p$. We consider the R/p-module $\operatorname{H}^i(X, \mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p)$, the universal property of the tilde-construction implies that we have a natural morphism

$$\mathrm{H}^{i}(X, \widetilde{\mathbf{L}} \otimes_{\mathbf{F}_{p}} \mathfrak{O}_{X}^{+}/p) \to \mathrm{R}^{i} \nu_{*}(\mathbf{L} \otimes_{\mathbf{F}_{p}} \mathfrak{O}_{X}^{+}/p)$$

In order to show that this map is an almost isomorphism, it is clearly sufficient to assume that $\mathfrak{X} = \operatorname{Spf} R$ is connected³³. Moreover, it is suffices to show that for any open formal subscheme $\operatorname{Spf} S = \mathfrak{U} \subset \mathfrak{X}$ with the adic generic fiber $U := \mathfrak{U}_C$, the restriction morphism

$$\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\mathbf{F}_{p}} \mathcal{O}_{X}^{+}/p) \otimes_{R/p} S/p \to \mathrm{H}^{i}(U, \mathbf{L} \otimes_{\mathbf{F}_{p}} \mathcal{O}_{X}^{+}/p)$$

is an almost isomorphism. This follows from Lemma 6.2.10.

Corollary 6.2.12. Let $\mathfrak{f}: \mathfrak{Y} \to \mathfrak{X}$ be an étale morphism of admissible formal schemes. Then the natural morphism

$$\mathfrak{f}^*\left(\mathrm{R}^i\nu_{\mathfrak{X}_C,*}(\mathbf{L}\otimes_{\mathbf{F}_p}\mathfrak{O}^+_{\mathfrak{X}_C}/p)\right)\to\mathrm{R}^i\nu_{\mathfrak{Y}_C,*}(\mathfrak{f}^*_C\mathbf{L}\otimes_{\mathbf{F}_p}\mathfrak{O}^+_{\mathfrak{Y}_C}/p)$$

is an almost isomorphism.

Proof. This follows from Theorem 6.2.11 and Lemma 6.2.10.

120

³³We use here that connected components of Spf R are clopen. This follows from the fact that the underlying topological space is noetherian.

6.3. The Complex $\mathbb{R}\nu_*(\mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p)$ is Almost Coherent for Smooth X. The main goal of this Section is to show that the complex $\mathbb{R}\nu_*(\mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p)$ has almost coherent cohomology sheaves for any admissible formal \mathcal{O}_C -scheme \mathfrak{X} with smooth adic generic fiber, and any \mathbf{F}_p -local system \mathbf{L} . We proceed in three steps: firstly, we show the result for the formal $\widehat{\mathbf{G}}_m^n$, then we deduce the result for affine formal schemes such that the adic generic fiber admits a map to a torus \mathbf{T}_C^n that is a composition of finite étale maps and rational embeddings. Finally, we show the result in general.

Let us mention the main content of each step. The first step really has been already done [Sch13, Lemma 5.5]. The second step was also sketched in [Sch13, Remark 5.7], but we do not need to show almost noetheriannes of the ring $\mathcal{O}_C/p[T_1, \ldots, T_n]$ as almost coherence of $\mathcal{O}_C/p[T_1, \ldots, T_n]$ is sufficient for our purposes. The last step uses Achinger's result ([Ach17, Proposition 6.6.1]) that any étale morphism g: Spa $(A, A^+) \rightarrow \mathbf{D}_C^n$ can be "modified" into a finite étale morphism provided that the residue characteristic of \mathcal{O}_C is p > 0. The proof of this result is given only for rigid spaces over the fraction field of a discrete valuation ring, but we need to apply it in the non-noethrian setup. So Appendix B provides the reader with a detailed proof of this result without any discreteness assumptions.

Now we consider the special case of a torus $\mathfrak{X} = \operatorname{Spf} \mathcal{O}_C \langle T_1^{\pm 1}, \ldots, T_n^{\pm 1} \rangle$. So we set $R^+ := \mathcal{O}_C \langle T_1^{\pm 1}, \ldots, T_n^{\pm 1} \rangle$ and $R_m^+ := \mathcal{O}_C \langle T_1^{\pm 1/p^m}, \ldots, T_n^{\pm 1/p^m} \rangle$. We note that the map $\operatorname{Spf} R_m^+ \to \operatorname{Spf} R^+$ defines a $\mu_{p^m}^n$ -torsor, thus $\mu_{p^m}^n$ continuously acts on R_m^+ by R^+ -linear automorphisms. Now we consider the *R*-algebra

$$R_{\infty}^{+} = \mathcal{O}_{C}\langle T_{1}^{\pm 1/p^{\infty}}, \dots, T_{n}^{\pm 1/p^{\infty}} \rangle = \left(\operatorname{colim}_{n} R_{m}^{+} \right)^{\widehat{}}$$

where $\widehat{}$ stands for the *p*-adic completion. Then we note that there is a continuous *R*-linear action of the group $\Delta_{\infty} := \mathbf{Z}_p(1)^n = T_p(\mu_{p^{\infty}})$ on R_{∞}^+ . We trivialize $\mathbf{Z}_p(1)$ by choosing some compatible system of p^i -th roots of unity $(\zeta_p, \zeta_{p^2}, \zeta_{p^3}, \ldots)$. In order to describe the action of Δ_{∞} on R_{∞}^+ we need the following definition:

Definition 6.3.1. For any $a \in \mathbb{Z}[1/p]$, we define ζ^a as $\zeta^{ap^l}_{p^l}$ whenever $ap^l \in \mathbb{Z}$. It is clear to see that this definition does not depend on a choice of l.

Then we see that k-th basis vector $\gamma_k \in \Delta_{\infty} \simeq \mathbf{Z}_p^n$ acts on R_{∞}^+ as

$$\gamma_k(T_1^{a_1}\dots T_n^{a_n}) = \zeta^{a_k} T_1^{a_1}\dots T_n^{a_n}$$

Lemma 6.3.2. [Sch13, Lemma 5.5] Let R^+ , R^+_{∞} and Δ_{∞} be as above. Then the cohomology groups $\mathrm{H}^i(\Delta_{\infty}, R^+_{\infty}/p)$ are almost coherent R^+/p -modules. And the natural map

$$\mathrm{H}^{i}_{cont}(\Delta_{\infty}, R^{+}_{\infty}/p) \otimes_{R^{+}/p} S^{+}/p \to \mathrm{H}^{i}_{cont}(\Delta_{\infty}, R^{+}_{\infty}/p \otimes_{R^{+}/p} S^{+}/p)$$

is an isomorphism for a *p*-torsionfree R^+ -algebra S^+ and $i \ge 0$.

Proof. We note that R/p is a coherent ring as it is finitely presented over \mathcal{O}_C/p by [FK18, Proposition 0.8.5.24 and Corollary 0.9.2.7]. Thus Theorem 2.6.11 reads that R/p is also almost coherent, so all almost finitely presented modules are almost coherent. Now [BMS18, Lemma 7.3] says that $\mathbf{R}\Gamma_{cont}(\Delta_{\infty}, R_{\infty}^+/p)$ is computed by means of the Koszul complex $K(R_{\infty}^+/p; \gamma_1 - 1, \ldots, \gamma_n - 1)$. Then, similarly to [Bha18, Lemma 4.6], we can write

$$K\left(R_{\infty}^{+}/p;\gamma_{1}-1,\ldots,\gamma_{n}-1\right) = K\left(R^{+}/p;0,0,\ldots,0\right) \oplus \bigoplus_{(a_{1},\ldots,a_{n})\in(\mathbf{Z}[1/p]\cap(0,1))^{n}} K\left(R^{+}/p;\zeta^{a_{1}}-1,\ldots,\zeta^{a_{n}}-1\right)$$

We observe that $\mathrm{H}^{i}(K(R^{+}/p;0,0,\ldots,0)) = \wedge^{i}(R^{+}/p)$ is a free finitely presented R^{+}/p -module. For each $(a_{1},\ldots,a_{n}) \in (\mathbf{Z}[1/p] \cap (0,1))^{n}$, we can assume that a_{1} has the minimal *p*-adic valuation

for the purpose of proving that $K(R_{\infty}^+/p; \gamma_1 - 1, \ldots, \gamma_n - 1)$ is almost coherent. Then [BMS18, Lemma 7.10] implies that $H^i(K(R^+/p; \zeta^{a_1} - 1, \ldots, \zeta^{a_n} - 1))$ is finitely presented over R^+/p and $\zeta^{a_1} - 1$ -torsion module. Note that

$$v_p(\zeta^{a_1} - 1) = v_p(\zeta_{p^l} - 1) = \frac{v(p)}{p^l - p^{l-1}} \to 0$$

where $a_1 = b/p^l$ with gcd(b, p) = 1. Moreover, for any $h \in \mathbb{Z}$, there are only finitely many indexes $(a_1, \ldots, a_n) \in (\mathbb{Z}[1/p] \cap (0, 1))^n$ with $v_p(a_j) \ge h$. This implies that

$$\mathrm{H}_{cont}^{i}(\Delta_{\infty}, R_{\infty}^{+}/p) = \mathrm{H}^{i}\left(K\left(R_{\infty}^{+}/p; \gamma_{1}-1, \ldots, \gamma_{n}-1\right)\right)$$

is a finitely presented R^+/p -module up to any p^{ε} -torsion. In particular, this module is almost finitely presented.

Now we show that $\operatorname{H}^{i}_{cont}(\Delta_{\infty}, R^{+}_{\infty}/p)$ commutes with base change for any \mathcal{O}_{C} -flat algebra S^{+} . In order to show this, we observe the $(R^{+}/p)[\Delta_{\infty}]$ -module R^{+}_{∞}/p comes as a tensor product $M \otimes_{\mathcal{O}_{C}/p} R^{+}/p$ for the $(\mathcal{O}_{C}/p)[\Delta_{\infty}]$ -module

$$M \coloneqq \bigoplus_{(a_1,\dots,a_n) \in (\mathbf{Z}[1/p] \cap [0,1))^n} (\mathfrak{O}_C/p) T_1^{a_1} \dots T_n^{a_n}$$

where the basis element γ_k acts by

$$\gamma_k(T_1^{a_1}\dots T_n^{a_n}) = \zeta^{a_k} T_1^{a_1}\dots T_n^{a_n}$$

Therefore, Lemma 6.2.9 implies that

$$\begin{aligned} \mathrm{H}^{i}_{cont}(\Delta_{\infty}, R^{+}_{\infty}/p) \otimes_{R^{+}/p} S^{+}/p &\simeq \left(\mathrm{H}^{i}_{cont}(\Delta_{\infty}, M) \otimes_{\mathbb{O}_{C}/p} R^{+}/p\right) \otimes_{R^{+}/p} S^{+}/p \\ &\simeq \mathrm{H}^{i}_{cont}(\Delta_{\infty}, M) \otimes_{\mathbb{O}_{C}/p} S^{+}/p \\ &\simeq \mathrm{H}^{i}_{cont}(\Delta_{\infty}, M \otimes_{\mathbb{O}_{C}/p} S^{+}/p) \\ &\simeq \mathrm{H}^{i}_{cont}(\Delta_{\infty}, R_{\infty}/p \otimes_{R^{+}/p} S^{+}/p) \end{aligned}$$

We will also need a slightly refined version of [Sch13, Lemma 4.5] specific to the situation of an étale morphism $\operatorname{Spa}(S, S^+) \to \mathbf{T}^n$. We define R (resp. R_m, R_∞) as $R^+[1/p]$ (resp. $R_m^+[1/p]$, $R_\infty^+[1/p]$). We note that the system $\mathbf{T}_\infty^n := \lim_m \operatorname{Spa}(R_m, R_m^+)^m \to \mathbf{T}^n = \operatorname{Spa}(R, R^+)$ is a pro-étale covering with $\widehat{\mathbf{T}}_\infty^n = \operatorname{Spa}(R_\infty, R_\infty^+)$ an affinoid perfectoid space.

Lemma 6.3.3. Let R be a topologically finitely presented \mathcal{O}_C -algebra, and P a topologically free R-module, i.e. $P = \bigoplus_{I} R$ for some set I. Then M is R-flat.

Proof. We start the proof by noting that [Sta21, Tag 00M5] guarantees that it suffices to show that $\operatorname{Tor}_{1}^{R}(P, M) = 0$ for any finitely presented *R*-module *M*. We choose a presentation

$$0 \to K \to R^n \to M \to 0$$

and observe that K is finitely presented because R is coherent. So vanishing of Tor₁ is equivalent to showing that

$$P \otimes_R K \to P \otimes_R R^n$$

is injective.

Now note that $K[p^{\infty}]$, $R^n[p^{\infty}]$, and $M[p^{\infty}]$ are bounded by [Bos14, Lemma 7.3/7], so the same holds for $\oplus_I K$, $\oplus_I R^n$, and $\oplus_I M$. Therefore, the usual *p*-adic completions of $\oplus_I K$, $\oplus_I R^n$ and $\oplus_I M$ coincide with their derived *p*-adic completions. Since derived *p*-adic completion is exact (in the

sense of triangulated categories) and coincides with the usual one on these modules, we get that the sequence

$$0 \to \widehat{\oplus}_I K \to \widehat{\oplus}_I R^n \to \widehat{\oplus}_I M \to 0$$

is exact.

Now we want to show that this short exact sequence is the same as the sequence

$$P \otimes_R K \to P \otimes_R R^n \to P \otimes_R M \to 0.$$

As a consequence, this will prove that $P \otimes_R K \to P \otimes_R R^n$ is injective.

For each R-module N, there is a canonical map

$$P \otimes_R N \to \widehat{\oplus}_I N.$$

So we have a morphism of sequences:

$$P \otimes_R K \longrightarrow P \otimes_R R^n \longrightarrow P \otimes_R M \longrightarrow 0$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$0 \longrightarrow \widehat{\oplus}_I K \longrightarrow \widehat{\oplus}_I R^n \longrightarrow \widehat{\oplus}_I M \longrightarrow 0.$$

The map $R^n \otimes_R P \to \widehat{\oplus}_I R^n$ is an isomorphism because $R^n \otimes_R P = P^n$ is already *p*-adically complete. This implies that the arrow

$$M \otimes_R P \to \widehat{\oplus}_I M$$

is surjective. But then

$$P \otimes_R K \to \widehat{\oplus}_I K$$

is surjective since M was an arbitrary finitely presented R-module. Now a diagram chase implies that

$$M \otimes_R P \to \widehat{\oplus}_I M$$

is also injective. And, therefore, it is an isomorphism. So

$$P \otimes_R K \to \widehat{\oplus}_I K$$

is also an isomorphism. Therefore, these two sequences are the same. In particular, $P \otimes_R K \rightarrow P \otimes_R R^n$ is injective.

Lemma 6.3.4. [Sch13, Lemma 4.5] Let $\text{Spa}(S, S^+) \to \text{Spa}(R, R^+) = \mathbf{T}^n$ be a morphism that is a composition of a finite étale maps and rational embeddings. Then the pullback system

$$\lim_{m} \operatorname{Spa}(S_m, S_m^+) \cong \mathbf{T}_{\infty}^n \times_{\mathbf{T}^n} \operatorname{Spa}(S, S^+) \to \operatorname{Spa}(S, S^+)$$

is a pro-étale cover of $\text{Spa}(S, S^+)$ by an affinoid perfectoid space. So, the Tate-Huber pair

$$(S_{\infty}, S_{\infty}^{+}) \coloneqq \left(\operatorname{colim}_{m}\left(S_{m}, S_{m}^{+}\right)\right)^{\gamma}$$

is a perfectoid pair. Moreover, for any $\varepsilon \in \mathbf{Q}_{>0}$, there exists m such that the morphism

$$S_m^+ \widehat{\otimes}_{R_m^+} R_\infty^+ \to S_\infty^+$$

is injective with cokernel annihilated by p^{ε} .

Proof. We note that [Sch13, Lemma 4.5] proves that " $\lim_m \operatorname{Spa}(S_m, S_m^+)'' \to \operatorname{Spa}(S, S^+)$ is a proétale cover by an affinoid perfectoid. We denote by A_m the *p*-adic completion of *p*-torsionfree quotient of $S_m^+ \widehat{\otimes}_{R_m^+} R_\infty^+$. Then [Sch13, Lemma 4.5] shows that, for any $\varepsilon \in \mathbf{Q}_{>0}$, there exists *m* such that the map $A_m \to S_\infty^+$ has cokernel annihilated by p^{ε} . Moreover, the map becomes an isomorphism after inverting *p*. We observe that this implies that $A_m \to S_\infty^+$ is injective as the kernel should be p^{∞} -torsion, but the *p*-adic completion of a *p*-torsionfree ring is *p*-torsionfree. Thus the only thing we need to show is that $S_m^+ \otimes_{R_m^+} R_\infty^+$ is already *p*-torsionfree for any *m*. We note that R_∞^+ is a topologically free as R_m^+ -module because

$$R_{\infty}^{+} = \mathcal{O}_{C}\langle T_{1}^{\pm 1/p^{\infty}}, \dots, T_{n}^{\pm 1/p^{\infty}} \rangle = \widehat{\bigoplus}_{(b_{1},\dots,b_{n})\in\mathbf{Z}^{n}\setminus m\mathbf{Z}^{n}} \mathcal{O}_{C}\langle T_{1}^{\pm 1/p^{m}},\dots, T_{n}^{\pm 1/p^{m}} \rangle T_{1}^{1/p^{b_{1}}}\dots T_{n}^{1/p^{b_{n}}}$$
$$= \widehat{\bigoplus}_{(b_{1},\dots,b_{n})\in\mathbf{Z}^{n}\setminus m\mathbf{Z}^{n}} R_{m}^{+} \cdot T_{1}^{1/p^{b_{1}}}\dots T_{n}^{1/p^{b_{n}}}.$$

Thus, R_{∞}^+ is R_m^+ -flat for any m by Lemma 6.3.3. Therefore, $S_m^+ \otimes_{R_m^+} R_{\infty}^+$ is flat over S_m^+ , so it is, in particular, \mathcal{O}_C -flat. As a consequence, it does not have any non-zero p-torsion. This finishes the proof.

Lemma 6.3.5. Let $X = \text{Spa}(S, S^+)$ be an affinoid smooth rigid *C*-space with an étale morphism $f: X \to \mathbf{T}^n = \text{Spa}(R, R^+)$ that factors as a composition of finite étale morphisms and rational embeddings. Then the cohomology groups

$$\mathrm{H}^{i}(X, \mathcal{O}_{X}^{+}/p)$$

are almost coherent S^+/p -modules for $i \ge 0$.

Proof. First of all, we note that [BGR84, §6.4.1, Corollary 5] implies that $S^+ = S^\circ$ is a flat, topologically finitely type \mathcal{O}_C -algebra. So S^+/p is coherent by [FK18, Proposition 0.8.5.24 and Corollary 0.9.2.7], and thus it is almost coherent by Lemma 2.6.11. So it is sufficient to show that $\mathrm{H}^i_{\mathrm{\acute{e}t}}(X, \mathcal{O}^+_X/p)$ is almost finitely presented over S^+/p .

We choose a pro-étale affinoid perfectoid cover $\mathbf{T}_{\infty}^n \to \mathbf{T}^n$ that is constructed before Lemma 6.3.4. We note that it is Δ_{∞} -torsor, so its base change

$$X_{\infty} = \lim_{m} \operatorname{Spa}(S_m, S_m^+)^{"} \coloneqq \mathbf{T}_{\infty}^n \times_{\mathbf{T}^n} \operatorname{Spa}(S, S^+) \to \operatorname{Spa}(S, S^+)$$

is also a Δ_{∞} -torsor. And X_{∞} is an affinoid perfectoid by Lemma 6.3.4. Thus [Sch13, Lemma 3.16 and Lemma 4.10] imply that the Čech complex for the covering " $\lim_{m} \operatorname{Spa}(S_m, S_m^+)$ " $\to X$ and the pro-étale sheaf \mathcal{O}_X^+/p is almost isomorphic to the complex

$$\mathbf{R}\Gamma_{cont}(\Delta_{\infty}, \Gamma(X_{\infty}, \mathbb{O}_X^+/p)) \cong^a \mathbf{R}\Gamma_{cont}(\Delta_{\infty}, S_{\infty}^+/p)$$

Moreover, the natural map $\mathbf{R}\Gamma_{cont}(\Delta_{\infty}, S_{\infty}^+/p) \to \mathbf{R}\Gamma(X, \mathcal{O}_X^+/p)$ is an almost isomorphism. So we reduce the problem to showing that the complex $\mathbf{R}\Gamma_{cont}(\Delta_{\infty}, S_{\infty}^+/p)$ has cohomology modules that are almost finitely presented over S/p.

Now we pick any $\varepsilon \in \mathbf{Q}_{>0}$ and use Lemma 6.3.4 to find m such that the map

$$S_m^+ \widehat{\otimes}_{R_m^+} R_\infty^+ \to S_\infty^+$$

is injective with cokernel killed by p^{ε} . Thus we conclude that the map

$$S_m^+/p \otimes_{R_m^+/p} R_\infty^+/p \to S_\infty^+/p$$

has kernel and cokernel annihilated by p^{ε} . Then it is clear that the induced map

$$\mathrm{H}^{i}_{cont}(\Delta_{\infty}, S^{+}_{m}/p \otimes_{R^{+}_{m}/p} R^{+}_{\infty}/p) \to \mathrm{H}^{i}_{cont}(\Delta_{\infty}, S^{+}_{\infty}/p)$$

has kernel and cokernel annihilated by $p^{2\varepsilon}$ for any $i \ge 0$. Therefore, Lemma 2.5.7 implies that it is sufficient to show $\mathrm{H}^{i}(\Delta_{\infty}, S_{m}^{+}/p \otimes_{R_{m}^{+}/p} R_{\infty}^{+}/p)$ is almost finitely presented over S^{+}/p for any $m \ge 0$ and any $i \ge 0$.

The trick now is to consider the subgroup $p^m \Delta_{\infty}$ that acts trivially on S_m^+/p to pull it out of cohomology group by Lemma 6.3.2. So we consider the Hochschild–Serre spectral sequence

$$\mathbf{E}_{2}^{i,j} = \mathbf{H}^{i}(\Delta_{\infty}/p^{m}\Delta_{\infty}, \mathbf{H}^{j}_{cont}(p^{m}\Delta_{\infty}, S^{+}_{m}/p \otimes_{R^{+}_{m}/p} R^{+}_{\infty}/p)) \Rightarrow \mathbf{H}^{i+j}_{cont}(\Delta_{\infty}, S^{+}_{m}/p \otimes_{R^{+}_{m}/p} R^{+}_{\infty}/p)$$

Now we recall that there is an explicit bar complex computing group cohomology for any finite group G acting on an S^+/p -module M. Namely, the complex looks like

$$C^0(G, M) \xrightarrow{d^0} C^1(G, M) \xrightarrow{d^1} \dots$$

where

$$\mathbf{C}^{i}(G,M) = \left\{ f: G^{i} \to M \right\} \simeq M^{\oplus i \cdot \# G}$$

and

$$d^{i}(f)(g_{0},g_{1},\ldots,g_{i}) = g_{0} \cdot f(g_{1},\ldots,g_{i}) + \sum_{j=1}^{i} (-1)^{j} f(g_{0},\ldots,g_{j-2},g_{j-1}g_{j},g_{j+1},\ldots,g_{i}) + (-1)^{i+1} f(g_{0},\ldots,g_{i-1}).$$

We note that all terms $C^i(G, M)$ have a natural structure of S^+/p -module, if G acts S^+/p -linearly on M. Moreover, they are finite direct sums of M as an S^+/p -module. In particular, they are almost coherent, if so is M. Thus Lemma 2.6.7 guarantees that all cohomology groups $H^i(G, M)$ are almost coherent over S^+/p provided that M is almost coherent over S^+/p .

We now apply it (together with Lemma 2.6.7) for

$$G = \Delta_{\infty}/p^m \Delta_{\infty}$$
 and $M = \mathrm{H}^{\mathfrak{I}}_{cont}(p^m \Delta_{\infty}, S^+_m/p \otimes_{R^+_m/p} R^+_{\infty}/p)$

to conclude that it is sufficient to show $\operatorname{H}_{cont}^{j}(p^{m}\Delta_{\infty}, S_{m}^{+}/p \otimes_{R_{m}^{+}/p} R_{\infty}^{+}/p)$ is almost coherent over S^{+}/p for any $j \geq 0, m \geq 0$. We note that S_{m}^{+} is finite over S^{+} by [BGR84, §6.4.1, Corollary 5]. Since both are \mathcal{O}_{C} -flat, we see that it is finitely presented by [Bos14, Theorem 7.3/4]. So Lemma 2.7.3 implies that it is enough to show that $\operatorname{H}_{cont}^{j}(p^{m}\Delta_{\infty}, S_{m}^{+}/p \otimes_{R_{m}^{+}/p} R_{\infty}^{+}/p)$ is almost coherent over S_{m}^{+}/p for $i \geq 0$ and $m \geq 0$. But now we can use Lemma 6.3.2 to write

$$\mathrm{H}^{j}_{cont}(p^{m}\Delta_{\infty}, S^{+}_{m}/p \otimes_{R^{+}_{m}/p} R^{+}_{\infty}/p) \simeq \mathrm{H}^{j}_{cont}(p^{m}\Delta_{\infty}, R^{+}_{\infty}/p) \otimes_{R^{+}_{m}/p} S^{+}_{m}/p$$

Moreover, Lemma 6.3.2 guarantees that $\mathrm{H}^{j}_{cont}(p^{m}\Delta_{\infty}, R^{+}_{\infty}/p)$ is almost finitely presented over R^{+}_{m}/p . Thus $\mathrm{H}^{j}_{cont}(p^{m}\Delta_{\infty}, R^{+}_{\infty}/p) \otimes_{R^{+}_{m}/p} S^{+}_{m}/p$ is almost finitely presented over S^{+}_{m}/p by Lemma 2.7.1. \Box

Corollary 6.3.6. Let $X = \text{Spa}(S, S^+)$ be as in Lemma 6.3.5, and let **L** be an \mathbf{F}_p -local system on X. Then the cohomology groups $\mathrm{H}^i(X, \mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$ are almost coherent over S^+/p .

Proof. We may and do assume that X is connected, so **L** has constant rank r. Then we choose some finite Galois cover $X' \to X$ that splits **L**. Denote its Galois group by G, then we have the Hochschild–Serre spectral sequence

$$\mathbf{E}_{2}^{i,j} = \mathbf{H}^{i}\left(G, \mathbf{H}^{j}\left(X', \left(\mathcal{O}_{X'}^{+}/p\right)^{r}\right)\right) \Rightarrow \mathbf{H}^{i+j}(X, \mathbf{L} \otimes_{\mathbf{F}_{p}} \mathcal{O}_{X}^{+}/p)$$

Similarly to the proof of Lemma 6.3.5, the argument with the explicit bar complex computing $H^{i}(G, -)$, [BGR84, §6.4.1, Corollary 5], Lemma 2.7.3 and Lemma 2.6.7 implies that it is sufficient to

show that $\mathrm{H}^{j}(X', \mathcal{O}^{+}_{X'}/p)$ is almost coherent over $\mathcal{O}^{+}_{X'}(X')/p$ for $j \geq 0$. But this is just Lemma 6.3.5.

Theorem 6.3.7. Let \mathfrak{X} be an admissible formal \mathcal{O}_C -scheme with the smooth adic generic fiber $X := \mathfrak{X}_C$ and the reduction scheme $X_0 := \mathfrak{X} \times_{\operatorname{Spf} \mathcal{O}_C} \operatorname{Spec} \mathcal{O}_C / p$. Then

$$\mathbf{R}\nu_*(\mathbf{L}\otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p) \in \mathbf{D}_{acoh}^+(X_0)$$

for any \mathbf{F}_p -local system \mathbf{L} .

Proof. Firstly, we note that \mathfrak{X} is rig-smooth in terminology of [BLR95, §3]. Thus, [BLR95, Proposition 3.7] states that there is an admissible blow-up $\pi: \mathfrak{X}' \to \mathfrak{X}$ and a covering of \mathfrak{X}' by open affine formal subschemes \mathfrak{U}'_i with "rig-étale" morphisms $g_i: \mathfrak{U}'_i \to \widehat{\mathbf{A}}^{n_i}_{\mathcal{O}_C}$, i.e. the adic generic fiber $g_{i,C}: \mathfrak{U}'_{i,C} \to \mathbf{D}^{n_i}$ is étale. We consider the commutative diagram



So we get that $\mathbf{R}\nu_*(\mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p) = \mathbf{R}\pi_* \circ \mathbf{R}\nu'_*(\mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$. The Almost Proper Mapping Theorem 5.1.3 implies that it is sufficient to show that $\mathbf{R}\nu'_*(\mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$ is almost coherent.

Theorem 6.2.11 states that $\mathbb{R}^{j}\nu'(\mathbf{L}\otimes_{\mathbf{F}_{p}} \mathbb{O}_{X}^{+}/p)$ is quasi-coherent for $j \geq 0$. And the restriction on \mathfrak{U}_{i}' is almost isomorphic to $\mathbb{H}^{j}(\mathcal{U}_{i}', \mathbf{L}\otimes_{\mathbf{F}_{p}} \mathbb{O}_{X}^{+}/p)$, where $\mathcal{U}_{i}' \coloneqq \mathfrak{U}_{i,C}'$ is the adic generic fiber. Thus the only thing we are left to show is that these cohomology groups are almost coherent over $\mathcal{O}(\mathfrak{U}_{i}')/p$. Thus we can and do assume that $\mathfrak{X} = \mathfrak{X}' = \operatorname{Spf} A$ is affine and admits a morphism $g \colon \mathfrak{X} \to \widehat{\mathbf{A}}_{\mathbb{O}_{C}}^{n}$ that is étale over the generic fiber.

Now we do the trick that allows us to reduce the question to Corollary 6.3.6. Namely, we use Theorem B.4 to see that we may assume that $g: \mathfrak{X} \to \widehat{\mathbf{A}}_{\mathbb{O}_C}^n$ is finite. In particular, the generic fiber $g_C: X \to \mathbf{D}^n$ is a finite étale morphism. Observe that the closed unit disc \mathbf{D}^n is a rational subset in the *n*-dimensional toris \mathbf{T}^n . Indeed, $\mathbf{T}^n(\frac{T_1-1}{p},\ldots,\frac{T_n-1}{p}) \simeq \mathbf{D}^n$, where T_i are the "standard" coordinates on \mathbf{T}^n . In particular, X admits an étale map to \mathbf{T}^n that factors as a composition of finite étale maps and rational embeddings. Thus Lemma 6.3.6 implies that $\mathrm{H}^i(X, \mathbf{L} \otimes_{\mathbf{F}_p} \mathbb{O}_X^+/p)$ is almost coherent over $\mathbb{O}_X^+(X)/p$. Finally, [BGR84, §6.4.1, Corollary 5] and [Bos14, Theorem 7.3/4] imply that $\mathbb{O}_X^+(X) = \mathbb{O}_X^\circ(X)$ is finitely presented over $\mathbb{O}_{\mathfrak{X}}(\mathfrak{X})$. Thus Lemma 2.7.3 guarantees that $\mathrm{H}^i(X, \mathbf{L} \otimes_{\mathbf{F}_p} \mathbb{O}_X^+/p)$ is almost coherent over $\mathbb{O}_{\mathfrak{X}}(\mathfrak{X})/p$. \Box

Corollary 6.3.8. Let $\text{Spa}(A, A^+)$ be an affinoid rigid space over $\text{Spa}(C, \mathcal{O}_C)$ and $f: X \to \text{Spa}(A, A^+)$ be a proper morphism with smooth X. Suppose **L** is an \mathbf{F}_p -local system on X. Then the cohomology groups $\text{H}^i(X, \mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$ are almost coherent over A_0/p for any topologically finite type ring of definition $A_0 \subset A^+$ and $i \geq 0$.

Proof. We choose some admissible formal model \mathfrak{X} of X with map $g: \mathfrak{X} \to \text{Spf } A_0$ such that $g_C = f$. This is possible by [BL93, Assertion (c) on p.307], moreover the map g is proper by [Lüt90, Lemma 2.6]. Now we can compute

 $\mathbf{R}\Gamma(X,\mathbf{L}\otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p) \simeq \mathbf{R}\Gamma(\mathfrak{X},\mathbf{R}\nu_*(\mathbf{L}\otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p))$

Theorem 6.3.7 reads that $\mathbf{R}\nu_*(\mathbf{L}\otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p) \in \mathbf{D}_{acoh}^+(X_0)$ as X is smooth. Theorem 5.1.3 implies that

$$\mathbf{R}\Gamma(X, \mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p) \simeq \mathbf{R}\Gamma(\mathfrak{X}, \mathbf{R}\nu_*(\mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p)) \in \mathbf{D}_{acoh}^+(A_0)$$

127

6.4. Proof of Theorem 6.1.1 for $\mathbf{R}\nu_*(\mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$. Theorems 6.2.11 and 6.3.7 almost give a proof of Theorem 6.1.1. We are left to show three things: $\mathbf{R}\nu_*(\mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$ is almost coherent even if the generic fiber X is not smooth, $\mathbf{R}\nu_*(\mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$ is concentrated in degrees $[0, 2 \dim X]$, and it is almost concentrated in degrees $[0, \dim X]$. It turns out that the hardest part is to show that this complex is almost concentrated in degrees $[0, \dim X]$. In order to achieve this result, we will need to use Scholze's theory of diamonds and the recent notion of perfectoidization developed in [BS19]. The proof is strongly motivated by [Guo19, Proposition 7.5.2]. The only new real input is that we use the embedding of the closed unit disc as a rational subset of a torus to get the result for *any* affinoid space.

We recall the definition of the diamond X^{\diamond} associated to an analytic adic space X. This is a functor on the category Perf of perfectoid spaces in characteristic p, whose functor of points is defined as

$$X^{\diamondsuit}(S) = \left\{ \left(\left(S^{\sharp}, \iota \right), f \colon S^{\sharp} \to X \right) \right\} / \text{isom}$$

where S^{\sharp} is a perfectoid space, and $\iota: (S^{\sharp})^{\flat} \to S$ is an identification of a S^{\sharp} as an until of S. It turns out that this defines a (locally spatial) diamond for any analytic adic space X. This roughly means that X^{\diamondsuit} is a sheaf in the big pro-étale topology³⁴, and it is pro-étale locally isomorphic a perfectoid space. However, it will be too big a digression to explain the definition of a diamond, instead we refer to [Sch17], and especially [Sch17, §11, 15], for an extensive discussion of this notion. The idea of the proof is that we can define the étale site of any diamond S in way that induces an equivalence $X_{\text{ét}} \simeq X_{\text{ét}}^{\diamondsuit}$. In particular, the sheaf $\mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p$ has its diamond counterpart $(\mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)^{\diamondsuit}$. It surprisingly turns out that it is easier to (almost) compute the right hand site of the equality

$$\mathrm{H}^{i}_{\mathrm{\acute{e}t}}(X, \mathbf{L} \otimes_{\mathbf{F}_{p}} \mathfrak{O}^{+}_{X}/p) \simeq \mathrm{H}^{i}_{\mathrm{\acute{e}t}}(X^{\diamondsuit}, (\mathbf{L} \otimes_{\mathbf{F}_{p}} \mathfrak{O}^{+}_{X}/p)^{\diamondsuit})$$
(6.1)

Even though this is not exactly what we are going to do, let us explain the main reason why the right hand side might be more trackable in some examples. The main observation is that the functor

 $(-)^{\diamond}$: {(Pre-)Adic Analytic Spaces} \rightarrow {Diamonds}

is not fully faithful. That will be the crux of our argument, we will roughly construct a certain pre-adic space that is not isomorphic to an affinoid perfectoid space, but its diamond is! So equation (6.1) will enable us to show that certain cohomology groups vanish even though the spaces is not necessarily perfectoid on the nose. In order to do make this work, we need the following recent theorem of Bhatt and Scholze:

Theorem 6.4.1. [BS19, Theorem 10.11] Let R be an integral perfectoid ring³⁵. Let $R \to S$ be the p-adic completion of an integral map. Then there exists an integral perfectoid ring S_{perfd} together with a map of R-algebras $S \to S_{\text{perfd}}$, such that it is initial among all of the R-algebra maps $S \to S'$ for S' being integral perfectoid.

Now we explain the main issue with that approach. The main difficulty is that it is rather hard to work with $X_{\text{ét}}^{\diamond}$ as it does not have a basis given by perfectoid spaces. It will be much more convenient to work with the so-called "v-topology" that we define in a moment.

 $^{^{34}}$ The definition of the big pro-étale topology is [Sch17, Definition 8.1]. We emphasize that this definition differs from [Sch13, Definition 3.9]

 $^{^{35}}$ We use [BMS18, Definition 3.5] as the definition for integral perfectoid rings here. This definition coincides with Definition 6.2.1 in the *p*-torsionfree case, but it is less restrictive in general

Definition 6.4.2. The category Perf is the category of κ -small perfectoid spaces of characteristic p.

A small v-sheaf is a v-sheaf Y on Perf such that there is a surjective map of v-sheaves $Y' \to Y$ for some (κ -small) perfectoid space Y'.

The *v*-site Y_v of a diamond Y is the site whose objects are all maps $Y' \to Y$ from small *v*-sheaves Y', with coverings given by families of jointly surjective maps.

We now define a morphism of sites³⁶ $\beta: X_v^{\diamond} \to X_{\text{pro\acute{e}t}}$. Firstly, we note that we have a functor $(-)^{\diamond}: X_{\acute{e}t} \to X_v^{\diamond}$ that is defined by

$$(Y \to X) \mapsto Y^{\diamondsuit} \to X^{\bigstar}$$

This functor is clearly continuous as it sends covers to covers by [Sch17, Lemma 15.6]. We claim that it commutes with finite limits.

Lemma 6.4.3. Let $(-)^{\diamond} \colon X_{\acute{e}t} \to X_v^{\diamond}$ be the functor as above. Then it commutes with finite limits.

Proof. It suffices to show that the functor commutes with fiber products. The essential input here is [Hub94, Proposition 3.7] that guarantees that, for any $U, V, W \in X_{\acute{e}t}$, the fiber product $U \times_V W$ is the fiber product in the category of adic spaces. Then it becomes straightforward to check that $(U \times_V W)^{\diamond} \simeq U^{\diamond} \times_{V^{\diamond}} W^{\diamond}$. Indeed, we have an identification

$$(U \times_V W)^{\diamondsuit}(S) = \left\{ \left(\left(S^{\sharp}, \iota \right), S^{\sharp} \to U \times_V W \right) \right\} / \text{isom} \\ = \left\{ \left(\left(S^{\sharp}, \iota \right), S^{\sharp} \to U \right) \right\} / \text{isom} \times_{\left\{ \left(\left(S^{\sharp}, \iota \right), S^{\sharp} \to W \right) \right\} / \text{isom}} \left\{ \left(\left(S^{\sharp}, \iota \right), S^{\sharp} \to W \right) \right\} / \text{isom} \\ = U^{\diamondsuit}(S) \times_{V^{\diamondsuit}(S)} W^{\diamondsuit}(S)$$

that is functorial in S.

From now on we suppose that the space X is quasi-compact and quasi-separated. Then we can define the site $X_{\text{proétqc}}$ consisting of quasi-compact objects in $X_{\text{proét}}$. This defines a site due to [Sch17, Proposition 3.12]. Moreover, as any étale morphism of strongly noetherian adic spaces is quasi-separated, we note that any object of $X_{\text{proétqc}}$ is quasi-compact and quasi-separated by [Sch17, Proposition 3.12]. Now we note that [Sch17, Lemma 12.17] says that X_v^{\diamond} contains all (κ' -small for any $\kappa' < \kappa$) limits of small v-sheaves with quasi-compact and quasi-separated transition maps³⁷. This allows us to extend the functor $(-)^{\diamond}: X_{\text{étqc}} \to X_v^{\diamond}$ to the functor $(-)^{\diamond}: X_{\text{proétqc}} \to X_v^{\diamond}$ as

$$U = "\lim_i U_i" \mapsto U^{\diamondsuit} \coloneqq \lim_i U_i^{\checkmark}$$

This functor preserves covers by [Sch17, Lemma 12.17]. Moreover, a straightforward argument using Lemma 6.4.3 shows that this functor preserves finite limits. Thus it defines a morphism of sites:

$$\alpha \colon X_v^{\diamondsuit} \to X_{\text{proetqc}}$$

for any quasi-compact and quasi-separated rigid C-space X. As the associated topoi of $X_{\text{pro\acute{e}t}}$ and $X_{\text{pro\acute{e}tgc}}$ are the same, we get a morphism of topoi:

$$(\alpha^{-1}, \alpha_*) \colon \mathbf{Shv}(X_v^{\diamondsuit}) \to \mathbf{Shv}(X_{\mathrm{pro\acute{e}t}})$$

³⁶We follows the convention of [Sta21, Tag 00X1]. So the morphism of cites $\mathcal{C} \to \mathcal{D}$ is a continuous functor $u: \mathcal{D} \to \mathcal{C}$ such that " u_s " is exact

 $^{^{37}}$ With some extra mild set-theoretic assumption that is automatic in our situation for an appropriate choice of κ

The last thing we observe is that this functor is quite explicit on affinoid perfectoid objects $U \in X_{\text{proétqc}}$. Namely, the affinoid perfectoid $U = \text{``lim}_i \operatorname{Spa}(A_i, A_i^+)$ '', with a perfectoid pair (A, A^+) being the *p*-adic completion $\operatorname{colim}_i(A_i, A_i^+)$, is sent to

$$U^{\diamondsuit} = \lim_{i} \operatorname{Spa}(A_i, A_i^+)^{\diamondsuit} = \lim_{i} \operatorname{Spd}(A_i, A_i^+) = \operatorname{Spd}(A, A^+) = \operatorname{Spa}(A^{\flat}, A^{+,\flat}) = \widehat{U}^{\flat}.$$

Now we define the "untilted integral structure sheaf" on X_v^{\diamond} following [Guo19]. We note that perfectoid spaces of characteristic p form a basis of v-topology by the definition of this topology. Thus any sheaf on the basis $\operatorname{Perf}_{X\diamond}$ uniquely extends to a v-sheaf on X^{\diamond} . We define the untilted structure sheaf $\widehat{\mathcal{O}}^+_{X\diamond}$ by

$$(S \to X^{\diamondsuit}) \mapsto \mathcal{O}^+_{S^{\sharp}}(S^{\sharp}) \text{ for } S \in \operatorname{Perf}$$

This functor is a sheaf for the v-topology by [Sch17, Theorem 8.7].

Lemma 6.4.4. Let X be a quasi-compact and quasi-separated rigid space over C, and let X^{\diamond} and α be as above. Then there is a canonical isomorphism $v: \widehat{\mathcal{O}}_X^+ \to \alpha_*(\widehat{\mathcal{O}}_{X\diamond}^+)$.

Proof. We recall that affinoid perfectoids form a basis for the (quasi-compact) pro-étale topology on X by [Sch13, Proposition 4.8]. So, it is sufficient to define r on these spaces in a functorial way. We pick an affinoid perfectoid $U = \lim_{n \to \infty} \operatorname{Spa}(A_i, A_i^+)^n$ with a perfectoid pair (A, A^+) being the p-adic completion $\operatorname{colim}_i(A_i, A_i^+)$. We recall that

$$U^{\diamondsuit} = \lim_{i} \operatorname{Spa}(A_i, A_i^+)^{\diamondsuit} = \lim_{i} \operatorname{Spd}(A_i, A_i^+) = \operatorname{Spd}(A, A^+) = \operatorname{Spa}(A^{\flat}, A^{+,\flat}) = \widehat{U}^{\flat}$$

So we see that

$$\widehat{\mathcal{O}}^+_{X\diamondsuit}(U\diamondsuit) = \widehat{\mathcal{O}}^+_X(\operatorname{Spa}(A^\flat, A^{+,\flat})) \simeq A^-$$

where the last isomorphism is the canonical isomorphism that comes from the Tilting Equivalence [Sch12, Proposition 6.17]. Namely, there is a unique up to a unique isomorphism until of the perfected pair $(A^{\flat}, A^{+,\flat})$ with the structure morphism from (C, \mathcal{O}_C) . Now we note that $A^+ \to \widehat{\mathcal{O}}^+_X(U)$ is also an isomorphism by [Sch13, Lemma 4.10]. So this allows us to define a morphism

$$\widehat{\mathcal{O}}_X^+(U) \to A^+ \to \widehat{\mathcal{O}}_{X^\diamondsuit}^+(U^\diamondsuit) = \alpha_*(\widehat{\mathcal{O}}_{X^\diamondsuit}^+)(U)$$

that is functorial in U and an isomorphism for any such U. This defines the morphism $\widehat{\mathbb{O}}_X^+ \to \alpha_*(\widehat{\mathbb{O}}_{X\diamond}^+)$, and this is an isomorphism as it is an isomorphism on the basis of affinoid perfectoids. \Box

Lemma 6.4.5. Let X be a quasi-compact and quasi-separated rigid space over C, and let X^{\diamond} and α be as above. Then there is an almost isomorphism $r: \mathcal{O}_X^+/p \to \alpha_*(\widehat{\mathcal{O}}_{X^{\diamond}}^+/p)$, i.e. the kernel and cokernel are annihilated by p^{ε} for any $\varepsilon \in \mathbf{Q}_{>0}$. Moreover, $\mathrm{R}^i \alpha_*(\widehat{\mathcal{O}}_{X^{\diamond}}^+/p)$ is almost zero (i.e. is annihilated by p^{ε} for any $\varepsilon \in \mathbf{Q}_{>0}$) for $i \geq 1$.

Proof. We recall that [Sch13, Lemma 4.1] guarantees that \mathcal{O}_X^+/p is canonically identified with $\widehat{\mathcal{O}}_X^+/p$. Thus, the isomorphism of Lemma 6.4.4 defines the morphism

$$\mathcal{O}_X^+/p \to \alpha_*(\mathcal{O}_{X^\diamond}^+/p)$$

We claim that this is an almost isomorphism. It is sufficient to check on the basis of affinoid perfectoids. We pick an affinoid perfectoid $U = \lim_n \operatorname{Spa}(A_i, A_i^+)$ with a perfectoid pair (A, A^+) being the *p*-adic completion $\operatorname{colim}_i(A_i, A_i^+)$. As perfectoid spaces also form a basis for the *v*topology on X^{\diamond} , we conclude that [Sch17, Proposition 8.8] and the Tilting Equivalence [Sch12, Proposition 6.17] imply that $\operatorname{H}^i_v(U^{\diamond}, \widehat{\mathbb{O}}^+_X) \cong^a 0$ for $i \geq 1$. This shows that the natural morphism

 $A^+/p \to \mathrm{H}^0_v(U^\diamondsuit, \widehat{\mathbb{O}}^+_X/p)$ is an almost isomorphism. Now note that $A^+/p \to \mathrm{H}^0(U, \mathbb{O}^+_X/p)$ is also an almost isomorphism by [Sch13, Lemma 4.10]. Thus, we have a commutative diagram



where the vertical maps are almost isomorphisms. Thus the lower horizontal map is an almost isomorphism as well. Since this holds for any element of the basis of affinoid perfectoids, we conclude that the map $\mathcal{O}_X^+/p \to \alpha_*(\widehat{\mathcal{O}}_{X^{\diamondsuit}}^+/p)$ is an almost isomorphism.

To show the second claim, we note that the $\mathbf{R}^i \alpha_* (\widehat{\mathcal{O}}^+_{X\diamond}/p)$ is defined as the sheafification of the presheaf

$$X_{\text{proetqc}} \ni U \to \mathrm{H}^{i}_{v}(U^{\diamondsuit}, \widehat{\mathrm{O}}^{+}_{X^{\diamondsuit}}/p)$$

So it suffices to show that this presheaf has almost zero values on affinoid perfectoids in X_{proetqc} for $i \geq 1$. Since U^{\diamond} is an affinoid perfectoid in this case, we reduce the question to show that $H_v^i(Y, \hat{\mathbb{O}}_{X\diamond}^+/p)$. As above, it follows from the fact that perfectoids form a basis for the v-topology on X^{\diamond} , [Sch17, Proposition 8.8] and the Tilting Equivalence [Sch12, Proposition 6.17].

Now we need a version of this result for any \mathbf{F}_p -local system \mathbf{L} on $X_{\text{\acute{e}t}}$. As before, we slightly abuse notations and denote by the same letter its "extension" $\lambda^{-1}\mathbf{L}^{38}$ on the pro-étale site. We denote $\alpha^{-1}\mathbf{L}$ as \mathbf{L}^{\diamond} .

The morphism $r: \mathcal{O}_X^+/p \to \alpha_*(\widehat{\mathcal{O}}_{X^{\diamondsuit}}^+/p)$ defines the morphism $\alpha^{-1}(r): \alpha^{-1}(\mathcal{O}_X^+/p) \to \widehat{\mathcal{O}}_{X^{\diamondsuit}}^+/p$ by the adjunction (α^{-1}, α_*) . Thus we obtain the morphism

$$\alpha^{-1}(\mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p) \simeq \mathbf{L}^{\diamondsuit} \otimes_{\mathbf{F}_p} \alpha^{-1}(\mathfrak{O}_X^+/p) \xrightarrow{\mathrm{Id} \otimes \alpha^{-1}(r)} \mathbf{L}^{\diamondsuit} \otimes_{\mathbf{F}_p} \widehat{\mathfrak{O}}_{X^{\diamondsuit}}^+/p$$

Using the adjunction once again we get the morphism

$$r_{\mathbf{L}} \colon \mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+ / p \to \alpha_* (\mathbf{L}^{\diamondsuit} \otimes_{\mathbf{F}_p} \widetilde{\mathfrak{O}}_{X^{\diamondsuit}}^+ / p).$$

Lemma 6.4.6. Let X be a quasi-compact and quasi-separated rigid C-space, and let X^{\diamondsuit} , **L**, and α be as above. Then the kernel and cokernel of $r_{\mathbf{L}}$ are annihilated by p^{ε} for any $\varepsilon \in \mathbf{Q}_{>0}$. Moreover, $\mathrm{R}^{i}\alpha_{*}(\mathbf{L}^{\diamondsuit} \otimes_{\mathbf{F}_{p}} \widehat{\mathbb{O}}^{+}_{X^{\diamondsuit}}/p)$ is almost zero (i.e. annihilated by p^{ε} for any $\varepsilon \in \mathbf{Q}_{>0}$) for $i \geq 1$.

Proof. Both claims are local on X, so we can check them étale locally. Then we may assume that X is connected and **L** is just the trivial \mathbf{F}_p -local system \mathbf{F}_p^n . In this case, both claims are just consequences of Lemma 6.4.5.

Now we are ready to give the first application of this formalism. Namely, we show that

$$\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\mathbf{F}_{P}} \mathfrak{O}_{X}^{+}/p)$$

is almost coherent over A_0/p for any rigid-analytic affinoid $X = \text{Spa}(A, A^+)$ with a topologically finite type ring of definition A_0 . The last ingredient we need is the notion of a hypercovering and cohomological descent. We briefly remind the reader the definition of this notion and refer to [Con] for a detailed discussion of this notion.

³⁸Recall that λ is the natural morphism $X_{\text{pro\acute{e}t}} \to X_{\acute{e}t}$.

Definition 6.4.7. Let \mathcal{C} be a category admitting finite limits. Let \mathbf{P} be a class of morphisms in \mathcal{C} which is stable under base change, preserved under composition (hence under products), and contains all isomorphisms. A simplicial object X_{\bullet} in \mathcal{C} is said to be a \mathbf{P} -hypercovering if, for all $n \geq 0$, the natural adjunction map

$$X_{\bullet} \to \operatorname{cosk}_n(\operatorname{sk}_n(X_{\bullet}))$$

induces a map $X_{n+1} \to (\cos k_n(sk_n(X_{\bullet})))_{n+1}$ in degree n+1 which is in **P**. If X_{\bullet} is an augmented simplicial complex, we make a similar definition but also require the case n = -1 (and we then say X_{\bullet} is a **P**-hypercovering of X_{-1}).

We refer to [Con, Definition 6.5] for a precise definition of a morphism of cohomological descent. What matters for us is that if $a: X_{\bullet} \to X$ is a morphism of cohomological descent, then we have a spectral sequence

$$\mathrm{E}_{1}^{i,j} = \mathrm{H}^{j}(X_{i}, a_{i}^{*}\mathcal{F}) \Rightarrow \mathrm{H}^{p+q}(X, \mathcal{F})$$

for \mathcal{F} a sheaf on X (in some topology on X) and $a_i: X_i \to X$ being the canonical map that comes from the augmentation $X_{\bullet} \to X$.

Theorem 6.4.8. Let $X = \text{Spa}(A, A^+)$ be an affinoid rigid-analytic space over $\text{Spa}(C, \mathcal{O}_C)$, and let **L** be an \mathbf{F}_p -local system on X. Suppose that $A_0 \subset A^+$ is a topologically finite type ring of definition. Then $\text{H}^i(X, \mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$ is almost coherent over A_0/p .

Proof. First of all, we note that quasi-compact rigid-analytic varieties over $\operatorname{Spa}(C, \mathcal{O}_C)$ admit resolution of singularities by [Tem12, Theorem 5.2.2]. Thus, the proof of [Con] carries over to show that there is a proper hypercovering³⁹ $a: X_{\bullet} \to X$ such that all X_i are smooth over $\operatorname{Spa}(C, \mathcal{O}_C)$. We now consider the simplicial object $a^{\diamond}: X_{\bullet}^{\diamond} \to X^{\diamond}$. We claim that this is a *v*-hypercovering⁴⁰. Indeed, the functor $(-)^{\diamond}$ commutes with finite limits on the category of rigid-analytic spaces over C by the same proof as in Lemma 6.4.3. So $((\operatorname{cosk}_n(\operatorname{sk}_n X_{\bullet}))_{n+1})^{\diamond} \simeq (\operatorname{cosk}_n(\operatorname{sk}_n X_{\bullet}^{\diamond}))_{n+1}$. Therefore, the only thing we need to check is that the morphism

$$X_{n+1}^{\diamondsuit} \to \left(\left(\operatorname{cosk}_n \left(\operatorname{sk}_n X_{\bullet} \right) \right)_{n+1} \right)^{\checkmark}$$

is a v-covering for any $n \ge -1$. Now we use [Sch17, Lemma 15.6] to see that topologically it is identified with

$$|X_{n+1}| \to |\left(\operatorname{cosk}_n\left(\operatorname{sk}_n X_{\bullet}\right)\right)_{n+1}|$$

that is certainly a v-cover as $X_{n+1} \to (\cosh_n(\operatorname{sk}_n X_{\bullet}))_{n+1}$ is proper. As in the discussion before the proof of Lemma 6.4.4 we have a commutative diagram of morphisms of topoi:

$$\begin{aligned} \mathbf{Shv}(X_{i,\mathrm{pro\acute{e}t}}) &\longleftrightarrow_{(\alpha_{i}^{-1},\alpha_{i,*})} \quad \mathbf{Shv}(X_{i,v}^{\diamondsuit}) \\ & \downarrow^{(a_{i}^{-1},a_{i,*})} \quad \qquad \downarrow^{((a_{i}^{\diamondsuit})^{-1},a_{i,*}^{\diamondsuit})} \\ \mathbf{Shv}(X_{\mathrm{pro\acute{e}t}}) &\longleftarrow_{(\alpha^{-1},\alpha_{*})} \quad \mathbf{Shv}(X_{v}^{\diamondsuit}) \end{aligned}$$

Then we note that

$$(a_i^{\diamondsuit})^{-1}(\mathbf{L}^{\diamondsuit} \otimes_{\mathbf{F}_p} \widehat{\mathcal{O}}_{X^{\diamondsuit}}^+/p) \simeq (a_i^{\diamondsuit})^{-1}(\mathbf{L}^{\diamondsuit}) \otimes_{\mathbf{F}_p} (\widehat{\mathcal{O}}_{X^{\diamondsuit}}^+/p)|_{X_{i,v}^{\diamondsuit}} \simeq (a_i^{-1}\mathbf{L})^{\diamondsuit} \otimes_{\mathbf{F}_p} \widehat{\mathcal{O}}_{X_i^{\diamondsuit}}^+/p$$

³⁹We consider this as a proper hypercover in the category of rigid-analytic spaces over C. This category has all finite limits and proper morphisms are preserved by base change and composition by [Tem00, Theorem 4.1], so it makes sense to speak about proper hypercoverings

 $^{^{40}}$ Similarly, the category of small *v*-sheaves has all finite limits. And *v*-covers are preserved by base change and compositions

Finally, we use [Con, Theorem 7.2 and Theorem 7.10] to conclude that $a^{\diamond} \colon X_{\bullet}^{\diamond} \to X^{\diamond}$ is a morphism of cohomological descent for the *v*-topology. So we can apply [Con, Theorem 6.11] to get the spectral sequence

$$\mathbf{E}_{1}^{i,j} = \mathbf{H}_{v}^{j}(X_{i}^{\Diamond}, (a_{i}^{\Diamond})^{-1}(\mathbf{L}^{\Diamond} \otimes_{\mathbf{F}_{p}} \widehat{\mathbf{O}}_{X^{\Diamond}}^{+}/p)) \Rightarrow \mathbf{H}_{v}^{i+j}(X^{\Diamond}, \mathbf{L}^{\Diamond} \otimes_{\mathbf{F}_{p}} \widehat{\mathbf{O}}_{X^{\Diamond}}^{+}/p)$$

that boils down to the spectral sequence:

$$\mathbf{E}_{1}^{i,j} = \mathbf{H}_{v}^{j}(X_{i}^{\diamondsuit}, (a_{i}^{-1}\mathbf{L})^{\diamondsuit} \otimes_{\mathbf{F}_{p}} \widehat{\mathcal{O}}_{X_{i}^{\diamondsuit}}^{+}/p) \Rightarrow \mathbf{H}_{v}^{i+j}(X^{\diamondsuit}, \mathbf{L}^{\diamondsuit} \otimes_{\mathbf{F}_{p}} \widehat{\mathcal{O}}_{X^{\diamondsuit}}^{+}/p)$$
(6.2)

We note that Lemma 6.4.6 implies that the natural map

$$r_i \colon a_i^{-1} \mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_{X_i}^+ / p \to \mathbf{R} \alpha_{i,*} \left(\left(a_i^{-1} \mathbf{L} \right)^{\diamondsuit} \otimes_{\mathbf{F}_p} \widehat{\mathfrak{O}}_{X_i^{\diamondsuit}}^+ / p \right)$$

is an almost isomorphism (i.e. cone is annihilated by p^{ε} for any $\varepsilon \in \mathbf{Q}_{>0}$). Thus, we have the following almost isomorphisms:

$$\mathbf{R}\Gamma_{v}\left(X_{i}^{\diamondsuit},\left(a_{i}^{-1}\mathbf{L}\right)^{\diamondsuit}\otimes_{\mathbf{F}_{p}}\widehat{\mathcal{O}}_{X_{i}^{\diamondsuit}}^{+}/p\right)\simeq\mathbf{R}\Gamma_{\mathrm{pro\acute{e}t}}\left(X_{i},\mathbf{R}\alpha_{i,*}\left(\left(a_{i}^{-1}\mathbf{L}\right)^{\diamondsuit}\otimes_{\mathbf{F}_{p}}\widehat{\mathcal{O}}_{X_{i}^{\diamondsuit}}^{+}/p\right)\right)$$
$$\cong^{a}\mathbf{R}\Gamma_{\mathrm{pro\acute{e}t}}\left(X_{i},a_{i}^{-1}\mathbf{L}\otimes_{\mathbf{F}_{p}}\mathcal{O}_{X_{i}}^{+}/p\right)$$

As each of X_i is smooth over $\text{Spa}(C, \mathcal{O}_C)$, we conclude that

$$\mathbf{R}\Gamma_{\text{pro\acute{e}t}}\left(X_p, a_q^{-1}\mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_{X_p}^+/p\right) \in \mathbf{D}_{acoh}(A_0/p)$$

by Corollary 6.3.8. In particular, each term of the spectral sequence 6.2 is almost coherent over A_0/p . The standard argument with (almost) coherence of A_0/p implies that each of $\mathbf{H}_v^{i+j}(X^{\diamondsuit}, \alpha^{-1}\mathbf{L}\otimes_{\mathbf{F}_p} \widehat{\mathcal{O}}^+_{X^{\diamondsuit}}/p)$ is also almost coherent over A_0/p . Then we use Lemma 6.4.6 once again to conclude that

$$\mathrm{H}^{i+j}_{\mathrm{pro\acute{e}t}}(X, \mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}^+_X/p) \cong^{a} \mathrm{H}^{i+j}_{v}(X^{\diamondsuit}, \alpha^{-1}\mathbf{L} \otimes_{\mathbf{F}_p} \widehat{\mathbb{O}}^+_{X^{\diamondsuit}}/p)$$

As a result, $\mathrm{H}^{n}_{\mathrm{pro\acute{e}t}}(X, \mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}^+_X/p)$ are indeed almost coherent over A_0/p for any $n \ge 0$.

Now we want to use similar ideas to show that cohomology groups $\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\mathbf{F}_{p}} \mathcal{O}_{X}^{+}/p)$ are almost concentrated in degrees $[0, \dim X]$ for any affinoid X. We need to recall the construction of the "standard" covering $\widehat{\mathbf{D}}_{\infty}^{d} \to \mathbf{D}^{d}$. We start with the "standard" $\Delta_{\infty} \simeq \mathbf{Z}_{p}(1)^{d}$ -torsor

$$\mathbf{T}_{\infty}^{d} \coloneqq \lim_{n} \operatorname{Spa}(C\langle T_{1}^{\pm 1/p^{n}}, \dots, T_{d}^{\pm 1/p^{n}}\rangle, \mathbb{O}_{C}\langle T_{1}^{\pm 1/p^{n}}, \dots, T_{d}^{\pm 1/p^{n}}\rangle)^{*} \to \mathbf{T}^{d}$$

with $\widehat{\mathbf{T}}_{\infty}^{d} \simeq \operatorname{Spa}(C\langle T_{1}^{\pm 1/p^{\infty}}, \ldots, T_{d}^{\pm 1/p^{\infty}} \rangle, \mathfrak{O}_{C}\langle T_{1}^{\pm 1/p^{\infty}}, \ldots, T_{d}^{\pm 1/p^{\infty}} \rangle)$ an affinoid perfectoid space. Then Lemma 6.3.4 implies that the restriction of this pro-étale cover on $\mathbf{D}^{d} = \mathbf{T}^{d}(\frac{T_{1}-1}{p}, \ldots, \frac{T_{n}-1}{p}) \subset \mathbf{T}^{d}$ defines a pro-étale covering

$$\mathbf{D}^d_\infty \coloneqq \mathbf{T}^d_\infty imes_{\mathbf{T}^d} \mathbf{D}^d o \mathbf{D}^d$$

with an affinoid perfectoid space $\widehat{\mathbf{D}}_{\infty}^{d}$. Now [Sch17, Proposition 15.4] implies that the map $\widehat{\mathbf{D}}_{\infty}^{d,\diamond} \rightarrow \mathbf{D}^{d,\diamond}$ is a *v*-cover that is $\underline{\Delta}_{\infty}$ -torsor, and $\widehat{\mathbf{D}}_{\infty}^{d,\diamond} \simeq \widehat{\mathbf{D}}_{\infty}^{d,\flat}$ is an affinoid perfectoid space of characteristic *p*. We show that this property is preserved under base change along finite morphisms.

Lemma 6.4.9. Let $g: X = \operatorname{Spa}(S, S^+) \to \mathbf{D}^d$ be a finite morphism of rigid *C*-spaces. Then the fiber product $X_{\infty} \coloneqq X^{\diamondsuit} \times_{\mathbf{D}^d} \widehat{\mathbf{D}}_{\infty}^{d,\diamondsuit}$ is isomorphic to an affinoid perfectoid space of characteristic p.

Proof. Let us say that $\mathbf{D}^d = \operatorname{Spa}(R, R^+)$ and $\widehat{\mathbf{D}}^d_{\infty} = \operatorname{Spa}(R_{\infty}, R^+_{\infty})$. We consider the map $g: X \to \mathbf{D}^d$ that defines a finite map $(R, R^+) \to (S, S^+)$ and we define $S^+_{\infty} \coloneqq R^+_{\infty} \widehat{\otimes}_{R^+} S^+$. Then we note that the map $R^+_{\infty} \to S^+_{\infty}$ is the *p*-adic completion of the integral map with R^+_{∞} an integral perfectoid ([BMS18, Lemma 3.20]). Thus Theorem 6.4.1 implies that there is an integral perfectoid *R*-algebra $(S^+_{\infty})_{\text{perfd}}$ that is initial among integral perfectoid⁴¹ *R*-algebras *S'* with a morphism $S^+_{\infty} \to S'$.

We define $S_{\text{perfd}}^{\dagger}$ as the *p*-torsionfree quotient of $(S_{\infty}^{+})_{\text{perfd}}$. We claim that $S_{\text{perfd}}^{\dagger}$ is a *p*-torsionfree integral perfectoid \mathcal{O}_{C} -algebra. The only non-trivial fact that we need to check is that it is *p*-adically complete. We use [BMS19, Proposition 4.19] to see that $(S_{\infty}^{+})_{\text{perfd}}$ has bounded p^{∞} -torsion. More precisely, we get that $(S_{\infty}^{+})_{\text{perfd}}[p^{\infty}] = (S_{\infty}^{+})_{\text{perfd}}[p]$. We note that any *p*-adically complete \mathbf{Z}_{p} -module is *p*-adically derived complete by [Sta21, Tag 091T]. In particular, both $(S_{\infty}^{+})_{\text{perfd}}[p]$ and $(S_{\infty}^{+})_{\text{perfd}}$ are *p*-adically derived complete. Thus the quotient

$$S_{\text{perfd}}^{\dagger} = \left(S_{\infty}^{+}\right)_{\text{perfd}} / \left(\left(S_{\infty}^{+}\right)_{\text{perfd}}[p]\right)$$

is *p*-adically derived complete by [Sta21, Tag 091U]. But now $S_{\text{perfd}}^{\dagger}$ is *p*-torsionfree, thus [Sta21, Tag 0BKG] implies that its derived *p*-adic completion coincides with the usual *p*-adic completion. So $S_{\text{perfd}}^{\dagger}$ is classically *p*-adically complete.

Now we define S_{perfd}^+ as the integral closure of S_{perfd}^\dagger in $S_{\text{perfd}}^\dagger [1/p]$. Then [BMS18, Lemma 3.21] implies that $(S_{\text{perfd}}^+[1/p], S_{\text{perfd}}^+)$ is a perfectoid pair, and it admits a continuous map $(S, S^+) \rightarrow (S_{\text{perfd}}^+[1/p], S_{\text{perfd}}^+)$, in particular, it is sheafy. Moreover, it is initial among maps to perfectoid pairs by its construction. [Hub94, Proposition 2.1] reads that we have a commutative diagram

that induces a commutative diagram

$$\begin{array}{ccc} Y^{\diamondsuit} & \longrightarrow & \widehat{\mathbf{D}}_{\infty}^{d,\diamondsuit} \\ \downarrow & & \downarrow \\ X^{\diamondsuit} & \longrightarrow & \mathbf{D}^{d,\diamondsuit} \end{array}$$

The universal property of fiber products induces a map $Y^{\diamond} \to X_{\infty}$. We claim that this map is an isomorphism. It suffices to check the equality on *T*-valued points for an affinoid Tate-perfectoid

 $^{^{41}}$ Similarly to Theorem 6.4.1 we need to use [BMS18, Definition 3.5] as the definition for integral perfectoid rings.

spaces T of characteristic p. So we have

$$\begin{split} Y^{\Diamond}(T) &= \left\{ \left(\left(T^{\sharp}, \iota \right), f \colon T^{\sharp} \to Y \right) \right\} / \text{isom} \\ &= \left\{ \left(\left(T^{\sharp}, \iota \right), f^{\#} \colon S^{+}_{\text{perfd}} \to \mathbb{O}^{+}(T^{\sharp}) \right) \right\} / \text{isom} \\ &= \left\{ \left(\left(T^{\sharp}, \iota \right), \colon S^{+}_{\infty} \widehat{\otimes}_{R^{+}} R^{+}_{\infty} \to \mathbb{O}^{+}(T^{\sharp}) \right) \right\} / \text{isom} \\ &= \left\{ \left(\left(T^{\sharp}, \iota \right), S^{+} \to \mathbb{O}^{+}(T^{\sharp}) \right) \right\} / \text{isom} \times_{\left\{ \left(\left(T^{\sharp}, \iota \right), R^{+} \to \mathbb{O}^{+}(T^{\sharp}) \right) \right\} / \text{isom}} \left\{ \left(\left(T^{\sharp}, \iota \right), R^{+}_{\infty} \to \mathbb{O}^{+}(T^{\sharp}) \right) \right\} / \text{isom} \\ &= X^{\Diamond}(T) \times_{\mathbf{D}^{d}, \Diamond(T)} \widehat{\mathbf{D}}^{d, \diamondsuit}_{\infty}(T) \\ &= X_{\infty}(T) \end{split}$$

Therefore, $X_{\infty} \simeq Y^{\diamondsuit} \simeq \text{Spa}(S_{\text{perfd}}^{+,\flat}[1/p], S_{\text{perfd}}^{+,\flat})$ is an affinoid perfectoid space of characteristic p.

Theorem 6.4.10. Let $X = \text{Spa}(S, S^+)$ be a rigid *C*-space of dimension *d*, and let **L** be an \mathbf{F}_p -local system on *X*. Then the cohomology groups $\mathrm{H}^i_{\text{ét}}(X, \mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}^+_X/p)$ are almost zero for any i > d.

Proof. First of all, we may and do assume that X is connected. Thus we can assume that **L** is an \mathbf{F}_p -local system of constant rank $\mathrm{rk}_{\mathbf{L}}$. Now we use the Noether Normalization [Bos14, Proposition 3.1.3] to find a finite morphism $f: X \to \mathbf{D}^d$ with $d = \dim X$. Now we consider the "standard" pro-étale cover $\mathbf{D}_{\infty}^d \to \mathbf{D}^d$. And consider the fiber product

$$X_{\infty} \coloneqq X^{\diamondsuit} \times_{\mathbf{D}^d} \widehat{\mathbf{D}}_{\infty}^{d,\diamondsuit}$$

Lemma 6.4.9 implies that $h: X_{\infty} \to X^{\diamond}$ is the *v*-cover of X^{\diamond} by an affinoid perfectoid space. Moreover, this a $\underline{\Delta}_{\infty}$ -torsor as the base change of $\widehat{\mathbf{D}}^{d,\diamond} \to \mathbf{D}^{d,\diamond}$.

As X is quasi-compact and quasi-separated, there is a morphism of topoi

$$(\alpha^{-1}, \alpha_*) \colon \mathbf{Shv}(X_v^{\diamondsuit}) \to X_{\mathrm{pro\acute{e}t}}$$

and Lemma 6.4.6 implies that

$$\mathbf{R}\Gamma_{\text{pro\acute{e}t}}(X, \mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p) \simeq^a \mathbf{R}\Gamma_v(X^\diamondsuit, \mathbf{L}^\diamondsuit \otimes_{\mathbf{F}_p} \widehat{\mathcal{O}}_{X^\diamondsuit}^+/p)$$

We now understand the right side of the equation.

Our first claim is that

$$\mathrm{H}^{i}_{v}(X_{\infty}, \mathbf{L}^{\diamondsuit} \otimes_{\mathbf{F}_{p}} \widehat{\mathrm{O}}^{+}_{X^{\diamondsuit}}/p) \cong^{a} 0 \text{ for } i > 0.$$

We choose some finite étale morphism $X' \to X$ that splits **L**. This corresponds to a finite étale morphism $X'^{\diamond} \to X^{\diamond}$ by [Sch17, Lemma 15.6]. Then X'^{\diamond} splits \mathbf{L}^{\diamond} . Therefore, $\mathbf{L}^{\diamond}|_{X_{\infty}} \in \mathbf{Shv}(X_{\infty})$ splits by a finite étale morphism

$$X'_{\infty} \coloneqq X'^{\diamondsuit} \times_{X^{\diamondsuit}} X_{\infty} \to X_{\infty}.$$

We note that X'_{∞} is an affinoid perfectoid space as it is finite étale over an affinoid perfectoid X_{∞}^{42} . The same holds for all selfproducts $X'_{\infty}^{j/X} := X'_{\infty} \times_{X_{\infty}} X'_{\infty} \times_{X_{\infty}} \cdots \times_{X_{\infty}} X'_{\infty}$. Therefore, [Sch17, Proposition 8.8] implies that

⁴²This is the definition of finite étale morphisms of diamonds [Sch17, Definition 10.1]

$$\mathrm{H}^{i}_{v}(X_{\infty}^{\prime j/X}, \mathbf{L}^{\diamondsuit} \otimes_{\mathbf{F}_{p}} \widehat{\mathbb{O}}^{+}_{X^{\diamondsuit}}/p) \simeq \mathrm{H}^{i}_{v}(X_{\infty}^{\prime j/X}, (\widehat{\mathbb{O}}^{+}_{X^{\diamondsuit}}/p)^{\mathrm{rk}_{\mathbf{L}}}) \cong^{a} 0 \text{ for } i > 0, j > 0$$

$$\mathrm{H}^{0}_{v}(X_{\infty}^{\prime j/X}, \mathbf{L}^{\diamondsuit} \otimes_{\mathbf{F}_{p}} \widehat{\mathrm{O}}_{X^{\diamondsuit}}^{+}/p) = \mathrm{H}^{0}_{v}(X_{\infty}^{\prime j/X}, (\widehat{\mathrm{O}}_{X^{\diamondsuit}}^{+}/p)^{\mathrm{rk}_{\mathbf{L}}}) \cong^{a} ((S^{\sharp, +}/p)^{\otimes_{R^{\sharp, +}/p^{j}}})^{\mathrm{rk}_{\mathbf{L}}} \text{ for } j > 0$$

where $X'_{\infty} = \operatorname{Spa}(S, S^+)$, $X_{\infty} = \operatorname{Spa}(R, R^+)$ and $(S^{\sharp}, S^{+,\sharp})$, $(R^{\sharp}, R^{+,\sharp})$ are their unique untilts over (C, \mathcal{O}_C) . This vanishing result implies that $\mathbf{R}\Gamma_v(X_{\infty}, \mathbf{L}^{\diamondsuit} \otimes_{\mathbf{F}_p} \widehat{\mathcal{O}}^+_{X^{\diamondsuit}}/p)$ is almost isomorphic to the Čech complex $\check{\mathbb{C}}(X'_{\infty}/X_{\infty}, \mathbf{L}^{\diamondsuit} \otimes_{\mathbf{F}_p} \widehat{\mathcal{O}}^+_{X^{\diamondsuit}}/p^{\flat})$.

The Almost Purity Theorem [Sch12, Theorem 7.9] implies that $S^{+,\sharp}$ is almost étale over $R^{+,\sharp}$, so it is almost flat over $R^{+,\sharp}$. We claim that $S^{+,\sharp}$ is almost faithfully flat over $R^{+,\sharp}$, i.e. the map Spec $S^{+,\sharp} \to \text{Spec } R^{+,\sharp}$ is surjective. We know that this map is integral by the definition of a finite étale morphism of perfectoid spaces. Thus, the Going-Up Theorem [Mat80, Theorem 9.4] implies that it is sufficient to show that the map $\text{Spec } S^{+,\sharp}[1/p] \to \text{Spec } R^{+,\sharp}[1/p]$ is surjective. But this follows from [Hub94, Lemma 1.4] as the map $X_{\infty}^{\prime\sharp} \to X_{\infty}^{\sharp}$ is a covering. Now the almost faithfully flat descent implies that $\check{\mathbb{C}}(X'_{\infty}/X_{\infty}, \mathbf{L}^{\diamondsuit} \otimes_{\mathbf{F}_{p}} \widehat{\mathbb{O}}^{+}_{X^{\diamondsuit}}/p)$ is almost isomorphic to the Amitsur complex for $\mathrm{H}_{v}^{0}(X^{\infty}, \mathbf{L}^{\diamondsuit} \otimes_{\mathbf{F}_{p}} \widehat{\mathbb{O}}^{+}_{X_{\infty}}/p)$ that is almost acyclic in positive degrees since $R^{+,\sharp}/p \to S^{+,\sharp}/p$ is almost faithfully flat because $R^{+,\sharp} \to S^{+,\sharp}$ is already almost faithfully flat.

Finally, we see that $X_{\infty} \to X^{\diamondsuit}$ is a $\underline{\Delta}_{\infty}$ -torsor such that $\mathrm{H}^{i}_{v}(X_{\infty}, \mathbf{L}^{\diamondsuit} \otimes_{\mathbf{F}_{p}} \widehat{\mathrm{O}}^{+}_{X^{\diamondsuit}}/p) \cong^{a} 0$ for $i \geq 1$. Moreover, this is a covering in the *v*-topology, so

$$\mathbf{R}\Gamma_{v}\left(X^{\diamondsuit},\mathbf{L}^{\diamondsuit}\otimes_{\mathbf{F}_{p}}\widehat{\mathbb{O}}_{X^{\diamondsuit}}^{+}/p\right)\cong^{a}\mathbf{R}\Gamma_{cont}\left(\Delta_{\infty},\mathrm{H}_{v}^{0}\left(X_{\infty},\mathbf{L}^{\diamondsuit}\otimes_{\mathbf{F}_{p}}\widehat{\mathbb{O}}_{X^{\diamondsuit}}^{+}/p\right)\right)\\\simeq\mathbf{R}\Gamma_{cont}\left(\Delta_{\infty},\mathrm{H}_{v}^{0}\left(X_{\infty},\mathbf{L}^{\diamondsuit}\otimes_{\mathbf{F}_{p}}\widehat{\mathbb{O}}_{X^{\diamondsuit}}^{+}/p\right)\right)\\\simeq\mathbf{R}\Gamma_{cont}\left(\mathbf{Z}_{p}^{d},\mathrm{H}_{v}^{0}\left(X_{\infty},\mathbf{L}^{\diamondsuit}\otimes_{\mathbf{F}_{p}}\widehat{\mathbb{O}}_{X^{\diamondsuit}}^{+}/p\right)\right)$$

Therefore, [BMS18, Lemma 7.3] implies that $\mathbf{R}\Gamma_{cont}(\mathbf{Z}_p^d, \mathrm{H}_v^0(X_\infty, \mathbf{L}^{\diamondsuit} \otimes_{\mathbf{F}_p} \widehat{\mathcal{O}}_{X^{\diamondsuit}}^+/p))$ is concentrated in degrees $[0, \dim X]$. The last step is to recall the almost isomorphism $\mathrm{H}_{\mathrm{\acute{e}t}}^i(X, \mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p) \cong^a \mathrm{H}_v^i(X^{\diamondsuit}, \mathbf{L}^{\diamondsuit} \otimes_{\mathbf{F}_p} \widehat{\mathcal{O}}_{X^{\diamondsuit}}^+/p)$ to finish the proof. \Box

Now we are ready to give a proof of Theorem 6.1.1 in the case of *p*-torsion local systems.

Theorem 6.4.11. Let \mathfrak{X} an admissible formal \mathcal{O}_C -scheme with the adic generic fiber $X := \mathfrak{X}_C$ of dimension d. Then

$$\mathbf{R}\nu_*(\mathbf{L}\otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p) \in \mathbf{D}_{acoh}^{[0,2d]}(X_0) \text{ and } \mathbf{R}\nu_*(\mathbf{L}\otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p) \in \mathbf{D}_{acoh}^{[0,d]}(X_0)^a$$

for an \mathbf{F}_p -local system **L**. More precisely, for an affine admissible $\mathfrak{X} = \operatorname{Spf} A$ with the adic generic fiber X, we the natural morphism

$$\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\mathbf{F}_{p}} \mathfrak{O}_{X}^{+}/p) \to \mathrm{R}^{i}\nu_{*}(\mathbf{L} \otimes_{\mathbf{F}_{p}} \mathfrak{O}_{X}^{+}/p)$$

is an almost isomorphism. Moreover, the formation of $\mathbb{R}^i \nu_*(\mathbf{L} \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p)$ almost commutes with étale base change, i.e., for any étale morphism $\mathfrak{f} \colon \mathfrak{X} \to \mathfrak{Y}$, the natural morphism

$$\mathfrak{f}_0^*\left(\mathrm{R}^i\nu_{\mathfrak{X}_C,*}(\mathbf{L}\otimes_{\mathbf{F}_p}\mathfrak{O}_{\mathfrak{X}_C}^+/p)\right)\to\mathrm{R}^i\nu_{\mathfrak{Y}_C,*}(\mathfrak{f}_C^*\mathbf{L}\otimes_{\mathbf{F}_p}\mathfrak{O}_{\mathfrak{Y}_C}^+/p)$$

is an almost isomorphism.

Proof. Theorem 6.2.11 claim that $\mathbb{R}^i \nu_*(\mathbf{L} \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p)$ is almost quasi-coherent for any $i \geq 0$. Moreover, the natural morphism

$$\mathrm{H}^{i}(X, \widetilde{\mathbf{L}} \otimes_{\mathbf{F}_{p}} \mathcal{O}_{X}^{+}/p) \to \mathrm{R}^{i}\nu_{*}(\mathbf{L} \otimes_{\mathbf{F}_{p}} \mathcal{O}_{X}^{+}/p)$$

is an almost isomorphism for $i \ge 0$. Moreover, Theorem 6.4.8 guarantees that

$$\mathrm{R}^{i}\nu_{*}(\mathbf{L}\otimes_{\mathbf{F}_{p}}\mathbb{O}_{X}^{+}/p)\in\mathbf{D}_{acoh}(X_{0})^{a}$$

for $i \geq 0$. Theorem 6.4.10 implies that $\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\mathbf{F}_{p}} \mathfrak{O}_{X}^{+}/p) \cong^{a}$ for i > d. Thus, we get that $\mathbf{R}\nu_{*}(\mathbf{L} \otimes_{\mathbf{F}_{p}} \mathfrak{O}_{X}^{+}/p) \in \mathbf{D}_{acoh}^{[0,d]}(X_{0})^{a}$. We note that [Hub96, Proposition 2.8.1] states that $\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\mathbf{F}_{p}} \mathfrak{O}_{X}^{+}/p) \cong 0$ for i > 2d. Therefore, $\mathbf{R}\nu_{*}(\mathbf{L} \otimes_{\mathbf{F}_{p}} \mathfrak{O}_{X}^{+}/p) \in \mathbf{D}_{acoh}^{[0,2d]}(X_{0})$. Finally, Corollary 6.2.12 guarantees that the formation of $\mathrm{R}^{i}\nu_{*}(\mathbf{L} \otimes_{\mathbf{F}_{p}} \mathfrak{O}_{X}^{+}/p)$ commutes with étale base change. \Box

6.5. **Proof of Theorem 6.1.1 for** $\mathbb{R}\nu_*(\mathbb{L}\otimes_{\widehat{\mathbb{Z}}_p}\widehat{\mathbb{O}}_X^+)$ **.** We prove the integral version of Theorem 6.1.1 in the section. The main idea is to get this result from "the finite level". The reduction to the finite level is very similar to that used in the proof of Theorem 5.1.6. We want to emphasize that it is possible to prove Theorem 6.1.1 for the trivial lisse $\widehat{\mathbb{Z}}_p$ -sheaf (see Definition C.7) in the way that is similar to our proof in the case of \mathcal{O}_X^+/p -coefficients. However, the case of a general lisse $\widehat{\mathbb{Z}}_p$ -sheaf \mathbb{L} seems harder because our proof of Theorem 6.4.11 in the case of \mathbb{F}_p local systems used existence of a finite étale cover that splits \mathbb{L} . We do not know at the moment how to adapt this argument to work for lisse $\widehat{\mathbb{Z}}_p$ -sheaves, so we use a limit argument. In order to make this work we firstly need to show the result for any locally constant $\mathbb{Z}/p^n\mathbb{Z}$ -sheaf. We refer to the Appendix C for the definition and main properties of these sheaves. In what follows we will freely use Theorem C.4.

We start the proof by generalizing Theorem 6.4.11 to the case of locally constant sheaf of $\mathbb{Z}/p^n\mathbb{Z}$ -modules.

Theorem 6.5.1. Let \mathfrak{X} an admissible formal \mathcal{O}_C -scheme with the adic generic fiber $X := \mathfrak{X}_C$ of dimension d. Then

$$\mathbf{R}\nu_*(\mathbf{L}\otimes_{\mathbf{Z}/p^n\mathbf{Z}}\mathfrak{O}_X^+/p^n) \in \mathbf{D}_{acoh}^{[0,2d]}(X_{n-1}) \text{ and } \mathbf{R}\nu_*(\mathbf{L}\otimes_{\mathbf{Z}/p^n\mathbf{Z}}\mathfrak{O}_X^+/p^n) \in \mathbf{D}_{acoh}^{[0,d]}(X_{n-1})^a$$

for $\mathbf{L} \in \operatorname{LocSys}_{\mathbf{Z}/p^n \mathbf{Z}, X_{\operatorname{pro\acute{e}t}}}$ and $X_{n-1} \coloneqq \mathfrak{X} \times_{\operatorname{Spf} \mathcal{O}_C} \operatorname{Spec} \mathcal{O}_C/p^n$. More precisely, for an affine admissible $\mathfrak{X} = \operatorname{Spf} A$ with the adic generic fiber X, the natural morphism

$$\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\mathbf{Z}/p^{n}\mathbf{Z}} \mathcal{O}_{X}^{+}/p^{n}) \to \mathrm{R}^{i}\nu_{*}(\mathbf{L} \otimes_{\mathbf{Z}/p^{n}\mathbf{Z}} \mathcal{O}_{X}^{+}/p^{n})$$

is an almost isomorphism. And, moreover, the formation of $\mathrm{R}^i \nu_*(\mathbf{L} \otimes_{\mathbf{Z}/p^n \mathbf{Z}} \mathcal{O}_X^+/p^n)$ almost commutes with étale base change, i.e., for any étale morphism $\mathfrak{f} \colon \mathfrak{Y} \to \mathfrak{X}$, the natural morphism

$$\mathfrak{f}_{n-1}^*\left(\mathrm{R}^i\nu_{\mathfrak{X}_C,*}(\mathbf{L}\otimes_{\mathbf{Z}/p^n\mathbf{Z}}\mathfrak{O}_{\mathfrak{X}_C}^+/p^n)\right)\to\mathrm{R}^i\nu_{\mathfrak{Y}_C,*}(\mathfrak{f}_C^*\mathbf{L}\otimes_{\mathbf{Z}/p^n\mathbf{Z}}\mathfrak{O}_{\mathfrak{Y}_C}^+/p^n)$$

is an almost isomorphism for every $i \ge 0$.

Proof. We prove the claim by induction on n. The case of n = 1 is done in Theorem 6.4.11. Now we suppose n > 1 and that the statement is proven for n - 1 and show it for n. The main idea is to use Corollary C.10 to reduce the claim for $\mathbf{L} \otimes_{\mathbf{Z}/p^n \mathbf{Z}} \mathcal{O}_X^+/p^n$ to the claims for $(p\mathbf{L}) \otimes_{\mathbf{Z}/p^{n-1}\mathbf{Z}} \mathcal{O}_X^+/p^{n-1}$ and $(\mathbf{L}/p\mathbf{L}) \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p$ that we can deal with by the induction hypothesis. More precisely, we have a short exact sequence

$$0 \to (p\mathbf{L}) \otimes_{\mathbf{Z}/p^{n-1}\mathbf{Z}} \mathfrak{O}_X^+/p^{n-1} \to \mathbf{L} \otimes_{\mathbf{Z}/p^n\mathbf{Z}} \mathfrak{O}_X^+/p^n \to (\mathbf{L}/p\mathbf{L}) \otimes_{\mathbf{F}_p} \mathfrak{O}_X^+/p \to 0$$
(6.3)

And we show that once we know the claim for the left and the right objects, then we know the result for the middle one. We proceed step by step.

Step 1. The sheaves $\mathbb{R}^i \nu_*(\mathbf{L} \otimes_{\mathbf{Z}/p^n \mathbf{Z}} \mathbb{O}_X^+/p^n)$ are almost quasi-coherent: Similarly to the proof of Theorem 6.2.11, it is sufficient to assume that $\mathfrak{X} = \operatorname{Spf} A$ is affine. Moreover, it suffices to show a stronger statement that the natural map

$$\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\mathbf{Z}/p^{n}\mathbf{Z}} \mathbb{O}_{X}^{+}/p^{n}) \to \mathrm{R}^{i}\nu_{*}(\mathbf{L} \otimes_{\mathbf{Z}/p^{n}\mathbf{Z}} \mathbb{O}_{X}^{+}/p^{n})$$

$$(6.4)$$

is an almost isomorphism. For any $f \in A$, we denote by \mathfrak{X}_f the principal open formal subscheme Spf $A_{\{f\}}$. We denote its generic fiber $\mathfrak{X}_{f,C}$ by X_f . Then the map (6.4) is an almost isomorphism if the morphism

$$\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\mathbf{Z}/p^{n}\mathbf{Z}} \mathbb{O}_{X}^{+}/p^{n}) \otimes_{A/p^{n}} A_{f}/p^{n} \to \mathrm{H}^{i}(X_{f}, \mathbf{L} \otimes_{\mathbf{Z}/p^{n}\mathbf{Z}} \mathbb{O}_{X}^{+}/p^{n})$$

is an almost isomorphism for every $f \in A$. We know that this holds for $(p\mathbf{L}) \otimes_{\mathbf{Z}/p^{n-1}\mathbf{Z}} \mathcal{O}_X^+/p^{n-1}$ and $(\mathbf{L}/p\mathbf{L}) \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p$ by the induction assumption. Thus it holds for $\mathbf{L} \otimes_{\mathbf{Z}/p^n\mathbf{Z}} \mathcal{O}_X^+/p^n$ by A/p^n -flatness of A_f/p^n and the induction hypothesis.

Step 2. The complex $\mathbb{R}\nu_*(\mathbb{L}\otimes_{\mathbb{Z}/p^n\mathbb{Z}} \mathcal{O}_X^+/p^n)$ is almost coherent: The question is again Zariski-local on \mathfrak{X} , so we may and do assume that $\mathfrak{X} = \operatorname{Spf} A$ is affine. Then Step 1 implies that it is sufficient to show that $\mathrm{H}^i(X, \mathbb{L}\otimes_{\mathbb{Z}/p^n\mathbb{Z}} \mathcal{O}_X^+/p^n)$ is almost coherent over A/p^n for any $i \geq 0$. Lemma 2.7.3 guarantees that an A/p^n -module is M is A-almost coherent if and only if it is A/p^n -almost coherent. Thus the induction hypothesis and Lemma 4.1.11 assert that

$$\mathrm{H}^{i}(X,(p\mathbf{L})\otimes_{\mathbf{Z}/p^{n-1}\mathbf{Z}} \mathbb{O}_{X}^{+}/p^{n-1}) \text{ and } \mathrm{H}^{i}(X,(\mathbf{L}/p\mathbf{L})\otimes_{\mathbf{F}_{p}} \mathbb{O}_{X}^{+}/p)$$

are almost coherent over A for $i \ge 0$. We write the long exact sequence associated with the short exact sequence 6.3 to obtain that the sequence

$$\begin{array}{c} \mathrm{H}^{i-1}(X, (\mathbf{L}/p\mathbf{L}) \otimes_{\mathbf{F}_{p}} \mathbb{O}_{X}^{+}/p) \xrightarrow{\delta^{i-1}} \mathrm{H}^{i}(X, (p\mathbf{L}) \otimes_{\mathbf{Z}/p^{n-1}\mathbf{Z}} \mathbb{O}_{X}^{+}/p^{n-1}) \longrightarrow \mathrm{H}^{i}(X, \mathbf{L} \otimes_{\mathbf{Z}/p^{n}\mathbf{Z}} \mathbb{O}_{X}^{+}/p^{n}) \\ \\ \end{array} \\ \xrightarrow{\mathrm{H}^{i}(X, (\mathbf{L}/p\mathbf{L}) \otimes_{\mathbf{F}_{p}} \mathbb{O}_{X}^{+}/p)} \underbrace{\overleftarrow{\delta^{i}}}_{\delta^{i}} \mathrm{H}^{i}(X, (p\mathbf{L}) \otimes_{\mathbf{Z}/p^{n-1}\mathbf{Z}} \mathbb{O}_{X}^{+}/p^{n-1})} \end{array}$$

is exact. We note that $\operatorname{Coker} \delta^{i-1}$ and $\ker \delta^i$ are both almost coherent by Lemma 2.6.7(4). Then we use Lemma 2.6.7(5) to conclude that $\operatorname{H}^i(X, \mathbf{L} \otimes_{\mathbf{Z}/p^n \mathbf{Z}} \mathcal{O}_X^+/p^n)$ is almost coherent over A.

Step 3. Boundedness of the complex $\mathbf{R}\nu_*(\mathbf{L} \otimes_{\mathbf{Z}/p^n\mathbf{Z}} \mathcal{O}_X^+/p^n)$: The only thing we are left to show is that

$$\mathbf{R}\nu_*(\mathbf{L}\otimes_{\mathbf{Z}/p^n\mathbf{Z}} \mathfrak{O}_X^+/p^n) \in \mathbf{D}^{[0,2d]}(X_{n-1}) \text{ and } \mathbf{R}\nu_*(\mathbf{L}\otimes_{\mathbf{Z}/p^n\mathbf{Z}} \mathfrak{O}_X^+/p^n) \in \mathbf{D}^{[0,d]}(X_{n-1})^a$$

These questions are Zariski-local on \mathfrak{X} , so we may and do assume that $\mathfrak{X} = \operatorname{Spf} A$ is affine. Then Step 1 says that what we need to show is that $\mathbf{R}\Gamma(X, \mathbf{L} \otimes_{\mathbf{Z}/p^n\mathbf{Z}} \mathfrak{O}_X^+/p^n)$ is concentrated in degrees [0, 2d] and almost concentrated in degrees [0, d]. Again, the same induction argument as in Step 2 shows this and finishes the proof.

Step 4. Formation of $\mathbb{R}^i \nu_*(\mathbf{L} \otimes_{\mathbf{Z}/p^n \mathbf{Z}} \mathfrak{O}_X^+/p^n)$ almost commutes with étale base change: The question is local on \mathfrak{X} and \mathfrak{Y} , so we can assume that $\mathfrak{X} = \operatorname{Spf} A$ and $\mathfrak{Y} = \operatorname{Spf} B$ are affine. Using

flatness of \mathfrak{f} and already established almost coherence of $\mathbf{R}\Gamma(\mathfrak{X}_C, \mathbf{L} \otimes_{\mathbf{Z}/p^n\mathbf{Z}} \mathfrak{O}_X^+/p^n)$, it suffices to show that the natural morphism

$$\mathbf{R}\Gamma(\mathfrak{X}_{C},\mathbf{L}\otimes_{\mathbf{Z}/p^{n}\mathbf{Z}}\mathbb{O}_{\mathfrak{X}_{C}}^{+}/p^{n})\otimes_{A/p^{n}A}^{L}B/p^{n}B\to\mathbf{R}\Gamma(\mathfrak{Y}_{C},\mathbf{L}\otimes_{\mathbf{Z}/p^{n}\mathbf{Z}}\mathbb{O}_{\mathfrak{Y}_{C}}^{+}/p^{n})$$

is an almost isomorphism. Then we note that

$$\mathbf{R}\Gamma(\mathfrak{X}_{C},(\mathbf{L}/p\mathbf{L})\otimes_{\mathbf{Z}/p^{n}\mathbf{Z}}\mathbb{O}_{\mathfrak{X}_{C}}^{+}/p^{n})\otimes_{A/p^{n}A}^{L}B/p^{n}B\simeq\mathbf{R}\Gamma(\mathfrak{X}_{C},(\mathbf{L}/p\mathbf{L})\otimes_{\mathbf{F}_{p}}\mathbb{O}_{\mathfrak{X}_{C}}^{+}/p)\otimes_{A/p^{n}A}^{L}B/p^{n}B\\\simeq\mathbf{R}\Gamma(\mathfrak{X}_{C},(\mathbf{L}/p\mathbf{L})\otimes_{\mathbf{F}_{p}}\mathbb{O}_{\mathfrak{X}_{C}}^{+}/p)\otimes_{A/p^{A}}^{L}B/pB$$

since B is A-flat. Similarly,

$$\mathbf{R}\Gamma(\mathfrak{X}_{C},\mathbf{L}[p]\otimes_{\mathbf{Z}/p^{n}\mathbf{Z}} \mathbb{O}_{\mathfrak{X}_{C}}^{+}/p^{n})\otimes_{A/p^{n}A}^{L}B/p^{n}B\simeq \mathbf{R}\Gamma(\mathfrak{X}_{C},\mathbf{L}[p]\otimes_{\mathbf{F}_{p}}\mathbb{O}_{\mathfrak{X}_{C}}^{+}/p)\otimes_{A/pA}^{L}B/pB$$

Therefore, one can use the short exact sequence (6.3) to reduce to the case n = 1 that was already done in Corollary 6.2.12.

Now we move to the proof of Theorem 6.1.1 for lisse $\widehat{\mathbf{Z}}_p$ -sheaves. The idea of the proof is firstly to show that $\mathbf{R}\nu_*(\mathbf{L}\otimes_{\widehat{\mathbf{Z}}_p}\widehat{\mathcal{O}}_X^+) \in \mathbf{D}^{[0,2\dim X+1]}(\mathfrak{X})$ for any $\mathbf{L} \in \mathrm{LocSys}_{\mathbf{Z}_p,X_{\mathrm{prooft}}}$. This, in turn, will allow us to imitate the proof of Theorem 5.1.6 to show that $\mathbf{R}\nu_*(\mathbf{L}\otimes_{\widehat{\mathbf{Z}}_p}\widehat{\mathcal{O}}_X^+) \in \mathbf{D}^{[0,2\dim X+1]}_{acoh}(\mathfrak{X})$. Finally, using the almost coherence of this complex as an input, we will show that the complex $\mathbf{R}\nu_*(\mathbf{L}\otimes_{\widehat{\mathbf{Z}}_p}\widehat{\mathcal{O}}_X^+)$ is concentrated in degrees $[0, 2\dim X]$ and almost concentrated in degrees $[0, \dim X]$.

Lemma 6.5.2. Let $X = \text{Spa}(A, A^+)$ be an affinoid rigid-analytic space over $\text{Spa}(C, \mathcal{O}_C)$ of dimension d, and let \mathbf{L} be a lisse \mathbf{Z}_p -sheaf on $X_{\text{pro\acute{e}t}}$. Then $\text{H}^i(X, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathcal{O}}_X^+) = 0$ for i > 2d + 1.

Proof. Lemma C.12 reads that we have the isomorphism

$$\mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathcal{O}}_X^+ \to \mathbf{R} \lim_n \left((\mathbf{L}/p^n \mathbf{L}) \otimes_{\mathbf{Z}/p^n \mathbf{Z}} \left(\mathcal{O}_X^+/p^n \right) \right)$$

As $\mathbf{R}\Gamma$ commutes with derived limits by [Sta21, Tag 0A07], we conclude that we have an isomorphism

$$\mathbf{R}\Gamma(X,\mathbf{L}\otimes_{\widehat{\mathbf{Z}}_p}\widehat{\mathcal{O}}_X^+)\simeq\mathbf{R}\lim_n\mathbf{R}\Gamma\left(X,(\mathbf{L}/p^n\mathbf{L})\otimes_{\mathbf{Z}/p^n\mathbf{Z}}\left(\mathcal{O}_X^+/p^n\right)\right)$$

Now we consider Milnor exact sequence applied to the right side:

$$0 \to \mathrm{R}^{1} \lim_{n} \mathrm{H}^{i-1} \left(X, \left(\mathbf{L}/p^{n} \mathbf{L} \right) \otimes \left(\mathcal{O}_{X}^{+}/p^{n} \right) \right) \to \mathrm{H}^{i} \left(X, \mathbf{L} \otimes \widehat{\mathcal{O}}_{X}^{+} \right) \to \lim_{n} \mathrm{H}^{i} \left(X, \left(\mathbf{L}/p^{n} \mathbf{L} \right) \otimes \left(\mathcal{O}_{X}^{+}/p^{n} \right) \right) \to 0$$

As each $\mathbf{L}/p^{n}\mathbf{L}$ comes from $\mathrm{LocSys}_{\mathbf{Z}/p^{n}\mathbf{Z},X_{\mathrm{\acute{e}t}}}$ by Theorem C.4, we conclude that

$$\mathrm{H}^{i}\left(X,\left(\mathbf{L}/p^{n}\mathbf{L}\right)\otimes\left(\mathcal{O}_{X}^{+}/p^{n}\right)\right)=0$$

for i > 2d. Therefore, we get that $\mathrm{H}^{i}\left(X, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathcal{O}}_{X}^{+}\right) = 0$ for i > 2d + 1.

Lemma 6.5.3. Let \mathfrak{X} an admissible formal \mathcal{O}_C -scheme with the adic generic fiber $X := \mathfrak{X}_C$ of dimension d. Then

$$\mathbf{R}\nu_*(\mathbf{L}\otimes_{\widehat{\mathbf{Z}}_p}\widehat{\mathbb{O}}_X^+)\in\mathbf{D}_{acoh}^{[0,2d]}(\mathfrak{X}),\ \mathbf{R}\nu_*(\mathbf{L}\otimes_{\widehat{\mathbf{Z}}_p}\widehat{\mathbb{O}}_X^+)\in\mathbf{D}_{acoh}^{[0,d]}(\mathfrak{X})^a$$

for $\mathbf{L} \in \operatorname{LocSys}_{\mathbf{Z}_p, X_{\operatorname{pro\acute{e}t}}}$. More precisely, for an affine admissible $\mathfrak{X} = \operatorname{Spf} A$ with the adic generic fiber X, the canonical morphism

$$\mathrm{H}^{i}(X,\mathbf{L}\otimes_{\widehat{\mathbf{Z}}_{p}}\widehat{\mathrm{O}}_{X}^{+})^{\Delta}\to\mathrm{R}^{i}\nu_{*}(\mathbf{L}\otimes_{\widehat{\mathbf{Z}}_{p}}\widehat{\mathrm{O}}_{X}^{+})$$

is an almost isomorphism. And, moreover, the formation of $\mathrm{R}^i \nu_*(\mathbf{L} \otimes_{\mathbf{Z}_p} \widehat{\mathcal{O}}_X^+)$ almost commutes with étale base change, i.e., for any étale morphism $\mathfrak{f}: \mathfrak{Y} \to \mathfrak{X}$, the natural morphism

$$\mathfrak{f}^*\left(\mathrm{R}^i\nu_{\mathfrak{X}_C,*}(\mathbf{L}\otimes_{\mathbf{Z}_p}\widehat{\mathcal{O}}^+_{\mathfrak{X}_C})\right)\to\mathrm{R}^i\nu_{\mathfrak{Y}_C,*}(\mathfrak{f}^*_C\mathbf{L}\otimes_{\mathbf{Z}_p}\widehat{\mathcal{O}}^+_{\mathfrak{Y}_C})$$

is an almost isomorphism for every $i \ge 0$.

Proof. The first claim is local on \mathfrak{X} , so we may and do assume that $\mathfrak{X} = \operatorname{Spf} A$ and $\mathfrak{Y} = \operatorname{Spf} B$ are affine and connected. Similarly to Lemma 5.1.8⁴³, we see that it is sufficient to show the three following claims:

- Claim 1: the A-module $\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathcal{O}}_{X}^{+})$ is almost coherent for any $i \geq 0$ and any $\mathbf{L} \in \mathrm{LocSys}_{\mathbf{Z}_{p}, X_{\mathrm{pro\acute{e}t}}}$.
- Claim 2: for any $g \in A$, the canonical map

$$\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathrm{O}}_{X}^{+}) \otimes_{A} A_{\{g\}} \to \mathrm{H}^{i}(X_{g}, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathrm{O}}_{X}^{+}),$$

where X_g is the adic generic fiber $(\operatorname{Spf} A_{\{g\}})_C$, is an almost isomorphism for any $i \ge 0$ and any $\mathbf{L} \in \operatorname{LocSys}_{\mathbf{Z}_p, X_{\operatorname{pro\acute{e}t}}}$.

- Claim 3: For any i > 2d, $\operatorname{H}^{i}(X, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathcal{O}}_{X}^{+}) = 0$. And, for any i > d, $\operatorname{H}^{i}(X, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathcal{O}}_{X}^{+}) \cong^{a} 0$.
- Claim 4: The natural morphism $\mathbf{R}\Gamma(\mathfrak{X}_C, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathcal{O}}^+_{\mathfrak{X}_C}) \otimes^L_A B \to \mathbf{R}\Gamma(\mathfrak{Y}_C, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathcal{O}}^+_{\mathfrak{Y}_C})$ is an almost isomorphism.

Now we start proving these three claims.

Step 1. Reduction to the case of "locally free" lisse sheaf \mathbf{L} : We consider the subsheaves $\mathbf{L}[p^m] \subset \mathbf{L}$ of p^m -torsion sections. As \mathbf{L} is locally given by $M \times_{\mathbf{Z}_p} \widehat{\mathbf{Z}}_p$ for some finitely generated \mathbf{Z}_p -module M, we conclude that there is some m such that $\mathbf{L}[p^m] = \mathbf{L}[p^{m+k}]$ for any $k \geq 0$. Then we see that $\mathbf{L}/\mathbf{L}[p^m]$ is p-torsion free, so pro-étale locally it is isomorphic to $\widehat{\mathbf{Z}}_p^r$ for some r. Moreover, we have a short exact sequence

$$0 \to \mathbf{L}[p^m] \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathbb{O}}_X^+ \to \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathbb{O}}_X^+ \to (\mathbf{L}/\mathbf{L}[p^m]) \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathbb{O}}_X^+ \to 0$$

Now we use that the maps $A \to A_{\{g\}}$ and $A \to B$ are flat to conclude that it suffices to show all four *Claims* separately for $\mathbf{L}[p^m]$ and $\mathbf{L}/\mathbf{L}[p^m]$. But $\mathbf{L}[p^m]$ comes from $\mathrm{LocSys}_{\mathbf{Z}/p^m\mathbf{Z},X_{\acute{e}t}}$ by Theorem C.4, so we know all four *Claims* for it by Theorem 6.5.1. Thus, it suffices to show the claim for $(\mathbf{L}/\mathbf{L}[p^m])$ that pro-étale locally is isomorphic to $\widehat{\mathbf{Z}}_p^r$ for some r.

Step 2. Claim 1 and Claim 2 for "locally free" lisse sheaf \mathbf{L} : We choose the universal covering space $\widetilde{X} \to X$ from Section 6.2. More precisely, we define here \widetilde{X} as the inverse system " $\lim_i \operatorname{Spa}(A_i, A_i^+)$ ", where A_i is roughly the system of all finite étale extensions of A[1/p] and A_i^+ is the integral closure of A in A_i . We refer to the discussion after Lemma 6.2.5 for the precise construction, but what we denoted there by $\widetilde{X} = \operatorname{Spa}(\widehat{\overline{A}}[1/p], \widehat{\overline{A}})$ is now the adic space $\widehat{\widetilde{X}}$ associated to the system \widetilde{X} . In any case, $\widetilde{X} \to X$ is a pro-étale covering of X by an affinoid perfectoid space. Moreover, it does not have any non-trivial finite étale covers in $X_{\text{proét}}$ by Lemma 6.2.5 and [Sch12, Lemma 7.5]. Thus each quotient $\mathbf{L}/p^n\mathbf{L}$ splits on \widetilde{X} , i.e. $\mathbf{L}/p^n\mathbf{L}|_{\widetilde{X}} \simeq (\mathbf{Z}/p^n\mathbf{Z})^r$. Therefore, we conclude that $\mathbf{L}|_{\widetilde{X}} \simeq \widehat{\mathbf{Z}}_p^r$.

We introduce the complex $K^{\prime \bullet} := \check{C}(\widetilde{X}/X, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathcal{O}}_X^+)$ that is just the Čech complex with respect to the pro-étale covering \widetilde{X}/X . Then our assumptions on \mathbf{L} and [Sch13, Lemma 4.10] imply that

 $^{^{43}}$ And using flatness of f and Lemma 4.7.3 for Claim 4

this complex almost computes cohomology of the sheaf $\mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathcal{O}}_X^+$. In particular, it is almost acyclic in degrees greater than 2d + 1 by Lemma 6.5.2. Moreover, loc.cit. guarantees that the quotient K'^{\bullet}/p^{k+1} almost computes cohomology groups of $(\mathbf{L}/p^{k+1}\mathbf{L}) \otimes_{\mathbf{Z}/p^{k+1}\mathbf{Z}} \mathcal{O}_X^+/p^{k+1}$, so it is almost acyclic in degrees greater than 2d by Theorem 6.4.10. We now define the complex $K^{\bullet} \coloneqq \mathfrak{m}K'^{\bullet}$ and $K_k^{\bullet} \coloneqq \mathfrak{m}K'^{\bullet}$, so we have a short exact sequence

$$0 \to K^{\bullet} \xrightarrow{p^{k+1}} K^{\bullet} \to K_k^{\bullet} \to 0$$

The above discussion implies that

$$\mathrm{H}^{i}(K^{\bullet}) \cong^{a} \mathrm{H}^{i}(X, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathrm{O}}_{X}^{+}), \ \mathrm{H}^{i}(K_{k}^{\bullet}) \cong^{a} \mathrm{H}^{i}(X, (\mathbf{L}/p^{k+1}\mathbf{L}) \otimes \mathrm{O}_{X}^{+}/p^{k+1}),$$

and both complexes are acyclic in degrees at least 2d+1. Each term K_i is isomorphic to $\widehat{\mathfrak{mO}}_X^+(X_{\infty}^{i/X})$ that is *p*-torsionfree and *p*-adically complete by [Sch13, Lemma 4.2]. We explain why it is *p*-adically complete in more detail: firstly, [FK18, Lemma 0.8.2.14] implies that the induced topology $\widehat{\mathfrak{mO}}_X^+(X_{\infty}^{i/X})$ coincides with the *p*-adic topology. Secondly, this submodule is closed in $\widehat{\mathcal{O}}_X^+(X_{\infty}^{i/X})$ as a kernel of a continuous morphism to a discrete module. Thus, it is complete in the induced topology, therefore it is complete in the *p*-adic topology. Finally, we know that *Claim 1* and *Claim 2* hold for cohomology of the complexes K_k^{\bullet} for all *k*. This follows from Theorem 6.5.1.

Once we verified all these properties of the complexes K^{\bullet} and K_k^{\bullet} , the arguments that go into proving Theorem 5.1.6 also apply mutatis mutandis to prove *Claim 1* and *Claim 2*. The only change is that we do not need to prove the *Claims* for all **L** simultaneously as $p^k \mathbf{L}$ is abstractly isomorphic to **L** in our situation.

Step 3. Claim 3 for "locally free" lisse sheaf L: Now we know that the complex $\mathbf{R}\Gamma(X, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathcal{O}}_X^+)$ is bounded and has almost coherent cohomology groups. So we consider a short exact sequence

$$0 \to \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathcal{O}}_X^+ \xrightarrow{p} \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathcal{O}}_X^+ \to (\mathbf{L}/p\mathbf{L}) \otimes_{\mathbf{F}_p} \mathcal{O}_X^+/p \to 0$$
(6.5)

Suppose that there is i > 2d such that $\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathbb{O}}_{X}^{+}) \neq 0$. Then the long exact sequence associated to the sequence (6.5), Theorem 6.4.11 and vanishing of $\mathrm{H}^{i}(X, (\mathbf{L}/p\mathbf{L}) \otimes_{\mathbf{F}_{p}} \mathbb{O}_{X}^{+}/p)$ for i > 2d show that

$$\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathrm{O}}_{X}^{+})/p \simeq 0$$

As the A-module $\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathcal{O}}_{X}^{+})$ is almost coherent by Claim 1, we conclude that $\mathrm{H}^{i}(X, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathcal{O}}_{X}^{+})$ must be zero by Lemma 2.5.15. The same proof shows that $\mathbf{R}\Gamma(X, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathcal{O}}_{X}^{+})$ is almost concentrated in degrees [0, d].

Step 4. Claim 4 for a "locally free" lisse $\widehat{\mathbf{Z}}_p$ -sheaf L: We already know that

$$\widetilde{\mathfrak{m}} \otimes \mathbf{R}\Gamma(\mathfrak{X}_C, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathfrak{O}}^+_{\mathfrak{X}_C}) \in \mathbf{D}^b_{qc,acoh}(\mathrm{Spf}\,A) \simeq \mathbf{D}^b_{acoh}(A)$$

by Step 2 and Theorem 4.6.15. Suppose

$$\widetilde{\mathfrak{m}} \otimes \mathbf{R}\Gamma(\mathfrak{X}_C, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathfrak{O}}^+_{\mathfrak{X}_C}) \simeq M$$

for some $M \in \mathbf{D}^b_{acoh}(A)$. Similarly, we conclude that

 $\widetilde{\mathfrak{m}} \otimes \mathbf{R} \Gamma(\mathfrak{Y}_C, \mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathcal{O}}_{\mathfrak{Y}_C}^+) \simeq N$

for some $N \in \mathbf{D}^b_{acoh}(B)$.

We want to show that the natural morphism

$$M \otimes^L_A B \to N$$

is an isomorphism. It suffices to show that its cone $C := \operatorname{cone}(M \otimes_A^L B \to N)$ is zero. We firstly note that $C \in \mathbf{D}_{acoh}(B)$ as a consequence of Corollary 2.7.2. And, therefore, Corollary 2.8.8 implies that M is derived p-adically complete modules.

Now Corollary 6.2.12 and local freeness of **L** imply that $C \otimes_B^L B/pB \simeq 0$. So the derived Nakayama Lemma [Sta21, Tag 0G1U] implies that $C \simeq 0$ finishing the proof.

Appendix

Appendix A. Complexes over Adhesive Rings

This section is just an elaboration of the section 2.8, but we stick ourselves to a more restrictive situation⁴⁴ to prove some more delicate statements.

Set-up A.1. We fix a ring R with an element ϖ and a compatible choice of roots $\varpi^{1/n}$ for each integer $n \ge 0$. Now we require R to be complete in the ϖ -adic topology and ϖ -adically adhesive. We define the ideal of almost mathematics as $\mathfrak{m} = \bigcup_n \varpi^{1/n} R$. It is clear that $\mathfrak{m}^2 = \mathfrak{m}$ and [GR03, Proposition 2.1.7] implies that $\widetilde{\mathfrak{m}} = \mathfrak{m} \otimes_R \mathfrak{m}$ is flat.

The basic example of such an algebra is an algebra R that is topologically finitely presented over a complete microbial valuation ring k^+ with algebraically closed fraction field. Then we pick ϖ to be a pseudo-uniformizer of k^+ and choose some compatible systems of roots $\varpi^{1/n}$ of ϖ using the fact that the fraction field of k^+ is algebraically closed. We note that R is adhesive by [FGK11, Theorem 7.3.2]

Lemma A.2. Let R be as in the Set-up A.1, and let $f: M \to N$ be a morphism of A-modules. Denote the cokernel by H, and suppose that

- M is ϖ -adically complete, and N is ϖ -adically separated
- H/ϖ is almost finitely generated.

Then H is almost finitely generated.

Proof. For any integer n we can find a map $\overline{\varphi} \colon \mathbb{R}^k \to H/\pi$ such that $Q \coloneqq \operatorname{Coker} \overline{\varphi}$ is killed by $\overline{\omega}^{1/n}$. Lift this map in an arbitrary way to a map $\varphi \colon \mathbb{R}^k \to H$ and denote the cokernel by $Q' \coloneqq \operatorname{Coker} \varphi$. We can also lift φ to some map $\Phi \colon \mathbb{R}^k \to N$ as $N \to H$ is surjective. We can summarize those constructions in the following commutative diagram:



Note that $\overline{\pi}: Q \to Q'$ is clearly surjective. If we can prove that this is an isomorphism, then it will imply that Q is $\overline{\omega}^{1/n}$ -torsion. Since n was arbitrary, it will imply that H is almost finitely generated over R. So we reduce the claim to show that $\overline{\pi}$ is injective.

We pick any $h \in H$ such that $\pi(h)$ lies in the image of $\overline{\varphi}$. In order to check that $\overline{\pi}$ is injective it suffices to show that h lives in the image of φ . However, we observe that the condition that $\pi(h) \in \text{Im}(\overline{\varphi})$ shows that there are elements $y_0 \in \mathbb{R}^k$ and $h_1 \in H$ such that

$$h = \varphi(y_0) + \varpi h_1 = \varphi(y_0) + \varpi^{1-1/n} \varpi^{1/n} h_1$$

 $^{^{44}}$ We do not know how to deal with the more general situation at the moment.

Now we use surjectivity of g to find $n_1 \in N$ such that $g(n_1) = h_1$. Here comes the crucial step, we note that the hypothesis that $\operatorname{Coker} \overline{\varphi}$ is annihilated by $\overline{\varphi}^{1/n}$ implies that

$$\varpi^{1/n} N \subset \Phi(R^k) + f(M) + \varpi N$$

In particular, we can find some $y_1 \in \mathbb{R}^k, m_1 \in M$ and $n_2 \in N$ such that

$$\varpi^{1/n} n_1 = \Phi(y_1) + f(m_1) + \varpi n_2 = \Phi(y_1) + f(m_1) + \varpi^{1-1/n} \varpi^{1/n} n_2$$

Applying the same procedure to $\varpi^{1/n} n_2$ we find $y_2 \in \mathbb{R}^k, m_2 \in M$ and $n_3 \in \mathbb{N}$ such that

$$\varpi^{1/n} n_2 = \Phi(y_2) + f(m_2) + \varpi n_3 = \Phi(y_2) + f(m_2) + \varpi^{1-1/n} \varpi^{1/n} n_3.$$

In particular, this implies that

$$\varpi^{1/n} n_1 = \Phi(y_1 + \varpi^{1-1/n} y_2) + f(m_1 + \varpi^{1-1/n} m_2) + \varpi^{2(1-1/n)} \varpi^{1/n} n_3.$$

We keep going this procedure to arrive at step r with a sequence of $y_1, \ldots, y_r \in \mathbb{R}^k, m_1, \ldots, m_2 \in M$ and $n_{r+1} \in N$ such that

$$\varpi^{1/n} n_1 = \Phi(y_1 + \varpi^{1-1/n} y_2 + \dots + \varpi^{(r-1)(1-1/n)} y_r) + f(m_1 + \varpi^{1-1/n} m_2 + \dots + \varpi^{(r-1)(1-1/n)} m_r) + \varpi^{r(1-1/n)} \varpi^{1/n} n_{r+1}.$$

We note that since N is ϖ -adically separated, we know that the limit of the sequence $\varpi^{r(1-1/n)} \varpi^{1/n} n_{r+1}$ is zero. Also, we recall that M and R^k are ϖ -complete by the assumption, thus each of the sums:

$$y_1 + \varpi^{1-1/n} y_2 + \dots + \varpi^{(r-1)(1-1/n)} y_r$$
 and $m_1 + \varpi^{1-1/n} m_2 + \dots + \varpi^{(r-1)(1-1/n)} m_r$

converge in \mathbb{R}^k and M, respectively. Denote those limits by $y \in \mathbb{R}^k$ and $m \in M$, then the discussion above guarantees that $\varpi^{1/n} n_1 = \Phi(y) + f(m)$. Thus we get

$$h = \varphi(y_0) + \varpi^{1-1/n} \varpi^{1/n} h_1 = \varphi(y_0) + \varpi^{1-1/n} g(n_1) = \varphi(y_0 + \varpi^{1-1/n} y).$$

deed in the image of ϕ .

So h is indeed in the image of ϕ .

Corollary A.3. Let R be as in the Set-up A.1, and let M be an ϖ -adically separated R-modules such that M/ϖ is almost finitely generated. Then M is almost finitely generated.

We also need two other statements that will be essential for our proof of the Almost Proper Mapping Theorem for formal schemes. Let us introduce the context. We consider a complex $(K^{\bullet}, d^{\bullet})$ of R-modules such that each term K^q is ϖ -adically complete. Then we can consider complexes $\varpi^{k+1} K^{\bullet} \subset K^{\bullet}$ and denote the quotient complex by K_k^{\bullet} :

$$0 \to \varpi^{k+1} K^{\bullet} \to K^{\bullet} \to K_k^{\bullet} \to 0$$

We define the filtration $F^{\bullet}H^{i}(K^{\bullet})$ as follows:

$$\mathbf{F}^{n}\mathbf{H}^{i}(K^{\bullet}) \coloneqq \mathrm{Im}(\mathbf{H}^{i}(\varpi^{n}K^{\bullet}) \to \mathbf{H}^{i}(K^{\bullet}))$$

The natural question to ask is whether this filtration defines the ϖ -adic topology on $H^i(K^{\bullet})$ for each i. It turns out that even more is true under the assumption that K^{q+1} has bounded ϖ^{∞} -torsion:

Lemma A.4. Let R and K^{\bullet} be as above. Suppose that $K^{q+1}[\varpi^{\infty}] = K^{q+1}[\varpi^n]$ for some n. Then for any $k \ge n$ we have

$$\varpi^k \mathrm{H}^q(K^{\bullet}) \subset \mathrm{F}^k \mathrm{H}^q(K^{\bullet}) \subset \varpi^{k-n} \mathrm{H}^q(K^{\bullet})$$

In particular, the filtration $F^{\bullet}H^{q}(K^{\bullet})$ defines the ϖ -adic topology on $H^{q}(K^{\bullet})$

Proof. We consider a short exact sequence

 $0 \to \ker d^q \to K^q \to \operatorname{Im} d^q \to 0$

and notice that $(\operatorname{Im} d^q)[\varpi^{\infty}] = (\operatorname{Im} d^q)[\varpi^n]$ as $\operatorname{Im} d^q$ is a submodule of a ϖ -adically complete module K^{q+1} . Therefore, [FK18, Lemma 0.8.2.14] implies that

$$\ker d^q \cap \varpi^{n+m} K^{q+1} = \varpi^m (\ker d^q \cap \varpi^n K^{q+1}) \text{ for any } m \ge 0$$

In particular, for any $k \ge n$ we have

$$\varpi^k \ker d^q \subset \ker d^q \cap \varpi^k K^q \subset \varpi^{k-n} \ker d^q$$

This, in turn, guarantees that

$$\varpi^k \mathrm{H}^q(K^{\bullet}) \subset \mathrm{F}^k \mathrm{H}^q(K^{\bullet}) \subset \varpi^{k-n} \mathrm{H}^q(K^{\bullet})$$

and finishes the proof.

Lemma A.5. Let R a ring as in the Set-up A.1 and let $R \to R'$ be a flat morphisms such that R' is π -adically complete and π -adically adhesive. Let K^{\bullet} (resp. L^{\bullet}) be a complex of R-modules (resp. R'-modules) such that each term is ϖ -adically complete, and let $K^{\bullet} \to L^{\bullet}$ be a morphism of R-complexes. Then this morphism induces a morphism of the corresponding long exact sequences⁴⁵:

$$\begin{array}{cccc} \mathrm{H}^{q}(\varpi^{k+1}K^{\bullet}) \otimes_{R} R' & \stackrel{a_{k}^{q}}{\longrightarrow} \mathrm{H}^{q}(K^{\bullet}) \otimes_{R} R' & \stackrel{b_{k}^{q}}{\longrightarrow} \mathrm{H}^{q}(K_{k}^{\bullet}) \otimes_{R} R' & \stackrel{c_{k}^{q}}{\longrightarrow} \mathrm{H}^{q+1}(\varpi^{k+1}K^{\bullet}) \otimes_{R} R' \\ & & \downarrow \varphi_{k}^{q} & & \downarrow \varphi_{k}^{q} & & \downarrow \varphi_{k}^{q+1} \\ \mathrm{H}^{q}(\varpi^{k+1}L^{\bullet}) & \stackrel{\alpha_{k}^{q}}{\longrightarrow} \mathrm{H}^{q}(L^{\bullet}) & \stackrel{\beta_{k}^{q}}{\longrightarrow} \mathrm{H}^{q}(L_{k}^{\bullet}) & \stackrel{\gamma_{k}^{q}}{\longrightarrow} \mathrm{H}^{q+1}(\varpi^{k+1}L^{\bullet}) \end{array}$$

Suppose that

- $\mathrm{H}^{q}(K^{\bullet})$ and $\mathrm{H}^{q}(L^{\bullet})$ are almost coherent R and R'-modules, respectively.
- $K^{q+1}[\varpi^{\infty}] = K^{q+1}[\varpi^n]$ and $L^{q+1}[\varpi^{\infty}] = L^{q+1}[\varpi^n]$ for some n.
- Φ_k^q is an isomorphism for any $k \ge n$.
- $\varphi_k^{\vec{q}}$ is injective for any $k \ge n$.

Then φ^q is an isomorphism.

Proof. We divide the proof into two part: we first show injectivity of the map φ^q and then surjectivity.

Injectivity of φ^q : We note that R' fits into the Set-up 2.8.3. Thus Lemma 2.7.1 and Lemma 2.8.7 imply that $\mathrm{H}^q(K^{\bullet}) \otimes_R R'$ is ϖ -adically complete. In particular, it is ϖ -adically separated. Therefore, it suffices to show that $\mathrm{ker}(\varphi^q)$ lies in $\varpi^m(\mathrm{H}^q(K^{\bullet}) \otimes_R R')$ for any $m \geq 0$. Now we write

$$\ker \varphi^q \subset \ker(\beta^q_k \circ \varphi^q) = \ker(b^q_k \circ \Phi^q_k) = \ker b^q_k = \operatorname{Im} \alpha^q_k$$

where the third equality comes from bijectivity of Φ_k^q . Now we note that *R*-flatness of R' implies that

$$\operatorname{Im} \alpha_k^q = \operatorname{F}^{k+1} \operatorname{H}^{\operatorname{q}}(K^{\bullet}) \otimes_R R' \subset \left(\varpi^{k+1-n} \operatorname{H}^{\operatorname{q}}(K^{\bullet}) \right) \otimes_R R' = \varpi^{k+1-n} \left(\operatorname{H}^{\operatorname{q}}(K^{\bullet}) \otimes_R R' \right)$$

where the second inequality comes from Lemma A.4. That implies that ker φ^q is actually zero by separatedness of $\mathrm{H}^q(K^{\bullet}) \otimes_R R'$.

144

⁴⁵The upper long sequence is exact by flatness of the map $R \to R'$
Surjectivity of φ^q : Note that the hypothesis that $\mathrm{H}^q(L^{\bullet})$ and $\mathrm{H}^q(K^{\bullet})$ are almost finitely generated implies that the modules $\mathrm{H}^q(L^{\bullet} \text{ and } \mathrm{H}^q(K^{\bullet}) \otimes_R R'$ are both ϖ -adically complete due to Lemma 2.7.1 and Lemma 2.8.7. Thus the standard argument with successive approximations implies that it is sufficient to show that φ^q is surjective modulo ϖ . However, an easy diagram chase shows that $\beta^q_k(\mathrm{Im}\,\varphi^q) = \mathrm{Im}\,\beta^q_k$, then φ^q induces a surjective morphism

$$\mathrm{H}^{q}(K^{\bullet}) \otimes_{R} R' \to \mathrm{H}^{q}(L^{\bullet})/\mathrm{F}^{k+1}\mathrm{H}^{q}(L^{\bullet})$$

Now we use Lemma A.4 to say that for k = n, we have an inclusion $F^{k+1}H^q(L^{\bullet}) \subset \varpi H^q(L^{\bullet})$. Thus the discussion above implies that the φ^q induces a surjective morphism

$$\mathrm{H}^{q}(K^{\bullet}) \otimes_{R} R' \to \mathrm{H}^{q}(L^{\bullet})/\varpi \mathrm{H}^{q}(L^{\bullet})$$

Therefore, it is surjective by the argument above.

Appendix B. Achinger's Result in the Non-Noetherian Case

Recall that Piotr Achinger proved a remarkable result [Ach17, Proposition 6.6.1] that says that an affinoid rigid space $X = \text{Spa}(A, A^+)$ that admits an étale map to a closed unit disc \mathbf{D}_K^n also admits a *finite* étale map to \mathbf{D}_K^n provided that K is the fraction field of a complete DVR R with residue field of characteristic p. This result is an analytic analogue of a more classical result of Kedlaya ([Ked05] and [Ach17, Proposition 5.2.1]) that an affine k-scheme X = Spec A that admits an étale map to an affine space \mathbf{A}_k^n also admits a *finite* étale to \mathbf{A}_k^n provided that k has characteristic p.

We generalize Achinger's result in the non-noetherian setting. The proof essentially follows the ideas of [Ach17], we only need to be slightly more careful at some places due to non-noetherian issues. We also show its formal counterpart.

Lemma B.1. Let k be a field of characteristic p, and let A be a finite type k-algebra such that dim $A \leq d$ for some integer d. Suppose that $x_1, \ldots, x_d \in A$ some elements of A, and m is any integer $m \geq 0$. Then there exist elements $y_1, \ldots, y_d \in A$ such that the map $f: k[T_1, \ldots, T_n] \to A$, defined as $f(T_i) = x_i + y_i^{p^m}$ is finite.

Proof. We extend the set x_1, \ldots, x_d to some set of generators $x_1, \ldots, x_d, \ldots, x_n$ of A as a k-algebra. This defines a presentation $A = k[T_1, \ldots, T_d, \ldots, T_n]/I$ for some ideal $I \subset k[T_1, \ldots, T_r, \ldots, T_n]$. We prove the claim by induction on n - d.

The case of n-d=0 is trivial as then the map $f: k[T_1, \ldots, T_d] \to A$, defined by $f(T_i) = x_i$, is surjective. Therefore, it is finite.

Now we do the induction argument, so we suppose that $n - d \ge 1$. We consider the elements

$$x'_i = x_i - x_n^{p^{im'}}, \ i = 1, \dots, n-1$$

for some integer $m' \ge m$. Now the assumption $n \ge d+1$ and Krull's principal ideal theorem imply that we can choose some non-zero element $g \in I$, thus we have an expression

$$g(x_1' + x_n^{p^{m'}}, x_2' + x_n^{p^{2m'}}, \dots, x_{n-1}' + x_n^{p^{(n-1)m'}}, x_n) = 0$$

Now [Mum99, §1] implies that there is some large m' such that this expression is a polynomial in x_n with coefficients in $k[x'_1, \ldots, x'_{n-1}]$ and a non-zero leading term. We may and do assume that this leading term is 1. So x_n is integral over a subring of R generated by x'_1, \ldots, x'_{n-1} , we denote this ring by R'. Since $x_i = x'_i + x_n^{p^{im'}}$, we conclude that R is integral over R'. Moreover, R is finite over R' as it is finite type over k. Now we note that [Mat86, Theorem 9.3] implies

BOGDAN ZAVYALOV

that dim $R' \leq \dim R \leq d$, and R' is generated by x'_1, \ldots, x'_{n-1} as a k-algebra. So we can use the induction hypothesis to find some elements

$$y'_1,\ldots,y'_d\in R'$$

such that the morphism $f': k[T_1, \ldots, T_d] \to R'$, defined as $f'(T_i) = x'_i + (y'_i)^{p^m}$, is finite. Therefore, the composite morphism

$$f: k[T_1, \ldots, T_d] \to R$$

is also finite. We now observe that

$$f(T_i) = x'_i + (y'_i)^{p^m} = x_i + x_n^{p^{im'}} + (y'_i)^{p^m} = x_i + (x_n^{p^{im'-m}} + y'_i)^{p^m}$$

Therefore, the set $(y_i \coloneqq x_n^{p^{im'-m}} + y'_i)_{i=1,\dots,d}$ does the job.

Lemma B.2. Let \mathcal{O} be a complete valuation ring of rank-1 with the maximal ideal \mathfrak{m} and the residue field k. Suppose that $f: A \to B$ is a morphism of topologically finitely generated \mathcal{O}_{K^-} algebras. Then f is finite if and only if $f \otimes_{\mathcal{O}} k: A \otimes_{\mathcal{O}} k \to B \otimes_{\mathcal{O}} k$ is finite.

Proof. The "only if" part is clear, so we only need to deal with the "if" part. We recall that [Mat80, Lemma (28.P), p. 212] says that $A \to B$ is finite if and only if $A/\pi \to B/\pi$ is finite for some pseudo-uniformizer $\pi \in \mathcal{O}$. So we only need to show that finiteness of $A \otimes_{\mathcal{O}} k \to B \otimes_{\mathcal{O}} k$ implies that there is a pseudo-uniformizer $\pi \in \mathcal{O}$ such that $A/\pi \to B/\pi$ is finite. Then we note that the maximal ideal \mathfrak{m} is a filtered colimit of its finitely generated subideal $\{I_j\}_{j\in J}$. Moreover, the valuation property of the ring \mathcal{O} implies that this colimit is actually direct and that $I_j = (\pi_j)$ is principal for any $j \in J$. We also observe that each π_j is a pseudo-uniformizer since \mathcal{O} is of rank-1. Thus we see that

$$A \otimes_{\mathbb{O}} k \to B \otimes_{\mathbb{O}} k = \operatorname{colim}_{i \in J} (A/\pi_i \to B/\pi_i)$$

and $A/\pi_j \to B/\pi_j$ is a finite type morphism by the assumption that both A and B are topologically finitely generated. Then [Sta21, Tag 07RG] implies that there is $j \in J$ such that $A/\pi_j \to B/\pi_j$ is finite. Therefore, $A \to B$ is finite as well.

Before going to the proof Theorem B.4, we need to show a result on dimension on rigid spaces spaces that seem to be missing in the literature. It seems that there is no generally accepted definition of a dimesion of an adic spaces. We define the dimension as $\dim X = \sup_{x \in X} \dim \mathcal{O}_{X,x}$, this is consistent with the definition of dimension in [FK18, Definition II.10.1.1]. We denote by $X^{\text{cl}} \subset X$ the set of all classical points of X.

Lemma B.3. Let $f: X = \operatorname{Spa}(B, B^+) \to Y = \operatorname{Spa}(A, A^+)$ be an étale morphism of rigid spaces over a complete rank-1 field K, then dim $B \ge \dim A$. If Y is equidimensional, i.e. dim $\mathcal{O}_{Y,y} =$ dim Y for any classical point $y \in Y^{\text{cl}}$, then we have an equality dim $B = \dim A$. In particular, if $f: \operatorname{Spa}(A, A^+) \to \mathbf{D}_K^d$ is étale, then dim A = d.

Proof. We note that [FK18, Proposition II.10.1.9 and Corollary II.10.1.10] imply that

$$\dim X = \dim B = \sup_{x \in X^{cl}} (\dim \mathcal{O}_{X,x}), \text{ and } \dim Y = \dim A = \sup_{y \in Y^{cl}} (\dim \mathcal{O}_{Y,y}).$$

Since f is topologically finite type, it sends classical points to classical points. Therefore, [Hub96, Lemma 1.6.4, Corollary 1.7.4, Proposition 1.7.9] imply that the map $\mathcal{O}_{Y,f(x)} \to \mathcal{O}_{X,x}$ is finite étale for any $x \in X^{\text{cl}}$. Thus we see that

$$\dim B = \sup_{x \in X^{\mathrm{cl}}} (\dim \mathcal{O}_{X,x}) = \sup_{x \in X^{\mathrm{cl}}} (\dim \mathcal{O}_{Y,f(x)}) \le \dim Y$$

It is also clear that this inequality becomes an equality, if Y is equidimensional.

146

Finally, we claim that $\mathbf{D}_{K}^{d} = \operatorname{Spa}(K\langle T_{1}, \ldots, T_{d} \rangle, \mathcal{O}_{K}\langle T_{1}, \ldots, T_{d} \rangle) = \operatorname{Spa}(A, A^{+})$ is equidimensional. Pick any classical point $x \in (\mathbf{D}_{K}^{d})^{\text{cl}}$ and a corresponding maximal ideal $\mathfrak{m}_{x} \in K\langle T_{1}, \ldots, T_{d} \rangle$. Then we know that $A_{\mathfrak{m}_{x}}$ and $\mathcal{O}_{\mathbf{D}_{K}^{d},x}$ are noetherian by [FK18, Proposition 0.9.3.9, Theorem II.8.3.6], and $\widehat{\mathcal{O}_{\mathbf{D}_{K}^{d},x}} \simeq \widehat{A_{\mathfrak{m}_{x}}}$ by [FK18, Proposition II.8.3.1]. Therefore, we get

$$\dim \mathfrak{O}_{\mathbf{D}_K^d, x} = \dim \widehat{\mathfrak{O}_{\mathbf{D}_K^d, x}} = \dim \widehat{A_{\mathfrak{m}_x}} = \dim A_{\mathfrak{m}_x} = d$$

where the last equality comes from [FK18, Proposition 0.9.3.9].

For the rest of the section we fix a complete rank-1 valuation ring \mathcal{O} with the fraction field Kand the characteristic p residue field k. We refer to [Hub96, §1.9] for the construction of the adic generic fiber of a topologically finitely generated formal \mathcal{O} -scheme. The only thing we mention here is that it sends an affine formal scheme Spf A to the affinoid adic space $\text{Spa}(A \otimes_{\mathcal{O}} K, A^+)$, where A^+ is the integral closure of the image $\text{Im}(A \to A \otimes_{\mathcal{O}} K)$.

Theorem B.4. In the notation as above, let $g: \operatorname{Spf} A \to \widehat{\mathbf{A}}^d_{\mathbb{O}}$ be a morphism of flat, topologically finitely generated formal O-schemes such that the rigid generic fiber $g_K: \operatorname{Spa}(A \otimes_{\mathbb{O}} K, A^+) \to \mathbf{D}^d_K$ is étale. Then there is a *finite* morphism $f: \operatorname{Spf} A \to \widehat{\mathbf{A}}^d_{\mathbb{O}}$ that is étale on rigid generic fibers.

Proof. First of all, we note that Lemma B.3 says that dim $A \otimes_0 K = d$. Now [FK18, Theorem 9.2.10] says that there exists an finite injective morphism

$$\varphi \colon \mathcal{O}\langle T_1, \dots, T_d \rangle \to A$$

with the \mathcal{O}_K -flat cokernel. This implies that $K\langle T_1, \ldots, T'_d \rangle \to A \otimes_{\mathcal{O}} K$ is finite and injective. Therefore, [Mat86, Theorem 9.3] implies that

$$d = \dim A \otimes_{\mathcal{O}} K = \dim K \langle T_1, \dots, T'_d \rangle = d'$$

Thus we get that d = d'. Flatness of Coker φ says that the map

$$k[T_1,\ldots,T_d] \to A \otimes_{\mathfrak{O}} k$$

is also finite and injective. Then the similar argument shows that dim $A \otimes_{\mathcal{O}} k = d$. Now we finish the proof in two slightly different ways depending on characteristic of K.

Case 1, char K = p: We consider the morphism $g^{\#} : \mathcal{O}(T_1, \ldots, T_d) \to A$ induced by g. We define $x_i := g^{\#}(T_i)$ for $i = 1, \ldots, d$. Since dim $A \otimes_{\mathcal{O}} k = d$ we can apply Lemma B.1 for the residue classes $\overline{x_1}, \ldots, \overline{x_d}$ and m = 1 to get elements $\overline{y_1}, \ldots, \overline{y_d} \in A \otimes_{\mathcal{O}} k$ such that the map

$$\overline{f^{\#}}: k[T_1, \dots, T_d] \to A \otimes_{\mathbb{O}} k$$
, defined as $\overline{f^{\#}}(T_i) = \overline{x_i} + \overline{y_i}^p$ for $i = 1, \dots, d$

is finite. We lift $\overline{y_i}$ in an arbitrary way to elements $y_i \in A$, and define

$$f^{\#} \colon \mathcal{O}\langle T_1, \dots, T_d \rangle \to A$$

as $f^{\#}(T_i) = x_i + y_i^p$ for any i = 1, ..., d. This map is finite by Lemma B.2.

Now we note that $X := \text{Spa}(A \otimes_0 K, A^+)$ is smooth over K, so [BLR95, Proposition 2.6] says that étaleness of $f_K \colon X \to \mathbf{D}_K^d$ is equivalent to bijectivity of the map

$$f_K^*\Omega^1_{\mathbf{D}^d_K/K} \to \Omega^1_{X/K}$$

This easily follows from étaleness of g_K and the fact that $d(x_i + y_i^p) = d(x_i)$ in characteristic p.

BOGDAN ZAVYALOV

Case 2, char K = 0: We denote Spf A by \mathfrak{X} and its adic generic fiber $\text{Spa}(A \otimes_0 K, A^+)$ by X. Then we use [BLR95, Proposition 2.6] once again to see that the map

$$g_K^*\Omega^1_{\mathbf{D}^d_K/K} \to \Omega^1_{X/K}$$

is an isomorphism. Since $(\widehat{\Omega^1}_{\mathfrak{X}/\mathfrak{O}})_K \simeq \Omega^1_{X/K}$ and the same for $\widehat{\mathbf{A}}^d_{\mathfrak{O}}$ and \mathbf{D}^d_K , we conclude that the fundamental short exact sequence ([FK18, Proposition I.3.6.3, Proposition I.5.2.5 and Theorem I.5.2.6])

$$g^*\widehat{\Omega}^1_{\widehat{\mathbf{A}}^d_{\mathcal{O}}/\mathcal{O}} \to \widehat{\Omega}^1_{\mathfrak{X}/\mathcal{O}} \to \widehat{\Omega}^1_{\mathfrak{X}/\widehat{\mathbf{A}}^d_{\mathcal{O}}} \to 0$$

implies that $\left(\widehat{\Omega}^{1}_{\mathfrak{X}/\widehat{\mathbf{A}}^{d}_{\mathcal{O}}}\right)_{K} = 0$. More precisely, we know that

$$\widehat{\Omega}^{1}_{\mathfrak{X}/\widehat{\mathbf{A}}^{d}_{\mathfrak{O}}} \cong \left(\widehat{\Omega}^{1}_{A/\mathfrak{O}\langle T_{1},...,T_{d}\rangle}\right)^{\Delta}$$

for a finite A-module $\widehat{\Omega}^{1}_{A/0\langle T_{1},...,T_{d}\rangle}$ ([FK18, Corollary I.5.1.11]). We denote this module by $\widehat{\Omega}^{1}_{g}$ for the rest of the proof, and recall that the condition $\left(\widehat{\Omega}^{1}_{\mathfrak{X}/\widehat{\mathbf{A}}^{d}_{0}}\right)_{K} = 0$ is equivalent to $\widehat{\Omega}^{1}_{g} \otimes_{0} K = 0$. Using finiteness of $\widehat{\Omega}^{1}_{g}$ and adhesiveness of A, we conclude that there is an integer k such that

$$p^k \widehat{\Omega}_g^1 = 0$$

as p is a pseudo-uniformizer in O. Now, similarly to the case of $\operatorname{char} K = p$, we consider the morphism

$$g^{\#} \colon \mathcal{O}\langle T_1, \ldots, T_d \rangle \to A$$

and define $x_i := g^{\#}(T_i)$ for i = 1, ..., d. Again, using that dim $A \otimes_{\mathbb{O}} k = d$ we can apply Lemma B.1 for the residue classes $\overline{x_1}, \ldots, \overline{x_d}$ and m = k + 1 to get elements $\overline{y_1}, \ldots, \overline{y_d} \in A \otimes_{\mathbb{O}} k$ such that the map

$$\overline{f^{\#}}: k[T_1, \dots, T_d] \to A \otimes_{\mathbb{O}} k$$
, defined as $\overline{f^{\#}}(T_i) = \overline{x_i} + \overline{y_i}^{p^{k+1}}$ for $i = 1, \dots, d$

is finite. We lift $\overline{y_i}$ to some elements $y_i \in A$ and define

$$f^{\#} \colon \mathfrak{O}\langle T_1, \dots, T_d \rangle \to A$$

by $f^{\#}(T_i) = x_i + y_i^{p^{k+1}}$. The map $f^{\#}$ is finite by Lemma B.2.

We are only left to show that the induced map

$$f: X \to \widehat{\mathbf{A}}^d_{\mathcal{O}}$$

is étale on adic generic fibers. We claim that $p^k(\widehat{\Omega}_f^1) = 0$. Indeed, we use [FK18, Proposition I.5.1.10] to trivialize $\widehat{\Omega}^1_{\mathcal{O}\langle T_1,...,T_d\rangle/\mathcal{O}} \simeq \bigoplus_{i=1}^d dT_i$, so we have the fundamental exact sequence

$$\bigoplus_{i=1}^{d} AdT_i \xrightarrow{dT_i \mapsto d(x_i + y_i^{p^{k+1}})} \widehat{\Omega}^1_{A/\mathcal{O}} \to \widehat{\Omega}^1_f \to \mathcal{O}_f$$

As $d(y_i^{p^{k+1}})$ is divisible by p^{k+1} . Therefore, we see that modulo p^{k+1} this sequence is equal to

$$\bigoplus_{i=1}^{d} A/p^{k+1} dT_i \xrightarrow{dT_i \to d(x_i)} \widehat{\Omega}^1_{A/\mathcal{O}}/p^{k+1} \to \widehat{\Omega}^1_f/p^{k+1} \to 0$$

Thus we see that $\widehat{\Omega}_f^1/p^{k+1} \simeq \widehat{\Omega}_g^1/p^{k+1}$. In particular,

$$\left(p^k\widehat{\Omega}_f^1\right)/p\left(p^k\widehat{\Omega}_f^1\right) = \left(p^k\widehat{\Omega}_g^1\right)/p\left(p^k\widehat{\Omega}_g^1\right) = 0$$

by the choice of k. Therefore, $p^k \widehat{\Omega}_f^1 = 0$ by [Mat80, Lemma (28.P), p. 212]. By passing to the adic generic fiber we get that $f_K \colon X \to \mathbf{D}_K^d$ such that the map

$$d(f_K) \colon f_K^* \Omega^1_{\mathbf{D}_K^d/K} \to \Omega^1_{X/K}$$

is surjective. However, we recall that X and \mathbf{D}_K^d are both smooth rigid spaces of (pure) dimension d. Thus $d_{f_K^*}$ is a surjective map of vector bundles of the same dimension d, so it must be an isomorphism. Finally, [BLR95, Proposition 2.6] implies that f_K is étale.

Corollary B.5. Let K be a complete rank-1 valuation field with a valuation ring \mathcal{O}_K , and the residue field k of characteristic p. Suppose that $g: X = \text{Spa}(A, A^+) \to \mathbf{D}_K^d$ is an étale morphism of rigid affinoid K-spaces. Then there exists a *finite* étale morphism $f: X \to \mathbf{D}_K^d$.

Proof. First of all, we note that [Hub94, Lemma 4.4] implies that $A^+ = A^\circ$. So the map g corresponds to the map

$$g^{\#} \colon (K\langle T_1, \dots, T_d \rangle, \mathcal{O}_K\langle T_1, \dots, T_d \rangle) \to (A, A^{\circ})$$

of Tate-Huber pairs. We note that it suffices to find a topologically finitely generated ring of definition $A_0 \subset A$ such that the map $\mathcal{O}_K \langle T_1, \ldots, T_d \rangle \to A^\circ$ factors through A_0 . Then Theorem B.4 will imply the corollary.

We choose some surjection $\varphi \colon K\langle X_1, \ldots, X_n \rangle \twoheadrightarrow A$ and consider a ring

$$A_0' \coloneqq \varphi(\mathfrak{O}_K \langle X_1, \dots, X_n \rangle)$$

This ring is open by the Banach Open Mapping Theorem ([Hub94, Lemma 2.4 (i)]). It is also bounded as any map of Tate rings is adic, so it preserves boundedness. Therefore, A'_0 is a ring of definition in A.

Now we use the universal property of Tate algebras ([Hub94, Lemma 3.3]) to get the unique K-linear continuous homomorphism

 $\psi \colon K\langle T_1, \ldots, T_d, X_1, \ldots, X_n \rangle \to A$

such that $\psi(T_i) = g^{\#}(T_i)$ and $\psi(X_j) = \varphi(X_j)$. Then a similar argument implies that

$$A_0 \coloneqq \psi(\mathcal{O}_K \langle T_1, \dots, T_d, X_1, \dots, X_n \rangle)$$

is a topologically finitely generated ring of definition in A such that the map $g^{+,\#} \colon \mathcal{O}_K \langle T_1, \ldots, T_d \to A^\circ$ factors through A_0 . We note that A_0 is \mathcal{O}_K flat as it is torsionfree. Therefore, we can apply Theorem B.4 to the map $\operatorname{Spf} A_0 \to \widehat{\mathbf{A}}^d_{\mathcal{O}_K}$ to construct a finite K-étale map $f \colon \operatorname{Spf} A_0 \to \widehat{\mathbf{A}}^d_{\mathcal{O}_K}$. Then the adic generic fiber $f_K \colon \operatorname{Spa}(A, A^\circ) \to \mathbf{D}^{d}_K^{46}$ is the desired finite étale map.

Appendix C. Locally Constant and Lisse Sheaves on the Pro-Étale Topology

We give two definitions of locally constant $\mathbf{Z}/p^n\mathbf{Z}$ -sheaves that eventually turn out to coincide, but this requires a proof. This is somewhat implicit in [Sch13, Proposition 8.3], but we prefer to give a precise proof. Then we discuss the notion of lisse $\hat{\mathbf{Z}}_p$ -sheaves on the pro-étale site. In what follows X is a rigid-analytic variety over $\operatorname{Spa}(C, \mathcal{O}_C)$ or, more generally, any locally strongly noetherian adic space over $\operatorname{Spa}(\mathbf{Q}_p, \mathbf{Z}_p)$.

⁴⁶Here we implicitly use [Hub94, Lemma 4.4] to say that $A_0^+ = A^\circ$.

BOGDAN ZAVYALOV

Definition C.1. A sheaf of $\mathbf{Z}/p^n \mathbf{Z}$ -modules \mathbf{L} on $X_{\text{\acute{e}t}}$ is called *locally constant* if, locally on $X_{\text{\acute{e}t}}$, it is isomorphic to the constant sheaf \underline{M} associated with a finitely generated $\mathbf{Z}/p^n \mathbf{Z}$ -module M. We denote this category by $\text{LocSys}_{\mathbf{Z}/p^n \mathbf{Z}, X_{\text{\acute{e}t}}}$.

We say that a sheaf of \mathbf{F}_p -modules \mathbf{L} on $X_{\text{\acute{e}t}}$ is a *local system*, if it is locally constant sheaf of \mathbf{F}_p -modules.

A sheaf of $\mathbb{Z}/p^n\mathbb{Z}$ -modules \mathbb{L} on $X_{\text{pro\acute{e}t}}$ is called *locally constant* if, locally on $X_{\text{pro\acute{e}t}}$, it is isomorphic to the constant sheaf \underline{M} associated with a finitely generated $\mathbb{Z}/p^n\mathbb{Z}$ -module M. We denote this category by $\text{LocSys}_{\mathbb{Z}/p^n\mathbb{Z},X_{\text{pro\acute{e}t}}}$.

Lemma C.2. Let \mathcal{F} be an abelian sheaf on $X_{\text{pro\acute{e}t}}$ such that $\mathcal{F}(U) = \operatorname{colim}_i \mathcal{F}(U_i)$ for any quasicompact and quasi-separated $U = \lim_i U_i \in X_{\text{pro\acute{e}t}}$. Then the natural adjunction map $\mathcal{F} \to \lambda^{-1}\lambda_*\mathcal{F}$ is an isomorphism.

Proof. As quasi-compact and quasi-separated objects form a basis for the pro-étale topology, it is sufficient to check that the adjunction map is an isomorphism for any qcqs $U = \lim_{i \to \infty} U_i \in X_{\text{proét}}$. Now we see that [Sch13, Lemma 3.16] guarantees that

$$\lambda^{-1}\lambda_*\mathcal{F}(U) = \operatorname{colim}_i \lambda_*\mathcal{F}(U_i) = \operatorname{colim}_i \mathcal{F}(U_i)$$

The same formula holds for $\mathcal{F}(U)$ by the assumption. That the map

$$\mathfrak{F}(U) \to (\lambda^{-1}\lambda_*\mathfrak{F})(U)$$

is an isomorphism for any qcqs $U \in X_{\text{pro\acute{e}t}}$.

Lemma C.3. Let $\mathbf{L} \in \operatorname{LocSys}_{\mathbf{Z}/p^n \mathbf{Z}, X_{\operatorname{pro\acute{e}t}}}$, then the adjunction $\mathbf{L} \to \lambda^{-1} \lambda_* \mathbf{L}$ is an isomorphism.

Proof. The question is local on X, so we may and do assume that X is quasi-compact and quasiseparated. Now we note that it is sufficient to show that the natural map $\operatorname{colim}_i \mathbf{L}(U_i) \to \mathbf{L}(U)$ is an isomorphism for any qcqs $U = \lim_i U_i^* \in X_{\operatorname{pro\acute{e}t}}$. Local constancy of **L** implies that there is a pro-étale covering $V \to X$ such that such $\mathbf{L}|_V$ is constant. Thus we see that for any quasi-compact, quasi-separated V' in the localized pro-étale site $X_{\operatorname{pro\acute{e}t}}/V$ we get an isomorphism $\operatorname{colim}_i \mathbf{L}(V_i') \to$ $\mathbf{L}(V)$ whenever $V' = \lim_i V_i'$ for a filtered system $V_i' \in X_{\operatorname{pro\acute{e}t}}/V$. Since quasi-compact, quasiseparated objects form a basis on $X_{\operatorname{pro\acute{e}t}}$, we can assume that V is qcqs.

Now we pick some quasi-compact and quasi-separated $U = \lim_{i \in I} U_i \in X_{\text{pro\acute{e}t}}$. And choose a presentation $V = \lim_{j \in J} V_j \in X_{\text{pro\acute{e}t}}$. The fiber product $W \coloneqq U \times V = \lim_{I \times J} U_i \times V_j = \lim_{I \to J} U_i \times V$ is quasi-compact and quasi-separated as we assume that so is X. Now we consider the commutative diagram:

We claim that the middle and the right vertical arrows are isomorphisms by the assumption on V. Since filtered colimits are exact, we conclude that both lines of the diagram C are exact. Thus the left vertical arrow must be also an isomorphism.

Theorem C.4. The functor $\lambda^{-1} \colon \mathcal{A}b(X_{\acute{e}t}) \to \mathcal{A}b(X_{\mathrm{pro\acute{e}t}})$ defines an equivalence of categories $\lambda^{-1} \colon \mathrm{LocSys}_{\mathbf{Z}/p^{n}\mathbf{Z}, X_{\acute{e}t}} \to \mathrm{LocSys}_{\mathbf{Z}/p^{n}\mathbf{Z}, X_{\mathrm{pro\acute{e}t}}}$ with the inverse given by λ_{*} .

Proof. We firstly note that the functor λ^{-1} clearly sends locally constant $\mathbf{Z}/p^{n}\mathbf{Z}$ -sheaves on $X_{\text{ét}}$ to locally constant $\mathbf{Z}/p^{n}\mathbf{Z}$ -sheaves on $X_{\text{pro\acute{e}t}}$. Moreover, [Sch13, Corollary 3.17] reads that the adjunction $\mathcal{F} \to \lambda_* \lambda^{-1} \mathcal{F}$ is an isomorphism for any $\mathcal{F} \in \mathcal{A}b(X_{\text{\acute{e}t}})$. Thus the functor λ^{-1} : LocSys $_{\mathbf{Z}/p^{n}\mathbf{Z},X_{\text{\acute{e}t}}} \to \text{LocSys}_{\mathbf{Z}/p^{n}\mathbf{Z},X_{\text{pro\acute{e}t}}}$ is fully faithful. Moreover, Lemma C.3 implies that the adjunction $\mathbf{L} \to \lambda^{-1}\lambda_*\mathbf{L}$ is an isomorphism for any $\mathbf{L} \in \text{LocSys}_{\mathbf{Z}/p^{n}\mathbf{Z},X_{\text{pro\acute{e}t}}}$. Thus the only thing we are left to show is that λ_* preserves local constancy of $\mathbf{Z}/p^{n}\mathbf{Z}$ -sheaves.

The problem is local on X, so we can assume that there is a quasi-compact and quasi-separated pro-étale covering $U = \lim_{I} U_i^{"} \to X$ such that $\phi \colon \underline{M}_U \to \mathbf{L}|_U$ for some finitely generated $\mathbf{Z}/p^n\mathbf{Z}$ module M. The proof of Lemma C.3 shows that the natural map $\operatorname{colim}_I \mathbf{L}(U_i) \to \mathbf{L}(U)$ is an isomorphism. Thus using the fact that any finitely generated $\mathbf{Z}/p^n\mathbf{Z}$ -module is finitely presented, we conclude there is some finite level $i \in I$ such that ϕ is defined U_i . More precisely, there is $i \in I$ and a morphism $\phi_i \colon \underline{M}_{U_i} \to \mathbf{L}|_{U_i}$ such that its pullback on U coincides with ϕ . We note that for any $j \ge i$, we have the morphism $\phi_j \colon \underline{M}_{U_j} \to \mathbf{L}|_{U_j}$ that is defined as the pullback of ϕ_i .

We claim that, possible after enlarging i, we can achieve ϕ_i to be an isomorphism. The fact that ϕ is surjective means that there are finitely many section $e_j \in \mathbf{L}(U)$ that globally generated \mathbf{L} as $\mathbf{Z}/p^n\mathbf{Z}$ -module. These sections are defined over some finite level. Thus after enlarging i, we can achieve that ϕ_i contains global sections of $\mathbf{L}(U_i)$ that generate $\mathbf{L}|_U$. In particular, they generate $\mathbf{L}|_{U_i}$ as surjectivity of a map of pro-étale sheaves can be checked pro-étale locally.

Now we want to make ϕ_i injective for large *i*. As the category $\operatorname{LocSys}_{\mathbf{Z}/p^n\mathbf{Z},X_{\operatorname{pro\acute{e}t}}}$ is clearly abelian, we conclude that ker ϕ_i is a locally constant $\mathbf{Z}/p^n\mathbf{Z}$ -sheaf. Moreover, we see that ker $\phi_i|_U \cong$ by the construction. Thus repeating the same argument for $\mathbf{L} = \ker \phi_i$, we get that there is a surjection $\underline{0}_{U_i} \to \ker \phi_i$ for some possibly large $i \in I$. This proves that ker ϕ_i becomes zero for large *i*.

All in all, this shows that there is some i such that $\mathbf{L}|_{U_i}$ is isomorphic to $\underline{M}|_{U_i}$ for some finitely generated $\mathbf{Z}/p^n\mathbf{Z}$ -module M. As $U_i \to X$ is étale by the construction, we conclude that $\lambda_* \mathcal{G} \in \mathrm{LocSys}_{\mathbf{Z}/p^n\mathbf{Z},X_{\mathrm{\acute{e}t}}}$. This finishes the proof.

Remark C.5. In the paper we freely identify pro-étale locally constant sheaves and étale locally constant sheaves. We usually slightly abuse notations and do not distinguish $\lambda^{-1}\mathbf{L}$ and \mathbf{L} for $\mathbf{L} \in \mathrm{LocSys}_{\mathbf{Z}/p^n\mathbf{Z},X_{\mathrm{\acute{e}t}}}$. This does not cause any real issues by Theorem C.4.

We also discuss the notion of lisse $\widehat{\mathbf{Z}}_p$ -sheaves in $X_{\text{proét}}$.

Definition C.6. The pro-étale sheaf $\widehat{\mathbf{Z}}_p$ is defined as $\widehat{\mathbf{Z}}_p \coloneqq \lim_n \mathbf{Z}/p^n \mathbf{Z}$ in $\mathbf{Shv}(X_{\text{proét}})$.

Definition C.7. A lisse $\widehat{\mathbf{Z}}_p$ -sheaf on $X_{\text{pro\acute{e}t}}$ is a sheaf \mathbf{L} of $\widehat{\mathbf{Z}}_p$ -modules on $X_{\text{pro\acute{e}t}}$, such that locally in $X_{\text{pro\acute{e}t}}$, \mathbf{L} is isomorphic to $\widehat{\mathbf{Z}}_p \otimes_{\mathbf{Z}_p} M$, where M is a finitely generated \mathbf{Z}_p -module.

We denote the category of lisse $\widehat{\mathbf{Z}}_p$ -sheaves by LocSys $_{\mathbf{Z}_p, X_{\text{pro\acute{t}}}}$.

Definition C.8. We say that a lisse $\widehat{\mathbf{Z}}_p$ -sheaf \mathbf{L} comes from $\operatorname{LocSys}_{\mathbf{Z}/p^n\mathbf{Z},X_{\acute{e}t}}$ if it is isomorphic to $\lambda^{-1}\mathbf{M}$ for some $\mathbf{M} \in \operatorname{LocSys}_{\mathbf{Z}/p^n\mathbf{Z},X_{\acute{e}t}}$.

Lemma C.9. Let $\mathbf{L} \in \text{LocSys}_{\mathbf{Z}_p, X_{\text{pro\acute{e}t}}}$, then the sheaves $\mathbf{L}/p^n \mathbf{L}$ come from $\text{LocSys}_{\mathbf{Z}/p^n \mathbf{Z}, X_{\acute{e}t}}$ for $n \ge 0$, and $p^n \mathbf{L}/p^{n+1} \mathbf{L}$ come from $\text{LocSys}_{\mathbf{Z}/p\mathbf{Z}, X_{\acute{e}t}}$ for $n \ge 0$.

Proof. It is clear that each $\mathbf{L}/p^{n}\mathbf{L}$ lies in $\operatorname{LocSys}_{\mathbf{Z}/p^{n}\mathbf{Z},X_{\operatorname{pro\acute{e}t}}}$. Theorem C.4 guarantees that its comes from $\operatorname{LocSys}_{\mathbf{Z}/p^{n}\mathbf{Z},X_{\operatorname{\acute{e}t}}}$. The same proof works for $p^{n}\mathbf{L}/p^{n+1}\mathbf{L}$.

Corollary C.10. Let $\mathbf{L} \in \operatorname{LocSys}_{\mathbf{Z}_p, X_{\operatorname{pro\acute{e}t}}}$, and let $\operatorname{Fil}^{\bullet}(\mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathbb{O}}_X^+)$ be the filtration defined by $\operatorname{Fil}^n = p^n(\mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathbb{O}}_X^+)$ for $n \ge 0$. Then the associated pieces $\operatorname{gr}^n(\mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathbb{O}}_X^+)$ has a form $\mathbf{L}_n \otimes_{\mathbf{F}_p} \mathbb{O}_X^+/p$ for some (étale) \mathbf{F}_p -local system \mathbf{L}_n . The quotients $(\mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathbb{O}}_X^+)/\operatorname{Fil}^n$ has the form $\mathbf{M}_n \otimes_{\mathbf{Z}/p^n \mathbf{Z}} \mathbb{O}_X^+/p^n$ for $\mathbf{M}_n \in \operatorname{LocSys}_{\mathbf{Z}/p^n \mathbf{Z}, X_{\operatorname{\acute{e}t}}}$.

Proof. First of all, we note that the sheaf $\widehat{\mathcal{O}}_X^+$ is flat over \mathbf{Z}_p by [Sch13, Lemma 4.2]. So $\widehat{\mathcal{O}}_X^+$ and \mathbf{L} are $\widehat{\mathbf{Z}}_p$ -tor independent for any lisse $\widehat{\mathbf{Z}}_p$ -sheaf \mathbf{L} . This implies that

$$gr^{n}(\mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathcal{O}}_{X}^{+}) \simeq p^{n}(\mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathcal{O}}_{X}^{+})/p^{n+1}(\mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathcal{O}}_{X}^{+})$$
$$\simeq (p^{n}\mathbf{L}/p^{n+1}\mathbf{L}) \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathcal{O}}_{X}^{+}$$
$$\simeq (p^{n}\mathbf{L}/p^{n+1}\mathbf{L}) \otimes_{\mathbf{F}_{p}} \mathcal{O}_{X}^{+}/p$$

So the result follows from Lemma C.9. The same argument works for $(\mathbf{L} \otimes_{\widehat{\mathbf{Z}}_n} \widehat{\mathbb{O}}_X^+)/\mathrm{Fil}^n$.

Now the last thing that we discuss is that $\mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathcal{O}}_X^+$ is always derived *p*-adically complete in $\mathbf{D}(X_{\text{pro\acute{e}t}}) \coloneqq \mathbf{D}(\mathcal{A}b(X_{\text{pro\acute{e}t}}))$. We refer to [Sta21, Tag 0995] for a self-contained discussion of this notion. We only briefly mention that a complex $K \in \mathbf{D}(X_{\text{pro\acute{e}t}})$ is *p*-adically derived complete if the morphism $K \to \mathbf{R} \lim_n [K/p^n K]$ is an isomorphism, where $[K/p^n K]$ stands for the cone of the p^n -multiplication map. This definition is equivalent to the definition given on Stacks Project by [Sta21, Tag 0A0E].

Lemma C.11. Let $K \in \mathbf{D}(X_{\text{pro\acute{e}t}})$ be a complex such that $\mathbf{R}\Gamma(U, K)$ is derived *p*-adically complete for any affinoid perfectoid U. Then K is derived *p*-adically complete.

Proof. Consider the derived completion $K \to \hat{K}$ and the associated distinguished triangle:

 $K \to \widehat{K} \to Q.$

We show that $Q \simeq 0$, it suffices to show that $\mathbf{R}\Gamma(U, Q) \simeq 0$ for any affinoid perfected U as affinoid perfected s form a basis in $X_{\text{pro\acute{e}t}}$. Now we use [Sta21, Tag 0BLX] to conclude that

$$\mathbf{R}\Gamma(U,\widehat{K})\simeq \mathbf{R}\widehat{\Gamma(U,K)},$$

so we get the distinguished triangle

$$\mathbf{R}\Gamma(U,K) \to \mathbf{R}\Gamma(U,K) \to \mathbf{R}\Gamma(U,Q).$$

Since $\mathbf{R}\Gamma(U,K)$ is derived *p*-adically complete by the assumption, so we see that the morphism

$$\mathbf{R}\Gamma(U,K) \to \mathbf{R}\Gamma(U,K)$$

is an isomorphism. Therefore, we conclude that $\mathbf{R}\Gamma(U,Q) \simeq 0$. This finishes the proof.

Lemma C.12. Let $\mathbf{L} \in \operatorname{LocSys}_{\mathbf{Z}_p, X_{\operatorname{pro\acute{e}t}}}$. Then the morphism

$$\mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathcal{O}}_X^+ \to \mathbf{R} \lim_n \left((\mathbf{L}/p^n \mathbf{L}) \otimes_{\mathbf{Z}/p^n \mathbf{Z}} \left(\mathcal{O}_X^+/p^n \right) \right)$$

is an isomorphism. In particular, $\mathbf{L} \otimes_{\widehat{\mathbf{Z}}_{p}} \widehat{\mathcal{O}}_{X}^{+}$ is derived *p*-adically complete.

Proof. Step 1. We show the claim for the sheaf $\widehat{\mathcal{O}}_X^+$: We note that [Sch13, Lemma 4.10] implies that $\mathbf{R}\Gamma(U,\widehat{\mathcal{O}}_X^+)$ is derived *p*-adically complete for any affinoid $U \in X_{\text{pro\acute{e}t}}$ since almost zero modules

REFERENCES

are clearly derived *p*-adically complete. Therefore, Lemma C.11 guarantees that $\widehat{\mathcal{O}}_X^+$ is derived *p*-adically complete.

Now we note that the sheaf $\widehat{\mathcal{O}}_X^+$ is flat over \mathbf{Z}_p by [Sch13, Lemma 4.2]. Thus, we see that $\mathcal{O}_X^+/p^n \simeq [\widehat{\mathcal{O}}_X^+/p^n]$. So derived completeness of $\widehat{\mathcal{O}}_X^+$ tells us that

$$\widehat{\mathcal{O}}_X^+ \simeq \mathbf{R} \lim_n [\widehat{\mathcal{O}}_X^+ / p^n] \simeq \mathbf{R} \lim_n \mathcal{O}_X^+ / p^n$$

Step 2. The general case: We can check that the map

$$\mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathcal{O}}_X^+ \to \mathbf{R} \lim_n \left((\mathbf{L}/p^n \mathbf{L}) \otimes_{\mathbf{Z}/p^n \mathbf{Z}} \left(\mathcal{O}_X^+/p^n \right) \right)$$

is an isomorphism pro-étale locally on X. So we may and do assume that **L** is given by $M \otimes_{\mathbf{Z}_p} \widehat{\mathbf{Z}}_p$ for some finitely generated \mathbf{Z}_p -module M. By the classification of finitely generated \mathbf{Z}_p -modules, we can write M as $M' \times \mathbf{Z}_p^r$ for some finitely generated torsion module M' and some non-negative integer r. Since derived limits commute with derived products, and finite derived products coincide with (naive) finite direct sums, we see that it suffices to show the claim separately for finite free modules and torsion modules. The case of finite free modules follows from Step 1. So we suppose that M is a finite torsion module. In particular, there is n such that $p^n M = 0$. This implies that the natural morphisms

$$\mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathcal{O}}_X^+ \to \mathbf{L}/p^m \mathbf{L} \otimes_{\mathbf{Z}/p^m \mathbf{Z}} \mathcal{O}_X^+/p^m \to \mathbf{L}/p^n \mathbf{L} \otimes_{\mathbf{Z}/p^n \mathbf{Z}} \mathcal{O}_X^+/p^n$$

are isomorphisms for $m \geq n$ and $\mathbf{L} = M \otimes_{\mathbf{Z}_p} \widehat{\mathbf{Z}}_p$. Thus the map

$$\mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathbb{O}}_X^+ \to \mathbf{R} \lim_n \left((\mathbf{L}/p^n \mathbf{L}) \otimes_{\mathbf{Z}/p^n \mathbf{Z}} \left(\mathbb{O}_X^+/p^n \right) \right)$$

is trivially an isomorphism in this case. Finally, we note that this implies that $\mathbf{L} \otimes_{\widehat{\mathbf{Z}}_p} \widehat{\mathcal{O}}_X^+$ is *p*-adically derived complete as derived complete objects are closed under derived limits.

References

[Ach17]	P. Achinger. "Wild ramification and $K(\pi, 1)$ spaces". In: Invent. Math. 210.2 (2017),
	pp. 453–499.
[AM69]	M. F. Atiyah and I. G. Macdonald. Introduction to commutative algebra. Addison-
	Wesley Publishing Co., Reading, MassLondon-Don Mills, Ont., 1969, pp. ix+128.
[BGR84]	S. Bosch, U. Güntzer, and R. Remmert. Non-Archimedean analysis. A systematic
	approach to rigid analytic geometry. Springer-Verlag, Berlin, 1984, pp. xii+436.
[Bha]	B. Bhatt. Lecture notes for a class on perfectoid spaces. http://www-personal.umich.edu/~bhatt
[Bha18]	B. Bhatt. "Specializing varieties and their cohomology from characteristic 0 to charac-
	teristic p". In: Algebraic geometry: Salt Lake City 2015. Vol. 97. Proc. Sympos. Pure
	Math. Amer. Math. Soc., Providence, RI, 2018, pp. 43–88.
[BL93]	S. Bosch and W. Lutkebohmert. "Formal and rigid geometry. I. Rigid spaces." In:
	Mathematische Annalen 295 (1993), pp. 291–318.
[BLR95]	S. Bosch, W. Lütkebohmert, and M. Raynaud. "Formal and rigid geometry. III. The
	relative maximum principle". In: Math. Ann. 302.1 (1995), pp. 1–29.
[BMS18]	B. Bhatt, M. Morrow, and P. Scholze. "Integral <i>p</i> -adic Hodge theory". In: <i>Publ. Math.</i>
	Inst. Hautes Études Sci. 128 (2018), pp. 219–397.
[BMS19]	B. Bhatt, M. Mathew, and P. Scholze. "Topological Hochschild homology and integral
	p-adic Hodge theory". In: Publ. Math. Inst. Hautes Études Sci. 129 (2019), pp. 199-
	310.

b/1

REFERENCES

[BNP17]	D. Ben-Zwi, D. Nadler, and A. Preygel. "Integral Transform For Coherent Sheaves"
	In: J. Eur. Math. Soc. (2017).

- [Bos14] S. Bosch. Lectures on formal and rigid geometry. Vol. 2105. Lecture Notes in Mathematics. Springer, Cham, 2014, pp. viii+254.
- [Bou98] N. Bourbaki. *Commutative algebra. Chapters 1–7.* Translated from the French, Reprint of the 1989 English translation. Springer-Verlag, Berlin, 1998, pp. xxiv+625.
- [BS19] B. Bhatt and P. Scholze. "Prisms and Prismatic Cohomology". In: (2019), pp. 1–105.
 URL: https://arxiv.org/abs/1905.08229.
- [Con] B. Conrad. Cohomological descent. http://math.stanford.edu/~conrad/papers/hypercover.pdf.
- [FGK11] K. Fujiwara, O. Gabber, and F. Kato. "On Hausdorff completions of commutative rings in rigid geometry". In: Journal Of Algebra 332 (2011), pp. 293–321.
- [FK18] K. Fujiwara and F. Kato. Foundations of rigid geometry. I. EMS Monographs in Mathematics. European Mathematical Society (EMS), Zürich, 2018, pp. xxxiv+829.
- [EGA I] A. Grothendieck and J. A. Dieudonné. Eléments de géométrie algébrique. I. Vol. 166. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 1971, pp. ix+466.
- [GR03] O. Gabber and L. Ramero. *Almost Ring Theory*. Lecture Notes in Mathematics. Springer, 2003, pp. viii+318.
- [SGA 1] A. Grothendieck. Revêtements étales et groupe fondamental. Vol. 1960/61. Séminaire de Géométrie Algébrique. Institut des Hautes Études Scientifiques, Paris, 1963, iv+143 pp.
- [EGA IV₃] A. Grothendieck. "Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. III". In: Inst. Hautes Études Sci. Publ. Math. 28 (1966), p. 255.
- [Guo19] H. Guo. Hodge-Tate decomposition for non-smooth spaces. https://arxiv.org/abs/1909.09917. 2019.
- [GZ67] P. Gabriel and M. Zisman. Calculus of fractions and homotopy theory. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 35. Springer-Verlag New York, Inc., New York, 1967, pp. x+168.
- [Hal18] J. Hall. GAGA Theorems. https://arxiv.org/abs/1804.01976. 2018.
- [Hub94] R. Huber. "A generalization of formal schemes and rigid analytic varieties". In: Math. Z. 217.4 (1994), pp. 513–551.
- [Hub96] R. Huber. Étale cohomology of rigid analytic varieties and adic spaces. Friedr. Vieweg & Sohn, Braunschweig, 1996, pp. x+450.
- [Ked05] K. S. Kedlaya. "More étale covers of affine spaces in positive characteristic". In: J. Algebraic Geom. 14.1 (2005), pp. 187–192.
- [Lim19] D. B. Lim. Grothendieck's Existence Theorem For Relatively Perfect Complexes On Algebraic Stacks. https://arxiv.org/abs/1907.05025. 2019.
- [Lur20] J. Lurie. Spectral Algebraic Geometry. https://www.math.ias.edu/~lurie/papers/SAG-rootfile. 2020.
- [Lüt90] Werner Lütkebohmert. "Formal-algebraic and rigid-analytic geometry". In: Math. Ann. 286.1-3 (1990), pp. 341–371.
- [Mat80] H. Matsumura. *Commutative algebra*. Second. Vol. 56. Mathematics Lecture Note Series. Benjamin/Cummings Publishing Co., Inc., Reading, Mass., 1980, pp. xv+313.
- [Mat86] H. Matsumura. *Commutative ring theory.* Vol. 8. Cambridge Studies in Advanced Mathematics. Translated from the Japanese by M. Reid. Cambridge University Press, Cambridge, 1986.

REFERENCES

- [Mum99] D. Mumford. The red book of varieties and schemes. expanded. Vol. 1358. Lecture Notes in Mathematics. Includes the Michigan lectures (1974) on curves and their Jacobians, With contributions by Enrico Arbarello. Springer-Verlag, Berlin, 1999, pp. x+306.
- [Ray70] M. Raynaud. Anneaux locaux henséliens. Vol. 169. Lecture Notes in Mathematics. Spinger-Verlag, 1970.
- [Sch12] P. Scholze. "Perfectoid spaces". In: Publ. Math. Inst. Hautes Études Sci. 116 (2012), pp. 245–313. ISSN: 0073-8301.
- [Sch13] P. Scholze. "*p*-adic Hodge theory for rigid-analytic varieties". In: 1 (2013), e1–77.
- [Sch16] P. Scholze. "*p*-adic Hodge theory for rigid-analytic varieties—corrigendum". In: *Forum* Math. Pi 4 (2016), e6, 4.
- [Sch17] P. Scholze. Étale Cohomology of diamonds. https://arxiv.org/abs/1709.07343. 2017.
- [Sch92] C. Scheiderer. "Quasi-augmented simplicial spaces, with an application to cohomological dimension". In: J. Pure Appl. Algebra 81.3 (1992), pp. 293–311.
- [Sta21] The Stacks project authors. The Stacks project. https://stacks.math.columbia.edu. 2021.
- [Tem00] M. Temkin. "On local properties of non-Archimedean analytic spaces". In: *Math. Ann* 318(3) (2000), pp. 585–607.
- [Tem12] M. Temkin. "Functorial desingularization of quasi-excellent schemes in characteristic zero: the nonembedded case". In: *Duke Math. J.* 161 (2012).
- [Wed19] T. Wedhorn. Adic Spaces. https://arxiv.org/abs/1910.05934. 2019.
- [Zav21] B. Zavyalov. "Mod-*p* Poincaré Duality in *p*-adic Analytic Geometry". In: *preparation* (2021).