MONOID OF LIOUVILLE SECTORS WITH CORNERS AND ITS INTRINSIC CHARACTERIZATION

YONG-GEUN OH

ABSTRACT. We provide a G-structure type characterization of Liouville sectors introduced in [GPS17] in terms of the characteristic foliation of the boundary, which we call *Liouville* σ -sectors. We extend this definition to the case with corners using the presymplectic geometry of null foliations of the coisotropic intersections of *clean coisotropic collection* of hypersurfaces which appear in the definition of Liouville sectors with corners. We give the definition of the structure of Liouville sectors with corners as a substructure of the monoid of manifolds with boundary and corners, and identify its automorphism group which enables us to give a natural definition of bundles of Liouville sectors. Then for a given Liouville σ -sector with corners (M, λ) , we introduce the class of gradientsectorial Lagrangian submanifolds and the notion of sectorial almost complex structures the pairs of which are amenable to the strong maximum principle. In particular the wrapped Fukaya category generated by gradient-sectorial Lagrangian branes on Liouville (σ -)sectors with corners becomes monoidal in the chain level under the monoidal product of manifolds with corners.

CONTENTS

	1. Introduction	2
	1.1. Sectional characterization of Liouville sectors	2
	1.2. Clean coisotropic collections and Liouville σ -sectors with corners	5
	1.3. Automorphism group of Liouville σ -sectors with corners	6
	1.4. Monoid of Liouville σ -sectors with corners	7
	1.5. Sectorial almost complex structures and gradient-sectorial Lagrangians	8
	2. Sectional characterization of sectorial hypersurfaces	11
	2.1. Definitions of σ -sectorial hypersurfaces and Liouville σ -sectors	12
	2.2. Preliminaries	13
	2.3. The leaf space is a topological manifold	16
	2.4. Smooth structure on the leaf space	20
	2.5. Symplectic structure on the leaf space	22
	2.6. Induced Liouville structure on the leaf space	24
	3. Geometry of clean coisotropic collections	25
	3.1. Gotay's coisotropic embedding theorem of presymplectic manifolds	25
	3.2. Structure of the null foliation of σ -sectorial corners	28
	3.3. Compatibility of null foliations of clean coisotropic intersections	31
4	4. Liouville σ -sectors and canonical splitting data	33

Date: October, 2021.

Key words and phrases. Monoid of Liouville σ -sectors with corners, end-profile functions, the (strong) maximum principle, sectorial almost complex structures, gradient-sectorial Lagrangian branes, presymplectic geometry.

This work is supported by the IBS project # IBS-R003-D1.

4.1. Definition of Liouville σ -sectors with corners	33
4.2. Integrable systems and canonical splitting data	34
5. Solution to [GPS17, Question 2.6] and convexity at infinity	38
5.1. Presymplectic geometry	38
5.2. Liouville geometry	41
5.3. Combining the two	43
6. Structure of Liouville σ -sectors and their automorphism groups	43
6.1. Some presymplectic geometry of ∂M	43
6.2. Automorphism group of Liouville σ -sectors	45
7. Monoid of Liouville σ -sectors and smoothing profiles	47
7.1. Monoid of manifolds with corners	47
7.2. Monoid of Liouville σ -sectors with corners	49
7.3. Smoothing profiles and their products	51
8. Sectorial almost complex structures and gradient-sectorial Lagrangians	54
8.1. Definition of sectorial almost complex structures	55
8.2. Gradient-sectorial Lagrangians and their products	56
8.3. Strong maximum principle and gradient-sectorial Lagrangian branes	58
9. Discussion and what to do	60
9.1. Pseudoconvex pairs and Z-invariant-at-infinity Lagrangian branes	60
9.2. Relationship with the Künneth-type formulae in Floer theory	61
9.3. Fukaya categories of Liouville manifolds of infinite type	62
References	63

1. INTRODUCTION

Unlike the set of manifolds with boundary, the set of manifolds with boundary and corners forms a monoid under the product of the enlarged category of manifolds with boundary and corners. (Obviously manifolds with boundary does not form a monoid: If X and Y are manifolds with nonempty boundary, then $X \times Y$ canonically becomes a manifold with boundary and corners.) For example, the topological boundary

$$\partial(X \times Y) = \partial X \times Y \coprod X \times \partial Y$$

itself is a manifold with corners but without boundary. For the simplicity of exposition, we will regard the boundary as a corner of codimension 1 and just call a manifold with boundary and corners a *manifold with corners*.

1.1. Sectional characterization of Liouville sectors. The original definition of Liouville sectors given in [GPS17] makes it somewhat awkward to identify the structure group of a bundle of Liouville sectors, as it is not a priori manifest (though it is verifiable) that the standard notion of Liouville automorphism of M is suitably compatible with certain sectorial data. It becomes even more awkward when one tries to define the bundle of Liouville sectors with *corners* as in [OT20]. which was the starting point of current investigation. In this paper, we introduce a more intrinsic definition but equivalent definition of Liouville sector which skirts this issue: We say it is more intrinsic in that our definition is closer to one in the sense of G-structures. (See [Che66] or [Ste83, Chapter VII] for a general introduction to G-structures.)

 $\mathbf{2}$

We start with our discussion of M for the case without corners.

Let (M, ω) be a symplectic manifold with boundary. The boundary ∂M (or more generally any coisotropic submanifold H) then carries a natural structure of a *presymplectic manifold* in the sense that the restriction two form

$$\omega_{\partial} := \iota^* \omega$$

has constant nullity. (See [Got82], [OP05] for some detailed explanation on presymplectic manifolds.) Here $\iota : \partial M \to M$ is the inclusion map.

Notation 1.1 $(\mathcal{D}_{\partial M}, \mathcal{N}_{\partial M} \text{ and } \pi : \partial M \to \mathcal{N}_{\partial M})$. We denote the characteristic distribution of $(\partial M, \omega_{\partial})$ by

$$\mathcal{D}_{\partial M} = \ker \omega_{\partial}.$$

With a slight abuse of notation, we also denote by $\mathcal{D}_{\partial M}$ the associated integrable foliation, and let $\pi_{\partial M} : \partial M \to \mathcal{N}_{\partial M}$ be its leaf map.

Now consider a Liouville manifold (M, λ) with boundary and denote by

 $(\partial_{\infty}M,\xi_{\infty})$

its ideal boundary as a contact manifold equipped with the contact distribution ξ_{∞} canonically induced by the Liouville form λ . (See [Gir17]. We recall that there is no contact form on $\partial_{\infty}M$ canonically induced from λ .)

Definition 1.2 (Liouville σ -sectors). We say a Liouville manifold with boundary (M, λ) is a *Liouville* σ -sector if the following hold:

- (a) The Liouville vector field Z of M is outward pointing along ∂M , and is tangent to ∂M at infinity.
- (b) $\partial_{\infty} M \cap \partial M$ is the boundary of $\partial_{\infty} M$, and is convex (as a hypersurface of the contact manifold $\partial_{\infty} M$).
- (c) The canonical projection map $\pi : \partial M \to \mathbb{N}_{\partial M}$ (to the leaf space of the characteristic foliation) admits a continuous section, and has fibers abstractly homeomorphic to \mathbb{R} .

The condition (c) in this definition is the difference from that of the *Liouville* sector of [GPS17] and is responsible for our naming of *Liouville* σ -sectors where σ stands either for 'section' or for 'sectional'. It can be replaced by the contractibility of fibers. (See Corollary 3.12.)

We also note that the details of λ matter only "at infinity"—the condition (c) depends only on the symplectic form $d\lambda$ (not on λ itself), while (a) and (b) depend only on the behavior of λ near the ideal contact boundary.

- **Remark 1.3.** (1) Our definition is closer to the one given in the spirit of G-structure (with integrability condition) [Che66], [Ste83]. (See also Corollary 5.2 for a similar characterization of convexity at infinity imposed in Definition (b).) In this sense, the choice of a section corresponds to a reduction of the structure group from Diff(\mathbb{R}) to Diff(\mathbb{R} , {0}) of the \mathbb{R} -bundle associated to the null foliation.
 - (2) It is worthwhile to mention that the presymplectic structure on $(\partial M, \omega_{\partial})$ uniquely determines a symplectic structure on the germ of a neighborhood up to symplectic diffeomorphism. (See [Got82].) Our definition of Liouville σ -sectors with corners is much based on Gotay's coisotropic embedding

theorem of presymplectic manifolds [Got82], applied to a germ of neighborhoods of the boundary ∂M or more generally of coisotropic submanifolds of $(M, d\lambda)$.

Note that a Liouville (σ -)sector M is a smooth manifold (possibly with noncompact corners) and the Liouville flow determines a well-defined contact manifold $\partial_{\infty}M$ "at infinity" (possibly with boundary). We will informally write

$$\partial_{\infty} M \cap \partial M$$

to mean the boundary of $\partial_{\infty} M$ and call it the *ceiling corner* of the Liouville sector. (When $\partial_{\infty} M$ has corners, "boundary" means the union of all boundary strata.)

Throughout this paper, by "near infinity," we mean "on the complement on some compact subset of M."

Theorem 1.4. Under the above definition of Liouville σ -sector, the following hold:

- (1) $\mathcal{N}_{\partial M}$ carries the structure of Hausdorff smooth manifold with corners such that π is a smooth submersion.
- (2) $\mathcal{N}_{\partial M}$ carries a canonical symplectic structure denoted by $\omega_{\mathcal{N}_{\partial M}}$ as a coisotropic reduction of $\partial M \subset M$.
- (3) $(\mathcal{N}_{\partial M}, \omega_{\mathcal{N}_{\partial M}})$ carries a canonical Liouville one-form $\lambda_{\mathcal{N}_{\partial M}}$ induced from the convexity hypothesis of $\partial_{\infty} M \cap \partial M \subset \partial_{\infty} M$.
- (4) We have a commutative diagram

where

- π is a smooth map which admits a smooth section $\sigma : \mathbb{N}_{\partial M} \to \partial M$ for which σ satisfies $\sigma^* \omega_{\partial} = \omega_{\mathbb{N}_{\partial M}}$,
- Setting $F := \text{Image } \sigma, \psi$ is a Liouville diffeomorphism between $(\mathcal{N}_{\partial M}, \lambda_{\mathcal{N}_{\partial M}})$ and the induced form $(F, \lambda|_F)$ defined by $\psi(\ell) = \sigma(\ell)$ for $\ell \in \mathcal{N}_{\partial M}$, and
- $\Psi : \partial M \to F \times \mathbb{R}$ is a diffeomorphism,

We refer to Section 2 for the precise description on the dependence of various structures and maps on the choice of section σ .

The following can be also derived in the course of proving the above theorem. (In fact the argument deriving this proposition is nearly identical to that of the proof of [GPS17, Lemma 2.5].)

Proposition 1.5. Let (M, λ) be a Liouville σ -sector. Then

- (1) Each choice of smooth section σ of π and a constant $0 < \alpha \leq 1$ canonically provides a smooth function $I : \partial M \to \mathbb{R}$ such that $Z(I) = \alpha I$,
- (2) There is a germ of neighborhood $Nbhd(\partial M)$ (unique up to a symplectomorphism fixing ∂M) on which the natural extension of I, still denoted by I, admits a unique function $R : Nbhd(\partial M) \to \mathbb{R}$ satisfying $\{R, I\} = 1$.

Another interesting consequence, when combined with Gotay's normal form theorem of neighborhoods of coisotropic submanifolds, is the following affirmative answer to a question raised by Ganatra-Pardon-Shende in [GPS17]. **Theorem 1.6** (Theorem 5.1; Question 2.6 [GPS17]). Suppose M is a Liouville manifold-with-boundary such that

- (1) the Liouville vector field is tangent to ∂M at infinity, and
- (2) there is a diffeomorphism $\partial M = F \times \mathbb{R}$ sending the characteristic foliation to the foliation by leaves $\mathbb{R} \times \{p\}$.

Then $\partial_{\infty} M \cap \partial M$ is convex in $\partial_{\infty} M$. In particular M is a Liouville sector in the sense of [GPS17].

The main task is to construct a contact vector field transversal to the ceiling corner

$$\partial_{\infty} M \cap \partial M =: F_{\infty}$$

in the contact manifold $\partial_{\infty} M$. This will be done by utilizing the aforementioned normal form theorem, whose details we refer readers to the proof of Theorem 5.1 in Section 5.

The following equivalence theorem is an immediate corollary of Theorem 1.4 and Theorem 1.6.

Theorem 1.7. Let (M, λ) be a Liouville manifold with boundary. Suppose the Liouville vector field Z of λ is tangent to ∂M at infinity. Then the followings are equivalent:

- (1) (M, λ) is a Liouville sector in the sense of [GPS17].
- (2) (M, λ) is a Liouville σ -sector.
- (3) There is a diffeomorphism $\partial M = F \times \mathbb{R}$ sending the characteristic foliation to the foliation by leaves $\mathbb{R} \times \{p\}$.

Remark 1.8 (Liouville sectors as a *G*-structure). In particular, the second characterization provides a natural characteristic of Liouville sectors in the spirit of *G*-structures purely in terms of the presymplectic geometry of ∂M in addition to a little bit of Liouville geometry of a neighborhood of the corner Nbhd $(\partial_{\infty} M \cap \partial M)$ in *M* requiring the Liouville vector field *Z* to be tangent to ∂M therein. Other defining data of Liouville sectors follow therefrom as 'properties'.

1.2. Clean coisotropic collections and Liouville σ -sectors with corners. The definition of Liouville σ -sector can be extended to the case with corners. The definition of Liouville σ -sectors with corners strongly relies on the general intrinsic geometry of the clean coisotropic collection. Study of this geometry in turn strongly relies on the coisotropic calculus and Gotay's coisotropic embedding theorem of general *presymplectic manifolds* [Got82].

Definition 1.9 (Clean coisotropic collection). Let (M, λ) be a Liouville manifold with corners. Let $H_1, \ldots, H_m \subset M$ be a collection of hypersurfaces Z-invariant at infinity, that satisfies

- (1) The H_i cleanly intersect,
- (2) All pairwise intersections $H_i \cap H_j$ are coisotropic.

Denote the associated codimension m corner by

$$C = H_1 \cap \dots \cap H_m$$

and by \mathbb{N}_C the leaf space of the null-foliation of the coisotropic submanifold C. Then we prove that for each choice of sections $\sigma = \{\sigma_1, \dots, \sigma_m\},\$

- there is a natural fiberwise \mathbb{R}^m -action on C which is a simultaneous linearization of the characteristic flows of the sectorial hypersurfaces H_i 's.
- each fiber is diffeomorphic to \mathbb{R}^m utilizing the standard construction of action-angle variables in the integrable system.

(See [Arn88] and Corollary 3.12 for the relevant discussion.) This leads us to the final definition of Liouville σ -sectors with corners.

Definition 1.10 (Liouville σ -sectors with corners). Let M be a manifold with corners equipped with a Liouville one-form λ . We call (M, λ) a Liouville σ -sector with corners if at each σ -sectorial corner δ of ∂M , the corner can be expressed as

$$C_{\delta} := H_{\delta,1} \cap \dots \cap H_{\delta,m}$$

for a clean coisotropic collection $\{H_{\delta,1}, \cdots, H_{\delta,m}\}$ such that each fiber of the canonical projection

$$\pi_{C_{\delta}}: C_{\delta} \to \mathcal{N}_C$$

is contractible. We call such a corner a σ -sectorial corner of codimension m.

We will show that each choice of σ will canonically provide an equivariant splitting data

$$(F, \{(R_i, I_i)\}_{i=1}^m), \quad d\lambda = \omega_F \oplus \sum_{i=1}^m dR \wedge dI$$

on $Nbhd(C_{\delta}) \cong F \times \mathbb{C}^{m}_{Re \geq 0}$ for σ -sectorial corners that is equipped with the Hamiltonian \mathbb{R}^{m} -action whose moment map is precisely the coordinate projection

 $\operatorname{Nbhd}(C) \to \mathbb{R}^m_{>0}; \quad x \mapsto (R_1(x), \dots, R_m(x)).$

(See Theorem 4.4 for the precise statement.)

We also prove the following equivalence result.

Theorem 1.11. Definition 1.10 is equivalent to that of Liouville sectors with corners from [GPS18].

We refer to Definition 3.1 for the comparison between Definition 1.10 and the definition of Liouville sectors with corners from [GPS18].

1.3. Automorphism group of Liouville σ -sectors with corners. Thanks to Theorem 1.4 or Theorem 1.7, our definition of Liouville σ -sectors with corners enables us to give a natural notion of Liouville automorphisms of Liouville sectors from [GPS17] which is similar to the case without boundary and which does not depend on choices of auxiliary defining function I that appear in the original definition of [GPS17, Definition 2.4].

We start with the following observation

Lemma 1.12 (Lemma 6.3). Fix a diffeomorphism $\phi : (M, \partial M) \to (M, \partial M)$ and suppose $\phi^* \lambda = \lambda + df$ for a function $f : M \to \mathbb{R}$, not necessarily compactly supported. Then the restriction $\phi|_{\partial M} = \phi_{\partial} : \partial M \to \partial M$ is a presymplectic diffeomorphism, i.e., satisfies $\phi^*_{\partial}\omega_{\partial} = \omega_{\partial}$. In particular, it preserves the characteristic foliation of ∂M .

Remark 1.13. Recall that a manifold with corners X is (pre)symplectic if there is a stratawise (pre)symplectic form ω , i.e., a collection of (pre)symplectic forms

that is compatible under the canonical inclusion map of strata

$$\iota_{\alpha\beta}: X_{\alpha} \hookrightarrow X_{\beta}, \quad \alpha < \beta$$

i.e., $\omega_{\alpha} = \iota_{\alpha\beta}^* \omega_{\beta}$. Here *I* is the POSET that indexes the strata of the stratified manifold *X*. By definition, a diffeomorphism between two manifolds with corners preserves dimensions of the strata.

Lemma 1.12 enables us to define the "structure" of Liouville σ -sectors (Definition 6.1), and to identity its automorphism group $\operatorname{Aut}(M, \lambda)$ in the same way as for the Liouville manifold case.

Definition 1.14 (Automorphisms group $\operatorname{Aut}(M, \lambda)$). Let (M, λ) be a Liouville σ -sector, possibly with corners. We call a diffeomorphism $\phi : (M, \partial M) \to (M, \partial M)$ a Liouville automorphism if ϕ satisfies the following:

 $\phi^*\lambda = \lambda + df$

for a compactly supported function $f: M \to \mathbb{R}$. We denote by $\operatorname{Aut}(M, \lambda)$ the set of automorphisms of (M, λ) .

Obviously $\operatorname{Aut}(M, \lambda)$ forms a topological group which is a subgroup of $\operatorname{Symp}(M, d\lambda)$, the group of symplectic diffeomorphisms of $(M, d\lambda)$.

1.4. Monoid of Liouville σ -sectors with corners. In [Oh21], the present author introduces the notion of λ -sectorial packages in the study of Floer theory on the Liouville sectors, where Lagrangian branes entering in the construction of Fukaya category are still assumed to be Z-invariant-at-infinity Lagrangian submanifolds as common in the literature such as [AS10], [GPS17].

In the present paper, we introduce a different class of Lagrangian submanifolds called gradient-sectorial Lagrangian submanifolds, with respect to which we can make this new Floer package go well along with the aforementioned monoidal structure of Liouville σ -sectors with corners. (See Section 7 for the detailed description of this class of Lagrangian branes.) We anticipate that this new package will facilitate the study of the Künneth-type formula in wrapped Fukaya category, the study of which we postpone elsewhere. (See [GPS17, Conjecture 3.40, Conjecture 4.39] for relevant conjectures concerning this monoidality issue.)

We show that the product $L_1 \times L_2$ of gradient-sectorial Lagrangians automatically are objects in the product $M_1 \times M_2$ of two Liouville (σ -)sectors with corners without making any deformation in our new framework. For example, we have a natural inclusion map

 $Ob(\mathsf{Fuk}(X,\omega_X)) \times Ob((\mathsf{Fuk}(Y,\omega_Y)) \hookrightarrow Ob(\mathsf{Fuk}(X \times Y,\omega_X \oplus \omega_Y)))$

on the nose. (See Definition 8.4 and Theorem 8.6.)

Remark 1.15. The standard definition of Z-invariant-at-infinity branes for the Liouville sectors is *not* monoidal in any obvious sense. For example, the product $L_1 \times L_2$ is not $(Z_X \oplus Z_Y)$ -invariant for the Z_X -invariant-brane-at-infinity L_1 of X and a Z_Y -invariant-at-infinity brane L_2 of Y. Because of this, one must take the following deformation process which has been used in the literature (See [Gro17, Gao17, GPS17, GPS18], especially [GPS18, Section 6] for a detailed explanation of this procedure in relation to a construction of Künneth-type embedding.) :

• take a corner smoothing of the corner $\partial_{\infty} M \cap \partial M$,

- take a deformation of the product Liouville form $\lambda_1 \times \lambda_2$ and the product vector field $Z_1 \times Z_2$ to the associated Liouville vector field,
- Then deform $L_1 \times L_2$ to a Z-invariant-at-infinity Lagrangian on $X \times Y$.

This whole deformation process will not be needed in our gradient-sectorial framework: This process is already subsumed in the sectorial package introduced in [Oh21] and augmented here: We modify the package from [Oh21] to one in the present paper so that it respects the aforementioned monoidal structure of manifolds with corners.

For the study of this monoidality property of the wrapped Fukaya category, our sectional characterization of Liouville sectors makes it natural to take the product of Liouville sectors with corners. The following is an easy consequence of our definition of Liouville σ -sectors with corners.

Proposition 1.16 (Proposition 7.11). The collection of Liouville σ -sectors with corners naturally forms a commutative monoid under direct product (with commutativity and associativity holding up to natural isomorphism).

The discussion in the present subsection and henceforth equally applies both with the definition of Liouville sectors of [GPS17] and with that of Liouville σ -sectors introduced in the present paper. We will mainly work with our definition of Liouville σ -sectors, unless otherwise mentioned.

1.5. Sectorial almost complex structures and gradient-sectorial Lagrangians. Now we propose a class of *gradient-sectorial* Lagrangian branes to form the object of a wrapped Fukaya category Fuk(M). (Our notation for Fuk(M) suppresses the dependence on λ and our choice working with gradient-sectorial Lagrangians instead of Z-invariant-at-infinity ones.)

For this purpose, we first equip (M, λ) with a splitting data

$$\operatorname{Nbhd}(\partial M) \cong F \times \mathbb{C}^k_{\operatorname{Re}>0}, \quad \{(R_i, I_i)\}$$

$$(1.2)$$

and an end-profile function

$$\mathfrak{s}:=\mathfrak{s}_{arphi}$$

which is introduced in [Oh21]. (See also Subsection 7.3.2 of the present paper for a brief description of \mathfrak{s}_{φ} .)

Remark 1.17. We will show in Theorem 4.4 that this data itself is canonically induced from the choice of sections $\sigma = \{\sigma_1, \ldots, \sigma_k\}$ for the clean coisotropic collection

$$\{H_1,\cdots,H_k\}$$

of σ -sectorial hypersurface H_i 's. We call this canonical splitting data the σ -splitting data.

Recall from [Oh21, Section 3] that \mathfrak{s}_{φ} is a collection of functions

$$s_{k_{\delta}+1,\varphi_{\delta}} = -\log\varphi(R_{\delta,1},\cdots,R_{\delta,k},e^{-s}) \tag{1.3}$$

associated to each sectorial corner δ of M which are glued by a partition of unity on M. Here $\varphi : \mathbb{R}_{\geq 0}^{k+1} \to \mathbb{R}_{\geq 0}$ is a convex corner-smoothing function. (See [FOOO20], [Oh21] for the details.) An upshot of this function \mathfrak{s}_{φ} is that it provides a pseudoconvex pair $(\mathfrak{s}_{\varphi}, J)$ on Nbhd $(\partial_{\infty} M \cup \partial M)$ for a sectorial complex structures J. (See [Oh21] for detailed discussion on the notion of pseudoconvex pairs.)

The splitting data given above also provides $Nbhd(\partial M)$ with a foliation \mathcal{F}_F whose leaves are given by

$$\mathcal{F}_F: F \times \{(R,I)\}, \quad (R,I) = (R_1 + \sqrt{-1}I_1, \cdots, R_k + \sqrt{-1}I_k) \in \mathbb{C}^k.$$
 (1.4)

Definition 1.18 (Sectorial almost complex structures). Let (M, λ) be a Liouville sector with boundary and corners equipped with a splitting data and an end-profile function \mathfrak{s}_{φ} . An ω -tame almost complex structure J on a Liouville sector is said to be *sectorial* (with respect to the given smoothing profile) if J satisfies the following:

(J1) [\mathcal{F}_F is *J*-complex] In a neighborhood of Nbhd(∂M) of ∂M , we require

$$J\left(T^*F \oplus 0_{\operatorname{span}\{dR_i, dI_i\}_{i=1}^k}\right) \subset T^*F \oplus 0_{\operatorname{span}\{dR_i, dI_i\}_{i=1}^k},\tag{1.5}$$

and J restricts to an almost complex structure of contact-type on F.

(J2) $[(\mathfrak{s}_{\varphi}, J)$ is a pseudoconvex pair] In a neighborhood $\operatorname{Nbhd}^{\mathbb{Z}}(\partial M) \cup \partial_{\infty})$ of $\partial M \setminus \operatorname{Nbhd}(\partial_{\infty} M)$, we have

$$-d(d\mathfrak{s}_{\varphi} \circ J) \ge 0$$

as a (1, 1)-current.

We denote by $\mathcal{J}_{\mathfrak{s}_{\varphi}}^{\mathrm{sec}} = \mathcal{J}_{\mathfrak{s}_{\varphi}}^{\mathrm{sec}}(M)$ the set of sectorial almost complex structures.

Remark 1.19. Obviously any almost complex structure J satisfying (J1) is sectorial if it satisfies

$$-d\mathfrak{s}_{\varphi}\circ J=\lambda+df$$

for some function f, not necessarily compactly supported. For example, the κ sectorial or λ -sectorial almost complex structures considered in [Oh21] are sectorial. This in particular shows that sectorial almost complex structures exist in abundance. The more complicated notion of λ -sectorial almost complex structures is needed to make the usual Z-invariant-at-infinity Lagrangians amenable to the strong maximum principle: Recall from [Oh21] that a pseudoconvex pair (ψ , J) is called Liouville-pseudoconvex if it satisfies the stronger condition

$$-d\psi \circ J = \lambda$$

in place of (J2). This duality requirement for $(-d\psi, \lambda)$ suffices to show that the Z-invariant-at-infinity Lagrangian submanifolds are amenable to the strong maximum principle. (See [Oh21, Section 11] for complete details of the construction of such a pair (ψ, J) .)

Now we introduce the notion of gradient-sectorial Lagrangians with respect to the end-profile function $\mathfrak{s} = \mathfrak{s}_{\varphi}$, which is amenable to the strong maximum principle with respect to sectorial almost complex structures for the given end-profile function \mathfrak{s} . We consider its normalized gradient vector field

$$Z_{\mathfrak{s}} := \frac{\operatorname{grad}\mathfrak{s}}{|\operatorname{grad}\mathfrak{s}|^2} \tag{1.6}$$

with respect to the usual metric

$$g_J(v,w) := \frac{d\lambda(v,Jw) + d\lambda(w,Jv)}{2}$$

Definition 1.20 (Gradient-sectorial Lagrangian branes). Let (M, λ) be a Liouville sector with corners. Let \mathfrak{s} be the end-profile function associated to a given smoothing profile. We say that an exact Lagrangian submanifold L of (M, ω) is gradient-sectorial if

- (1) $L \subset \operatorname{Int} M \setminus \partial M$ and $\operatorname{dist}(L, \partial M) > 0$.
- (2) There exists a sufficiently large $r_0 > 0$ such that $L \cap \mathfrak{s}^{-1}([r_0,\infty))$ is $Z_{\mathfrak{s}}$ invariant, i.e., $Z_{\mathfrak{s}}$ is tangent to $L \cap \mathfrak{s}^{-1}([r_0, \infty))$.

Then in Theorem 8.7 we prove the confinement results -hence Gromov compactnessfor J-holomorphic curves with boundaries on gradient-sectorial Lagrangians for sectorial almost-complex structures by proving that such J-holomorphic curves are amenable to the strong maximum principle in terms of the end-profile function \mathfrak{s}_{φ} .

We establish the product Lagrangian $L_1 \times L_2$ itself of two gradient-sectorial Lagrangian submanifolds L_1 and L_2 (not just up to isotopy as in [GPS17, GPS18, Gao17) can be used as boundary conditions for holomorphic curves—in other words, as objects for an appropriate wrapped Fukaya category.

Theorem 1.21 (Theorem 8.6). Let (X, ω_X) and (Y, ω_Y) be Liouville σ -sectors. Then $L_1 \times L_2$ is gradient-sectorial if both L_1 and L_2 are gradient-sectorial.

We refer readers to Section 7 for more detailed discussion on the product.

Acknowledgments: The present work is supported by the IBS project IBS-R003-D1. We would like to thank Hiro Lee Tanaka for his collaboration on the study of Liouville sectors and for useful comments on the preliminary draft of the present work.

Conventions:

- Hamiltonian vector field X_H : $X_H \rfloor \omega = dH$,
- Canonical one-form θ_0 on T^*Q : $\theta_0 = \sum_{i=1}^n p_i dq_i$,

- Canonical symplectic form ω₀ on T*Q: ω₀ = d(-θ) = Σⁿ_{i=1} dq_i ∧ dp_i,
 Liouville one-form on (T*Q, ω₀): λ = -θ = -Σⁿ_{i=1} p_idq_i,
 Symplectization SC of contact manifold (C, θ): SC = C × ℝ with ω = $d(e^{s}\theta)$. Here note that we write the \mathbb{R} -factor after the C-factor.

Notations:

- $\partial_{\infty} M$: the asymptotic boundary of M.
- \overline{M} : the completion of M which is $\partial_{\infty} M [] \partial M$.
- DM: the union $\partial_{\infty} M \cup \partial M$ in \overline{M} .
- $\partial_{\infty}^{\text{Liou}} M$: the ideal boundary of a Liouville manifold M (or sector).
- Aut (M, λ) : The group of Liouville diffeomorphisms of Liouville σ -sector $(M, \lambda).$
- $\omega_{\partial} = d\lambda_{\partial}$: The induced presymplectic form on ∂M with $\lambda_{\partial} := \iota^* \lambda$.
- Aut (M, λ_{∂}) : The group of pre-Liouville diffeomorphisms of exact presymplectic manifolds $(M, d\lambda_{\partial})$.
- H : a σ -sectorial hypersurface $H \subset M$.

2. Sectional characterization of sectorial hypersurfaces

We start with the case without corners but with nonempty boundary ∂M , postpointing the study of the case with corners till Section 3.

For the comparison, we recall the definition of Liouville sectors in [GPS17]. In fact we will consider the definition of sectorial hypersurfaces in [GPS18, Definition 9.2] and restrict that to the sectorial boundary of a Liouville domain.

To facilitate our exposition, we utilize Giroux's notion of the *ideal completion* of the Liouville domain (M, λ) .

Definition 2.1 (Ideal completion \overline{M}). [Gir17]

- (1) An *ideal Liouville domain* (W, ω) is a domain endowed with an ideal Liouville structure ω .
- (2) The *ideal Liouville structure* is an exact symplectic form on Int W admitting a primitive θ such that: For some (and then any) function $u: W \to \mathbb{R}_{\geq 0}$ with regular level set $\partial_{\infty}W = \{u = 0\}$, the product $u\theta$ extends to a smooth one-form on W which induces a contact form on ∂W .
- (3) When a Liouville manifold (M, λ) is Liouville isomorphic to $(\operatorname{Int} W, \theta)$, we call W the ideal completion of M and denote it by \overline{M} .

Remark 2.2. First, this definition provides a natural topology and smooth structure on the completion \overline{M} and a Liouville structure on $M(= \operatorname{Int} W)$ as an open Liouville manifold. Secondly it also provides a natural class of Liouville diffeomorphisms on M as the restriction of diffeomorphisms of $\overline{M} = W$. (See [Gir17].)

For a (noncompact) Liouville manifold (M, λ) (without boundary) its ideal boundary, denoted by $\partial_{\infty} M$, is defined to be the set of asymptotic rays of Liouville vector field Z. Then the *ideal completion* is the coproduct

$$\overline{M} = M \coprod \partial_{\infty} M$$

equipped with the obvious topology. When (M, λ) is a Liouville sector with boundary ∂M , its ideal boundary is still well-defined by the Z-invariance requirement at infinity put on ∂M in the definition of Liouville sectors [GPS17] and so is its completion \overline{M} . Then we have the formula for the topological boundary

$$\partial \overline{M} = \partial_{\infty} M \cup \partial M$$

To ease our exposition, we often abuse our notation

$$\partial_{\infty}(\partial M) = \partial_{\infty} M \cup \partial M$$

for the coproduct $\partial_{\infty} M \coprod \partial M$ after the present section, as long as there is no danger of confusion. Likewise we also abuse the notation like

$$\partial_{\infty}H = \partial_{\infty}M \cap H$$

where the intersection is actually taken as a subset of M.

This being said, the following definition is nothing but when the sectorial collection there in [GPS18, Definition 9.2] is a single element.

Definition 2.3 (See Definition 9.2 [GPS17]). A sectorial hypersurface $H \subset M$ is a hypersurface Liouville manifold-with-boundary M satisfying the following equivalent definitions:

• For some $\alpha > 0$, there exists a function $I : H \to \mathbb{R}$ with $ZI = \alpha I$ and $dI_{\mathcal{D}} > 0$.

- For every $\alpha > 0$, there exists a function $I : H \to \mathbb{R}$ with $ZI = \alpha I$ and $dI_{\mathcal{D}} > 0$.
- The ideal boundary $\partial_{\infty} M \cap H = \partial_{\infty} H$ is convex and there is a diffeomorphism $H = F \times \mathbb{R}$ sending the characteristic foliation of H to the foliation of $F \times \mathbb{R}$ by leaves $\{p\} \times \mathbb{R}$.

2.1. Definitions of σ -sectorial hypersurfaces and Liouville σ -sectors. Here we give another more intrinsic definition of sectorial hypersurface. The definition is intrinsic in that it is closer to one in the spirit of *G*-structures (with integrability condition), not involving the defining function *I*: Existence of the data of function *I* appearing in the definition of Liouville sectors in [GPS17] is now a 'property', not a 'defining data', of Liouville σ -sector in our definition. (We refer readers to [Ste83, Chapter VII] for a nice introduction to the geometry of *G*-structures.)

Definition 2.4 (σ -sectorial hypersurface). Let (M, λ) be a Liouville manifold with boundary (without corners). Let $H \subset M$ be a smooth hypersurface such that its completion \overline{H} has the union

$$(\partial_{\infty}M \cap \overline{H}) \cup (\overline{H} \cap \partial M) =: \partial_{\infty}H \cup \partial \overline{H}$$

as its (topological) boundary. H is a σ -sectorial hypersurface if it satisfies the following:

- (1) Z is tangent to H at infinity,
- (2) $F_{\infty} := \partial_{\infty} M \cap H \subset \partial_{\infty} M$ is a convex hypersurface of the contact manifold $\partial_{\infty} M$,
- (3) The canonical projection map $\pi : H \to \mathcal{N}_H$ has a continuous section and each of its fiber is homeomorphic to \mathbb{R} .

To avoid some confusion with the corners in ∂M , we call the intersection

$$\partial_{\infty} M \cap \partial \overline{M}$$

the *ceiling corner*. This is the corner of the ideal completion \overline{M} of M of codimension 2. (We will call the genuine corners of M the *sectorial corners* in Section 3 when we consider the Liouville sectors with corners.)

By applying the notion of σ -sectorial hypersurface to the boundary $\partial M \subset M$, we obtain the following definition.

Definition 2.5 (Liouville σ -sector). Let M be a noncompact manifold with boundary such that its completion \overline{M} has (topological) boundary given by the union

$$\partial_{\infty} M \cup \partial M = DM$$

and $\partial_{\infty} M \cap \partial M$ is the codimension two corner of \overline{M} . M is called an *Liouville* σ -sector if its boundary $\partial M \subset M$ is a σ -sectorial hypersurface in the sense of Definition 2.4.

We start with the well-known fact that each hypersurface $H \subset M$ in a symplectic manifold (M, ω) carries the canonical characteristic foliation \mathcal{D} . The definition of this foliation is based on the fact that any hypersurface S of (M, ω) is a *coisotropic* submanifold in that

(1) We have

$$(T_x H)^{\omega_x} \subset T_x H,$$

for any $x \in H$, where $(T_x H)^{\omega_x}$ is the ω_x -orthogonal complement
 $(T_x H)^{\omega_x} := \{ v \in T_x M \mid \omega_x(v, w) = 0 \,\forall w \in T_x H \}.$

(2) Let
$$\iota_H : H \to M$$
 be the inclusion map and

 $\ker \iota_H^* \omega_x := \{ v \in T_x H \mid \omega_x(v, w) = 0 \,\forall w \in T_x H \}$

has constant rank 1 for all $x \in H$.

Then we denote $\mathcal{D} = \ker \iota_H^* \omega$ which defines a 1-dimensional (integrable) distribution of H, and call it the characteristic distribution or the null distribution of H. We denote by \mathcal{N}_H the leaf space of the associated foliation. It is also well-known that \mathcal{D} carries a transverse symplectic structure which induces one on the leaf space

$$\mathcal{N}_H := H/\sim \tag{2.1}$$

chart-wise. With slight abuse of notation, we will also denote by $\mathcal D$ the associated foliation.

Of course, the quotient topology of a leaf space may not be Hausdorff in general. We will show that under the conditions laid out in Definition 1.2, the aforementioned transverse symplectic form, as well as its smooth structure, descends to the leaf space. For the rest of this section, we always assume M is a Liouville σ -sector as in Definition 1.2, unless otherwise said.

2.2. Preliminaries.

2.2.1. Orientations. We write

$$F_{\infty} = F_{H,\infty} := \partial_{\infty} M \cap H.$$

At each point $x \in H \cap Nbhd(\partial_{\infty}M)$, we have a natural exact sequence

$$0 \to \mathcal{D}_x \to T_x H \to T_x H / \mathcal{D}_x \to 0. \tag{2.2}$$

The quotient carries a canonical symplectic bilinear form and so carries a natural symplectic orientation.

Choice 2.6 (Orientation of \mathcal{D}). Let $H \subset M$ be a σ -sectorial hypersurface. Make a choice of orientation on the trivial line bundle $\mathcal{D} \to H$.

Definition 2.7 (Presymplectic orientation on H). Let $\mathcal{D} \to H$ be given an orientation $o_{\mathcal{D}}$ on a neighborhood of F_{∞} in $\partial_{\infty}M$. We call the orientation on $TH|_{F_{\infty}}$ given by the direct sum orientation

$$T_xH|_{F_{\infty}} = (T_xH/\mathcal{D}_x) \oplus \mathcal{D}_x$$

the presymplectic orientation of H relative to $o_{\mathcal{D}}$.

When we are given a section $\sigma : \mathcal{N}_H \to H$ such that Image $\sigma = F_N$ far out close to $\partial_{\infty} M$, then this orientation coincides with that of $T_x H$ induced by the exact sequence

$$0 \to T_x F_N \to T_x H \to \mathcal{D}_x \to 0 \tag{2.3}$$

as oriented vector spaces. (We alert readers that the presymplectic orientation of H is *not* the one naturally given by the exact sequence (2.2), rather arising from (2.3).)

Definition 2.8 (Asymptotic boundary orientation of $F_{\infty} \subset \overline{H}$). Regard

$$F_{\infty} := \partial_{\infty} M \cap H$$

as the asymptotic boundary of H where H is equipped with the *presymplectic* orientation of H relative to $o_{\mathcal{D}}$. We call this orientation of F_{∞} the asymptotic boundary orientation in H.

The orientation on F_{∞} given in Definition 2.8 may or may not coincide with the symplectic orientation of F_{∞} .

Definition 2.9. Equip \mathcal{D} with an orientation $o_{\mathcal{D}}$ and in turn orient H by the presymplectic orientation. The intersection F_{∞} is decomposed into

$$F_{\infty} = F_{\infty}^{+} \coprod F_{\infty}^{-}$$

such that the asymptotic boundary orientation of $F_{\infty} \subset \overline{H}$ coincides with the symplectic orientation on F_{∞}^+ and not on F_{∞}^- .

This discussion leads us to the following:

Definition 2.10. Let M be a Liouville σ -sector and H be a σ -sectorial hypersurface.

- (1) We call $F_{\infty} = \partial_{\infty} M \cap H$ the *end* of *H*.
- (2) Let \mathcal{D} be equipped with an orientation $o_{\mathcal{D}}$. We call F_{∞}^{\pm} the positive (resp. the negative) end of H and the positive (resp. the negative) boundary of $\partial_{\infty} M$ with respect to $o_{\mathcal{D}}$.

We illustrate the decomposition in Definition 2.9 in the case of sectorial boundary ∂M of a Liouville sector M.

Example 2.11 (Boundary orientation for $H = \partial M$). In the special case of the boundary $H = \partial M$ of a Liouville sector, ∂M itself carries the canonical boundary orientation of the symplectic $(M, d\lambda)$, and hence a natural orientation on $\mathcal{D}_{\partial M}$ induced by the short exact sequence (2.3),

$$0 \to T_x F_\infty \to T_x(\partial M)|_{F_\infty} \to \mathcal{D}_{\partial M}|_x \to 0.$$

This in turn induces the presymplectic orientation on ∂M and the asymptotic boundary orientation on F_{∞} .

Example 2.12 $(F_{\infty}^{\pm} \text{ on } T^*[0,1])$. Now consider the case of the cotangent bundle $M = T^*[0,1]$ of the closed interval [0,1] equipped with the Liouville form

$$\lambda = -p \, dq. \tag{2.4}$$

(This is the negative of the standard Liouville one-form pdq in the cotangent bundle.) The standard orientation of the interval induces a diffeomorphism $M \cong [0,1]_q \times \mathbb{R}_p$ which carries the symplectic orientation induced by the symplectic form

$$dq \wedge dp$$
.

(We alert the readers that this is the negative of the convention $dp \wedge dq$ used by [GPS17].) The boundary $\partial M \cong \{0,1\} \times \mathbb{R}_p$ has 2 connected components. The characteristic foliation's orientation is compatible with the vector field $\frac{\partial}{\partial p}$. Note that the Liouville vector field of the Liouville form (2.4) on $T^*[0,1] \cong [0,1]_q \times \mathbb{R}_p$ is given by the Euler vector field

$$\vec{E} := p \frac{\partial}{\partial p} \tag{2.5}$$

on T^*M which vanishes at p = 0. So each leaf $\{q\} \times \mathbb{R}_p$ of the foliation consists of 3 different orbit sets of the Liouville vector field

$$\mathbb{R}_+ = (0, \infty), \quad \{0\}, \mathbb{R}_- = (-\infty, 0).$$

We may identify $\partial_{\infty} M$ with two disjoint copies of [0, 1] at " $p = \pm \infty$." F_{∞} consists of four points, which we will denote by $(0, \pm \infty)$ and $(1, \pm \infty)$ again using the informal notation allowing p to attain $\pm \infty$. Under this notation, we have that

$$F_{\infty}^{+} = \{(0, -\infty), (1, \infty)\},$$
 and $F_{\infty}^{-} = \{(0, \infty), (1, -\infty)\}.$

Example 2.13. More generally, let $Q = Q^n$ be a connected *n*-manifold with boundary and let $M = T^*Q$. The inclusion $T(\partial Q) \hookrightarrow TQ$ induces a quotient map $T^*Q|_{\partial Q} \to T^*(\partial Q)$ of bundles on ∂Q ; the kernel induces the characteristic foliation on

$$T^*Q|_{\partial Q} = \partial M.$$

Informally: At a point $(q, p) \in \partial M$, the oriented vector defining the characteristic foliation is the symplectic dual to an inward vector normal to ∂Q . For example, identifying Q near ∂Q with the right half plane with final coordinate p_n , in standard Darboux coordinate (q, p), the characteristic foliation is generated by $\frac{\partial}{\partial p_n}$.

If dim $Q \ge 2$, we have that $F_{\infty}^+ = F_{\infty}$ (and so $F_{\infty}^- = \emptyset$), and F_{∞} is identified with the restriction of ∂Q of the cosphere-at-infinity bundle of T^*Q .

2.2.2. Convexity of $\partial_{\infty} M \cap \partial M$ and choice of contact vector field. Recall that $\partial_{\infty} M$ is naturally oriented as the ideal boundary of symplectic manifold M with Z pointing outward along $\partial_{\infty} M$. We take a contact-type hypersurface $S_0 \subset M$ that is transverse to ∂M and identify a neighborhood Nbhd $(\partial_{\infty} M)$ with the (half) of the symplectization $S(S_0)$ of the contact manifold $(S_0, \iota_{S_0}^* \lambda)$

$$S_+(S_0) := S_0 \times [0,\infty)$$

and decompose M into

$$M = (M \setminus Nbhd(\partial_{\infty} M) \cup (S_0 \times [0, \infty))$$

so that $Z = \frac{\partial}{\partial s}$ for the symplectization form $d(e^s \iota_{S_0}^* \lambda)$ of the contact manifold $(S_0, \iota_{S_0}^* \lambda)$ on Nbhd $(\partial_{\infty} M) = S_0 \times [0, \infty)$.

Next, by the convexity hypothesis of $\partial_{\infty} M \cap H$ in $\partial_{\infty} M =: F_{\infty}$, there exists a contact vector field η of the contact structure $(\partial_{\infty} M, \xi_{\infty})$ on a neighborhood of F_{∞} in $\partial_{\infty} M$ that is transverse to F_{∞} . This naturally equips F_{∞} with the reduced symplectic form and so the symplectic orientation thereon. On the other hand, a choice of contact vector field η transverse to F_{∞} in $\partial_{\infty} M$, also gives rise to an isomorphism

$$(T(\partial_{\infty} M)/\xi)|_{F_{\infty}} \cong \operatorname{span}_{\mathbb{R}}\{\eta\}.$$

In particular we have an exact sequence

$$0 \to TF_{\infty} \to T(\partial_{\infty}M)|_{F_{\infty}} \to \operatorname{span}_{\mathbb{R}}\{\eta\} \to 0.$$
(2.6)

Choice 2.14 (Choice of contact vector field η). Equip the asymptotic boundary $\partial_{\infty} M$ of the symplectic manifold $(M, d\lambda)$ with the (asymptotic) boundary orientation, i.e., the one oriented by the Liouville vector field Z on M which is outward pointing along $\partial_{\infty} M$. We also equip F_{∞} the symplectic orientation. Then we make the choice of the contact vector field η so that the sequence (2.6) becomes an exact sequence of oriented vector spaces.

Example 2.15. (1) The choice of contact vector field η in Choice 2.14 is consistent with the convention of writing the symplectization as $S_0 \times [0, \infty)$,

i.e., writing the \mathbb{R} -factor after the S_0 -factor. On the symplectization of general contact manifold (C, θ) in general, we have the splitting

$$T(SC) = \xi_C \oplus \mathbb{R} \langle R_\theta \rangle \oplus \mathbb{R} \langle Z \rangle.$$

Then $\eta = \pm R_{\theta}$ so that $\{Z, \eta\}$ forms the positively oriented basis of the complex orientation of $\mathbb{C} \cong \mathbb{R}\langle R_{\theta} \rangle \oplus \mathbb{R}\langle Z \rangle$.

(2) The choice is also compatible with the writing of the splitting data in [GPS17]

$$\mathcal{N}bhd(\partial M) = F \times \mathbb{C}_{Re>0}$$

as oriented manifolds. In this case, the orientation given by $\{Z,\eta\}$ above corresponds to the orientation

$$\left\{\frac{\partial}{\partial p}, -\frac{\partial}{\partial q}\right\} \cong \left\{\frac{\partial}{\partial q}, \frac{\partial}{\partial p}\right\}$$

where the contact vector field $\eta = -\frac{\partial}{\partial q} = X_p$ on $s^{-1}(N)$ is pointing *outward* of $\mathbb{C}_{\operatorname{Re}\geq 0}$, when we take the symplectization radial function $s = \log |p|$ at infinity of $T^*\mathbb{R} \cong \mathbb{C}$.

In the remaining section and henceforth, we will always assume that $H \subset M$ is a σ -sectorial hypersurface (Definition 1.2) without further mentioning, unless otherwise mentioned.

2.3. The leaf space is a topological manifold. Let $H \subset M$ be a σ -sectorial hypersurface of a Liouville σ -sector (M, λ) . Equip the leaf space \mathcal{N}_H with the quotient topology induced by the projection $\pi = \pi_H : H \to \mathcal{N}_H$. Before providing a smooth atlas on \mathcal{N}_H , our first order of business is to prove:

Proposition 2.16. The leaf space \mathcal{N}_H is a topological manifold. (In particular, \mathcal{N}_H is second countable and Hausdorff.)

The proof of Proposition 2.16 occupies the rest of this subsection. We consider the given continuous section $\sigma_{\text{ref}} : \mathcal{N}_H \to H$ guaranteed by Definition 1.2. We write

$$F_{\rm ref} = F_{H,\rm ref} := {\rm Image}\,\sigma_{\rm ref} \subset H. \tag{2.7}$$

We choose a contact-type hypersurface $Y \subset M$ far out near infinity so that Y is transversal to the characteristic foliation of H. (Recall that the Liouville vector field Z is tangent to the boundary H near infinity by definition of σ -sectorial hypersurface.) This induces a natural inclusion maps $\iota_{\infty}^{\pm} : F_{\infty}^{\pm} \to H$ and the composition map $\pi_{\infty}^{\pm} : F_{\infty}^{\pm} \to \mathcal{N}_{H}$ given by $\pi_{\infty}^{+} = \pi \circ \iota_{\infty}^{+}$. We denote by

$$\mathrm{Nbhd}^{\pm}(\partial_{\infty}M \cap H) := \mathrm{Nbhd}(F_{\infty}^{\pm})$$

respectively.

Lemma 2.17. There exists a pair of smooth functions

$$h_{\pm} : \operatorname{Nbhd}^{\pm}(\partial_{\infty}M \cap H) \to \mathbb{R}$$
 (2.8)

satisfying $Z[h_{\pm}] = h_{\pm}$ and h_{\pm} are submersions along the characteristic leaves of ∂M .

Proof. By a defining datum of Liouville σ -sectors, Z is tangent to H at infinity.

Since F_{∞}^{\pm} are convex in $\partial_{\infty}M$, there exists a contact vector field η_{\pm} on a neighborhood of F_{∞}^{\pm} in ∂_{∞} such that $\eta_{\pm} \pitchfork F_{\infty}^{\pm}$ respectively. We can choose η_{\pm} so that it points outward of ∂M . In particular, we have

$$dh_{\pm}(X) = d\lambda(X_{h_{\pm}}, X) = d\lambda(\eta_{\pm}, X) \neq 0$$

for any nonzero vector $X \in \ker d\lambda_H$ with $\lambda_H = \iota_H^* \lambda$: We derive the nonvanishing by combining the following:

- $d\lambda(X, TH) = 0$ by definition of ker $d\lambda_H$,
- $\eta \notin TH$ and so $TM = TH + \operatorname{span}\{\eta\}$, and
- nondegeneracy of $d\lambda$.

This finishes the proof.

In particular the level set of h_+ is a smooth submanifold that is transverse to H, We take a symplectization radial function s so that

$$\operatorname{Nbhd}^+(\partial_\infty M) = s^{-1}(0) \times \mathbb{R}_+$$

and $Z = \frac{\partial}{\partial s}$ and the Liouville form

$$\lambda = e^s \pi^* \theta, \quad \theta := \iota_{S_0}^* \lambda$$

where $\pi: S_0 \times \mathbb{R} \to S_0$ with $S_0 := s^{-1}(0)$. Denote

$$F_N^+ := s^{-1}(N) \cap H$$
 (2.9)

for a sufficiently large N > 0. We similarly define F_N^- by considering $N \ll 0$.

We also have the continuous maps

$$\pi_N^{\pm}: F_N^{\pm} \to \mathcal{N}_H \tag{2.10}$$

given as the restrictions of π .

We orient \mathcal{D} so that it points upward in terms of h_+ on F_N^+ , i.e.,

$$dh_+(\mathcal{D}) = d\lambda(X_{h_+}, \mathcal{D}) > 0.$$

Then both Z and D point upward: By suitably adjusting the contact-type hypersurface $h_+^{-1}(N)$ near H we may choose the symplectization end radial function s so that $s = \frac{1}{\alpha} \log h_+ \pm C$ (for any given $0 < \alpha \leq 1$) in a neighborhood of Nbhd⁺($\partial_{\infty} \cap H$) and then clearly we have $Z[h_+] > 0$.

Lemma 2.18. The canonical Z-flow projection provides a natural diffeomorphism between the ideal boundary F_{∞}^+ and F_N^+ .

Proof. First note

$$Z = \frac{1}{\alpha} \operatorname{grad}_g \log h_+$$

for a metric g on M of bounded geometry that is cylindrical on $h_+^{-1}([N,\infty)) \subset H$. Explicitly, the cylindrical metric g has the form

$$g = g_{\infty} \oplus ds^2$$

for a metric g_{∞} on $\partial_{\infty}M$. In particular, the flow of Z—which is the same as the gradient flow of s on $h_{+}^{-1}([N,\infty))$ —defines an obvious diffeomorphism

$$\varphi_N^+: h_+^{-1}([N,\infty)) \cap H \to F_N^+ \times [N,\infty).$$

The gradient flow also induces a diffeomorphism between F_N^+ and $F_\infty^+ \subset F_\infty$. \Box

The same discussion applies to F_{∞}^{-} . We denote the resulting diffeomorphism by $\phi_{N}^{\pm}: F_{N}^{\pm} \to F_{\infty}^{\pm}.$ (2.11)

Proposition 2.19. The map π_N^+ in (2.10) is a homeomorphism.

Proof. The domain of π_N^+ is compact, while the codomain is Hausdorff because of the existence of the continuous section σ_{ref} . Thus it suffices to show that π_N^+ is a bijection. Recall the convexity hypothesis on F_∞ in $\partial_\infty M$ and the choice of orientation on \mathcal{D} above imply $dh_+(\mathcal{D}) > 0$ on F_N^+ for sufficiently large N > 0. Therefore since each leaf is connected, $y_1, y_2 \in F_N^+$ cannot be contained in same leaf of H if $y_1 \neq y_2$. This shows that π_N^+ is a one-one map. (In fact, consider the restriction of π to $h_+^{-1}([N,\infty)) \cap H$. Then we see that π is nothing but the assignment $y \mapsto \ell_y \in \mathcal{N}_H$ where ℓ_y is the leaf of H through $y \in F_N^+$.)

We next show that the map π_N^+ is also surjective. This is the key step in the proof of this proposition. To prove surjectivity, we need to rule out possibilities of the following phenomena:

- appearance of closed leaves,
- appearance of limit cycles or more generally of non-proper leaves.

For this purpose, we consider the identifications (via the aforementioned gradient flow)

$$\begin{aligned} & \operatorname{Nbhd}^+(\partial_{\infty}M) \cap H &\cong F_N^+ \times (N-\delta,\infty), \\ & \operatorname{Nbhd}^-(\partial_{\infty}M) \cap H &\cong F_N^- \times (-\infty, -N+\delta) \end{aligned}$$

for a sufficiently small $\delta > 0$. (Note that these are codimension 0 open subsets of H.) We consider a smooth vector field \widetilde{Z} on H such that

(1) On $F_N^+ \times [N, \infty)$, $\widetilde{Z} \equiv Z_0^+$, and on $F_N^- \times (-\infty, -N]$, $\widetilde{Z} \equiv Z_0^-$ for some smooth vector fields Z_0^{\pm} respectively such that

$$dh_{\pm}(Z_0^{\pm}) = h_{\pm}.$$
 (2.12)

(Without loss of generality, we set $\alpha = 1$ in the rest of the proof for the simplicity of exposition.) In particular, vector fields Z_0^+ (resp. Z_0^-) are forward-complete (resp. back-ward complete).

(2) Z is tangent to the foliation \mathcal{D} and is compatible with the orientation (2.2) of the leaves on

$$H_{\mathrm{mid}} := H \setminus \left(\operatorname{Nbhd}^+(\partial_\infty M) \cup \operatorname{Nbhd}^-(\partial_\infty M) \right).$$

(Indeed, it is the orientability of \mathcal{D} that guarantees the existence of \widetilde{Z} .) In particular, \widetilde{Z} is nowhere vanishing.

Definition 2.20 (Leaf-generating vector field Z' of \mathcal{D}_H). We consider the interpolating smooth vector field

$$Z' = (1 - \chi)Z_0 + \chi \widetilde{Z} \tag{2.13}$$

where $\{1 - \chi, \chi\}$ is a partition of unity on H subordinate to the covering

 $\{(\operatorname{Nbhd}^+(\partial_{\infty}M)\cap H)\cup (\operatorname{Nbhd}^-(\partial_{\infty}M)\cap H), H_{\operatorname{mid}}\}.$

Note that by taking δ to be small enough, we can guarantee that Z' is nowhere vanishing. Since Z_0^{\pm} are complete forward or backward on

$$\operatorname{Nbhd}^+(\partial_{\infty}M) \cup \operatorname{Nbhd}^-(\partial_{\infty}M)$$

respectively, and H_{mid} is compact, the vector field Z' is complete and so defines a global flow on H. Now the proof of surjectivity of π_N^+ relies on the following lemma:

Lemma 2.21. Any trajectory of Z' eventually exits from H_{mid} both forward and backward. Moreover every leaf is a flow orbit of Z' and vice versa.

Proof. It is a standard fact that each leaf is second countable because the manifold M is assumed to be second countable. (This rules out the possibility for a leaf becomes a 'Long line' [Ste95, pp. 71-72].) Note that since Z' is regular, each leaf of H of the characteristic foliation is a flow line of the regular vector field Z'. (See [CC00, Section 2.1].) Furthermore no leaf can be a point. By the condition stated in Definition 1.2 (d), Z' cannot have a nontrivial periodic orbit either. Therefore each flow map from \mathbb{R} to H is one-one and so there is a uniquely defined $T \in \mathbb{R}$ such that $\phi_{Z'}^T(\sigma_{\text{ref}}(\pi(x)) = x$ for each $x \in \partial X$.

Combining this discussion with the aforementioned completeness, we can define a flow map

 $\Psi_{\text{ref}}: H \to F_{\text{ref}} \times \mathbb{R}; \quad \Psi_{\text{ref}}(x) = (\sigma_{\text{ref}}(\pi(x)), T(x))$

where T(x) is the time for $\sigma_{\text{ref}}(\pi(x))$ to reach the point x along the flow of Z'. We define a continuous function $T: M \to \mathbb{R}$ by

T(x) := "the reaching time of the flow of Z' issued at $\sigma_{ref}(\pi(x))$ "

By definition, H is an increasing union

$$H = \bigcup_{N \in \mathbb{N}} T^{-1}(-N, N)$$

of open subset $T^{-1}(-N, N) =: U_N$. Since H_{mid} is compact, there is some $N_0 \in \mathbb{N}$ such that $H_{\text{mid}} \subset U_{N_0}$. Therefore any point $y \in F_{\text{ref}}$ has its forward (resp., backward) flow point $x_+ \in \text{Nbhd}^+(F_{\infty})$ (resp., $x_- \in \text{Nbhd}^-(F_{\infty})$. Once the flow reaches there, it just follows the flow of Z_0^{\pm} forward and backward respectively which escape to infinity. This finishes the proof of Lemma 2.21.

We now wrap up the proof of Proposition 2.19. Lemma 2.21 implies that any trajectory of Z' is an extension of some trajectories Z_0^{\pm} , one from each of $\mathrm{Nbhd}_{\pm}(\partial_{\infty}M)$. On the other hand, if $\phi_{Z'}(t) \in \mathrm{Nbhd}_{\pm}(\partial_{\infty}M)$, we have $ds_{\pm}(Z') = ds_{\pm}(Z_0^{\pm}) = 1$. In particular we have Z'[T] = 1 and

$$T(x) = s_{\pm}(x) + C'_{\pm},$$

on each of Nbhd[±]($\partial_{\infty} M$) respectively for some constant $C'_{\pm} = C'_{\pm}(\mathcal{D}_x)$ depending only on the leaf \mathcal{D}_x containing x, and hence $T(x) \to \pm \infty$ if $x \to \partial_{\infty} M \cap H$ along the leaf $\pi(x(\ell))$.

This proves that the function $x \mapsto T(x)$ restricts to a homeomorphism from the leaf $\pi(x)$ to \mathbb{R} . This in particular implies that any leaf of \mathcal{N}_H is an extension of the asymptotic rays of F_N^+ and so the map π_N^+ is also surjective. \Box

Lemma 2.22. \mathcal{N}_H equipped with the quotient topology is homeomorphic to F_{ref} .

Proof. As we mentioned above, any leaf of \mathcal{N}_H is an extension of the asymptotic rays of F_N^+ . Therefore by Proposition 2.19 we have a homeomorphism

$$\Psi_{\text{ref}}: H \to F_{\text{ref}} \times \mathbb{R}, \qquad \Psi_{\text{ref}}(x) = (\sigma_{\text{ref}}(\pi(x)), T(x)) \tag{2.14}$$

for a continuous map $T : H \to \mathbb{R}$. In particular \mathcal{N}_M equipped with the quotient topology of \mathcal{N}_H is homeomorphic to F_{ref} .

Proposition 2.23. F_{ref} with the subspace topology of H is Hausdorff and locally Euclidean (and in particular, locally compact).

Proof. Since the function $T : H \to \mathbb{R}$ is continuous and $F_{\text{ref}} = T^{-1}(0)$, F_{ref} is a closed subset of a smooth manifold H. In particular F_{ref} with the subspace topology of H is Hausdorff.

Furthermore since Z'[T] > 0, T is monotonically increasing. To see the locally Euclidean property of F_{ref} , let $x_0 \in F_{\text{ref}}$ be any given point. We have only to note that (2.14) induces a homeomorphism

$$U/ \sim \to F_{\mathrm{ref}} \cap U$$

for a sufficiently small foliation chart U containing x_0 where \sim is the orbit equivalence with respect to Z'. Since U/\sim is homeomorphic to \mathbb{R}^{2n-1} , so is $F_{\text{ref}} \cap U$. This proves that $F_{\text{ref}} \cap U$ is locally Euclidean.

Wrap up of the proof of Proposition 2.16. We have only to combine Proposition 2.23 and Lemma 2.22. $\hfill \Box$

The following corollary of the above proof will be useful for the study of smooth and symplectic structures of the leaf space.

Corollary 2.24 (Section σ_N^+). The definition (2.9) defines another section σ_N^+ defined by

$$\sigma_N^+(\ell) = \ell \cap F_N^+ \subset \operatorname{Nbhd}(\partial_\infty M \cap \partial M) \tag{2.15}$$

for $\ell \in \mathcal{N}_H$.

2.4. Smooth structure on the leaf space. The goal of this section is to prove the first item of Theorem 1.4. We start with the following proposition.

Proposition 2.25. The leaf space \mathcal{N}_H carries a canonical smooth manifold structure such that

- (1) $\pi: H \to \mathcal{N}_H$ is a smooth submersion, and
- (2) there is a smooth diffeomorphism $\Psi : H \to \mathcal{N}_H \times \mathbb{R}$ which makes the following diagram commute



Actually, when the leaf space is Hausdorff and locally Euclidean, the well-known construction of coisotropic reduction (or symplectic reduction) applies to prove existence of smooth structure and the symplectic structure on the leaf space. (See [AM78] for example.) Since we also need to construct the map Ψ and will also use the details of the proof later, we provide the full details of the existence proofs of both structures below along the way for readers' convenience.

We follow the standard notation of [CC00] in our discussion of foliations. It follows from a well-known result in foliation theory that the foliation \mathcal{F} is determined by its holonomy cocycle $\gamma = \{\gamma_{\alpha\beta}\}_{\alpha,\beta\in\mathfrak{U}}$ with

$$\gamma_{\alpha\beta}: y_{\beta}(U_{\alpha} \cap U_{\beta}) \to y_{\alpha}(U_{\alpha} \cap U_{\beta}).$$

arising from the transverse coordinate map $y_{\alpha}: U_{\alpha} \to \mathbb{F}^{2n-2} = \mathbb{R}^{2n-2}$ or \mathbb{H}^{2n} .

Each y_{α} is a submersion and $\gamma_{\alpha\beta}$ is given by $y_{\alpha} = y_{\alpha}(y_{\beta})$ in coordinates. (See e.g., [CC00, Definition 1.2.12].) Furthermore for the null foliation \mathcal{F} of the coisotropic submanifold H, we can choose a foliated chart $\mathcal{U} = \{(U_{\alpha}, \varphi_{\alpha})\}_{\alpha \in \mathfrak{U}}$ so that the associated cocycle elements $\gamma_{\alpha\beta}$ become symplectic, i.e., the foliation \mathcal{F} carries a transverse symplectic structure. We refer readers to the proof of Proposition 2.25 below for the details.

Remark 2.26. When *H* has corners, the foliated chart $B = B_{\tau} \times B_{\uparrow}$ means that the *tangential factor* B_{τ} of the foliated chart has no boundary but the *transverse factor* B_{\uparrow} has a boundary. (See e.g., [CC00, Definition 1.1.18] for the definition.)

Proof of Proposition 2.25. We will show that the above holonomy cocycle naturally descends to a smooth atlas on \mathcal{N}_H under the defining condition of σ -sectorial hypersurface above, especially in the presence of a continuous section of the projection $\pi_H: H \to \mathcal{N}_H$.

For this purpose, we consider a coherent regular foliated atlas $\{\varphi_{\alpha} : U_{\alpha} \rightarrow \mathbb{R}^{2n-1}\}$, and its associated foliation cocycle $\gamma = \{\gamma_{\alpha\beta}\}$ (see e.g., [CC00, Section 1.2.A]). Let $\sigma_N^+ : \mathcal{N}_H \rightarrow H$ be the section given by (2.15). We take the subcollection $\{U_{\alpha'}\}$ that covers F_N^+ and such that $U_{\alpha'} \cap F_N^+ \neq \emptyset$. Furthermore since $F_N^+ = h^{-1}(N)$ is a level set of the function h in (2.8) and we

Furthermore since $F_N^+ = h^{-1}(N)$ is a level set of the function h in (2.8) and we may assume that each plaque in a small $Nbhd(F_N^+)$ is the gradient trajectory of h: By considering a refinement $\{U_{\alpha'}\}$ of the given covering, we can choose a collection of foliated charts $\varphi_{\alpha'}: U_{\alpha'} \to \mathbb{R} \times \mathbb{R}^{2n-2}$ of the form

$$\varphi_{\alpha'} = (h, y_1, \cdots, y_{2n-2})$$

whose transversal coordinates (y_1, \cdots, y_{2n-2}) satisfy

$$dy_i(Z) = 0.$$

We take a maximal such collection of F_N^+ which we denote by

$$\mathcal{O}' = \{(\varphi_{\alpha'}, U_{\alpha'})\}. \tag{2.17}$$

By the definition of transverse coordinates (y_1, \dots, y_{2n-2}) of the foliated chart, it follows that the collection thereof defines a *smooth* atlas of \mathcal{N}_H . We write the resulting atlas of \mathcal{N}_H by

$$[\mathfrak{O}'] := \{ [\varphi_{\alpha'}] : [U_{\alpha'}] \to \mathbb{R}^{2n-2} \}.$$

Lemma 2.27. The projection map $\pi: H \to \mathcal{N}_H$ is a smooth submersion.

Proof. To show smoothness of π , we will show that for any smooth function $f : \mathcal{N}_H \to \mathbb{R}$ the composition $f \circ \pi$ is smooth.

For this purpose, at any point x, we consider the foliated chart $\varphi_{\alpha} : U_{\alpha} \to \mathbb{R}^{2n-1}$ of the form

$$(h', y_1, \dots, y_{2n-2})$$
 (2.18)

whose transversal coordinate (y_1, \ldots, y_{2n-2}) satisfies

$$dh'(Z') \equiv 1, \quad dy_i(Z') = 0, \quad i = 1..., 2n-2.$$
 (2.19)

Let $f : \mathcal{N}_H \to \mathbb{R}$ be any smooth function on \mathcal{N}_H . With respect to the aforementioned foliated atlas of H, we will show that $f \circ \pi$ is smooth at every point $x \in H$.

If x is contained in $U_{\alpha'}$, we have

$$(f \circ \pi) \circ (\varphi_{\alpha'})^{-1}(h', y_1, \cdots, y_{2n-2}) = f \circ [\varphi_{\alpha'}]^{-1}(y_1, \cdots, y_{2n-2})$$

The right hand side is smooth in the variables y_1, \dots, y_{2n-2} by the hypothesis on f, and does not depend on h'-variable. This in particular implies that the left hand map $(f \circ \pi) \circ (\varphi_{\alpha'})^{-1}$ is smooth at x.

Otherwise, let $(\varphi_{\beta}, U_{\beta})$ be a foliation chart at x. We take a flow map $\phi_{Z'}^T$ satisfying $y := \phi_{Z'}^T(x) \in U'_{\beta}$ for some chart $(\varphi_{\beta'}, U_{\beta'}) \in \mathcal{O}'$ at y given by

$$(U_{\beta'} = \phi_{Z'}^T (U_{\beta}), \quad \varphi_{\beta'} = \varphi_{\beta} \circ (\phi_{Z'}^T)^{-1})$$

which is contained in O' by the maximality of the collection O'.

Therefore the map $(f \circ \pi) \circ \varphi_{\beta'}^{-1}$ is smooth at $y = \phi_{Z'}^T(x) \in U_{\beta'}$. We note

$$f \circ \pi = \left((f \circ \pi) \circ \varphi_{\beta'}^{-1} \right) \circ \left((\varphi_{\beta'}^{-1} \circ \phi_{Z'}^T |_{U_\beta}) \right)$$

which is a composition of two smooth maps and so smooth at x. This implies $f \circ \pi$ is smooth at x again. This finishes the proof of smoothness $\pi \circ f$ for all smooth function $f : \mathcal{N}_H \to \mathbb{R}$. This proves that π is smooth.

Submersivity of π is obvious by the above construction.

Now a closer examination of the above proof also shows the following whose proof we leave to the readers.

Corollary 2.28. Let \mathcal{N}_H be equipped with this smooth structure.

(1) The smooth structure on \mathcal{N}_H constructed above does not depend on the choice of the section (2.15), rewritten as

$$\sigma_N^+(\ell) := \ell \cap h^{-1}(N), \tag{2.20}$$

does not depend on the choice of such σ_N^+ .

- (2) The map π_N^+ in (2.10) is a smooth submersion with respect to the smooth structure on \mathcal{N}_H with which we have just equipped \mathcal{N}_H .
- (3) The map σ_N^+ is a smooth section of the submersion π_N^+ .

2.5. Symplectic structure on the leaf space. Now we turn to the construction of symplectic structure. We fix the section $\sigma_N^+ : \mathcal{N}_H \to H$ constructed before.

When we choose the above used coherent atlas, we can choose them so that the associated cocycle $\gamma_{\alpha\beta}$ becomes symplectic by requiring the chart $(U_{\alpha}, \varphi_{\alpha})$ also to satisfy the defining equation

$$y^*_{\alpha}\omega_0 = \iota^*_H\omega, \quad \omega = d\lambda$$
 (2.21)

of the general coisotropic reduction (see [AM78, Theorem 5.3.23] for example) where $\iota_H : H \to M$ is the inclusion map and ω_0 is the standard symplectic from on \mathbb{R}^{2n-2} . (See also [Got82], [OP05].) By using such a foliated chart satisfying (2.21), the associated holonomy cycles define symplectic atlas and so a symplectic structure on \mathcal{N}_H , when the holonomy is trivial as in our case where we assume the presence of smooth section. This will then finish construction of reduced symplectic structures on \mathcal{N}_H . (We refer to [OP05, Section 5] for a detailed discussion on the construction of transverse symplectic structure for the null foliation of general coisotropic submanifolds.)

Now it remains to construct a diffeomorphism $\Psi^{\sigma} : H \to \mathcal{N}_H \times \mathbb{R}$ which makes the diagram (2.16) commute. For this, it follows from the above discussions that we can define another map

$$\Psi: H \to \mathcal{N}_H \times \mathbb{R}; \quad \Psi(x) = (\pi(x), t(x)) \tag{2.22}$$

by replacing the original defining section σ_{ref} by the new *smooth* section σ_N^+ defined by (2.20): Here we put

$$t(x) :=$$
 the unique time determined by $\phi_{Z'}^{t(x)}(\sigma_N^+(\pi(x))) = x$ (2.23)

which is clearly a smooth function. Furthermore, we have shown $\pi_{\text{ref}} = \pi_N^+$ which also shows that π_{ref} is smooth with respect to the smooth structure just given to \mathcal{N}_H .

Its inverse map $\Phi : \mathcal{N}_H \times \mathbb{R} \to H$ is given by

$$\Phi(\ell, t) = \phi_{Z'}^t(\sigma_N^+(\ell))$$

which is obviously smooth. This finishes the proof of Proposition 2.25.

An immediate corollary of the above proof is the following construction of the defining function $I : \partial M \to \mathbb{R}$ appearing in the definition of Liouville sectors in [GPS17].

Corollary 2.29. Let t be the function used in the definition of the map (2.22). For each $\alpha > 0$, we define the function $I : \partial M \to \mathbb{R}$ by

$$I(x) = \pm e^{\alpha t(x)}.$$

Then we have $Z[I] = \alpha I$ on $Nbhd(\partial_{\infty} M) \cap H$.

This proves Proposition 1.5, i.e., that any Liouville σ -sector is a Liouville sector in the sense of [GPS17].

Remark 2.30. On the other hand, the converse is almost a tautological statement in that [GPS17, Lemma 2.5] shows that any of their three defining conditions given in [GPS17, Definition 2.4] is equivalent to the condition

• There exists a diffeomorphism $\Psi : H \to F \times \mathbb{R}$ making (2.16) commute Once this is in our disposal, Ψ induces a diffeomorphism $[\Psi] : \mathcal{N}_H \to F$. Therefore we can choose a continuous section $\sigma_{\text{ref}} : \mathcal{N}_H \to H$ required for the definition of σ -sectorial hypersurface to be

$$\sigma_{\mathrm{ref}}(\ell) := [\Psi]^{-1}(\ell), \quad \ell \in \mathcal{N}_H.$$

Now we wrap up the proof of Theorem 1.4.

Proof of Theorem 1.4. Proposition 2.25 and Lemma 2.27 prove (1) of the theorem. Let

$$\iota_F = \iota_{F_N} : F_N \to H$$

be the inclusion map and consider the diagram

$$\begin{array}{c} H \xrightarrow{\iota_H} M \\ \downarrow_{\pi_H} \\ \aleph_H \end{array}$$

Then the reduced symplectic form $\omega_{\mathcal{N}_H}$ is characterized by

$$(\pi_H)^* \omega_{\mathcal{N}_H} = (\iota_H)^* \omega_H, \quad \omega_H := \iota_H^*(d\lambda)$$
(2.24)

as an example of coisotropic reduction. This proves (2).

Now we take $\sigma = \sigma_N^+$ and $F = F_N^+ := \text{Image } \sigma_N^+$. By pulling back the two form $(\iota_H)^* \omega_H$ by the inclusion $F \hookrightarrow H$, we have obtained a two-form $(\iota_{F_N})^* \omega$ on F_N

which is symplectic for any sufficiently large N > 0 by the convexity hypothesis of F_{∞}^+ in $\partial_{\infty} M$. This finishes the proof of (4). Finally setting $I = \pm e^{\alpha t(x)}$, we have proved Proposition 1.5.

It is useful for the later study of intrinsic characterization of Liouville sectors with corners to keep in mind the following corollary of the above proof.

Corollary 2.31. The line bundle $\mathcal{D}_H \to H$ of characteristic distribution is trivial for any σ -sectorial hypersurface.

Of course this is a tautological property with the original definition of Liouville sectors from [GPS17].

2.6. Induced Liouville structure on the leaf space. Finally we extract some consequences on the above constructed symplectic structure on \mathcal{N}_H derived from the given property of the characteristic foliation \mathcal{D} at infinity. By composing the two symplectic diffeomorphisms

$$\mathfrak{N}_H \xrightarrow{\sigma_N^+} F_N^+ \xrightarrow{\phi_N} F_\infty^+ \hookrightarrow \partial_\infty M$$

we have a smooth map σ_{∞}^+ : $\mathcal{N}_H \to \partial_{\infty} M$ which is a diffeomorphism onto the convex hypersurface (F_{∞}^+, ω^+) of the contact manifold $(\partial_{\infty} M, \xi)$. By the convexity hypothesis on F_{∞} , we have a contact vector field ν on $\partial_{\infty} M$ that is transverse to F_{∞}^+ .

Lemma 2.32. The symplectic manifold $(\mathcal{N}_H, \omega_{\mathcal{N}_H})$ is exact.

Proof. Note that the symplectic form on F_N^+ is nothing but the exact symplectic form given by $d\lambda_N^+$ where

$$\lambda_N^+ = (\iota_N^+)^* \lambda$$

for the inclusion map $\iota_N^+ : F_N^+ \hookrightarrow H \hookrightarrow (M, \lambda)$. Therefore it follows from (2.24) and $\pi_F^* \circ \sigma_N^+ = id_{\mathcal{N}_H}$

$$\omega_{\mathcal{N}_H} = (\pi_{F_N^+} \circ \sigma_N^+)^* \omega_{\mathcal{N}_H} = (\sigma_N^+)^* (\pi_{F_N^+}^* \omega_{\mathcal{N}_H})$$
$$= (\sigma_N^+)^* (\iota_{F_N^+}^* d\lambda) = (\sigma_N^+)^* d\lambda_N^+ = d((\sigma_N^+)^* \lambda_N^+)$$

which proves exactness of $\omega_{\mathcal{N}_H}$: Here the third equality follows from the defining equation (2.24) and the equalities

$$\pi_{F_N^+} = \pi_H \circ \iota_{F_N^+H}, \quad \iota_{F_N^+} = \iota_H \circ \iota_{F_N^+H}.$$

We next prove that the one-form $(\sigma_N^+)^* \lambda_N^+$ on \mathcal{N}_H appearing in the above proof does not depend on N.

Lemma 2.33. We have

$$(\sigma_N^+)^*\lambda_N^+ = (\sigma_{N'}^+)^*\lambda_{N'}^+$$

on \mathcal{N}_H for all $N, N' \geq 0$. We denote this common one-form by $\lambda_{\mathcal{N}_H}$.

Proof. The flow map $\phi_{NN'}$ of the vector field Z' (2.13) between F_N^+ and $F_{N'}^+$ is the same as

$$(\phi_{N'}^+)^{-1} \circ \phi_N^+ : F_N^+ \to F_{N'}^+$$

Therefore it intertwines two maps ϕ_N^+ and $\phi_{N'}^+$ and satisfies

$$\phi_{NN'}^+(\lambda_{N'}^+) = \lambda_N^+$$

for all large $N, N' \geq 0$. This finishes the proof.

Definition 2.34 (Reduced Liouville structure). We call the primitive $\lambda_{\mathcal{N}_H}$ of $\omega_{\mathcal{N}_{\partial H}}$ defined as above the canonical Liouville structure on $(\mathcal{N}_H, \omega_{\mathcal{N}_H})$.

Remark 2.35. We remark that while we have used the existence of section σ_{ref} to equip \mathcal{N}_H with a smooth structure for which there exists a diffeomorphism $\Psi : H \to \mathcal{N}_H \times \mathbb{R}$, construction of the symplectic structure does not depend on the choice of the section, but depends only on the Liouville geometry of the neighborhood $\mathcal{N}bhd(\partial_{\infty}M \cap \partial M)$.

3. Geometry of clean coisotropic collections

Recall that [GPS18] imposes the following restriction on the boundary strata when studying Liouville sectors:

Definition 3.1 (Definition 9.2 & Lemma 9.4 & Definition 9.14 [GPS18]). A sectorial collection is a collection of m hypersurfaces $H_1, \ldots, H_m \subset M$, cylindrical at infinity, such that:

- (S1) The H_i cleanly intersect,
- (S2) All pairwise intersections $H_i \cap H_j$ are coisotropic, and
- (S3) There exist functions I_i : Nbhd $(\partial M) \to \mathbb{R}$, linear near infinity, satisfying the following on the characteristic foliations \mathcal{D}_i of H_i :

 $dI_i|_{\mathcal{D}_i} \neq 0, \, dI_i|_{\mathcal{D}_i} = 0 \quad \text{for } i \neq j, \quad \{I_i, I_j\} = 0.$ (3.1)

A Liouville sector (M, λ) with corners is a Liouville manifold-with-corners whose boundary strata form a sectorial collection.

We will introduce another definition of sectorial collection by replacing Condition (S3) in the spirit of Definition 2.4.

For this purpose, we need some preparations. We start with introducing the following definition

Definition 3.2 (Clean coisotropic collection). Let (M, λ) be a Liouville manifold with boundary and corners. Let $H_1, \ldots, H_m \subset M$ be a collection hypersurfaces cylindrical at infinity, that satisfies Conditions (S1), (S2) of Definition 3.1.

In the remaining section, we first study the underlying geometry and prove a general structure theorem of such a collection. In the next section, based on the theorem, we will provide an intrinsic characterization of the sectorial collection and Liouville sectors with corners above purely in terms of geometry of coisotropic submanifolds. We call the resulting structure the structure of *Liouville* σ -sectors with corners.

3.1. Gotay's coisotropic embedding theorem of presymplectic manifolds. For a finer study of the neighborhood structure of the sectorial corner C, we first recall below some basic properties of the coisotropic submanifolds and the coisotropic embedding theorem of Gotay [Got82]. See also [Wei79], [OP05] for relevant material on the geometry of coisotropic submanifolds. We will mostly adopt the notations used in [Got82], [OP05, Section 3].

Let (Y, ω_Y) be any presymplectic manifold. The null distribution on Y is the vector bundle

 $E := (TY)^{\omega_Y} \subset TY, \quad E_y = \ker \omega_Y|_y.$

This distribution is integrable since ω_Y is closed. We call the corresponding foliation the *null foliation* on Y and denote it by

 $\mathcal{F} = \mathcal{F}_{\mathbf{V}}.$

(Then E is nothing but the total space of the foliation tangent bundle $T\mathcal{F}$.) We now consider the dual bundle $\pi: E^* \to Y$ which is the foliation cotangent bundle

$$E^* = T^* \mathcal{F}.$$

The tangent bundle TE^* of the total space E^* has its restriction to the zero section $Y \hookrightarrow E^*$; this restriction carries a canonical decomposition

$$TE^*|_Y \cong TY \oplus E^*$$

Example 3.3. A typical example of a presymplectic manifold is given by

$$(Y, \omega_Y) = (H, \omega_H), \quad \omega_H := \iota_H^* \omega$$

arising from any coisotropic submanifold $H \subset^{\iota_H} (X, \omega)$. Then $E = \mathcal{D}_H$, the null distribution of (H, ω_H) . It is easy to check that the isomorphism

$$TX \to T^*X$$

maps TY^{ω} to the conormal $N^*Y \subset T^*X$, and induces an isomorphism between $NY = (TX)|_Y/TY$ and E^* .

Gotay [Got82] takes a transverse symplectic subbundle G of TY and associates to each splitting

$$\Gamma: \quad TY = G \oplus E, \quad E = T\mathcal{F} \tag{3.2}$$

the zero section map

 $\Phi_{\Gamma}: Y \hookrightarrow T^* \mathcal{F} = E^*$

as a coisotropic embedding with respect to a 'canonical' two-form ω_{E^*} on E^* which restricts to a symplectic form on a neighborhood of the zero section of E^* such that

$$\omega_Y = \Phi_{\Gamma}^* \omega_{E^*}.$$

Remark 3.4. When $\omega_Y = 0$, Gotay's embedding theorem reduces to the wellknown Weinstein's neighborhood theorem of Lagrangian submanifolds L in which case $E^* = T^*L$ with Y = L.

We now describe the last symplectic form closely following [Got82].

We denote the aforementioned neighborhood by

$$V \subset T^* \mathcal{F} = E^*.$$

Using the splitting Γ , which may be regarded as an 'Ehresmann connection' of the 'fibration'

$$T\mathcal{F} \to Y \to \mathcal{N}_Y,$$

we can explicitly write down a symplectic form ω_{E^*} as follows.

First note that as a vector bundle, we have a natural splitting

$$TE^*|_Y \cong TY \oplus E^* \cong G \oplus E \oplus T^*\mathcal{P}$$

on Y, which can be extended to a neighborhood V of the zero section $Y \subset E^*$ via the 'connection of the fibration' $T^*\mathcal{F} \to Y$. (We refer readers to [OP05] for a complete discussion on this.)

We denote

$$p_{\Gamma}: TY \to T\mathcal{F}$$

the (fiberwise) projection to $E = T\mathcal{F}$ over Y with respect to the splitting (3.2). We have the bundle map

$$TE^* \xrightarrow{T\pi} TY \xrightarrow{p_{\Gamma}} E$$

over Y.

Definition 3.5 (Canonical one-form θ_{Γ} on E^*). Let $\zeta \in E^*$ and $\xi \in T_{\zeta}E^*$. We define the one form θ_{Γ} on E^* whose value is to be the linear functional

$$\theta_{\Gamma}|_{\zeta} \in T^*_{\zeta}E^*$$

at ζ that is determined by its value

$$\theta_{\Gamma}|_{\zeta}(\xi) := \zeta(p_{\Gamma} \circ T\pi(\xi)) \tag{3.3}$$

against $\xi \in T_{\zeta}(T^*\mathcal{F}).$

(We remark that this is reduced to the canonical Liouville one-form θ on the cotangent bundle T^*L for the case of Lagrangian submanifold L in which case $\omega_Y = 0$ and the splitting is trivial and not needed.)

Then we define the closed (indeed exact) two form on $E^* = T^* \mathcal{F}$ by

 $-d\theta_{\Gamma}$.

Together with the pull-back form $\pi^* \omega_Y$, we consider the closed two-form $\omega_{E^*,\Gamma}$ defined by

$$\omega_{E^*,\Gamma} := \pi^* \omega_Y - d\theta_\Gamma \tag{3.4}$$

on $E^* = T^* \mathcal{F}$. It is easy to see that $\omega_{E^*,\Gamma}$ is non-degenerate in a neighborhood $V \subset E^*$ of the zero section (See the coordinate expression [OP05, Equation (6.6)] of $d\theta_{\Gamma}$ and ω_V .)

Definition 3.6 (Gotay's symplectic form [Got82]). We denote the restriction of $\omega_{E^*,\Gamma}$ to V by ω_V , i.e.,

$$\omega_V := (\pi^* \omega_Y - d\theta_\Gamma)|_V.$$

We call this two-form Gotay's symplectic form on $V \subset E^*$.

The following theorem ends the description of Gotay's normal form for the neighborhood of a coisotropic submanifold $C \subset (M, \omega)$ of any symplectic manifold (M, ω) as a neighborhood V of the zero section of $T^*\mathcal{F}_C$ of its null foliation \mathcal{F}_C on C equipped with the symplectic form.

Theorem 3.7 (See [Got82, OP05]). Let $Y \subset (X, \omega_X)$ be any coisotropic submanifold. Fix a splitting Γ in (3.2). Then there is a neighborhood $Nbhd(Y) := U \subset X$ and a diffeomorphism

$$\Phi_{\Gamma}: U \to V \subset E^*$$

such that the following hold:

- (1) $\omega_X = \Phi_{\Gamma}^* \omega_{E^*,\Gamma}$ on $U \subset X$.
- (2) For two different choices, Γ and Γ' , of splitting of TY, the associated two forms $\omega_{E^*,\Gamma}$ and $\omega_{E^*,\Gamma'}$ are diffeomorphic relative to the zero section $Y \subset E^*$, on a possibly smaller neighborhood $V' \subset E^*$ of Y.

Proof. The first statement is proved in [Got82]. Statement (2) is then proved in [OP05, Theorem 10.1]. \Box

We have the natural projection map

$$\widetilde{\pi}_Y : \operatorname{Nbhd}(Y) \to Y$$
 (3.5)

defined by

$$\widetilde{\pi}_Y := \pi_{E^*} \circ \Phi_\Gamma \circ \iota_Y, \tag{3.6}$$

for the inclusion map $\iota_Y : Y \hookrightarrow \operatorname{Nbhd}(Y) =: U \subset X$, which is induced by restricting the canonical projection $E^* \to Y$ to the neighborhood $V \subset E^*$ of the zero section Y. In particular, we have

$$\ker d_x \pi_Y = E_x = \mathcal{D}_Y|_x.$$

3.2. Structure of the null foliation of σ -sectorial corners. We apply the discussion in the previous subsection to general clean coisotropic collection

$$\{H_1,\cdots,H_m\}$$

For any given subset $I \subset \{1, \dots, m\}$, we denote

$$H_I = \bigcap_{i \in I} H_i$$

and $\pi_{H_I}: H_I \to \mathcal{N}_{H_I}$ be the canonical projection. We also denote the full intersection by

$$C = \bigcap_{i=1}^{m} H_i.$$

Furthermore, by the clean intersection property of the coisotropic collection, we can choose the collection $\{\sigma_{C,1}, \ldots, \sigma_{C,m}\}$ to have the complete intersection property in that their images form a collection of clean intersection. More precisely, we fix the following choice of smooth sections for a finer study of the neighborhood structure of further constructions we will perform

Choice 3.8 (Choice of sections $\sigma_i : \mathcal{N}_{H_i} \to H_i$). For each $i = 1, \ldots, m$, we choose a smooth section

$$\sigma_i: \mathcal{N}_{H_i} \to H_i$$

for each i = 1, ..., n. Denote the set of sections $\sigma_i : \mathcal{N}_{H_i} \to H_i$ by

$$\sigma = \{\sigma_1, \dots, \sigma_m\}. \tag{3.7}$$

Recall from Section 2 that for each i a choice of smooth section

$$\sigma_i: \mathcal{N}_{H_i} \to H_i$$

provides the trivialization map

$$\Psi_i^{\sigma_i}: H_i \to \mathcal{N}_{H_i} \times \mathbb{R}, \quad \Psi_i^{\sigma_i}(x) = (\pi_{H_i}(x), t_i^{\sigma_i}(x))$$

given in (2.22). We choose each σ_i to be $\sigma_i = \sigma_{H_i}^+$ as defined in (2.15). For the given choice of $\sigma = \{\sigma_1, \ldots, \sigma_m\}$, we collectively write

$$\Psi_i^{\sigma} := \Psi_i^{\sigma_i}, \quad i = 1, \dots, m.$$
(3.8)

The following theorem is the generalization of Theorem 1.4 whose proof also extends the one used in Section 2 to the case with corners. The main task for this extension is to establish compatibility of the null foliations of various coisotropic intersections arising from taking a sub-collection $I \subset \{1, \ldots, m\}$: This compatibility

condition and construction of relevant strata is in the same spirit as the combinatorial construction of a toric variety out of its associated fan. (See [Ful93] for example.)

Theorem 3.9. Let (M, λ) be a Liouville σ -sector with corners, and let Z be the Liouville vector field of (M, λ) . Let

$$\sigma = \{\sigma_1, \cdots, \sigma_m\}$$

be a collection of sections $\sigma_i : \mathbb{N}_{H_i} \to H_i$ for $i = 1, \ldots, m$. Then the leaf space \mathbb{N}_C carries a canonical structure $\lambda_{\mathbb{N}_C}$ of a Liouville manifold with boundary and corners.

We also define the function $t_i^{C,\sigma}:C\to\mathbb{R}$ to be the restriction

$$t_i^{C,\sigma} = t_i^{\sigma_i}|_C \tag{3.9}$$

where $t_i^{\sigma_i}$ is the function appearing in (2.22). The collection $\sigma = \{\sigma_i\}$ also induces a surjective map $\Psi_C : C \to \mathcal{N}_C \times \mathbb{R}^m$,

$$\Psi_C^{\sigma}(x) := \left(\pi_C(x), \left(t_1^{C,\sigma}(x), \dots, t_m^{C,\sigma}(x)\right)\right)$$
(3.10)

which is also smooth with respect to the induced smooth structure on \mathcal{N}_C . (The functions $t_i^{C,\sigma}$ correspond to t_i appearing in [Arn88, Section 49] in the discussion following below.)

Proposition 3.10. There is an \mathbb{R}^m -action on C that is free, proper and discontinuous such that C is foliated by the \mathbb{R}^m -orbits. In particular the map

$$\Psi_C^{\sigma}: C \to \mathcal{N}_C \times \mathbb{R}^{d}$$

is an \mathbb{R}^m -equivariant diffeomorphism with respect to the \mathbb{R}^m -action on C and that of linear translations on \mathbb{R}^m .

Proof. Let (s_1, \ldots, s_m) be the standard coordinates of \mathbb{R}^m . We set

$$Z_i := (\Psi_C^{\sigma})^* \left(\vec{0}_{\mathcal{N}_C} \oplus \frac{\partial}{\partial s_i} \right).$$
(3.11)

Then $Z_i \in \mathcal{D}_C$, and $[Z_i, Z_j] = 0$ since $[\frac{\partial}{\partial s_i}, \frac{\partial}{\partial s_j}] = 0$. On C, we also have

$$t_j^{C,\sigma}(Z_i) = d(s_j \circ \Psi_C^{\sigma}) \left((\Psi_C^{\sigma})^* \left(\vec{0}_{\mathcal{N}_C} \oplus \frac{\partial}{\partial s_i} \right) \right) = ds_j(\frac{\partial}{\partial s_i}) = \delta_{ij}.$$

In particular Z_i is tangent to all level sets of $t_i^{C,\sigma}$ with $j \neq i$, and is transversal to the level sets of $t_i^{C,\sigma}$ for each *i*.

The so-constructed global frame $\{Z_1, \dots, Z_m\}$ of TC on C are commuting vector fields. Therefore we have an \mathbb{R}^m -action on C induced by the flows of commuting vector fields $\{Z_1, \dots, Z_m\}$.

Lemma 3.11. This \mathbb{R}^m -action is also proper and discontinuous. In particular, its isotropy subgroup is a discrete subgroup of \mathbb{R}^m .

Proof. The Liouville vector field Z is tangent to every H_i at infinity. Since Z is tangent to H_i for all i at infinity, the flag

$$H_1 \cap \dots \cap H_m \subset H_1 \cap \dots \cap H_{m-1} \subset \dots \subset H_1$$

is Z-invariant at infinity, and in particular we have

 $Z\in TC$

at infinity of C. Since Z[s] = 1, Z is also transversal to $s^{-1}(r)$ for all sufficiently large r > 0. Therefore the \mathbb{R}^m -action induces a free \mathbb{R}^m/\mathbb{R} -action on the set $\partial_{\infty}C =$ $\partial_{\infty}M \cap C$ of asymptotic Liouville rays tangent to C. Since the latter set is compact, it follows that the \mathbb{R}^m/\mathbb{R} -action is proper and discontinuous. Since the flow of Zor the \mathbb{R} -action induced by Z moves the level of s by 1 as time varies by 1, we conclude that the \mathbb{R}^m -action on C is proper and discontinuous.

Once the action is proved to be proper and discontinuous, the second statement of the lemma follows e.g. from the proof in [Arn88, Section 49, Lemma 3], to which we refer. This finishes the proof. \Box

With Lemma 3.11 in our disposal, the standard argument in the construction of action-angle coordinates proves that each orbit of the \mathbb{R}^m -action is homeomorphic to $\mathbb{R}^{n_1} \times T^{n_2}$ for some n_1 , n_2 with $n_1 + n_2 = n$. (See [Arn88, Section 49, Lemma 3] and its proof.) Since we are assuming that fibers are contractible, we immediate conclude the following.

Corollary 3.12. Suppose $\pi_C : C \to \mathcal{N}_C$ has contractible fibers. Then

- (1) The \mathbb{R}^m -action is free and its fiber is naturally diffeomorphic to \mathbb{R}^m , i.e., it is a principle \mathbb{R}^m bundle over \mathcal{N}_C .
- (2) The map Ψ is an \mathbb{R}^m -equivariant diffeomorphism with respect to the translations of \mathbb{R}^m .

The inverse of Ψ_C^{σ} denoted by

$$\Phi_C^{\sigma}: \mathcal{N}_C \times \mathbb{R}^m \to C \tag{3.12}$$

is also easy to explicitly write down as follows. First we note

$$t_i^{C,\sigma}(\sigma_{C,i}(\pi_C(x))) = 0$$

for all $i = 1, \ldots, m$ by the definitions of $\sigma_{C,i}$ and $t_i^{C,\sigma}$. Now let a point

$$(\ell, (t_1, \ldots, t_m)) \in \mathbb{N}_C \times \mathbb{R}^m$$

be given. Then there is a unique point $x \in C$ satisfying

$$\begin{cases} \pi_C(x) = \ell \\ x = \bigcap_{i=1}^n (t_i^{C,\sigma})^{-1}(t_i). \end{cases}$$
(3.13)

(See (2.23) for the definition of $t_i^{C,\sigma}$ and (2.13) for the definition of the vector field Z'_i respectively.) Then we define $\Phi^{\sigma}_C(\ell, (t_1, \ldots, t_m))$ to be this unique point. It is easy to check from definition that Φ^{σ}_C is indeed the inverse of Ψ^{σ}_C . This finishes the proof of Proposition 3.10.

By applying the above proof and Proposition 3.10 to any sub-collection $I \subset \{1, \dots, m\}$ including the full collection itself, we also obtain the following stronger form of Theorem 3.9

Theorem 3.13. Let $I \subset \{1, \dots, m\}$ be any sub-collection, and define

$$H_I = \bigcap_{i \in I} H_i$$

Assume $\pi_{H_I} : H_I \to \mathcal{N}_{H_I}$ has contractible fibers. Let $\lambda_{\mathcal{N}_{H_I}}$ be the canonical induced Liouville form as before. Then the following hold:

(1) There is an $\mathbb{R}^{|I|}$ -action on H_I that is free, proper and discontinuous such that H_I is foliated by the $\mathbb{R}^{|I|}$ -orbits. In particular the map

$$\Psi_{H_I}^{\sigma}: H_I \to \mathcal{N}_{H_I} \times \mathbb{R}^{|I|}$$

is an $\mathbb{R}^{|I|}$ -equivariant diffeomorphism with respect to the $\mathbb{R}^{|I|}$ -action on H_I and that of linear translations on $\mathbb{R}^{|I|}$.

(2) The leaf space \mathcal{N}_{H_I} carries a canonical structure of Liouville manifold with boundary and corners.

By applying the above to the full collection $C = H_{\{1,...,m\}}$, we have finished the proof of Theorem 3.9.

3.3. Compatibility of null foliations of clean coisotropic intersections. Let $C_{\delta} = C$ as in the previous section and let $\{\sigma_1, \dots, \sigma_m\}$ a collection of sections $\sigma_i : \mathcal{N}_{H_i} \to H_i$ made in Choice 3.8. For each subset $I \subset \{1, \dots, m\}$, we have the following section

$$\sigma_I: \mathcal{N}_{H_I} \to H_I$$

defined by

$$\sigma_I([\ell]) := \Phi^{\sigma}_{H_I}([\ell], (0, \cdots, 0)) = (\Psi^{\sigma}_{H_I})^{-1}([\ell], (0, \cdots, 0))$$
(3.14)

for the diffeomorphism Φ_{H_I} given in (3.13) applied to $C = H_I$.

Then for each pair of subsets $I \subset J$ of $\{1, \dots, n\}$, we have the map

$$\psi_{JI}^{\sigma}: \mathcal{N}_{H_J} \to \mathcal{N}_{H_J}$$

given by

$$\psi_{JI}^{\sigma}([\ell]) := \pi_{\mathcal{N}_{H_{J}}}(\Phi_{H_{J}}^{\sigma}([\ell], (0, \cdots, 0)).$$
(3.15)

In particular consider the cases with $I = \{i\}$, $J = \{i, j\}$ and $K = \{i, j, k\}$. Then we prove the following compatibility of the collection of maps ψ_{IJ} : For each $i \neq j$, we consider the maps

$$\psi_{ij,i}^{\sigma}: \mathcal{N}_{H_i \cap H_j} \to \mathcal{N}_H$$

defined by $\psi_{ij,i}^{\sigma} := \psi_{\{ij\}\{i\}}$, and the inclusion maps

$$\iota_{ij,i}: H_i \cap H_j \to H_i.$$

Proposition 3.14. Let $\{H_1, \ldots, H_m\}$ be a collection of hypersurfaces satisfying only (S1) and (S2). Then the maps $\psi_{ij,i}^{\sigma}$ satisfy the following:

- (1) They are embeddings.
- (2) The diagram

$$\begin{array}{c} H_i \cap H_j \xrightarrow{\iota_{ij,i}} H_i \\ \pi_{ij} \bigvee \qquad & \downarrow \\ \mathcal{N}_{H_i \cap H_j} \xrightarrow{\psi_{ij,i}^{\sigma}} \mathcal{N}_{H_i} \end{array}$$
(3.16)

commutes for all pairs $1 \leq i, j \leq n$.

(3) The diagrams are compatible in the sense that we have

$$\psi_{ij,i}^{\sigma} \circ \psi_{ijk,ij}^{\sigma} = \psi_{ijk,i}^{\sigma}$$

for all triples $1 \leq i, j, k \leq n$.

Proof. We first show that the map $\psi_{ij,i}^{\sigma}$ is an embedding. Let ℓ_1, ℓ_2 be two leaves of the null-foliation of $H_i \cap H_j$ such that

$$\ell_1 \cap H_i = \ell_2 \cap H_i$$

By definition of leaves, we have only to show that $\ell_i \cap \ell_j \neq \emptyset$.

Let $x \in H_i$ in the above two common intersection which obviously implies

$$x \in \ell_1 \cap \ell_2 \subset H_i \cap H_j.$$

This proves $\psi_{ij,i}^{\sigma}$ is a one-one map. Then smoothness and the embedding property of $\psi_{ij,i}^{\sigma}$ follow from the definition of smooth structures given on the leaf spaces.

For the commutativity, we first note

$$\psi_{ij,i}^{\sigma}(\pi_{ij}(x)) = \pi_i(\Phi_{ij}^{\sigma}((\pi_{ij}(x), 0, 0)))$$
(3.17)

by the definition of the maps $\psi_{ij,i}^{\sigma}$. But by the definition (3.13) of Φ_{ij}^{σ} , the point

$$y := \Phi_{ij}^{\sigma}((\pi_{ij}(x), 0, 0))$$

is the intersection point

$$y \in \operatorname{Image} \sigma_i \cap \operatorname{Image} \sigma_j$$

Since $x \in H_i \cap H_j$, we can express it as

$$x = \Phi^{\sigma}_{ij}(\pi_{ij}(x), t_1, t_2)$$

for some $t_1, t_2 \in \mathbb{R}$. In other words, it is obtained from y by the characteristic flows of H_i and H_j by definition of Φ_{ij}^{σ} in (3.13). In particular, we have

$$\pi_i(\iota_{ij,i}(x)) = \pi_i(y)$$

On the other hand, the definition of the null foliation of \mathcal{N}_{H_i} implies

$$\pi_i(y) = \psi^{\sigma}_{ij,i}(\pi_{ij}(x)) \tag{3.18}$$

for all $x \in H_i \cap H_j$. Combining the last two equalities and commutativity of the diagram $\pi_i \circ \iota_{ij,i} = \psi_{ij,i}^{\sigma} \circ \pi_{ij}$, we have proved the commutativity of (3.16).

Finally we show that $\psi_{i,i}^{\sigma}$ is a symplectic map. Consider the pull-back

$$\omega_{ij}^{\sigma} := (\psi_{ij,i}^{\sigma})^* (\omega_{\mathcal{N}_{H_i}})$$

We will show that ω_{ij}^{σ} satisfies the defining property

$$\pi^*_{H_i \cap H_j} \omega^{\sigma}_{ij} = \iota^*_{H_i \cap H_j} \omega, \quad \omega = d\lambda$$

of the reduced form on $\mathcal{N}_{H_i \cap H_j}$ under the coisotropic reduction on the coisotropic submanifolds $H_i \cap H_j \subset M$. We compute

$$\pi_{H_i \cap H_j}^* \omega_{ij}^{\sigma} = \pi_{H_i \cap H_j}^* (\psi_{ij,i}^{\sigma})^* (\omega_{\mathcal{N}_{H_i}})$$

$$= (\psi_{ij,i}^{\sigma} \circ \pi_{H_i \cap H_j})^* (\omega_{\mathcal{N}_{H_i}})$$

$$= (\pi_{H_i} \circ \iota_{H_i \cap H_j, H_i})^* \omega_{\mathcal{N}_{H_i}}$$

$$= (\iota_{H_i \cap H_j, H_i})^* (\pi_{H_i}^* \omega_{\mathcal{N}_{H_i}})$$

$$= (\iota_{H_i \cap H_j, H_i})^* (\iota_{H_i}^* \omega) = \iota_{H_i \cap H_j}^* \omega$$

where we use the defining condition of the reduced form $\omega_{\mathcal{N}_{H_i}}$ of $\omega_{\partial H_i}$

$$\pi_{H_i}^* \omega_{\mathcal{N}_{H_i}} = \iota_{H_i}^* \omega_{\mathcal{N}_{H_i}}$$

for the penultimate equality. Therefore we have proved

$$\pi^*_{H_i \cap H_i} \omega^{\sigma}_{ij} = \iota^*_{H_i \cap H_i} \omega.$$

This shows that the form ω_{ij}^{σ} satisfies the defining equation (2.24) of the reduced form $\omega_{H_i \cap H_j}$. Then by the uniqueness of the reduced form, we have derived

$$\omega_{ij}^o = \omega_{H_i \cap H_j}$$

This proves $(\psi_{ij,i}^{\sigma})^* \omega_{H_i} = \omega_{H_i \cap H_j}$, which finishes the proof of Statement (1).

Statement (2) also follows by a similar argument this time from the naturality of the *coisotropic reduction by stages*: Consider H_i , H_j , H_k in the given coisotropic collection and consider the two flags

$$H_i \cap H_j \cap H_k \subset H_i \cap H_j \subset H_i \tag{3.19}$$

and

$$H_i \cap H_j \cap H_k \subset H_i. \tag{3.20}$$

The composition $\psi_{ij,i}^{\sigma} \circ \psi_{ijk,ij}^{\sigma}$ is the map obtained by the coisotropic reductions in two stages and $\psi_{ijk,i}^{\sigma}$ is the one obtained by the one stage reduction performed in the proof of Statement 1 with the replacement of the pair $(H_i \cap H_j, H_i)$ by $(H_i \cap H_j \cap H_k, H_i)$. Then by the naturality of the coisotropic reduction, we have proved Statement 2. This finishes the proof of the proposition. \Box

The following is an immediate corollary of the above proposition and its proof. (See Remark 1.13 for the relevant remark on the stratified presymplectic manifolds.)

Corollary 3.15. The collection of maps

$$\{\psi_I\}_{I \subset \{1,...,m\}}$$

are compatible in that the leaf space \mathcal{N}_{H_I} carries the structure of symplectic manifold with boundary and corners.

4. Liouville σ -sectors and canonical splitting data

Let $\{H_1, \dots, H_m\}$ be a clean coisotropic collection as in Definition 3.2. We denote their intersection by

$$C = H_1 \cap \dots \cap H_m$$

as before, which is a coisotropic submanifold of codimension m associated thereto.

4.1. Definition of Liouville σ -sectors with corners. Denote by $\iota_{CH_i} : C \to H_i$ the inclusion map, and $\sigma = \{\sigma_1, \ldots, \sigma_m\}$ be the collection as before. This induces the diagram

$$C \xrightarrow{\iota_{CH_i}} H_i$$

$$\pi_C \bigvee_{\mathcal{N}_C} \xrightarrow{\psi_{CH_i}^{\sigma}} \mathcal{N}_{H_i}$$

$$(4.1)$$

for all i which are compatible in the sense of Statement (2) of Proposition 3.14. In fact, we have

$$\mathcal{D}_C = \mathcal{D}_{H_1}|_C + \mathcal{D}_{H_2}|_C + \dots + \mathcal{D}_{H_m}|_C \tag{4.2}$$

which canonically induces the leaf map $\psi^{\sigma}_{CH_i}$ in the bottom arrow that makes the diagram commute.

With these preparations, we are finally ready to provide the sectional characterization of Liouville sectors with corners. **Definition 4.1** (Liouville σ -sectors with corners). Let M be a manifold with corners equipped with a Liouville one-form λ . We call (M, λ) a Liouville σ -sector with corners if at each sectorial corner δ of ∂M , the corner can be expressed as

$$C_{\delta} := H_{\delta,1} \cap \cdots \cap H_{\delta,m}$$

for a clean coisotropic collection

$$\{H_{\delta,1},\cdots,H_{\delta,m}\}$$

of σ -sectorial hypersurfaces such that fibers of the map

$$\pi_{C_{\delta}}: C_{\delta} \to \mathcal{N}_{C_{\delta}}$$

are contractible. We call such a corner C_{δ} a σ -sectorial corner of codimension m.

In the remaining section, we will derive the consequences of this definition.

4.2. Integrable systems and canonical splitting data. By applying Theorem 3.7 to the coisotropic submanifold C, we will obtain a neighborhood $Nbhd(C) \subset M$ and the projection

$$\widetilde{\pi}_C$$
: $\mathrm{Nbhd}(C) \to C.$

Choice 4.2 (Splitting Γ_C^{σ}). Let $\sigma = \{\sigma_1, \dots, \sigma_m\}$ be a choice of sections of clean coisotropic collection $\{H_1, \dots, H_m\}$. Then we associate the splitting

$$\Gamma = \Gamma_C^{\sigma} : \quad TC = G_C^{\sigma} \oplus \mathcal{D}_C \tag{4.3}$$

thereto given by the transversal symplectic subspace

$$G_C^{\sigma}|_x := (d\Psi_C^{\sigma}|_x)^{-1} (T_{\pi_C(x)} \mathcal{N}_C \oplus \{0\}_{\mathbb{R}^m}).$$
(4.4)

Applying Theorem 3.7, we obtain a diffeomorphism

$$\Psi^{\sigma}_{\Gamma} : \mathrm{Nbhd}(C) \to V \subset E^* = T^* \mathcal{F}_C$$

where \mathcal{F}_C is the null foliation of C. Furthermore the pushforward of symplectic form $d\lambda$ on U is given by the canonical Gotay's symplectic form on $V \subset E^*$

$$(\Psi_{\Gamma}^{\sigma})_*(d\lambda) = \pi^* \omega_C - d\theta$$

for the presymplectic form $\omega_C = \iota_C^*(d\lambda)$ on C. (See Theorem 3.7.)

Note that we have

$$\mathcal{D}_C|_x = \operatorname{span}_{\mathbb{R}} \{ Z_1(x), \cdots, Z_m(x) \}$$

by definition of Z_i above. Therefore the aforementioned \mathbb{R}^m -action induces an \mathbb{R}^m -equivariant bundle isomorphism

$$\mathcal{D}_C \cong C \times \mathbb{R}^m$$

over C. (This isomorphism does not depend on the choice of σ but depends only on the Liouville geometry of Nbhd $(C \cap \partial_{\infty} M)$. See Remark 2.35)

Then we have made the aforementioned splitting $TC = G_C^{\sigma} \oplus \mathcal{D}_C$ given in (4.4) \mathbb{R}^m -equivariant. In other words, for each group element $\mathbf{t} = (t_1, \ldots, t_m) \in \mathbb{R}^m$, we have the equality

$$d\mathsf{t}(G_x^{\sigma}) = G_{\mathsf{t}\cdot x}^{\sigma}$$

For a fixed $\alpha > 0$, we put

$$I_i^{\sigma} = \pm e^{\alpha t_i^{C,\sigma}} \tag{4.5}$$

which then satisfies $dI_i^{\sigma}(Z_i) = \alpha I_i^{\sigma}$ on C.

Noting that the induced \mathbb{R}^m -action on TC preserves the subbundle

$$T\mathcal{F}_C = \mathcal{D}_C \subset TC$$

the canonically induced action on T^*C also preserves the subbundle

 $\mathcal{D}_C^\perp \subset T^*C$

for which we have the isomorphism

 $T^* \mathcal{F} \cong \mathcal{D}_C^\perp$.

Therefore the \mathbb{R}^m -action on C can be lifted to $T^*\mathcal{F}$ which is the restriction of the canonical induced action on T^*C of the one on C.

Lemma 4.3. We can lift the vector fields Z_j 's to \widetilde{Z}_j on $T^*\mathcal{F}$ which are the generators of the induced \mathbb{R}^m -action such that

(1)
$$Z_j|_C = Z_j$$
,

(2) The collection $\{\widetilde{Z}_i\}$ are commuting.

Proof. Let $\phi_{Z_j}^t$ be the flow of Z_j on C. Since the \mathbb{R}^m -action is abelian, the vector fields Z_j 's are pairwise commuting. Then lifting \widetilde{Z}_j is nothing but the vector field generating the isotopy of canonical derivative maps

$$((d\phi_{Z_{i}}^{t})^{*})^{-1}: T^{*}C \to T^{*}C$$

on T^*C . Since the flows $\phi_{Z_j}^t$ are commuting, their derivatives are also commuting. Then obviously their dual flows $((d\phi_{Z_j}^t)^*)^{-1}$ on T^*C are also commuting and hence \widetilde{Z}_j 's too. The first condition also follows since any derivative maps zero vector to a zero vector. This finishes the proof.

We now define

$$\widetilde{I}_i^{\sigma} = I_i^{\sigma} \circ \pi_{T^* \mathcal{F}}.$$

Then $\{dI_1^{\sigma}, \dots, dI_m^{\sigma}\}\$ are linearly independent on a neighborhood of the zero section of $T^*\mathcal{F}$ if we choose the neighborhood small enough. This is because $\{dI_1^{\sigma}, \dots, dI_m^{\sigma}\}\$ are linearly independent on C. By suitably adjusting the parametrization $t_i^{C,\sigma}$ of the \mathbb{R}^m -action, we can make the equation

$$d\widetilde{I}_i^{\sigma}(\widetilde{Z}_j) = \alpha \delta_{ij} \widetilde{I}_i^{\sigma} \tag{4.6}$$

hold.

This is precisely the situation of completely integrable system to which we can apply the standard construction of action-angle coordinates. (See [Arn88, Section 49] for example.) Therefore, regarding $\{\tilde{I}_1^{\sigma}, \ldots, \tilde{I}_m^{\sigma}\}$ as the (fiberwise) angle coordinates, we can find a unique choice of (fiberwise) action coordinates

$$\{\widetilde{R}_1^{\sigma},\cdots,\widetilde{R}_m^{\sigma}\}$$

over \mathcal{N}_C satisfying

$$\{\widetilde{R}_i^{\sigma}, \widetilde{I}_i^{\sigma}\} = \delta_{ij}, \quad \widetilde{R}_i^{\sigma} \circ \Phi_C^{\sigma}|_{H_i} = 0$$

on a neighborhood $V \subset T^* \mathcal{F}_C$ of the zero section $0_{T^* \mathcal{F}_C} \cong C$. Now we define the pull-back functions

$$R_i^{\sigma} := \widetilde{R}_i^{\sigma} \circ \Phi_C^{\sigma}, \quad I_j^{\sigma} := \widetilde{I}_j^{\sigma} \circ \Phi_C^{\sigma}$$

on U = Nbhd(C). We also pull-back the vector fields Z_j to Nbhd(C) by Φ_C^{σ} and denote them by Z_j . (Note that the notations I_j^{σ} and Z_j are consistent in that their

restrictions to C are nothing but the above already given I_j^{σ} or Z_j respectively on C.) Furthermore, we have the relationship

$$Z_j = X_{R_j^{\sigma}}.$$

(See the definition (3.11) of Z_i on C.)

Then we have

$$\{R_i^{\sigma}, R_j^{\sigma}\} = \omega(X_{R_i^{\sigma}}, X_{R_j^{\sigma}}) = \omega(Z_i, Z_j) = 0$$

on Nbhd(C). Since $Z_i := X_{R_i^{\sigma}}$, we have

$$Z_i \rfloor \omega = dR_i^\sigma \tag{4.7}$$

on $U = \operatorname{Nbhd}^Z(C)$. This is precisely the defining equation of the moment map $\phi_{G,C}^{\sigma} : \operatorname{Nbhd}(C) \to \mathfrak{g}^* \cong \mathbb{R}^m$ with $G = \mathbb{R}^m$ given by

$$\phi_{G,C}^{\sigma}(x) = (R_1^{\sigma}(x), \cdots, R_m^{\sigma}(x))$$

for the above $G = \mathbb{R}^m$ -action. Recall that the hypersurfaces H_i are Z-invariant at infinity. Therefore we can choose the neighborhood Nbhd(C) so that it is Zinvariant at infinity. Then by the requirement put on the Liouville vector field Z which is pointing outward along ∂M , we can choose the whole neighborhood Nbhd(C) Z-invariant. Together with the normalization condition of R_i 's

$$R_i^{\sigma}|_{H_i} = \tilde{R}_i \circ \Phi_C^{\sigma}|_{H_i} = 0,$$

it also implies $R_i^{\sigma} \ge 0$ on Nbhd(C) for all *i*. We now take the neighborhood $U \subset M$ to be this Z-invariant neighborhood

$$U = \operatorname{Nbhd}^Z(C).$$

The content of the above discussion can be summarized into the following intrinsic derivation of the splitting data.

Theorem 4.4 (σ -Splitting data). Let $C \subset \partial M$ be a sectorial corner of codimension n associated to the sectorial coisotropic collection $\{H_1, \ldots, H_m\}$ on ∂M . Then for each choice

$$\sigma = \{\sigma_1, \cdots, \sigma_m\}$$

of sections $\sigma_i : \mathcal{N}_{H_i} \to H_i$ of π_{H_i} for $i = 1, \dots, n$, there is a diffeomorphism

$$\Psi^{\sigma}_C : \mathrm{Nbhd}^Z(C) \cap \partial M \to F \times \mathbb{R}^m$$

and

$$\psi_C^{\sigma}: \mathcal{N}_C \to F_C^{\sigma}$$

such that

- (1) $F_C^{\sigma} = \text{Image } \sigma_1 \cap \cdots \cap \text{Image } \sigma_m$,
- (2) $(\Psi_C^{\sigma})_*\omega_{\partial} = \pi_F^*\omega_F,$
- (3) The following diagram

$$\frac{\partial M|_C \longrightarrow F_C^{\sigma} \times \mathbb{R}^m}{\bigvee_{\pi_{\partial M}} \pi_{F_C^{\sigma}}}$$

$$\frac{\chi_{\pi_{\partial M}}}{\bigvee_{\pi_{F_C^{\sigma}}}} F_C^{\sigma}.$$

$$(4.8)$$

commutes for the map

$$\Psi_C^{\sigma} = (\sigma_C \circ \pi_{F_C}, (I_1^{\sigma}, \cdots, I_m))$$

(4) The G-action with $G = \mathbb{R}^m$ has the moment map $\phi_{G,C}^{\sigma} : \operatorname{Nbhd}_{\epsilon}^Z(C) \to \mathbb{R}^m$ given by

$$\phi^{\sigma}_{G,C} = (R^{\sigma}_1, \cdots, R^{\sigma}_m)$$

for a collection of Poisson-commuting R_i 's satisfying the simultaneous normalization condition

$$R_i^{\sigma}|_{H_i} = 0, \quad R_i^{\sigma} \ge 0$$

for all i on $\operatorname{Nbhd}^Z(C)$.

(5) The map $\operatorname{Nbhd}(\partial M) \to F_C \times \mathbb{C}^m_{\operatorname{Re} \geq 0}$ is given by the formula

$$\widetilde{\Psi}_C^{\sigma}(x) = \left(\sigma_C(\pi_{F_C}(x)), R_1^{\sigma}(x) + \sqrt{-1}I_1^{\sigma}(x), \dots, R_m^{\sigma}(x) + \sqrt{-1}I_m^{\sigma}(x)\right).$$
(4.9)
such that

such that

$$(\widetilde{\Psi}_C^{\sigma})_*\omega = \pi_F^*\omega_{F_C} + \sum_{i=1}^m dR_i^{\sigma} \wedge dI_i^{\sigma}.$$
(4.10)

We call these data a σ -splitting data of Nbhd(C) associated to the choice σ = $\{\sigma_1, \cdots, \sigma_m\}$ of sections $\sigma_i : \mathcal{N}_{H_i} \to H_i$.

We also gather the following consequences of the above discussion separately. The first one, in particular, states that Proposition 1.5 still holds for the Liouville σ -sectors with corners.

Theorem 4.5. (1) Each Liouville σ -sector with corners is a Liouville sector in the sense of Definition 3.1.

(2) The leaf space $\mathcal{N}_{C_{\delta}}$ carries a natural structure of manifold with corners at each sectorial corner δ such that the map $\pi_{C_{\delta}} : \partial M \to \mathcal{N}_{C_{\delta}}$ is a morphism of manifolds with corners.

Proof. We have already constructed a diffeomorphism

$$\Psi^{\sigma}_{\delta}: \partial M|_{C_{\delta}} \to F^{\sigma}_{\delta} \times \mathbb{R}^{r}$$

given by

$$\Psi^{\sigma}_{\delta}(x) = (\pi_{F^{\sigma}_{\delta}}(x), I^{\sigma}_1(x), \dots, I^{\sigma}_m(x)).$$

Each I_i^{σ} defined on ∂M is extended to the function $\widetilde{I}_i^{\sigma} \circ \Phi_{C_{\delta}}^{\sigma}$ on a symplectic neighborhood $U_{\delta} := \operatorname{Nbhd}^Z(C_{\delta}) \subset M$ via Gotay's coisotropic neighborhood map

$$\Phi^{\sigma}_{C_{\delta}} : \mathrm{Nbhd}(C_{\delta}) \hookrightarrow T^* \mathfrak{F}_{C_{\delta}}$$

where the function \widetilde{I}_i^{σ} is canonically defined on a neighborhood

$$V \subset E^* = T^* \mathcal{F}_{C_\delta}$$

This diffeomorphism $\Phi_{C_{\delta}}$ onto $V_{\delta} \subset T^* \mathcal{F}$ also induces a splitting of the tangent bundle TC_{δ}

$$\Gamma_{C_{\delta}}^{\sigma}: \quad TC_{\delta} = G_{\delta}^{\sigma} \oplus T\mathcal{F}_{C_{\delta}} = G_{\delta}^{\sigma} \oplus \mathcal{D}_{C_{\delta}}$$

such that G^{σ}_{δ} is a transverse symplectic subbundle of TC_{δ}

$$G^{\sigma}_{\delta}|_{x} := d\Psi^{-1}\left(T_{\pi_{F^{\sigma}_{\delta}}(x)}F^{\sigma}_{\delta} \oplus \{0\}\right)$$

at each $x \in C_{\delta}$. Theorem 4.4 then finishes the construction of the data laid out in Definition 3.1.

For the proof of Statement (2), we start with the observation that for each $H = H_i$ the canonical smooth structure on \mathcal{N}_H carries the natural structure of a manifold with boundary and corners through a choice of smooth section made in Choice 3.8, whose existence relies on the defining hypothesis of σ -sectorial hypersurfaces that the projection map $\pi_H : H \to \mathcal{N}_H$ admits a continuous section. For each choice of smooth section, by the same construction as in Subsection 2.6, we have a symplectic structure $(\mathcal{N}_H, \omega_{\mathcal{N}_H})$, and a smooth map $\sigma_{\infty}^+ : \mathcal{N}_H \to \partial_{\infty} M$ which is a symplectic diffeomorphism onto the convex hypersurface (F_{∞}^+, ω^+) of the contact manifold $(\partial_{\infty} M, \xi)$. For two different choices of splittings, the resulting structures are diffeomorphic.

Finally it remains to verify the property of \mathcal{N}_C carrying the structure of the Liouville manifolds with corners. But this immediately follows from the compatibility result, Proposition 3.14: The moment map $\phi_{G,\delta}^{\sigma}$: $\mathcal{N}bhd^Z(C_{\delta}) \to \mathbb{R}^m_+$ provides local description of the codimension k-corner of $\mathcal{N}_{C_{\delta}}$. This finishes the proof. \Box

5. Solution to [GPS17, Question 2.6] and convexity at infinity

As an application of our arguments used to derive the canonical splitting data, we can now provide the affirmative answer to a question raised by Ganatra-Pardon-Shende in [GPS17].

Theorem 5.1 (Question 2.6 [GPS17]). Suppose (M, λ) is a Liouville manifoldwith-boundary that satisfies the following:

- (1) Its Liouville vector field Z is tangent to ∂M at infinity.
- (2) There is a diffeomorphism $\partial M = F \times \mathbb{R}$ sending the characteristic foliation to the foliation by leaves $\mathbb{R} \times \{p\}$.

Then M is a Liouville σ -sector.

The proof will be divided into two parts: we first examine the presymplectic geometry component of the proof, and then combine the discussion with that of the Liouville geometry.

In the mean time, the following is an immediate corollary of Theorem 5.1.

Corollary 5.2. In the presence of other conditions, the convexity condition (b) in Definition (1.2) is equivalent to the existence of a diffeomorphism

$$\partial M \cap \mathcal{N}bhd(\partial_{\infty}M) \cong F_0 \times [N,\infty)$$

sending the characteristic foliation to the foliation by leaves $\{p\} \times [N, \infty)$ for a sufficiently large N > 0.

5.1. **Presymplectic geometry.** Denote by $\iota_{\partial M} : \partial M \to M$ the inclusion map. Then the one-form $\lambda_{\partial} := \iota_{\partial M}^* \lambda$ induces the structure of presymplectic manifold

 $(\partial M, d\lambda_{\partial}).$

By definition, $\mathcal{D}_{\partial M} = \ker d\lambda_{\partial}$. Denote by $\Psi : \partial M \to F \times \mathbb{R}$ the diffeomorphism entering in the hypothesis. Then the hypothesis implies that we have a commutative diagram

where $\psi := [\Psi] : \mathcal{N}_{\partial M} \to F$ the obvious quotient map, which becomes a diffeomorphism. Obviously the map $\sigma : \mathcal{N}_{\partial M} \to \partial M$ defined by

$$\sigma(\ell) := \Psi^{-1}(\psi(\ell), 0) \tag{5.2}$$

defines a continuous section of $\pi_{\partial M} : \partial M \to \mathcal{N}_{\partial M}$, one of the defining data of Liouville σ -sectors.

Next, by Condition (1), we have

$$\partial_{\infty} M \cap \partial M = \partial_{\infty} (\partial M).$$

Therefore it remains to show convexity of $\partial_{\infty} M \cap \partial M$ in $\partial_{\infty} M$, i.e., that there exists a contact vector field defined on Nbhd $(\partial_{\infty} M \cap \partial M) \subset \partial_{\infty} M$ that is transversal to the hypersurface

$$\partial_{\infty} M \cap \partial M$$

We denote the reduced symplectic form on $\mathcal{N}_{\partial M}$ of the presymplectic form $d\lambda_{\mathcal{N}_{\partial M}}$ by

 $\omega_{\mathcal{N}_{\partial M}}.$

Next we prove

Lemma 5.3. Suppose that Z is tangent to ∂M outside a compact subset $K \subset M$. Consider the pull-back $\lambda_{\partial} := \iota_{\partial} M^* \lambda$ which is a presymplectic form. Let X be a vector field tangent to ker $\iota_{\partial} M^* \lambda = \mathcal{D}_{\partial} M$. Then we have

$$\mathcal{L}_X \lambda_\partial = 0$$

on $\partial M \cap (M \setminus K)$.

Proof. Let Z be tangent to $\partial M \cap (M \setminus K)$ for a compact set $K \subset M$. Since X spans the characteristic distribution of $(\partial M, \lambda_{\partial})$, we have

$$X | d\lambda_{\partial} = 0$$

on ∂M . On the other hand, since Z is tangent to $\partial M \cap (M \setminus K)$ and $X \in \ker \omega_{\partial} = d\lambda_{\partial}$, we also have

$$0 = d\lambda_{\partial}(Z, X) = \lambda_{\partial}(X)$$

where the second equality follows by definition of Liouville vector field Z. Therefore on $\partial M \cap (M \setminus K)$, we compute

$$\mathcal{L}_X \lambda_\partial = (d(X \rfloor \lambda) + X \rfloor d\lambda)|_{\partial M} = 0$$

which finishes the proof.

We push-forward the presymplectic structure on ∂M to $F \times \mathbb{R}$ by Ψ and write the resulting presymplectic form by

$$\lambda^{\operatorname{pre}} := \Psi_*(\lambda_\partial)$$

on $F \times \mathbb{R}$.

Lemma 5.4. $\mathcal{L}_{\frac{\partial}{\partial t}}\lambda^{\text{pre}} = 0$ on $F \times [N, \infty)$ and so

$$\lambda^{\rm pre} = \pi_F^* \lambda_F$$

for some one-form λ_F on F, where $\pi_F : F \times \mathbb{R} \to F$ is the projection.

Proof. Let X be the pull-back vector field defined by $X = \Psi^*(\frac{\partial}{\partial I})$ on ∂M . Suppose that Z is tangent to ∂M on $(M \setminus K) \cap \partial M$ for a compact subset $K \subset M$. Since K is compact, there exists a sufficiently large N > 0 such that

$$\Psi(K) \subset F \times [-N, N].$$

In particular, we have $F \times [N, \infty) \subset \Psi(\partial M \setminus K)$ for a sufficiently large $N \in \mathbb{R}$. Then Lemma 5.3 applied to $H = \partial M$ implies

$$\mathcal{L}_{\frac{\partial}{\partial t}}\lambda^{\mathrm{pre}} = 0$$

on $F \times [N, \infty)$, i.e., it is $\frac{\partial}{\partial t}$ -invariant and hence there exists a one-form λ_F on F such that $\pi_F^* \lambda_F = \lambda^{\text{pre}}$ thereon. This finishes the proof.

We denote by

$$(Y, d\lambda^{\text{pre}}), \quad Y := F \times \mathbb{R}$$

the resulting presymplectic manifold $(F \times \mathbb{R}, d\lambda^{\text{pre}})$.

Using Condition (1) of Theorem 5.1, we can choose a smooth section defined as in Lemma 2.33. From now on, we fix $\sigma = \sigma_N^+$ to be such a smooth section, which also induces the natural Liouville form $\lambda_{N_{\partial M}}$ from λ_{∂} as constructed in Lemma 2.33, which satisfies

$$\lambda_{\partial}|_{\{I \ge N\}} = \pi_{\partial M}^* \lambda_{\mathcal{N}_{\partial M}} \tag{5.3}$$

by choosing N so large that the form λ_{∂} on $\{I \geq N\}$ is projectable where $I = t \circ \pi_{\mathbb{C}}$ for the coordinate function t of \mathbb{R} . (Similar discussion also applies on $\{I \leq -N\}$ with N > 0.) Then we have

$$\psi_*(\omega_{\mathcal{N}_{\partial M}}) = d\lambda_F := \omega_F$$

which is also the reduced symplectic form of $d\lambda^{\text{pre}}$ of $F \times \mathbb{R}$ relative to the choice of section

$$\sigma_F := \Psi \circ \sigma \circ \circ \psi^{-1} \tag{5.4}$$

of the trivial fibration $F \times \mathbb{R} \to F$.

By applying Theorem 4.4 for n = 1, we can extend the presymplectic map $\Psi : \partial M \to F \times \mathbb{R}$ to a symplectic thickening

$$\Psi : \mathrm{Nbhd}(\partial M) \to F \times \mathbb{C}.$$

We equip $V = Nbhd(F \times \mathbb{R}) \subset F \times \mathbb{C}$ with the pushforward symplectic form

$$\omega_V = \Psi_*(d\lambda). \tag{5.5}$$

We have the canonical isomorphism

$$T^* \mathfrak{F} \cong Y \times \mathbb{R} = F \times \mathbb{R} \times \mathbb{R} \cong F \times \mathbb{C}$$

and H is contained in $V \subset T^* \mathcal{F}$ as the zero section of $T^* \mathcal{F}$. We denote by

$$\mathrm{pr}: V \subset T^* \mathcal{F} \to Y$$

the projection.

Lemma 5.5. Let $s + \sqrt{-1t}$ be the standard coordinates of \mathbb{C} . We define

$$R = s \circ \pi_{\mathbb{C}} \circ \widetilde{\Psi}, \quad I = t \circ \pi_{\mathbb{C}} \circ \widetilde{\Psi}$$

on $F \times \mathbb{C}$. Then there is a neighborhood $V' \subset V$ and a diffeomorphism

$$\Upsilon: V' \cap \{I > N'\} \to V$$

onto its image that preserves $(Y, d\lambda^{\text{pre}})$ and so that

$$\omega_{V'} := \Upsilon^* \omega_V = \pi_F^* \omega_F + dR \wedge dI \tag{5.6}$$

on $\{I > N'\} \cap V'$ for some sufficiently large N' > 0.

Proof. Recall from (5.3) that

$$\mathrm{pr}^*\lambda^{\mathrm{pre}} = \pi_F^*\lambda_F$$

on $\{I \ge N\}$ for a sufficiently large N > 0. Then the inclusion

$$(F \times \{I \ge N\}, d\lambda^{\text{pre}}) \hookrightarrow (F \times \mathbb{C}, \pi_F^* \omega_F + dR \wedge dI)$$

is a coisotropic embedding and the lemma follows by the uniqueness of Gotay's coisotropic embedding theorem. $\hfill \Box$

Based on this lemma, we may and will assume

$$\Psi_* d\lambda = \omega_V = \pi_F^* \omega_F + dR \wedge dI, \quad \omega_F = \psi_* \omega_{\mathcal{N}_{\partial M}} \tag{5.7}$$

on $Nbhd(F \times \mathbb{R}) \subset F \times \mathbb{C}_{Re \geq 0}$. From now on, we will work with the trivial fibration $F \times \mathbb{R} =: H$ as a hypersurface in the Liouville manifold (V, ω_V) with $V \subset F \times \mathbb{C} \cong T^* \mathcal{F}$.

5.2. Liouville geometry. Now we take the Liouville form of (V, ω_V) given by

$$\lambda_V = \widetilde{\Psi}_* \lambda$$

and let $Z_V = \widetilde{\Psi}_* Z$ be its associated Liouville vector field. By definition, Z_V satisfies

$$Z_V \rfloor d\lambda_V = \lambda_V \tag{5.8}$$

where we have

$$d\lambda_V = \pi_F^* \omega_F + dR \wedge dI$$

from (5.7). By decomposing the Liouville vector field Z_V into

$$Z_V = X_F + a\frac{\partial}{\partial R} + b\frac{\partial}{\partial I}$$
(5.9)

in terms of the splitting $TV = TF \oplus T\mathbb{C}$, we compute

$$Z_V \rfloor d\lambda_V = X_F \rfloor \pi_F^* \omega_F + a \, dI - b \, dR$$

for some coefficient functions a = a(y, R, I), b = b(y, R, I) for $(y, R, I) \in F \times \mathbb{C}$. Then (5.8) becomes

$$X_F \rfloor \pi_F^* \omega_F + a \, dI - b \, dR = \lambda_V. \tag{5.10}$$

Proposition 5.6. Regard $\{R = 0\} =: H$ as a hypersurface of V. Let b be the coefficient function appearing in (5.9). Then we have $b \neq 0$ on $V' \cap \{I > C'\}$ on a possibly smaller neighborhood $V' \subset V$ of H for a sufficiently large constant C' > 0.

Proof. We denote by $\iota_H : H \to V$ the inclusion map. We first recall from Lemma 5.3 that $X = \Psi^* \frac{\partial}{\partial I} \in \ker d\lambda_\partial$. Therefore we have

$$\frac{\partial}{\partial I} \rfloor \Psi_* d\lambda_\partial = 0$$

on $F \times \mathbb{R}$, since $\Psi_* \lambda_\partial = \lambda^{\text{pre}} = \iota_H^* \lambda_V$. Since $\omega_F = d\lambda_F$ and $d\lambda_V = \pi_F^* d\lambda_F + dR \wedge dI$, we have

$$d(\lambda_V - \pi_F^* \lambda_F - IdR) = 0.$$

Since the choice of σ made in Lemma 2.32 implies

$$\pi_F^* \lambda_F = \Psi_* \lambda_\partial = \iota_H^* \lambda_V$$

we have $\iota_H^*(\lambda_V - \pi_F^*\lambda_F - IdR) = 0$. In particular the form $\lambda_V - \pi_F^*\lambda_F - IdR$ is exact on any neighborhood V of $\{R = 0\}$ which deformation retracts to $\{R = 0\}$. Therefore we have

$$\lambda_V - \pi_F^* \lambda_F - IdR = dh_V$$

on such a neighborhood V for some function $h_V: V \to \mathbb{R}$, i.e.,

$$\lambda_V = \pi_F^* \lambda_F + I dR + dh_V. \tag{5.11}$$

Since ker $\iota_H^*(d\lambda_V) = \operatorname{span}\{\frac{\partial}{\partial I}\}$, we have

$$\frac{\partial}{\partial I} \rfloor d\lambda_V = 0$$

on H. Then since Z is tangent to H at infinity, we have

$$\lambda_V\left(\frac{\partial}{\partial I}\right) = d\lambda\left(Z, \frac{\partial}{\partial I}\right) = 0$$

Obviously we have $(\pi_F^* \lambda_F + IdR)(\frac{\partial}{\partial I}) = 0$. Therefore we have derived

$$\left. \frac{\partial h_V}{\partial I} \right|_{R=0} = 0$$

by evaluating (5.11) against $\frac{\partial}{\partial I}$. Therefore $h_V|_{\{R=0\}}$ does not depend on I. In particular, we have

$$||h_V|_{\{R=0\}}||_{C^1} \le C$$

for some constant C > 0. In turn, since h_V is a smooth function, we have

$$\|h_V\|_{C^1} \le C$$

by precompactness of V/\mathbb{R} on a sufficient small neighborhood $V' \subset V$ of $H = \{R = 0\}$ for some constant C' choosing C' larger, if necessary. In particular we have

$$\left\|\frac{\partial h_V}{\partial R}\right\|_{C^0;V'} \le C' \tag{5.12}$$

Substituting (5.11) into (5.10), we obtain the equation

$$X_F \rfloor d\lambda_V + a \, dI - b \, dR = \pi_F^* \lambda_F + I dR + dh_V.$$

By evaluating this equation against $\frac{\partial}{\partial R}$, we obtain

$$b = -\left(I + \frac{\partial h_V}{\partial I}\right)$$

on V. Therefore we have $b(y, R, I) \neq 0$ for all $(y, R, I) \in V'$, if |I| > C'. This finishes the proof.

From (5.9), we have derived

$$Z_V[I] = b(y, R, I)$$

In particular $Z_V[I] \neq 0$ on

$$I^{-1}(N) \cap V$$

for any $N \ge C'$ and hence any such level set is a contact-type hypersurface in V.

5.3. Combining the two. The following lemma then will finish the proof of Theorem 5.1.

Lemma 5.7. The Hamiltonian vector field X_I induces a contact vector field on the contact-type hypersurface $I^{-1}(N)$ that is transversal to $F_N := I^{-1}(N) \cap \partial M$ for all $N \geq C'$.

Proof. By the expression of symplectic form in (5.6) we have

$$\frac{\partial}{\partial R} = X_I, \quad \frac{\partial}{\partial I} = -X_R.$$

This implies

$$1 = d\lambda(X_R, X_I) = -b \, d\lambda(Z, X_I) = -b \, \lambda(X_I)$$

Since X_I is tangent to the level set $I^{-1}(N)$, X_I defines a contact vector field thereon which is transversal to the contact distribution of $I^{-1}(N)$ induced by the contact form $\theta_N := \iota_N^* \lambda$.

The same discussion applies to F_{∞}^{-} , which finishes the proof.

This finishes the proof of Theorem 5.1.

6. Structure of Liouville σ -sectors and their automorphism groups

Our definition of Liouville σ -sectors with corners enables us to give a natural notion of Liouville automorphisms which is the same as the case without boundary and which does not depend on the choice of splitting data.

We first recall the following well-known definition of automorphisms of Liouville manifold (without boundary)

Definition 6.1. Let (M, λ) be an Liouville manifold without boundary. We call a diffeomorphism $\phi : M \to M$ a Liouville automorphism if ϕ satisfies

 $\phi^*\lambda = \lambda + df$

for a compactly supported function $f: M \to \mathbb{R}$. We denote by $\operatorname{Aut}(M)$ the set of automorphisms of (M, λ) .

It is easy to check that $\operatorname{Aut}(M)$ forms a topological group. Now we would like extend this definition of automorphisms to the case of Liouville σ -sectors. For this purpose, we need some preparations by examining the universal geometric structures inherent on the boundary ∂M of a Liouville manifold with boundary and corners.

6.1. Some presymplectic geometry of ∂M . We start with the observation that $(\partial M, \omega_{\partial M})$ carries the structure of *presymplectic manifolds* as usual for any coisotropic submanifold mentioned as before. We first introduce automorphisms of presymplectic manifolds (Y, ω) in general context.

Definition 6.2. Let (Y, ω) and (Y', ω') be two presymplectic manifolds. A diffeomorphism $\phi : Y \to Y'$ is called *presymplectic* if $\phi^* \omega' = \omega$. We denote by $\mathcal{P}Symp(Y, \omega)$ the set of presymplectic diffeomorphisms.

(We refer to [OP05] for some detailed discussion on the geometry of presymplectic manifolds and their automorphisms and their application to the deformation problem of coisotropic submanifolds.)

Then we note that any diffeomorphism $\phi: (M, \partial M) \to (M, \partial M)$ satisfying

$$\phi^* \lambda = \lambda + df \tag{6.1}$$

for some function f, not necessarily compactly supported, induces a presymplectic diffeomorphism

$$\phi_{\partial} := \phi|_{\partial M}$$

on ∂M equipped with the presymplectic form

$$\omega_{\partial} := d\lambda_{\partial}, \quad \lambda_{\partial} := \iota^* \lambda$$

for the inclusion map $\iota : \partial M \to M$.

Lemma 6.3. The presymplectic diffeomorphism $\phi_{\partial} : \partial M \to \partial M$ preserves the characteristic foliation of ∂M .

Proof. We have

$$\mathcal{D}_{\partial M} = \ker \omega_{\partial}$$

Since any Liouville automorphism ϕ of $(M, \partial M)$ satisfies (6.1), we have

$$\phi_{\partial}^*\omega_{\partial} = \omega_{\partial}$$

Therefore we have

 $\phi_*(\mathcal{D}_{\partial M}) = \mathcal{D}_{\partial M}$

which finishes the proof.

In fact, for the current case of our interest $Y = \partial M$, the presymplectic form ω_{∂} is exact in that

$$\omega_{\partial} = d\lambda_{\partial}, \quad \lambda_{\partial} := \iota^* \lambda_{\partial}$$

Furthermore (6.1) implies that ϕ actually restricts to an exact presymplectic diffeomorphism

$$\phi_{\partial}: (\partial M, \omega_{\partial}) \to (\partial M, \omega_{\partial})$$

on ∂M in that

$$\phi_{\partial}^* \lambda_{\partial} - \lambda_{\partial} = dh, \quad h = f \circ d$$

where the function $h: \partial X \to \mathbb{R}$ is not necessarily compactly supported.

We have a natural restriction map

$$\operatorname{Aut}(M,\lambda) \to \operatorname{\mathscr{P}Symp}(\partial M,\omega_{\partial}); \quad \phi \mapsto \phi_{\partial}.$$
 (6.2)

Definition 6.4 (Pre-Liouville automorphism group Aut $(\partial M, \lambda_{\partial})$). We call a diffeomorphism $\phi : (\partial M, \lambda_{\partial}) \to (\partial M, \lambda_{\partial})$ a pre-Liouville diffeomorphism if the form $\phi^* \lambda_{\partial} - \lambda_{\partial}$ is exact. We say ϕ is a pre-Liouville automorphism if it satisfies

$$\phi^*\lambda_\partial = \lambda_\partial + dh$$

for a compactly supported function $h : \partial M \to \mathbb{R}$. We denote by $\operatorname{Aut}(\partial M, \lambda_{\partial})$ the set of pre-Liouville automorphisms of $(\partial M, \lambda_{\partial})$.

The following is an immediate consequence of the definition.

Corollary 6.5. The restriction map (6.2) induces a canonical group homomorphism

$$\operatorname{Aut}(M,\lambda) \to \operatorname{Aut}(\partial M,\lambda_{\partial}).$$

We recall that ∂M carries a canonical transverse symplectic structure arising from the presymplectic form $d\lambda_{\partial}$. (See [OP05, Section 4].)

44

Proposition 6.6. The induced pre-Liouville automorphism $\phi_{\partial} := \phi|_{\partial M} : \partial M \to \partial M$ descends to a (stratawise) symplectic diffeomorphism

$$\phi_{\mathcal{N}_{\partial M}}:\mathcal{N}_{\partial M}\to\mathcal{N}_{\partial M}$$

and satisfies

$$\pi_{\partial M} \circ \phi_{\partial} = \phi_{\mathcal{N}_{\partial M}} \circ \pi_{\partial M}$$

when we regard both ∂M and $\mathcal{N}_{\partial M}$ as manifolds with corners.

6.2. Automorphism group of Liouville σ -sectors. Now we are ready give the geometric structure of *Liouville* σ -sectors.

Definition 6.7 (Structure of Liouville σ -sectors). We say two Liouville σ -sectors (M, λ) and (M', λ') are isomorphic, it there exists a diffeomorphism $\psi : M \to M'$ (as a manifold with corners) such that $\psi^*\lambda' = \lambda + df$ for some compactly supported function $f: M \to \mathbb{R}$. A structure of Liouville σ -sectors is defined to be an isomorphism class of Liouville σ -sectors.

With this definition of the structure of Liouville σ -sectors in our disposal, the following is an easy consequence of the definition and Proposition 6.6, which shows that the definition of an automorphism of a Liouville sector (M, λ) is in the same form as the case of Liouville manifold given by the defining equation

$$\psi^*\lambda = \lambda + df$$

for some compactly supported function $f: M \to \mathbb{R}$, except that ψ is a self diffeomorphism of M as a stratified manifold and the equality of the above equation as in the sense of Remark 1.13.

Theorem 6.8 (Automorphism group). Let (M, λ) be a Liouville σ -sector. Suppose a diffeomorphism $\psi : M \to M$ satisfies

$$\psi^* \lambda = \lambda + df \tag{6.3}$$

for some compactly supported function $f: M \to \mathbb{R}$. Then ψ is an automorphism of the *structure of Liouville* σ -sectors.

Proof. We first discuss how the action of diffeomorphisms ψ satisfying

$$\psi^*\lambda = \lambda + dj$$

affects the structure of Liouville σ -sectors, when the function f is compactly supported. In particular we have

- $\psi^* d\lambda = d\lambda$,
- $\psi^* \lambda = \lambda$ at infinity.

Then ψ restricts to a presymplectic diffeomorphism $\psi_{\partial} : \partial M \to \partial M$ which is also pre-Liouville, i.e., satisfies

$$(\psi|_{\partial M})^* \lambda_{\partial} = \lambda_{\partial} + dh$$

for a *compactly supported* function h on ∂M .

We need to show that the *structure* of Liouville σ -sectors with respect to

$$(M, \psi^* \lambda) = (M, \lambda + df)$$

is isomorphic to that of (M, λ) . For this, we make a choice of $\sigma = \{\sigma_1, \dots, \sigma_m\}$ associated to a clean coisotropic collection $\{H_1, \dots, H_m\}$ for each sectorial corner δ of M with

$$C_{\delta} = H_1 \cap \dots \cap H_m.$$

Such a collection exists by definition for (M, λ) being a Liouville σ -sector.

$$\{H'_1, \cdots, H'_m\} := \{\psi(H_1), \dots, \psi(H_m)\}.$$

Since smooth diffeomorphisms between two manifolds with corners preserve strata dimensions by definition, we work with the defining data of $(M, \psi^* \lambda)$ stratawise of the fixed dimensional strata.

We first need to show that each H'_i is σ -sectorial hypersurface by finding a collection

$$\sigma' = \{\sigma'_1, \ldots, \sigma_m\}$$

where each σ'_i is a section o H'_i respectively. For this purpose, we prove the following

Lemma 6.9. Choose the sections σ_i s so that

Image
$$\sigma_i \subset M \setminus \operatorname{supp} df$$
.

Then there exists a neighborhood $Nbhd(\partial_{\infty}M)$ such that the following hold:

- (1) The map ψ : $\mathbb{N}bhd(\partial_{\infty}M) \cap H_i \to H_i$ descends to a diffeomorphism $[\psi]$: $\mathcal{N}_{H_i} \to \mathcal{N}_{H_i}.$ (2) The map $\sigma_i^{\psi} : \mathcal{N}_{H_i} \to \psi(H_i)$ defined by

$$\sigma_i^{\psi} := \psi \circ \sigma_i \circ [\psi]^{-1}$$

is a section of the projection $\psi(H_i) \to \mathcal{N}_{\psi(H_i)} = \mathcal{N}_{H_i}$.

Proof. Since Image $\sigma_i \subset M \setminus \operatorname{supp} df$, we have

$$\psi^*\lambda = \lambda$$

on Image $\sigma_i := F_i$. In particular, the projection $\pi_{H_i} : H_i \to \mathcal{N}_{H_i}$ restricts to a bijective map on F_i . Furthermore since $\psi^* \lambda = \lambda$ on Nbhd $(\partial_{\infty} M)$, the associated Liouville vector field Z_{λ} of λ satisfies

$$\psi_* Z_\lambda = Z_\lambda$$

thereon. Recall that ψ restricts to a diffeomorphism on ∂M (as a map on manifold with corners). Then the equality $\psi^* \lambda = \lambda$ implies $\psi^*_{\partial} d\lambda_{\partial} = d\lambda_{\partial}$ and hence

$$d\psi_{\partial}(\ker d\lambda_{\partial}) = \ker d\lambda_{\partial}$$

on $\mathcal{N}bhd(\partial M) \cap H_i$. Therefore ψ descends to a diffeomorphism $[\psi] : \mathcal{N}_{H_i} \to \mathcal{N}_{H_i}$ so that we have the commutative diagram

$$\begin{array}{c} H_i & \stackrel{\psi}{\longrightarrow} \psi(H_i) \\ \downarrow^{\pi_{H_i}} & \downarrow^{\pi_{\psi(H_i)}} \\ \mathcal{N}_{H_i} & \stackrel{[\psi]}{\longrightarrow} \mathcal{N}_{H_i}. \end{array}$$

By composing $\sigma'_i = \psi \circ \sigma_i$ with $\pi_{\psi(H_i)}$ to the left, we obtain

$$\pi_{\psi(H_i)}\sigma'_i = \pi_{\psi(H_i)} \circ \psi \circ \sigma_i = [\psi] \circ \pi_{H_i} \circ \sigma_i = [\psi]$$

which is a diffeomorphism. Therefore the map

$$\sigma_i^{\psi} := \psi \circ \sigma_i' = \psi \circ \sigma_i \circ [\psi]^{-1}$$

is a section of the projection $H'_I \to \mathcal{N}_{H'_i}$. This finishes the proof.

Clearly any diffeomorphism preserves the clean intersection property. This proves that any diffeomorphism ψ satisfying $\psi^* \lambda = \lambda + df$ with compactly supported fis an automorphism of the *structure of Liouville* σ -sectors. (See Definition 3.1 and 6.7.) This finishes the proof of the theorem.

Based on this discussion, we will unambiguously denote by $\operatorname{Aut}(M)$ the automorphism group of Liouville σ -sector (M, λ) as in the case of Liouville manifolds.

- **Remark 6.10.** (1) The above proof shows that the group $\operatorname{Aut}(M, \lambda)$ is manifestly the automorphism group of the structure of Liouville σ -sectors. We alert the readers that this is not manifest in the original definition of Liouville sectors from [GPS17], [GPS18].
 - (2) This simple characterization of the automorphism groups of Liouville σ sectors with corners enables one to define the bundle of Liouville sectors
 with corners in the same way for the case of Liouville manifolds (with
 boundary) without corners. See [OT20] for the usage of such bundles in the
 construction of continuous actions of Lie groups on the wrapped Fukaya
 category of Liouville sectors (with corners).
 - (3) Recall that the Liouville structure λ on M induces a natural contact structure on its ideal boundary $\partial_{\infty} M$. We denote the associated contact structure by ξ_{∞} . Then we have another natural map

$$\operatorname{Aut}(M,\lambda) \to \operatorname{Cont}(\partial_{\infty}M,\xi_{\infty})$$

where $(\partial_{\infty} M, \xi_{\infty})$ is the group of *contactomorphisms of the contact mani*fold $(\partial_{\infty} M, \xi_{\infty})$. (See [Gir17], [OT] for the details.)

(4) The different geometric nature of $(\partial_{\infty} M, \xi_{\infty})$ and $(\partial M, \lambda_{\partial})$ is partially responsible for the difficulty of the constructions of a pseudoconvex pair (ψ, J) in a neighborhood

 $\mathcal{N}bhd(\partial_{\infty} M \cup \partial M)$

such that the almost complex structures J is amenable to the (strong) maximum principle) for the (perturbed) pseudoholomorphic maps into the Liouville sectors as manifested in [Oh21].

7. Monoid of Liouville σ -sectors and smoothing profiles

As mentioned before, the enlarged set of manifolds with corners forms a monoid under the product. We now show that this monoidal structure restricts to the set of *Liouville* σ -sectors with corners by defining the product as a Liouville σ -sector with corners.

In this section, when we say 'sectorial', it always means σ -sectorial and will not mention this throughout the section, unless necessary.

7.1. Monoid of manifolds with corners. We start with some discussion on the boundary of manifolds with corners. We will regard the boundaries themselves as corners of codimension one and do not separate them from other corners, and call manifolds with boundary and corners just *manifolds with corners* for the simplicity of naming.

For a noncompact manifold with corner M, we have the standard definition of the *ideal boundary*, also called as the asymptotic boundary. In the present paper, to facilitate our exposition on the ideal boundary of the product of Liouville sectors later, we reserve the name 'ideal boundary' for the ideal boundary for the Liouville manifolds (with corners) and denote by

 $\partial_{\infty}^{\text{Liou}} M$

for the latter meaning. We reserve the name 'asymptotic boundary' for the ideal boundary in the context of topological spaces and denote them by

 $\partial_{\infty}M$

which is defined as follows.

We recall some basic definitions of the end and the asymptotic boundary (or also called the ideal boundary) of a noncompact manifold M. We adopt the definitions from [Ric63] where it is applied to a surface which however applies equally to general topological spaces.

Definition 7.1. Let M be a noncompact topological space. An *end* of M is an equivalence class of nested sequences $p = \{P_1 \supset P_2 \supset ...\}$ of connected unbounded regions in M such that

- (1) The boundary of P_i in M is compact for every i.
- (2) For any bounded subset A of M, $P_i \cap A = \emptyset$ for sufficiently large i

Two ends $p = \{P_1 \supset P_2 \supset ...\}$ and $q = \{Q_1 \supset Q_2 \supset ...\}$ are equivalent if for any n there is a corresponding integer N such that $P_n \subset Q_N$ holds and vice versa. We say an equivalence class

$$[\{P_1 \supset P_2 \supset \dots\}]$$

an end of M.

When M is a manifold of finite type, we assume that each end is cylindrical, i.e., we may assume the sequence $P_1 \supset P_2 \supset \cdots$ is topologically stable and assume that P_i is diffeomorphic to $\partial P_i \times [0, \infty)$ and ∂P_i is compact. (In the term of [Ric63] in two dimensions, such an end is called a *planar end.*) So from now on, we assume M is of finite type.

Then we define the notion of the *asymptotic boundary* of M as follows.

Definition 7.2 (Asymptotic boundary). The asymptotic boundary, denoted by $\partial_{\infty} M$ of M is a topological space equipped with the topology induced from ∂P_j for sufficiently large j.

Now we introduce the notion of *ideal completion*.

Definition 7.3 (Ideal completion). Let M be as above. The ideal completion denoted by \overline{M} is the coproduct

$$\overline{M} = M \coprod \partial_{\infty} M$$

equipped with the finest topology for which both inclusion maps $\partial_{\infty} M, M \to \overline{M}$ are continuous.

Now we restrict ourselves to the case of noncompact manifolds with boundary and corners.

It is also useful to consider the definition of the full boundary of the ideal completion in the study of pluri-subharmonic exhaustion functions of Int M.

Definition 7.4 (Full boundary of \overline{M}). We define the DM to be the boundary of \overline{M} which is the union of ∂M and $\partial_{\infty} M$

 $DM := \partial \overline{M} = \partial M \cup \partial_{\infty} M.$

Recall that for given two manifolds with corners X and Y, we have

$$\partial(X \times Y) = \partial X \times Y \bigcup X \times \partial Y$$

as a manifold with corners. Then we have the following

Lemma 7.5. Let (X, λ_X) and (Y, λ_Y) be two Liouville sectors (with corners). Then

$$\partial_{\infty}(X \times Y) := (\partial_{\infty}X \times \overline{Y}) \cup (\overline{X} \times \partial_{\infty}Y), D(X \times Y) := DX \times \overline{Y} \cup (\overline{X} \times DY).$$

Proof. The proof is a straightforward calculation and so omitted.

Example 7.6. Let (M, λ) be a Liouville sector and consider the splitting

$$\mathcal{N}bhd(\partial M) \cong F \times \mathbb{C}_{Re>0}$$

as the product of manifolds with boundary and corners. Then we have

$$\partial_{\infty}(F \times \mathbb{C}_{\operatorname{Re} \ge 0}) = (\partial_{\infty}F \times \overline{\mathbb{C}_{\operatorname{Re} \ge 0}}) \cup (\overline{F} \times \partial_{\infty}\mathbb{C}_{\operatorname{Re} \ge 0}).$$

We also have

$$D(F \times \mathbb{C}_{\text{Re} \ge 0}) = (DF \times \overline{\mathbb{C}_{\text{Re} \ge 0}}) \cup (\overline{F} \times D(\mathbb{C}_{\text{Re} \ge 0}))$$

which carries a natural structure of manifold with corners.

We also have the following which is one of the main reasons why the Floer moduli spaces arising in the wrapped Fukaya category and the symplectic cohomology are monoidal. (See Subsection 8.2 below.)

Corollary 7.7. Let ψ be an exhaustion function for a neighborhood Nbhd(DX) of the Liouville sector X. The level sets $\psi^{-1}(r)$ smoothly approximate $D(F \times \mathbb{C}_{\text{Re} \ge 0})$ as $r \to \infty$ in the obvious sense as a manifold with corners.

7.2. Monoid of Liouville σ -sectors with corners. In this section, Liouville sectors mean Liouville sectors with corners omitting 'with corners'.

We start with recalling the following standard definition of the ideal boundary of Liouville manifolds (M, λ) .

Definition 7.8 (Asymptotic Liouville rays). Let (M, λ) be a Liouville manifold and denote by Z its Liouville vector field. We call a Liouville trajectory escaping to infinity an *asymptotic Liouville rays* as time flows to $+\infty$. When dim M = 2, we also define a positive (resp. negative) asymptotic ray as it escapes to infinity as time flows to $\pm\infty$ respectively, as described in Section 2.

Definition 7.9 (Ideal boundary of Liouville manifolds). The ideal boundary $\partial_{\infty}^{\text{Liou}} M$ is the set of Liouville equivalence classes of asymptotic Liouville rays of Z_M .

The following is easy to check whose proof is omitted.

Lemma 7.10. Let (M, λ) be a Liouville sector. Then we have

$$\partial_{\infty}^{\text{Liou}} M = \partial_{\infty} M$$

We consider the product of Liouville sectors. We now show that the monoidal structure of the category of manifolds with corners restricts to the set of (intrinsic) *Liouville sectors with corners* by defining the product as a Liouville sector with corners.

Proposition 7.11. The product of Liouville σ -sectors (X, λ_X) and (Y, λ_Y)

$$(X \times Y, \pi_X^* \lambda_X + \pi_Y^* \lambda_Y)$$

is canonically a Liouville σ -sector with corners. In particular the set of Liouville σ -sectors with corners is a submonoid of the monoid of manifolds with boundary and corners under the product.

Proof. First we check that the vector field

$$Z_{X\times Y} := Z_X \oplus Z_Y$$

satisfies

$$Z_{X \times Y} \rfloor d(\pi_X^* \lambda_X + \pi_Y^* \lambda_Y) = (Z_X \oplus 0) \rfloor d(\pi_X^* \lambda_X) \oplus (0 \oplus Z_Y) \rfloor d(\pi_Y^* \lambda_Y)$$
$$= (Z_X \rfloor d\lambda_X) \oplus (Z_Y \rfloor d\lambda_Y) = Z_X \oplus Z_Y = Z_{X \times Y}$$

on $\operatorname{Int}(X \times Y)$. This shows that $Z_{X \times Y}$ is the Liouville vector field of the Liouville form $\pi_X^* \lambda_X + \pi_Y^* \lambda_Y$ on $\operatorname{Int}(X \times Y)$. Furthermore it is tangent to $\partial(X \times Y)$ near infinity if Z_X and Z_Y are tangent to $X \times \{y\}$ and $\{x\} \times Y$ near infinity respectively. This proves that $Z_{X \times Y}$ is tangent to the boundary

$$\partial(X \times Y) \setminus \partial X \times \partial Y$$

at infinity, where $\partial X \times \partial Y$ is a corner of $X \times Y$ of codimension 2.

Therefore a Liouville trajectory $\gamma : \mathbb{R} \to X \times Y$ is given by

$$\gamma(t) = (\gamma_X(t), \gamma_Y(t))$$

which is complete in the sense that either γ is forward complete or exits $\partial(X \times Y)$ in finite time.

Next we examine the σ -sectorial structure of the product $X \times Y$. Note that

$$\partial(X \times Y) = \partial X \times Y \bigcup X \times \partial Y.$$

And the characteristic foliation of $\partial(X \times Y)$ is given by $C \times \{y\}$ on $\partial X \times Y$ and $\{x\} \times C'$ respectively. Obviously they are homeomorphic to \mathbb{R} if C and C' are. Therefore we have the defining continuous section $\sigma_{\text{ref}} : \mathcal{N}_{\partial(X \times Y)} \to \partial(X \times Y)$ given by

$$\sigma_{\rm ref}(\ell) = \begin{cases} (\sigma_{\rm ref}^X(\pi_X(\ell)), \pi_Y(\ell)) & \text{on } \partial X \times Y \\ (\pi_X(\ell), \sigma_{\rm ref}^Y(\pi_Y(\ell))) & \text{on } X \times \partial Y \end{cases}$$

where σ_{ref}^X and σ_{ref}^Y the defining sections of the Liouville σ -sectors X, Y respectively. These prove that the pair (H_1, H_2) of

$$H_1 := \partial X \times Y, \quad H_2 := X \times \partial Y$$

is a sectorial collection. This finishes the proof.

Corollary 7.12. We have

$$\partial_{\infty}^{\text{Liou}}(X \times Y) = \partial_{\infty}^{\text{Liou}}X \times \overline{Y} \bigcup \overline{X} \times \partial_{\infty}^{\text{Liou}}Y$$
$$D(X \times Y) = DX \times \overline{Y} \bigcup \overline{X} \times DY.$$

50

Proof. The lemma immediately follows from Lemma 7.5 and Proposition 7.10. \Box

Remark 7.13. One can define the *compact* σ -sectorial domain with corners W in the same way as one does the Liouville domain, except that we need to add one requirement of the existence of *outer collaring* of ∂W which reads that the outward pointing Liouville vector fields along the various codimension zero strata of ∂M are compatible: Then can be smoothly interpolated to one another along the *outer collar* of the intersections of the codimension zero strata of ∂M . Our canonical description, Theorem 4.4 of the σ -splitting data at the sectorial corners enables us to prove the existence of such an outer collaring. Since we do not use this in the present paper, we do not elaborate this further here postponing its full explanation elsewhere.

7.3. Smoothing profiles and their products.

7.3.1. Compatible corner smoothing of $[0, \infty)^n$. In this subsection, we borrow the construction from [FOOO20, Section 18.5] and [Oh21, Section 3] which provides a family of local models of corner smoothing of

$$\mathbb{R}_{>0}^m = [0,\infty)^n$$

that is compatible with various m and with the S_m -symmetry for the coordinate swapping. Combination of constructions given in [FOOO20, Condition 18.21] and in [Oh21, Section 3] provides a family of compatible convex corner smoothing functions as follows.

Definition 7.14 (Convex smoothing functions of $\mathbb{R}^m_{\geq 0}$). A symmetric convex smoothing function on \mathbb{R}^k is a function $\varphi : \mathbb{R}^k \to \mathbb{R}$ satisfying the following:

- (CV1) The restriction $\varphi|_{\mathbb{R}^J}$ is $\mathsf{S}_{|J|}$ -invariant for all subsets $J \subset \underline{n} = \{1, \ldots, n\}$. Here $\mathbb{R}^J \subset \mathbb{R}^n$ is the obvious copy of $\mathbb{R}^{|J|}$.
- (CV2) $\text{Hess}(\varphi)$ is positive semi-definite everywhere.

(CV3) Hess($\varphi|_{\mathbb{R}^J}$) is compactly supported on \mathbb{R}^J for all subsets $J \subset \underline{n}$ for $|J| \ge 1$.

We denote the set thereof by $\mathfrak{Conv}_{\mathfrak{sm}}^{\mathsf{S}_k}(\mathbb{R}^k)$.

It is shown in [Oh21, Section 3] that the set is nonempty, which are also convex and so contractible.

We now fix a constant $\epsilon_0 > 0$ which will measure the size of Nbhd (∂M) once and for all. We will utilize a two-parameter family of such convex smoothing functions parameterized by $\epsilon > 0$ and $T_0 > 0$ as follows.

Definition 7.15 (Convex smoothing functions of $\mathbb{R}^m_{\geq 0}$). Let $\epsilon > 0$ be a sufficiently small constant and $T_0 > 0$ a sufficiently large constant such that

$$\epsilon_0 < 2T_0\sqrt{\epsilon} < \frac{3}{2}\epsilon_0.$$

Then we consider a family of functions $\varphi_m^{\epsilon} : \mathbb{R}_{\geq 0}^m \to \mathbb{R}_{\geq 0}$ that satisfies the following:

(1) We have

$$\varphi_m^{\epsilon}(x_1, \dots, x_m) = x_i \quad \text{when } x_i \ge 2T_0\sqrt{\epsilon}, \text{ and } 0 \le x_j \le \frac{\sqrt{\epsilon}}{4} \text{ for } j \ne i \qquad (7.1)$$
(2) $d\varphi(x_1, \dots, x_m) = 0$ if and only if $(x_1, \dots, x_m) = 0$.

7.3.2. Smoothing profiles. Recall from Definition 4.1 that ∂M is a union of a collection of cleanly intersecting hypersurfaces $H_1, \ldots, H_m \subset M$ (cylindrical at infinity) near each sectorial corner of ∂M of codimension k for some k with

$$C = H_1 \cap \dots \cap H_m.$$

The σ -splitting data given in Theorem 4.4 is the collection of maps

$$\widetilde{\Psi}_C : \mathrm{Nbhd}(\partial M) \to F_C \times \mathbb{R}^m_{>0}$$

associated to the corners satisfying

$$(\widetilde{\Psi}_C)_*(d\lambda) = \omega_{F_C} + \sum_{i=1} dR_i \wedge dI_i$$

with a canonically given symplectic from ω_{F_C} on F_C .

Definition 7.16 (Sectorial corner smoothing). Let C be a sectorial corner of M. Define the function

$$s_{m,\varphi} = -\log\varphi \circ \Psi_C^{\sigma} = -\log\varphi(R_1, \dots, R_m)$$
(7.2)

which we call a sectorial corner smoothing function.

Then we fix a contact-type hypersurface $S_0 \subset M$ and a Liouville embedding $S \times [0, \infty) \hookrightarrow M$ that equips a symplectization end of M. We denote by $s = s_{S_0}$ the associated radial function.

Definition 7.17 (End-profile function). Consider the function

$$s_{m_{\delta}+1,\varphi_{\delta}} := -\log\varphi_2\left(e^{-s_{m_{\delta},\varphi_{\delta}}}, e^{-s}\right) \tag{7.3}$$

at each sectorial corner C_{δ} . We glue these functions of the corners by a partition of unity and denote by

$$\mathfrak{s}_{\varphi}, \quad \varphi := \{\varphi_{\delta}\}$$

the resulting function, which we call an *end-profile function* of the end of M, $\partial_{\infty} M \cup \partial M$.

Condition 7.18 (Smoothing profile). (1) We fix a contact-type hypersurface $S_0 \subset M$ and the associated decomposition

$$M = W \cup_{\partial W} S_0 \times [0, \infty), \quad \partial W = S_0.$$

and the associated radial function $s = s_{S_0}$.

- (2) At each sectorial corner C_{δ} of $Nbhd(\partial_{\infty} M \cup \partial M)$, we fix the following data:
 - a splitting data $(F_{\delta}, \{(R_{\delta}, I_{\delta})\})$ with $F_{\delta} = (R_{\delta}, I_{\delta})^{-1}(0, 0)$,
 - convex smoothing function

$$\varphi_{\delta} = \varphi_{k_{\delta}+1} \in \mathfrak{Conv}_{\mathfrak{sm}}(\mathbb{R}^{k_{\delta}+1}).$$

and its associated sectorial corner smoothing function

$$s_{k_{\delta},\varphi}: \mathrm{Nbhd}(C_{\delta}) \to \mathbb{R}.$$

(3) An end profile function \mathfrak{s}_{φ} defined as in Definition 7.17.

7.3.3. Product of smoothing profiles. Let (X, λ_X) and (Y, λ_Y) be Liouville σ -sectors with corners.

The following shows that the σ -splitting data is monoidal whose proof is straightforward and so is omitted.

Lemma 7.19. For given σ -splitting data $(F_{\alpha}^{X}, \{(R_{\alpha}^{X}, I_{\alpha}^{X})\})$ of X along a sectorial corner C_{α} of codimension k_{α} and $(F_{\beta}^{Y}, \{(R_{\beta}^{Y}, I_{\beta}^{Y})\})$ of Y along C_{β} of codimension k_{β} , their product

$$(F_{\alpha,\beta}^{X\times Y}, \{R_{\alpha,\beta}^{X\times Y}, I_{\alpha,\beta}^{X\times Y}\})$$

:= $\left(F_{\alpha}^{X} \times Y \bigcup X \times F_{\beta}^{Y}, \left(\{(\pi_{X}^{*}R_{\alpha}^{X}, \pi_{X}^{*}I_{\alpha}^{X})\} \bigcup \{(\pi_{Y}^{*}R_{\beta}^{Y}, \pi_{Y}^{*}I_{\beta}^{Y})\}\right)\right)$ (7.4)

is a σ -splitting data for $X \times Y$.

We also have

$$D(X \times Y) = \partial_{\infty}^{\text{Liou}}(X \times Y) \bigcup \partial(X \times Y)$$
$$= DX \times \overline{Y} \bigcup \overline{X} \times DY.$$

Remark 7.20. An upshot of this function \mathfrak{s}_{φ} is that it is *J*-convex for sectorial complex structures *J* to be introduced later in that $-d(d\mathfrak{s}_{\varphi} \circ J) \geq 0$ as a (1,1)-current.

Next equip each of X and Y with an end-profile function

$$\mathfrak{s}_X := \mathfrak{s}_{\varphi_X}, \quad \mathfrak{s}_Y := \mathfrak{s}_{\varphi_Y}$$

defined as in Definition 7.17.

Equip X, Y with end-profile functions \mathfrak{s}_X and \mathfrak{s}_Y associated to the collections

$$\{s_{k_{\alpha}+1,\varphi_{\alpha}}, \{s_{k_{\beta}+1,\varphi_{\beta}}\}.$$

with

$$s_{k_{\alpha}+1,\varphi_{\alpha}} = -\log \varphi_2(e^{-s_{k_{\alpha},\varphi_{\alpha}}}, e^{-s})\}$$

$$s_{k_{\beta}+1,\varphi_{\beta}} = -\log \varphi_2(e^{-s_{k_{\alpha},\varphi_{\alpha}}}, e^{-s})$$
(7.5)

for convex symmetric smoothing functions $\varphi_{k_{\alpha}+1} : \mathbb{R}^{k_{\alpha}+1}_+ \to \mathbb{R}$ and $\varphi_{k_{\beta}+1} : \mathbb{R}^{k_{\beta}+1}_+ \to \mathbb{R}$.

Next we state the following lemma whose proof is immediate by definition and so left to the readers.

Lemma 7.21. Take symplectization radial functions s^X and s^Y of X and Y respectively. Then the function $-\log \varphi_2(e^{-s^X}, e^{-s^Y})$ is transversal to the product Liouville vector fields $Z^X \times Z^Y$ on $s_{X \times Y}^{-1}(r)$ for all sufficiently large r > 0. We denote

$$s^{X \times Y} := -\log \varphi_2(e^{-s^X}, e^{-s^Y})$$

and call it the *(smoothed) product radial coordinate* of s^X and s^Y . Furthermore, we have

$$s^{X \times Y} = \begin{cases} \pi_X^* s^X & s^X \ge R^+, \ e^{-s^Y} \ge \epsilon_\varphi \\ \pi_Y^* s^Y & s^Y \ge R^+, \ e^{-s^X} \ge \epsilon_\varphi \end{cases}$$
(7.6)

for a fixed small constant $\epsilon_{\varphi} > 0$ depending on φ and for a sufficiently large constant $R^+ > 0$.

Definition 7.22 (Product end-profile function). We define the product end-profile function

$$\mathfrak{s}_X *_{\varphi} \mathfrak{s}_Y : X \times Y \to \mathbb{R}$$

by taking the union of the local convex interpolation

$$\{s_{k_{\alpha}+k_{\beta}+1,\varphi}: \mathbb{N}bhd(C_{\alpha} \times C_{\beta}) \to \mathbb{R}\}$$

defined by

$$s_{k_{\alpha}+k_{\beta}+1,\varphi} := -\log\varphi_3\left(e^{-s_{k_{\alpha},\varphi_{\alpha}}}, e^{-s_{k_{\alpha},\varphi_{\alpha}}}, e^{-s^{X\times Y}}\right)$$

followed by taking a partition of unity thereof on $X \times Y$.

We unravel by definition the arguments inside parenthesis into

$$s_{k_{\alpha},\varphi_{\alpha}} = -\log\left(\varphi_{k_{\alpha}}(R_{\alpha,1},\ldots,R_{\alpha,k_{\alpha}})\right)$$

$$s_{k_{\beta},\varphi_{\beta}} = -\log\left(\varphi_{k_{\beta}}(R_{\beta,1},\ldots,R_{\beta,k_{\beta}})\right)$$

and hence we have

$$:= -\log\varphi_3\left(\varphi_{k_{\alpha}}(R_{\alpha,1},\ldots,R_{\alpha,k_{\alpha}}),\varphi_{k_{\beta}}(R_{\beta,1},\ldots,R_{\beta,k_{\beta}}),\varphi_2\left(e^{-s^X},e^{-s^Y}\right)\right)$$
(7.7)

Lemma 7.23. The function

$$\mathfrak{s}_X *_{\varphi} \mathfrak{s}_Y : X \times Y \to \mathbb{R}$$

is an exhaustion function of $Nbhd(D(X \times Y))$ and its level sets smoothly approximate the full boundary $D(X \times Y)$.

Proof. We first note that the function

$$-\log\varphi_3(e^{-x}, e^{-y}, e^{-z})$$

is a convex three-term interpolation of the functions x, y and z on $\mathbb{R}^3_{\geq 0}$ whose level sets smoothly approximates of the boundary $\partial \mathbb{R}^3_{\geq 0}$. This being said, the first statement follows from definition of the convex smoothing functions φ_k , and the second is obvious by construction.

8. Sectorial almost complex structures and gradient-sectorial Lagrangians

In this section, we assume that we are given the splitting data and end-profile functions

$$\mathfrak{s}_X = \mathfrak{s}_{\varphi^X}, \quad \mathfrak{s}_Y := \mathfrak{s}_{\varphi^Y}$$

respectively for X and Y. Then we consider a product end-profile function

$$\mathfrak{s}_{X\times Y}:=\mathfrak{s}_X*_\varphi\mathfrak{s}_Y$$

as defined before. Note that this product depends on the choice of convex functions φ which is, however, a contractible choice.

8.1. **Definition of sectorial almost complex structures.** Note that a splitting data

$$\mathcal{N}bhd(\partial M) \cong F \times \mathbb{C}^k_{\text{Re} \ge 0}, \quad \{(R^{\sigma}_i, I^{\sigma}_i)\}_{i=1}^k$$

provides a foliation, denoted by \mathcal{F}_F , of symplectic hypersurfaces whose leaves are given by

$$F \times \{(x,y)\}_{(x,y) \in \mathbb{C}_{\mathrm{Re} \ge 0}}$$

We now introduce the following class of almost complex structures.

Definition 8.1 (Sectorial almost complex structures). Let (M, λ) be a Liouville sector with boundary and corners equipped with a smoothing profile whose associated splitting is given by

$$\mathbb{N}bhd(\partial M) \cong F \times \mathbb{C}^k_{\mathrm{Re}>0}, \quad \{(R^{\sigma}_i, I^{\sigma}_i)\}_{i=1}^k$$

and whose associated end-profile function is \mathfrak{s}_{φ} .

An ω -tame almost complex structure J on a Liouville sector is said to be *sectorial* (with respect to the given smoothing profile) if J satisfies the following:

(J1) [\mathcal{F}_F is *J*-complex] In a neighborhood of Nbhd^Z(∂M) of ∂M , we require

$$J\left(T^*F \oplus 0_{\operatorname{span}\left\{dR_i^{\sigma}, dI_i^{\sigma}\right\}_{i=1}^k}\right) \subset T^*F \oplus 0_{\operatorname{span}\left\{dR_i^{\sigma}, dI_i^{\sigma}\right\}_{i=1}^k},\tag{8.1}$$

and J restricts to an almost complex structure of contact-type on F.

(J2) [($\mathfrak{s}_{\varphi}, J$) is a pseudoconvex pair] In a neighborhood $\operatorname{Nbhd}^{\mathbb{Z}}(\partial M) \cup \partial_{\infty}$) of $\partial M \setminus \operatorname{Nbhd}(\partial_{\infty} M)$, we have

$$-d(d\mathfrak{s}_{\varphi}\circ J)\geq 0$$

is a positive (1, 1)-current.

We denote by $\mathcal{J}_{\mathfrak{s}_{\omega}}^{\mathrm{sec}} = \mathcal{J}_{\mathfrak{s}_{\omega}}^{\mathrm{sec}}(M)$ the set of sectorial almost complex structures.

Such an almost complex structures is also studied in [Oh21] where the common asymptotically Z-invariant Lagrangians are still adopted as branes of the wrapped Fukaya category. Obviously any almost complex structure of the form

$$-d\mathfrak{s}_{\varphi}\circ J=\lambda+df$$

satisfying (J1) is sectorial, which includes both the class of κ -sectorial almost complex structures and that of λ -sectorial almost complex structures from [Oh21]. In particular $\mathcal{J}_{\mathfrak{s}_{\alpha}}^{\operatorname{sec}}$ is nonempty and contractible.

The following proposition is the reason why we consider the sectorial almost complex structures instead of λ -sectorial ones from [Oh21]. Such a natural inclusion does not exist for the set of λ -sectorial almost complex structures thereform.

Proposition 8.2. Consider the product $J_1 \times J_2$ for $J_1 \in \mathcal{J}_{\mathfrak{s}_X}^{\text{sect}}$ and $J_2 \in \mathcal{J}_{\mathfrak{s}_Y}^{\text{sect}}$. Then $(\mathfrak{s}_{X \times Y}, J_1 \times J_2)$ satisfies the following:

(1) $-d\mathfrak{s}_{X\times Y} \circ (J_1 \times J_2) = f_X \pi_1^* \lambda_1 + f_Y \pi_2^* \lambda_2$ for positive function $f_X, f_Y : X \times Y \to \mathbb{R}$ of the form

$$f_X = \psi_1(\mathfrak{s}_X, \mathfrak{s}_Y), \qquad f_Y = \psi_2(\mathfrak{s}_X, \mathfrak{s}_Y)$$

where $\psi_1, \psi_2 > 0$.

(2) $(\mathfrak{s}_{X\times Y}, J_1 \times J_2)$ is a pseudoconvex pair.

In particular we have

$$\mathcal{J}_{\mathfrak{s}_X}^{\mathrm{sec}} \times \mathcal{J}_{\mathfrak{s}_Y}^{\mathrm{sec}} \subset \mathcal{J}_{\mathfrak{s}_X \times Y}^{\mathrm{sec}}.$$

Proof. We compute

$$-d\mathfrak{s}_{X\times Y}\circ(J_1\times J_2) = -\frac{\partial\varphi_2}{\partial x}(\mathfrak{s}_{X\times Y})d\mathfrak{s}_X\circ J_1 - \frac{\partial\varphi_2}{\partial y}(\mathfrak{s}_{X\times Y})d\mathfrak{s}_Y\circ J_2$$
$$= \frac{\partial\varphi_2}{\partial x}(\mathfrak{s}_{X\times Y})\pi_X^*\lambda_1 + \frac{\partial\varphi_2}{\partial y}(\mathfrak{s}_{X\times Y})\pi_Y^*\lambda_2.$$

By setting

$$f_X := \frac{\partial \varphi_2}{\partial x}(\mathfrak{s}_{X \times Y}), \quad f_Y := \frac{\partial \varphi_2}{\partial y}(\mathfrak{s}_{X \times Y})$$

we have finished the first statement.

To prove the second statement, we keep computing

$$-d(d\mathfrak{s}_{X\times Y}\circ(J_1\times J_2)) = \frac{\partial^2\varphi_2}{\partial x^2}(\mathfrak{s}_{X\times Y})(d\mathfrak{s}_X\circ J_1)\wedge d\mathfrak{s}_X +\frac{\partial\varphi_2}{\partial x}(\mathfrak{s}_{X\times Y})d\lambda_1 + \frac{\partial\varphi_2}{\partial y}(\mathfrak{s}_{X\times Y})d\lambda_2 +\frac{\partial^2\varphi_2}{\partial y^2}(\mathfrak{s}_{X\times Y})(d\mathfrak{s}_Y\circ J_2)\wedge d\mathfrak{s}_Y.$$

The first and the third summand are nonnegative currents since $d\mathfrak{s}_X \circ J_1 \wedge d\mathfrak{s}_X$ and $d\mathfrak{s}_Y \circ J_2 \wedge d\mathfrak{s}_Y$ are nonnegative (1, 1)-currents and

$$\frac{\partial^2 \varphi_2}{\partial x^2}, \quad \frac{\partial^2 \varphi_2}{\partial x^2} \ge 0$$

On the other hand, the second summand is nonnegative because $d\lambda$ is a positive (1, 1)-current and

$$\frac{\partial \varphi_2}{\partial x}, \quad \frac{\partial \varphi_2}{\partial y} \ge 0.$$

This finishes the proof.

As usual, we denote by g_J the ω -tame metric given by

$$g_J(v,w) := \frac{\omega(v,Jw) + \omega(w,Jv)}{2}$$

for each given sectorial almost complex structure J.

8.2. Gradient-sectorial Lagrangians and their products. As mentioned before, the enlarged set of manifolds with corners forms a monoid under the product.

We next identify the set of objects for the wrapped Fukaya category of a Liouville sectors with corners. We omit other brane data such as the orientation or the spin structure since they are not changed from the usual description in the literature and do not enter the discussion of the current paper.

We introduce the following normalized gradient vector field on a neighborhood Nbhd(DM)

$$Z_{\mathfrak{s}} := \frac{\operatorname{grad}\mathfrak{s}}{|\operatorname{grad}\mathfrak{s}|^2} \tag{8.2}$$

where the gradient is taken with respect to the metric $g = g_J$. By definition, we have $Z_{\mathfrak{s}}[\mathfrak{s}] = 1 > 0$ on Nbhd(DM).

The following is the key lemma towards our proof of monoidality of the class of gradient-sectorial Lagrangian submanifolds we introduce below. Its proof is an immediate consequence of the definition of convex smoothing function φ_2 from [Oh21, Section 3.2], more specifically by (7.6).

Lemma 8.3. For any given $\epsilon_0 > 0$, there exists sufficiently large $r_0 > 0$ such that

$$\mathfrak{s}_{X \times Y} = s^{X \times Y} = -\log \varphi_2(e^{-s^X}, e^{-s^Y})$$

whenever dist $(x, \partial(X \times Y)) \ge \epsilon_0$ and $\mathfrak{s}_{X \times Y}(x) \ge r_0$.

Now we are ready to introduce the notion of gradient-sectorial Lagrangians with respect to the end-profile function \mathfrak{s}_{φ} . This a variation of sectorial Lagrangians introduced in [Oh21]. Equip M with any tame metric such as g_J given above for a sectorial J.

Definition 8.4 (Gradient-sectorial Lagrangian branes). Let (M, λ) be a Liouville sector with corners. Let \mathfrak{s} be the end-profile function associated to a given smoothing profile. We say an exact Lagrangian submanifold L of (M, ω) is gradient-sectorial if

- (1) $L \subset \operatorname{Int} M \setminus \partial M$, and $\operatorname{dist}(L, \partial M) > 0$.
- (2) There exist a sufficiently large $r_0 > 0$ such that $L \cap \mathfrak{s}^{-1}([r_0, \infty))$ is $Z_{\mathfrak{s}}$ -invariant.

Remark 8.5 (Comparison with sectorial Lagrangians [Oh21]). The vector field $Z_{\mathfrak{s}}$ coincides with the Liouville vector field Z on the symplectization end $[0,\infty) \times S_0$ of M if we set $\mathfrak{s} = s$ and J is Z-invariant so that

$$\frac{\partial}{\partial s} = Z$$

in the Z-invariant splitting

$$TM \cong \operatorname{span}_{\mathbb{R}} \{Z, R_{\theta}\} \oplus \xi_{\{s=0\}}$$

with respect $\omega = d(e^s \theta)$, where R_{θ} is the Reeb vector field of the contact-type hypersurface S_0 equipped with the contact form $\theta = \iota_{S_0}^* \lambda$. In this regard, the notion of gradient-sectorial Lagrangian branes is a generalization of Z-invariantat-infinity Lagrangian submanifolds. However while the latter definition does not respect the product operation of Liouville manifolds, the definition of gradientsectorial Lagrangian does under the product operations of the manifold together with that of product operation of end-profile functions

$$(\mathfrak{s}_X,\mathfrak{s}_Y)\mapsto\mathfrak{s}_X*_{\varphi}\mathfrak{s}_Y=:\mathfrak{s}_{X\times Y}.$$

The gradient-sectorial Lagrangians have the following fundamental monoidal property under the product operation. Such monoidality fails for the commonly adopted Z-invariant Lagrangians in the study of wrapped Fukaya category which results in much nuisance in the construction of Künneth-type mappings. (See [Gao17, GPS17, GPS18] for example.) Here crucially enters the special interpolation property of the end-profile function \mathfrak{s}_{φ} having the expression

$$s_{k+1,\varphi} = -\log\varphi(R_1,\cdots,R_k,e^{-s})$$

on the ceiling of each sectorial corner of $Nbhd(\partial M)$ slightly away from the corner $\partial_{\infty} M \cap \partial M$.

Theorem 8.6. Let (X, ω_X) and (Y, ω_Y) be Liouville sectors with corners equipped with a smoothing profile. Let $\mathfrak{s}_{X \times Y}$ be the product end-profile function given in Definition 7.22. Then for any gradient-sectorial Lagrangian branes L_1 and L_2 , the product $L_1 \times L_2$ is also gradient-sectorial with respect to $\mathfrak{s}_{X \times Y}$.

Proof. Obviously we have

 $L_1 \times L_2 \subset \operatorname{Int}(X \times Y)$

if $L_1 \subset \operatorname{Int} X$ and $L_2 \subset \operatorname{Int} Y$, and

$$\operatorname{dist}(L_1 \times L_2, \partial(X \times Y)) \ge \min\{\operatorname{dist}(L_1, \partial X), \operatorname{dist}(L_2, \partial Y)\} > 0.$$

This shows that Definition 8.4 (1) is satisfied for $L_1 \times L_2$ This also implies

$$(L_1 \times L_2) \cap \mathfrak{s}_{X \times Y}^{-1}([r_0, \infty)) \cap R^{-1}((0, e^{-r_0}]) = \emptyset$$

for all $R = R_{\delta_X,i}$ (resp. $R = R_{\delta_Y,j}$) at any sectorial corner of X (resp. of Y), because we have $L_X \cap R^{-1}(0, e^{-r_0}]) = \emptyset$ (resp. $L_Y \cap R^{-1}(0, e^{-r_0}]) = \emptyset$), whenever

$$e^{-r_0} < \min\{\operatorname{dist}(L_1, \partial X), \operatorname{dist}(L_2, \partial Y)\})$$

or equivalently whenever

$$\max\{-\log \operatorname{dist}(L_1, \partial X), -\log \operatorname{dist}(L_2, \partial Y)\} > r_0.$$

Therefore from the definition of $\mathfrak{s}_{X \times Y}$ in Definition 7.22, (7.6) and the definition of sectorial Lagrangians above, we obtain

$$\mathfrak{s}_{X \times Y} = \begin{cases} \pi_X^* s^X & \text{on } (L_1 \times L_2) \cap \{s^Y \ge r_0\} \\ \pi_Y^* s^Y & \text{on } (L_1 \times L_2) \cap \{s^X \ge r_0\}. \end{cases}$$

This and the splitting property of $J_1 \times J_2$ give rise to

$$\operatorname{grad}_{g_{J_1 \times J_2}} \mathfrak{s}_{X \times Y} = \operatorname{grad}_{g_{J_1}} \mathfrak{s}^X \oplus 0$$

on $(L_1 \times L_2) \cap \{s^Y \ge r_0\}$ and equal $0 \oplus Z_{\mathfrak{s}^Y}$ on $(L_1 \times L_2) \cap \{s^X \ge r_0\}$ respectively.

Therefore it follows from $Z_{\mathfrak{s}_X}$ -invariance of L_1 and $Z_{\mathfrak{s}_Y}$ -invariance of L_2 at infinity that $Z_{\mathfrak{s}^X \times Y}$ is tangent to

$$(L_1 \times L_2) \cap \mathfrak{s}_{X \times Y}^{-1}([r,\infty))$$

for all sufficiently large r > 0. We remark that the open subset $\mathfrak{s}_{X \times Y}^{-1}((r, \infty))$ is a neighborhood of $\partial_{\infty}(X \times Y)$ and that on any given neighborhood

$$\operatorname{Nbhd}(\partial_{\infty}(X \times Y)) = \operatorname{Nbhd}(\partial_{\infty}X \times Y \cup X \times \partial_{\infty}Y)$$

we can choose r sufficiently large so that

$$(L_1 \times L_2) \cap \mathfrak{s}_{X \times Y}^{-1}([r, \infty)) \subset \operatorname{Nbhd}(\partial_\infty(X \times Y))$$

This proves that $L_1 \times L_2$ satisfies Condition 2 of Definition 8.4 and hence the proof.

8.3. Strong maximum principle and gradient-sectorial Lagrangian branes. In this subsection, we explain how the sectorial almost complex structure nicely pairs with gradient-sectorial Lagrangian branes to become amenable to the strong maximum principle and hence gives rise to fundamental confinement results for various Floer-type equations. We will just illustrate this for the Floer equation which produces the structure maps of the associated A_{∞} category, leaving the proofs for other cases to [Oh21] which handles the more subtle case of λ -sectorial almost complex structures and Z-invariant-at-infinity Lagrangian branes.

Let J be a sectorial almost complex structure of the Liouville sectors with corners M. Consider a (k + 1)-tuple (L_0, \ldots, L_k) of gradient-sectorial Lagrangian submanifolds. We denote

$$\Sigma = D^2 \setminus \{z_0, \dots, z_k\}$$

and equip Σ with strip-like coordinates (τ, t) with $\pm \tau \in [0, \infty)$ and $t \in [0, 1]$ near each z_i .

Then for a given collection of intersection points $p_i \in L_i \cap L_{i+1}$ for i = 0, ..., k, we wish to study maps $u : \Sigma \to M$ satisfying the Cauchy-Riemann equation

$$\begin{cases} \overline{\partial}_J u = 0\\ u(\overline{z_i z_{i+1}}) \subset L_i & i = 0, \dots k\\ u(\infty_i, t) = p_i, & i = 0, \dots k. \end{cases}$$

$$(8.3)$$

The following theorem will establish both vertical and horizontal confinement results simultaneously.

Theorem 8.7. Suppose that there exists some $\delta > 0$ such that

$$\operatorname{dist}(L_i, \partial M) \ge \delta$$

for all i = 0, ..., k. Let u be a solution to (8.3). Then there exists a sufficiently large r > 0 such that

Image
$$u \subset (\mathfrak{s}_{\varphi})^{-1}((-\infty, r])$$
 (8.4)

Proof. First all L_i 's are contained in $\{R > \delta\} \subset \text{Int } M$ since we have

$$\min\{d(L_i, \partial M) \mid i = 0, \dots, k\} > \delta$$

by the hypothesis.

Then by definition of sectorial almost complex structure J, J is associated to a given splitting data and the end-profile function \mathfrak{s}_{φ} . Since a neighborhood of $\partial_{\infty} M \cup \partial M$ is exhausted by the family of hypersurfaces

$$(\mathfrak{s}_{\varphi})^{-1}(r)$$

for $r \geq 0$, it is enough to prove

$$\operatorname{Image} u \subset (\mathfrak{s}_{\varphi})^{-1}((-\infty, r]) \tag{8.5}$$

for some r > 0. We first recall that du is *J*-holomorphic and satisfies $-d(d\mathfrak{s}_{\varphi} \circ J) \ge 0$ from the definition of pseudoconvex pair $(\mathfrak{s}_{\varphi}, J)$. Since u is *J*-holomorphic, we obtain

$$d\left(\mathfrak{s}_{\varphi}\circ u\right)\circ j=d\mathfrak{s}_{\varphi}\circ J\circ du=u^{*}(d\mathfrak{s}_{\varphi}\circ J)$$

By taking the differential of the equation, we derive

$$-d\left(d\left(\mathfrak{s}_{\varphi}\circ u\right)\circ j\right) = -u^{*}\left(d(d\mathfrak{s}_{\varphi}\circ J)\right) \geq 0.$$

In particular, the function $\mathfrak{s}_{\varphi} \circ u$ is a subharmonic function and cannot carry an interior maximum on $\mathbb{R} \times [0, 1]$ by the maximum principle.

Next we will show by the strong maximum principle that u cannot have a boundary maximum in a neighborhood of $\partial_{\infty} M \cup \partial M$ either. This will then enable us to obtain a C^0 confinement result

Image
$$u \subset \{\mathfrak{s}_{\varphi} \leq r_0\}$$

for any finite energy solution u with fixed asymptotics given in (8.3) provided r_0 is sufficiently large.

Now suppose to the contrary that $\mathfrak{s}_{\varphi} \circ u$ has a boundary local maximum point $z' \in \partial D^2 \setminus \{z_0, \ldots, z_k\}$. By the strong maximum principle, we must have

$$0 < \frac{\partial}{\partial \nu}(\mathfrak{s}_{\varphi}(u(z'))) = d\mathfrak{s}_{\varphi}\left(\frac{\partial u}{\partial \nu}(z')\right)$$
(8.6)

for the outward unit normal $\frac{\partial}{\partial \nu}|_{z'}$ of $\partial \Sigma$. unless $\mathfrak{s}_{\varphi} \circ u$ is a constant function in which case there is nothing to prove. Let (r, θ) be an isothermal coordinate of a neighborhood of $z' \in \partial \Sigma$ in (Σ, j) adapted to $\partial \Sigma$, i.e., such that $\frac{\partial}{\partial \theta}$ is tangent to $\partial \Sigma$ and $|dz|^2 = (dr)^2 + (d\theta)^2$ for the complex coordinate $z = r + i\theta$ and

$$\frac{\partial}{\partial \nu} = \frac{\partial}{\partial r} \tag{8.7}$$

along the boundary of Σ . Since u is J-holomorphic, we also have

$$\frac{\partial u}{\partial r} + J \frac{\partial u}{\partial \theta} = 0.$$

Therefore we derive

$$d\mathfrak{s}_{\varphi}\left(\frac{\partial u}{\partial \nu}(z')\right) = d\mathfrak{s}_{\varphi}\left(-J\frac{\partial u}{\partial \theta}(z')\right).$$

By the \mathfrak{s}_{φ} -gradient sectoriality of L and the boundary condition $u(\partial \Sigma) \subset L$, both $Z_{\mathfrak{s}_{\varphi}}(u(z'))$ and $\frac{\partial u}{\partial \theta}(z')$ are contained in $T_{u(z')}L$, which is a $d\lambda$ -Lagrangian subspace. Therefore we have

$$0 = d\lambda \left(Z_{\mathfrak{s}_{\varphi}}(u(z')), \frac{\partial u}{\partial \theta}(z') \right) = d\lambda \left(Z_{\mathfrak{s}_{\varphi}}(u(z')), J\frac{\partial u}{\partial \nu}(z') \right)$$
$$= g_J \left(Z_{\mathfrak{s}_{\varphi}}(z'), \frac{\partial u}{\partial \nu}(z') \right) = \frac{1}{|Z_{\mathfrak{s}_{\varphi}}(u(z'))|^2} d\mathfrak{s}_{\varphi} \left(\frac{\partial u}{\partial \nu}(z') \right)$$

where the last equality follows from the definition of normalized gradient vector field $Z_{\mathfrak{s}_{\varphi}}$. This is a contradiction to (8.6) (unless $\mathfrak{s}_{\varphi} \circ u$ is constant) and hence the function $\mathfrak{s}_{\varphi} \circ u$ cannot have a boundary maximum either. This then implies

$$\max \mathfrak{s}_{\varphi} \circ u \leq \max \{ \mathfrak{s}_{\varphi}(p_i) \mid i = 0, \dots, k \}$$

By setting

$$r_0 = \max\{\mathfrak{s}_{\varphi}(p_i) \mid i = 0, \dots, k\} + 1,$$

we have finished the proof.

We remark that the constant $\max\{\mathfrak{s}_{\varphi}(p_i) \mid i = 0, ..., k\}$ (and so r_0) depends only on the intersection set

$$\bigcup_{i=0}^{k} L_i \cap L_{i+1} (\mod k)$$

and not on the maps u.

9. DISCUSSION AND WHAT TO DO

9.1. **Pseudoconvex pairs and** Z-invariant-at-infinity Lagrangian branes. A construction of sectorial almost complex structures is given in [Oh21] whose details we refer readers thereto. A subtle difficulty to overcome in the construction of a pseudoconvex pair (ψ, J) in [Oh21] lies in the fact that for given Liouville sector with boundary and corners the asymptotic boundary $\partial_{\infty} M$ is of contact type and does not form a coisotropic collection when it is added to the coisotropic collection

$$\{H_1, H_2, \cdots, H_m\}$$

associated to a sectorial corner δ of the boundary ∂M . (Compare the definitions of $\mathfrak{s}_{k,\varphi}$ and $\mathfrak{s}_{k+1,\varphi}$: The former one involves interpolations between the sectorial corners only, while the latter involves both sectorial corners and the ceiling corner.)

This destroys the contact-type property of the union

$$DM = \partial_{\infty} M [] \partial M$$

in the sense that it may not be approximated by the hypersurfaces of contact-type, unless the sector is sufficiently expanded in the horizontal direction. (See [GPS17, Lemma 2.31].) However [Oh21, Theorem 1.2.3] shows that it admits an exhaustion function ψ that becomes *J*-convex for some ω -tame almost complex structures *J* which also satisfies

$$-d\psi \circ J = \lambda$$

Such a pair is called a *Liouville-pseudoconvex pair*, or more specifically a λ -pseudoconvex pair. It is shown in [Oh21] that for any Liouville-pseudoconvex pair (ψ, J) , the function $\psi \circ u$ satisfies the (interior) maximum principle as well as the strong maximum principle when a *J*-holomorphic curve u satisfies the boundary condition attached to a *Z*-invariant-at-infinity Lagrangian submanifold. The pairs constructed in [Oh21] are nothing but those consisting of the pair (ψ, J) with

 $\mathfrak{s}_{\varphi,\kappa} = \kappa$ -wiggled end-profile function, $J = \lambda$ -sectorial almost complex structures (9.1)

precise definitions of which we refer to [Oh21]. The aforementioned difficulty is what was overcome in [Oh21] as a byproduct of the existence result of λ -sectorial almost complex structures introduced therein. The main task in [Oh21] then is to interpolate the two requirements on the intersection Nbhd $(\partial_{\infty} M \cap \partial M)$ in the way that relevant maximum and strong maximum principle are still applicable. This construction of J in [Oh21] requires to unveil background geometry of Liouville sectors with corners and to go through a careful pointwise consideration of almost complex structures near the corner $\partial_{\infty} M \cap \partial M$ to reveal what presents the obstruction to interpolating the aforementioned two geometric structures, presymplectic geometry of ∂M and the Liouville geometry of Nbhd $(\partial_{\infty} M)$, near the corner so that Z-invariant Lagrangian boundary condition becomes amenable to the strong maximum principle.

9.2. Relationship with the Künneth-type formulae in Floer theory. One main consequence of Lemma 7.23 combined with the usage of sectorial almost complex structures and gradient-sectorial Lagrangians in the present paper is the following monoidality of the various Floer moduli spaces under the product of Liouville sectors.

Let K = K(t, x) be a sectorial Hamiltonian with respect to the end-profile function \mathfrak{s}_X , i.e., $K = \rho \circ \mathfrak{s}_X$ of the function $\rho : \mathbb{R} \to \mathbb{R}$ with $\rho' \ge 0$ and $\operatorname{supp} \rho'$ having compact support. We denote by

$$\mathcal{M}(M; J, K), \quad \mathcal{M}(M, L_0, L_1; J, K)$$

the moduli spaces of the Hamiltonian-perturbed Floer trajectories for the closed string and for the open string cases respectively, and by

$$\mathcal{M}(M,\mathcal{L};J), \quad \mathcal{L}=(L_0,\cdots,L_k)$$

the moduli spaces of the *J*-holomorphic polygons entering in the construction of (wrapped) Fukaya category $\mathsf{Fuk}(M)$.

The following, especially Statement (2), is an immediate consequence of Lemma 7.23 and Theorem 8.7 applied to the pseudoconvex pair

$$(\mathfrak{s}_{M_1} *_{\varphi} \mathfrak{s}_{M_2}, J_1 \times J_2)$$

under the boundary condition of product gradient-sectorial Lagrangians.

Corollary 9.1. Let (\mathfrak{s}_{M_i}, J) be a pair of end-profile function of Liouville sector (M_i, λ_i) and J_i its associated sectorial almost complex structure, respectively for i = 1, 2. Then we have the following natural inclusion maps:

(1) For any sectorial pair K_1 and K_2 of Hamiltonians on M_1 and M_2 respectively, we have

 $\mathcal{M}(M_1, L_1; J_1, K_1) \times \mathcal{M}(M_1, L_1; J_1, K_1) \to \mathcal{M}(M_1 \times M_2, L_1 \times L_2; J_1 \oplus J_2, K_1 \oplus K_2)$

(2) For any tuple of Lagrangians $\mathcal{L}_i = (L_1^1, L_1^2, \dots, L_1^k)$ for i = 1, 2, we have

 $\mathcal{M}(M_1,\mathcal{L}_1) \times \mathcal{M}(M_2,\mathcal{L}_2) \to \mathcal{M}(M_1 \times M_2,\mathcal{L}_1 \times \mathcal{L}_2).$

Similar proof can be given to prove Statement (1) whose details we refer to [Oh21] which deals with the more subtle case of the *Liouville-pseudoconvex pairs* ($\mathfrak{s}_{\varphi,\kappa}, J$) of (9.1) under the boundary condition of *Z*-invariant Lagrangian submanifolds.

An immediate consequence of this corollary will be that all the Künneth-type maps in the studies of symplectic cohomology (from Statement (1) above) and of the Hochschild homology (and cohomology) (from Statement (2) above) of the wrapped Fukaya category have chain-level monoidal property under the product operation of the Liouville sectors with corners, which follow rather straightforwardly by the algebraic arguments in homological algebra given in [Sei08] or in its references or in the one as summarized in [GPS18, Appendix B]. We will elaborate this remark in a sequel to the present paper. We also refer readers to [Amo17, Amo16], [Fuk17] for the relevant study of tensor products in the filtered setting of compact symplectic manifolds.

9.3. Fukaya categories of Liouville manifolds of infinite type. As explained in [Ric63], a topological space M of infinite type that has infinite number of ends can have *non-cylindrical* ends such as in the case of a Riemann surface with infinite type with *non-planar* ends. In this context, the standard setting in the literature using Liouville manifolds with cylindrical ends and cylindrical-at-infinity Lagrangian submanifolds as the objects of the Fukaya category cannot be applied. The standard approach of defining the Fukaya category as the (homotopy) colimit of the Fukaya category of finite type by considering the increasing union

$$M = \bigcup_{N=1}^{\infty} M_N$$

where each M_N is a compact Liouville domain of finite type is not necessarily the only way of defining the Fukaya category of such Liouville manifolds.

We propose to adopt the gradient-sectorial Lagrangian submanifolds with respect to a suitably chosen exhaustion function ψ as the objects of the Fukaya category. In adjunction with this choice, we then consider the class of almost complex structures J for which (ψ, J) forms a pseudoconvex pair as defined in the present paper and [Oh21], and the associated sectorial Hamiltonians H as defined in [Oh21] as the relevant wrapping Hamiltonians. We refer to [Cho] for the study of the Fukaya category for the infinite-type Riemann surfaces in this point of view. It remains to be seen whether this new definition of the Fukaya category is quasi-isomorphic to the colimit definition of the Fukaya category or not, even for the case of Riemann surfaces.

Generalizing this construction to higher dimensional cases and its applications is a subject of future research.

References

- [AM78] R. Abraham and J. Marsden, Foundations of Mechanics, Benjamin/Cummings, Reading, MA, 1978, 2nd Edition.
- [Amo16] Lino Amorim, Tensor product of filtered A_{∞} -algebras, J. Pure Appl. Algebra **220** (2016), no. 12, 3984–4016.
- [Amo17] _____, The Künneth theorem for the Fukaya algebra of a product of Lagrangians, Internat. J. Math. 28 (2017), no. 4, 1750026, 38 pp.
- [Arn88] V. I. Arnol'd, Mathematical Methods of Classical Mechanics, Graduate Texts in Mathematics, vol. 60, Springer-Verlag, New York, NY, 1988, 2nd Edition.
- [AS10] Mohammed Abouzaid and Paul Seidel, An open string analogue of Viterbo functoriality, Geom. Topol. 14 (2010), no. 2, 627–718. MR 2602848 (2011g:53190)
- [CC00] Alberto Candel and Lawrence Conlon, *Foliations I*, Graduate Studies in Mathematics, 23., American Mathematical Socienty, 2000.
- [Che66] Shiing-Shen Chern, The geometry of G-structures, Bulletin of the American Mathematical Society 72 (1966), no. 2, 167–219.
- [Cho] Jaeyoung Choi, Wrapped Fukaya category of infinite-type surfaces, Thesis in POSTECH (in preparation).
- [FOOO20] Kenji Fukaya, Yong-Geun Oh, Hiroshi Ohta, and Kaoru Ono, Kuranishi structures and virtual fundamental chains, Springer Monographs in Mathematics, Springer, Singapore, 2020, 638 pp.
- [Fuk17] Kenji Fukaya, Unobstructed immersed Lagrangian correspondence and filtered A_{∞} -functor, arXiv:1706.02131v1, 2017.
- [Ful93] William Fulton, Introduction to toric varieties, Annals of Mathematics Studies, vol. 131, Princeton University Press, Princeton, NJ, 1993, The William H. Roever Lectures in Geometry.
- [Gao17] Yuan Gao, Functors of wrapped Fukaya categories from Langrangian correspondence, arXiv:1712.00225, 2017.
- [Gir17] E. Giroux, Ideal Liouville domains a cool gadget, arXiv:1708.08855, 2017.
- [Got82] Mark J Gotay, On coisotropic imbeddings of presymplectic manifolds, Proc. Amer. Math. Soc. 84 (1982), no. 1, 111–114.
- [GPS17] Sheel Ganatra, John Pardon, and Vivek Shende, Covariantly functorial wrapped Floer theory on Liouville sectors, arXiv:1706.03152, 2017.
- [GPS18] _____, Sectorial descent for wrapped Fukaya categories, arXiv:1809.03427v2, 2018.
- [Gro17] Yoel Groman, Floer theory on open manifolds, preprint, arXiv:1510.04265v5.
- [Oh21] Yong-Geun Oh, Geometry of Liouville sectors and the maximum principle, preprint, 2021.
- [OP05] Yong-Geun Oh and Jae-Suk Park, Deformations of coisotropic submanifolds and strong homotopy Lie algebroids, Invent. Math. 161 (2005), no. 2, 287–360. MR 2180451 (2006g:53152)
- [OT] Yong-Geun Oh and Hiro Lee Tanaka, Smooth approximation for classifying spaces of diffeomorphism groups, Algebraic & Geometric Topology, (to appear), arxiv:2003.06033.
- [OT20] _____, Holomorphic curves and continuation maps in Liouville bundles, arXiv:2003.04977, 2020.
- [Ric63] Ian Richards, On the classification of noncompact Riemann surfaces, Trans. Amer. Math. Soc. 106 (1963), 259–269.
- [Sei08] Paul Seidel, Fukaya Categories and Picard-Lefschetz Theory, Zurich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich, 2008. MR 2441780 (2009f:53143)
- [Ste83] Shlomo Sternberg, Lectures on Differential Geometry, 2nd ed. ed., Chelsea Publishing Co, New York, 1983.
- [Ste95] J. Arthur Jr. Steen, Lynn Arthur; Seebach, Counterexamples in Topology, Springer-Verlag, Berlin, New York, 1995, Dover reprint of 1978 ed.

[Wei79] A. Weinstein, Lectures on symplectic manifolds, CBMS Regional Conference Series in Mathematics, vol. 29, American Mathematical Society, Providence, R.I., 1979, Corrected reprint, ii + 48 pp.

Center for Geometry and Physics, Institute for Basic Science (IBS), 77 Cheongamro, Nam-gu, Pohang-si, Gyeongsangbuk-do, Korea 790-784 & POSTECH, Gyeongsangbuk-do, Korea

 $Email \ address: \ {\tt yongohl@postech.ac.kr}$