

GREEN-LAZARSFELD INDEX OF SQUARE-FREE MONOMIAL IDEALS AND THEIR POWERS

MOHAMMAD FARROKHI D. G., YASIN SADEGH, AND ALI AKBAR YAZDAN POUR

ABSTRACT. Let \mathbb{K} be a field and I be a square-free monomial ideal in the polynomial ring $\mathbb{K}[x_1, \dots, x_n]$. The Green-Lazarsfeld index, $\text{index}(I)$, counts the number of steps to reach to a syzygy minimally generated by a nonlinear form in a graded minimal free resolution of I . In this paper, we study this invariant for I and its powers from a combinatorial point of view. We characterize all square-free monomial ideals I generated in degree 3 such that $\text{index}(I) > 1$. Utilizing this result, we also characterize all square-free monomial ideals generated in degree 3 such that $\text{index}(I) > 1$ and $\text{index}(I^2) = 1$. In case $n \leq 5$, it is shown that $\text{index}(I^k) > 1$ for all k if I is any square-free monomial ideal with $\text{index}(I) > 1$.

INTRODUCTION

Let \mathbb{K} be a field and $S = \mathbb{K}[x_1, \dots, x_n]$ be the polynomial ring in n variables endowed with the standard grading (i.e. $\deg(x_i) = 1$ for all $1 \leq i \leq n$). Let $I \neq 0$ be a homogeneous ideal in S generated by elements of degree d . A classical problem in algebraic geometry and commutative algebra is to study equations of syzygies of I and determine when these equations are linear forms. The ideal I is said to satisfy the $N_{d,p}$ -property ($p > 0$), if

$$\beta_{i,i+j}(I) = 0, \quad \text{for all } i < p, \text{ and } j > d.$$

The quantity

$$\text{index}(I) = \sup \{p : I \text{ satisfies } N_{d,p}\text{-property}\}$$

is called the *Green-Lazarsfeld index*, or simply *index* of I . The $N_{d,p}$ notation defined in [10] comes essentially from the notation N_p of Green and Lazarsfeld in [15, 14] (see also [8]). This notation indicates when the minimal equations defining the syzygies are in the simplest form (that is of the linear form). The Green-Lazarsfeld index of ideals is very difficult to compute in general. Important conjectures, such as Green's conjecture [9, Chapter 9], predicts the value of this invariant for certain families of varieties. In [4] the authors study the Green-Lazarsfeld index of the Veronese embeddings $\nu_c: \mathbb{P}^{n-1} \rightarrow \mathbb{P}^N$ of degree c of projective spaces and, more generally, of the Veronese embeddings of arbitrary varieties (see also [13, 17, 20, 21, 23] in this matter). In their interesting paper [10], Eisenbud, Green, Hulek, and Popescu provide some nice examples and conjectures about $N_{2,p}$ -property. As a remarkable result, they characterize the property $N_{2,p}$ for monomial ideals in degree 2 [10, Theorem 2.1, Proposition 2.3]. It turns out that the edge ideal $I(G)$ of a graph G satisfies $N_{2,p}$ -property if and only if every induced cycle in G has length $\geq p + 3$. In particular,

$$\text{index}(I(G)) = \inf \{ |C| - 3 : C \text{ is an induced cycle in } \bar{G} \text{ of length } > 3 \}.$$

2010 *Mathematics Subject Classification.* Primary: 13D02, 05E40; Secondary: 13P20.

Key words and phrases. Green-Lazarsfeld index, linear syzygy, powers of ideal, clutter.

Though we have such a good combinatorial formulation for the index of square-free monomials generated in degree 2, the precise value of index of powers of such ideals is still mysterious. In this regard, we have the following result due to Bigdeli, Herzog, and Zaare-Nahandi [3, Theorem 2.1].

Theorem 1. *Let $I = I(G)$ be the edge ideal of a graph G , where G may have some loops. The following conditions are equivalent:*

- (a) G is gap-free, i.e. no induced subgraph of G consists of two disjoint edges;
- (b) $\text{index}(I^k) > 1$, for all k ;
- (c) $\text{index}(I^k) > 1$, for some k .

The above theorem implies in particular that for a monomial ideal generated in degree 2 we have $\text{index}(I) = 1$ if and only if $\text{index}(I^k) = 1$ for all k , while simple examples show that this fails if I is not generated in degree 2 (see Theorem 3.3).

In this paper, we consider the square-free monomial ideals I generated in degree 3 and study the index of I and its powers. Note that such ideals are in one-to-one correspondence with edge ideals of 3-uniform clutters. Recall that a 3-uniform clutter \mathcal{C} on vertex set $V = \{v_1, \dots, v_n\}$ is a collection of 3-subsets of V such that $V = \cup \mathcal{C}$, and the edge ideal $I(\mathcal{C})$ of \mathcal{C} is the ideal generated by all monomials $x_i x_j x_k$ such that $\{v_i, v_j, v_k\} \in \mathcal{C}$. In view of the above theorem, it is reasonable to ask whether for the edge ideal $I = I(\mathcal{C})$ of a 3-uniform clutter \mathcal{C} , the property $\text{index}(I) > 1$ is translated to a combinatorial property of \mathcal{C} . To this end, we interpret the above theorem as follows: If $I = I(G)$ is the edge ideal of a graph G , then the followings are equivalents:

- (i) the complement \bar{G} of G is C_4 -free, where C_4 stands for the cycle of length 4;
- (ii) $\text{index}(I) > 1$;
- (iii) $\text{index}(I^k) > 1$, for some k .

Here we consider C_4 as the minimal triangulation of 1-sphere that is not the triangle. Note that the equivalence of (i) and (ii) has been proved first in [7, Corollary 2.9]. We show in Theorem 2.2 and Proposition 2.4 that if $I = I(\mathcal{C})$ is the edge ideal of a 3-uniform clutter \mathcal{C} , then the followings are equivalent:

- (i) the complement $\bar{\mathcal{C}}$ of \mathcal{C} is \mathcal{C} -free;
- (ii) $\text{index}(I) > 1$;
- (iii) $\text{index}(I^k) > 1$, for some k ,

where \mathcal{C} refers to a set of four 3-uniform clutters whose three of them come from the minimal triangulation of 2-sphere that is not the tetrahedron (see Remark 1 for precise definition of \mathcal{C}). Indeed, as we will see later, the condition that $\bar{\mathcal{C}}$ is \mathcal{C} -free has strong implications for the lcm-lattice. As a byproduct of the above result, it follows that $\text{index}(I(\mathcal{C})) > 1$ if \mathcal{C} is a 3-uniform clutter with $|\mathcal{C}| > \binom{|V(\mathcal{C})|}{3} - 6$. We will see in Proposition 2.3 that indeed $\text{index}(I(\mathcal{C})) = \infty$ if \mathcal{C} is a d -uniform clutter with $|\mathcal{C}| > \binom{|V(\mathcal{C})|}{d} - 2d$.

If I is a (square-free) monomial ideal generated in degree 2 with $\text{index}(I) > 1$, then Theorem 1 implies that $\text{index}(I^k) > 1$ for all k . Examples show that the same statement does not hold for ideals generated in degree greater than 2. However, in Theorem 2.5 we show that for any square-free monomial ideal $I \subseteq \mathbb{K}[x_1, \dots, x_5]$ with $\text{index}(I) > 1$ we have $\text{index}(I^k) > 1$ for all k . Moreover, in Theorem 3.3 we characterize all square-free monomial ideals generated in degree 3 such that $\text{index}(I) > 1$ and $\text{index}(I^2) = 1$. It turns out that if $I = I(\mathcal{C})$ is the edge ideal of a 3-uniform clutter \mathcal{C} and $\text{index}(I) > 1$, then $\text{index}(I^2) > 1$ if and only if \mathcal{C} is \mathcal{D} -free where \mathcal{D} is the family of clutters defined by Figures 1, 2, 3.

1. PRELIMINARIES

Throughout, let \mathbb{K} be a field and $S = \mathbb{K}[x_1, \dots, x_n]$ be the polynomial ring over n variables. In this section, we quickly review basic notions and preliminaries that we will meet in the sequel.

1.1. Multigraded Betti numbers. Let $I \subset S$ be a monomial ideal in the polynomial ring S . Then I is a multigraded S -module and so it admits a minimal multigraded free S -resolution

$$\cdots \rightarrow F_2 \rightarrow F_1 \rightarrow F_0 \rightarrow I \rightarrow 0,$$

where $F_i = \bigoplus_{\mathbf{a} \in \mathbb{Z}^n} S(-\mathbf{a})^{\beta_{i,\mathbf{a}}(I)}$. The numbers $\beta_{i,\mathbf{a}}(I) = \dim_{\mathbb{K}} \operatorname{Tor}_i^S(\mathbb{K}, I)_{\mathbf{a}}$ are called the *multigraded Betti numbers* of I . For $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{Z}^n$, let $|\mathbf{a}| = \sum_i a_i$. Define

$$\beta_{i,j}(I) = \sum_{\substack{\mathbf{a} \in \mathbb{Z}^n \\ |\mathbf{a}|=j}} \beta_{i,\mathbf{a}}(I).$$

The numbers $\beta_{i,j}(I)$ are the *graded Betti numbers* of I with respect to the standard grading on S (i.e. $\deg(x_i) = 1$). A monomial ideal $I \subset S$ is said to have a *d -linear resolution* if $\beta_{i,j}(I) = 0$, for all i, j with $j - i \neq d$.

1.2. Green-Lazarsfeld index. The Green-Lazarsfeld index (or simply index) counts the number of linear steps in the graded minimal free resolution of an ideal. The monomial ideal I is *r -step linear* if I has a linear resolution up to homological degree r ; in other words, if I is generated by homogeneous elements in degree d and $\beta_{i,j}(I) = 0$ for all pairs (i, j) with $0 \leq i \leq r$ and $j - i > d$. The quantity

$$\operatorname{index}(I) = \sup\{r : I \text{ is } r\text{-step linear}\} + 1$$

is called the *index* of I . In particular, I has a linear resolution if and only if $\operatorname{index}(I) = \infty$. A monomial ideal I is called *linearly presented* if $\operatorname{index}(I) > 1$.

1.3. Clutters and their associated ideals. A *clutter* \mathcal{C} on vertex set $V = \{v_1, \dots, v_n\}$ is a collection of subsets of V , called *circuits* of \mathcal{C} , such that $V = \cup \mathcal{C}$ and if F_1 and F_2 are distinct circuits, then $F_1 \not\subseteq F_2$. A clutter \mathcal{C} is called *d -uniform* if every circuit of \mathcal{C} has d vertices. Let $\mathbf{x}_{\emptyset} = 0$ and $\mathbf{x}_F = \prod_{v_i \in F} x_i$ for any non-empty subset F of V . For a non-empty clutter \mathcal{C} we define the ideal $I(\mathcal{C})$ to be

$$I(\mathcal{C}) = (\mathbf{x}_T : T \in \mathcal{C}),$$

and we set $I(\emptyset) = 0$. The ideal $I(\mathcal{C})$ is called the *edge ideal* of \mathcal{C} . If \mathcal{C} is a d -uniform clutter on V , then the *complement* $\bar{\mathcal{C}}$ of \mathcal{C} is defined to be

$$\bar{\mathcal{C}} = \{F \subset V : |F| = d, F \notin \mathcal{C}\}.$$

Let \mathcal{C} be a clutter and let e be a subset of V . By $\mathcal{C} \setminus e$ we mean the clutter $\{F \in \mathcal{C} : e \not\subseteq F\}$.

1.4. Simplicial complexes and Stanley-Reisner ideals. A *simplicial complex* Δ on the vertex set $V = \{v_1, \dots, v_n\}$ is a collection of subsets of V , called *faces*, such that $\{v_i\} \in \Delta$ for all i and, $F \in \Delta$ implies that all subsets of F are also in Δ . Maximal faces of Δ are called *facets* of Δ . A simplicial complex with facets F_1, \dots, F_m is denoted by $\langle F_1, \dots, F_m \rangle$. The *induced* subcomplex $\Delta[W]$ of Δ is the simplicial complex $\Delta[W] = \{F \in \Delta : F \subseteq W\}$. Also, the Stanley-Reisner ideal I_{Δ} is the square-free monomial ideal generated by all \mathbf{x}_F with $F \notin \Delta$.

Let \mathcal{C} be a d -uniform clutter on the vertex set $[n] := \{1, \dots, n\}$. A subset F of $[n]$ is a *clique* in \mathcal{C} if either $|F| < d$ or else all d -subsets of F belong to \mathcal{C} . The *clique complex*

$\Delta(\mathcal{C})$ of \mathcal{C} is the simplicial complex whose faces are cliques of \mathcal{C} . It is easy to see that $I(\mathcal{C}) = I_{\Delta(\bar{\mathcal{C}})}$.

1.5. Gasharov-Peeva-Welker formula. In the following, by a *poset* we mean a partially ordered set. A subset C of a poset P is called a *chain* in P if C is a totally ordered subset of P with respect to the induced order.

Let I be a monomial ideal and $\mathcal{G}(I) = \{u_1, \dots, u_m\}$ be the unique minimal set of monomial generators of I . We denote by $L(I)$ the *lcm-lattice* of I , that is a poset whose elements are labeled by the least common multiples of subsets of monomials in $\mathcal{G}(I)$ ordered by divisibility. For any $u \in L(I)$ we denote by $(1, u)$ the open interval of $L(I)$, which is by definition

$$(1, u) = \{v \in L(I) : 1 < v < u\}.$$

Furthermore, we denote by $\Delta_I(1, u)$ the order complex of the poset $(1, u)$, that is the simplicial complex whose faces are the chains in the poset $(1, u)$.

Theorem 1.1 ([12, Theorem 2.1]). *For all $i \geq 0$ and all $u \in L(I)$,*

$$\beta_{i,u}(I) = \dim_{\mathbb{K}} \tilde{H}_{i-1}(\Delta_I(1, u); \mathbb{K}).$$

2. INDEX OF EQUIGENERATED SQUARE-FREE MONOMIAL IDEALS AND THEIR POWERS

Let \mathcal{C} be a clutter on vertex set V and W be a subset of V . The *induced* subclutter $\mathcal{C}[W]$ of \mathcal{C} is the clutter $\mathcal{C}[W] = \{F \in \mathcal{C} : F \subseteq W\}$. Assume that \mathcal{F} is a family of clutters. We say that \mathcal{C} is \mathcal{F} -free if \mathcal{C} does not contain an induced subclutter isomorphic to any element in \mathcal{F} . Let G be a graph and $I = I(G)$ be the edge ideal of G . It follows from [7, Corollary 2.9] that \bar{G} is C_4 -free if and only if $\text{index}(I) > 1$. We intend to present a similar result for 3-uniform clutters (Theorem 2.2). To do this, we need the following setting: Let I be a monomial ideal whose minimal generators are all of degree d . In [3] the authors define a graph G_I whose vertex set is $\mathcal{G}(I)$ and for which $\{u, v\}$ is an edge of G_I if and only if $\deg(\text{lcm}(u, v)) = d + 1$. For all $u, v \in \mathcal{G}(I)$, let $G_I^{u,v}$ be the induced subgraph of G_I with vertex set

$$\{w \in \mathcal{G}(I) : w \text{ divides } \text{lcm}(u, v)\}.$$

The following result obtained simply from the proofs of [3, Proposition 1.1 and Corollary 1.2] and also Theorem 1.1 plays a crucial role in this paper.


Theorem 2.1. *Let I be a monomial ideal generated in degree d . Then*

- (i) *if $u, v \in \mathcal{G}(I)$ are such that $\deg \text{lcm}(u, v) > d + 1$ and $\Delta_I(1, \text{lcm}(u, v))$ is disconnected, then u, v belong to distinct connected components of $G_I^{u,v}$;*
- (ii) *if $\Delta_I(1, \text{lcm}(u, v))$ is connected for all $u, v \in \mathcal{G}(I)$, then so is $G_I^{u,v}$ for all $u, v \in \mathcal{G}(I)$.*

In particular, $\text{index}(I) > 1$ if and only if for all $u, v \in \mathcal{G}(I)$ there is a path in $G_I^{u,v}$ connecting u and v .

Remark 1. Let \mathcal{B} be the following bi-pyramid, $\mathcal{B}_1 = \mathcal{B} \cup \{125\}$, and $\mathcal{B}_2 = \mathcal{B} \cup \{125, 135\}$. Since $\langle \mathcal{B} \rangle$ is a triangulation of the 2-sphere, $\beta_{1,5}(I(\bar{\mathcal{B}})) = \dim_{\mathbb{K}} \tilde{H}_2(\Delta(\bar{\mathcal{B}}); \mathbb{K}) = 1$. Hence, by [2, Theorem 2.1], we have

$$1 = \beta_{1,5}(I(\bar{\mathcal{B}})) = \beta_{1,5}(I(\bar{\mathcal{B}}_1)) = \beta_{1,5}(I(\bar{\mathcal{B}}_2)).$$



On the other hand, if $\mathcal{B}' = \binom{[6]}{3} \setminus \{123, 456\}$ then one may easily check that $\beta_{1,6}(I(\bar{\mathcal{B}}')) = 1$. Let \mathcal{C} be the family $\{\mathcal{B}, \mathcal{B}_1, \mathcal{B}_2, \mathcal{B}'\}$ of clutters. The following theorem shows that this family of clutters tends to a characterization of all square-free monomial ideals I such that $\text{index}(I) > 1$.

Theorem 2.2. *Let \mathcal{C} be a 3-uniform clutter and let $I = I(\mathcal{C})$ be its edge ideal. Let $\bar{\mathcal{C}}$ be the family of clutters as in the above remark. The following conditions are equivalent:*

- (a) $\bar{\mathcal{C}}$ is \mathcal{C} -free;
- (b) $\text{index}(I) > 1$, that is I is linearly presented;
- (c) $\beta_{1,5}(I) = \beta_{1,6}(I) = 0$.

Proof. $a \Rightarrow b$: Suppose that $\bar{\mathcal{C}}$ is \mathcal{C} -free. Using Theorem 2.1, it is enough to show that for all $u, v \in \mathcal{G}(I)$ there is a path in $G := G_I^{u,v}$ between u and v . Let $u = x_i x_j x_k$ and $v = x_r x_s x_t$.

If $\text{deg lcm}(u, v) = 4$, then $\{u, v\} \in E(G)$ and we have nothing to prove. Suppose that $\text{deg lcm}(u, v) = 5$. Without loss of generality, we may assume that $u = x_1 x_2 x_3$ and $v = x_1 x_4 x_5$. Suppose on the contrary that there is no path in G between u and v . Let $X = \{124, 125, 134, 135\}$. If \mathbf{x}_F is a vertex of G for some $F \in X$, then $u - \mathbf{x}_F - v$ is a path in G between u and v , a contradiction. Thus $X \subseteq \bar{\mathcal{C}}$. Now put

$$\begin{aligned} A_1 &= \{x_2 x_3 x_4, x_2 x_4 x_5\}, & A_2 &= \{x_2 x_3 x_4, x_3 x_4 x_5\}, \\ A_3 &= \{x_2 x_3 x_5, x_2 x_4 x_5\}, & A_4 &= \{x_2 x_3 x_5, x_3 x_4 x_5\}. \end{aligned}$$

Observe that $A_i \not\subseteq V(G)$, for $i = 1, \dots, 4$ for otherwise $A_i \cup \{u, v\}$ yields a path between u and v in G . Hence $\bar{\mathcal{C}} \cap A_i \neq \emptyset$, for $i = 1, \dots, 4$. A simple verification shows that the clutter induced by X and those circuits $F \in \bar{\mathcal{C}}$ with $\mathbf{x}_F \in A_1 \cup \dots \cup A_4$ contains one of the elements of \mathcal{C} contradicting the \mathcal{C} -freeness of $\bar{\mathcal{C}}$.

Next assume that $\text{deg lcm}(u, v) = 6$. Without loss of generality, assume that $u = x_1 x_2 x_3$ and $v = x_4 x_5 x_6$. We claim that there is a path in G connecting u to v (of length 3). Suppose on the contrary that there is no path between u and v in G . Let $B_{ijab} = \{x_i x_j x_a, x_i x_a x_b\}$ for all $1 \leq i < j \leq 3$ and $4 \leq a < b \leq 6$. Then $B_{ijab} \not\subseteq V(G)$ because otherwise $B_{ijab} \cup \{u, v\}$ is a path of length 3 in G between u and v , contradicting the assumption. The group $H = \langle (1\ 2\ 3), (1\ 2), (4\ 5\ 6), (4\ 5) \rangle$ acts naturally on the set

$$\Omega := \{x_i x_j x_a : 1 \leq i < j \leq 3, 4 \leq a \leq 6\}$$

by $(x_i x_j x_a)^\pi = x_{i^\pi} x_{j^\pi} x_{a^\pi}$ for all $\pi \in H$ and $x_i x_j x_a \in \Omega$. Suppose $X := \Omega \cap V(G)$ is a k -subset of Ω , for some $0 \leq k \leq 9$. Then X belongs to an orbit of H on $\binom{\Omega}{k}$, the set of all k -subsets of Ω . Since all elements of an orbit share the same properties, we can restrict ourselves to a fixed representative of any orbit. Observe that G has 1, 1, 3, 6, 7, 7, 6, 3, 1, 1 orbits in action on $\binom{\Omega}{k}$, for $k = 0, \dots, 9$, respectively, so that we have 36 cases to consider. For any such case, we have $iab \in \bar{\mathcal{C}}$ (resp. $ija \in \bar{\mathcal{C}}$) whenever

$x_i x_j x_a \in X$ (resp. $x_i x_j x_a \in \Omega \setminus X$) because $B_{ijab} \not\subseteq V(G)$. What remains is to consider all possibilities $iab \in \mathcal{C}$ or $iab \in \bar{\mathcal{C}}$ for all iab satisfying $x_i x_j x_a \in X$ whose the number of them is bounded above by $2^5 = 32$ except for the case where $X = \emptyset$. A simple computer program shows that we have totally 105 cases and, in any case, $\bar{\mathcal{C}}$ has an element of \mathcal{C} as an induced subclutter. This contradicts the fact that $\bar{\mathcal{C}}$ is \mathcal{C} -free.

(b) \Rightarrow (c): This implication is clear by the definition of the index of an ideal.

(c) \Rightarrow (a): Suppose on the contrary that $\bar{\mathcal{C}}$ is not \mathcal{C} -free. So $\bar{\mathcal{C}}$ contains an induced subclutter \mathcal{D} isomorphic to one of the elements of \mathcal{C} . If $\mathcal{D} \not\cong \binom{[6]}{3} \setminus \{123, 456\}$, then in view of Remark 1, we obtain

$$\beta_{1,5}(I) \geq \beta_{1,5}(I(\bar{\mathcal{D}})) = 1,$$

which contradicts to our assumption. Thus $\mathcal{D} \cong \binom{[6]}{3} \setminus \{123, 456\}$. Without loss of generality, assume that $\mathcal{D} = \binom{[6]}{3} \setminus \{123, 456\}$. Then by setting $\alpha = x_1 \cdots x_6$, the interval $(1, \alpha)$ consists of only two points $x_1 x_2 x_3$ and $x_4 x_5 x_6$. Thus $\Delta(1, \alpha)$ is disconnected. It follows that $\beta_{1,6}(I) \geq \beta_{1,\alpha}(I) = \dim \tilde{H}_0(\Delta(1, \alpha); \mathbb{K}) = 1$, which is a contradiction. \square

Theorem 2.2 implies that $\text{index}(I(\mathcal{C})) > 1$ for any 3-uniform clutter \mathcal{C} on a vertex set $[n]$ with $|\mathcal{C}| > \binom{n}{3} - 6$. In the following, we show that an stronger result holds under the same assumptions. This result is not restricted only to 3-uniform clutters.

Proposition 2.3. *Let $d > 1$ be a positive integer and κ_d be the smallest size of a d -uniform clutter \mathcal{C} such that $I(\bar{\mathcal{C}})$ does not have linear resolution. Then $\kappa_d = 2d$. In particular, if \mathcal{C} is a d -uniform clutter with $|\mathcal{C}| > \binom{|V(\mathcal{C})|}{d} - 2d$, then $I(\mathcal{C})$ has a linear resolution.*

Proof. First we show that $\kappa_d \leq 2d$. Let

$$\mathcal{C}_d = \left(\binom{\{1, \dots, d+1\}}{d} \cup \binom{\{2, \dots, d+2\}}{d} \right) \setminus \{2, \dots, d+1\}.$$

The simplicial complex $\langle \mathcal{C}_d \rangle$ is a triangulation of the $(d-1)$ -sphere \mathbb{S}^{d-1} so that

$$\tilde{H}_{d-1}(\Delta(\mathcal{C}_d); \mathbb{K}) = \tilde{H}_{d-1}(\langle \mathcal{C}_d \rangle; \mathbb{K}) = H_{d-1}(\mathbb{S}^{d-1}; \mathbb{K}) \neq 0.$$

Thus $\beta_{1,d+2}(I(\bar{\mathcal{C}}_d)) \neq 0$ by Hochster's formula [16, Theorem 5.1], and hence $I(\bar{\mathcal{C}}_d)$ does not have linear resolution showing that $\kappa_d \leq 2d$.

To complete the proof, let \mathcal{C} be a d -uniform clutter such that $|\mathcal{C}| < 2d$. Let $\Delta = \Delta(\mathcal{C})$ be the clique complex of \mathcal{C} . It is obvious that $d-1 \leq \dim \Delta \leq d$. Suppose $\partial_{d-1}(\sum_{i=1}^m c_i F_i) = 0$ for some circuits F_1, \dots, F_m of \mathcal{C} and put $\mathcal{C}' = \{F_1, \dots, F_m\}$. Clearly, every $(d-1)$ -subset of F_i is contained in F_j for some $j \neq i$. Let $X = \cup \mathcal{C}'$. We show that any t -subset of X is contained in some circuit of \mathcal{C}' provided that $t \leq d$. Clearly, the result holds for $t = 1$. Assume the claim is true for some $t < d$ but not for t . Let X' be a t -subset of X . By assumptions, every $(t-1)$ -subset of X' is contained in at least $d-t+2$ circuits $F_{k_1}^{X'}, \dots, F_{k_{d-t+2}}^{X'}$ of \mathcal{C}' for X' is contained in some circuit F and F has $d-t+2$ subsets of size $d-1$ containing X' . Since $F_{k_i}^{X'}$ and $F_{k_j}^{X''}$ are distinct for $X' \neq X''$ or $i \neq j$, it follows that

$$t(d-t+2) = \# \left\{ F_{k_i, X'} : X' \in \binom{X}{t-1}, 1 \leq i \leq d-t+2 \right\} \leq |\mathcal{C}'| < 2d.$$

This yields $d < t$, which is a contradiction. It follows that $\mathcal{C}' = \binom{X}{d}$, hence $X \in \Delta$ and $|X| = d+1$. This shows that $\sum_{i=1}^m c_i F_i$ is a scalar multiple of $\partial_d(X)$ and consequently $\tilde{H}_{d-1}(\Delta[W], \mathbb{K}) = 0$ for all $W \subseteq V(\mathcal{C})$.

If $\dim \Delta = d - 1$, then $\tilde{H}_i(\Delta[W]; \mathbb{K}) = 0$ by the argument above for all $W \subseteq V(\mathcal{C})$ and $i \geq d - 1$. Hence $I(\bar{\mathcal{C}}) = I_{\Delta(\mathcal{C})}$ has a linear resolution (see e.g. [24, Proposition 3.1]). So, assume that $\dim \Delta = d$. In this case, \mathcal{C} has a unique clique F of size $d + 1$. Indeed, if F and F' are two distinct cliques in \mathcal{C} of size $d + 1$, then $|F \cap F'| \leq d$ so that $F \cup F'$ contains at least $2d + 1$ circuits of \mathcal{C} contradicting the fact that $|\mathcal{C}| < 2d$. Since $\dim \Delta = d$, it follows that $\tilde{H}_d(\Delta; \mathbb{K}) = \ker \partial_d = 0$, where ∂_d is the chain map from $\mathbb{K}F$ into $\bigoplus_{e \in \mathcal{C}} \mathbb{K}e$. Consequently, $\tilde{H}_i(\Delta[W]; \mathbb{K}) = 0$ for all $W \subseteq V(\mathcal{C})$ and $i \geq d - 1$. Thus $I(\bar{\mathcal{C}})$ has a linear resolution in this case too. \square

If G is a graph and $\text{index}(I(G)^k) > 1$ for some $k \geq 1$, then \bar{G} is C_4 -free (Theorem 1). Here we prove an analogue result for 3-uniform clutters.

Proposition 2.4. *Let \mathcal{C} be a 3-uniform clutter and $I = I(\mathcal{C})$ be the corresponding edge ideal. The following conditions are equivalent:*

- (a) $\bar{\mathcal{C}}$ is \mathcal{C} -free;
- (b) $\text{index}(I^k) > 1$, for some $k \geq 1$.

Proof. (a) \Rightarrow (b). If $\bar{\mathcal{C}}$ is \mathcal{C} -free, then by Theorem 2.2, $\text{index}(I) > 1$.

(b) \Rightarrow (a). If $k = 1$ the result holds in view of Theorem 2.2, so assume that $k > 1$ and $\text{index}(I^k) > 1$. Suppose on the contrary that $\bar{\mathcal{C}}$ has an induced subclutter isomorphic to an element of \mathcal{C} . First assume that $\bar{\mathcal{C}}$ contains the bi-pyramid

$$\mathcal{B} = \{123, 124, 134, 235, 345, 245\}$$

as induced subclutter. It follows that $x_1x_2x_5, x_1x_3x_5, x_1x_4x_5, x_2x_3x_4 \in I$. Note that the monomials $(x_2x_3x_4)^k, (x_1x_2x_5)(x_2x_3x_4)^{k-1}, (x_1x_3x_5)(x_2x_3x_4)^{k-1}$, and $(x_1x_4x_5)(x_2x_3x_4)^{k-1}$ are atoms in the lcm-lattice of I^k . Clearly $\alpha := x_1x_2^kx_3^kx_4^kx_5$ covers these four atoms and the open interval $(1, \alpha)$ consists of the four atoms. Hence $\Delta(1, \alpha)$ is disconnected. Consequently,

$$\beta_{1,3k+2}(I^k) \geq \beta_{1,\alpha}(I^k) = \dim_{\mathbb{K}} \left(\tilde{H}_0(\Delta(1, \alpha); \mathbb{K}) \right) \neq 0.$$

This implies that $\text{index}(I^k) = 1$, a contradiction. If $\mathcal{B}_1 = \mathcal{B} \cup \{125\}$ or $\mathcal{B}_2 = \mathcal{B} \cup \{125, 135\}$ is an induced subclutter of $\bar{\mathcal{C}}$, then the same discussion as above leads us to a contradiction. Finally, suppose that $\bar{\mathcal{C}}$ contains $\binom{[6]}{3} \setminus \{123, 456\}$ as induced subclutter. In this case, put $\alpha = (x_1x_2x_3)^k(x_4x_5x_6)$. Then the open interval $(1, \alpha)$ consists of two atoms $(x_1x_2x_3)^{k-1}(x_4x_5x_6)$ and $(x_1x_2x_3)^k$. Thus $\Delta(1, \alpha)$ is disconnected and

$$\beta_{1,3k+3}(I^k) \geq \beta_{1,\alpha}(I^k) = \dim_{\mathbb{K}} \left(\tilde{H}_0(\Delta(1, \alpha); \mathbb{K}) \right) \neq 0.$$

This implies that $\text{index}(I^k) = 1$, which is again a contradiction. \square

If $I = I(G)$ is the edge ideal of a graph G and $\text{index}(I) > 1$, then Theorem 1 implies that $\text{index}(I^k) > 1$ for all $k \geq 1$. Examples show that this statement is not true if I is generated by (square-free) monomials generated in degree greater than 2 (see Theorem 3.3). The next theorem shows that this result still holds for square-free monomial ideals in a polynomial ring with at most five variables. Theorem 3.3 shows that we can not extend this result for polynomial rings in more than five variables.

Theorem 2.5. *Let $I \subseteq \mathbb{K}[x_1, \dots, x_n]$ be a square-free monomial ideal with $\text{index}(I) > 1$, where $n \leq 5$. Then $\text{index}(I^k) > 1$ for all $k \geq 1$.*

Proof. Suppose $I \subseteq \mathbb{K}[x_1, \dots, x_n]$ is a square-free monomial ideal generated in degree d and that $x_1 \cdots x_n \mid \text{lcm}(\mathcal{G}(I))$. If $d = 1$, then there is nothing to prove. The case $d = 2$ follows from Theorem 1. If $d = n - 1$ or $d = n$, then I is a polymatroidal ideal and hence all powers of I have linear resolutions (see [6, Theorems 5.2 and 5.3]). So, the only case we need to consider is $n = 5$ and $d = 3$. Suppose on the contrary that $\text{index}(I^m) = 1$ for some m , and that m is minimum with this property. Then $\text{index}(I^i) > 1$, for all $i = 1, \dots, m - 1$. By Theorem 2.1, there exist $u, v \in \mathcal{G}(I^m)$ such that u and v belong to different connected components of $G_{I^m}^{u,v}$. Let $u = u_1 \cdots u_m$ and $v = v_1 \cdots v_m$ with $u_i, v_i \in \mathcal{G}(I)$, for $i = 1, \dots, m$. Let $U = \{u_1, \dots, u_m\}$ and $V = \{v_1, \dots, v_m\}$. It turns out that $U \cap V = \emptyset$ for $G_{I^{m-1}}^{u/u_i, v/v_j}$ is connected when $u_i = v_j$, for some $1 \leq i, j \leq m$.

For a monomial u let $\text{supp}(u)$ be the set $\{i: x_i \mid u\}$. First assume that $\text{supp}(u_i) \cap \text{supp}(v_j)$ is a singleton for all $1 \leq i, j \leq m$. Without loss of generality, suppose that $u_1 = x_1x_2x_3$ and $v_1 = x_1x_4x_5$. Clearly, $U \subseteq \{x_1x_2x_3, x_2x_3x_4, x_2x_3x_5\}$ and $V \subseteq \{x_1x_4x_5, x_2x_4x_5, x_3x_4x_5\}$. Since $\text{index}((u_1, X)) = 1$ for any $\emptyset \neq X \subseteq V$ (see Theorem 2.2), we conclude that $\mathcal{G}(I) \setminus (\{u_1\} \cup V)$ (and similarly $\mathcal{G}(I) \setminus (\{v_1\} \cup U)$) is non-empty. Let $x_a x_b x_c \in \mathcal{G}(I) \setminus (\{u_1\} \cup V)$. If $x_a x_b x_c \notin U$, then we may assume w.l.o.g. that $x_a x_b x_c = x_1x_2x_4$. Let $w_1 = (u/x_3)x_4$ and $w_2 = (v/x_5)x_2$. Then $w_1, w_2 \in \mathcal{G}(I^m)$ and divide $\text{lcm}(u, v)$. Since $w_1 = x_1x_2x_4(u/u_1)$ and $w_2 = x_1x_2x_4(v/v_1)$ share the same element $x_1x_2x_4 \in \mathcal{G}(I)$ and the graph $G_{I^{m-1}}^{w_1/x_1x_2x_4, w_2/x_1x_2x_4}$ is connected, there exists a path between w_1 and w_2 in $G_{I^m}^{u,v}$. Since $u \sim w_1$ and $w_2 \sim v$, it follows that u and v are connected via a path in $G_{I^m}^{u,v}$, which is a contradiction. Thus $\mathcal{G}(I) \subseteq U \cup V$. Clearly, $|\mathcal{G}(I) \cap U|, |\mathcal{G}(I) \cap V| \geq 2$. W.l.o.g. we can assume that $x_2x_3x_4 \in \mathcal{G}(I) \cap U$ and $x_2x_4x_5 \in \mathcal{G}(I) \cap V$. Let $w_1 = (u/x_1)x_4$, $w_2 = (u/x_1x_3)x_4x_5$, and $w_3 = (v/x_1)x_2$. Clearly, $w_1, w_2, w_3 \in \mathcal{G}(I^m)$ and divide $\text{lcm}(u, v)$. Notice that $w_2 = x_2x_4x_5(u/u_1)$ and $w_3 = x_2x_4x_5(v/v_1)$ share the same element $x_2x_4x_5$ of $\mathcal{G}(I)$. Since $u \sim w_1 \sim w_2$, $w_3 \sim v$, and $w_2/x_2x_4x_5$ and $w_3/x_2x_4x_5$ are connected via a path in $G_{I^{m-1}}^{w_2/x_2x_4x_5, w_3/x_2x_4x_5}$, it follows that u and v are connected via a path in $G_{I^m}^{u,v}$. This contradiction shows that there exist i, j such that u_i and v_j have two variables in common, say $i = j = 1$, $u_1 = x_1x_2x_3$, and $v_1 = x_1x_2x_4$. Let $u = x_1^{\alpha_1} \cdots x_5^{\alpha_5}$ and $v = x_1^{\beta_1} \cdots x_5^{\beta_5}$. We claim

(*) If $u_s = x_i x_j x_k$ and $v_t = x_i x_j x_l$, then $\alpha_k \leq \beta_k$ and $\alpha_l \geq \beta_l$.

If not, we may assume $\alpha_k > \beta_k$. Then $v \sim (v/v_t)u_s$, $(v/v_t)u_s$ divides $\text{lcm}(u, v)$, and u/u_s and v/v_t are connected via a path in $G_{I^{m-1}}^{u/u_s, v/v_t}$. Hence u and v are connected via a path in $G_{I^m}^{u,v}$, which contradicts our assumption.

Since $u_1 = x_1x_2x_3$ and $v_1 = x_1x_2x_4$ we get $\alpha_4 \geq \beta_4 > 0$ and $0 < \alpha_3 \leq \beta_3$. This shows that $x_4 \mid u$ and $x_3 \mid v$. By symmetry, we have three cases to consider:

Case 1. $x_4x_5 \nmid u_i$ and $x_3x_5 \nmid v_i$, for $i = 1, \dots, m$. Since $x_4 \mid u$ and $x_3 \mid v$, we can assume that $u_2 = x_1x_3x_4$ and $v_2 = x_2x_3x_4$. As $\alpha_3 \leq \beta_3 < m$ and $\beta_4 \leq \alpha_4 < m$, there exist $3 \leq i, j \leq m$ such that $x_3 \nmid u_i$ and $x_4 \nmid v_j$. From the inequalities $u_i \neq x_1x_2x_4$ and $v_j \neq x_1x_2x_3$, we observe that $x_5 \mid u_i, v_j$. By assumption, $x_4 \nmid u_i$ and $x_3 \nmid v_j$. Thus $u_i = v_j = x_1x_2x_5$, which is a contradiction.

Case 2. $x_4x_5 \mid u_i$ and $x_3x_5 \mid v_j$ for some $1 \leq i, j \leq m$. W.l.o.g. assume that $i = j = 2$. By symmetry of 1 and 2 in u_1 and v_1 , either $u_2 = x_1x_4x_5$ or $u_2 = x_3x_4x_5$. First assume that $u_2 = x_1x_4x_5$. Then $\alpha_2 \geq \beta_2$ and $\alpha_5 \leq \beta_5$ by (*). We have three cases for v_2 . If $v_2 = x_1x_3x_5$, then we get $\alpha_i = \beta_i$ for all $1 \leq i \leq 5$ by (*) and hence $u = v$, a contradiction. If $v_2 = x_2x_3x_5$, then $\alpha_5 = \beta_5$ by (*). Since $\alpha_1 \leq \beta_1 < m$, $u_i \mid x_2x_3x_4x_5$ for some $i > 2$. Analogously, $v_j \mid x_1x_3x_4x_5$ for some $j > 2$ as $\beta_2 \leq \alpha_2 < m$. As $U \cap V = \emptyset$, we deduce that

$x_4 \mid u_i$ and $x_3 \mid v_j$, from which it follows that $\alpha_4 = \beta_4$ and $\alpha_3 = \beta_3$ by (*), respectively. If $x_2 \mid u_i$ (resp. $x_1 \mid v_1$), then (*) yields $\alpha_1 = \beta_1$ (resp. $\alpha_2 = \beta_2$) and hence $\alpha_2 = \beta_2$ (resp. $\alpha_1 = \beta_1$) as $\deg(u) = \deg(v)$. Thus $u = v$, which is a contradiction. Hence $x_2 \nmid u_i$ and $x_1 \nmid v_j$, which implies that $u_i = v_j = x_3x_4x_5$ contradicting the fact that $U \cap V = \emptyset$. The case $v_3 = x_3x_4x_5$ is ruled out with the same discussion as in the case $v_2 = x_2x_3x_5$. Now, assume $u_2 = x_3x_4x_5$. By symmetry, either $v_2 = x_1x_3x_5$ or $v_2 = x_3x_4x_5$. The case $v_2 = x_1x_3x_5$ leads us to a contradiction as in the case $u_2 = x_1x_4x_5$. Hence $v_2 = x_3x_4x_5$, which is a contradiction as $U \cap V = \emptyset$.

Case 3. $x_4x_5 \mid u_i$ for some $2 \leq i \leq m$ but $x_3x_5 \nmid v_j$, for all $2 \leq j \leq m$. W.l.o.g. assume that $i = 2$. By symmetry of x_1 and x_2 in u_1 and v_1 , either $u_2 = x_1x_4x_5$ or $u_2 = x_3x_4x_5$. Since $\beta_3 \geq \alpha_3 > 0$, $x_3 \mid v_j$ for some $2 \leq j \leq m$, say $x_3 \mid v_2$. As $x_5 \nmid v_2$ and $v_2 \neq x_1x_2x_3$, it follows that $x_4 \mid v_2$. Thus $v_2 = x_1x_3x_4$ or $v_2 = x_2x_3x_4$. First assume that $u_2 = x_1x_4x_5$. Then $\alpha_2 \geq \beta_2$ and $\alpha_5 \leq \beta_5$ by (*). From $\beta_5 \geq \alpha_5 > 0$, it follows that $x_5 \mid v_j$ for some $3 \leq j \leq m$, say $x_5 \mid v_3$. As $x_3 \nmid v_3$ and $v_3 \neq x_1x_4x_5$, we must have $x_2 \mid v_3$. Hence $v_3 = x_1x_2x_5$ or $v_3 = x_2x_4x_5$. If $v_2 = x_1x_3x_4$, then by swapping u_1 with u_2 , and v_2 with v_3 , we are at the position of Case 2, which leads us to a contradiction. Hence $v_2 = x_2x_3x_4$ and consequently $\alpha_1 \leq \beta_1$ by (*). If $v_3 = x_1x_2x_5$, then by replacing v_1 and v_2 by v_2 and v_3 , respectively, we get a contradiction as we are at the position of Case 2 once again. Thus $v_3 = x_2x_4x_5$. From the inequalities $\alpha_1 \leq \beta_1 < m$, we conclude that $x_1 \nmid u_i$ for some $3 \leq i \leq m$, say $x_1 \nmid u_3$. Since $U \cap V = \emptyset$, it follows that $x_3x_5 \mid u_3$. Now by replacing u_1, u_2, v_1, v_2 by v_1, v_3, u_1, u_3 , we are at the position of Case 2 leading us to a contradiction. This shows that $u_2 \neq x_1x_4x_5$, that is we must have $u_2 = x_3x_4x_5$. Then by swapping u_1 with u_2 , and v_1 with v_2 , we arrive at the case where $u_2 = x_1x_4x_5$ after suitable relabeling of the variables, which ruled out before. The proof is complete. \square

Remark 2. Theorem 2.5 is not valid for $n > 5$, or monomial ideals with $n \leq 5$ that are not square-free. In Theorem 3.3 we present (all possible) square-free monomial ideals $I \subseteq \mathbb{K}[x_1, \dots, x_6]$ with $\text{index}(I) > 1$ and $\text{index}(I^2) = 1$. As an instance of a monomial ideal I satisfying $\text{index}(I) > 1$ and $\text{index}(I^2) = 1$ in $n \leq 5$ variables, we can refer to Conca's example $I = (x_1^2x_2, x_1^2x_3, x_1x_3^2, x_2x_3^2, x_1x_3x_4) \subseteq \mathbb{K}[x_1, x_2, x_3, x_4]$ (see [5, Example 2.5]).

3. INDEX OF THE SQUARE OF LINEARLY PRESENTED CUBIC SQUARE-FREE MONOMIAL IDEALS

In the rest of the paper, we characterize all cubic square-free monomial ideals I such that $\text{index}(I) > 1$ and $\text{index}(I^2) = 1$. To this end, we need some preparations we mention here.

Lemma 3.1. *Let I be a monomial ideal, $k \geq 1$, and $u = u_1 \cdots u_k$ and $v = v_1 \cdots v_k$ be monomials in $\mathcal{G}(I^k)$, where $u_1, \dots, u_k, v_1, \dots, v_k \in \mathcal{G}(I)$. Let $A, B \subseteq [k]$ be non-empty sets such that $|A| + |B| = k$ and $\prod_{i \in A} u_i \prod_{j \in B} v_j$ divides $\text{lcm}(u, v)$. If $\text{index}(I^{|A|}) > 1$ and $\text{index}(I^{|B|}) > 1$, then u and v are connected via a path in $G_{I^k}^{u, v}$.*

Proof. Without loss of generality, we may assume that $A = \{1, \dots, r\}$ and $B = \{r+1, \dots, k\}$. Let $u' := u_1 \cdots u_r$, $u'' := u_{r+1} \cdots u_k$, $v' := v_1 \cdots v_r$, and $v'' := v_{r+1} \cdots v_k$. Suppose $u' = \alpha_0, \alpha_1, \dots, \alpha_t = v'$ is a path in $G_{I^{|A|}}^{u', v'}$ and $u'' = \beta_0, \beta_1, \dots, \beta_s = v''$ is a path in $G_{I^{|B|}}^{u'', v''}$. Then

$$u = u' \beta_0, u' \beta_1, \dots, u' \beta_s = \alpha_0 v'', \alpha_1 v'', \dots, \alpha_t v'' = v$$

is a path in $G_{I^k}^{u,v}$, as required. \square

Corollary 3.2. *Let I be a monomial ideal and $k \geq 1$ be such that $\text{index}(I^i) > 1$ for $i = 1, \dots, k-1$. Let $u = u_1 \cdots u_k$ and $v = v_1 \cdots v_k$ be monomials in $\mathcal{G}(I^k)$, where $u_1, \dots, u_k, v_1, \dots, v_k \in \mathcal{G}(I)$. If u, v belong to distinct connected components of $G_{I^k}^{u,v}$, then $\prod_{i \in A} u_i \prod_{j \in B} v_j$ does not divide $\text{lcm}(u, v)$ for any two non-empty subsets A and B of $[k]$ with $|A| + |B| = k$.*

Let \mathcal{D}_i^n be the 3-clutter associated to the i -th vertex in Figure 1, 3, or 2 when $n = 6$, $n = 7$, or $n = 8$, respectively, where

$$\begin{aligned} \mathcal{D}_1^6 &= \{123, 246, 145, 356, 134, 136, 146, 346\}, \\ \mathcal{D}_6^7 &= \{123, 124, 127, 145, 147, 247, 267, 347\}, \\ \mathcal{D}_{48}^7 &= \{123, 124, 136, 145, 146, 147, 167, 246, 267, 346, 347\}, \\ \mathcal{D}_1^8 &= \{123, 124, 125, 145, 147, 246, 248, 258, 456\}, \end{aligned}$$

and an arrow with label abc from i -th vertex to j -th vertex means $\mathcal{D}_j^n = \mathcal{D}_i^n \cup \{abc\}$. Notice that all parallel edges in Figures 1, 2, 3 have the same labels and that all clutters \mathcal{D}_i^n are uniquely determined by diagram chasing. Let \mathcal{D} denote the set of all \mathcal{D}_i^n for $n = 6, 7, 8$. The ideal associated to the clutter \mathcal{D}_1^6 , introduced by Sturmfels [22] in 2000, is the first example of an ideal with linear resolution (quotients) whose square does not have linear resolution.




FIGURE 1. $n = 6$




FIGURE 2. $n = 8$




FIGURE 3. $n = 7$

Let I be a monomial ideal in a polynomial ring generated in degree d . It follows from [24, Theorem 2.1] that $\text{index}(I) > 1$ if and only if $\beta_{1,d+2}(I) = \dots = \beta_{1,2d}(I) = 0$. In particular, if I is generated in degree 3, then $\text{index}(I^2) > 1$ if and only if $\beta_{1,8}(I^2) = \dots = \beta_{1,12}(I^2) = 0$. The following theorem shows surprisingly that if $I = I(\mathcal{C})$ is the edge ideal of a 3-uniform clutter \mathcal{C} with $\text{index}(I) > 1$, then $\text{index}(I^2) > 1$ if and only if $\beta_{1,8}(I^2) = 0$.

Theorem 3.3. *Let \mathcal{C} be a 3-uniform clutter and let $I = I(\mathcal{C})$ be such that $\text{index}(I) > 1$. The following conditions are equivalent:*

- (a) \mathcal{C} is \mathcal{D} -free;
- (b) $\text{index}(I^2) > 1$;
- (c) $\beta_{1,8}(I^2) = 0$.

Proof. Let \mathcal{C}' be a 3-uniform clutter on $[n]$ and $J = I(\mathcal{C}')$ be such that $\text{index}(J) > 1$. Let $u, v \in \mathcal{G}(J^2)$ be monomials such that $\text{supp}(uv) = [n]$. We show that \mathcal{C}' is isomorphic to an element of \mathcal{D} if and only if u, v belong to distinct connected components of $G := G_{J^2}^{u,v}$. Suppose u and v belong to distinct connected components of G . Let $u = u_1u_2$ and $v = v_1v_2$ with $u_1, u_2, v_1, v_2 \in \mathcal{G}(J)$. Corollary 3.2 implies that $u_iv_j \nmid \text{lcm}(u, v)$ for $i, j \in \{1, 2\}$. As a result, $\text{gcd}(u_i, v_j) \neq 1$ for $i, j \in \{1, 2\}$ from which it follows that $n \leq 9$. Notice that $\{u_1, u_2\} \cap \{v_1, v_2\} = \emptyset$ otherwise $u_i = v_j$ for some i, j , say $u_1 = v_1$ and consequently $u_1u_2 = u_2v_1 \mid \text{lcm}(u, v)$, a contradiction. If $n = 9$, then $u_1 = x_1x_2x_3$, $u_2 = x_1x_4x_5$, $v_1 = x_1x_6x_7$, and $v_2 = x_1x_8x_9$ after a relabeling of the variables. Then $u_1v_1 \mid \text{lcm}(u, v)$,

which is a contradiction. On the other hand, Theorem 2.5 in conjunction with Theorem 2.1 reveal that $n \geq 6$. Thus $6 \leq n \leq 8$.

By invoking Corollary 3.2 and a computer search presented by Algorithm 1, we arrive at the following cases after a suitable relabeling of variables:

$$\begin{aligned} u_1 = x_1x_2x_3, u_2 = x_2x_4x_6, v_1 = x_1x_4x_5, v_2 = x_3x_5x_6, & \quad \text{if } n = 6, \\ u_1 = x_1x_2x_3, u_2 = x_3x_4x_7, v_1 = x_1x_4x_5, v_2 = x_2x_6x_7, & \quad \text{if } n = 7, \\ u_1 = x_1x_2x_3, u_2 = x_4x_5x_6, v_1 = x_1x_4x_7, v_2 = x_2x_5x_8, & \quad \text{if } n = 8. \end{aligned}$$

For a given fixed $n \in \{6, 7, 8\}$, let $\mathcal{C}_1^n := \{F: \mathbf{x}_F \in \{u_1, u_2, v_1, v_2\}\}$, and define $\mathcal{C}_2^n, \mathcal{C}_3^n$, and \mathcal{C}_4^n as follows:

- \mathcal{C}_2^n is the set of all triples F such that for $w_1 := \mathbf{x}_F$ there exists $w_2 \in \{u_1, u_2\}$ satisfying $w_1w_2 \in V(G)$, $u \sim w_1w_2$, and $v_iw_j \mid \text{lcm}(v, w_1w_2)$ as well as all triples F such that for $w_1 := \mathbf{x}_F$ there exists $w_2 \in \{v_1, v_2\}$ satisfying $w_1w_2 \in V(G)$, $v \sim w_1w_2$, and $u_iw_j \mid \text{lcm}(u, w_1w_2)$.
- $\mathcal{C}_3^n = A^n \cup B^n \cup C^n$, where A^n, B^n , and C^n are as follows: A^n is the set of all 2-sets $\{F_1, F_2\}$ such that $\mathbf{x}_{F_1}\mathbf{x}_{F_2} \in V(G)$ and $u \sim \mathbf{x}_{F_1}\mathbf{x}_{F_2} \sim v$; B^n is the set of all 2-sets $\{F_1, F_2\}$ such that $\mathbf{x}_{F_1}w_1, \mathbf{x}_{F_2}w_2 \in V(G)$ and $u \sim \mathbf{x}_{F_1}w_1 \sim \mathbf{x}_{F_2}w_2 \sim v$ for some $w_1, w_2 \in \{u_1, u_2, v_1, v_2\}$; and C^n is the set of all 2-sets $\{F_1, F_2\}$ such that either $u = \mathbf{x}_{F_1}\mathbf{x}_{F_2}$ and $\mathbf{x}_{F_i}v_j \mid \text{lcm}(u, v)$, or $v = \mathbf{x}_{F_1}\mathbf{x}_{F_2}$ and $\mathbf{x}_{F_i}u_j \mid \text{lcm}(u, v)$, for some i, j .
- $\mathcal{C}_4^n := \binom{[n]}{3} \setminus (\mathcal{C}_1^n \cup \mathcal{C}_2^n \cup \mathcal{C}_3^n)$.

Observe that $\mathcal{C}_2^n \cap \mathcal{C} = \emptyset$ and $\{F_1, F_2\} \not\subseteq \mathcal{C}$ for all $\{F_1, F_2\} \in \mathcal{C}_3^n$. Let G_n be the graph obtained from \mathcal{C}_3^n as a set of edges. Then $\mathcal{C} \cap V(G_n)$ is an independent set in G_n . It follows that, the possible candidates for \mathcal{C}' are those of the form $\mathcal{C}_1^n \cup X \cup Y$, where X is an independent set of G_n and Y is a subset of \mathcal{C}_4^n . In view of Algorithm 2, analysing all these cases yields 6, 48, and 6 possible 3-clutters \mathcal{C}' , namely \mathcal{D}_i^n 's, for which $\text{index}(J) > 1$ and u, v belong to different connected components of G for $n = 6, 7$, and 8 , respectively. Hence \mathcal{C}' is isomorphic to one of the elements of \mathcal{D} . Since the vertices u, v belong to distinct connected components of $G_{I(\mathcal{D}_i^n)}^{u,v}$ by the argument above, the converse also holds.

Now let $I = I(\mathcal{C})$ be the edge ideal of a 3-uniform clutter \mathcal{C} such that $\text{index}(I) > 1$. Also, let $I_{u,v}$ be the edge ideal of $\mathcal{C}[\text{supp}(uv)]$ for all $u, v \in \mathcal{G}(I^2)$. Theorem 2.1 states that $\text{index}(I^2) > 1$ if and only if $\text{index}(I_{u,v}^2) > 1$ for all monomials $u, v \in \mathcal{G}(I^2)$ or equivalently u, v belong to the same connected component of $G_{I_{u,v}^2}^{u,v}$ for all monomials $u, v \in \mathcal{G}(I^2)$. Our discussions above show that (a) and (b) are equivalent. Clearly, (b) implies (c). It remains to prove that (c) implies (a). Note that

$$\begin{aligned} \beta_{1,8}(I^2) &= \sum_{\substack{w \in L(I^2) \\ \deg w = 8}} \beta_{1,w}(I^2) = \sum_{\substack{w \in L(I^2) \\ \deg w = 8}} \dim_{\mathbb{K}}(\tilde{H}_0(\Delta_{I^2}(1, w); \mathbb{K})) \\ &\geq \dim_{\mathbb{K}}(\tilde{H}_0(\Delta_{I^2}(1, \text{lcm}(u, v)); \mathbb{K})) \\ &= \dim_{\mathbb{K}}(\tilde{H}_0(\Delta_{I_{u,v}^2}(1, \text{lcm}(u, v)); \mathbb{K})) \end{aligned}$$

for all $u, v \in \mathcal{G}(I^2)$. If \mathcal{C} contains an induced subclutter isomorphic to some \mathcal{D}_i^n of \mathcal{D} , then there exist $u, v \in \mathcal{G}(I^2)$ such that $\mathcal{D}_i^n \cong \mathcal{C}[\text{supp}(uv)]$ and u, v belong to distinct connected components of $G_{I_{u,v}^2}^{u,v} = G_{I^2}^{u,v}$. Hence, by Theorem 2.1, $\Delta_{I_{u',v'}^2}(1, \text{lcm}(u', v')) = \Delta_{I^2}(1, \text{lcm}(u', v'))$ is disconnected for some $u', v' \in \mathcal{G}(I^2)$, which implies that $\beta_{1,8}(I^2) \neq 0$. The proof is complete. \square

Let G_F^n be the stabilizer of $\{\{1, 2, 3\}, F\}$ in S_n on its action on $2^{\binom{[n]}{3}}$ for $n \in \{6, 7, 8\}$, and $F = \{1, 2, 4\}$ or $F = \{1, 4, 5\}$. It is easy to see that $G_{\{1, 2, 4\}}^n = \langle (1\ 2), S_{\{4, \dots, n\}} \rangle$ and $G_{\{1, 4, 5\}}^n = \langle (2\ 3), (4\ 5), S_{\{6, \dots, n\}} \rangle$. The following algorithms are used in the proof of Theorem 3.3. The source codes of algorithms in GAP are available in [11].

Algorithm 1

Require: Positive integer $n \in \{6, 7, 8\}$

Ensure: All quadruples of cubic square-free monomials (u_1, u_2, v_1, v_2) such that $\deg \text{lcm}(u_1 u_2, v_1 v_2) > 8$ and $u_i v_j \nmid \text{lcm}(u_1 u_2, v_1 v_2)$ for all $i, j \in \{1, 2\}$ modulo relabeling of indices

```

1:  $\mathcal{S} \leftarrow \binom{[n]}{3}$ 
2:  $U_1 \leftarrow \{1, 2, 3\}$ 
3: for  $U_1 \in \{\{1, 2, 4\}, \{1, 4, 5\}\}$  do
4:    $\mathcal{T}_{V_1} \leftarrow \emptyset$ 
5:   for  $U_2 \in \mathcal{S}$  do
6:     for  $V_2 \in \mathcal{S}$  do
7:        $u \leftarrow \mathbf{x}_{U_1} \mathbf{x}_{U_2}$  and  $v \leftarrow \mathbf{x}_{V_1} \mathbf{x}_{V_2}$ 
8:       if  $\mathbf{x}_{U_i} \mathbf{x}_{V_j} \nmid \text{lcm}(u, v)$  for all  $i, j \in \{1, 2\}$  then
9:          $\mathcal{T}_{V_1} \leftarrow \mathcal{T}_{V_1} \cup \{(\{U_1, U_2\}, \{V_1, V_2\})\}$ 
10:      end if
11:    end for
12:  end for
13:   $\mathcal{O}_{V_1} \leftarrow$  the set of representatives of orbits of  $G_{V_1}^n$  on  $\mathcal{T}_{V_1}$ 
14:   $\mathcal{O}_{V_1} \leftarrow \mathcal{O}_{V_1} \setminus \{(\{U_1, U_2\}, \{V_1, V_2\}) : |(U_1 \cup U_2) \cap (V_1 \cup V_2)| = 5\}$ 
15: end for
16: return  $\{(\mathbf{x}_{U_1}, \mathbf{x}_{U_2}, \mathbf{x}_{V_1}, \mathbf{x}_{V_2}) : (\{U_1, U_2\}, \{V_1, V_2\}) \in \mathcal{O}_{\{1, 2, 4\}} \cup \mathcal{O}_{\{1, 4, 5\}}\}$ 

```

Algorithm 2

Require: Any (u_1, u_2, v_1, v_2) returned by Algorithm 1 for a positive integer $n \in \{6, 7, 8\}$

Ensure: All 3-uniform clutters \mathcal{D} on $[n]$ such that $J = I(\mathcal{D})$ contains u_1, u_2, v_1, v_2 and satisfies $\text{index}(J) > 1$ and $\text{index}(J^2) = 1$

```

1:  $u \leftarrow u_1 u_2$  and  $v \leftarrow v_1 v_2$ 
2: Compute  $\mathcal{C}_2^n, \mathcal{C}_3^n, \mathcal{C}_4^n$ 
3:  $G_n \leftarrow$  simple graph on  $\binom{[n]}{3}$  with edge set  $\mathcal{C}_3^n$ 
4:  $i \leftarrow 0$ 
5: for  $X$  an independent set of  $G_n$  do
6:   for  $Y \subseteq \mathcal{C}_4^n$  do
7:      $\mathcal{C}' \leftarrow \mathcal{C}_1^n \cup X \cup Y$ 
8:      $J \leftarrow I(\mathcal{C}')$ 
9:     if  $\bar{\mathcal{C}}'$  is  $\mathcal{C}$ -free then
10:      if  $u, v$  are in distinct connected components of  $G_{J^2}^{u, v}$  then
11:         $i \leftarrow i + 1$ 
12:         $\mathcal{D}_i^n \leftarrow \mathcal{C}'$ 
13:      end if
14:    end if
15:  end for
16: end for
17: return  $\mathcal{D}_1^n, \dots, \mathcal{D}_i^n$ 

```

Let G be a graph and $I = I(G)$ be the edge ideal of G . If \bar{G} is C_4 -free then $\text{index}(I^k) > 1$ for all k (Theorem 1). Francisco, Hà, and Van Tuyl raised the question whether in this

case $\text{index}(I^k) = \infty$ for all $k \geq 2$? (see [19, Question 1.7]). Nevo and Peeva show by an example that this statement is not true in general (see [19, Counterexample 1.10]). Nevo [18] shows that $\text{index}(I^2) = \infty$ if G is both gap-free and claw-free. Also, Banerjee [1, Theorem 1.2] shows that $\text{index}(I^k) = \infty$ for all k if G is both gap-free and cricket-free. In this regard, we pose the following question in the case of 3-uniform clutters instead of graphs.

Question 1. Let \mathcal{C} be a 3-uniform clutter and $I = I(\mathcal{C})$ be the edge ideal of \mathcal{C} . Assume that $\bar{\mathcal{C}}$ is \mathcal{C} -free and \mathcal{C} is \mathcal{D} -free.

- (i) Under which conditions I^2 has a linear resolution?
- (ii) Under which conditions I^k has a linear resolution for all k ?

For non-negative integers $d \geq 1$ and $k \geq 0$, let $\Omega_{d,k}$ denote the set of all d -uniform clutters \mathcal{C} satisfying the following conditions:

- (i) $\text{index}(I(\mathcal{C})), \dots, \text{index}(I(\mathcal{C})^{k-1}) > 1$,
- (ii) $\text{index}(I(\mathcal{C})^k) = 1$,
- (iii) no proper induced sub-clutter of \mathcal{C} satisfies (i) and (ii).

Also, let $\Omega_{d,k}(n)$ be the number of isomorphism classes of d -uniform clutters of $\Omega_{d,k}$ with n vertices, for all $n \geq 1$. It turns out that,

$$\begin{aligned} \{\Omega_{2,1}(n)\} &= 0, 0, 0, 1, 0, 0, \dots, \\ \{\Omega_{3,1}(n)\} &= 0, 0, 0, 0, 3, 1, 0, 0, \dots, \\ \{\Omega_{3,2}(n)\} &= 0, 0, 0, 0, 0, 6, 48, 6, 0, 0, \dots \end{aligned}$$

Conjecture 3.4. *With the notation as above, the sequence $\{\Omega_{d,k}(n)\}_n$ is unimodal.*

REFERENCES

- [1] A. Banerjee, The regularity of powers of edge ideals, *J. Algebraic Combin.* **41** (2015), 303–321.
- [2] M. Bigdeli, A. A. Yazdan Pour, and R. Zaare-Nahandi, Stability of Betti numbers under reduction processes: Towards chordality of clutters, *J. Combin. Theory Ser. A* **145** (2017), 129–149.
- [3] M. Bigdeli, J. Herzog, and R. Zaare-Nahandi, On the index of powers of edge ideals, *Comm. Algebra* **46**(3) (2018), 1080–1095.
- [4] W. Bruns, A. Conca, and T. Römer, Koszul homology and syzygies of Veronese subalgebras, *Math. Ann.* **351** (2011), 761–779.
- [5] A. Conca, Regularity jumps for powers of ideals, *Commutative algebra*, 21–32, Lect. Notes Pure Appl. Math., 244, *Chapman & Hall/CRC, Boca Raton, FL*, 2006.
- [6] A. Conca and J. Herzog, Castelnuovo-Mumford regularity of products of ideals, *Collect. Math.* **54**(2) (2003), 137–152.
- [7] H. Dao, C. Huneke, and J. Schweig, Bounds on the regularity and projective dimension of ideals associated to graphs, *J. Algebraic Combin.* **38** (2013), 37–55.
- [8] L. Ein and R. Lazarsfeld, Syzygies and Koszul cohomology of smooth projective varieties of arbitrary dimension, *Invent. Math.* **111** (1993), 51–67.
- [9] D. Eisenbud, *The Geometry of Syzygies: A Second Course in Commutative Algebra and Algebraic Geometry*, Graduate Texts in Mathematics **229**, Springer 2005.
- [10] D. Eisenbud, M. Green, K. Hulek, and S. Popescu, Restricting linear syzygies: algebra and geometry, *Compos. Math.* **14**(6) (2005), 1460–1478.
- [11] M. Farrokhi D. G., Source codes of Algorithms 1 and 2, Available at: <https://iasbs.ac.ir/~farrokhi/gap/gli.txt>.
- [12] V. Gasharov, I. Peeva, and V. Welker, The lcm-lattice in monomial resolutions, *Math. Res. Lett.* **6** (1990), 521–532.
- [13] M. L. Green, Koszul cohomology and the geometry of projective varieties, *J. Differential Geom.* **19** (1984), 125–171.

- [14] M. L. Green and R. Lazarsfeld, On the projective normality of complete linear series on an algebraic curve, *Invent. Math.* **83** (1986), 73–90.
- [15] M. L. Green and R. Lazarsfeld, Some results on the syzygies of finite sets and algebraic curves, *Compos. Math.* **67** (1988), 301–314.
- [16] M. Hochster, Cohen-Macaulay rings, combinatorics, and simplicial complexes. Ring theory, II (*Proc. Second Conf., Univ. Oklahoma, Norman, Okla., 1975*), pp. 171–223. Lecture Notes in Pure and Appl. Math., Vol. 26, Dekker, New York, 1977.
- [17] T. Jozefiak, P. Pragacz, and J. Weyman, Resolutions of determinantal varieties and tensor complexes associated with symmetric and antisymmetric matrices, *Astérisque* **87-88** (1981), 109–189.
- [18] E. Nevo, Regularity of edge ideals of C_4 -free graphs via the topology of the lcm-lattice, *J. Combin. Theory Ser. A* **118** (2011), 491–501.
- [19] E. Nevo and I. Peeva, C_4 -free edge ideals, *J. Algebraic Combin.* **37** (2013), 243–248.
- [20] G. Ottaviani and R. Paoletti, Syzygies of Veronese embeddings, *Compos. Math.* **125** (2011), 31–37.
- [21] E. Rubei, A result on resolutions of Veronese embeddings, *Ann. Univ. Ferrara Sez. VII (N. S.)* **50** (2004), 151–165.
- [22] B. Sturmfels, Four counterexamples in combinatorial algebraic geometry, *J. Algebra* **230** (2000), 282–294.
- [23] J. Weyman, *Cohomology of Vector Bundles and Syzygies*, Cambridge University Press 2003.
- [24] A. A. Yazdan Pour, Candidates for nonzero Betti numbers of monomial ideals, *Commun. Algebra* **45**(4) (2017), 1483–1492.

Email address: m.farrokhi.d.g@gmail.com, farrokhi@iasbs.ac.ir

RESEARCH CENTER FOR BASIC SCIENCES AND MODERN TECHNOLOGIES (RBST), INSTITUTE FOR ADVANCED STUDIES IN BASIC SCIENCES (IASBS), ZANJAN 45137-66731, IRAN

Email address: yassin.sadegh@yahoo.com, y.sadegh@uma.ac.ir

DEPARTMENT OF MATHEMATICS, FACULTY OF MATHEMATICAL SCIENCES, UNIVERSITY OF MOHAGHEGH ARDABIL, 56199-11367, ARDABIL, IRAN.

Email address: yazdan@iasbs.ac.ir

DEPARTMENT OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDIES IN BASIC SCIENCES (IASBS), ZANJAN 45137-66731, IRAN