On the average hitting times of the squares of cycles

Yoshiaki Doi, Norio Konno, Tomoki Nakamigawa, Tadashi Sakuma, Etsuo Segawa, Hidehiro Shinohara, Shunya Tamura, Yuuho Tanaka, Kosuke Toyota,

Abstract

The exact formula for the average hitting time (HT, as an abbreviation) of simple random walks from one vertex to any other vertex on the square C_N^2 of an N-vertex cycle graph C_N was given by N. Chair [Journal of Statistical Physics, 154 (2014) 1177-1190]. In that paper, the author gives the expression for the even N case and the expression for the odd N case separately. In this paper, by using an elementary method different from Chair (2014), we give a much simpler single formula for the HT's of simple random walks on C_N^2 . Our proof is considerably short and fully combinatorial, in particular, has no-need of any spectral graph theoretical arguments. Not only the formula itself but also intermediate results through the process of our proof describe clear relations between the HT's of simple random walks on C_N^2 and the Fibonacci numbers.

Keywords: simple random walk, hitting time, the square of a cycle, Fibonacci number, Kirchhoff index

1 Introduction

A simple random walk on a graph G is a discrete stochastic model such that a random walker on a vertex $u \in V(G)$ moves to a vertex v adjacent to the vertex u at the next step with the probability of $\frac{1}{\deg_G(u)}$, where $\deg_G(u)$ denotes the degree of the vertex u of G. The number of steps required until the random walker starting from a vertex p of G will first arrive at a vertex q of G is called the hitting time from p to q of the simple random walk on G. The average hitting time (HT, as an abbreviation) from p to q on G which is denoted by h(G; p, q), means the expected value of the hitting times from p to q of simple random walks on G. The exact formula for the average hitting time from one vertex to any other vertex is far from available in general. For some very special graph classes with high symmetries, it

^{*}Sakata City Daini Junior High School, Shinbashi, Sakata 998-0864, Japan, y.doi6532@gmail.com

[†]Department of Applied Mathematics, Faculty of Engineering, Yokohama National University, Hodogaya, Yokohama, 240-8501, Japan, konno-norio-bt@ynu.ac.jp

[‡]Department of Information Science Shonan Institute of Technology, Tsujido-Nishikaigan, Fujisawa, 251-8511, Japan, nakami@info.shonan-it.ac.jp

[§]Faculty of Science, Yamagata University, Kojirakawa, Yamagata, 990–8560, Japan, sakuma@sci.kj.yamagata-u.ac.jp

 $[\]P$ Graduate School of Education Center and Graduate School of Environment Information Sciences, Yokohama National University, Hodogaya, Yokohama, 240-8501, Japan, segawa-etsuo-tb@ynu.ac.jp

Institute for Excellence in Higher Education, Tohoku University, Aoba, Sendai, 980-8576, Japan, shino.setsystem@gmail.com

^{**}Graduate School of Science and Engineering, Yokohama National University, Hodogaya, Yokohama, 240-8501, Japan, tamura-shunya-kj@ynu.jp

^{††}Graduate School of Information Sciences, Tohoku University Aoba, Sendai, 980-8579, Japan, tanaka.yuuho.t4@dc.tohoku.ac.jp

^{‡‡}Graduate School of Science and Engineering (Science), Yamagata University, Kojirakawa, Yamagata, 990–8560, Japan, s201001m@st.yamagata-u.ac.jp

may be possible to obtain such exact formulas. For example, as far as the authors know, only in the case of $i \in \{1,2\}$, exact formulas for the i-th power C_N^i of an N-vertex cycle graph C_N are obtained. Furthermore, it was not until 2014 that such a formula for the case of i=2 was given by N. Chair [1]. In that paper, by calculating the exact value of Wu's formula [9] of the effective resistances in terms of the eigenvalues and the eigenvectors of the Laplacian matrices of C_N^2 , the author gives the expression for the even N case and the expression for the odd N case separately. Let us define $V(C_N^2) := \mathbb{Z}_N$ and $E(C_N^2) := \{\{u,v\} \mid u-v=\pm 1, \pm 2\}$. Let us denote by $h_N(k,l)$ the average hitting time for simple random walks from the vertex k to the vertex k of C_N^2 . Note that k0,00 = 0. Since the dihedral group D_N acts on C_N^2 , we have that $\forall k,l \in \mathbb{Z}_N, h_N(0,l) = h_N(k,k+l) = h_N(k+l,k)$. Let F_i denote the i-th Fibonacci number.

Theorem 1.1 (Chair, 2014 [1]). The exact formula for the HT's of simple random walks on $C_N^2(N \ge 5)$ is as follows:

1. For the case of N=2n

$$h_N(0,l) = \frac{2}{5}l(N-l) + (-1)^{l+1} \frac{2N}{\sqrt{5}} F_l^2 \left(\frac{1 + \left(\frac{3-\sqrt{5}}{2}\right)^N}{1 - \left(\frac{3-\sqrt{5}}{2}\right)^N} \right) + (-1)^l \frac{2N}{5} F_{2l}.$$

2. For the case of N = 2n + 1,

$$h_N(0,l) = \frac{2}{5}l\left(N-l\right) + (-1)^{l+1}\frac{2N}{\sqrt{5}}F_l^2\left(\frac{1-\left(\frac{3-\sqrt{5}}{2}\right)^N}{1+\left(\frac{3-\sqrt{5}}{2}\right)^N}\right) + (-1)^l\frac{2N}{5}F_{2l}.$$

In this paper, by using an elementary method different from Chair [1], we give a simpler single formula for HT's of simple random walks on C_N^2 , as follows:

Theorem 1.2. The exact formula for the HT's of simple random walks on $C_N^2(N \ge 5)$ is,

$$h_N(0,l) = \frac{2}{5} \left(l (N-l) + 2N \frac{F_l F_{N-l}}{F_N} \right).$$

Our proof is considerably short and fully combinatorial. In particular, it has no-need of any spectral graph theoretical arguments. Not only the formula itself but also intermediate results through the process of our proof describe clear relations between the HT's of simple random walks on C_N^2 and the Fibonacci numbers.

2 Preliminary

In this section, we prepare some well-known formulas on Fibonacci numbers, which will be used throughout this paper. For proofs of these formulas, please refer to the appropriate literature on Fibonacci numbers (e.g., [7, 8]).

$$F_{n+1}F_{n-1} - F_n^2 = (-1)^n. (2.1)$$

$$F_{2n} - 3F_{2(n-1)} + F_{2(n-2)} = 0. (2.2)$$

$$F_{2n+1} = F_{n+1}^2 + F_n^2. (2.3)$$

$$F_{m-n} = (-1)^{n+1} F_{m-1} F_n + (-1)^n F_m F_{n-1}.$$
(2.4)

$$\frac{1}{F_{2n-1}F_{2n+1}} = \frac{F_{2n+2}}{F_{2n+1}} - \frac{F_{2n}}{F_{2n-1}}. (2.5)$$

$$\sum_{l=1}^{l} (-1)^{l-k} F_{2k-1} = F_l^2. \tag{2.6}$$

$$F_l^2 = \frac{1}{5}(F_{2l-1} + F_{2l+1}) + \frac{2}{5}(-1)^{l-1}.$$
 (2.7)

Proof of Theorem 1.2 3

Let L denote the Laplacian matrix of C_N^2 and let L' denote the matrix obtained from L by deleting its first row and column. Let \vec{h} be the column vector whose i-th entry is $h_N(0,i)$. Let $\vec{1}$ be a column vector of proper dimension whose entries are all 1. For the case of a random walk on C_N^2 , a random walker moves, with the probability of $\frac{1}{4}$, to an arbitrary vertex adjacent to the vertex where the walker is. Hence we have that $\forall l \in \mathbb{Z}_N, \ h_N(0,l) = \frac{1}{4} \left((1 + h_N(-2,l)) + (1 + h_N(-1,l)) + (1 + h_N(1,l)) + (1 + h_N(2,l)) \right),$ and hence $\forall l \in \mathbb{Z}_N, -h_N(0,l-2) - h_N(0,l-1) + 4h_N(0,l) - h_N(0,l+1) - h_N(0,l+2) = 4.$ Thus, we have $L'\tilde{h}=4\tilde{1}$. Combining this with the fact that $h_N(0,l)=h_N(0,N-l)$ for all $l \in \mathbb{Z}_N$, we can halve the number of variables of our problem. Let L'(i,j) denote the (i,j)-th entry of L'. Let H_N be the $\lfloor N/2 \rfloor \times \lfloor N/2 \rfloor$ -matrix whose (i,j)-th entry is L'(i,j) if $j = \frac{N}{2}(N = 2n)$, and L'(i, j) + L'(i, N - j) otherwise. Let \vec{x} be an $\lfloor N/2 \rfloor$ -dimensional variable column vector whose i-th entry is corresponding to $h_N(0,i)$. Then we have $H_N\vec{x}=4\vec{1}$. For our problem, it suffices to solve this matrix equation. As a technical issue, in the case of N=2n, our matrix H_{2n} is not symmetric. However, if we multiply the last row of this matrix by $\frac{1}{2}$, then the resultant matrix H'_{2n} will be symmetric. Hence, only in the case of N=2n, we redefine $H_{2n}:=H'_{2n}$ and change our matrix equation to $H_{2n}\vec{x}=^t(4,\ldots,4,2)$.

Let U_N be the upper triangular matrix whose (i, j)-th entry is 1 if $j \geq i$.

Let W_N be the following matrix:

$$W_N := \begin{bmatrix} 1 & 0 & 0 & \cdots & 0 & 0 \\ \frac{F_1}{F_3} & 1 & 0 & \cdots & 0 & 0 \\ 0 & \frac{F_3}{F_5} & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 0 \\ 0 & 0 & 0 & \cdots & \frac{F_2 \lfloor \frac{N}{2} \rfloor - 3}{F_2 \lfloor \frac{N}{2} \rfloor - 1} & 1 \end{bmatrix}.$$

Then its inverse W_N^{-1} is as follows:

$$W_N^{-1} = \begin{bmatrix} 1 & 0 & 0 & \cdots & 0 & 0 \\ -\frac{F_1}{F_3} & 1 & 0 & \cdots & 0 & 0 \\ \frac{F_1}{F_5} & -\frac{F_3}{F_5} & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ (-1)^{\left\lfloor \frac{N}{2} \right\rfloor - 2} \frac{F_1}{F_2 \left\lfloor \frac{N}{2} \right\rfloor - 3} & (-1)^{\left\lfloor \frac{N}{2} \right\rfloor - 3} \frac{F_3}{F_2 \left\lfloor \frac{N}{2} \right\rfloor - 3} & (-1)^{\left\lfloor \frac{N}{2} \right\rfloor - 3} \frac{F_5}{F_2 \left\lfloor \frac{N}{2} \right\rfloor - 3} & \cdots & 1 & 0 \\ (-1)^{\left\lfloor \frac{N}{2} \right\rfloor - 1} \frac{F_1}{F_2 \left\lfloor \frac{N}{2} \right\rfloor - 1} & (-1)^{\left\lfloor \frac{N}{2} \right\rfloor - 2} \frac{F_3}{F_2 \left\lfloor \frac{N}{2} \right\rfloor - 1} & (-1)^{\left\lfloor \frac{N}{2} \right\rfloor - 3} \frac{F_5}{F_2 \left\lfloor \frac{N}{2} \right\rfloor - 1} & \cdots & -\frac{F_2 \left\lfloor \frac{N}{2} \right\rfloor - 3}{F_2 \left\lfloor \frac{N}{2} \right\rfloor - 1} & 1 \end{bmatrix}.$$

Let D_N be the following matrix:

$$D_N := \begin{bmatrix} \frac{F_3}{F_1} & 0 & 0 & \cdots & 0 & 0\\ 0 & \frac{F_5}{F_3} & 0 & \cdots & 0 & 0\\ 0 & 0 & \frac{F_7}{F_5} & \cdots & 0 & 0\\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots\\ 0 & 0 & 0 & \cdots & \frac{F_2 \left\lfloor \frac{N}{2} \right\rfloor - 1}{F_2 \left\lfloor \frac{N}{2} \right\rfloor - 3} & 0\\ 0 & 0 & 0 & \cdots & 0 & \frac{F_N}{F_2 \left\lfloor \frac{N}{2} \right\rfloor - 1} \end{bmatrix}.$$

Then we have the following:

Theorem 3.1. $H_N = U_N^{-1} W_N D_N^{\ t} W_N^{\ t} U_N^{-1}$.

Proof of Theorem 3.1.

Let S_N denote the matrix $U_N^{-1}W_ND_N{}^tW_N{}^tU_N^{-1}$. By simple matrix-computation, the entries of S_N can be calculated as follows.

$$S_N(i,j) = \begin{cases} 0 & (|i-j| \geq 3), \\ -1 & (|i-j| = 2), \\ 2 - \frac{F_{2i-3} + F_{2i+1}}{F_{2i-1}} & (j-i=1, j \neq \left\lfloor \frac{N}{2} \right\rfloor), \\ 1 - \frac{F_{2\lfloor \frac{N}{2} \rfloor - 3} + F_N}{F_{2\lfloor \frac{N}{2} \rfloor - 1}} & ((i,j) = (\left\lfloor \frac{N}{2} \right\rfloor - 1, \left\lfloor \frac{N}{2} \right\rfloor)), \\ \frac{F_{2\lfloor \frac{N}{2} \rfloor - 1} + F_{2i+1}}{F_{2i-1}} + \frac{F_{2i-1} + F_{2i+3}}{F_{2i+1}} - 2 & (i,j) = (1,1), \\ \frac{F_{2\lfloor \frac{N}{2} \rfloor - 1} + F_{2\lfloor \frac{N}{2} \rfloor - 3}}{F_{2\lfloor \frac{N}{2} \rfloor - 3}} + \frac{F_{2\lfloor \frac{N}{2} \rfloor - 3} + F_N}{F_{2\lfloor \frac{N}{2} \rfloor - 1}} - 2 & ((i,j) = (\left\lfloor \frac{N}{2} \right\rfloor - 1, \left\lfloor \frac{N}{2} \right\rfloor)), \\ \frac{F_{2\lfloor \frac{N}{2} \rfloor - 3} + F_N}{F_{2\lfloor \frac{N}{2} \rfloor - 3}} + \frac{F_{2\lfloor \frac{N}{2} \rfloor - 3} + F_N}{F_{2\lfloor \frac{N}{2} \rfloor - 1}} - 2 & ((i,j) = (\left\lfloor \frac{N}{2} \right\rfloor - 1, \left\lfloor \frac{N}{2} \right\rfloor - 1)), \\ \frac{F_{2\lfloor \frac{N}{2} \rfloor - 3} + F_N}{F_{2\lfloor \frac{N}{2} \rfloor - 1}} & ((i,j) = (\left\lfloor \frac{N}{2} \right\rfloor - 1, \left\lfloor \frac{N}{2} \right\rfloor)). \end{cases}$$
In the case that $|i-i| = 1, i \neq \lfloor \frac{N}{2} \rfloor$ we have

In the case that $|i-j|=1, j\neq \lfloor \frac{N}{2} \rfloor$, we have

$$S_N(i,j) = 2 - \frac{F_{2l-1} + F_{2l+3}}{F_{2l+1}}$$

$$= 2 - \frac{3F_{2l+1}}{F_{2l+1}}$$

$$= -1.$$
 (by (2.2))

In the case that $i = j \notin \{1, \lfloor \frac{N}{2} \rfloor - 1, \lfloor \frac{N}{2} \rfloor \}$, we have

$$S_N(i,j) = \frac{F_{2l-1} + F_{2l+3}}{F_{2l+1}} + \frac{F_{2l+1} + F_{2l+5}}{F_{2l+3}} - 2 = \frac{3F_{2l+1}}{F_{2l+1}} + \frac{3F_{2l+3}}{F_{2l+3}} - 2 \qquad \text{(by (2.2))}$$

$$= 4.$$

In the case of $S_N(1,1)$, we have

$$S_N(1,1) = \frac{F_3}{F_1} + \frac{F_1 + F_5}{F_3} - 2 = 2 + \frac{1+5}{2} - 2 = 3.$$

We will show $S_N(\left\lfloor \frac{N}{2} \right\rfloor - 1, \left\lfloor \frac{N}{2} \right\rfloor - 1) = H_N(\left\lfloor \frac{N}{2} \right\rfloor - 1, \left\lfloor \frac{N}{2} \right\rfloor - 1)$. In the case that N is odd,

$$\begin{split} S_N\left(\left\lfloor \frac{N}{2} \right\rfloor - 1, \left\lfloor \frac{N}{2} \right\rfloor - 1\right) &= \frac{F_{2\lfloor \frac{N}{2} \rfloor - 1} + F_{2\lfloor \frac{N}{2} \rfloor - 5}}{F_{2\lfloor \frac{N}{2} \rfloor - 3}} + \frac{F_{2\lfloor \frac{N}{2} \rfloor - 3} + F_N}{F_{2\lfloor \frac{N}{2} \rfloor - 1}} - 2 \\ &= \frac{F_{2\lfloor \frac{N}{2} \rfloor - 1} + F_{2\lfloor \frac{N}{2} \rfloor - 5}}{F_{2\lfloor \frac{N}{2} \rfloor - 3}} + \frac{F_{2\lfloor \frac{N}{2} \rfloor - 3} + F_{2\lfloor \frac{N}{2} \rfloor + 1}}{F_{2\lfloor \frac{N}{2} \rfloor - 1}} - 2 \\ &= \frac{3F_{2\lfloor \frac{N}{2} \rfloor - 3}}{F_{2\lfloor \frac{N}{2} \rfloor - 3}} + \frac{3F_{2\lfloor \frac{N}{2} \rfloor - 1}}{F_{2\lfloor \frac{N}{2} \rfloor - 1}} - 2 \\ &= 4. \end{split} \tag{by (2.2)}$$

In the case that N is even,

$$\begin{split} S_{N}\left(\left\lfloor\frac{N}{2}\right\rfloor-1,\left\lfloor\frac{N}{2}\right\rfloor-1\right) &= \frac{F_{2\lfloor\frac{N}{2}\rfloor-1}+F_{2\lfloor\frac{N}{2}\rfloor-5}}{F_{2\lfloor\frac{N}{2}\rfloor-3}} + \frac{F_{2\lfloor\frac{N}{2}\rfloor-3}+F_{N}}{F_{2\lfloor\frac{N}{2}\rfloor-1}} - 2\\ &= \frac{F_{2\lfloor\frac{N}{2}\rfloor-1}+F_{2\lfloor\frac{N}{2}\rfloor-5}}{F_{2\lfloor\frac{N}{2}\rfloor-3}} + \frac{F_{2\lfloor\frac{N}{2}\rfloor-3}+F_{2\lfloor\frac{N}{2}\rfloor}}{F_{2\lfloor\frac{N}{2}\rfloor-1}} - 2\\ &= \frac{F_{2\lfloor\frac{N}{2}\rfloor-1}+F_{2\lfloor\frac{N}{2}\rfloor-5}}{F_{2\lfloor\frac{N}{2}\rfloor-3}} + \frac{F_{2\lfloor\frac{N}{2}\rfloor-3}+F_{2\lfloor\frac{N}{2}\rfloor+1}-F_{2\lfloor\frac{N}{2}\rfloor-1}}{F_{2\lfloor\frac{N}{2}\rfloor-1}} - 2\\ &= \frac{3F_{2\lfloor\frac{N}{2}\rfloor-3}}{F_{2\lfloor\frac{N}{2}\rfloor-3}} + \frac{2F_{2\lfloor\frac{N}{2}\rfloor-1}}{F_{2\lfloor\frac{N}{2}\rfloor-1}} - 2 & \text{ (by (2.2))} \\ &= 3 \end{split}$$

We will show $S_N(\lfloor \frac{N}{2} \rfloor - 1, \lfloor \frac{N}{2} \rfloor) = H_N(\lfloor \frac{N}{2} \rfloor - 1, \lfloor \frac{N}{2} \rfloor)$. In the case that N is odd,

$$S_N\left(\left\lfloor \frac{N}{2} \right\rfloor - 1, \left\lfloor \frac{N}{2} \right\rfloor\right) = 1 - \frac{F_{2\lfloor \frac{N}{2} \rfloor - 3} + F_N}{F_{2\lfloor \frac{N}{2} \rfloor - 1}}$$

$$= 1 - \frac{F_{2\lfloor \frac{N}{2} \rfloor - 3} + F_{2\lfloor \frac{N}{2} \rfloor + 1}}{F_{2\lfloor \frac{N}{2} \rfloor - 1}}$$

$$= 1 - \frac{3F_{2\lfloor \frac{N}{2} \rfloor - 1}}{F_{2\lfloor \frac{N}{2} \rfloor - 1}} \qquad \text{(by (2.2))}$$

$$= -2.$$

In the case that N is even,

$$S_{N}\left(\left\lfloor \frac{N}{2} \right\rfloor - 1, \left\lfloor \frac{N}{2} \right\rfloor\right) = 1 - \frac{F_{2\lfloor \frac{N}{2} \rfloor - 3} + F_{N}}{F_{2\lfloor \frac{N}{2} \rfloor - 1}}$$

$$= 1 - \frac{F_{2\lfloor \frac{N}{2} \rfloor - 3} + F_{2\lfloor \frac{N}{2} \rfloor}}{F_{2\lfloor \frac{N}{2} \rfloor - 1}}$$

$$= 1 - \frac{F_{2\lfloor \frac{N}{2} \rfloor - 3} + F_{2\lfloor \frac{N}{2} \rfloor + 1} - F_{2\lfloor \frac{N}{2} \rfloor - 1}}{F_{2\lfloor \frac{N}{2} \rfloor - 1}}$$

$$= 1 - \frac{2F_{2\lfloor \frac{N}{2} \rfloor - 1}}{F_{2\lfloor \frac{N}{2} \rfloor - 1}} \qquad \text{(by (2.2))}$$

$$= -1$$

Last, we will show $S_N(\left\lfloor \frac{N}{2} \right\rfloor, \left\lfloor \frac{N}{2} \right\rfloor) = H_N(\left\lfloor \frac{N}{2} \right\rfloor, \left\lfloor \frac{N}{2} \right\rfloor)$ In the case that N is odd,

$$\begin{split} S_N\left(\left\lfloor\frac{N}{2}\right\rfloor, \left\lfloor\frac{N}{2}\right\rfloor\right) &= \frac{F_2\lfloor\frac{N}{2}\rfloor - 3 + F_N}{F_2\lfloor\frac{N}{2}\rfloor - 1} \\ &= \frac{F_2\lfloor\frac{N}{2}\rfloor - 3 + F_2\lfloor\frac{N}{2}\rfloor + 1}{F_2\lfloor\frac{N}{2}\rfloor - 1} \\ &= \frac{3F_2\lfloor\frac{N}{2}\rfloor - 1}{F_2\lfloor\frac{N}{2}\rfloor - 1} \\ &= 3. \end{split} \tag{by (2.2)}$$

In the case that N is even,

$$S_{N}\left(\left\lfloor \frac{N}{2} \right\rfloor, \left\lfloor \frac{N}{2} \right\rfloor\right) = \frac{F_{2\lfloor \frac{N}{2} \rfloor - 3} + F_{N}}{F_{2\lfloor \frac{N}{2} \rfloor - 1}}$$

$$= \frac{F_{2\lfloor \frac{N}{2} \rfloor - 3} + F_{2\lfloor \frac{N}{2} \rfloor}}{F_{2\lfloor \frac{N}{2} \rfloor - 1}}$$

$$= \frac{F_{2\lfloor \frac{N}{2} \rfloor - 3} + F_{2\lfloor \frac{N}{2} \rfloor + 1} - F_{2\lfloor \frac{N}{2} \rfloor - 1}}{F_{2\lfloor \frac{N}{2} \rfloor - 1}}$$

$$= \frac{2F_{2\lfloor \frac{N}{2} \rfloor - 1}}{F_{2\lfloor \frac{N}{2} \rfloor - 1}}$$

$$= 2$$
(by (2.2))

Therefore, we have $S_N = H_N$.

Now let us define the new variable vector $\vec{y} = {}^tU_N^{-1}\vec{x}$. Then both the matrix equation $H_N\vec{x} = 4\vec{1}$ in the case of N = 2n + 1 and the matrix equation $H_N\vec{x} = {}^t(4, 4, \dots, 2)$ in the case of N = 2n can be expressed by the following single matrix equation.

$$\vec{y} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_l \\ \vdots \\ y_{\left \lfloor \frac{N}{2} \right \rfloor} \end{bmatrix} := \begin{bmatrix} x_1 \\ x_2 - x_1 \\ \vdots \\ x_l - x_{l-1} \\ \vdots \\ x_{\left \lfloor \frac{N}{2} \right \rfloor} - x_{\left \lfloor \frac{N}{2} \right \rfloor - 1} \end{bmatrix} = 2 \ ^t W_N^{-1} D_N^{-1} W_N^{-1} \begin{bmatrix} N - 1 \\ N - 3 \\ \vdots \\ N - 2l + 1 \\ \vdots \\ N - 1 - 2(\left \lfloor \frac{N}{2} \right \rfloor - 1) \end{bmatrix}.$$

Again, let us define the new variable vector $\vec{z} := {}^tW_N\vec{y}$. Then we have

$$\vec{z} = \begin{bmatrix} z_1 \\ z_2 \\ \vdots \\ z_{\left\lfloor \frac{N}{2} \right\rfloor - 1} \\ z_{\left\lfloor \frac{N}{2} \right\rfloor} \end{bmatrix} = 2D_N^{-1}W_N^{-1} \begin{bmatrix} N - 1 \\ N - 3 \\ \vdots \\ N - 1 - 2(\left\lfloor \frac{N}{2} \right\rfloor - 2) \\ N - 1 - 2(\left\lfloor \frac{N}{2} \right\rfloor - 1) \end{bmatrix}.$$

By solving this matrix equation, we obtain the following.

Lemma 3.1.

$$1 \leq \forall l \leq \left\lfloor \frac{N}{2} \right\rfloor - 1, \quad z_l = \frac{2}{5} \left(N - 2l + 1 + \frac{(N - 2l - 1)F_{2l - 1} + 2N(-1)^{l - 1}}{F_{2l + 1}} \right).$$

$$z_{\left\lfloor \frac{N}{2} \right\rfloor} = \frac{2}{5F_N} \left(\left(N - 2 \left\lfloor \frac{N}{2} \right\rfloor \right) \left(F_{2\left\lfloor \frac{N}{2} \right\rfloor + 1} + F_{2\left\lfloor \frac{N}{2} \right\rfloor - 1} \right) + F_{2\left\lfloor \frac{N}{2} \right\rfloor} + 2N(-1)^{\left\lfloor \frac{N}{2} \right\rfloor - 1} \right).$$

 $Proof\ of\ Lemma\ 3.1.$

For the case that $1 \leq \forall l \leq \lfloor \frac{N}{2} \rfloor - 1$, we have

$$z_{l} = 2D_{N}^{-1}W_{N}^{-1t}\left[N-1, N-3, \cdots, N-1-2(\left\lfloor\frac{N}{2}\right\rfloor-1)\right]$$

$$= \frac{2}{F_{2l+1}}\sum_{k=1}^{l}(-1)^{l-k}(N-2k+1)F_{2k-1}$$

$$= \frac{2}{F_{2l+1}}\left((N+1)\sum_{k=1}^{l}(-1)^{l-k}F_{2k-1}-2\sum_{k=1}^{l}kF_{2k-1}\right)$$

$$= \frac{2}{F_{2l+1}}\left((N+1)F_{l}^{2}-\frac{2}{5}F_{2l-1}+\frac{2}{5}(-1)^{l-1}(2l-1)-2lF_{l}^{2}\right) \qquad (by (2.6))$$

$$= \frac{2}{5F_{2l+1}}\left(5(N-2l+1)F_{l}^{2}-2F_{2l-1}+2(-1)^{l-1}(2l-1)\right)$$

$$= \frac{2}{5F_{2l+1}}\left((N-2l+1)(F_{2l-1}+F_{2l+1}+2(-1)^{l-1})-2F_{2l-1}+2(-1)^{l-1}(2l-1)\right) \qquad (by (2.7))$$

$$= \frac{2}{5F_{2l+1}}\left((N-2l-1)F_{2l-1}+(N-2l+1)F_{2l+1}+2N(-1)^{l-1}\right)$$

$$= \frac{2}{5}\left((N-2l+1)+\frac{(N-2l-1)F_{2l-1}+2N(-1)^{l-1}}{F_{2l+1}}\right).$$

In the case of $z_{\lfloor \frac{N}{2} \rfloor}$, we have

$$\begin{split} z_{\lfloor \frac{N}{2} \rfloor} &= \frac{2}{F_N} \sum_{k=1}^{\lfloor \frac{N}{2} \rfloor} (-1)^{\lfloor \frac{N}{2} \rfloor - k} (N - 2k + 1) F_{2k-1} \\ &= \frac{2}{F_N} \left(\sum_{k=1}^{\lfloor \frac{N}{2} \rfloor} (-1)^{\lfloor \frac{N}{2} \rfloor - k} F_{2k-1} - 2 \sum_{k=1}^{\lfloor \frac{N}{2} \rfloor} k F_{2k-1} \right) \\ &= \frac{2}{F_N} \left((N+1) F_{\lfloor \frac{N}{2} \rfloor}^2 - \frac{2}{5} F_{2\lfloor \frac{N}{2} \rfloor - 1} + \frac{2}{5} (-1)^{\lfloor \frac{N}{2} \rfloor - 1} \left(2 \lfloor \frac{N}{2} \rfloor - 1 \right) - 2 \lfloor \frac{N}{2} \rfloor F_{\lfloor \frac{N}{2} \rfloor}^2 \right) \quad \text{(by (2.6))} \\ &= \frac{2}{5F_N} \left(5 (N - 2 \lfloor \frac{N}{2} \rfloor + 1) F_{\lfloor \frac{N}{2} \rfloor}^2 - 2 F_{2\lfloor \frac{N}{2} \rfloor - 1} + 2 (-1)^{\lfloor \frac{N}{2} \rfloor - 1} \left(2 \lfloor \frac{N}{2} \rfloor - 1 \right) \right) \\ &= \frac{2}{5F_N} \left(\left(N - 2 \lfloor \frac{N}{2} \rfloor + 1 \right) \left(F_{2\lfloor \frac{N}{2} \rfloor - 1} + F_{2\lfloor \frac{N}{2} \rfloor + 1} + 2 (-1)^{\lfloor \frac{N}{2} \rfloor - 1} \right) \right) \\ &- 2 F_{2\lfloor \frac{N}{2} \rfloor - 1} + (-1)^{\lfloor \frac{N}{2} \rfloor - 1} \left(4 \lfloor \frac{N}{2} \rfloor - 2 \right) \right) \\ &= \frac{2}{5F_N} \left(\left(N - 2 \lfloor \frac{N}{2} \rfloor \right) \left(F_{2\lfloor \frac{N}{2} \rfloor - 1} + F_{2\lfloor \frac{N}{2} \rfloor + 1} \right) + F_{2\lfloor \frac{N}{2} \rfloor} + (-1)^{\lfloor \frac{N}{2} \rfloor - 1} 2 N \right). \end{split}$$

Combining the above lemma with the equation $\vec{y} = {}^tW_N^{-1}\vec{z},$ we have the following.

Lemma 3.2.

$$y_l = \frac{2}{5} \left((N - 2l + 1) + 2N \frac{(-1)^{l-1} F_{N-2l+1}}{F_N} \right).$$

Proof of Lemma 3.2.

$$\begin{split} y_{l} &= F_{2l-1} \sum_{i=l}^{\left \lfloor \frac{N}{2} \right \rfloor - 1} (-1)^{l+i} \frac{z_{i}}{F_{2i-1}} + (-1)^{l+\left \lfloor \frac{N}{2} \right \rfloor} \frac{F_{2l-1}}{F_{2\left \lfloor \frac{N}{2} \right \rfloor - 1}} z_{\left \lfloor \frac{N}{2} \right \rfloor} \\ &= \frac{2}{5} F_{2l-1} \Biggl(\sum_{i=l}^{\left \lfloor \frac{N}{2} \right \rfloor - 1} (-1)^{l+i} \frac{(N-2i+1)F_{2i+1} + (N-2i-1)F_{2i-1} + 2N(-1)^{i-1}}{F_{2i-1}F_{2i+1}} \\ &+ \frac{(-1)^{l+\left \lfloor \frac{N}{2} \right \rfloor}}{F_{N}F_{2\left \lfloor \frac{N}{2} \right \rfloor - 1}} \left(\left(N-2 \left \lfloor \frac{N}{2} \right \rfloor + 1 \right) F_{2\left \lfloor \frac{N}{2} \right \rfloor + 1} + \left(N-2 \left \lfloor \frac{N}{2} \right \rfloor - 1 \right) F_{2\left \lfloor \frac{N}{2} \right \rfloor - 1} + 2N(-1)^{\left \lfloor \frac{N}{2} \right \rfloor - 1} \right) \right) \\ &= \frac{2}{5} F_{2l-1} \Biggl(\sum_{i=l}^{\left \lfloor \frac{N}{2} \right \rfloor - 1} (-1)^{l+i} \left(\frac{(N-2i-1)(F_{2i+1} + F_{2i-1}) + 2F_{2i+1} - 2N(-1)^{i}}{F_{2i-1}F_{2i+1}} \right) \\ &+ \frac{(-1)^{l+\left \lfloor \frac{N}{2} \right \rfloor}}{F_{N}F_{2\left \lfloor \frac{N}{2} \right \rfloor - 1}} \left((N-2 \left \lfloor \frac{N}{2} \right \rfloor + 1) F_{2\left \lfloor \frac{N}{2} \right \rfloor + 1} + (N-2 \left \lfloor \frac{N}{2} \right \rfloor - 1) F_{2\left \lfloor \frac{N}{2} \right \rfloor - 1} + 2N(-1)^{\left \lfloor \frac{N}{2} \right \rfloor - 1} \right) \right) \right) \\ &= \frac{2}{5} F_{2l-1} \left(\frac{N-2l+1}{F_{2l-1}} + 2N(-1)^{l-1} \sum_{i=l}^{\left \lfloor \frac{N}{2} \right \rfloor - 1} \frac{1}{F_{2i-1}F_{2i+1}} + \frac{2N(-1)^{l-1}}{F_{N}F_{2\left \lfloor \frac{N}{2} \right \rfloor - 1}} \right) \right) \end{split}$$

$$= \frac{2}{5}F_{2l-1} \left(\frac{N-2l+1}{F_{2l-1}} + 2N(-1)^{l-1} \sum_{i=l}^{\left\lfloor \frac{N}{2} \right\rfloor - 1} \left(\frac{F_{2i+2}}{F_{2i+1}} - \frac{F_{2i}}{F_{2i-1}} \right) + \frac{2N(-1)^{l-1}}{F_N F_{2\left\lfloor \frac{N}{2} \right\rfloor - 1}} \right)$$

$$= \frac{2}{5}F_{2l-1} \left(\frac{N-2l+1}{F_{2l-1}} + 2N(-1)^{l-1} \left(\frac{F_{2\left\lfloor \frac{N}{2} \right\rfloor}}{F_{2\left\lfloor \frac{N}{2} \right\rfloor - 1}} - \frac{F_{2l}}{F_{2l-1}} \right) + 2N(-1)^{l-1} \left(\frac{F_{N+1}}{F_N} - \frac{F_{2\left\lfloor \frac{N}{2} \right\rfloor}}{F_{2\left\lfloor \frac{N}{2} \right\rfloor - 1}} \right) \right)$$

$$= \frac{2}{5} \left(N - 2l + 1 + 2N(-1)^{l-1} \left(\frac{-F_N F_{2l} + F_{N+1} F_{2l-1}}{F_N} \right) \right)$$

$$= \frac{2}{5} \left(N - 2l + 1 + 2N \frac{(-1)^{l-1} F_{N-2l+1}}{F_N} \right).$$
 (by (2.4))

Combining Lemmas 3.1 and 3.2 and the equation $x_l = \sum_{i=1}^l y_i$, we obtain the exact formula for the variables x_l as follows:

$$h_{N}(0,l) = x_{l} = \sum_{i=1}^{l} \frac{2}{5} \left((N - 2i + 1) + 2N \frac{(-1)^{i-1} F_{N-2i+1}}{F_{N}} \right)$$

$$= \frac{2}{5} l(N+1) - \frac{4}{5} \sum_{i=1}^{l} i - \frac{4N}{5F_{N}} \sum_{i=1}^{l} (-1)^{i} (F_{N-1} F_{2i-1} - F_{N} F_{2i-2})$$

$$= \frac{2}{5} l(N-l) - \frac{4N}{5F_{N}} \left(F_{N-1} \sum_{i=1}^{l} (-1)^{i} F_{2i-1} - F_{N} \sum_{i=1}^{l} (-1)^{i} F_{2i-2} \right)$$

$$= \frac{2}{5} l(N-l) + \frac{4N}{5F_{N}} \left((-1)^{l-1} F_{N-1} F_{l}^{2} + (-1)^{l} F_{N} F_{l-1} F_{l} \right) \qquad (by (2.6))$$

$$= \frac{2}{5} l(N-l) + \frac{4N}{5F_{N}} F_{l} F_{N-l} = \frac{2}{5} \left(l(N-l) + 2N \frac{F_{l} F_{N-l}}{F_{N}} \right). \qquad \Box$$

From Theorem 3.1 we have $H_N^{-1} = {}^tU_N\,{}^tW_N^{-1}D_N^{-1}W_N^{-1}\,U_N$. Hence the entries of H_N^{-1} also can be calculated as follows: For $1 \le i, j \le \left\lfloor \frac{N}{2} \right\rfloor$,

$$H_N^{-1}(i,j) = \begin{cases} \frac{F_i F_j F_{N-i} F_{N-j}}{F_N F_{N-1}} + \sum_{k=1}^{i-1} \frac{F_{i-k} F_{j-k} F_{N-i-k} F_{N-j-k}}{F_{N-2k+1} F_{N-2k-1}} & \text{if } i \leq j, \\ \frac{F_i F_j F_{N-i} F_{N-j}}{F_N F_{N-1}} + \sum_{k=1}^{j-1} \frac{F_{i-k} F_{j-k} F_{N-i-k} F_{N-j-k}}{F_{N-2k+1} F_{N-2k-1}} & \text{otherwise.} \end{cases}$$

Combining the above with the equation $H_N \vec{x} = 4\vec{1}$, we have the following.

Proposition 3.1.

$$h_N(0,1) = \frac{2}{F_N} \sum_{i=0}^{N} F_i F_{N-i}.$$

Proof.

$$h_N(0,1) = 4 \sum_{j=1}^{\left\lfloor \frac{N}{2} \right\rfloor} H_N^{-1}(1,j) = 4 \sum_{j=1}^{\left\lfloor \frac{N}{2} \right\rfloor} \frac{F_1 F_j F_{N-1} F_{N-j}}{F_N F_{N-1}} = 4 \sum_{j=1}^{\left\lfloor \frac{N}{2} \right\rfloor} \frac{F_j F_{N-j}}{F_N} = \frac{2}{F_N} \sum_{j=0}^{N} F_j F_{N-j}.$$

Remark 3.1. We also obtain the counterpart of Theorem 3.1 for the case of C_N^3 [2], which convinced us that our method is always efficient for the case of general C_N^m , although their calculations get much complicated as the parameter m get larger.

4 Effective Resistance, Kirchhoff Index, and Graph Complexity

The effective resistance between a pair of vertices $\{p,q\}$ of a connected graph G, denoted by r(G;p,q), is the electrical resistance measured across p and q when the graph G represents an electrical circuit such that each edge of G is a unit resistor.

In 1959, Nash-Williams [6] proved the following surprising formula between the effective resistances and the average hitting times.

Theorem 4.1 (Nash-Williams, 1959 [6]).

$$r(G; x, y) = \frac{h(G; x, y) + h(G; y, x)}{2|E(G)|}.$$

The Kirchhoff index Kf(G) of the graph G is defined as $Kf(G) := \sum_{i < j} r(G; v_i, v_j)$.

Theorem 4.2.

$$Kf(C_N^2) = \frac{N(N-1)(5N+17)}{300} + \frac{2N^2}{25} \frac{F_{N-1}}{F_N}.$$

Proof. The following equation is derived immediately from Theorem 1.2 and Proposition 3.1.

$$\sum_{i=0}^{N} F_i F_{N-i} = \frac{1}{5} ((N-1)F_N + 2NF_{N-1}). \tag{4.1}$$

Then we have:

$$Kf(C_N^2) = \sum_{x < y} r(C_N^2; x, y) = \sum_{i=1}^{N-1} (N - i) r(C_N^2; 0, i) = \sum_{i=1}^{N-1} (N - i) \frac{h(C_N^2; 0, i) + h(C_N^2; i, 0)}{2|E(C_N^2)|}$$

$$= \sum_{i=1}^{N-1} (N - i) \frac{h(C_N^2; 0, i)}{2N} = \sum_{i=1}^{N-1} \frac{(N - i)}{5N} \left(i(N - i) + 2N \frac{F_i F_{N-i}}{F_N} \right)$$

$$= \frac{1}{5} \left(N \sum_{i=1}^{N-1} i - \sum_{i=1}^{N-1} i^2 + \frac{2N}{F_N} \sum_{i=1}^{N-1} F_i F_{N-i} \right) - \frac{1}{5N} \left(N \sum_{i=1}^{N-1} i^2 - \sum_{i=1}^{N-1} i^3 + \frac{2N}{F_N} \sum_{i=1}^{N-1} i F_i F_{N-i} \right)$$

$$= \frac{1}{5} \left(N \sum_{i=1}^{N-1} i - 2 \sum_{i=1}^{N-1} i^2 + \frac{2N}{F_N} \sum_{i=1}^{N-1} F_i F_{N-i} \right) - \frac{1}{5N} \left(- \sum_{i=1}^{N-1} i^3 + \frac{N^2}{F_N} \sum_{i=1}^{N-1} F_i F_{N-i} \right)$$

$$= \frac{1}{5} \left(N \sum_{i=1}^{N-1} i - 2 \sum_{i=1}^{N-1} i^2 + \frac{1}{N} \sum_{i=1}^{N-1} i^3 + \frac{N}{F_N} \sum_{i=1}^{N-1} F_i F_{N-i} \right)$$

$$= \frac{1}{5} \left(\frac{N^2(N-1)}{2} - \frac{N(N-1)(2N-1)}{3} + \frac{N(N-1)^2}{4} + \frac{N}{5F_N} ((N-1)F_N + 2NF_{N-1}) \right)$$
 (by (4.1))

$$\begin{split} &=\frac{1}{5}\left(\frac{N(N-1)(N+1)}{12}+\frac{N}{5F_N}(2NF_{N-1}+(N-1)F_N)\right)\\ &=\frac{1}{5}\left(\frac{N(N-1)(N+1)}{12}+\frac{2N^2F_{N-1}}{5F_N}+\frac{N(N-1)}{5}\right)\\ &=\frac{1}{5}\left(\frac{N(N-1)(5N+17)}{60}+\frac{2N^2F_{N-1}}{5F_N}\right)\\ &=\frac{N(N-1)(5N+17)}{300}+\frac{2N^2}{25}\frac{F_{N-1}}{F_N}. \end{split}$$

The graph complexity of a graph G, denoted by t(G), means the number of spanning trees in G. Then let t(G; x, y) denote the graph complexity of the graph obtained by identifying two vertices x and y of G. The following famous formula between the effective resistances and the graph complexities was proven in Kirchhoff [4].

Theorem 4.3 (G. Kirchhoff, 1847 [4]).

$$r(G; x, y) = \frac{t(G; x, y)}{t(G)}.$$

Furthermore, D. J. Kleitman and B. Golden [5] proved the following beautiful formula for the graph complexity of C_N^2 .

Theorem 4.4 (D. J. Kleitman and B. Golden, 1975 [5]).

$$t(C_N^2) = NF_N^2.$$

Combining Theorems 1.2, 4.1, 4.3, and 4.4, we have the following.

Proposition 4.1.

$$t(C_N^2; 0, l) = \frac{F_N}{5} \left(l(N - l)F_N + 2NF_l F_{N-l} \right).$$

Proof.

$$t(C_N^2; 0, l) = r(C_N^2; 0, l)t(C_N^2) = \frac{h_N(0, l)F_N^2}{2} = \frac{F_N^2}{5} \left(l(N - l) + 2N \frac{F_l F_{N-l}}{F_N} \right)$$
$$= \frac{F_N}{5} \left(l(N - l)F_N + 2N F_l F_{N-l} \right).$$

5 Asymptotic Behavior

Theorem 1.2 is useful for calculating the asymptotic behavior of HT's of simple random walks on C_N^2 . Let $\phi = \frac{1+\sqrt{5}}{2}$.

Lemma 5.1 ([3]).

$$\lim_{n \to \infty} \frac{F_n}{F_{n-1}} = \phi.$$

Theorem 5.1.

$$\lim_{N \to \infty} \frac{h_N(0, l) - \frac{2}{5}l(N - l)}{N} = \frac{4}{5\sqrt{5}} \qquad (l = 0, 1, 2, \dots).$$

Proof. By Theorem 1.2 and Lemma 5.1, we have

$$h_N(0,l) = \frac{2}{5} \left(l(N-l) + 2N \frac{F_l \cdot F_{N-l}}{F_N} \right)$$

$$= \frac{2}{5} \left(l(N-l) + 2N \frac{1}{\sqrt{5}} \left(\frac{\phi^l \cdot \phi^{N-l}}{\phi^N} + O\left(\frac{1}{N}\right) \right) \right)$$

$$= \frac{2}{5} \left(l(N-l) + \frac{2N}{\sqrt{5}} \left(1 + O\left(\frac{1}{N}\right) \right) \right).$$

Then,

$$\lim_{N \to \infty} \frac{h_N(0, l) - \frac{2}{5}l(N - l)}{N} = \lim_{N \to \infty} \frac{4}{5\sqrt{5}} \left(1 + O\left(\frac{1}{N}\right) \right) = \frac{4}{5\sqrt{5}}.$$

Theorem 5.2.

$$\lim_{\substack{N \to \infty \\ \frac{l}{N} \to x}} \frac{h_N(0, l)}{N^2} = \frac{2}{5}x(1 - x).$$

Proof.

$$\lim_{\substack{N \to \infty \\ \frac{N}{N} \to x}} \frac{h_N(0, l)}{N^2} = \lim_{\substack{N \to \infty \\ \frac{N}{N} \to x}} \frac{2}{5} \left(\frac{l}{N} \left(1 - \frac{l}{N} \right) + \frac{2}{N} \cdot \frac{1}{\sqrt{5}} \left(1 + O\left(\frac{1}{N}\right) \right) \right)$$
$$= \frac{2}{5} x (1 - x).$$

References

- N. Chair.: The Effective Resistance of the N-Cycle Graph with Four Nearest Neighbors, Journal of Statistical Physics, 154 (2014) 1177-1190.
- [2] Y. Doi, T. Miezaki, T. Nakamigawa, T. Sakuma, E. Segawa, H. Shinohara, S. Tamura, Y. Tanaka, and K. Toyota: Manuscript in preparation.
- [3] S. Jacob.: Ein new und wohlgegründt Rechenbuch, auf den Linien und Ziffern, Sampt der Welschen Practica ..., Frankfurt am Meyn, Christian Egenolff, 1571.
- [4] G. Kirchhoff.: Uber die Auflösung der Gleichungen, auf welche man bei der Untersuchung der linearen Verteilung galvanischer Ströme geführt wird, Annalen Physik und Chemie, 72 (1847) 497-508
- [5] D. J. Kleitman, B. Golden.: Counting Trees in a Certain Class of Graphs, The American Mathematical Monthly, 82 No.1 (1975) 40-44
- [6] C. S. J. A. Nash-Williams.: Random walk and electric currents in networks. Proceedings of the Cambridge Philosophical Society. Mathematical and Physical Sciences, 55 (1959) 181-194.
- [7] T. Koshy.: Fibonacci and Lucas Numbers with Applications, Volume 1, Pure and Applied Mathematics: A Wiley Series of Texts, Monographs and Tracts, 2018.
- [8] T. Koshy.: Fibonacci and Lucas Numbers with Applications, Volume 2, Pure and Applied Mathematics: A Wiley Series of Texts, Monographs and Tracts, 2019.
- [9] F. Y. Wu.: Theory of resistor networks: the two-point resistance, *Journal of Physics A: Mathematical and General*, **37**(26) (2004) 6653-6673.