

Maximal 3-wise Intersecting Families

József Balogh^{*†} Ce Chen^{*} Kevin Hendrey[‡] Ben Lund[‡] Haoran Luo^{*§}
Casey Tompkins[¶] Tuan Tran^{||}

Abstract

A family \mathcal{F} on ground set $[n] := \{1, 2, \dots, n\}$ is *maximal k -wise intersecting* if every collection of at most k sets in \mathcal{F} has non-empty intersection, and no other set can be added to \mathcal{F} while maintaining this property. In 1974, Erdős and Kleitman asked for the minimum size of a maximal k -wise intersecting family. We answer their question for $k = 3$ and sufficiently large n . We show that the unique minimum family is obtained by partitioning the ground set $[n]$ into two sets A and B with almost equal sizes and taking the family consisting of all the proper supersets of A and of B .

1 Introduction

A central topic in extremal set theory is intersection properties of families of finite sets. This topic began with classical results of Erdős, Ko and Rado [14] determining the maximum size of an intersecting family of subsets of an n -element set, for both arbitrary subsets and for subsets of a given size.

A family \mathcal{F} of sets is *maximal* (or *saturated*) with respect to some property if it satisfies the property, but no family properly containing \mathcal{F} satisfies the property. The problem of finding the smallest object which is maximal with respect to some property has an extensive literature. In the setting of graphs the topic was initiated by Erdős, Hajnal and Moon [12], who determined the minimum number of edges in a maximal n -vertex graph not containing a clique of size k (see [16] for an extensive survey on graph saturation). For set systems consisting of sets of a fixed size r , Erdős and Lovász [15] suggested the problem of finding a maximal intersecting family of minimum size. For this problem, the best known lower bound of $3r$ (when there is a projective plane of order $r - 1$) was proved by Dow, Drake, Füredi and Larson [7], and the best known upper bound of $r^2/2 + O(r)$ is due to Boros, Füredi and Kahn [5]. Kahn [22] conjectured that $O(r)$ is an upper bound. Other properties for which saturation results have been obtained for set systems include the

^{*}Department of Mathematics, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA. E-mail: {jobal, cechen4, haoranl8}@illinois.edu.

[†]Research is partially supported by NSF Grant DMS-1764123, NSF RTG grant DMS 1937241, Arnold O. Beckman Research Award (UIUC Campus Research Board RB 22000), and the Langan Scholar Fund (UIUC).

[‡]Discrete Mathematics Group, Institute for Basic Science (IBS), Daejeon, Republic of Korea. E-mail: {kevinhendrey, benlund}@ibs.re.kr. This work was supported by the Institute for Basic Science (IBS-R029-C1).

[§]Research is partially supported by UIUC Campus Research Board RB 22000.

[¶]Alfréd Rényi Institute of Mathematics, Hungarian Academy of Sciences. E-mail: casey@renyi.hu. This work was supported by NKFIH grant K135800.

^{||}School of Mathematical Sciences, University of Science and Technology of China, China. E-mail: trantuan@ustc.edu.cn. This work was supported by the Institute for Basic Science (IBS-R029-Y1), and the Outstanding Young Talents Program (Overseas) of the National Natural Science Foundation of China.

k -Sperner [18], equal disjoint union free [8], VC-dimension at most k [17] and k -matching free [6] properties.

In this paper we study a natural saturation problem concerning an intersection property. To state the problem we need to introduce some notation. For a positive integer n , denote by $[n]$ the set $\{1, 2, \dots, n\}$. Given a set S , we write 2^S for the power set of S , which is the family of all subsets of S . For $\mathcal{F} \subseteq 2^{[n]}$, let $\bar{\mathcal{F}} := \{F^c : F \in \mathcal{F}\}$ where F^c denotes $[n] \setminus F$. We use the notation $\mathcal{F} \Delta \mathcal{G}$ for the symmetric difference $(\mathcal{F} \setminus \mathcal{G}) \cup (\mathcal{G} \setminus \mathcal{F})$ of two set families \mathcal{F} and \mathcal{G} .

We say that a family $\mathcal{F} \subseteq 2^{[n]}$ is *maximal k -wise intersecting* if the intersection of every collection of at most k sets in \mathcal{F} is non-empty, and no set from $2^{[n]} \setminus \mathcal{F}$ can be added to \mathcal{F} while preserving this property. In the case $k = 2$, it is well-known that every maximal 2-wise intersecting family has the same size, namely 2^{n-1} . For $k \geq 3$, an old question of Erdős and Kleitman [13] from 1974 asks for the size of the smallest maximal k -wise intersecting family. We answer this question for the case $k = 3$ and n sufficiently large.

We now present our construction, which we call *a balanced pair of linked cubes*. A balanced pair of linked cubes is a set family of the form $\{A : S \subsetneq A \subseteq [n]\} \cup \{B : S^c \subsetneq B \subseteq [n]\}$ for some $S \subseteq [n]$ with $|S| = \lfloor n/2 \rfloor$. It is not hard to check that a balanced pair of linked cubes is a maximal 3-wise intersecting family of size $2^{\lfloor n/2 \rfloor} + 2^{\lceil n/2 \rceil} - 3$.

Our main result is that for sufficiently large n , the smallest maximal 3-wise intersecting families are balanced pairs of linked cubes.

Theorem 1.1. *If \mathcal{F} is a maximal 3-wise intersecting family of minimum size on ground set $[n]$, where n is sufficiently large, then \mathcal{F} is a balanced pair of linked cubes.*

Remark. The even case of Theorem 1.1 was first proved by the third, fourth, sixth and seventh authors [20]. In [4], the first, second and fifth authors used the method of [20] together with some new ideas to establish the odd case. The present paper provides a unified treatment of both cases.

A key ingredient in the proof of Theorem 1.1 is the following stability result, which shows that a maximal 3-wise intersecting family of size at most $(1 + o(1))(2^{\lfloor n/2 \rfloor} + 2^{\lceil n/2 \rceil})$ must be close in structure to a balanced pair of linked cubes.

Theorem 1.2. *For every $\varepsilon > 0$ there exists $\delta > 0$ such that, if \mathcal{F} is a maximal 3-wise intersecting family on $[n]$ of size $|\mathcal{F}| \leq (1 + \delta)(2^{\lfloor n/2 \rfloor} + 2^{\lceil n/2 \rceil})$, where n is sufficiently large, then there exists a balanced pair of linked cubes \mathcal{F}_0 such that $|\mathcal{F} \Delta \mathcal{F}_0| \leq \varepsilon 2^{\lfloor n/2 \rfloor}$.*

Organization and notation: The rest of the paper is organized as follows. In Section 2 we prove Theorem 1.2, in Section 3 we deduce Theorem 1.1 from Theorem 1.2 using a perturbation argument, and in Section 4 we discuss maximal k -wise intersecting families when $k \geq 4$.

We use standard asymptotic notation throughout. For functions $f = f(n)$ and $g = g(n)$ we write $f = o(g)$, $f \ll g$, or $g \gg f$ to mean $f/g \rightarrow 0$ as $n \rightarrow \infty$. All asymptotics are as $n \rightarrow \infty$.

2 Proof of Theorem 1.2

2.1 Proof Overview

In the proof of Theorem 1.2, we find it more convenient to work with $\bar{\mathcal{F}}$.

Definition. A *balanced pair of cubes on $[n]$* is a set system of the form $\{A : A \subsetneq S\} \cup \{B : B \subsetneq S^c\}$, where $S \subseteq [n]$ with $|S| = \lfloor \frac{n}{2} \rfloor$. More generally, a *balanced series of k cubes on $[n]$* , denoted by $\mathcal{F}_{n,k}$, is a set system of the form $\bigcup_{i=1}^k \{A : A \subsetneq S_i\}$, where S_1, \dots, S_k is a partition of $[n]$ with $\lfloor \frac{n}{k} \rfloor \leq |S_i| \leq \lceil \frac{n}{k} \rceil$ for each $i \in [k]$.

We also need the following key concepts, whose relevance will be revealed in Lemma 2.1 below.

Definition. A set system $\mathcal{H} \subseteq 2^{[n]}$ is a $(1 - \varepsilon)$ - k -generator for $[n]$ if all but at most $\varepsilon 2^n$ subsets of $[n]$ can be expressed as a union of at most k disjoint sets of \mathcal{H} .

Definition. Given a set system $\mathcal{H} \subseteq 2^{[n]}$, denote by $\text{dp}(\mathcal{H})$ the number of disjoint pairs in \mathcal{H} , i.e., $\text{dp}(\mathcal{H}) = |\{\{A, B\} : A, B \in \mathcal{H}, A \cap B = \emptyset\}|$.

The following lemma is central to the proof of Theorem 1.2.

Lemma 2.1. *If \mathcal{F} is a maximal 3-wise intersecting family on $[n]$, then*

$$\text{dp}(\bar{\mathcal{F}}) \geq 2^n - |\mathcal{F}|.$$

Moreover, if $|\mathcal{F}| \leq \varepsilon 2^n$, then $\bar{\mathcal{F}}$ is a $(1 - \varepsilon)$ -2-generator for $[n]$.

Proof. Let $\mathcal{F}^c := 2^{[n]} \setminus \mathcal{F}$, then $|\mathcal{F}^c| = 2^n - |\mathcal{F}|$. For every $A \in \mathcal{F}^c$, there exist $B, C \in \mathcal{F}$ such that $B \cap C \subseteq A^c$, since \mathcal{F} is a maximal 3-wise intersecting family. Notice that \mathcal{F} is upward closed, i.e., if $F \in \mathcal{F}$ and $F \subseteq F' \subseteq [n]$, then $F' \in \mathcal{F}$. Thus, we may select these sets $B, C \in \mathcal{F}$ so that $A^c = B \cap C$ and $B \cup C = [n]$, which means $B^c \cap C^c = \emptyset$. This (not necessarily unique) choice of B and C defines an injective map from \mathcal{F}^c into disjoint pairs of sets in $\bar{\mathcal{F}}$ (where \emptyset is mapped to $\{\emptyset, \emptyset\}$). Therefore,

$$\text{dp}(\bar{\mathcal{F}}) \geq |\mathcal{F}^c| = 2^n - |\mathcal{F}|.$$

Now assume that $|\mathcal{F}| \leq \varepsilon 2^n$. Since $|\mathcal{F}^c| = 2^n - |\mathcal{F}| \geq (1 - \varepsilon)2^n$ and every set in \mathcal{F}^c can be expressed as a union of at most two disjoint sets of $\bar{\mathcal{F}}$, $\bar{\mathcal{F}}$ is a $(1 - \varepsilon)$ -2-generator for $[n]$. \square

The following result together with Lemma 2.1 immediately implies Theorem 1.2. We will prove it in Section 2.3.

Theorem 2.2. *For every $\varepsilon' > 0$, there exists $\delta = \delta(\varepsilon') > 0$ such that for sufficiently large n , if $\mathcal{H} \subseteq 2^{[n]}$ is a $(1 - \delta)$ -2-generator for $[n]$ with $|\mathcal{H}| \leq (1 + \delta)(2^{\lfloor n/2 \rfloor} + 2^{\lceil n/2 \rceil})$, then there exists a balanced pair of cubes $\mathcal{F}_{n,2}$ such that $|\mathcal{H} \Delta \mathcal{F}_{n,2}| \leq \varepsilon' 2^{\lfloor n/2 \rfloor}$.*

Remark. Theorem 2.2 for n even was first proved by Ellis and Sudakov [9, Proposition 9]. Our proof of Theorem 2.2 uses their method.

Proof of Theorem 1.2 assuming Theorem 2.2. Let $\varepsilon > 0$ and n be a sufficiently large integer. Let $\delta = \delta(\varepsilon) > 0$ be obtained from Theorem 2.2. Let \mathcal{F} be a maximal 3-wise intersecting family on $[n]$ of size $|\mathcal{F}| \leq (1 + \delta)(2^{\lfloor n/2 \rfloor} + 2^{\lceil n/2 \rceil})$, then $|\mathcal{F}| \leq \delta 2^n$, when n is sufficiently large. Since $|\bar{\mathcal{F}}| = |\mathcal{F}| \leq \delta 2^n$, $\bar{\mathcal{F}}$ is a $(1 - \delta)$ -2-generator for $[n]$ by Lemma 2.1. Applying Theorem 2.2 to $\mathcal{H} := \bar{\mathcal{F}}$, we conclude that there exists a balanced pair of cubes $\mathcal{F}_{n,2}$ with $|\bar{\mathcal{F}} \Delta \mathcal{F}_{n,2}| \leq \varepsilon 2^{\lfloor n/2 \rfloor}$. Taking the complements of the sets in $\bar{\mathcal{F}}$ and $\mathcal{F}_{n,2}$ yields the desired result. \square

Lemma 2.1 also leads to another classic question posed by Erdős [11]: How many disjoint pairs of sets can there be in a set system of given size? A lower bound comes from $\mathcal{F}_{n,k}$, a balanced series of k cubes, i.e. $\mathcal{F}_{n,k} = \cup_{i=1}^k \{A : A \subseteq S_i\}$, where S_1, \dots, S_k is a partition of $[n]$ with $\lfloor \frac{n}{k} \rfloor \leq |S_i| \leq \lceil \frac{n}{k} \rceil$ for each $i \in [k]$. With the additional assumption that k divides n , it is easy to see that $|\mathcal{F}_{n,k}| = k 2^{n/k} - 2k + 1$ and $\text{dp}(\mathcal{F}_{n,k}) > (1 - 1/k) \binom{|\mathcal{F}_{n,k}|}{2}$. Solving a conjecture of Daykin and Erdős [19], Alon and Frankl [1] proved that $\mathcal{F}_{n,k}$ has asymptotically the maximum number of disjoint pairs, given its size.

Theorem 2.3 (Alon–Frankl). *For every positive integer k , there exists $\beta = \beta(k) > 0$ such that, if \mathcal{H} is a set system on $[n]$ of size $m := |\mathcal{H}| \geq 2^{(1/(k+1)+\varepsilon)n}$, where $\varepsilon > 0$, then*

$$\text{dp}(\mathcal{H}) \leq \left(1 - \frac{1}{k}\right) \binom{m}{2} + O\left(m^{2-\beta\varepsilon^2}\right).$$

We strengthen Theorem 2.3 by proving a stability result, showing families with close to $(1 - \frac{1}{k}) \binom{m}{2}$ disjoint pairs must resemble a balanced series of k cubes.

Theorem 2.4. *For every $\varepsilon > 0$ and integer $k \geq 2$, there exists an $\eta > 0$ such that for every sufficiently large n , if a set system \mathcal{H} on $[n]$ of size $m := |\mathcal{H}| \geq (1-\eta)k2^{n/k}$ has at least $(1 - \frac{1}{k} - \eta)\frac{m^2}{2}$ disjoint pairs, then k divides n , and there exists a balanced series of k cubes $\mathcal{F}_{n,k}$ with $|\mathcal{H} \Delta \mathcal{F}_{n,k}| \leq \varepsilon 2^{\lfloor n/k \rfloor}$.*

The proof of Theorem 2.4, given in Section 2.2, borrows some ideas of Alon and Frankl [1], and Alon, Das, Glebov and Sudakov [2]. The even case of Theorem 2.2 is proved using Theorem 2.4.

Proof of Theorem 2.2 for even n assuming Theorem 2.4. Let η be as given by Theorem 2.4 for $k = 2$ and $\varepsilon > 0$, and let $\delta = \eta^2$. Let $\mathcal{H}' \supseteq \mathcal{H}$ such that $|\mathcal{H}'| = \lfloor (1 + \delta)2^{(n+2)/2} \rfloor$, and every element of $\mathcal{H}' \setminus \mathcal{H}$ has size greater than $n/2$. This is possible because n is large and the number of sets of size greater than $n/2$ is nearly 2^{n-1} . Since $\mathcal{H}' \supseteq \mathcal{H}$ and \mathcal{H} is a $(1 - \delta)$ -2-generator for $[n]$, we have

$$|\mathcal{H}'| + \text{dp}(\mathcal{H}') \geq |\mathcal{H}| + \text{dp}(\mathcal{H}) \geq (1 - \delta)2^n.$$

Hence $\text{dp}(\mathcal{H}') \geq (1 - \delta)2^n - (1 + \delta)2^{(n+2)/2} \geq (1 - 2\delta)2^n$ for n sufficiently large. On the other hand,

$$\left(\frac{1}{2} - \eta\right) \frac{|\mathcal{H}'|^2}{2} \leq \left(\frac{1}{2} - \eta\right) (1 + \eta)^2 2^{n+1} = (1 - 3\eta^2 - 2\eta^3)2^n \leq (1 - 3\delta)2^n.$$

It follows that $\text{dp}(\mathcal{H}') \geq (1/2 - \eta)|\mathcal{H}'|^2/2$, and the hypotheses of Theorem 2.4 are satisfied for \mathcal{H}' . Theorem 2.4 gives a balanced pair of cubes $\mathcal{F}_{n,2}$ with $|\mathcal{H}' \Delta \mathcal{F}_{n,2}| \leq \varepsilon 2^{n/2}$. By the second property of \mathcal{H}' , $\mathcal{H}' \setminus \mathcal{H}$ is disjoint from $\mathcal{F}_{n,2}$. Hence, $|\mathcal{H} \Delta \mathcal{F}_{n,2}| \leq |\mathcal{H}' \Delta \mathcal{F}_{n,2}| \leq \varepsilon 2^{n/2}$. \square

Before starting the proofs of Theorems 2.2 and 2.4, we introduce the so-called disjointness graph, which is the main object that we analyze in Sections 2.2 and 2.3.

Definition. For a set system \mathcal{H} , the *disjointness graph* $G_{\mathcal{H}}$ is the graph with vertex set \mathcal{H} and edge set $\{\{A, B\} \subseteq \mathcal{H} : A \cap B = \emptyset\}$. For two (not necessarily disjoint) set families $\mathcal{H}_1, \mathcal{H}_2$, the *disjointness bipartite graph* $G_{\mathcal{H}_1, \mathcal{H}_2}$ is the bipartite graph with classes $(\mathcal{H}_1, \mathcal{H}_2)$, where there is an edge between $A \in \mathcal{H}_1$ and $B \in \mathcal{H}_2$ if and only if $A \cap B = \emptyset$.

2.2 Proof of Theorem 2.4

The heart of the proof of Theorem 2.4 is the following lemma, which shows that the disjointness graph of a large family has only few cliques of size $k + 1$. This lemma essentially appears in [1]. For completeness, we include its proof here.

Lemma 2.5. *For every $\varepsilon > 0$, $\gamma > 0$ and integer $k \geq 2$, and sufficiently large integer n , if \mathcal{H} is a set system on $[n]$ of size $m := |\mathcal{H}| \geq 2^{(1/(k+1)+\varepsilon)n}$, then $G_{\mathcal{H}}$ contains at most $\gamma \binom{m}{k+1}$ copies of K_{k+1} .*

We denote by $K_r(t)$ the complete r -partite graph with parts of size t . The following proposition is standard and follows from a result of Erdős [10] by a simple averaging argument (see for instance [2, Proposition 3.2]).

Proposition 2.6. *For integers $r \geq 2$, $t \geq 1$ and any real $\gamma > 0$, there exists $\delta_{2.6} = \delta_{2.6}(r, t, \gamma) > 0$, such that if m is sufficiently large and G is a graph on m vertices with at least $\gamma \binom{m}{r}$ copies of K_r , then G contains at least $\delta_{2.6} \binom{m}{rt}$ copies of $K_r(t)$.*

We will use a probabilistic argument to derive Lemma 2.5 from Proposition 2.6.

Proof of Lemma 2.5. Let $t = \lceil \frac{2}{\varepsilon} \rceil + k$. Select t sets $A_1, \dots, A_t \in \mathcal{H}$ independently uniformly at random, with repetitions allowed. The probability that $|A_1 \cup A_2 \cup \dots \cup A_t| \leq \frac{n}{k+1}$ is bounded above by

$$\sum_{\substack{S \subseteq [n] \\ |S| = \lfloor n/(k+1) \rfloor}} \Pr [A_i \subseteq S, i = 1, \dots, t] \leq 2^n \left(2^{n/(k+1)} / m \right)^t \leq m^{-k},$$

since $m \geq 2^{(1/(k+1)+\varepsilon)n}$, where 2^n estimates the number of choices of S , and S has $2^{\lfloor n/(k+1) \rfloor}$ many subsets, each could be chosen as A_i , if they are in \mathcal{H} .

Let $\mathbf{A} = (A_1, A_2, \dots, A_{(k+1)t})$ be a random sample (chosen independently uniformly at random, allowing repetition) of $(k+1)t$ vertices of $G_{\mathcal{H}}$. It follows from the discussion above that the probability that $|A_{(i-1)t+1} \cup A_{(i-1)t+2} \cup \dots \cup A_{it}| \leq \frac{n}{k+1}$ for some $i \in [k+1]$ is at most $(k+1)m^{-k}$. On the other hand, if our random sample \mathbf{A} gives a copy of $K_{(k+1)}(t)$ with $\{A_{(i-1)t+1}, \dots, A_{it}\}$ being the i -th vertex class for every $i \in [k+1]$, then $A_1 \cup \dots \cup A_t, \dots, A_{kt+1} \cup \dots \cup A_{(k+1)t}$ are $k+1$ disjoint subsets of $[n]$. Hence, in this case, we must have $|A_{(i-1)t+1} \cup A_{(i-1)t+2} \cup \dots \cup A_{it}| \leq \frac{n}{k+1}$ for some $i \in [k+1]$. Moreover, the probability that our random sample gives such a copy of $K_{k+1}(t)$ is precisely

$$m^{-(k+1)t} \cdot (k+1)!(t!)^{k+1} \cdot \# \text{ copies of } K_{k+1}(t) \text{ in } G_{\mathcal{H}}.$$

Therefore, the number of copies of $K_{k+1}(t)$ in $G_{\mathcal{H}}$ is at most

$$\frac{m^{(k+1)t-k}}{k!(t!)^{k+1}} \leq \delta_{2.6}(k+1, t, \gamma) \binom{m}{(k+1)t}$$

for m sufficiently large. Hence, by Proposition 2.6, $G_{\mathcal{H}}$ has at most $\gamma \binom{m}{k+1}$ copies of K_{k+1} . This completes the proof of Lemma 2.5. \square

We will make use of the following theorem of Balogh, Bushaw, Collares, Liu, Morris and Sharifzadeh [3, Theorem 1.2].

Theorem 2.7. *For every $m, k, t \in \mathbb{N}$, the following holds. Suppose G is graph on m vertices with at most $\frac{m^{k-1}}{e^{2k} \cdot k!} \left(e(G) + t - \left(1 - \frac{1}{k}\right) \frac{m^2}{2} \right)$ copies of K_{k+1} . Then there is a partition of the vertex set of G as $V(G) = V_1 \cup V_2 \cup \dots \cup V_k$ with $\sum_{i=1}^k e(V_i) \leq t$.*

We also use the following well-known estimate on the size of a set system in terms of the binary entropy function.

Lemma 2.8. *Let \mathcal{F} be a set system on $[n]$. Denote by p_i the fraction of sets in \mathcal{F} that contain i . Then*

$$|\mathcal{F}| \leq 2^{\sum_{i=1}^n h(p_i)},$$

where $h(p) = -p \log_2 p - (1-p) \log_2 (1-p)$ is the binary entropy function.

We are now ready to prove Theorem 2.4. The proof closely follows the approach from [2].

Proof of Theorem 2.4. Let \mathcal{H} be a family of subsets of $[n]$ such that $m := |\mathcal{H}| \geq (1 - o(1))k2^{n/k}$ and $\text{dp}(\mathcal{H}) \geq (1 - 1/k - o(1))\frac{m^2}{2}$. By Lemma 2.5, $G_{\mathcal{H}}$ contains at most $o(m^{k+1})$ copies of K_{k+1} . Thus, applying Theorem 2.7 with $t = o(m^2)$, we conclude that $G_{\mathcal{H}}$ has a k -partite subgraph with $(1 - 1/k - o(1))\frac{m^2}{2}$ edges. In order to contain this many edges, it is clear that the vertex classes must all have size $(1 - o(1))m/k > (1 - o(1))2^{n/k}$. Furthermore, again by the edge density, all but at most $o(m)$ vertices must have at least $(1 - 1/k - o(1))m$ neighbors. Thus, we may take a k -partite subgraph G of $G_{\mathcal{H}}$ consisting of parts of size at least $(1 - o(1))2^{n/k}$ such that each vertex has at least $(1 - 1/k - o(1))k2^{n/k}$ neighbors, and thus at least $(1 - o(1))2^{n/k}$ neighbors in every other class.

Let $\varepsilon > 0$ be a sufficiently small constant with respect to k . Let $\mathcal{H}_1, \mathcal{H}_2, \dots, \mathcal{H}_k$ be the sets from \mathcal{H} corresponding to the k color classes of G , respectively. By the above discussion,

$$|\mathcal{H}_i| \geq (1 - \varepsilon)2^{n/k} \quad \text{for all } i \in [k]. \quad (1)$$

We will construct a balanced series of k cubes containing $\bigcup_{i=1}^k \mathcal{H}_i$. To do this we begin with a partition of the ground set $[n]$ into $2k + 1$ disjoint sets: $A_1, B_1, \dots, A_k, B_k, R$. For $i \in [k]$, let A_i and B_i be the set of those elements of $[n] \setminus \bigcup_{j=1}^{i-1} (A_j \cup B_j)$ occurring in more than $\varepsilon|\mathcal{H}_i|$ members of \mathcal{H}_i and between 1 and $\varepsilon|\mathcal{H}_i|$ members of \mathcal{H}_i , respectively. Let R be the set of remaining elements of $[n]$.

By the definition of the partition, if for some $i \in [k]$ we have $x \in B_i$, then there is a set $F \in \mathcal{H}_i$ such that $x \in F$. It follows from the density properties of G discussed above that for $j \neq i$, x is not contained in at least $(1 - \varepsilon)|\mathcal{H}_j|$ members of \mathcal{H}_j . If $x \in A_i$, then for $j \neq i$, x does not appear in any member of \mathcal{H}_j , since each member of \mathcal{H}_j is disjoint from at least $(1 - \varepsilon)|\mathcal{H}_i|$ members of \mathcal{H}_i . It follows that no set $F \in \mathcal{H}$ has vertices from $\bigcup_{j=1}^{i-1} A_j$. Furthermore, each element $x \in \bigcup_{j=1}^i B_j$ is contained in at most $\varepsilon|\mathcal{H}_i|$ sets in \mathcal{H}_i . Using Lemma 2.8 we have

$$|\mathcal{H}_i| \leq 2^{|A_i| + h(\varepsilon)(\sum_{j < i} |B_j|)}.$$

We thus get

$$\begin{aligned} 2^{n-\frac{1}{2}} < (1 - \varepsilon)^k 2^n &\leq \prod_{i=1}^k |\mathcal{H}_i| \leq 2^{\sum_{i \in [k]} (|A_i| + h(\varepsilon)(k-i+1)|B_i|)} \\ &= 2^{n-|R| - \sum_{i \in [k]} (1-h(\varepsilon)(k-i+1))|B_i|} \leq 2^{n-|R| - \frac{3}{4} \sum_{i \in [k]} |B_i|}, \end{aligned}$$

where in the first equality we used the fact that $n = |R| + \sum_{i \in [k]} (|A_i| + |B_i|)$, and the last inequality holds since $1 - h(\varepsilon)k \geq 3/4$ for ε sufficiently small with respect to k . This implies that $R = B_1 = \dots = B_k = \emptyset$. As $\mathcal{H}_i \subseteq 2^{A_i \cup B_i}$, by the definition of A_i and B_i , we have that $|\mathcal{H}_i| \leq 2^{|A_i| + |B_i|} = 2^{|A_i|}$. Together with the lower bound $|\mathcal{H}_i| \geq (1 - \varepsilon)2^{n/k}$ from (1), we must have $|A_i| > (n - 1)/k$, and so $|A_i| \geq n/k$. Since $\sum_{i \in [k]} |A_i| = n$, we in fact have $|A_i| = n/k$ for all $i \in [k]$. It follows that the family $\bigcup_{i=1}^k \mathcal{H}_i$ is contained in the balanced series of k cubes A_1, \dots, A_k . Since \mathcal{H} contains at most $\varepsilon|\mathcal{H}|$ members not in $\bigcup_{i=1}^k \mathcal{H}_i$, the proof is complete. \square

2.3 Proof of Theorem 2.2

Before stating the proof of Theorem 2.2, we need some preparation.

Definition. Given set systems $\mathcal{H}_1, \mathcal{H}_2$ and a bipartite subgraph $B \subseteq G_{\mathcal{H}_1}$ with bipartition $(\mathcal{X}, \mathcal{Y})$, we say B (and sometimes we say $E(B)$) *generates* \mathcal{H}_2 if every $H \in \mathcal{H}_2$ can be expressed as a disjoint union of some $X \in \mathcal{X}$ and $Y \in \mathcal{Y}$, i.e., every $H \in \mathcal{H}_2$ corresponds to an edge of B .

Definition. For a set system \mathcal{H} and $i \in [n]$, let $\mathcal{H}_i^- := \{H \in \mathcal{H} : i \notin H\}$ be the subfamily of \mathcal{H} , consisting of sets not containing i and $\mathcal{H}_i^+ := \{H \setminus \{i\} : i \in H \in \mathcal{H}\}$. Note that $|\mathcal{H}_i^+| + |\mathcal{H}_i^-| = |\mathcal{H}|$.

We will use the following result by Ellis and Sudakov [9].

Lemma 2.9 ([9, Proposition 18]). *Let $c > 0$. Then, for every set family $\mathcal{H} \subseteq 2^{[n]}$ with $|\mathcal{H}| \geq c2^{n/2}$, the disjointness graph $G_{\mathcal{H}}$ can be made bipartite by deleting at most $o(|\mathcal{H}|^2)$ edges.*

We are now ready to prove Theorem 2.2.

Proof of Theorem 2.2. We have already proved the even case of Theorem 2.2, hence from now on we assume that $n = 2\ell + 1$ is a sufficiently large odd integer. Let $\varepsilon' \gg \varepsilon \gg \delta > 0$ be sufficiently small. Suppose that $\mathcal{H} \subseteq 2^{[n]}$ is a $(1 - \delta)$ -2-generator for $[n]$ with $|\mathcal{H}| \leq (1 + \delta)(2^{\lfloor n/2 \rfloor} + 2^{\lceil n/2 \rceil}) = 3(1 + \delta)2^\ell$. Then, the number of ways to choose at most two disjoint sets (whose unions are all different) from \mathcal{H} is at least $(1 - \delta)2^n$, by the definition of a $(1 - \delta)$ -2-generator. Hence, $|\mathcal{H}|^2 \geq (1 - \delta)2^n$, implying

$$|\mathcal{H}| \geq \sqrt{1 - \delta} \cdot 2^{n/2}.$$

Moreover, let G_0 be the disjointness graph of \mathcal{H} , then

$$1 + |\mathcal{H}| + e(G_0) \geq (1 - \delta)2^n,$$

which implies that

$$e(G_0) \geq (1 - \delta)2^{2\ell+1} - 3(1 + \delta)2^\ell - 1.$$

We conclude that G_0 has edge-density

$$\frac{e(G_0)}{\binom{|\mathcal{H}|}{2}} \geq \frac{(1 - \delta)2^{2\ell+1} - 3(1 + \delta)2^\ell - 1}{9(1 + \delta)^2 2^{2\ell-1}} \geq \frac{4 - 5\delta}{9(1 + \delta)^2},$$

where the last inequality comes from $3(1 + \delta)2^\ell + 1 \leq \delta 2^{2\ell-1}$. Applying Lemma 2.9 to \mathcal{H} with $c = \sqrt{1 - \delta}$, we conclude that one can delete at most

$$o(|\mathcal{H}|^2) = o(2^{2\ell+1})$$

edges from G_0 and obtain a bipartite graph $G = (\mathcal{X}, \mathcal{Y})$ with $\mathcal{X} \cup \mathcal{Y} = \mathcal{H}$. Note that G generates all but at most

$$\delta 2^{2\ell+1} + 1 + |\mathcal{H}| + o(2^{2\ell+1})$$

subsets of $[n]$. Since $|\mathcal{H}| \leq 3(1 + \delta)2^\ell = o(2^{2\ell+1})$ and $\delta \ll \varepsilon$, G generates all but at most $\varepsilon 2^{2\ell+1}$ subsets of $[n]$. In particular,

$$e(G) \geq (1 - \varepsilon)2^{2\ell+1}. \tag{2}$$

Let $\alpha := |\mathcal{X}|/2^\ell$ and $\beta := |\mathcal{Y}|/2^\ell$. Since $|\mathcal{X}| + |\mathcal{Y}| = |\mathcal{H}| \leq 3(1 + \delta)2^\ell$, we have

$$\alpha + \beta \leq 3(1 + \delta). \tag{3}$$

Therefore, we have

$$\alpha\beta \leq \frac{9}{4}(1 + \delta)^2. \tag{4}$$

Since $\alpha\beta 2^{2\ell} = |\mathcal{X}||\mathcal{Y}| \geq e(G) \geq (1-\varepsilon)2^{2\ell+1}$, we also have $\alpha\beta \geq 2-2\varepsilon$. Combining with (3), we get

$$1-3\varepsilon < \alpha, \beta < 2+3\varepsilon, \quad (5)$$

where we use $\varepsilon \gg \delta$.

Let

$$\mathcal{X}_{1/3} := \{i \in [n] : |\mathcal{X}_i^+| \geq |\mathcal{X}|/3\} \quad \text{and} \quad \mathcal{Y}_{1/3} := \{i \in [n] : |\mathcal{Y}_i^+| \geq |\mathcal{Y}|/3\}. \quad (6)$$

Letting $A := \mathcal{X}_{1/3}$ and $B := \mathcal{Y}_{1/3}$, we will show that A, B forms an equipartition of $[n]$, and that \mathcal{X} and \mathcal{Y} are not too far from 2^A and 2^B , respectively, thereby proving Theorem 2.2. We will achieve this goal step by step via the following lemma and a series of claims.

Lemma 2.10. *It cannot happen that all the following equations hold at the same time:*

$$|\mathcal{X}|, |\mathcal{Y}| = (3/2 - o(1))2^\ell, \quad |\mathcal{X}_n^+|, |\mathcal{Y}_n^+| = (1 - o(1))2^{\ell-1} \quad \text{and} \quad |\mathcal{X}_n^-|, |\mathcal{Y}_n^-| = (1 - o(1))2^\ell. \quad (7)$$

Proof. Suppose for a contradiction that (7) holds. Since all but at most $\varepsilon 2^n = 2\varepsilon \cdot 2^{n-1}$ sets in $[n]$ can be expressed as a union of at most two disjoint sets in \mathcal{H} , we have $\mathcal{H}_n^- = \mathcal{X}_n^- \cup \mathcal{Y}_n^-$ is a $(1-2\varepsilon)$ -2-generator for $[n-1]$. By the assumption that (7) holds, $|\mathcal{H}_n^-| = (2 - o(1))2^\ell = (2 - o(1))2^{(n-1)/2}$. As $|\mathcal{F}_{n-1,2}| = 2 \cdot 2^{(n-1)/2} - 1$, we can apply the even case of Theorem 2.2 to \mathcal{H}_n^- and conclude that there exists an equipartition $S_1 \cup S_2 = [n-1]$ such that each of $\mathcal{X}_n^-, \mathcal{Y}_n^-$ contains at least $(1 - o(1))2^\ell$ sets in $2^{S_1}, 2^{S_2}$. Define $\mathcal{U} := \{F \in \mathcal{X} : F \cap S_2 = \emptyset\}, \mathcal{V} := \{F \in \mathcal{Y} : F \cap S_1 = \emptyset\}$. Note that $\mathcal{U}_n^- = \mathcal{X}_n^- \cap 2^{S_1}$ and $\mathcal{V}_n^- = \mathcal{Y}_n^- \cap 2^{S_2}$. We have $|\mathcal{U}_n^-|, |\mathcal{V}_n^-| = (1 - o(1))2^\ell$, implying

$$|\mathcal{X}_n^- \setminus \mathcal{U}_n^-|, |\mathcal{Y}_n^- \setminus \mathcal{V}_n^-| = o(2^\ell). \quad (8)$$

Now we prove that

$$|\mathcal{X}_n^+ \setminus \mathcal{U}_n^+|, |\mathcal{Y}_n^+ \setminus \mathcal{V}_n^+| = o(2^\ell).$$

In fact, for every $X \in \mathcal{X}_n^+ \setminus \mathcal{U}_n^+$, we have $X \cap S_2 \neq \emptyset$ by the definition of \mathcal{U} , thus $X \cup \{n\}$ is disjoint from at most $2^{\ell-1}$ subsets of S_2 . Since $|\mathcal{Y}_n^- \setminus 2^{S_2}| = |\mathcal{Y}_n^- \setminus \mathcal{V}_n^-| = o(2^\ell)$, the set $X \cup \{n\}$ is disjoint from at most $2^{\ell-1} + o(2^\ell)$ sets in \mathcal{Y} . Similarly, for every $Y \in \mathcal{Y}_n^+ \setminus \mathcal{V}_n^+$, the set $Y \cup \{n\}$ is disjoint from at most $2^{\ell-1} + o(2^\ell)$ sets in \mathcal{X} . Let e_n be the number of edges $XY \in E(G)$ such that $n \in X \cup Y$. Since G generates all but at most $\varepsilon 2^n$ subsets of $[n]$, at least $2^{2\ell} - \varepsilon 2^n = (1-2\varepsilon)2^{2\ell}$ sets containing n correspond to edges of G , which implies that

$$e_n \geq (1-2\varepsilon)2^{2\ell} = (1-o(1))2^{2\ell}.$$

Let $\phi := |\mathcal{U}_n^+|/|\mathcal{X}_n^+|$ and $\theta := |\mathcal{V}_n^+|/|\mathcal{Y}_n^+|$, then $\phi, \theta \in [0, 1]$. Combining with (7), we have

$$\begin{aligned} e_n &\leq |\mathcal{U}_n^+||\mathcal{Y}_n^-| + |\mathcal{X}_n^+ \setminus \mathcal{U}_n^+| \left(2^{\ell-1} + o(2^\ell) \right) + |\mathcal{V}_n^+||\mathcal{X}_n^-| + |\mathcal{Y}_n^+ \setminus \mathcal{V}_n^+| \left(2^{\ell-1} + o(2^\ell) \right) \\ &= \phi(1-o(1))2^{2\ell-1} + (1-\phi)(1-o(1))2^{2\ell-2} + \theta(1-o(1))2^{2\ell-1} + (1-\theta)(1-o(1))2^{2\ell-2} \\ &= (2 + \phi + \theta - o(1))2^{2\ell-2}. \end{aligned}$$

Hence,

$$(1-o(1))2^{2\ell} \leq e_n \leq (2 + \phi + \theta - o(1))2^{2\ell-2},$$

which implies that $\phi, \theta = 1 - o(1)$. Therefore, $|\mathcal{X}_n^+ \setminus \mathcal{U}_n^+| = o(|\mathcal{X}_n^+|) = o(2^\ell)$, and $|\mathcal{Y}_n^+ \setminus \mathcal{V}_n^+| = o(|\mathcal{Y}_n^+|) = o(2^\ell)$, as desired.

Now we have

$$|\mathcal{X} \setminus \mathcal{U}| + |\mathcal{Y} \setminus \mathcal{V}| = |\mathcal{X}_n^- \setminus \mathcal{U}_n^-| + |\mathcal{X}_n^+ \setminus \mathcal{U}_n^+| + |\mathcal{Y}_n^- \setminus \mathcal{V}_n^-| + |\mathcal{Y}_n^+ \setminus \mathcal{V}_n^+| = o(2^\ell).$$

For a set $F \in \mathcal{H}$, notice that $F \in (\mathcal{X} \setminus \mathcal{U}) \cup (\mathcal{Y} \setminus \mathcal{V})$ if and only if $F \in \mathcal{X}$ satisfies $F \cap S_2 \neq \emptyset$ or $F \in \mathcal{Y}$ satisfies $F \cap S_1 \neq \emptyset$. For $S \subseteq S_1$, if $S \cup \{n\} \in \mathcal{X}$, then $S \in \mathcal{X}_n^+$ by definition. Hence, there are at least $2^{|S_1|} - |\mathcal{X}_n^+| = 2^\ell - (1 - o(1))2^{\ell-1} = (1 + o(1))2^{\ell-1}$ sets $S \subseteq S_1$ satisfying $S \cup \{n\} \notin \mathcal{X}$. Similarly, there are at least $(1 + o(1))2^{\ell-1}$ sets $S' \subseteq S_2$ satisfying $S' \cup \{n\} \notin \mathcal{Y}$. Thus, there are at least $(1 + o(1))2^{2\ell-2}$ sets of the form $S \cup S' \cup \{n\} \subseteq [n]$ satisfying $S \subseteq S_1, S \cup \{n\} \notin \mathcal{X}$ and $S' \subseteq S_2, S' \cup \{n\} \notin \mathcal{Y}$. Denote the family of sets of this form by \mathcal{S} . Since G generates all but at most $\varepsilon 2^n$ subsets of $[n]$, at least $(1 + o(1))2^{2\ell-2} - \varepsilon 2^{2\ell+1} = (1 - o(1))2^{2\ell-2}$ sets in \mathcal{S} correspond to edges of G . If $F \in \mathcal{S}$ can be expressed as a disjoint union of $X \in \mathcal{X}$ and $Y \in \mathcal{Y}$, then either $X \cap S_2 \neq \emptyset$ or $Y \cap S_1 \neq \emptyset$, implying that either X or Y is in $(\mathcal{X} \setminus \mathcal{U}) \cup (\mathcal{Y} \setminus \mathcal{V})$. Hence, the number of choices for F is at most $o(2^\ell)|\mathcal{H}| \leq o(2^\ell) \cdot 3(1 + \delta)2^\ell \ll (1 - o(1))2^{2\ell-2}$, a contradiction. \square

Claim 2.11. $A \cup B = [n]$.

Proof. Suppose for a contradiction that $A \cup B \neq [n]$. We may assume without loss of generality that $n \notin A \cup B$. Let $x := |\mathcal{X}_n^+|/|\mathcal{X}|$ and $y := |\mathcal{Y}_n^+|/|\mathcal{Y}|$, then $x, y < 1/3$ by the definitions of A and B . Recalling that $e_n \geq (1 - 2\varepsilon)2^{2\ell}$ is the number of disjoint pairs $X \in \mathcal{X}, Y \in \mathcal{Y}$ such that $n \in X \cup Y$, we have

$$\begin{aligned} (1 - 2\varepsilon)2^{2\ell} \leq e_n &\leq |\mathcal{X}_n^+||\mathcal{Y}_n^-| + |\mathcal{Y}_n^+||\mathcal{X}_n^-| = |\mathcal{X}_n^+|(|\mathcal{Y}| - |\mathcal{Y}_n^+|) + |\mathcal{Y}_n^+|(|\mathcal{X}| - |\mathcal{X}_n^+|) \\ &= x\alpha 2^\ell(\beta 2^\ell - y\beta 2^\ell) + y\beta 2^\ell(\alpha 2^\ell - x\alpha 2^\ell) = (x + y - 2xy)\alpha\beta 2^{2\ell}. \end{aligned} \quad (9)$$

Define the function $f(x, y) := x + y - 2xy$. On $0 \leq x, y \leq 1/3$, the function $f(x, y)$ attains its maximum value $4/9$, when $x = y = 1/3$. Combining with (9), we have

$$1 - 2\varepsilon \leq f(x, y)\alpha\beta \leq \frac{4}{9}\alpha\beta.$$

Recalling that $\alpha + \beta \leq 3(1 + \delta)$ by (3), we then get

$$\frac{3}{2} - 4\sqrt{\frac{\varepsilon}{2}} \leq \alpha, \beta \leq \frac{3}{2} + 4\sqrt{\frac{\varepsilon}{2}}.$$

Additionally, $\alpha\beta \leq \frac{9}{4}(1 + \delta)^2$ by (4), implying that $1 - 2\varepsilon \leq f(x, y)\alpha\beta \leq \frac{9}{4}(1 + \delta)^2 f(x, y)$, so

$$\frac{1}{3} - 3\varepsilon \leq x, y \leq \frac{1}{3}.$$

In summary, we have

$$|\mathcal{X}|, |\mathcal{Y}| = (3/2 - o(1))2^\ell, \quad |\mathcal{X}_n^+|, |\mathcal{Y}_n^+| = (1 - o(1))2^{\ell-1} \quad \text{and} \quad |\mathcal{X}_n^-|, |\mathcal{Y}_n^-| = (1 - o(1))2^\ell,$$

where $\varepsilon = o(1)$. Therefore, Claim 2.11 follows from Lemma 2.10. \square

Claim 2.12. $A \cap B = \emptyset$.

Proof. Suppose for a contradiction that $A \cap B \neq \emptyset$. We may assume without loss of generality that $n \in A \cap B$. Let $x := |\mathcal{X}_n^+|/|\mathcal{X}|$ and $y := |\mathcal{Y}_n^+|/|\mathcal{Y}|$, then $x, y \geq 1/3$ by the definitions of A and B . Notice that

$$(2 - 2\varepsilon)2^{2\ell} \leq e(G) \leq |\mathcal{X}||\mathcal{Y}| - |\mathcal{X}_n^+||\mathcal{Y}_n^+| = (1 - xy)\alpha\beta 2^{2\ell}.$$

Hence,

$$2 - 2\varepsilon \leq (1 - xy)\alpha\beta \leq \frac{8}{9}\alpha\beta.$$

Recalling that $\alpha + \beta \leq 3(1 + \delta)$ by (3), we then get

$$\frac{3}{2} - 2\sqrt{\varepsilon} \leq \alpha, \beta \leq \frac{3}{2} + 2\sqrt{\varepsilon}.$$

Additionally, $\alpha\beta \leq \frac{9}{4}(1 + \delta)^2$ by (4), implying that $2 - 2\varepsilon \leq (1 - xy)\alpha\beta \leq \frac{9}{4}(1 + \delta)^2(1 - xy)$, so

$$\frac{1}{3} \leq x, y \leq \frac{1}{3} + 3\varepsilon.$$

In summary, we have

$$|\mathcal{X}|, |\mathcal{Y}| = (3/2 - o(1))2^\ell, \quad |\mathcal{X}_n^+|, |\mathcal{Y}_n^+| = (1 - o(1))2^{\ell-1} \quad \text{and} \quad |\mathcal{X}_n^-|, |\mathcal{Y}_n^-| = (1 - o(1))2^\ell,$$

where again $\varepsilon = o(1)$. By Lemma 2.10, we have completed the proof of Claim 2.12. \square

By Claims 2.11 and 2.12, we now know that $A \cup B$ is a partition of $[n]$. It remains to show that $A \cup B$ is in fact an equipartition of $[n]$ and \mathcal{X}, \mathcal{Y} are close to $2^A, 2^B$, respectively. The following observation is simple but will be useful hereafter.

Observation 2.13. *If $F \in \mathcal{X} \setminus 2^A$, then F has at most $2|\mathcal{Y}|/3$ neighbors in \mathcal{Y} . Similarly, if $F \in \mathcal{Y} \setminus 2^B$, then F has at most $2|\mathcal{X}|/3$ neighbors in \mathcal{X} .*

Proof. By symmetry, it suffices to prove the first part. Suppose $F \in \mathcal{X} \setminus 2^A$, then there exists $i \in [n]$ such that $i \in F \cap B$. By the definition of $B = \mathcal{Y}_{1/3}$ (see (6)), there are at least $|\mathcal{Y}|/3$ sets in \mathcal{Y} containing i , which therefore have non-empty intersection with F . By the definition of G , we conclude that F has at most $|\mathcal{Y}| - |\mathcal{Y}|/3 = 2|\mathcal{Y}|/3$ neighbors in \mathcal{Y} , as desired. \square

Claim 2.14. *We have $|\mathcal{X} \cap 2^A| \geq (2/3 - 3\varepsilon)|\mathcal{X}|$ and $|\mathcal{Y} \cap 2^B| \geq (2/3 - 3\varepsilon)|\mathcal{Y}|$. Additionally, $[n] = A \cup B$ is an equipartition.*

Proof. Let $\theta := \frac{|\mathcal{X} \cap 2^A|}{|\mathcal{X}|}$ and $\phi := \frac{|\mathcal{Y} \cap 2^B|}{|\mathcal{Y}|}$. By Observation 2.13 and (4), we have

$$(2 - 2\varepsilon)2^{2\ell} \leq e(G) \leq |\mathcal{X} \cap 2^A| \cdot |\mathcal{Y}| + |\mathcal{X} \setminus 2^A| \cdot \frac{2}{3}|\mathcal{Y}| = \frac{2 + \theta}{3}\alpha\beta 2^{2\ell} \leq \frac{2 + \theta}{3} \cdot \frac{9}{4}(1 + \delta)^2 2^{2\ell}.$$

Hence $\frac{2 + \theta}{3} \cdot \frac{9}{4}(1 + \delta)^2 \geq 2 - 2\varepsilon$, which implies that $\theta \geq 2/3 - 3\varepsilon$, as desired. Similarly, we can prove $\phi \geq 2/3 - 3\varepsilon$.

If $|A| \leq \ell - 1$, then

$$|\mathcal{X}| = \frac{|\mathcal{X} \cap 2^A|}{\theta} \leq \frac{|2^A|}{\theta} \leq \frac{2^{\ell-1}}{2/3 - 3\varepsilon} < (1 - 2\varepsilon)2^\ell,$$

a contradiction to (5), so we have $|A| \geq \ell$. Similarly, we have $|B| \geq \ell$. Therefore, $[n] = A \cup B$ is an equipartition. \square

According to Claim 2.14, we can assume from now on that $|A| = \ell$ and $|B| = \ell + 1$. We claim that

$$\alpha \leq \frac{3}{2} + 8\varepsilon, \quad \beta \geq \frac{3}{2} - 7\varepsilon, \tag{10}$$

which are better bounds for α, β than (5). Notice that we only need to show $\beta \geq 3/2 - 7\varepsilon$, as $\alpha + \beta \leq 3(1 + \delta)$ will then imply $\alpha \leq 3(1 + \delta) - \beta \leq 3/2 + 8\varepsilon$. Suppose that $\beta = 3/2 - \gamma$ with some $\gamma \geq 0$, then $\alpha \leq 3/2 + 3\delta + \gamma$ since $\alpha + \beta \leq 3(1 + \delta)$ by (3). By (2) and Observation 2.13, we have

$$\begin{aligned} (2 - 2\varepsilon)2^{2\ell} &\leq e(G) \leq |\mathcal{X} \cap 2^A||\mathcal{Y}| + |\mathcal{X} \setminus 2^A| \cdot \frac{2}{3}|\mathcal{Y}| = |\mathcal{X} \cap 2^A||\mathcal{Y}| + (|\mathcal{X}| - |\mathcal{X} \cap 2^A|) \cdot \frac{2}{3}|\mathcal{Y}| \\ &= \left(\frac{1}{3}|\mathcal{X} \cap 2^A| + \frac{2}{3}|\mathcal{X}|\right)|\mathcal{Y}| \leq \left(\frac{1}{3} \cdot 2^\ell + \frac{2}{3} \left(\frac{3}{2} + 3\delta + \gamma\right) 2^\ell\right) \left(\frac{3}{2} - \gamma\right) 2^\ell \leq \left(2 + 3\delta - \frac{1}{3}\gamma\right) 2^{2\ell}, \end{aligned}$$

which implies that $2 - 2\varepsilon \leq 2 + 3\delta - \gamma/3$. Therefore, $\gamma \leq 7\varepsilon$.

The final three claims show that \mathcal{X} and \mathcal{Y} are not too far from 2^A and 2^B , respectively.

Claim 2.15. (i) $|2^A \setminus \mathcal{X}| \leq 24\varepsilon 2^\ell$.

(ii) $|\mathcal{Y} \setminus 2^B| \leq (\sqrt{\varepsilon} + 3\varepsilon)2^\ell$.

Proof. Let

$$D := \min\{|2^B \setminus \mathcal{Y}|, |\mathcal{Y} \setminus 2^B|\},$$

then $D \leq |\mathcal{Y} \setminus 2^B| \leq (1/3 + 3\varepsilon)|\mathcal{Y}| < 2|\mathcal{Y}|/3$ by Claim 2.14. Define \mathcal{Y}' from \mathcal{Y} by adding D sets in $2^B \setminus \mathcal{Y}$ and deleting D sets in $\mathcal{Y} \setminus 2^B$. Thus, $|\mathcal{Y}'| = |\mathcal{Y}| \leq (2 + 3\varepsilon)2^\ell$ by (5). Note that \mathcal{X} and \mathcal{Y}' are not necessarily disjoint. Let $G_1 = G_{\mathcal{X}, \mathcal{Y}'}$. If $D = |2^B \setminus \mathcal{Y}| \leq |\mathcal{Y} \setminus 2^B|$, then $2^B \subseteq \mathcal{Y}'$ and $|\mathcal{Y}' \setminus 2^B| = |\mathcal{Y}'| - 2^{\ell+1} \leq 2\varepsilon 2^{\ell+1}$; if $D = |\mathcal{Y} \setminus 2^B| \leq |2^B \setminus \mathcal{Y}|$, then $\mathcal{Y}' \subseteq 2^B$ and $|\mathcal{Y}' \setminus 2^B| = 0$. In both cases, we have $|\mathcal{Y}' \cap 2^B| \geq |\mathcal{Y} \cap 2^B|$ and

$$|\mathcal{Y}' \setminus 2^B| \leq 2\varepsilon 2^{\ell+1}. \quad (11)$$

We now compare $e(G_1)$ and $e(G)$. Every deleted $Y \in \mathcal{Y} \setminus 2^B$ has at most $2|\mathcal{X}|/3$ neighbors in \mathcal{X} by Observation 2.13. On the other hand, every added $S \in 2^B \setminus \mathcal{Y}$ is disjoint from every set in $2^A \cap \mathcal{X}$, thus has at least $|2^A \cap \mathcal{X}| \geq (2/3 - 3\varepsilon)|\mathcal{X}|$ neighbors in \mathcal{X} by Claim 2.14. Therefore, by (3),

$$e(G) - e(G_1) \leq D \left(\frac{2}{3}|\mathcal{X}| - \left(\frac{2}{3} - 3\varepsilon \right) |\mathcal{X}| \right) \leq \frac{2}{3}|\mathcal{Y}| \cdot 3\varepsilon|\mathcal{X}| = 2\varepsilon\alpha\beta 2^{2\ell} \leq 2\varepsilon \cdot \frac{9}{4}(1 + \delta)2^{2\ell} \leq 3\varepsilon 2^{2\ell+1},$$

which, with (2), implies that

$$e(G_1) \geq e(G) - 3\varepsilon 2^{2\ell+1} \geq (1 - \varepsilon)2^{2\ell+1} - 3\varepsilon 2^{2\ell+1} = (1 - 4\varepsilon)2^{2\ell+1}.$$

Similarly, we define another bipartite graph G_2 . Let

$$C := \min\{|2^A \setminus \mathcal{X}|, |\mathcal{X} \setminus 2^A|\}.$$

Define \mathcal{X}' from \mathcal{X} by adding C sets in $2^A \setminus \mathcal{X}$ and deleting C sets in $\mathcal{X} \setminus 2^A$. Thus, $|\mathcal{X}'| = |\mathcal{X}| \geq (1 - 3\varepsilon)2^\ell$ by (5). Note that \mathcal{X}' and \mathcal{Y}' are not necessarily disjoint. Let $G_2 = G_{\mathcal{X}', \mathcal{Y}'}$. If $C = |2^A \setminus \mathcal{X}| \leq |\mathcal{X} \setminus 2^A|$, then $2^A \subseteq \mathcal{X}'$ and $|\mathcal{X}' \cap 2^A| = |2^A| = 2^\ell$; if $C = |\mathcal{X} \setminus 2^A| \leq |2^A \setminus \mathcal{X}|$, then $\mathcal{X}' \subseteq 2^A$ and $|\mathcal{X}' \cap 2^A| = |\mathcal{X}'| \geq (1 - 3\varepsilon)2^\ell$. In both cases, we have $|\mathcal{X}' \cap 2^A| \geq (1 - 3\varepsilon)2^\ell$. We now compare $e(G_2)$ and $e(G_1)$. Every deleted $X \in \mathcal{X} \setminus 2^A$ intersects B , thus has at most 2^ℓ neighbors in 2^B . By (11), X has at most $2\varepsilon 2^{\ell+1}$ neighbors in $\mathcal{Y}' \setminus 2^B$, thus has at most $(1 + 4\varepsilon)2^\ell$ in \mathcal{Y}' . On the other hand, every added $S \in 2^A \setminus \mathcal{X}$ is disjoint from every set in $2^B \cap \mathcal{Y}'$, thus has at least

$$|2^B \cap \mathcal{Y}'| = |\mathcal{Y}'| - |\mathcal{Y}' \setminus 2^B| \geq (3/2 - 7\varepsilon)2^\ell - (2\varepsilon)2^{\ell+1} = (3/2 - 11\varepsilon)2^\ell$$

neighbors by (10) and (11). Therefore,

$$e(G_2) \geq e(G_1) + C \left(\frac{3}{2} - 11\varepsilon - (1 + 4\varepsilon) \right) 2^\ell \geq (1 - 4\varepsilon)2^{2\ell+1} + C \left(\frac{1}{2} - 15\varepsilon \right) 2^\ell. \quad (12)$$

If $|\mathcal{X}'| \leq 2^\ell$, then by (3), we have that $e(G_2) \leq |\mathcal{X}'||\mathcal{Y}'| \leq |\mathcal{X}'|(3(1 + \delta)2^\ell - |\mathcal{X}'|)$ attains its maximum value $(1 + \frac{3}{2}\delta)2^{2\ell+1}$ when $|\mathcal{X}'| = 2^\ell$. If $|\mathcal{X}'| > 2^\ell$, then let $|\mathcal{X}'| = (1 + \gamma)2^\ell$ for some $\gamma > 0$. Since $|\mathcal{X}'| + |\mathcal{Y}'| = |\mathcal{X}| + |\mathcal{Y}| \leq 3(1 + \delta)2^\ell$ by (3), we have $|\mathcal{Y}'| \leq (2 + 3\delta - \gamma)2^\ell$. Recalling $|\mathcal{Y}'| = |\mathcal{Y}| \geq (3/2 - 7\varepsilon)2^\ell$ by (10), we have $\gamma \leq 1/2 + 7\varepsilon + 3\delta$. By the definition of \mathcal{X}' , we have

$\mathcal{X}' \supseteq 2^A$, so $|\mathcal{X}' \setminus 2^A| = |\mathcal{X}'| - |2^A| = \gamma 2^\ell$. Every $X \in \mathcal{X}' \setminus 2^A$ intersects B , thus has at most 2^ℓ neighbors in $\mathcal{Y}' \cap 2^B$, so it has at most $(1 + 4\varepsilon)2^\ell$ neighbors in \mathcal{Y}' by (11). Therefore,

$$\begin{aligned} e(G_2) &\leq |\mathcal{X}' \setminus 2^A|(1 + 4\varepsilon)2^\ell + |\mathcal{X}' \cap 2^A||\mathcal{Y}'| \leq \gamma 2^\ell(1 + 2\varepsilon)2^\ell + 2^\ell(2 + 3\delta - \gamma)2^\ell \\ &= (1 + \varepsilon\gamma + \frac{3}{2}\delta)2^{2\ell+1} \leq (1 + \varepsilon)2^{2\ell+1}. \end{aligned}$$

In both cases, we have

$$e(G_2) \leq (1 + \varepsilon)2^{2\ell+1}. \quad (13)$$

Combining (2), (12) and (13), we conclude that

$$e(G_1) - e(G) \leq e(G_2) - e(G) \leq (1 + \varepsilon)2^{2\ell+1} - (1 - \varepsilon)2^{2\ell+1} = 2\varepsilon 2^{2\ell+1}. \quad (14)$$

Now we prove (i). By (12) and (13), we have

$$C \leq \frac{10\varepsilon}{1/2 - 15\varepsilon}2^\ell \leq 21\varepsilon 2^\ell, \quad (15)$$

so if $C = |2^A \setminus \mathcal{X}|$, then we are done. Assume $C = |\mathcal{X} \setminus 2^A|$. Then, by (5) and (15), we have

$$|2^A \setminus \mathcal{X}| = |2^A| - |2^A \cap \mathcal{X}| = 2^{|A|} - (|\mathcal{X}| - |\mathcal{X} \setminus 2^A|) \leq 2^\ell - (1 - 3\varepsilon)2^\ell + 21\varepsilon 2^\ell = 24\varepsilon 2^\ell,$$

as desired.

For (ii), we show that it suffices to prove

$$D \leq \sqrt{\varepsilon}2^\ell. \quad (16)$$

Indeed, if $D = |\mathcal{Y} \setminus 2^B|$, then we are done. Assume $D = |2^B \setminus \mathcal{Y}|$. Then, by (5), we have

$$|\mathcal{Y} \setminus 2^B| = |\mathcal{Y}| - |2^B \cap \mathcal{Y}| = |\mathcal{Y}| - (|2^B| - |2^B \setminus \mathcal{Y}|) \leq (2 + 3\varepsilon)2^\ell - 2^{2\ell+1} + \sqrt{\varepsilon}2^\ell = (\sqrt{\varepsilon} + 3\varepsilon)2^\ell,$$

as desired.

Now suppose for a contradiction that $D > \sqrt{\varepsilon}2^\ell$. We claim that there exists some $Y \in \mathcal{Y} \setminus 2^B$ having at least $2|\mathcal{X}|/3 - 8\sqrt{\varepsilon}2^\ell$ neighbors in \mathcal{X} . Otherwise, recall that every $S \in \mathcal{Y}' \setminus \mathcal{Y} \subseteq 2^B \setminus \mathcal{Y}$ has at least $(2/3 - 3\varepsilon)|\mathcal{X}|$ neighbors in \mathcal{X} and $|\mathcal{X}| \leq (3/2 + 8\varepsilon)2^\ell$ by (10). Then,

$$e(G_1) - e(G) \geq D \left(\left(\frac{2}{3} - 3\varepsilon \right) |\mathcal{X}| - \left(\frac{2|\mathcal{X}|}{3} - 8\sqrt{\varepsilon}2^\ell \right) \right) \geq \sqrt{\varepsilon} \left(8\sqrt{\varepsilon} - \frac{9}{2}\varepsilon - 24\varepsilon^2 \right) 2^{2\ell} > 2\varepsilon 2^{2\ell+1},$$

a contradiction to (14). Fix some $Y \in \mathcal{Y} \setminus 2^B$ having at least $2|\mathcal{X}|/3 - 8\sqrt{\varepsilon}2^\ell$ neighbors in \mathcal{X} . Note that there exists $i \in [n]$ such that $i \in Y \cap A$. We may assume without loss of generality that $i = n$. By the definition of G , at most $|\mathcal{X}| - (2|\mathcal{X}|/3 - 8\sqrt{\varepsilon}2^\ell) = (1/3 + o(1))|\mathcal{X}|$ sets in \mathcal{X} contain n , i.e., $|\mathcal{X}_n^+|/|\mathcal{X}| \leq 1/3 + o(1)$. On the other hand, since $|2^A \setminus \mathcal{X}| \leq 24\varepsilon 2^\ell$ by (i), at least $2^{\ell-1} - 24\varepsilon 2^\ell = (1 - o(1))2^{\ell-1}$ subsets of A containing n are contained in \mathcal{X} , so $|\mathcal{X}_n^+| \geq (1 - o(1))2^{\ell-1}$ and hence $|\mathcal{X}| \geq (3/2 - o(1))2^\ell$. Recalling that $|\mathcal{X}| \leq (3/2 + o(1))2^\ell$ by (10), we have $|\mathcal{X}| = (3/2 - o(1))2^\ell$, which then implies that $|\mathcal{X}_n^+| = (1 - o(1))2^{\ell-1}$ and $|\mathcal{Y}| = (3/2 - o(1))2^\ell$. By the definition of B , we have $|\mathcal{Y}_n^+|/|\mathcal{Y}| \leq 1/3$, since $n \in A$. Recall (9), where we have now $x, y \leq 1/3 + o(1)$ and $\alpha, \beta \leq 3/2 + o(1)$. We get $y = 1/3 + o(1)$ and hence $|\mathcal{Y}_n^+| = (1 - o(1))2^{\ell-1}$. Therefore, $|\mathcal{X}_n^-|, |\mathcal{Y}_n^-| = (1 - o(1))2^\ell$. Again, by Lemma 2.10, we obtain a contradiction and complete the proof of Claim 2.15. \square

Claim 2.16. $|2^B \setminus \mathcal{Y}| \leq 6\sqrt{\varepsilon}2^\ell$.

Proof. For every $e \in E(G)$, call e *bad* if e has an endpoint in $\mathcal{Y} \setminus 2^B$ and *good* otherwise. By Claim 2.15 (ii) and (10), G has at most

$$|\mathcal{Y} \setminus 2^B| |\mathcal{X}| \leq (\sqrt{\varepsilon} + 3\varepsilon) 2^\ell \cdot (3/2 + 8\varepsilon) 2^\ell \leq 2\sqrt{\varepsilon} 2^{2\ell} \quad (17)$$

bad edges.

Fix $S' \in 2^B \setminus \mathcal{Y}$ and choose a set $F \subseteq [n]$ of the form $F = S \cup S'$, where $S \subseteq A$. If F corresponds to a good edge of G , assume that $F = X \cup Y$ with $X \in \mathcal{X}, Y \in \mathcal{Y} \cap 2^B, X \cap Y = \emptyset$. By the assumption that $S' \in 2^B \setminus \mathcal{Y}$ and $Y \in \mathcal{Y} \cap 2^B$, we have $Y \neq S'$ and hence $Y \subsetneq S'$. Hence, $X \cap B \neq \emptyset$, so $X \in \mathcal{X} \setminus 2^A$. Furthermore, $X \cap A = S$, which implies that S is determined by X . By (10) and Claim 2.15 (i), we have

$$|\mathcal{X} \setminus 2^A| = |\mathcal{X}| - |\mathcal{X} \cap 2^A| = |\mathcal{X}| - (|2^A| - |2^A \setminus \mathcal{X}|) \leq (3/2 + 8\varepsilon) 2^\ell - (1 - 24\varepsilon) 2^\ell = (1/2 + 32\varepsilon) 2^\ell.$$

Hence, for every fixed $S' \in 2^B \setminus \mathcal{Y}$, there are at most $(1/2 + 32\varepsilon) 2^\ell$ sets F of the form $F = S \cup S'$, where $S \subseteq A$, corresponding to good edges of G . There are $2^\ell |2^B \setminus \mathcal{Y}|$ sets of the form $S \cup S'$ with $S \subseteq A, S' \in 2^B \setminus \mathcal{Y}$ in total, so there are at least $|2^B \setminus \mathcal{Y}| (2^\ell - (1/2 + 32\varepsilon) 2^\ell)$ sets of the form $S \cup S'$ with $S \subseteq A, S' \in 2^B \setminus \mathcal{Y}$ that correspond to bad edges of G or do not correspond to edges of G . Recalling that \mathcal{H} is a $(1 - \varepsilon)$ -2-generator for $[n]$ and using (17), we have

$$|2^B \setminus \mathcal{Y}| \left(2^\ell - (1/2 + 32\varepsilon) 2^\ell \right) \leq 2\sqrt{\varepsilon} 2^{2\ell} + \varepsilon 2^n = (2\sqrt{\varepsilon} + 2\varepsilon) 2^{2\ell}.$$

Therefore, we get $|2^B \setminus \mathcal{Y}| \leq 6\sqrt{\varepsilon} 2^\ell$. □

Claim 2.17. $|\mathcal{X} \setminus 2^A| \leq 7\sqrt{\varepsilon} 2^\ell$.

Proof. By Claims 2.15 (i) and 2.16, we have

$$\begin{aligned} 3(1 + \delta) 2^\ell \geq |\mathcal{H}| &= |\mathcal{X}| + |\mathcal{Y}| = |\mathcal{X} \setminus 2^A| + (|2^A| - |2^A \setminus \mathcal{X}|) + |\mathcal{Y} \setminus 2^B| + (|2^B| - |2^B \setminus \mathcal{Y}|) \\ &\geq |\mathcal{X} \setminus 2^A| + (1 - 24\varepsilon) 2^\ell + (2 - 6\sqrt{\varepsilon}) 2^\ell = |\mathcal{X} \setminus 2^A| + 3 \cdot 2^\ell - (6\sqrt{\varepsilon} + 24\varepsilon) 2^\ell, \end{aligned}$$

implying $|\mathcal{X} \setminus 2^A| \leq 7\sqrt{\varepsilon} 2^\ell$. □

By Claim 2.14, $[n] = A \cup B$ is an equipartition. By Claims 2.15, 2.16 and 2.17, we have

$$|\mathcal{H} \Delta (2^A \cup 2^B)| \leq (14\sqrt{\varepsilon} + 27\varepsilon) 2^\ell \leq \varepsilon' 2^\ell,$$

completing the proof of Theorem 2.2. □

3 Proof of Theorem 1.1

In this section, we will employ a perturbation argument to deduce Theorem 1.1 from Theorem 1.2. Let n be a sufficiently large integer and \mathcal{F} be a maximal 3-wise intersecting family on $[n]$ of size at most $2^{\lfloor n/2 \rfloor} + 2^{\lfloor n/2 \rfloor} - 3$. Let $\mathcal{H} := \bar{\mathcal{F}} = \{F^c : F \in \mathcal{F}\}$ and fix some $\varepsilon \in (0, 1/4)$. By Theorem 1.2, there exists $S \subseteq [n]$ of size $\lfloor n/2 \rfloor$ such that $\mathcal{F}_0 := \{A : A \subseteq S\} \cup \{B : B \subseteq S^c\}$ satisfies $|\mathcal{H} \Delta \mathcal{F}_0| \leq \varepsilon 2^{\lfloor n/2 \rfloor}$.

Recall that \mathcal{H} is downward closed since \mathcal{F} is upward closed, so $\emptyset \in \mathcal{H}$. We first prove that $S \notin \mathcal{H} = \bar{\mathcal{F}}$. Suppose for a contradiction that $S^c \in \mathcal{F}$. Since $|\mathcal{H} \Delta \mathcal{F}_0| \leq \varepsilon 2^{\lfloor n/2 \rfloor}$, among the $2^{\lfloor n/2 \rfloor}$ subsets of S^c , there exists some $A \subseteq S^c$ such that both A and $S^c \setminus A$ are contained in \mathcal{H} . However, this would imply that $S^c, A^c, (S^c \setminus A)^c = S \cup A$ are in \mathcal{F} . As $S^c \cap A^c \cap (S \cup A) = \emptyset$, this contradicts that \mathcal{F} is a 3-wise intersecting family. It can be proved similarly that $S^c \notin \mathcal{H}$.

We work with the following partition $\mathcal{H} = \mathcal{H}_1 \cup \mathcal{H}_2 \cup \mathcal{H}_3 \cup \{\emptyset\}$, where

$$\mathcal{H}_1 := \{A \in \mathcal{H} : \emptyset \subsetneq A \subsetneq S\}, \quad \mathcal{H}_2 := \{B \in \mathcal{H} : \emptyset \subsetneq B \subsetneq S^c\}, \quad \mathcal{H}_3 := \mathcal{H} \setminus \mathcal{F}_0.$$

Claim 3.1. $|2^{[n]} \setminus (\mathcal{F} \cup \mathcal{F}_0)| \leq |\mathcal{H}_1||\mathcal{H}_2| + |\mathcal{H}_3| \cdot \varepsilon 2^{\lfloor n/2 \rfloor}$.

Proof. If $A \notin \mathcal{F}$, then as stated in the proof of Lemma 2.1, there exist $B, C \in \mathcal{H}$ such that $A = B \cup C$ with $B \cap C = \emptyset$. There exists a pair $\{B, C\}$ such that

$$g(\{B, C\}) := \min\{|B \setminus S| + |C \setminus S^c|, |B \setminus S^c| + |C \setminus S|\}$$

attains its minimum value over all such choices of B and C . Therefore, we can define an injection f on $2^{[n]} \setminus (\mathcal{F} \cup 2^S \cup 2^{S^c})$ by mapping A to a pair of sets $\{B, C\}$ in \mathcal{H} , which has minimum $g(\{B, C\})$ given $A = B \cup C$ and $B \cap C = \emptyset$. Since $\emptyset \in \mathcal{H}$, $S, S^c \notin \mathcal{H}$ and \mathcal{H} is downward closed, we have that $f(A) = \{B, C\}$ must be one of the following two types: $|\{B, C\} \cap \mathcal{H}_1| = |\{B, C\} \cap \mathcal{H}_2| = 1$; $|\{B, C\} \cap \mathcal{H}_3| \geq 1$. The number of pairs $\{B, C\}$ satisfying $|\{B, C\} \cap \mathcal{H}_1| = |\{B, C\} \cap \mathcal{H}_2| = 1$ is at most $|\mathcal{H}_1||\mathcal{H}_2|$, so it suffices to prove that if $f(A) = \{B, C\}$ with $B \in \mathcal{H}_3$, then there are at most $\varepsilon 2^{\lfloor n/2 \rfloor}$ choices for C .

Suppose $B = X \cup Y$, where $\emptyset \neq X \subsetneq S, \emptyset \neq Y \subsetneq S^c$. Then, $g(\{B, C\}) > 0$ and $Y \in \mathcal{H}$, since $Y \subseteq B \in \mathcal{H}$ and \mathcal{H} is downward closed. There are three possibilities for C .

- (1) $C \subseteq S$: Define $B' := X \cup C, C' := Y$, then $B' \cup C' = B \cup C = A$. Since $g(\{B', C'\}) = 0 < g(\{B, C\})$ and $C' = Y \in \mathcal{H}$, we have that $B' \notin \mathcal{H}$ by the definition of f . Note that $B' = X \cup C$ is determined by C , so the number of choices for $C \subseteq S$ is at most $|2^S \setminus \mathcal{H}|$.
- (2) $C \subseteq S^c$: Similarly, the number of choices for C is at most $|2^{S^c} \setminus \mathcal{H}|$.
- (3) $C \in \mathcal{H} \setminus \mathcal{F}_0$: The number of choices for such C is at most $|\mathcal{H} \setminus \mathcal{F}_0|$.

In summary, the number of choices for C is at most

$$|2^S \setminus \mathcal{H}| + |2^{S^c} \setminus \mathcal{H}| + |\mathcal{H} \setminus \mathcal{F}_0| = |\mathcal{H} \Delta \mathcal{F}_0| \leq \varepsilon 2^{\lfloor n/2 \rfloor},$$

as desired. \square

Claim 3.2. $|\mathcal{H}_1||\mathcal{H}_2| + |\mathcal{H}_3| \cdot \varepsilon 2^{\lfloor n/2 \rfloor} \leq (2^{\lfloor n/2 \rfloor} - 2)(2^{\lceil n/2 \rceil} - 2)$. Equality holds if and only if $\mathcal{H}_1 = \{A: \emptyset \subsetneq A \subsetneq S\}$, $\mathcal{H}_2 = \{B: \emptyset \subsetneq B \subsetneq S^c\}$ and $\mathcal{H}_3 = \emptyset$.

Proof. By the definitions of \mathcal{H}_1 and \mathcal{H}_2 , we have

$$|\mathcal{H}_1||\mathcal{H}_2| \leq (2^{\lfloor n/2 \rfloor} - 2)(2^{\lceil n/2 \rceil} - 2). \quad (18)$$

Notice that $\min\{|\mathcal{H}_1|, |\mathcal{H}_2|\} \geq 2^{\lfloor n/2 \rfloor} - |\mathcal{F}_0 \setminus \mathcal{H}| \geq 2^{\lfloor n/2 \rfloor} - |\mathcal{F}_0 \Delta \mathcal{H}| \geq (1 - \varepsilon)2^{\lfloor n/2 \rfloor}$. Define the function $\varphi(h_1, h_2, h_3) := h_1 h_2 + h_3 \cdot \varepsilon 2^{\lfloor n/2 \rfloor}$. Given that $\min\{h_1, h_2\} \geq (1 - \varepsilon)2^{\lfloor n/2 \rfloor}$, we claim that $\varphi(h'_1, h_2, h'_3) > \varphi(h_1, h_2, h_3)$ if $h'_1 = h_1 + 1$ and $h'_3 = h_3 - 1$. In fact, we have

$$\varphi(h'_1, h_2, h'_3) - \varphi(h_1, h_2, h_3) = h_2 - \varepsilon 2^{\lfloor n/2 \rfloor} \geq (1 - 2\varepsilon)2^{\lfloor n/2 \rfloor} > 0.$$

Therefore, fixing $h_1 + h_2 + h_3$ and stipulating $h_1, h_2 \geq (1 - \varepsilon)2^{\lfloor n/2 \rfloor}$ and $h_3 \geq 0$, the function $\varphi(h_1, h_2, h_3)$ attains its maximum when $h_3 = 0$. Take $h_i = |\mathcal{H}_i|$ for $i \in [3]$. Combining with (18), we have

$$|\mathcal{H}_1||\mathcal{H}_2| + |\mathcal{H}_3| \cdot \varepsilon 2^{\lfloor n/2 \rfloor} \leq (2^{\lfloor n/2 \rfloor} - 2)(2^{\lceil n/2 \rceil} - 2),$$

where equality holds if and only if $\mathcal{H}_1 = \{A: \emptyset \subsetneq A \subsetneq S\}$, $\mathcal{H}_2 = \{B: \emptyset \subsetneq B \subsetneq S^c\}$ and $\mathcal{H}_3 = \emptyset$. \square

By Claims 3.1 and 3.2, we have

$$2^n - (2^{\lfloor n/2 \rfloor} + 2^{\lceil n/2 \rceil} - 1) - |\mathcal{F}| \leq |2^{[n]} \setminus (\mathcal{F} \cup \mathcal{F}_0)| \leq (2^{\lfloor n/2 \rfloor} - 2)(2^{\lceil n/2 \rceil} - 2) = 2^n - 2 \cdot 2^{\lfloor n/2 \rfloor} - 2 \cdot 2^{\lceil n/2 \rceil} + 4,$$

which implies that $|\mathcal{F}| \geq 2^{\lfloor n/2 \rfloor} + 2^{\lceil n/2 \rceil} - 3$. By our assumption that $|\mathcal{F}| \leq 2^{\lfloor n/2 \rfloor} + 2^{\lceil n/2 \rceil} - 3$, equality has to hold in Claim 3.2. Hence, we have $\mathcal{H} = \mathcal{F}_0$, which means that \mathcal{F} is a balanced pair of linked cubes. This completes our proof of Theorem 1.1.

4 The Case when $k \geq 4$

Denote by $f(n, k)$ the minimum possible size of a maximal k -wise intersecting family on $[n]$. For the case when $k \geq 4$, we remark that our method also gives the following general bounds for $f(n, k)$.

Proposition 4.1. *For every $k \geq 4$ there exist positive constants c_k and d_k such that for every positive integer n , we have*

$$c_k \cdot 2^{n/(k-1)} \leq f(n, k) \leq d_k \cdot 2^{n/\lceil k/2 \rceil}.$$

Proof. The proof of Lemma 2.1 easily generalizes to give a lower bound for $f(n, k)$. Indeed, there is an injective map from \mathcal{F}^c to $\binom{\mathcal{F}}{k-1}$, such that each set $A \in \mathcal{F}^c$ is sent to a $(k-1)$ -tuple $\{B_1, B_2, \dots, B_{k-1}\}$ of sets in \mathcal{F} such that $B_1 \cap B_2 \cap \dots \cap B_{k-1} = A^c$ (unless A is so small that fewer than $k-1$ sets contain A^c). This observation immediately leads to the lower bound.

The linked cubes construction also generalizes. In more detail, suppose that ℓ divides n , and let S_1, S_2, \dots, S_ℓ be a partition of $[n]$ such that $|S_i| = n/\ell$ for each i . Let $\mathcal{F}_i = \{A: [n] \setminus S_i \subsetneq A \subseteq [n]\}$, and let $\mathcal{F} = \mathcal{F}_1 \cup \mathcal{F}_2 \cup \dots \cup \mathcal{F}_\ell$. For any set \mathcal{G} of $2\ell - 1$ elements of \mathcal{F} , there is an i such that $|\mathcal{G} \cap \mathcal{F}_i| \leq 1$. The sets in $\mathcal{G} - \mathcal{F}_i$ all contain S_i , and the set in $\mathcal{G} \cap \mathcal{F}_i$ contains at least one element of S_i , which element is in every set of \mathcal{G} . Hence, \mathcal{G} is $(2\ell - 1)$ -wise intersecting. On the other hand, if $A \notin \mathcal{F}$, then either A is missing elements from two distinct sets S_i , or A is missing all of the elements from some S_i . In either case, it is easy to find a non-intersecting $(2\ell - 1)$ -tuple of elements in $\mathcal{F} \cup \{A\}$.

For the case when k is even, let \mathcal{F}' be a maximal $(k-1)$ -intersecting family on $[n-1]$. Then the family $\mathcal{F} = \{A \cup \{n\} : A \in \mathcal{F}'\} \cup \{[n-1]\}$ is a maximal k -wise intersecting family.

These constructions give the upper bound. \square

Very recently, Janzer [21] showed that the lower bound in Proposition 4.1 is the correct order of magnitude of $f(n, k)$, by constructing a maximal k -wise intersecting family of size $O(2^{n/(k-1)})$ for every $k \geq 3$ and n . Note that in the special case $k = 3$, Theorem 1.1 matches Janzer's [21] construction.

Assume that n is sufficiently large. Let \mathcal{F} be a maximal k -wise intersecting family on $[n]$ with minimum size. Similarly to the case $k = 3$, one can show that $\bar{\mathcal{F}}$ is a $(1 - \varepsilon)$ - $(k-1)$ -generator for $[n]$, where $\varepsilon = o(1)$. A modification of the method in this paper could be used to determine the structure of $\bar{\mathcal{F}}$, if $|\bar{\mathcal{F}} \setminus \mathcal{F}_{n, k-1}|$ was small. However, Janzer [21] showed that $|\bar{\mathcal{F}} \setminus \mathcal{F}_{n, k-1}|$ cannot be small.

Theorem 4.2 ([21, Lemma 1.2]). *For every $k \geq 4$, there exist $d = d(k) > 0$, $c = c(k) > 0$ and $n_0 = n_0(k) > 0$ such that the following holds when $n \geq n_0$. Let $S_1 \cup \dots \cup S_{k-1}$ be a partition of $[n]$, where $\frac{n}{k-1} - d \leq |S_i| \leq \frac{n}{k-1} + d$ for every $i \in [k-1]$. Let $\mathcal{F}_0 = 2^{S_1} \cup \dots \cup 2^{S_{k-1}}$. For every set system $\mathcal{F} \subseteq 2^{[n]}$ with $|\mathcal{F} \setminus \mathcal{F}_0| \leq c \cdot 2^{n/(k-1)}$, $\bar{\mathcal{F}}$ cannot be maximal k -wise intersecting.*

Combining our method with Theorem 4.2, we obtain the following result.

Proposition 4.3. *For every $k \geq 4$, there exists $c = c(k) > 0$ such that*

$$f(n, k) \geq (1 + c)|\mathcal{F}_{n, k-1}| = (1 + c) \left((k-1)2^{\frac{n}{k-1}} - k + 2 \right),$$

when n is divisible by $k-1$.

Therefore, a maximal k -wise intersecting family of minimum size will necessarily have a more complex structure when $k \geq 4$. It is worth mentioning that the exact value of the upper bound on Janzer's construction [21] is $(k-1)2^{k-3}2^{n/(k-1)} - (k-2)(2^{k-1} - 1)$, which is about $2^{k-3}|\mathcal{F}_{n, k-1}|$.

Acknowledgment

The authors are grateful for Jingwei Xu, Simon Piga and Andrew Treglown, who participated in fruitful discussions at the beginning of the project. Simon Piga and Andrew Treglown's visit to University of Illinois was partially supported by NSF RTG grant DMS 1937241.

References

- [1] N. Alon and P. Frankl, *The maximum number of disjoint pairs in a family of subsets*, Graphs and Combinatorics, **1**, (1985): 13–21.
- [2] N. Alon, S. Das, R. Glebov and B. Sudakov, *Comparable pairs in families of sets*, Journal of Combinatorial Theory, Series B, **115**, (2015): 164–185.
- [3] J. Balogh, N. Bushaw, M. Collares, H. Liu, R. Morris and M. Sharifzadeh, *The typical structure of graphs with no large cliques*, Combinatorica, **37**, (2017): 617–632.
- [4] J. Balogh, C. Chen and H. Luo, *Maximal 3-wise Intersecting Families with Minimum Size: the Odd Case*, arXiv:2206.09334 (2022).
- [5] E. Boros, Z. Füredi and J. Kahn, *Maximal intersecting families and affine regular polygons in $PG(2, q)$* , Journal of Combinatorial Theory, Series A, **52**(1), (1989): 1–9.
- [6] M. Bucić, S. Letzter, B. Sudakov and T. Tran, *Minimum saturated families of sets*, Bulletin of the London Mathematical Society, **50**(4), (2018): 725–732.
- [7] S. J. Dow, D. A. Drake, Z. Füredi and J. A. Larson, *A lower bound for the cardinality of a maximal family of mutually intersecting sets of equal size*, Proceedings of the 16th Southeastern international conference on combinatorics, graph theory and computing, **48**, (1985): 47–48.
- [8] P. Dukes and L. Howard, *Small maximally disjoint union-free families*, Discrete Mathematics, **308**(18), (2008): 4272–4275.
- [9] D. Ellis and B. Sudakov, *Generating all subsets of a finite set with disjoint unions*, Journal of Combinatorial Theory, Series A, **118**(8), (2011): 2319–2345.
- [10] P. Erdős, *On extremal problems of graphs and generalized graphs*, Israel Journal of Mathematics, **2**(3), (1964): 183–190.
- [11] P. Erdős, *Problem sessions*, I. Rival (Ed.), Ordered Sets, Proceedings of the NATO Advanced Study, Reidel, Dordrecht, (1981): 860–861.
- [12] P. Erdős, A. Hajnal and J. W. Moon, *A problem in graph theory*, The American Mathematical Monthly, **71**(1), (1964): 1107–1110.
- [13] P. Erdős and D. J. Kleitman, *Extremal problems among subsets of a set*, Discrete Mathematics, **8**, (1974): 281–294.
- [14] P. Erdős, C. Ko and R. Rado, *Intersection theorems for systems of finite sets*, Quarterly Journal of Mathematics, Oxford Second Series, **12**, (1961): 313–320.

- [15] P. Erdős and L. Lovász, *Problems and results on 3-chromatic hypergraphs and some related questions*, Colloquia Mathematica Societatis János Bolyai 10. Infinite and Finite Sets Keszthely (Hungary), (1973).
- [16] J. Faudree, R. Faudree and J. R. Schmitt, *A survey of minimum saturated graphs*, The Electronic Journal of Combinatorics, **18**, (2011), Dynamic Survey 19, 36 pp.
- [17] N. Frankl, S. Kiselev, A. Kupavskii and B. Patkós, *VC-saturated set systems*, European Journal of Combinatorics, **104**, (2022), Paper No. 103528, 10 pp.
- [18] D. Gerbner, B. Keszegh, N. Lemons, C. Palmer, D. Pálvölgyi and B. Patkós, *Saturating Sperner families*, Graphs and Combinatorics, **29**(5), (2013): 1355–1364.
- [19] R. K. Guy, *A miscellany of Erdős problems*, The American Mathematical Monthly, **90**, (1983): 118–120.
- [20] K. Hendrey, B. Lund, C. Tompkins and T. Tran, *Maximal 3-wise intersecting families*, arXiv:2110.12708v1 (2021).
- [21] B. Janzer, *A note on saturation for k -wise intersecting families*, Combinatorial Theory, **2**(2), (2022): Paper No. 11, 5 pp.
- [22] J. Kahn, *On a problem of Erdős and Lovász*, Journal of the American Mathematical Society, **7**(1), (1994): 125–143.