

Model-Independent Constraints on Non-Unitary Neutrino Mixing from High-Precision Long-Baseline Experiments

Sanjib Kumar Agarwalla,^{a,b,c,d} Sudipta Das,^{a,b} Alessio Giarnetti,^e Davide Meloni^e

^a*Institute of Physics, Sachivalaya Marg, Sainik School Post, Bhubaneswar 751005, India*

^b*Homi Bhabha National Institute, Training School Complex, Anushakti Nagar, Mumbai 400094, India*

^c*International Centre for Theoretical Physics, Strada Costiera 11, 34151 Trieste, Italy*

^d*Department of Physics & Wisconsin IceCube Particle Astrophysics Center, University of Wisconsin, Madison, WI 53706, U.S.A.*

^e*Dipartimento di Matematica e Fisica, Università di Roma Tre
Via della Vasca Navale 84, 00146 Rome, Italy*

E-mail: sanjib@iopb.res.in (ORCID:0000-0002-9714-8866),
sudipta.d@iopb.res.in (ORCID:0000-0002-5508-7751),
giarnetti.alessio@uniroma3.it (ORCID:0000-0001-8487-8045),
davide.meloni@uniroma3.it (ORCID:0000-0001-7680-6957)

ABSTRACT: Our knowledge on the active 3ν mixing angles (θ_{12} , θ_{13} , and θ_{23}) and the CP phase δ_{CP} is becoming accurate day-by-day enabling us to test the unitarity of the leptonic mixing matrix with utmost precision. Future high-precision long-baseline experiments are going to play an important role in this direction. In this work, we study the impact of possible non-unitary neutrino mixing (NUNM) in the context of next-generation long-baseline experiments DUNE and T2HKK/JD+KD having one detector in Japan (T2HK/JD) and a second detector in Korea (KD). We estimate the sensitivities of these setups to place direct, model-independent, and competitive constraints on various NUNM parameters. We demonstrate the possible correlations between the NUNM parameters, θ_{23} , and δ_{CP} . Our numerical results obtained using only far detector data and supported by simple approximate analytical expressions of the oscillation probabilities in matter, reveal that JD+KD has better sensitivities for $|\alpha_{21}|$ and α_{22} as compared to DUNE, due to its larger statistics in the appearance channel and less systematic uncertainties in the disappearance channel, respectively. For $|\alpha_{31}|$, $|\alpha_{32}|$, and α_{33} , DUNE gives better constraints as compared to JD+KD, due to its larger matter effect and wider neutrino energy spectrum. For α_{11} , both DUNE and JD+KD give similar bounds. We also show how much the bounds on the NUNM parameters can be improved by combining the prospective data from DUNE and JD+KD setups. We find that due to zero-distance effects, the near detectors alone can also constrain α_{11} , $|\alpha_{21}|$, and α_{22} in both these setups. Finally, we observe that the ν_τ appearance sample in DUNE can improve the constraints on $|\alpha_{32}|$ and α_{33} .

KEYWORDS: Neutrino, Non-unitary neutrino mixing, Long-baseline, DUNE, T2HK, T2HKK

ARXIV EPRINT: [2111.00329](https://arxiv.org/abs/2111.00329)

Contents

1	Introduction and motivation	1
2	Oscillation probabilities in matter with non-unitary neutrino mixing	4
3	Key features of the experiments	9
3.1	DUNE	9
3.2	T2HK (JD) and T2HKK (JD+KD)	10
4	Simulation details and discussion at the event level	11
5	Numerical results	15
5.1	Correlations in test $(\theta_{23} - \alpha_{ij})$ and test $(\delta_{\text{CP}} - \alpha_{ij})$ planes	15
5.2	Constraints on non-unitary neutrino mixing parameters	17
6	Benefits of having near detectors	21
7	Improvement due to the ν_τ sample in DUNE	25
8	Summary and conclusions	27
A	Analytical expressions of the transition probabilities in the regime of small matter effects	29
B	Impact of marginalization over the other NUNM parameters while showing the constraints on α_{ij}	31

1 Introduction and motivation

The Standard Model (SM) of particle physics is the most successful gauge theory which has been rigorously tested in several laboratory experiments, including those which are being performed at the Large Hadron Collider at CERN in Geneva [1]. In spite of this huge success, the observed input parameters of the SM, such as the Higgs mass, the fermion masses and mixings, and the QCD theta term are quite offbeat and certainly require further explanations [2, 3]. These pressing issues point towards the existence of a microscopic theory beyond the Standard Model which should address the electroweak hierarchy problem [4, 5], the flavor puzzle, and the strong CP problem [6–8]. Various next-generation high-precision experiments in the energy, intensity, and cosmic frontiers are expected to provide crucial information on these issues. In the intensity frontier, several high-precision neutrino oscillation experiments are currently running and also in the pipeline to measure the mass-mixing

parameters with unprecedented precision. Marvelous data from several pioneering neutrino experiments like Super-K [9–11], IceCube-DeepCore [12], ANTARES [13], Daya Bay [14], RENO [15], MINOS [16], Tokai-to-Kamioka (T2K) [17], and NuMI Off-axis ν_e Appearance (NO ν A) [18] have been decisive for our understanding of neutrino flavor mixing. Global fit analyses of the world neutrino data [19–22] have already been able to determine the values of the neutrino oscillation parameters with reasonable accuracy. At present, the relative 1σ precision on the mixing angles θ_{23} , θ_{13} , and θ_{12} lies in the range of $\mathcal{O}(3 - 7)\%$. For the mass-squared differences, the achieved relative 1σ precision is around $\mathcal{O}(1 - 3)\%$. Another important result that is being emerged from these global fit studies is that now we have an overall 1.6σ hint for leptonic CP-violation ($\sin \delta_{\text{CP}} < 0$) [20, 22].

The upcoming medium-baseline reactor experiment Jiangmen Underground Neutrino Observatory (JUNO) [23, 24] is going to address the issue of neutrino mass hierarchy at high confidence level. JUNO will also improve our knowledge on the mixing angle θ_{12} and mass-squared differences Δm_{21}^2 and Δm_{31}^2 . The Fermilab-based Deep Underground Neutrino Experiment (DUNE) [25–28] is expected to resolve the issue of neutrino mass hierarchy and leptonic CP violation with a confidence level never achieved before. DUNE will measure the value of CP phase δ_{CP} and atmospheric oscillation parameters θ_{23} and Δm_{31}^2 with high precision using its wide band neutrino beam, which provides information on oscillation parameters at several L/E values. Another major candidate in the race of next-generation high-precision long-baseline neutrino oscillation experiments is Tokai to Hyper-Kamiokande (T2HK) with one detector in Japan, which we refer as JD in the present work [29, 30]. The possibility of having a second detector in Korea (KD) is also being explored actively. The combination of these two detectors (one in Japan and other in Korea) exposed to a common high-intensity, off-axis (2.5°), narrow-band beam from J-PARC, is widely known as T2HKK [31], which we refer as JD+KD setup in the present study. The JD setup with a relatively shorter baseline as compared to DUNE and high statistics is going to address the issue of leptonic CP violation with unmatched sensitivity and measure the value of δ_{CP} quite precisely without facing the issue of fake CP violation due to matter effect. On the other hand, the KD setup having a baseline roughly around four times that of JD contains some information on Earth matter effect and measure δ_{CP} around the second oscillation maximum with reasonable precision.

With the excellent foreseen precision on the neutrino mass-mixing parameters in the coming years, it will be interesting to explore whether tiny new physics effects beyond the standard three-flavor framework are present in the scenario and what are their impact on the neutrino oscillation probabilities [32]. In the present paper, in the context of high-precision long-baseline experiments, we study the possible consequences of the fact that the 3×3 active neutrino mixing matrix N no longer respects the unitarity condition $N^\dagger N = I$. Theoretically, there are various neutrino mass models in the literature [33–36], which allows the possibility of non-unitary neutrino mixing (NUNM). The most appealing ones are the so-called see-saw models, in which new heavy neutral leptons and/or scalars are introduced in the basic Standard Model to explain tiny neutrino masses. In these models, the standard 3×3 active neutrino mixing matrix becomes a non-unitary sub-matrix of the larger mixing matrix (see discussion in Appendix of Ref. [37]). If the heavy lepton is very massive in the

see-saw model [38–40], then it can cause a departure from the unitarity of the order of 10^{-3} . The amount of non-unitarity can be even larger in the low-scale see-saw models [41, 42].

In this work, our aim is to study the sensitivity of various long-baseline experiments to probe the non-unitarity of the standard 3×3 active neutrino mixing matrix via oscillation, which allows us to constrain the NUNM parameters considering one at a time or all of them at the same time. Here, we adopt a completely model-independent approach to invoke this possible deviation from unitarity without relying on the underlying mechanism.

In the literature, many theoretical/phenomenological studies have been performed to investigate the possible signatures of non-unitarity of the three-flavor neutrino mixing matrix in the context of neutrino oscillation experiments. Considerable efforts have been made to study the possible impact of NUNM on the measurement of leptonic CP phase δ_{CP} , determination of the neutrino mass-hierarchy [43–48], and in estimating the performance of current and future long-baseline experiments to constrain the various NUNM parameters [37, 39, 49–58].

In the present work, we study the potential of two next-generation long-baseline experiments DUNE and T2HKK (JD+KD), to constrain the NUNM parameters in a complete model-independent approach. We present the expected bounds, as obtained from these experiments individually as well as from their combination, assuming no priors in the fit procedure. Compared to other similar studies, we differ in several aspects; first of all, to carry on our numerical simulations, we use the most recent configuration of the DUNE [27] and T2HKK [31] experiments, also investigating the role of near detectors (ND) and τ neutrino detection in the case of DUNE. Then, for a better understanding of the obtained bounds as well as of the observed correlations between the standard oscillation parameters θ_{23} and δ_{CP} with the various NUNM parameters, we provide simple and useful analytical expressions of the relevant transition probabilities in the regime of small matter effects, which is a correct approximation for the neutrino facilities under investigation.

The manuscript is organized as follows. In Sec. 2, we discuss the formalism and the parameterization of non-unitary neutrino mixing used in this work, and derive simple approximate analytical expressions of the oscillation probability in $\nu_\mu \rightarrow \nu_e$ and $\nu_\mu \rightarrow \nu_\mu$ oscillation channels in the presence of standard matter interaction. In Sec. 3, we give a brief description of the future long-baseline experiments discussed in this work. Sec. 4 provides the details of the numerical simulations performed in our analysis and some results on the expected signal events. The main results of our study are presented in Sec. 5, where we illustrate the correlation of the standard oscillation parameters θ_{23} and δ_{CP} with various NUNM parameters and give the expected constraints achievable by DUNE and T2HKK separately as well as their combination. In Sec. 6, we show the possible improvements in the bounds due to the presence of near detectors. Section 7 shows the improved constraints on the NUNM parameters when we add ν_τ events sample in the DUNE simulation. Our concluding remarks are discussed in Sec. 8. Two appendices complete our work: in Appendix A, we compare our analytical expressions of the oscillation probabilities and those computed numerically in the presence of non-unitarity. We also provide the analytical expressions of the oscillation probabilities for all the remaining channels. In Appendix B, we discuss in detail the impact on the bounds discussed in the main text of

α_{11}	α_{22}	α_{33}	$ \alpha_{21} $	$ \alpha_{31} $	$ \alpha_{32} $
< 0.031	< 0.005	< 0.110	< 0.013	< 0.033	< 0.009

Table 1: 90% confidence level (C.L.) limits on the NUNM parameters using data from various short-baseline and long-baseline experiments, as obtained from the Ref. [53].

the simultaneous marginalization over all NUNM parameters.

2 Oscillation probabilities in matter with non-unitary neutrino mixing

The non-unitarity of the PMNS matrix can be parameterized in different ways [49, 56, 59–62]. One possibility, which turned out to be very useful in oscillation analysis, consists of factorizing the deviation from unitarity into a matrix α in such a way that the non-unitary neutrino mixing matrix N is expressed as¹:

$$N = (I + \alpha) U_{PMNS}. \quad (2.1)$$

In order to compare our numerical results with those already presented in the literature, we adhere here at the widely used lower triangular structure of the matrix α , containing nine free parameters organized as follows:

$$\alpha = \begin{pmatrix} \alpha_{11} & 0 & 0 \\ |\alpha_{21}|e^{i\phi_{21}} & \alpha_{22} & 0 \\ |\alpha_{31}|e^{i\phi_{31}} & |\alpha_{32}|e^{i\phi_{32}} & \alpha_{33} \end{pmatrix}. \quad (2.2)$$

This parameterization simplifies the oscillation probabilities and let the parameter α_{ij} to be the main source of non-unitarity for the oscillation channel $\nu_i \rightarrow \nu_j$ ($i, j = e, \mu, \tau$).

Bounds on the α_{ij} parameters have been recently computed, among others, in Ref. [53] and reported for convenience in Table 1. These results have been obtained using data from the short-baseline experiments NOMAD and NuTeV, and the long-baseline experiments MINOS/MINOS+, T2K, and NO ν A. For the off-diagonal NUNM parameters, the authors also used the triangular inequalities² $\alpha_{ij} \leq \sqrt{1 - (1 + \alpha_{ii})^2} \sqrt{1 - (1 + \alpha_{jj})^2}$ [40], which appear due to the assumption that the standard 3 \times 3 active neutrino mixing matrix is a non-unitary sub-matrix of a larger unitary mixing matrix. Note that in the present study, we do not take into account these inequalities in order to study the capability of long-baseline experiments alone to put bounds on these NUNM parameters in a model-independent fashion.

¹We choose the convention in which the matrix α , which invokes non-unitarity, is added to the identity matrix. Note that other phenomenological studies adopt the relation $N = (1 - \alpha) U_{PMNS}$ [39]. Our results can be compared to the others just changing the sign in front of the diagonal elements.

²Note that in this model, the diagonal parameters can only be negative.

θ_{23}	θ_{13}	θ_{12}	δ_{CP}	$\Delta m_{21}^2 [\text{eV}^2]$	$\Delta m_{31}^2 [\text{eV}^2]$
45°	8.61°	33.6°	-90°	7.39×10^{-5}	2.52×10^{-3}

Table 2: The benchmark values of the oscillation parameters used in our analysis. These values are consistent with the present best-fit values as obtained in various global fit studies [19–22]. We assume normal mass hierarchy (NH) of neutrino throughout this work.

Taking into account the expression in Eq. 2.1, the complete effective neutrino propagation Hamiltonian in the mass-eigenstate is:

$$H = \frac{1}{2E_\nu} \left[\begin{pmatrix} 0 & 0 & 0 \\ 0 & \Delta m_{21}^2 & 0 \\ 0 & 0 & \Delta m_{31}^2 \end{pmatrix} + N^\dagger \begin{pmatrix} a_e + a_n & 0 & 0 \\ 0 & a_n & 0 \\ 0 & 0 & a_n \end{pmatrix} N \right]. \quad (2.3)$$

As usual, the matter potential parameters are given by $a_e = 2\sqrt{2}E_\nu G_F N_e$ and $a_n = -\sqrt{2}E_\nu G_F N_n$ where, N_e and N_n are the electron and the neutron number densities, respectively. Note that in this framework, the neutral current (NC) matter potential is necessary since the non-unitarity of the matrix N does not allow the subtraction of an identity matrix proportional to a_n . From the Schrodinger equation, the transition probability at a given baseline L is obtained from the following expression:

$$P_{\alpha\beta} = |(N e^{-iHL} N^\dagger)_{\beta\alpha}|^2. \quad (2.4)$$

The relevant probabilities for long-baseline experiments are the $\nu_\mu \rightarrow \nu_e$ appearance and $\nu_\mu \rightarrow \nu_\mu$ disappearance channels. We will discuss these probabilities here in details, while for the sake of completeness, we also quote the τ appearance and electron disappearance into the Appendix A.

In order to get approximate analytical expressions for the transition probabilities, we observe that the vacuum approximation cannot be sufficiently precise in experiments like DUNE, since the matter effects can modify the appearance probability up to about 10%. For this reason, we derive approximate analytical expressions in the presence of matter. We use perturbation theory in the small expansion parameters (r , s , and a) defined as follows:

$$\sin \theta_{13} = \frac{r}{\sqrt{2}}, \quad \sin \theta_{12} = \frac{1}{\sqrt{3}}(1 + s), \quad \sin \theta_{23} = \frac{1}{\sqrt{2}}(1 + a), \quad (2.5)$$

where, r , s , and a represent the deviation from the tri-bimaximal mixing values of the neutrino mixing parameters, namely, $\sin \theta_{13} = 0$, $\sin \theta_{23} = 1/\sqrt{2}$, and $\sin \theta_{12} = 1/\sqrt{3}$ [63, 64]. Given the recent global fit of neutrino oscillation data, we can assume that $r, s, a \sim \mathcal{O}(0.1)$ and we can further expand them up to the second order [19–22]. To further simplify the notation, we also introduce $\Delta_{31} = \Delta m_{31}^2 L/4E_\nu$, $\Delta_e = a_e L/4E_\nu$ and $\Delta_n = a_n L/4E_\nu$; at the DUNE peak energy, namely, $E_\nu = 2.5$ GeV, $\Delta_e \sim 0.36$ and $\Delta_n \sim 0.18$, we can further expand in the small matter potentials up to the first order. Note that for the other experimental facilities discussed in this paper, this approximation is even better; in fact, for the Tokai to Hyper-Kamiokande (T2HK) setup with a far detector in Japan

(JD), at beam energy of $E_\nu = 0.6$ GeV, we have $\Delta_e \sim 0.08$ and $\Delta_n \sim 0.04$, while with a second in Korea (KD) at a distance of 1100 km from the source with same energy, we get $\Delta_e \sim 0.30$ and $\Delta_n \sim 0.15$. Also, we use one mass scale dominance (OMSD) approximation ($\Delta_{31} \gg \Delta_{21}$, where $\Delta_{21} = \Delta m_{21}^2 L / 4E_\nu$) in our derivation; which is a valid approximation in the atmospheric regime.

In the case of the $\nu_\mu \rightarrow \nu_e$ appearance probability, we thus obtain:

$$\begin{aligned}
P_{\mu e} = & \left(\frac{r^2}{\Delta_{31}} \right) \sin \Delta_{31} [(\Delta_{31} + 2\Delta_e) \sin \Delta_{31} - 2\Delta_{31}\Delta_e \cos \Delta_{31}] + \\
& \left(\frac{2|\alpha_{31}|r}{\Delta_{31}} \right) \Delta_n \sin \Delta_{31} [\cos(\delta_{\text{CP}} - \phi_{31}) \sin \Delta_{31} - \Delta_{31} \cos(\delta_{\text{CP}} + \Delta_{31} - \phi_{31})] + \\
& \left(\frac{|\alpha_{21}|r}{\Delta_{31}} \right) \{ \sin \Delta_{31} [2\Delta_{31}(\Delta_e + \Delta_n) \cos(\delta_{\text{CP}} + \Delta_{31} - \phi_{21}) - \Delta_n \sin(\delta_{\text{CP}} - \Delta_{31} - \phi_{21})] + \\
& \sin(\delta_{\text{CP}} + \Delta_{31} - \phi_{21}) [(-2\Delta_{31} - 2\Delta_e + \Delta_n) \sin \Delta_{31} + 2\Delta_{31}\Delta_e \cos \Delta_{31}] \} + \\
& \left(\frac{|\alpha_{21}||\alpha_{31}|}{\Delta_{31}} \right) \Delta_n [-2\Delta_{31} \sin(\phi_{21} - \phi_{31}) + \cos(2\Delta_{31} - \phi_{21} + \phi_{31}) - \cos(\phi_{21} - \phi_{31})] + \\
& \left(\frac{|\alpha_{21}|^2}{\Delta_{31}} \right) [\Delta_{31} - \Delta_n(1 - \cos 2\Delta_{31})] . \tag{2.6}
\end{aligned}$$

From the above expression, it is clear that the $\nu_\mu \rightarrow \nu_e$ appearance probability strongly depends on $|\alpha_{21}|$ and $|\alpha_{31}|$. The parameter $|\alpha_{21}|$ survives in the vacuum case while $|\alpha_{31}|$ always appears with the matter potential Δ_n . This essentially means that an experiment in which the matter effect is not negligible is able to put strong bounds also to $|\alpha_{31}|$ which, otherwise, would not be accessible by $P_{\mu e}$. Note that due to the loss of unitarity property of the neutrino mixing, some terms remain non-zero in $\nu_\mu \rightarrow \nu_e$ appearance probability expression even when the neutrino propagation length L is zero, which is known as zero-distance effect:

$$P_{\mu e}^{L=0} \sim |\alpha_{21}|^2 . \tag{2.7}$$

So, it is clear that even the near detectors (ND) of long-baseline experiments could contribute to the bounds of non-unitarity parameters, as it will be discussed in Sec. 6. Finally, we point out that the vacuum limit of Eq. 2.6 assumes a particularly simple expression:

$$P_{\mu e}^{\text{vac}} = |\alpha_{21}|^2 + r^2 \sin^2 \Delta_{31} - 2|\alpha_{21}|r \sin \Delta_{31} \sin(\delta_{\text{CP}} + \Delta_{31} - \phi_{21}) , \tag{2.8}$$

which, in the limit of vanishing solar mass difference, agrees with the results presented in Ref. [37].

In order to assess the modifications in the $\nu_\mu \rightarrow \nu_e$ appearance probabilities caused by the presence of non-unitary neutrino mixing, in Fig. 1, we show the exact $\nu_\mu \rightarrow \nu_e$ appearance probabilities as a function of energy, obtained numerically using the General long-baseline Experiment Simulator (GLoBES) software [65, 66]. In the extreme left column, we consider a baseline of 1300 km for DUNE. In the middle column, we show the results for the JD baseline of 295 km. In the extreme right column, we deal with the KD setup having a baseline of 1100 km. Note that for both JD and KD, we consider the




Figure 1: $\nu_\mu \rightarrow \nu_e$ appearance probability as a function of energy in the presence of off-diagonal NUNM parameters. Left, middle, and right columns correspond to the baselines of 1300 km (DUNE), 295 km (JD), and 1100 km (KD), respectively. The four colored curves correspond to four benchmark values of the phases associated with off-diagonal NUNM parameters: 0° , 90° , 180° , and -90° . We consider $\delta_{\text{CP}} = 0^\circ$ and $\sin^2 \theta_{23} = 0.5$. The values of the other oscillation parameters are taken from Table 2.

neutrino energy range of 0 to 1.5 GeV having a peak around 0.6 GeV. In every rows, we switch-on one off-diagonal NUNM parameters at a time, while maintaining others to zero. The effect of $|\alpha_{21}|$ is shown in the top panels, $|\alpha_{31}|$ in the middle panels, and $|\alpha_{32}|$ in the bottom panels. In each panel, the solid black curves correspond to the probabilities in the unitary neutrino mixing (UNM) case, while the colored curves correspond to the NUNM cases with four benchmark values of the phases associated with each off-diagonal NUNM parameters, as reported in the legend. As we can see, the impact of the NUNM parameter $|\alpha_{21}|$ (top panels) is comparatively larger than the other two off-diagonal NUNM parameters $|\alpha_{31}|$ and $|\alpha_{32}|$ even though the strength of the $|\alpha_{21}|$ is much smaller than the other two. This feature is more clear for JD in the middle columns. In fact, from the approxi-




Figure 2: $\nu_\mu \rightarrow \nu_\mu$ disappearance probabilities as a function of neutrino energy in the presence of the NUNM parameters α_{22} and $|\alpha_{32}|$ assuming $\phi_{32} = 0$ one at a time. Left, middle, and right columns correspond to the baselines of 1300 km (DUNE), 295 km (JD), and 1100 km (KD), respectively. We consider $\delta_{\text{CP}} = 0^\circ$ and $\sin^2 \theta_{23} = 0.5$. The values of the other oscillation parameters are taken from Table 2.

mated analytical expression of $\nu_\mu \rightarrow \nu_e$ appearance probability in Eqs. 2.6 and 2.8, we see that only terms containing $|\alpha_{21}|$ survive in vacuum, while the effect of $|\alpha_{31}|$ and $|\alpha_{32}|$ is linked to Δ_e and Δ_n which are very small at the considered baseline. This is not the case for DUNE ($L \simeq 1300$ km) where, given the largest baseline under consideration, Δ_e and Δ_n are no longer negligible and the impact of $|\alpha_{31}|$ and $|\alpha_{32}|$ on $P_{\mu e}$ is of the same order as $|\alpha_{21}|$.

For the $\nu_\mu \rightarrow \nu_\mu$ disappearance channel, we get:

$$\begin{aligned}
P_{\mu\mu} = & \cos^2 \Delta_{31} (1 + 4\alpha_{22}) - 2|\alpha_{32}| \Delta_n \sin 2\Delta_{31} \cos \phi_{32} + 4a^2 \sin^2 \Delta_{31} + \\
& 2|\alpha_{21}|^2 \cos \Delta_{31} [2 \sin \Delta_{31} (\Delta_e + \Delta_n) + \cos \Delta_{31}] + 6\alpha_{22}^2 \cos^2 \Delta_{31} - \\
& 2|\alpha_{21}| r \sin(2\Delta_{31}) (\Delta_e + \Delta_n) \cos(\delta_{\text{CP}} - \phi_{21}) + \\
& \left(\frac{8a}{\Delta_{31}} \right) (\alpha_{22} - \alpha_{33}) \Delta_n \sin \Delta_{31} (\sin \Delta_{31} - \Delta_{31} \cos \Delta_{31}) \\
& - 2|\alpha_{21}| |\alpha_{31}| \Delta_n \sin 2\Delta_{31} \cos(\phi_{21} - \phi_{31}) - 8\alpha_{22} |\alpha_{32}| \Delta_n \sin 2\Delta_{31} \cos \phi_{32} \\
& - 2|\alpha_{32}| \alpha_{33} \Delta_n \sin 2\Delta_{31} \cos \phi_{32}, \tag{2.9}
\end{aligned}$$

from which we learn that all the NUNM parameters α_{ij} but α_{11} enter into the probability expression. Also, in this case, we get the zero-distance expression of the $\nu_\mu \rightarrow \nu_\mu$ survival probability, given by:

$$P_{\mu\mu}^{L=0} \sim 1 + 2|\alpha_{21}|^2 + 6\alpha_{22}^2 + 4\alpha_{22}, \tag{2.10}$$

and the vacuum approximation (also in agreement with Ref. [37] in the limit of vanishing Δm_{21}^2):

$$P_{\mu\mu}^{\text{vac}} = \cos^2 \Delta_{31} (1 + 2|\alpha_{21}|^2 + 4\alpha_{22} + 6\alpha_{22}^2) + 4a^2 \sin^2 \Delta_{31}. \tag{2.11}$$

In Fig. 2, we show the exact $\nu_\mu \rightarrow \nu_\mu$ oscillation probabilities as a function of energy for the baseline lengths corresponding to DUNE (left panel), JD (middle panel), and KD (right panel) setups. In each panel, the black solid curves correspond to the UNM case, while the red and blue curves show the presence of α_{22} and α_{32} , respectively with strength reported in the legend³, one at a time. The impact of these two NUNM parameters can be understood from our approximated analytical expressions in Eq. 2.9. When matter effects are negligible (for example, in the middle panel of Fig. 2), we expect that the parameter α_{22} dominates the deviation from UNM since it appears already at first order in $P_{\mu\mu}$. This remains true when matter parameters are switched-on; the relevant difference compared to the vacuum case relies on the fact that also $|\alpha_{32}|$ enter at first order, although suppressed by Δ_n . Thus, we expect that for DUNE and KD, one can see deviation from the UNM predictions, as visible in Fig. 2. Note that the impact of α_{32} is amplified by the larger benchmark value compared to the choice for α_{22} .

3 Key features of the experiments

Long-baseline experiments provide a clear environment to study the phenomenology of neutrino oscillations. Indeed, the possibility to fix the baseline and have a focused neutrino beam allow us to have different features of the experiments under control. The next-generation long-baseline experiments will increase the neutrino events statistics with the help of high precision detectors. Two experiments currently under construction are DUNE and T2HK. For the latter is being taken into account the possibility of a second far detector placed approximately at the second oscillation maximum. These two complementary experiments, which will be described in more details in the following subsections, are going to improve the measurements of the oscillation parameters and the searches of new physics effects in a significant way. In the non-unitarity framework, the unprecedented statistics collected by the detectors of the two experiments, as well as the possibility to look for ν_μ disappearance, ν_e appearance but also ν_τ appearance in DUNE, will provide the chance to bound at a very good level all the non-standard parameters.

3.1 DUNE

DUNE (Deep Underground Neutrino Experiment) is a next-generation long-baseline experiment which will play an important role in solving the existing puzzles in neutrino oscillation as well as in increasing the precision on the measured neutrino oscillation parameters. It will use an on-axis, high-intensity, wide-band neutrino beam traveling a distance of 1284 km from Fermilab to South Dakota. For this baseline, they consider an average matter density of 2.85 g/cm^3 . The far detector (FD) is a liquid argon time projection chamber (LArTPC) of 40 kt fiducial mass placed underground at the Homestake mine. According to the recent Technical Design Report (TDR) [27], DUNE will use a proton beam having 1.2 MW power which will deliver 1.1×10^{21} proton on target (P.O.T.) per year. DUNE considers a total run-time of 10 years with 5 years each in neutrino and antineutrino modes.

³Even though the analytical expression of $P_{\mu\mu}$ reported in Eq. 2.9 shows the presence of other NUNM parameters, we have numerically checked that they do not have any significant impact.

However, following most of the existing studies related to DUNE, we present our results using a total run-time of 7 years with 3.5 years each in neutrino and antineutrino modes. Beside the far detector, the possibility to have three modules of near detectors (ND) has been proposed [67]. The first one would be a liquid Argon Time Projection chamber (TPC) situated at a distance of 574 m from the source. It will measure the flux and cross-section of the neutrinos. The second one would be a multi purpose detector (MPD) equipped with a magnetic spectrometer with one ton High-Pressure Gaseous Argon TPC, which will be useful to study possible new physics signals. The third one would be the System for On-Axis Neutrino Detection (SAND). It will be made up of a former KLOE magnet and a calorimeter which will track the outgoing particles.

3.2 T2HK (JD) and T2HKK (JD+KD)

T2HK (Tokai to Hyper-Kamiokande) is another promising next-generation long-baseline experiment which will play a very important role in δ_{CP} measurement as well as in the study of various BSM physics [68–70]. In the T2HK setup [29, 71, 72], an intense neutrino beam from the J-PARC proton synchrotron facility will be detected at the Hyper-Kamiokande (HK) detector situated at a distance of 295 km from the source. The power of the proton beam at the source is 1.3 MW which will produce 27×10^{21} P.O.T. in its total 10 years run-time. The detector is a water Cherenkov (WC) detector with 187 kt of fiducial mass. To have an equal contribution from the neutrino and the antineutrino signal events, the proposed running time of the experiment is 2.5 years (25% exposure) in neutrino mode and 7.5 years (75% exposure) in antineutrino mode. The HK detector will be placed at 2.5° off-axis angle from the neutrino beam-line to receive a narrow band beam with energy peaked at around the first oscillation maximum (~ 0.6 GeV).

There is also a proposal [73] to have another identical detector which will be situated in Korea, 1100 km far from J-PARC. We assume that this detector will also be placed at an off-axis angle of 2.5° from the neutrino beamline (similar to JD) and operate around the second oscillation maximum with a baseline of 1100 km and an average neutrino energy of 0.6 GeV. Combination of the T2HK setup along with the detector in Korea is known as T2HKK. However, in our work, we call the Hyper-Kamiokande as the Japanese detector (JD) and the detector in Korea as the Korean detector (KD). For both JD and KD baselines, we assume an average matter density of 2.8 g/cm^3 . In the following, we will show our numerical results on the sensitivity to the various NUNM parameters separately for each detectors as well as for their combination.

In addition to these two detectors, two more near detectors, namely, the ND280 detector and Intermediate Water Cherenkov detector (IWCD), have been proposed. ND280 will be located at 280 m from the source with the same off-axis angle as the far detectors. It will be helpful to measure the flux of the unoscillated neutrino beam and study the neutrino cross-sections [74]. IWCD is a water Cherenkov detector of mass of 1 kt and possibly located at a distance of 1 km from the source [75, 76]. One advantage of this detector is that it can be moved vertically to take data at different off-axis angles. In Table 3, we summarize the relevant information of all the experimental setups discussed in this section.

	DUNE	JD/KD
Detector Mass	40 kt LArTPC	187 kt WC (each)
Baseline	1300 km	295/1100 km
Proton Energy	120 GeV	80 GeV
Beam type	Wide-band, on-axis	Narrow-band, off-axis (2.5°)
Beam power	1.2 MW	1.3 MW
P.O.T./year	1.1×10^{21}	2.7×10^{21}
Run time ($\nu + \bar{\nu}$)	3.5 yrs + 3.5 yrs	2.5 yrs + 7.5 yrs

Table 3: Essential features of DUNE [27] and JD/KD [31] experiments used in our simulation.

4 Simulation details and discussion at the event level

In order to perform our numerical simulations, we use the General long-baseline Experiment Simulator (GLOBES) package [65, 66] along with the plug-in MonteCUBES [39]. For the standard oscillation parameters, we use the value as given in Table 2. For the simulation of the DUNE experiment, we use the most recent GLOBES configuration files as given in Ref. [27]. The simulation details of T2HK and T2HKK setups are taken from Ref. [31]. In Table 3, we summarize the relevant information of the two experiments considered in our simulation. Note that while showing the sensitivity of the FDs in constraining various NUNM parameters, we do not take into account the NDs in our simulation. The only exception is in Sec. 6, where we discuss the possible improvement in the bounds due to presence of near detectors without considering any possible correlations among FDs and NDs (see Sec. 6 for further details). Both DUNE and T2HKK are expected to have access to two oscillation channels, namely the ν_μ ($\bar{\nu}_\mu$) disappearance and the ν_e ($\bar{\nu}_e$) appearance channels. In particular:

- for DUNE, backgrounds to the appearance channel are the ν_e beam contamination and the misidentified ν_μ , ν_τ , and NC events. The signal systematic normalization error have been chosen to be 2%. For the disappearance channel, background to the signal are misidentified ν_τ and NC events. The signal error is 5%. Efficiency functions as well as smearing matrices have been provided by the DUNE collaboration in Ref. [27];
- for the two T2HKK far detectors JD and KD, background in the appearance channels are the ν_e from the beam contamination and misidentified ν_μ and NC events. Signal systematic uncertainties are 5% normalization and 5% calibration errors. In the disappearance channel, backgrounds to the signal are misidentified ν_e and NC events. Systematic uncertainties are 3.5% normalization and 5% calibration errors. Efficiencies and energy resolutions are taken from the Ref. [31].

The number of expected signal events for both channels simulated here are summarized in Tables 4 and 5, where the cases of UNM and NUNM (for some benchmark values of the α_{ij} parameters) are reported. The impact of NUNM parameters on the number of events

	ν_e appearance			$\bar{\nu}_e$ appearance		
	DUNE	JD	KD	DUNE	JD	KD
UNM	1259	1836	169	221	767	31
NUNM	DUNE	JD	KD	DUNE	JD	KD
$ \alpha_{21} $ (= 0.025)	1328	1893	169	232	756	29
$ \alpha_{31} $ (= 0.1)	1420	1893	187	264	754	30
$ \alpha_{32} $ (= 0.25)	1300	1855	172	213	756	30
α_{11} (= -0.02)	1203	1761	162	214	738	29
α_{22} (= -0.015)	1223	1782	164	215	744	30
α_{33} (= -0.15)	1208	1817	168	227	779	30

Table 4: Comparison of the total signal rate for the ν_e and $\bar{\nu}_e$ appearance channels in DUNE, JD, and KD setups in UNM case as well as in presence of various NUNM parameters. The relevant features of these facilities are given in Table 3. The values of the standard oscillation parameters used to calculate event rate are quoted in Table 2. The phases associated with the off-diagonal NUNM parameters are considered to be zero.

is fully in agreement with our analytical discussions. First of all, as shown in Eq. 2.6, the appearance channel is mainly influenced by $|\alpha_{21}|$, even in vacuum. This reflects in an enhancement of the number of events by roughly 5% in both JD and DUNE. On the other hand, the NUNM parameters $|\alpha_{31}|$ and α_{33} are also relevant but they are coupled to the matter potentials, so we expect them to be relevant primarily for DUNE, where matter effects are more important: in fact, $|\alpha_{31}|$ causes an increase in the number of events up to 10%, while α_{33} provokes a small but visible reduction of the order of 4%. Finally, some impact on the number of signal events is also given by α_{11} , even though it only appears at higher orders in our perturbative expansion and has not been displayed (but it present in the vacuum probabilities reported in Ref. [37]). Note that the number of ν_e and $\bar{\nu}_e$ events in KD is only slightly influenced by the NUNM parameters due to fact that the experiment works close to the second oscillation maximum of the atmospheric oscillation ($\nu_\mu \rightarrow \nu_\tau$), where the $\nu_\mu \rightarrow \nu_e$ appearance probability approached one of its minima and the effects of new physics are suppressed.

For the disappearance channel, the parameter α_{22} , which enters at the first perturbative order in Eq. 2.9, produces a reduction of about 6% in the number of events for all three detectors. This can be roughly understood from the fact that the standard disappearance probability is multiplied by $4\alpha_{22} = 0.06$, which causes a reduction by a similar factor in the number of events. The other relevant NUNM parameter is $|\alpha_{32}|$ which, being coupled to matter potential in Eq. 2.9, can cause a $\sim 6\%$ increase of events especially in DUNE. The other parameters at their benchmark values only have a negligible impact on the number of disappearance events.

In Fig. 3, we show the relation between the appearance events in neutrino and antineutrino modes at the three detectors discussed in this paper, namely DUNE (left panel),

	ν_μ disappearance			$\bar{\nu}_\mu$ disappearance		
	DUNE	JD	KD	DUNE	JD	KD
UNM	10359	9064	1266	6034	8625	1144
NUNM	DUNE	JD	KD	DUNE	JD	KD
$ \alpha_{21} $ (= 0.025)	10371	9074	1264	6045	8640	1149
$ \alpha_{31} $ (= 0.1)	10351	9062	1261	6035	8627	1168
$ \alpha_{32} $ (= 0.25)	10978	9203	1255	6005	8467	1158
α_{11} (= -0.02)	10359	9064	1266	6034	8625	1145
α_{22} (= -0.015)	9748	8531	1192	5681	8120	1077
α_{33} (= -0.15)	10406	9077	1268	6040	8619	1145

Table 5: Comparison of the total signal rate for the ν_μ and $\bar{\nu}_\mu$ disappearance channels in DUNE, JD, and KD setups in UNM case as well as in presence of various NUNM parameters. The relevant features of these facilities are given in Table 3. The values of the standard oscillation parameters used to calculate event rate are quoted in Table 2. The phases associated with the off-diagonal NUNM parameters are considered to be zero.

JD (middle panel), and KD (right panel). The cyan (black) ellipses in each panel correspond to the standard interaction case (obtained varying the CP phase δ_{CP} in the range $[-180^\circ, 180^\circ]$), referring to the normal (inverted) mass hierarchy. The red (blue) colored blobs in the upper panels show the bi-events in the presence of the non-unitarity parameters $|\alpha_{21}|$ with strength 0.025 for the NH (IH) case. The lower panels show the same but in the presence of $|\alpha_{31}|$ fixed to 0.1. In both cases, CP phase δ_{CP} and the phase associated with corresponding off-diagonal NUNM parameter are varied in the range -180° to 180° . The plots show that DUNE (left panels) is in principle expected to be able to distinguish the mass hierarchy even in presence of NUNM. On the other hand, the two standard model curves and the two shadowed regions overlap in the case of JD (center panels) and KD (right panels). For these, the amplitude of the shadowed regions when ϕ_{31} is varied is much smaller than the DUNE one since, as it is clear from the analytical formulas, $|\alpha_{31}|$ is always coupled to matter effects, which are smaller for the two J-PARC based experiments.

In our numerical simulations on the sensitivity to the NUNM parameters, the true values of the standard oscillation parameters have been chosen as in Table 2, while the true values of the NUNM parameters are set to zero. Since the presence of non-unitarity has basically the same effect on the number of events in the case of NH and IH, as shown in Fig. 3, we consider only the NH case. The computation of the $\Delta\chi^2$ is based on the pull method [77–79] implemented in the GLoBES software. We study the NUNM parameters by fixing the mixing angles θ_{12} , θ_{13} , and two mass-squared differences Δm_{21}^2 , and Δm_{31}^2 both in data and theory at their best fit values as given in Table 2. We check that the marginalization over the atmospheric mass-squared difference Δm_{31}^2 does not have any significant effect on our analysis. On the other hand, the only notable effect of the marginalization over the reactor mixing angle θ_{13} (which has a very small experimental




Figure 3: Bi-event plots for the DUNE (left panel), JD (middle panel), and KD (right panel). The ellipses in each panel are obtained by varying the CP phase δ_{CP} in the range $[-180^\circ, 180^\circ]$ assuming $\alpha_{ij} = 0$. The colored blobs in the upper panels show the bi-events in the presence of $|\alpha_{21}|$ with a strength of 0.025, while the lower panels depict the impacts of $|\alpha_{31}|$ with a strength of 0.1. To obtain the blobs with non-zero α_{ij} , we vary the associated NUNM phases and the CP phase δ_{CP} in their allowed range of -180° to 180° . The values of the other standard three-flavor oscillation parameters are taken from Table 2.

uncertainty of 3%), is the worsening of the α_{11} bound at the level of 15%. This is due to the fact that there is a correlation between these two parameters, which appear in a term proportional to $\alpha_{11}^2 \sin^2 2\theta_{13}$ in the $\nu_\mu \rightarrow \nu_e$ transition probability as shown in Ref. [37]. Finally, we marginalized θ_{23} in its current 3σ allowed range [22], which is approximately $[40^\circ, 50^\circ]$ and the CP phase δ_{CP} in its entire possible range $[-180^\circ, 180^\circ]$. We keep both these parameters with true values as in Table 2. Moreover, we consider one NUNM parameter at a time, *i.e.*, when a parameter is taken into account the others are considered to be zero. In Appendix B, we also discuss in detail how our results would be affected by the marginalization over multiple NUNM parameters. Finally, the phases associated with each off-diagonal NUNM parameters are marginalized over the entire possible range from -180° to 180° .

5 Numerical results

In this section, we provide a detailed discussion of the numerical results obtained with the GLOBES simulations. We study the correlations between the NUNM parameters and the two less constrained standard oscillation parameters θ_{23} and δ_{CP} . Then, we determine the expected bounds on all the non-standard parameters at 90% confidence level (C.L.) when the expected data from DUNE, JD, KD, and their various combinations are taken into account.

5.1 Correlations in test $(\theta_{23} - \alpha_{ij})$ and test $(\delta_{\text{CP}} - \alpha_{ij})$ planes




Figure 4: 1σ (2 d.o.f.) C.L. contours in the $(\theta_{23} - \alpha_{ij})$ planes for DUNE (red curves) and JD+KD (blue curves). The true values of θ_{23} considered here are $\theta_{23} = 40^\circ$, 45° , and 50° (shown by the black dots in each panel). True value of δ_{CP} is considered to be -90° . In the fit, δ_{CP} and the phases associated with off-diagonal NUNM parameters (see lower panels) are marginalized over the range of $[-180^\circ, 180^\circ]$. The other oscillation parameters are fixed at their best fit values (see Table 2).

In Fig. 4, we show the correlation between various NUNM parameters and the atmospheric mixing angle θ_{23} . Three different true values of θ_{23} (shown by the black dot in each panel) have been analyzed, namely, $\theta_{23} = 40^\circ$ ($\sin^2 \theta_{23} = 0.413$) in the lower octant, the




Figure 5: 1σ (2 d.o.f.) C.L. contours in the $(\delta_{\text{CP}} - \alpha_{ij})$ planes for DUNE (red curves) and JD+KD (blue curves). The true values of δ_{CP} considered here are $\delta_{\text{CP}} = 0^\circ, 90^\circ, 180^\circ, -90^\circ$ (shown by the black dots in each panel). The true value of θ_{23} is 45° , while its test value is kept free in the range $[40^\circ, 50^\circ]$. In the fit, the phases associated with off-diagonal NUNM parameters (see lower panels) are marginalized over the range of $[-180^\circ, 180^\circ]$. The other oscillation parameters are fixed at the values as given in Table 2.

maximal value $\theta_{23} = 45^\circ$ ($\sin^2 \theta_{23} = 0.5$), and $\theta_{23} = 50^\circ$ ($\sin^2 \theta_{23} = 0.586$) in the upper octant. We keep the true value of δ_{CP} as -90° . Red contours correspond to 1σ (2 d.o.f.) C.L. obtained using the DUNE setup, while the blue contours show the same for the JD+KD combination. We observe that the correlation between $\sin^2 \theta_{23}$ and α_{11} (top left panel) is almost the same for both DUNE and JD+KD setups. In the $(\theta_{23} - \alpha_{22})$ and $(\theta_{23} - \alpha_{21})$ planes, the allowed regions for JD+KD are smaller than the ones corresponding to DUNE, which suggest that the J-PARC based experiments will have a better sensitivity on these parameters (see the following subsection). For the other three NUNM parameters, DUNE shows much better sensitivity than JD+KD, due to the larger matter effects which couples to the NUNM parameters α_{33} , $|\alpha_{31}|$, and $|\alpha_{32}|$. In particular, for maximal value of θ_{23} , there are no lower limits on α_{33} from JD+KD. One interesting point to be noted is that, in the presence of non-unitary mixing, $\sin^2 \theta_{23}$ can be constrained better for non-maximal true values of θ_{23} .

In Fig. 5, we show the correlation of various NUNM parameters with the CP phase δ_{CP} . In each panel, the black dots correspond to the true values of $\delta_{\text{CP}} = 0^\circ, 90^\circ, 180^\circ,$ and -90° and the benchmark value chosen for the α_{ij} under consideration. The red contours refer to the sensitivity at 1σ (2 d.o.f.) C.L. expected to be obtained by DUNE setup and the blue curve shows the same for JD+KD combination. The plots follow a similar behavior already seen in Fig. 4. In particular, the allowed regions for the two experiments are almost the same in the case of α_{11} , smaller for JD+KD in the case of α_{22} and $|\alpha_{21}|$ and, finally, much smaller for DUNE in the case of the NUNM parameters α_{33} , $|\alpha_{31}|$, and $|\alpha_{32}|$. Note that when true $\delta_{\text{CP}} = 90^\circ, 180^\circ,$ and -90° , the J-PARC based detectors will not be able to set any lower limit on α_{33} .

5.2 Constraints on non-unitary neutrino mixing parameters

In this subsection, we present our numerical results showing the expected constraints on the six NUNM parameters (α_{ij}) that DUNE, JD, KD, and JD+KD setups can place. We derive bounds on α_{ij} following the simulation method as discussed in Sec. 4. The statistical significance with which we can constrain the NUNM parameters (α_{ij}) in a given experiment is defined as

$$\Delta\chi^2 = \min_{(\theta_{23}, \delta_{\text{CP}}, \phi_{ij}, \lambda_1, \lambda_2)} \left[\chi^2(\alpha_{ij} \neq 0) - \chi^2(\alpha_{ij} = 0) \right], \quad (5.1)$$

where, $\chi^2(\alpha_{ij} \neq 0)$ and $\chi^2(\alpha_{ij} = 0)$ are calculated by fitting the prospective data assuming NUNM ($\alpha_{ij} \neq 0$) and UNM ($\alpha_{ij} = 0$). Note that $\chi^2(\alpha_{ij} = 0) \approx 0$ because the statistical fluctuations are suppressed to obtain the median sensitivity of a given experiment in the frequentist approach [80]. While estimating the constraints, we marginalize over the most uncertain oscillation parameters ($\theta_{23}, \delta_{\text{CP}}$) and the phases associated with the off-diagonal NUNM parameters (ϕ_{ij}) in the fit. We also minimize over the systematic pulls on signal (λ_1) and background (λ_2).

In Fig. 6, we plot the $\Delta\chi^2$ function for the six NUNM parameters analyzed in our paper considering only the far detectors in a given setup. Upper (Lower) panels show the results for the diagonal (off-diagonal) NUNM parameters. The red curves in each panel refer to the sensitivities obtained for the DUNE setup considering a total of 336 kt-MW-yrs exposure, corresponding to a total 7 years of data collection with equal run-time in neutrino and antineutrino modes. The green curves show the results for JD for which we consider a total exposure of 2431 kt-MW-yrs with 10 years of total run-time (2.5 years in neutrino mode and 7.5 years in antineutrino mode). The magenta curves correspond to KD assuming the same exposure. We also estimate the results for JD+KD as shown by the blue curves. From the upper left panel, we observe that DUNE and JD+KD place similar constraints on α_{11} . The sensitivity to this parameter comes from two contributions: disappearance of intrinsic ν_e beam and the ν_e appearance. For both of these contributing channels, DUNE has better systematics and JD+KD has more statistics. As a result, limits on α_{11} is found to be almost the same for the two setups. However, for α_{22} (upper middle panel), JD+KD has significantly better sensitivity compared to DUNE setup. This is because α_{22} is mainly constrained by the disappearance channels, which due to the large statistics, is primarily




Figure 6: Expected limits on the NUNM parameters from DUNE (red curves), JD (green curves), KD (magenta curves), and JD+KD (blue curves). The upper (lower) panels are for the diagonal (off-diagonal) NUNM parameters one at a time. True values of θ_{23} and δ_{CP} are 45° and -90° , respectively. For the diagonal NUNM parameters, we marginalize over θ_{23} in the range $[40^\circ, 50^\circ]$ and δ_{CP} in the range $[-180^\circ, 180^\circ]$ in the fit. For the off-diagonal NUNM parameters, apart from θ_{23} and δ_{CP} , we also marginalize over the associated NUNM phases in the range of -180° to 180° .

limited by the systematic uncertainties. Since the normalization error for this channel is 3.5% (5%) for JD+KD (DUNE), it is clear that JD+KD can put better limit than DUNE. We have checked that if we consider the same amount of systematic uncertainties for both the setups, DUNE shows slightly better sensitivity than T2HKK.

For $|\alpha_{21}|$, DUNE and JD+KD have comparable sensitivities (see lower left panel). The slightly better limit on $|\alpha_{21}|$ achieved in the case of JD+KD as compared to DUNE, is due to the fact that JD+KD has larger statistics in the appearance channels. For the other three parameters $|\alpha_{31}|$, $|\alpha_{32}|$ and α_{33} , which enter the $\nu_\mu \rightarrow \nu_e$ appearance channel through matter parameters Δ_e and Δ_n (see Eq. 2.6), DUNE outperforms JD+KD setups because of its large matter effects.

We summarize our results in Table 6, where we give the bounds on the six NUNM parameters at 90% C.L. for the various long-baseline experimental setups discussed in

	DUNE	JD	KD	JD+KD	JD+KD+DUNE	T2K+NO ν A
α_{11}	[-0.020, 0.020]	[-0.025, 0.025]	[-0.040, 0.040]	[-0.022, 0.022]	[-0.017, 0.017]	[-0.06, 0.06]
α_{22}	[-0.014, 0.014]	[-0.0087, 0.0087]	[-0.013, 0.013]	[-0.007, 0.007]	[-0.006, 0.006]	[-0.02, 0.02]
α_{33}	[-0.2, 0.17]	< 0.6	< 0.63	< 0.476	[-0.17, 0.17]	< 0.64
$ \alpha_{21} $	< 0.022	< 0.015	< 0.10	< 0.016	< 0.012	< 0.06
$ \alpha_{31} $	< 0.15	< 0.48	< 0.70	< 0.34	< 0.11	< 2.20
$ \alpha_{32} $	< 0.33	< 1.2	< 0.85	< 0.71	< 0.27	< 1.4

Table 6: Bounds on the NUNM parameters at 90% C.L. (1 d.o.f.) using DUNE (second column), JD (third column), KD (fourth column), and JD+KD (fifth column). Sixth column shows the results for the combination of DUNE and JD+KD. Last column depicts results using the full exposure of T2K and NO ν A.

Parameter	DUNE (3.5 yrs+3.5 yrs)	DUNE (5 yrs+5 yrs)
α_{11}	[-0.020, 0.020]	[-0.020, 0.018]
α_{22}	[-0.014, 0.014]	[-0.013, 0.013]
α_{33}	[-0.2, 0.17]	[-0.19, 0.15]
$ \alpha_{21} $	< 0.022	< 0.016
$ \alpha_{31} $	< 0.15	< 0.12
$ \alpha_{32} $	< 0.33	< 0.31

Table 7: 90% C.L. (1 d.o.f.) limits on the NUNM parameters considering two different exposures of DUNE: total run-time of 7 years (see second column) and 10 years (see third column) equally divided in neutrino and antineutrino modes.

this paper. As clear from our previous discussion, the expected constraints on NUNM parameters from DUNE is better than the other two experiments JD and KD (and their combination) except for the parameters α_{22} and $|\alpha_{21}|$, where JD has better sensitivity than DUNE. Finally, in the sixth column of the Table, we give the final constraints on the NUNM parameters by combining the expected results from DUNE and JD+KD setups. As we have anticipated, the bounds experience a general improvements by $\sim 20\%$, with the precise magnitude depending on the parameter under consideration.

For a comparison with the ongoing long-baseline experiments, we also add the expected constraints from the combination of the T2K and NO ν A setup in the last column. For T2K, we consider a total exposure of 84.4 kt-MW-yrs with 22.5 kt detector mass, 750 kW proton beam power, 5 years run-time (2.5 years each for neutrino and antineutrino mode). For NO ν A, the considered exposure is 58.8 kt-MW-yrs with 14 kt detector mass, 700 kW proton beam power, and 6 years for total run-time (3 years each for neutrino and antineutrino modes). Due to the limited statistics, we observe that the expected constraints from T2K+NO ν A setup is worse than the DUNE or JD+KD setup.

Note that if some information coming from the near detector (for an example, measurement of the initial neutrino flux) are used to analyze the far detector data then the constraints on α_{11} and α_{22} may be modified (see Sec. 6 for a detailed discussion). However,

in this section, we adopt a different strategy, where we simulate the far detector data alone to set limits on the NUNM parameters. In principle, this approach is valid if the initial flux can be predicted by some theoretical calculation or measured at an experiment which is insensitive to neutrino oscillation phenomena. In fact, the MINOS/MINOS+ experiment adopted this strategy where the oscillation data at the far detector was analyzed using information from the MINERvA flux predictions [81]. In future, if somehow we can apply this approach for DUNE and T2HKK, then in principle, one can estimate the limits on the NUNM parameters using only the far detector data.

At the same time, we understand that the assumptions that we take in our paper for the systematic uncertainties at the far detector may be too optimistic if we do not use the near detector to measure the initial flux. To have a better understanding on this issue, we perform some study and find that limits on α_{11} and α_{22} are mainly governed by the systematic uncertainties. In other words, the expected constraints on these two NUNM parameters are proportional to the systematic uncertainties that we consider in our simulation. Therefore, it may be possible to predict what would be the limits on α_{11} and α_{22} for a given assumptions on systematic uncertainties.

We compare our results summarized in Table 6 with the bounds reported in Table 1⁴. We observe that the bound we achieve from the DUNE+JD+KD (DUNE+T2HKK) setup for the diagonal α_{11} is $\sim 80\%$ better than the bound quoted in Ref. [53]. In the α_{22} case, the two results are comparable, with a slightly better limit when the global neutrino data analysis is performed. For the remaining diagonal parameter α_{33} , NC data from MINOS/MINOS+ give a 60% stronger bound [53] compared to the one expected from the DUNE+JD+KD setup. As for α_{21} , the authors of the Ref. [53] make use of the triangular inequality as well as the data from the short-baseline experiments; this allows to constrain the mentioned parameter very tightly. However, due to the large statistics and good systematics of DUNE and JD+KD setups, we can achieve an almost similar bound without using any external hypothesis on the relations between the α_{ij} . On the other hand, constraints on α_{31} and α_{32} in Table 1 are substantially better than the ones we obtain from DUNE+JD+KD setup. Also, in these cases, the triangular inequalities which link them to the diagonal NUNM parameters play an important role, together with the short-baseline experiments limits on the ν_τ appearance. However, it is important to stress that all our results are obtained in a complete model independent fashion, relying only on the expected data from DUNE and T2HKK. We check that for our best setup, namely DUNE+T2HKK, the only parameter whose limit gets improved when we consider these inequalities is $|\alpha_{32}|$ because of the stringent bound on α_{22} . Since $|\alpha_{21}|$ is already tightly constrained, we do not see any improvement in its limit using these inequalities. As far as the bound on $|\alpha_{31}|$ is concerned, since the limit on the diagonal parameter α_{33} is very poor, we also do not see any improvement.

⁴In order to get their results, the authors of Ref. [40] left free the standard oscillation parameters θ_{23} , δ_{CP} , and Δm_{31}^2 and the NUNM parameters α_{11} , α_{21} , α_{22} . Conversely, in our work we marginalize over δ_{CP} and θ_{23} only, but we have checked that the marginalization over Δm_{31}^2 does not have any significant impact. In Appendix B, we also show that the marginalization over α_{11} , α_{21} , and α_{22} does not cause remarkable change in the results.

Recently, the DUNE collaboration [27] exploited the possibility of increasing the exposure of the experiment from 336 kt-MW-yrs to 480 kt-MW-yrs (corresponding to an increase of the data taking time from 7 years to 10 years with 5 years in neutrino mode and 5 years in antineutrino mode). In Table 7, we compare our previous constraints from the DUNE experiment, Table 6, with those obtained in the (5 + 5) years configuration. We observe that the constraints on all six NUNM parameters improve by small amount except for $|\alpha_{21}|$, which shows a significant improvement. This happens because the higher run-time increases statistics of the $\nu_\mu \rightarrow \nu_e$ appearance channel, which is the one driving the α_{21} sensitivity. On the other hand, the $\nu_\mu \rightarrow \nu_\mu$ disappearance channel is almost already saturated by systematics after 3.5 years + 3.5 years of running. This leads to only small improvements on the other NUNM parameters sensitivities.

6 Benefits of having near detectors

Near detectors (ND) are a fundamental component for long-baseline neutrino experiments. Indeed, a detector placed very close to the beam source (from hundreds of meters to a few kilometers) is able to monitor the neutrino beam, measuring the flavor composition, and the total number of neutrinos emitted from the source. Near detectors are not expected to improve any of the standard oscillation parameter measurements, since at such short distances, oscillations do not develop for neutrinos with energies in the GeV range. However, in some new physics scenarios, in which, oscillation probabilities contain zero-distance terms, near detectors can be used to constrain non-standard parameters in a very straightforward way. This is the case of the non-unitarity framework under discussion where, as already mentioned in Sec. 2, at vanishing distances we have zero-distance terms in case $\nu_\mu \rightarrow \nu_e$ appearance channel: $P_{\mu e}^{L=0} \sim |\alpha_{21}|^2$, and $\nu_\mu \rightarrow \nu_\mu$ disappearance channel: $P_{\mu\mu}^{L=0} \sim 1 + 2|\alpha_{21}|^2 + 6\alpha_{22}^2 + 4\alpha_{22}$. Thus, we can expect that T2HKK and DUNE near detectors would be able to constrain two parameters $|\alpha_{21}|$ and α_{22} from $\nu_\mu \rightarrow \nu_e$ appearance and $\nu_\mu \rightarrow \nu_\mu$ disappearance channel, respectively, but also α_{11} considering the ν_e beam contamination (see Eq. A1). So, in this section, we analytically infer the order of magnitude of bounds implied by ND measurements. Let us consider the total number of events of a given channel as [82]:

$$N = N_0 P_{\alpha\beta}(\alpha_{ij}), \quad (6.1)$$

where, the normalization factor N_0 includes all the detector properties. For an oscillation channel $\nu_\alpha \rightarrow \nu_\beta$, N_0 can be defined as:

$$N_0 = \int_{E_\nu} dE_\nu \sigma_\beta(E_\nu) \frac{d\phi_\alpha}{dE_\nu}(E_\nu) \varepsilon_\beta(E_\nu), \quad (6.2)$$

where, σ_β denotes the production cross-section of the β lepton, ε_β represents the detector efficiency, and ϕ_α stands for the initial neutrino flux of flavor α . If we want to put bounds on new physics parameters, we can use a simple χ^2 test with a gaussian χ^2 defined as

$$\chi^2 = \frac{(N_{obs} - N_{fit})^2}{\sigma^2}, \quad (6.3)$$

Expt.	α_{11}		α_{22}	
	w/o norm.	w/ norm.	w/o norm.	w/ norm.
DUNE	[-0.02, 0.02]	[-0.043, 0.034]	[-0.014, 0.014]	[-0.036, 0.048]
JD+KD	[-0.022, 0.022]	[-0.048, 0.040]	[-0.007, 0.007]	[-0.038, 0.050]
DUNE+JD+KD	[-0.017, 0.017]	[-0.036, 0.026]	[-0.006, 0.006]	[-0.026, 0.039]

Table 8: 90% C.L. (1 d.o.f.) limits on the NUNM parameters α_{11} and α_{22} for the two setups, DUNE, JD+KD, and combination of them. Second and fourth column correspond to the constraints assuming only far detector. Third and fifth columns correspond to the constraints using the FD and ND correlation (or with normalization factor in the oscillation probability).

where, σ represents the uncertainty on the number of events; in this case, neglecting the backgrounds, we get:

$$\chi^2 = \frac{N_0^2}{\sigma^2} \left[\delta_{\alpha\beta} - P_{\alpha\beta}(\alpha_{ij}^{fit}) \right]^2. \quad (6.4)$$

For the $\nu_\mu \rightarrow \nu_\mu$ disappearance channel, the leading term of the probability is $P_{\mu\mu}^{L=0} = 1 + 4\alpha_{22}$. Therefore, the χ^2 assumes the form:

$$\chi^2 = \frac{16N_0^2\alpha_{22}^2}{\sigma^2}. \quad (6.5)$$

At a chosen confidence level, represented by a cut at χ_0^2 , it is possible to exclude the region satisfying:

$$|\alpha_{22}| > \frac{\sqrt{\chi_0^2}\sigma}{4N_0}. \quad (6.6)$$

Since the disappearance channel is expected to produce a huge number of events at the near detector, one can consider the uncertainty to be dominated by systematic errors σ_{sys} . Thus, it is possible to approximate $\sigma \sim N_0\sigma_{sys}$, where N_0 represents the number of events in absence of zero-distance effects, being the disappearance probability in that case equal to 1. This allows to simplify Eq. 6.6 as follows:

$$|\alpha_{22}| > \frac{\sqrt{\chi_0^2}\sigma_{sys}}{4}, \quad (6.7)$$

which tells us that, neglecting backgrounds effects, the near detector limits would be of the order of the chosen systematic uncertainty. A similar approach can be used for the $\nu_e \rightarrow \nu_e$ oscillation channel (see Eq. A1), which arises from the ν_e beam contamination, obtaining an inequality for $|\alpha_{11}|$ of the similar form:

$$|\alpha_{11}| > \frac{\sqrt{\chi_0^2}\sigma_{sys}^{\nu_e}}{4}, \quad (6.8)$$

where $\sigma_{sys}^{\nu_e}$ refers to the systematic uncertainty on the $\nu_e \rightarrow \nu_e$ transition.

For the appearance channel, the zero-distance probability reads $P_{\mu e}^{L=0} = |\alpha_{21}|^2$; the χ^2 function can therefore be written as:

$$\chi^2 = \frac{N_0^2|\alpha_{21}|^4}{\sigma^2}, \quad (6.9)$$




Figure 7: Upper panels show the improvements in the sensitivities to α_{11} , α_{22} , and $|\alpha_{21}|$ due to the presence of 67 tons LArTPC near detector placed at distance of 574 meters from the neutrino source for DUNE. Lower panels portray the same for JD+KD having a 1 kt water Cherenkov near detector placed at a distance of 1 km from J-PARC. The blue curves show the performance with only near detectors. The red curves represent the combined sensitivities due to both near and far detectors. True values of the standard oscillation parameters are taken from Table 2. We obtain our results marginalizing over δ_{CP} in the range $[-180^\circ, 180^\circ]$ and θ_{23} in the range $[40^\circ, 50^\circ]$ in the fit. We also marginalize over the associated phase ϕ_{21} in the range $[-180^\circ, 180^\circ]$ for the off-diagonal NUNM parameter $|\alpha_{21}|$.

and the excluded region is expected to be determined by the following relation:

$$|\alpha_{21}| < \sqrt[4]{\frac{\chi_0^2 \sigma^2}{N_0^2}}. \quad (6.10)$$

Since the number of events at the near detector is in principle very small (being only caused by new physics), the uncertainty is dominated by statistics. Thus, given a certain number of observed events, $\sigma \sim \sqrt{N_{obs}}$ and the excluded values of $|\alpha_{21}|$ reduced to:

$$|\alpha_{21}| < \sqrt[4]{\frac{\chi_0^2 N_{obs}}{N_0^2}}, \quad (6.11)$$

Parameter	DUNE (ND)	DUNE FD w/ norm.	IWCD	JD+KD w/ norm.
α_{11}	[-0.020, 0.024]	[-0.043, 0.034]	[-0.029, 0.033]	[-0.048, 0.040]
α_{22}	[-0.033, 0.037]	[-0.036, 0.048]	[-0.019, 0.020]	[-0.038, 0.050]
$ \alpha_{21} $	< 0.007	< 0.022	< 0.01	< 0.015

Table 9: 90% C.L. (1 d.o.f.) bounds on the NUNM parameters α_{11} , α_{22} , and $|\alpha_{21}|$ obtained with and without near detectors in DUNE and JD+KD. Note that IWCD is the near detector for JD+KD setup.

suggesting that the bounds are very sensitive to the number of events and to the running time of the experiment.

Both DUNE and T2HKK will have near detectors [55, 67, 83] which may play a crucial role to probe various new physics scenarios including the possibility of non-unitarity of the PMNS matrix which is the main thrust of this work. In our analysis, for DUNE, we consider a 67 tons LArTPC near detector placed at a baseline of 574 meters from Fermilab [67]. For JD+KD, we consider a 1 kt water Cherenkov near detector located at a baseline of 1 km from J-PARC which is known as IWCD [75, 76]. In order to simulate their responses, we scale the far detector fluxes for ND baselines and take into account their fiducial masses. We follow a very conservative approach as far as the systematic uncertainties at the near detectors are concerned. We multiply the FD systematic uncertainties by a factor of three and consider them as inputs for the ND. In DUNE near detector, we expect $\mathcal{O}(10^7)$ ν_μ and $\bar{\nu}_\mu$ events, which provide bounds on α_{22} . DUNE can place stringent constraints on α_{11} and α_{21} using $\mathcal{O}(10^6)$ ν_e and $\bar{\nu}_e$ events at ND, which stem from both intrinsic ν_e ($\bar{\nu}_e$) beam contamination and via $\nu_\mu \rightarrow \nu_e$ ($\bar{\nu}_\mu \rightarrow \bar{\nu}_e$) appearance caused due to zero-distance effect. For the NDs, we consider their appropriate baselines, fiducial masses, and systematic uncertainties which we assume to be larger than the systematic uncertainties considered for the FDs.

Before discussing the limits that the near detectors would be able to set using their own data, we want to study the effect of the ND flux measurements on the far detector constraints. Indeed, if the initial neutrino flux is measured at the near detector and then extrapolated to the far detector, the probability which could be inferred at the far detector is the effective probability defined as

$$P_{\alpha\beta}^{\text{eff}} = \frac{P_{\alpha\beta}}{P_{\alpha\alpha}^{L=0}}. \quad (6.12)$$

The $P_{\alpha\alpha}^{L=0}$ term that appears in the denominator is the survival probability initial neutrino flavor at the source or the zero-distance term which act as a normalization factor. If we normalize the $\nu_\mu \rightarrow \nu_\mu$ survival probability at the far detector using the zero-distance term in Eq. 2.10, it is observed that the contribution from α_{22} gets canceled at the leading order. As a result, sensitivity to the parameter α_{22} is worsened for a given setup. The same happens for α_{11} , whose contribution in the effective $\nu_e \rightarrow \nu_e$ disappearance probability is canceled at the leading order (see Eq. A1 and A4). Since the sensitivity to this parameter arises partially due to the intrinsic ν_e that we have in the beam to begin with, the near detector normalization causes a deterioration of α_{11} limits. In Table 8, we show how

the constraints on α_{11} and α_{22} would be modified when taking into account the FD and ND correlation for the three setups namely, DUNE, JD+KD, and DUNE+JD+KD. We observe that the bound on α_{11} is increased by a factor of almost two when we consider the correlation between the FD and ND. For α_{22} , the bound is deteriorated at least three times compared to the FD case only. We have checked that no other NUNM parameter is affected significantly if we consider the FD and ND correlation. Indeed, the non-diagonal parameters and α_{33} can be constrained using appearance channels (for which we do not have full cancellations in the effective probabilities) or using the interplay with matter effects, which are not developed at the near site.

The bounds obtained using the above-mentioned near detectors are shown in Fig. 7 for DUNE and T2HKK together with the results we got using the FD data with effective probabilities. For α_{11} , NDs of the two setups can put bounds better than the one set by the FDs considering the ND normalization due to the very high statistics and the strong α_{11} dependence of the zero-distance probability. The improvement is roughly a factor of two for DUNE and 60% for T2HKK (see Table 9). Note that the obtained limits are in agreement with the predictions deduced from Eq. 6.8, once we insert a normalization uncertainty of 6% (15%) for DUNE (JD+KD).

For the second diagonal parameter α_{22} we also observe a similar situation, in which the ND alone can put more stringent bounds than the far detector when the normalization is considered, despite of the increased systematics. The improvement can be quantified as roughly 25% in DUNE and a factor of two in T2HKK. Once again, the analytical predictions from Eq. 6.8 are sufficiently recovered by the numerical simulations, considering that the near detectors normalization systematics are 15% for DUNE and 10.5% for T2HKK. Note that the bounds from the far detector data alone would be considerably better than the near detector ones.

Finally, for the NUNM parameter $|\alpha_{21}|$ the near detectors bounds are considerably better than the far detector ones, due to the zero-distance effect outlined in Eq. 2.7. In particular, the limits are ~ 3 times smaller than the one set by the far detector in the DUNE facility and $\sim 70\%$ smaller in the case of T2HKK. Considering a number of observed events of $\mathcal{O}(10)$, and taking into account that we expect $N_0 \sim 10^6$ per year [67], our analytic estimate for $|\alpha_{21}|$ is comparable with the numerical results.

7 Improvement due to the ν_τ sample in DUNE

The ν_τ production at accelerator experiments is very challenging since the charged current interactions of such particles with nuclei have an energy threshold of 3.1 GeV. Thus, many proposed long-baseline experiments are not able to detect such neutrinos⁵. However, the DUNE neutrino spectra will have peak at around 2.5 GeV (differently from the T2HKK where $E_\nu \simeq 0.6$ GeV) and the most energetic neutrinos of the beam will have enough energy to produce τ leptons. Recently, different studies [51, 85–87] take into account the possibility of including the ν_τ ($\bar{\nu}_\tau$) sample in the DUNE analysis.

⁵In this respect, the OPERA setup provided neutrinos with an average energy close to 13 GeV [84].

Parameter	w/o ν_τ appearance	w/ ν_τ appearance
α_{33}	[-0.2, 0.17]	[-0.16, 0.15]
$ \alpha_{32} $	< 0.33	< 0.19

Table 10: 90% C.L. limits on the NUNM parameters α_{33} and $|\alpha_{32}|$ from the DUNE setup. Second (third) column shows the results without (with) τ in the analysis.

The recognition of such events could be in principle possible due to the imaging capabilities of LArTPC detectors. Because of a relatively small number of neutrinos with an energy above the production threshold and of the short lifetime of the τ leptons which could make the recognition of the τ interaction and the decay vertices a difficult task, the ν_τ appearance channel is not really useful to constrain the standard oscillation parameters. However, when new physics affects the oscillation probabilities, it has been shown in Refs. [51, 86, 88, 89] that the NUNM parameters α_{33} and $|\alpha_{32}|$ constraints can be improved even by the small number of ν_τ events.




Figure 8: Comparison between the DUNE sensitivities on $|\alpha_{32}|$ (left panel) and α_{33} (right panel) when ν_τ appearance channel is included in the analysis (red lines) and the case where no τ events are analyzed (blue lines). True values of the standard oscillation parameters are taken from Table 2. All results have been obtained marginalizing over δ_{CP} in the range $[-180^\circ, 180^\circ]$ and θ_{23} in the range $[40^\circ, 50^\circ]$. For $|\alpha_{32}|$ (left panel), we also marginalize over ϕ_{32} in the range $[-180^\circ, 180^\circ]$.

In case of non-unitary neutrino mixing, the $\nu_\mu \rightarrow \nu_\tau$ oscillation probability mainly depends on the three parameters $|\alpha_{32}|$, α_{33} and α_{22} (see Eq. A3 in Appendix A). While we expect that the sensitivity to α_{22} will not be improved by the small number of ν_τ events, the other two poorly constrained NUNM parameters $|\alpha_{32}|$ and α_{33} could take advantage of $\nu_\mu \rightarrow \nu_\tau$ oscillation channel. We include τ events in our analysis in the following fashion.

- For the hadronic decays of τ events having a branching ratio of 65%, a 30% signal efficiency has been assumed and 10% of the NC events are considered as background [51].
- For τ decaying to electron with a branching ratio of 17.4%, we assume a signal efficiency of 30% considering ν_e events as possible background. We consider the signal to background ratio of 2.45 in our analysis [86].

The muonic decays have not been taken into account since the discrimination of the number of background events would be too large compared to the signal events [51, 86]. The total number of ν_τ ($\bar{\nu}_\tau$) events in DUNE is expected to be roughly 72 (37) per year. The normalization error for the signal is taken to be 20%. The results of our analysis for α_{33} and $|\alpha_{32}|$ are shown in Fig. 8 and Table 10. The allowed range of α_{33} , which appears only at the second order in the probability, is reduced of $\sim 13\%$ by the inclusion of the new oscillation channel, and the new limits are set into the range $[-0.16, 0.15]$ (see Table 10). On the other hand, the sensitivity to $|\alpha_{32}|$, which impacts linearly the ν_τ appearance probability, is significantly improved: in this case, the new upper bound is roughly 60% smaller than the one set by the standard oscillation channels, namely $|\alpha_{32}| < 0.19$.

8 Summary and conclusions

Neutrino oscillation has been one of the most important discoveries over the last few decades and a large number of pioneering neutrino experiments have confirmed this phenomena of mass-induced flavor transition. Excellent data from various neutrino oscillation experiments having different baselines and energies have enabled us to achieve remarkable precision on almost all the three-flavor oscillation parameters. However, some of the features of neutrino mass-mixing parameters, like, neutrino mass hierarchy, the value of the CP-violating phase δ_{CP} , and octant of θ_{23} are still poorly known and next-generation neutrino experiments will play an important role to address these issues with high confidence level.

Given the magnificent precision on neutrino mixing angles and rapidly increasing knowledge on δ_{CP} , it seems quite natural to ask if there is any violation of the unitary property of 3×3 PMNS mixing matrix, which may be related to the existence of new mass eigenstates of neutrinos. In this context, there exists one well-known parameterization in the literature, which takes into account the non-unitary neutrino mixing (NUNM) by introducing a lower triangular matrix with three real and three complex parameters, α_{ij} .

To study the impact of these NUNM parameters on various oscillation channels, for the first time, we derive simple approximate analytical expressions for the oscillation probabilities in matter in the atmospheric regime ($\Delta_{31} \gg \Delta_{21}$). Our perturbative expansions are valid up to second order in the small deviations from the mixing angles by their tri-bimaximal values, second order in the NUNM parameters, and first order in the matter potential. We have shown that the $\nu_\mu \rightarrow \nu_e$ appearance probability (see Eq. 2.6) mainly depends on $|\alpha_{21}|$, but when matter potential is large, the impact of $|\alpha_{31}|$ and $|\alpha_{32}|$ can also be significant. On the other hand, the $\nu_\mu \rightarrow \nu_\mu$ disappearance probability (see Eq. 2.9) mainly relies on α_{22} , but sub-leading dependencies on $|\alpha_{21}|$ in vacuum and α_{33} , $|\alpha_{31}|$, and $|\alpha_{32}|$ in matter are also present. The only parameter that does not appear in our formulae

is α_{11} , which is only relevant at the higher-orders in our perturbative expansions as shown in Ref. [37] for the vacuum case.

In this work, we analyze in detail, the impact of possible NUNM in the context of long-baseline experiments DUNE and T2HK having one detector in Japan (JD) and a second detector in Korea (KD), and the combination of these two detectors, popularly known as T2HKK or JD+KD. First, we show how the various NUNM parameters (α_{ij}) are correlated with the oscillation parameters θ_{23} and δ_{CP} for these setups. Then, we estimate in detail the sensitivities of these experiments to place direct, model-independent, competitive constraints on the six NUNM parameters at 90% C.L. (see Fig. 6 and Table 6). The wide-band neutrino beam in DUNE encompassing both first and second oscillation maxima allows us to measure the NUNM parameters at several L/E values in the presence of significant matter effect due to its 1300 km baseline. Indeed, DUNE shows better sensitivity than JD+KD in constraining the NUNM parameters, which are influenced by matter effects, namely, α_{33} , $|\alpha_{31}|$, and $|\alpha_{32}|$. We observe that JD+KD will provide a stringent constraint on α_{22} as compared to DUNE because it has less systematic uncertainties in the disappearance channel. Also, due to the larger statistics in the appearance channel, JD+KD will give significantly better limit on the NUNM parameter $|\alpha_{21}|$ in comparison to DUNE. Lastly, α_{11} is expected to be constrained almost in the same way by these two experiments. We show how the limits on the NUNM parameters get improved in case of DUNE, if the total exposure is increased from 336 kt-MW-yrs to 480 kt-MW-yrs (corresponding to an increase in the total run-time from 7 years to 10 years), as proposed in the recent TDR [27]. We also estimate how much the sensitivities can be improved by adding the prospective data from DUNE and JD+KD. Finally, we compare our results with the constraints that can be achieved using the full exposure of currently running experiments T2K and NO ν A.

Due to the so-called zero-distance effects which are induced by the non-unitary neutrino mixing in neutrino oscillation probabilities, the prospective data from near detectors in both DUNE and JD+KD experiments could be in principle used to bound the three NUNM parameters $|\alpha_{21}|$, α_{11} and α_{22} . However, the zero-distance effect should also be taken into account if the near detectors data will be used to measure the initial neutrino flux for both experiments. This would lead to a substantial deterioration of the limits that the far detectors could set on α_{11} and α_{22} , as summarized in Table 8.

Moreover, in DUNE, the expected limits on $|\alpha_{32}|$ and α_{33} get improved by $\mathcal{O}(20)\%$ when we also add the ν_τ appearance sample in our analysis.

Long-baseline experiments have contributed significantly in our journey to establish the three-neutrino paradigm on a strong footing. Future high-precision long-baseline experiments such as DUNE and T2HKK (JD+KD) will continue this legacy and rigorously test the unitarity of the PMNS matrix by measuring the mixing angles and CP phase δ_{CP} with remarkable precision. We hope our present work will provide further boost along this direction.

Acknowledgments

We would like to thank M. Blennow, E. Fernandez Martinez, S. Roy, S. Sahoo, M. Singh, and M. Tortola for useful discussions. S.K.A. is supported by the INSPIRE Faculty Re-

search Grant [IFA-PH-12] from the Department of Science and Technology (DST), Govt. of India. S.K.A. acknowledges the financial support from the Swarnajayanti Fellowship Research Grant (No. DST/SJF/PSA-05/2019-20) provided by the DST, Govt. of India and the Research Grant (File no. SB/SJF/2020-21/21) from the Science and Engineering Research Board (SERB) under the Swarnajayanti Fellowship by the DST, Govt. of India. S.K.A would like to thank the United States-India Educational Foundation for providing the financial support through the Fulbright-Nehru Academic and Professional Excellence Fellowship (Award No. 2710/F-N APE/2021). The numerical simulations are carried out using SAMKHYA: High-Performance Computing Facility at Institute of Physics, Bhubaneswar.

A Analytical expressions of the transition probabilities in the regime of small matter effects

Exact vs. approximate probabilities

In this appendix, we provide a check of the goodness of our expanded probabilities, Eqs. 2.6 to 2.9. To this aim, in the upper panels of Fig. A1, we show the $\nu_\mu \rightarrow \nu_e$ appearance probabilities as a function of neutrino energy considering the NUNM parameters $|\alpha_{21}|$ and $|\alpha_{31}|$ one at a time. We portray the same in the lower panels for $\nu_\mu \rightarrow \nu_\mu$ disappearance channel in presence of the diagonal NUNM parameter α_{22} . We consider three different baselines, corresponding to DUNE (left panels), JD (middle panels), and KD (right panels). In each panel, solid curves are obtained numerically from the full (not expanded) Hamiltonian, while dashed curves correspond to the oscillation probabilities obtained analytically using Eq. 2.6 and Eq. 2.9. The value of the CP phase δ_{CP} and phases ϕ_{21} , ϕ_{31} are fixed to zero, while the values of the other standard oscillation parameters are taken from Table 2. As we can see, the agreement between the full solid and the analytical dashed curves is generally very good for both probabilities.

Perturbative expansion of P_{ee} , $P_{e\tau}$, and $P_{\mu\tau}$

For the sake of completeness, we give here the perturbative expressions of the probabilities P_{ee} , $P_{e\tau}$ and $P_{\mu\tau}$, using the same approximations discussed in Sec. 2. We start with the $\nu_e \rightarrow \nu_e$ disappearance channel:

$$\begin{aligned}
P_{ee} = & 1 + 4\alpha_{11} + 6\alpha_{11}^2 - \left(\frac{4r}{\Delta_{31}}\right) \Delta_n [|\alpha_{21}| \cos(\delta - \phi_{21}) + \alpha_{31} \cos(\delta - \phi_{31})] \sin^2 \Delta_{31} + \\
& - \left(\frac{2r}{\Delta_{31}}\right) \sin \Delta_{31} [(\Delta_{31} + 2\Delta_e) \sin \Delta_{31} - 2\Delta_{31}\Delta_e \cos \Delta_{31}] . \tag{A1}
\end{aligned}$$




Figure A1: Upper panels show the $\nu_\mu \rightarrow \nu_e$ appearance probability in presence of the NUNM parameters $|\alpha_{21}| = 0.025$ and $|\alpha_{31}| = 0.1$. Lower panels depict the $\nu_\mu \rightarrow \nu_\mu$ disappearance probability for $\alpha_{22} = -0.015$. Left, middle, and right columns correspond to the baselines of 1300 km (DUNE), 295 km (JD), and 1100 km (KD), respectively. Solid curves in each panel show the oscillation probabilities obtained numerically, while the dashed curves are the probabilities obtained analytically using Eq. 2.6 and Eq. 2.9. We consider the values of δ_{CP} and the phases ϕ_{21}, ϕ_{31} equal to zero. The values of the other standard oscillation parameters are taken from Table 2.

The $\nu_e \rightarrow \nu_\tau$ transition is governed by the following expression:

$$\begin{aligned}
P_{e\tau} = & \left(\frac{r^2}{\Delta_{31}} \right) \sin \Delta_{13} [(\Delta_{31} + 2\Delta_e) \sin \Delta_{31} - 2\Delta_{31}\Delta_e \cos \Delta_{31}] \\
& \left(\frac{|\alpha_{31}|^2}{\Delta_{31}} \right) [\Delta_{31} - \Delta_n(1 - \cos 2\Delta_{31})] + \\
& \left(\frac{2|\alpha_{21}|r}{\Delta_{31}} \right) \Delta_n \sin \Delta_{31} [\sin \Delta_{31} \cos(\delta_{\text{CP}} - \phi_{21}) - \Delta_{31} \cos(\delta_{\text{CP}} - \Delta_{31} - \phi_{21})] + \quad (\text{A2}) \\
& \left(\frac{|\alpha_{31}|r}{\Delta_{31}} \right) [2\Delta_{31} \sin \Delta_{31} (\Delta_e + \Delta_n) \cos(\delta_{\text{CP}} - \Delta_{13} - \phi_{31}) + \\
& \sin(\delta_{\text{CP}} - \Delta_{31} - \phi_{31}) (\sin \Delta_{31} (2\Delta_{31} + 2\Delta_e - \Delta_n) - 2\Delta_{31}\Delta_e \cos \Delta_{31}) + \\
& \Delta_n \sin \Delta_{31} \sin(\delta + \Delta_{31} - \phi_{31})] + \\
& \left(\frac{|\alpha_{21}||\alpha_{31}|}{\Delta_{31}} \right) \Delta_n [-2\Delta_{31} \sin(\phi_{21} - \phi_{31}) + \cos(2\Delta_{31} - \phi_{21} + \phi_{31}) - \cos(\phi_{21} - \phi_{31})].
\end{aligned}$$

Eventually, the ν_τ appearance from a ν_μ beam is regulated by the following probability expression:

$$\begin{aligned}
P_{\mu\tau} = & \sin^2 \Delta_{31} (1 + 2\alpha_{22} + 2\alpha_{33} - 4a^2 + \alpha_{22}^2 + \alpha_{33}^2 + 4\alpha_{22}\alpha_{33}) + \\
& |\alpha_{32}| \sin 2\Delta_{13} [2\Delta_n \cos \phi_{32} - \sin \phi_{32}] + \\
& \left(\frac{r^2}{\Delta_{31}} \right) \sin(\Delta_{31}) [2\Delta_{31}\Delta_e \cos \Delta_{31} - \sin \Delta_{31}(\Delta_{31} + 2\Delta_e)] + \\
& \left(\frac{2}{\Delta_{31}} \right) (|\alpha_{21}|^2 + |\alpha_{31}|^2) \Delta_n \sin \Delta_{31} [\sin \Delta_{31} - \Delta_{31} \cos \Delta_{31}] + \\
& \left(\frac{|\alpha_{32}|^2}{\Delta_{31}} \right) [\Delta_n \sin 2\phi_{32}(\sin 2\Delta_{31} - 2\Delta_{31} \cos 2\Delta_{31}) + \Delta_{31} \cos^2 \Delta_{13}] + \\
& \left(\frac{2|\alpha_{21}|r}{\Delta_{31}} \right) \sin \Delta_{31} [\sin \Delta_{31} \cos(\delta_{\text{CP}} - \phi_{21})(\Delta_{31} + \Delta_e - \Delta_n) - \Delta_{31}\Delta_e \cos(\delta_{\text{CP}} + \Delta_{31} - \phi_{21}) + \\
& \quad \Delta_{31}\Delta_n \cos(\delta_{\text{CP}} - \Delta_{31} - \phi_{21})] + \\
& \left(\frac{2|\alpha_{31}|r}{\Delta_{31}} \right) [\sin \Delta_{31}(\sin \Delta_{31} \cos(\delta_{\text{CP}} - \phi_{31})(\Delta_{31} + \Delta_e - \Delta_n) - \Delta_{31}\Delta_e \cos(\delta_{\text{CP}} - \Delta_{31} - \phi_{31}) + \\
& \quad \Delta_{31}\Delta_n \cos(\delta_{\text{CP}} + \Delta_{31} - \phi_{31}))] + \left(\frac{8a}{\Delta_{31}} \right) (\alpha_{22} - \alpha_{33}) \Delta_n \sin \Delta_{31}(\Delta_{31} \cos \Delta_{31} - \sin \Delta_{31}) + \\
& 4a|\alpha_{32}| \sin^2 \Delta_{31} \cos \phi_{32} + \\
& \left(\frac{2|\alpha_{21}||\alpha_{31}|}{\Delta_{31}} \right) \sin \Delta_{31} [2\Delta_{31}(\Delta_e + \Delta_n) \cos(\Delta_{31} - \phi_{21} + \phi_{31}) + 2\Delta_n \sin \Delta_{31} \cos(\phi_{21} - \phi_{31}) - \\
& \quad \Delta_{31} \sin(\Delta_{31} - \phi_{21} + \phi_{31})] + \\
& 4\alpha_{22}|\alpha_{32}| \sin \Delta_{31} \left(\frac{\Delta_n \cos \phi_{32}(\sin \Delta_{31} + 2\Delta_{31} \cos \Delta_{31})}{\Delta_{31}} - \cos \Delta_{31} \sin \phi_{32} \right) + \\
& - \left(\frac{2|\alpha_{32}|\alpha_{33}}{\Delta_{31}} \right) \sin \Delta_{31} [2\Delta_n \cos \phi_{32}(\sin \Delta_{31} - 3\Delta_{31} \cos \Delta_{31}) + \Delta_{31} \cos \Delta_{31} \sin \phi_{32}] . \tag{A3}
\end{aligned}$$

The zero-distance effects are as follows:

$$\begin{aligned}
P_{ee}^{L=0} & \sim 1 + 4\alpha_{11} + 6\alpha_{11}^2, \\
P_{e\tau}^{L=0} & \sim |\alpha_{31}|^2, \\
P_{\mu\tau}^{L=0} & \sim |\alpha_{32}|^2.
\end{aligned} \tag{A4}$$

It is easy to check that, when all α parameters are set to zero, we recover the unitary relation $P_{\mu e} + P_{\mu\mu} + P_{\mu\tau} = 1$.

B Impact of marginalization over the other NUNM parameters while showing the constraints on α_{ij}

In this appendix, we derive the constraints on the NUNM parameters obtained, differently from the procedure adopted in the main text, marginalizing over the other NUNM parameters α_{ij} . Our numerical results are shown in Fig. B1, where we show the impact of the marginalization over various NUNM parameters on the sensitivity of the DUNE+JD+KD




Figure B1: The combined sensitivity of DUNE and JD+KD setups to the NUNM parameters. True values of the standard oscillation parameters are given in Table 2. We marginalize over θ_{23} and δ_{CP} in the fit (see text for details). Black curves show the sensitivity when only one NUNM parameter is considered at a time, while the others are taken to be zero. Red curves in all the panels correspond to the case when all other NUNM parameters are kept free in the fit.

setup. In each panel, black curves show the sensitivity for a particular NUNM parameter, when all other NUNM parameters are fixed in the fit. The red curves show the situation when all un-displayed NUNM parameters are marginalized in the fit with no priors. The 90% C.L. (1 d.o.f.) constraints on various NUNM parameters are summarized in Table B1. We observe that the marginalization over all other NUNM parameters worsen the bounds on all of them but α_{22} . This is due to the fact that, as it can be seen in Eq. 2.9, α_{22} appears at the leading order, with no correlations to the other α_{ij} . The other diagonal parameter α_{11} shows only a marginal deterioration at the level of 15% since, as before, its correlation with the other NUNM parameters are mild in ν_e appearance probability as shown in Ref. [37]. We see from Table B1 that there is a considerable deterioration in the sensitivities for α_{33} , $|\alpha_{21}|$, $|\alpha_{31}|$, and $|\alpha_{32}|$ when we marginalize over the un-displayed NUNM parameters in the fit. This happens because all these four parameters are strongly correlated among them and with other two NUNM parameters α_{11} and α_{22} (see Eqs. 2.6

Parameter	Other α_{ij} 's zero	Other α_{ij} 's free
α_{11}	[-0.017, 0.017]	[-0.02, 0.02]
α_{22}	[-0.006, 0.006]	[-0.006, 0.006]
α_{33}	[-0.20, 0.17]	[-0.35, 0.33]
$ \alpha_{21} $	< 0.012	< 0.017
$ \alpha_{31} $	< 0.11	< 0.18
$ \alpha_{32} $	< 0.27	< 0.40

Table B1: 90% C.L.(1 d.o.f.) limits on various NUNM parameters. Second column shows the constraints considering only one NUNM parameter at a time, while other NUNM parameters are assumed to be zero in the fit. Third column depicts the bounds on a given NUNM parameter when all other NUNM parameters and the phases associated with the off-diagonal parameters are kept free in the fit. True values of the standard oscillation parameters are taken from Table 2. We marginalize over θ_{23} and δ_{CP} in the fit (see text for details).

and 2.9). Note that $|\alpha_{21}|$ does not have any correlation with the other NUNM parameters in $\nu_\mu \rightarrow \nu_e$ appearance channel but it is strongly correlated with α_{22} , $|\alpha_{31}|$, and $|\alpha_{32}|$ in $\nu_\mu \rightarrow \nu_\mu$ disappearance channel which reduces its sensitivity by roughly 40%, when we keep the other NUNM parameter free in the fit. Similar correlations among the NUNM parameters are also responsible for worsening the bounds on $|\alpha_{31}|$ by 60%, $|\alpha_{32}|$ by 45%, and by a factor of two for α_{33} when we marginalize over the other NUNM parameters in the fit.

References

- [1] L. R. Evans and P. Bryant, *LHC Machine*, *JINST* **3** (2008) S08001. 164 p. This report is an abridged version of the LHC Design Report (CERN-2004-003).
- [2] M. Dine, *TASI lectures on the strong CP problem*, in *Theoretical Advanced Study Institute in Elementary Particle Physics (TASI 2000): Flavor Physics for the Millennium*, pp. 349–369, 6, 2000. [hep-ph/0011376](#).
- [3] M. C. Gonzalez-Garcia and Y. Nir, *Neutrino Masses and Mixing: Evidence and Implications*, *Rev. Mod. Phys.* **75** (2003) 345–402, [[hep-ph/0202058](#)].
- [4] F. Jegerlehner, *The hierarchy problem of the electroweak Standard Model revisited*, [arXiv:1305.6652](#).
- [5] G. 't Hooft, *Naturalness, chiral symmetry, and spontaneous chiral symmetry breaking*, *NATO Sci. Ser. B* **59** (1980) 135–157.
- [6] M. McCullough, *Lectures on Physics Beyond the Standard Model.*, in *6th Tri-Institute Summer School on Elementary Particles*, 2018.
- [7] R. D. Peccei, *The Strong CP problem and axions*, *Lect. Notes Phys.* **741** (2008) 3–17, [[hep-ph/0607268](#)].
- [8] R. D. Peccei and H. R. Quinn, *CP conservation in the presence of pseudoparticles*, *Phys. Rev. Lett.* **38** (Jun, 1977) 1440–1443.
- [9] **Super-Kamiokande** Collaboration, K. Abe et al., *Solar neutrino results in Super-Kamiokande-III*, *Phys. Rev. D* **83** (2011) 052010, [[arXiv:1010.0118](#)].
- [10] **Super-Kamiokande** Collaboration, K. Abe et al., *Atmospheric neutrino oscillation analysis with external constraints in Super-Kamiokande I-IV*, *Phys. Rev. D* **97** (2018), no. 7 072001, [[arXiv:1710.09126](#)].
- [11] **Super-Kamiokande** Collaboration, M. Jiang et al., *Atmospheric Neutrino Oscillation Analysis with Improved Event Reconstruction in Super-Kamiokande IV*, *PTEP* **2019** (2019), no. 5 053F01, [[arXiv:1901.03230](#)].
- [12] **IceCube** Collaboration, M. G. Aartsen et al., *Measurement of Atmospheric Neutrino Oscillations at 6–56 GeV with IceCube DeepCore*, *Phys. Rev. Lett.* **120** (2018), no. 7 071801, [[arXiv:1707.07081](#)].
- [13] **ANTARES** Collaboration, A. Albert et al., *Measuring the atmospheric neutrino oscillation parameters and constraining the 3+1 neutrino model with ten years of ANTARES data*, *JHEP* **06** (2019) 113, [[arXiv:1812.08650](#)].
- [14] **Daya Bay** Collaboration, D. Adey et al., *Measurement of the Electron Antineutrino Oscillation with 1958 Days of Operation at Daya Bay*, *Phys. Rev. Lett.* **121** (2018), no. 24 241805, [[arXiv:1809.02261](#)].
- [15] **RENO** Collaboration, G. Bak et al., *Measurement of Reactor Antineutrino Oscillation Amplitude and Frequency at RENO*, *Phys. Rev. Lett.* **121** (2018), no. 20 201801, [[arXiv:1806.00248](#)].
- [16] **MINOS** Collaboration, P. Adamson et al., *Measurement of Neutrino and Antineutrino Oscillations Using Beam and Atmospheric Data in MINOS*, *Phys. Rev. Lett.* **110** (2013), no. 25 251801, [[arXiv:1304.6335](#)].

- [17] **T2K** Collaboration, K. Abe et al., *Constraint on the matter–antimatter symmetry-violating phase in neutrino oscillations*, *Nature* **580** (2020), no. 7803 339–344, [[arXiv:1910.03887](#)]. [Erratum: *Nature* 583, E16 (2020)].
- [18] **NOvA** Collaboration, M. A. Acero et al., *An Improved Measurement of Neutrino Oscillation Parameters by the NOvA Experiment*, [arXiv:2108.08219](#).
- [19] NuFIT 5.0 (2020), <http://www.nu-fit.org/>.
- [20] I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, T. Schwetz, and A. Zhou, *The fate of hints: updated global analysis of three-flavor neutrino oscillations*, *JHEP* **09** (2020) 178, [[arXiv:2007.14792](#)].
- [21] P. F. de Salas, D. V. Forero, S. Gariazzo, P. Martínez-Miravé, O. Mena, C. A. Ternes, M. Tórtola, and J. W. F. Valle, *2020 global reassessment of the neutrino oscillation picture*, *JHEP* **02** (2021) 071, [[arXiv:2006.11237](#)].
- [22] F. Capozzi, E. Di Valentino, E. Lisi, A. Marrone, A. Melchiorri, and A. Palazzo, *Unfinished fabric of the three neutrino paradigm*, *Phys. Rev. D* **104** (2021), no. 8 083031, [[arXiv:2107.00532](#)].
- [23] **JUNO** Collaboration, F. An et al., *Neutrino Physics with JUNO*, *J. Phys. G* **43** (2016), no. 3 030401, [[arXiv:1507.05613](#)].
- [24] **JUNO** Collaboration, A. Abusleme et al., *JUNO Physics and Detector*, [arXiv:2104.02565](#).
- [25] **DUNE** Collaboration, B. Abi et al., *Deep Underground Neutrino Experiment (DUNE), Far Detector Technical Design Report, Volume I Introduction to DUNE*, *JINST* **15** (2020), no. 08 T08008, [[arXiv:2002.02967](#)].
- [26] **DUNE** Collaboration, B. Abi et al., *Long-baseline neutrino oscillation physics potential of the DUNE experiment*, *Eur. Phys. J. C* **80** (2020), no. 10 978, [[arXiv:2006.16043](#)].
- [27] **DUNE** Collaboration, B. Abi et al., *Experiment Simulation Configurations Approximating DUNE TDR*, [arXiv:2103.04797](#).
- [28] **DUNE** Collaboration, A. A. Abud et al., *Low exposure long-baseline neutrino oscillation sensitivity of the DUNE experiment*, [arXiv:2109.01304](#).
- [29] **Hyper-Kamiokande Proto-** Collaboration, K. Abe et al., *Physics potential of a long-baseline neutrino oscillation experiment using a J-PARC neutrino beam and Hyper-Kamiokande*, *PTEP* **2015** (2015) 053C02, [[arXiv:1502.05199](#)].
- [30] **Hyper-Kamiokande** Collaboration, K. Abe et al., *Hyper-Kamiokande Design Report*, [arXiv:1805.04163](#).
- [31] **Hyper-Kamiokande** Collaboration, K. Abe et al., *Physics potentials with the second Hyper-Kamiokande detector in Korea*, *PTEP* **2018** (2018), no. 6 063C01, [[arXiv:1611.06118](#)].
- [32] C. S. Fong, H. Minakata, and H. Nunokawa, *A framework for testing leptonic unitarity by neutrino oscillation experiments*, *JHEP* **02** (2017) 114, [[arXiv:1609.08623](#)].
- [33] R. N. Mohapatra and J. W. F. Valle, *Neutrino Mass and Baryon Number Nonconservation in Superstring Models*, *Phys. Rev. D* **34** (1986) 1642.
- [34] R. N. Mohapatra and G. Senjanovic, *Neutrino Mass and Spontaneous Parity Nonconservation*, *Phys. Rev. Lett.* **44** (1980) 912.

- [35] E. K. Akhmedov, M. Lindner, E. Schnapka, and J. W. F. Valle, *Dynamical left-right symmetry breaking*, *Phys. Rev. D* **53** (1996) 2752–2780, [[hep-ph/9509255](#)].
- [36] M. Malinsky, T. Ohlsson, and H. Zhang, *Non-unitarity effects in a realistic low-scale seesaw model*, *Phys. Rev. D* **79** (2009) 073009, [[arXiv:0903.1961](#)].
- [37] F. J. Escrihuela, D. V. Forero, O. G. Miranda, M. Tortola, and J. W. F. Valle, *On the description of nonunitary neutrino mixing*, *Phys. Rev. D* **92** (2015), no. 5 053009, [[arXiv:1503.08879](#)]. [Erratum: *Phys.Rev.D* 93, 119905 (2016)].
- [38] A. de Gouvêa and A. Kobach, *Global Constraints on a Heavy Neutrino*, *Phys. Rev. D* **93** (2016), no. 3 033005, [[arXiv:1511.00683](#)].
- [39] M. Blennow, P. Coloma, E. Fernandez-Martinez, J. Hernandez-Garcia, and J. Lopez-Pavon, *Non-Unitarity, sterile neutrinos, and Non-Standard neutrino Interactions*, *JHEP* **04** (2017) 153, [[arXiv:1609.08637](#)].
- [40] F. J. Escrihuela, D. V. Forero, O. G. Miranda, M. Tórtola, and J. W. F. Valle, *Probing CP violation with non-unitary mixing in long-baseline neutrino oscillation experiments: DUNE as a case study*, *New J. Phys.* **19** (2017), no. 9 093005, [[arXiv:1612.07377](#)].
- [41] E. K. Akhmedov, M. Lindner, E. Schnapka, and J. W. F. Valle, *Left-right symmetry breaking in NJL approach*, *Phys. Lett. B* **368** (1996) 270–280, [[hep-ph/9507275](#)].
- [42] M. Malinsky, T. Ohlsson, Z.-z. Xing, and H. Zhang, *Non-unitary neutrino mixing and CP violation in the minimal inverse seesaw model*, *Phys. Lett. B* **679** (2009) 242–248, [[arXiv:0905.2889](#)].
- [43] Y.-F. Li and S. Luo, *Neutrino Oscillation Probabilities in Matter with Direct and Indirect Unitarity Violation in the Lepton Mixing Matrix*, *Phys. Rev. D* **93** (2016), no. 3 033008, [[arXiv:1508.00052](#)].
- [44] S. Parke and M. Ross-Lonergan, *Unitarity and the three flavor neutrino mixing matrix*, *Phys. Rev. D* **93** (2016), no. 11 113009, [[arXiv:1508.05095](#)].
- [45] S.-F. Ge, P. Pasquini, M. Tortola, and J. W. F. Valle, *Measuring the leptonic CP phase in neutrino oscillations with nonunitary mixing*, *Phys. Rev. D* **95** (2017), no. 3 033005, [[arXiv:1605.01670](#)].
- [46] O. G. Miranda, M. Tortola, and J. W. F. Valle, *New ambiguity in probing CP violation in neutrino oscillations*, *Phys. Rev. Lett.* **117** (2016), no. 6 061804, [[arXiv:1604.05690](#)].
- [47] D. Dutta, P. Ghoshal, and S. Roy, *Effect of Non Unitarity on Neutrino Mass Hierarchy determination at DUNE, NO ν A and T2K*, *Nucl. Phys. B* **920** (2017) 385–401, [[arXiv:1609.07094](#)].
- [48] H. Päs and P. Sicking, *Discriminating sterile neutrinos and unitarity violation with CP invariants*, *Phys. Rev. D* **95** (2017), no. 7 075004, [[arXiv:1611.08450](#)].
- [49] S. Antusch, C. Biggio, E. Fernandez-Martinez, M. B. Gavela, and J. Lopez-Pavon, *Unitarity of the Leptonic Mixing Matrix*, *JHEP* **10** (2006) 084, [[hep-ph/0607020](#)].
- [50] L. S. Miranda, P. Pasquini, U. Rahaman, and S. Razzaque, *Searching for non-unitary neutrino oscillations in the present T2K and NO ν A data*, *Eur. Phys. J. C* **81** (2021), no. 5 444, [[arXiv:1911.09398](#)].
- [51] A. De Gouvêa, K. J. Kelly, G. V. Stenico, and P. Pasquini, *Physics with Beam Tau-Neutrino Appearance at DUNE*, *Phys. Rev. D* **100** (2019), no. 1 016004, [[arXiv:1904.07265](#)].

- [52] E. Fernandez-Martinez, J. Hernandez-Garcia, and J. Lopez-Pavon, *Global constraints on heavy neutrino mixing*, *JHEP* **08** (2016) 033, [[arXiv:1605.08774](#)].
- [53] D. V. Forero, C. Giunti, C. A. Ternes, and M. Tortola, *Nonunitary neutrino mixing in short and long-baseline experiments*, *Phys. Rev. D* **104** (2021), no. 7 075030, [[arXiv:2103.01998](#)].
- [54] S. A. R. Ellis, K. J. Kelly, and S. W. Li, *Current and Future Neutrino Oscillation Constraints on Leptonic Unitarity*, *JHEP* **12** (2020) 068, [[arXiv:2008.01088](#)].
- [55] P. Coloma, J. López-Pavón, S. Rosauero-Alcaraz, and S. Urrea, *New physics from oscillations at the DUNE near detector, and the role of systematic uncertainties*, [arXiv:2105.11466](#).
- [56] Z. Hu, J. Ling, J. Tang, and T. Wang, *Global oscillation data analysis on the 3ν mixing without unitarity*, *JHEP* **01** (2021) 124, [[arXiv:2008.09730](#)].
- [57] Y. Abe, Y. Asano, N. Haba, and T. Yamada, *Heavy neutrino mixing in the T2HK, the T2HKK and an extension of the T2HK with a detector at Oki Islands*, *Eur. Phys. J. C* **77** (2017), no. 12 851, [[arXiv:1705.03818](#)].
- [58] D. Dutta and S. Roy, *Non-Unitarity at DUNE and T2HK with Charged and Neutral Current Measurements*, *J. Phys. G* **48** (2021), no. 4 045004, [[arXiv:1901.11298](#)].
- [59] Z.-z. Xing, *Towards testing the unitarity of the 3×3 lepton flavor mixing matrix in a precision reactor antineutrino oscillation experiment*, *Phys. Lett. B* **718** (2013) 1447–1453, [[arXiv:1210.1523](#)].
- [60] W. Flieger, J. Gluza, and K. Porwit, *New limits on neutrino non-unitary mixings based on prescribed singular values*, *JHEP* **03** (2020) 169, [[arXiv:1910.01233](#)].
- [61] K. Bielas, W. Flieger, J. Gluza, and M. Gluza, *Neutrino mixing, interval matrices and singular values*, *Phys. Rev. D* **98** (2018), no. 5 053001, [[arXiv:1708.09196](#)].
- [62] S. A. R. Ellis, K. J. Kelly, and S. W. Li, *Leptonic Unitarity Triangles*, *Phys. Rev. D* **102** (2020), no. 11 115027, [[arXiv:2004.13719](#)].
- [63] S. F. King, *Parametrizing the lepton mixing matrix in terms of deviations from tri-bimaximal mixing*, *Phys. Lett. B* **659** (2008) 244–251, [[arXiv:0710.0530](#)].
- [64] S. Pakvasa, W. Rodejohann, and T. J. Weiler, *Unitary parametrization of perturbations to tribimaximal neutrino mixing*, *Phys. Rev. Lett.* **100** (2008) 111801, [[arXiv:0711.0052](#)].
- [65] P. Huber, M. Lindner, and W. Winter, *Simulation of long-baseline neutrino oscillation experiments with GLOBES (General Long Baseline Experiment Simulator)*, *Comput. Phys. Commun.* **167** (2005) 195, [[hep-ph/0407333](#)].
- [66] P. Huber, J. Kopp, M. Lindner, M. Rolinec, and W. Winter, *New features in the simulation of neutrino oscillation experiments with GLOBES 3.0: General Long Baseline Experiment Simulator*, *Comput. Phys. Commun.* **177** (2007) 432–438, [[hep-ph/0701187](#)].
- [67] **DUNE** Collaboration, A. Abed Abud et al., *Deep Underground Neutrino Experiment (DUNE) Near Detector Conceptual Design Report*, [arXiv:2103.13910](#).
- [68] K. J. Kelly, *Searches for new physics at the Hyper-Kamiokande experiment*, *Phys. Rev. D* **95** (2017), no. 11 115009, [[arXiv:1703.00448](#)].
- [69] S. K. Agarwalla, S. S. Chatterjee, and A. Palazzo, *Signatures of a Light Sterile Neutrino in T2HK*, *JHEP* **04** (2018) 091, [[arXiv:1801.04855](#)].
- [70] S. Choubey, D. Dutta, and D. Pramanik, *Measuring the Sterile Neutrino CP Phase at DUNE and T2HK*, *Eur. Phys. J. C* **78** (2018), no. 4 339, [[arXiv:1711.07464](#)].

- [71] K. Abe et al., *Letter of Intent: The Hyper-Kamiokande Experiment — Detector Design and Physics Potential* —, [arXiv:1109.3262](#).
- [72] **Hyper-Kamiokande Working Group** Collaboration, K. Abe et al., *A Long Baseline Neutrino Oscillation Experiment Using J-PARC Neutrino Beam and Hyper-Kamiokande*, 12, 2014. [arXiv:1412.4673](#).
- [73] S.-H. Seo, *Neutrino Telescope at Yemilab, Korea*, [arXiv:1903.05368](#).
- [74] **T2K** Collaboration, K. Abe et al., *T2K ND280 Upgrade - Technical Design Report*, [arXiv:1901.03750](#).
- [75] **J-PARC E61** Collaboration, E. Drakopoulou, *An intermediate water Cherenkov detector for the Hyper-Kamiokande experiment: overview and status*, *PoS ICRC2017* (2018) 1021.
- [76] J. R. Wilson, *The Hyper-K Near Detector Programme*, *J. Phys. Conf. Ser.* **1342** (2020), no. 1 012053.
- [77] P. Huber, M. Lindner, and W. Winter, *Superbeams versus neutrino factories*, *Nucl. Phys. B* **645** (2002) 3–48, [[hep-ph/0204352](#)].
- [78] G. L. Fogli, E. Lisi, A. Marrone, D. Montanino, and A. Palazzo, *Getting the most from the statistical analysis of solar neutrino oscillations*, *Phys. Rev. D* **66** (2002) 053010, [[hep-ph/0206162](#)].
- [79] M. C. Gonzalez-Garcia and M. Maltoni, *Atmospheric neutrino oscillations and new physics*, *Phys. Rev. D* **70** (2004) 033010, [[hep-ph/0404085](#)].
- [80] M. Blennow, P. Coloma, P. Huber, and T. Schwetz, *Quantifying the sensitivity of oscillation experiments to the neutrino mass ordering*, *JHEP* **03** (2014) 028, [[arXiv:1311.1822](#)].
- [81] **MINERvA** Collaboration, L. Aliaga et al., *Neutrino Flux Predictions for the NuMI Beam*, *Phys. Rev. D* **94** (2016), no. 9 092005, [[arXiv:1607.00704](#)]. [Addendum: *Phys.Rev.D* 95, 039903 (2017)].
- [82] A. Giarnetti and D. Meloni, *Probing source and detector nonstandard interaction parameters at the DUNE near detector*, *Phys. Rev. D* **104** (2021), no. 1 015027, [[arXiv:2005.10272](#)].
- [83] O. G. Miranda, P. Pasquini, M. Tórtola, and J. W. F. Valle, *Exploring the Potential of Short-Baseline Physics at Fermilab*, *Phys. Rev. D* **97** (2018), no. 9 095026, [[arXiv:1802.02133](#)].
- [84] **OPERA** Collaboration, N. Agafonova et al., *Observation of a first ν_τ candidate in the OPERA experiment in the CNGS beam*, *Phys. Lett. B* **691** (2010) 138–145, [[arXiv:1006.1623](#)].
- [85] P. Machado, H. Schulz, and J. Turner, *Tau neutrinos at DUNE: New strategies, new opportunities*, *Phys. Rev. D* **102** (2020), no. 5 053010, [[arXiv:2007.00015](#)].
- [86] A. Ghoshal, A. Giarnetti, and D. Meloni, *On the role of the ν_τ appearance in DUNE in constraining standard neutrino physics and beyond*, *JHEP* **12** (2019) 126, [[arXiv:1906.06212](#)].
- [87] I. Martinez-Soler and H. Minakata, *Measuring tau neutrino appearance probability via unitarity*, [arXiv:2109.06933](#).
- [88] P. B. Denton and J. Gehrlein, *New tau neutrino oscillation and scattering constraints on unitarity violation*, [arXiv:2109.14575](#).

- [89] P. B. Denton, *Tau Neutrino Identification in Atmospheric Neutrino Oscillations Without Particle Identification or Unitarity*, [arXiv:2109.14576](https://arxiv.org/abs/2109.14576).