

Quantum-classical correspondence principle for heat distribution in quantum Brownian motion

Jin-Fu Chen,¹ Tian Qiu,¹ and H. T. Quan^{1,2,3,*}

¹*School of Physics, Peking University, Beijing, 100871, China*

²*Collaborative Innovation Center of Quantum Matter, Beijing 100871, China*

³*Frontiers Science Center for Nano-optoelectronics,
Peking University, Beijing, 100871, China*

(Dated: November 23, 2021)

Abstract

Quantum Brownian motion, described by the Caldeira-Leggett model, brings insights to understand phenomena and essence of quantum thermodynamics, especially the quantum work and heat associated with their classical counterparts. By employing the phase-space formulation approach, we study the heat distribution of a relaxation process in the quantum Brownian motion model. The analytical result of the characteristic function of heat is obtained at any relaxation time with an arbitrary friction coefficient. By taking the classical limit, such a result approaches the heat distribution of the classical Brownian motion described by the Langevin equation, indicating the quantum-classical correspondence principle for heat distribution. We also demonstrate that the fluctuating heat at any relaxation time satisfies the exchange fluctuation theorem of heat, and its long-time limit reflects complete thermalization of the system. Our research brings justification for the definition of the quantum fluctuating heat via two-point measurements.

* Corresponding author: htquan@pku.edu.cn

I. INTRODUCTION

In the past few decades, the discovery of fluctuation theorems [1–4] and the establishment of the framework of stochastic thermodynamics [5–7] deepened our understanding about the fluctuating nature of thermodynamic quantities (such as work, heat and entropy production) in microscopic systems [8–13]. Among various fluctuation theorems, the non-equilibrium work relation [2] sharpens our understanding of the second law of thermodynamics by presenting an elegant and precise equality associating the free energy change with the fluctuating work. Such a relation was later extended to the quantum realm based on the two-point measurement definition of the quantum fluctuating work [14, 15], soon after its discovery in the classical regime. The work statistics has been widely studied in various microscopic classical and quantum systems [16–26]. Historically, the quantum-classical correspondence principle played an essential role in the development of the theory of quantum mechanics and the interpretation of the transition from quantum to classical world [27, 28]. In Refs. [19, 22, 24], it is demonstrated that the existence of the quantum-classical correspondence principle for work distribution brings justification for the definition of quantum fluctuating work via two-point measurements.

Compared to work statistics, heat statistics relevant to thermal transport associated with a nonequilibrium stationary state has been extensively studied [29–38], but the heat statistics in a finite-time quantum thermodynamic process [39–41] and its quantum-classical correspondence have been less explored. A challenge is that the precise description of the bath dynamics requires handling a huge number of degrees of freedom of the heat bath. Different approaches have been proposed to calculate the quantum fluctuating heat and its statistics, such as the non-equilibrium Green’s function approach to quantum thermal transport [29, 32, 36, 42–44] and the path-integral approach to quantum thermodynamics [45–49]. However, very few analytical results about the heat statistics have been obtained for the relaxation processes in open quantum systems. These analytical results are limited to either the relaxation dynamics described by the Lindblad master equation [39, 40] or the long-time limit independent of the relaxation dynamics [50]. On the other hand, some results about the heat statistics in the classical Brownian motion model have been reported [51–59]. How the quantum and the classical heat statistics (especially associated with the relaxation dynamics in finite time) are related to each other has not been explored so far, probably

due to the difficulty in studying the heat statistics in open quantum systems [60–62].

In this article, we study the heat statistics of a quantum Brownian motion model described by the Caldeira-Leggett Hamiltonian [48, 63–68], where the heat bath is modeled as a collection of harmonic oscillators. Although it is well known that the dynamics of such an open quantum system can approach that of the classical Brownian motion in the classical limit $\hbar \rightarrow 0$ [64], less is known about the heat statistics of this model during the finite-time relaxation process. We here focus on the relaxation process without external driving (the Hamiltonian of the system is time-independent), and the quantum fluctuating heat can thus be defined as the difference of the system energy between the initial and the final measurements [69]. Under the Ohmic spectral density, the dynamics of the composite system is exactly solvable in the continuum limit of the bath oscillators [70]. By employing the phase-space formulation approach [71–73], we obtain analytical results of the characteristic function of heat for the Caldeira-Leggett model at any relaxation time τ with an arbitrary friction coefficient κ . Previously, such an approach was employed to study the quantum corrections to work [74–76] and entropy [77, 78]. Analytical results of the heat statistics bring important insights to understand the fluctuating property of heat. By taking the classical limit $\hbar \rightarrow 0$, the heat statistics of the Caldeira-Leggett model approaches that of the classical Brownian motion model. Thus, our results verify the quantum-classical correspondence principle for heat distribution, and provide justification for the definition of the quantum fluctuating heat via two-point measurements. We also verify from the analytical results that the heat statistics satisfies the exchange fluctuation theorem of heat [4].

The rest of this article is organized as follows. In Sec. II, we introduce the Caldeira-Leggett model and define the quantum fluctuating heat. In Sec. III, the analytical results of the characteristic function of heat are obtained by employing the phase-space formulation approach. We show the quantum-classical correspondence of the heat distribution, and discuss the heat distribution in the long-time limit or with the extremely weak or strong coupling strength. The conclusion is given in Sec. IV.

II. THE CALDEIRA-LEGGETT MODEL AND THE HEAT STATISTICS

A. The Caldeira-Leggett model

The quantum Brownian motion is generally described by the Caldeira-Leggett model [64, 65], where the system is modeled as a single particle moving in a specific potential, and the heat bath is a collection of harmonic oscillators. For simplicity, we choose the harmonic potential for the system [66, 79–81], where the dynamics of such an open quantum system can be solved analytically. The system will relax to the equilibrium state at the temperature of the heat bath. We study the heat distribution of such a quantum relaxation process, and obtain analytically the characteristic function of heat and its classical correspondence based on the phase-space formulation of quantum mechanics.

The total Hamiltonian of the composite system is $H_{\text{tot}} = H_S + H_B + H_{SB}$ with each term

$$H_S = \frac{1}{2} \frac{\hat{p}_0^2}{m_0} + \frac{1}{2} m_0 \omega_0^2 \hat{q}_0^2 \quad (1)$$

$$H_B = \sum_{n=1}^N \left(\frac{1}{2} \frac{\hat{p}_n^2}{m_n} + \frac{1}{2} m_n \omega_n^2 \hat{q}_n^2 \right) \quad (2)$$

$$H_{SB} = -\hat{q}_0 \sum_{n=1}^N (C_n \hat{q}_n) + \sum_{n=1}^N \left(\frac{C_n^2}{2m_n \omega_n^2} \hat{q}_0^2 \right), \quad (3)$$

where $m_0, \omega_0, \hat{q}_0, \hat{p}_0$ ($m_n, \omega_n, \hat{q}_n, \hat{p}_n$ with $n = 1, 2, 3, \dots, N$) are the mass, frequency, position and momentum of the system (the n -th bath harmonic oscillator), and C_n is the coupling strength between the system and the n -th bath harmonic oscillator. The counter-term $\sum_n [C_n^2 / (2m_n \omega_n^2)] \hat{q}_0^2$ is included in the interaction Hamiltonian H_{SB} to cancel the frequency shift of the system.

The spectral density is defined as $J(\omega) := \sum_n [C_n^2 / (2m_n \omega_n)] \delta(\omega - \omega_n)$. We adopt an Ohmic spectral density with the Lorentz-Drude cutoff [67]

$$J(\omega) = \frac{m_0 \kappa}{\pi} \omega \frac{\Omega_0^2}{\Omega_0^2 + \omega^2}, \quad (4)$$

where κ is the friction coefficient. A sufficiently large cutoff frequency Ω_0 ($\Omega_0 \gg \omega_0$) is applied to ensure a finite counter-term, and the dynamics with the timescale exceeding $1/\Omega_0$ is Markovian. Under such a spectral density, the dissipation dynamics of the Caldeira-Leggett

model with a weak coupling strength $\kappa \ll \omega_0$ reproduces that of the classical underdamped Brownian motion when taking the classical limit $\hbar \rightarrow 0$ [64].

We assume the initial state to be a product state of the system and the heat bath

$$\rho(0) = \rho_S(0) \otimes \rho_B^G, \quad (5)$$

which makes it possible to define the quantum fluctuating heat via two-point measurements. Here, $\rho_S(0)$ is the initial state of the system, and $\rho_B^G = \exp(-\beta H_B)/Z_B(\beta)$ is the Gibbs distribution of the heat bath with the inverse temperature β and the partition function $Z_B(\beta) = \text{Tr}[\exp(-\beta H_B)]$.

B. The quantum fluctuating heat in the relaxation process

We study the heat distribution of the relaxation process based on the two-point measurement definition of the quantum fluctuating heat. When no external driving is applied to the system, the Hamiltonian of the system is time-independent. Since no work is performed during the relaxation process, the quantum fluctuating heat can be defined as

$$Q_{\nu l} = E_{\nu'}^S - E_l^S, \quad (6)$$

where E_l^S ($E_{\nu'}^S$) is the eigenenergy of the system corresponding to the outcome l (ν') at the initial (final) time $t = 0$ ($t = \tau$). The two-point measurements over the heat bath can be hardly realized due to a huge number of degrees of freedom of the heat bath [20], while the measurements over the small quantum system are much easier in principle. The positive sign corresponds to the energy flowing from the heat bath to the system.

For the system prepared in an equilibrium state, no coherence exists in the initial state, and the initial density matrix of the system commutes with the Hamiltonian of the system, $[\rho(0), H_S] = 0$. The probability of observing the transition from l and ν' is

$$p_{\tau, \nu l} = \gamma_{\tau, \nu l} p_l, \quad (7)$$

with the conditional transition probability $\gamma_{\tau, \nu l} = \text{Tr} \left[(\hat{P}_{\nu'}^S \otimes I_B) U_{\text{tot}}(\tau) (\hat{P}_l^S \otimes \rho_B^G) U_{\text{tot}}^\dagger(\tau) \right]$ and the initial probability $p_l = \text{Tr}[\rho(0) \hat{P}_l^S]$. Here, $\hat{P}_l^S = |l\rangle \langle l|$ is the projection operator corresponding to the outcome l . The heat distribution is defined as

$$P_\tau(q) := \sum_{\nu,l} \delta(q - Q_{\nu l}) p_{\tau,\nu l}. \quad (8)$$

The characteristic function of heat $\chi_\tau(\nu)$ is defined as the Fourier transform of the heat distribution $\chi_\tau(\nu) := \sum_{\nu,l} \exp[i\nu(E_\nu^S - E_l^S)] p_{\tau,\nu l}$, which can be rewritten explicitly as

$$\chi_\tau(\nu) = \text{Tr} \left[e^{i\nu H_S} U_{\text{tot}}(\tau) (e^{-i\nu H_S} \rho(0)) U_{\text{tot}}^\dagger(\tau) \right], \quad (9)$$

where $U_{\text{tot}}(\tau) = \exp(-iH_{\text{tot}}\tau/\hbar)$ is the unitary time-evolution operator of the composite system.

Our goal is to analytically calculate the characteristic function $\chi_\tau(\nu)$. Previously, the quantum-classical correspondence principle for heat statistics has been analyzed with the path-integral approach to quantum thermodynamics [48], yet the explicit result of the characteristic function (or generating function) of heat has not been obtained so far. We employ the phase-space formulation approach to solve this problem, and rewrite the characteristic function Eq. (9) into

$$\chi_\tau(\nu) = \text{Tr} \left[e^{i\nu H_S^H(\tau)} \eta(0) \right], \quad (10)$$

where the system Hamiltonian in the Heisenberg picture is

$$H_S^H(\tau) = U_{\text{tot}}^\dagger(\tau) H_S U_{\text{tot}}(\tau), \quad (11)$$

and the density-matrix-like operator $\eta(0)$ is

$$\eta(0) = [e^{-i\nu H_S} \rho_S(0)] \otimes \rho_B^G. \quad (12)$$

We express Eq. (10) with the phase-space formulation of quantum mechanics [71–76]

$$\chi_\tau(\nu) = \frac{1}{(2\pi\hbar)^{N+1}} \int d\mathbf{z} \left[e^{i\nu H_S^H(\tau)} \right]_w(\mathbf{z}) \cdot P(\mathbf{z}), \quad (13)$$

where \mathbf{z} represents a point $\mathbf{z} = [\mathbf{q}, \mathbf{p}] = [q_0, \dots, q_N, p_0, \dots, p_N]$ in the phase space of the composite system, and the integral is performed over the whole phase space. The subscript “ w ” indicates the Weyl symbol of the corresponding operator, and $P(\mathbf{z})$ is the Weyl symbol of the operator $\eta(0)$, which is explicitly defined as [71]

$$P(\mathbf{z}) := \int d\mathbf{y} \left\langle \mathbf{q} - \frac{\mathbf{y}}{2} \left| \eta(0) \right| \mathbf{q} + \frac{\mathbf{y}}{2} \right\rangle e^{\frac{i\mathbf{p}\cdot\mathbf{y}}{\hbar}}. \quad (14)$$

In the following, we will calculate the heat statistics Eq. (13) by employing the phase-space formulation approach.

III. RESULTS OF THE CHARACTERISTIC FUNCTION OF HEAT

We show a sketch of the derivation of the heat statistics $\chi_\tau(\nu)$ with the details left in Appendix A. We specifically consider the system is initially prepared at an equilibrium state $\rho_S(0) = \exp(-\beta' H_S)/Z_S(\beta')$ with the inverse temperature β' and the partition function $Z_S(\beta') = 1/[2 \sinh(\beta' \hbar \omega_0/2)]$. The heat bath is at the inverse temperature β , which is different from β' . In Eq. (13), the two Weyl symbols $\left[e^{i\nu H_S^H(\tau)} \right]_w(\mathbf{z})$ and $P(\mathbf{z})$ are obtained as

$$\left[e^{i\nu H_S^H(\tau)} \right]_w(\mathbf{z}) = \frac{1}{\cos\left(\frac{\nu \hbar \omega_0}{2}\right)} \exp\left[\frac{i}{2\hbar} \mathbf{z}^T \tilde{\mathbf{\Lambda}}_{\nu z}(\tau) \mathbf{z}\right], \quad (15)$$

and

$$P(\mathbf{z}) = \frac{2 \sinh\left(\frac{\beta' \hbar \omega_0}{2}\right)}{\cosh\left[\frac{(\beta' + i\nu) \hbar \omega_0}{2}\right]} \cdot \left[\prod_{n=1}^N 2 \tanh\left(\frac{\beta \hbar \omega_n}{2}\right) \right] \cdot \exp\left(-\frac{1}{2\hbar} \mathbf{z}^T \mathbf{\Lambda}_{\beta z} \mathbf{z}\right), \quad (16)$$

where the explicit expressions of the matrices $\tilde{\mathbf{\Lambda}}_{\nu z}(\tau)$ and $\mathbf{\Lambda}_{\beta z}$ are given in Eqs. (A10) and (A36), respectively.

Substituting Eqs. (15) and (16) into Eq. (13), the characteristic function of heat at any relaxation time τ with an arbitrary friction coefficient κ is finally obtained as

$$\chi_\tau(\nu) = \left\{ \left[(1 + i\Xi)(1 - i\Theta\Xi) - i\Xi(1 - \Theta - i\Theta\Xi) \frac{\kappa^2 \cos(2\hat{\omega}_0\tau) - 4\omega_0^2}{(\kappa^2 - 4\omega_0^2)e^{\kappa\tau}} \right]^2 + \Xi^2(1 - \Theta - i\Theta\Xi)^2 \left[\left(\frac{\kappa^2 \cos(2\hat{\omega}_0\tau) - 4\omega_0^2}{(\kappa^2 - 4\omega_0^2)e^{\kappa\tau}} \right)^2 - e^{-2\kappa\tau} \right] \right\}^{\frac{1}{2}}, \quad (17)$$

where the quantities Ξ and Θ are

$$\Xi = \frac{\tan\left(\frac{\nu \hbar \omega_0}{2}\right)}{\tanh\left[\frac{(\beta' + i\nu) \hbar \omega_0}{2}\right] - i \tan\left(\frac{\nu \hbar \omega_0}{2}\right)}, \quad (18)$$

$$\Theta = \frac{\tanh\left[\frac{(\beta' + i\nu) \hbar \omega_0}{2}\right] - i \tan\left(\frac{\nu \hbar \omega_0}{2}\right)}{\tanh\left(\frac{\beta \hbar \omega_0}{2}\right)}. \quad (19)$$

Induced by the friction, the frequency of the system harmonic oscillator is shifted to $\hat{\omega}_0 = \sqrt{\omega_0^2 - \kappa^2/4}$.

From the analytical results of the heat statistics Eq. (17), the average heat $\langle Q \rangle(\tau) = -i\partial_\nu[\ln \chi_\tau(\nu)]|_{\nu=0}$ is immediately obtained as

$$\langle Q \rangle(\tau) = \frac{\omega_0 \hbar}{2} \left[\coth \left(\frac{\beta \omega_0 \hbar}{2} \right) - \coth \left(\frac{\beta' \omega_0 \hbar}{2} \right) \right] \left[1 - \frac{\kappa^2 \cos(2\hat{\omega}_0 \tau) - 4\omega_0^2}{(\kappa^2 - 4\omega_0^2)e^{\kappa\tau}} \right], \quad (20)$$

and the variance $\text{Var}(Q)(\tau) = -\partial_\nu^2[\ln \chi_\tau(\nu)]|_{\nu=0}$ is

$$\text{Var}(Q)(\tau) = \text{I} + \text{II} \cdot e^{-\kappa\tau} + \text{III} \cdot e^{-2\kappa\tau}, \quad (21)$$

with

$$\text{I} = \frac{\omega_0^2 \hbar^2 \left[\text{csch}^2 \left(\frac{\beta \omega_0 \hbar}{2} \right) + \text{csch}^2 \left(\frac{\beta' \omega_0 \hbar}{2} \right) \right]}{4}, \quad (22)$$

$$\text{II} = \frac{\kappa^2 \cos(2\hat{\omega}_0 \tau) - 4\omega_0^2}{2\hat{\omega}_0^2} \cdot \frac{\omega_0^2 \hbar^2 \left[\coth^2 \left(\frac{\beta \omega_0 \hbar}{2} \right) + \text{csch}^2 \left(\frac{\beta' \omega_0 \hbar}{2} \right) - \coth \left(\frac{\beta \omega_0 \hbar}{2} \right) \coth \left(\frac{\beta' \omega_0 \hbar}{2} \right) \right]}{4} \quad (23)$$

$$\text{III} = \frac{\kappa^4 \cos(4\hat{\omega}_0 \tau) + 8\omega_0^2 \kappa^2 [1 - 2 \cos(2\hat{\omega}_0 \tau)] + 16\omega_0^4}{16\hat{\omega}_0^4} \cdot \frac{\omega_0^2 \hbar^2 \left[\coth \left(\frac{\beta \omega_0 \hbar}{2} \right) - \coth \left(\frac{\beta' \omega_0 \hbar}{2} \right) \right]^2}{4}. \quad (24)$$

Similarly, one can calculate the higher cumulants from the analytical results of the heat statistics. In the following, we will examine the properties of the heat statistics of the quantum Brownian motion.

A. Quantum-classical correspondence principle for heat statics and the exchange fluctuation theorem of heat

We further take the classical limit $\hbar \rightarrow 0$, or more rigorously $\beta \hbar \omega_0 \rightarrow 0$. The two quantities approach $\Xi \rightarrow \nu/\beta'$ and $\Theta \rightarrow \beta'/\beta$, and the characteristic function of heat [Eq. (17)] becomes

$$\chi_\tau^{\text{cl}}(\nu) = \left\{ \left[\left(1 + i \frac{\nu}{\beta'} \right) \left(1 - i \frac{\nu}{\beta} \right) - \frac{i\nu(\beta - \beta' - i\nu)}{\beta\beta'} \frac{\kappa^2 \cos(2\hat{\omega}_0 \tau) - 4\omega_0^2}{(\kappa^2 - 4\omega_0^2)e^{\kappa\tau}} \right]^2 + \nu^2 \left(\frac{\beta - \beta' - i\nu}{\beta\beta'} \right)^2 \left[\left(\frac{\kappa^2 \cos(2\hat{\omega}_0 \tau) - 4\omega_0^2}{(\kappa^2 - 4\omega_0^2)e^{\kappa\tau}} \right)^2 - e^{-2\kappa\tau} \right] \right\}^{-\frac{1}{2}}, \quad (25)$$

which is consistent with the results obtained from the classical Brownian motion described by the Kramers equation (see Ref. [58] or Appendix C). The average heat is

$$\langle Q^{\text{cl}} \rangle (\tau) = \frac{\beta' - \beta}{\beta\beta'} \left[1 - \frac{\kappa^2 \cos(2\hat{\omega}_0\tau) - 4\omega_0^2}{(\kappa^2 - 4\omega_0^2)e^{\kappa\tau}} \right], \quad (26)$$

and the variance $\text{Var}(Q^{\text{cl}})(\tau) = -\partial_\nu^2[\ln \chi_\tau^{\text{cl}}(\nu)]|_{\nu=0}$ is

$$\text{Var}(Q^{\text{cl}})(\tau) = \text{I}^{\text{cl}} + \text{II}^{\text{cl}} \cdot e^{-\kappa\tau} + \text{III}^{\text{cl}} \cdot e^{-2\kappa\tau}, \quad (27)$$

with

$$\text{I}^{\text{cl}} = \frac{\beta^2 + \beta'^2}{\beta^2\beta'^2} \quad (28)$$

$$\text{II}^{\text{cl}} = \frac{\kappa^2 \cos(2\hat{\omega}_0\tau) - 4\omega_0^2}{2\hat{\omega}_0^2} \cdot \frac{\beta^2 - \beta\beta' + \beta'^2}{\beta^2\beta'^2} \quad (29)$$

$$\text{III}^{\text{cl}} = \frac{\kappa^4 \cos(4\hat{\omega}_0\tau) + 8\omega_0^2\kappa^2[1 - 2\cos(2\hat{\omega}_0\tau)] + 16\omega_0^4}{16\hat{\omega}_0^4} \cdot \frac{(\beta - \beta')^2}{\beta^2\beta'^2}. \quad (30)$$

From Eq. (17) [or the classical counterpart Eq. (25)], one can see the characteristic function of heat exhibits the following symmetry

$$\chi_\tau(\nu) = \chi_\tau[-i(\beta - \beta') - \nu], \quad (31)$$

which shows the heat distribution satisfies the exchange fluctuation theorem of heat in the differential form $P_\tau(Q)/P_\tau(-Q) = \exp[-(\beta - \beta')Q]$ [4]. By setting $\nu = 0$, we obtain the relation $\chi_\tau[-i(\beta - \beta')] = \chi_\tau(0) = 1$, which is exactly the exchange fluctuation theorem of heat in the integral form $\langle \exp[(\beta - \beta')Q] \rangle = 1$.

B. Long-time limit

In the long-time limit $\tau \rightarrow \infty$, the characteristic functions of heat [Eqs. (17) and (25)] become

$$\chi_\infty(\nu) = \frac{(1 - e^{-\beta'\omega_0\hbar})(1 - e^{-\beta\omega_0\hbar})}{(1 - e^{-(\beta'+i\nu)\omega_0\hbar})(1 - e^{-(\beta-i\nu)\omega_0\hbar})}, \quad (32)$$

and

$$\chi_{\infty}^{\text{cl}}(\nu) = \frac{\beta' \beta}{(\beta' + i\nu)(\beta - i\nu)}. \quad (33)$$

Such results, independent of the relaxation dynamics, are in the form

$$\chi_{\text{th}}(\nu) = \frac{Z_S(\beta' + i\nu)Z_S(\beta - i\nu)}{Z_S(\beta')Z_S(\beta)}, \quad (34)$$

reflecting complete thermalization of the system [53]. For example, the relaxation of a harmonic oscillator governed by the quantum-optical master equation gives the identical characteristic function of heat in the long-time limit [39]. In Appendix D, we demonstrate that the characteristic function of heat for any relaxation processes with complete thermalization is always in the form of Eq. (34). With the simple expressions (32) and (33) of the characteristic functions, the heat distributions are obtained from the inverse Fourier transform as

$$P_{\infty}(q) = \begin{cases} \frac{(1-e^{-\beta'\omega_0\hbar})(1-e^{-\beta\omega_0\hbar})}{1-e^{-(\beta'+\beta)\omega_0\hbar}} \sum_{j=0}^{\infty} \delta(q - j\omega_0\hbar) e^{-\beta q} & q \geq 0 \\ \frac{(1-e^{-\beta'\omega_0\hbar})(1-e^{-\beta\omega_0\hbar})}{1-e^{-(\beta'+\beta)\omega_0\hbar}} \sum_{j=1}^{\infty} \delta(q + j\omega_0\hbar) e^{\beta' q} & q < 0 \end{cases}, \quad (35)$$

and

$$P_{\infty}^{\text{cl}}(q) = \begin{cases} \frac{\beta' \beta}{\beta' + \beta} e^{-\beta q} & q \geq 0 \\ \frac{\beta' \beta}{\beta' + \beta} e^{\beta' q} & q < 0 \end{cases}, \quad (36)$$

which are exactly the same as the long-time results obtained in Ref. [39].

C. Weak/Strong-coupling limit in finite time

In the weak-coupling limit $\kappa \ll \omega_0$, the characteristic function of heat [Eq. (17)] becomes

$$\chi_{\tau}^{\text{w}}(\nu) = \frac{1}{(1 + i\nu\Xi)(1 - i\nu\Xi\Theta)(1 - e^{-\kappa\tau}) + e^{-\kappa\tau}}. \quad (37)$$

There is only one relaxation timescale associated to κ . Such situation corresponds to the highly underdamped regime of the classical Brownian motion, and a systematic method has been proposed to study the heat distribution [56] as well as the work distribution under an external driving [82, 83].




Figure 1. The cumulative heat distribution function $\Pr(Q < q)$. The choices of the parameters are given in the main text. We compare the results of the Caldeira-Leggett model (blue solid, orange dotted and green dot-dashed curves) in Eq. (17) and those of the classical Brownian motion (black dashed curve) in Eq. (25). The rescaled relaxation time is $\tilde{\tau} = \kappa\tau = 1$ in the upper subfigures and $\tilde{\tau} = 10$ in the lower subfigures. The left, middle and right subfigures illustrate the results for the weak ($\kappa = 0.01$), intermediate ($\kappa = 1$) and strong coupling strength ($\kappa = 100$).

In the strong coupling limit $\kappa \gg \omega_0$, the characteristic function of heat [Eq. (17)] becomes

$$\chi_\tau^s(\nu) = \frac{1}{\sqrt{(1 + i\Xi)(1 - i\Xi\Theta)(1 - e^{-2\kappa\tau}) + e^{-2\kappa\tau}}} \times \frac{1}{\sqrt{(1 + i\Xi)(1 - i\Xi\Theta) \left(1 - e^{-\frac{2\omega_0^2}{\kappa}\tau}\right) + e^{-\frac{2\omega_0^2}{\kappa}\tau}}}. \quad (38)$$

The relaxation timescales of the momentum (the first factor) and the coordinate (the second factor) are separated. The long-time limits of both Eqs. (37) and (38) are equal to Eq. (32). In classical thermodynamics, the usual overdamped approximation neglects the motion of the momentum, hence the heat statistics derived under such an approximation is incomplete [52]. Actually, the momentum degree of freedom also contributes to the heat statistics.




Figure 2. The evolution of the mean value $\langle Q \rangle(\tau)$ (upper subfigures) and the variance $\text{Var}(Q)(\tau)$ (lower subfigures) of the heat statistics as functions of the rescaled time $\tilde{\tau} = \kappa\tau$.

D. Numerical results

In Fig. 1, we show the cumulative heat distribution function $\text{Pr}(Q < q) := \int_{-\infty}^q P_{\tau}(q')dq'$ with different friction coefficients $\kappa = 0.01, 1$ and 100 at the rescaled relaxation time $\tilde{\tau} = \kappa\tau = 1$ and 10 . We set the mass $m_0 = 1$ and the frequency $\omega_0 = 1$ for the system harmonic oscillator, the inverse temperatures $\beta = 1$ and $\beta' = 2$ for the initial equilibrium states of the heat bath and the system, respectively. The Planck constant is set to be $\hbar = 1, 0.5, 0.1$. With the decrease of \hbar , the quantum results [Eq. (17)] approaches the classical result [Eq. (25)]. The quantum-classical correspondence of the heat distribution is thus demonstrated for generic values of the friction coefficient κ .

For $\kappa = 0.01$ and 1 , complete thermalization is achieved at $\tilde{\tau} = 10$. The left-lower and middle-lower subfigures show the identical distribution characterized by Eqs. (35) and (36). For $\kappa = 100$, the momentum degree of freedom has been thermalized [$\exp(-2\tilde{\tau}) \approx 0$ in Eq. (38)] while the coordinate degree of freedom remains frozen [$\exp[-2(\omega_0^2/\kappa^2)\tilde{\tau}] \approx 1$ in Eq. (38)]. Thus, the distribution in the right-lower subfigure is different from the middle-lower subfigure.

In Fig. 2, we illustrate the results of the mean value $\langle Q \rangle(\tau)$ and the variance $\text{Var}(Q)(\tau)$ with different friction coefficients $\kappa = 0.01, 1$ and 100 . The parameters are the same as those in Fig. 1. The quantum results approach the classical results with the decrease of

\hbar . For $\kappa = 0.01$ and 1 (left and middle subfigures), complete thermalization is reached when $\tilde{\tau} > 5$. The mean value and the variance approach $\lim_{\tau \rightarrow \infty} \langle Q^{\text{cl}} \rangle (\tau) = 1/\beta - 1/\beta'$ and $\lim_{\tau \rightarrow \infty} \text{Var} (Q^{\text{cl}}) (\tau) = 1/\beta^2 + 1/\beta'^2$ (gray horizontal lines). For $\kappa = 100$ (right subfigures), only the momentum degree of freedom is thermalized at this timescale. Thus, the mean value and the variance take half value of their long-time limits. When the coordinate degree of freedom is also thermalized in the long-time limit ($\tilde{\tau} \gg \kappa^2/\omega_0^2 = 10^4$), the mean value and the variance are expected to approach the same values as those in the middle subfigures.

IV. CONCLUSION

Previously, the heat statistics of the relaxation processes has been studied analytically in open quantum systems described by the Lindblad master equation [39, 40, 50]. However, due to the rotating wave approximation and other approximations. Such quantum systems do not possess a well-defined classical counterpart. Hence, the quantum-classical correspondence principle for heat distribution has not been well established.

In this paper, we study the heat statistics of the quantum Brownian motion model described by the Caldeira-Leggett Hamiltonian, in which the bath dynamics is explicitly considered. By employing the phase-space formulation approach, we obtain the analytical expressions of the characteristic function of heat at any relaxation time τ with an arbitrary friction coefficient κ . Analytical results of heat statistics bring important insights to the studies of quantum thermodynamics. For example, in the classical limit, our results approach the heat statistics of the classical Brownian motion. Thus, the quantum-classical correspondence principle for heat statistics is verified in this model. Our analytical results provide justification for the definition of quantum fluctuating heat via two-point measurements.

We have also discussed the characteristic function of heat in the long-time limit or with the extremely weak/strong coupling strength. In the long-time limit, the form of the characteristic function of heat reflects complete thermalization of the system. In addition, from the analytical expressions of the heat statistics, we can immediately verify the exchange fluctuation theorem of heat. The phase-space formulation can be further utilized to study the joint statistics of work and heat in a driven open quantum system, which will be beneficial to explore the fluctuations of power and efficiency in finite-time quantum heat engines.

ACKNOWLEDGMENTS

H. T. Quan acknowledges support from the National Natural Science Foundation of China under Grants No. 11775001, No. 11534002, and No. 11825001. This paper is dedicated to Prof. Wojciech Zurek on the occasion of his 70th birthday for his kind and generous supports to one of the authors (H. T. Quan), and for his many insightful contributions to our understanding about the quantum to classical transition.

Appendix A: Derivation to the characteristic function of heat (17)

We show the detailed derivation to the characteristic function of heat $\chi_\tau(\nu)$. We first calculate the two Wigner functions $\left[e^{i\nu H_S^H(t)} \right]_w(\mathbf{z})$ and $P(\mathbf{z})$. Then the final result Eq. (17) is obtained from Eq. (13).

1. $\left[e^{i\nu H_S^H(t)} \right]_w(\mathbf{z})$

With the quadratic Hamiltonian $H_S^H(t)$, the Wigner function $\left[e^{i\nu H_S^H(t)} \right]_w(\mathbf{z})$ is [78, 81]

$$\left[e^{i\nu H_S^H(t)} \right]_w(\mathbf{z}) = \frac{1}{\cos\left(\frac{\omega_0 \hbar \nu}{2}\right)} \exp \left[i \frac{m_0 \omega_0}{\hbar} \tan\left(\frac{\omega_0 \hbar \nu}{2}\right) q_0^2(t) + i \frac{1}{m_0 \hbar \omega_0} \tan\left(\frac{\omega_0 \hbar \nu}{2}\right) p_0^2(t) \right] \quad (\text{A1})$$

$$= \frac{1}{\cos\left(\frac{\omega_0 \hbar \nu}{2}\right)} \exp \left[\frac{i}{2\hbar} \mathbf{z}^T(t) \mathbf{\Lambda}_{\nu z} \mathbf{z}(t) \right], \quad (\text{A2})$$

where $\mathbf{z}(t)$ gives the trajectory in the phase space determined by the initial point $\mathbf{z}(0) = \mathbf{z}$, and $\mathbf{\Lambda}_{\nu z}$ is a rank-2 diagonal matrix

$$\mathbf{\Lambda}_{\nu z} = \begin{pmatrix} 2m_0\omega_0 \tan\left(\frac{\omega_0 \hbar \nu}{2}\right) & & & \\ & \mathbf{O} & & \\ & & \frac{2}{m_0\omega_0} \tan\left(\frac{\omega_0 \hbar \nu}{2}\right) & \\ & & & \mathbf{O} \end{pmatrix}, \quad (\text{A3})$$

with an $N \times N$ zero matrix \mathbf{O} . The unlisted elements are zeros. The trajectory $\mathbf{z}(t)$ satisfies the classical equation of motion (also the equation of motion in the Heisenberg picture)

We rewrite the quadratic form into $\mathbf{z}^T(t)\mathbf{\Lambda}_{\nu z}\mathbf{z}(t) = \mathbf{z}^T(0)\tilde{\mathbf{\Lambda}}_{\nu z}(t)\mathbf{z}(0)$ with

$$\tilde{\mathbf{\Lambda}}_{\nu z}(t) = \exp(\mathbf{L}^T t) \mathbf{\Lambda}_{\nu z} \exp(\mathbf{L} t). \quad (\text{A10})$$

We next carry out every element in Eq. (A9) through the Laplace transforms of Eqs. (A4)-(A7)

$$s\tilde{q}_0(s) - q_0(0) = \frac{\tilde{p}_0(s)}{m_0}, \quad (\text{A11})$$

$$s\tilde{q}_n(s) - q_n(0) = \frac{\tilde{p}_n(s)}{m_n}, \quad (\text{A12})$$

$$s\tilde{p}_0(s) - p_0(0) = -m_0\tilde{\omega}_0^2\tilde{q}_0(s) + \sum_n C_n\tilde{q}_n(s), \quad (\text{A13})$$

$$s\tilde{p}_n(s) - p_n(0) = -m_n\omega_n^2\tilde{q}_n(s) + C_n\tilde{q}_0(s). \quad (\text{A14})$$

Representing $\tilde{q}_n(s)$ and $\tilde{p}_n(s)$ with $\tilde{q}_0(s)$ and the initial conditions, we obtain

$$\left\{ s^2 + \tilde{\omega}_0^2 - \sum_n \left[\frac{C_n^2}{m_0 m_n (s^2 + \omega_n^2)} \right] \right\} \tilde{q}_0(s) = \dot{q}_0(0) + sq_0(0) + \sum_n \frac{C_n}{m_0} \left[\frac{\dot{q}_n(0) + sq_n(0)}{s^2 + \omega_n^2} \right]. \quad (\text{A15})$$

Under the Ohmic spectral density [Eq. (4)], the above equation is simplified to

$$(s^2 + \kappa s + \omega_0^2)\tilde{q}_0(s) = \dot{q}_0(0) + sq_0(0) + \sum_n \frac{C_n}{m_0} \left[\frac{\dot{q}_n(0) + sq_n(0)}{s^2 + \omega_n^2} \right], \quad (\text{A16})$$

where the summation on the left-hand side of Eq. (A15) can be approximately expressed as

$$\sum_n \left[\frac{C_n^2}{m_0 m_n (s^2 + \omega_n^2)} \right] \approx -\kappa s + \sum_n \frac{C_n^2}{m_0 m_n \omega_n^2}, \quad (\text{A17})$$

with a large cutoff frequency Ω_0 . The inverse Laplace transform gives the differential equation of $q_0(t)$ as

$$\ddot{q}_0(t) + \kappa\dot{q}_0(t) + \omega_0^2 q_0(t) = \underbrace{-\kappa q_0(0)\delta(t)}_{\text{initial velocity change}} + \underbrace{\sum_n \frac{C_n}{m_0} \left[\dot{q}_n(0) \frac{\sin(\omega_n t)}{\omega_n} + q_n(0) \cos(\omega_n t) \right]}_{\text{stochastic force}}. \quad (\text{A18})$$

On the right-hand side, the second term presents the stochastic force induced by the heat bath; the first term indicates an abrupt velocity change $-\kappa q_0(0)$ of the system particle at the

initial time $t = 0$ [63, 84, 85]. The sudden change of velocity occurs for the system harmonic oscillator when the coupling between the system and the heat bath is switched on. Such an initial slippage is caused by the assumption of the initial product state. To avoid such an initial discontinuous problem, we drop the first term by considering the particle motion as starting at $t = 0+$ [70]. Under such a modification, the Caldeira-Leggett model can reproduce the complete Langevin equation with an arbitrary friction coefficient κ for both the underdamped and the overdamped regimes, and the heat distribution of the Caldeira-Leggett model approaches that of the classical Brownian motion described by the Kramers equation [86]. In Appendix B, for the classical counterpart of the Caldeira-Leggett model, we show the initial slippage can be naturally eliminated by choosing another initial state.

After dropping the first term, Eq. (A16) becomes

$$(s^2 + \kappa s + \omega_0^2)\tilde{q}_0(s) = \dot{q}_0(0) + (\kappa + s)q_0(0) + \sum_n \frac{C_n}{m_0} \left[\frac{\dot{q}_n(0) + s q_n(0)}{s^2 + \omega_n^2} \right]. \quad (\text{A19})$$

The solutions to $\tilde{q}_0(s)$ and $\tilde{q}_n(s)$ follow immediately as

$$\tilde{q}_0(s) = \frac{\dot{q}_0(0) + (\kappa + s)q_0(0) + \sum_n \frac{C_n}{m_0} \frac{\dot{q}_n(0) + s q_n(0)}{s^2 + \omega_n^2}}{s^2 + \kappa s + \omega_0^2}, \quad (\text{A20})$$

$$\tilde{q}_n(s) = \frac{\dot{q}_n(0) + s q_n(0)}{s^2 + \omega_n^2} + \frac{C_n}{m_n} \cdot \frac{\dot{q}_0(0) + (\kappa + s)q_0(0) + \sum_l \frac{C_l}{m_0} \frac{\dot{q}_l(0) + s q_l(0)}{s^2 + \omega_l^2}}{(s^2 + \omega_n^2)(s^2 + \kappa s + \omega_0^2)}. \quad (\text{A21})$$

With the inverse Laplace transform, the elements in the matrix $\exp(\mathbf{L}t)$ [Eq. (A9)] are determined by

$$\begin{pmatrix} q_0(t) \\ q_1(t) \\ \dots \\ q_N(t) \end{pmatrix} = \begin{pmatrix} \alpha_0 & \alpha_1 & \alpha_2 & \dots & \alpha_N & \beta_0 & \beta_1 & \beta_2 & \dots & \beta_N \\ \gamma_1 & \Lambda_{11} & \Lambda_{12} & \dots & \Lambda_{1N} & \xi_1 & \Delta_{11} & \Delta_{12} & \dots & \Delta_{1N} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \gamma_N & \Lambda_{N1} & \Lambda_{N2} & \dots & \Lambda_{NN} & \xi_N & \Delta_{N1} & \Delta_{N2} & \dots & \Delta_{NN} \end{pmatrix} \begin{pmatrix} q_0(0) \\ q_1(0) \\ \dots \\ q_N(0) \\ \dot{q}_0(0) \\ \dot{q}_1(0) \\ \dots \\ \dot{q}_N(0) \end{pmatrix}, \quad (\text{A22})$$

where the elements in the matrix of the right-hand side are explicitly solved as [70]

$$\alpha_0 = e^{-\frac{\kappa t}{2}} \left[\cos(\hat{\omega}_0 t) + \frac{\kappa}{2\hat{\omega}_0} \sin(\hat{\omega}_0 t) \right], \quad (\text{A23})$$

$$\beta_0 = \frac{e^{-\frac{\kappa t}{2}}}{\hat{\omega}_0} \sin(\hat{\omega}_0 t), \quad (\text{A24})$$

$$\alpha_n = \frac{C_n}{m_0} f_n(t), \quad (\text{A25})$$

$$\beta_n = \frac{C_n}{m_0} g_n(t), \quad (\text{A26})$$

$$\gamma_n = \frac{C_n}{m_n} [f_n(t) + \kappa g_n(t)], \quad (\text{A27})$$

$$\xi_n = \frac{C_n}{m_n} g_n(t), \quad (\text{A28})$$

$$\Lambda_{nl} = \delta_{nl} \cos(\omega_n t) + \frac{C_n C_l}{m_n m_0} F_{nl}(t), \quad (\text{A29})$$

$$\Delta_{nl} = \frac{\delta_{nl}}{\omega_n} \sin(\omega_n t) + \frac{C_n C_l}{m_n m_0} G_{nl}(t). \quad (\text{A30})$$

The functions $f_n(t)$, $g_n(t)$, $F_{nl}(t)$ and $G_{nl}(t)$ are explicitly

$$f_n(t) = \mathcal{L}^{-1} \left[\frac{s}{(s^2 + \kappa s + \omega_0^2)(s^2 + \omega_n^2)} \right], \quad (\text{A31})$$

$$g_n(t) = \mathcal{L}^{-1} \left[\frac{1}{(s^2 + \kappa s + \omega_0^2)(s^2 + \omega_n^2)} \right], \quad (\text{A32})$$

$$F_{nl}(t) = \mathcal{L}^{-1} \left[\frac{s}{(s^2 + \kappa s + \omega_0^2)(s^2 + \omega_n^2)(s^2 + \omega_l^2)} \right], \quad (\text{A33})$$

$$G_{nl}(t) = \mathcal{L}^{-1} \left[\frac{1}{(s^2 + \kappa s + \omega_0^2)(s^2 + \omega_n^2)(s^2 + \omega_l^2)} \right], \quad (\text{A34})$$

where $\mathcal{L}^{-1}(\cdot)$ denotes the inverse Laplace transform with $\mathcal{L}(\cdot) = \int_0^\infty (\cdot) e^{-st} dt$.

2. $P(\mathbf{z})$

$P(\mathbf{z})$ is the Wigner function of the state $\eta(0)$ for the composite system [78, 81]

$$P(\mathbf{z}) = \frac{2 \sinh\left(\frac{\beta' \hbar \omega_0}{2}\right)}{\cosh\left[\frac{(\beta' + i\nu)\hbar \omega_0}{2}\right]} \cdot \left[\prod_{n=1}^N 2 \tanh\left(\frac{\beta \hbar \omega_n}{2}\right) \right] \cdot \exp\left[-\frac{1}{2\hbar} \mathbf{z}^T \mathbf{\Lambda}_{\beta z} \mathbf{z}\right], \quad (\text{A35})$$

where $\mathbf{\Lambda}_{\beta z}$ is a $(2N + 2) \times (2N + 2)$ diagonal matrix

$$\mathbf{\Lambda}_{\beta z} = \text{diag}(\lambda_{\beta' q_0}, \theta_1, \dots, \theta_N, \lambda_{\beta' p_0}, \mu_1, \dots, \mu_N), \quad (\text{A36})$$

with the elements

$$\theta_n = 2m_n\omega_n \tanh\left(\frac{\beta\hbar\omega_n}{2}\right), \quad (\text{A37})$$

$$\mu_n = \frac{2}{m_n\omega_n} \tanh\left(\frac{\beta\hbar\omega_n}{2}\right), \quad (\text{A38})$$

$$\lambda_{\beta'q_0} = 2m_0\omega_0 \tanh\left[\frac{(\beta' + i\nu)\hbar\omega_0}{2}\right], \quad (\text{A39})$$

$$\lambda_{\beta'p_0} = \frac{2}{m_0\omega_0} \tanh\left[\frac{(\beta' + i\nu)\hbar\omega_0}{2}\right]. \quad (\text{A40})$$

3. Calculation of the integral

With the explicit expressions of $\left[e^{i\nu H_S^H(\tau)}\right]_w(\mathbf{z})$ and $P(\mathbf{z})$, we perform the integral in Eq. (13), and obtain the result of the characteristic function of heat

$$\chi_\tau(\nu) = \sqrt{\frac{\det(\mathbf{\Lambda}_{\beta z} - i\mathbf{\Lambda}_{\nu z})}{\det[\mathbf{\Lambda}_{\beta z} - i\tilde{\mathbf{\Lambda}}_{\nu z}(\tau)]}}. \quad (\text{A41})$$

We have used the following integral formula

$$\int d\mathbf{x} e^{-\frac{1}{2}\mathbf{x}^T \mathbf{T} \mathbf{x}} = \sqrt{\frac{(2\pi)^{\dim(\mathbf{T})}}{\det(\mathbf{T})}}, \quad (\text{A42})$$

where all the eigenvalues of \mathbf{T} have positive real parts.

By introducing a diagonal matrix $\mathbf{A} = \mathbf{\Lambda}_{\beta z} - i\mathbf{\Lambda}_{\nu z}$, we rewrite Eq. (A41) as

$$\chi_\tau(\nu) = \sqrt{\frac{1}{\det\left(\mathbf{I} + i\sqrt{\mathbf{A}^{-1}}\left[\mathbf{\Lambda}_{\nu z} - \tilde{\mathbf{\Lambda}}_{\nu z}(\tau)\right]\sqrt{\mathbf{A}^{-1}}\right)}}. \quad (\text{A43})$$

Since $\tilde{\mathbf{\Lambda}}_{\nu z}(\tau)$ is a rank-2 matrix, we rewrite it as

$$\tilde{\mathbf{\Lambda}}_{\nu z}(\tau) = 2m_0\omega_0 \tan\left(\frac{\omega_0\hbar\nu}{2}\right) \left[\mathbf{v}_{q_0}(\tau)\mathbf{v}_{q_0}^T(\tau) + \frac{1}{m_0^2\omega_0^2}\mathbf{v}_{p_0}(\tau)\mathbf{v}_{p_0}^T(\tau) \right], \quad (\text{A44})$$

with the vectors

$$\mathbf{v}_{q_0}(\tau) = \left(\alpha_0, \alpha_1, \dots, \alpha_N, \frac{\beta_0}{m_0}, \frac{\beta_1}{m_1}, \dots, \frac{\beta_N}{m_N} \right)^T, \quad (\text{A45})$$

$$\mathbf{v}_{p_0}(\tau) = \left(m_0\dot{\alpha}_0, m_0\dot{\alpha}_1, \dots, m_0\dot{\alpha}_N, \dot{\beta}_0, \frac{m_0}{m_1}\dot{\beta}_1, \dots, \frac{m_0}{m_N}\dot{\beta}_N \right)^T. \quad (\text{A46})$$

Here, the evolution time t in the terms α_n and β_n is set to τ . We rewrite the matrix in the determinant [see Eq. (A43)] as

$$\sqrt{\mathbf{A}^{-1}} \left(\mathbf{\Lambda}_{\nu z} - \tilde{\mathbf{\Lambda}}_{\nu z}(\tau) \right) \sqrt{\mathbf{A}^{-1}} = \mathbf{M}\mathbf{M}^T, \quad (\text{A47})$$

with the matrix

$$\mathbf{M}^T = \begin{pmatrix} \sqrt{2m_0\omega_0 \tan \frac{\omega_0\hbar\nu}{2}} \mathbf{v}_{q_0}^T(0) \\ \sqrt{\frac{2}{m_0\omega_0} \tan \frac{\omega_0\hbar\nu}{2}} \mathbf{v}_{p_0}^T(0) \\ i\sqrt{2m_0\omega_0 \tan \frac{\omega_0\hbar\nu}{2}} \mathbf{v}_{q_0}^T(\tau) \\ i\sqrt{\frac{2}{m_0\omega_0} \tan \frac{\omega_0\hbar\nu}{2}} \mathbf{v}_{p_0}^T(\tau) \end{pmatrix} \sqrt{\mathbf{A}^{-1}}. \quad (\text{A48})$$

The determinant in Eq. (A43) can be simplified to

$$\det \left(\mathbf{I} + i\sqrt{\mathbf{A}^{-1}} \left[\mathbf{\Lambda}_{\nu z} - \tilde{\mathbf{\Lambda}}_{\nu z}(\tau) \right] \sqrt{\mathbf{A}^{-1}} \right) = \det \left(\mathbf{I}_4 + i\mathbf{M}^T\mathbf{M} \right), \quad (\text{A49})$$

where the right-hand side is the determinant of a 4×4 matrix, and \mathbf{I}_4 is the 4×4 identity matrix. Notice that the initial values of the two vectors are

$$\mathbf{v}_{q_0}(0) = (1, 0, \dots, 0, 0, 0, \dots, 0)^T, \quad (\text{A50})$$

$$\mathbf{v}_{p_0}(0) = (0, 0, \dots, 0, 1, 0, \dots, 0)^T. \quad (\text{A51})$$

The explicit result of $\mathbf{M}^T\mathbf{M}$ is obtained as

$$\mathbf{M}^T\mathbf{M} = \Xi \begin{pmatrix} 1 & 0 & i\alpha_0 & i\dot{\alpha}_0/\omega_0 \\ 0 & 1 & i\omega_0\beta_0 & i\dot{\beta}_0 \\ i\alpha_0 & i\omega_0\beta_0 & -h_{11}(\tau) & -h_{12}(\tau) \\ i\dot{\alpha}_0/\omega_0 & i\dot{\beta}_0 & -h_{12}(\tau) & -h_{22}(\tau) \end{pmatrix}, \quad (\text{A52})$$

where the elements are functions of the final time τ , and the functions $h_{11}(\tau)$, $h_{12}(\tau)$ and $h_{22}(\tau)$ are

$$h_{11}(\tau) = \frac{2m_0\omega_0}{\Xi} \tan\left(\frac{\omega_0\hbar\nu}{2}\right) \mathbf{v}_{q_0}^T(\tau) \mathbf{A}^{-1} \mathbf{v}_{q_0}(\tau), \quad (\text{A53})$$

$$h_{22}(\tau) = \frac{2}{m_0\omega_0\Xi} \tan\left(\frac{\omega_0\hbar\nu}{2}\right) \mathbf{v}_{p_0}^T(\tau) \mathbf{A}^{-1} \mathbf{v}_{p_0}(\tau), \quad (\text{A54})$$

$$h_{12}(\tau) = \frac{2}{\Xi} \tan\left(\frac{\omega_0\hbar\nu}{2}\right) \mathbf{v}_{q_0}^T(\tau) \mathbf{A}^{-1} \mathbf{v}_{p_0}(\tau), \quad (\text{A55})$$

with

$$\mathbf{v}_{q_0}^T(t) \mathbf{A}^{-1} \mathbf{v}_{q_0}(t) = \frac{\alpha_0^2 + \omega_0^2 \beta_0^2}{2m_0\omega_0 \left\{ \tanh\left[\frac{(\beta' + i\nu)\hbar\omega_0}{2}\right] - i \tan\left(\frac{\omega_0\hbar\nu}{2}\right) \right\}} + \sum_{n=1}^N \left(\frac{\alpha_n^2}{\theta_n} + \frac{1}{\mu_n} \frac{\beta_n^2}{m_n^2} \right), \quad (\text{A56})$$

$$\mathbf{v}_{p_0}^T(t) \mathbf{A}^{-1} \mathbf{v}_{p_0}(t) = \frac{m_0 \left(\dot{\alpha}_0^2 + \omega_0^2 \dot{\beta}_0^2 \right)}{2\omega_0 \left\{ \tanh\left[\frac{(\beta' + i\nu)\hbar\omega_0}{2}\right] - i \tan\left(\frac{\omega_0\hbar\nu}{2}\right) \right\}} + m_0^2 \sum_{n=1}^N \left(\frac{\dot{\alpha}_n^2}{\theta_n} + \frac{1}{\mu_n} \frac{\dot{\beta}_n^2}{m_n^2} \right), \quad (\text{A57})$$

$$\mathbf{v}_{q_0}^T(t) \mathbf{A}^{-1} \mathbf{v}_{p_0}(t) = \frac{\frac{d}{dt} (\alpha_0^2 + \omega_0^2 \beta_0^2)}{4\omega_0 \left\{ \tanh\left[\frac{(\beta' + i\nu)\hbar\omega_0}{2}\right] - i \tan\left(\frac{\omega_0\hbar\nu}{2}\right) \right\}} + \frac{m_0}{2} \sum_{n=1}^N \frac{d}{dt} \left(\frac{\alpha_n^2}{\theta_n} + \frac{1}{\mu_n} \frac{\beta_n^2}{m_n^2} \right). \quad (\text{A58})$$

The summations are replaced by the integral with the Ohmic spectral density, and every element in Eq. (A52) is carried out as

$$\alpha_0(\tau) = e^{-\frac{\kappa\tau}{2}} \left[\cos(\hat{\omega}_0\tau) + \frac{\kappa \sin(\hat{\omega}_0\tau)}{2\hat{\omega}_0} \right], \quad (\text{A59})$$

$$\beta_0(\tau) = \frac{e^{-\frac{\kappa\tau}{2}} \sin(\hat{\omega}_0\tau)}{\hat{\omega}_0}, \quad (\text{A60})$$

$$h_{11}(\tau) = \Theta + e^{-\kappa\tau} \left[\frac{\omega_0^2}{\hat{\omega}_0^2} + \frac{\kappa \sin(2\hat{\omega}_0\tau)}{2\hat{\omega}_0} - \frac{\kappa^2 \cos(2\hat{\omega}_0\tau)}{4\hat{\omega}_0^2} \right] (1 - \Theta), \quad (\text{A61})$$

$$h_{22}(\tau) = \Theta + e^{-\kappa\tau} \left[\frac{\omega_0^2}{\hat{\omega}_0^2} - \frac{\kappa \sin(2\hat{\omega}_0\tau)}{2\hat{\omega}_0} - \frac{\kappa^2 \cos(2\hat{\omega}_0\tau)}{4\hat{\omega}_0^2} \right] (1 - \Theta), \quad (\text{A62})$$

$$h_{12}(\tau) = \frac{\kappa\omega_0 e^{-\kappa\tau}}{2\hat{\omega}_0^2} (\Theta - 1) [1 - \cos(2\hat{\omega}_0\tau)]. \quad (\text{A63})$$

Then, Eq. (17) is obtained by directly calculating the determinant of a 4×4 matrix in Eq. (A49).

Appendix B: Classical Caldeira-Leggett model

We consider the classical Caldeira-Leggett model, where coordinates and momenta commute with each other. To eliminate the initial slippage, the initial state is amended as a coupled state

$$\rho^{\text{cl}}(\mathbf{z}; 0) = \frac{e^{-\beta' H_S(0) - \beta [H_B(0) + H_{SB}(0)]}}{Z^{\text{cl}}(\beta', \beta)}, \quad (\text{B1})$$

which represents the probability density in the phase space of the composite system. The classical partition function is obtained by performing the integral in the phase space

$$Z^{\text{cl}}(\beta', \beta) = \iint e^{-\beta' H_S(0) - \beta [H_B(0) + H_{SB}(0)]} dq_0 dq_1 \dots dq_N dp_0 dp_1 \dots dp_N \quad (\text{B2})$$

$$= \frac{2\pi}{\beta' \omega_0} \prod_{n=1}^N \left(\frac{2\pi}{\beta \omega_n} \right), \quad (\text{B3})$$

which is independent of the interaction (notice that the partition function of the quantum model relies on the interaction strength [68, 87]).

We also define the classical fluctuating heat as the energy difference of the initial and the final system energy. For classical dynamics, the initial and the final states are directly represented by the points in the phase space, and the measurements over the system can be applied without disturbing the composite system. Therefore, the characteristic function of heat is

$$\chi_\tau^{\text{cl}}(\nu) = \frac{\iint e^{i\nu H_S(\tau) - (\beta' + i\nu) H_S(0) - \beta [H_B(0) + H_{SB}(0)]} dq_0 dq_1 \dots dq_N dp_0 dp_1 \dots dp_N}{Z^{\text{cl}}(\beta', \beta)}, \quad (\text{B4})$$

where the energy of the system $H_S(t) = [p_0(t)]^2/(2m_0) + m_0\omega_0^2[q_0(t)]^2/2$ is determined by $p_0(t)$ and $q_0(t)$ associated with the initial point \mathbf{z} . We choose a new set of initial variables $q_0, \mathbf{q}_n := q_n - C_n q_0/(m_n \omega_n^2), p_0$ and p_n in the following calculation.

We rewrite the evolution of the coordinate $q_0(t)$ of the system [Eq. (A16)] as

$$(s^2 + \kappa s + \omega_0^2) \tilde{q}_0(s) = \dot{q}_0(0) + \left(s + \sum_n \frac{C_n^2}{m_0 m_n \omega_n^2} \frac{s}{s^2 + \omega_n^2} \right) q_0(0) + \sum_n \frac{C_n}{m_0} \frac{\dot{q}_n(0) + s \mathbf{q}_n}{s^2 + \omega_n^2}. \quad (\text{B5})$$

For the Ohmic spectral density, the summation in the second term is

$$\sum_n \frac{C_n^2}{m_0 m_n \omega_n^2} \frac{s}{s^2 + \omega_n^2} = \kappa. \quad (\text{B6})$$

Thus, Eq. (B5) naturally leads to Eq. (A19) by substituting $q_n(0)$ into \mathbf{q}_n . The initial slippage is rationally eliminated by choosing a coupled initial state [Eq. (B1)]. In reality, the interaction between the system and the heat bath always exist, and one cannot prepare the initial state of the composite system without the influence of the interaction. The initial state of the composite system is more likely in the coupled form [Eq. (B1)]. The heat bath encodes partial information of the system due to the interaction.

Similar to Eq. (A10), the system energy at time t can be represented by the dynamical map as

$$H_S(t) = \frac{1}{2} \mathbf{z}^T(t) \mathbf{\Lambda}_{H_S} \mathbf{z}(t) \quad (\text{B7})$$

$$= \frac{1}{2} \mathbf{z}^T(0) \tilde{\mathbf{\Lambda}}_{H_S}(t) \mathbf{z}(0), \quad (\text{B8})$$

with the matrices [88]

$$\mathbf{\Lambda}_{H_S} = \begin{pmatrix} m_0 \omega_0^2 & & & \\ & \mathbf{O} & & \\ & & \frac{1}{m_0} & \\ & & & \mathbf{O} \end{pmatrix}, \quad (\text{B9})$$

and

$$\tilde{\mathbf{\Lambda}}_{H_S}(t) = \exp(\mathbf{L}^T t) \mathbf{\Lambda}_{H_S} \exp(\mathbf{L} t). \quad (\text{B10})$$

The initial vector is now amended to

$$\mathbf{z}(0) = (q_0(0), \mathbf{q}_1, \dots, \mathbf{q}_n, p_0(0), \dots, p_N(0))^T. \quad (\text{B11})$$

The initial Hamiltonians $H_S(0)$ and $H_B(0) + H_{SB}(0)$ are

$$H_S(0) = \frac{1}{2} \mathbf{z}^T(0) \mathbf{\Lambda}_{H_S} \mathbf{z}(0), \quad (\text{B12})$$

$$\begin{aligned} H_B(0) + H_{SB}(0) &= \sum_{n=1}^N \left(\frac{1}{2} \frac{p_n^2}{m_n} + \frac{1}{2} m_n \omega_n^2 \mathbf{q}_n^2 \right) \\ &= \frac{1}{2} \mathbf{z}^T(0) \mathbf{\Lambda}_{H_B} \mathbf{z}(0), \end{aligned} \quad (\text{B13})$$

with the matrix

$$\mathbf{\Lambda}_{H_B} = \text{diag}(0, m_1 \omega_1^2, \dots, m_N \omega_N^2, 0, \frac{1}{m_1}, \dots, \frac{1}{m_N}). \quad (\text{B14})$$

According to the integral formula (A42), we carry out the characteristic function Eq. (B4) into

$$\chi_\tau^{\text{cl}}(\nu) = \sqrt{\frac{\det[\beta' \mathbf{\Lambda}_{H_S} + \beta \mathbf{\Lambda}_{H_B}]}{\det[\beta' \mathbf{\Lambda}_{H_S} + \beta \mathbf{\Lambda}_{H_B} - i\nu (\tilde{\mathbf{\Lambda}}_{H_S}(\tau) - \mathbf{\Lambda}_{H_S})]}}. \quad (\text{B15})$$

For the classical limit $\hbar \rightarrow 0$, we can verify

$$\lim_{\hbar \rightarrow 0} \frac{\mathbf{\Lambda}_{\beta z}}{\hbar} = (\beta' + i\nu) \mathbf{\Lambda}_{H_S} + \beta \mathbf{\Lambda}_{H_B}, \quad (\text{B16})$$

$$\lim_{\hbar \rightarrow 0} \frac{\mathbf{\Lambda}_{\nu z}}{\hbar} = \nu \mathbf{\Lambda}_{H_S}, \quad (\text{B17})$$

$$\lim_{\hbar \rightarrow 0} \frac{\tilde{\mathbf{\Lambda}}_{\nu z}(t)}{\hbar} = \nu \tilde{\mathbf{\Lambda}}_{H_S}(t), \quad (\text{B18})$$

and obtain

$$\lim_{\hbar \rightarrow 0} \chi_\tau(\nu) = \chi_\tau^{\text{cl}}(\nu), \quad (\text{B19})$$

with $\chi_\tau(\nu)$ given in Eq. (A41). The final result [Eq. (B15)] is the same as Eq. (25). Hence, we use the same notation.

Appendix C: The characteristic function of heat for the classical Brownian motion

We derive the characteristic function of heat for the classical Brownian motion. For an underdamped Brownian particle moving in a potential $V(x)$, the stochastic dynamics is described by the complete Langevin equation

$$\ddot{x} + \kappa \dot{x} + \frac{1}{m} \frac{\partial V}{\partial x} = \frac{1}{m} F_{\text{fluc}}(t). \quad (\text{C1})$$

The fluctuating force is a Gaussian white noise satisfying the fluctuation-dissipation relation

$$\langle F_{\text{fluc}}(t) F_{\text{fluc}}(t') \rangle = 2m\kappa k_B T \delta(t - t'). \quad (\text{C2})$$

The evolution of the system state is characterized by the probability density function $\rho(x, p; t)$ in the phase space. The stochastic dynamics is then described by the Kramers equation [86]

$$\frac{\partial \rho}{\partial t} = \mathcal{L}[\rho], \quad (\text{C3})$$

with the Liouville operator

$$\mathcal{L}[\rho] = -\frac{\partial}{\partial x}\left(\frac{p}{m}\rho\right) + \frac{\partial}{\partial p}\left[\kappa p\rho + \frac{\partial V(x)}{\partial x}\rho + \frac{\kappa m}{\beta}\frac{\partial \rho}{\partial p}\right]. \quad (\text{C4})$$

Similarly in the phase space, we calculate the characteristic function of heat for the classical Brownian motion

$$\chi_\tau^{\text{cl}}(\nu) = \iint dx dp e^{i\nu\left[\frac{p^2}{2m}+V(x)\right]}\eta(x, p; \tau), \quad (\text{C5})$$

where a probability-density-like function $\eta(x, p; t)$ also satisfies the dynamic equation (C3) with the initial condition

$$\eta(x, p; 0) = e^{-i\nu\left[\frac{p^2}{2m}+V(x)\right]}\rho(x, p; 0). \quad (\text{C6})$$

We consider the system potential as a harmonic potential $V(x) = m\omega_0^2 x^2/2$ and the initial system state as an equilibrium state

$$\rho(x, p; 0) = \frac{1}{Z_S^{\text{cl}}(\beta')} e^{-\beta'\left(\frac{p^2}{2m} + \frac{1}{2}m\omega_0^2 x^2\right)}, \quad (\text{C7})$$

with the inverse temperature β' and the classical partition function $Z_S^{\text{cl}}(\beta') = 2\pi/(\beta'\omega_0)$. Under such conditions, the probability-density-like function $\eta(x, p; t)$ is always in a quadratic form, assumed as

$$\eta(x, p; t) = \frac{1}{Z_S^{\text{cl}}(\beta')} e^{-\left[a(t)\frac{p^2}{2m} + b(t)\frac{1}{2}m\omega_0^2 x^2 + c(t)\omega_0 x p + \Lambda(t)\right]}. \quad (\text{C8})$$

The Kramers equation (C3) for $\eta(x, p; t)$ leads to the following ordinary differential equations

$$\dot{\Lambda} = -\kappa\left(1 - \frac{a}{\beta}\right), \quad (\text{C9})$$

$$\dot{a} = 2\kappa a\left(1 - \frac{a}{\beta}\right) - 2\omega_0 c, \quad (\text{C10})$$

$$\dot{b} = 2c\left(\omega_0 - \frac{\kappa}{\beta}c\right), \quad (\text{C11})$$

$$\dot{c} = \omega_0(a - b) + \kappa c - 2\frac{\kappa}{\beta}ac, \quad (\text{C12})$$

with the initial conditions $a(0) = b(0) = \beta' + i\nu$, $c(0) = 0$ and $\Lambda(0) = 0$. According to the conservation of the probability $\iint \eta(x, p; t) dx dp = \text{const}$, the coefficient $\Lambda(t)$ is obtained as

$$e^{-\Lambda(t)} = \frac{\sqrt{a(t)b(t) - c(t)^2}}{\beta' + i\nu}. \quad (\text{C13})$$

Substituting Eq. (C8) into Eq. (C5), we obtain the characteristic function for the classical Brownian motion as

$$\chi_\tau^{\text{cl}}(\nu) = \frac{\beta'}{\beta' + i\nu} \sqrt{\frac{a(\tau)b(\tau) - c(\tau)^2}{[a(\tau) - i\nu][b(\tau) - i\nu] - c(\tau)^2}}. \quad (\text{C14})$$

To solve the nonlinear differential equations (C10)-(C12), we introduce a new set of variables

$$A = \frac{a}{ab - c^2}, \quad (\text{C15})$$

$$B = \frac{b}{ab - c^2}, \quad (\text{C16})$$

$$C = \frac{c}{ab - c^2}, \quad (\text{C17})$$

and obtain the linear differential equations

$$\frac{dA}{dt} = -2\omega_0 C, \quad (\text{C18})$$

$$\frac{dB}{dt} = 2\omega_0 C - 2\kappa B + 2\frac{\kappa}{\beta}, \quad (\text{C19})$$

$$\frac{dC}{dt} = \omega_0(A - B) - \kappa C, \quad (\text{C20})$$

with the initial conditions $A(0) = B(0) = 1/(\beta' + i\nu)$ and $C(0) = 0$. The characteristic function Eq. (C14) becomes

$$\chi_\tau^{\text{cl}}(\nu) = \frac{\beta'}{\beta' + i\nu} \sqrt{\frac{1}{1 - i\nu(A + B) - \nu^2(AB - C^2)}}. \quad (\text{C21})$$

The solutions to Eqs. (C18)-(C20) are

$$A(t) = -\frac{e^{-\kappa t} [\kappa^2 \cos(2\hat{\omega}_0 t) - 2\hat{\omega}_0 \kappa \sin(2\hat{\omega}_0 t) - 4\omega_0^2] \beta - \beta' - i\nu}{4\beta\hat{\omega}_0^2} \frac{1}{\beta' + i\nu} + \frac{1}{\beta}, \quad (\text{C22})$$

$$B(t) = -\frac{e^{-\kappa t} [\kappa^2 \cos(2\hat{\omega}_0 t) + 2\hat{\omega}_0 \kappa \sin(2\hat{\omega}_0 t) - 4\omega_0^2] \beta - \beta' - i\nu}{4\beta\hat{\omega}_0^2} \frac{1}{\beta' + i\nu} + \frac{1}{\beta}, \quad (\text{C23})$$

$$C(t) = -\frac{2e^{-\kappa t} \kappa \omega_0 [\cos(2\hat{\omega}_0 t) - 1] \beta - \beta' - i\nu}{4\beta\hat{\omega}_0^2} \frac{1}{\beta' + i\nu}, \quad (\text{C24})$$

with $\hat{\omega}_0 = \sqrt{\omega_0^2 - \kappa^2/4}$. Plugging the solutions into Eq. (C21), we immediately obtain Eq. (25). We remark that the heat distribution of the classical Brownian motion has been obtained by the path-integral method in Ref. [58], but they only consider the initial temperature of the system to be the same as that of the bath.

1. The long-time limit

After sufficiently long relaxation time, the solutions $a(t)$, $b(t)$ and $c(t)$ to Eqs. (C10)-(C12) eventually approach $a(\infty) = b(\infty) = \beta$ and $c(\infty) = 0$. The long-time limit of Eq. (C14) reproduces Eq. (33).

2. The underdamped limit

In the underdamped limit $\kappa/\omega_0 \rightarrow 0$, the differential equations (C10)-(C12) are reduced to

$$\dot{a} = \kappa a \left(1 - \frac{a}{\beta} \right), \quad (\text{C25})$$

with $b = a$ and $c = 0$. The solution is

$$a(t) = \frac{\beta(\beta' + i\nu)}{\beta' + i\nu + (\beta - \beta' - i\nu)e^{-\kappa t}}, \quad (\text{C26})$$

and Eq. (C14) becomes

$$\chi_\tau^{\text{w,cl}}(\nu) = \frac{\beta\beta'}{(\beta - i\nu)(\beta' + i\nu)(1 - e^{-\kappa\tau}) + \beta\beta'e^{-\kappa\tau}}. \quad (\text{C27})$$

It can be checked that Eq. (37) reproduces Eq. (C27) in the classical limit $\hbar \rightarrow 0$.

3. The overdamped limit

In the overdamped limit $\kappa/\omega_0 \rightarrow \infty$, the differential equations (C10)-(C12) are reduced to

$$\dot{a} = 2\kappa a \left(1 - \frac{a}{\beta}\right), \quad (\text{C28})$$

$$\dot{b} = 2\omega_0 c \left(1 - \frac{\kappa}{\beta\omega_0} c\right), \quad (\text{C29})$$

$$0 = (\kappa c + \omega_0 a) - 2\frac{\kappa}{\beta} ac - b\omega_0. \quad (\text{C30})$$

Eliminating c in Eq. (C30), Eq. (C29) becomes

$$\dot{b} = \frac{2\omega_0^2 (\beta - a - b) (b - a)}{\kappa (\beta - 2a) \left(1 - 2\frac{a}{\beta}\right)}. \quad (\text{C31})$$

Notice that in the overdamped limit, the relaxation timescales of the momentum and the coordinate are separated. We can substitute $a = \beta$ in Eq. (C31) and obtain

$$\dot{b} = \frac{2\omega_0^2}{\kappa} b \left(1 - \frac{b}{\beta}\right). \quad (\text{C32})$$

With the initial condition $a(0) = \beta' + i\nu$ and $b(0) = \beta' + i\nu$, the solutions are

$$a(t) = \frac{(\beta' + i\nu) \beta}{\beta' + i\nu + (\beta - \beta' - i\nu) e^{-2\kappa t}}, \quad (\text{C33})$$

$$b(t) = \frac{(\beta' + i\nu) \beta}{\beta' + i\nu + (\beta - \beta' - i\nu) e^{-\frac{2\omega_0^2}{\kappa} t}}. \quad (\text{C34})$$

We substitute Eqs. (C33), (C34) and $c(t) \approx 0$ into Eq. (C14) and obtain

$$\begin{aligned} \chi_\tau^{\text{s,cl}}(\nu) &= \frac{\beta\beta'}{\sqrt{(\beta - i\nu)(\beta' + i\nu) (1 - e^{-2\kappa\tau}) + \beta\beta' e^{-2\kappa\tau}}} \\ &\times \frac{1}{\sqrt{(\beta - i\nu)(\beta' + i\nu) \left(1 - e^{-\frac{2\omega_0^2}{\kappa}\tau}\right) + \beta\beta' e^{-\frac{2\omega_0^2}{\kappa}\tau}}}. \end{aligned} \quad (\text{C35})$$

It can be checked that Eq. (38) reproduces Eq. (C35) in the classical limit $\hbar \rightarrow 0$.

Appendix D: The characteristic function of heat for complete thermalization

We derive the characteristic function of heat for a complete thermalization process [Eq. (34)] in the main content. For a complete thermalization process (typically with infinite relaxation time), the information of the initial state is completely forgotten, and the final state is always an equilibrium state at the inverse temperature β of the heat bath

$$\gamma_{\text{th},l'l} = p_{l'}^{\text{eq}}, \quad (\text{D1})$$

regardless of the initial state l . Therefore, the characteristic function of heat for complete thermalization is $\chi_{\text{th}}(\nu) = \sum_{l',l} \exp[i\nu(E_{l'}^S - E_l^S)] p_l p_{l'}^{\text{eq}}$. We immediately obtain Eq. (34) by plugging into the initial distribution $p_l = \exp(-\beta' E_l^S)/Z_S(\beta')$ and the final distribution $p_{l'}^{\text{eq}} = \exp(-\beta E_{l'}^S)/Z_S(\beta)$, where $Z_S(\beta') = \sum_l \exp(-\beta' E_l^S)$ is the partition function of the system at the inverse temperature β' .

-
- [1] G. Gallavotti and E. G. D. Cohen, *Phys. Rev. Lett.* **74**, 2694 (1995).
 - [2] C. Jarzynski, *Phys. Rev. Lett.* **78**, 2690 (1997).
 - [3] G. E. Crooks, *Phys. Rev. E* **60**, 2721 (1999).
 - [4] C. Jarzynski and D. K. Wójcik, *Phys. Rev. Lett.* **92**, 230602 (2004).
 - [5] C. Jarzynski, *Annu. Rev. Condens. Matter Phys.* **2**, 329 (2011).
 - [6] K. Sekimoto, *Stochastic Energetics* (Springer Berlin Heidelberg, 2010).
 - [7] U. Seifert, *Rep. Prog. Phys.* **75**, 126001 (2012).
 - [8] U. Seifert, *Phys. Rev. Lett.* **95**, 040602 (2005).
 - [9] M. Esposito, U. Harbola, and S. Mukamel, *Rev. Mod. Phys.* **81**, 1665 (2009).
 - [10] M. Campisi, P. Hänggi, and P. Talkner, *Rev. Mod. Phys.* **83**, 771 (2011).
 - [11] R. Klages, *Nonequilibrium statistical physics of small systems : fluctuation relations and beyond* (Wiley-VCH, Weinheim, Germany, 2013).
 - [12] M. Horodecki and J. Oppenheim, *Nat Commun* **4**, 2059 (2013).
 - [13] S. Ciliberto, *Phys. Rev. X* **7**, 021051 (2017).
 - [14] H. Tasaki, [cond-mat/0009244](#).
 - [15] J. Kurchan, (2000), [arXiv:cond-mat/0007360 \[cond-mat.stat-mech\]](#).

- [16] P. Talkner, E. Lutz, and P. Hänggi, *Phys. Rev. E* **75**, 050102 (2007).
- [17] S. Deffner and E. Lutz, *Phys. Rev. E* **77**, 021128 (2008).
- [18] F. Liu, *Phys. Rev. E* **90**, 032121 (2014).
- [19] L. Zhu, Z. Gong, B. Wu, and H. T. Quan, *Phys. Rev. E* **93**, 062108 (2016).
- [20] K. Funo and H. Quan, *Phys. Rev. Lett.* **121**, 040602 (2018).
- [21] D. S. P. Salazar and S. A. Lira, *Phys. Rev. E* **99**, 062119 (2019).
- [22] C. Jarzynski, H. Quan, and S. Rahav, *Phys. Rev. X* **5**, 031038 (2015).
- [23] S. Deffner, J. P. Paz, and W. H. Zurek, *Phys. Rev. E* **94**, 010103(R) (2016).
- [24] I. García-Mata, A. J. Roncaglia, and D. A. Wisniacki, *Phys. Rev. E* **95**, 050102 (2017).
- [25] Z. Fei and H. Quan, *Phys. Rev. Lett.* **124**, 240603 (2020).
- [26] T. Qiu, Z. Fei, R. Pan, and H. T. Quan, *Phys. Rev. E* **101**, 032111 (2020).
- [27] W. H. Zurek, *Phys. Today* **44**, 36 (1991).
- [28] W. H. Zurek, *Rev. Mod. Phys.* **75**, 715 (2003).
- [29] K. Saito and A. Dhar, *Phys. Rev. Lett.* **99**, 180601 (2007).
- [30] Y. Dubi and M. D. Ventra, *Rev. Mod. Phys.* **83**, 131 (2011).
- [31] J. Thingna, J. L. García-Palacios, and J.-S. Wang, *Phys. Rev. B* **85**, 195452 (2012).
- [32] J.-S. Wang, B. K. Agarwalla, H. Li, and J. Thingna, *Front. Phys.* **9**, 673 (2014).
- [33] J. Thingna, D. Manzano, and J. Cao, *Sci. Rep.* **6**, 28027 (2016).
- [34] D. He, J. Thingna, J.-S. Wang, and B. Li, *Phys. Rev. B* **94**, 155411 (2016).
- [35] D. Segal and B. K. Agarwalla, *Ann. Phys. Chem.* **67**, 185 (2016).
- [36] M. Kilgour, B. K. Agarwalla, and D. Segal, *J. Chem. Phys.* **150**, 084111 (2019).
- [37] C. Wang, J. Ren, and J. Cao, *Phys. Rev. A* **95**, 023610 (2017).
- [38] E. Aurell, B. Donvil, and K. Mallick, *Phys. Rev. E* **101**, 052116 (2020).
- [39] T. Denzler and E. Lutz, *Phys. Rev. E* **98**, 052106 (2018).
- [40] D. S. P. Salazar, A. M. S. Macêdo, and G. L. Vasconcelos, *Phys. Rev. E* **99**, 022133 (2019).
- [41] M. Popovic, M. T. Mitchison, A. Strathearn, B. W. Lovett, J. Goold, and P. R. Eastham, *PRX Quantum* **2**, 020338 (2021).
- [42] M. B. Karsten Balzer, *Nonequilibrium Green's Functions Approach to Inhomogeneous Systems* (Springer Berlin Heidelberg, 2012).
- [43] M. Esposito, M. A. Ochoa, and M. Galperin, *Phys. Rev. Lett.* **114**, 080602 (2015).
- [44] C. A. Polanco, *Nanoscale Microscale Thermophys. Eng.* **25**, 1 (2021).

- [45] C. Aron, G. Biroli, and L. F. Cugliandolo, *J. Stat. Mech.* **2010**, P11018 (2010).
- [46] K. Mallick, M. Moshe, and H. Orland, *J. Phys. A* **44**, 095002 (2011).
- [47] M. Carrega, P. Solinas, A. Braggio, M. Sasseti, and U. Weiss, *New J. Phys.* **17**, 045030 (2015).
- [48] K. Funo and H. T. Quan, *Phys. Rev. E* **98**, 012113 (2018).
- [49] J. Yeo, *Phys. Rev. E* **100**, 062107 (2019).
- [50] H. C. Fogedby, *J. Stat. Mech.: Theory Exp.* **2020**, 083208 (2020).
- [51] R. van Zon and E. G. D. Cohen, *Phys. Rev. E* **69**, 056121 (2004).
- [52] A. Imparato, L. Peliti, G. Pesce, G. Rusciano, and A. Sasso, *Phys. Rev. E* **76**, 050101 (2007).
- [53] H. C. Fogedby and A. Imparato, *J. Phys. A: Math. Theor.* **42**, 475004 (2009).
- [54] D. Chatterjee and B. J. Cherayil, *Phys. Rev. E* **82**, 051104 (2010).
- [55] J. R. Gomez-Solano, A. Petrosyan, and S. Ciliberto, *Phys. Rev. Lett.* **106**, 200602 (2011).
- [56] D. S. P. Salazar and S. A. Lira, *J. Phys. A: Math. Theor.* **49**, 465001 (2016).
- [57] A. Pagare and B. J. Cherayil, *Phys. Rev. E* **100**, 052124 (2019).
- [58] P. V. Paraguassú, R. Aquino, and W. A. M. Morgado, (2021), [arXiv:2102.09115](https://arxiv.org/abs/2102.09115) [[cond-mat.stat-mech](https://arxiv.org/abs/2102.09115)].
- [59] D. Gupta and D. A. Sivak, (2021), [arXiv:2103.09358](https://arxiv.org/abs/2103.09358) [[cond-mat.stat-mech](https://arxiv.org/abs/2103.09358)].
- [60] M. Esposito, M. A. Ochoa, and M. Galperin, *Phys. Rev. B* **92**, 235440 (2015).
- [61] P. Talkner and P. Hänggi, *Phys. Rev. E* **94**, 022143 (2016).
- [62] P. Talkner and P. Hänggi, *Rev. Mod. Phys.* **92**, 041002 (2020).
- [63] W. Bez, *Z. Phys. B* **39**, 319 (1980).
- [64] A. Caldeira and A. Leggett, *Phys. A* **121**, 587 (1983).
- [65] A. Caldeira and A. Leggett, *Ann. Phys.* **149**, 374 (1983).
- [66] W. G. Unruh and W. H. Zurek, *Phys. Rev. D* **40**, 1071 (1989).
- [67] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, 2007).
- [68] U. Weiss, *Quantum Dissipative Systems* (World Scientific Publishing Company, 2008).
- [69] Usually the quantum fluctuating heat is defined via two-point measurements over the heat bath. When the Hamiltonian of the system is time-independent, the internal energy change of the system is completely caused by the heat exchange. The quantum fluctuating heat can thus be alternatively defined via two-point measurements over the system, whose number of

degrees of freedom is much smaller than that of the heat bath. Hence, the calculation of the heat statistics can be significantly simplified under this definition.

- [70] L. H. Yu and C.-P. Sun, *Phys. Rev. A* **49**, 592 (1994).
- [71] E. Wigner, *Phys. Rev.* **40**, 749 (1932).
- [72] M. Hillery, R. O’Connell, M. Scully, and E. Wigner, *Phys. Rep.* **106**, 121 (1984).
- [73] A. Polkovnikov, *Ann. Phys.* **325**, 1790 (2010).
- [74] Z. Fei, H. T. Quan, and F. Liu, *Phys. Rev. E* **98**, 012132 (2018).
- [75] Y. Qian and F. Liu, *Phys. Rev. E* **100**, 062119 (2019).
- [76] O. Brodier, K. Mallick, and A. M. O. de Almeida, *J. Phys. A: Math. Theor.* **53**, 325001 (2020).
- [77] T. Qiu, Z. Fei, R. Pan, and H. T. Quan, *Phys. Rev. E* **101**, 032113 (2020).
- [78] T. Qiu and H.-T. Quan, *Commun. Theor. Phys.* **73**, 095602 (2021).
- [79] B. L. Hu, J. P. Paz, and Y. Zhang, *Phys. Rev. D* **45**, 2843 (1992).
- [80] R. Karrlein and H. Grabert, *Phys. Rev. E* **55**, 153 (1997).
- [81] G. W. Ford and R. F. O’Connell, *Phys. Rev. D* **64**, 105020 (2001).
- [82] D. S. P. Salazar, *Phys. Rev. E* **101**, 030101 (2020).
- [83] Y. H. Chen, J.-F. Chen, Z. Fei, and H. T. Quan, (2021), [arXiv:2108.04128](https://arxiv.org/abs/2108.04128) [cond-mat.stat-mech].
- [84] J. S. Cañizares and F. Sols, *Phys. A* **212**, 181 (1994).
- [85] K.-K. Ju, C.-X. Guo, and X.-Y. Pan, *Chin. Phys. Lett.* **34**, 010301 (2017).
- [86] H. A. Kramers, *Physica* **7**, 284 (1940).
- [87] H. Grabert, U. Weiss, and P. Talkner, *Z. Phys. B* **55**, 87 (1984).
- [88] Strictly, the substitution requires to amend $\gamma_n, \Lambda_{nm}, \xi_n, \Delta_{nm}$ accordingly, but we only require $q_0(t)$ and $p_0(t)$ to calculate the characteristic function of heat, so we skip the further amendment.