

GEOMETRIC CLASSIFICATION OF TOTAL STABILITY CONDITIONS

YU QIU AND XIAOTING ZHANG

Dedicated to Alastair King on the occasion of his sixtieth birthday

ABSTRACT. We construct a geometric model for the root category $\mathcal{D}_\infty(Q)/[2]$ of any Dynkin quiver Q , which is an h_Q -gon \mathbf{V}_Q with cores, where h_Q is the Coxeter number and $\mathcal{D}_\infty(Q) = \mathcal{D}^b(Q)$ is the bounded derived category associated to Q . As an application, we describe all spaces $\text{ToSt}\mathcal{D}$ of total stability conditions on triangulated categories \mathcal{D} , where \mathcal{D} must be of the form $\mathcal{D}_\infty(Q)$. More precisely, we prove that $\text{ToSt}\mathcal{D}_\infty(Q)/[2]$ is isomorphic to a suitable moduli space of stable h_Q -gons of type Q .

In particular, an h_Q -gon \mathbf{V} of type D_n is a (centrally) symmetric doubly punctured $2(n-1)$ -gon. \mathbf{V} is stable if it is positively convex and the punctures are inside the level- $(n-2)$ diagonal-gon. Another interesting case is E_6 , where the (stable) h_Q -gon (dodecagon) can be realized as a pair of planar tiling pattern.

Key words: stability conditions, root system, Dynkin diagram, geometric model

CONTENTS

1. Introduction	2
2. Preliminaries	8
3. Prototype of ToSt (i.e. type A_n)	12
4. ToSt of type D_n	15
5. Exceptional type: Preview	23
6. ToSt of exceptional type E_6	24
7. ToSt of exceptional type E_7	30
8. ToSt of exceptional type E_8	33
9. Non simply-laced types via folding	37
References	38
Appendix A. Tikz Art Gallery by Qy	40

Date: June 28, 2022.

1. INTRODUCTION

The notion of stability conditions on a triangulated category \mathcal{D} is introduced by Bridgeland [B1], whose motivation is Π -stability in string theory. They measure certain stability structure, as the name suggested, in physics as well as in mathematics. A stability condition $\sigma = (Z, \mathcal{P})$ on \mathcal{D} consists of a group homomorphism $Z: K\mathcal{D} \rightarrow \mathbb{C}$, known as the central charge, and an \mathbb{R} -collection of abelian subcategories $\mathcal{P}(\phi)$, known as the slicing. The main theorem in [B1] states that all stability conditions on a triangulated category form a complex manifold. Moreover, one recent breakthrough in this area is the identification of stability spaces with the moduli spaces of quadratic differentials, cf. [BS, HKK]. Although there are already many studies, the known examples of global structures of stability spaces are still very limited.

1.1. Total stability. One key concept in the theory of stability conditions is stable object, i.e., simple object in some $\mathcal{P}(\phi)$, $\phi \in \mathbb{R}$. It goes back to geometric invariant theory and King's θ -stability, which plays an important role in the study of Donaldson-Thomas invariant, as well as cluster theory, cf. [K]. When passing to the dynamical system side, i.e. identifying a stability condition with a quadratic differential on some Riemann/marked surface, stable objects correspond to non-broken geodesics (connecting zeroes) or saddles. An interesting question proposed in [Q1] is to find stability conditions that make all indecomposable objects in the triangulated category stable. Such stability conditions are called total stability condition. The abelian version of this question is proposed by Reineke [R] when studying quantum dilogarithm identities of the module category $\text{mod } \mathbf{k}Q$ of an ADE quiver Q .

Motivated by q -deformation of stability conditions in [IQ1, IQ2], Ikeda-Qiu introduce the notion of global dimension ($\mathbb{R}_{\geq 0}$ -valued) function gldim of a stability condition σ to measure how stable a stability condition is. Qiu [Q2] shows that σ is totally stable if and only if $\text{gldim } \sigma < 1$, which is very rare. In fact, such an existence of σ implies that \mathcal{D} must be the bounded derived category $\mathcal{D}_{\infty}(Q) = \mathcal{D}^b(Q)$ of a Dynkin diagram Q (cf. [KOT, Q3]). In this paper, we classify all spaces $\text{ToSt}(Q)$ of total conditions on the Dynkin category $\mathcal{D}_{\infty}(Q)$ by constructing their geometric models.

1.2. Geometric model for Dynkin categories. For type A_n , [Q2] gives a geometric description of $\text{ToSt } \mathcal{D}_{\infty}(A_n)$, i.e. as the moduli space of positively convex $(n+1)$ -gon. It can be viewed as a variation/consequence of the fact that $\mathcal{D} = \mathcal{D}_{\infty}(Q)$ is the topological Fukaya category of $(n+1)$ -gon as follows.

Let \mathbf{S}^{λ} be a graded marked surface, that is a topological surface \mathbf{S} with marked points on its boundary $\partial\mathbf{S}$ and a grading $\lambda \in H_1(\mathbb{P}\mathbf{T}\mathbf{S})$. Given an ' ∞ -angulation' \mathbf{A} of \mathbf{S}^{λ} , one obtains a graded gentle algebra $\Lambda_{\mathbf{A}}$ and the topological Fukaya category $\mathcal{D}_{\infty}(\mathbf{S}^{\lambda})$ can be constructed as the bounded derived category $\mathcal{D}^b(\Lambda_{\mathbf{A}})$ of $\Lambda_{\mathbf{A}}$. The arcs correspond to indecomposable objects in $\mathcal{D}^b(\Lambda_{\mathbf{A}})$, cf. [HKK]. We have $\mathcal{D}_{\infty}(\mathbf{S}^{\lambda}) = \mathcal{D}_{\infty}(A_n)$ for a disk \mathbf{S}^{λ} with $(n+1)$ marked points. A geometric model for module category \mathcal{H} of an A_n quiver (with any orientation) is given in [BGMS] (and they construct a Tost with heart \mathcal{H} as a consequence).

If \mathbf{S}^λ has punctures, which carry additional \mathbb{Z}_2 -symmetry see [QZZ], cf. [FST, S, QZ], the story still works. Then $\Lambda_{\mathbf{A}}$ is a graded skew-gentle algebra. In particular, $\mathcal{D}_\infty(\mathbf{S}^\lambda) = \mathcal{D}_\infty(D_n)$ for a once-punctured disk \mathbf{S} with $n - 1$ marked points on $\partial\mathbf{S}$.

In this paper, we introduce another model for $\mathcal{D}_\infty(D_n)$, featuring double cover, which is similar, but different from the ones in [AP, AB]. More precisely, they take the double cover branching at punctures but we take the double cover of a once-punctured disk branching at a point other than the puncture. What we get is a doubly punctured disk with $2(n - 1)$ marked points on its boundary. As the other models mentioned above, we can also realized objects as arcs (without tagging). Moreover the two punctures in our model naturally correspond to the two ways of tagging in [FST, QZ, QZZ]. For instance, the tagged(-switching) rotation (introduced in [BQ]) corresponding to the Auslander-Reiten (AR) translation $\tau \in \text{Aut } \mathcal{D}_\infty(Q)$ becomes the puncture-switching rotation in our setting. See Figure 1 that $\tau(B_-V_3) = B_+V_2$ in both cases, where the subscripts $+/-$ denote untagged/tagged respectively in the left picture.




FIGURE 1. tagged/puncture-switching rotation as τ , type D_4

Moreover, after straightening the model and making it (centrally) symmetric (then the geometric center coincide with the branching point), the (oriented) arcs corresponding to the objects naturally becomes their central charges. Furthermore, we can easily describe total stability using such a model, which we will mention in Section 1.3.

One step further, we manage to find similar geometric models (to describe objects, central charge and total stability conditions) for all exceptional type categories, i.e. $\mathcal{D}_\infty(E_{6,7,8})$ (see Remark 5.1). Such models shed lights on understanding the geometry of $\text{Stab } \mathcal{D}_\infty(Q)$ and its connection to the root system/Kleinian singularities/Calabi-Yau categories.

1.3. Summary of notations and results. A (labelled) h -gon \mathbf{V} (in \mathbb{C}) with vertices $V_0, V_1, \dots, V_h = V_0$ and (oriented) edges $0 \neq z_j = V_{j-1}V_j, \forall j \in \mathbb{Z}_h$. We write VW for the vector (or oriented edges) \overrightarrow{VW} . The h -gon \mathbf{V} is *positively convex* if all other V_i is on the left hand side of the edge $z_j = V_{j-1}V_j$. A polygon is symmetric if it is centrally symmetric, which implies the number of vertices/edges is even.

For $1 \leq s \leq h/2$, the (oriented) *length- s diagonals* of an h -gon are $V_j V_{j+s}$. For instance, length-1 diagonals are just edges. For a positively convex h -gon \mathbf{V} , the *level- s diagonal-gon* is the convex polygon bounded by its length- s diagonals (i.e. on the left hand side of). See the hexagon bounded by the orange diagonals in Figure 3 as a level-2 diagonal-gon.

Let Q be a Dynkin quiver.

Definition 1.1. An h -gon of type Q is defined respectively as

A_n : an $(n + 1)$ -gon.

B_n : a symmetric $2n$ -gon with one puncture at its geometric center.

C_n : a symmetric $2n$ -gon.

D_n : a symmetric doubly punctured $2(n - 1)$ -gon.

E_6 : a 12-gon satisfying (4 triangle relations and 3 square relations):

$$\begin{cases} z_j + z_{j+4} + z_{j+8} = 0, \\ z_j - z_{j-3} + z_{j-6} - z_{j-9} = 0, \end{cases} \quad \forall j \in \mathbb{Z}_{12}. \quad (1.1)$$

Note that the rank of the 7 relation equations is actually 6.

E_7 : a symmetric 18-gon satisfying (3 hexagon relations):

$$z_j + z_{j+1} + z_{j+6} + z_{j+7} + z_{j+12} + z_{j+13} = 0, \quad \forall j \in \mathbb{Z}_{18}. \quad (1.2)$$

Note that after setting $z_{j+9} = -z_j$ (by the central symmetry), the rank of the 3 relation equations is actually 2.

E_8 : a symmetric 30-gon satisfying (5 triangle relations and 3 pentagon relations):

$$\begin{cases} z_j + z_{j+10} + z_{j+20} = 0, \\ z_j + z_{j+6} + z_{j+12} + z_{j+18} + z_{j+24} = 0, \end{cases} \quad \forall j \in \mathbb{Z}_{30}. \quad (1.3)$$

Note that after setting $z_{j+15} = -z_j$ (by the central symmetry), the rank of the 8 relation equations is actually 7.

F_4 : a symmetric 12-gon satisfying (1.1). Note that after setting $z_{j+6} = -z_j$ (by the central symmetry), the rank of the 7 relation equations in (1.1) further reduces to 4.

G_2 : a symmetric 6-gon satisfying (2 triangle relations)

$$z_j + z_{j+2} + z_{j+4} = 0, \quad \forall j \in \mathbb{Z}_6. \quad (1.4)$$

Note that after setting $z_{j+3} = -z_j$ (by the central symmetry), the two relation equations are equivalent.

Note that the edge vectors also need to satisfy the condition $\sum_{j=1}^h z_h = 0$, which is implied by the symmetric condition or the equations above except for type A_n . In each of the types except D_n , one can choose/fix n linear independent $0 \neq z_j \in \mathbb{C}$ to be the local coordinates and thus makes the space of h_Q -gons of Q a complex submanifold of \mathbb{C}^n . In type D_n , one can freely choose all $n - 1$ $\{z_j = -z_{j+n-1} \mid 1 \leq j \leq n - 1\}$'s together with one vector (say $V_0 B_+$ to determine the position of the two punctures) to form the $2(n - 1)$ -gon of type D_n . Thus, the space of h_Q -gons of Q is still a complex submanifold of \mathbb{C}^n in this case.

In particular, an h -gon of type Q is an h_Q -gon (satisfying extra conditions), where h_Q is the Coxeter number associated to the Dynkin diagram Δ .

For an exceptional case (or F_4 case), the relation (1.1), (1.2) or (1.3) induces a pair of $h_Q/2$ -gons, called the *ice* and *fire cores*. See Figure 7 for the two cores in type E_6 , the upper picture of Figure 9 for the ice core in type E_7 and Figure 10 for the ice core in type E_8 . The fire core is the symmetric mirror of the ice core in type E_7/E_8 .

Definition 1.2. An h_Q -gon of type Q is *stable* if it is positively convex and moreover:

- D_n : the punctures are inside the level- $(n-2)$ diagonal-gon.
- E_n : the ice and fire cores are inside the level- $(n-3)$ diagonal-gon, for $n \in \{6, 7, 8\}$.
- F_4 : same as E_6 . Note that the pair of 6-gons are symmetric to each other.

In particular, a stable h -gon means a positively convex h -gon without referring the types.

Two key observations are: (I) the central charges of objects in any τ -orbit of the root category form an h_Q -gon (see Proposition 3.6) and (II) there are distinguished τ -orbits, called the far-end τ -orbits (cf. Lemma 2.5, as well as the mid-end/near-end τ -orbits). We always fix a far-end τ -orbit and call the corresponding h_Q -gon the far-end one.

Definition 1.3. Denote by $\text{Stgon}(Q)$ the moduli space of stable h_Q -gons of type Q , where two stable h_Q -gons are equivalent if and only if they are parallel, i.e. are related by a translation of \mathbb{C} .

Note that the stability condition for an h_Q -gon is an open condition. Hence, $\text{Stgon}(Q)$ is an open subspace of the moduli space of h_Q -gon, which is a complex manifold of dimension n . For instance, in type E_6 , $\{z_j \mid 1 \leq j \leq 6\}$ is a choice of local coordinate for $\text{Stgon}(E_6)$ (satisfying certain inequalities).

Theorem 1. *An h_Q -gon of type Q provides a geometric model for the root category $\mathcal{D}_\infty(Q)/[2]$, in the sense that it induces a central charge naturally. Moreover, we have the isomorphism (between complex manifolds)*

$$Z_h: \text{ToSt}(Q)/[2] \cong \text{Stgon}(Q), \quad (1.5)$$

sending a total stability condition to the far-end stable h_Q -gon (of type Q).

If further quotient by $\mathbb{C}^* = \mathbb{C}/[2]$, where \mathbb{C} here is the natural \mathbb{C} -action on Stab , then we have

$$Z_h: \text{ToSt}(Q)/\mathbb{C} \cong \text{Sth}(Q),$$

where $\text{Sth}(Q)$ denotes the space of stable h -gon up to similarity.

A side product is that we have a simple description of the global dimension function on $\text{ToSt}(Q)$ (Theorem 3.7).

1.4. Connection to root systems. In this subsection, we explain the connection between $\text{ToSt}(Q)$, the root system $\Lambda(Q)$ and the space $\text{Stab } \mathcal{D}_2(Q)$ of stability conditions on $\mathcal{D}_2(Q)$. Here, $\mathcal{D}_2(Q)$ is the finite dimensional derived category of the Calabi-Yau-2 Ginzburg dg algebra/derived preprojective algebra associated to Q . It can also be

constructed from coherent sheaves for the Kleinian singularity \mathbb{C}^2/G , where G is the finite subgroup of $\mathrm{SL}_2(\mathbb{C})$ corresponding to Q (i.e. the McKay correspondence).

Let \mathfrak{g} be the finite-dimensional complex simple Lie algebra associated to Q , with Cartan subalgebra \mathfrak{h} and $\Lambda(Q)$ the corresponding root system. Let $\mathfrak{h}^{\mathrm{reg}}$ be the complement of the root hyperplanes in \mathfrak{h} :

$$\mathfrak{h}^{\mathrm{reg}} = \{v \in \mathfrak{h} \mid v(\alpha) \neq 0, \forall \alpha \in \Lambda(Q)\}.$$

The Weyl group W , generated by reflections of the root hyperplanes, acts freely on $\mathfrak{h}^{\mathrm{reg}}$. In [B2], Bridgeland shows

$$\mathrm{Stab} \mathcal{D}_2(Q) / \mathrm{Br}_Q \cong \mathfrak{h}^{\mathrm{reg}} / W_Q.$$

On the other hand, one expects that


$$\mathrm{Stab} \mathcal{D}_\infty(Q) \cong \mathfrak{h} / W_Q \tag{1.6}$$

and it is confirmed in [HKK] for type A . Clearly, there is a close relation between these two results. For instance, one can find a detailed case study in [BQS] for A_2 case. For all Dynkin case, there is also a conjectural description on the (almost) Frobenius structure on $\mathrm{Stab} \mathcal{D}_?(Q)$, which has been proved in [IQ2] for type A . More precisely, [IQ1] introduces the Calabi-Yau- \mathbb{X} category to link the Calabi-Yau- ∞ category and the Calabi-Yau-2 one. By the induction-reduction procedure there, each stability condition σ in $\mathrm{ToSt}(Q)$ induces a stability condition in $\mathrm{Stab} \mathcal{D}_2(Q)$.

Another connection between $\mathrm{ToSt}(Q)$ and the root system is the Gepner point σ_G , a stability condition with extra symmetry. In the Dynkin case, such a point is the (unique up to \mathbb{C} -action) solution to the equation (found by [KST])

$$\tau(\sigma) = (-2/h_Q) \cdot \sigma, \tag{1.7}$$

where $(-2/h_Q)$ is the \mathbb{C} -action on the right hand side. The stability condition σ_G is also the most stable one, in the sense that it is the minimal point of the global dimension function gldim , cf. [Q2, Thm. 4.7]. Interestingly, the central charge at the Gepner point for a Dynkin quiver Q is given by the projection of the root system on the Coxeter plane (Lemma 2.7). This was pointed out by Lutz Hille to Qy and Alastair King in Oberwolfach, Jan. 2020. We draw many pictures in Appendix A showing such projections (together with features of characterization of total stability conditions, i.e. the stable h_Q -gon). Here is the trailer:



1.5. Further studies. One of the applications of our description of $\text{ToSt}(Q)$ is to prove Reineke's conjecture.

Conjecture 1.4. [R, K] *For any Dynkin quiver Q , there is a stability function Z on $\mathcal{H}_Q = \text{mod } \mathbf{k}Q$ such that all indecomposable objects are stable.*

Note that a total stability condition induces a stability function on its heart that makes all indecomposable objects in the heart stable. Thus, to prove this conjecture, we only need to understand the heart of a total stability condition and construct a stable h_Q -gon so that its heart is equivalent to any given \mathcal{H}_Q . This will be done in the forthcoming sequel [CQZ].

Another follow-up work is to attack the following conjecture (we states for $\mathcal{D}_\infty(Q)$ but one could consider general case).

Conjecture 1.5. [Q2, Q3] *For any Dynkin quiver Q , the global dimension function gldim is a piece-wise Morse function on $\text{Stab } \mathcal{D}_\infty(Q)/\mathbb{C}$. Its only critical point is the Gepner point and hence induces a contractible flow on $\text{Stab } \mathcal{D}_\infty(Q)$.*

Note that (cf. [Q2]) we have $\text{ToSt}(Q) = \text{gldim}^{-1}[1 - 2/h_Q, 1)$. So $\text{ToSt}(Q)$ is the core of $\text{Stab } \mathcal{D}_\infty(Q)$. The conjecture above decomposes into two parts:

- 1°. gldim is piece-wise Morse on $\text{Stab } \mathcal{D}_\infty(Q) \setminus \text{ToSt}(Q)$ and thus $\text{Stab } \mathcal{D}_\infty(Q)$ contracts to $\text{ToSt}(Q)$.
- 2°. gldim is piece-wise Morse on $\text{ToSt}(Q)$.

The first part is proved in [Q3] for graded gentle algebras (and in particular for type A). Presumably, the method there should generalize to graded skew-gentle algebras (with extra effort to dealing with punctures, and in particular for type D) once the geometric model sets up (e.g. [QZZ]). Now with our geometric model for all Dynkin type, one may try to prove the first part for type E as well. Combining the expectation (1.6), the first part says that there is a canonical contraction: $\mathfrak{h}/W_Q \rightarrow \text{ToSt}(Q)$.

The second part is much more tricky, even in type A. The statement (say in type A) is easy to understand: the derivative of the function gldim gives a canonical way to deform any convex polygon into a regular polygon. This part is specific to the Dynkin case, while the phenomenon of the first part is much more general (e.g. for coherent sheaves on complex line in [Q2] and for coherent sheaves on complex plane for [FLLQ]).

Another potential application of our geometric model is to study tilted algebras of type E. The tilted algebras of type A and D lead to gentle and skew-gentle algebras, where the corresponding geometric models can be glued together to produce the topological Fukaya categories of (graded marked) surfaces, as mentioned above. It would be interesting to see how could these work in type E case.

Finally, Tost corresponds to $\text{gldim} < 1$ and one may want to study the ToSS =total semi-stability case, which corresponds to $\text{gldim} \leq 1$.

Acknowledgments. Qy would like to thank Alastair King for pushing him think carefully about total stability and suggesting the abbreviation ToSt among numerous comments. We would also like to thank Yu Zhou and Shiquan Ruan for pointing out

references for Coxeter elements and Wen Chang for proofreading. Qy is supported by National Key R&D Program of China (No. 2020YFA0713000), Beijing Natural Science Foundation (Grant No.Z180003) and National Natural Science Foundation of China (Grant No.12031007). Zx is supported by National Natural Science Foundation of China (Grant No. 12101422).

2. PRELIMINARIES

2.1. Stability conditions. Following [B1], we recall the notion of stability conditions on a triangulated category. In this paper, \mathcal{D} is a triangulated category with Grothendieck group $K\mathcal{D}$ and assume that $K\mathcal{D} \cong \mathbb{Z}^n$ for some n . Denote by $\text{Ind } \mathcal{D}$ the set of (isomorphism classes of) indecomposable objects in \mathcal{D} .

Definition 2.1. A *stability condition* $\sigma = (Z, \mathcal{P})$ on \mathcal{D} consists of a group homomorphism $Z: K\mathcal{D} \rightarrow \mathbb{C}$, called the *central charge*, and a family of full additive subcategories $\mathcal{P}(\phi) \subset \mathcal{D}$ for $\phi \in \mathbb{R}$, called the *slicing*, satisfying the following conditions:

- (a) if $0 \neq E \in \mathcal{P}(\phi)$, then $Z(E) = m(E) \exp(i\pi\phi)$ for some $m(E) \in \mathbb{R}_{>0}$,
- (b) for all $\phi \in \mathbb{R}$, $\mathcal{P}(\phi + 1) = \mathcal{P}(\phi)[1]$,
- (c) if $\phi_1 > \phi_2$ and $A_i \in \mathcal{P}(\phi_i)$ ($i = 1, 2$), then $\text{Hom}(A_1, A_2) = 0$,
- (d) for each object $0 \neq E \in \mathcal{D}$, there is a finite sequence of real numbers

$$\phi_1 > \phi_2 > \cdots > \phi_l \tag{2.1}$$

and a collection of exact triangles (known as the *HN-filtration*)

$$0 = E_0 \longrightarrow E_1 \longrightarrow E_2 \longrightarrow \cdots \longrightarrow E_{l-1} \longrightarrow E_l = E$$

with $A_i \in \mathcal{P}(\phi_i)$ for all i .

- (e) a technique condition, known as the support property, which holds automatically in our setting.

The categories $\mathcal{P}(\phi)$ are then abelian. Their non zero objects are called *semistable of phase ϕ* and simple objects *stable of phase ϕ* . For a semistable object $E \in \mathcal{P}(\phi)$, denote by $\phi_\sigma(E) := \phi$ its phase.

There is a natural \mathbb{C} -action on the set $\text{Stab } \mathcal{D}$ of all stability conditions on \mathcal{D} , namely:

$$s \cdot (Z, \mathcal{P}) = (Z \cdot e^{-i\pi s}, \mathcal{P}_{\text{Re}(s)}),$$

where $\mathcal{P}_x(\phi) = \mathcal{P}(\phi + x)$ and $s \in \mathbb{C}$. There is also a natural action on $\text{Stab } \mathcal{D}$ by the group of autoequivalences $\text{Aut } \mathcal{D}$, namely:

$$\Phi(Z, \mathcal{P}) = (Z \circ \Phi^{-1}, \Phi(\mathcal{P})),$$

where $\Phi \in \text{Aut } \mathcal{D}$.

The famous result in [B1] states that $\text{Stab } \mathcal{D}$ is a complex manifold with dimension $\text{rank } K\mathcal{D}$ and the local coordinate is provided by the central charge Z .

2.2. Total (semi)stability via global dimension function. Let $\sigma = (Z, \mathcal{P}) \in \text{Stab } \mathcal{D}$.

Definition 2.2. The *global dimension* of σ is defined as

$$\text{gldim } \mathcal{P} = \sup\{\phi_2 - \phi_1 \mid \text{Hom}(\mathcal{P}(\phi_1), \mathcal{P}(\phi_2)) \neq 0\} \in \mathbb{R}_{\geq 0} \cup \{+\infty\}, \quad (2.2)$$


which is a continuous function and an invariant under both the \mathbb{C} -action and $\text{Aut } \mathcal{D}$. The global dimension of \mathcal{D} is defined to be the inf of $\text{gldim } \sigma$ for all $\sigma \in \text{Stab } \mathcal{D}$.

Note that the notion generalizes the global dimension of an algebra/abelian category.

We recall the notion of total (semi)stability on triangulated categories, whose abelian version is due to Reineke [R], cf. [Q1, Conjecture 7.13] and comments there.

Definition 2.3. A stability condition σ is called *totally (semi)stable*, if any indecomposable object in \mathcal{D} is (semi)stable w.r.t. σ . We will call it a total stability condition for short.

2.3. Dynkin diagrams and KOT-Q classification. A *simply-laced Dynkin quiver* Q is an oriented graph whose underlying graph Δ is one of ADE Dynkin diagram. Explicitly, Δ is of the form $T_{p,q,r}$:



(2.3)

with $1 \leq p \leq q \leq r$ and

$$\frac{1}{p} + \frac{1}{q} + \frac{1}{r} > 1.$$

Then we have ($n = p + q + r - 2$)

- Q is of type A_n if $\Delta = T_{1,q,r}$ for $q + r = n - 1$.
- Q is of type D_n if $\Delta = T_{2,2,n-2}$ for $n \geq 4$.
- Q is of type E_n if $\Delta = T_{2,3,n-3}$ for $n \in \{6, 7, 8\}$.

Definition 2.4. For a simply-laced Dynkin quiver Q , we call its leaves (i.e. univalent vertices) the *boundary vertices*. Moreover, we call

- the leaf at the end of length p branch of Q the *near-end vertex*.
- the leaf at the end of length q branch of Q the *mid-end vertex*.
- the leaf at the end of length r branch of Q the *far-end vertex*.

Note that there is a choice involved for fixing the far-end vertex in type A_n , D_4 and E_6 .

Denote by $\mathcal{D}_\infty(Q) = \mathcal{D}^b(\mathbf{k}Q)$ the bounded derived category of the path algebra $\mathbf{k}Q$ for a simply-laced Dynkin quiver Q . Denote by $\text{AR } \mathcal{D}_\infty(Q)$ the *AR quiver* of $\mathcal{D}_\infty(Q)$, which is isomorphic to $\mathbb{Z}Q$. Each vertex of Q canonically corresponds to a τ -orbit (containing the corresponding projective). The τ -orbit that corresponds to an xx vertex is call xx τ -orbit, for xx being boundary/near-end/mid-end/far-end.

The following is well-known.

Lemma 2.5. *For type A_n and E_6 , there are two choices of far-end τ -orbits of $\text{AR } \mathcal{D}_\infty(Q)$, which are shift [1] of each other. For type D_4 , there are three choices of far-end τ -orbit of $\text{AR } \mathcal{D}_\infty(Q)$ (i.e. any boundary τ -orbit). For other cases, there is a unique choice of the far-end τ -orbit.*


For the orbit categories of $\mathcal{D}_\infty(Q)$ (e.g. the root category $\mathcal{D}_\infty(Q)/[2]$ we are about to mention in particular), we keep the same notions. We will always choose a preferred far-end vertex/ τ -orbit in each case.

A non-simply-laced Dynkin quiver Q , denoted by R^ι , is defined to be a simply-laced Dynkin quiver R together with an automorphism ι of R . All possible cases are listed as follows (we omit the orientations but they should be compatible with ι):

B_n : R is of type D_{n+1} and R^ι is of type B_n while ι exchanges the black bullets in the same column.



C_n : R is of type A_{2n-1} and R^ι is of type C_n while ι exchanges the black bullets in the same column.



F_4 : R is of type E_6 and R^ι is of type F_4 while ι exchanges the black bullets in the same column.



G_2 : R is of type D_4 and R^ι is of type G_2 while ι rotates the black bullets in the same column.



The label below any black bullet in R^t is its weight (i.e. number of vertices in that orbit).

In the non-simply-laced case, $\mathcal{D}_\infty(Q)$ means the ι -stable category $\mathcal{D}_\infty(R)^t$ and the boundary/far-end vertices (of Q) and τ -orbits (of $\text{AR } \mathcal{D}_\infty(Q)$) are induced from R . Lemma 2.5 implies that there is a unique far-end τ -orbit in any non-simply-laced case.

Denote by $\text{ToSt } \mathcal{D}$ the subspace of $\text{Stab } \mathcal{D}$ consisting of all total stability conditions on a triangulated category \mathcal{D} . The classification theorem below is a combination of the following two results:

- [Q2, Prop. 3.5]: σ is totally stable if and only if $\text{gldim } \sigma < 1$.
- [Q3, Thm. 3.2]: \mathcal{D} admits a stability condition σ with $\text{gldim } \sigma < 1$ if and only if \mathcal{D} equals $\mathcal{D}_\infty(Q)$ for a Dynkin diagram Δ .

Note that the second result is essentially due to Kikuta -Ouchi-Takahashi, where they impose a mild condition that excludes the non simply-laced case.

Theorem 2.6. *ToSt \mathcal{D} is non-empty if and only if \mathcal{D} equals $\mathcal{D}_\infty(Q)$ for a Dynkin quiver Q . In such a case, we write $\text{ToSt}(Q)$ for $\text{ToSt } \mathcal{D}_\infty(Q)$.*

2.4. Root system. Let \mathfrak{g} be the complex simple Lie algebra of type Q and \mathfrak{h} its Cartan subalgebra. Denote by $\Lambda(Q)$ the associated *root system* and h_Q the *Coxeter number*.

The *Coxeter element* w is the product of all simple reflections. Although it depends on the order of the product, all such elements are conjugate to each other. Thus, up to symmetry, there is essentially one Coxeter element. After fix the Coxeter element w , there is a unique plane P_w , known as the *Coxeter plane*, on which w acts by rotation by $2\pi/h_Q$.

For instance, the projection of the root system on the Coxeter plane P_w of type D_5 , E_6 , E_7 and E_8 are shown in Figures 13, 14, 15 and 16 respectively.

2.5. Root categories. Note that the dimension function \dim depends on the orientation of Q but the root category does not. Recall the famous Gabriel's theorem that the root category categorifies the root system, in the sense that there is a bijection

$$\dim: \text{Ind } \mathcal{D}_\infty(Q)/[2] \rightarrow \Lambda(Q). \quad (2.4)$$

Set $\bar{\tau} = \tau^{-1}$. It is well-known that

$$\bar{\tau}^{h_Q} = [2]$$

and $\mathbb{S} = [1] \circ \tau$ is the Serre functor. The following observation is due to L. Hille.

Lemma 2.7. *Under the bijection in (2.4), the projection of the root system on the Coxeter plane gives the central charge of the Gepner point σ_G of $\mathcal{D}_\infty(Q)$.*

Proof. As the Coxeter element w acts by rotation by $2\pi/h_Q$, the roots with the same length spread evenly in P_w . This matches the Gepner equation (1.7) and implies the lemma. See the top pictures in Figures 14 and 15. \square

3. PROTOTYPE OF ToSt (I.E. TYPE A_n)3.1. The h_Q -gons induced by τ -orbits.

Lemma 3.1. $\text{id} - \tau$ (or $\text{id} + \mathbb{S}$) is a non-singular linear map on $\mathbb{R}^n = K\mathcal{D}_\infty(Q) \otimes \mathbb{R}$.

Proof. Taking the basis $\{\dim P_i \mid i \in Q_0\}$ of \mathbb{R}^n for P_i being the projectives of Q . As $\tau(P_i) = I_i[-1]$ for I_i being the injectives, the matrix presentation $C(Q)$ of τ , as a linear transformation on $K\mathcal{D}_\infty(Q)$ is the so-called *Coxeter transformation* of Q . It is well-known that the eigenvalues of the Coxeter transformation is not 1 for Dynkin type (cf. [L]). Thus the lemma follows. \square

For any M in $\text{Ind } \mathcal{D}_\infty(Q)/[2]$, define

$$g(M) := \sum_{j=1}^{h_Q} [\tau^j M],$$

to be the sum of all objects in the τ -orbit of M in $K\mathcal{D}_\infty(Q)$.

Proposition 3.2. *The function $g \equiv 0$.*

Proof. Note that τ is a group automorphism of $K\mathcal{D}_\infty(Q)$. Then $\tau^{h_Q} = [-2] = \text{id}$ implies

$$(\text{id} - \tau)(\text{id} + \tau + \cdots + \tau^{h_Q-1}) = 0.$$

Then Lemma 3.1 implies the proposition.

An alternating categorical proof is as follows, which fits better in our content. We claim that any $g(P_i), i \in Q_0$ are proportional. Note that we only need to check this for the simply laced case, as the other cases can be obtained from folding and hence hold also well. Then

$$\sum_{i \in Q_0} g(P_i) = \sum_{M \in \text{Ind } \mathcal{D}_\infty(Q)/[2]} [M] = \sum_{M \in \text{Ind } \mathcal{H}_Q} ([M] + [M[1]]) = 0$$

implies the proposition. For the claim, let i be a leaf of Q (e.g. the boundary vertex, cf. Definition 2.4) and j be its neighbour vertex. Summing all the mesh relations in $K\mathcal{D}_\infty(Q)$ at the τ -orbit of P_i , which are triangle relations, we have

$$2g(P_i) = g(P_j).$$

If j is not the tri-valent vertex in (2.3), let k be its neighbour vertex other than i . Summing all the mesh relation in $K\mathcal{D}_\infty(Q)$ at the τ -orbit of P_i , which are square relations, we have

$$2g(P_j) = g(P_i) + g(P_k)$$

and hence $g(P_k) = 3g(P_i)$. Do all such calculations, we see the claim holds. \square

Let Z be any central charge and denote by $\bar{\tau} = \tau^{-1}$. Then we have the following fact:

Corollary 3.3. *For any $M \in \text{Ind } \mathcal{D}_\infty(Q)$, the vectors $z_1 = Z(\bar{\tau}M), z_2 = Z(\bar{\tau}^2M), \dots, z_{h_Q} = Z(\bar{\tau}^{h_Q}M)$ form an h_Q -gon.*

Definition 3.4. The h_Q -gon (w.r.t. to any given central charge Z) induced by a boundary/?-end τ -orbit (of $\text{AR } \mathcal{D}_\infty(Q)/[2]$) is called the *boundary/?-end h_Q -gon*, for ? being mid/near/far.

Once we choose our favorite far-end vertex/ τ -orbit, we will have the corresponding far-end h_Q -gon. It will not make much difference in type A_n and E_6 . But as we will explain in Section 4.4, the choice matters in type D_4 .

3.2. Positive convexity as stability for h_Q -gons.

Lemma 3.5. *Let $\sigma = (Z, \mathcal{P}) \in \text{ToSt}(Q)$. If there is a (non-trivial) path from M to L in the AR quiver of $\mathcal{D}_\infty(Q)$, then $\phi_\sigma(M) < \phi_\sigma(L)$.*

Proof. Any path in the AR-quiver consists of irreducible morphism (which are non-zero). Hence, the total stability implies that the phases of (stable) objects in the path are increasing. \square

Proposition 3.6. *If $\sigma = (Z, \mathcal{P}) \in \text{ToSt}(Q)$, then, for any $M \in \text{Ind } \mathcal{D}_\infty(Q)$, the vectors $Z(\bar{\tau}M), Z(\bar{\tau}^2M), \dots, Z(\bar{\tau}^{h_Q}M)$ form a positively convex h_Q -gon.*

Proof. In the A_1 case, the statement is trivial. Assume that $n = |Q_0| \geq 2$ in the following.

In the AR quiver $\mathcal{D}_\infty(Q)$, there is a path from M to $\bar{\tau}M$ and hence $\phi_\sigma(M) < \phi_\sigma(\bar{\tau}M)$ by Lemma 3.5. Then

$$\phi_\sigma(M) < \phi_\sigma(\bar{\tau}M) < \dots < \phi_\sigma(\bar{\tau}^{h_Q-1}M) < \phi_\sigma(\bar{\tau}^{h_Q}M) = \phi_\sigma(M[2]) = \phi_\sigma(M) + 2.$$

Let $z_i = Z(\bar{\tau}^i M), \forall i \in \mathbb{Z}_{h_Q}$. Then $\arg z_0 = \pi \cdot \phi_\sigma(M)$. So we have

$$\arg z_0 < \arg z_1 < \dots < \arg z_{h_Q-1} < \arg z_{h_Q} + 2\pi = \arg z_0 + 2\pi,$$

where \arg takes values in $[\pi \cdot \phi_\sigma(M), \pi \cdot \phi_\sigma(M) + 2\pi)$. It implies the h_Q -gon with edges z_i is indeed positively convex. \square


Up to translation, each τ -orbit induces one positively convex h_Q -gon for a chosen $\sigma \in \text{ToSt}(Q)$. We call all such h_Q -gons the σ -induced positively convex h_Q -gons.

A direct consequence is the following characterization of gldim on $\text{ToSt}(Q)$, which generalizes [Q2, Prop. 3.6] for type A_n to all Dynkin cases.

Theorem 3.7. *If $\sigma \in \text{ToSt}(Q)$, then $\pi \cdot \text{gldim } \sigma$ equals the maximal angle among any interior angles of σ -induced positively convex h_Q -gons.*

Proof. By AR duality, we have $\text{Hom}(E, F) = D \text{Hom}(F, \tau(E[1]))$. So in particular, $\text{Hom}(E, \tau(E[1])) = D \text{Hom}(E, E) \neq 0$. Moreover, by hammock property of AR-quiver of $\mathcal{D}_\infty(Q)$, we know that $\text{Hom}(E, F) \neq 0$ implies that there is path from E to F to $\tau(E)[1]$ in the AR-quiver, for any indecomposable E, F . By Lemma 3.5,

$$\phi_\sigma(E) \leq \phi_\sigma(F) \leq \phi_\sigma(\tau(E[1])),$$

FIGURE 2. AR quiver of $\mathcal{D}_\infty(A_n)/[2]$

and thus $\phi_\sigma(F) - \phi_\sigma(E) \leq \phi_\sigma(\tau(E[1])) - \phi_\sigma(E)$. Hence,

$$\text{gldim } \sigma = \max\{\phi_\sigma(\tau(E[1])) - \phi_\sigma(E) \mid E \in \text{Ind } \mathcal{D}_\infty(Q)\}.$$

Finally, the statement follows from that $\pi \cdot (\phi_\sigma(\tau(E[1])) - \phi_\sigma(E))$ is indeed an interior angle of the h_Q -gons induced by the τ -orbit containing E (with respect to σ). \square

3.3. ToSt of type A_n . Although the orientation of the quiver does not matter when considering $\text{ToSt}(Q)$, we will choose our favorite one for convenience (e.g. to describe objects).

Set-up 3.8. Take the A_n quiver with straight orientation

$$1 \longleftarrow 2 \longleftarrow \cdots \longleftarrow n \quad (3.1)$$

We declare vertex 1 as the fixed far-end vertex and label the indecomposable objects in $\mathcal{D}_\infty(A_n)$ by

$$P_i^j := \bar{\tau}^j P_i, \quad j \in \mathbb{Z}, 1 \leq i \leq n, \quad (3.2)$$

where each P_i is the projective corresponding to vertex i . The AR quiver of $\mathcal{D}_\infty(A_n)/[2]$ is illustrated as in Figure 2, where $j(j+i)$ denotes P_i^j and $(j+i)j$ denotes $P_i^j[1]$ for $1 \leq i \leq n$ and $0 \leq j \leq n-i$.


Note that the oriented diagonals of a $(n+1)$ -gon is a model for indecomposable objects in the root category of type A_n is well-known, cf. e.g. [BM].

Recall that $\text{Stgon}(A_n)$ denotes the moduli space of stable $(n+1)$ -gons on \mathbb{C} up to translation. The following proposition is a \mathbb{C}^* -covering version of [Q2, Prop. 3.6]. But the proof is essentially the same.

Proposition 3.9. *There is a natural isomorphism*

$$Z_h: \text{ToSt}(A_n)/[2] \rightarrow \text{Stgon}(A_n)$$

sending a total stability condition σ to the far-end $(n+1)$ -gon.


 FIGURE 3. The 6-gon of type A_5 with level-2 diagonal-gon shaded


More precisely, we have $z_j = Z(P_1^{j-1})$ for $j \in \mathbb{Z}_{n+1}$ for the edges of the far-end $(n+1)$ -gon and

$$Z(P_i^j) = V_j V_{j+i}, \quad 1 \leq i \leq n, 0 \leq j \leq n-i.$$

A study of total stability for the module category \mathcal{H} of type A is in [Ki], which can be deduced from the description above after fixing the heart of a total stability condition to be \mathcal{H} .

4. ToSt OF TYPE D_n

Set-up 4.1. Set $m = n - 1$ and then $h_Q = 2m$ for $Q = D_n$. We choose an orientation as follows:



Moreover, we fix 1 as our favorite far-end vertex for $n = 4$ and it is the unique far-end vertex for $n \geq 5$. Label the indecomposable objects in the $\mathcal{D}_\infty(D_n)$ still by (3.2). Thus the i^{th} - τ -orbit consisting of objects $\{P_i^j\}_{j \in \mathbb{Z}}$ is the one corresponding to vertex i .

As $\bar{\tau}^{2m} = [2]$, the indecomposables in the root category are $\{P_i^j \mid 1 \leq i \leq n, j \in \mathbb{Z}_{2m}\}$. For instance, see the AR quiver $\text{AR } \mathcal{D}_\infty(D_5)/[2]$ in the low picture of Figure 4, where $j(j+i)$ denotes P_i^j , $j \varrho(j)$ denotes P_m^j and $j \varrho(j+1)$ denotes P_n^j . Here

$$\varrho(x) = \text{sign}(-1)^x: \mathbb{Z} \rightarrow \{\pm\}$$

is the parity function.

4.1. Symmetric doubly punctured h -gons.

Definition 4.2. A *symmetric h -gon \mathbf{V}* is an h -gon (cf. notations in Section 1.3) such that h is even and

$$z_{j+h/2} = -z_j. \tag{4.1}$$

Let O be its geometric center. It is *doubly punctured* if there is a pair of punctures B_{\pm} such that O is the middle point of them.




FIGURE 4. The 8-gon of type D_5 and AR quiver of $\mathcal{D}_{\infty}(D_5)$.

Given a central charge $Z: K\mathcal{D}_\infty(D_n) \rightarrow \mathbb{C}$, we construct a symmetric doubly punctured $2(n-1)$ -gon (\mathbf{V}_Z, B_\pm) as follows:

$$\begin{cases} V_j = -\left(\sum_{t=j}^{j+m-1} Z(P_1^t)\right)/2, \\ B_+ = (Z(P_m) - Z(P_n))/2 \\ B_- = (Z(P_n) - Z(P_m))/2. \end{cases} \quad (4.2)$$

So $z_j = V_{j-1}V_j = Z(P_1^{j-1})$ noticing $Z(P_1^{j+m-1}) = -Z(P_1^{j-1})$. More precisely,

- $\mathbf{V} = \mathbf{V}_Z$ is the far-end $2(n-1)$ -gon, which is indeed symmetric since

$$\bar{\tau}^m = [1]$$

holds on the far-end τ -orbit $\{P_1^t \mid t \in \mathbb{Z}_{2m}\}$ that implies (4.1).

- Moreover, the geometric center of \mathbf{V} is at origin as

$$\begin{cases} V_j = -V_{j+m}, & j \in \mathbb{Z}_{2m} \\ B_+ = -B_-. \end{cases} \quad (4.3)$$

Definition 4.3. An h -gon of type D_n is a symmetric doubly punctured $2(n-1)$ -gon. It is *stable* (of type D_n) if it is positively convex and the punctures are inside the level- $(n-2)$ diagonal-gon. Denote by $\text{Stgon}(D_n)$ the moduli space of stable $2(n-1)$ -gon of type D_n up to translation, which has complex dimension n .


Next we show that total stability implies the stability of $2(n-1)$ -gon of type D_n .

Proposition 4.4. *If Z is the central charge of some $\sigma \in \text{ToSt}(D_n)$, then the symmetric doubly punctured $2(n-1)$ -gon (\mathbf{V}, B_\pm) defined above is a stable $2(n-1)$ -gon of type D_n . We call it the far-end stable $2(n-1)$ -gon with respect to σ .*

Proof. We need to show that the total stability of σ implies the stability of the corresponding far-end h -gon \mathbf{V} together with punctures B_\pm .

Firstly, the positive convexity of \mathbf{V} follows from Proposition 3.6.

Secondly, we will check that B_\pm is inside the level- $(m-1)$ diagonal-gon. Namely, for any $j \in \mathbb{Z}_{2m}$, it suffices to show that B_\pm is on the left hand side of $V_{j+1}V_{j+m}$:



Note that we have AR triangles ($i \in \{m, n\}$)

$$P_i^j \rightarrow P_{n-2}^{j+1} \rightarrow P_i^{j+1} \rightarrow P_i^j[1]$$

with central charges

$$V_j B_\pm = B_\mp V_{j+m}, \quad V_{j+1} V_{j+m}, \quad V_{j+1} B_\mp \quad \text{and} \quad -V_j B_\pm$$

Here $\pm = \varrho(j)$ if $i = m$ and $\pm = \varrho(j+1)$ if $i = n$ to be more precise. Since all P_i^j are stable objects, their phases are increasing. Therefore we have

$$\arg B_\mp V_{j+m} < \arg V_{j+1} V_{j+m} < \arg V_{j+1} B_\mp < \arg B_\mp V_{j+m} + \pi,$$

where \arg takes values in $[\pi \cdot \phi_\sigma(P_i^j), \pi \cdot \phi_\sigma(P_i^j) + 2\pi)$. This completes the proof. \square

4.2. Geometric model for root category of type D_n . Suppose that we have a $2(n-1)$ -gon (\mathbf{V}, B_\pm) of type D_n (with vertices V_j and punctures B_\pm). Up to translation, we may assume its geometric center is at the origin, i.e. (4.3) holds. Then we have

$$\begin{cases} V_j V_k = V_{k+m} V_{j+m}, \\ V_j B_+ = B_- V_{j+m}. \end{cases} \quad (4.4)$$

Theorem 4.5. *A $2(n-1)$ -gon of type D_n is a geometric model for the root category $\mathcal{D}_\infty(D_n)/[2]$ in the sense that by setting*

$$\begin{cases} Z(P_i^j) = V_j V_{j+i} = V_{j+i+m} V_{j+m}, & j \in \mathbb{Z}_{2m}, 1 \leq i \leq m-1, \\ Z(P_m^j) = V_j B_{\varrho(j)} = B_{\varrho(j+1)} V_{j+m}, & j \in \mathbb{Z}_{2m}, \\ Z(P_n^j) = V_j B_{\varrho(j+1)} = B_{\varrho(j)} V_{j+m}, & j \in \mathbb{Z}_{2m}, \end{cases} \quad (4.5)$$

we obtain a central charge $Z: K \mathcal{D}_\infty(D_n) \rightarrow \mathbb{C}$.

Proof. To show that Z is a group homomorphism, we only need to check all the mesh relations in the root category still hold after applying Z . To start with, notice that

$$Z(P_m^j) + Z(P_n^j) = V_j V_{j+m} \quad (4.6)$$

by (4.4). The rest of the proof is just a direct checking:

1°. At the τ -orbit of $i = 1$ the mesh relation has the form

$$[P_1^j] + [P_1^{j+1}] = [P_2^j]$$

and indeed we have

$$V_j V_{j+1} + V_{j+1} V_{j+2} = V_j V_{j+2}.$$

2°. At the τ -orbit of $1 < i < n-2$ the mesh relation has the form

$$[P_i^j] + [P_i^{j+1}] = [P_{i-1}^{j+1}] + [P_{i+1}^j]$$

and indeed we have

$$V_j V_{j+i} + V_{j+1} V_{j+i+1} = V_{j+1} V_{j+i} + V_j V_{j+i+1}.$$

3°. At the τ -orbit of $i = n - 2$ the mesh relation has the form

$$[P_{n-2}^j] + [P_{n-2}^{j+1}] = [P_{n-3}^{j+1}] + [P_m^j] + [P_n^j]$$

and indeed we have, using (4.6),

$$V_j V_{j+n-2} + V_{j+1} V_{j+n-1} = V_{j+1} V_{j+n-2} + V_j V_{j+n-1}.$$

4°. At the τ -orbit of $i \in \{m, n\}$, the mesh relation has the form

$$[P_i^j] + [P_i^{j+1}] = [P_{n-2}^{j+1}]$$

and indeed we have, using (4.4),

$$V_j B_{\pm} + V_{j+1} B_{\mp} = B_{\mp} V_{j+m} + V_{j+1} B_{\mp} = V_{j+1} V_{j+m}.$$

Note that here the sign \pm depends on $i \in \{m, n\}$ and the parity function ρ .

Note that all these calculations can be easily checked in Figure 4 for case $n = 5$. \square

Example 4.6 (Type $Q = D_5$). In Figure 4, we have

- The objects drawn in blue/violet/green circle, in the τ -orbit of $\text{AR } \mathcal{D}_{\infty}(Q)$, correspond to length-1/2/3 diagonals drawn in blue/violet/green respectively as in the upper 8-gons.
- The objects drawn in orange circle, in the upper/lower τ -orbit of $\text{AR } \mathcal{D}_{\infty}(Q)$, correspond to solid/dashed orange line segments respectively as in the right upper 8-gon.

4.3. From stable h -gon to total stability conditions of type D_n . Given a stable $2(n-1)$ -gon \mathbf{V} of type D_n , we can construct a total stability condition $\sigma = (Z, \mathcal{P})$ with Z defined as in Theorem 4.5.

Construction 4.7. Let us construct a slicing \mathcal{P} as follows.

Step 1: Assign a real number $\phi(M)$ for each object M in the far-end τ -orbit of $\text{AR } \mathcal{D}_{\infty}(D_n)$:

- Let

$$\phi(P_1[k]): = \arg Z(P_1)/\pi + k,$$

for any $k \in \mathbb{Z}$, where \arg takes values in $[0, 2\pi)$.

- For $1 \leq j \leq m-1$ and $k \in \mathbb{Z}$, let

$$\phi(P_1^j[k]): = \arg Z(P_1^j)/\pi + k,$$

where \arg takes values in $[\arg Z(P_1), Z(P_1) + 2\pi)$.

So we have the monotonicity and periodicity:

$$\begin{cases} \phi(P_1^j) < \phi(P_1^{j+1}), \\ \phi(P_1^{j+m}) = \phi(P_1^j[1]) = \phi(P_1^j) + 1, \end{cases} \quad \forall j \in \mathbb{Z}. \quad (4.7)$$

Step 2: For any $2 \leq i \leq n-2 < m$ and $j \in \mathbb{Z}$, we have

$$Z(P_i^j) = V_j V_{j+i} = \sum_{t=j}^{j+i-1} V_t V_{t+1} = Z(P_1^j) + \dots + Z(P_1^{j+i-1}).$$

Note that by Step 1, the positive convexity of \mathbf{V} gives

$$\phi(P_1^j) < \phi(P_1^{j+1}) < \dots < \phi(P_1^{j+i-1}) < \phi(P_1^{j+m}) = \phi(P_1^j[1]) = \phi(P_1^j) + 1$$

Then let

$$\phi(P_i^j[k]): = \arg Z(P_i^j)/\pi + k, \quad (4.8)$$

where \arg takes values in $[\pi \cdot \phi(P_1^j), \pi \cdot \phi(P_1^j) + 2\pi)$. It is straightforward to check that the monotonicity and periodicity of (4.7) are inherited:

$$\begin{cases} \phi(P_i^j) < \phi(P_i^{j+1}), \\ \phi(P_i^{j+m}) = \phi(P_i^j[1]) = \phi(P_i^j) + 1, \end{cases} \quad \forall j \in \mathbb{Z}.$$

Note that, locally, the additive subcategory of $\text{AR } \mathcal{D}_\infty(D_n)$ generated by

$$\begin{cases} P_1^j, \dots, P_1^{j+i-2}, P_1^{j+i-1} \\ P_2^j, \dots, P_2^{j+i-2} \\ \dots \\ P_i^j \end{cases} \quad (4.9)$$

is isomorphic to $\text{mod } \mathbf{k}A_i$ for an A_i quiver with straight orientation. Inductively, the positive convexity implies that

$$\begin{cases} \phi(P_t^s) < \phi(P_{t-1}^{s+1}) < \phi(P_t^{s+1}) & 1 < t \leq i, j \leq s < j + i - t \\ \phi(P_t^s) < \phi(P_{t+1}^s) < \phi(P_t^{s+1}) & 1 \leq t < i, j \leq s < j + i - t. \end{cases} \quad (4.10)$$

Step 3: For $i \in \{m, n\}$ and $j \in \mathbb{Z}$, let

$$\phi(P_i^j): = \arg Z(P_i^j)/\pi, \quad (4.11)$$

where \arg takes value in $[\pi \cdot \phi(P_{n-2}^j), \pi \cdot \phi(P_{n-2}^j) + 2\pi)$. Note that

$$\phi(P_{n-2}^{j+m}) = \phi(P_{n-2}^j[1]) = \phi(P_{n-2}^j) + 1,$$

and

$$\begin{cases} \phi(P_m^{j+m}) = \phi(P_m^j[1]) = \phi(P_m^j) + 1, \\ \phi(P_n^{j+m}) = \phi(P_n^j[1]) = \phi(P_n^j) + 1, \end{cases}$$

if m is even and

$$\begin{cases} \phi(P_m^{j+m}) = \phi(P_n^j[1]) = \phi(P_n^j) + 1, \\ \phi(P_n^{j+m}) = \phi(P_m^j[1]) = \phi(P_m^j) + 1, \end{cases}$$

if m is odd. This completes the assigning ϕ .

Step 4: Define

$$\mathcal{P}(\varphi) = \text{Add} \left(\bigoplus_{\substack{P \in \text{Ind } \mathcal{D}_\infty(D_n) \\ \phi(P) = \varphi}} P \right) \quad (4.12)$$

Proposition 4.8. $\sigma = (Z, \mathcal{P})$ defined as above is a total stability condition on $\mathcal{D}_\infty(D_n)$.

Proof. By construction, we already have


- all indecomposable objects are in some $\mathcal{P}(\phi)$ for $\phi \in \mathbb{R}$ and
- \mathcal{P} is compatible with the central charge Z as well as the shift $[1]$.

Thus, what is left to show is $\phi(M) < \phi(L)$ whenever there is a non-zero map from M to L for any indecomposable objects M, L . This amounts to check that, for any arrow $M \rightarrow L$ in $\text{AR } \mathcal{D}_\infty(D_n)$, we have $\phi(M) < \phi(L)$. There are three cases:

- If the arrow is between the far-end τ -orbit (corresponding to vertex 1) and the double-trivalent τ -orbit (corresponding to vertex $n - 2$), then it is a type A_n issue, cf. (4.9), and (4.10) gives the inequality.
- If the arrow is from the double-trivalent τ -orbit to the boundary τ -orbits (corresponding to vertex m or n), then (4.11) implies the inequality.
- If the arrow is from the boundary τ -orbits (corresponding to vertex m or n) to the double-trivalent τ -orbit, then we need to examine the triangle

$$\begin{array}{ccccc}
 & & P_{n-3}^{j+1} & & \\
 & \nearrow & & \searrow & \\
 P_{n-2}^j & \longrightarrow & P_m^j & \longrightarrow & P_{n-2}^{j+1} \\
 & \searrow & & \nearrow & \\
 & & P_n^j & &
 \end{array}$$

that corresponds to



Then, from the stability condition of (\mathbf{V}, B_{\pm}) , we know that B_{\pm} is bounded by solid lines in the above picture. By (4.8) and (4.11), we deduce that both $\phi(P_i^j)$, $i \in \{m, n\}$, and $\phi(P_{n-2}^{j+1})$ are in $[\phi(P_{n-2}^j), \phi(P_{n-2}^j) + 2)$. Thus we have

$$\begin{aligned}
 \phi(P_{n-2}^j) &= \frac{\arg V_j V_{j+m-1}}{\pi} \\
 &< \frac{\arg V_j B_{\pm}}{\pi} = \phi(P_i^j) \\
 &< \frac{\arg V_{j+1} V_{j+m}}{\pi} = \phi(P_{n-2}^{j+1})
 \end{aligned}$$

where \pm in the second row depends on $i \in \{m, n\}$ and the parity function ρ . Note that \arg takes values in $[\pi \cdot \phi(P_{n-2}^j), \pi \cdot \phi(P_{n-2}^j) + 2\pi)$.

In all, the first case uses positive convexity of \mathbf{V} and the last two cases uses the extra condition of stability of (\mathbf{V}, B_{\pm}) . \square

Remark 4.9. Alternatively, one can define \mathcal{P} as follows:

- For any $1 \leq i \leq n$, let

$$\phi(P_i) = \arg Z(P_i)/\pi,$$

where \arg takes value in $[\arg Z(P_1), Z(P_1) + 2\pi)$.

- For fixed $1 \leq i \leq n$, let

$$\phi(P_i^j) = \arg Z(P_i^j)/\pi$$

for $1 \leq j \leq m$, where \arg takes values in $[\arg Z(P_i), Z(P_i) + 2\pi)$.

- Define \mathcal{P} as in (4.12).

But the simplicity of this construction will trade off more complexity of the proof above.

Theorem 4.10. *There is a natural isomorphism*

$$Z_h: \text{ToSt}(D_n)/[2] \rightarrow \text{Stgon}(D_n),$$

sending a total stability condition σ to the far-end stable $2(n-1)$ -gon.

Proof. It follows from combining Propositions 4.4 and 4.8, noticing that in the slices can shift two simultaneously in Construction 4.7. \square




FIGURE 5. Three far-end 6-gons of type D_4 for a given central charge

4.4. Example: D_4 with three far-end stable 6-gons. As mentioned above couple of times, there are three choices of far-end vertices/ τ -orbits/6-gons. The isomorphism in Theorem 4.10 depends on such a choice. Here, we give an example, as shown in Figure 5, of how the three (far-end) stable 6-gons look like (3 hexagons on the left), with respect to a fixed $\sigma \in \text{ToSt}(D_4)$ (whose central charge is on the right).

5. EXCEPTIONAL TYPE: PREVIEW

For an exceptional quiver $Q = E_n$, where $n \in \{6, 7, 8\}$, we fix the following orientation

$$\begin{array}{ccccccc} & & & 3 & & & \\ & & & \uparrow & & & \\ 1 & \leftarrow & 2 & \leftarrow & 4 & \longrightarrow & \cdots \longrightarrow n \end{array} \quad (5.1)$$

and still label the indecomposable objects in the $\mathcal{D}_\infty(E_n)$ by (3.2). The objects in the boundary τ -orbits play the key role in constructing geometric model and we label them as

$$\begin{cases} 1^{\text{st}}\text{-}\tau\text{-orbit: Denote } C_j = P_1^{j-1}. & \text{This is the mid-end } \tau\text{-orbit.} \\ 3^{\text{rd}}\text{-}\tau\text{-orbit: Denote } M_j = P_3^{j-1}. & \text{This is the near-end } \tau\text{-orbit.} \\ n^{\text{th}}\text{-}\tau\text{-orbit: Denote } B_j = P_n^{j-1}. & \text{This is the far-end } \tau\text{-orbit.} \end{cases} \quad (5.2)$$

In Figure 6, 9 and 11, the objects in the mid-end/near-end/far-end orbits will be drawn in green/yellow/blue in the AR-quiver, respectively.

Moreover, we will use the following family of triangles

$$B_j \rightarrow C_{j+2} \rightarrow B_{j+n-2} \rightarrow B_j[1] \quad (5.3)$$

and octahedral diagrams

$$\begin{array}{ccccccc} & & B_j & \xlongequal{\quad} & B_j & & \\ & & \downarrow & & \downarrow & & \\ C_j & \longrightarrow & M_{j+1} & \longrightarrow & B_{j+n-3} & \longrightarrow & C_j[1] \\ \parallel & & \downarrow & & \downarrow & & \parallel \\ C_j & \longrightarrow & C_{j+3} & \longrightarrow & L_j & \longrightarrow & C_j[1] \\ & & \downarrow & & \downarrow & & \\ & & B_j[1] & \xlongequal{\quad} & B_j[1] & & \end{array} \quad (5.4)$$

in $\mathcal{D}_\infty(Q)$, for

$$L_j = \begin{cases} M_{j+4} & n = 6, \\ C_{j+6} & n = 7, \\ B_{j+10} & n = 8. \end{cases} \quad (5.5)$$

Remark 5.1. We will construct a geometric model \mathbf{V} for $\mathcal{D}_\infty(Q)/[2]$ of a type E_n quiver Q consisting of the following data:

- \mathbf{V} is an h_Q gon whose edges are given by some central charge of the indecomposable objects in the far-end τ -orbit of $\mathcal{D}_\infty(Q)$.
- its ice/fire core is an $h_Q/2$ gon whose edges are given by the (same) central charge of half of the indecomposable objects in the mid-end τ -orbit of $\mathcal{D}_\infty(Q)$.

Such an h_Q -gon is stable if and only if both the ice and fire cores are inside the length- $(n-3)$ diagonal-gon of \mathbf{V} . Moreover, the key of realizing the root category is knowing the central charges of objects in the boundary τ -orbits, i.e. we will prove:

$$\begin{cases} Z(P_1^{j-1}) = Z(C_j) = W_{j-1}W_{j+1} \\ Z(P_3^{j-1}) = Z(M_j) = V_{j-2}W_j = W_{j+1}V_{j+n-4}, \\ Z(P_n^{j-1}) = Z(B_j) = V_{j-1}V_j. \end{cases} \quad (5.6)$$

in the following sections. We will also show that the edges of the ice/fire cores are given by

$$w_{j+2} := z_j + z_{j+n-2}, \quad j \in \mathbb{Z}_h \quad (5.7)$$

for $h = h_Q$.

Furthermore, the usual types (A and D) can also be thought as a degeneration of the model above, in the sense that:

- for $Q = D_n$, the ice/fire core shrinks to the puncture B_\pm .
- for $Q = A_n$, the ice/fire core vanishes.

6. ToSt OF EXCEPTIONAL TYPE E_6

6.1. The h -gon of type E_6 .

Definition 6.1. An h -gon \mathbf{V} of type E_6 is a 12-gon satisfying (1.1), i.e. the equations in Figure 6.

A direct calculation shows that:

Lemma 6.2. *The set of 7 equations (1.1) has rank 6. So the space of 12-gons of type E_6 has complex dimension $12 - 6 = 6$.*

By (1.1), we have the following hexagon relations.

$$z_j + z_{j+1} + z_{j+2} + z_{j+6} + z_{j+7} + z_{j+8} = 0, \quad \forall j \in \mathbb{Z}_{12}. \quad (6.1)$$

Construction 6.3. Using the triangle relations in (1.1), we can draw triangles

$$\mathbf{T}_j := V_{j-1}V_jW_j \quad \forall j \in \mathbb{Z}_{12}$$

with edges


$$V_{j-1}V_j = z_j, \quad V_jW_j = z_{j+4}, \quad W_jV_{j-1} = z_{j+8}.$$

Note that we have 4 sets of 3 parallel triangles, see Figure 6,

$$\{\mathbf{T}_{j+4k} \mid k = 0, 1, 2\} \quad \text{for } j \in \mathbb{Z}_4,$$

drawn in orange/blue/green/violet respectively. Moreover, the square relations in (1.1) correspond to the squares

$$S_j := V_{j-1}V_jW_{j+1}W_{j-1} \quad \forall j \in \mathbb{Z}_{12}$$



$$z_j + z_{j+4} + z_{j+8} = 0, \quad \forall j \in \mathbb{Z}_{12}$$

$$z_j - z_{j-3} + z_{j-6} - z_{j-9} = 0, \quad \forall j \in \mathbb{Z}_{12}$$




FIGURE 6. The 12-gon of type E_6 and AR quiver of $\mathcal{D}_\infty(E_6)$

with edges

$$V_{j-1}V_j = z_j, \quad V_jW_{j+1} = -z_{j-3}, \quad W_{j+1}W_{j-1} = z_{j-6}, \quad W_{j-1}V_{j-1} = -z_{j-9}.$$

Note that we have 3 sets of 4 parallel/anti-parallel squares

$$\{\mathbf{S}_{j+3k} \mid k = 0, 1, 2, 3\} \quad \text{for } j \in \mathbb{Z}_3,$$

see the squares in Figure 7.

Thus it is better to have two copies of \mathbf{V} (called *ice* and *fire*, cf. Figure 7):

\mathbf{V}_{ice} : the one with 6 triangles $\{\mathbf{T}_{2j} \mid j \in \mathbb{Z}_6\}$ and 6 squares $\{\mathbf{S}_{2j+1} \mid j \in \mathbb{Z}_6\}$. Then we have a hexagon *ice core* $\mathbf{V}_{\text{ice}}^{\odot}$ of \mathbf{V} with vertices W_{2j} .

\mathbf{V}_{fire} : the one with 6 triangles $\{\mathbf{T}_{2j+1} \mid j \in \mathbb{Z}_6\}$ and 6 squares $\{\mathbf{S}_{2j} \mid j \in \mathbb{Z}_6\}$.

Then we have a hexagon *fire core* $\mathbf{V}_{\text{fire}}^{\odot}$ of \mathbf{V} with vertices W_{2j+1} .

Set w_j as (5.7) for $n = 6$ and $h = 12$. By the triangle relation in (1.1), we have $w_{j+2} = -z_{j+8}$.

Remark 6.4. An interesting feature is that both ice and fire configurations induce planar tiling patterns, see Figure 17 in Appendix A. In other words, a 12-gon of type E_6 is equivalent to ‘A Tiling of Ice and Fire’ as shown there.

6.2. Geometric model for the root category of type E_6 .

Set-up 6.5. Take the orientation in (5.1) with $n = 6$. Note that 1/3/6 are the mid-end/near-end/far-end vertices in this case. The lower picture in Figure 6 is part of $\text{AR}\mathcal{D}_{\infty}(E_6)$ and the labelling is given by

1st- τ -orbit: $j + 6/j + 5$ denotes $C_j := P_1^{j-1}$ (drawn in green).

2nd- τ -orbit: $j + 7/j + 5$ denotes P_2^{j-1} .

3rd- τ -orbit: Let $M_j := P_3^{j-1}$ (drawn in yellow).




FIGURE 7. Configurations Ice and Fire

- 4th- τ -orbit:** $j - 1/j + 2$ denotes P_4^{j-1} .
5th- τ -orbit: $j - 1/j + 1$ denotes P_5^{j-1} .
6th- τ -orbit: $j - 1/j$ denotes $B_j := P_6^{j-1}$ (drawn in blue).

for $1 \leq j \leq 6$.

Given any central charge $Z: K\mathcal{D}_\infty(E_6) \rightarrow \mathbb{C}$. Let \mathbf{V}_Z be the far-end 12-gon of $\mathcal{D}_\infty(E_6)$ with edges $z_j = Z(B_j)$ for $1 \leq j \leq 12$.

Lemma 6.6. *The set $\{[B_j] \mid 1 \leq j \leq 6\}$ spans $K\mathcal{D}_\infty(E_6)$ (and in fact is a basis). Moreover, \mathbf{V}_Z is a 12-gon of type E_6 and its ice/fire core is formed by (the odd/even) half of*

$$\{Z(C_j) \mid 1 \leq j \leq 12\}$$

for $Z(C_j) = w_j$.

A coincidence in this case is $w_j = -z_{j+6}$ as mentioned above.

Proof. The first statement can be checked directly, e.g. via dimension vectors.

Now we will show that all z_j satisfy (1.1). On one hand, by (5.3) for $n = 6$ we have

$$[B_j] + [B_{j+4}] = [C_{j+2}]$$

and thus $Z(C_{j+2}) = z_j + z_{j+4} = w_{j+2}$ by (5.7). Due to $B_{j+8} = C_{j+2}[1]$, we have

$$[B_j] + [B_{j+4}] + [B_{j+8}] = 0.$$

Thus the triangle relations in (1.1) holds. On the other hand, triangles in the second row/column of (5.4) for $n = 6$ implies that

$$[B_j] + [C_{j+3}] = [M_{j+1}] = [C_j] + [B_{j+3}].$$

Noticing that $C_{j+6} = B_j[1]$ and $B_{j+3} = B_{j-9}[2]$, the above equation becomes

$$[B_j] - [B_{j-3}] = -[B_{j-6}] + [B_{j-9}],$$

again using. Thus the square relations in (1.1) holds.

In particular, we see that the edges of the ice/fire core are also w_j 's. \square

In this subsection, we will construct a central charge from a 12-gon \mathbf{V} of type E_6 . Recall that \mathbf{V} has edges $z_j = V_{j-1}V_j$, $j \in \mathbb{Z}_{12}$, as shown in the upper picture of Figure 6. Moreover, we have the associated ice/fire configuration/core according to Construction 6.3.

Theorem 6.7. *A 12-gon \mathbf{V} of type E_6 is a geometric model for the root category $\mathcal{D}_\infty(E_6)/[2]$ in the sense that by setting (for $n = 6$)*

$$\begin{cases} Z(P_1^j) = W_j W_{j+2} \\ Z(P_2^j) = V_{j-1} W_{j+2} = W_{j+1} V_{j+n-3} \\ Z(P_3^j) = V_{j-1} W_{j+1} = W_{j+2} V_{j+n-3}, \\ Z(P_i^j) = V_j V_{j+1+(n-i)}, \end{cases} \quad 4 \leq i \leq n, \quad (6.2)$$

we obtain a central charge $Z: K\mathcal{D}_\infty(E_6) \rightarrow \mathbb{C}$.

In particular, (5.6) holds (for mid-end/near-end/far-end τ -orbits). Also, central charges of objects in the mid-end τ -orbits are given by the edges of ice/fire core of \mathbf{V} and central charges of objects in the i^{th} - τ -orbits are given by length- $(7-i)$ diagonals of \mathbf{V} for $4 \leq i \leq n$.

Remark 6.8. In fact, there are several ways to realize the indecomposable objects in the τ -orbits. For instance, for any object $E \in \text{Ind mod } \mathbf{k}E_6$ that is not in the 3^{rd} - τ -orbit, it admits a labeling j/k as in Set-up 6.5, which is also the labelling corresponding to its image under the central charge, namely

$$Z(E) = V_j V_k \quad \text{and} \quad Z(E[1]) = V_k V_j;$$

for each M_j in the 3^{rd} - τ -orbits, it can be realized in two more ways:

$$W_{j+h/2} V_{j-2+h/2} = V_{j-2} W_j = W_{j+1} V_{j+n-4} = V_{j+n-4+h/2} W_{j+1+h/2} \quad (6.3)$$

for $h = 12$. See the thick black line segments in Figure 8 for $j = 2$.

In type E_6 , the far-end and mid-end τ -orbits are shift of each other which enable us to realize the objects in many ways. However, in types E_7 and E_8 , this is not the case. Therefore, we write the statement in the above theorem of the form that can be generalized to types E_7 and E_8 .

Proof of Theorem 6.7. The proof follows the same way as in the proof of Theorem 4.5. One has to show that the central charge Z preserves the mesh relations of $\text{AR } \mathcal{D}_\infty(E_6)/[2]$.




FIGURE 8. The triangles/squares for E_6

Here we only point out the reason that $Z(M_j)$ can be realized as in (6.3) is due to the triangles in the second row/column of (5.4) and Construction 6.3. The remaining calculations are left to the reader. \square

6.3. Stability of 12-gon for E_6 . Suppose that \mathbf{V} is a positively convex 12-gon of type E_6 . We have the following observations.

- Its ice/fire core is convex. This follows from the fact that the edges of ice/fire core are exactly half of the edges of the 12-gon induced by the mid-end τ -orbit.
- Any length-3 diagonal $V_{j-1}V_{j+2}$ is a long diagonal of a narrow hexagon

$$\mathbf{H}_j = V_{j-1}V_jV_{j+1}V_{j+2}W_{j+2}W_j$$

that corresponds to (6.1). Note that \mathbf{H}_j and \mathbf{H}_{j+6} are related by a translation $V_{j+2}V_{j+5}$. Hence they inherit the convexity of \mathbf{V} .

- See the left picture in Figure 7, the configuration ice contains 6 narrow hexagons $\{\mathbf{H}_{2j}\}$ and the corresponding length-3 diagonals (blue dashed lines in the left picture) bound another hexagon, we call the *ice-boundary*, which contains the ice core automatically due to the convexity of the narrow hexagons.
- Dually, see the right picture in Figure 7, the configuration fire contains 6 narrow hexagons $\{\mathbf{H}_{2j+1}\}$ and the corresponding length-3 diagonals (red dashed lines in the right picture) bound another hexagon, we call the *fire-boundary*, which also contains the fire core automatically.

Definition 6.9. A 12-gon \mathbf{V} of type E_6 is *stable* if it is positively convex and both its ice and fire cores are inside the level-3 diagonal-gon.

Denote by $\text{Stgon}(E_6)$ the moduli space of stable 12-gon of type E_6 up to translation. By Lemma 6.2, the complex dimension of $\text{Stgon}(E_6)$ is 6.

Using the above observations, one can simplify the stability condition of 12-gons a bit as in the lemma below.

Lemma 6.10. *A positively convex 12-gon \mathbf{V} of type E_6 is stable if and only if its ice/fire core is inside its fire/ice boundary respectively.*

Proof. As We know, the positive convexity of \mathbf{V} implies the convexity of the narrow hexagons. Then, the ice/fire core is automatically inside ice/fire boundary respectively. Moreover, the level-3 diagonal-gon is the intersection of the ice and fire boundaries. Thus, the stability condition of \mathbf{V} can be simplified to the condition stated in the lemma, provided the positive convexity of \mathbf{V} . \square

Now we proceed to show that total stability of σ deduces stability of its far-end 12-gon.

Proposition 6.11. *If $\sigma = (Z, \mathcal{P}) \in \text{ToSt}(E_6)$, then its far-end 12-gon is a stable 12-gon of type E_6 .*

Proof. By Proposition 3.6, the far-end 12-gon \mathbf{V} is positively convex. Since the equation (1.1) holds due to Lemma 6.6, we can follow Construction 6.3 to obtain its ice/fire core with vertices $W_j, j \in \mathbb{Z}_{12}$ via triangles (cf. Figure 6). Then we only need to show that

each W_j are bounded by length-3-diagonals of \mathbf{V} . By Lemma 6.10, it suffices to check that W_{2j} is bounded by the fire boundary and W_{2j+1} is bounded by the ice boundary. By the observation of the narrow hexagons, it is equivalent to prove that each W_{j+1} is on the left side of the diagonal $V_{j-1}V_{j+2}$.

Consider the triangle $V_{j-1}W_{j+1}V_{j+2}$ with edges

$$V_{j-1}W_{j+1} = Z(P_3^j), \quad W_{j+1}V_{j+2} = Z(P_3^{j-1}), \quad V_{j-1}V_{j+2} = Z(P_4^{j-1}).$$

These edges correspond to the central charges of the terms in the following AR triangle:

$$P_3^{j-1} \rightarrow P_4^{j-1} \rightarrow P_3^j \rightarrow P_3^{j-1}[1]. \quad (6.4)$$

See the yellow triangle (and its mirror in lighter yellow) in Figure 8 for $j = 1$. And the total stability implies

$$\phi_\sigma(P_3^{j-1}) < \phi_\sigma(P_4^{j-1}) < \phi_\sigma(P_3^j) < \phi_\sigma(P_3^{j-1}) + 1$$

By taking arg in $[\pi \cdot \phi_\sigma(P_3^{j-1}), \pi \cdot \phi_\sigma(P_3^{j-1}) + 2\pi)$, we conclude that each W_{j+1} is indeed on the left side of $Z(P_4^{j-1}) = V_{j-1}V_{j+2}$. \square

6.4. From stable 12-gon to total stability conditions for E_6 . Finally, we would like to construct a slicing \mathcal{P} from a stable 12-gon \mathbf{V} , which is compatible with the central charge defined in Theorem 6.7 and the shift [1]. Such a construction is basically taking arg in appropriate length 2π intervals for central charges of each indecomposables. It follows exactly the same line of work as in Section 4.3. Thus, we have a similar statement in type E_6 .

Theorem 6.12. *There is a natural isomorphism*

$$Z_h: \text{ToSt}(E_6)/[2] \rightarrow \text{Stgon}(E_6),$$

sending a total stability condition σ to the far-end 12-gon.

7. TOSt OF EXCEPTIONAL TYPE E_7

7.1. The h -gon of type E_7 .

Definition 7.1. An h -gon \mathbf{V} of type E_7 is a symmetric 18-gon satisfying (1.2), i.e. the equation in Figure (9).

A direct calculation shows that:

Lemma 7.2. *After setting $z_{j+9} = -z_j$ for $j \in \mathbb{Z}_{18}$, the set of 3 equations (1.2) has rank 2. So the space of 18-gons of type E_7 has complex dimension $9 - 2 = 7$.*

Construction 7.3. Using the hexagon relations in (1.2), we can draw hexagons

$$\mathbf{L}_{2j}: = V_{2j-1}V_{2j}V_{2j+1}W_{2j+1}U_{2j}W_{2j-1}$$

with edges

$$\begin{aligned} V_{2j-1}V_{2j} &= z_{2j}, & V_{2j}V_{2j+1} &= z_{2j+1}, & V_{2j+1}W_{2j+1} &= z_{2j+6}, \\ W_{2j+1}U_{2j} &= z_{2j+7}, & U_{2j}W_{2j-1} &= z_{2j+12}, & W_{2j-1}V_{2j-1} &= z_{2j+13} \end{aligned}$$

as shown in the upper picture of Figure 9. Note that we have 3 sets of 3 parallel hexagons

$$\{\mathbf{L}_{2j+6k} \mid k = 0, 1, 2\} \quad \text{for } j \in \mathbb{Z}_3,$$

drawn in blue/green/orange respectively.




FIGURE 9. The (regular) 18-gon of type E_7 and AR-quiver of $\mathcal{D}_\infty(E_7)$

Set w_j as in (5.7) for $n = 7$ and $h = 18$. As $z_{j+9} = -z_j$, we also have $w_{j+9} = -w_j$. Moreover, each hexagon decomposes into 3 triangles corresponding to the above equation and 1 triangle corresponding to the relation

$$w_j + w_{j+6} + w_{j+12} = 0.$$

So there is a 9-gon $\mathbf{V}_{\text{ice}}^{\odot}$, called the *ice core* of \mathbf{V} , with vertices $\{W_{2j+1} \mid j \in \mathbb{Z}_9\}$ and edges

$$w_{2j} = W_{2j-1}W_{2j+1}. \quad (7.1)$$

Remark 7.4. We take nine \mathbf{L}_{2j} with even indices to obtain $\mathbf{V}_{\text{ice}}^{\odot}$. One can also take the other nine \mathbf{L}_{2j+1} with odd indices to obtain another 9-gon $\mathbf{V}_{\text{fire}}^{\odot}$, called the *fire core* of \mathbf{V} , with vertices $\{W_{2j} \mid j \in \mathbb{Z}_9\}$ and edges $w_{2j+1} = W_{2j}W_{2j+2}$. However, since \mathbf{V} is symmetric, these two cores are also symmetric to each other.

7.2. Geometric model for the root category of type E_7 .

Set-up 7.5. Take the orientation in (5.1) with $n = 7$. Note that 1/3/7 are the mid-end/near-end/far-end vertices in this case. The lower picture in Figure 9 is part of $\text{AR } \mathcal{D}_{\infty}(E_7)$ and for simplicity, we only label the objects in the boundary τ -orbits, i.e. $C_j/M_j/B_j$ in (5.2).

Given any central charge $Z: K\mathcal{D}_{\infty}(E_7) \rightarrow \mathbb{C}$. Let \mathbf{V}_Z be the far-end 18-gon of $\mathcal{D}_{\infty}(E_7)$ with edges $z_j = Z(B_j)$ for $1 \leq j \leq 18$.

Lemma 7.6. *The set $\{[B_j] \mid 1 \leq j \leq 9\}$ spans $K\mathcal{D}_{\infty}(E_7)$. Moreover, \mathbf{V}_Z is a 18-gon of type E_7 and its ice/fire core is formed by (the odd/even) half of*

$$\{Z(C_j) \mid 1 \leq j \leq 18\}$$

for $Z(C_j) = w_j$.

Proof. The first statement can be checked directly, e.g. via dimension vectors.

Now we will show that all z_j satisfy (1.2). By (5.3) for $n = 7$ we have

$$[C_{j+2}] = [B_j] + [B_{j+5}]$$

and thus $Z(C_{j+2}) = z_j + z_{j+5} = w_{j+2}$ as we have the triangle relation (5.7).

Moreover, triangles in the second row/column of (5.4) for $n = 7$ implies that

$$[B_j] + [C_{j+3}] = [M_{j+1}] = [C_j] + [B_{j+4}]$$

and thus $z_j + w_{j+3} = w_j + z_{j+4}$. Substitute (5.7) to kill w_j , we obtain

$$z_j + (z_{j+1} + z_{j+6}) = (z_{j-2} + z_{j+3}) + z_{j+4}.$$

Noticing that $z_{k+9} = -z_k$, the above equation becomes (1.2).

Thanks to (5.7), the last statement is clear. \square

Remark 7.7. The triangle relation (7.1) corresponds to the triangle in the third row of (5.4) as $L_j = C_{j+6}$ in this case.

Now we describe a geometric model in E_7 case.

Theorem 7.8. *An 18-gon \mathbf{V} of type E_7 is a geometric model for the root category $\mathcal{D}_\infty(E_7)/[2]$ in the sense that by setting (6.2) for $n = 7$, we obtain a central charge $Z: K\mathcal{D}_\infty(E_7) \rightarrow \mathbb{C}$.*

As in type E_6 , for each M_j in the $3^{\text{rd}}\text{-}\tau$ -orbits, it can be realized by (6.3) for $h = 18$. For instance, the (edges of the) yellow triangle in the top picture of Figure 9 corresponds to the (central charges of) triangle

$$M_2 \rightarrow X \rightarrow M_3 \rightarrow M_2[1] \quad (7.2)$$

in $\mathcal{D}_\infty(E_7)$, similar to the yellow triangle in the case of E_6 .

7.3. Stability of 18-gon for E_7 .

Definition 7.9. An h -gon \mathbf{V} of type E_7 is *stable* if it is positively convex and its ice/fire core is inside the level-4 diagonal-gon.

Note that the positive convexity of \mathbf{V} will be inherited by its ice and fire cores as in the E_6 case. Denote by $\text{Stgon}(E_7)$ the moduli space of stable 18-gon of type E_7 up to translation. By Lemma 7.2, the complex dimension of $\text{Stgon}(E_7)$ is 7.

Proposition 7.10. *If $\sigma = (Z, \mathcal{P}) \in \text{ToSt}(E_7)$, then its far-end 18-gon is a stable 18-gon of type E_7 .*

Proof. To prove the statement, one needs to use the AR triangle (7.2) similar as in type E_6 , whose central charges of its terms form the yellow triangle in Figure 9. Then the rest argument follows the same way as in Proposition 6.11. \square

Theorem 7.11. *There is a natural isomorphism*

$$Z_h: \text{ToSt}(E_7)/[2] \rightarrow \text{Stgon}(E_7).$$

sending a total stability condition σ to the far-end 18-gon.

8. TOSt OF EXCEPTIONAL TYPE E_8

8.1. The h -gon of type E_8 .


Definition 8.1. An h -gon \mathbf{V} of type E_8 is a symmetric 30-gon satisfying (1.3), i.e. the equation in Figure (10).

A direct calculation shows that:

Lemma 8.2. *After setting $z_{j+15} = -z_j$ for $j \in \mathbb{Z}_{30}$, the set of 8 equations (1.3) has rank 7. So the space of 30-gons of type E_8 has complex dimension $15 - 7 = 8$.*

Construction 8.3. Using the triangle/pentagon relations in (1.3), we can draw pentagons

$$\mathbf{P}_{2j}: = V_{2j}V_{2j+1}W_{2j+1}U_{2j}W_{2j-1} \quad \forall j \in \mathbb{Z}_{15}$$

FIGURE 10. The (regular) 30-gon of type E_8

with edges

$$\begin{cases} V_{2j}V_{2j+1} = z_{2j+1}, & V_{2j+1}W_{2j+1} = z_{2j+7}, & W_{2j+1}U_{2j} = z_{2j+13}, \\ U_{2j}W_{2j-1} = z_{2j+19}, & W_{2j-1}V_{2j} = z_{2j+25} \end{cases}$$

and triangles

$$\mathbf{T}_{2j} := V_{2j-1}V_{2j}W_{2j-1} \quad \forall j \in \mathbb{Z}_{15}$$

with edges

$$V_{2j-1}V_{2j} = z_{2j}, \quad V_{2j}W_{2j-1} = z_{2j+10}, \quad W_{2j-1}V_{2j-1} = z_{2j+20},$$

as shown in Figure 10. Note that we have 3 sets of 5 parallel pentagons

$$\{\mathbf{P}_{2j+6k} \mid k = 0, 1, 2, 3, 4\}, \quad \text{for } j \in \mathbb{Z}_3$$

drawn in orange/green/blue respectively and 5 sets of 3 parallel triangles

$$\{\mathbf{T}_{2j+10k} \mid k = 0, 1, 2\}, \quad \text{for } j \in \mathbb{Z}_5$$

drawn in violet with different opacity respectively.

So there is a 15-gon $\mathbf{V}_{\text{ice}}^{\odot}$, called the *ice core* of \mathbf{V} with vertices $\{W_{2j-1} \mid j \in \mathbb{Z}_{15}\}$ and edges $w_{2j} = W_{2j-1}W_{2j+1}$.

Remark 8.4. We take fifteen \mathbf{P}_{2j} and fifteen \mathbf{T}_{2j} with even indices to obtain $\mathbf{V}_{\text{ice}}^{\odot}$. One can also take the other fifteen \mathbf{P}_{2j+1} and fifteen \mathbf{T}_{2j+1} with odd indices to obtain another 15-gon $\mathbf{V}_{\text{fire}}^{\odot}$, called the *fire core* of \mathbf{V} , with vertices $\{W_{2j} \mid j \in \mathbb{Z}_{15}\}$ and edges $w_{2j+1} = W_{2j}W_{2j+2}$. However, since \mathbf{V} is symmetric, these two cores are also symmetric to each other.

Note that as $W_{j+1}W_{j+3} = W_{j+1}U_{j+2} + U_{j+2}W_{j+3}$, we have formula (5.7) for $n = 8$ and $h = 30$. As $z_{j+15} = -z_j$, we also have $w_{j+15} = -w_j$.

8.2. Geometric model for the root category of type E_8 .

Set-up 8.5. Take the orientation in (5.1) with $n = 8$. Note that $1/3/8$ are the mid-end/near-end/far-end vertices in this case. The picture in Figure 11 is part of $\text{AR } \mathcal{D}_{\infty}(E_8)$ and for simplicity, we only label the objects in the boundary τ -orbits, i.e. $C_j/M_j/B_j$ in (5.2).

Given any central charge $Z: K\mathcal{D}_{\infty}(E_8) \rightarrow \mathbb{C}$. Let \mathbf{V}_Z be the far-end 30-gon of $\mathcal{D}_{\infty}(E_8)$ with edges $z_j = Z(B_j)$ for $1 \leq j \leq 30$.

Lemma 8.6. *The set $\{[B_j] \mid 1 \leq j \leq 15\}$ spans $K\mathcal{D}_{\infty}(E_8)$. Moreover, \mathbf{V}_Z is a 30-gon of type E_8 and its ice/fire core is formed by (the odd/even) half of*

$$\{Z(C_j) \mid 1 \leq j \leq 30\}$$

for $Z(C_j) = w_j$.

Proof. The first statement can be checked directly, e.g. via dimension vectors.

Now we will show that all z_j satisfy (1.3). By (5.3) for $n = 8$ we have

$$[C_{j+2}] = [B_j] + [B_{j+6}]$$

and thus $Z(C_{j+2}) = z_j + z_{j+6} = w_{j+2}$ as we have the triangle relation (5.7).

Moreover, triangles in the third row/column of (5.4) for $n = 8$ implies that


$$\begin{cases} z_j - z_{j+5} + z_{j+10} = 0, \\ w_j - w_{j+3} + z_{j+10} = 0, \end{cases} \quad \forall j \in \mathbb{Z}_{30} \quad (8.1)$$

as $L_j = B_{j+10}$ in this case. Noticing $-z_{j+5} = z_{j+20}$, the first equation of (8.1) becomes the triangle relation in (1.3). Substitute (5.7) to kill w 's in the second equation of (8.1), we have

$$z_{j-2} + z_{j+4} - z_{j+1} - z_{j+7} + z_{j+10} = 0, \quad \forall j \in \mathbb{Z}_{30}.$$

Noticing $z_k = -z_{k+15}$, the above equation becomes the pentagon relation in (1.3). \square

Now we describe a geometric model in type E_8 .

FIGURE 11. The AR-quiver of $\mathcal{D}_\infty(E_8)$

Theorem 8.7. A 30-gon \mathbf{V} of type E_8 is a geometric model for the root category $\mathcal{D}_\infty(E_8)/[2]$ in the sense that by setting (6.2) for $n = 8$ we obtain a central charge $Z: K\mathcal{D}_\infty(E_8) \rightarrow \mathbb{C}$.

As in types E_6 and E_7 , for each M_j in the $3^{\text{rd}}\text{-}\tau$ -orbits, it can be realized by (6.3) for $h = 30$. For instance, the (edges of the) yellow triangle in the top picture of Figure 10 corresponds to the (central charges of) triangle $M_2 \rightarrow X \rightarrow M_3 \rightarrow M_2[1]$ in $\mathcal{D}_\infty(E_8)$.

8.3. Stability of 30-gon for E_8 .

Definition 8.8. A 30-gon \mathbf{V} of type E_8 is *stable* if it is positively convex and its ice/fire core is inside the level-5 diagonal-gon.

Denote by $\text{Stgon}(E_8)$ the moduli space of stable 30-gon of type E_8 up to translation. By Lemma 8.2, the complex dimension of $\text{Stgon}(E_8)$ is 8.

As above in type E_7 , we have the following proposition and theorem for E_8 .

Proposition 8.9. If $\sigma = (Z, \mathcal{P}) \in \text{ToSt}(E_8)$, then its far-end 30-gon is a stable 30-gon of type E_8 .

Theorem 8.10. *There is a natural isomorphism*

$$Z_h: \text{ToSt}(E_8)/[2] \rightarrow \text{Stgon}(E_8).$$

sending a total stability condition σ to the far-end 30-gon.

Remark 8.11. An interesting numerical observation is that the triangle/square relations in type E_6 correspond to x^4 and y^3 , and triangle/pentagon relations in type E_6 correspond to x^5 and y^3 , respectively, in the corresponding simple (surface) singularity

$$f(x, y, z) = \begin{cases} x^{n+1} + y^2 + z^2, & Q = A_n (n \geq 1); \\ x^{n-1} + xy^2 + z^2, & Q = D_n, (n \geq 4); \\ x^4 + y^3 + z^2, & Q = E_6; \\ x^3y + y^3 + z^2, & Q = E_7; \\ x^5 + y^3 + z^2, & Q = E_8. \end{cases}$$

These equations are used to define $\mathcal{D}_\infty(Q)$ via (graded) matrix factorizations in [KST]. The correspondence in type E_7 is not as nice as the other two.

9. NON SIMPLY-LACED TYPES VIA FOLDING

The non-simply-laced case is obtained by folding the corresponding simply-laced case, see Section 2.3. The derived category/stability conditions for R^l are the ι -stable part of the one for Q (cf. [CQ] for more details). Thus, the stable h_Q -gons for R^l are just ι -stable h -gons, i.e. satisfying extra symmetry. More precisely, we have the following

- B_n : It is obtained by folding the corresponding quiver of type D_{n+1} . Thus the extra symmetry condition on the corresponding doubly punctured $2n$ -gon is the two punctures coincide (and thus at the geometric center of the $2n$ -gon).
- C_n : It is obtained by folding the corresponding quiver of type A_{2n-1} . Thus the extra symmetry condition on the corresponding $2n$ -gon is central symmetry.
- F_4 : It is obtained by folding the corresponding quiver of type E_6 . Thus the extra symmetry condition on the corresponding 12-gon is central symmetry.
- G_2 : It is obtained by folding the corresponding quiver of type D_4 . Thus the extra condition is: concision of the punctures, central symmetry and (1.4).

So Theorem 1 also holds in these cases.

Example 9.1. In Figure 12, we show that how to deform the central charge and associated far-end stable h_Q -gon of type D_4 into the ones of type B_3 and of type G_2 , respectively.

From D_4 to B_3 , it requires the (orange/violet) central charges of objects in the two boundary (besides the chosen far-end) τ -orbits coincides correspondingly. Equivalently, the two punctures coincide as mentioned above.

From D_4 to G_2 , it requires the (orange/violet/blue) central charges of objects in all boundary τ -orbits coincides correspondingly. Equivalently, the long diagonals of the $h_Q = 6$ -gon intersect at its geometric center that divide it into six parallel/anti-parallel triangles.




FIGURE 12. Deforming stable h -gon and central charge of D_4 to B_3 to G_2

REFERENCES

- [AB] C. Amiot, and T. Brüstle, Derived equivalences between skew-gentle algebras, [arXiv:1912.04367](#)
- [AP] C. Amiot, and P-G. Plamondon, The cluster category of a surface with punctures via group actions, *Adv. Math.* **389** (2021) Paper No. 107884, 63 pp.
- [BGMS] E. Barnard, E. Gunawan, E. Meehan and R. Schiffler, Cambrian combinatorics on quiver representations (type A_n), [arXiv:1912.02840](#)
- [BM] K. Baur and R.-Marsh, Categorification of a frieze pattern determinant, [arXiv:1008.5329](#)
- [B1] T. Bridgeland, Stability conditions on triangulated categories, *Ann. Math.* **166** (2007) 317–345. ([arXiv:0212237](#))
- [B2] T. Bridgeland, Stability conditions and Kleinian singularities, *Int. Math. Res. Not.* **21** (2009) 4142–4157. ([arXiv:0508257](#))
- [BQS] T. Bridgeland, Y. Qiu, and T. Sutherland, Stability conditions and the A_2 quiver, [arXiv:1406.2566](#).
- [BS] T. Bridgeland and I. Smith, Quadratic differentials as stability conditions, *Publ. Math. de l’IHÉS* **121** (2015) 155–278. ([arXiv:1302.7030](#))
- [BQ] T. Brüstle and Y. Qiu, Tagged main class group: Auslander-Reiten translations, *Math. Zeit.* **279** (2015) 1103–1120. ([arXiv:1212.0007](#))
- [CQ] W. Chang and Y. Qiu, Folding quivers and numerical stability conditions, [arXiv:1210.0243](#)
- [CQZ] W. Chang, Y. Qiu and X. Zhang, Geometric model for module categories of Dynkin quivers via hearts of total stability conditions, in preparation.
- [FLLQ] Y. Fan, C. Li, W. Liu and Y. Qiu, Contractibility of space of stability conditions on the projective plane via global dimension function, [arXiv:2001.11984](#)

- [FST] S. Fomin, M. Shapiro and D. Thurston, Cluster algebras and triangulated surfaces, part I: Cluster complexes, *Acta Mathematica* **201** (2008) 83–146.
- [HKK] F. Haiden, L. Katzarkov and M. Kontsevich, Stability in Fukaya categories of surfaces, *Publ. Math. de l’IHÉS* **126** (2017) 247–318. ([arXiv:1409.8611](#))
- [IQ1] A. Ikeda and Y. Qiu, q -Stability conditions on Calabi-Yau- \mathbb{X} categories and twisted periods, [arXiv:1807.00469](#).
- [IQ2] A. Ikeda and Y. Qiu, q -Stability conditions via q -quadratic differentials, [arXiv:1812.00010](#).
- [KST] H. Kajiura, K. Saito and A. Takahashi, Matrix factorizations and representations of quivers II: Type ADE case, *Adv. Math.* **211** (2007) 327–362. ([arxiv:0511155](#))
- [K] B. Keller, On cluster theory and quantum dilogarithm. [arXiv:1102.4148v4](#).
- [KOT] K. Kikuta, G. Ouchi and A. Takahashi, Serre dimension and stability conditions, [arXiv:1907.10981](#).
- [Ki] R. Kinser, Total stability functions for type A quivers, [arXiv:2002.12396](#)
- [L] H. Lenzing, Coxeter Transformations associated with Finite Dimensional Algebras, *Computational methods for representations of groups and algebras* (1999) 287–308.
- [Q1] Y. Qiu, Stability conditions and quantum dilogarithm identities for Dynkin quivers, *Adv. Math.* **269** (2015) 220–264. ([arXiv:1111.1010](#))
- [Q2] Y. Qiu, Global dimension function, Gepner equations and q -stability conditions, [arXiv:1807.00010](#).
- [Q3] Y. Qiu, Contractible flow of stability conditions via global dimension function, [arXiv:2008.00282](#).
- [QW] Y. Qiu and J. Woolf, Contractible stability spaces and faithful braid group actions, *Geom. Topol.* **22** (2018) 3701–3760. ([arXiv:1407.5986](#))
- [QZ] Y. Qiu and Y. Zhou, Cluster categories for marked surfaces: punctured case, *Compos. Math.* **153** (2017) 1779–1819. ([arXiv:1311.0010](#))
- [QZZ] Y. Qiu, C. Zhang and Y. Zhou, Two geometric models for graded skew-gentle algebras, in preparation.
- [R] M. Reineke, The Harder-Narasimhan system in quantum groups and cohomology of quiver moduli, *Invent. Math.* 152 (2003), no. 2, 349–368. ([arXiv:math/0204059v1](#))
- [S] R. Schiffler, A geometric model for cluster categories of type D_n , *J. Algebraic Combin.* **27** (2008) 1–21.

QY: YAU MATHEMATICAL SCIENCES CENTER AND DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, 100084 BEIJING, CHINA. & BEIJING INSTITUTE OF MATHEMATICAL SCIENCES AND APPLICATIONS, YANQI LAKE, BEIJING, CHINA

Email address: yu.qiu@bath.edu

ZX: BEIJING ADVANCED INNOVATION CENTER FOR IMAGING THEORY AND TECHNOLOGY, ACADEMY FOR MULTIDISCIPLINARY STUDIES, CAPITAL NORMAL UNIVERSITY, BEIJING 100048, CHINA

Email address: xiaoting.zhang09@hotmail.com

APPENDIX A. TIKZ ART GALLERY BY QY

In this appendix, we collection figures of stable h -gons at Gepner points that interact with the corresponding projection of the root systems in the Coxeter plans, as well as a tilting of Ice and Fire of type E_6 .




FIGURE 13. The (Coxeter projection of the) root system of type D_6 and the stable h -gon




FIGURE 14. The central charges, root system and stable h -gons of type E_6




FIGURE 15. The central charges, root system and stable h -gons of type E_7




FIGURE 16. The root system and (two) stable h -gons of type E_8

FIGURE 17. A Tiling of Ice and Fire

