

The Hydrogen-Poor Superluminous Supernovae from the Zwicky Transient Facility Phase-I Survey: I. Light Curves and Measurements

Z. H. CHEN,¹ LIN YAN,² T. KANGAS,³ R. LUNNAN,⁴ S. SCHULZE,⁵ J. SOLLERMAN,⁴ D. A. PERLEY,⁶ T.-W. CHEN,⁴ K. TAGGART,⁷ K. R. HINDS,⁶ A. GAL-YAM,⁸ X. F. WANG,^{1,9} I. ANDREONI,¹⁰ E. BELLM,¹¹ J. S. BLOOM,^{12,13} K. BURDGE,¹⁴ A. BURGOS,¹⁵ D. COOK,¹⁶ A. DAHIWALE,¹⁴ K. DE,¹⁰ R. DEKANY,² A. DUGAS,¹⁷ S. FREDERIK,¹⁸ C. FREMLING,¹⁴ M. GRAHAM,¹⁰ M. HANKINS,¹⁹ A. HO,²⁰ J. JENCSON,²¹ V. KARAMBELKAR,¹⁰ M. KASLIWAL,¹⁴ S. KULKARNI,¹⁴ R. LAHER,¹⁶ B. RUSHOLME,¹⁶ Y. SHARMA,¹⁴ F. TADDIA,²² L. TARTAGLIA,²³ B. P. THOMAS,²⁴ A. TZANIDAKIS,¹⁴ J. VAN ROESTEL,¹⁴ R. WALTER,¹⁰ Y. YANG,²⁵ Y. H. YAO,¹⁰ AND O. YARON⁸

¹Physics Department and Tsinghua Center for Astrophysics (THCA), Tsinghua University, Beijing, 100084, China

²Caltech Optical Observatories, California Institute of Technology, Pasadena, CA 91125, USA

³KTH Royal Institute of Technology, Stockholm, Sweden

⁴The Oskar Klein Centre, Department of Astronomy, Stockholm University, AlbaNova, SE-106 91 Stockholm, Sweden

⁵The Oskar Klein Centre, Department of Physics, Stockholm University, AlbaNova, SE-106 91 Stockholm, Sweden

⁶Astrophysics Research Institute, Liverpool John Moores University, 146 Brownlow Hill, Liverpool L3 5RF, UK

⁷Department of Astronomy and Astrophysics, University of California, Santa Cruz, CA 95064, USA

⁸Department of particle physics and astrophysics, Weizmann Institute of Science, 76100 Rehovot, Israel

⁹Beijing Planetarium, Beijing Academy of Sciences and Technology, Beijing, 100044, China

¹⁰Cahill Center for Astrophysics, California Institute of Technology, 1200 E. California Blvd. Pasadena, CA 91125, USA

¹¹DIRAC Institute, Department of Astronomy, University of Washington, 3910 15th Avenue NE, Seattle, WA 98195, USA

¹²Department of Astrophysics, University of California, Berkeley, CA 94720-3411, USA

¹³Lawrence Berkeley National Laboratory, 1 Cyclotron Road, MS 50B-4206, Berkeley, CA 94720, USA

¹⁴Division of Physics, Mathematics, and Astronomy, California Institute of Technology, Pasadena, CA 91125, USA

¹⁵Instituto de Astrofísica de Canarias, E-38 200 La Laguna, Tenerife, Spain

¹⁶IPAC, California Institute of Technology, 1200 E. California Blvd, Pasadena, CA 91125, USA

¹⁷Department of Physics and Astronomy, Watanabe 416, 2505 Correa Road, Honolulu, HI 96822, USA

¹⁸Department of Astronomy, University of Maryland, MD 20742-2421, USA

¹⁹Arkansas Tech University, Russellville, AR 72801, USA

²⁰Department of Astronomy and Miller Institute for Basic Research in Science, University of California, Berkeley, CA, USA

²¹Steward Observatory, University of Arizona, 933 North Cherry Avenue, Tucson, AZ 85721-0065, USA

²²Department of Physics and Astronomy, Aarhus University, Ny Munkegade 120, DK-8000 Aarhus C, Denmark

²³INAF-Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, I-35122 Padova, Italy

²⁴Department of Astronomy, University of Texas at Austin, 2515 Speedway, Stop C1400 Austin, TX 78712-1205, USA

²⁵Department of Astronomy, University of California, Berkeley, CA 94720-3411, USA

(Accepted Nov. 1, 2022)

Submitted to APJ

ABSTRACT

During the Zwicky Transient Facility (ZTF) Phase-I operation, 78 hydrogen-poor superluminous supernovae (SLSNe-I) were discovered in less than three years, making up the largest sample from a single survey. This paper (Paper I) presents the data, including the optical/ultraviolet light curves and classification spectra, while Paper II in this series will focus on the detailed analysis of the light curves and modeling. Our photometry is primarily taken by the ZTF in the g, r, i bands, and with additional data from other ground-based facilities and *Swift*. The events of our sample cover a redshift range of $z = 0.06 - 0.67$, with a median and 1σ error (16% and 84% percentiles) $z_{\text{med}} = 0.265^{+0.143}_{-0.135}$. The peak luminosity covers $-22.8 \text{ mag} \leq M_{g,\text{peak}} \leq -19.8 \text{ mag}$, with a median value of $-21.48^{+1.13}_{-0.61} \text{ mag}$. Their

light curves evolve slowly with the mean rest-frame rise time of $t_{\text{rise}} = 41.9 \pm 17.8$ days. The luminosity and time scale distributions suggest that low luminosity SLSNe-I with peak luminosity ~ -20 mag or extremely fast rising events (< 10 days) exist but are rare. We confirm previous findings that slowly rising SLSNe-I also tend to fade slowly. The rest-frame color and temperature evolution show large scatters, suggesting that the SLSN-I population may have diverse spectral energy distributions. The peak rest-frame color shows a moderate correlation with the peak absolute magnitude, *i.e.* brighter SLSNe-I tend to have bluer colors. With optical and ultraviolet photometry, we construct bolometric luminosity and derive a bolometric correction relation generally applicable for converting g, r -band photometry to bolometric luminosity for SLSNe-I.

Keywords: Stars: supernovae: general

1. INTRODUCTION

Superluminous supernovae (SLSNe) are a rare class of stellar explosions first discovered over fifteen years ago (*i.e.* SN 2005ap, Quimby et al. 2007). Their peak luminosities (10^{43-44} erg s $^{-1}$) are 10 – 100 times higher than those of normal Type Ia and core-collapse supernovae (SNe). Their light curves (LCs) usually evolve rather slowly, with rise times of $\sim 20 - 100$ days. The combination of these two features can not be explained by conventional SN models, *i.e.* standard radioactive decay (Kasen 2017). With the discovery of the first several SLSNe (SN 2005ap, SN 2006gy, SN 2007bi & SN 2008es), it was quickly recognized that, like normal SNe, SLSNe can be divided into two spectroscopic subclasses, one with hydrogen emission lines (SLSNe-II, Miller et al. 2009; Gezari et al. 2009; Inserra et al. 2018b) and the other without hydrogen (SLSNe-I, Quimby et al. 2007; Ofek et al. 2007; Gal-Yam et al. 2009; Smith et al. 2007; Gal-Yam 2012). In recent years, the subclass of H-poor but Helium-rich SLSNe (SLSNe-Ib) were first identified by Quimby et al. (2018), and later a sample by (Yan et al. 2020).

Three popular models have been proposed to explain the extraordinary radiative power of SLSNe. One involves energy injection from a central engine, such as the spin-down of a fast-rotating neutron star (magnetar, Kasen & Bildsten 2010; Woosley 2010). Alternatively, the interaction between the SN ejecta and dense circumstellar material (CSM) can efficiently convert kinetic energy into radiation (Chevalier & Irwin 2011; Chatzopoulos et al. 2013). Finally, some SLSNe could be powered by massive amounts of ^{56}Ni synthesised in a pair-instability supernova explosion (PISN) of low-metallicity stars with masses $> 140 M_{\odot}$ (Woosley et al. 2007; Kasen et al. 2011). It is commonly accepted that most SLSNe-II are analogous to Type IIn SNe (Schlegel 1990; Filippenko 1997), primarily powered by ejecta interactions with dense CSM (Ofek et al. 2007; Miller et al. 2010; Chevalier & Irwin 2011), while a small fraction show broad H α features with no signs of strong inter-

action in spectra, *e.g.* SN 2008es, SN 2013hx, PS15br (Miller et al. 2009; Gezari et al. 2009; Inserra et al. 2018b), but interactions are still likely required in these events (Kangas et al. 2022).

Between 2005 and 2009, a handful of SLSNe were discovered by several untargeted transient surveys which were not specifically targeting bright nearby galaxies. This small number of luminous events sparked a flurry of studies in both theory and observation of SLSNe. The next big advance in this field came between 2009 and 2016 when large area, untargeted transient surveys started operating. For example, the Palomar Transient Factory (PTF, Law et al. 2009), the Pan-STARRS1 Medium Deep Survey (PS1 MDS, Chambers et al. 2016), the Catalina Real-time Transient Survey (CRTS, Drake et al. 2009), the All-Sky Automated Survey for SuperNovae (ASAS-SN, Shappee 2014), the Dark Energy Survey (DES, The Dark Energy Survey Collaboration 2005) and the Gaia Photometric Science Alerts (Gaia, Hodgkin et al. 2021) made major contributions to the discoveries of several dozen SLSNe at both low ($z \sim 0.2$) and high redshift ($z \sim 1$) (Nicholl et al. 2015; De Cia et al. 2018; Quimby et al. 2018; Lunnan et al. 2018; Angus et al. 2019). However, with over ~ 90 discovered SLSNe-I by the end of 2017, many questions regarding their physical nature still remained unclear. For example, the SLSN volumetric rates are poorly constrained, with only estimates from small SLSN-I samples (Quimby et al. 2013; McCrum et al. 2015; Prajs et al. 2017; Frohmaier et al. 2021). Attempts to examine the statistical distributions, such as luminosity functions, were also quite limited due to small number statistics.

Assembling a large sample of low- z SLSNe with a well-defined survey volume and cadence is one of the goals of the Zwicky Transient Facility (ZTF, Graham et al. 2019; Bellm et al. 2019a; Masci et al. 2019). ZTF utilizes a 600-megapixel camera mounted on the Palomar Samuel Oschin 48 inch Schmidt telescope to reach a 47 deg 2 field of view (Dekany et al. 2020). ZTF can cover the full

Northern Sky in 3 days down to a 5σ limiting magnitude of 20.5 – 20.8 magnitude, which is about 3.5 and 0.5 magnitudes deeper than that of ASAS-SN (Shappee 2014), and the Asteroid Terrestrial-impact Last Alert System (ATLAS, Tonry et al. 2018), respectively. The ZTF survey offers several advantages for discovering rare transient events such as SLSNe. It is an untargeted, all sky, and moderately high cadence survey, probing large volumes with its large area coverage and deep sensitivity limits. Its well-defined survey strategy – area coverage and cadence – also makes it possible to quantify the survey efficiency.

ZTF conducted several surveys with different cadences (ranging from minutes to days) and area coverage during the phase-I operation (Bellm et al. 2019b). Among them, a particularly important one for extragalactic transient studies is the Northern Sky Public Survey. ZTF covered roughly the entire northern sky accessible from Palomar, corresponding to a total sky area of $\sim 23,675 \text{ deg}^2$. In every 3 days, each field was observed once in g band and once in r band, with an interval of at least 30 minutes between observations.

Between March 17, 2018 and October 31, 2020, ZTF Phase-I¹ discovered and spectroscopically confirmed 85 SLSNe-I, 6 SLSNe-I.5 (classified as SLSNe-I but showing H lines after the peak) and 61 SLSNe-II (defined as SNe II with peak magnitudes brighter than -20.0 mag). The number of SLSNe discovered by ZTF from 2018 to 2020 (about 60 per year) is roughly 5 – 7 times higher than that detected in any previous year. The SLSN-I sample will be the focus of a series of three papers. Paper I (this paper) presents the observational data and analysis of the overall observational properties. Paper II (Chen et al. 2022) discusses the LC modeling and analysis of the LC morphology. Paper III (Yan et al. in preparation) will present the derived SLSN-I volumetric rates and luminosity functions at $z \leq 0.7$. Several additional papers based on some individual SLSNe discovered during the ZTF Phase-I operation have been recently published. Lunnan et al. (2020) showcased the first four SLSNe-I discovered by ZTF during its science commissioning phase. Yan et al. (2020) presented the discovery of six He-rich SLSNe-I (SLSNe-Ib), revealing additional constraints on the progenitor mass-loss history.

This paper is organized as follows. Section 2 introduces the selection and classification of this sample. Sec-

tion 3 presents the photometry from ZTF and other facilities. Section 4 discusses our methodology with various photometric corrections and the calculations of peak absolute magnitudes. The measurements of time scales, colors, black-body temperatures and bolometric luminosities are presented in Section 5. Section 6 summarizes the conclusions. Throughout the paper, all magnitudes are in the AB system unless explicitly noted otherwise. We adopt a Λ CDM cosmology with $H_0 = 70.0 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.3$ and $\Omega_\Lambda = 0.7$.

2. THE SLSN-I SAMPLE FROM ZTF-I

During the phase-I survey, ZTF discovered 85 SLSNe-I. This paper focuses on 78 of these 85 events whose LCs have turned over from the peak by October 31, 2020, enabling better LC modeling. Of these 78 SLSNe-I, seven can be classified as He-rich SLSNe-Ib, including six published by Yan et al. (2020) and one by Tereran et al. (2020). For completeness, this sample paper also includes the four sources published in Lunnan et al. (2020). In addition, Schulze et al. (in preparation) will focus on an extremely slow and peculiar SLSN-I, ZTF18acenqto (SN 2018ibb), and provided the derived parameters to include in our catalog.

Table A1 compiles the metadata for each of the targets, including the internal ZTF name, IAU name, right ascension (RA), declination (DEC), redshift, Galactic extinction $E(B - V)$, discovery group, and additional information on spectral classification. Our sample covers the redshift of $z \sim 0.06 - 0.67$ with a median of $z_{\text{med}} = 0.265^{+0.143}_{-0.135}$. All redshifts in our sample are determined using the narrow emission lines from the host galaxy, except for nine events without host lines. The redshifts for these nine events are estimated from template matching by running *superfit* (Howell et al. 2006) over a range of z values. These redshifts are less accurate and marked with \star in Table A1. There are two additional events, ZTF19abcvwrz (SN 2019aamx) and ZTF19aawsqsc (SN 2019hno), which also have less accurate redshifts because of the low signal-to-noise ratios of Mg II $\lambda\lambda 2796, 2803$ absorption lines in their host-galaxy spectra. Figure 1 displays the redshift distribution of the full sample, including those derived from template fitting. To avoid possible biases caused by the choice of histogram grids, we apply kernel density estimation on all the histograms in this paper using a Gaussian kernel offered by the machine learning package Scikit-learn (Pedregosa et al. 2011), as shown in Figure 1.

Several SLSN-I samples from different surveys, including PS1, DES and PTF, as well as samples collected from the literature by Nicholl et al. (2015) and Inserra et al. (2018a), have revealed many important properties

¹ The ZTF public survey transitioned to a 2-day cadence on October 1, 2020, however the official start of the ZTF Phase II is December 1, 2020. We picked our date range for the convenience of the sources analysed in our sample.

of SLSNe-I. Compared with these previous samples, our sample size is significantly larger and the observing cadence is also better, as shown in Table 1. These two

key features allow us to investigate the LC properties of SLSNe with much better statistics.

Table 1. SLSN samples

Source	Candidates	Redshift range	Observing cadence ^a (rest-frame days)	Reference
Literature	25 ^b (14)	0.10 - 1.19	-	Nicholl et al. (2015); Inserra et al. (2018a)
PS1	17	0.32 - 1.57	2.30	Lunnan et al. (2018)
DES	22	0.22 - 2.00	3.65	Angus et al. (2019)
PTF	26	0.06 - 0.74	2.26	De Cia et al. (2018)
ZTF	78	0.06 - 0.67	1.45	This paper

^aThe median value of the observing cadence in the rest frame.

^bIncluding 11 SLSNe-I from PTF and PS1 samples and 14 independent SLSNe-I.

2.1. Photometric Selection

Here we briefly describe the photometric selection system of our SLSN-I candidates. Daily ZTF alerts are ingested into a dynamic science portal called the GROWTH Marshal (Kasliwal et al. 2019). A filter is implemented within the Marshal to select SLSN candidates. The candidates passing the filter do not get automatically saved; instead, each week, a human scanner has to visually examine the LC of each candidate selected by the filter and make a decision if it is worth being saved. The subsequent spectral follow up is based on the human-saved candidates. This filter adopts several cutoff conditions, including: [1] Not moving; the same alert is detected in two consecutive epochs with $\Delta t > 0.02$ days. [2] Not a star, based on the SDSS star-galaxy scores (Tachibana & Miller 2018). [3] Excluding bogus alerts based on scores constructed by Duev et al. (2019). [4] Not in the galactic plane, with Galactic latitude $|b| > 7^\circ$. [5] Variability was not detected at this location more than a year prior to the alert.

In addition, we assign numerical scores to several properties, including: [1] slowly rising events; [2] faint blue hosts; [3] spatial location of the transient relative to the center of the host (against nuclear transient); [4] brightness of the candidate relative to the host brightness (against faint transients); [5] the time interval between the first detection and the last one (against very long-lived transients like active galactic nucleus). This candidate filter gives high scores (thus preferences) to slowly rising events with faint blue hosts. For example, a candidate rising at least 20 - 25 days with a faint host

galaxy will have a high score and pass the filter and be also saved by the human scanner.

We note that the ZTF collaboration has multiple transient groups which also perform daily alert stream scanning, and most transient candidates can be saved by multiple groups, for example, the ZTF Bright Transient Survey (BTS, Fremling et al. 2020), the ZTF Census of the Local Universe (CLU, De et al. 2020), the fast transient group, the stripped envelop SN group and the ZTF Nuclear Transient group. Therefore, the classification of a specific SLSN-I candidate can also be drawn from other classification efforts, most noticeably the BTS. A few classifications are taken from the Transient Name Server (TNS) reported by external groups. But almost all of these externally-classified sources were also identified as candidates by our filter. The filtering method will be described in detail in the forthcoming work.

With the above selection criteria, on the order of 50 candidates per week are saved and get another round of vetting when spectral classification observations are planned. Any candidates brighter than 18.5 mag are classified by the ZTF BTS. Their classification efficiency at ≤ 18.5 mag is very high, close to 95% (Fremling et al. 2020; Perley et al. 2020). We anticipate that our catalog is almost entirely complete to SLSNe within the ZTF footprint peaking at magnitudes brighter than 18.5 mag, and highly complete to 19.0 mag. For magnitudes fainter than 19.0 (44 out of 78 events in total), there may be biases primarily related to the nature of the host and/or the rising phase of the light curve that will be examined in future work. The detailed structure of the light curve, *e.g.* the presence of bumps, was not a crucial factor in selection or triggering follow up.




Figure 1. Distribution of redshifts for the sample of 78 SLSNe-I presented in this paper. Other SLSN-I samples are plotted as hollow bars for comparison. The dashed line and the shaded area mark the median value and 1σ error (16% and 84% percentiles) of ZTF sample, $z_{\text{med}} = 0.265^{+0.143}_{-0.135}$. The black solid line shows the kernel density estimation of the distribution.

Our effort for spectroscopic classification is primarily focused on SLSN candidates fainter than 18.5 mag, using the Spectral Energy Distribution Machine (SEDM, Blagorodnova et al. 2018) and the Double Beam Spectrograph (DBSP, Oke & Gunn 1983) mounted on the Palomar 60 inch (P60) and 200 inch (P200) telescope, respectively. Additional facilities include the Low Resolution Imaging Spectrometer (LRIS, Oke et al. 1995) on the Keck I telescope, the Alhambra Faint Object Spectrograph and Camera (ALFOSC) on the 2.56m Nordic Optical Telescope (NOT), SPectrograph for the Rapid Acquisition of Transients (SPRAT) on the 2m Liverpool Telescope (LT), and the Intermediate-dispersion Spectrograph and Imaging System (ISIS) on the 4.2 m William Herschel Telescope (WHT). The basic information of classification spectra is listed in Table A2. The spectral reduction is performed using various standard reduction pipelines. This includes the SEDM automated pipeline (Rigault et al. 2019), the pyraf-dbsp package (Bellm & Sesar 2016) and DBSP_DRP (Roberson et al. 2022) pipelines for the DBSP data, and the LPipe package (Perley et al. 2019) for the LRIS data.

Finally, we note that the SLSN project has at least 0.5 to 1 night of DBSP time on P200 per month for spectral classification (PI: Yan). More quantitative discussions on the completeness of spectral classification will be included in Paper III.

2.2. Spectral Classification

Every event in our sample has at least one spectrum and most have multi-epoch spectra. The hallmark spectral features for SLSNe-I are the five O II absorption features in the wavelength range of 3737 – 4650 Å in pre-peak and/or near-peak optical spectra. These were identified and discussed in Quimby et al. (2011, 2018). We utilize the large SLSN-I spectral template library assembled by Quimby et al. (2018), and update the library by adding the missing phase information to some of the templates. Our classification relies on matching with the spectral templates using SNID (Blondin & Tonry 2007) and *superfit*.

To determine the best matched spectral templates, we run both *superfit* and SNID on the smoothed spectra with host-galaxy emission lines removed and with redshifts fixed for most sources. If the spectrum of a SLSN-I has a good match with that of a SNIc but its g -band peak luminosity is higher than -20.0 mag, we classify this candidate as a SLSN-I since many SLSNe-I develop spectra similar to those of SNeIc after the peak (Pastorello et al. 2010; Quimby et al. 2018; Gal-Yam 2019).

In Appendix A, Figures A7 – A9 present the best-fit spectral template for each event of our sample, with the information of event name, phase and template information labeled after the spectra. The phases are measured relative to the rest-frame g -band peak in this paper. We note that the phase differences between the

observed spectra of our SLSNe-I and the templates in the library are not zero, but generally within 30 days (rest frame). This is not surprising since SLSNe-I can have similar photospheric spectra but different LC evolution time scales (see Kangas et al. 2017; Quimby et al. 2018).

The above classification procedure can sometimes give ambiguous results for a small number of events, *i.e.* two events for our sample. In these cases, we rely on additional LC information, such as rise time and peak luminosity to break the degeneracy. For example, ZTF19aacxrab (SN 2019J) and ZTF19aaqrime (SN 2019kwt) are almost equally well-matched with the spectral templates of SNIa and SLSN-I. In the case of SN 2019J, although the overall spectral features broadly match with those of SNe Ia, its spectrum lacks S II $\lambda\lambda$ 5433, 5606 commonly seen in SNe Ia. In addition, because of their slow rising LCs and high peak luminosities (*i.e.* SN 2019kwt has $M_g \sim -22.8$ mag), we adopt the SLSN-I classifications, corresponding to the template spectra of SN 2007bi and PTF09cnd, respectively.

In summary, the 78 sources listed in Table A1 can be classified as SLSNe-I according to their features of spectra and light curves. The classification spectra are made available to the public as part of electronic data at the Journal website and will be uploaded to the Weizmann Interactive Supernova Data Repository (WiSeREP, Yaron & Gal-Yam 2012)².

3. OBSERVATIONS AND DATA

3.1. ZTF data and forced photometry

The bulk of the photometric data comes from ZTF, including the data from the public survey with a 3-day cadence, the ZTF partnership and Caltech surveys with a faster cadence (≤ 2 days) over smaller areas (Bellm et al. 2019b).

The IPAC ZTF pipeline produces reference-subtracted images using the ZOGY algorithm (Zackay et al. 2016) and aperture photometry for all transients detected at $\geq 5\sigma$. However, there are two issues with the LCs produced by this pipeline. One is that the reference images built in 2018 may contain signals from transients that exploded during the same period, for which the photometric offset needs to be corrected. The second issue is that the upper limits prior to the first detection are aperture photometry and based on the image noise measured over the entire quadrant. This can significantly underestimate transient signals.

To fix these two issues, we perform forced PSF photometry using the software provided by IPAC³. With a code from Yao et al. (2019), we refine the astrometric position of each event by using only the images around the peak phase. The very early and late-time forced photometry without transient signals allow us to compute baseline offsets to the LCs. In addition, we reject bad-quality data if: [1] the image processing and instrumental calibration fail to meet predefined quality criteria; [2] the robust estimate of 1σ value of the spatial noise per pixel in the image is over 25; [3] the seeing is larger than $5''$, similar to what was used in Yao et al. (2019). The photometry of the same transient observed with different CCD quadrants can have a systematic offset. This problem gets fixed by our photometric reprocessing as well. For the final photometric collections, a detection is defined as 4σ (3σ for *Swift* data) above the background and an upper limit is computed at 3σ .

The ZTF astrometric and photometric systems are calibrated using the *Gaia* DR1 data and the Pan-STARRS 1 catalogs, respectively (for details see Masci et al. 2019). The output magnitudes are in the AB system. The airmass and color term corrections are included when the photometry is calibrated to the PS1 system. The color information for color corrections ($g-r$ for g, r bands and $r-i$ for i band) is obtained from the g, r, i photometry taken at the same epoch or neighboring observations.

3.2. Supplemental photometry

3.2.1. P60, LT and P200 photometry

When ZTF is not able to take observations due to scheduling conflicts, bad weather, or when the transient is fainter than the ZTF detection limits, additional data are taken with SEDM on the P60 (Blagorodnova et al. 2018), the Optical imager of the Infrared-Optical suite of instruments (IO:O) on the LT (Steele et al. 2004) or the Wafer-Scale camera for Prime (WASP)⁴ on the P200. All of the P60, LT and P200 photometric results are obtained via PSF fitting, which are calibrated with the PS1 and SDSS (for u -band and part of P60 data only) standard stars. For the P60 data, the systematic errors between PS1 and SDSS systems may introduce an additional uncertainty of 0.01 mag in r and i bands and a larger error (up to 0.25 mag) in g band. The airmass and color-term corrections are included in all the P60/LT/P200 data. The P60 data are processed using the software Fpipe (Fremming et al. 2016) and image

² <https://www.wiserep.org/>

³ <http://web.ipac.caltech.edu/staff/fmasci/ztf/forcedphot.pdf>

⁴ <https://sites.astro.caltech.edu/palomar/observer/200inchResources/waspmmanual.html>

subtraction uses the SDSS references. While LT photometric data are processed with the software specifically built for IO:O (Steele et al. 2004; Fremling et al. 2016, Taggart et al. 2022 in prep.). The IO:O collects data in SDSS u , g , r , i and z bands and image subtraction is performed using reference images from the PS1 or the SDSS (for u -band only). The P200 data are processed using the software AutoPhOT (Brennan & Fraser 2022) and the PS1 reference images are used for image subtractions before performing photometry.

3.2.2. *Swift* data

27 of 78 events in our sample have observations from the UV/Optical Telescope (UVOT, Roming et al. 2005) aboard the *Neil Gehrels Swift Observatory* (Gehrels et al. 2004). We retrieved the UVOT data from the NASA *Swift* Data Archive⁵, and used the standard UVOT data analysis software distributed with HEASoft version 6.19⁶ (Nasa High Energy Astrophysics Science Archive Research Center 2014), along with the standard calibration data. We manually define sky background apertures devoid of any sources. Four events in our sample, namely ZTF18aavrmcg (SN 2018bgv), ZTF18acslpji (SN 2018hti), ZTF19aawfbtg (SN 2019hge) and ZTF19abpbopt (SN 2019neq), have bright host galaxies. We therefore requested host-galaxy images at very late phases when the SNe had faded. The photometry for these four objects are obtained by subtracting out the host galaxy fluxes. For most of the other events, their host galaxies are very faint and contribute no more than 10% of the SN fluxes in the UV-bands, so no host-galaxy subtractions are applied.

All the photometry data are listed in Table A3, and are available to the public in electronic format at the journal website.

4. THE OBSERVED LIGHT CURVES

The observed LCs for the full sample are presented in Figures A1 - A6, where the left Y-axis is apparent magnitude and the X-axis is the rest-frame days relative to the g -band peak phase. For a small number of events without a g -band peak phase, we use the r -band peak phase as the reference. Table A3 contains the complete upper limits over much wider time ranges (early and late times) than those shown in the plots. For better display, we plot only a small number of photometric upper limits.

One striking feature apparent in Figures A1 - A6 is that a large fraction of the LCs in our sample show significant undulations, and some have multiple peaks. In

Paper II, we find 18 – 44% of the well-sampled SLSNe-I have LC undulations. Among a small sub-set of SLSNe-I with early phase coverage, 6 – 44% show early double peak LCs. The quantitative analysis of the LCs and the discussions of these features are included in Paper II.

4.1. *Magnitude corrections*

4.1.1. *Extinction corrections*

The Galactic reddening $E(B - V)$ is taken from the Schlafly & Finkbeiner (2011) dust map using the NASA/IPAC InfraRed Science Archive (IRSA) database. The extinction corrections at different wavelengths are computed using the empirical dust extinction laws of Fitzpatrick & Massa (2007) with $R_V = 3.1$.

Previous studies have shown that SLSN-I hosts are mostly low-mass, metal-poor dwarf galaxies (Lunnan et al. 2014, 2015; Leloudas et al. 2015; Perley et al. 2016; Angus et al. 2016; Chen et al. 2017a; Schulze et al. 2018). This is the case for most of the events in our sample. Because their LCs do not show particularly red colors at pre-peak phases, we do not make any host-galaxy reddening corrections to these events. However, a small fraction of SLSN-I hosts are massive ($10^9-10 M_\odot$, Perley et al. 2016; Chen et al. 2017b), and the host-galaxy reddening in such cases can be large. We find that seven events in our sample (Table 2) have non-negligible host-galaxy reddening. They either reside in bright hosts ($M_r \lesssim -18.5$ mag) or have redder ($g - r$) colors at peak than average (see Figure 8), which is presumably due to host-galaxy reddening.

Table 2. Host-galaxy Reddening

Name	$E(B - V)_{\text{host}}$ (mag)	Method
SN 2018don	0.4 ^a	spec.temp
SN 2018kyt	0.11	line ratio
SN 2019kwt	0.22	line ratio
SN 2019aamx	0.07	spec.temp
SN 2019aamu	0.23	spec.temp
SN 2019stc	0.18	line ratio
SN 2020xkv	0.24	spec.temp

^aEstimated in Lunnan et al. (2020).

The host-galaxy reddening is difficult to measure accurately, but can be roughly estimated using two different methods. The first one uses Balmer line ratio measured from the host-galaxy spectra. Without dust attenuation, the predicted line ratio between $H\alpha$ and $H\beta$ re-

⁵ <https://heasarc.gsfc.nasa.gov/cgi-bin/W3Browse/swift.pl>

⁶ <https://heasarc.gsfc.nasa.gov/lheasoft/download.html>

mains constant in a wide range of temperature and electron density in the nebular regions (*e.g.* $H\alpha/H\beta \approx 2.86$, Osterbrock & Ferland 2006). The attenuation caused by dust scattering is wavelength-dependent and stronger at shorter wavelengths. This method is affected by both stellar and supernova absorption features. We remove the stellar Balmer absorption using *FIREFLY* (Wilkinson et al. 2017) before we measure the line flux, but ignore the influence of supernova features. This method is applied to the host spectra of ZTF18acyxnyw (SN 2018kyt) and ZTF19acbonaa (SN 2019stc). Note that the SN location may not coincide with the H II region responsible for the Balmer lines, so the real extinction at the SN site could be different from the inferred value. But such an estimation is at least an indicator of whether the galaxy has significant dust.

The second method is to infer the host-galaxy reddening by matching the observed spectrum (including spectral continuum slope and absorption features) with a SLSN-I template with negligible extinction and at a similar phase. For this method, the assumption is that the observed red spectral color is due to the host-galaxy reddening. The results from this method are very uncertain, but offer crude extinction estimates for those events without host emission lines. This method is applied to ZTF18aajqcue (SN 2018don), SN 2019aamx and ZTF19acvxquk (SN 2019aamu), whose spectra lack distinct Balmer emission lines from host galaxies. For example, to match the near-peak spectrum of SN 2018don, the $t \sim +54$ day template spectrum of SN 2007bi is required to be reddened by $E(B - V) \sim 0.4$ mag (Lunnan et al. 2020). Similarly, the +9 day spectrum of SN 2019aamu matches well with the -7 day template spectrum of PTF12gty after considering a host-galaxy reddening of $E(B - V) = 0.23$ mag. And the +13 day spectrum of SN 2019aamx matches well with the +22 day template spectrum of PTF09cwl with $E(B - V) = 0.07$ mag (shown in Figures A7 – A9). Note that both PTF12gty and PTF09cwl have a faint, dwarf host, so their host-galaxy reddenings are assumed to be negligible (Quimby et al. 2018; Perley et al. 2016).

For SN 2019kwt and ZTF20abzaacf (SN 2020xkv), we applied both of the above methods, since their spectra contain strong supernova signals and we are not able to remove the stellar absorption nor supernova features. The derived $E(B - V)$ for SN 2019kwt is 0.22 mag from the host-galaxy spectrum and 0.18 mag from template matching, which appear to be consistent. In the case of SN 2020xkv, the $E(B - V)$ derived from the spectral matching ranges from 0.22 – 0.26 mag, in contrast to the lower value (0.17 mag) inferred from the Balmer decrement. Its spectrum has very low signal-to-noise ratio at

$H\beta$, and with both highly uncertain measurements, we adopt the average value of 0.24 mag from the template matching method.

4.1.2. The K-corrections

One of the key parameters of SLSNe-I is the absolute g -band peak magnitudes. Proper estimate of this value requires proper K-corrections to derive reliable rest-frame values, either in g -band for sources at $z \leq 0.17$ or in r -band for sources at $z > 0.17$. This switch is chosen because the observed r -band LC at $z > 0.17$ is closer to the rest-frame g -band in wavelength.

Most events in our sample have at least one spectrum around the LC peak (*i.e.* $\lesssim 20$ days in rest frame), which allow us to compute the K-corrections using the formula described in Hogg et al. (2002). For those without spectra near the peak phases, we apply a constant K-correction of $-2.5 \times \log(1 + z)$. This approximation is not too far from the spectral estimates, as shown below.

Figure 2 compares the calculated K-corrections based on observed spectra (blue dots) and an assumed 10^4 K blackbody spectral energy distribution (SED) (green line) with the constant correction (black line). The constant value of $-2.5 \times \log(1 + z)$ appears to capture most of the corrections within a range of ± 0.1 mag (shaded region) for SLSNe-I at $0.06 < z < 0.67$. We list the K-corrections for each event in Table A4.




Figure 2. K-corrections from the observed band to the rest-frame g band for SLSNe-I around the peak. Blue points represent the corrections computed from the observed spectra. The black line shows the constant correction of $-2.5 \times \log(1 + z)$ with an uncertainty of 0.1 mag shown by the gray shaded area. The green line shows the correction calculated assuming a blackbody spectrum with a temperature of 10^4 K. The break at $z = 0.17$ is caused by the change of observed band (from g to r band).

4.1.3. The S-corrections

As the g -, r - and i -band photometry of SLSNe-I presented in this work are obtained from four different telescopes, the errors caused by filter differences need to be evaluated. Since most of the data are from the ZTF itself, we choose to correct the LT/P200/P60 photometry to that of the ZTF filters. Following the method of Stritzinger et al. (2002, 2005), Pignata et al. (2008) and Wang et al. (2009), the correction between different instruments (known as the S-correction) can be computed using

$$Sc_{\lambda 1} = M_{\lambda 1} - m_{\lambda 1} - CT_{\lambda 1}(m_{\lambda 1} - m_{\lambda 2}) - ZP_{\lambda 1}, \quad (1)$$

where, $M_{\lambda 1}$ is the SN synthetic magnitude computed with the response functions of ZTF $\lambda 1$ filter, and $m_{\lambda 1}$ and $m_{\lambda 2}$ are the SN synthetic magnitudes computed with the LT/P200/P60 filters. $CT_{\lambda 1}$ is the color term and $ZP_{\lambda 1}$ is a zero point, which are measured by convolving the ZTF filters with a large sample of spectrophotometric Landolt standard stars from Stritzinger et al. (2005). The response function above includes filter transmission, detector quantum efficiency and atmospheric transmission.

As S-corrections are spectrum-dependent and accurate measurements at any given epoch require far more SLSN-I spectra than we have in this work. We use the S-correction values computed from the spectra as an additional 1σ error in photometry, and add them to the existing errors of LT/P200/P60 photometry in quadrature. The additional 1σ errors due to the S-corrections are listed in Table 3.

Table 3. The estimated error caused by the S-correction.

Telescopes	Filters		
	g (mag)	r (mag)	i (mag)
LT	0.018	0.044	0.030
P60	0.042	0.030	0.026
P200	0.023	0.030	0.025

4.2. Empirical LC Fitting Method

The LC analysis requires estimates of various parameters such as peak magnitude, rise time and rise rate, which all involve numerical interpolation and fitting. We adopt a machine learning algorithm, Gaussian Process (GP) regression which has various kernel functions. The GP interpolation can reduce the influence of outliers and

gives robust error estimates. We use a composite kernel which is the sum of Matérn kernel with a white noise kernel. We tested Matérn kernels with ν parameter of 3/2 (Matérn 3/2) (Inserra et al. 2018a; Angus et al. 2019; Lunnan et al. 2020) as well as $\nu = 5/2$ (Matérn 5/2). When we compute peak magnitudes and phases, GP fit with Matérn 3/2 is performed in the flux space. The Python package `george` (Ambikasaran et al. 2016) and the Scikit-learn give comparable results.

4.3. LCs in Absolute g -band magnitudes

To set an approximate luminosity scale, we also plot the absolute magnitude on the right Y-axis of Figures A1 - A6. This is calculated by assuming a constant K-correction of $-2.5 \times \log(1+z)$. In these figures, the LCs have been corrected for the Galactic and host-galaxy reddening whenever possible.

To calculate the apparent peak magnitude and phase, we run the GP to interpolate the LCs in flux space. The errors are shown as 1σ uncertainty. The g -band absolute peak magnitudes are derived using $M_g = m_g - \mu - KC$, where μ is the distance modulus, KC is the K-correction, and m_g is the apparent g -band magnitudes corrected for both the host (in rest frame) and the Galactic extinction (in observed frame). The rest-frame g -band absolute peak magnitudes and the peak dates are tabulated in Table A4. The peaks of some events are not well constrained due to the poor sampling around the peak. Figure 3 shows the distribution of the g -band absolute peak magnitudes (M_g) for the ZTF sample, ranging from -19.8 mag to -22.8 mag with a median and 1σ error of $M_{g,\text{med}} = -21.48_{-0.61}^{+1.13}$ mag. Both our median value and dispersion are consistent with those of previous samples.

The weak bimodal distribution in the peak absolute magnitudes should not be over-interpreted because this is the raw distribution function without applying corrections for various selection biases (*e.g.* Malmquist bias, preference of our selection system for slowly-rising events). Selection biases could also have a strong impact on the distribution of the timescales (*e.g.* Figure 5), peak bolometric luminosities (*e.g.* Figure 16) and other photometric properties. The correct interpretation of these distributions requires a well-defined selection function of our sample, the detailed simulation of which, however, is well beyond the scope of this paper. Further studies (*e.g.* Paper III) are required to confirm these distributions.

5. LIGHT CURVE PARAMETERS

5.1. Rise/decay time scales – fast and slow SLSN-I events




Figure 3. The distribution of the rest-frame g -band absolute peak magnitudes. The dashed line and shaded region mark the median value and the 1σ dispersion, $-21.48_{-0.61}^{+1.13}$ mag. The black solid line shows the kernel density estimation of the distribution. Other SLSN-I samples are plotted in different hatching patterns for comparison. DES and PS1 samples are measured at 400 nm, which is bluer than the g band (472 nm).

Traditional rise times, defined as the interval between the explosion date and the peak, are notoriously difficult to measure, especially for distant transients such as SLSNe-I. Although the ZTF detection can go down to 20.5–21.0 magnitude, at the median redshift of our sample, it only probes an absolute magnitude of ~ -20 mag. The upper limits before the first detection usually can not place strong constraints on the LC evolution in very early phases and only about one-third of our sample have high-quality early time data. In addition, Anderson et al. (2018) shows that a SLSN-I (SN 2018bsz) can have a long, slowly rising “plateau” before a steeper, faster rise to the peak, which makes it harder to speculate explosion dates without enough deep, early data.

Instead of using the traditional rise time, we define $t_{\text{rise/decay},x}$ as the time interval between the peak and the flux when it is at a fraction x of the peak value. Here the x factor can be 10% ($\Delta\text{mag} = 2.5$), or $1/e$ ($\Delta\text{mag} \approx 1.09$). Time dilation is corrected for all time scale measurements. The derived $t_{\text{rise},10\%}$ and $t_{\text{rise/decay},1/e}$ are listed in Table A4. The poorly constrained time scales are not included in the following analysis.

Figure 4 shows a direct proportionality between $t_{\text{rise},1/e}$ and $t_{\text{rise},10\%}$ in the rest-frame g band. Linear fit gives $t_{\text{rise},1/e} = 0.80 t_{\text{rise},10\%} - 1.73$ days with 1σ uncertainty as small as ~ 2.84 days. The small scatter implies that the LC rise rates at the very early phases do not have significant differences in our sample. The rise

times $t_{\text{rise},10\%}$ cover a wide distribution, ranging from 10–90 days, with the mean (median) value and a standard deviation of $\bar{t}_{\text{rise},10\%} = 41.9(38.3) \pm 17.8$ days.




Figure 4. Rise time measured at 10% of the peak flux $t_{\text{rise},10\%}$ versus that measured at $1/e$ of the peak flux $t_{\text{rise},1/e}$. The vertical solid gray line marks the mean value of $t_{\text{rise},10\%}$, the dashed and dotted lines represent the 1σ and 2σ uncertainties, respectively. The black solid line and shaded area show the linear fit to the measured time scales ($t_{\text{rise},1/e} = 0.80 t_{\text{rise},10\%} - 1.73$ days) and the 1σ dispersion. The events with extreme time scales are highlighted in different colors.

Of the total 56 SLSNe-I with $t_{\text{rise},10\%}$ measurements, the two slowest rising events SN 2018ibb and ZTF20aapaecd (SN 2020fyq) stand out in Figure 4, with time scales longer than $\bar{t}_{\text{rise},10\%} + 2\sigma = 78$ days.

For another slowly-evolving event ZTF20aadzbcf (SN 2020fvm), its $t_{\text{rise},1/e}$ reaches 91 days, suggestive of an unusually long rise time, although $t_{\text{rise},10\%}$ is not well constrained. SN 2020fvm has two LC peaks and we set the second, also the brighter one as its main peak. If the first one is set as the main peak, its $t_{\text{rise},1/e}$ is around 48 days. The fastest event in our sample is SN 2018bgv (Lunnan et al. 2020), which rises to the peak in less than 10 days. Such a fast and luminous event is very rare and it may be associated with the Fast Blue Optical Transients (FBOT, Drout et al. 2014; Ho et al. 2021). Compared with previous SLSN-I samples, our sample contains slightly more fast-evolving events. For instance, 9 out of 56 events (9-26%, calculated at a confidence level of 95% based on Gehrels 1986) are found to have $t_{\text{rise},1/e} \lesssim 15$ days, while only 5 in 55 events (4-18%) were reported in the previous samples.

Outside the $\pm 1\sigma$ range, we have 10 fast-evolving events with $t_{\text{rise},10\%} \leq \bar{t}_{\text{rise},10\%} - 1\sigma = 24$ days and 10 slow events with $t_{\text{rise},10\%} \geq \bar{t}_{\text{rise},10\%} + 1\sigma = 60$ days. The $t_{\text{rise},10\%}$ shows a continuous distribution and can not be

divided into two separate fast and slow subclasses, as indicated by previous studies (Nicholl et al. 2015; De Cia et al. 2018).

Figure 5 shows comparison of the rise and decay time scales measured at $1/e$ of the peak flux for 48 events. Both rise/decay time scales show a distribution centered at $\sim 25 - 30$ days with an extended tail. For the whole sample, the rise and decay time scales show a strong positive correlation with a Spearman correlation coefficient of $\rho = 0.73$ and null probability $p < 10^{-8}$, *i.e.* slowly-rising events tend to decay slowly. Applying a linear fit, we find $t_{\text{decay},1/e} = (1.47 \pm 0.07) t_{\text{rise},1/e} + (0.35 \pm 2.50)$ days, which is similar to what was found in previous studies (Nicholl et al. 2015; De Cia et al. 2018).

It is interesting to note that four of the five SLSNe-Ib in the Yan et al. (2020) sample, namely SN 2018kyt, SN 2019hge, ZTF19acgjpgh (SN 2019unb) and ZTF20ablkuio (SN 2020qef), have rise time scales which are 9 – 32 days longer than that of the rise-decay time correlation. The last SLSN-Ib, ZTF19aamhhiz (SN 2019kws), in the Yan et al. (2020) sample has a short rise time but much longer decay time. As noted in Yan et al. (2020), these SLSNe-Ib may have He-rich CSM and CSM interaction may affect the LC evolution before or after the peak. Detailed modeling and discussions of their LCs are given in Paper II.

Although SNe Ic and SLSNe-I have similar post-peak spectra, they have very different peak luminosity and time scales. Figure 6 compares the peak M_g and rise times between normal SNe Ic, broad-lined SNe Ic (SNe Ic-BL) and SLSNe-I. The SLSNe-I are from our sample, while the normal SNe Ic sample and the SNe Ic-BL sample are taken from Barbarino et al. (2021) and Taddia et al. (2019), respectively. We exclude two bright SNe Ic from literature, iPTF12gty and iPTF15eov, which can be better classified as SLSNe-I as suggested by their authors. Both SNe Ic samples are obtained from the (intermediate) PTF, which has similar observation depth to ZTF. The overall distribution in Figure 6 is similar to that shown in De Cia et al. (2018). It should be noted that correction for observational selection biases and volumetric correction are required to interpret the distribution correctly. The peak M_g of SLSNe-I is about 4 and 3 magnitude brighter than those of normal SNe Ic and SNe Ic-BL, respectively. All SLSNe-Ib have low luminosities and moderate rise times compared with normal SLSNe-I. SLSNe-I have significantly longer rise times and wider dispersion compared to those of SNe Ic. First, this is physically related to the fact that they have larger ejecta masses and more massive progenitor stars (see Paper II). Second, SNe Ic are primarily powered by radioactive decay which has a constant time scale




Figure 5. Time scales measured at $1/e$ of the peak flux for the rising and declining portions of LCs. Normal SLSNe-I are shown in blue circles while He-rich SLSNe-Ib are highlighted in red. The black dashed line and the shaded area show the best linear fit and 1-sigma error, $t_{\text{decay}} = 1.47 t_{\text{rise}} + 0.35$ days. The histograms along the horizontal and vertical axes show the distributions of the rise and decay time scales, respectively. The black solid lines show the kernel density estimation of the distributions. The time scales from other SLSN-I samples are plotted with smaller gray points for comparison. The data from PTF are measured at 1 mag below peak magnitude and is slightly shorter than the $1/e$ -maximum (1.09 mag) time scales. And those from the literature (Nicholl et al. 2015) and the PS1 are measured from bolometric LCs and may have biases due to redshift and the SED.

of energy injection. In the magnetar model proposed for SLSNe-I, the time scale of energy injection depends on magnetic field B ($\propto B^{-2}$) and spin period P ($\propto P^2$). The LC width expands with the decrease of B (see Figure 2 of Kasen & Bildsten 2010). The large spread in their rise times could suggest that the magnetic field strengths of the neutron stars can vary over a wide range as well.

5.2. Color and blackbody temperature

Understanding the variation and uniformity of the SLSN-I SEDs as a function of time should shed light on the physical nature of this population of stellar explosions. Although the complete characterization of the SLSN-I SEDs is beyond the scope of this paper, we can infer some basic properties by examining the distributions of $(g-r)$ colors and blackbody temperatures since both parameters are determined by the transient SEDs.




Figure 6. Rise times $t_{\text{rise},10\%}$ versus g -band absolute peak magnitudes M_g for SLSNe-I (this paper), normal SNe Ic (from Barbarino et al. 2021) and broad-lined SNe Ic (SNe Ic-BL, from Taddia et al. 2019). The M_g of the SNe Ic and SNe Ic-BL samples is computed from the r -band magnitudes using a color correction of ~ 0.36 mag (Taddia et al. 2015; Prentice et al. 2016). The rise times of SNe Ic and SNe Ic-BL are measured from the explosion date in r band, which are slightly longer than $t_{\text{rise},10\%}$. The SLSNe-Ib in our sample are highlighted.

Figure 7 shows the observed color evolution as a function of time. All of the colors have been corrected for reddening but not for K-corrections, since that requires better spectral coverage than we have. The overall $(g-r)$ color trend evolves from blue (*i.e.* ~ -0.3 mag, hotter temperatures) at early phases to red and reaches $(g-r) \sim +0.8$ mag at about 2 months after the peak. In our sample, four events have peculiar color (temperature) evolution, and do not follow the general trend. We discuss these outliers below. We further examine the distribution of the observed $(g-r)$ color obtained at the maximum light in Figure 8. At the peak phases, the median observed color is close to zero, $(g-r)_{\text{med}} = -0.03^{+0.12}_{-0.11}$ mag.

Figures 7 and 8 show large scatters. The observed colors are not easy to interpret because they sample different parts of the SEDs, depending on the transient redshifts. The proper measurements are the rest-frame color tracks, however, reliable results would require many more spectra than our sample has. Here we compute only the rest-frame $(g-r)$ colors at the peak phase using the color corrections computed from the near-peak spectra. The rest-frame $(g-r)$ colors are corrected from the observed $(g-r)$ for the events at $z \leq 0.17$ and $(i-r)$ for $z > 0.17$, if photometry data and spectra are available. All the peak colors are listed in Table A5. The green histogram in Figure 8 shows that

the median rest-frame $(g-r)_{\text{rest,med}}$ is $-0.21^{+0.19}_{-0.12}$ mag, which is consistent with -0.27 mag calculated from the PTF SLSNe-I sample (De Cia et al. 2018). The large scatter in the peak rest-frame $(g-r)$ indicates that the SEDs of SLSNe-I may show a diverse shape and hence a wide range of blackbody temperatures.

Figure 9 illustrates a moderate correlation ($\rho = 0.52$, $p < 10^{-3}$) between the peak rest-frame $(g-r)$ colors and the g -band absolute peak magnitudes $M_{g,\text{peak}}$, *i.e.* brighter SLSNe-I tend to have bluer color, which was also previously found by Inserra & Smartt (2014) and De Cia et al. (2018) in smaller samples. If combined with the PTF sample from De Cia et al. (2018), the correlation becomes stronger ($\rho = 0.56$, $p \approx 2 \times 10^{-5}$), and can be described by a linear function, $M_{g,\text{peak}} = (11.5 \pm 2.6) \times (g-r) - (18.9 \pm 0.5)$ mag, with 1σ error of 1.7 mag. This correlation can be used in cosmological searches of SLSNe, *e.g.* Inserra et al. (2021). However, this is beyond the scope of this paper.

To fit the blackbody temperature, we adopt a modified blackbody function, defined by $f_\lambda = \max[0, 1 - A \times (1.0 - \lambda/3000.0)] \times B_\lambda$, with B_λ being the Planck function, and A being the scaling factor. The modified blackbody function aims to quantitatively capture the variations in the UV spectral suppression for different events, as shown by the HST UV spectra of SLSNe-I (Yan et al. 2017, 2018). The scaling factor A is derived from the fitting at $\lambda \leq 3000 \text{ \AA}$, set to zero at $\lambda \geq 3000 \text{ \AA}$ and fit in a range of 0–3. Larger scaling factors represent stronger suppression in UV and $A = 1$ represents the SED function used in Nicholl et al. (2017). The error associated with the temperature is estimated using the Markov Chain Monte Carlo (MCMC) method.

Of the 78 events in our sample, only 15 have at least 3 epochs of *Swift* UV photometry for properly computing the blackbody temperatures. In Figure 10, the left panel shows the temperature evolution tracks for 12 events, and the right panel shows the rest three with peculiar color/temperature evolution. Although the statistics is not large, Figure 10 indicates the large temperature spread at any given phase, especially at pre-peak and peak phases. In the left panel, some SLSNe-I have high temperatures at ~ 15000 K at $t \sim -10$ days, and cool down to ~ 9000 K at $+20$ days. Interestingly, there are also two cooler SLSNe-I (*i.e.* SN 2019hge and SN 2019unb) with the peak temperatures less than ~ 10000 K. Consistently, neither of them shows clear O II absorption lines in their classification spectra, which require a high ionization temperature (*i.e.* $T \sim 15000$ K, Quimby et al. 2018). We perform a polynomial fit to the temperature tracks and derive $T = 0.090201 t^3 - 3.4306 t^2 - 180.30 t + 13395$ K and




Figure 7. The observed $(g-r)$ color evolution tracks with time. We highlight four events whose $(g-r)$ color tracks do not follow the general trend shown in gray.

$T = -0.052720t^3 + 0.3319t^2 - 15.84t + 9007$ K for the high and low temperature tracks, respectively. These are shown as blue and red shaded regions with the $\pm 1\sigma$ uncertainty in Figure 10. Consistent with the temperature evolution, the $(g-r)$ colors of low-temperature events are $\sim 0.2 - 0.3$ mag redder than those of high-temperature ones at the peak but become indistinguishable from that of the full sample after +15 days after the peak.

Moreover, the color distribution plots suggest that there are more low-temperature SLSNe-I. For instance, 10 out of 35 (29%) events are found to have redder peak rest-frame colors than the two low-temperature events (~ 0.10 mag). However these 10 events have no UV photometry available, thus no temperature measurements. It is possible that these 10 red events could also have low peak temperatures.

Combining the temperature measurements from the PS1 sample, we do see more low-temperature events. To better quantify the temperature distribution of SLSNe-I, we choose three typical epochs (*i.e.* $t \sim -10$ days, 0 days and +20 days relative to the rest-frame g -band peak) when sufficient data are available from the ZTF and PS1 sample. We show the temperature distribution at these epochs in Figure 11. At $t \sim -10$ days, the temperature of SLSNe-I ranges from 7000 to 23000 K for different events, while this range is 6000 – 20000 K around the peak and 6000 – 12000 K (excluding one special event SN 2019szu) at $t \sim +20$ days. As SLSNe-I evolve, both the median value and the spread of temperature become

smaller. This conclusion will still hold true if we expand the temporal range from -20 to $+40$ days, according to the temperature trend shown in Figure 10. Furthermore, the temperatures at any epochs in Figure 11 are found to show flat and unimodal distributions, which indicates that SLSNe-I may have a continuous temperature distribution.

We also measure the blackbody radius of the photosphere. As shown in the top panel of Figure 12, most events show a linearly expanding photosphere from explosion to 10 – 20 days around the peak. This is due to that the recession of the photosphere is negligible during this phase. We apply a linear fit to measure the photosphere velocity, V_{phot} . In paper II, we measured the velocity from the Fe II and O II absorption lines in spectra. The velocity implied from the species V_{ion} are expected to be higher than V_{phot} since only the line features formed at higher velocity and lying outside the photosphere can be observed. As shown in the bottom panel of Figure 12, the distribution of the $\Delta V = V_{\text{phot}} - V_{\text{ion}}$ proves that the most events have a negative ΔV and their V_{phot} is on average lower than V_{ion} by about $2000 - 3000 \text{ km s}^{-1}$. Two outliers, SN 2018bgv and SN 2018kvt, are found to have significantly higher V_{phot} than V_{ion} . SN 2018bgv is the fastest-evolving event in our sample, while SN 2018kvt also exhibits relatively fast-evolving behaviors. Thus the inconsistency between V_{phot} and V_{ion} may be due to that V_{phot} is the average velocity measured in a period of 10 – 20 days while V_{ion} is fast-evolving and measured at




Figure 8. The distribution of the $(g-r)$ colors at peak. *Top panel:* The red dotted line represents the original colors of the events with host-galaxy reddening and the red area shows their corrected observed color. The blue area represents the observed colors of the events with only Galactic extinction correction. After host-galaxy reddening correction, the observed $(g-r)$ color at peak has a median value with the 1σ dispersion of $-0.03^{+0.12}_{-0.11}$ mag, marked by the blue dashed line. The blue solid line shows the kernel density estimation of the distribution. *Bottom panel:* The green area represents the rest-frame colors of the events with K-correction. The rest-frame $(g-r)$ color at peak has a median value with the 1σ dispersion of $-0.21^{+0.19}_{-0.12}$ mag, which is represented by the green dashed line. The green and blue solid lines show the kernel density estimations of the rest-frame colors and the observed colors (normalized to the number of the rest-frame colors), respectively.

a single epoch. All of the measured temperature and radius values are listed in Table A6.

In summary, our measurements show that the SLSN-I have a wide range of temperatures, especially at early phases. While most SLSNe-I from ZTF have temperatures over 11000 K at the peak and cool down




Figure 9. The correlation between the g -band absolute peak magnitudes $M_{g,\text{peak}}$ and the rest-frame $(g-r)$ colors. The linear fit and 1σ error are shown by the black line and the shaded area. The SLSN-I sample from this paper is marked in blue while that from PTF (De Cia et al. 2018) is in orange.

rapidly, there are also many lower-temperature and slowly-evolving events. This indicates that the SEDs of SLSNe-I may have diverse shapes and different evolutionary tracks.

Finally, we discuss four peculiar events that have extraordinary color and temperature evolution as shown in Figures 7 and 10.

SN 2018hti is well sampled in UV and shows notably higher temperatures and much bluer $(g-r)$ colors compared with other events. Its LC can be well reproduced by a magnetar model (Lin et al. 2020). This event may represent some of the population with higher temperatures and bluer SEDs. However, we caution that this event has a very high Galactic extinction $E(B-V) = 0.4$ mag. The dust extinction corrections in the UV-bands are highly uncertain.

SN 2020fvm has two almost equally bright LC peaks (at phase ~ -60 and 0 days, respectively) and the longest $1/e$ -maxima rise time scale of 91 days. Both its color and temperature evolutions are peculiar. The $(g-r)$ color around two peaks shows similar evolutionary trends, *i.e.* initially evolving from red to blue before the peak and then turning red after the peak. During the phase from -75 days to $+35$ days, its temperature remains almost constant of ~ 10000 K, which could be due to the absence of sampling. Such a long time scale and double-peak evolution are unexpected for a simple radiative cooling or magnetar model. Dessart (2019) examined the color evolution for various configurations of magnetar models, and found that almost all have fairly blue colors in the early phase and none shows the color change of ~ 0.7 mag from red to blue. The ejecta inter-




Figure 10. Blackbody temperature evolution as a function of time. For comparison, we plot the temperature measurements from the PS1 sample in light gray. **Left panel:** We include 12 of 15 events with at least 3 epochs of UV data. We apply third-order polynomial fits to the temperature evolution of two low-temperature events, SN 2019hge and SN 2019unb, and show the result with the 1σ error in the red shaded area. Similarly, the fit of 10 high-temperature events is shown by the blue shaded area. **Right panel:** We highlight the three extraordinary events and plot the 12 normal events in dark gray for comparison.




Figure 11. The temperature distributions (including both ZTF and PS1 samples) at -10 days, 0 days and $+20$ days relative to the rest-frame g -band peak. The dashed lines mark the median temperature at different epochs, *i.e.* $T \sim 12600$ K at -10 days, $T \sim 12000$ K at peak and $T \sim 9200$ K at $+20$ days. The solid lines show the kernel density estimation of the distribution.

action with an extended H-poor CSM could be a possible explanation.

ZTF19aanestg (SN 2019cdt) has much redder color at $+20$ days after the peak and it evolves much faster in

comparison with other objects of our sample. No temperature evolution is computed due to the lack of UV data. This event is somewhat similar to SN 2018bgv, both belonging to the fast-evolving and CSM-model-favored events (see LC modeling in Paper II), though it has redder ($g - r$) color. Such a red color and fast LC evolution could also be consistent with a magnetar model with high kinetic energy, as shown in Figure 13 of Dessart (2019). Nevertheless, the Fe II velocity and the kinetic energy of the ejecta derived for SN 2019cdt are ~ 14400 km s $^{-1}$ and 7.6×10^{51} erg, respectively, which are both higher than the average value (see Paper II).

Finally, unlike any other SLSN-I, ZTF19acfwynw (SN 2019szu) shows unusually high temperatures which are rising as the luminosity declines. At the early phases, its colors are blue, consistent with having high temperatures. Although the black-body temperatures at late time have large errors due to contamination of increasingly strong nebular emission lines, its spectral sequence has revealed a clear excess of UV continuum emission well past the peak phase. One possible explanation is CSM interaction, which could offer an additional heating source, boosting the emission at shorter wavelength. Another possibility is that the ejecta could be ionized by the ionizing flux from a long-lived central source (*e.g.* a magnetar, Margutti et al. 2017). Ionization can increase the optical opacity dominated by the electron scattering




Figure 12. *Top panel:* The photosphere radius evolution as a function of time. Our measurements are plotted as open points while those for PS1 sample (Lunnan et al. 2018) are shown in grey lines for comparison. For the events showing a linearly expanding photosphere before or at the peak, we apply a linear fit for them and plot the results with solid lines. *Bottom panel:* The distribution of the velocity difference between those derived from photosphere radius (V_{phot}) and those measured from the absorption lines in spectra (V_{ion}). Blue and orange bars represent the value measured from Fe II and O II, respectively. The solid lines show the kernel density estimation of the distributions.

and decrease the UV opacity dominated by line transitions of metals, which leads to a shift of the peak of the SED from optical to UV frequencies.

It is worth noting that SN 2020fvm and SN 2019cdt are poorly fit by simple magnetar models. It is possible that CSM interaction plays significant roles in these two systems. Additional modeling and analysis are presented in Paper II.

5.3. Bolometric correction and bolometric LC

Bolometric light curve is an important indicator of the total radiative energy of a transient and sets constraints on the possible explosion mechanisms. Due to the high photospheric temperatures of SLSNe-I, UV ra-

diation contributes to a large fraction of their emission (Yan et al. 2017, 2018). Since only a small portion of the sample has UV data, we need to first derive an empirical bolometric correction (BC) relation, which can tie g - and r -band photometry to the bolometric luminosity, with $\log L_{\text{bol}} = \log L_{gr} + \text{BC}$ ⁷. Here L_{gr} is the sum of the g - and r -band luminosities.

Lyman et al. (2013) derived bolometric corrections for a sample of core-collapse SNe at low redshifts using their well-observed SEDs. We adopt a similar method and apply it to our sample. The basic concept is following. Bolometric correction is strongly influenced by the redshift and the temperature (slope of the spectra) at each epoch. As the observed $(g - r)$ color is also largely determined by the same two parameters, it is expected to see a correlation between the BC and the $(g - r)$ color. This correlation can serve as the basis of constructing bolometric LCs with only optical data.

We first compute the bolometric luminosity at the epochs when both UV and optical data are available, using the modified blackbody SED fit described in §5.2. The UV scaling factor A is derived by fitting the UV to optical SEDs. For the epochs without UV data, we set $A = 1.55$, which is the median value of the epochs with UV data. The bolometric luminosities are shown in Figure 14 (only 15 events with at least three-epoch UV photometry) and listed in Table A6.

Figure 13 shows the observed $(g - r)$ color versus the bolometric correction, *i.e.* $\log L_{\text{bol}}/L_{gr}$, the luminosity ratio. We apply a third-order polynomial fit by optimizing a Gaussian likelihood function and derive an empirical relation as $\log(L_{\text{bolo}}/L_{gr}) = -1.093x^3 + 1.244x^2 - 0.261x + 0.410$, where $x = (g - r)$. Each point is weighted by the errors of both color and luminosity ratio. This equation can be used to compute the BC for low- z SLSNe-I with similar $(g - r)$ color evolution as our sample. Note that our fit is applicable for $-0.4 < (g - r) < +0.9$ mag, the phase range of -74 days $< \text{phase} < +173$ days and $0.06 < z < 0.57$.

As inferred from the residuals, the systematic error for the derived bolometric luminosity is about 19%. The BC uncertainty (shown as the shaded area in Figure 13) is the combined errors from MCMC estimates and the systematic error. This error is the dominant one compared with other error sources like host-galaxy reddening and redshifts. In Figure 14, the bolometric luminosities constructed with both UV and optical photometry are con-

⁷ Another definition of bolometric correction is $M_{\text{bolo}} = M_K + \text{BC}_K$ for K -band absolute magnitude M_K , where M_{bolo} is bolometric magnitude. Our correction is applied to two bands (g and r), and we use the definition introduced in §5.3.




Figure 13. Bolometric corrections (ratio between bolometric luminosity and the gr -band luminosity) derived from the events with both UV and gr -band photometry. X axis represents the $(g-r)$ color in the observed frame, where the Galactic and host-galaxy reddening have been corrected. The color bar on the right indicates the redshift of each data point. The black solid line shows the result of third-polynomial fit and the gray area shows 1σ uncertainty.

sistent with the ones derived from g - and r -band photometry. This illustrates the reliability of our method.

The bolometric LCs derived from g - and r -band photometry are shown in Figure 15. The errors of both photometry and bolometric correction are combined together in quadrature to represent the error of the bolometric LCs. ZTF19aauiresf (SN 2019fiy) and SN 2019aamu are excluded in this figure due to the bad photometry quality. Figure 15 illustrates a diversity in both peak luminosities and LC widths. Bumpy LCs are commonly seen and we perform detailed analysis in Paper II.

The peak bolometric luminosities are tabulated in Table A4 and the distribution is shown in Figure 16. The peak bolometric luminosities show a similar distribution as the absolute peak magnitudes. These are the raw distribution functions without any corrections for selection biases. Further studies are required to confirm such bimodal distributions. The peak bolometric luminosity spans from 3.22×10^{43} to $7.80 \times 10^{44} \text{ erg s}^{-1}$, with a median value and the 1σ dispersion of $L_{\text{bolo,med}} = 2.00_{-1.44}^{+1.97} \times 10^{44} \text{ erg s}^{-1}$. Our measurements are consistent with the results from the PTF and PS1 sample. Note that the DES sample tends to have lower luminosities, which could be due to that they are pseudo bolometric luminosities constructed from trapezoidal in-

tegration of photometry. Compared to the median peak luminosity of SNe Ib/c ($\sim 2 \times 10^{42} \text{ erg s}^{-1}$, Prentice et al. 2016), SLSNe-I are about 100 times more luminous, indicating different energy sources and/or explosion mechanisms from normal ccSNe.

6. CONCLUSIONS

The large sky coverage, high sensitivity and uniform cadence have made ZTF a very efficient discovery machine for superluminous supernovae. During the phase-I operation of ZTF survey, a total of 85 SLSNe-I were discovered. This paper presents 78 SLSNe-I whose LCs cover both pre- and post-peak phases. The other 7 SLSNe-I were still rising before October 30, 2021, and are not included in current analysis.

This sample represents the largest sample of SLSNe-I at $z < 1$ discovered from a single survey. Compared with the previous SLSN-I samples, our sample also has a better observing cadence on average. The large sample size and relative good observing cadence make it a good SLSN-I sample for statistical and detailed studies on SLSN-I light curves (*e.g.* precursor peak and post-peak bump features investigated in Paper II).

Based on this sample, we derive a bolometric correction relation for SLSNe-I, which allows a simple conversion between the optical g - and r -band photometry to the bolometric luminosity for low- z SLSNe-I. The bolometric correction follows, $\log(L_{\text{bolo}}/L_{gr}) = -1.093x^3 + 1.244x^2 - 0.261x + 0.410$, where x is the observed color $(g-r)$.

Our other findings are summarized as follows.

1. The rise time $t_{\text{rise},10\%}$ of our SLSN-I events has a mean value of 41.9 ± 17.8 days. Compared with SNe Ic, SLSNe-I have significantly longer rise times and wider dispersion (Nicholl et al. 2015; De Cia et al. 2018). In our sample, there is one fast-evolving SLSN-I, SN 2018gbv, with $t_{\text{rise},10\%}$ shorter than 10 days; while there are only three slowly-evolving events with $t_{\text{rise},10\%} > 78$ days, which is about 5% in our sample. Compared with previous samples, the observed ratio of fast-rising ($t_{\text{rise},1/e} \lesssim 15$ days) events in our sample is slightly higher, despite the target selection criteria disfavor the capture of fast-rising SNe.

We confirm that the $t_{\text{rise},10\%}$ shows a continuous distribution and can not be divided into two clearly detached subclasses (De Cia et al. 2018). And as proven by many studies (Nicholl et al. 2015, 2017; De Cia et al. 2018), the $1/e$ -maxima rise and decay time scales show a positive correlation, *i.e.* slowly-rising events tend to decay slowly,




Figure 14. Bolometric LCs for the events with at least three-epoch UV photometry. The black points present the bolometric luminosities constructed with both UV and optical photometry. The bolometric LCs derived from g - and r -band photometry are shown in solid lines with 1σ error marked by the shaded area. The breaks in LCs are due to the lack of data (*e.g.* SN 2019szu) or that the $(g-r)$ color is 0.1 mag beyond the range where the bolometric correction is reliable (*e.g.* SN 2018bgv).

and the decay time scales are about 1.4–1.6 times the rise time scales.

2. The observed $(g-r)$ color at peak has a median value of $-0.03^{+0.12}_{-0.11}$ mag and the rest-frame $(g-r)$ color $-0.21^{+0.19}_{-0.12}$ mag. The majority of our SLSNe-I follows a wide color trend that evolves from blue ($(g-r) \sim -0.3$ mag) at early phases to red, and reaches $(g-r) \sim +0.8$ mag at about 2 months after the peak. As [Inserra & Smartt \(2014\)](#) and [De Cia et al. \(2018\)](#) have proposed, we confirm that the peak rest-frame $(g-r)$ color is moderately corre-

lated with the g -band absolute peak magnitudes, *i.e.* brighter SLSNe-I tend to have bluer color.

3. SLSNe-I have a wide range of temperatures at any given phase from $t \sim -20$ to $+40$ days relative to the peak, especially at early phases, and the spread becomes smaller as SLSNe-I evolve. Based on the measurements from our sample and the PS1 sample, we suggest that the temperature of SLSNe-I at a given phase can have a continuous distribution.
4. We find four peculiar events that have extraordinary temperature and $(g-r)$ color evolution.




Figure 15. The derived bolometric LCs from g - and r -band photometry for our whole sample. Due to the LC quality, SN 2019fy and SN 2019aamu are excluded. The LCs are plotted in different colors, line styles and widths.




Figure 16. The distribution of the peak bolometric luminosities. The black dashed line and the gray region show the median value and the 1σ dispersion of $2.00^{+1.97}_{-1.44} \times 10^{44} \text{ erg s}^{-1}$. The black solid line shows the kernel density estimation of the distribution. Other SLSN-I samples are plotted in different hatching patterns for comparison.

SN 2018hti shows a notably high temperature and blue color. SN 2020fvm remains almost at a constant temperature of 10000 K for over 100 days and has a double-peak color evolution like its LC. SN 2019szu shows an unusual temperature evolution, which slowly rises from 13000 K at -20 day to 20000 K at $+170$ day. The color of SN 2019cdt turns red much faster than that of any other event.

5. The absolute peak magnitudes of our SLSN-I sample are $-22.8 \text{ mag} \leq M_g \leq -19.8 \text{ mag}$, with a median value and the 1σ error of $-21.48^{+1.13}_{-0.61} \text{ mag}$. On average, the peak M_g of SLSNe-I is around 4 mag and 3 mag brighter than normal SNeIc and SNeIc-BL, respectively. The peak bolometric luminosities of our sample distribute from 3.22×10^{43} to $7.80 \times 10^{44} \text{ erg s}^{-1}$ and have a median value with the 1σ dispersion of $2.00^{+1.97}_{-1.44} \times 10^{44} \text{ erg s}^{-1}$.

ACKNOWLEDGMENTS

We acknowledge very helpful discussions about data processing of P200 photometry with Shengyu Yan from Tsinghua University.

Based on observations obtained with the Samuel Oschin Telescope 48-inch and the 60-inch Telescope at the Palomar Observatory as part of the Zwicky Transient Facility project. ZTF is supported by the National Science Foundation under Grant No. AST-1440341

and a collaboration including Caltech, IPAC, the Weizmann Institute for Science, the Oskar Klein Center at Stockholm University, the University of Maryland, the University of Washington, Deutsches Elektronen-Synchrotron and Humboldt University, Los Alamos National Laboratories, the TANGO Consortium of Taiwan, the University of Wisconsin at Milwaukee, and Lawrence Berkeley National Laboratories. Operations are conducted by COO, IPAC, and UW. SED Machine is based upon work supported by the National Science Foundation under Grant No. 1106171. The ZTF forced-photometry service was funded under the Heising-Simons Foundation grant #12540303 (PI: Graham).

The Liverpool Telescope is operated on the island of La Palma by Liverpool John Moores University in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias with financial support from the UK Science and Technology Facilities Council. The Nordic Optical Telescope is owned in collaboration by the University of Turku and Aarhus University, and operated jointly by Aarhus University, the University of Turku and the University of Oslo, representing Denmark, Finland and Norway, the University of Iceland and Stockholm University at the Observatorio del Roque de los Muchachos, La Palma, Spain, of the Instituto de Astrofísica de Canarias. This research has made use of data obtained through the High Energy Astrophysics Science Archive Research Center Online Service, provided by the NASA/Goddard Space Flight Center. This work was supported by the GROWTH project funded by the National Science Foundation under Grant No. 1545949.

Z. Chen acknowledges support from the China Scholarship Council. T. K. acknowledges support from the Swedish National Space Agency and the Swedish Research Council. S. Schulze acknowledges support from the G.R.E.A.T research environment, funded by *Vetenskapsrådet*, the Swedish Research Council, project number 2016-06012. R. L. acknowledges support from a Marie Skłodowska-Curie Individual Fellowship within the Horizon 2020 European Union (EU) Framework Programme for Research and Innovation (H2020-MSCA-IF-2017-794467). A. Gal-Yam acknowledges support from the EU via ERC grant No. 725161, the ISF GW excellence center, an IMOS space infrastructure grant and BSF/Transformative and GIF grants, as well as the André Deloro Institute for Advanced Research in Space and Optics, the Schwartz/Reisman Collaborative Science Program and the Norman E Alexander Family M Foundation ULTRASAT Data Center Fund, Minerva and Yeda-Sela. The work of X. Wang

is supported by the National Natural Science Foundation of China (NSFC grants 12288102, 12033003 and 11633002), the Major State Basic Research Development Program (grant 2016YFA0400803), the Scholar Program of Beijing Academy of Science and Technology (DZ:BS202002), and the Tencent XPLOERER Prize.

Facilities: PO:1.2m, PO:1.5m, PO:Hale, Liverpool:2m, NOT:2.56m, Keck:I, WHT:4.2m

Software: Scikit-learn (Pedregosa et al. 2011), *FIREFLY* (Wilkinson et al. 2017), *george* (Ambikasaran et al. 2015), *HEASoft* (Nasa High Energy Astrophysics Science Archive Research Center 2014, <https://heasarc.gsfc.nasa.gov/lheasoft/download.html>), SEDM pipeline (Rigault et al. 2019), *pyraf-dbsp* (Bellm & Sesar 2016), *DBSP_DRP* (Roberson et al. 2022), *LPipe* (Perley et al. 2019), *Fpipe* (Fremling et al. 2016), *AutoPhOT* (Brennan & Fraser 2022).

APPENDIX

A. INFORMATION ON THE ZTF SLSN-I SAMPLE

Table A1. The ZTF SLSN-I Sample

ZTF Name	IAU Name	RA ^a	DEC ^a	Redshift	$E(B - V)^b$ (mag)	Discovery group	Template	Spectrum Phase ^c	Template Phase ^c	Template Type
ZTF18aaisyyp	SN 2018avk	13:11:27.72	+65:38:16.7	0.132	0.011	Gaia	PTF12gty	-13.5	-7	SLSN-I
ZTF18aajqcue	SN 2018don	13:55:08.65	+58:29:42.0	0.0735	0.0089	PS1	SN2007gr	-2.1	-7	Ic
ZTF18aapgrxo	SN 2018bym	18:43:13.42	+45:12:28.2	0.2744	0.0517	ATLAS	PTF13ajg	+5.4	+8	SLSN-I
ZTF18aavrmcg	SN 2018bgv	11:02:30.29	+55:35:55.8	0.0795	0.0074	Gaia	SN2011ke	+17.8	+20	SLSN-I
ZTF18aazgrfl	SN 2018lzv	12:44:02.32	+56:01:44.5	0.434	0.0081	ZTF	PTF09atu	-19.1	-20	SLSN-I
ZTF18abjwagv	SN 2018gbw	15:55:38.02	+28:21:38.0	0.3454	0.0386	PS1	PTF13ajg	-4.1	-4	SLSN-I
ZTF18abmasep	SN 2018fcg	21:09:36.78	+33:28:59.6	0.1011	0.1435	ZTF	PTF12dam	-4.4	-22	SLSN-I
ZTF18abrzcbp	SN 2018lzw	07:39:32.76	+27:44:02.7	0.3198	0.0373	ZTF	PTF10uhf	+27.6	+15	SLSN-I
ZTF18abshezu	SN 2018gft	23:57:17.95	-15:37:53.3	0.2320	0.0256	ZTF	PTF09cnd	-38.4	-14	SLSN-I
ZTF18abszecem	SN 2018lzx	22:29:27.24	+13:10:39.8	0.4373	0.0529	ZTF	PTF09cnd	+19.2	+37	SLSN-I
ZTF18abvgjyl	SN 2018gkz	07:58:11.54	+19:31:07.9	0.2405	0.036	ZTF	SN2007bi	+10.6	+50	SLSN-I
ZTF18acapyww	SN 2018hpb	18:28:41.24	+75:48:47.3	0.124	0.0868	Gaia	PTF10nmn	+25.0	max	SLSN-I
ZTF18acenqto	SN 2018ibb	04:38:56.93	-20:39:44.2	0.166	0.0275	ATLAS	PTF12dam	-4.0	+7	SLSN-I
ZTF18acqyvag	SN 2018lfe	09:33:29.56	+00:03:08.4	0.3505	0.0286	PS1	SN2011ke	+59.6	+53	SLSN-I
ZTF18acsplji	SN 2018hti	03:40:53.77	+11:46:37.9	0.0613	0.3983	ATLAS	PTF12dam	+21.2	+7	SLSN-I
ZTF18acxgqxq	SN 2018lfd	23:14:59.32	+48:45:27.6	0.2686	0.1506	ZTF	PTF12dam	-7.5	-22	SLSN-I
ZTF18acyxnyw	SN 2018kyt	12:27:56.23	+56:23:35.6	0.1080	0.0091	ZTF	PTF10hgi	+45.5	+47	SLSN-I
ZTF19aacxrab	SN 2019J	10:03:46.78	+06:46:24.7	0.1346	0.0229	PS1	PTF10gvb	+8.8	-6	SLSN-I
ZTF19aajwogx	SN 2019cca	12:02:50.91	-16:39:53.6	0.4103	0.0462	ZTF	PTF10uhf	+28.5	+15	SLSN-I
ZTF19aaknqmp	SN 2019bgu	09:57:15.34	+32:00:05.6	0.1480	0.0123	ATLAS	PTF12gty	+16.6	-7	SLSN-I
ZTF19aalbrph	SN 2019kwq	17:07:58.84	+58:42:03.9	0.49*	0.0257	ZTF	PTF09cnd	+68.9	+37	SLSN-I
ZTF19aamhast	SN 2019dgr	09:45:32.68	+04:56:02.2	0.3815	0.0348	ATLAS	PTF13ajg	+13.2	+8	SLSN-I
ZTF19aamhhiz	SN 2019kws	14:15:04.46	+50:39:06.8	0.1977	0.0142	ZTF	PTF12gty	+21.7	-7	SLSN-I
ZTF19aanesgt	SN 2019cdt	08:17:53.90	+65:28:46.7	0.153	0.0441	ZTF	PTF10nmn	+11.6	+29	SLSN-I
ZTF19aantokv	SN 2019aamp	14:37:49.27	+20:18:16.6	0.4040	0.0237	ZTF	PTF13ajg	+8.4	+8	SLSN-I
ZTF19aaohuwc	SN 2019dlr	11:17:34.18	+00:30:02.6	0.26*	0.0309	ZTF	SN2004aw	+57.7	+11	Ic
ZTF19aapaeye	SN 2019cwu	14:51:37.29	+48:59:13.7	0.32*	0.0186	ZTF	SN2011ke	+27.6	+20	SLSN-I
ZTF19aaqrime	SN 2019kwt	19:39:22.59	+78:45:43.8	0.3562	0.0775	ZTF	SN2007bi	+11.3	+50	SLSN-I
ZTF19aarpwhc	SN 2019eot	18:00:29.95	+50:17:43.3	0.3057	0.0366	ZTF	PTF13ajg	-13.8	-4	SLSN-I
ZTF19aaruijx	SN 2019kwu	13:57:39.77	+64:21:18.6	0.60*	0.0138	ZTF	PTF10vqv	+28.4	+11	SLSN-I
ZTF19aasdvfr	SN 2019gqi	14:21:11.98	+28:54:05.9	0.3642	0.0129	ATLAS	PTF11rks	+21.3	+7	SLSN-I
ZTF19aauiref	SN 2019fiy	14:05:46.73	+33:27:38.3	0.67*	0.0142	PS1	PTF09atu	+15.6	+28	SLSN-I
ZTF19aauvzyh	SN 2019gam	10:19:18.32	+17:12:42.6	0.1235	0.0272	ATLAS	SN2011ke	+17.6	+29	SLSN-I
ZTF19aavouyw	SN 2019gfm	15:35:46.59	+24:03:45.0	0.1816	0.0465	PS1	PTF11rks	+16.8	+18	SLSN-I
ZTF19aawfbtg	SN 2019hge	22:24:21.20	+24:47:17.1	0.0866	0.058	ZTF	PTF12dam	-25.7	+2	SLSN-I
ZTF19aawsqsc	SN 2019hno	19:39:12.95	+62:43:41.0	0.26	0.0583	ATLAS	PTF09cnd	+1.6	+9	SLSN-I
ZTF19aayclnm	SN 2019aamq	20:55:36.14	-08:40:31.4	0.386	0.0636	ZTF	PTF10bjp	+3.8	+8	SLSN-I
ZTF19abaeyqw	SN 2019kcy	14:08:19.78	+08:58:01.0	0.399	0.0227	ZTF	PTF13ajg	-19.3	-4	SLSN-I
ZTF19abcvwrz	SN 2019aamx	15:57:48.27	+27:28:03.5	0.41	0.0362	ZTF	PTF09cwl	+13.0	+22	SLSN-I
ZTF19abdlzyq	SN 2019aamr	15:29:23.55	+38:06:12.6	0.42*	0.0118	ZTF	PTF09cwl	+8.0	+22	SLSN-I
ZTF19abfvnns	SN 2019lsq	00:04:40.58	+42:52:11.3	0.1295	0.0829	ATLAS	PTF12dam	-3.3	-22	SLSN-I
ZTF19abkfhj	SN 2019otl	02:52:21.63	-17:48:12.4	0.500	0.0241	ZTF	PTF13ajg	+11.2	-4	SLSN-I
ZTF19abnacvf	SN 2019nhs	00:52:01.44	+07:36:59.7	0.189	0.0514	PS1	PTF13ajg	-8.6	-4	SLSN-I
ZTF19abnqqdp	SN 2019aams	23:43:36.16	+12:29:01.0	0.6360	0.0448	ZTF	PTF13ajg	+0.8	-4	SLSN-I
ZTF19abpbopt	SN 2019neq	17:54:26.76	+47:15:40.6	0.1060	0.0334	ZTF	PTF11rks	+16.1	+7	SLSN-I

Table A1 *continued*

Table A1 (*continued*)

ZTF Name	IAU Name	RA ^a	DEC ^a	Redshift	$E(B - V)^b$	Discovery	Template	Spectrum	Template	Template
					(mag)			group	Phase ^c	Phase ^c
ZTF19abrbrsvm	SN 2019obk	22:33:54.08	-02:09:42.3	0.1656	0.048	PS1	PTF12gty	+12.8	-7	SLSN-I
ZTF19abuolvj	SN 2019qgk	22:29:57.55	-04:06:02.2	0.3468	0.054	ZTF	SN2010gx	+47.2	+29	SLSN-I
ZTF19abuyuwa	SN 2019sgg	01:01:11.77	+14:01:35.4	0.5726	0.0373	ZTF	PTF13ajg	-18.6	-4	SLSN-I
ZTF19abzoyeg	SN 2019aamt	21:15:08.00	+32:43:01.3	0.2138	0.126	ZTF	PTF12dam	+15.2	+7	SLSN-I
ZTF19abzqmau	SN 2019sgh	01:12:39.42	+36:28:24.8	0.3436	0.0599	ZTF	PTF10uhf	+39.8	+15	SLSN-I
ZTF19acbonaa	SN 2019stc	06:54:23.10	+17:29:31.4	0.1178	0.072	ZTF	PTF10hgi	+25.6	+47	SLSN-I
ZTF19acfwynw	SN 2019szu	00:10:13.14	-19:41:32.4	0.2120	0.018	ATLAS	PTF12dam	-15.2	+2	SLSN-I
ZTF19acjjpg	SN 2019unb	09:47:57.02	+00:49:36.0	0.0635	0.1045	ZTF	PTF10hgi	-7.5	+15	SLSN-I
ZTF19ackjrru	SN 2019ujb	09:03:15.18	+40:14:32.6	0.2008	0.0165	ZTF	PTF11rks	+25.7	+18	SLSN-I
ZTF19acsajxn	SN 2019xdy	08:24:51.33	+22:10:46.0	0.2206	0.0425	ZTF	PTF10vwg	+31.6	+22	SLSN-I
ZTF19acucxij	SN 2019vvc	09:13:30.13	+44:46:26.2	0.3314	0.0115	ATLAS	PTF09cwl	+16.0	+22	SLSN-I
ZTF19acvxquk	SN 2019aamu	02:55:08.89	+11:27:22.4	0.2590	0.1925	ZTF	PTF12gty	+8.6	-7	SLSN-I
ZTF19adaivcf	SN 2019zbv	10:15:01.10	+43:24:53.6	0.3785	0.0119	ZTF	PTF13ajg	+10.4	+8	SLSN-I
ZTF20aahbfmf	SN 2020ank	08:16:14.65	+04:19:26.9	0.2485	0.0193	ZTF	PTF13ajg	-0.3	-4	SLSN-I
ZTF20aaifybu	SN 2020aau	16:34:12.51	+13:05:51.9	0.280	0.0565	ZTF	PTF13ajg	-2.4	+8	SLSN-I
ZTF19acujysi	SN 2019aamw	23:48:54.54	+24:59:59.8	0.22*	0.0526	ZTF	PTF10vwg	+137.6	+147	SLSN-I
ZTF20aadzbcf	SN 2020fvm	14:12:45.93	+34:44:16.2	0.2428	0.0129	ZTF	PTF10uhf	-1.4	+15	SLSN-I
ZTF20aagikvv	SN 2019aamv	12:45:01.65	+33:33:14.1	0.3996	0.0127	ZTF	PTF13ajg	-11.4	+8	SLSN-I
ZTF20aahrxgw	SN 2020aup	13:09:44.44	+12:29:13.4	0.31*	0.0208	ZTF	PTF09cwl	+16.2	+22	SLSN-I
ZTF20aaqwpo	SN 2020dlb	08:08:34.14	+34:44:12.9	0.398	0.0327	ZTF	SN2005ap	+0.2	+4	SLSN-I
ZTF20aapaecd	SN 2020fyq	14:46:10.44	+23:48:02.0	0.1765	0.0325	PS1	SN2011ke	+27.5	+53	SLSN-I
ZTF20aattyuz	SN 2020exj	14:42:40.01	+30:14:39.1	0.1216	0.0096	ATLAS	PTF10hgi	+44.2	+47	SLSN-I
ZTF20aaouudz	SN 2020htd	17:44:17.28	+38:55:30.4	0.3515	0.026	PS1	PTF09cnd	+7.4	+13	SLSN-I
ZTF20aavfbqz	SN 2020iyj	09:15:36.64	+53:27:32.0	0.3690	0.0155	ZTF	PTF09cwl	+1.9	+1	SLSN-I
ZTF20aavqrzc	SN 2020kox	11:06:04.97	+26:17:28.7	0.456	0.0166	ATLAS	PTF13ajg	+2.6	-4	SLSN-I
ZTF20aawfxdt	SN 2020jii	15:34:55.31	+02:51:11.5	0.396	0.0427	PS1	PTF13ajg	-1.8	-4	SLSN-I
ZTF20aawkgxa	SN 2020afah	10:20:18.32	+53:19:21.4	0.3754	0.0086	ZTF	PTF09cnd	-41.8	-16	SLSN-I
ZTF20abisijg	SN 2020afag	00:15:46.25	+47:00:08.5	0.3815	0.0898	ZTF	PTF09cwl	+6.7	+22	SLSN-I
ZTF20abjwrx	SN 2020onb	14:23:00.61	+49:10:40.7	0.16*	0.0208	ZTF	SN2004aw	+24.8	+4	Ic
ZTF20ablkuiu	SN 2020qef	22:56:10.53	+28:45:53.3	0.1831	0.0459	ATLAS	PTF12gty	+24.4	-7	SLSN-I
ZTF20abpuwxi	SN 2020rmv	00:40:00.19	-14:35:25.1	0.2621	0.0185	ATLAS	PTF10uhf	+13.2	+15	SLSN-I
ZTF20abzaacf	SN 2020xkv	22:37:46.00	+23:31:37.4	0.2410	0.0303	ATLAS	SN2011ke	-8.7	+20	SLSN-I
ZTF20aceqspy	SN 2020xgd	00:19:45.83	+05:08:18.7	0.455	0.0145	PS1	PTF13ajg	-4.4	-4	SLSN-I

^aIn J2000.^bFrom Schlafly & Finkbeiner (2011).^cIn rest-frame days.* means the redshift is fit by *superfit*.**Table A2.** Classification spectra

Name	Observing date	Telescope + instrument	Exposure time ^a (s)
SN 2018avk	2018-05-04	NOT+ALFOSC	1800
SN 2018don	2018-06-09	P200+DBSP	2 × 600/2 × 600
SN 2018bym	2018-06-12	P200+DBSP	600/600
SN 2018bgv	2018-06-04	NOT+ALFOSC	2 × 2400
SN 2018lzv	2018-07-14	WHT+ISIS	600
SN 2018gbw	2018-08-13	P200+DBSP	600/600
SN 2018fcg	2018-08-21	P200+DBSP	600/600
SN 2018lzw	2018-10-06	P200+DBSP	610/400

^aSlash-separated values indicate the exposure time of blue and red sides, respectively. (This table is available in its entirety in machine-readable form.)

Table A3. Photometry data

Name	MJD (days)	Filter	Mag(AB) (mag)	Error _{mag} (mag)	Label ^a	Flux _{ratio} ^b (10 ⁻⁹)	Tel.+Ins. ^c
SN 2018avk	58202.30	g	20.48	0.21	F	6.40	P48+ZTF
SN 2018avk	58202.32	g	20.64	0.18	F	5.53	P48+ZTF
SN 2018avk	58202.32	g	20.74	0.16	F	5.04	P48+ZTF
SN 2018avk	58202.34	g	20.78	0.19	F	4.86	P48+ZTF
SN 2018avk	58202.36	g	20.92	0.14	F	4.27	P48+ZTF
SN 2018avk	58202.36	g	20.80	0.21	F	4.77	P48+ZTF
SN 2018avk	58203.35	g	20.54	0.15	F	6.06	P48+ZTF
SN 2018avk	58204.39	g	20.10	0.19	F	9.09	P48+ZTF

^aF means real detection and T means upper limit.^bThe ratio of observed flux and flux zero point.^cTelescope+Instrument.

(This table is available in its entirety in machine-readable form.)

Table A4. LC properties

Name	K-correction ^a (mag)	M_{peak} (mag)	L_{peak} (10 ⁴³ erg s ⁻¹)	t_{peak} (MJD)	$t_{\text{rise},1/e}$ (days)	$t_{\text{decay},1/e}$ (days)	$t_{\text{rise},10\%}$ (days)
SN 2018avk	-0.00	-20.27 ± 0.03	5.5 ± 1.0	58257.29 ^{+5.45} _{-4.35}	41.94 ^{+4.84} _{-3.88}	76.83 ^{+4.70} _{-5.44}	50.27 ^{+4.99} _{-3.91}
SN 2018don	0.09	-20.05 ± 0.04	3.4 ± 0.6	58280.21 ^{+3.68} _{-3.49}	48.70 ^{+3.55} _{-3.29}	49.04 ^{+3.91} _{-4.31}	-
SN 2018bym	-0.22	-22.03 ± 0.03	33.2 ± 5.9	58274.14 ^{+1.82} _{-1.87}	24.05 ^{+2.33} _{-1.81}	43.10 ^{+1.50} _{-1.46}	-
SN 2018bgv	-0.09	-20.61 ± 0.10	(14.9 ± 3.6) ^b	58253.74 ^{+1.30} _{-1.76}	8.40 ^{+1.20} _{-1.63}	19.39 ^{+1.66} _{-1.22}	9.99 ^{+1.20} _{-1.63}
SN 2018lzv	-0.39	-22.23 ± 0.10	52.6 ± 16.1	58340.35 ^{+12.83} _{-9.63}	40.47 ^{+8.97} _{-6.75}	67.81 ^{+7.07} _{-9.56}	47.35 ^{+8.98} _{-6.75}
SN 2018gbw	-0.33	-22.10 ± 0.07	(41.1 ± 7.6)	58348.56 ^{+4.55} _{-3.23}	22.58 ^{+3.53} _{-2.62}	-	32.78 ^{+3.60} _{-2.77}
SN 2018fcg	-0.13	-20.35 ± 0.06	(6.0 ± 1.1)	58355.83 ^{+2.93} _{-2.54}	13.33 ^{+2.67} _{-2.32}	18.00 ^{+2.33} _{-2.68}	17.75 ^{+2.76} _{-2.35}
SN 2018lzw	-0.30*	(-21.68 ± 0.10)	(22.7 ± 4.6)	(58360.51 ^{+3.57} ₋₀)	-	(56.57 ^{+1.40} _{-2.99})	-
SN 2018gft	-0.18	-22.22 ± 0.02	40.1 ± 7.2	58420.27 ^{+7.66} _{-5.97}	37.70 ^{+6.22} _{-4.85}	-	49.95 ^{+6.26} _{-4.88}
SN 2018lzx	-0.44	-22.01 ± 0.04	32.6 ± 5.9	58437.41 ^{+11.79} _{-9.64}	60.45 ^{+8.24} _{-6.75}	108.79 ^{+10.04} _{-13.22}	76.50 ^{+8.55} _{-7.05}
SN 2018gkz	-0.24	(-21.81 ± 0.03)	(21.3 ± 4.3)	(58383.86 ^{+9.92} _{-9.14})	-	(86.65 ^{+8.11} _{-8.60})	-
SN 2018hpg	-0.13*	-20.10 ± 0.03	4.8 ± 0.9	58427.85 ^{+0.93} _{-1.31}	20.37 ^{+1.88} _{-3.47}	-	31.92 ^{+1.41} _{-1.99}
SN 2018lfe	-0.33*	-21.43 ± 0.09	27.4 ± 9.1	58465.45 ^{+3.90} _{-6.95}	17.12 ^{+3.21} _{-5.34}	26.03 ^{+6.25} _{-5.42}	-
SN 2018hti	-0.09	-22.09 ± 0.07	55.1 ± 14.9	58460.47 ^{+8.51} _{-5.88}	28.11 ^{+8.02} _{-5.55}	-	35.73 ^{+8.03} _{-5.55}
SN 2018lfd	-0.19	-21.98 ± 0.11	38.6 ± 8.5	58496.49 ^{+23.40} _{-9.15}	35.45 ^{+18.61} _{-7.50}	-	-
SN 2018kyl	-0.13	-20.51 ± 0.08	9.2 ± 0.4 * *	58506.63 ^{+4.10} _{-6.14}	29.07 ^{+3.75} _{-5.63}	26.42 ^{+5.69} _{-4.76}	33.35 ^{+3.70} _{-5.54}
SN 2019J	-0.05	-19.86 ± 0.13	3.7 ± 0.9	58546.97 ^{+9.61} _{-18.26}	44.98 ^{+8.70} _{-16.29}	30.58 ^{+16.26} _{-8.77}	54.62 ^{+8.60} _{-16.20}
SN 2019cca	-0.37*	-21.99 ± 0.10	(33.3 ± 7.8)	58559.82 ^{+6.71} _{-10.28}	35.36 ^{+10.88} _{-9.31}	39.88 ^{+9.58} _{-5.97}	-
SN 2019bgu	-0.14	-20.27 ± 0.15	5.5 ± 1.4	58559.92 ^{+5.53} _{-11.12}	-	24.74 ^{+9.98} _{-5.20}	-
SN 2019kwq	-0.43*	-21.99 ± 0.05	32.2 ± 5.8	58563.33 ^{+5.87} _{-8.38}	45.37 ^{+8.32} _{-8.52}	63.96 ^{+6.57} _{-5.02}	63.10 ^{+12.81} _{-10.97}
SN 2019dgr	-0.37	-21.89 ± 0.10	33.0 ± 9.8	58581.83 ^{+5.11} _{-7.84}	25.06 ^{+3.99} _{-5.84}	-	36.56 ^{+3.87} _{-6.08}
SN 2019kws	-0.31	-20.05 ± 0.04	5.1 ± 0.9	58573.97 ^{+4.64} _{-3.92}	22.70 ^{+3.97} _{-3.40}	53.26 ^{+3.36} _{-3.99}	-
SN 2019cdt	-0.04	-20.85 ± 0.05	7.6 ± 1.4	58583.63 ^{+2.01} _{-6.10}	17.59 ^{+1.89} _{-5.34}	14.24 ^{+5.29} _{-1.77}	26.66 ^{+5.70} _{-5.86}
SN 2019aamp	-0.40	-21.92 ± 0.04	30.5 ± 5.7	58588.27 ^{+6.54} _{-6.84}	-	43.93 ^{+5.98} _{-5.34}	-
SN 2019dlr	-0.24	-21.11 ± 0.07	18.1 ± 3.7	58593.30 ^{+6.51} _{-7.17}	28.54 ^{+5.67} _{-6.04}	42.00 ^{+9.39} _{-6.08}	38.58 ^{+8.43} _{-6.61}
SN 2019cwu	-0.30	-21.27 ± 0.04	17.8 ± 3.4	58599.56 ^{+4.14} _{-3.73}	17.43 ^{+3.15} _{-2.84}	48.64 ^{+3.14} _{-3.32}	20.92 ^{+3.20} _{-2.86}
SN 2019kwt	-0.34	-22.74 ± 0.05	65.1 ± 11.9	58650.73 ^{+5.03} _{-4.48}	51.21 ^{+4.08} _{-3.67}	49.09 ^{+4.94} _{-4.72}	70.50 ^{+8.91} _{-4.56}
SN 2019eot	-0.28	-22.13 ± 0.03	40.1 ± 7.1	58644.97 ^{+2.46} _{-2.33}	31.28 ^{+1.90} _{-1.80}	37.61 ^{+2.04} _{-2.11}	40.87 ^{+2.04} _{-1.90}
SN 2019kwu	-0.51*	-22.32 ± 0.11	50.8 ± 9.4	58620.64 ^{+5.06} _{-3.19}	14.54 ^{+3.18} _{-2.02}	27.67 ^{+2.62} _{-3.50}	20.29 ^{+3.27} _{-2.19}
SN 2019gqi	-0.34*	-21.43 ± 0.08	19.3 ± 4.8	58636.92 ^{+4.27} _{-5.71}	26.80 ^{+3.32} _{-4.30}	-	34.99 ^{+3.61} _{-4.53}
SN 2019fny	-0.69	-22.71 ± 0.03	-	58640.96 ^{+5.24} _{-5.82}	23.49 ^{+3.21} _{-3.54}	-	32.24 ^{+4.75} _{-4.20}
SN 2019gam	0.32	(-19.88 ± 0.12)	(2.6 ± 0.7)	(58648.17 ⁺⁰ _{-4.25})	(26.13 ^{+1.91} _{-4.49})	-	(38.28 ^{+1.07} _{-3.92})
SN 2019gfm	-0.02	-20.99 ± 0.05	13.7 ± 2.7	58645.17 ^{+3.58} _{-2.24}	13.80 ^{+3.19} _{-2.03}	26.18 ^{+2.32} _{-3.23}	20.53 ^{+3.51} _{-2.68}

Table A4 continued

Table A4 (continued)

Name	K-correction ^a (mag)	M_{peak} (mag)	L_{peak} (10^{43} erg s ⁻¹)	t_{peak} (MJD)	$t_{\text{rise},1/e}$ (days)	$t_{\text{decay},1/e}$ (days)	$t_{\text{rise},10\%}$ (days)
SN 2019hge	-0.11	-19.82 ± 0.05	3.6 ± 0.2 **	58693.95 ^{+3.00} _{-7.76}	46.63 ^{+2.79} _{-7.15}	39.35 ^{+7.26} _{-2.92}	57.21 ^{+14.29} _{-7.24}
SN 2019hno	-0.21	-21.05 ± 0.04	13.4 ± 2.5	58663.92 ^{+7.75} _{-4.02}	22.89 ^{+6.17} _{-3.24}	29.89 ^{+3.51} _{-6.25}	29.45 ^{+6.30} _{-3.30}
SN 2019aamq	-0.36	-21.76 ± 0.03	32.9 ± 11.8	58690.78 ^{+11.60} _{-10.21}	50.47 ^{+9.01} _{-7.92}	-	-
SN 2019kcy	-0.41	-22.12 ± 0.09	46.6 ± 11.8	58693.95 ^{+16.31} _{-11.69}	33.01 ^{+11.69} _{-8.40}	-	42.58 ^{+11.82} _{-8.45}
SN 2019aamx	-0.33	-21.80 ± 0.07	21.7 ± 4.8	58709.67 ^{+9.10} _{-12.60}	42.97 ^{+6.58} _{-9.03}	45.91 ^{+10.48} _{-8.31}	50.92 ^{+6.51} _{-8.98}
SN 2019aamr	-0.41	-21.36 ± 0.10	20.5 ± 4.3	58684.58 ^{+6.37} _{-3.66}	14.04 ^{+4.50} _{-2.62}	27.09 ^{+3.89} _{-5.38}	17.70 ^{+4.50} _{-2.60}
SN 2019lsq	-0.22	-20.87 ± 0.04	12.5 ± 2.3	58707.77 ^{+1.48} _{-1.44}	21.29 ^{+1.36} _{-1.33}	43.02 ^{+1.50} _{-1.80}	31.31 ^{+1.65} _{-1.46}
SN 2019otl	-0.46	-22.41 ± 0.06	(61.2 ± 14.4)	58715.21 ^{+12.61} _{-13.82}	-	-	-
SN 2019nhs	-0.05	-21.44 ± 0.06	18.6 ± 0.9 **	58732.19 ^{+3.96} _{-4.79}	22.77 ^{+3.36} _{-4.06}	27.57 ^{+4.16} _{-3.42}	30.59 ^{+3.70} _{-4.18}
SN 2019aams	-0.60	-22.11 ± 0.08	(47.2 ± 13.6)	58730.76 ^{+8.27} _{-4.61}	25.18 ^{+6.30} _{-3.55}	23.86 ^{+3.23} _{-5.82}	33.92 ^{+7.83} _{-5.12}
SN 2019neq	-0.24	-21.30 ± 0.03	20.1 ± 0.4 **	58730.16 ^{+0.80} _{-0.83}	14.69 ^{+0.73} _{-0.76}	19.97 ^{+0.88} _{-0.83}	21.97 ^{+0.74} _{-0.76}
SN 2019obk	-0.10	-19.80 ± 0.11	3.8 ± 1.1	58735.13 ^{+10.43} _{-10.59}	37.49 ^{+10.91} _{-10.12}	-	56.10 ^{+10.52} _{-12.16}
SN 2019qgk	-0.32*	(-21.55 ± 0.11)	(21.2 ± 5.5)	(58744.36 ^{+9.70} _{-7.55})	-	(43.96 ^{+3.91} _{-9.03})	-
SN 2019sgg	-0.42	(-22.32 ± 0.07)	56.6 ± 12.0	(58791.33 ⁺⁰ _{-8.53})	(42.77 ^{+0.95} _{-5.51})	-	(51.83 ^{+1.60} _{-5.61})
SN 2019aamt	-0.24	-20.74 ± 0.05	17.0 ± 7.1	58773.51 ^{+6.45} _{-7.02}	24.61 ^{+5.32} _{-5.79}	-	33.31 ^{+5.37} _{-5.82}
SN 2019sgh	-0.34	-21.32 ± 0.05	19.3 ± 3.5	58758.54 ^{+7.89} _{-3.86}	13.96 ^{+5.89} _{-2.90}	32.03 ^{+6.33} _{-10.59}	18.16 ^{+6.34} _{-3.15}
SN 2019stc	-0.02	-20.55 ± 0.06	5.8 ± 1.0	58783.42 ^{+12.67} _{-5.51}	26.13 ^{+12.49} _{-5.24}	38.22 ^{+6.25} _{-11.77}	-
SN 2019szu	-0.06	-21.19 ± 0.06	22.3 ± 5.2	58829.39 ^{+6.34} _{-11.76}	54.57 ^{+11.65} _{-11.07}	-	-
SN 2019unb	-0.03	-20.13 ± 0.07	4.1 ± 0.1 **	58843.01 ^{+4.50} _{-4.87}	-	20.80 ^{+4.64} _{-4.30}	-
SN 2019ujb	-0.05	-21.13 ± 0.08	12.2 ± 2.6	58818.10 ^{+5.57} _{-7.89}	26.06 ^{+4.67} _{-6.58}	50.81 ^{+6.58} _{-4.66}	35.07 ^{+8.06} _{-6.93}
SN 2019xdy	-0.22*	-20.72 ± 0.06	7.9 ± 4.2	58833.43 ^{+3.14} _{-5.67}	22.84 ^{+6.08} _{-8.07}	32.42 ^{+4.65} _{-2.59}	37.94 ^{+3.82} _{-6.22}
SN 2019vvc	-0.31*	-22.06 ± 0.02	44.7 ± 18.1	58859.72 ^{+3.61} _{-3.91}	40.06 ^{+2.86} _{-3.09}	-	53.55 ^{+5.18} _{-4.85}
SN 2019aamu	-0.25*	(-21.64 ± 0.07)	-	(58861.15 ⁺⁰ _{-7.55})	(45.20 ^{+1.81} _{-6.33})	-	(65.18 ^{+1.02} _{-6.05})
SN 2019zvb	-0.39	-21.98 ± 0.04	33.4 ± 6.0	58866.72 ^{+3.55} _{-3.40}	26.50 ^{+2.77} _{-2.60}	54.00 ^{+3.13} _{-3.16}	-
SN 2020ank	-0.22	-21.75 ± 0.06	24.6 ± 4.5	58893.40 ^{+2.91} _{-2.06}	15.56 ^{+2.37} _{-1.69}	32.76 ^{+2.06} _{-3.09}	22.81 ^{+3.48} _{-2.41}
SN 2020auv	-0.21	-21.85 ± 0.09	31.2 ± 6.7	58884.04 ^{+5.22} _{-6.59}	-	26.77 ^{+5.23} _{-4.16}	-
SN 2019aamw	-0.22*	-20.49 ± 0.07	6.4 ± 1.3	58854.17 ^{+8.75} _{-11.51}	43.66 ^{+7.27} _{-9.51}	57.18 ^{+18.25} _{-11.62}	63.85 ^{+7.30} _{-9.62}
SN 2020fvm	-0.19	-21.39 ± 0.05	15.1 ± 2.7	58999.69 ^{+4.88} _{-2.82}	91.12 ^{+4.02 c} _{-2.41}	-	-
SN 2019aamv	-0.38	-21.37 ± 0.06	19.6 ± 3.9	58919.90 ^{+6.07} _{-3.72}	51.73 ^{+5.17} _{-3.39}	58.20 ^{+4.38} _{-5.00}	66.80 ^{+19.86} _{-6.96}
SN 2020aup	-0.28	-21.48 ± 0.12	24.6 ± 4.8	58882.70 ^{+5.31} _{-4.11}	14.87 ^{+4.18} _{-3.27}	28.44 ^{+4.82} _{-4.58}	20.82 ^{+4.66} _{-3.70}
SN 2020dlb	-0.36*	-22.61 ± 0.07	78.0 ± 23.9	58948.76 ^{+3.86} _{-3.88}	30.31 ^{+2.78} _{-2.80}	32.26 ^{+3.08} _{-3.00}	41.57 ^{+3.21} _{-2.93}
SN 2020fyq	-0.18*	-19.89 ± 0.06	3.2 ± 0.6	58965.65 ^{+4.40} _{-33.19}	70.06 ^{+4.04} _{-28.25}	62.83 ^{+28.28} _{-4.70}	86.77 ^{+4.71} _{-28.37}
SN 2020exj	-0.22	-20.08 ± 0.07	5.2 ± 1.0	58946.37 ^{+3.63} _{-2.43}	19.70 ^{+3.51} _{-2.53}	21.99 ^{+3.44} _{-3.24}	26.57 ^{+3.44} _{-2.61}
SN 2020htd	-0.33*	-21.22 ± 0.05	14.3 ± 2.6	58987.98 ^{+4.85} _{-4.94}	50.90 ^{+4.35} _{-4.93}	47.26 ^{+5.44} _{-3.99}	65.69 ^{+4.72} _{-4.30}
SN 2020iyj	-0.35	-21.73 ± 0.03	23.3 ± 4.3	58995.41 ^{+6.73} _{-7.58}	30.48 ^{+4.96} _{-5.57}	68.36 ^{+17.54} _{-27.95}	39.22 ^{+5.07} _{-5.65}
SN 2020kox	-0.36	-21.78 ± 0.11	(44.6 ± 12.7)	58994.23 ^{+9.02} _{-8.89}	26.52 ^{+6.30} _{-6.20}	-	34.57 ^{+6.46} _{-6.33}
SN 2020jii	-0.40	-21.86 ± 0.07	44.2 ± 13.9	59000.44 ^{+4.02} _{-3.77}	30.36 ^{+3.27} _{-3.00}	32.84 ^{+2.83} _{-3.01}	39.44 ^{+3.27} _{-3.22}
SN 2020afah	-0.35*	-21.84 ± 0.05	25.0 ± 5.3	59055.47 ^{+19.11} _{-7.13}	61.40 ^{+13.90} _{-5.20}	54.31 ^{+6.56} _{-17.50}	69.96 ^{+13.91} _{-5.21}
SN 2020afag	-0.37	-21.67 ± 0.04	23.6 ± 4.3	59065.68 ^{+3.48} _{-2.84}	36.29 ^{+3.23} _{-2.59}	49.05 ^{+3.39} _{-4.45}	49.71 ^{+6.04} _{-4.63}
SN 2020onb	-0.11	-21.07 ± 0.12	9.8 ± 1.8	59078.29 ^{+2.54} _{-5.48}	28.91 ^{+2.25} _{-4.75}	-	40.04 ^{+2.32} _{-4.78}
SN 2020qef	-0.26	-20.03 ± 0.05	4.3 ± 0.8	59083.09 ^{+2.01} _{-2.18}	36.52 ^{+1.99} _{-2.06}	31.05 ^{+9.62} _{-2.30}	47.82 ^{+2.28} _{-2.37}
SN 2020rmv	-0.18	-21.60 ± 0.05	22.1 ± 3.9	59117.41 ^{+7.57} _{-6.28}	39.81 ^{+6.23} _{-5.49}	-	54.13 ^{+7.81} _{-5.25}
SN 2020xkv	-0.36	-21.02 ± 0.02	10.7 ± 2.0	59144.80 ^{+9.81} _{-8.07}	46.48 ^{+7.94} _{-6.55}	-	71.62 ^{+8.08} _{-6.72}
SN 2020xgd	-0.45	-21.83 ± 0.09	34.7 ± 6.7	59149.39 ^{+4.78} _{-3.70}	23.67 ^{+3.31} _{-2.58}	21.92 ^{+4.10} _{-3.77}	29.58 ^{+3.85} _{-2.78}

M_{peak} , t_{peak} , $t_{\text{rise},1/e}$, $t_{\text{decay},1/e}$ and $t_{\text{rise},10\%}$ are measured in the rest-frame g -band LCs.

^aK-correction to rest-frame g -band at the peak. It equals $K_{g \rightarrow g}$ for $z \leq 0.17$ and $K_{r \rightarrow g}$ for $z > 0.17$.

^bThe value in parentheses means they are not well-constrained due to the lack of data.

^cSN 2020fvm has two LC peaks and its $t_{\text{rise},1/e}$ is 47.52^{+6.66}_{-6.88} days if the first (and the fainter) one is set as the main peak.

* means the K-correction is calculated by $-2.5 \times \log(1+z)$.

** means the L_{peak} is calculated from multi-band data instead of corrected from g - and r -band flux.

Table A5. $(g - r)$ colors at peak

Name	Observed (mag)	Rest frame (mag)
SN 2018avk	-0.03 ± 0.03	-0.21 ± 0.03
SN 2018don	0.15 ± 0.02	-0.01 ± 0.02
SN 2018bym	-0.11 ± 0.01	-0.32 ± 0.06
SN 2018bgv	-0.23 ± 0.03	-0.25 ± 0.03
SN 2018lzv	-0.03 ± 0.07	-
SN 2018gbw	-0.12 ± 0.04	-
SN 2018fcg	-0.16 ± 0.04	-0.18 ± 0.04
SN 2018gft	-0.14 ± 0.03	-0.21 ± 0.05
SN 2018lzx	0.02 ± 0.04	-
SN 2018hpq	-0.11 ± 0.02	-

(This table is available in its entirety in machine-readable form.)

Table A6. Temperatures and bolometric luminosities

Name	Phase (days)	Temperature (10^4 K)	L_{bolo} (10^{43} erg s $^{-1}$)	Radius (10^{10} km)
SN 2018avk	-4.23	$1.05^{+0.15}_{-0.11}$	$5.14^{+0.58}_{-0.56}$	$2.46^{+0.51}_{-0.71}$
SN 2018avk	16.97	$0.74^{+0.05}_{-0.04}$	$4.24^{+0.43}_{-0.40}$	$4.39^{+0.54}_{-0.63}$
SN 2018bym	1.06	$1.21^{+0.07}_{-0.06}$	$29.67^{+1.13}_{-1.24}$	$4.43^{+0.42}_{-0.50}$
SN 2018bym	9.70	$1.14^{+0.13}_{-0.10}$	$25.89^{+2.09}_{-2.29}$	$4.66^{+0.81}_{-1.08}$
SN 2018bym	33.24	$0.86^{+0.03}_{-0.03}$	$12.23^{+0.58}_{-0.57}$	$5.64^{+0.44}_{-0.46}$
SN 2018bgv	6.26	$1.39^{+0.32}_{-0.19}$	$7.44^{+0.68}_{-0.64}$	$1.67^{+0.46}_{-0.77}$
SN 2018bgv	9.04	$1.17^{+0.10}_{-0.09}$	$5.98^{+0.34}_{-0.34}$	$2.12^{+0.32}_{-0.37}$
SN 2018bgv	11.82	$0.94^{+0.12}_{-0.09}$	$4.47^{+0.50}_{-0.51}$	$2.83^{+0.55}_{-0.73}$
SN 2018bgv	17.38	$0.73^{+0.57}_{-0.09}$	$3.43^{+0.84}_{-0.70}$	$4.07^{+1.15}_{-6.33}$
SN 2018bgv	20.16	$0.64^{+0.05}_{-0.04}$	$2.64^{+0.34}_{-0.31}$	$4.63^{+0.68}_{-0.82}$
SN 2018bgv	22.94	$0.61^{+0.18}_{-0.05}$	$2.29^{+0.42}_{-0.49}$	$4.87^{+0.85}_{-2.98}$

(This table is available in its entirety in machine-readable form.)




Figure A1. The LCs of our sample. All magnitudes are in the AB system and have been corrected for Milky Way extinction. The absolute magnitude y-axis on the right side is calculated by assuming a constant K-correction of $-2.5 \times \log(1+z)$. The rings mark the data measured from the combined image of multiple nearby images. The symbol ‘S’ at the top of each panel shows the epoch of the closest spectrum to the peak, which is used to calculate accurate K-corrections and color corrections. X-axis shows the rest-frame days relative to the observed *g*-band peak. SN 2018gbw, SN 2019cca, SN 2019fiy, SN 2019xdy, SN 2019vvc, SN 2019aamu and SN 2020ank are relative to *r*-band peak.




Figure A2. Similar to Figure A1.




Figure A3. Similar to Figure A1.




Figure A4. Similar to Figure A1.




Figure A5. Similar to Figure A1.




Figure A6. Similar to Figure A1.




Figure A7. Spectral classification. This figure shows the comparison between our spectrum and their best-matched templates (shown in gray) from *superfit*. Event name, the phase of our spectrum (rest-frame days), the best-matched templates, the template phase and the template type are labeled after each spectrum in the same color.




Figure A8. Similar to Figure A7




Figure A9. Similar to Figure A7

REFERENCES

- Ambikasaran, S., Foreman-Mackey, D., Greengard, L., Hogg, D. W., & O’Neil, M. 2015, *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 38, 252, doi: [10.1109/TPAMI.2015.2448083](https://doi.org/10.1109/TPAMI.2015.2448083)
- Ambikasaran, S., Foreman-Mackey, D., Greengard, L., Hogg, D. W., & O’Neil, M. 2016, *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 38, 252, doi: [10.1109/TPAMI.2015.2448083](https://doi.org/10.1109/TPAMI.2015.2448083)
- Anderson, J. P., Pessi, P. J., Dessart, L., et al. 2018, ArXiv e-prints. <https://arxiv.org/abs/1806.10609>
- Angus, C. R., Levan, A. J., Perley, D. A., et al. 2016, *MNRAS*, 458, 84, doi: [10.1093/mnras/stw063](https://doi.org/10.1093/mnras/stw063)
- Angus, C. R., Smith, M., Sullivan, M., et al. 2019, *Monthly Notices of the Royal Astronomical Society*, 487, 2215–2241, doi: [10.1093/mnras/stz1321](https://doi.org/10.1093/mnras/stz1321)
- Barbarino, C., Sollerman, J., Taddia, F., et al. 2021, *A&A*, 651, A81, doi: [10.1051/0004-6361/202038890](https://doi.org/10.1051/0004-6361/202038890)
- Bellm, E. C., & Sesar, B. 2016, pyraf-dbsp: Reduction pipeline for the Palomar Double Beam Spectrograph. <http://ascl.net/1602.002>
- Bellm, E. C., Kulkarni, S. R., Graham, M. J., et al. 2019a, *PASP*, 131, 018002, doi: [10.1088/1538-3873/aaecbe](https://doi.org/10.1088/1538-3873/aaecbe)
- Bellm, E. C., Kulkarni, S. R., Barlow, T., et al. 2019b, *PASP*, 131, 068003, doi: [10.1088/1538-3873/ab0c2a](https://doi.org/10.1088/1538-3873/ab0c2a)
- Blagorodnova, N., Neill, J. D., Walters, R., et al. 2018, *PASP*, 130, 035003, doi: [10.1088/1538-3873/aaa53f](https://doi.org/10.1088/1538-3873/aaa53f)
- Blondin, S., & Tonry, J. L. 2007, *ApJ*, 666, 1024, doi: [10.1086/520494](https://doi.org/10.1086/520494)
- Brennan, S. J., & Fraser, M. 2022, arXiv e-prints, arXiv:2201.02635. <https://arxiv.org/abs/2201.02635>
- Chambers, K. C., Magnier, E. A., Metcalfe, N., et al. 2016, arXiv e-prints, arXiv:1612.05560. <https://arxiv.org/abs/1612.05560>
- Chatzopoulos, E., Wheeler, J. C., Vinko, J., Horvath, Z. L., & Nagy, A. 2013, *ApJ*, 773, 76, doi: [10.1088/0004-637X/773/1/76](https://doi.org/10.1088/0004-637X/773/1/76)
- Chen, T.-W., Smartt, S. J., Yates, R. M., et al. 2017a, *MNRAS*, 470, 3566, doi: [10.1093/mnras/stx1428](https://doi.org/10.1093/mnras/stx1428)
- Chen, T.-W., Schady, P., Xiao, L., et al. 2017b, *ApJL*, 849, L4, doi: [10.3847/2041-8213/aa8f40](https://doi.org/10.3847/2041-8213/aa8f40)
- Chen, Z. H., Yan, L., Kangas, T., et al. 2022, arXiv e-prints, arXiv:2202.02060. <https://arxiv.org/abs/2202.02060>
- Chevalier, R. A., & Irwin, C. M. 2011, *ApJL*, 729, L6, doi: [10.1088/2041-8205/729/1/L6](https://doi.org/10.1088/2041-8205/729/1/L6)
- De, K., Kasliwal, M. M., Tzanidakis, A., et al. 2020, *ApJ*, 905, 58, doi: [10.3847/1538-4357/abb45c](https://doi.org/10.3847/1538-4357/abb45c)
- De Cia, A., Gal-Yam, A., Rubin, A., et al. 2018, *ApJ*, 860, 100, doi: [10.3847/1538-4357/aab9b6](https://doi.org/10.3847/1538-4357/aab9b6)
- Dekany, R., Smith, R. M., Riddle, R., et al. 2020, *PASP*, 132, 038001, doi: [10.1088/1538-3873/ab4ca2](https://doi.org/10.1088/1538-3873/ab4ca2)
- Dessart, L. 2019, *A&A*, 621, A141, doi: [10.1051/0004-6361/201834535](https://doi.org/10.1051/0004-6361/201834535)
- Drake, A. J., Djorgovski, S. G., Mahabal, A., et al. 2009, *ApJ*, 696, 870, doi: [10.1088/0004-637X/696/1/870](https://doi.org/10.1088/0004-637X/696/1/870)
- Drout, M. R., Chornock, R., Soderberg, A. M., et al. 2014, *ApJ*, 794, 23, doi: [10.1088/0004-637X/794/1/23](https://doi.org/10.1088/0004-637X/794/1/23)
- Duev, D. A., Mahabal, A., Masci, F. J., et al. 2019, *MNRAS*, 489, 3582, doi: [10.1093/mnras/stz2357](https://doi.org/10.1093/mnras/stz2357)
- Filippenko, A. V. 1997, *ARA&A*, 35, 309, doi: [10.1146/annurev.astro.35.1.309](https://doi.org/10.1146/annurev.astro.35.1.309)
- Fitzpatrick, E. L., & Massa, D. 2007, *The Astrophysical Journal*, 663, 320–341, doi: [10.1086/518158](https://doi.org/10.1086/518158)
- Fremling, C., Sollerman, J., Taddia, F., et al. 2016, *A&A*, 593, A68, doi: [10.1051/0004-6361/201628275](https://doi.org/10.1051/0004-6361/201628275)
- Fremling, C., Miller, A. A., Sharma, Y., et al. 2020, *ApJ*, 895, 32, doi: [10.3847/1538-4357/ab8943](https://doi.org/10.3847/1538-4357/ab8943)
- Frohmaier, C., Angus, C. R., Vincenzi, M., et al. 2021, *MNRAS*, 500, 5142, doi: [10.1093/mnras/staa3607](https://doi.org/10.1093/mnras/staa3607)
- Gal-Yam, A. 2012, *Science*, 337, 927, doi: [10.1126/science.1203601](https://doi.org/10.1126/science.1203601)
- . 2019, *ARA&A*, 57, 305, doi: [10.1146/annurev-astro-081817-051819](https://doi.org/10.1146/annurev-astro-081817-051819)
- Gal-Yam, A., Mazzali, P., Ofek, E. O., et al. 2009, *Nature*, 462, 624, doi: [10.1038/nature08579](https://doi.org/10.1038/nature08579)
- Gehrels, N. 1986, *ApJ*, 303, 336, doi: [10.1086/164079](https://doi.org/10.1086/164079)
- Gehrels, N., Chincarini, G., Giommi, P., et al. 2004, *ApJ*, 611, 1005, doi: [10.1086/422091](https://doi.org/10.1086/422091)
- Gezari, S., Halpern, J. P., Grupe, D., et al. 2009, *ApJ*, 690, 1313, doi: [10.1088/0004-637X/690/2/1313](https://doi.org/10.1088/0004-637X/690/2/1313)
- Graham, M. J., Kulkarni, S. R., Bellm, E. C., et al. 2019, *PASP*, 131, 078001, doi: [10.1088/1538-3873/ab006c](https://doi.org/10.1088/1538-3873/ab006c)
- Ho, A. Y. Q., Perley, D. A., Gal-Yam, A., et al. 2021, arXiv e-prints, arXiv:2105.08811. <https://arxiv.org/abs/2105.08811>
- Hodgkin, S. T., Harrison, D. L., Breedt, E., et al. 2021, *A&A*, 652, A76, doi: [10.1051/0004-6361/202140735](https://doi.org/10.1051/0004-6361/202140735)
- Hogg, D. W., Baldry, I. K., Blanton, M. R., & Eisenstein, D. J. 2002, ArXiv Astrophysics e-prints
- Howell, D. A., Sullivan, M., Nugent, P. E., et al. 2006, *Nature*, 443, 308, doi: [10.1038/nature05103](https://doi.org/10.1038/nature05103)
- Inserra, C., Prajs, S., Gutierrez, C. P., et al. 2018a, *The Astrophysical Journal*, 854, 175, doi: [10.3847/1538-4357/aaaaaa](https://doi.org/10.3847/1538-4357/aaaaaa)
- Inserra, C., & Smartt, S. J. 2014, *ApJ*, 796, 87, doi: [10.1088/0004-637X/796/2/87](https://doi.org/10.1088/0004-637X/796/2/87)

- Inserra, C., Smartt, S. J., Gall, E. E. E., et al. 2018b, *Monthly Notices of the Royal Astronomical Society*, 475, 1046–1072, doi: [10.1093/mnras/stx3179](https://doi.org/10.1093/mnras/stx3179)
- Inserra, C., Sullivan, M., Angus, C. R., et al. 2021, *MNRAS*, 504, 2535, doi: [10.1093/mnras/stab978](https://doi.org/10.1093/mnras/stab978)
- Kangas, T., Blagorodnova, N., Mattila, S., et al. 2017, *MNRAS*, 469, 1246, doi: [10.1093/mnras/stx833](https://doi.org/10.1093/mnras/stx833)
- Kangas, T., Yan, L., Schulze, S., et al. 2022, *MNRAS*, doi: [10.1093/mnras/stac2218](https://doi.org/10.1093/mnras/stac2218)
- Kasen, D. 2017, *Unusual Supernovae and Alternative Power Sources*, ed. A. W. Alsabti & P. Murdin, 939, doi: [10.1007/978-3-319-21846-5_32](https://doi.org/10.1007/978-3-319-21846-5_32)
- Kasen, D., & Bildsten, L. 2010, *ApJ*, 717, 245, doi: [10.1088/0004-637X/717/1/245](https://doi.org/10.1088/0004-637X/717/1/245)
- Kasen, D., Woosley, S. E., & Heger, A. 2011, *ApJ*, 734, 102, doi: [10.1088/0004-637X/734/2/102](https://doi.org/10.1088/0004-637X/734/2/102)
- Kasliwal, M. M., Cannella, C., Bagdasaryan, A., et al. 2019, *PASP*, 131, 038003, doi: [10.1088/1538-3873/aafbc2](https://doi.org/10.1088/1538-3873/aafbc2)
- Law, N. M., Kulkarni, S. R., Dekany, R. G., et al. 2009, *PASP*, 121, 1395, doi: [10.1086/648598](https://doi.org/10.1086/648598)
- Leloudas, G., Schulze, S., Krühler, T., et al. 2015, *MNRAS*, 449, 917, doi: [10.1093/mnras/stv320](https://doi.org/10.1093/mnras/stv320)
- Lin, W. L., Wang, X. F., Li, W. X., et al. 2020, *MNRAS*, 497, 318, doi: [10.1093/mnras/staa1918](https://doi.org/10.1093/mnras/staa1918)
- Lunnan, R., Chornock, R., Berger, E., et al. 2014, *ApJ*, 787, 138, doi: [10.1088/0004-637X/787/2/138](https://doi.org/10.1088/0004-637X/787/2/138)
- . 2015, *ApJ*, 804, 90, doi: [10.1088/0004-637X/804/2/90](https://doi.org/10.1088/0004-637X/804/2/90)
- . 2018, *ApJ*, 852, 81, doi: [10.3847/1538-4357/aa9fla](https://doi.org/10.3847/1538-4357/aa9fla)
- Lunnan, R., Yan, L., Perley, D. A., et al. 2020, *ApJ*, 901, 61, doi: [10.3847/1538-4357/abaeeec](https://doi.org/10.3847/1538-4357/abaeeec)
- Lyman, J. D., Bersier, D., & James, P. A. 2013, *Monthly Notices of the Royal Astronomical Society*, 437, 3848–3862, doi: [10.1093/mnras/stt2187](https://doi.org/10.1093/mnras/stt2187)
- Margutti, R., Metzger, B. D., Chornock, R., et al. 2017, *ApJ*, 836, 25, doi: [10.3847/1538-4357/836/1/25](https://doi.org/10.3847/1538-4357/836/1/25)
- Masci, F. J., Laher, R. R., Rusholme, B., et al. 2019, *PASP*, 131, 018003, doi: [10.1088/1538-3873/aae8ac](https://doi.org/10.1088/1538-3873/aae8ac)
- McCrum, M., Smartt, S. J., Rest, A., et al. 2015, *MNRAS*, 448, 1206, doi: [10.1093/mnras/stv034](https://doi.org/10.1093/mnras/stv034)
- Miller, A. A., Chornock, R., Perley, D. A., et al. 2009, *ApJ*, 690, 1303, doi: [10.1088/0004-637X/690/2/1303](https://doi.org/10.1088/0004-637X/690/2/1303)
- Miller, A. A., Silverman, J. M., Butler, N. R., et al. 2010, *MNRAS*, 404, 305, doi: [10.1111/j.1365-2966.2010.16280.x](https://doi.org/10.1111/j.1365-2966.2010.16280.x)
- Nasa High Energy Astrophysics Science Archive Research Center. 2014, HEASoft: Unified Release of FTOOLS and XANADU. <http://ascl.net/1408.004>
- Nicholl, M., Guillochon, J., & Berger, E. 2017, *ArXiv e-prints*. <https://arxiv.org/abs/1706.00825>
- Nicholl, M., Smartt, S. J., Jerkstrand, A., et al. 2015, *MNRAS*, 452, 3869, doi: [10.1093/mnras/stv1522](https://doi.org/10.1093/mnras/stv1522)
- Ofek, E. O., Cameron, P. B., Kasliwal, M. M., et al. 2007, *ApJL*, 659, L13, doi: [10.1086/516749](https://doi.org/10.1086/516749)
- Oke, J. B., & Gunn, J. E. 1983, *ApJ*, 266, 713, doi: [10.1086/160817](https://doi.org/10.1086/160817)
- Oke, J. B., Cohen, J. G., Carr, M., et al. 1995, *PASP*, 107, 375, doi: [10.1086/133562](https://doi.org/10.1086/133562)
- Osterbrock, D. E., & Ferland, G. J. 2006, *Astrophysics of gaseous nebulae and active galactic nuclei*
- Pastorello, A., Smartt, S. J., Botticella, M. T., et al. 2010, *ApJL*, 724, L16, doi: [10.1088/2041-8205/724/1/L16](https://doi.org/10.1088/2041-8205/724/1/L16)
- Pedregosa, F., Varoquaux, G., Gramfort, A., et al. 2011, *Journal of Machine Learning Research*, 12, 2825
- Perley, D. A., Quimby, R. M., Yan, L., et al. 2016, *ApJ*, 830, 13, doi: [10.3847/0004-637X/830/1/13](https://doi.org/10.3847/0004-637X/830/1/13)
- Perley, D. A., Mazzali, P. A., Yan, L., et al. 2019, *MNRAS*, 484, 1031, doi: [10.1093/mnras/sty3420](https://doi.org/10.1093/mnras/sty3420)
- Perley, D. A., Fremling, C., Sollerman, J., et al. 2020, *ApJ*, 904, 35, doi: [10.3847/1538-4357/abbd98](https://doi.org/10.3847/1538-4357/abbd98)
- Pignata, G., Benetti, S., Mazzali, P. A., et al. 2008, *MNRAS*, 388, 971, doi: [10.1111/j.1365-2966.2008.13434.x](https://doi.org/10.1111/j.1365-2966.2008.13434.x)
- Prajs, S., Sullivan, M., Smith, M., et al. 2017, *MNRAS*, 464, 3568, doi: [10.1093/mnras/stw1942](https://doi.org/10.1093/mnras/stw1942)
- Prentice, S. J., Mazzali, P. A., Pian, E., et al. 2016, *MNRAS*, 458, 2973, doi: [10.1093/mnras/stw299](https://doi.org/10.1093/mnras/stw299)
- Quimby, R. M., Aldering, G., Wheeler, J. C., et al. 2007, *ApJL*, 668, L99, doi: [10.1086/522862](https://doi.org/10.1086/522862)
- Quimby, R. M., Yuan, F., Akerlof, C., & Wheeler, J. C. 2013, *MNRAS*, 431, 912, doi: [10.1093/mnras/stt213](https://doi.org/10.1093/mnras/stt213)
- Quimby, R. M., Kulkarni, S. R., Kasliwal, M. M., et al. 2011, *Nature*, 474, 487, doi: [10.1038/nature10095](https://doi.org/10.1038/nature10095)
- Quimby, R. M., De Cia, A., Gal-Yam, A., et al. 2018, *ApJ*, 855, 2, doi: [10.3847/1538-4357/aaac2f](https://doi.org/10.3847/1538-4357/aaac2f)
- Rigault, M., Neill, J. D., Blagorodnova, N., et al. 2019, *A&A*, 627, A115, doi: [10.1051/0004-6361/201935344](https://doi.org/10.1051/0004-6361/201935344)
- Roberson, M., Fremling, C., & Kasliwal, M. 2022, *The Journal of Open Source Software*, 7, 3612, doi: [10.21105/joss.03612](https://doi.org/10.21105/joss.03612)
- Roming, P. W. A., Kennedy, T. E., Mason, K. O., et al. 2005, *SSRv*, 120, 95, doi: [10.1007/s11214-005-5095-4](https://doi.org/10.1007/s11214-005-5095-4)
- Schlafly, E. F., & Finkbeiner, D. P. 2011, *The Astrophysical Journal*, 737, 103, doi: [10.1088/0004-637x/737/2/103](https://doi.org/10.1088/0004-637x/737/2/103)
- Schlegel, E. M. 1990, *MNRAS*, 244, 269
- Schulze, S., Krühler, T., Leloudas, G., et al. 2018, *MNRAS*, 473, 1258, doi: [10.1093/mnras/stx2352](https://doi.org/10.1093/mnras/stx2352)
- Shappee, B. J. 2014, PhD thesis, The Ohio State University
- Smith, N., Li, W., Foley, R. J., et al. 2007, *ApJ*, 666, 1116, doi: [10.1086/519949](https://doi.org/10.1086/519949)
- Steele, I. A., Smith, R. J., Rees, P. C., et al. 2004, in *Proc. SPIE*, Vol. 5489, *Ground-based Telescopes*, ed. J. M. Oschmann, Jr., 679–692, doi: [10.1117/12.551456](https://doi.org/10.1117/12.551456)

- Stritzinger, M., Suntzeff, N. B., Hamuy, M., et al. 2005, *PASP*, 117, 810, doi: [10.1086/431468](https://doi.org/10.1086/431468)
- Stritzinger, M., Hamuy, M., Suntzeff, N. B., et al. 2002, *AJ*, 124, 2100, doi: [10.1086/342544](https://doi.org/10.1086/342544)
- Tachibana, Y., & Miller, A. A. 2018, *PASP*, 130, 128001, doi: [10.1088/1538-3873/aae3d9](https://doi.org/10.1088/1538-3873/aae3d9)
- Taddia, F., Sollerman, J., Leloudas, G., et al. 2015, *A&A*, 574, A60, doi: [10.1051/0004-6361/201423915](https://doi.org/10.1051/0004-6361/201423915)
- Taddia, F., Sollerman, J., Fremling, C., et al. 2019, *A&A*, 621, A71, doi: [10.1051/0004-6361/201834429](https://doi.org/10.1051/0004-6361/201834429)
- Terreran, G., Blanchard, P., DeMarchi, L., et al. 2020, *The Astronomer's Telegram*, 13970, 1
- The Dark Energy Survey Collaboration. 2005, arXiv e-prints, astro. <https://arxiv.org/abs/astro-ph/0510346>
- Tonry, J. L., Denneau, L., Heinze, A. N., et al. 2018, *PASP*, 130, 064505, doi: [10.1088/1538-3873/aabadf](https://doi.org/10.1088/1538-3873/aabadf)
- Wang, X., Li, W., Filippenko, A. V., et al. 2009, *ApJ*, 697, 380, doi: [10.1088/0004-637X/697/1/380](https://doi.org/10.1088/0004-637X/697/1/380)
- Wilkinson, D. M., Maraston, C., Goddard, D., Thomas, D., & Parikh, T. 2017, *MNRAS*, 472, 4297, doi: [10.1093/mnras/stx2215](https://doi.org/10.1093/mnras/stx2215)
- Woosley, S. E. 2010, *ApJL*, 719, L204, doi: [10.1088/2041-8205/719/2/L204](https://doi.org/10.1088/2041-8205/719/2/L204)
- Woosley, S. E., Blinnikov, S., & Heger, A. 2007, *Nature*, 450, 390, doi: [10.1038/nature06333](https://doi.org/10.1038/nature06333)
- Yan, L., Perley, D. A., De Cia, A., et al. 2018, *ApJ*, 858, 91, doi: [10.3847/1538-4357/aabad5](https://doi.org/10.3847/1538-4357/aabad5)
- Yan, L., Quimby, R., Gal-Yam, A., et al. 2017, *ApJ*, 840, 57, doi: [10.3847/1538-4357/aa6b02](https://doi.org/10.3847/1538-4357/aa6b02)
- Yan, L., Perley, D. A., Schulze, S., et al. 2020, *ApJL*, 902, L8, doi: [10.3847/2041-8213/abb8c5](https://doi.org/10.3847/2041-8213/abb8c5)
- Yao, Y., Miller, A. A., Kulkarni, S. R., et al. 2019, *ApJ*, 886, 152, doi: [10.3847/1538-4357/ab4cf5](https://doi.org/10.3847/1538-4357/ab4cf5)
- Yaron, O., & Gal-Yam, A. 2012, *PASP*, 124, 668, doi: [10.1086/666656](https://doi.org/10.1086/666656)
- Zackay, B., Ofek, E. O., & Gal-Yam, A. 2016, *ApJ*, 830, 27, doi: [10.3847/0004-637X/830/1/27](https://doi.org/10.3847/0004-637X/830/1/27)