Post-Newtonian Feasibility of the Closed String Massless Sector

Kang-Sin Choi*

Scranton Honors Program, Ewha Womans University, Seodaemun-gu, Seoul 03760, Korea Institute of Mathematical Sciences, Ewha Womans University, Seodaemun-gu, Seoul 03760, Korea

Jeong-Hyuck Park[†]

Department of Physics, Sogang University, 35 Baekbeom-ro, Mapo-gu, Seoul 04107, Korea

We perform post Newtonian analysis of Double Field Theory as a test of string theory in gravitational sector against observations. We identify the Eddington–Robertson–Schiff parameters β_{PPN} , γ_{PPN} with the charges of electric *H*-flux and dilaton respectively, and further relate them to stress-energy tensor. We show $\beta_{\text{PPN}} = 1$ from weak energy condition and argue that the observation of $\gamma_{\text{PPN}} \simeq 1$ signifies the ultrarelativistic equation of state in baryons, or the suppression of gluon condensate.

Introduction.—What is the gravitational theory that string theory predicts? The conventional answer has been General Relativity (GR) on account of a metric $g_{\mu\nu}$ appearing in closed string quantisation. However, the metric is only one segment of the closed string massless sector that includes two additional fields, a two-form potential $B_{\mu\nu}$ and a scalar dilaton ϕ . Given the fact that the O(D, D) symmetry of T-duality [1–6] transforms the trio to one another, it may be not unreasonable to regard the whole sector as gravitational. This idea has come true in recent years. The O(D, D) manifest formulation of supergravity [7–12], dubbed Double Field Theory (DFT), has matured into stringy gravity which is based on a beyond-Riemannian geometry [13–16] and has its own version of 'Einstein field equations' carrying $\mathbf{O}(D, D)$ indices [17],

$$G_{AB} = T_{AB} \,. \tag{1}$$

As off-shell and on-shell conserved $\mathbf{O}(D, D)$ tensors, the LHS and RHS of the equality represent stringy curvature and matter respectively. Parametrising the fundamental variables of DFT in terms of the trio $\{g, B, \phi\}$ (*c.f.* [18– 25] for non-Riemannian alternatives), the above single formula is decomposed into

$$R_{\mu\nu} + 2\nabla_{\mu}(\partial_{\nu}\phi) - \frac{1}{4}H_{\mu\rho\sigma}H_{\nu}^{\ \rho\sigma} = K_{(\mu\nu)},$$
$$\frac{1}{2}e^{2\phi}\nabla^{\rho}\left(e^{-2\phi}H_{\rho\mu\nu}\right) = K_{[\mu\nu]}, \quad (2)$$

$$R + 4\Box \phi - 4\partial_{\mu}\phi\partial^{\mu}\phi - \frac{1}{12}H_{\lambda\mu\nu}H^{\lambda\mu\nu} = T_{(0)}.$$

While $K_{(\mu\nu)}, K_{[\mu\nu]}, T_{(0)}$ are *a priori* the components of the DFT stress-energy tensor T_{AB} in (1), each line of (2) may be identified—at least on-shell—as the equation of motion of $g_{\mu\nu}, B_{\mu\nu}$ and a DFT-dilaton *d* absorbing $\sqrt{-g}$ as $d \equiv \phi - \frac{1}{2} \ln \sqrt{-g}$. The $\mathbf{O}(D, D)$ symmetry principle, once taken as the working hypothesis, fixes their coupling to matters completely. For example, when coupled to the Standard Model [26], spin-half fermions (diffeomorphically half-unit-weighted) respond to the H-flux [15] and gauge bosons do so to the dilaton [27]:

$$\int d^4x \,\bar{\psi} \big[i\gamma^{\mu} (D_{\mu} + \frac{1}{24} H_{\mu\nu\rho} \gamma^{\nu\rho}) - m \big] \psi - \frac{1}{4} e^{-2d} \operatorname{Tr}(F_{\mu\nu} F^{\mu\nu}) \,,$$
(3)

which gives [15, 17] $T_{\scriptscriptstyle (0)} = \frac{1}{4} \text{Tr}(F_{\mu\nu}F^{\mu\nu})$ and

$$K_{\mu\nu} = \frac{-ie^{2d}}{4} \bar{\psi} \left(\gamma_{\mu} \vec{D}_{\nu} - \vec{D}_{\nu} \gamma_{\mu} + \frac{1}{4} H_{\nu\rho\sigma} \gamma_{\mu}{}^{\rho\sigma} \right) \psi + \frac{1}{2} \operatorname{Tr} \left(F_{\mu\rho} F_{\nu}{}^{\rho} \right)$$
(4)

Having stated the above, to the best of our knowledge, the gravitational coupling in quantum field theories has never been tested experimentally. All the observations are based on the geodesic motion (and deviation) of point particles including photon. The $\mathbf{O}(D, D)$ symmetry then enforces the particles to couple to $g_{\mu\nu}$ only in the usual manner, $-m_* \int d\tau \sqrt{-g_{\mu\nu} \dot{y}^{\mu} \dot{y}^{\nu}}$, which gives $T_{(0)} = 0$ and

$$K^{\mu\nu}(x) = \frac{1}{2}m_* \int d\tau \ \dot{y}^{\mu} \dot{y}^{\nu} \ e^{2d} \delta^{(4)}(x - y(\tau)) \ . \tag{5}$$

That is to say [28], the strong equivalence principle is saved in the string frame. While the sources can be stringy, the probes are still point-like up to now.

Yet, given the augmented stress-energy tensor, the gravitational physics of DFT should be richer than that of GR. To develop some physical intuition on the governing equations (2), we may take the following three steps. *i*) The second formula dictates how $K_{[\mu\nu]}$ shapes the *H*-flux. *ii*) Subtracting the third from the trace of the first, one solves for the dilaton:

$$\Box \left(e^{-2\phi} \right) = \left(K_{\mu}{}^{\mu} - T_{(0)} + \frac{1}{6} H_{\lambda\mu\nu} H^{\lambda\mu\nu} \right) e^{-2\phi} \,. \tag{6}$$

iii) The metric is lastly determined by the first formula.

It is the purpose of the present Letter to address the feasibility of the closed string massless sector, or D = 4 DFT as *stringy gravity* alternative to GR, within the parametrised post Newtonian (PPN) formalism [29–31]. Keeping astrophysical systems in mind rather than the entire Universe, we shall neglect any large scale contribution to $T_{(0)}$ such as a (non-critical string) cosmological constant [32–35]. We postulate the spacetime

^{*} kangsin@ewha.ac.kr

[†] park@sogang.ac.kr

to be asymptotically flat and static. The Eddington–Robertson–Schiff parameters $\beta_{\text{PPN}}, \gamma_{\text{PPN}}$ are then defined in an isotropic coordinate system,

$$ds^{2} = -\left(1 - \frac{2MG_{\rm N}}{r} + \frac{2\beta_{\rm PPN}(MG_{\rm N})^{2}}{r^{2}} + \cdots\right) dt^{2} + \left(1 + \frac{2\gamma_{\rm PPN}MG_{\rm N}}{r} + \cdots\right) dx^{i} dx^{j} \delta_{ij},$$

$$(7)$$

where $r = \sqrt{x^i x^j \delta_{ij}}$ is the isotropic radius and G_N is the Newtonian constant of gravitation (setting " $\alpha_{PPN} \equiv 1$ ").

The current stringent observational bounds for the solar gravity are $\gamma_{\rm PPN} - 1 = (2.1 \pm 2.3) \times 10^{-5}$ (Shapiro timedelay by Cassini spacecraft) [31, 36] and adopting this value $\beta_{\rm PPN} - 1 = (-4.1 \pm 7.8) \times 10^{-5}$ or $(0.4 \pm 2.4) \times 10^{-4}$ (perihelion shifts of Mercury or Mars) [31, 37]. Other observations include $4\beta_{\rm PPN} - \gamma_{\rm PPN} - 3 = (4.44 \pm 4.5) \times 10^{-4}$ for the Earth gravity [38] and $\gamma_{\rm PPN} = 0.98 \pm 0.07$ on galactic size scales [39]. These all agree with the GR prediction $\beta_{\rm PPN} = \gamma_{\rm PPN} = 1$ and rule out various alternative gravities [40–42]. For example, a class of scalar-tensor theories, notably Brans–Dicke, gives

$$\gamma_{\rm PPN} = \frac{1 + \omega_{\rm BD}}{2 + \omega_{\rm BD}} \,, \tag{8}$$

and thus the observations compel the coupling constant $\omega_{\rm BD}$ unnaturally huge. Truncating $K_{[\mu\nu]}$, $T_{(0)}$, and the *H*-flux consistently, the leftover components of (2) coincide with the Brans–Dicke field equations for $\omega_{\rm BD} = -1$. This has been often regarded as a "smoking gun" to necessarily eliminate the string dilaton ϕ . Below, analysing (2) with care, we show that spherically symmetric static objects in stringy gravity (henceforth collectively "stars" though applicable to the Earth) have $\beta_{\rm PPN} = 1$ and make $\gamma_{\rm PPN}$ depend on equation-of-state parameters. As a byproduct, we shall point out that the well-known relation (8) is an artifact of restricting to 'pressureless' stars. It can generalise to $\gamma_{\rm PPN} = 3p/\rho$ for $\omega_{\rm BD} = -1$, and thus becomes phenomenologically viable with ultrarelativistic pressure.

Post Newtonian Double Field Theory.—Following the three steps i), ii), iii) above, it is possible to solve the linearised version of (2), using the Green function method with retarded time $t' = t - |\mathbf{x} - \mathbf{x}'|$ [44],

$$H_{\lambda\mu\nu}(x) \simeq -\frac{1}{2\pi} \int d^3 x' \frac{3\partial'_{[\lambda}K_{\mu\nu]}(x')}{|\mathbf{x}-\mathbf{x}'|},$$

$$\phi(x) \simeq \frac{1}{8\pi} \int d^3 x' \frac{[K_{\mu}{}^{\mu} - T_{(0)} + \frac{1}{6}H_{\lambda\mu\nu}H^{\lambda\mu\nu}](x')}{|\mathbf{x}-\mathbf{x}'|},$$

$$g_{\mu\nu}(x) \simeq \eta_{\mu\nu} + \frac{1}{2\pi} \int d^3 x' \frac{[K_{(\mu\nu)} + \frac{1}{4}H_{\mu\rho\sigma}H_{\nu}{}^{\rho\sigma}](x')}{|\mathbf{x}-\mathbf{x}'|}.$$
(9)

Mapping this to the isotropic coordinate system, we can read off MG_N and γ_{PPN} , but not β_{PPN} which would require higher order analysis. Instead, we turn to a known D = 4 three-parameter exact solution to (2) [43]. It has vanishing sources ($T_{AB} = 0$) and can be identified as the most general, asymptotically flat, static, and spherically symmetric vacuum geometry ($G_{AB} = 0$) [28]. For our purposes, we perform a radial coordinate transformation, $r_{\text{there}} = r + \frac{a^2 + b^2}{16r} + \frac{\alpha_{\text{there}} - \beta_{\text{there}}}{2}$ from [28], and put the geometry decisively in the isotropic (as well as often spherical) coordinate system,

$$e^{2\phi} = \gamma_{+} \left(\frac{4r - \sqrt{a^{2} + b^{2}}}{4r + \sqrt{a^{2} + b^{2}}}\right)^{\frac{2b}{\sqrt{a^{2} + b^{2}}}} + \gamma_{-} \left(\frac{4r + \sqrt{a^{2} + b^{2}}}{4r - \sqrt{a^{2} + b^{2}}}\right)^{\frac{2b}{\sqrt{a^{2} + b^{2}}}},$$

$$H_{(3)} = h \sin \vartheta \, \mathrm{d}t \wedge \mathrm{d}\vartheta \wedge \mathrm{d}\varphi = h \, \mathrm{d}t \wedge \left(\frac{\epsilon_{ijk} x^{i} \, \mathrm{d}x^{j} \wedge \mathrm{d}x^{k}}{2r^{3}}\right),$$

$$\mathrm{d}s^{2} = g_{tt}(r) \, \mathrm{d}t^{2} + g_{rr}(r) \left(\mathrm{d}r^{2} + r^{2} \mathrm{d}\Omega^{2}\right),$$

$$(10)$$

where $d\Omega^2 = d\vartheta^2 + \sin^2 \vartheta d\varphi^2$, and

$$g_{tt}(r) = -e^{2\phi(r)} \left(\frac{4r - \sqrt{a^2 + b^2}}{4r + \sqrt{a^2 + b^2}}\right)^{\frac{2a}{\sqrt{a^2 + b^2}}},$$

$$g_{rr}(r) = e^{2\phi(r)} \left(\frac{4r + \sqrt{a^2 + b^2}}{4r - \sqrt{a^2 + b^2}}\right)^{\frac{2a}{\sqrt{a^2 + b^2}}} \left(1 - \frac{a^2 + b^2}{16r^2}\right)^2.$$

The three "free" parameters are $\{a, b, h \mid b^2 \ge h^2\}$, which set $\gamma_{\pm} = \frac{1}{2} (1 \pm \sqrt{1 - h^2/b^2})$.

The geometry (10) can be identified as the outer geometry of a (stringy) compact star. Crucially, from the full Einstein equations (2), it becomes possible to ascribe the three parameters to the star's stress-energy tensor or $\{K_t^t, K_r^r, K_\vartheta^{\ \vartheta} = K_\varphi^{\ \varphi}, K_{tr}, K_{[\vartheta\varphi]}\}$ [17], such as

$$a = \frac{1}{4\pi} \int d^3x \ e^{-2d} \left(K_i{}^i - K_t{}^t - T_{(0)} + \frac{1}{6} H_{ijk} H^{ijk} \right) .$$
(11)

Inside the star, while the electric H-flux is persistently of the rigid form (10), the magnetic flux can be nontrivial,

$$H^{r\vartheta\varphi} = -2e^{2d} \int_{r}^{r_{\star}} \mathrm{d}r' \ e^{-2d} K^{[\vartheta\varphi]} \,, \tag{12}$$

where the upper limit of the integral r_{\star} denotes the star's finite radius. As implied by the first of (2) for the isotropic metric, the electric and the magnetic fluxes should meet

$$H_{t\vartheta\varphi}H_r^{\ \vartheta\varphi} = -2K_{(tr)}\,,\tag{13}$$

while the second of (2) gives $K_{[tr]} = 0$.

Now, from the 1/r-expansion of the outer geometry (10), we are able to identify the Newtonian mass [28],

$$MG_{\rm N} = \frac{1}{2} \left(a + b\sqrt{1 - h^2/b^2} \right),$$
 (14)

and the two post Newtonian parameters,

$$(\beta_{\rm PPN}-1)(MG_{\rm N})^2 = \frac{\hbar^2}{4}, \ (\gamma_{\rm PPN}-1)MG_{\rm N} = -b\sqrt{1-\frac{\hbar^2}{b^2}}.$$
(15)

Since the inverse relations hold,

$$h = \pm 2\sqrt{\beta_{\text{PPN}} - 1} MG_{\text{N}}, \qquad a = (\gamma_{\text{PPN}} + 1)MG_{\text{N}},$$
$$b = (1 - \gamma_{\text{PPN}})\sqrt{1 + 4\frac{\beta_{\text{PPN}} - 1}{(\gamma_{\text{PPN}} - 1)^2}} MG_{\text{N}},$$

the outer geometry (10) is also completely determinable by the triple of $\{MG_{\rm N}, \beta_{\rm PPN}, \gamma_{\rm PPN}\}$, such as

$$\phi \simeq \frac{(\gamma_{\rm PPN} - 1)MG_{\rm N}}{2r} + \frac{(\beta_{\rm PPN} - 1)(MG_{\rm N})^2}{r^2},$$

$$H_{(3)} = \pm \sqrt{\beta_{\rm PPN} - 1}MG_{\rm N} dt \wedge \left(\frac{\epsilon_{ijk}x^i dx^j \wedge dx^k}{r^3}\right),$$

$$g_{rr} \simeq 1 + \frac{2\gamma_{\rm PPN}MG_{\rm N}}{r} + \frac{(6\beta_{\rm PPN} + 7\gamma_{\rm PPN}^2 - 7)(MG_{\rm N})^2}{r^2}.$$
(16)

The expansion of ϕ suggests that $(\gamma_{\text{PPN}}-1)MG_{\text{N}}$ is the dilaton charge. Indeed, multiplying $\sqrt{-g}$ to (6) and using $\sqrt{-g}\Box(e^{-2\phi}) = \partial_{\mu}(\sqrt{-g}\partial^{\mu}e^{-2\phi})$, we can compute the dilaton charge from the stress-energy tensor of the star,

$$(\gamma_{\rm PPN} - 1)MG_{\rm N} = \frac{1}{4\pi} \oint_{\infty} \mathrm{d}S_i \,\partial^i e^{-2\phi}$$
$$= \frac{1}{4\pi} \int \mathrm{d}^3 x \, e^{-2d} \left(K_{\mu}{}^{\mu} - T_{(0)} + \frac{1}{6} H_{\lambda\mu\nu} H^{\lambda\mu\nu} \right)$$
(17)

where the *H*-flux has been fixed by (12) and (13). Similarly, from (16) we identify $\sqrt{\beta_{\text{PPN}}-1}MG_{\text{N}}$ as the *electric H*-flux charge which is through (12) and (13) also related to the stress-energy tensor,

$$\sqrt{\beta_{\rm PPN} - 1} M G_{\rm N} = \left| \oint_{\infty} \frac{\mathrm{d}S_i}{16\pi} \epsilon^{ijk} H_{tjk} \right| = \left| \frac{K_{(tr)} g^{rr} (e^{-2d} / \sin\vartheta)}{2 \int_r^{T_*} \mathrm{d}r' (e^{-2d} K^{[\vartheta\varphi]})} \right|$$
(18)

For consistency, the fractional quantity at the end of (18) must be independent of r, while the ϑ -dependency is trivially canceled since $K_{[\vartheta \varphi]} \propto \sin \vartheta$, $K^{[\vartheta \varphi]} \propto 1/\sin \vartheta$, and

$$e^{-2d} = e^{-2\phi(r)}\sqrt{-g_{tt}(r)g_{rr}^3(r)}r^2\sin\vartheta$$
. (19)

Lastly, from (11), (15), (17), the Newtonian mass (14) has its own integral expression [17],

$$M = \frac{1}{4\pi G_{\rm N}} \int d^3x \ e^{-2d} \left(-K_t^{\ t} - \frac{1}{2} H_{t\mu\nu} H^{t\mu\nu} \right) \ . \tag{20}$$

Substituting the expression of $-K_t^t > 0$ for particles (5) into (20), we identify m_* in terms of the rest mass,

$$m_* = 8\pi G_{\rm N} m \,, \tag{21}$$

and further confirm that, like in GR, the Newtonian mass M is given by the sum of the energy $m\dot{t} = m/\sqrt{1-v^2}$ rather than the rest mass. However, in contrast to GR, it is a priori $-4K_t^{\ t}$ rather than the conventional energy, or $\propto -2K_t^{\ t}+T_{(0)}$, that enters the mass formula. This difference is significant for gauge bosons, in particular gluons: as seen from (4) only the electric field F_{ti} contributes to the mass, but the magnetic field F_{ij} does not. Since e^{-2d} is the rightful integral measure in DFT, all

Since e^{-2d} is the rightful integral measure in DFT, all the integral expressions above are 'proper', *i.e.* invariant under static (radial) diffeomorphisms, which is rather notoriously not the case for the Schwarzschild mass in GR.

 $\beta_{\rm PPN} = 1$ from a weak energy condition.—The exact expressions of the Newtonian mass (20) and the magnetic H-flux (12) are in good agreement with the linearised general solutions (9). The former (20) is obvious and the latter (12) is due to a shell theorem. However, the electric *H*-flux given through (15) and (18) appears highly nonperturbative. In fact, for the linearised expression (9), when $K_{[\mu\nu]}$ therein is static and spherical, we should have $\partial_t K_{[\mu\nu]} = 0$ and $K_{[ti]} = x_i f(r)$ for some radial function f(r). Consequently $\partial_{[t} K_{ij]}$ vanishes and the linear version of the electric H-flux must be trivial. We resolve this discrepancy by arguing that, the electric H-flux in the exact solution should be trivial too. With (19), its contribution to the Newtonian mass (20) is an $1/r^2$ integral, $\int_0^\infty dr \, e^{-2\phi} h^2/(G_N\sqrt{-g_{tt}g_{rr}}r^2)$, which—provided the geometry at the centre of the star is non-singular would diverge unless h = 0. Since the mass M is finite and a weak energy condition $-K_t^t > 0$ should hold, the electric *H*-flux must be inevitably trivial. Besides, the last expression of (18) ought to be independent of r. In its small r-limit, from (19) the numerator vanishes. Hence we arrive at the same conclusion h = 0. Consequently, from (13) $K_{(tr)} = 0$ and from (15) $\beta_{PPN} = 1$.

 $\gamma_{\rm PPN} \simeq 1$ from subhadronic pressure.—With the vanishing electric *H*-flux (h = 0), the volume integrals of the total mass (20) and $\gamma_{\rm PPN}$ (17) are now all restricted to the star's interior,

$$\gamma_{\rm PPN} = 1 - \frac{\int_{\rm star} d^3 x \ e^{-2d} \left(K_{\mu}{}^{\mu} - T_{(0)} + \frac{1}{6} H_{ijk} H^{ijk} \right)}{\int_{\rm star} d^3 x \ e^{-2d} K_t^{\ t}} , \quad (22)$$

where $\frac{1}{6}H_{ijk}H^{ijk} = H_{r\vartheta\varphi}H^{r\vartheta\varphi}$ is set by $K^{[\vartheta\varphi]}$ from (12). Clearly, γ_{PPN} depends on the stress-energy tensor for which we introduce volume-averaged, equation-of-state parameters (*c.f.* [45]),

$$w_{K} = \frac{\int_{\text{star}} d^{3}x \ e^{-2d} \ \frac{1}{3}K_{i}^{i}}{\int_{\text{star}} d^{3}x \ e^{-2d} \left(-K_{t}^{t}\right)}, \quad w_{T} = \frac{\int_{\text{star}} d^{3}x \ e^{-2d} T_{(0)}}{\int_{\text{star}} d^{3}x \ e^{-2d} \left(-K_{t}^{t}\right)},$$
(23)

and further for the magnetic H-flux part,

$$\delta_{H-\text{flux}} = \frac{16\pi \int_{0}^{r_{\star}} dr \ r^{2} e^{2\phi} \sqrt{g_{rr}^{3}/(-g_{tt})} \left(\int_{r}^{r_{\star}} dr' \ e^{-2d} K^{[\vartheta\varphi]}\right)^{2}}{\int_{\text{star}} d^{3}x \ e^{-2d} \left(-K_{t}^{\ t}\right)}.$$
(24)

Note from the identification of the usual energymomentum tensor $\mathcal{T}_{\mu\nu} = e^{-2\phi} \left(2K_{(\mu\nu)} - g_{\mu\nu}T_{(0)} \right) c.f.$ (27), the conventional equation-of-state parameter reads

$$w = \frac{\mathcal{T}_i^i}{-3\mathcal{T}_t^t} = \frac{p}{\rho} = \frac{w_K - \frac{1}{2}w_T}{1 + \frac{1}{2}w_T}.$$
 (25)

The post Newtonian parameter (22) amounts to

$$\gamma_{\rm PPN} = 3w_K - w_T + \delta_{H-\rm flux} \,. \tag{26}$$

For the ideal gas of particles, we have $w_T = \delta_{H-\text{flux}} = 0$ and hence simply $\gamma_{\text{PPN}} = 3w$. In terms of the averaged speed v, this is equivalent to $\gamma_{\text{PPN}} \simeq v^2$ since from (5) we have $K_{\mu}{}^{\mu}/K_t{}^t = -1/(g_{tt}\dot{t}^2) = -g^{tt}(1-v^2) \simeq 1-v^2$.

If a star were composed of non-relativistic pressureless dust, we would get $\gamma_{\rm PPN} = 0$, which certainly fails to explain the observations. This is actually the case with the usual analysis of the Brans–Dicke theory having $\omega_{\rm BD} = -1$ (8) and non-relativistic sources. More realistic stars would exert pressure along the radial direction to balance the gravitational force. In the Newtonian gravity, the pressure is fixed to meet $\frac{dp(r)}{dr} = -M(r)G_N\rho/r^2$ and leads to $w = \frac{\langle p \rangle}{\rho} = \frac{MG_N}{2r_\star}$ (volume-averaged). The Sun would have then merely $\gamma_{\rm PPN} = 3w \sim 3 \times 10^{-6}$ which is far less than unity. In GR, the outer Schwarzschild geometry of a star depends on the total mass only, being inert to the pressure. In DFT, the outer geometry (10)depends on the equation-of-state parameters (23), yet the Einstein equations thereof (1) still do not impose any restriction on them, as seen e.q. from (17) and (20). In flat spacetime, the conservation of the (ordinary) stressenergy tensor implies that the volume integral of the pressure vanishes, which is known as the von Laue condition [46]. To correctly include the stringy gravitational effect, we call the on-shell conservation $\nabla_A T^A{}_B = 0$ of (1) [17] (c.f. [47, 48]) which becomes more concretly $\nabla_{\mu} \left(e^{-2\phi} K^{[\mu\nu]} \right) = 0$ and

$$\nabla^{\rho} \left(e^{-2\phi} K_{(\rho\mu)} \right) + \frac{1}{2} e^{-2\phi} \left(H_{\mu\rho\sigma} K^{[\rho\sigma]} - \partial_{\mu} T_{(0)} \right) = 0.$$
 (27)

For the spherically symmetric static case of our interest where $K_{tr} = 0 = H_{t\vartheta\varphi}$ and (12) holds, writing the covariant derivative in (27) explicitly as

$$\begin{split} r\sqrt{-g}\nabla^{\rho} \left(e^{-2\phi}K_{(\rho r)}\right) &= \frac{\mathrm{d}}{\mathrm{d}r} \left[re^{-2d} \left(K_{r}^{\ r} - \frac{1}{3}K_{i}^{\ i}\right)\right] \\ &+ \frac{1}{3}r^{3}\frac{\mathrm{d}}{\mathrm{d}r} \left(r^{-2}e^{-2d}K_{i}^{\ i}\right) - \frac{1}{2}re^{-2d} \left(\frac{\dot{g}_{tt}}{g_{tt}}K_{t}^{\ t} + \frac{\dot{g}_{rr}}{g_{rr}}K_{i}^{\ i}\right), \end{split}$$

we relax the von Laue condition rather inevitably,

$$\int d^{3}x \ e^{-2d} \left(K_{i}^{i} - \frac{3}{2}T_{(0)} + \frac{1}{8}H_{ijk}H^{ijk} \right)$$

$$= \int d^{3}x \ e^{-2d} \left[\frac{r\dot{g}_{tt}}{2g_{tt}} \left(-K_{t}^{t} + \frac{1}{2}T_{(0)} + \frac{1}{24}H_{ijk}H^{ijk} \right) - \frac{r\dot{g}_{rr}}{2g_{rr}} \left(K_{i}^{i} - \frac{3}{2}T_{(0)} + \frac{1}{8}H_{ijk}H^{ijk} \right) - r\dot{\phi} \left(T_{(0)} + \frac{1}{12}H_{ijk}H^{ijk} \right) \right]$$
(28)

While $\frac{\dot{g}_{tt}}{2g_{tt}}(-K_t^{\ t})$ corresponds to the gravitational force, the presence of the self-reflective term of $-\frac{r\dot{g}_{rr}}{2g_{rr}}(K_i^{\ i}-\frac{3}{2}T_{(0)}+\frac{1}{8}H_{ijk}H^{ijk})$ invalidates the aforementioned Newtonian underestimation of the total pressure. In fact, (27) and (28) are identities for any matter and geometric data $\{K_{\mu\nu}, T_{(0)}, g_{\mu\nu}, B_{\mu\nu}, \phi\}$ that satisfy the Einstein equations (2), since $\nabla_A G^A{}_B = 0$ is a 'Bianchi' identity in DFT [48, 49]. For $\gamma_{\rm PPN} \simeq v^2$ to be close to unity within the stringent observational bound of 2.1×10^{-5} [31, 36], the constituting particles—if not strings—should be ultrarelativistic. In terms of temperature $3k_{\rm B}T = m(\frac{1}{\sqrt{1-v^2}}-1)$, we get $\gamma_{\rm PPN} \simeq 1 - (\frac{m}{m+3k_{\rm B}T})^2$. Assuming $T = 1.57 \times 10^7$ K at the center of the Sun [50], from $(\frac{m}{m+3k_{\rm B}T})^2 < 2.1 \times 10^{-5}$, the particle mass m should not exceed 20 eV. Certainly no such a light atom exists. While so, a recent experiment has revealed a pressure inside protons as high as 10^{35} Pa ~ 0.005 GeV⁴ [51] comparable to their mass density 10^{18} kg/m³ ~ 0.004 GeV⁴ ~ $(\Lambda_{\rm QCD})^4$ [52]. Henceforth we look into the energy and pressure inside baryons, or the QCD matter of (3). The deviation $\gamma_{\rm PPN}-1$ (22) is then given by the sum of $\delta_{H-\rm flux}$ (24) and an integral,

$$\frac{1}{4\pi M G_{\rm N}} \int_{\rm star}^{\rm d} 3x \, \frac{1}{4} e^{-2d} {\rm Tr}(F_{\mu\nu}F^{\mu\nu}) - \frac{1}{2}\bar{\psi}(m+i\frac{1}{12}\gamma^{ijk}H_{ijk})\psi \,.$$
⁽²⁹⁾

In our normalisation (2), from (21), $K_{\mu\nu}, T_{(0)}$ are order of $8\pi G_N$. Hence the magnetic *H*-flux (12) gives a subleading contribution and may be ignored. Then, (29) boils down faithfully to the integral of the gluon and quark condensates (vacuum expectation value) [53, 54]. According to [55–63], the condensates have local support restricted to the hadrons' interior. Put differently, the energy, pressure, and condensates specified by the components of $K_{\mu\nu}, T_{(0)}$ are all confined. Although the condensates may have the natural order of magnitudes $(\Lambda_{\rm QCD})^4$ and $m \times (\Lambda_{\rm QCD})^3$ respectively, while the latter can be chirally rotated and averaged away, the empirical measurements of the former [64–68] show significant scatter and even differences in sign [58], allowing zero as $0.006 \pm 0.012 \,\text{GeV}^4$ [64]. Perhaps, at (fourdimensional) thermal equilibrium, the equal number of electric and magnetic gluon field-strength-squared cancel each other to suppress the gluon condensate normalized by the baryon mass density, c.f. [60]. This ratio essentially estimates (29) and the observation $\gamma_{\rm PPN} \simeq 1$ is consistent with the suppression.

Discussion.—To conclude, DFT sets $\beta_{\text{PPN}} = 1$ and lets γ_{PPN} depend on the equation-of-state parameters (26). Rather than ruling out the theory, applied to baryons' interior where the energy and pressure are both confined, the apparently universal observations $\gamma_{\text{PPN}} \simeq 1$ including the Sun and the Earth [31, 36, 38] signify the ultrarelativistic equation of state inside baryons, $w_K \simeq \frac{1}{3}, w_T \simeq 0$, through the suppression of the (massless) gluon condensate (29). We call for both theoretical and experimental verifications of this rather drastic conclusion of ordinary matter being ultrarelativistic at a subhadronic level.

Some comments are in order. In terms of the adiabatic index c_P/c_V , one gets $w = c_P/c_V - 1$ which becomes 1/3 for ultrarelativistic ideal fluid having $c_P = 4k_B$ and $c_V = 3k_B$, e.g. [69].

Any two-derivative potential-free effective scalar the-

ory, if O(D, D)-symmetric, should be of the form,

$$S_{\text{eff.}} = -\int d^4 x \ e^{-2d} g^{\mu\nu} \partial_\mu \Phi^I \partial_\nu \Phi^J \mathcal{G}_{IJ}(\Phi) \,, \qquad (30)$$

which gives $K_{\mu}{}^{\mu} = g^{\mu\nu}\partial_{\mu}\Phi^{I}\partial_{\nu}\Phi^{J}\mathcal{G}_{IJ} = T_{(0)}, \ K^{[\vartheta\varphi]} = 0,$ and thus rather precisely $\gamma_{\text{PPN}} = 1.$

The *B*-field is dual to a scalar axion in four dimensions. Recently, a static Kähler axion has been suggested to make γ_{PPN} close to unity (for w = 0 and $\omega_{\text{BD}} = \frac{3}{2}$) [70]. In our case, the axion dual to the magnetic *H*-flux would be time, rather than radial, dependent.

Since $\delta_{H-\text{flux}} \geq 0$, if any star were ever found to feature $\gamma_{\text{PPN}} > 1$, it should be stringy. From the superextensive multi-integrals appearing in $\delta_{H-\text{flux}}$ (24), bigger stringy stars or galaxies would have larger value of γ_{PPN} .

The equations of state $w_K \simeq \frac{1}{3}$, $w_T \simeq 0$ for $\gamma_{\text{PPN}} \simeq 1$ prevents also ϕ from the cosmological time evolution [45, 71]. This may satisfy an extremely tight bound on the time variation of the fine structure constant [72], since e^{-2d} is an overall factor hence affecting the coupling constant of the gauge boson Lagrangian (3).

Acknowledgments.—We wish to thank Robert Brandenberger, Minkyoo Kim, Bum-Hoon Lee, Chang-Hwan Lee, Hocheol Lee, Taekoon Lee, Wonwoo Lee, Shinji Mukohyama, Sang-Jin Sin, and Clifford Will for useful comments and correspondences. This work is supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) Grants, NRF-2016R1D1A1B01015196, NRF-2018R1A2B2007163, and NRF-2020R1A6A1A03047877 (Center for Quantum Space Time).

- T. H. Buscher, "A Symmetry of the String Background Field Equations," Phys. Lett. B **194** (1987) 59.
 T. H. Buscher, "Path Integral Derivation of Quantum
- [2] T. H. Buscher, "Path Integral Derivation of Quantum Duality in Nonlinear Sigma Models," Phys. Lett. B 201 (1988) 466.
- [3] A. Giveon, E. Rabinovici and G. Veneziano, "Duality in String Background Space," Nucl. Phys. B **322** (1989), 167-184 doi:10.1016/0550-3213(89)90489-6
- [4] M. Duff, "Duality Rotations in String Theory," Nucl. Phys. B 335 (1990), 610
- [5] A. A. Tseytlin, "Duality Symmetric Formulation Of String World Sheet Dynamics," Phys. Lett. B 242, 163 (1990).
- [6] A. A. Tseytlin, "Duality Symmetric Closed String Theory And Interacting Chiral Scalars," Nucl. Phys. B 350, 395 (1991).
- [7] W. Siegel, "Two vierbein formalism for string inspired axionic gravity," Phys. Rev. D 47 (1993) 5453 [hep-th/9302036].
- [8] W. Siegel, "Superspace duality in low-energy superstrings," Phys. Rev. D 48 (1993) 2826 [hep-th/9305073].
- [9] C. Hull and B. Zwiebach, "Double Field Theory," JHEP 0909 (2009) 099 [arXiv:0904.4664 [hep-th]].
- [10] C. Hull and B. Zwiebach, JHEP 0909 (2009) 090 [arXiv:0908.1792 [hep-th]].
- [11] O. Hohm, C. Hull and B. Zwiebach, "Background independent action for double field theory," JHEP 1007 (2010) 016 [arXiv:1003.5027 [hep-th]].
- [12] O. Hohm, C. Hull and B. Zwiebach, "Generalized metric formulation of double field theory," JHEP 08 (2010), 008 [arXiv:1006.4823 [hep-th]].
- [13] I. Jeon, K. Lee and J. H. Park, "Differential geometry with a projection: Application to double field theory," JHEP **1104** (2011) 014 doi:10.1007/JHEP04(2011)014 [arXiv:1011.1324 [hep-th]].
- [14] I. Jeon, K. Lee and J. H. Park, "Stringy differential geometry, beyond Riemann," Phys. Rev. D 84 (2011) 044022 doi:10.1103/PhysRevD.84.044022 [arXiv:1105.6294 [hepth]].
- [15] I. Jeon, K. Lee and J. H. Park, "Incorporation of fermions into double field theory," JHEP 11 (2011), 025

doi:10.1007/JHEP11(2011)025 [arXiv:1109.2035 [hep-th]].

- [16] J. H. Park, "Comments on double field theory and diffeomorphisms," JHEP 06 (2013), 098 doi:10.1007/JHEP06(2013)098 [arXiv:1304.5946 [hepth]].
- [17] S. Angus, K. Cho and J. H. Park, "Einstein Double Field Equations," Eur. Phys. J. C **78** (2018) no.6, 500 doi:10.1140/epjc/s10052-018-5982-y [arXiv:1804.00964 [hep-th]]. We have mapped $G_{AB} = 8\pi G_{\text{there}} T_{AB}^{\text{there}}$ of [17] to (1) by

putting $16\pi G_{\text{there}} = 1$ and $\frac{1}{2}T_{AB}^{\text{there}} = T_{AB}$. The physically observable Newton constant G_{N} appears through the post Newtonian parametrisation of the metric (7).

- [18] K. Lee and J. H. Park, "Covariant action for a string in doubled yet gauged spacetime," Nucl. Phys. B 880 (2014) 134 [arXiv:1307.8377 [hep-th]].
- [19] S. M. Ko, C. Melby-Thompson, R. Meyer and J. H. Park, "Dynamics of Perturbations in Double Field Theory & Non-Relativistic String Theory," JHEP **12** (2015), 144 doi:10.1007/JHEP12(2015)144 [arXiv:1508.01121 [hepth]].
- [20] K. Morand and J. H. Park, "Classification of non-Riemannian doubled-yet-gauged spacetime," Eur. Phys. J. C 77 (2017) no.10, 685 [arXiv:1707.03713 [hep-th]].
- [21] D. S. Berman, C. D. A. Blair and R. Otsuki, "Non-Riemannian geometry of M-theory," JHEP 07 (2019), 175 doi:10.1007/JHEP07(2019)175 [arXiv:1902.01867 [hep-th]].
- [22] C. D. A. Blair, "A worldsheet supersymmetric Newton-Cartan string," JHEP 10 (2019), 266 doi:10.1007/JHEP10(2019)266 [arXiv:1908.00074 [hepth]].
- [23] K. Cho and J. H. Park, "Remarks on the non-Riemannian sector in Double Field Theory," Eur. Phys. J. C 80 (2020) no.2, 101 doi:10.1140/epjc/s10052-020-7648-9 [arXiv:1909.10711 [hep-th]].
- [24] J. H. Park and S. Sugimoto, "String Theory and non-Riemannian Geometry," Phys. Rev. Lett. **125** (2020) no.21, 211601 doi:10.1103/PhysRevLett.125.211601 [arXiv:2008.03084 [hep-th]].

- [25] A. D. Gallegos, U. Gürsoy, S. Verma and N. Zinnato, "Non-Riemannian gravity actions from double field theory," [arXiv:2012.07765 [hep-th]].
- [26] K. S. Choi and J. H. Park, "Standard Model as a Double Field Theory," Phys. Rev. Lett. **115** (2015) no.17, 171603 doi:10.1103/PhysRevLett.115.171603 [arXiv:1506.05277 [hep-th]].
- [27] I. Jeon, K. Lee and J. H. Park, "Double field formulation of Yang-Mills theory," Phys. Lett. B 701 (2011), 260-264 doi:10.1016/j.physletb.2011.05.051 [arXiv:1102.0419 [hep-th]].
- [28] S. M. Ko, J. H. Park and M. Suh, "The rotation curve of a point particle in stringy gravity," JCAP 06 (2017), 002 doi:10.1088/1475-7516/2017/06/002 [arXiv:1606.09307 [hep-th]].
- [29] C. M. Will and K. Nordtvedt, Jr., "Conservation Laws and Preferred Frames in Relativistic Gravity. I. Preferred-Frame Theories and an Extended PPN Formalism," Astrophys. J. 177 (1972), 757 doi:10.1086/151754
- [30] T. Damour and G. Esposito-Farese, "Tensor multiscalar theories of gravitation," Class. Quant. Grav. 9 (1992), 2093-2176 doi:10.1088/0264-9381/9/9/015
- [31] C. M. Will, "The Confrontation between General Relativity and Experiment," Living Rev. Rel. 17 (2014), 4 doi:10.12942/lrr-2014-4 [arXiv:1403.7377 [gr-qc]].
- [32] T. R. Taylor and G. Veneziano, "Dilaton Couplings at Large Distances," Phys. Lett. B 213 (1988), 450-454 doi:10.1016/0370-2693(88)91290-7
- [33] T. Damour and K. Nordtvedt, "General relativity as a cosmological attractor of tensor scalar theories," Phys. Rev. Lett. **70** (1993), 2217-2219 doi:10.1103/PhysRevLett.70.2217
- [34] T. Damour and K. Nordtvedt, "Tensor scalar cosmological models and their relaxation toward general relativity," Phys. Rev. D 48 (1993), 3436-3450 doi:10.1103/PhysRevD.48.3436
- [35] T. Damour, F. Piazza and G. Veneziano, "Runaway dilaton and equivalence principle violations," Phys. Rev. Lett. 89 (2002),081601 doi:10.1103/PhysRevLett.89.081601 [arXiv:gr-qc/0204094 [gr-qc]].
- [36] B. Bertotti, L. Iess and P. Tortora, "A test of general relativity using radio links with the Cassini spacecraft," Nature 425 (2003), 374-376 doi:10.1038/nature01997.
- [37] A. Konopliv, S. Asmar, W. Folkner, O. Karatekin, D. Nunes, S. Smrekar, C. Yoder and M. Zuber, "Mars high resolution gravity fields from MRO, Mars seasonal gravity, and other dynamical parameters," Icarus 211 issue 1 (2011), 401-428 doi:10.1016/j.icarus.2010.10.004.
- [38] J. G. Williams, S. G. Turyshev and D. H. Boggs, "Progress in lunar laser ranging tests of relativistic gravity," Phys. Rev. Lett. **93** (2004), 261101 doi:10.1103/PhysRevLett.93.261101 [arXiv:gr-qc/0411113 [gr-qc]].
- [39] A. S. Bolton, S. Rappaport and S. Burles, "Constraint on the Post-Newtonian Parameter gamma on Galactic Size Scales," Phys. Rev. D 74 (2006), 061501(R) doi:10.1103/PhysRevD.74.061501 [arXiv:astro-ph/0607657 [astro-ph]].
- [40] N. Wei-Tou, "Theoretical frameworks for testing relativistic gravity. iv. a compendium of metric theories of gravity and their post-newtonian limits," Astrophys. J. 176 (1972), 769-796 doi:10.1086/151677
- [41] S. Weinberg, "Gravitation and Cosmology: Principles

and Applications of the General Theory of Relativity," John Wiley & Sons (1972), ISBN: 9780471925675, 9780471925675 (Print).

- [42] C. W. Misner, K. S. Thorne and J. A. Wheeler, "Gravitation," Princeton University Press (1973), ISBN: 0716703440, 9780716703440.
- [43] C. P. Burgess, R. C. Myers and F. Quevedo, "On spherically symmetric string solutions in four-dimensions," Nucl. Phys. B 442 (1995), 75-96 doi:10.1016/S0550-3213(95)00090-9 [arXiv:hep-th/9410142 [hep-th]].
- [44] K. Cho, K. Morand and J. H. Park, "Stringy Newton Gravity with *H*-flux," Phys. Rev. D 101 (2020) no.6, 064020 doi:10.1103/PhysRevD.101.064020 [arXiv:1912.13220 [hep-th]].
- [45] S. Angus, K. Cho, G. Franzmann, S. Mukohyama and J. H. Park, "O(D, D) completion of the Friedmann equations," Eur. Phys. J. C 80 (2020) no.9, 830 doi:10.1140/epjc/s10052-020-8379-7 [arXiv:1905.03620 [hep-th]].
- [46] M. Laue, "Zur Dynamik der Relativitätstheorie," Annalen Phys. **340** (1911) no.8, 524-542 doi:10.1002/andp.19113400808
- [47] C. D. A. Blair, "Conserved Currents of Double Field Theory," JHEP 04 (2016), 180 doi:10.1007/JHEP04(2016)180 [arXiv:1507.07541 [hepth]].
- [48] J. H. Park, S. J. Rey, W. Rim and Y. Sakatani, "O(D, D) covariant Noether currents and global charges in double field theory," JHEP **1511** (2015) 131 [arXiv:1507.07545 [hep-th]].
- [49] O. Hohm and B. Zwiebach, "On the Riemann Tensor in Double Field Theory," JHEP 05 (2012), 126 doi:10.1007/JHEP05(2012)126 [arXiv:1112.5296 [hepth]].
- [50] "NASA/Marshall Solar Physics", Marshall Space Flight Center (2007), https://solarscience.msfc.nasa.gov/.
- [51] V. D. Burkert, L. Elouadrhiri and F. X. Girod, "The pressure distribution inside the proton," Nature 557 (2018) no.7705, 396-399 doi:10.1038/s41586-018-0060-z
- [52] X. D. Ji, "A QCD analysis of the mass structure of the nucleon," Phys. Rev. Lett. **74** (1995), 1071-1074 doi:10.1103/PhysRevLett.74.1071 [arXiv:hep-ph/9410274 [hep-ph]].
- [53] A. I. Vainshtein, V. I. Zakharov and M. A. Shifman, "A Possible mechanism for the Delta T = 1/2 rule in nonleptonic decays of strange particles," JETP Lett. **22** (1975), 55-56
- [54] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, "Light Quarks and the Origin of the Delta I = 1/2 Rule in the Nonleptonic Decays of Strange Particles," Nucl. Phys. B **120** (1977), 316-324 doi:10.1016/0550-3213(77)90046-3
- [55] A. Casher and L. Susskind, "Chiral magnetism (or magnetohadrochironics)," Phys. Rev. D 9 (1974), 436-460 doi:10.1103/PhysRevD.9.436
- [56] P. Maris, C. D. Roberts and P. C. Tandy, "Pion mass and decay constant," Phys. Lett. B 420 (1998), 267-273 doi:10.1016/S0370-2693(97)01535-9 [arXiv:nucl-th/9707003 [nucl-th]].
- [57] P. Maris and C. D. Roberts, "Pi- and K meson Bethe-Salpeter amplitudes," Phys. Rev. C 56 (1997), 3369-3383 doi:10.1103/PhysRevC.56.3369 [arXiv:nucl-th/9708029 [nucl-th]].
- [58] S. J. Brodsky and R. Shrock, "On Condensates in

Strongly Coupled Gauge Theories," [arXiv:0803.2541 [hep-th]].

- [59] S. J. Brodsky and R. Shrock, "Standard-Model Condensates and the Cosmological Constant," Science 108 (2011), 45-50 [arXiv:0803.2554 [hep-th]].
- [60] S. J. Brodsky and R. Shrock, "Condensates in Quantum Chromodynamics and the Cosmological Constant," Proc. Nat. Acad. Sci. 108 (2011), 45-50 doi:10.1073/pnas.1010113107 [arXiv:0905.1151 [hep-th]].
- [61] S. J. Brodsky, C. D. Roberts, R. Shrock and P. C. Tandy, "Essence of the vacuum quark condensate," Phys. Rev. C 82 (2010), 022201(R) doi:10.1103/PhysRevC.82.022201 [arXiv:1005.4610 [nucl-th]].
- [62] L. Chang, C. D. Roberts and P. C. Tandy, "Expanding the concept of in-hadron condensates," Phys. Rev. C 85 (2012), 012201(R) doi:10.1103/PhysRevC.85.012201 [arXiv:1109.2903 [nucl-th]].
- [63] S. J. Brodsky, C. D. Roberts, R. Shrock and P. C. Tandy, "Confinement contains condensates," Phys. Rev. C 85 (2012), 065202 doi:10.1103/PhysRevC.85.065202 [arXiv:1202.2376 [nucl-th]].
- [64] R. Barate et al. [ALEPH], "Studies of quantum chromodynamics with the ALEPH detector," Phys. Rept. 294 (1998), 1-165 doi:10.1016/S0370-1573(97)00045-8
- [65] B. V. Geshkenbein, B. L. Ioffe and K. N. Zybablyuk, "The Check of QCD based on the tau - decay data analysis in the complex q**2 - plane," Phys. Rev. D 64 (2001), 093009 doi:10.1103/PhysRevD.64.093009 [arXiv:hep-ph/0104048 [hep-ph]].
- [66] B. L. Ioffe and K. N. Żyablyuk, "Gluon condensate in

charmonium sum rules with three loop corrections," Eur. Phys. J. C **27** (2003), 229-241 doi:10.1140/epjc/s2002-01099-8 [arXiv:hep-ph/0207183 [hep-ph]].

- [67] M. Davier, A. Hocker and Z. Zhang, "ALEPH Tau Spectral Functions and QCD," Nucl. Phys. B Proc. Suppl. 169 (2007), 22-35 doi:10.1016/j.nuclphysbps.2007.02.109 [arXiv:hep-ph/0701170 [hep-ph]].
- [68] M. Davier, S. Descotes-Genon, A. Hocker, B. Malaescu and Z. Zhang, "The Determination of alpha(s) from Tau Decays Revisited," Eur. Phys. J. C 56 (2008), 305-322 doi:10.1140/epjc/s10052-008-0666-7 [arXiv:0803.0979 [hep-ph]].
- [69] J. H. Park and S. W. Kim, "Existence of a critical point in the phase diagram of the ideal relativistic neutral Bose gas," New J. Phys. **13** (2011), 033003 doi:10.1088/1367-2630/13/3/033003 [arXiv:1001.1823 [cond-mat.quantgas]].
- [70] C. P. Burgess and F. Quevedo, "Axion Homeopathy: Screening Dilaton Interactions," [arXiv:2110.10352 [hepth]].
- [71] A. A. Tseytlin and C. Vafa, "Elements of string cosmology," Nucl. Phys. B **372** (1992), 443-466 doi:10.1016/0550-3213(92)90327-8 [arXiv:hep-th/9109048 [hep-th]].
- [72] J. K. Webb, V. V. Flambaum, C. W. Churchill, M. J. Drinkwater and J. D. Barrow, "Evidence for time variation of the fine structure constant," Phys. Rev. Lett. 82 (1999), 884-887 doi:10.1103/PhysRevLett.82.884 [arXiv:astro-ph/9803165 [astro-ph]].