COMPLETENESS OF BETHE ANSATZ FOR GAUDIN MODELS ASSOCIATED WITH $\mathfrak{gl}(1|1)$

KANG LU

ABSTRACT. We study the Gaudin models associated with $\mathfrak{gl}(1|1)$. We give an explicit description of the algebra of Hamiltonians (Gaudin Hamiltonians) acting on tensor products of polynomial evaluation $\mathfrak{gl}(1|1)[t]$ -modules. It follows that there exists a bijection between common eigenvectors (up to proportionality) of the algebra of Hamiltonians and monic divisors of an explicit polynomial written in terms of the highest weights and evaluation parameters. In particular, our result implies that each common eigenspace of the algebra of Hamiltonians has dimension one. Therefore, we confirm [HMVY19, Conjecture 8.3]. We also give dimensions of the generalized eigenspaces. Moreover, we express the generating pseudo-differential operator of Gaudin transfer matrices associated to antisymmetrizers in terms of the quadratic Gaudin transfer matrix and the center of $U(\mathfrak{gl}(1|1)[t])$.

Keywords: Gaudin models, Bethe ansatz, pseudo-differential operators.

1. INTRODUCTION

Supersymmetric integrable models have been studied extensively for the last 4 decades since their introduction back to 1980s [KS82, Kul85]. However, the vast majority of work are done for spin chains while there is far less information about Gaudin models associated to Lie superalgebras.

This paper is devoted to the study of the Gaudin models associated to the simplest Lie superalgebra $\mathfrak{gl}(1|1)$. Our main motivation is to prove the Bethe ansatz conjecture for the $\mathfrak{gl}(1|1)$ Gaudin models defined on tensor products of polynomial modules, which is formulated in the form of [HMVY19, Conjecture 8.3]. We expect the results of this paper is important to understand the completeness of Bethe ansatz for $\mathfrak{gl}(m|n)$ Gaudin models, see [HMVY19, Conjecture 8.2] and cf. [CLV20, MNV20].

Our main tool is the statements of Bethe ansatz for $\mathfrak{gl}(1|1)$ Gaudin models that are established in the generic situation, see [MVY15]. For generic parameters or more specially in the case when the Bethe ansatz equation has no multiple roots, the Bethe ansatz describes completely the spectrum of the Gaudin Hamiltonians, and the Gaudin Hamiltonians are simultaneously diagonalizable with simple spectrum. The more subtle case is when the Bethe ansatz equation has roots of non-trivial multiplicity, then the Gaudin Hamiltonians develop Jordan blocks. To attack this, we describe the image of algebra of Hamiltonians (Bethe algebra) explicitly and show that the Gaudin model for $\mathfrak{gl}(1|1)$ is perfectly integrable, see [Lu20]. The perfect integrability introduced in [Lu20] is motivated by [MTV09] to study the completeness of Bethe ansatz. Here by perfect integrability, we mean the algebra of Hamiltonians acts on the Hilbert space \mathbb{V} cyclically and the image of the algebra of Hamiltonians in End(\mathbb{V}) is a Frobenius algebra.

Let us discuss our findings in more detail.

We consider tensor products $L_{\Lambda}(\boldsymbol{b}) = \bigotimes_{s=1}^{k} L_{\lambda^{(s)}}(b_s)$ of polynomial evaluation modules of $U(\mathfrak{gl}(1|1)[t])$, where $\Lambda = (\lambda^{(1)}, \ldots, \lambda^{(k)})$ is a sequence of polynomial $\mathfrak{gl}(1|1)$ -weights and $\boldsymbol{b} = (b_1, \ldots, b_k)$ is a sequence of distinct complex numbers. Set $\lambda^{(s)} = (\alpha_s, \beta_s)$.

Define the Bethe algebra to be the unital subalgebra of $U(\mathfrak{gl}(1|1)[t])$ generated by the coefficients of the (quadratic) Gaudin transfer matrix and the center of $U(\mathfrak{gl}(1|1)[t])$. Here the Gaudin transfer matrix, see (2.4), can be thought as the generating function of Gaudin Hamiltonians, see (2.5). We need to find the spectrum of the Gaudin transfer matrix acting on the subspace $(L_{\Lambda}(b))_{(n-l,l)}^{\text{sing}}$ of singular vectors in $L_{\Lambda}(b)$ of weight (n-l,l) where $n = \sum_{s=1}^{k} (\alpha_s + \beta_s)$ and $0 \leq l < k$. Indeed, we obtain more than that. Let us list our main results about completeness of Bethe ansatz, see Theorem 4.8.

- We prove that the Bethe algebra acts on $(L_{\Lambda}(\boldsymbol{b}))_{(n-l,l)}^{\text{sing}}$ cyclically and hence its image in $\operatorname{End}((L_{\Lambda}(\boldsymbol{b}))_{(n-l,l)}^{\text{sing}})$ has dimension $\binom{k-1}{l}$.
- We further show that the image of the Bethe algebra in $\operatorname{End}((L_{\Lambda}(b))_{(n-l,l)}^{\operatorname{sing}})$ is isomorphic to

$$\mathbb{C}[w_1,\ldots,w_{k-1}]^{\mathfrak{S}_l\times\mathfrak{S}_{k-l-1}}/\langle n\prod_{i=1}^k(x-w_i)-\varphi_{\mathbf{\Lambda},\mathbf{b}}(x)\rangle,$$

where

$$\varphi_{\mathbf{\Lambda},\mathbf{b}}(x) := \prod_{s=1}^{k} (x - b_s) \sum_{s=1}^{k} \frac{\alpha_s + \beta_s}{x - b_s}.$$

In particular, the image of the Bethe algebra is a Frobenius algebra. Thus, we establish the perfect integrability of the Gaudin models for $\mathfrak{gl}(1|1)$ in the sense of [Lu20].

 Consequently, we obtain that the eigenvectors (up to proportionality) of the Bethe algebra in (L_Λ(b))^{sing}_(n-l,l) are in a bijective correspondence with the monic polynomials of degree l which divide φ_{Λ,b}(x). And to each monic divisor y we have exactly one eigenvector of the Bethe algebra and a generalized eigenspace of dimension given by the product of binomial coefficients

$$\prod_{a \in \mathbb{C}} \binom{\operatorname{Mult}_a(\varphi_{\mathbf{\Lambda}, \mathbf{b}}(x))}{\operatorname{Mult}_a(y(x))},$$

where $\operatorname{Mult}_{a}(f)$ denotes the order of zero of f(x) at x = a.

The proof of aforementioned statements are based on the idea of [MTV09], cf. [LM21a]. Note that the perfect integrability for the inhomogeneous $\mathfrak{gl}(m|n)$ Gaudin models with diagonal twists defined on tensor products of symmetric powers of the vector representations was established in [HM20, Corollary 5.3] by studying duality between $\mathfrak{gl}(m|n)$ and $\mathfrak{gl}(k)$ Gaudin models and using the known results from [MTV08].

Finally, we give a description of the eigenvalues/eigenvectors of the Bethe algebra in terms of "opers" under the philosophy of the standard geometric Langlands. Given a monic divisor y of $\varphi_{\Lambda,b}(x)$, or, in other words, an eigenvector v_y of the Bethe algebra, following [HMVY19], we have a pseudo-differential operator \mathcal{D}_y , see the right hand side of (4.9). From the explicit formula for the

3

eigenvalue, one sees that the coefficients of the pseudo-differential operator in this case are essentially eigenvalues of the first two Gaudin transfer matrices acting on v_y . Again, we improve on that as follows. Let $L(x) = ((-1)^{|i|}e_{ij}(x))_{i,j=1,2}$ be the generating matrix of the algebra $U(\mathfrak{gl}(1|1)[t])$. By [MR14], the Berezinian $\operatorname{Ber}(\partial_x - L(x))$ is a generating function for the Gaudin transfer matrices, see (4.7) and (4.8). By $\operatorname{Ber}(\partial_x - L(x))v_y = \mathcal{D}_y v_y$, see Theorem 4.11, it follows that there is a universal formula for the pseudo-differential operator in terms of the first two Gaudin transfer matrices, which produces \mathcal{D}_y when applied to the vector v_y for all y, see Corollary 4.14.

The theory of Bethe ansatz has been developed for Gaudin models associated with $\mathfrak{osp}(1|2)$ in [KM01]. We plan to further study the completeness of Bethe ansatz for $\mathfrak{osp}(1|2)$ Gaudin models, cf. [LM21c].

The paper is organized as follows. In Section 2, we fix notations and discuss basic facts of the algebraic Bethe ansatz for $\mathfrak{gl}(1|1)$ Gaudin models. Then we study the space $\mathcal{V}^{\mathfrak{S}}$ and Weyl modules, and their properties in Section 3. Section 4 contains the main theorems where we also discuss the higher Gaudin transfer matrices and the relations between higher Gaudin transfer matrices and the first two Gaudin transfer matrices. Section 5 is dedicated to the proofs of main theorems.

Acknowledgments. The author thanks C.-L. Huang and E. Mukhin for interesting discussions.

2. PRELIMINARIES

2.1. Lie superalgebra $\mathfrak{gl}(1|1)$ and its representations. A vector superspace $V = V_{\bar{0}} \oplus V_{\bar{1}}$ is a \mathbb{Z}_2 -graded vector space. Elements of $V_{\bar{0}}$ are called *even*; elements of $V_{\bar{1}}$ are called *odd*. We write $|v| \in \{\bar{0}, \bar{1}\}$ for the parity of a homogeneous element $v \in V$. Set $(-1)^{\bar{0}} = 1$ and $(-1)^{\bar{1}} = -1$.

Consider the vector superspace $\mathbb{C}^{1|1}$, where $\dim(\mathbb{C}_{\bar{0}}^{1|1}) = 1$ and $\dim(\mathbb{C}_{\bar{1}}^{1|1}) = 1$. We choose a homogeneous basis v_1, v_2 of $\mathbb{C}^{1|1}$ such that $|v_1| = \bar{0}$ and $|v_2| = \bar{1}$. For brevity we shall write their parities as $|v_i| = |i|$. Denote by $E_{ij} \in \operatorname{End}(\mathbb{C}^{1|1})$ the linear operator of parity |i| + |j| such that $E_{ij}v_r = \delta_{jr}v_i$ for i, j, r = 1, 2.

The Lie superalgebra $\mathfrak{gl}(1|1)$ is spanned by elements e_{ij} , i, j = 1, 2, with parities $e_{ij} = |i| + |j|$ and the supercommutator relations are given by

$$[e_{ij}, e_{rs}] = \delta_{jr} e_{is} - (-1)^{(|i|+|j|)(|r|+|s|)} \delta_{is} e_{rj}.$$

Let \mathfrak{h} be the commutative Lie subalgebra of $\mathfrak{gl}(1|1)$ spanned by e_{11}, e_{22} . Denote the universal enveloping algebras of $\mathfrak{gl}_{1|1}$ and \mathfrak{h} by $U(\mathfrak{gl}_{1|1})$ and $U(\mathfrak{h})$, respectively.

We call a pair $\lambda = (\lambda_1, \lambda_2)$ of complex numbers a $\mathfrak{gl}(1|1)$ -weight. Set $|\lambda| = \lambda_1 + \lambda_2$. A $\mathfrak{gl}(1|1)$ -weight λ is non-degenerate if $\lambda_1 + \lambda_2 \neq 0$.

Let M be a $\mathfrak{gl}(1|1)$ -module. A non-zero vector $v \in M$ is called *singular* if $e_{12}v = 0$. Denote the subspace of all singular vectors of M by $(M)^{\text{sing}}$. A non-zero vector $v \in M$ is called *of weight* $\lambda = (\lambda_1, \lambda_2)$ if $e_{11}v = \lambda_1 v$ and $e_{22}v = \lambda_2 v$. Denote by $(M)_{\lambda}$ the subspace of M spanned by vectors of weight λ . Set $(M)^{\text{sing}}_{\lambda} = (M)^{\text{sing}} \cap (M)_{\lambda}$.

Let $\Lambda = (\lambda^{(1)}, \dots, \lambda^{(k)})$ be a sequence of $\mathfrak{gl}(1|1)$ -weights. Set $|\Lambda| = \sum_{s=1}^{k} |\lambda^{(s)}|$.

Denote by L_{λ} the irreducible $\mathfrak{gl}(1|1)$ -module generated by an even singular vector v_{λ} of weight λ . Then L_{λ} is two-dimensional if λ is non-degenerate and one-dimensional otherwise. Clearly, $\mathbb{C}^{1|1} \cong L_{\omega_1}$, where $\omega_1 = (1, 0)$.

A $\mathfrak{gl}(1|1)$ -module M is called a *polynomial module* if M is a submodule of $(\mathbb{C}^{1|1})^{\otimes n}$ for some $n \in \mathbb{Z}_{\geq 0}$. We say that λ is a *polynomial weight* if L_{λ} is a polynomial module. Weight $\lambda = (\lambda_1, \lambda_2)$ is a polynomial weight if and only if $\lambda_1, \lambda_2 \in \mathbb{Z}_{\geq 0}$ and either $\lambda_1 > 0$ or $\lambda_1 = \lambda_2 = 0$. We also write $L_{(\lambda_1, \lambda_2)}$ for L_{λ} .

For non-degenerate polynomial weights $\lambda = (\lambda_1, \lambda_2)$ and $\mu = (\mu_1, \mu_2)$, we have

$$L_{(\lambda_1,\lambda_2)} \otimes L_{(\mu_1,\mu_2)} = L_{(\lambda_1+\mu_1,\lambda_2+\mu_2)} \oplus L_{(\lambda_1+\mu_1-1,\lambda_2+\mu_2+1)}$$

2.2. Current superalgebra $\mathfrak{gl}(1|1)[t]$. Denote by $\mathfrak{gl}(1|1)[t]$ the Lie superalgebra $\mathfrak{gl}(1|1) \otimes \mathbb{C}[t]$ of $\mathfrak{gl}_{1|1}$ -valued polynomials with the point-wise supercommutator. Call $\mathfrak{gl}(1|1)[t]$ the *current superal-gebra* of $\mathfrak{gl}(1|1)$. We identify $\mathfrak{gl}(1|1)$ with the subalgebra $\mathfrak{gl}(1|1) \otimes 1$ of constant polynomials in $\mathfrak{gl}(1|1)[t]$.

We write $e_{ij}[r]$ for $e_{ij} \otimes t^r$, $r \in \mathbb{Z}_{\geq 0}$. A basis of $\mathfrak{gl}(1|1)[t]$ is given by $e_{ij}[r]$, i, j = 1, 2 and $r \in \mathbb{Z}_{\geq 0}$. They satisfy the supercommutator relations

$$[e_{ij}[r], e_{kl}[s]] = \delta_{jk} e_{il}[r+s] - (-1)^{(|i|+|j|)(|k|+|l|)} \delta_{il} e_{kj}[r+s].$$

In particular, one has

$$(e_{12}[r])^2 = (e_{21}[r])^2 = 0, \quad e_{21}[r]e_{21}[s] = -e_{21}[s]e_{21}[r]$$
 (2.1)

in the universal enveloping superalgebra $U(\mathfrak{gl}(1|1)[t])$. The universal enveloping superalgebra $U(\mathfrak{gl}(1|1)[t])$ is a Hopf superalgebra with the coproduct given by

 $\Delta(X) = X \otimes 1 + 1 \otimes X, \quad \text{for } X \in \mathfrak{gl}(1|1)[t].$

Let $e_{ij}(x) = \sum_{r=0}^{\infty} e_{ij}[r]x^{-r-1}$, where x is a formal variable. For each $a \in \mathbb{C}$, there exists an automorphism of $U(\mathfrak{gl}(1|1)[t])$, $\rho_a : e_{ij}(x) \to e_{ij}(x-a)$. Given a $\mathfrak{gl}(1|1)[t]$ -module M, denote by M(a) the pull-back of M through the automorphism ρ_a .

For each $a \in \mathbb{C}$, we have the evaluation map $ev_a : e_{ij}(x) \mapsto e_{ij}/(x-a)$. For a $\mathfrak{gl}(1|1)$ -module L, denote by L(a) the $\mathfrak{gl}(1|1)[t]$ -module obtained by pulling back L through the evaluation map ev_a . We call L(a) an *evaluation module* at a.

Given any series $\zeta(x) \in x^{-1}\mathbb{C}[x^{-1}]$, we have the one-dimensional $\mathfrak{gl}(1|1)[t]$ -module generated by an even vector v satisfying $e_{ij}(x)v = \delta_{ij}(-1)^{|j|}\zeta(x)v$. We denote this module by \mathbb{C}_{ζ} .

If b_1, \ldots, b_n are pairwise distinct complex numbers and L_1, \ldots, L_n are finite-dimensional irreducible $\mathfrak{gl}(1|1)$ -modules, then the $\mathfrak{gl}(1|1)[t]$ -module $\bigotimes_{s=1}^n L_s(b_s)$ is irreducible.

There is a natural $\mathbb{Z}_{\geq 0}$ -gradation on $U(\mathfrak{gl}(1|1)[t])$ such that $\deg(e_{ij}[r]) = r$. Let M be a $\mathbb{Z}_{\geq 0}$ -graded space with finite-dimensional homogeneous components. Let $M_j \subset M$ be the homogeneous component of degree j. We call the formal power series in variable q,

$$\operatorname{ch}(M) = \sum_{j=0}^{\infty} \dim(M_j) q^j$$
(2.2)

the graded character of M.

2.3. Gaudin Hamiltonians and Bethe ansatz. In this section, we recall the Gaudin Hamiltonians and the corresponding Bethe ansatz from [MVY15].

Let $\Lambda = (\lambda^{(1)}, \ldots, \lambda^{(k)})$ be a sequence of polynomial $\mathfrak{gl}(1|1)$ -weights and $b = (b_1, \ldots, b_k)$ a sequence of distinct complex numbers, where $\lambda^{(s)} = (\alpha_s, \beta_s)$. Set $n = |\Lambda| = \sum_{s=1}^k (\alpha_s + \beta_s)$ and $L_{\Lambda} = \bigotimes_{s=1}^k L_{\lambda^{(s)}}$. The quadratic Gaudin Hamiltonians are the linear maps $\mathcal{H}_r \in \operatorname{End}(L_{\Lambda})$ given by

$$\mathcal{H}_r := \sum_{s=1}^k \frac{e_{11}^{(r)} e_{11}^{(s)} - e_{12}^{(r)} e_{21}^{(s)} + e_{21}^{(r)} e_{12}^{(s)} - e_{22}^{(r)} e_{22}^{(s)}}{b_r - b_s}, \quad 1 \le r \le k.$$
(2.3)

where $e_{ab}^{(r)} = 1^{\otimes (r-1)} \otimes e_{ab} \otimes 1^{\otimes (k-r)}$.

Lemma 2.1 ([MVY15, Proposition 3.1]). The Gaudin Hamiltonians \mathcal{H}_r

- (i) are mutually commuting: $[\mathcal{H}_r, \mathcal{H}_s] = 0$ for all r, s;
- (ii) commute with the action of $\mathfrak{gl}(1|1)$: $[\mathcal{H}_r, X] = 0$ for all r and $X \in \mathfrak{gl}(1|1)$.

Instead of working on Gaudin Hamiltonians \mathcal{H}_s , we work on the generating function of Gaudin Hamiltonians,

$$\mathcal{H}(x) := \sum_{r=2}^{\infty} \mathcal{H}_r x^{-r} = \frac{1}{2} \sum_{a,b=1}^{2} e_{ab}(x) e_{ba}(x) (-1)^{|b|}.$$
(2.4)

The operator $\mathcal{H}(x)$ acts on the tensor product of the evaluation $\mathfrak{gl}(1|1)[t]$ -modules

$$L_{\mathbf{\Lambda}}(\mathbf{b}) := \bigotimes_{s=1}^{k} L_{\mathbf{\lambda}^{(s)}}(b_s).$$

Note that $L_{\Lambda}(b)$ and L_{Λ} are isomorphic as $\mathfrak{gl}(1|1)$ -modules via the identity map, then we have

$$\mathcal{H}(x) = \frac{1}{2} \sum_{s=1}^{k} \frac{\alpha_s(\alpha_s - 1) - \beta_s(\beta_s + 1)}{(x - b_s)^2} \mathrm{Id} + \sum_{s=1}^{k} \frac{1}{x - b_s} \mathcal{H}_s,$$
(2.5)

as operators in $\operatorname{End}(L_{\Lambda}) = \operatorname{End}(L_{\Lambda}(b))$. We call $\mathcal{H}(x)$ the Gaudin transfer matrix.

We are interested in finding the eigenvalues and eigenvectors of the Gaudin transfer matrix in $L_{\Lambda}(b)$. To be more precise, we call

$$\xi(x) = \sum_{r=2}^{\infty} \xi_r x^{-r}, \qquad \xi_r \in \mathbb{C},$$
(2.6)

an *eigenvalue* of $\mathcal{H}(x)$ if there exists a non-zero vector $v \in L_{\Lambda}(\mathbf{b})$ such that $\mathcal{H}_r v = \xi_r v$ for all $r \in \mathbb{Z}_{>1}$. If $\xi(x)$ is a rational function, we consider it as a power series in x^{-1} as (2.6). The vector v is called an *eigenvector* of $\mathcal{H}(x)$ corresponding to eigenvalue $\xi(x)$. We also define the *eigenspace* of $\mathcal{H}(x)$ in $L_{\Lambda}(\mathbf{b})$ corresponding to eigenvalue $\xi(x)$ as $\bigcap_{r=2}^{\infty} \ker(\mathcal{H}_r|_{L_{\Lambda}(\mathbf{b})} - \xi_r)$.

It is sufficient to consider L_{Λ} with $\beta_s = 0$ for all s. Indeed, if $L_{\Lambda}(b)$ is an arbitrary tensor product and

$$\xi(x) = \sum_{s=1}^{k} \frac{\beta_s}{x - b_s},$$

then

$$L_{\Lambda}(\boldsymbol{b}) \otimes \mathbb{C}_{\xi} \cong L_{\widetilde{\Lambda}}(\boldsymbol{b}), \quad \boldsymbol{\lambda}^{(s)} = (\alpha_s + \beta_s, 0)$$

Identify $L_{\Lambda}(\mathbf{b}) \otimes \mathbb{C}_{\xi}$ with $L_{\Lambda}(\mathbf{b})$ as vector spaces. Then $\mathcal{H}(x)$ acting on $L_{\Lambda}(\mathbf{b}) \otimes \mathbb{C}_{\xi}$ coincides with $\mathcal{H}(x) + 2\zeta(x)(e_{11}(x) + e_{22}(x))$ acting on $L_{\Lambda}(\mathbf{b})$. Note that the coefficients of $e_{11}(x) + e_{22}(x)$ are central in $U(\mathfrak{gl}(1|1)[t])$ and hence acts on $L_{\Lambda}(\mathbf{b})$ by scalars, therefore the problem of diagonalization of the Gaudin transfer matrix in $L_{\Lambda}(\mathbf{b})$ is reduced to diagonalization of the Gaudin transfer matrix in $L_{\widetilde{\Lambda}}(\mathbf{b})$.

Since L_{λ} is one-dimensional if λ is degenerate, similarly, it suffices to consider the case that all participant $\mathfrak{gl}(1|1)$ -weights are non-degenerate. Hence, we shall always assume throughout the paper that $\lambda^{(s)}$ are non-degenerate for all $1 \leq s \leq k$.

The main method to find eigenvalues and eigenvectors of the Gaudin transfer matrix in L_{Λ} is the algebraic Bethe ansatz which we recall from [MVY15].

Fix a non-negative integer l. Let $t = (t_1, \ldots, t_l)$ be a sequence of complex numbers. Define the polynomial $y_t = \prod_{i=1}^l (x - t_i)$. We say that polynomial y_t represents t.

Set

$$\zeta_{\mathbf{\Lambda},\mathbf{b}}(x) := \sum_{s=1}^{k} \frac{\alpha_s + \beta_s}{x - b_s}.$$
(2.7)

A sequence of complex numbers t is called a *solution to the Bethe ansatz equation associated to* Λ , b, l if

$$y_t(x)$$
 divides the polynomial $\varphi_{\mathbf{\Lambda}, \mathbf{b}}(x) := \zeta_{\mathbf{\Lambda}, \mathbf{b}}(x) \prod_{s=1}^k (x - b_s).$ (2.8)

We do not distinguish solutions which differ by a permutation of coordinates (that is represented by the same polynomial).

Let v_s be the highest weight vector of $L_{\lambda^{(s)}}$, and set $|0\rangle = v_1 \otimes \cdots \otimes v_k$. We call $|0\rangle$ the vacuum vector.

Define the *off-shell Bethe vector* $\mathbb{B}_l(t) \in (L_{\Lambda})_{(n-l,l)}$ by

$$\mathbb{B}_{l}(\boldsymbol{t}) = e_{12}(t_{1}) \cdots e_{12}(t_{l}) |0\rangle.$$
(2.9)

Since $e_{12}(x)e_{12}(u) = -e_{12}(u)e_{12}(x)$, the order of t_i is not important. Moreover, the off-shell Bethe vector is zero if $t_i = t_j$ for some $1 \le i \ne j \le l$.

If t is a solution of the Bethe ansatz equation (2.8), we call $\mathbb{B}_l(t)$ an *on-shell Bethe vector*.

Let t be a solution of the Bethe ansatz equation associated to Λ , b, l. The following statements are known, see [MVY15, Section VI].

Theorem 2.2 ([MVY15]). If the on-shell Bethe vector $\mathbb{B}_l(t)$ is non-zero, then $\mathbb{B}_l(t)$ is an eigenvector of the Gaudin transfer matrix $\mathfrak{H}(x)$ with the corresponding eigenvalue

$$\mathcal{E}_{y_{t},\Lambda,b}(x) = \frac{1}{2}\zeta'_{\Lambda,b}(x) - \zeta_{\Lambda,b}(x)\frac{y'_{t}(x)}{y_{t}(x)} + \sum_{r,s=1}^{k} \frac{\alpha_{r}\alpha_{s} - \beta_{r}\beta_{s}}{2(x-b_{r})(x-b_{s})}.$$
(2.10)

where $\zeta_{\Lambda,b}(x)$ is given by (2.7).

Consider another Gaudin transfer matrix, see Example 4.9,

$$\mathcal{T}(x) = \frac{1}{2} \left(\dot{e}_{11}(x) + \dot{e}_{22}(x) \right) + \frac{1}{2} \left(e_{11}(x) + e_{22}(x) \right)^2 - \mathcal{H}(x), \tag{2.11}$$

where $\dot{e}_{ii}(x) = \partial_x(e_{ii}(x))$, i = 1, 2. Then the eigenvalue of $\mathcal{T}(x)$ acting on the on-shell Bethe vector $\mathbb{B}_l(t)$ is

$$\mathcal{E}_{y_{t},\Lambda,b}(x) = \zeta_{\Lambda,b}(x) \frac{y_{t}'(x)}{y_{t}(x)} + \sum_{r,s=1}^{k} \frac{\alpha_{r}\beta_{s} + \alpha_{s}\beta_{r} + 2\beta_{r}\beta_{s}}{2(x-b_{r})(x-b_{s})}$$

$$= \zeta_{\Lambda,b}(x) \Big(\frac{y_{t}'(x)}{y_{t}(x)} + \sum_{s=1}^{k} \frac{\beta_{s}}{x-b_{s}}\Big).$$
(2.12)

Proposition 2.3 ([MVY15]). *The on-shell Bethe vector* $\mathbb{B}_l(t)$ *is singular.*

It is important to know if the on-shell Bethe vectors are non-zero.

Proposition 2.4 ([MVY15]). Suppose the polynomial $\varphi_{\Lambda,b}(x)$ only has simple roots, then the onshell Bethe vector $\mathbb{B}_l(t)$ is nonzero.

The conjecture of completeness of Bethe ansatz for Gaudin models associated with $\mathfrak{gl}(1|1)$ was reformulated in [HMVY19, Conjecture 8.2] as follows.

Conjecture 2.5. Suppose all weights $\lambda^{(s)}$, $1 \leq s \leq k$, are polynomial $\mathfrak{gl}(1|1)$ -weights. Then the Gaudin transfer matrix $\mathfrak{H}(x)$ has a simple spectrum in $(L_{\Lambda}(\mathbf{b}))^{\text{sing}}$. There exists a bijectively correspondence between the monic divisors y of the polynomial $\varphi_{\Lambda,b}$ and the eigenvectors v of the Gaudin transfer matrices (up to multiplication by a non-zero constant). Moreover, this bijection is such that $\mathfrak{H}(x)v = \mathcal{E}_{y,\Lambda,b}(x)v$, where $\mathcal{E}_{y,\Lambda,b}(x)$ is given by (2.10).

By simple spectrum we mean that if v_1 , v_2 are eigenvectors of $\mathcal{H}(x)$ and $v_1 \neq cv_2$, $c \in \mathbb{C}^{\times}$, then the eigenvalues of $\mathcal{H}(x)$ on v_1 and v_2 are different.

The conjecture follows from Theorem 4.8 proved in Section 5.3.

The conjecture is clear for the case when $\varphi_{\Lambda,b}$ only has simple roots which we also recall from [MVY15]. Note that dim $L_{\Lambda}(b) = 2^k$ and dim $(L_{\Lambda}(b))^{\text{sing}} = 2^{k-1}$. If the polynomial $\varphi_{\Lambda,b}$ has no multiple roots, then $\varphi_{\Lambda,b}$ has the desired number of distinct monic divisors. Therefore, we have desired number of on-shell Bethe vectors which are also nonzero by Proposition 2.4. By Theorem 2.2, it implies that we do have an eigenbasis of the Gaudin transfer matrix consisting of on-shell Bethe vectors in $(L_{\Lambda}(b))^{\text{sing}}$, see Proposition 2.3, with different eigenvalues. Thus the algebraic Bethe ansatz works well for this situation.

Theorem 2.6 ([MVY15]). Suppose all weights $\lambda^{(s)}$, $1 \leq s \leq k$, are polynomial $\mathfrak{gl}(1|1)$ -weights. If the polynomial $\varphi_{\Lambda,b}$ has no multiple roots, then the Gaudin transfer matrix $\mathfrak{H}(x)$ is diagonalizable and the Bethe ansatz is complete. In particular, for any given Λ and generic \mathbf{b} , the Gaudin transfer matrix $\mathfrak{H}(x)$ is diagonalizable and the Bethe ansatz is complete. \Box

3. Space $\mathcal{V}^{\mathfrak{S}}$ and Weyl modules

In this section, we discuss the super-analog of \mathcal{V}^S in [MTV09, Section 2.5], cf. [LM21a, Section 3].

The symmetric group \mathfrak{S}_n acts naturally on $\mathbb{C}[z_1, \ldots, z_n]$ by permuting variables. Denote by $\sigma_i(\boldsymbol{z})$ the *i*-th elementary symmetric polynomial in z_1, \ldots, z_n . The algebra of symmetric polynomials $\mathbb{C}[z_1, \ldots, z_n]^{\mathfrak{S}}$ is freely generated by $\sigma_1(\boldsymbol{z}), \ldots, \sigma_n(\boldsymbol{z})$.

Fix $\ell \in \{0, 1, ..., n\}$. We have a subgroup $\mathfrak{S}_{\ell} \times \mathfrak{S}_{n-\ell} \subset \mathfrak{S}_n$. Then \mathfrak{S}_{ℓ} permutes the first ℓ variables while $\mathfrak{S}_{n-\ell}$ permutes the last $n-\ell$ variables. Denote by $\mathbb{C}[z_1, ..., z_n]^{\mathfrak{S}_{\ell} \times \mathfrak{S}_{n-\ell}}$ the subalgebra of $\mathbb{C}[z_1, ..., z_n]$ consisting of $\mathfrak{S}_{\ell} \times \mathfrak{S}_{n-\ell}$ -invariant polynomials. It is known that $\mathbb{C}[z_1, ..., z_n]^{\mathfrak{S}_{\ell} \times \mathfrak{S}_{n-\ell}}$ is a free $\mathbb{C}[z_1, ..., z_n]^{\mathfrak{S}}$ -module of rank $\binom{n}{\ell}$.

3.1. **Definition of** $\mathcal{V}^{\mathfrak{S}}$. Let $V = (\mathbb{C}^{1|1})^{\otimes n}$ be the tensor power of the vector representation of $\mathfrak{gl}(1|1)$. The $\mathfrak{gl}(1|1)$ -module V has weight decomposition

$$V = \bigoplus_{\ell=0}^{n} (V)_{(n-\ell,\ell)}.$$

Let \mathcal{V} be the space of polynomials in variables $\boldsymbol{z} = (z_1, z_2, \dots, z_n)$ with coefficients in V,

$$\mathcal{V} = V \otimes \mathbb{C}[z_1, z_2, \dots, z_n].$$

The space V is identified with the subspace $V \otimes 1$ of constant polynomials in \mathcal{V} . The space \mathcal{V} has a natural grading induced from the grading on $\mathbb{C}[z_1, \ldots, z_n]$ with $\deg(z_i) = 1$. Namely, the degree of an element $v \otimes p$ in \mathcal{V} is given by the degree of the polynomial p, $\deg(v \otimes p) = \deg p$. Clearly, the space $\operatorname{End}(\mathcal{V})$ has a gradation structure induced from that on \mathcal{V} .

Let $P^{(i,j)}$ be the graded flip operator which acts on the *i*-th and *j*-th factors of *V*. Let s_1, s_2, \ldots , s_{n-1} be the simple permutations of the symmetric group \mathfrak{S}_n . Define the \mathfrak{S}_n -action on \mathcal{V} by the rule:

$$s_i: \boldsymbol{f}(z_1,\ldots,z_n) \mapsto P^{(i,i+1)}\boldsymbol{f}(z_1,\ldots,z_{i+1},z_i,\ldots,z_n),$$

for $f(z_1, \ldots, z_n) \in \mathcal{V}$. Note that the \mathfrak{S}_n -action respects the gradation on \mathcal{V} . Denote the subspace of all vectors in \mathcal{V} invariant with respect to the \mathfrak{S}_n -action by $\mathcal{V}^{\mathfrak{S}}$.

Clearly, the $\mathfrak{gl}(1|1)$ -action on \mathcal{V} commutes with the \mathfrak{S}_n -action on \mathcal{V} and preserves the grading. Therefore, $\mathcal{V}^{\mathfrak{S}}$ is a graded $\mathfrak{gl}(1|1)$ -module. Hence we have the weight decomposition for both $\mathcal{V}^{\mathfrak{S}}$ and $(\mathcal{V}^{\mathfrak{S}})^{\operatorname{sing}}$,

$$\mathcal{V}^{\mathfrak{S}} = \bigoplus_{\ell=0}^{n} (\mathcal{V}^{\mathfrak{S}})_{(n-\ell,\ell)}, \qquad (\mathcal{V}^{\mathfrak{S}})^{\operatorname{sing}} = \bigoplus_{\ell=0}^{n} (\mathcal{V}^{\mathfrak{S}})_{(n-\ell,\ell)}^{\operatorname{sing}}$$

Note that $(\mathcal{V}^{\mathfrak{S}})_{(n-\ell,\ell)}$ and $(\mathcal{V}^{\mathfrak{S}})_{(n-\ell,\ell)}^{\operatorname{sing}}$ are also graded $\mathbb{C}[z_1,\ldots,z_n]^{\mathfrak{S}}$ -modules. The space \mathcal{V} is a $\mathfrak{gl}(1|1)[t]$ -module where $e_{ij}[r]$ acts by

$$e_{ij}[r](p(z_1,\ldots,z_n)w_1\otimes\cdots\otimes w_n)$$

= $p(z_1,\ldots,z_n)\sum_{s=1}^n (-1)^{(|w_1|+\cdots+|w_{s-1}|)(|i|+|j|)} z_s^r w_1\otimes\cdots\otimes e_{ij}w_s\otimes\cdots\otimes w_n,$ (3.1)

for $p(z_1, \ldots, z_n) \in \mathbb{C}[z_1, \ldots, z_n]$ and $w_s \in \mathbb{C}^{1|1}$.

The following lemma is straightforward.

Lemma 3.1. The $\mathfrak{gl}(1|1)[t]$ -action on \mathcal{V} commutes with the \mathfrak{S}_n -action on \mathcal{V} . Both \mathcal{V} and $\mathcal{V}^{\mathfrak{S}}$ are graded $\mathfrak{gl}(1|1)[t]$ -modules.

3.2. Properties of $\mathcal{V}^{\mathfrak{S}}$ and $(\mathcal{V}^{\mathfrak{S}})^{\text{sing}}$. In this section, we recall properties of $\mathcal{V}^{\mathfrak{S}}$ and $(\mathcal{V}^{\mathfrak{S}})^{\text{sing}}$ from [LM21a, Section 3].

Lemma 3.2. The space $(\mathcal{V}^{\mathfrak{S}})_{(n-\ell,\ell)}$ is a free $\mathbb{C}[z_1, \ldots, z_n]^{\mathfrak{S}}$ -module of rank $\binom{n}{\ell}$. In particular, the space $\mathcal{V}^{\mathfrak{S}}$ is a free $\mathbb{C}[z_1, \ldots, z_n]^{\mathfrak{S}}$ -module of rank 2^n .

Set $v^+ = v_1^{\otimes n} = v_1 \otimes \cdots \otimes v_1$.

Lemma 3.3. The $\mathfrak{gl}(1|1)[t]$ -module $\mathcal{V}^{\mathfrak{S}}$ is a cyclic module generated by v^+ .

Lemma 3.4. The set

$$\{e_{21}[r_1]e_{21}[r_2]\cdots e_{21}[r_\ell]v^+ \mid 0 \leqslant r_1 < r_2 < \cdots < r_\ell \leqslant n-1\}$$
(3.2)

is a free generating set of $(\mathcal{V}^{\mathfrak{S}})_{(n-\ell,\ell)}$ over $\mathbb{C}[z_1,\ldots,z_n]^{\mathfrak{S}}$.

Lemma 3.5. The space $(\mathcal{V}^{\mathfrak{S}})_{(n-\ell,\ell)}^{\operatorname{sing}}$ is a free $\mathbb{C}[z_1,\ldots,z_n]^{\mathfrak{S}}$ -module of rank $\binom{n-1}{\ell}$ with a free generating set given by

$$\{e_{12}[0]e_{21}[0]e_{21}[r_1]\cdots e_{21}[r_\ell]v^+, \quad 1 \le r_1 < r_2 < \cdots < r_\ell \le n-1\}.$$
(3.3)

In particular, the space $(\mathcal{V}^{\mathfrak{S}})^{\text{sing}}$ is a free $\mathbb{C}[z_1, \ldots, z_n]^{\mathfrak{S}}$ -module of rank 2^{n-1} .

Set $(q)_r = \prod_{i=1}^r (1 - q^i)$.

Proposition 3.6. We have

$$\operatorname{ch}((\mathcal{V}^{\mathfrak{S}})_{(n-\ell,\ell)}) = \frac{q^{\ell(\ell-1)/2}}{(q)_{\ell}(q)_{n-\ell}}, \qquad \operatorname{ch}((\mathcal{V}^{\mathfrak{S}})_{(n-\ell,\ell)}^{\operatorname{sing}}) = \frac{q^{\ell(\ell+1)/2}}{(q)_{\ell}(q)_{n-1-\ell}(1-q^n)}$$

Given $\mathbf{a} = (a_1, \ldots, a_n) \in \mathbb{C}^n$, let $I_{\mathbf{a}}$ be the ideal of $\mathbb{C}[z_1, \ldots, z_n]^{\mathfrak{S}}$ generated by $\sigma_i(\mathbf{z}) - \mathbf{a}$, $i = 1, \ldots, n$. Then for any \mathbf{a} , by Lemmas 3.1 and 3.2, the quotient space $\mathcal{V}^{\mathfrak{S}}/I_{\mathbf{a}}\mathcal{V}^{\mathfrak{S}}$ is a $\mathfrak{gl}(1|1)[t]$ -module of dimension 2^n over \mathbb{C} . Denote by \overline{v}^+ be the image of v^+ under this quotient.

3.3. Weyl modules. In this section, we discuss a special family of Weyl modules. We are not ware of references about Weyl modules of $U(\mathfrak{gl}(1|1)[t])$, cf. [CLS19], except the quantum affine case [Zha17].

Let $\eta(x)$ be a monic polynomial of degree m, where $m \in \mathbb{Z}_{\geq 0}$,

$$\eta(x) = \sum_{i=0}^{m} \gamma_i x^i, \qquad \gamma_m = 1.$$

Denote by W_n the $\mathfrak{gl}(1|1)[t]$ -module generated by an even vector w subject to the relations:

$$e_{11}(x)w = \eta'(x)/\eta(x)w, \qquad e_{22}(x)w = e_{12}(x)w = 0,$$
 (3.4)

$$\sum_{i=0}^{m} \gamma_i e_{21}[i]w = 0.$$
(3.5)

It is convenient to write (3.5) as $(e_{21} \otimes \eta(t))w = 0$.

Clearly, we have dim $W_{\eta} \leq 2^m$ by PBW theorem and (2.1), (3.5). The module W_{η} is the universal $\mathfrak{gl}(1|1)[t]$ -module satisfying (3.4), (3.5) which we call a *Weyl module*.

If $\eta(x) = (x - b)^m$, we write W_η as $W_m(b)$.

Lemma 3.7. Let $a = (0, ..., 0) \in \mathbb{C}^n$. Then $\mathcal{V}^{\mathfrak{S}}/I_a \mathcal{V}^{\mathfrak{S}}$ is isomorphic to $W_n(0)$ as $\mathfrak{gl}(1|1)[t]$ -modules.

Proof. It is clear that \bar{v}^+ satisfies the relations (3.4) for $\eta(x) = x^n$. In addition, it follows from the equation

$$e_{21}[n]v^{+} = \sum_{i=1}^{n} (-1)^{i-1} \sigma_{i}(\boldsymbol{z}) e_{21}[n-i]v^{+}$$

that \bar{v}^+ also satisfies the relation (3.5) for $\eta(x) = x^n$. Therefore, we have a surjective $\mathfrak{gl}(1|1)[t]$ -module homomorphism $W_n(0) \twoheadrightarrow \mathcal{V}^{\mathfrak{S}}/I_a\mathcal{V}^{\mathfrak{S}}$. By $\dim \mathcal{V}^{\mathfrak{S}}/I_a\mathcal{V}^{\mathfrak{S}} = 2^n \ge \dim W_n(0)$, we obtain that it is also an isomorphism.

In particular, we have dim $W_m(b) = 2^m$.

Lemma 3.8. Let $\eta(x) = \prod_{s=1}^{k} (x-b_s)^{n_s}$, where $b_s \neq b_r$ for $1 \leq s \neq r \leq k$. Then W_η is isomorphic to $\bigotimes_{s=1}^{k} W_{n_s}(b_s)$ as $\mathfrak{gl}(1|1)[t]$ -modules.

Proof. For $1 \leq s \leq k$, let w_s be the highest weight vector of $W_{n_s}(b_s)$ and set $\eta_s(x) = (x - b_s)^{n_s}$. Similar to Lemma 3.7, since dim $W_\eta \leq \dim \bigotimes_{s=1}^k W_{n_s}(b_s)$, it suffices to show that $\bigotimes_{s=1}^k W_{n_s}(b_s)$ is generated by $w^+ := \bigotimes_{s=1}^k w_s$ and w^+ satisfies the relations (3.4), (3.5).

The proof of the fact that $\bigotimes_{s=1}^{k} W_{n_s}(b_s)$ is generated by w^+ is similar to the even case, see e.g. [CP01]. The relations (3.4) are obvious. By $[e_{11}[r], e_{21}[i]] = -e_{21}[r+i]$ and relations (3.4), (3.5), we have

 $(e_{21} \otimes \eta_s(t)t^r)w_s = 0, \qquad r \ge 0, \qquad 1 \le s \le k.$

Therefore, we also have $(e_{21} \otimes \eta(t))w_s = 0$ for all $1 \leq s \leq k$. Now the relation (3.5) follows immediately.

Given sequences $\mathbf{n} = (n_1, \ldots, n_k)$ of nonnegative integers and $\mathbf{b} = (b_1, \ldots, b_s)$ of distinct complex numbers, by Lemma 3.8, we call $\bigotimes_{s=1}^k W_{n_s}(b_s)$ the Weyl module associated with \mathbf{n} and \mathbf{b} .

Given $a = (a_1, \ldots, a_n) \in \mathbb{C}^n$, define $k \in \mathbb{Z}_{>0}$, $b_s \in \mathbb{C}$, and $n_s \in \mathbb{Z}_{>0}$ for $1 \leq s \leq k$ by

$$x^{n} + \sum_{i=1}^{n} (-1)^{i} a_{i} x^{n-i} = \prod_{s=1}^{k} (x - b_{s})^{n_{s}}.$$
(3.6)

Note that $n = \sum_{s=1}^{k} n_s$.

Lemma 3.9. The $\mathfrak{gl}(1|1)[t]$ -module $\mathcal{V}^{\mathfrak{S}}/I_{\boldsymbol{a}}\mathcal{V}^{\mathfrak{S}}$ is isomorphic to the Weyl module $\bigotimes_{s=1}^{k} W_{n_s}(b_s)$.

Proof. Similar to the proof of Lemma 3.7, the statement follows from Lemma 3.8 by checking relations and comparing dimensions. \Box

Lemma 3.10. We have the following properties for $W_m(0)$.

- (i) The module $W_m(0)$ has a unique grading such that $W_m(0)$ is a graded $\mathfrak{gl}(1|1)[t]$ -module and such that the degree of the highest weight vector w equals 0.
- (ii) As a $\mathfrak{gl}(1|1)$ -module, $W_m(0)$ is isomorphic to $(\mathbb{C}^{1|1})^{\otimes m}$.
- (iii) A $\mathfrak{gl}(1|1)[t]$ -module M is an irreducible subquotient of $W_m(0)$ if and only if M has the form $L_{\lambda}(0)$, where λ is a polynomial weight such that $|\lambda| = m$.

Proof. The first two statements follows from Lemma 3.7 and the construction of $\mathcal{V}^{\mathfrak{S}}/I_a\mathcal{V}^{\mathfrak{S}}$. The last statement follows from the first two.

Lemma 3.11. Let $b \in \mathbb{C}$. We have the following properties for $W_m(b)$.

- (i) As a $\mathfrak{gl}(1|1)$ -module, $W_m(b)$ is isomorphic to $(\mathbb{C}^{1|1})^{\otimes m}$.
- (ii) A $\mathfrak{gl}(1|1)[t]$ -module M is an irreducible subquotient of $W_m(b)$ if and only if M has the form $L_{\lambda}(b)$, where λ is a polynomial weight such that $|\lambda| = m$.

Proof. It follows from Lemma 3.10.

Corollary 3.12. A $\mathfrak{gl}(1|1)[t]$ -module M is an irreducible subquotient of $\bigotimes_{s=1}^{k} W_{n_s}(b_s)$ if and only if M has the form $\bigotimes_{s=1}^{k} L_{\lambda^{(s)}}(b_s)$, where $\lambda^{(s)}$ is a polynomial weight such that $|\lambda^{(s)}| = n_s$ for each $1 \leq s \leq k$.

Proof. It follows from part (ii) of Lemma 3.11 and the irreducibility of $\bigotimes_{s=1}^{k} L_{\lambda^{(s)}}(b_s)$, and the Jordan-Hölder theorem.

4. MAIN THEOREMS

4.1. The algebra \mathcal{O}_l . Let Ω_l be the *n*-dimensional affine space with coordinates $f_1, \ldots, f_l, g_1, \ldots, g_{n-l-1}$ and Σ_n . Introduce two polynomials

$$f(x) = x^{l} + \sum_{i=1}^{l} f_{i} x^{l-i}, \quad g(x) = x^{n-l-1} + \sum_{i=1}^{n-l-1} g_{i} x^{n-l-i-1}.$$
(4.1)

Denote by \mathcal{O}_l the algebra of regular functions on Ω_l , namely

 $\mathcal{O}_l = \mathbb{C}[f_1, \ldots, f_l, g_1, \ldots, g_{n-l-1}, \Sigma_n].$

Define the degree function by

$$\deg f_i = i, \qquad \deg g_j = j, \qquad \deg \Sigma_n = n,$$

for all i = 1, ..., l and j = 1, ..., n - l - 1. The algebra O_l is graded with the graded character given by

$$\operatorname{ch}(\mathcal{O}_l) = \frac{1}{(q)_l(q)_{n-l-1}(1-q^n)}.$$
(4.2)

Let $\Sigma_1, \ldots, \Sigma_{n-1}$ be the elements of \mathcal{O}_l such that

$$nf(x)g(x) = nx^{n-1} + \sum_{i=1}^{n-1} (-1)^i (n-i) \Sigma_i x^{n-1-i}.$$
(4.3)

The homomorphism

$$\pi_l : \mathbb{C}[z_1, \dots, z_n]^{\mathfrak{S}} \to \mathcal{O}_l, \qquad \sigma_i(\boldsymbol{z}) \mapsto \Sigma_i, \qquad i = 1, \dots, n,$$
(4.4)

is injective and induces a $\mathbb{C}[z_1, \ldots, z_n]^{\mathfrak{S}}$ -module structure on \mathcal{O}_l .

Express f'(x)g(x) as follows,

$$nf'(x)g(x) = nlx^{n-2} + \sum_{i=1}^{n-2} G_i x^{n-2-i},$$
(4.5)

where $G_i \in \mathcal{O}_l$.

Lemma 4.1. The elements G_i and Σ_j , i = 1, ..., n-2, j = 1, ..., n, generate the algebra \mathcal{O}_l . \Box

Lemma 4.2. The elements G_i and Σ_j are homogeneous of degrees i and j, respectively, for $i = 1, \ldots, n-2, j = 1, \ldots, n$.

4.2. Bethe algebra. We call the unital subalgebra of $U(\mathfrak{gl}(1|1)[t])$ generated by the coefficients of

$$e_{11}(x) + e_{22}(x), \quad 2\mathcal{H}(x) = e_{11}(x)e_{11}(x) - e_{12}(x)e_{21}(x) + e_{21}(x)e_{12}(x) - e_{22}(x)e$$

the *Bethe algebra*. We denote the Bethe algebra by \mathcal{B} . Note that the coefficients of $e_{11}(x) + e_{22}(x)$ generate the center of $U(\mathfrak{gl}(1|1)[t])$.

Lemma 4.3 ([MR14]). *The Bethe algebra* \mathcal{B} *is commutative. The Bethe algebra* \mathcal{B} *commutes with the subalgebra* $U(\mathfrak{gl}(1|1)) \subset U(\mathfrak{gl}(1|1)[t])$.

Being a subalgebra of $U(\mathfrak{gl}(1|1)[t])$, the Bethe algebra \mathcal{B} acts on any $\mathfrak{gl}(1|1)[t]$ -module M. Since \mathcal{B} commutes with $U(\mathfrak{gl}(1|1))$, the Bethe algebra preserves the subspace $(M)^{\text{sing}}_{\lambda}$ for any weight λ . If $K \subset M$ is a \mathcal{B} -invariant subspace, then we call the image of \mathcal{B} in End(K) the *Bethe algebra associated with* K.

Let $\boldsymbol{a} = (a_1, \ldots, a_n) \in \mathbb{C}^n$. Define $k \in \mathbb{Z}_{>0}$, a sequence of positive integers $\boldsymbol{n} = (n_1, \ldots, n_k)$, and a sequence of distinct complex numbers $\boldsymbol{b} = (b_1, \ldots, b_k)$ by (3.6). Let $\boldsymbol{\Lambda} = (\boldsymbol{\lambda}^{(1)}, \ldots, \boldsymbol{\lambda}^{(k)})$ be a sequence of polynomial $\mathfrak{gl}(1|1)$ -weights such that $|\boldsymbol{\lambda}^{(s)}| = n_s$.

We study the action of the Bethe algebra \mathcal{B} on the following \mathcal{B} -modules:

$$\mathcal{M}_{l} = (\mathcal{V}^{\mathfrak{S}})_{(n-l,l)}^{\operatorname{sing}}, \quad \mathcal{M}_{l,\boldsymbol{a}} = (\bigotimes_{s=1}^{k} W_{n_{s}}(b_{s}))_{(n-l,l)}^{\operatorname{sing}}, \quad \mathcal{M}_{l,\boldsymbol{\Lambda},\boldsymbol{b}} = (\bigotimes_{s=1}^{k} L_{\boldsymbol{\lambda}^{(s)}}(b_{s}))_{(n-l,l)}^{\operatorname{sing}}.$$

Denote the Bethe algebras associated with \mathcal{M}_l , $\mathcal{M}_{l,a}$, $\mathcal{M}_{l,\Lambda,b}$ by \mathcal{B}_l , $\mathcal{B}_{l,a}$, $\mathcal{B}_{l,\Lambda,b}$, respectively. For any element $X \in \mathcal{B}$, we denote by X(z), $X(\alpha)$, $X(\Lambda, b)$ the respective linear operators.

Since by Lemma 3.3 the $\mathfrak{gl}(1|1)[t]$ -module $\mathcal{V}^{\mathfrak{S}}$ is generated by $v_1^{\otimes n} = v_1 \otimes \cdots \otimes v_1$, the series $e_{11}(x) + e_{22}(x)$ acts on $\mathcal{V}^{\mathfrak{S}}$ by multiplication by the series

$$\sum_{i=1}^{n} \frac{1}{x - z_i} = \sum_{i=1}^{n} \sum_{j=0}^{\infty} z_i^j x^{-j-1}.$$

Therefore there exist unique central elements C_1, \ldots, C_n of $U(\mathfrak{gl}(1|1)[t])$ of minimal degrees such that each C_i acts on $\mathcal{V}^{\mathfrak{S}}$ by multiplication by $\sigma_i(\mathbf{z})$.

Define $B_i \in \mathcal{B}$ by

$$\left(x^{n} + \sum_{i=1}^{n} (-1)^{i} C_{i} x^{n-i}\right) \mathcal{T}(x) = x^{n} \sum_{i=2}^{\infty} B_{i} x^{-i},$$
(4.6)

where $\mathcal{T}(x)$ is defined in (2.11).

Lemma 4.4. We have $B_i(z) = 0$ for i > n and $B_2(z) = nl$.

Proof. Let $V(\mathbf{c}) = \bigotimes_{i=1}^{n} \mathbb{C}^{1|1}(c_i)$, where $c_i \in \mathbb{C}$. Note that $B_i(\mathbf{z})$ is a polynomial in \mathbf{z} with values in $\operatorname{End}((V)_{(n-l,l)}^{\operatorname{sing}})$. For any sequence of complex numbers $\mathbf{c} = (c_1, \ldots, c_n)$, we can evaluate $B_i(\mathbf{z})$ at $\mathbf{z} = \mathbf{c}$ to an operator on $(V(\mathbf{c}))_{(n-l,l)}^{\operatorname{sing}}$. By Theorem 2.6, the Gaudin transfer matrix $\mathcal{H}(x)$ is diagonalizable and the Bethe ansatz is complete for $(V(\mathbf{c}))_{(n-l,l)}^{\operatorname{sing}}$ when $\mathbf{c} \in \mathbb{C}^n$ is generic. Hence by (2.12) and (4.6) that $(x^n + \sum_{i=1}^n (-1)^i C_i x^{n-i}) \mathcal{T}(x)$ acts on $(V(\mathbf{c}))_{(n-l,l)}^{\operatorname{sing}}$ as a polynomial in x for generic \mathbf{c} . In particular, it implies that B_i , i > n, acts on $(V(\mathbf{c}))_{(n-l,l)}^{\operatorname{sing}}$ by zero for generic \mathbf{c} . Therefore $B_i(\mathbf{z})$, i > n, is identically zero.

By the same reasoning, one shows that $B_2(z) = nl$. Alternatively, it also follows from $B_2 = (e_{11} + e_{22})e_{22} - e_{21}e_{12}$.

Lemma 4.5. The elements $B_i(z)$ and $C_j(z)$, i = 3, ..., n, j = 1, ..., n, generate the algebra \mathcal{B}_l .

Proof. It follows from the definition of \mathcal{B} , (4.6), and Lemma 4.4.

Lemma 4.6. The elements B_i and C_j are homogeneous of degrees i - 2 and j, respectively, for $i = 3, 4, \ldots$ and $j = 1, \ldots, n$.

4.3. **Main theorems.** Recall from Proposition 3.6 that there exists a unique vector (up to proportionality) of degree l(l+1)/2 in \mathcal{M}_l explicitly given by

$$\mathfrak{u}_l := e_{12}[0]e_{21}[0]e_{21}[1]\cdots e_{21}[l]v^+,$$

see Lemma 3.5.

Any commutative algebra \mathcal{A} is a module over itself induced by left multiplication. We call it the *regular representation of* \mathcal{A} . The dual space \mathcal{A}^* is naturally an \mathcal{A} -module which is called the *coregular representation*. A bilinear form $(\cdot|\cdot) : \mathcal{A} \otimes \mathcal{A} \to \mathbb{C}$ is called *invariant* if (ab|c) =(a|bc) for all $a, b, c \in \mathcal{A}$. A finite-dimensional commutative algebra \mathcal{A} admitting an invariant nondegenerate symmetric bilinear form $(\cdot|\cdot) : \mathcal{A} \otimes \mathcal{A} \to \mathbb{C}$ is called a *Frobenius algebra*. The regular and coregular representations of a Frobenius algebra are isomorphic.

Let M be an \mathcal{A} -module and $\mathcal{E} : \mathcal{A} \to \mathbb{C}$ a character, then the \mathcal{A} -eigenspace associated to \mathcal{E} in M is defined by $\bigcap_{a \in \mathcal{A}} \ker(a|_M - \mathcal{E}(a))$. The generalized \mathcal{A} -eigenspace associated to \mathcal{E} in M is defined by $\bigcap_{a \in \mathcal{A}} \left(\bigcup_{m=1}^{\infty} \ker(a|_M - \mathcal{E}(a))^m \right)$.

Theorem 4.7. The action of the Bethe algebra \mathcal{B}_l on \mathcal{M}_l has the following properties.

- (i) The map $\eta_l : G_i \mapsto B_{i+2}(z), \Sigma_j \mapsto C_j(z), i = 1, ..., n-2, j = 1, ..., n$, extends uniquely to an isomorphism $\eta_l : \mathcal{O}_l \to \mathcal{B}_l$ of graded algebras. Moreover, the isomorphism η_l is an isomorphism of $\mathbb{C}[z_1, ..., z_n]^{\mathfrak{S}}$ -modules.
- (ii) The map $\rho_l : \mathcal{O}_l \mapsto \mathcal{M}_l, F \mapsto \eta_l(F)\mathfrak{u}_l$, is an isomorphism of graded vector spaces identifying the \mathcal{B}_l -module \mathcal{M}_l with the regular representation of \mathcal{O}_l .

Theorem 4.7 is proved in Section 5.

Let $\boldsymbol{a} = (a_1, \ldots, a_n) \in \mathbb{C}^n$. Define $k \in \mathbb{Z}_{>0}$, a sequence of positive integers $\boldsymbol{n} = (n_1, \ldots, n_k)$, and a sequence of distinct complex numbers $\boldsymbol{b} = (b_1, \ldots, b_k)$ by (3.6). Let $\boldsymbol{\Lambda} = (\boldsymbol{\lambda}^{(1)}, \ldots, \boldsymbol{\lambda}^{(k)})$ be a sequence of non-degenerate polynomial weights such that $|\boldsymbol{\lambda}^{(s)}| = n_s$ for each $1 \leq s \leq k$.

Theorem 4.8. The action of the Bethe algebra $\mathcal{B}_{l,\Lambda,b}$ on $\mathcal{M}_{l,\Lambda,b}$ has the following properties.

(i) The Bethe algebra $\mathcal{B}_{l,\Lambda,b}$ is isomorphic to

$$\mathbb{C}[w_1,\ldots,w_{k-1}]^{\mathfrak{S}_l\times\mathfrak{S}_{k-l-1}}/\langle\sigma_i(\boldsymbol{w})-\varepsilon_i\rangle_{i=1,\ldots,k-1},$$

where ε_i are given by

$$\varphi_{\mathbf{\Lambda}, \mathbf{b}}(x) := \prod_{s=1}^{k} (x - b_s) \sum_{s=1}^{k} \frac{n_s}{x - b_s} = n \left(x^{k-1} + \sum_{i=1}^{k-1} (-1)^i \varepsilon_i x^{k-1-i} \right)$$

and $\sigma_i(\boldsymbol{w})$ are elementary symmetric functions in w_1, \ldots, w_{k-1} .

- (ii) The Bethe algebra $\mathcal{B}_{l,\Lambda,b}$ is a Frobenius algebra. Moreover, the $\mathcal{B}_{l,\Lambda,b}$ -module $\mathcal{M}_{l,\Lambda,b}$ is isomorphic to the regular representation of $\mathcal{B}_{l,\Lambda,b}$.
- (iii) The Bethe algebra $\mathcal{B}_{l,\Lambda,b}$ is a maximal commutative subalgebra in $\mathcal{M}_{l,\Lambda,b}$ of dimension $\binom{k-1}{l}$.
- (iv) Every \mathcal{B} -eigenspace in $\mathcal{M}_{l,\Lambda,b}$ has dimension one.
- (v) The \mathcal{B} -eigenspaces in $\mathcal{M}_{l,\Lambda,b}$ bijectively correspond to the monic degree l divisors y(x) of the polynomial $\varphi_{\Lambda,b}(x)$. Moreover, the eigenvalue of $\mathcal{H}(x)$ corresponding to the monic divisor y is described by $\mathcal{E}_{y,\Lambda,b}(x)$, see (2.10).
- (vi) Every generalized \mathcal{B} -eigenspace in $\mathcal{M}_{l,\Lambda,b}$ is a cyclic \mathcal{B} -module.
- (vii) The dimension of the generalized \mathcal{B} -eigenspace associated to $\mathcal{E}_{y,\Lambda,b}(x)$ is

$$\prod_{a\in\mathbb{C}} \binom{\operatorname{Mult}_a(\varphi_{\mathbf{\Lambda},\boldsymbol{b}})}{\operatorname{Mult}_a(y)},$$

1--- (

where $Mult_a(p)$ is the multiplicity of a as a root of the polynomial p.

Theorem 4.8 is proved in Section 5.

Note that the results of is quite parallel to that of XXX spin chains, see [LM21a, Theorem 4.11].

4.4. Higher Gaudin transfer matrices. To define higher Gaudin transfer matrices, we first recall basics about pseudo-differential operators. Let \mathcal{A} be a differential superalgebra with an even derivation $\partial : \mathcal{A} \to \mathcal{A}$. For $r \in \mathbb{Z}_{>0}$, denote the *r*-th derivative of $a \in \mathcal{A}$ by $a_{[r]}$. Define the *superalgebra* of pseudo-differential operators $\mathcal{A}((\partial^{-1}))$ as follows. Elements of $\mathcal{A}((\partial^{-1}))$ are Laurent series in ∂^{-1} with coefficients in \mathcal{A} , and the product is given by

$$\partial \partial^{-1} = \partial^{-1} \partial = 1, \quad \partial^r a = \sum_{s=0}^{\infty} \binom{r}{s} a_{[s]} \partial^{r-s}, \quad r \in \mathbb{Z}, \quad a \in \mathcal{A},$$

where

$$\binom{r}{s} = \frac{r(r-1)\cdots(r-s+1)}{s!}.$$

Let

$$\mathcal{A}_{x}^{m|n} = \mathcal{U}(\mathfrak{gl}(1|1)[t])((x^{-1})) = \left\{ \sum_{r=-\infty}^{s} g_{r}x^{r}, \ r \in \mathbb{Z}, \ g_{r} \in \mathcal{U}(\mathfrak{gl}(1|1)[t]) \right\}$$

Consider the operator in $\operatorname{End}(\mathbb{C}^{1|1}) \otimes \mathcal{A}_x^{m|n}((\partial_x^{-1}))$,

$$\mathfrak{Z}(x,\partial_x) := \sum_{a,b=1}^2 E_{ab} \otimes \left(\delta_{ab}\partial_x - e_{ab}(x)(-1)^{|a|}\right).$$

which is a Manin matrix, see [MR14, Lemma 3.1] and [HM20, Lemma 4.2]. Define the *Berezinian*, see [Naz91], of $\mathfrak{Z}(x, \partial_x)$ by

$$\operatorname{Ber}(\mathfrak{Z}(x,\partial_x)) = (\partial_x - e_{11}(x)) \Big(\partial_x + e_{22}(x) + e_{21}(x) \big(\partial_x - e_{11}(x)\big)^{-1} e_{12}(x)\Big)^{-1}.$$
 (4.7)

Denote the Berezinian by $\mathfrak{D}(x,\partial_x)$ and expand it as an element in $\mathcal{A}_x^{m|n}((\partial_x^{-1}))$,

$$\mathfrak{D}(x,\partial_x) = \sum_{r=0}^{\infty} (-1)^r \mathcal{G}_r(x) \partial_x^{-r}.$$
(4.8)

We call the series $\mathcal{G}_r(x) \in \mathcal{A}_x^{m|n}$, $r \in \mathbb{Z}_{\geq 0}$, the higher Gaudin transfer matrices. In particular, we call $\mathcal{G}_1(x)$ and $\mathcal{G}_2(x)$ the first and second Gaudin transfer matrices, respectively.

Example 4.9. We have $\mathcal{G}_0(x) = 1$,

$$\mathcal{G}_1(x) = e_{11}(x) + e_{22}(x), \quad \mathcal{G}_2(x) = (e_{11}(x) + e_{22}(x))e_{22}(x) - e_{21}(x)e_{12}(x).$$

Moreover, we have

$$\mathcal{H}(x) = \frac{1}{2} \left(\mathcal{G}_1(x) \right)^2 - \mathcal{G}_2(x) + \frac{1}{2} \partial_x \mathcal{G}_1(x), \quad \mathcal{T}(x) = \mathcal{G}_2(x),$$

see (2.11).

Remark 4.10. In principal, the Bethe algebra should be the unital subalgebra of $U(\mathfrak{gl}(1|1)[t])$ generated by coefficients $\mathcal{G}_r(x)$, $r \in \mathbb{Z}_{>0}$, cf. [MM15]. However, it turns out that the first two transfer matrices already give (almost) complete information about the Bethe algebra, see the discussion below.

Now we describe the eigenvalues of higher Gaudin transfer matrices acting on the on-shell Bethe vector.

Let $\Lambda = (\lambda^{(1)}, \ldots, \lambda^{(k)})$ be a sequence of $\mathfrak{gl}(1|1)$ -weights and $\boldsymbol{b} = (b_1, \ldots, b_k)$ a sequence of distinct complex numbers, where $\lambda^{(s)} = (\alpha_s, \beta_s)$. Let $\boldsymbol{t} = (t_1, \ldots, t_l)$, where $0 \leq l < k$. Suppose that y_t divides the polynomial $\varphi_{\Lambda, \boldsymbol{b}}$ (namely \boldsymbol{t} satisfies the Bethe ansatz equation), see (2.8).

Theorem 4.11. If $t_i \neq t_j$ for $1 \leq i < j \leq l$, then

$$\mathfrak{D}(x,\partial_x)\mathbb{B}_l(t) = \mathbb{B}_l(t)\Big(\partial_x - \sum_{s=1}^k \frac{\alpha_s}{x - b_s} + \frac{y_t'}{y_t}\Big)\Big(\partial_x + \sum_{s=1}^k \frac{\beta_s}{x - b_s} + \frac{y_t'}{y_t}\Big)^{-1}.$$
(4.9)

This is a differential analog of [LM21a, Theorem 6.4]. Note that the pseudo-differential operator in the right hand side of (4.9), denoted by $\mathfrak{D}_{y,\Lambda,b}$, was introduced [HMVY19, Section 5.3]. This theorem can be generalized to the $\mathfrak{gl}(m|n)$ case where on the right hand side the pseudodifferential operator describing the eigenvalues of higher Gaudin transfer matrices should be replaced by the pseudo-differential operator in [HMVY19, Equation (6.5)]. This generalization is a classical limit of [LM21b, Conjecture 5.15] which connects the rational difference operator introduced in [HLM19, Equation (5.6)] with the eigenvalues of higher transfer matrices on the on-shell Bethe vector for XXX spin chains associated with $\mathfrak{gl}(m|n)$.

The proof of Theorem 4.11 can be obtained from [LM21a, Theorem 6.4] by taking classical limit, see [MTV06] and cf. [HLM19, Remark 5.4]. We shall provide a proof of [LM21b, Conjecture 5.15] and the generalization of of Theorem 4.11 to the $\mathfrak{gl}(m|n)$ case in a further publication using nested Bethe ansatz, see [KR83, MTV06, BR08]. Hence the proof of Theorem 4.11 is omitted here.

Remark 4.12. As shown in [HMVY19, Lemma 5.7], the odd reflections of $\mathfrak{D}_{y,\Lambda,b}$ cf. [HMVY19, equation (3.1)], which come from the study of the fermionic reproduction procedure of the Bethe ansatz equation, are compatible with the odd reflections in Lie superalgebras. The difference analog of this fact has been used in [Lu21] to investigate the odd reflections of super Yangian of type A and the fermionic reproduction procedure of the Bethe ansatz equation for XXX spin chains.

We conclude this section by discussing the connections between $\mathcal{G}_i(x)$, $i \ge 3$, and $\mathcal{G}_1(x)$, $\mathcal{G}_2(x)$. Let

$$\mu(x) = \sum_{s=1}^{k} \frac{\alpha_s}{x - b_s} - \frac{y'_t}{y_t}, \quad \nu(x) = \sum_{s=1}^{k} \frac{\beta_s}{x - b_s} + \frac{y'_t}{y_t}.$$

For simplicity, we do not write the dependence of $\mu(x)$ and $\nu(x)$ on Λ, b, t explicitly. Then the eigenvalue of $\mathfrak{D}(x, \partial_x)$ acting on $\mathbb{B}_l(t)$ is given by

$$(\partial_x - \mu(x))(\partial_x + \nu(x))^{-1} = 1 - (\mu(x) + \nu(x))(\partial_x + \nu(x))^{-1}.$$
(4.10)

Hence the eigenvalues of $\mathcal{G}_i(x)$ are essentially determined only by $\mu(x) + \nu(x)$ and $\nu(x)$. Comparing (4.8) and the expansion of (4.10), we have

$$\mathcal{G}_1(x)\mathbb{B}_l(\boldsymbol{t}) = (\mu(x) + \nu(x))\mathbb{B}_l(\boldsymbol{t}), \quad \mathcal{G}_2(x)\mathbb{B}_l(\boldsymbol{t}) = (\mu(x) + \nu(x))\nu(x)\mathbb{B}_l(\boldsymbol{t}), \quad (4.11)$$

see also (2.12). Therefore, the spectrum of all higher transfer matrices are determined simply by that of the first two transfer matrices which justifies our definition of Bethe algebra.

Lemma 4.13. Let the complex parameters c_1, \ldots, c_m and the positive integer m vary. Then the kernels of the representations $\bigotimes_{i=1}^m \mathbb{C}^{1|1}(c_i)$ of $U(\mathfrak{gl}(1|1)[t])$ have the zero intersection.

Proof. The proof is contained in the proof of [Naz20, Proposition 1.7].

Corollary 4.14. We have

$$\mathfrak{D}(x,\partial_x) = \left(\partial_x - \mathcal{G}_1(x) + \frac{\mathcal{G}_2(x)}{\mathcal{G}_1(x)}\right) \left(\partial_x + \frac{\mathcal{G}_2(x)}{\mathcal{G}_1(x)}\right)^{-1}.$$
(4.12)

Proof. By Lemma 4.13, it suffices to check that the left hand side and the right hand side of (4.12) act identically on a basis of $\bigotimes_{i=1}^{m} \mathbb{C}^{1|1}(c_i)$ for all $m \in \mathbb{Z}_{>0}$ and generic $c = (c_1, \ldots, c_m)$.

By Theorem 2.6, there is a basis of $\bigotimes_{i=1}^{m} \mathbb{C}^{1|1}(c_i)$ consisting of on-shell Bethe vectors for generic c. Therefore, the statement follows from Theorem 4.11, (4.10), (4.11).

5. PROOF OF MAIN THEOREMS

In this section, we prove the main theorems.

5.1. The first isomorphism.

Proof of Theorem 4.7. We first show the homomorphism defined by η_l is well-defined.

Consider the tensor product $V(c) = \bigotimes_{i=1}^{n} \mathbb{C}^{1|1}(c_i)$, where $c_i \in \mathbb{C}$, and the corresponding Bethe ansatz equation associated to weight (n - l, l). Let t be a solution with distinct coordinates and $\mathbb{B}_l(t)$ the corresponding on-shell Bethe vector. Denote $\mathcal{E}_{i,t}$ the eigenvalues of B_i acting on $\mathbb{B}_l(t)$, see Theorem 2.2 and (2.12).

Define a character $\pi : \mathcal{O}_l \to \mathbb{C}$ by sending

$$f(x) \mapsto y_t(x), \qquad g(x) \mapsto \frac{1}{ny_t(x)} \prod_{i=1}^n (x - c_i) \sum_{i=1}^n \frac{1}{x - c_i}, \qquad \Sigma_n \mapsto \prod_{i=1}^n c_i.$$

Then

$$\pi(\Sigma_i) = \sigma_i(\boldsymbol{c}), \qquad \pi(G_i) = \mathcal{E}_{i,t}, \tag{5.1}$$

by (4.3) and by (2.10), (2.12), (4.5), respectively.

Let now $P(G_i, \Sigma_j)$ be a polynomial in G_i, Σ_j such that $P(G_i, \Sigma_j)$ is equal to zero in \mathcal{O}_l . It suffices to show $P(B_i(\boldsymbol{z}), C_j(\boldsymbol{z}))$ is equal to zero in \mathcal{B}_l .

Note that $P(B_i(\boldsymbol{z}), C_j(\boldsymbol{z}))$ is a polynomial in z_1, \ldots, z_n with values in $End((V)_{(n-l,l)}^{sing})$. For any sequence \boldsymbol{c} of complex numbers, we can evaluate $P(B_i(\boldsymbol{z}), C_j(\boldsymbol{z}))$ at $\boldsymbol{z} = \boldsymbol{c}$ to an operator on $(V(\boldsymbol{c}))_{(n-l,l)}^{sing}$. By Theorem 2.6, the transfer matrix $\mathcal{T}(x)$ is diagonalizable and the Bethe ansatz is complete for $(V(\boldsymbol{c}))_{(n-l,l)}^{sing}$ when $\boldsymbol{c} \in \mathbb{C}^n$ is generic. Hence by (5.1) the value of $P(B_i(\boldsymbol{z}), C_j(\boldsymbol{z}))$ at $\boldsymbol{z} = \boldsymbol{c}$ is also equal to zero for generic \boldsymbol{c} . Therefore $P(B_i(\boldsymbol{z}), C_j(\boldsymbol{z}))$ is identically zero and the map η_l is well-defined.

Let us now show that the map η_l is injective. Let $P(G_i, \Sigma_j)$ be a polynomial in G_i, Σ_j such that $P(G_i, \Sigma_j)$ is non-zero in \mathcal{O}_l . Then the value at a generic point of Ω_l (e.g. the non-vanishing points of $P(G_i, \Sigma_j)$ such that f and g are relatively prime and have only simple zeros) is not equal to zero. Moreover, at those points the transfer matrix $\mathcal{T}(x)$ is diagonalizable and the Bethe ansatz is

complete again by Theorem 2.6. Therefore, again by (5.1), the polynomial $P(B_i(z), C_j(z))$ is a non-zero element in \mathcal{B}_l . Thus the map η_l is injective.

The surjectivity of η_l follows from Lemma 4.5. Hence η_l is an isomorphism of algebras.

The fact that η_l is an isomorphism of graded algebra respecting the gradation follows from Lemmas 4.2 and 4.6. This completes the proof of part (i).

The kernel of ρ_l is an ideal of \mathcal{O}_l . If we identify $\sigma_i(z)$ with Σ_i , then the algebra \mathcal{O}_l contains the algebra $\mathbb{C}[z_1, \ldots, z_n]^{\mathfrak{S}}$, see (4.4). The kernel of ρ_l intersects $\mathbb{C}[z_1, \ldots, z_n]^{\mathfrak{S}}$ trivially. Therefore the kernel of ρ_l is trivial as well. Hence ρ_l is an injective map. Comparing (4.2) and Proposition 3.6, we have $\operatorname{ch}(\mathcal{M}_l) = q^{l(l+1)/2} \operatorname{ch}(\mathcal{O}_l)$. Thus ρ_l is an isomorphism of graded vector spaces which shifts the degree by l(l+1)/2, completing the proof of part (ii).

5.2. The second isomorphism. Let $a = (a_1, \ldots, a_n)$ be a sequence of complex numbers. Define $k \in \mathbb{Z}_{>0}$, a sequence of positive integers $n = (n_1, \ldots, n_k)$, and a sequence of distinct complex numbers $b = (b_1, \ldots, b_k)$ by (3.6). Let $I_{l,a}^{\mathcal{O}}$ be the ideal of \mathcal{O}_l generated by the elements $\Sigma_i - a_i$, $i = 1, \ldots, n$, where $\Sigma_1, \ldots, \Sigma_{n-1}$ are defined in (4.3). Let $\mathcal{O}_{l,a}$ be the quotient algebra

$$\mathcal{O}_{l,oldsymbol{a}} = \mathcal{O}_l / I_{l,oldsymbol{a}}^\mathcal{O}$$

Let $I_{l,a}^{\mathcal{B}}$ be the ideal of \mathcal{B}_l generated by $C_i(z) - a_i, i = 1, \ldots, n$. Consider the subspace

$$I_{l,\boldsymbol{a}}^{\mathcal{M}} = I_{l,\boldsymbol{a}}^{\mathcal{B}} \mathcal{M}_{l} = (I_{\boldsymbol{a}} \mathcal{V}^{\mathfrak{S}})_{(n-l,l)}^{\operatorname{sing}},$$

where I_a as before is the ideal of $\mathbb{C}[z_1, \ldots, z_n]^{\mathfrak{S}}$ generated by $\sigma_i(z) - a_i, i = 1, \ldots, n$.

Lemma 5.1. We have

$$\eta_l(I_{l,\boldsymbol{a}}^{\mathcal{O}}) = I_{l,\boldsymbol{a}}^{\mathcal{B}}, \quad \rho_l(I_{l,\boldsymbol{a}}^{\mathcal{O}}) = I_{l,\boldsymbol{a}}^{\mathcal{M}}, \quad \mathcal{B}_{l,\boldsymbol{a}} = \mathcal{B}_l/I_{l,\boldsymbol{a}}^{\mathcal{B}}, \quad \mathcal{M}_{l,\boldsymbol{a}} = (\mathcal{V}^{\mathfrak{S}})_{(n-l,l)}^{\mathrm{sing}}/I_{l,\boldsymbol{a}}^{\mathcal{M}}.$$

Proof. The lemma follows from Theorem 4.7 and Lemma 3.9.

By Lemma 5.1, the maps η_l and ρ_l induce the maps

$$\eta_{l,\boldsymbol{a}}:\mathcal{O}_{l,\boldsymbol{a}}\to\mathcal{B}_{l,\boldsymbol{a}},\qquad \rho_{l,\boldsymbol{a}}:\mathcal{O}_{l,\boldsymbol{a}}\to\mathcal{M}_{l,\boldsymbol{a}}.$$

The map $\eta_{l,a}$ is an isomorphism of algebras. Since $\mathcal{B}_{l,a}$ is finite-dimensional, by e.g. [MTV09, Lemma 3.9], $\mathcal{O}_{l,a}$ is a Frobenius algebra, so is $\mathcal{B}_{l,a}$. The map $\rho_{l,a}$ is an isomorphism of vector spaces. Moverover, it follows from Theorem 4.7 and Lemma 5.1 that $\rho_{l,a}$ identifies the regular representation of $\mathcal{O}_{l,a}$ with the $\mathcal{B}_{l,a}$ -module $\mathcal{M}_{l,a}$.

The statement of this section implies, by e.g. [Lu20, Lemma 1.3], the following. Set

$$\zeta_{n,b}(x) = \sum_{s=1}^{k} \frac{n_s}{x - b_s}, \qquad \psi_{n,b}(x) := \zeta_{n,b}(x) \prod_{r=1}^{k} (x - b_r)^{n_s}.$$

Theorem 5.2. Suppose $\mathbf{b} = (b_1, \ldots, b_k)$ is a sequence of distinct complex numbers. Then the Gaudin transfer matrix $\mathfrak{H}(x)$ has a simple spectrum in $(\bigotimes_{s=1}^k W_{n_s}(b_s))^{\text{sing}}$. There exists a bijective correspondence between the monic divisors y of the polynomial $\psi_{n,\mathbf{b}}$ and the eigenvectors v_y of the

Gaudin transfer matrix $\mathcal{H}(x)$ (up to multiplication by a non-zero constant). Moreover, this bijection is such that

$$\mathcal{H}(x)v_y = \left(\frac{1}{2}\zeta'_{\boldsymbol{n},\boldsymbol{b}}(x) - \zeta_{\boldsymbol{n},\boldsymbol{b}}(x)\frac{y'(x)}{y(x)} + \frac{1}{2}\left(\zeta_{\boldsymbol{n},\boldsymbol{b}}(x)\right)^2\right)v_y.$$

Remark 5.3. Fix $l \in \mathbb{Z}_{\geq 0}$ and set $\boldsymbol{t} = (t_1, \ldots, t_l)$. Let \boldsymbol{y}_t represent \boldsymbol{t} . Then the Bethe ansatz equation for $(\bigotimes_{s=1}^k W_{n_s}(b_s))^{\text{sing}}$ is

 $y_t(x)$ divides the polynomial $\psi_{n,b}(x)$.

Note that in this case, y_t may have multiple roots. If there are multiple roots in y_t , then the corresponding on-shell Bethe vector is zero. Therefore an actual eigenvector should be obtained via an appropriate derivative as pointed out in [HMVY19, Section 8.2].

5.3. The third isomorphism. Recall from Section 2.3, that without loss of generality, we can assume that $\beta_s = 0, 1 \le s \le k$. In this case, $\alpha_s = n_s, 1 \le s \le k$.

Lemma 5.4. There exists a surjective $\mathfrak{gl}(1|1)[t]$ -module homomorphism from $\bigotimes_{s=1}^{k} W_{n_s}(b_k)$ to $\bigotimes_{s=1}^{k} L_{\lambda^{(s)}}(b_k)$ which maps vacuum vector to vacuum vector.

Proof. It follows from Lemma 3.11 and our assumption that $\beta_s = 0$ for all $1 \leq s \leq k$.

By Lemma 3.9, the surjective $\mathfrak{gl}(1|1)[t]$ -module homomorphism

$$\bigotimes_{s=1}^{k} W_{n_s}(b_k) \twoheadrightarrow \bigotimes_{s=1}^{k} L_{\boldsymbol{\lambda}^{(s)}}(b_k)$$

induces a surjective $\mathfrak{gl}(1|1)[t]$ -module homomorphism

$$\mathcal{V}^{\mathfrak{S}} \twoheadrightarrow \bigotimes_{s=1}^{k} L_{\boldsymbol{\lambda}^{(s)}}(b_k).$$

The second map then induces a projection of the Bethe algebras $\mathcal{B}_l \twoheadrightarrow \mathcal{B}_{l,\Lambda,b}$. We describe the kernel of this projection. We consider the corresponding ideal in the algebra \mathcal{O}_l .

Suppose $l \leq k - 1$. Define the polynomial h(x) by

$$h(x) = \prod_{s=1}^{k} (x - b_s)^{n_s - 1}.$$

Divide the polynomial g(x) in (4.1) by h(x) and let

$$p(x) = x^{k-l-1} + p_1 x^{k-l-2} + \dots + p_{k-l-2} x + p_{k-l-1},$$
(5.2)

$$r(x) = r_1 x^{n-k-1} + r_2 x^{n-k-2} + \dots + r_{n-k-1} x + r_{n-k}$$
(5.3)

be the quotient and the remainder, respectively. Clearly, $p_i, r_j \in \mathcal{O}_l$.

Denote by $I_{l,\Lambda,b}^{\mathcal{O}}$ the ideal of \mathcal{O}_l generated by $r_1, \ldots, r_{n-k}, \Sigma_n - a_n$, and the coefficients of polynomial

$$\varphi_{\mathbf{\Lambda},\mathbf{b}}(x) - np(x)f(x) = \prod_{s=1}^{k} (x - b_s) \sum_{s=1}^{k} \frac{n_s}{x - b_s} - np(x)f(x).$$

Let $\mathcal{O}_{l,\Lambda,b}$ be the quotient algebra

$$\mathcal{O}_{l, \boldsymbol{\Lambda}, \boldsymbol{b}} = \mathcal{O}_l / I_{l, \boldsymbol{\Lambda}, \boldsymbol{b}}^{\mathcal{O}}$$

Clearly, if $\mathcal{O}_{l,\Lambda,b}$ is finite-dimensional, then it is a Frobenius algebra.

Let $I_{l,\Lambda,b}^{\mathcal{B}}$ be the image of $I_{l,\Lambda,b}^{\mathcal{O}}$ under the isomorphism η_l .

Lemma 5.5. The ideal $I_{l,\Lambda,b}^{\mathcal{B}}$ is contained in the kernel of the projection $\mathcal{B}_{l} \twoheadrightarrow \mathcal{B}_{l,\Lambda,b}$.

Proof. We treat $\boldsymbol{b} = (b_1, \ldots, b_k)$ as variables. Note that the elements of $I_{l,\Lambda,\boldsymbol{b}}^{\mathcal{B}}$ act on $\mathcal{M}_{l,\Lambda,\boldsymbol{b}}$ as polynomials in \boldsymbol{b} with values in $\operatorname{End}((L_{\Lambda})_{(n-l,l)}^{\operatorname{sing}})$. Therefore it suffices to show it for generic \boldsymbol{b} . Let $\mathfrak{f}(x)$ be the image of f(x) under η_l . The condition that $I_{l,\Lambda,\boldsymbol{b}}^{\mathcal{B}}$ vanishes is equivalent to the condition that $\varphi_{\Lambda,\boldsymbol{b}}(x)$ is divisible by $\mathfrak{f}(x)$.

By Theorem 2.6, there exists an eigenbasis of the operator $\mathcal{T}(x)$ in $\mathcal{M}_{l,\Lambda,b}$ for generic *b*. Clearly, a solution of Bethe ansatz equation associated to Λ, b, l is also a solution to Bethe ansatz equation for $\mathcal{M}_{l,a}$, see Theorem 5.2 and Remark 5.3. Moreover, the expressions of corresponding on-shell Bethe vectors coincide (with different vacuum vectors). By Lemma 5.4 and Theorems 2.2, 5.2, $\varphi_{\Lambda,b}(x)$ is divisible by $\mathfrak{f}(x)$ for generic *b* since the eigenvalue of $\mathfrak{f}(x)$ corresponds to $y_t(x)$ in (2.10). Therefore $I_{l,\Lambda,b}^{\mathcal{B}}$ vanishes for generic *b*, thus completing the proof.

Therefore, we have the epimorphism

$$\mathcal{O}_{l,\Lambda,b} \cong \mathcal{B}_l / I_{l,\Lambda,b}^{\mathcal{B}} \twoheadrightarrow \mathcal{B}_{l,\Lambda,b}.$$
(5.4)

We claim that the surjection in (5.4) is an isomorphism by checking dim $\mathcal{O}_{l,\Lambda,b} = \dim \mathcal{B}_{l,\Lambda,b}$.

Lemma 5.6. We have dim $\mathcal{O}_{l,\Lambda,b} = \binom{k-1}{l}$.

Proof. Note that $\mathbb{C}[p_1, \ldots, p_{k-l-1}, r_1, \ldots, r_{n-k}] \cong \mathbb{C}[g_1, \ldots, g_{n-l-1}]$, where p_i and r_j are defined in (5.2) and (5.3). It is not hard to check that

$$\mathcal{O}_{l,\mathbf{\Lambda},\mathbf{b}} \cong \mathbb{C}[f_1,\ldots,f_l,p_1,\ldots,p_{k-l-1}]/\tilde{I}^{\mathcal{O}}_{l,\mathbf{\Lambda},\mathbf{b}},\tag{5.5}$$

where $\tilde{I}_{l,\Lambda,b}^{\mathcal{O}}$ is the ideal of $\mathbb{C}[f_1, \ldots, f_l, p_1, \ldots, p_{k-l-1}]$ generated by the coefficients of the polynomial $\varphi_{\Lambda,b}(x) - np(x)f(x)$.

Introduce new variables $\boldsymbol{w} = (w_1, \ldots, w_{k-1})$ such that

$$f(x) = \prod_{i=1}^{l} (x - w_i), \quad p(x) = \prod_{i=1}^{k-l-1} (x - w_{l+i})$$

Let $\boldsymbol{\varepsilon} = (\varepsilon_1, \dots, \varepsilon_{k-1})$ be complex numbers such that

$$\varphi_{\mathbf{\Lambda},\mathbf{b}}(x) = \prod_{s=1}^{k} (x-b_s) \sum_{s=1}^{k} \frac{n_s}{x-b_s} = n \Big(x^{k-1} + \sum_{i=1}^{k-1} (-1)^i \varepsilon_i x^{k-1-i} \Big).$$

Then

$$\mathbb{C}[f_1,\ldots,f_l,p_1,\ldots,p_{k-l-1}]/\tilde{I}^{\mathcal{O}}_{l,\mathbf{\Lambda},\mathbf{b}} \cong \mathbb{C}[w_1,\ldots,w_{k-1}]^{\mathfrak{S}_l \times \mathfrak{S}_{k-l-1}}/\langle \sigma_i(\boldsymbol{w}) - \varepsilon_i \rangle_{i=1,\ldots,k-1}.$$
(5.6)

The lemma now follows from the fact that $\mathbb{C}[w_1, \ldots, w_{k-1}]^{\mathfrak{S}_l \times \mathfrak{S}_{k-l-1}}$ is a free $\mathbb{C}[w_1, \ldots, w_{k-1}]^{\mathfrak{S}_-}$ module of rank $\binom{k-1}{l}$.

$$\dim \mathcal{B}_{l,\mathbf{\Lambda},\boldsymbol{b}} = \dim \mathcal{M}_{l,\mathbf{\Lambda},\boldsymbol{b}} = \binom{k-1}{l}.$$
(5.7)

Proof of Theorem 4.8. Part (i) follows from Lemma 5.6 and (5.4), (5.5), (5.6), (5.7). Clearly, we have $\mathcal{B}_{l,\Lambda,b} \cong \mathcal{O}_{l,\Lambda,b}$ is a Frobenius algebra. Moreover, the map ρ_l from Theorem 4.7 induces a map

$$\rho_{l,\Lambda,b}: \mathcal{O}_{l,\Lambda,b} \to \mathcal{M}_{l,\Lambda,b}$$

which identifies the regular representation of $\mathcal{O}_{l,\Lambda,b}$ with the $\mathcal{B}_{l,\Lambda,b}$ -module $\mathcal{M}_{l,\Lambda,b}$. Therefore part (ii) is proved.

Since $\mathcal{B}_{l,\Lambda,b}$ is a Frobenius algebra, the regular and coregular representations of the algebra $\mathcal{B}_{l,\Lambda,b}$ are isomorphic to each other. Parts (iii)–(vi) follow from the general facts about the coregular representations, see e.g. [MTV09, Section 3.3] or [Lu20, Lemma 1.3].

Due to part (iv), it suffices to consider the algebraic multiplicity of every eigenvalue. It is well known that roots of a polynomial depend continuously on its coefficients. Hence the eigenvalues of $\mathcal{T}(x)$ depend continuously on **b**. Part (vii) follows from the deformation argument and Theorem 2.6.

REFERENCES

- [BR08] S. Belliard, E. Ragoucy, *The nested Bethe ansatz for 'all' closed spin chains*, J. Phys A: Math. and Theor. 41 (2008), 295202.
- [CLS19] L. Calixto, J. Lemay, A. Savage, Weyl modules for Lie superalgebras Proc. Amer. Math. Soc. 147 (2019), no. 8, 3191–3207.
- [CLV20] D. Chernyak, S. Leurent, D. Volin, Completeness of Wronskian Bethe equations for rational $\mathfrak{gl}(m|n)$ spin chain, arXiv:2004.02865.
- [CP01] V. Chari, A. Pressley, *Weyl Modules for Classical and Quantum Affine algebras*, Represent. Theory **5** (2001), 191–223.
- [HLM19] C.-L. Huang, K. Lu, E. Mukhin, Solutions of $\mathfrak{gl}(m|n)$ XXX Bethe ansatz equation and rational difference operators, J. Phys. A **52** (2019), 375204, 31pp.
- [HM20] C.-L. Huang, E. Mukhin, The duality of $\mathfrak{gl}(m|n)$ and \mathfrak{gl}_k Gaudin models, J. Algebra 548 (2020), 1–24.
- [HMVY19] C.-L. Huang, E. Mukhin, B. Vicedo, C. Young, *The solutions of* $\mathfrak{gl}(m|n)$ *Bethe ansatz equation and rational pseudodifferential operators*, Sel. Math. New Ser. (2019) **25**:52.
- [Kul85] P. Kulish, Integrable graded magnets, Zap. Nauchn. Sem. LOMI 145 (1985), 140–163.
- [KM01] P. Kulish, N. Manojlovic, *Bethe vectors of the* $\mathfrak{osp}(1|2)$ *Gaudin model*, Lett. Math. Phys. **55** (2001), no. 1, 77–95.
- [KR83] P. Kulish, N.Yu.Reshetikhin, *Diagonalization of* GL(N) *invariant transfer-matrices and quantum* N-wave *system (Lee model)*, J. Phys A: Math. Theor. **15** (1983), L591–L596.
- [KS82] P. Kulish, E. Sklyanin, On solutions of the Yang-Baxter equation, Zap. Nauchn. Sem. LOMI 95 (1980), 129–160; Engl. transl.: J. Soviet Math. 19 (1982) 19–56.
- [Lu20] K. Lu, Perfect integrability and Gaudin models, SIGMA 16 (2020), 132, 10 pages.
- [Lu21] K. Lu, A note on odd reflections of super Yangian and Bethe ansatz, arXiv:2111.10655.
- [LM21a] K. Lu, E. Mukhin, On the supersymmetric XXX spin chains associated with gl_{1|1}, Commun. Math. Phys. 386 (2021), 711–747.

- [LM21b] K. Lu, E. Mukhin, Jacobi-Trudi identity and Drinfeld functor for super Yangian, Int. Math. Res. Not. IMRN, 2021 (2021), no. 21, 16749–16808.
- [LM21c] K. Lu, E. Mukhin, *Bethe ansatz equations for orthosymplectic Lie superalgebra and self-dual superspaces*, Ann. Henri Poincaré **22** (2021), 4087–4130.
- [MM15] A. Molev, E. Mukhin, *Invariants of the vacuum module associated with the Lie superalgebra* gl(1|1), J. Phys A: Math. Theor. **48** (2015), no. 31, 314001, 20 pp.
- [MNV20] J. M. Maillet, G. Niccoli, L. Vignoli, Separation of variables bases for integrable $gl_{\mathcal{M}|\mathcal{N}}$ and Hubbard models, SciPost Phys. 9, 060 (2020), 55 pages.
- [MR14] A. Molev, E. Ragoucy, *The MacMahon Master Theorem for right quantum superalgebras and higher Sugawara operators for* $\widehat{\mathfrak{gl}}(m|n)$, Moscow Math. J., **14** (2014), no. 1, 83–119.
- [MTV06] E. Mukhin, V. Tarasov, A. Varchenko, *Bethe eigenvectors of higher transfer matrices*, J. Stat. Mech. Theor. Exp. (2006) P08002.
- [MTV08] E. Mukhin, V. Tarasov, A. Varchenko, *Spaces of quasi-exponentials and representations of* \mathfrak{gl}_N , J. Phys A: Math. Theor. **41** (2008), no. 19, 194017, 28 pp.
- [MTV09] E. Mukhin, V. Tarasov, A. Varchenko, *Schubert calculus and representations of general linear group*, J. Amer. Math. Soc. **22** (2009), no. 4, 909–940.
- [MVY15] E. Mukhin, B. Vicedo, C. Young, *Gaudin models for* $\mathfrak{gl}(m|n)$, J. Math. Phys. **56** (2015), no. 5, 051704, 30 pp,
- [Naz91] M. Nazarov, Quantum Berezinian and the classical capelli identity, Lett. Math. Phys. 21 (1991), 123–131.
- [Naz20] M. Nazarov, Yangian of the General Linear Lie Superalgebra, SIGMA Symmetry Integrability Geom. Methods Appl. 16 (2020), Paper No. 112, 24 pages.
- [Zha17] H. Zhang, Fundamental representations of quantum affine superalgebras and R-matrices, Transform. Groups 22 (2017), no. 2, 559–590.
 - K.L.: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DENVER,

2390 S. York St., Denver, CO 80208, USA

Email address: kang.lu@du.edu