

Search for the Gravitational-wave Background from Cosmic Strings with the Parkes Pulsar Timing Array Second Data Release

Zu-Cheng Chen,^{1,2,*} Yu-Mei Wu,^{3,4,†} and Qing-Guo Huang^{3,4,5,‡}

¹*Department of Astronomy, Beijing Normal University, Beijing 100875, China*

²*Advanced Institute of Natural Sciences, Beijing Normal University, Zhuhai 519087, China*

³*CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China*

⁴*School of Physical Sciences, University of Chinese Academy of Sciences, No. 19A Yuquan Road, Beijing 100049, China*

⁵*School of Fundamental Physics and Mathematical Sciences, Hangzhou Institute for Advanced Study, UCAS, Hangzhou 310024, China*

(Dated: May 17, 2022)

We perform a direct search for an isotropic stochastic gravitational-wave background (SGWB) produced by cosmic strings in the Parkes Pulsar Timing Array second data release. We find no evidence for such an SGWB, and therefore place 95% confidence level upper limits on the cosmic string tension, $G\mu$, as a function of the reconnection probability, p , which can be less than 1 in the string-theory-inspired models. The upper bound on the cosmic string tension is $G\mu \lesssim 5.1 \times 10^{-10}$ for $p = 1$, which is about five orders of magnitude tighter than the bound derived from the null search of individual gravitational wave burst from cosmic string cusps in the PPTA DR2.

I. INTRODUCTION

Over the past few years, the gravitational wave (GW) community witnessed the detection of a population of GW events [1–4] with ground-based interferometers that are sensitive in frequencies from Hz to kHz. A pulsar timing array (PTA) [5–7] offers a unique opportunity of extending the GW observations to the very low frequencies from nHz to μ Hz, by regularly monitoring the time of arrivals (TOAs) of radio pulses from an array of highly stable millisecond pulsars in the Milky Way. There are three major PTA projects, namely the European PTA (EPTA) [8], the North American Nanohertz Observatory for GWs (NANOGrav) [9], and the Parkes PTA (PPTA) [10]. These PTA projects have been monitoring the TOAs from dozens of pulsars with a weekly to monthly cadence for more than a decade. These PTAs along with the Indian PTA (InPTA) [11], the Chinese PTA (CPTA) [12] and the MeerKAT interferometer [13], support the International PTA (IPTA) [14, 15]. Potential GW sources in the PTA frequency band include a variety of physical phenomena such as supermassive black hole binaries (SMBHBs) [16–18], scalar-induced GWs [19–24], first-order phase transition [25, 26], and cosmic strings [27–30].

Recently, the NANOGrav [31], PPTA [32], EPTA [33] and IPTA [34] successively reported strong evidence for a stochastic common-spectrum process modeled by a power-law spectrum in their latest data sets. However, there is no significant evidence for the tensor transverse spatial correlations, which are necessary to claim a detection of stochastic GW background (SGWB) predicted

by general relativity. The origin of this spatially uncorrelated common-spectrum process (UCP) is still unknown. Further investigations indicate that the UCP can possibly come from various physical processes such as phase transitions [35–40], domain walls [37, 41, 42], cosmic strings [37, 41, 43, 44], or the non-tensorial polarization modes from alternative gravity theories [45–48].

Cosmic strings are linear topological defects that can either form in the early Universe from symmetry-breaking phase transitions at high energies [49–52] or be the fundamental strings of superstring theory (or one-dimensional D-branes) stretched out to astrophysical lengths [52, 53]. After their formation, the intersection between cosmic strings can lead to reconnections and form loops, which will then decay due to relativistic oscillation and emit GWs. PTA observations may detect a cosmic string network either through GW bursts emitted at cusps or through the SGWB superposed by radiation from all loops existing through cosmic history. A null detection of the individual GW burst from cosmic strings in PPTA DR2 has been reported in [30], thus placing a 95% upper limit on the cosmic string tension to be $G\mu \lesssim 10^{-5}$. In this work, we perform the first direct search for the SGWB produced from a network of cosmic strings in the PPTA DR2 data set. The remainder of this paper is organized as follows. In Sec. II, we review the SGWB energy spectrum produced by the cosmic strings. In Sec. III, we describe the data set and methodology used in our analyses. Finally, we summarize the results and give some discussion in Sec. IV.

II. SGWB FROM COSMIC STRINGS

We now review the SGWB produced by cosmic strings following [54]. A cosmic string network consists of both long (or “infinite”) strings that are longer than the horizon size and loops formed from smaller strings. When

* zucheng.chen@bnu.edu.cn

† Corresponding author: wuyumei@itp.ac.cn

‡ Corresponding author: huangqg@itp.ac.cn

two cosmic strings meet one another, they can exchange partners with a reconnection probability p , and form loops. Once loops are formed, they oscillate and decay through the emission of GWs [50], shrinking in size. A cosmic string network grows along with the cosmic expansion and evolves toward the scaling regime in which all the fundamental properties of the system scale with the cosmic time. The scaling regime can be achieved through the formation and subsequent decay of loops. The GW spectrum created by a cosmic string network is exceptionally broadband, depending on the size of the loops created.

We describe the GW energy spectrum of cosmic strings in terms of the dimensionless tension, $G\mu$, and the reconnection probability, p . Even though $p = 1$ for classical strings, it can be less than 1 in the string-theory-inspired models. In this work, we adopt the convention that the speed of light $c = 1$. The dimensionless GW energy density parameter per logarithm frequency as the fraction of the critical energy density is [54]

$$\Omega_{\text{gw}}(f) = \frac{8\pi Gf}{3H_0^2 p} \rho_{\text{gw}}(t_0, f), \quad (1)$$

where t_0 is cosmic time today, and ρ_{gw} is the GW energy density per unit frequency that can be computed by

$$\rho_{\text{gw}}(t, f) = G\mu^2 \sum_{n=1}^{\infty} C_n P_n, \quad (2)$$

with

$$C_n(f) = \int_0^{t_0} \frac{dt}{(1+z)^5} \frac{2n}{f^2} n(l, t). \quad (3)$$

Here, P_n is the radiation power spectrum of each loop, and $n(l, t)$ is the density of loops per unit volume per unit range of loop length l existing at time t . The SGWB from a network of cosmic strings has been computed in [54] with a complete end-to-end method by (i) simulating the long string network to extract a representative sample of loop shapes; (ii) using a smoothing model to estimate loop shape deformations due to gravitational backreaction; (iii) computing GW spectrum for each loop; (iv) evaluating the distribution of loops over redshift by integrating over cosmological time; (v) integrating the GW spectrum of each loop over the redshift-dependent loop distribution to get the overall emission spectrum; and at last (vi) integrating the overall emission spectrum over cosmological time to get the current SGWB. The simulations span a large parameter space of $G\mu \in [10^{-25}, 10^{-8}]$, and $f \in [10^{-15}, 10^{10}]$; and the output of the expected energy density spectra has been made publicly available¹.

III. THE DATA SET AND METHODOLOGY

The PPTA DR2 [55] data set includes pulse TOAs from high-precision timing observations for 26 pulsars collected with the 64-m Parkes radio telescope in Australia. The data were acquired between 2003 and 2018, spanning about 15 years, with observations taken at a cadence of approximately three weeks [55]. Details of the observing systems and data processing procedures are described in [10, 55].

To search for the GW signal from the PTA data, one needs to provide a comprehensive description of the stochastic processes that can induce the arrival time variations. The stochastic processes can be categorized as being correlated (red) or uncorrelated (white) in time. A careful analysis of the noise processes for individual pulsars in the PPTA sample has been performed in [56], showing that the PPTA data sets contain a wide variety of noise processes, including instrument dependent or band-dependent processes. Similar to [47], we adopt the noise model developed in [56] to characterize the noise processes. After subtracting the timing model of the pulsar from the TOAs, the timing residuals $\delta\mathbf{t}$ for each single pulsar can be decomposed into (see *e.g.* [57])

$$\delta\mathbf{t} = M\boldsymbol{\epsilon} + \delta\mathbf{t}_{\text{RN}} + \delta\mathbf{t}_{\text{DET}} + \delta\mathbf{t}_{\text{WN}} + \delta\mathbf{t}_{\text{CP}}. \quad (4)$$

The first term $M\boldsymbol{\epsilon}$ in the above equation accounts for the inaccuracies in the subtraction of timing model [58], where M is the timing model design matrix obtained from TEMPO2 [59, 60] through `libstempo`² interface, and $\boldsymbol{\epsilon}$ is a small offset vector denoting the difference between the true parameters and the estimated parameters of timing model. The second term $\delta\mathbf{t}_{\text{RN}}$ is the stochastic contribution from red noise [56], including achromatic spin noise (SN) [61]; frequency-dependent dispersion measure (DM) noise [62]; frequency-dependent chromatic noise (CN) [63]; achromatic band noise (BN) and system (“group”) noise (GN) [57]. We use 30 frequency components for the red noise of the individual pulsar. The third term $\delta\mathbf{t}_{\text{DET}}$ represents deterministic noise [56], including chromatic exponential dips [57], extreme scattering events [62], and annual dispersion measure variations [64]. The fourth term $\delta\mathbf{t}_{\text{WN}}$ represents white noise (WN), including a scale parameter on the TOA uncertainties (EFAC), an added variance (EQUAD), and a per-epoch variance (ECORR) for each backend/receiver system [65]. The last term $\delta\mathbf{t}_{\text{CP}}$ is the stochastic contribution due to the common-spectrum process (such as an SGWB), which is described by the cross-power spectral density [66]

$$S_{IJ}(f) = \frac{H_0^2}{16\pi^4 f^5} \Gamma_{IJ}(f) \Omega_{\text{GW}}(f), \quad (5)$$

¹ <http://cosmos.phy.tufts.edu/cosmic-string-spectra/>

² <https://vallis.github.io/libstempo>

Parameter	Description	Prior	Comments
White noise			
E_k	EFAC per backend/receiver system	U[0, 10]	single pulsar analysis only
Q_k [s]	EQUAD per backend/receiver system	log-U[-8.5, -5]	single pulsar analysis only
J_k [s]	ECORR per backend/receiver system	log-U[-8.5, -5]	single pulsar analysis only
Red noise (including SN, DM and CN)			
\mathcal{A}_{RN}	red noise power-law amplitude	log-U[-20, -8]	one parameter per pulsar
γ_{RN}	red noise power-law index	U[0, 10]	one parameter per pulsar
Band/System noise			
$\mathcal{A}_{\text{BN/GN}}$	band/group-noise power-law amplitude	log-U[-20, -8]	one parameter per band/system
$\gamma_{\text{BN/GN}}$	band/group-noise power-law index	U[0, 10]	one parameter per band/system
Deterministic noise			
\mathcal{A}_{E}	exponential dip amplitude	log-U[-10, -2]	one parameter per exponential dip event
t_{E} [MJD]	time of the event	U[57050, 57150] for PSR J1643 U[56100, 56500] for PSR J2145	one parameter per exponential dip event
$\log_{10}\tau_{\text{E}}$ [MJD]	relaxation time for the dip	U[log ₁₀ 5, 2]	one parameter per exponential-dip event
\mathcal{A}_{G}	Gaussian bump amplitude	log-U[-6, -1]	one parameter per Gaussian bump event
t_{G} [MJD]	time of the bump	U[53710, 54070]	one parameter per Gaussian bump event
σ_{G} [MJD]	width of the bump	U[20, 140]	one parameter per Gaussian bump event
\mathcal{A}_{Y}	annual variation amplitude	log-U[-10, -2]	one parameter per annual event
ϕ_{Y}	phase of the annual variation	U[0, 2 π]	one parameter per annual event
SGWB from cosmic string			
$G\mu$	cosmic string tension	log-U[-15, -8]	one parameter per PTA
p	reconnection probability	log-U[-3, 0]	one parameter per PTA

TABLE I. Model parameters and their prior distributions used in the Bayesian inference.

where Γ_{IJ} is the Hellings & Downs coefficients [67] measuring the spatial correlations of the pulsars I and J . Following [68], we use 5 frequency components for the common process among all of the pulsars.

We perform the Bayesian parameter inferences based on the methodology in [68, 69] to search for the SGWB from cosmic strings in the PPTA DR2 data set. Since it is challenging to obtain a complete noise model for pulsar J0437-4715 and pulsar J1713+0747 [56], we do not include these two pulsars. A summary of the noise model for the 24 pulsars used in our analyses can be found in Table 1 of [47]. We quantify the model selection scores by the Bayes factor defined as

$$\mathcal{BF} \equiv \frac{\Pr(\mathcal{D}|\mathcal{M}_2)}{\Pr(\mathcal{D}|\mathcal{M}_1)}, \quad (6)$$

where $\Pr(\mathcal{D}|\mathcal{M})$ measures the evidence that the data \mathcal{D} are produced under the hypothesis of model \mathcal{M} . Model \mathcal{M}_2 is preferred over \mathcal{M}_1 if the Bayes factor is sufficiently large. As a rule of thumb, $\mathcal{BF} \leq 3$ implies the evidence supporting the model \mathcal{M}_2 over \mathcal{M}_1 is “not worth more than a bare mention” [70]. In practice, we estimate the Bayes factor using the *product-space method* [71–74]. Our analyses are based on the latest JPL solar system ephemeris (SSE) DE438 [75]. We first infer the parameters of each single pulsar without including the common-spectrum process δt_{CP} in Eq. (4), and then fix the white

noise parameters to their max likelihood values from single-pulsar analysis to reduce the computational costs as was commonly done in literature (see *e.g.* [68, 69]). We use the open-source software packages `enterprise` [76] and `enterprise_extension` [77] to calculate the likelihood and Bayes factors and use `PTMCMCSampler` [78] package to do the Markov chain Monte Carlo sampling. Similar to [68, 79], we use draws from empirical distributions based on the posteriors obtained from the single-pulsar Bayesian analysis to sample the parameters of red noise and deterministic noise. Using empirical distributions can reduce the number of samples needed for the chains to burn in. The model parameters and their prior distributions are listed in Table I.

IV. RESULTS AND DISCUSSION

We first consider a model in which both the cosmic string tension $G\mu$ and the reconnection probability p are free parameters. Fig. 1 shows the posterior distributions of the $G\mu$ and p parameters obtained from the Bayesian search. The Bayes factor of the model including both the UCP and CS signal versus the model including only the UCP is $\mathcal{BF}_{\text{UCP}}^{\text{UCP}+\text{CS}} = 0.591 \pm 0.008$, indicating no evidence for an SGWB signal produced by the cosmic string in the PPTA DR2.




FIG. 1. One- and two-dimensional marginalized posterior distributions for the cosmic string tension, $G\mu$, and the reconnection probability, p . We show both the 1σ and 2σ contours in the two-dimensional plot.




FIG. 2. **Top panel:** the 95% CL upper limits on the cosmic string tension, $G\mu$, as a function of the reconnection probability, p , from the PPTA DR2. **Bottom panel:** the corresponding Bayes factor, \mathcal{BF} , as a function of the reconnection probability, p .

We also consider models in which the reconnection probability p is fixed to a specific value while the cosmic string tension $G\mu$ is allowed to vary. The lower panel of Fig. 2 shows the Bayes factor as a function of reconnection probability. For all the values of $p \in [10^{-3}, 1]$, we have $\mathcal{BF}_{\text{UCP}}^{\text{UCP}+\text{CS}} \lesssim 3$, confirming that there is no evidence for an SGWB produced by cosmic strings in the PPTA DR2. We, therefore, place 95% confidence level upper limit on cosmic string tension $G\mu$ as a function of reconnection probability p as shown in Fig. 2. The blue shaded region indicates parameter space that is excluded by the PPTA DR2. For $p = 1$, the upper bound on the cosmic string tension is $G\mu \lesssim 5.1 \times 10^{-10}$, which is about five orders of magnitude tighter than the bound of $G\mu \lesssim 10^{-5}$ [30] derived from the null search of individual gravitational wave burst from cosmic string cusps in the PPTA DR2. Note that Ω_{gw} is enhanced for $p < 1$, and therefore tighter constraints on $G\mu$ are obtained for $p < 1$.

To sum up, we have searched for the SGWB produced by a cosmic string network in the PPTA DR2 in the work. We find no evidence for such SGWB signal, and therefore place 95% upper limit on cosmic string tension as a function of reconnection probability.

ACKNOWLEDGMENTS

We thank Xing-Jiang Zhu, Zhi-Qiang You, Xiao-Jin Liu, Zhu Yi, and Shen-Shi Du for useful discussions. We acknowledge the use of HPC Cluster of ITP-CAS and HPC Cluster of Tianhe II in National Supercomputing Center in Guangzhou. This work is supported by the National Key Research and Development Program of China Grant No.2020YFC2201502, grants from NSFC (grant No. 11975019, 11991052, 12047503), Key Research Program of Frontier Sciences, CAS, Grant NO. ZDBS-LY-7009, CAS Project for Young Scientists in Basic Research YSBR-006, the Key Research Program of the Chinese Academy of Sciences (Grant NO. XDPB15).

[1] B. P. Abbott *et al.* (LIGO Scientific, Virgo), “GWTC-1: A Gravitational-Wave Transient Catalog of Compact Binary Mergers Observed by LIGO and Virgo during

the First and Second Observing Runs,” *Phys. Rev. X* **9**, 031040 (2019), [arXiv:1811.12907 \[astro-ph.HE\]](https://arxiv.org/abs/1811.12907).

[2] R. Abbott *et al.* (LIGO Scientific, Virgo), “GWTC-2:

- Compact Binary Coalescences Observed by LIGO and Virgo During the First Half of the Third Observing Run,” *Phys. Rev. X* **11**, 021053 (2021), arXiv:2010.14527 [gr-qc].
- [3] R. Abbott *et al.* (LIGO Scientific, VIRGO), “GWTC-2.1: Deep Extended Catalog of Compact Binary Coalescences Observed by LIGO and Virgo During the First Half of the Third Observing Run,” (2021), arXiv:2108.01045 [gr-qc].
- [4] R. Abbott *et al.* (LIGO Scientific, VIRGO, KAGRA), “GWTC-3: Compact Binary Coalescences Observed by LIGO and Virgo During the Second Part of the Third Observing Run,” (2021), arXiv:2111.03606 [gr-qc].
- [5] M. V. Sazhin, “Opportunities for detecting ultralong gravitational waves,” *Soviet Astronomy* **22**, 36–38 (1978).
- [6] Steven L. Detweiler, “Pulsar timing measurements and the search for gravitational waves,” *Astrophys. J.* **234**, 1100–1104 (1979).
- [7] R. S. Foster and D. C. Backer, “Constructing a Pulsar Timing Array,” *Astrophys. J.* **361**, 300 (1990).
- [8] Michael Kramer and David J. Champion, “The European Pulsar Timing Array and the Large European Array for Pulsars,” *Class. Quant. Grav.* **30**, 224009 (2013).
- [9] Maura A. McLaughlin, “The North American Nanohertz Observatory for Gravitational Waves,” *Class. Quant. Grav.* **30**, 224008 (2013), arXiv:1310.0758 [astro-ph.IM].
- [10] R. N. Manchester *et al.*, “The Parkes Pulsar Timing Array Project,” *Publ. Astron. Soc. Austral.* **30**, 17 (2013), arXiv:1210.6130 [astro-ph.IM].
- [11] Bhal Chandra Joshi *et al.*, “Precision pulsar timing with the ORT and the GMRT and its applications in pulsar astrophysics,” (2018), 10.1007/s12036-018-9549-y.
- [12] K. J. Lee, “Prospects of Gravitational Wave Detection Using Pulsar Timing Array for Chinese Future Telescopes,” in *Frontiers in Radio Astronomy and FAST Early Sciences Symposium 2015*, Astronomical Society of the Pacific Conference Series, Vol. 502, edited by L. Qain and D. Li (2016) p. 19.
- [13] M. Bailes *et al.*, “The MeerKAT telescope as a pulsar facility: System verification and early science results from MeerTime,” *Publ. Astron. Soc. Austral.* **37**, e028 (2020), arXiv:2005.14366 [astro-ph.IM].
- [14] G. Hobbs *et al.*, “The international pulsar timing array project: using pulsars as a gravitational wave detector,” *Gravitational waves. Proceedings, 8th Edoardo Amaldi Conference, Amaldi 8, New York, USA, June 22-26, 2009*, *Class. Quant. Grav.* **27**, 084013 (2010), arXiv:0911.5206 [astro-ph.SR].
- [15] R. N. Manchester, “The International Pulsar Timing Array,” *Class. Quant. Grav.* **30**, 224010 (2013), arXiv:1309.7392 [astro-ph.IM].
- [16] Andrew H. Jaffe and Donald C. Backer, “Gravitational waves probe the coalescence rate of massive black hole binaries,” *Astrophys. J.* **583**, 616–631 (2003), arXiv:astro-ph/0210148.
- [17] Alberto Sesana, Alberto Vecchio, and Carlo Nicola Colacino, “The stochastic gravitational-wave background from massive black hole binary systems: implications for observations with Pulsar Timing Arrays,” *Mon. Not. Roy. Astron. Soc.* **390**, 192 (2008), arXiv:0804.4476 [astro-ph].
- [18] A. Sesana, A. Vecchio, and M. Volonteri, “Gravitational waves from resolvable massive black hole binary systems and observations with Pulsar Timing Arrays,” *Mon. Not. Roy. Astron. Soc.* **394**, 2255 (2009), arXiv:0809.3412 [astro-ph].
- [19] Ryo Saito and Jun’ichi Yokoyama, “Gravitational wave background as a probe of the primordial black hole abundance,” *Phys. Rev. Lett.* **102**, 161101 (2009), [Erratum: *Phys. Rev. Lett.* 107, 069901 (2011)], arXiv:0812.4339 [astro-ph].
- [20] Chen Yuan, Zu-Cheng Chen, and Qing-Guo Huang, “Probing Primordial-Black-Hole Dark Matter with Scalar Induced Gravitational Waves,” (2019), arXiv:1906.11549 [astro-ph.CO].
- [21] Chen Yuan, Zu-Cheng Chen, and Qing-Guo Huang, “Log-dependent slope of scalar induced gravitational waves in the infrared regions,” (2019), arXiv:1910.09099 [astro-ph.CO].
- [22] Zu-Cheng Chen, Chen Yuan, and Qing-Guo Huang, “Pulsar Timing Array Constraints on Primordial Black Holes with NANOGrav 11-Year Dataset,” *Phys. Rev. Lett.* **124**, 251101 (2020), arXiv:1910.12239 [astro-ph.CO].
- [23] Chen Yuan, Zu-Cheng Chen, and Qing-Guo Huang, “Scalar Induced Gravitational Waves in Different Gauges,” (2019), arXiv:1912.00885 [astro-ph.CO].
- [24] Zu-Cheng Chen, Chen Yuan, and Qing-Guo Huang, “Confronting the primordial black hole scenario with the gravitational-wave events detected by LIGO-Virgo,” *Phys. Lett. B* **829**, 137040 (2022), arXiv:2108.11740 [astro-ph.CO].
- [25] Edward Witten, “Cosmic Separation of Phases,” *Phys. Rev. D* **30**, 272–285 (1984).
- [26] C.J. Hogan, “Gravitational radiation from cosmological phase transitions,” *Mon. Not. Roy. Astron. Soc.* **218**, 629–636 (1986).
- [27] L. Lentati *et al.*, “European Pulsar Timing Array Limits On An Isotropic Stochastic Gravitational-Wave Background,” *Mon. Not. Roy. Astron. Soc.* **453**, 2576–2598 (2015), arXiv:1504.03692 [astro-ph.CO].
- [28] Jose J. Blanco-Pillado, Ken D. Olum, and Xavier Siemens, “New limits on cosmic strings from gravitational wave observation,” *Phys. Lett. B* **778**, 392–396 (2018), arXiv:1709.02434 [astro-ph.CO].
- [29] Z. Arzoumanian *et al.* (NANOGrav), “The NANOGrav 11-year Data Set: Pulsar-timing Constraints On The Stochastic Gravitational-wave Background,” *Astrophys. J.* **859**, 47 (2018), arXiv:1801.02617 [astro-ph.HE].
- [30] N. Yonemaru *et al.*, “Searching for gravitational wave bursts from cosmic string cusps with the Parkes Pulsar Timing Array,” *Mon. Not. Roy. Astron. Soc.* **501**, 701–712 (2021), arXiv:2011.13490 [gr-qc].
- [31] Zaven Arzoumanian *et al.* (NANOGrav), “The NANOGrav 12.5-year Data Set: Search For An Isotropic Stochastic Gravitational-Wave Background,” (2020), arXiv:2009.04496 [astro-ph.HE].
- [32] Boris Goncharov *et al.*, “On the Evidence for a Common-spectrum Process in the Search for the Nanohertz Gravitational-wave Background with the Parkes Pulsar Timing Array,” *Astrophys. J. Lett.* **917**, L19 (2021), arXiv:2107.12112 [astro-ph.HE].
- [33] S. Chen *et al.*, “Common-red-signal analysis with 24-yr high-precision timing of the European Pulsar Timing Array: inferences in the stochastic gravitational-wave background search,” *Mon. Not. Roy. Astron. Soc.* **508**, 4970–4993 (2021), arXiv:2110.13184 [astro-ph.HE].
- [34] J. Antoniadis *et al.*, “The International Pulsar Timing

- Array second data release: Search for an isotropic gravitational wave background,” *Mon. Not. Roy. Astron. Soc.* **510**, 4873–4887 (2022), [arXiv:2201.03980 \[astro-ph.HE\]](#).
- [35] Andrea Addazi, Yi-Fu Cai, Qingyu Gan, Antonino Marciano, and Kaiqiang Zeng, “NANOGrav results and dark first order phase transitions,” *Sci. China Phys. Mech. Astron.* **64**, 290411 (2021), [arXiv:2009.10327 \[hep-ph\]](#).
- [36] Wolfram Ratzinger and Pedro Schwaller, “Whispers from the dark side: Confronting light new physics with NANOGrav data,” *SciPost Phys.* **10**, 047 (2021), [arXiv:2009.11875 \[astro-ph.CO\]](#).
- [37] Ligong Bian, Rong-Gen Cai, Jing Liu, Xing-Yu Yang, and Ruiyu Zhou, “Evidence for different gravitational-wave sources in the NANOGrav dataset,” *Phys. Rev. D* **103**, L081301 (2021), [arXiv:2009.13893 \[astro-ph.CO\]](#).
- [38] Shou-Long Li, Lijing Shao, Puxun Wu, and Hongwei Yu, “NANOGrav signal from first-order confinement-deconfinement phase transition in different QCD-matter scenarios,” *Phys. Rev. D* **104**, 043510 (2021), [arXiv:2101.08012 \[astro-ph.CO\]](#).
- [39] Xiao Xue *et al.*, “Constraining Cosmological Phase Transitions with the Parkes Pulsar Timing Array,” *Phys. Rev. Lett.* **127**, 251303 (2021), [arXiv:2110.03096 \[astro-ph.CO\]](#).
- [40] Zaven Arzoumanian *et al.* (NANOGrav), “Searching for Gravitational Waves from Cosmological Phase Transitions with the NANOGrav 12.5-Year Dataset,” *Phys. Rev. Lett.* **127**, 251302 (2021), [arXiv:2104.13930 \[astro-ph.CO\]](#).
- [41] Deng Wang, “Novel Physics with International Pulsar Timing Array: Axionlike Particles, Domain Walls and Cosmic Strings,” (2022), [arXiv:2203.10959 \[astro-ph.CO\]](#).
- [42] Ricardo Z. Ferreira, Alessio Notari, Oriol Pujolas, and Fabrizio Rompineve, “Gravitational Waves from Domain Walls in Pulsar Timing Array Datasets,” (2022), [arXiv:2204.04228 \[astro-ph.CO\]](#).
- [43] Simone Blasi, Vedran Brdar, and Kai Schmitz, “Has NANOGrav found first evidence for cosmic strings?” *Phys. Rev. Lett.* **126**, 041305 (2021), [arXiv:2009.06607 \[astro-ph.CO\]](#).
- [44] John Ellis and Marek Lewicki, “Cosmic String Interpretation of NANOGrav Pulsar Timing Data,” *Phys. Rev. Lett.* **126**, 041304 (2021), [arXiv:2009.06555 \[astro-ph.CO\]](#).
- [45] Zu-Cheng Chen, Chen Yuan, and Qing-Guo Huang, “Non-tensorial gravitational wave background in NANOGrav 12.5-year data set,” *Sci. China Phys. Mech. Astron.* **64**, 120412 (2021), [arXiv:2101.06869 \[astro-ph.CO\]](#).
- [46] Zaven Arzoumanian *et al.* (NANOGrav), “The NANOGrav 12.5-year Data Set: Search for Non-Einsteinian Polarization Modes in the Gravitational-wave Background,” *Astrophys. J. Lett.* **923**, L22 (2021), [arXiv:2109.14706 \[gr-qc\]](#).
- [47] Yu-Mei Wu, Zu-Cheng Chen, and Qing-Guo Huang, “Constraining the Polarization of Gravitational Waves with the Parkes Pulsar Timing Array Second Data Release,” *Astrophys. J.* **925**, 37 (2022), [arXiv:2108.10518 \[astro-ph.CO\]](#).
- [48] Zu-Cheng Chen, Yu-Mei Wu, and Qing-Guo Huang, “Searching for Isotropic Stochastic Gravitational-Wave Background in the International Pulsar Timing Array Second Data Release,” (2021), [arXiv:2109.00296 \[astro-ph.CO\]](#).
- [49] T. W. B. Kibble, “Topology of Cosmic Domains and Strings,” *J. Phys. A* **9**, 1387–1398 (1976).
- [50] A. Vilenkin, “Gravitational radiation from cosmic strings,” *Phys. Lett. B* **107**, 47–50 (1981).
- [51] Alexander Vilenkin, “Cosmic Strings and Domain Walls,” *Phys. Rept.* **121**, 263–315 (1985).
- [52] Gia Dvali and Alexander Vilenkin, “Formation and evolution of cosmic D strings,” *JCAP* **03**, 010 (2004), [arXiv:hep-th/0312007](#).
- [53] Edmund J. Copeland, Robert C. Myers, and Joseph Polchinski, “Cosmic F and D strings,” *JHEP* **06**, 013 (2004), [arXiv:hep-th/0312067](#).
- [54] Jose J. Blanco-Pillado and Ken D. Olum, “Stochastic gravitational wave background from smoothed cosmic string loops,” *Phys. Rev. D* **96**, 104046 (2017), [arXiv:1709.02693 \[astro-ph.CO\]](#).
- [55] Matthew Kerr *et al.*, “The Parkes Pulsar Timing Array project: second data release,” *Publ. Astron. Soc. Austral.* **37**, e020 (2020), [arXiv:2003.09780 \[astro-ph.IM\]](#).
- [56] Boris Goncharov *et al.*, “Identifying and mitigating noise sources in precision pulsar timing data sets,” *Mon. Not. Roy. Astron. Soc.* **502**, 478–493 (2021), [arXiv:2010.06109 \[astro-ph.HE\]](#).
- [57] L. Lentati *et al.*, “From Spin Noise to Systematics: Stochastic Processes in the First International Pulsar Timing Array Data Release,” *Mon. Not. Roy. Astron. Soc.* **458**, 2161–2187 (2016), [arXiv:1602.05570 \[astro-ph.IM\]](#).
- [58] Sydney J. Chamberlin, Jolien D. E. Creighton, Xavier Siemens, Paul Demorest, Justin Ellis, Larry R. Price, and Joseph D. Romano, “Time-domain Implementation of the Optimal Cross-Correlation Statistic for Stochastic Gravitational-Wave Background Searches in Pulsar Timing Data,” *Phys. Rev. D* **91**, 044048 (2015), [arXiv:1410.8256 \[astro-ph.IM\]](#).
- [59] George Hobbs, R. Edwards, and R. Manchester, “Tempo2, a new pulsar timing package. 1. overview,” *Mon. Not. Roy. Astron. Soc.* **369**, 655–672 (2006), [arXiv:astro-ph/0603381](#).
- [60] Russell T. Edwards, G. B. Hobbs, and R. N. Manchester, “Tempo2, a new pulsar timing package. 2. The timing model and precision estimates,” *Mon. Not. Roy. Astron. Soc.* **372**, 1549–1574 (2006), [arXiv:astro-ph/0607664](#).
- [61] Ryan M. Shannon and James M. Cordes, “Assessing the Role of Spin Noise in the Precision Timing of Millisecond Pulsars,” *Astrophys. J.* **725**, 1607–1619 (2010), [arXiv:1010.4794 \[astro-ph.SR\]](#).
- [62] M. J. Keith *et al.*, “Measurement and correction of variations in interstellar dispersion in high-precision pulsar timing,” *Mon. Not. Roy. Astron. Soc.* **429**, 2161 (2013), [arXiv:1211.5887 \[astro-ph.GA\]](#).
- [63] Andrew G. Lyne and Francis Graham-Smith, *Pulsar astronomy*, 48 (Cambridge University Press, 2012).
- [64] W. A. Coles *et al.*, “Pulsar Observations of Extreme Scattering Events,” *Astrophys. J.* **808**, 113 (2015), [arXiv:1506.07948 \[astro-ph.SR\]](#).
- [65] Z. Arzoumanian *et al.* (NANOGrav), “The NANOGrav Nine-year Data Set: Limits on the Isotropic Stochastic Gravitational Wave Background,” *Astrophys. J.* **821**, L3 (2016), [arXiv:1508.03024 \[astro-ph.GA\]](#).
- [66] Eric Thrane and Joseph D. Romano, “Sensitivity curves for searches for gravitational-wave backgrounds,” *Phys. Rev. D* **88**, 124032 (2013), [arXiv:1310.5300 \[astro-ph.IM\]](#).

- [67] R. w. Hellings and G. s. Downs, “UPPER LIMITS ON THE ISOTROPIC GRAVITATIONAL RADIATION BACKGROUND FROM PULSAR TIMING ANALYSIS,” *Astrophys. J.* **265**, L39–L42 (1983).
- [68] Zaven Arzoumanian *et al.* (NANOGrav), “The NANOGrav 12.5 yr Data Set: Search for an Isotropic Stochastic Gravitational-wave Background,” *Astrophys. J. Lett.* **905**, L34 (2020), [arXiv:2009.04496 \[astro-ph.HE\]](#).
- [69] Z. Arzoumanian *et al.* (NANOGrav), “The NANOGrav 11-year Data Set: Pulsar-timing Constraints On The Stochastic Gravitational-wave Background,” *Astrophys. J.* **859**, 47 (2018), [arXiv:1801.02617 \[astro-ph.HE\]](#).
- [70] Robert E. Kass and Adrian E. Raftery, “Bayes factors,” *Journal of the American Statistical Association* **90**, 773–795 (1995).
- [71] Bradley P. Carlin and Siddhartha Chib, “Bayesian model choice via markov chain monte carlo methods,” *Journal of the Royal Statistical Society. Series B (Methodological)* **57**, 473–484 (1995).
- [72] Simon J. Godsill, “On the relationship between markov chain monte carlo methods for model uncertainty,” *Journal of Computational and Graphical Statistics* **10**, 230–248 (2001).
- [73] Sonke Hee, Will Handley, Mike P. Hobson, and Anthony N. Lasenby, “Bayesian model selection without evidences: application to the dark energy equation-of-state,” *Mon. Not. Roy. Astron. Soc.* **455**, 2461–2473 (2016), [arXiv:1506.09024 \[astro-ph.CO\]](#).
- [74] Stephen R. Taylor, Rutger van Haasteren, and Alberto Sesana, “From Bright Binaries To Bumpy Backgrounds: Mapping Realistic Gravitational Wave Skies With Pulsar-Timing Arrays,” *Phys. Rev. D* **102**, 084039 (2020), [arXiv:2006.04810 \[astro-ph.IM\]](#).
- [75] William M. Folkner and Ryan S. Park, “Planetary ephemeris DE438 for Juno,” Tech. Rep. IOM392R-18-004, Jet Propulsion Laboratory, Pasadena, CA (2018).
- [76] Justin A. Ellis, Michele Vallisneri, Stephen R. Taylor, and Paul T. Baker, “Enterprise: Enhanced numerical toolbox enabling a robust pulsar inference suite,” Zenodo (2020).
- [77] Stephen R. Taylor, Paul T. Baker, Jeffrey S. Hazboun, Joseph Simon, and Sarah J. Vigeland, “enterprise_extensions,” (2021), v2.2.0.
- [78] Justin Ellis and Rutger van Haasteren, “jellis18/ptmcmcsampler: Official release,” (2017).
- [79] K. Aggarwal *et al.*, “The NANOGrav 11-Year Data Set: Limits on Gravitational Waves from Individual Supermassive Black Hole Binaries,” *Astrophys. J.* **880**, 2 (2019), [arXiv:1812.11585 \[astro-ph.GA\]](#).