Testing quantum gravity with primordial gravitational waves

Gianluca Calcagni,^a Leonardo Modesto^{b,c}

^aInstituto de Estructura de la Materia, CSIC, Serrano 121, 28006 Madrid, Spain

^bDepartment of Physics, Southern University of Science and Technology, Shenzhen 518055, China

 $^c\mathrm{Dipartimento}$ di Fisica, Università di Cagliari, Cittadella Universitaria, 09042 Monserrato, Italy

E-mail: g.calcagni@csic.es, leonardo.modesto@unica.it

Abstract. We propose a testable alternative to inflation directly built in a very general class of ultraviolet complete theories of quantum gravity enjoying Weyl conformal invariance. After the latter is spontaneously broken, logarithmic quantum corrections to the action make both the primordial tensor spectrum (from graviton fluctuations) and the scalar spectrum (from thermal fluctuations) quasi scale invariant. We predict a scalar spectral index $n_{\rm s}$ which only depends on two parameters and is consistent with observations, a tensor index $n_{\rm t} = 1 - n_{\rm s} > 0$, a tensor-to-scalar ratio $r_{0.05} \approx 0.01$ detectable by BICEP Array and LiteBIRD in the immediate or near future, respectively, and a blue-tilted stochastic gravitational-wave background observable by DECIGO in the further future.

Keywords: Alternatives to inflation, Cosmology of theories beyond the SM, Quantum gravity phenomenology, Primordial gravitational waves (theory)

Contents

| 1 | Introduction | 1 |
|----------|--|----|
| 2 | Summary of the setting | 2 |
| 3 | Primordial spectra | 5 |
| | 3.1 Flat spacetime approximation | 5 |
| | 3.2 Tensor perturbations | 9 |
| | 3.3 Scalar perturbations | 10 |
| | 3.4 Parameter space | 13 |
| | 3.5 Tensor-to-scalar ratio | 15 |
| | 3.6 Primordial GW background | 15 |
| 4 | Conclusions | 16 |
| A | Tree-level graviton propagator | 18 |
| | A.1 Tree-level propagator in the theory (2.1) | 18 |
| | A.2 One-loop graviton propagator | 20 |
| | A.3 One-loop linearized equations of motion | 24 |
| в | Spatial correlation function and power spectrum | 25 |
| С | Two-point correlation function of the energy-momentum tensor | 26 |

1 Introduction

The early universe is an ideal laboratory where to test theories beyond Einstein gravity and the Standard Model of quantum interactions. Microscopic scales can inflate to cosmological ones in the blink of an eye; extreme curvature regimes can dominate near the big bang or a big bounce; exotic particles can show their invisible presence through the cosmic evolution of the large-scale structures; and so on. Unfortunately, in general, one has to strike a compromise between theoretical control and predictivity, and the more rigorous and assumption-free is the theory, the less it makes contact with observations, and vice versa. For example, quantum gravity has made many of its advances to phenomenology in the context of inflation (see, e.g., [1] for an overview) but, until now, it has been challenging to overcome theoretical difficulties and, at the same time, extract predictions that could be falsified in the near future.

In its purest form as a single scalar field, the inflationary mechanism requires a considerable amount of engineering, from the choice of potential to the embedding in a particle-physics or quantum-gravity framework. It is significant that the most successful scenario of early-universe acceleration to date [2], Starobinsky inflation [3], is actually a model of gravity where the scalar degree of freedom is hidden, and that such model is not ultraviolet (UV) complete *per se* but requires an embedding in a larger theory of quantum gravity, such as those attempted in supergravity [4–6], nonlocal gravity [7–11], or string theory [12]. Considering also that string theory is facing a swampland crisis [13–15] and it seems difficult to realize single-field inflation therein, the drive to look for alternatives might be felt even more urgently by some.

In this paper, instead of trying to get inflation from a fundamental theory, we propose a candidate with fundamental nonlocality which contains, basically for free, the ingredients needed to solve the singularity, horizon and flatness problems of the hot big bang [16] and to explain the anisotropies of the cosmic microwave background (CMB), all without inflation. These achievements are due to an anomaly-free symmetry, Weyl (or conformal, or scale) invariance, and its spontaneous breaking; all the other details of the theory are unessential. As a phenomenological mechanism, conformal invariance as an alternative to inflation is not new [17-20] but, to date, it was not known how to make it work in a fundamental theory due to the challenge of obtaining UV finiteness. The novelty here lies, on one hand, in the fact that we provide a concrete setting where quantum conformal invariance is realized and, on the other hand, in the extraction of rigid predictions on the tensor sector which can be not only verified, but even falsified by present- and next-generation gravitationalwave (GW) observatories. Such rigidity comes from the fact that the cosmological model discussed below is derived *directly* from the full theory under assumptions that can hardly be relaxed without spoiling the self-consistency of the theory. For these reasons, this is arguably one of the sharpest instances, compared with few others [11, 21-24], where the observation of the gravity-wave universe is opening a robust window on microphysics.

The paper is organized as follows. In section 2, we briefly recall the setting of nonlocal quantum gravity, the transition from a Planckian conformally invariant phase to a post-Planckian nonconformal one and the resolution of the hot big-bang problems detailed in the companion paper [16]. In section 3, we compute the primordial tensor and scalar spectra generated during the post-Planckian phase, their indices, the tensor-to-scalar ratio and the GW background, showing that the theory yields characteristic predictions in a wide range of frequencies that, once combined, can be discriminated from inflationary models. Conclusions are in section 4. Appendices contain technical material and calculations.

2 Summary of the setting

A theory of gravity nonminimally coupled with matter was introduced in [25–27]. Thanks to the presence of certain nonlocal operators (i.e., terms with infinitely many derivatives), the quantum properties of this proposal are very similar to those of a more general class of nonlocal theories [28–37] and include perturbative unitarity (absence of classical instabilities or quantum ghost modes), same nonlinear stability properties as in Einstein gravity, same treelevel scattering amplitudes, same macro-causality properties, and super-renormalizability or finiteness at the quantum level. In four dimensions, the action is

$$S = \int d^4x \sqrt{|g|} \left[\mathcal{L}_{\rm loc} + E_i F^{ij}(\mathbf{\Delta}) E_j + V(E_i) \right], \qquad \mathcal{L}_{\rm loc} = \frac{M_{\rm Pl}^2}{2} R[g] + \mathcal{L}_{\rm m} + \mathcal{L}_2^{\rm loc}, \quad (2.1)$$

where $\Delta_{ki} \coloneqq \delta^2 S_{\text{loc}} / \delta \Phi_k \delta \Phi_i$ is the local Hessian for all gravitational and matter fields Φ_i , E_i give the local equations of motion $E_i = 0$, and $V(E_i)$ is a collection of local operators at least cubic in E that secure the finiteness of the theory in even dimensions [27, 33]. The local extremals are $E_{\mu\nu} \coloneqq (M_{\text{Pl}}^2 G_{\mu\nu} - T_{\mu\nu})/2$, where $M_{\text{Pl}}^2 \coloneqq (8\pi G)^{-1}$ is the reduced Planck mass, $G_{\mu\nu} = R - g_{\mu\nu}R/2$, and $T_{\mu\nu}$ is the energy-momentum tensor derived from the matter Lagrangian \mathcal{L}_{m} . The term $\mathcal{L}_2^{\text{loc}}$ will be explained shortly. The nonlocal equations of motions of the theory (2.1) in the gravitational sector are

$$\left[e^{\bar{\mathrm{H}}(\boldsymbol{\Delta})}\right]^{\sigma\tau}_{\mu\nu}E_{\sigma\tau} + O(E^2_{\mu\nu}) = 0.$$
(2.2)

where $\overline{\mathrm{H}}(\Delta) := \mathrm{H}(\Delta) - \mathrm{H}(0)$ is an entire analytic function of the Hessian entering the definition of the form factor F:

$$2\Delta F(\Delta) \equiv e^{\bar{\mathrm{H}}(\Delta)} - 1 = e^{\gamma_{\mathrm{E}} + \Gamma[0, p(z)]} p(z) - 1, \qquad (2.3a)$$

$$H(\mathbf{\Delta}) = \gamma_{\rm E} + \Gamma[0, p(z)] + \ln p(z), \qquad z = \frac{4\mathbf{\Delta}}{M_{\rm Pl}^2 \Lambda_*^2},$$
 (2.3b)

where Λ_* is an energy scale, $\gamma_{\rm E} \approx 0.577$ is the Euler–Mascheroni constant, Γ is the upper incomplete gamma function, and p(z) is the quartic polynomial

$$p(z) = b_0 - b z + z^4$$
, (2.3c)

where b > 0. Note that, with the choice (2.3c), H(0) = 0 when $b_0 = 0$ ($\bar{H} = H$); later on we will drop b_0 altogether, since it is negligible in the energy regimes we will consider. The form factor (2.3) simplifies previous proposals [29, 31] and guarantees a well-defined classical problem of initial conditions [38], the absence of extra poles in the propagators, a unique Lorentzian limit [39–41], unitarity, and power-counting super-renormalizability in four dimensions [27] (which is essentially given by the highest-order operators $\mathcal{R}^2 \to h \Box^2 h$ of Stelle gravity [42], where h is the graviton and $\mathcal{R} = R_{\mu\nu}, R$). In fact, in the UV the form factor is asymptotically polynomial:

$$e^{\mathrm{H}(z)} \stackrel{z \gg 1}{\simeq} e^{\gamma_{\mathrm{E}}} p(z) ,$$
 (2.4)

and one can perform the usual power counting of divergences in Feynman diagrams. Note for later that, if $b \gtrsim O(1)$, then there exists an intermediate UV regime where the linear term dominates over the quartic one.

Finally, the metric in eq. (2.1) is defined as $g_{\mu\nu} =: (\phi/M_{\rm Pl})^2 \hat{g}_{\mu\nu}$, where ϕ is the dilaton field. All matter fields are rescaled similarly, so that the theory is classically invariant under Weyl transformations

$$\hat{g}'_{\mu\nu} = \Omega^2(x)\,\hat{g}_{\mu\nu}\,,\qquad \phi' = \Omega^{-1}(x)\,\phi\,,\qquad\dots.$$
 (2.5)

Most importantly, the theory is also conformally invariant at the quantum level, since it is finite, all beta functions vanish, and the conformal anomaly is zero [27].

As a consequence, the theory has no physical energy scales, since all couplings are rescaled by powers of the dilaton field $\phi \equiv M_{\rm Pl}/\sqrt{2} + \varphi$. Still, one can define a UV and an infrared (IR) regime when comparing the energy component of the momentum and the scale Λ_* appearing in the form factors, which then become physical after conformal symmetry is broken [16]. At UV energy scales $E := k^0 \gg \Lambda_*$, the scattering amplitudes are suppressed [43]. At IR energy scales $E \ll \Lambda_*$, the quantum theory acquires logarithmic corrections just like Stelle gravity:

$$\mathcal{L}_Q = \beta_R R \ln\left(\frac{-\Box}{\Lambda_*^2 \delta_0^2}\right) R + \beta_{\mathrm{Ric}} R_{\mu\nu} \ln\left(\frac{-\Box}{\Lambda_*^2}\right) R^{\mu\nu}, \qquad (2.6)$$

where δ_0 is a constant to be determined by an explicit calculation we do not need here. At energy scales $E \sim M_{\rm Pl}$, the theory is manifestly Weyl invariant, there is no absolute notion of spacetime distance, and the correlation functions of fields with nonzero conformal weight are identically zero as a consequence of conformal symmetry [44]. In this conformal phase, the quantum corrections (2.6) always feature three or more fields and, therefore, do not enter perturbative two-point functions.

Spontaneous symmetry breaking occurs via a conformal-invariant term $\mathcal{L}_2^{\text{loc}}$ in the local Lagrangian \mathcal{L}_{loc} which provides a potential for the dilaton with minima. Two possibilities [16] are the quartic potential arising from a cosmological constant term $\sqrt{|g|} \Lambda_{\text{cc}} = \sqrt{|\hat{g}|} \Lambda_{\text{cc}} \phi^4$ or, more interestingly, from the Higgs-dilaton potential proposed in [45, 46]:

$$\mathcal{L}_2^{\text{loc}} = \lambda (\mathfrak{h}^{\dagger} \mathfrak{h} - \alpha \phi^2)^2 + \lambda' \phi^4 , \qquad (2.7)$$

where \mathfrak{h} is the Higgs field. At the minima, the dilaton acquires the expectation value $\phi = M_{\rm Pl}/\sqrt{2}$, determined empirically by the measurement of Newton's constant, and conformal symmetry is broken. When $\varphi \ll M_{\rm Pl}$ and the dilaton sits in its expectation value, the gravitational sector becomes local and quadratic in the Ricci tensor around energies $E \sim \Lambda_{\rm hd}$ [16],

$$\mathcal{L}_{\Lambda_{\rm hd}} = \frac{M_{\rm Pl}^2}{2} R + \mathcal{L}_{\rm m} + E_{\mu\nu} E^{\mu\nu} + V(E_{\mu\nu}) + \mathcal{L}_Q \,, \qquad (2.8)$$

where

$$\Lambda_{\rm hd} \coloneqq \sqrt{\frac{\max(1, b_0)}{b}} \Lambda_* \,, \tag{2.9}$$

(actually, it will turn out that we can drop b_0 so that $\Lambda_{\rm hd} = \Lambda_*/\sqrt{b}$), $E_{\mu\nu}E^{\mu\nu} \propto R_{\mu\nu}R^{\mu\nu}$ in vacuum in D = 4 dimensions (the expression with matter is [16, eq. (A.5)]), and the quantum corrections (2.6) contribute to the graviton propagator. In this sub-Planckian *Higgs phase*, a background is selected and one can consider cosmological perturbations thereon. At yet lower energies, $\mathcal{L} \sim R$ and Einstein gravity is recovered.

In [16], we started the description of the cosmology stemming from this theory. To recapitulate, there we concentrated on the background, leaving perturbations to the present paper. Conformal invariance provides an extremely simple, natural, and model-independent solution to the problems of hot-big-bang cosmology. All these issues are addressed in the conformal phase of the theory, while the quasi scale-invariant primordial temperature spectrum arises in the Higgs phase. For details, see [16, sections 3.3 and 4].

The big-bang problem is solved because the Borde–Guth–Vilenkin theorem, establishing geodesic incompleteness in a very general class of gravitational theories [47], does not apply here. In fact, the average expansion condition at the core of the theorem is not a conformally invariant statement and cannot be met in a theory where the dynamics is conformally invariant. Vice versa, a conformal transformation can always make an expanding background static or even contracting and any singular metric can be rescaled into a regular one. In particular, the Friedmann–Lemaître–Robertson–Walker (FLRW) metric can be mapped into asymptotically flat metrics.

The flatness problem is dispensed with by pure symmetry arguments. Conformal invariance forces all regions of the Universe to be in causal contact, which eventually lead to select homogeneous and isotropic metrics (i.e., Minkowski and FLRW with intrinsic curvature $\kappa = 0, \pm 1$) at the onset of the Higgs phase. The well-know observation that all FLRW metrics are physically equivalent to the FLRW spacetime with zero intrinsic curvature in a Weyl×Diff invariant theory further reduces the possibilities to Minkowski and FLRW with $\kappa = 0$.

The horizon problem does not arise because horizons and distances do not have any absolute meaning in the conformal phase; at the onset of the Higgs phase, a horizon forms which is much larger than the horizon of FLRW in Einstein's gravity. The monopole problem is avoided because in nonlocal quantum gravity all the couplings run to zero in the UV, similarly to the scenario of [48]. In the conformal phase, there are no causally disconnected domains where the Higgs field of a grand unification theory could take different expectation values.

Finally, the trans-Planckian problem does not occur because, while in standard inflation primordial perturbations with sub-Planckian wavelengths eventually become observable, here the conformal phase does not admit a comparison of physical lengths with any given scale. From the point of view of quantum field theory, the higher-derivative operators $E_i(\Box/M_{\rm Pl}^2)^n E^i$, important at or above the Planck energy, are quadratic in E_i and do not contribute to the linearized nonlocal equations of motion

$$e^{\bar{\mathrm{H}}(\mathbf{\Delta})^{(0)}} E_i^{(1)} + E_i^Q + O(E_i^2) = 0, \qquad (2.10)$$

where the superscripts denote, respectively, the background and the first order in field perturbations and E^Q is the contribution from the quantum corrections (2.6). In particular, all background solutions of the Einstein's equations $E_{\mu\nu} = 0$ (including FLRW metrics) are exact solutions of the theory (2.1). Examples of backgrounds, meaningful only after spontaneous symmetry breaking, that are exactly flat and geodesically complete are given in [16].

3 Primordial spectra

While the conformal phase can address the classic problems of the hot-big-bang model, it does not explain the origin of the primordial spectra and their almost scale invariance. It turns out that these are generated during the post-Planckian phase, where conformal invariance is spontaneously broken. The one-loop quantum corrections to the classical action are directly responsible for the small deviation from Harrison–Zel'dovich spectra.

In this section, after describing some basic approximations needed for physical and technical reasons, we derive the quasi-scale-invariant spectrum of primordial tensor and scalar perturbations. To this aim, we need the graviton propagator for the theory (2.1) with form factor (2.3). The tree-level expression can be found in Appendix A.1, while the graviton propagator from the one-loop quantum effective Lagrangian (2.8) is calculated in Appendix A.2.

3.1 Flat spacetime approximation

According to the discussion in [16, sections 2 and 3] and the results derived in Appendix A.2, the computations we are going to make in this section rely on two assumptions for the energy regime in which we will consider the theory (2.1):

(i) Quadratic-gravity or higher-derivative (hd) regime [16]

$$\Lambda_{\rm hd} \lesssim E \lesssim \Lambda_* \,, \tag{3.1}$$

In this range, the theory has the local limit (2.8) which captures its main UV properties.¹ The reason why one should consider the quadratic-curvature regime of the theory is not just convenience (the theory in the fully nonlocal regime may be very difficult

¹More generally, what follows holds also for Stelle gravity with a nonzero coefficient of the R^2 operator.

to handle cosmologically), but also empirical. We know that early-universe spectra are quasi-scale-invariant. Then, if they are generated by curvature (as they should in this model, since we do not have matter fields sustaining inflation), in four dimensions they must proceed from quadratic curvature terms, as is clear by the dimensional scaling of the graviton two-point function. Thus, if the theory can account for the primordial cosmological spectra, it must do so in the Stelle-like regime. Quantum corrections naturally break exact scale invariance, as described below.

(ii) Perturbative (one-loop) approximation (A.30), corresponding to $E \gtrsim \Lambda_*/10$. It guarantees that one-loop quantum corrections in the propagator are subdominant with respect to the classical part consistently with the quantum field theory perturbative expansion. This condition also permits some analytic simplifications (eqs. (A.25) and (A.26)).

Therefore, if $\max(1, b_0)/b < O(10^{-2})$, then $\Lambda_{\rm hd} < \Lambda_*/10$ and the conditions (i) and (ii) imply that $\Lambda_*/10 \leq E \leq \Lambda_*$, which boils down to pushing the energy to the lower limit:

$$E \simeq \frac{\Lambda_*}{10} \,. \tag{3.2}$$

If we make the identification $\Lambda_* = M_{\text{Pl}}$, then the regime (3.2) falls within the post-Planckian or Higgs phase where conformal invariance is spontaneously broken.

In [16, section 3.1], we argued that $\Lambda_* \leq M_{\rm Pl}$ only using conformal invariance. Here, we end up with a weaker statement using (3.2) and the fact that the energy of particles propagating in the Higgs phase must be smaller than the Planck mass, $E < M_{\rm Pl}$:

$$\Lambda_* < 10 M_{\rm Pl}, \qquad \ell_* > \frac{\ell_{\rm Pl}}{10}.$$
 (3.3)

where $\ell_* \coloneqq \Lambda_*^{-1}$ and $\ell_{\text{Pl}} \coloneqq M_{\text{Pl}}^{-1}$.

In order to compute the primordial scalar and tensor spectrum, we first show that we can approximate the asymptotic FLRW metric with the Minkowski one. Then, we can evaluate the correlation functions in flat spacetime and avoid the complications of quantum field theory on curved spaces. This approximation is not a novelty in certain scenarios of conformal cosmology [17, 18] and cosmologies from quantum gravity [49, 50] where the correct primordial spectra are generated subhorizon. The condition that has to be fulfilled is to consider perturbations within the Hubble horizon,

$$\lambda = \frac{1}{E} \ll R_H = H^{-1} \,, \tag{3.4}$$

where λ is the wavelength of the perturbation. There are three important moments in the evolution of the Universe: the instant t_{Λ_*} when $\lambda = \ell_*$ and the Universe exits the conformal phase, the time (or, more precisely, a narrow range of time scales t_{λ} when $\lambda \simeq 10\ell_*$ and the wavelength has the same scale of validity of the Stelle regime (see eq. (3.2)), and the instant $t_{\Lambda_{\rm hd}}$ when $\lambda = \Lambda_{\rm hd}^{-1}$ before which the theory can be approximated by the local quadratic action (2.8) and after which it starts to recover Einstein gravity. Apart from these, there is another time instant of interest t_* which signals the moment after which the cosmic expansion becomes almost indistinguishable from the one in Einstein gravity. As we will see shortly, the recovery of the background-independent Einstein action does not happen exactly at the

same time of the recovery of the standard cosmic expansion, and $t_{\Lambda_{\rm hd}} \neq t_{\star}$. In fact, since the class of FLRW metrics is an exact solution of the full theory ($E_{\mu\nu} = 0$, which solves eq. (2.2)), the standard cosmic expansion is expected to be reached some time before $t_{\Lambda_{\rm hd}}$, so that $t_{\Lambda_{\rm hd}} > t_{\star}$. This is indeed the case and t_{\star} is determined as follows.

Since we are in the regime (3.2), i.e., out of the conformally invariant phase, we have to consider a metric being asymptotically FLRW at late times when Einstein gravity is recovered. The most natural setting in all scenarios alternative to inflation is a radiationdominated Universe described by the scale factor

$$a(t) \simeq \sqrt{\frac{t}{t_0}} \,. \tag{3.5}$$

In Einstein gravity, a perturbation living in this background would cross the horizon at a time t_{\star} such that $\lambda(t_{\star}) = R_H(t_{\star})$. At the instant t, the wavelength is given by $\lambda = a(t) \lambda_{\text{com}}$, while $H = \dot{a}/a \simeq 1/(2t)$ and $R_H = H^{-1} \simeq 2t$. Therefore, at $t = t_{\star}$ we have

$$\sqrt{\frac{t_{\star}}{t_0}} \lambda_{\rm com} \simeq 2t_{\star} \implies t_{\star} \simeq \frac{\lambda_{\rm com}^2}{4t_0},$$
(3.6)

where $\lambda_{\rm com} = 2\pi/|\mathbf{k}|$ is the comoving wavelength and $|\mathbf{k}|$ is the comoving wavenumber. In conformal nonlocal gravity, if the condition (3.4) holds, then there is no horizon crossing during the production of primordial perturbations and t_{\star} has the physical interpretation advanced above, as one can see in Fig. 1. At later points in the evolution of the universe, perturbations $\lambda(t > t_{\star})$ are still generated *inside* the actual horizon $R_H > 2t$ calculated from the exact metric and the condition (3.4) holds also beyond t_{\star} .

Let us now find the instants t_{Λ_*} , t_{λ} , and $t_{\Lambda_{hd}}$ and provide the chronological order of the events for which eq. (3.4) is reliable. We take the asymptotic form (3.5) for the scale factor, which is sufficient to establish the hierarchy of time scales. The instant t_{Λ_*} is given by

$$t_{\Lambda_*}: \sqrt{\frac{t_{\Lambda_*}}{t_0}} \lambda_{\rm com} \simeq \ell_* \qquad \Longrightarrow \qquad t_{\Lambda_*} \simeq \frac{\ell_*^2 t_0}{\lambda_{\rm com}^2} = \frac{\ell_*^2}{4t_\star}.$$
 (3.7)

The instant t_{λ} is given by

$$t_{\lambda} \colon \sqrt{\frac{t_{\lambda}}{t_0}} \lambda_{\rm com} \simeq 10 \,\ell_* \qquad \Longrightarrow \qquad t_{\lambda} \simeq 10^2 \,t_{\Lambda_*} \,.$$
 (3.8)

The instant $t_{\Lambda_{\rm hd}}$ is given by

$$t_{\Lambda_{\rm hd}}: \ \sqrt{\frac{t_{\Lambda_{\rm hd}}}{t_0}} \lambda_{\rm com} \simeq \sqrt{\frac{b}{\max(1,b_0)}} \ell_* \qquad \Longrightarrow \qquad t_{\Lambda_{\rm hd}} \simeq \frac{b}{\max(1,b_0)} t_{\Lambda_*} \,. \tag{3.9}$$

Since $b/\max(1, b_0) \gg 1$, we have the hierarchy

$$t_{\Lambda_*} \ll t_{\lambda} < t_{\Lambda_{\rm hd}} \,, \tag{3.10}$$

which is the time analogue of the interval (3.1).

The condition (3.4) is satisfied in the regime (3.2) when

$$\lambda \simeq 10 \,\ell_* \ll H^{-1} \simeq 2t_\lambda \qquad \Longrightarrow \qquad t_\lambda \gg 5\ell_* \,. \tag{3.11}$$




Figure 1. The cosmological scenario proposed in this article. The black curve with support on the whole t line is the scale factor $a = \sqrt{A + t/t_0} e^{\Gamma(0,t/t_{\rm Pl})}$ [16, eq. (5.9a) and Fig. 4a], which is chosen only for definiteness among any of the other solutions described in [16] reducing to FLRW at late times. The black curve with support on the t > 0 half-line is the FLRW radiation-dominated scale factor (3.5). The red curve and red straight line are the Hubble radius associated with the above scale factors, respectively. For $t < t_{\Lambda_*}$, the Universe is in the trans-Planckian conformal phase described by the theory (2.1). For $t \gtrsim t_{\Lambda_*}$, the conformal symmetry is spontaneously broken. In the epoch $t_{\Lambda_*} < t \ll t_{\Lambda_{\rm hd}}$, the Universe is well described by quadratic gravity (Stelle's theory) plus quantum corrections. For $t > t_*$, the background solution is indistinguishable from a radiation-dominated Universe in Einstein gravity, although the Einstein-Hilbert action is recovered only at later times $t > t_{\Lambda_{\rm hd}}$. In Einstein gravity, t_* would represent the time of horizon crossing of the perturbation (k = aH), eq. (3.6), but notice that all regions of the Universe are in causal contact at any time in the plot for nonlocal quantum gravity (the Hubble radius is always larger than the scale factor). t_{λ} is when the perturbation is evaluated after the spontaneous breaking of the Weyl symmetry, consistently with the perturbative approximation (3.2). The CMB is formed at the decoupling time $t_{\rm dec}$.

Plugging eq. (3.6) into (3.11), we find

$$t_{\lambda} \gg 5\ell_* \simeq 10\sqrt{t_{\star} t_{\Lambda_*}},$$

in which we now replace $t_{\Lambda_*} \simeq t_{\lambda}/10^2$:

$$t_{\star} \ll t_{\lambda} \,. \tag{3.12}$$

Since t_{\star} marks an event posterior to the onset time of the Higgs phase, we also have $t_{\Lambda_*} < t_{\star}$ which, combined with (3.10) and (3.12), yields the interval

$$t_{\Lambda_*} < t_{\star} \ll t_{\lambda} < t_{\Lambda_{\rm hd}} \,. \tag{3.13}$$

The cosmic evolution (not just the scale factor a(t)) becomes standard at times $t > t_{\Lambda_{hd}}$, so that the time range of production of primordial perturbations in this cosmological model is roughly $t_{\Lambda_*} < t < t_{\Lambda_{hd}}$. The conformal and the Higgs phase as well as the scale hierarchy (3.13) and the local UV limit of the theory are depicted in Fig. 1.

3.2 Tensor perturbations

After the spontaneous breaking of conformal invariance, distances acquire a meaning, correlation functions become nonvanishing, and one can consider metric perturbations. In this noninflationary early-Universe scenario, tensor perturbations are the subhorizon metric quantum fluctuations corresponding to the graviton, in a regime where one can ignore the curvature of the background and the cosmological expansion. In contrast, in standard inflation metric fluctuations are induced by the quantum fluctuations of a scalar field and the spectrum is calculated at subhorizon scales but evaluated at horizon crossing.

The tensor spectrum in the regime (3.2) is readily obtained from the graviton propagator (A.31) derived in Appendix A.2. The two-point correlation function for transverse and traceless tensors in four dimensions reads [18]

$$\begin{aligned}
G_{ijkl}^{(2)}(x,x') &\coloneqq \langle h_{ij}(x)h_{kl}(x') \rangle \\
&\stackrel{(\mathbf{A}.31)}{=} -\mathbf{C} \int \frac{d^4k}{(2\pi)^4} e^{ik \cdot (x-x')} \frac{P_{ijkl}^{(2)}(k)}{k^4 \left(\frac{k}{\Lambda_*}\right)^{2\epsilon_2}} \\
&= -\mathbf{C}\Lambda_*^{2\epsilon_2} P_{ijkl}^{(2)}(\partial_x) \int \frac{d^4k}{(2\pi)^4} \frac{e^{ik \cdot (x-x')}}{k^{4+2\epsilon_2}},
\end{aligned} \tag{3.14}$$

where Latin indices run over spatial directions, the constants C and ϵ_2 are given by eqs. (A.31) and (A.19), respectively,

$$C \coloneqq \frac{4\Lambda_*^2}{b\,e^{\tilde{\gamma}_{\rm E}}M_{\rm Pl}^2}, \qquad \epsilon_2 = -\frac{2\beta_{\rm Ric}\Lambda_*^2}{b\,e^{\tilde{\gamma}_{\rm E}}M_{\rm Pl}^2}, \qquad \tilde{\gamma}_{\rm E} \coloneqq \gamma_{\rm E} - {\rm H}(0) \tag{3.15}$$

and $P_{ijkl}^{(2)}(k)$ are the spatial components of the spin-2 projector in eq. (A.3) [51–53]. The angular brackets in the first line of eq. (3.14) are defined via the Lorentzian path integral and the final expression is also Lorentzian. To perform the integral explicitly, however, we now move to Euclidean momenta as described in [54]. We recall that the two-point correlation function in *D*-dimensional Euclidean momentum space is obtained through the Fourier transform

$$(x^{2})^{-s} = B_{D}(2s) \int \frac{d^{D}k}{(2\pi)^{D}} e^{ik \cdot x} (k^{2})^{s - \frac{D}{2}}, \qquad (3.16a)$$

$$B_D(2s) \coloneqq \frac{2^{D-2s} \pi^{\frac{D}{2}} \Gamma\left(\frac{D}{2} - s\right)}{\Gamma(s)}.$$
(3.16b)

Therefore, comparing eqs. (3.14) and (3.16) with D = 4, we see that $s = -\epsilon_2$, so that the propagator in position space reads

$$G_{ijkl}^{(2)}(x,x') = -\frac{C\Lambda_*^{2\epsilon_2}}{B_4(-2\epsilon_2)} P_{ijkl}^{(2)}(\partial_x)|x-x'|^{2\epsilon_2}.$$
(3.17)

For $x'_0 = x_0$ and using again eq. (3.16) with $s = -\epsilon_2$, but now in D-1 dimensions,

$$G_{ijkl}^{(2)}(\boldsymbol{x}, \boldsymbol{x}') = -\frac{C\Lambda_*^{2\epsilon_2}}{B_4(-2\epsilon_2)} P_{ijkl}^{(2)}(\partial_x) |\boldsymbol{x} - \boldsymbol{x}'|^{2\epsilon_2} = -\frac{C\Lambda_*^{2\epsilon_2}B_3(-2\epsilon_2)}{B_4(-2\epsilon_2)} P_{ijkl}^{(2)}(\partial_x) \int \frac{d^3\boldsymbol{k}}{(2\pi)^3} \frac{e^{i\boldsymbol{k}\cdot(\boldsymbol{x}-\boldsymbol{x}')}}{(\boldsymbol{k}^2)^{\epsilon_2+\frac{3}{2}}}.$$

From the second formula in eq. (3.16), we can compute the ratio $B_3(-2\epsilon_2)/B_4(-2\epsilon_2)$:

$$\frac{B_{D-1}(2s)}{B_D(2s)} = \frac{1}{2\sqrt{\pi}} \frac{\Gamma\left(\frac{D-1}{2}\right)}{\Gamma\left(\frac{D}{2}\right)} \stackrel{D=4}{=} \frac{1}{4}.$$
(3.18)

Finally, according to the definitions in Appendix B,

$$G_{ijkl}^{(2)}(r) = -\frac{C\Lambda_{*}^{2\epsilon_{2}}}{4}P_{ijkl}^{(2)}(\partial_{x})\int \frac{d^{3}\mathbf{k}}{(2\pi)^{3}}\frac{e^{i\mathbf{k}\cdot\mathbf{r}}}{k^{2\epsilon_{2}+3}}$$

$$\stackrel{(\mathbf{B}.1)}{=} -P_{ijkl}^{(2)}(\partial_{x})\int_{0}^{+\infty} dk \,\frac{k^{2}}{2\pi^{2}} P_{h}(k) \,\frac{\sin(kr)}{kr}$$

$$\stackrel{(\mathbf{B}.3)}{=} -P_{ijkl}^{(2)}(\partial_{x})\int_{0}^{+\infty} \frac{dk}{k} \,\Delta_{h}^{2}(k) \,\frac{\sin(kr)}{kr} , \qquad (3.19)$$

where $\boldsymbol{r} = \boldsymbol{x} - \boldsymbol{x}', \ r = |\boldsymbol{r}|, \ k = |\boldsymbol{k}|, \ \text{and}$

$$P_h(k) = \frac{C\Lambda_*^{2\epsilon_2}}{4k^{2\epsilon_2+3}}, \qquad \Delta_h^2(k) = \frac{C}{8\pi^2} \left(\frac{\Lambda_*}{k}\right)^{2\epsilon_2}.$$
(3.20)

Taking into account the two polarization modes of the graviton, the final result for the tensor power spectrum \mathcal{P}_t is

$$\mathcal{P}_{t}(k) \coloneqq 2\Delta_{h}^{2}(k) = \frac{4\Lambda_{*}^{2}}{b\left(2\pi\right)^{2} e^{\tilde{\gamma}_{E}} M_{Pl}^{2}} \left(\frac{k}{\Lambda_{*}}\right)^{-2\epsilon_{2}}.$$
(3.21)

Therefore, using the definition (A.20), the tensor spectrum reads

$$\mathcal{P}_{\rm t}(k) = \frac{n_{\rm t}}{(2\pi)^2 \,\beta_{\rm Ric}} \,(\ell_* k)^{n_{\rm t}} \,\,, \tag{3.22}$$

where the tensor spectral index is

$$n_{\rm t} \coloneqq \frac{d\ln \mathcal{P}_{\rm t}}{d\ln k} = -2\epsilon_2 = \frac{4\beta_{\rm Ric}}{b\,e^{\tilde{\gamma}_{\rm E}}}\frac{\Lambda_*^2}{M_{\rm Pl}^2}\,,\tag{3.23}$$

which is positive because $\beta_{\text{Ric}} > 0$.

3.3 Scalar perturbations

Scalar perturbations are thermal fluctuations of the radiation filling the Universe. They are not induced by the quantum fluctuations of a scalar field as in inflation. In order to convert the power spectrum of primordial density perturbations to the spectrum of fluctuations in the CMB at large angular separations, we follow the standard treatment of the Sachs–Wolfe effect relating temperature deviations $\delta T/T$ to the gravitational potential Φ at the last-scattering surface [55, 56]:

$$\frac{\delta T(\boldsymbol{x})}{T} = -\frac{1+3w}{3(1+w)} \Phi(\boldsymbol{x}), \qquad (3.24)$$

where w is the constant barotropic index defining the equation of state $p = w\rho$ of the perfect fluid assumed to dominate, where p is the pressure and ρ is the energy density. For radiation, w = 1/3 and the front coefficient in eq. (3.24) is -1/2. At the last-scattering surface, the theory (2.1) is well approximated by Einstein gravity. Therefore, the potential is related to density perturbations by the standard Poisson equation

$$\nabla^2 \Phi(\boldsymbol{x}) = \frac{1}{2M_{\rm Pl}^2} \delta \rho(\boldsymbol{x}) \,. \tag{3.25}$$

Regarding the potential, at the last-scattering surface the leading contribution to Φ comes from the Einstein–Hilbert action because, at that time, the energy scale is much smaller than the Planck energy [17]. Moreover, we remind the reader that the FLRW solutions of Einstein's equations are exact solutions also of the theory (2.1).

Equations (3.24) and (3.25) imply

$$rac{\delta T(m{x})}{T} = -rac{1+3w}{3(1+w)} \, rac{1}{2M_{
m Pl}^2} rac{1}{
abla^2} \delta
ho(m{x}) \, ,$$

so that the two-point function of CMB temperature fluctuations reads

$$\left\langle \frac{\delta T(\boldsymbol{x})}{T} \frac{\delta T(\boldsymbol{y})}{T} \right\rangle = \left[\frac{1+3w}{3(1+w)} \right]^2 \left\langle \Phi(\boldsymbol{x}) \Phi(\boldsymbol{y}) \right\rangle, \qquad (3.26)$$

$$\left\langle \Phi(\boldsymbol{x})\Phi(\boldsymbol{y})\right\rangle = \frac{1}{4M_{\rm Pl}^4} \left\langle \frac{1}{\nabla_{\mathbf{x}}^2} \delta\rho(\boldsymbol{x}) \frac{1}{\nabla_{\mathbf{y}}^2} \delta\rho(\boldsymbol{y}) \right\rangle, \qquad (3.27)$$

where we formally inverted eq. (3.25).

On the other hand, both the density perturbations and their two-point correlation function have to be evaluated at the energy scale $E \simeq \Lambda_*/10$ right after the conformal phase. Therefore, we need to express the perturbations of the energy-momentum tensor in terms of the gravitational perturbations through the linearized equations of motion for the theory (2.1) at the scale (3.2). Afterwards, such density perturbations affect the fluctuations of the gravitational potential, which satisfies the 00 component (3.25) of Einstein's equations at the decoupling time t_{dec} . In turn, the potential affects the temperature of the CMB. Schematically,

$$\delta \rho \Big|_{E \simeq \frac{\Lambda_*}{10}} \underset{(\overline{\mathbf{3.25}})}{\Longrightarrow} \Phi \Big|_{\text{last scattering}} \Longrightarrow \frac{\delta T}{T} \Big|_{\text{last scattering}}.$$

In order to relate the density perturbations to the graviton two-point correlation function, we need the linearized equations of motion for the theory (2.1) including the quantum corrections, i.e.,

$$\left[e^{\mathbf{H}^{(0)}}\right]_{\mu\nu}^{\sigma\tau} \left[\frac{M_{\rm Pl}^2 G_{\sigma\tau}^{(1)} - T_{\sigma\tau}^{(1)}}{2}\right] + E_{\mu\nu}^{Q(1)} = 0, \qquad (3.28)$$

where E^Q are the quantum corrections to the equations of motion coming from the Lagrangian (2.6). The linearized equations are given in Appendix A.3 by eq. (A.35), written in compact notation as (A.36).

Now we are ready to evaluate the two-point correlation function of the energy momentum tensor. The details of this tedious computation are given in Appendix C and the final result is eq. (C.2). However, we are only interested in the two-point correlation function of the energy-density perturbation $\delta \rho \coloneqq T_{00}^{(1)}$, which is given by eq. (C.9). Let us focus on the first contribution proportional to 5/12, later showing that the other term is negligible:

$$\langle \delta \rho(x) \delta \rho(y) \rangle \simeq -\frac{4M_{\rm Pl}^2 \Lambda_*^2}{b \, e^{\tilde{\gamma}_{\rm E}}} \int \frac{d^4 k}{(2\pi)^4} \, e^{ik \cdot (x-y)} \frac{5}{12} \left(\frac{k}{\Lambda_*}\right)^{2\epsilon_2}$$

$$\stackrel{(3.16)}{=} -\frac{5M_{\rm Pl}^2 \Lambda_*^{2(1-\epsilon_2)}}{3b \, e^{\tilde{\gamma}_{\rm E}}} \, \frac{(|x-y|^2)^{-\epsilon_2-2}}{B_4(2\epsilon_2+4)} \,.$$

$$(3.29)$$

On the spatial section $x_0 = y_0$, this becomes

$$\begin{aligned} \langle \delta \rho(\boldsymbol{x}) \delta \rho(\boldsymbol{y}) \rangle &= -\frac{5M_{\rm Pl}^2 \Lambda_*^{2(1-\epsilon_2)}}{3b \, e^{\tilde{\gamma}_{\rm E}}} \, \frac{(|\boldsymbol{x} - \boldsymbol{y}|^2)^{-\epsilon_2 - 2}}{B_4(2\epsilon_2 + 4)} \\ &= -\frac{5M_{\rm Pl}^2 \Lambda_*^{2(1-\epsilon_2)}}{3b \, e^{\tilde{\gamma}_{\rm E}}} \, \frac{B_3(2\epsilon_2 + 4)}{B_4(2\epsilon_2 + 4)} \int \frac{d^3 \boldsymbol{k}}{(2\pi)^3} \, e^{i\boldsymbol{k}\cdot(\boldsymbol{x} - \boldsymbol{y})} (k^2)^{\epsilon_2 + 2 - \frac{3}{2}} \,, \quad (3.30) \end{aligned}$$

where in the last equality we used again eq. (3.16) for D = 3 and $s = \epsilon_2 + 2$. Using the second formula in eq. (3.16), we can compute the ratio

$$\frac{B_3(2\epsilon_2+4)}{B_4(2\epsilon_2+4)} = \frac{\Gamma\left(-\epsilon_2-\frac{1}{2}\right)}{2\sqrt{\pi}\Gamma(-\epsilon_2)} \simeq \epsilon_2, \qquad (3.31)$$

where we assumed $\epsilon_2 \ll 1$. Finally,

$$\langle \delta \rho(\boldsymbol{x}) \delta \rho(\boldsymbol{y}) \rangle \simeq -\frac{5M_{\rm Pl}^2 \Lambda_*^{2(1-\epsilon_2)}}{3b \, e^{\tilde{\gamma}_{\rm E}}} \, \epsilon_2 \int \frac{d^3 \boldsymbol{k}}{(2\pi)^3} \, e^{i \boldsymbol{k} \cdot (\boldsymbol{x}-\boldsymbol{y})} k^{2\epsilon_2+1}. \tag{3.32}$$

At this point, we can compute eq. (3.27) employing the definitions of Appendix B. In particular, we apply eq. (B.1) to $\psi(\mathbf{x}) = \Phi(\mathbf{x})$:

$$\langle \Phi(\boldsymbol{x})\Phi(\boldsymbol{y})\rangle = \frac{1}{4M_{\rm Pl}^4} \left[-\frac{5M_{\rm Pl}^2\Lambda_*^{2(1-\epsilon_2)}}{3b\,\epsilon^{\tilde{\gamma}_{\rm E}}} \right] \epsilon_2 \int \frac{d^3\boldsymbol{k}}{(2\pi)^3} e^{i\boldsymbol{k}\cdot(\boldsymbol{x}-\boldsymbol{y})} \frac{1}{k^4} k^{2\epsilon_2+1}$$

$$= -\frac{5\Lambda_*^{2(1-\epsilon_2)}}{12b\,\epsilon^{\tilde{\gamma}_{\rm E}}M_{\rm Pl}^2} \frac{\epsilon_2}{2\pi^2} \int_0^{+\infty} \frac{dk}{k} \frac{\sin(kr)}{kr} k^{2\epsilon_2}$$

$$= \int_0^{+\infty} \frac{dk}{k} \Delta_{\Phi}^2(k) \frac{\sin(kr)}{kr} ,$$

$$(3.33)$$

where, according to eq. (B.2), we introduced the power spectrum

$$\Delta_{\Phi}^{2}(k) = -\frac{5\Lambda_{*}^{2}}{6b \, e^{\tilde{\gamma}_{\rm E}} M_{\rm Pl}^{2}} \frac{\epsilon_{2}}{(2\pi)^{2}} \left(\frac{k}{\Lambda_{*}}\right)^{2\epsilon_{2}} \stackrel{(3.15)}{=} \frac{5}{12} \frac{\epsilon_{2}^{2}}{(2\pi)^{2} \beta_{\rm Ric}} \left(\ell_{*}k\right)^{2\epsilon_{2}}, \qquad (3.34)$$

which is positive since β_{Ric} is positive.

In order to get the full result including the second term in eq. (C.9) proportional to 5/24, we simply have to add to eq. (3.34) a similar contribution obtained replacing 5/12 with

5/24 and ϵ_2 with ϵ_0 . Hence, the total contribution to the power spectrum is

$$\Delta_{\Phi}^{2}(k) = \frac{5}{12} \frac{\epsilon_{2}^{2}}{(2\pi)^{2} \beta_{\text{Ric}}} (\ell_{*}k)^{2\epsilon_{2}} + \frac{5}{24} \frac{\epsilon_{0}^{2}}{(2\pi)^{2} (\beta_{\text{Ric}} + 3\beta_{R})} (\ell_{*}k)^{2\epsilon_{0}}$$

$$= \frac{5}{12} \frac{\epsilon_{2}^{2}}{(2\pi)^{2} \beta_{\text{Ric}}} (\ell_{*}k)^{2\epsilon_{2}} + \frac{5}{24} \frac{\epsilon_{2}^{2} (\beta_{\text{Ric}} + 3\beta_{R})}{(2\pi)^{2} \beta_{\text{Ric}}^{2}} (\ell_{*}k)^{2\epsilon_{0}}$$

$$= \frac{5}{12} \frac{\epsilon_{2}^{2}}{(2\pi)^{2} \beta_{\text{Ric}}} \left[(\ell_{*}k)^{2\epsilon_{2}} + \frac{\beta_{\text{Ric}} + 3\beta_{R}}{2\beta_{\text{Ric}}} (\ell_{*}k)^{2\epsilon_{0}} \right].$$
(3.35)

However, according to eq. (A.24), the coefficient of the second term is $\ll 1$. Therefore, eq. (3.34) is a good approximation of the scalar power spectrum.

The primordial scalar spectrum $\mathcal{P}_{\rm s}$ is defined by the spectrum of the gauge-invariant curvature perturbation on uniform density slices ζ , $\mathcal{P}_{\rm s}(k) \coloneqq k^3 P_{\zeta}(k)/(2\pi^2)$. In the presence of one barotropic fluid with constant w and ignoring the anisotropic stress, one can show that the relation between the potential Φ and ζ is (see, e.g., [57, Chap. 8])

$$\Phi \simeq -\frac{3+3w}{5+3w}\,\zeta\,.\tag{3.36}$$

In particular, for radiation $\Phi = -(2/3)\zeta$, so that the final form for the scalar power spectrum is

$$\mathcal{P}_{\rm s}(k) = \frac{9}{4} \Delta_{\Phi}^2(k) = \frac{15}{64} \frac{(n_{\rm s} - 1)^2}{(2\pi)^2 \beta_{\rm Ric}} \left(\ell_* k\right)^{n_{\rm s} - 1},\tag{3.37}$$

since the scalar spectral index is

$$n_{\rm s} - 1 \coloneqq \frac{d\ln \mathcal{P}_{\rm s}}{d\ln k} = 2\epsilon_2 = -\frac{4\beta_{\rm Ric}\Lambda_*^2}{b\,e^{\tilde{\gamma}_{\rm E}}M_{\rm Pl}^2},\tag{3.38}$$

which is negative definite. Here we used eq. (3.15).

Comparing eqs. (3.23) and (3.38), we get the prediction

$$n_{\rm t} = 1 - n_{\rm s} > 0$$
. (3.39)

This relation between the tensor and the scalar index is typical of noninflationary scenarios where scalar perturbations are sustained by thermal fluctuations. Examples are string-gas cosmology $[58-60]^2$ and the new ekpyrotic model [63-65].

3.4 Parameter space

Let us now discuss the parameter space of the theory compatible with data.

The energy scale $\Lambda_* = \ell_*^{-1}$ is free and the only constraints are that it be larger than $\sim 10 \text{ TeV}$ (LHC center-of-mass scale) and smaller than or equal to the Planck mass $M_{\rm Pl}$. Its actual value does not change the main prediction of the theory (section 3.5) and here we can marginalized with $\Lambda_* = M_{\rm Pl}$ to have the nonlocality scale as high as the Planck scale.

 $^{^{2}}$ A noninflationary model coming from matrix theory may realize string-gas cosmology from first principles [61, 62]. Its cosmological spectra are under construction.

This is a sort of "worst-case" scenario from the phenomenological point of view, where one would naively expect that the higher the fundamental energy, the more difficult to extract observable effects.

According to the PLANCK Legacy Release [66], at the CMB pivot scale $k_0 = 0.05 \text{ Mpc}^{-1}$ and assuming no running, the scalar index is $n_s = 0.9649 \pm 0.0042$ at 68% confidence level, assuming $dn_s/d\ln k = 0$. Plugging the observed value of n_s into eqs. (3.22) and (3.37) and taking $\ell_* = \ell_{\text{Pl}}$, we find

$$\mathcal{P}_{\rm t}(k_0) \simeq \frac{8.266 \times 10^{-6}}{\beta_{\rm Ric}}, \qquad \mathcal{P}_{\rm s}(k_0) \simeq \frac{7.867 \times 10^{-4}}{\beta_{\rm Ric}}.$$
 (3.40)

Therefore, in order to get the observed amplitude $\mathcal{P}_{\rm s}(k_0) \approx 2.2 \times 10^{-9}$, from eq. (3.40) it follows that $\beta_{\rm Ric} \approx 3.6 \times 10^5$. Such a large value of this coefficient can be justified by noting that it depends on the number of fundamental particles in the matter sector, in which case one might consider to populate the theory with superfields. Since $\beta_{\rm Ric} \propto N_{\rm fields}$ (Appendix A.2 and [67, 68]), where $N_{\rm fields}$ is the total number of fields in the theory, the latter requires a large number of new fundamental particles. This assumption may be consistent with a particle-physics solution of the problem of dark matter, interpreted as the contribution of exotic particles, while at the same time it could also populate the "desert" lying between the mass of the Higgs boson and the Planck scale.

The theory does not predict the value of the free mass scale $M_{\rm Pl}/\sqrt{b}$ that appears in $n_{\rm t}$, but we can argue that it takes an intriguing value. From eqs. (3.23) and (3.39) with $\Lambda_* = M_{\rm Pl}$,

$$b = \frac{4\beta_{\rm Ric}}{(1-n_{\rm s})\,e^{\tilde{\gamma}_{\rm E}}}\,.\tag{3.41}$$

For $\beta_{\rm Ric} \approx 3.6 \times 10^5$ and $\tilde{\gamma}_{\rm E} = \gamma_{\rm E}$, we get $b \approx 2.3 \times 10^7$ and an effective intermediate mass scale $M_{\rm Pl}/\sqrt{b} = 5 \times 10^{14} \,\text{GeV}$, which looks like low-end grand-unification scale and is about one order of magnitude larger than the Starobinsky mass for the curvaton. Grand unification scenarios are much easier to realize in the presence of supersymmetry, which would naturally generate a large $\beta_{\rm Ric}$.

The scenario just described is not the only possibility to explain the observed value of the scalar spectral amplitude. Remaining within the non-supersymmetric Standard Model of particle physics, another mechanism could be rooted in the choice of a different background in the post-Planckian phase, in particular, the interpolating metric [16]

$$g'_{\mu\nu} = -\left(A + B \tanh\frac{\tau}{\tau_{\rm Pl}}\right)\eta_{\mu\nu}, \qquad \lim_{\tau \to \pm\infty} a^2 = A \pm B, \qquad A \ge B, \qquad (3.42)$$

where $\tau = \int dt/a(t)$ is conformal time. When performing a Weyl rescaling $\eta_{\mu\nu} = \Omega^{-2}g'_{\mu\nu}$ with $\Omega = a(\tau)$ from the Minkowski metric $\eta_{\mu\nu}$ to the spacetime (3.42), the two-point correlation function must be rescaled, too. Since the scalar potential Φ is proportional to the metric perturbation \boldsymbol{h} and $\boldsymbol{h}' = \Omega^2 \boldsymbol{h}$, one has $\langle \Phi(\boldsymbol{x})\Phi(\boldsymbol{y})\rangle' = \Omega^4 \langle \Phi(\boldsymbol{x})\Phi(\boldsymbol{y})\rangle$ and

$$\mathcal{P}_{\rm s}' = \Omega^4 \mathcal{P}_{\rm s} \,. \tag{3.43}$$

The same rescaling is applied to the amplitude $\mathcal{P}_{\rm t}$, but the ratio $r := \mathcal{P}_{\rm t}/\mathcal{P}_{\rm s}$ is conformally invariant. Therefore, evaluating Ω^2 at the Planck conformal time, i.e., for $\tau \gtrsim \tau_{\rm Pl}$, $\Omega^2(\tau_{\rm Pl}) = a^2(\tau_{\rm Pl}) \simeq A + B$, we get an amplitude $\mathcal{P}'_{\rm s}(k_0)$ consistent with observations if, for instance, $\beta_{\rm Ric} = 1$ and $A + B = 1.7 \times 10^{-3}$, or $\beta_{\rm Ric} = 40$ and $A + B = 10^{-2}$. Therefore, in this case the level of fine tuning is very modest.

3.5 Tensor-to-scalar ratio

Using eqs. (3.22) and (3.37), we can compute the tensor-to-scalar ratio at the pivot scale k_0 :

$$r \coloneqq \frac{\mathcal{P}_{\rm t}(k_0)}{\mathcal{P}_{\rm s}(k_0)} = \frac{64}{15(1-n_{\rm s})} \left(\ell_* k_0\right)^{2(1-n_{\rm s})}.$$
(3.44)

Remarkably, this observable only depends on the scale ℓ_* , while the other parameters of the model, the constants β_{Ric} and b, are hidden in the scalar index. This implies that, despite the need of more understanding about the values of these constants, the theory is already falsifiable. In fact, from eq. (3.3) and for $0.1 \leq \ell_*/\ell_{\text{Pl}} \leq 1$, we obtain the very narrow range

$$0.009 \leqslant r \leqslant 0.011$$
, $k_0 = 0.05 \,\mathrm{Mpc}^{-1}$, (3.45)

with the value

$$r_{0.05} = 0.011 \tag{3.46}$$

being the most natural one ($\ell_* = \ell_{\rm Pl}$). Not only is this prediction three times larger than the one of Starobinsky inflation, $r_{\rm Starobisky} = 0.0037$ for $\mathcal{N} = 57$ e-folds [2, 3] or Weyl-squared quantum gravity [69], but it is also within reach of present observations on primordial gravitational waves [70, 71]. BICEP Array, the imminent upgrade of Keck Array and BICEP3, is projected to reach a sensitivity allowing to determine the tensor-to-scalar ratio with an uncertainty $\sigma(r) \sim 0.003$ within four years [70]. Therefore, this instrument will be able to detect any signal with r > 0.009, establish an implication for $r \sim 0.006-0.009$, and rule out any model predicting r > 0.006 in the case of no signal being found. The satellite LiteBIRD will further reduce the uncertainty in case of detection [72]. In contrast, Starobinsky inflation will have to wait LiteBIRD to be verified, which will establish an upper bound of r < 0.002at the 95% confidence level in case of no detection.

3.6 Primordial GW background

Since

$$\epsilon_2 = \frac{n_{\rm s} - 1}{2} \approx -0.01755, \qquad k_0 = 0.05 \,{\rm Mpc}^{-1},$$
(3.47)

the primordial tensor spectrum is blue-tilted with a spectral index

$$n_{\rm t} \approx 0.0351$$
. (3.48)

This can give rise to an observable gravitational-wave background. The theory reduces to Einstein gravity soon after the conformal symmetry breaking and by the time cosmological perturbations re-enter the horizon. Therefore, one can use the standard dimensionless spectral shape

$$\Omega_{\rm GW}(k,\tau_0) \coloneqq \frac{1}{\rho_{\rm crit}} \frac{d\rho_{\rm GW}}{d\ln k} = \frac{k^2}{12a_0^2 H_0^2} \mathcal{P}_{\rm t}(k) \,\mathcal{T}^2(k,\tau_0),\tag{3.49}$$

where $\rho_{\rm crit} = 3M_{\rm Pl}^2 H_0^2$ is the critical energy density, $\rho_{\rm GW}$ is the energy density of gravitational waves (spatial average of the kinetic energy \dot{h}^2 of the perturbation), the subscript 0 denotes quantities evaluated today, τ is conformal time, and \mathcal{T} is a transfer function describing how the primordial spectrum evolved since its generation [73–77]. Its form, valid for any theory where the observed perturbations came from outside the horizon (independently of where they were generate, in- or outside) can be found, e.g., in [11]. This formula applies also to our non-inflationary scenario where the universe has always been radiation dominated and there is no reheating, a situation which can be assimilated to instantaneous reheating at the Planck temperature $T_{\rm reh} = T_{\rm Pl}$. Then, as shown in Fig. 2, the stochastic background will be able to reach the sensitivity curve of the planned DECIGO interferometer [78], thus providing an independent test of the theory.




Figure 2. Gravitational-wave background (3.49) with primordial tensor spectrum (3.22) compared with the sensitivity curves (dashed) of LIGO-Virgo-KAGRA (LVK), SKA, LISA, Einstein Telescope (ET), and DECIGO. Here $\ell_* = \ell_{Pl}$; other choices close to the Planck scale do not change the curve appreciably. The single-frequency CMB constraint has been given thickness to make it more visible in the plot. Also shown are the big-bang nucleosynthesis (BBN) exclusion region and, as a thick red segment, a model of the signal (compatible with a population of supermassive black-hole binaries) recently detected by the international network of Pulsar Timing Arrays [79–83]. Stochastic backgrounds weaker than the observed signal would be drowned by it at those frequencies (shaded red region). However, the high-frequency cutoff of this signal is much smaller than the DECIGO range.

4 Conclusions

In this paper, we have continued the analysis of the cosmological scenario of the early universe arising in nonminimally coupled nonlocal quantum gravity started in [16]. In this model, generated directly in the full theory under a minimal set of approximations, we have

calculated the tensor and scalar primordial spectra, their indices, the tensor-to-scalar ratio, and the gravitational-wave background. These findings rely on the property of finiteness of the theory, that secures Weyl conformal invariance at the classical as well as at the quantum level. Once the conformal symmetry is spontaneously broken, two-point correlation functions get perturbative logarithmic quantum corrections that make primordial spectra deviate from exact scale invariance.

The main theoretical results of the paper are the tensor spectrum \mathcal{P}_{t} (3.22), the tensor index n_{t} (3.23), the scalar spectrum \mathcal{P}_{s} (3.37), the scalar index n_{s} (3.38), the relation (3.39) between the tensor and the scalar index, and the tensor-to-scalar ratio r (3.44). The tensor spectral index n_{t} is positive and equal to $1 - n_{s}$ and the tensor-to-scalar ratio r only depends on n_{s} and the fundamental length scale ℓ_{*} .

The main phenomenological results are the tight range of values (3.45) for r (and its most natural value (3.46)) and the primordial gravitational-wave background shown in Fig. 2. These are unique, falsifiable predictions of the theory.

The only free parameters in the model are the constants β_{Ric} (or, actually, the number of matter fields in the spectrum of the theory) and b (or, equivalently, n_{s} , which is known from observations). However, β_{Ric} can only affect the amplitude of the perturbations, but not n_{t} or r, that are completely fixed in our model regardless of the details of the theory in the UV regime. The whole scenario is summarized in Fig. 1.

If observations failed to detect a large tensor-to-scalar ratio at the level predicted by our model, the latter should be ruled out because there are no really adjustable parameters in eq. (3.44), since the scale ℓ_* cannot vary more than one order of magnitude within the allowed range. Indeed, inequalities (3.3) are not imposed for convenience but are dictated by theoretical arguments during the conformal phase and by the self-consistency of the perturbative treatment of the quantum field theory. As a consequence, the value of r cannot change by more than 10%, as seen in (3.45).

One cannot give up any of the assumptions and approximations without destroying the whole set of predictions. These approximations are only three: quadratic Stelle-like limit, validity of the perturbative expansion, and flat spacetime regime. Any attempt to relax our construction to something else adapting to future observations (for instance, a lower or undetected r) would inevitably lead to a more complicated model with more assumptions, or to the necessity to include an inflaton. This rigidity is due to using perturbation theory in a well-defined way, which has undergone several cross-checks in our treatment. We do not see how one could possibly relax (3.3) and similar constraints while keeping the rest of the construction standing. Giving up the higher-derivative regime would not only make the problem possibly intractable, but it would also break the quasi-scale invariance of the primordial spectra, for which quadratic curvature operators are necessary in the absence of inflatonic fields. Modifying the model for the sake of saving it would possibly be theoretically unacceptable and, even if one could manage to do so, it would jeopardize the status of the full theory. The latter would not be ruled out *per se*, but it would be removed as a parsimonious contender among other top-down approaches having a say on the early universe.

The blue-tilted GW background, not achievable in single-field inflation, can help to discriminate our model from others with a high-r prediction, such as α -attractors and $V \propto \phi^{4/3}$ inflation [2]. In parallel, none of the other quantum-gravity-related models with a similar prediction of the GW background has the same level of r [11]. Therefore, having two independent observations of the GW background at two different frequency ranges is already sufficient to discriminate among models predicting a similar amplitude in either frequency range.

In order to complete the cosmological picture and further test the predictions of our model, one should look at other aspects of the evolution of the early as well as of the late universe. In the first case, at an observational level, scenarios based on conformal invariance are known to be able to generate an observable level of primordial non-Gaussianity [17, 18, 84]. In particular, in [18, 84] it was found that the level of primordial local non-Gaussianity is $|f_{\rm NL}^{\rm local}| = O(1)$, i.e., larger than in standard inflation (where $|f_{\rm NL}^{\rm local}| = O(\epsilon)$ and ϵ is the first slow-roll parameter) but small enough to be still compatible with the current 1σ constraint $f_{\rm NL}^{\rm local} = -0.9 \pm 5.1$ [85]. Since conformal invariance washes away many of the fine details of the underlying model, it is quite possible that also our model will produce a level of non-Gaussianity of such a magnitude. None of the models in [18, 84] predict a high r, so that non-Gaussianity can provide a genuine cross-check of our model against high-r inflationary scenarios. The calculation of the bispectrum is conceptually straightforward but can be quite involved, and we leave it for future work.

At the level of late-time cosmology, in this paper we have said nothing about the cosmological constant problem, which could be another field of application of the symmetries of the theory. Also, any late-time modification of the Λ CDM model should be able to explain the H_0 and the σ_8 tensions and satisfy certain general conditions outlined in [86–88].

The theory of quantum gravity proposed here directly generates a cosmological model with very few free parameters and rigid, clearly falsifiable predictions in the tensor sector. These will be tested by data on primordial gravitational waves, both in the present (~ 0-5 years, BICEP Array) and in the future (≤ 10 years, LiteBIRD; ≥ 20 years, DECIGO). Already before next-generation observatories become operational, gravitational-wave astronomy can arise as one of the most powerful tools to probe gravity at the interface with the quantum.

Acknowledgments

The authors are supported by grant PID2020-118159GB-C41 funded by the Spanish Ministry of Science and Innovation MCIN/AEI/10.13039/501100011033. L.M. was also supported by the Basic Research Program of the Science, Technology, and Innovation Commission of Shenzhen Municipality (grant no. JCYJ20180302174206969).

A Tree-level graviton propagator

In this appendix, we calculate the tree-level graviton propagator for the theory (2.1) in vacuum, after recalling some results in the literature of quantum gravity.

A.1 Tree-level propagator in the theory (2.1)

The graviton propagator for the theory (2.1) has been computed in [25]. Here, we recall the form of the kinetic operator and the gauge-invariant components of the propagator. In D dimensions, the expansion of the vacuum action at second order in the graviton perturbation

is

$$S^{(2)}[h] = \frac{1}{2} \int d^{D}x \left\{ h^{\alpha\beta} \left(\frac{\delta^{2} S_{\text{loc}}}{\delta h^{\alpha\beta} \delta h^{\gamma\sigma}} \right) \left[e^{\bar{\mathcal{H}}(\Delta)} \right]_{\mu\nu}^{\gamma\sigma} h^{\mu\nu} \right\}$$

$$= \frac{M_{\text{Pl}}^{2}}{8} \int d^{D}x h^{\alpha\beta} \left\{ \left[P^{(2)} - (D-2)P^{(0)} \right] \Box \right\}_{\alpha\beta}^{\gamma\sigma} \left\{ e^{\bar{\mathcal{H}} \left[\frac{P^{(2)} - (D-2)P^{(0)}}{\Lambda_{*}^{2}} \Box \right]} \right\}_{\gamma\sigma,\mu\nu} h^{\mu\nu} (A.1)$$

$$= \frac{M_{\text{Pl}}^{2}}{8} \int d^{D}x h^{\alpha\beta} \left(\left\{ P^{(2)} e^{\bar{\mathcal{H}} \left(\frac{\Box}{\Lambda_{*}^{2}} \right)} - (D-2)P^{(0)} e^{\bar{\mathcal{H}} \left[-\frac{(D-2)\Box}{\Lambda_{*}^{2}} \right]} \right\} \Box \right)_{\alpha\beta,\mu\nu} h^{\mu\nu}$$

$$=: \frac{M_{\text{Pl}}^{2}}{8} \int d^{D}x h^{\alpha\beta} \left\{ \left[P^{(2)} e^{\mathcal{H}_{2}} - (D-2)P^{(0)} e^{\mathcal{H}_{0}} \right] \Box \right\}_{\alpha\beta,\mu\nu} h^{\mu\nu}$$

$$= \frac{1}{2} \int d^{D}x h^{\alpha\beta} \mathcal{O}_{\alpha\beta\mu\nu}^{K} h^{\mu\nu} , \qquad (A.2)$$

where $\overline{H}(z) = H(z) - H(0)$ was defined in eq. (2.3a), the entire function H(z) is defined in eq. (2.3b), in the second and third equalities we wrote explicitly the dimensionless arguments of H, and we have used the projectors

$$P^{(2)}_{\mu\nu\rho\sigma} \coloneqq \frac{1}{2} \left(\Theta_{\mu\rho} \Theta_{\nu\sigma} + \Theta_{\mu\sigma} \Theta_{\nu\rho} \right) - \frac{1}{D-1} \Theta_{\mu\nu} \Theta_{\rho\sigma} , \qquad (A.3a)$$

$$P^{(0)}_{\mu\nu\rho\sigma} \coloneqq \frac{1}{D-1} \Theta_{\mu\nu} \Theta_{\rho\sigma} , \qquad \Theta_{\mu\nu} \coloneqq \eta_{\mu\nu} - \omega_{\mu\nu} .$$
(A.3b)

Therefore, the propagator can be obtained from the propagator of the Einstein–Hilbert theory under the replacements

$$\Box P^{(2)} \rightarrow \Box e^{\overline{\mathrm{H}}\left(\frac{\Box}{\Lambda_{*}^{2}}\right)} P^{(2)} =: \Box e^{\mathrm{H}_{2}} P^{(2)}, \qquad (A.4)$$

$$\Box P^{(0)} \rightarrow \Box e^{\overline{\mathrm{H}} \left[-\frac{(D-2)\Box}{\Lambda_*^2} \right]} P^{(0)} =: \Box e^{\mathrm{H}_0} P^{(0)} .$$
 (A.5)

These rules are useful when computing the propagator for the action (2.1) including the logarithmic one-loop quantum corrections (see eq. (A.10)). Augmenting the kinetic operator \mathcal{O}^{K} with a gauge-fixing term \mathcal{O}_{gf} , from $\mathcal{O} = \mathcal{O}^{K} + \mathcal{O}_{gf}$ we get the tree-level graviton propagator up to gauge-dependent contributions [54, 89]:

$$\mathcal{O}^{-1} = \frac{4}{M_{\rm Pl}^2} \left\{ \frac{P^{(2)}}{\prod_{e} \bar{\mathrm{H}}\left(\frac{\Box}{\Lambda_*^2}\right)} - \frac{P^{(0)}}{\prod_{e} (D-2)e^{\bar{\mathrm{H}}\left[-\frac{(D-2)\Box}{\Lambda_*^2}\right]}} \right\} = \frac{4}{M_{\rm Pl}^2} \left[\frac{P^{(2)}}{\prod_{e} e^{\mathrm{H}_2}} - \frac{P^{(0)}}{(D-2)\Box_{e} e^{\mathrm{H}_0}} \right].$$
(A.6)

According to eq. (2.4), in the UV the propagator becomes polynomial:

$$\mathcal{O}^{-1} \stackrel{\text{\tiny UV}}{\simeq} \frac{4}{M_{\text{Pl}}^2 e^{\tilde{\gamma}_{\text{E}}}} \left\{ \frac{P^{(2)}}{\Box p\left(\frac{\Box}{\Lambda_*^2}\right)} - \frac{P^{(0)}}{(D-2)\Box p\left[-\frac{(D-2)\Box}{\Lambda_*^2}\right]} \right\},\tag{A.7}$$

where $\tilde{\gamma}_{\rm E}$ is defined in eq. (3.15). Writing explicitly the scale in the argument of the polynomial (2.3c),

$$p\left(\frac{\Box}{\Lambda_*^2}\right) = b_0 - b \frac{\Box}{\Lambda_*^2} + \left(\frac{\Box}{\Lambda_*^2}\right)^4, \qquad (A.8)$$

$$p\left[-\frac{(D-2)\Box}{\Lambda_*^2}\right] = b_0 + (D-2)b\frac{\Box}{\Lambda_*^2} + \left[(D-2)\frac{\Box}{\Lambda_*^2}\right]^4.$$
 (A.9)

When $b_0 \neq 0$, this constant can be dropped from the polynomial at the energy scale $E \gtrsim \Lambda_{\rm hd}$ defined in eq. (2.9).

A.2 One-loop graviton propagator

In this section, we calculate the graviton propagator for the quantum effective Lagrangian (2.8). Taking the sum of eq. (A.2) and the quadratic expansion in the graviton $h_{\mu\nu}$ of eq. (2.6), the quantum effective action reads

$$\Gamma^{(2)}[h] = \int d^{D}x \left(\mathcal{L} + \mathcal{L}^{Q}\right) = \frac{1}{2} \int d^{D}x h_{\mu\nu} \left[\mathcal{O}^{K} + \mathcal{O}^{Q}\right]^{\mu\nu\rho\sigma} h_{\rho\sigma}$$

$$= \frac{M_{\rm Pl}^{2}}{8} \int d^{D}x h_{\mu\nu} \Box \left(\left(e^{\rm H_{2}} + \Box\gamma_{2}^{Q}\right) P^{(2)} - \left\{(D-2)e^{\rm H_{0}} - \Box \left[4(D-1)\gamma_{0}^{Q} + D\gamma_{2}^{Q}\right]\right\} P^{(0)}\right)^{\mu\nu\rho\sigma} h_{\rho\sigma}, \qquad (A.10)$$

where γ_2^Q and γ_0^Q are the quantum corrections to the kinetic operator. According to eq. (2.6), the quantum form factors are

$$\gamma_2^Q \coloneqq \frac{2\beta_{\rm Ric}}{M_{\rm Pl}^2} \ln\left(-\frac{\Box}{\Lambda_*^2}\right), \qquad \gamma_0^Q \coloneqq \frac{2\beta_R}{M_{\rm Pl}^2} \ln\left(-\frac{\Box}{\Lambda_*^2\delta_0^2}\right). \tag{A.11}$$

The linearized equations of motion that we get from eq. (A.10) are

$$0 = \Box \left(\left(e^{H_2} + \Box \gamma_2^Q \right) P^{(2)} - \left\{ (D-2)e^{H_0} - \Box \left[4(D-1)\gamma_0^Q + D\gamma_2^Q \right] \right\} P^{(0)} \right)_{\mu\nu}^{\rho\sigma} h_{\rho\sigma}$$

$$= \Box \left\{ e^{H_2}P^{(2)} - (D-2)e^{H_0}P^{(0)} + \Box \gamma_2^Q P^{(2)} + \Box \left[4(D-1)\gamma_0^Q + D\gamma_2^Q \right] P^{(0)} \right\}_{\mu\nu}^{\rho\sigma} h_{\rho\sigma}$$

$$= \Box \left\{ e^{\bar{H}} \left[P^{(2)} - (D-2)P^{(0)} \right] + \Box \gamma_2^Q P^{(2)} + \Box \left[4(D-1)\gamma_0^Q + D\gamma_2^Q \right] P^{(0)} \right\}_{\mu\nu}^{\rho\sigma} h_{\rho\sigma}$$

$$= \Box \left(e^{\bar{H}} \left\{ \left[P^{(2)} - (D-2)P^{(0)} \right] + e^{-H_2} \Box \gamma_2^Q P^{(2)} + e^{-H_0} \Box \left[4(D-1)\gamma_0^Q + D\gamma_2^Q \right] P^{(0)} \right\}_{\mu\nu}^{\rho\sigma} h_{\rho\sigma}, \qquad (A.12)$$

where the operator \overline{H} acting on the first term contains the projectors according to eq. (A.1). In the last two steps of eq. (A.12), we used the completeness relations of the projectors [51–53]:

$$P^{(2)^2} = P^{(2)}, \qquad P^{(0)2} = P^{(0)}, \qquad P^{(2)}P^{(0)} = 0.$$
 (A.13)

From the action (A.10) augmented by the gauge-fixing term, the one-loop one-particleirreducible propagator for the quantum effective Lagrangian (2.8) reads

$$\mathcal{O}^{-1}(k) = -\frac{4}{M_{\rm P1}^2} \left(\frac{\mathbf{P}^{(2)}}{k^2 \left[e^{\mathrm{H}_2} - k^2 \gamma_2^Q(k^2) \right]} - \frac{\mathbf{P}^{(0)}}{k^2 \left\{ (D-2)e^{\mathrm{H}_0} + k^2 \left[4(D-1)\gamma_0^Q(k^2) + D\gamma_2^Q(k^2) \right] \right\}} \right).$$
(A.14)

In the high-energy limit, we can replace the form factors with the polynomial as we have done in eq. (A.7), and, thus, the propagator (A.14) turns into

$$\mathcal{O}^{-1} \stackrel{\text{UV}}{\simeq} -\frac{4}{M_{\text{Pl}}^2} \left(\frac{\mathbf{P}^{(2)}}{k^2 \left[e^{\tilde{\gamma}_{\text{E}}} p\left(-\frac{k^2}{\Lambda_*^2} \right) - k^2 \gamma_2^Q(k^2) \right]} - \frac{\mathbf{P}^{(0)}}{k^2 \left\{ (D-2) e^{\tilde{\gamma}_{\text{E}}} p\left[\frac{(D-2)k^2}{\Lambda_*^2} \right] + k^2 \left[4(D-1)\gamma_0^Q(k^2) + D\gamma_2^Q(k^2) \right] \right\}} \right)$$
(A.15)

Assuming now to be in the energy regime (3.1), which is still well within what we consider the UV regime, the action reduces to Stelle's theory with a certain choice of coefficients [16] and the dominant contribution comes from the first monomial in eqs. (A.8) and (A.9), since at energy scales $E \gtrsim \Lambda_{\rm hd}$ we can forget the constant b_0 in the polynomial:

$$e^{\tilde{\gamma}_{\rm E}} p\left(-\frac{k^2}{\Lambda_*^2}\right) \simeq \frac{e^{\tilde{\gamma}_{\rm E}} b \, k^2}{\Lambda_*^2} \,, \qquad e^{\tilde{\gamma}_{\rm E}} p\left[\frac{(D-2)k^2}{\Lambda_*^2}\right] \simeq -\frac{e^{\tilde{\gamma}_{\rm E}} (D-2) \, b \, k^2}{\Lambda_*^2} \,. \tag{A.16}$$

The propagator turns into:

$$\begin{split} \mathcal{O}^{-1}(k) &\simeq -\frac{4}{M_{\rm Pl}^2} \left(\frac{P^{(2)}}{k^4 \left[\frac{e^{\tilde{\gamma}_{\rm E}b}}{\Lambda_*^2} - \gamma_2^Q(k^2) \right]} \\ &- \frac{P^{(0)}}{k^4 \left\{ -(D-2) \frac{e^{\tilde{\gamma}_{\rm E}}(D-2)b}{\Lambda_*^2} + \left[4(D-1)\gamma_0^Q(k^2) + D\gamma_2^Q(k^2) \right] \right\}} \right) \\ &= -\frac{4}{e^{\tilde{\gamma}_{\rm E}}b} \frac{\Lambda_*^2}{M_{\rm Pl}^2} \left(\frac{P^{(2)}}{k^4 \left[1 - \frac{\Lambda_*^2}{e^{\tilde{\gamma}_{\rm E}b}} \gamma_2^Q(k^2) \right]} \\ &+ \frac{P^{(0)}}{k^4 \left\{ (D-2)^2 - \frac{\Lambda_*^2}{e^{\tilde{\gamma}_{\rm E}b}} \left[4(D-1)\gamma_0^Q(k^2) + D\gamma_2^Q(k^2) \right] \right\}} \right). \end{split}$$

Replacing the quantum corrections (A.11) in the above expression, we get

$$\mathcal{O}^{-1}(k) = -\frac{4}{e^{\tilde{\gamma}_{\rm E}}b} \frac{\Lambda_{*}^{2}}{M_{\rm Pl}^{2}} \left(\frac{\mathbf{P}^{(2)}}{k^{4} \left[1 - \frac{2\Lambda_{*}^{2}}{e^{\tilde{\gamma}_{\rm E}}b} \frac{\beta_{\rm Ric}}{M_{\rm Pl}^{2}} \ln\left(\frac{k^{2}}{\Lambda_{*}^{2}}\right) \right]} + \frac{\mathbf{P}^{(0)}}{k^{4} \left\{ (D-2)^{2} - \frac{\Lambda_{*}^{2}}{e^{\tilde{\gamma}_{\rm E}}b} \left[8(D-1) \frac{\beta_{R}}{M_{\rm Pl}^{2}} \ln\left(\frac{k^{2}}{\Lambda_{*}^{2}\delta_{0}^{2}}\right) + 2D \frac{\beta_{\rm Ric}}{M_{\rm Pl}^{2}} \ln\left(\frac{k^{2}}{\Lambda_{*}^{2}}\right) \right] \right\}} \right)}{e^{-\frac{4}{e^{\tilde{\gamma}_{\rm E}}b}} \frac{\Lambda_{*}^{2}}{M_{\rm Pl}^{2}k^{4}} \left\{ \frac{\mathbf{P}^{(2)}}{1 + \epsilon_{2}\ln\left(\frac{k^{2}}{\Lambda_{*}^{2}}\right)} + \frac{\mathbf{P}^{(0)}}{(D-2)^{2} \left[1 + \epsilon_{0}\ln\left(\frac{k^{2}}{\Lambda_{*}^{2}}\right) + \frac{8(D-1)\Lambda_{*}^{2}\beta_{R}}{(D-2)^{2}e^{\tilde{\gamma}_{\rm E}}bM_{\rm Pl}^{2}} \ln \delta_{0}^{2} \right]} \right\}}{e^{-\frac{4}{e^{\tilde{\gamma}_{\rm E}}b}} \frac{\Lambda_{*}^{2}}{M_{\rm Pl}^{2}k^{4}} \left[\frac{\mathbf{P}^{(2)}}{1 + Q_{2}} + \frac{\mathbf{P}^{(0)}}{(D-2)^{2} \left(1 + Q_{0} + Q_{b} \right)} \right], \qquad (A.17)$$

where we defined

$$Q_2 \coloneqq \epsilon_2 \ln \frac{k^2}{\Lambda_*^2}, \qquad Q_0 \coloneqq \epsilon_0 \ln \frac{k^2}{\Lambda_*^2}, \qquad Q_b \coloneqq \frac{8(D-1)\Lambda_*^2 \beta_R}{(D-2)^2 e^{\tilde{\gamma}_{\rm E}} b M_{\rm Pl}^2} \ln \delta_0^2, \qquad (A.18)$$

$$\epsilon_2 \coloneqq -\frac{2\beta_{\mathrm{Ric}}\Lambda_*^2}{b\,e^{\tilde{\gamma}_{\mathrm{E}}}M_{\mathrm{Pl}}^2}, \qquad \epsilon_0 \coloneqq -\frac{2[D\beta_{\mathrm{Ric}} + 4(D-1)\beta_{\mathrm{R}}]\Lambda_*^2}{(D-2)^2\,b\,e^{\tilde{\gamma}_{\mathrm{E}}}M_{\mathrm{Pl}}^2}.$$
(A.19)

In D = 4 dimensions, eq. (A.19) reads

$$\epsilon_2 = -\frac{2\beta_{\rm Ric}\Lambda_*^2}{b\,e^{\tilde{\gamma}_{\rm E}}M_{\rm Pl}^2}, \qquad \epsilon_0 = -\frac{2(\beta_{\rm Ric}+3\beta_{\rm R})\Lambda_*^2}{b\,e^{\tilde{\gamma}_{\rm E}}M_{\rm Pl}^2}, \qquad \epsilon_0 - \epsilon_2 = -\frac{6\Lambda_*^2}{b\,e^{\tilde{\gamma}_{\rm E}}M_{\rm Pl}^2}\beta_R = 3\epsilon_2\frac{\beta_R}{\beta_{\rm Ric}}.$$
(A.20)

A comparison between the one-loop coefficients (which are not beta functions!) in the nonlocal theory (2.1) and the beta functions $\beta_{2,3}$ of Stelle gravity gives [16]

$$\beta_{\rm Ric} = \frac{\beta_2}{(4\pi)^2}, \qquad \beta_R = \frac{3\beta_3 - 2\beta_2}{6(4\pi)^2}.$$
 (A.21)

Therefore,

$$\beta_{\rm Ric} + 3\beta_R = \frac{3}{2} \frac{\beta_3}{(4\pi)^2} \,, \tag{A.22}$$

and

$$\epsilon_0 = \frac{3\epsilon_2}{2} \frac{\beta_3}{\beta_2} \ll \epsilon_2 , \qquad \epsilon_2 - \epsilon_0 = \epsilon_2 \frac{2\beta_2 - 3\beta_3}{2\beta_2} \simeq \epsilon_2 , \qquad (A.23)$$

where in eq. (A.23) we assumed that $\beta_3 \ll \beta_2$, as is usually the case since β_3 depends on the number of fields N_{fields} , while β_2 does not (see [67, formulæ (9.1) and (9.8)] and [68, formula (4.137)]). From eq. (A.21), this implies

$$\frac{\beta_{\rm Ric} + 3\beta_R}{\beta_{\rm Ric}} \ll 1\,,\tag{A.24}$$

which can happen because $\beta_R < 0$.

In order to manipulate the propagator analytically, at this point we make the following approximations:

$$1 + Q_2 = 1 + \ln\left(\frac{k^2}{\Lambda_*^2}\right)^{\epsilon_2} \simeq \left(\frac{k^2}{\Lambda_*^2}\right)^{\epsilon_2} =: u^{2\epsilon_2}, \qquad (A.25)$$

$$1 + Q_0 = 1 + \ln\left(\frac{k^2}{\Lambda_*^2}\right)^{\epsilon_0} \simeq \left(\frac{k^2}{\Lambda_*^2}\right)^{\epsilon_0} = u^{2\epsilon_0} \,. \tag{A.26}$$

According to eq. (A.23), $\epsilon_0 \ll \epsilon_2$, so that it is sufficient to focus on ϵ_2 , the largest of these two parameters. For $\epsilon_2 = -0.0175$ (observed value), the approximation (A.25) has an accuracy of 10% or better if

$$u^{2\epsilon_2} - (1+Q_2) < 0.01 \implies 0.02 < u < 60.$$
 (A.27)

In the main text, we show that our cosmological model fully lies in this range of values. The other approximation we have to take into account is the self-consistency of the one-loop perturbative computation, namely,

$$\left|\ln u^{2\epsilon_2}\right| \ll 1, \qquad \left|\ln u^{2\epsilon_0}\right| \ll 1.$$
 (A.28)

Again, for $\epsilon_2 = -0.0175$ and a quantum correction smaller than 10% of the classical part, we found

$$\left| \ln u^{2\epsilon_2} \right| < 0.1 \implies 0.06 < u < 17.$$
 (A.29)

This lower bound on u is larger than the one in (A.27) and we can take it as a reference. For simplicity and to be even more conservative, we approximate it to

$$u \gtrsim 0.1$$
, (A.30)

corresponding to an energy scale $E \gtrsim \Lambda_*/10$, which is within the conformally invariant trans-Planckian regime if we identify Λ_* with $M_{\rm Pl}$.

Coming back to the calculation of the propagator, making use of the replacements (A.25) and (A.26) into eq. (A.17) we get

$$\mathcal{O}^{-1}(k) \simeq -\frac{4}{e^{\tilde{\gamma}_{\rm E}}b} \frac{\Lambda_*^2}{M_{\rm Pl}^2 k^4} \left\{ \frac{\mathbf{P}^{(2)}}{\left(\frac{k}{\Lambda_*}\right)^{2\epsilon_2}} + \frac{\mathbf{P}^{(0)}}{(D-2)^2 \left[\left(\frac{k}{\Lambda_*}\right)^{2\epsilon_0} + Q_b\right]} \right\}$$
$$= -\frac{4}{e^{\tilde{\gamma}_{\rm E}}b} \frac{\Lambda_*^{2(1+\epsilon_2)}}{M_{\rm Pl}^2 k^{4+2\epsilon_2}} \left[\mathbf{P}^{(2)} + \frac{\mathbf{P}^{(0)}}{(D-2)^2 \left(\frac{k}{\Lambda_*}\right)^{2(\epsilon_0-\epsilon_2)}} \right]. \tag{A.31}$$

Here, we assumed that $Q_b \ll 1$. Indeed, in D = 4 dimensions

$$Q_b = \frac{6\Lambda_*^2 \beta_R}{e^{\tilde{\gamma}_{\rm E}} b M_{\rm Pl}^2} \ln \delta_0^2 = (\epsilon_2 - \epsilon_0) \ln \delta_0^2, \qquad (A.32)$$

which is much smaller than 1 for $\epsilon_2 - \epsilon_0 \simeq \epsilon_2 \ll 1$ (see eq. (A.23)) or $\delta_0 \sim 1$. This analysis justifies the assumption $Q_b = 0$ in the main text.

A.3 One-loop linearized equations of motion

On the same vein, we can write the equations of motion (A.12) in vacuum as

$$0 = \Box \left(e^{\bar{H}} \left\{ \left[P^{(2)} - (D-2)P^{(0)} \right] + \frac{\frac{2\beta_{\rm Ric}}{M_{\rm Pl}^2} \Box \ln \left(-\frac{\Box}{\Lambda_*^2} \right)}{\exp H_2} P^{(2)} + \frac{\Box \left[8(D-1)\frac{\beta_R}{M_{\rm Pl}^2} \ln \left(-\frac{\Box}{\Lambda_*^2 \delta_0^2} \right) + 2D\frac{\beta_{\rm Ric}}{M_{\rm Pl}^2} \ln \left(-\frac{\Box}{\Lambda_*^2} \right) \right]}{\exp H_0} P^{(0)} \right\} \right)_{\mu\nu}^{\rho\sigma} h_{\rho\sigma} ,$$

where \overline{H} is taken at the zero order in the perturbation and, for $\Lambda_{hd} \leq E \ll \Lambda_*$, we can make the replacements (A.8) and (A.9):

$$\begin{split} 0 &= \Box \left(e^{\bar{\mathrm{H}}} \left\{ \left[P^{(2)} - (D-2)P^{(0)} \right] + \frac{\frac{2\beta_{\mathrm{Ric}}}{M_{\mathrm{Pl}}^2} \Box \ln \left(-\frac{\Box}{\Lambda_*^2} \right)}{e^{\tilde{\gamma}_{\mathrm{E}}} \left(-b\frac{\Box}{\Lambda_*^2} \right)} P^{(2)} \right. \\ &+ \frac{\Box \left[8(D-1)\frac{\beta_R}{M_{\mathrm{Pl}}^2} \ln \left(-\frac{\Box}{\Lambda_*^2 \delta_0^2} \right) + 2D\frac{\beta_{\mathrm{Ric}}}{M_{\mathrm{Pl}}^2} \ln \left(-\frac{\Box}{\Lambda_*^2} \right) \right]}{e^{\tilde{\gamma}_{\mathrm{E}}} b(D-2)\frac{\Box}{\Lambda_*^2}} P^{(0)} \\ &= \Box \left(e^{\bar{\mathrm{H}}} \left\{ \left[P^{(2)} - (D-2)P^{(0)} \right] - \frac{2\beta_{\mathrm{Ric}}\Lambda_*^2 \ln \left(-\frac{\Box}{\Lambda_*^2} \right)}{b \, e^{\tilde{\gamma}_{\mathrm{E}}} M_{\mathrm{Pl}}^2} P^{(2)} \right. \\ &+ \Lambda_*^2 \frac{8(D-1)\beta_R \ln \left(-\frac{\Box}{\Lambda_*^2 \delta_0^2} \right) + 2D\beta_{\mathrm{Ric}} \ln \left(-\frac{\Box}{\Lambda_*^2} \right)}{b \, e^{\tilde{\gamma}_{\mathrm{E}}} (D-2)M_{\mathrm{Pl}}^2} P^{(0)} \\ \end{split} \right\} \right)_{\mu\nu}^{\rho\sigma} h_{\rho\sigma} \, . \end{split}$$

Using again the completeness relations for the projectors (A.3), we can factorize the operator $P^{(2)} - (D-2)P^{(0)}$ and get

$$0 = \Box \left\{ e^{\bar{H}} \left[P^{(2)} - (D-2)P^{(0)} \right] \left[\mathbb{1} - \frac{2\beta_{\rm Ric}\Lambda_*^2 \ln\left(-\frac{\Box}{\Lambda_*^2}\right)}{b \, e^{\tilde{\gamma}_{\rm E}}M_{\rm Pl}^2} P^{(2)} - \Lambda_*^2 \frac{8(D-1)\beta_R \ln\left(-\frac{\Box}{\Lambda_*^2}\delta_0^2\right) + 2D\beta_{\rm Ric} \ln\left(-\frac{\Box}{\Lambda_*^2}\right)}{b \, e^{\tilde{\gamma}_{\rm E}}(D-2)^2 M_{\rm Pl}^2} P^{(0)} \right] \right\}_{\mu\nu}^{\rho\sigma} h_{\rho\sigma} ,$$

where $\mathbb{1}_{\mu\nu,\rho\sigma} := (\eta_{\mu\rho}\eta_{\nu\sigma} + \eta_{\mu\sigma}\eta_{\nu\rho})/2$ is the identity tensor [89]. Finally, for $Q_b \ll 1$ we write the linearized quantum equations of motion as

$$0 = \Box \left\{ e^{\bar{H}} \left[P^{(2)} - (D-2)P^{(0)} \right] \left[\mathbb{1} + \epsilon_2 \ln \left(-\frac{\Box}{\Lambda_*^2} \right) P^{(2)} + \epsilon_0 \ln \left(-\frac{\Box}{\Lambda_*^2} \right) P^{(0)} \right] \right\}_{\mu\nu}^{\rho\sigma} h_{\rho\sigma}$$

= $\Box \left\{ e^{\bar{H}} \left[P^{(2)} - (D-2)P^{(0)} \right] \left[\mathbb{1} + Q_2 P^{(2)} + Q_0 P^{(0)} \right] \right\}_{\mu\nu}^{\rho\sigma} h_{\rho\sigma}.$ (A.33)

In the presence of matter, the equations of motion are (3.28). Expanding them at the linear level around the Minkowski background (indeed, the metric $\eta_{\mu\nu}$ is an exact solutions of

the equations of motion (3.28) also including the quantum corrections E^Q), the contribution of the nonlocal form factor is of zero order in the fluctuations. Hence, it is diagonal and we can write the modified Einstein's equations (3.28) as

$$\left[e^{\mathbf{H}^{(0)}}\right]_{\mu\nu}^{\sigma\tau}G_{\sigma\tau}^{(1)} + \frac{2E_{\mu\nu}^{Q(1)}}{M_{\mathrm{Pl}}^2} = \frac{1}{M_{\mathrm{Pl}}^2}\left[e^{\mathbf{H}^{(0)}}\right]_{\mu\nu}^{\sigma\tau}T_{\sigma\tau}^{(1)},\tag{A.34}$$

where $^{(0)}$ and $^{(1)}$ stand for, respectively, the zero-order and the first-order expansion in the metric or matter perturbations. There is no H⁽¹⁾ contribution, as shown in [37]. The left-hand side of eq. (A.34) at the linear level is given by eq. (A.12). Hence, the equations of motion (A.34) at the linear level read

$$\Box \left(e^{\mathbf{H}^{(0)}} \left\{ \left[P^{(2)} - (D-2)P^{(0)} \right] + \frac{\Box \gamma_2^Q}{e^{\mathbf{H}_2^{(0)}}} P^{(2)} + \frac{\Box \left[4(D-1)\gamma_0^Q + D\gamma_2^Q \right]}{e^{\mathbf{H}_0^{(0)}}} P^{(0)} \right\} \right)_{\mu\nu}^{\sigma\tau} h_{\sigma\tau}$$
$$= \frac{1}{M_{\rm Pl}^2} \left[e^{\mathbf{H}^{(0)}} \right]_{\mu\nu}^{\sigma\tau} T_{\sigma\tau}^{(1)},$$

and, factoring out the exponential form factor $\exp H^{(0)}$ on both sides (which can be done since $\exp \bar{H}$ is entire),

$$\frac{T_{\mu\nu}^{(1)}}{M_{\rm Pl}^2} = \Box \left\{ \left[P^{(2)} - (D-2)P^{(0)} \right] + e^{-H_2^{(0)}} \Box \gamma_2^Q P^{(2)} + e^{-H_0^{(0)}} \Box \left[4(D-1)\gamma_0^Q + D\gamma_2^Q \right] P^{(0)} \right\}_{\mu\nu}^{\sigma\tau} h_{\sigma\tau} ,$$
(A.35)

where one should not confuse the labels $^{(0)}$ and $^{(2)}$ of the projectors with the linearization subscripts. Finally, using the result (A.33) and a compact vectorial notation, we can replace the quantum form factors in eq. (A.35) to end up with

$$\frac{\boldsymbol{T}^{(1)}}{M_{\rm Pl}^2} = \Box \left[\boldsymbol{P}^{(2)} - (D-2)\boldsymbol{P}^{(0)} \right] \left[\mathbf{1} + Q_2 \boldsymbol{P}^{(2)} + Q_0 \boldsymbol{P}^{(0)} \right] \boldsymbol{h} \,. \tag{A.36}$$

As stated in [16], the linearized quantum equations of motion (A.36) reduce to the linearized Einstein's equations in the classical limit $\hbar \to 0$, i.e., for $Q_2 = Q_0 = 0$.

B Spatial correlation function and power spectrum

The spatial correlation function for a generic field $\Psi(x)$ is defined by

$$\begin{split} \langle \Psi(\boldsymbol{x})\Psi(\boldsymbol{y})\rangle &= \langle \Psi(\boldsymbol{x})\Psi(\boldsymbol{x}-\boldsymbol{r})\rangle \\ &=: \int \frac{d^{3}\boldsymbol{k}}{(2\pi)^{3}} P_{\Psi}(k)e^{i\boldsymbol{k}\cdot\boldsymbol{r}} \\ &= \int_{0}^{2\pi} d\phi \int_{0}^{+\infty} \frac{dk \, k^{2}}{(2\pi)^{3}} P_{\Psi}(k) \int_{-1}^{1} du \, e^{ikru} \\ &= \int_{0}^{+\infty} \frac{dk \, k^{2}}{(2\pi)^{2}} P_{\Psi}(k) \int_{-1}^{1} du \, e^{ikru} \\ &= \int_{0}^{+\infty} \frac{dk \, k^{2}}{(2\pi)^{2}} P_{\Psi}(k) \frac{2\sin(kr)}{kr} \\ &= \int_{0}^{+\infty} \frac{dk \, k^{2}}{2\pi^{2}} P_{\Psi}(k) \frac{\sin(kr)}{kr} \,, \end{split}$$
(B.1)

where $\mathbf{r} = \mathbf{x} - \mathbf{y}$, $r = |\mathbf{r}|$, and $k = |\mathbf{k}|$. It is common to define the dimensionless power spectrum as

$$\Delta_{\Psi}^2(k) \coloneqq \frac{k^3}{2\pi^2} P_{\Psi}(k) \,, \tag{B.2}$$

so that eq. (B.1) turns into

$$\langle \Psi(\boldsymbol{x})\Psi(\boldsymbol{y})\rangle = \int_0^{+\infty} \frac{dk}{k} \,\Delta_{\Psi}^2(k) \,\frac{\sin(kr)}{kr} \,. \tag{B.3}$$

C Two-point correlation function of the energy-momentum tensor

In this section, we provide all the details about the computation of the two-point correlation function of the perturbed energy-momentum tensor $T^{(1)}_{\mu\nu}$. For the sake of simplicity, we omit the indices, which are hidden in the boldface notation. Hence, from the linearized equations of motion (A.36) we can get the two-point function

$$\frac{1}{M_{\rm Pl}^{4}} \langle \mathbf{T}^{(1)}(x) \, \mathbf{T}^{(1)}(y) \rangle = \Box_{x} \left[\mathbf{P}_{x}^{(2)} - (D-2) \mathbf{P}_{x}^{(0)} \right] \left[\mathbf{1} + Q_{2} \mathbf{P}_{x}^{(2)} + Q_{0} \mathbf{P}_{x}^{(0)} \right] \\
\times \Box_{y} \left[\mathbf{P}_{y}^{(2)} - (D-2) \mathbf{P}_{y}^{(0)} \right] \left[\mathbf{1} + Q_{2} \mathbf{P}_{y}^{(2)} + Q_{0} \mathbf{P}_{y}^{(0)} \right] \langle \mathbf{h}(x) \mathbf{h}(y) \rangle \\
= \int \frac{d^{D}k}{(2\pi)^{D}} e^{ik \cdot (x-y)} \mathcal{O}^{-1}(k) \, k^{4} \\
\times \left[\mathbf{P}^{(2)} - (D-2) \mathbf{P}^{(0)} \right] \left[\mathbf{1} + Q_{2} \mathbf{P}^{(2)} + Q_{0} \mathbf{P}^{(0)} \right] \\
\times \left[\mathbf{P}^{(2)} - (D-2) \mathbf{P}^{(0)} \right] \left[\mathbf{1} + Q_{2} \mathbf{P}^{(2)} + Q_{0} \mathbf{P}^{(0)} \right] \\
\overset{(\mathbf{A}.\mathbf{17})}{=} \int \frac{d^{D}k}{(2\pi)^{D}} e^{ik \cdot (x-y)} \, k^{4} \left[\mathbf{P}^{(2)} - (D-2) \mathbf{P}^{(0)} \right] \left[\mathbf{1} + Q_{2} \mathbf{P}^{(2)} + Q_{0} \mathbf{P}^{(0)} \right] \\
\times \left[\mathbf{P}^{(2)} - (D-2) \mathbf{P}^{(0)} \right] \left[\mathbf{1} + Q_{2} \mathbf{P}^{(2)} + Q_{0} \mathbf{P}^{(0)} \right] \\
\times \left[\mathbf{P}^{(2)} - (D-2) \mathbf{P}^{(0)} \right] \left[\mathbf{1} + Q_{2} \mathbf{P}^{(2)} + Q_{0} \mathbf{P}^{(0)} \right] \\
\times \left(-\frac{4}{b \, e^{\tilde{\gamma}_{E}}} \frac{\Lambda_{*}^{2}}{M_{\rm Pl}^{2} k^{4}} \right) \left[\frac{\mathbf{P}^{(2)}}{\mathbf{1} + Q_{2}} + \frac{\mathbf{P}^{(0)}}{(D-2)^{2} (\mathbf{1} + Q_{0} + Q_{b})} \right]. \quad (C.1)$$

Making iterated use of the properties (A.13) in the product of the first and third brackets with the last bracket, we can further simplify eq. (C.1),

$$\langle \mathbf{T}^{(1)}(x) \, \mathbf{T}^{(1)}(y) \rangle = -\frac{4M_{\rm Pl}^2 \Lambda_*^2}{b \, e^{\tilde{\gamma}_{\rm E}}} \int \frac{d^D k}{(2\pi)^D} e^{ik \cdot (x-y)} \left[\mathbf{1} + Q_2 \mathbf{P}^{(2)} + Q_0 \mathbf{P}^{(0)} \right] \times \left[\mathbf{1} + Q_2 \mathbf{P}^{(2)} + Q_0 \mathbf{P}^{(0)} \right] \left[\frac{\mathbf{P}^{(2)}}{1+Q_2} + \frac{\mathbf{P}^{(0)}}{1+Q_0} \right] = -\frac{4M_{\rm Pl}^2 \Lambda_*^2}{b \, e^{\tilde{\gamma}_{\rm E}}} \int \frac{d^D k}{(2\pi)^D} e^{ik \cdot (x-y)} \left[\mathbf{1} + (2Q_2 + Q_2^2) \mathbf{P}^{(2)} + (2Q_0 + Q_0^2) \mathbf{P}^{(0)} \right] \times \left[\frac{\mathbf{P}^{(2)}}{1+Q_2} + \frac{\mathbf{P}^{(0)}}{1+Q_0} \right] = -\frac{4M_{\rm Pl}^2 \Lambda_*^2}{b \, e^{\tilde{\gamma}_{\rm E}}} \int \frac{d^D k}{(2\pi)^D} e^{ik \cdot (x-y)} \left[\frac{(1+Q_2)^2 \mathbf{P}^{(2)}}{1+Q_2} + \frac{(1+Q_0)^2 \mathbf{P}^{(0)}}{1+Q_0} \right] = -\frac{4M_{\rm Pl}^2 \Lambda_*^2}{b \, e^{\tilde{\gamma}_{\rm E}}} \int \frac{d^D k}{(2\pi)^D} e^{ik \cdot (x-y)} \left[(1+Q_2) \mathbf{P}^{(2)} + (1+Q_0) \mathbf{P}^{(0)} \right].$$
(C.2)

Let us now display the indices and also introduce the explicit form of the projectors (A.3) in the two-point function (C.2):

$$\langle T_{\mu\nu}^{(1)}(x) T_{\rho\sigma}^{(1)}(y) \rangle = -\frac{4M_{\rm Pl}^2 \Lambda_*^2}{b \, e^{\tilde{\gamma}_{\rm E}}} \int \frac{d^D k}{(2\pi)^D} e^{ik \cdot (x-y)} \left[(1+Q_2) P_{\mu\nu\rho\sigma}^{(2)} + (1+Q_0) P_{\mu\nu\rho\sigma}^{(0)} \right]$$

$$\stackrel{(\mathbf{A}.3)}{=} -\frac{4M_{\rm Pl}^2 \Lambda_*^2}{b \, e^{\tilde{\gamma}_{\rm E}}} \int \frac{d^D k}{(2\pi)^D} e^{ik \cdot (x-y)}$$

$$\times \left\{ (1+Q_2) \left[\frac{1}{2} \left(\Theta_{\mu\rho} \Theta_{\nu\sigma} + \Theta_{\mu\sigma} \Theta_{\nu\rho} \right) - \frac{1}{D-1} \Theta_{\mu\nu} \Theta_{\rho\sigma} \right]$$

$$+ (1+Q_0) \frac{1}{D-1} \Theta_{\mu\nu} \Theta_{\rho\sigma} \right\}$$

$$= -\frac{4M_{\rm Pl}^2 \Lambda_*^2}{b \, e^{\tilde{\gamma}_{\rm E}}} \int \frac{d^D k}{(2\pi)^D} e^{ik \cdot (x-y)}$$

$$\times \left((1+Q_2) \left\{ \frac{1}{2} \left[\left(\eta_{\mu\rho} - \frac{k_{\mu}k_{\rho}}{k^2} \right) \left(\eta_{\nu\sigma} - \frac{k_{\nu}k_{\sigma}}{k^2} \right) \right]$$

$$+ \left(\eta_{\mu\sigma} - \frac{k_{\mu}k_{\sigma}}{k^2} \right) \left(\eta_{\nu\rho} - \frac{k_{\rho}k_{\sigma}}{k^2} \right) \right\}$$

$$+ (1+Q_0) \frac{1}{D-1} \left(\eta_{\mu\nu} - \frac{k_{\mu}k_{\nu}}{k^2} \right) \left(\eta_{\rho\sigma} - \frac{k_{\rho}k_{\sigma}}{k^2} \right) \right) . \quad (C.3)$$

Expanding the products in the brackets:

$$\langle T_{\mu\nu}^{(1)}(x) T_{\rho\sigma}^{(1)}(y) \rangle = -\frac{4M_{\rm Pl}^2 \Lambda_*^2}{b \, e^{\tilde{\gamma}_{\rm E}}} \int \frac{d^D k}{(2\pi)^D} e^{ik \cdot (x-y)} \times \left\{ (1+Q_2) \left[\frac{1}{2} \left(\eta_{\mu\rho} \eta_{\nu\sigma} - \eta_{\mu\rho} \frac{k_{\nu} k_{\sigma}}{k^2} - \frac{k_{\mu} k_{\rho}}{k^2} \eta_{\nu\sigma} + \frac{k_{\mu} k_{\rho} k_{\nu} k_{\sigma}}{k^4} \right) \right. \left. + \eta_{\mu\sigma} \eta_{\nu\rho} - \eta_{\mu\sigma} \frac{k_{\nu} k_{\rho}}{k^2} - \frac{k_{\mu} k_{\sigma}}{k^2} \eta_{\nu\rho} + \frac{k_{\mu} k_{\sigma} k_{\nu} k_{\rho}}{k^4} \right) \right] \left. - \frac{1}{D-1} \left(\eta_{\mu\nu} \eta_{\rho\sigma} - \eta_{\mu\nu} \frac{k_{\rho} k_{\sigma}}{k^2} - \frac{k_{\mu} k_{\nu}}{k^2} \eta_{\rho\sigma} + \frac{k_{\mu} k_{\nu} k_{\rho} k_{\sigma}}{k^4} \right) \right] \left. + (1+Q_0) \frac{1}{D-1} \left(\eta_{\mu\nu} \eta_{\rho\sigma} - \eta_{\mu\nu} \frac{k_{\rho} k_{\sigma}}{k^2} - \frac{k_{\mu} k_{\nu}}{k^2} \eta_{\rho\sigma} + \frac{k_{\mu} k_{\nu} k_{\rho} k_{\sigma}}{k^4} \right) \right\}.$$
(C.4)

According to Lorentz invariance, the following identities hold:

$$\int d^{D}k f(k) k_{\mu} k_{\nu} = \frac{1}{D} \eta_{\mu\nu} \int d^{D}k f(k) k^{2}, \qquad (C.5)$$

$$\int d^{D}k f(k) k_{\mu} k_{\nu} k_{\nu} k_{\nu} = \frac{1}{D} (n_{\nu} n_{\nu} + n_{\nu} n_{\nu} + n_{\nu} n_{\nu}) \int d^{D}k f(k) k^{4} (C.6)$$

$$\int d^D k f(k) k_{\mu} k_{\nu} k_{\rho} k_{\sigma} = \frac{1}{D(D+2)} \left(\eta_{\mu\nu} \eta_{\rho\sigma} + \eta_{\mu\rho} \eta_{\nu\sigma} + \eta_{\mu\sigma} \eta_{\nu\rho} \right) \int d^D k f(k) k^4 , \quad (C.6)$$

which can be easily verified contracting the left-hand sides with the Minkowski metric. Applying these identities to eq. (C.4), we obtain

$$\langle T_{\mu\nu}^{(1)}(x) T_{\rho\sigma}^{(1)}(y) \rangle = -\frac{4M_{\rm Pl}^2 \Lambda_*^2}{b \, e^{\tilde{\gamma}_{\rm E}}} \int \frac{d^D k}{(2\pi)^D} e^{ik \cdot (x-y)} \\ \times \left\{ (1+Q_2) \frac{(D-2)(D+1)}{(D+2)(D-1)} \left[\frac{1}{2} \left(\eta_{\mu\rho} \eta_{\nu\sigma} + \eta_{\mu\sigma} \eta_{\nu\rho} \right) - \frac{1}{D} \eta_{\mu\nu} \eta_{\rho\sigma} \right] \\ + (1+Q_0) \frac{1}{D(D+2)(D-1)} \left[(D^2-3) \eta_{\mu\nu} \eta_{\rho\sigma} + \eta_{\mu\rho} \eta_{\nu\sigma} + \eta_{\mu\sigma} \eta_{\nu\rho} \right] \right\}.$$
(C.7)

Therefore, the two-point correlation function of the perturbation of the energy density in D = 4 dimensions is

$$\begin{aligned} \langle \delta \rho(x) \delta \rho(y) \rangle &= \langle T_{00}^{(1)}(x) T_{00}^{(1)}(y) \rangle \\ &= -\frac{4M_{\rm Pl}^2 \Lambda_*^2}{b \, e^{\tilde{\gamma}_{\rm E}}} \int \frac{d^D k}{(2\pi)^D} \, e^{ik \cdot (x-y)} \frac{D+1}{D(D+2)} \left[(D-2)(1+Q_2) + (1+Q_0) \right] \\ &\sum_{=}^{D=4} -\frac{4M_{\rm Pl}^2 \Lambda_*^2}{b \, e^{\tilde{\gamma}_{\rm E}}} \int \frac{d^4 k}{(2\pi)^4} \, e^{ik \cdot (x-y)} \frac{5}{24} \left[2(1+Q_2) + (1+Q_0) \right]. \end{aligned}$$
(C.8)

Replacing eqs. (A.25) and (A.26) into eq. (C.8), we find

$$\left\langle \delta\rho(x)\delta\rho(y)\right\rangle = -\frac{4M_{\rm Pl}^2\Lambda_*^2}{b\,e^{\tilde{\gamma}_{\rm E}}} \int \frac{d^4k}{(2\pi)^4}\,e^{ik\cdot(x-y)}\frac{5}{24}\left[2\left(\frac{k}{\Lambda_*}\right)^{2\epsilon_2} + \left(\frac{k}{\Lambda_*}\right)^{2\epsilon_0}\right].\tag{C.9}$$

References

- [1] G. Calcagni, *Classical and Quantum Cosmology*, Springer, Switzerland (2017).
- [2] Y. Akrami et al. [PLANCK Collaboration], Planck 2018 results. X. Constraints on inflation, Astron. Astrophys. 641 (2020) A10 [arXiv:1807.06211].
- [3] A.A. Starobinsky, A new type of isotropic cosmological models without singularity, Phys. Lett. B 91 (1980) 99.
- [4] S. Cecotti, Higher derivative supergravity is equivalent to standard supergravity coupled to matter, Phys. Lett. B 190 (1987) 86.
- J. Ellis, D.V. Nanopoulos, and K.A. Olive, No-scale supergravity realization of the Starobinsky model of inflation, Phys. Rev. Lett. 111 (2013) 111301; Phys. Rev. Lett. 111 (2013) 129902
 [arXiv:1305.1247].
- [6] R. Kallosh and A. Linde, Superconformal generalizations of the Starobinsky model, J. Cosmol. Astropart. Phys. 1306 (2013) 028 [arXiv:1306.3214].
- [7] F. Briscese, A. Marcianò, L. Modesto, and E.N. Saridakis, Inflation in (super-)renormalizable gravity, Phys. Rev. D 87 (2013) 083507 [arXiv:1212.3611].
- [8] F. Briscese, L. Modesto, and S. Tsujikawa, Super-renormalizable or finite completion of the Starobinsky theory, Phys. Rev. D 89 (2014) 024029 [arXiv:1308.1413].
- [9] A.S. Koshelev, L. Modesto, L. Rachwał, and A.A. Starobinsky, Occurrence of exact R² inflation in non-local UV-complete gravity, J. High Energy Phys. 1611 (2016) 067 [arXiv:1604.03127].
- [10] A.S. Koshelev, K.S. Kumar, and A.A. Starobinsky, R² inflation to probe non-perturbative quantum gravity, J. High Energy Phys. 1803 (2018) 071 [arXiv:1711.08864].
- [11] G. Calcagni and S. Kuroyanagi, Stochastic gravitational-wave background in quantum gravity, J. Cosmol. Astropart. Phys. 2103 (2021) 019 [arXiv:2012.00170].
- [12] R. Blumenhagen, A. Font, M. Fuchs, D. Herschmann, and E. Plauschinn, Towards axionic Starobinsky-like inflation in string theory, Phys. Lett. B 746 (2015) 217 [arXiv:1503.01607].
- [13] H. Ooguri and C. Vafa, On the geometry of the string landscape and the swampland, Nucl. Phys. B 766 (2007) 21 [arXiv:hep-th/0605264].
- [14] G. Obied, H. Ooguri, L. Spodyneiko, and C. Vafa, de Sitter space and the swampland, arXiv:1806.08362.
- [15] E. Palti, The swampland: introduction and review, Fortsch. Phys. 67 (2019) 1900037 [arXiv:1903.06239].
- [16] L. Modesto and G. Calcagni, Early universe in quantum gravity, arXiv:2206.06384.
- [17] I. Antoniadis, P.O. Mazur, and E. Mottola, Conformal invariance and cosmic background radiation, Phys. Rev. Lett. 79 (1997) 14 [arXiv:astro-ph/9611208].
- [18] I. Antoniadis, P.O. Mazur, and E. Mottola, Conformal invariance, dark energy, and CMB non-Gaussianity, J. Cosmol. Astropart. Phys. 1209 (2012) 024 [arXiv:1103.4164].
- [19] G. Amelino-Camelia, M. Arzano, G. Gubitosi, and J. Magueijo, Dimensional reduction in momentum space and scale-invariant cosmological fluctuations, Phys. Rev. D 88 (2013) 103524 [arXiv:1309.3999].
- [20] G. Amelino-Camelia, M. Arzano, G. Gubitosi, and J. Magueijo, Gravity as the breakdown of conformal invariance, Int. J. Mod. Phys. D 24 (2015) 1543002 [arXiv:1505.04649].
- [21] G. Calcagni, Quantum gravity and gravitational-wave astronomy, in: C. Bambi, S. Katsanevas, and K.D. Kokkotas (Eds.), Handbook of Gravitational Wave Astronomy (Springer, Singapore, 2021) [arXiv:2012.08251].

- [22] A. Addazi et al. Quantum gravity phenomenology at the dawn of the multi-messenger era—A review, Prog. Part. Nucl. Phys. **125** (2022) 103948 [arXiv:2111.05659].
- [23] P. Auclair et al. [LISA Cosmology Working Group], Cosmology with the Laser Interferometer Space Antenna, Living Rev. Relativ. 26 (2023) 5 [arXiv:2204.05434].
- [24] K.G. Arun et al. [LISA Fundamental Physics Working Group], New horizons for fundamental physics with LISA, Living Rev. Relativ. 25 (2022) 4 [arXiv:2205.01597].
- [25] L. Modesto, Nonlocal spacetime-matter, arXiv:2103.04936.
- [26] L. Modesto, The Higgs mechanism in nonlocal field theory, J. High Energy Phys. 2106 (2021) 049 [arXiv:2103.05536].
- [27] G. Calcagni, B.L. Giacchini, L. Modesto, T. de Paula Netto, and L. Rachwał, *Renormalizability of nonlocal quantum gravity coupled to matter*, J. High Energy Phys. 2309 (2023) 034 [arXiv:2306.09416].
- [28] N.V. Krasnikov, Nonlocal gauge theories, Theor. Math. Phys. 73 (1987) 1184 [Teor. Mat. Fiz. 73 (1987) 235].
- [29] Yu.V. Kuz'min, The convergent nonlocal gravitation, Sov. J. Nucl. Phys. 50, 1011 (1989) [Yad. Fiz. 50, 1630 (1989)].
- [30] E.T. Tomboulis, Super-renormalizable gauge and gravitational theories, arXiv:hep-th/9702146.
- [31] L. Modesto, Super-renormalizable quantum gravity, Phys. Rev. D 86 (2012) 044005 [arXiv:1107.2403].
- [32] T. Biswas, E. Gerwick, T. Koivisto, and A. Mazumdar, Towards singularity and ghost free theories of gravity, Phys. Rev. Lett. 108 (2012) 031101 [arXiv:1110.5249].
- [33] L. Modesto and L. Rachwał, Super-renormalizable and finite gravitational theories, Nucl. Phys. B 889 (2014) 228 [arXiv:1407.8036].
- [34] L. Modesto and L. Rachwał, Universally finite gravitational and gauge theories, Nucl. Phys. B 900 (2015) 147 [arXiv:1503.00261].
- [35] P. Donà, S. Giaccari, L. Modesto, L. Rachwał, and Y. Zhu, Scattering amplitudes in super-renormalizable gravity, J. High Energy Phys. 1508 (2015) 038 [arXiv:1506.04589].
- [36] S. Giaccari and L. Modesto, *Causality in nonlocal gravity*, arXiv:1803.08748.
- [37] L. Modesto and G. Calcagni, Tree-level scattering amplitudes in nonlocal field theories, J. High Energy Phys. 2110 (2021) 169 [arXiv:2107.04558].
- [38] G. Calcagni, L. Modesto, and G. Nardelli, Non-perturbative spectrum of non-local gravity, Phys. Lett. B 795 (2019) 391 [arXiv:1803.07848].
- [39] G.V. Efimov, Analytic properties of Euclidean amplitudes, Sov. J. Nucl. Phys. 4, 309 (1967)
 [Yad. Fiz. 4, 432 (1966)].
- [40] R. Pius and A. Sen, Cutkosky rules for superstring field theory,
 J. High Energy Phys. 1610 (2016) 024; Erratum-ibid. 1809 (2018) 122 [arXiv:1604.01783].
- [41] F. Briscese and L. Modesto, Cutkosky rules and perturbative unitarity in Euclidean nonlocal quantum field theories, Phys. Rev. D 99 (2019) 104043 [arXiv:1803.08827].
- [42] K.S. Stelle, Renormalization of higher-derivative quantum gravity, Phys. Rev. D 16 (1977) 953.
- [43] J.F. Liu, F. Briscese, G. Calcagni, L. Qiang, and L. Modesto, Amplitudes and unitarity in nonlocal field theory, In preparation.
- [44] P. Di Francesco, P. Mathieu, and D. Senechal, *Conformal Field Theory*, Springer-Verlag, New York, N.Y. (1997).

- [45] I. Bars, Standard model of particles and forces in the framework of two-time physics, Phys. Rev. D 74 (2006) 085019 [arXiv:hep-th/0606045].
- [46] I. Bars, P. Steinhardt, and N. Turok, Local conformal symmetry in physics and cosmology, Phys. Rev. D 89 (2014) 043515 [arXiv:1307.1848].
- [47] A. Borde, A.H. Guth, and A. Vilenkin, Inflationary spacetimes are incomplete in past directions, Phys. Rev. Lett. 90 (2003) 151301 [arXiv:gr-qc/0110012].
- [48] S.P. Robinson and F. Wilczek, Gravitational correction to running of gauge couplings, Phys. Rev. Lett. 96 (2006) 231601 [arXiv:hep-th/0509050].
- [49] A. Bonanno and M. Reuter, Cosmology of the Planck era from a renormalization group for quantum gravity, Phys. Rev. D 65 (2002) 043508 [arXiv:hep-th/0106133].
- [50] A. Bonanno and M. Reuter, Entropy signature of the running cosmological constant, J. Cosmol. Astropart. Phys. 0708 (2007) 024 [arXiv:0706.0174].
- [51] R.J. Rivers, Lagrangian theory for neutral massive spin-2 fields, Nuovo Cim. 34 (1964) 386.
- [52] K.J. Barnes, Lagrangian theory for the second?rank tensor field, J. Math. Phys. 6 (1965) 788.
- [53] P. Van Nieuwenhuizen, On ghost-free tensor Lagrangians and linearized gravitation, Nucl. Phys. B 60 (1973) 478.
- [54] G. Calcagni and L. Modesto, Path integral and conformal instability in nonlocal quantum gravity, arXiv:2402.14785.
- [55] R.K. Sachs and A.M. Wolfe, Perturbations of a cosmological model and angular variations of the microwave background, Astrophys. J. 147 (1967) 73.
- [56] M.J. White and W. Hu, The Sachs-Wolfe effect, Astron. Astrophys. 321 (1997) 8 [arXiv:astro-ph/9609105].
- [57] D.H. Lyth and A.R. Liddle, *The Primordial Density Perturbation*, Cambridge University Press, Cambridge, U.K. (2009).
- [58] R. Brandenberger and C. Vafa, Superstrings in the early universe, Nucl. Phys. B 316 (1989) 391.
- [59] R.H. Brandenberger, String gas cosmology after Planck, Classical Quantum Gravity 32 (2015) 234002 [arXiv:1505.02381].
- [60] H. Bernardo, R. Brandenberger, and G. Franzmann, Solution of the size and horizon problems from classical string geometry, J. High Energy Phys. 2010 (2020) 155 [arXiv:2007.14096].
- [61] S. Brahma, R. Brandenberger, and S. Laliberte, Emergent cosmology from matrix theory, J. High Energy Phys. 2203 (2022) 067 [arXiv:2107.11512].
- [62] S. Brahma, R. Brandenberger, and S. Laliberte, Emergent early universe cosmology from BFSS matrix theory, arXiv:2205.06016.
- [63] R. Brandenberger and Z. Wang, Nonsingular ekpyrotic cosmology with a nearly scale-invariant spectrum of cosmological perturbations and gravitational waves, *Phys. Rev.* D 101 (2020) 063522 [arXiv:2001.00638].
- [64] R. Brandenberger and Z. Wang, Ekpyrotic cosmology with a zero-shear S-brane, Phys. Rev. D 102 (2020) 023516 [arXiv:2004.06437].
- [65] R. Brandenberger, K. Dasgupta, and Z. Wang, *Reheating after S-brane ekpyrosis*, *Phys. Rev.* D 102 (2020) 063514 [arXiv:2007.01203].
- [66] N. Aghanim et al. [PLANCK Collaboration], Planck 2018 results. VI. Cosmological parameters, Astron. Astrophys. 641 (2020) A6; Erratum-Ibid. 652 (2021) C4 [arXiv:1807.06209].

- [67] I.L. Buchbinder, S.D. Odintsov, and I.L. Shapiro, Effective Action in Quantum Gravity, IOP, Bristol, U.K. (1992).
- [68] I.G. Avramidi, *Heat Kernel and Quantum Gravity*, Springer, New York, NY (2000).
- [69] D. Anselmi, E. Bianchi, and M. Piva, Predictions of quantum gravity in inflationary cosmology: effects of the Weyl-squared term, J. High Energy Phys. **2007** (2020) 211 [arXiv:2005.10293].
- [70] P.A.R. Ade et al. [BICEP/Keck Collaboration], Improved constraints on primordial gravitational waves using Planck, WMAP, and BICEP/Keck observations through the 2018 observing season, Phys. Rev. Lett. 127 (2021) 151301 [arXiv:2110.00483].
- [71] M. Tristram et al. Improved limits on the tensor-to-scalar ratio using BICEP and Planck, Phys. Rev. D 105 (2022) 083524 [arXiv:2112.07961].
- [72] E. Allys et al. [LiteBIRD Collaboration], Probing cosmic inflation with the LiteBIRD cosmic microwave background polarization survey, Prog. Theor. Exp. Phys. 2022 (2022) ptac150 [arXiv:2202.02773].
- [73] M.S. Turner, M.J. White, and J.E. Lidsey, Tensor perturbations in inflationary models as a probe of cosmology, Phys. Rev. D 48 (1993) 4613 [arXiv:astro-ph/9306029].
- [74] Y. Watanabe and E. Komatsu, Improved calculation of the primordial gravitational wave spectrum in the standard model, Phys. Rev. D 73 (2006) 123515 [arXiv:astro-ph/0604176].
- [75] S. Kuroyanagi, T. Chiba, and N. Sugiyama, Precision calculations of the gravitational wave background spectrum from inflation, Phys. Rev. D 79 (2009) 103501 [arXiv:0804.3249].
- [76] K. Nakayama, S. Saito, Y. Suwa, and J. Yokoyama, Probing reheating temperature of the universe with gravitational wave background, J. Cosmol. Astropart. Phys. 0806 (2008) 020 [arXiv:0804.1827].
- [77] S. Kuroyanagi, T. Takahashi, and S. Yokoyama, Blue-tilted tensor spectrum and thermal history of the universe, J. Cosmol. Astropart. Phys. 1502 (2015) 003 [arXiv:1407.4785].
- [78] S. Kawamura et al., Current status of space gravitational wave antenna DECIGO and B-DECIGO, Prog. Theor. Exp. Phys. 2021 (2021) 05A105 [arXiv:2006.13545].
- [79] G. Agazie et al. [NANOGrav], The NANOGrav 15-year data set: evidence for a gravitational-wave background, Astrophys. J. Lett. 951 (2023) L8 [arXiv:2306.16213].
- [80] J. Antoniadis et al., The second data release from the European Pulsar Timing Array III. Search for gravitational wave signals, Astron. Astrophys. 678 (2023) A50 [arXiv:2306.16214].
- [81] D.J. Reardon et al. [PPTA], Search for an isotropic gravitational-wave background with the Parkes Pulsar Timing Array, Astrophys. J. Lett. **951** (2023) L6 [arXiv:2306.16215].
- [82] H. Xu et al. [CPTA], Searching for the nano-Hertz stochastic gravitational wave background with the Chinese Pulsar Timing Array Data Release I, Res. Astron. Astrophys. 23 (2023) 075024 [arXiv:2306.16216].
- [83] G. Agazie et al. [International Pulsar Timing Array], Comparing recent PTA results on the nanohertz stochastic gravitational wave background, arXiv:2309.00693.
- [84] P. Agrawal, S. Gukov, G. Obied, and C. Vafa, Topological gravity as the early phase of our universe, arXiv:2009.10077.
- [85] Y. Akrami et al. [PLANCK Collaboration], Planck 2018 results. IX. Constraints on primordial non-Gaussianity, Astron. Astrophys. 641 (2020) A9 [arXiv:1905.05697].
- [86] L. Heisenberg, H. Villarrubia-Rojo, and J. Zosso, Can late-time extensions solve the H_0 and σ_8 tensions?, arXiv:2202.01202.
- [87] L. Heisenberg, H. Villarrubia-Rojo, and J. Zosso, Simultaneously solving the H_0 and σ_8 tensions with late dark energy, arXiv:2201.11623.

- [88] B.-H. Lee, W. Lee, E. Ó Colgáin, M.M. Sheikh-Jabbari and S. Thakur, Is local H₀ at odds with dark energy EFT?, J. Cosmol. Astropart. Phys. 2204 (2022) 004 [arXiv:2202.03906].
- [89] A. Accioly, A. Azeredo, and H. Mukai, Propagator, tree-level unitarity and effective nonrelativistic potential for higher-derivative gravity theories in D dimensions, J. Math. Phys. (N.Y.) 43 (2002) 473.