

Random Tensor Networks with Nontrivial Links

Newton Cheng¹, Cécilia Lancien², Geoff Penington^{1,3}, Michael Walter^{4,5,6}, and Freek Witteveen⁴

¹Center for Theoretical Physics and Department of Physics, University of California, Berkeley, USA

²Institut Fourier & CNRS, Université Grenoble Alpes, Gières, France

³Institute for Advanced Study, Princeton, USA

⁴Korteweg-de Vries Institute for Mathematics and QuSoft, University of Amsterdam, The Netherlands

⁵Institute for Theoretical Physics and ILLC, University of Amsterdam, The Netherlands

⁶Faculty of Computer Science, Ruhr University Bochum, Germany

Abstract

Random tensor networks are a powerful toy model for understanding the entanglement structure of holographic quantum gravity. However, unlike holographic quantum gravity, their entanglement spectra are flat. It has therefore been argued that a better model consists of random tensor networks with link states that are not maximally entangled, i.e., have nontrivial spectra. In this work, we initiate a systematic study of the entanglement properties of these networks. We employ tools from free probability, random matrix theory, and one-shot quantum information theory to study random tensor networks with bounded and unbounded variation in link spectra, and in cases where a subsystem has one or multiple minimal cuts. If the link states have bounded spectral variation, the limiting entanglement spectrum of a subsystem with two minimal cuts can be expressed as a free product of the entanglement spectra of each cut, along with a Marchenko-Pastur distribution. For a class of states with unbounded spectral variation, analogous to semiclassical states in quantum gravity, we relate the limiting entanglement spectrum of a subsystem with two minimal cuts to the distribution of the minimal entanglement across the two cuts. In doing so, we draw connections to previous work on split transfer protocols, entanglement negativity in random tensor networks, and Euclidean path integrals in quantum gravity.

Contents

1	Introduction	2
1.1	Notation and conventions	7
2	Random tensor network states	8
2.1	The replica trick for random tensor networks	9
2.2	Maximally entangled link states and minimal cuts	10
2.3	The replica trick for general background states	11
3	Link states with bounded spectral variation	14
3.1	Random matrices, free probability and non-crossing partitions	14
3.2	Entanglement spectrum of random tensor network states as a free product	18
3.3	Nontrivial link states and entanglement negativity	24
4	Link states with unbounded spectral variation	28
4.1	One-shot entropies	28
4.2	Recovery isometries	30
4.3	One minimal cut	35

4.4	Two competing minimal cuts	37
4.5	Computing entropies with two minimal cuts	53
Acknowledgments		59
A Euclidean gravity path integrals and entanglement spectra		60
A.1	The replica trick and Euclidean path integrals	60
A.2	States at a minimal surface phase transition	62
A.3	Replica wormholes and JT gravity	64
B Random tensor network states and split transfer protocols		67
B.1	Split transfer and recovery in holography	68
C Joint smoothing of link states		70
D A family of metrics on the symmetric group		78
References		79

1 Introduction

More than twenty years after its discovery, the AdS/CFT correspondence [Mal99] remains the only known example of a theory of quantum gravity.¹ A crucial feature of this correspondence is that the emergence of a (classical) spacetime is closely related to the entanglement structure of the boundary theory. Tensor networks appear to provide useful toy models for this aspect of AdS/CFT, mirroring many of its expected properties in a setting that can be made completely mathematically rigorous [Swi12b, Swi12a, PYHP15, HNQ⁺16]. A particularly powerful model is given by *random tensor networks*, which have the advantage of being highly analytically tractable, while exhibiting remarkably precise agreement with gravitational calculations (even including certain exponentially small corrections) [YHQ16, HNQ⁺16, QYY17, QY18, PSSY19, DQW21, KFNR21, QSY21]. Random tensors and tensor networks also arise in a number of other fields of physics, including quantum information, where they have been used to explore generic entanglement properties of quantum states [HLW06, CNŽ10, Aub12, ASY12, AN12, CGGPG13, CNŽ13, CDKW14, CN16, AS17, Has17, NW20, WW21, MB21, LPG21, AFLR21] and condensed matter physics, e.g. in the study of random circuits and measurements [YYQ18, VPYL19, LPWV20, NRSR21, MVS21, YLFC21, LC21, LVFL21].


The most basic version of a random tensor network is characterized by a choice of bond dimension D and a graph $G = (V, E)$, where the vertices $V = V_b \sqcup V_\partial$ of G are partitioned into “bulk” vertices V_b and “boundary” vertices V_∂ . To each edge $e \in E$, we associate a maximally entangled state

$$\frac{1}{\sqrt{D}} \sum_{i=1}^D |ii\rangle \tag{1.1}$$


on two D -dimensional Hilbert spaces, one of which is associated to each endpoint of e ; each vertex $v \in V$ is therefore associated with a Hilbert space \mathcal{H}_v of dimension $D^{\deg(v)}$. Finally, we project each *bulk* vertex $v_b \in V_b$ onto a Haar random state $|\psi_{v_b}\rangle \in \mathcal{H}_{v_b}$. The resulting “random tensor network state” lives in the Hilbert space $\mathcal{H}_\partial = \otimes_{v_\partial} \mathcal{H}_{v_\partial}$ associated to the boundary vertices $v_\partial \in V_\partial$, as shown in Figure 1. Such states, obtained by projecting maximally entangled edge states onto (not necessarily random) bulk vertex states, are also known as projected entangled pair states (PEPS) in the condensed matter literature [VC04, VMC08, CPGSV21].

To characterize the typical entanglement structure of random tensor network states, we can compute the von Neumann entropy $H(\rho_A)$ of the reduced density matrix ρ_A on a subset $A \subset V_\partial$ of the boundary vertices. In the limit where the bond dimension D is very large, this entropy can be shown to converge with high

¹Here, we are requiring any putative theory of quantum gravity to (a) be defined nonperturbatively and (b) have strong evidence for the existence of a semiclassical limit consisting of Einstein gravity coupled to quantum field theory.



(a) The skeleton of a random tensor network. We take the green-colored vertices to be the boundary vertices V_∂ and the remaining, orange-colored vertices to be the bulk vertices V_b . We use this network for illustration throughout the remainder of this paper.



(b) A close-up picture of a random tensor network. We first associate each vertex $v \in V$ with a Hilbert space \mathcal{H}_v of dimension $D^{\deg(v)}$ (here, $\deg(v) = 3$). Then maximally entangled states are distributed according to the graph's edges, after which the state at each bulk vertex $v_b \in V_b$ is projected onto a Haar random state $|\psi_{v_b}\rangle \in \mathcal{H}_v$.

Figure 1: The basic structure of a random tensor network.

probability to $\log(D)|\gamma_A|$, where γ_A is the set of edges crossing the minimal cut (for the moment, assumed to be the unique such cut) in the graph separating A from its boundary complement $V_\partial \setminus A$ (see [Figure 2a](#)). This formula is closely analogous to the Ryu-Takayanagi (RT) formula and its generalizations in AdS/CFT, in which entropies are given by the area of minimal surfaces homologous to a subregion of the conformal boundary [[RT06b](#), [RT06a](#), [HRT07](#), [LM13](#), [FLM13](#), [EW15](#)]. Indeed, this connection is one of the primary reasons for studying tensor networks as a toy model of quantum gravity. However, if one goes beyond the von Neumann entropy and studies finer details of the entanglement spectrum of random tensor network states, significant divergences from holography begin to appear, as we will see shortly.

When studying entanglement in either random tensor networks or quantum gravity, or more generally in quantum field theory, it is often convenient to study k -th Rényi entropies $H_k(\rho_A) = (1 - k)^{-1} \log \text{tr}[\rho_A^k]$. For integer $k > 1$, these are more amenable to direct computation than the von Neumann entropy, and one can extract the von Neumann entropy by analytic continuation to $k = 1$. The computation of Rényi entropies in random tensor network models is very similar to holographic computations. In both cases, the idea is to use the *replica trick* – essentially, this is the observation that $\text{tr}[\rho_A^k] = \text{tr}[\tau \rho_A^{\otimes k}]$ where τ is an operator which permutes the k copies of A cyclically. In the holographic computation, this can be written as a path integral, on k copies of the theory, glued together in an appropriate way. By the holographic dictionary, this path integral can then be computed by the action of a bulk geometry with certain boundary conditions [[LM13](#)]. For random tensor networks, one finds that $\text{tr}[\rho_A^k]$ concentrates around its expectation, and can be computed as the partition function of a classical spin model on the bulk vertices, with boundary conditions dictated by the choice of boundary subsystem [[HNQ⁺16](#)]. This computation will be explained in detail in [Section 2.1](#). From these computations, one finds that holographic CFT states and random tensor network states behave quite differently when $k \neq 1$. For random tensor network states, the Rényi entropies are approximately independent of k in the large D limit, meaning their entanglement spectrum is close to “flat;” the boundary state ρ_A is approximately maximally-mixed within a certain subspace. On the other hand, CFT states that are dual to semiclassical spacetime geometries have Rényi entropies that vary non-trivially with k , meaning their entanglement spectrum contains a wide range of eigenvalues that contribute significantly to the state. Recently, it has been argued that the class of “fixed-area states” in AdS/CFT *do* have flat spectra, and more generally have an entanglement structure that closely matches random tensor network states [[AR19](#), [DHM19](#), [BPSW19](#), [MWW20](#), [DQW21](#)]. Fixed-area states have a well-defined semiclassical

geometry associated to a fixed spatial slice; however, thanks to the uncertainty principle, they cannot describe a single semiclassical *spacetime* geometry [BPSW19]. We discuss the connection between these states and random tensor networks in more detail in Appendix A.

In the random tensor network model, the flatness of the spectrum can be traced to the maximally-entangled states used as “link states” (see Eq. (1.1) and Figure 1b) on the edges of the graph, which themselves have flat entanglement spectra. To take results about random tensor networks beyond the fixed-area state regime, it is natural – see, e.g., discussion in [HNQ⁺16, BPSW19] – to replace the maximally entangled link states by general states

$$|\phi_e\rangle = \sum_{i=1}^D \sqrt{\lambda_{e,i}} |ii\rangle. \quad (1.2)$$

The variation in the entanglement spectrum $\lambda_{e,i}$ represents the fluctuations in area in semiclassical gravitational states. We introduce this model in Section 2. The goal of this paper will be to understand the entanglement spectra of random tensor networks with such *nontrivial link states* and spectra in a number of different regimes. In fact, a number of our results apply in an even more general setting, where the product of link states $\otimes_e |\phi_e\rangle$ is replaced by a completely general “background state” density matrix ϕ_V . From a quantum gravity perspective, random tensor networks with general background states are needed to model bulk quantum fields in AdS/CFT, which yield significant physical consequences when the bulk entropies are large [AP20, AP22]. Beyond holography, they also play a central role in the quantum information processing tasks of multipartite state merging *split transfer* [DH10], a connection we elaborate on in Appendix B.

If one considers a random tensor network with non-trivial link states, if there is a *single* minimal cut γ_A for a subsystem A , then the resulting density matrix ρ_A will have an entanglement spectrum that converges to that of $|\gamma_A|$ copies of the link state along the minimal cut as $D \rightarrow \infty$; indeed, this was implicit in [HNQ⁺16]. A more complex question, and the main focus of this work, is the case where there are *two* minimal cuts, as in Figure 2b. This situation is motivated by questions in holography: it can be used to study the phase transition at the point where there are two competing minimal surfaces [MWW20, AP20], which has been relevant to recent advances on the black hole information paradox [Pen20, AEMM19, PSSY19, AHM⁺20, MM20].

For our first main result, in Section 3, we consider a family of link states with increasing bond dimension D as in Eq. (1.2). For each D , the link state has an associated distribution

$$\mu_e^{(D)} = \frac{1}{D} \sum_{i=1}^D \delta_{D\lambda_{e,i}},$$

where $\lambda_{e,i} = \lambda_{e,i}^{(D)}$ also depends on D . We then require that the moments


$$m_k^{(D)} = D^{k-1} \sum_{i=1}^D \lambda_{e,i}^k$$

of the distributions $\mu_e^{(D)}$ converge to a finite limit as $D \rightarrow \infty$ for all positive integer k . We refer to this as the *bounded spectral variation* limit. This means, in particular, that we must have $\lambda_{e,i} = \mathcal{O}(1/D)$ for all but a vanishing fraction of the eigenvalues $\lambda_{e,i}$. If we let γ_A denote a minimal cut for a boundary domain A , then we may similarly define the associated distribution

$$\mu_{\gamma_A}^{(D)} = \frac{1}{D^{|\gamma_A|}} \sum_{\{i_e\}_{e \in \gamma_A}} \delta_{D^{|\gamma_A|} \prod_{e \in \gamma_A} \lambda_{e,i_e}}.$$

By assumption, the moments of the distribution $\mu_{\gamma_A}^{(D)}$ converge to a finite limit (because those of each distribution $\mu_e^{(D)}$ do), implying that $\mu_{\gamma_A}^{(D)}$ converges weakly to some distribution μ_{γ_A} . Now consider the empirical distribution of the spectrum of reduced state ρ_A , which is the (random) distribution

$$\mu_A^{(D)} = \frac{1}{D^{|\gamma_A|}} \sum_{\lambda \in \text{spec}(\rho_A)} \delta_{D^{|\gamma_A|} \lambda}.$$



(a) The boundary domain A has a unique minimal cut Γ_A . The set of edges crossing this minimal cut is denoted by γ_A .

(b) The boundary domain A has two minimal cuts $\Gamma_{A,i}$ with cut-sets $\gamma_{A,i}$. In the notation of Section 2.2, the red region is $\Gamma_{A,1} = V_1$, the blue region is $\Gamma_{A,2} \setminus \Gamma_{A,1} = V_2$ and the green region is $V \setminus \Gamma_{A,2} = V_3$. Note that $\Gamma_{A,1}$ has two connected components, while $\Gamma_{A,2}$ is connected.

Figure 2: Tensor networks with one and two minimal cuts.

In the case where there are two non-intersecting minimal cuts $\gamma_{A,1}$ and $\gamma_{A,2}$, we find that $\mu_A^{(D)}$ converges weakly, in probability, to a limiting distribution μ_A given by a *free product* $\text{MP}(1) \boxtimes \mu_{\gamma_{A,1}} \boxtimes \mu_{\gamma_{A,2}}$, a notion from the theory of free probability. Here, $\text{MP}(1)$ is the Marchenko-Pastur distribution of parameter 1. The situation is summarized by our first main result, which we state more precisely as Theorem 3.4:

Theorem (Informal). *Consider a family of link states in the bounded spectral variation limit. If there is a unique minimal cut γ_A for a boundary subsystem A , then $\mu_A^{(D)}$ converges weakly, in probability, to μ_{γ_A} , while if there are exactly two non-intersecting minimal cuts $\gamma_{A,1}$ and $\gamma_{A,2}$, it converges to $\text{MP}(1) \boxtimes \mu_{\gamma_{A,1}} \boxtimes \mu_{\gamma_{A,2}}$.*

In Section 3.3 we briefly discuss the closely related problem of computing the entanglement negativity spectrum in the same regime.

For our second main result, in Section 4, we investigate a different regime, in which link states are allowed to have *unbounded spectral variation* in the large D limit. This is the more relevant regime for holography, where fluctuations in the area of a surface (in Planck units) grow sublinearly but without bound in the semiclassical limit. When the spectral variation is unbounded, there still exists a reasonable notion of a minimal cut that determines the entanglement spectrum of the boundary state, but the key difference is that minimality must now be defined entropically, rather than geometrically. In fact, the underlying graph essentially plays no role in this regime. A sensible way to formalize this would be to use one-shot entropies: we might say that a cut is “minimal” if the rank of the state along the cut is smaller than the inverse of the largest element in the entanglement spectrum along any other cut. This condition, while intuitive, is a little too restrictive, and one can use smooth conditional entropies to get a weaker, but still meaningful, condition. In Definition 4.6, following [AP20] we introduce the notion of a *unique (ε, K) -minimal cut* Γ_A , where $\Gamma_A \subseteq V$ is a subset of vertices such that the ε -smooth conditional min-entropy of Γ_A compared to competing cuts either contained in Γ_A or containing Γ_A is lower bounded by K . Intuitively, we want K to be as large as possible, and indeed, we show in Theorem 4.7 that the spectrum of a reduced density matrix ρ_A will be close to the spectrum along a unique (ε, K) -minimal cut with an error exponentially small in K .

The situation is more complicated if there are two non-intersecting (ε, K) -minimal cuts, defined in

Definition 4.10 as both cuts satisfying an (ε, K) -minimality property for the same ε and K . Here, we need to impose a regularity condition on the link states, motivated by the example of a link state that is a (large) number of copies of a fixed state:

$$|\phi_\varepsilon\rangle = |\phi_0\rangle^{\otimes n} \quad (1.3)$$

where $|\phi_0\rangle$ is a bipartite state with local dimension d , so the total bond dimension is $D = d^n$. In this case, if ϕ_0 is not maximally entangled, the measure $\mu_e^{(D)}$ will not converge with increasing n as the spectrum is not concentrated around $\frac{1}{D}$. However, using a different measure, the entanglement spectrum of $|\phi_\varepsilon\rangle$ satisfies a central limit theorem. Namely, if $|\phi_\varepsilon\rangle$ has Schmidt coefficients $\{\lambda_i\}$, and $|\phi_0\rangle$ has entanglement entropy H_0 , then the distribution of the random variable $X^{(n)}$ which takes values $\frac{1}{\sqrt{n}}(\log(\frac{1}{\lambda_i}) - nH_0)$ with probability λ_i , converges weakly to a centered Gaussian distribution as $n \rightarrow \infty$. Since we subtracted the entropy nH_0 , the random variable $X^{(n)}$ has expectation zero. Its variance can be thought of as a measure of the fluctuation of $\log(\frac{1}{\lambda_i})$ around the entropy, and is relevant for second-order asymptotic rates in quantum information processing tasks [TH13]. We take this central limit theorem as motivation for a regularity condition on the spectra of general link and background states, and we allow the states to have varying bond dimensions $D(n)$, e.g. $D(n) \sim d^n$. To be more precise, we define the following measure along a cut γ_A :

$$\nu_{\gamma_A}^{(n)} = \sum_{\{i_e\}_{e \in \gamma_A}} \prod_{e \in \gamma_A} \lambda_{e, i_e} \delta_{\frac{1}{\sqrt{n}} \left(\sum_{e \in \gamma_A} \log \frac{1}{\lambda_{e, i_e}} - H(n) \right)},$$

where $H(n)$ is a function of n (which one can think of as being approximately equal to the entanglement entropy along the cut γ_A), and we assume that $\nu_{\gamma_A}^{(n)}$ converges weakly to a *continuous* distribution. Note that the distribution described above reduces to the distribution of $X^{(n)}$ if the link state is of the form in Eq. (1.3), and is very different from the distribution $\mu_A^{(D)}$ we study for link states with bounded spectral variation. Similarly, we let

$$\nu_A^{(n)} = \sum_{\lambda \in \text{spec}(\rho_A)} \lambda \delta_{\frac{1}{\sqrt{n}}(\log(\frac{1}{\lambda}) - H(n))}$$

be the corresponding (random) distribution for the boundary spectrum. Knowledge of this distribution allows computation of the entropy of ρ_A (and fluctuations) as a correction to $H(n)$. In the random tensor network setting, we find that in a situation with two competing minimal cuts, the random tensor network will ‘select’ the minimal parts of each cut, in the following sense:

Theorem (Informal). *Assume that we have a family of (states with) two non-intersecting $(\varepsilon(n), K(n))$ -minimal cuts $\gamma_{A,1}$ and $\gamma_{A,2}$, as defined in Definition 4.10. Suppose the entanglement spectra along the two minimal cuts are such that $\nu_{\gamma_{A,1}}^{(n)}$ and $\nu_{\gamma_{A,2}}^{(n)}$ converge weakly to continuous measures ν_1 and ν_2 respectively as $n \rightarrow \infty$. Then $\nu_A^{(n)}$ converges weakly, in probability, to $\min_*(\nu_1, \nu_2)$ which is the pushforward of ν_1 and ν_2 along the function $\min: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$. In other words, for any bounded continuous function $f \in C_b(\mathbb{R})$*

$$\sum_{\lambda \in \text{spec}(\rho_A)} \lambda f\left(\frac{\log(\frac{1}{\lambda}) - H(n)}{\sqrt{n}}\right) \rightarrow \int \int f(\min(x_1, x_2)) d\nu_1(x_1) d\nu_2(x_2) \quad (1.4)$$

in probability.

We also show in Corollary 4.14 that, up to an error of size $\mathcal{O}(\log(n))$, this allows us to compute the entropy of ρ_A .

An analogous statement to (1.4) was previously conjectured in [MWW20] to be valid in quantum gravity and justified at a physics level of rigour in [AP20]. Our proof of (1.4) is closely related to the arguments in [AP20], but converting the physical arguments into a mathematical proof and carefully controlling the relevant sources of error requires significant technical work, and constitutes the majority of Section 4.

In [Appendix A](#), we relate our results to the study of holographic gravity computations, particularly in situations with competing minimal surfaces. This is not needed to understand the results of this work, but provides additional motivation for the relevance of our results and clarifies the way in which random tensor networks provide a useful toy model for holographic quantum gravity. Then, in [Appendix B](#), we discuss the relation of our results to split transfer and its relevance in holography. This is again not needed to understand the main text, but rather pointing at topics our work is connected to. Next, in [Appendix C](#), we prove two technical lemmas on joint smoothing that are necessary to analyze competing minimal cuts in the unbounded spectral variation regime. Finally, in [Appendix D](#), we prove that a certain function on the symmetric group is a metric; this is not directly used in the current work, but may be of independent interest for the study of random tensor networks.

After the completion of this manuscript, we became aware of independent work by Jinzhao Wang [[Wan22](#)] on the use of free products to describe entanglement in the toy model of quantum gravity introduced in [[PSSY19](#)] that has strong overlap with the ideas in [Section 3.2](#) and [Appendix A](#).

1.1 Notation and conventions

For $k \in \mathbb{N}$, we let $[k] = \{1, \dots, k\}$, and we denote by S_k the group of permutations of this set. We denote by $\|a\|_p$ the ℓ_p -norm of a vector a , defined by $\|a\|_p^p = \sum_i |a_i|^p$. If a and b are vectors of different dimension, we extend the shorter vector by zeros and still write $\|a - b\|_p$ for their distance. For example, if $a \in \mathbb{C}^{d_1}$ and $b \in \mathbb{C}^{d_2}$ with $d_2 > d_1$, we write $\|a - b\|_1 = \sum_{i=1}^{d_1} |a_i - b_i| + \sum_{i=d_1+1}^{d_2} |b_i|$. We also denote by $\|A\|_p$ the Schatten p -norm of an operator A , defined as the ℓ_p -norm of the singular values $\{s_i\}$ of A ; it can also be computed by $\|A\|_p^p = \text{tr}((A^\dagger A)^{p/2})$. The operator norm is given by $\|A\|_\infty = \max\{s_i\}$. If \mathcal{H} is a Hilbert space, we introduce the notation $\mathcal{P}(\mathcal{H})$ for the set of positive semidefinite operators on \mathcal{H} . We often refer to positive semidefinite operators as ‘density operators’ or ‘states’, *without* requiring them to be normalized to unit trace. We write $\mathcal{P}_=(\mathcal{H})$ for the set of $\rho \in \mathcal{P}(\mathcal{H})$ with unit trace, $\text{tr}[\rho] = 1$, and we denote by $\mathcal{P}_\leq(\mathcal{H})$ the set of subnormalized states, that is, $\rho \in \mathcal{P}(\mathcal{H})$ with $\text{tr}[\rho] \leq 1$. We use the convention that for a vector $|\phi\rangle$, we denote the corresponding pure state by ϕ , so $\phi = |\phi\rangle\langle\phi|$. Given a positive semidefinite operator ρ , we denote by $\text{spec}(\rho)$ the vector containing its spectrum in non-increasing order, and we write $\text{spec}_+(\rho)$ for the nonzero part of the spectrum. It is a well-known fact that

$$\|\text{spec}(\rho) - \text{spec}(\sigma)\|_1 = \|\text{spec}_+(\rho) - \text{spec}_+(\sigma)\|_1 \leq \|\rho - \sigma\|_1 \quad (1.5)$$

(in the second expression we use the convention for the distance of vectors of possibly different dimension introduced above). If A is a quantum system with Hilbert space \mathcal{H}_A , we write $\mathcal{P}(A) = \mathcal{P}(\mathcal{H}_A)$, $\mathcal{P}_=(A) = \mathcal{P}_=(\mathcal{H}_A)$, and $\mathcal{P}_\leq(A) = \mathcal{P}_\leq(\mathcal{H}_A)$, and we use subscripts, e.g. $\rho_A \in \mathcal{P}(A)$, to indicate which system and Hilbert space a quantum state is associated with. For a bipartite state $\rho_{AB} \in \mathcal{P}(AB)$ defined on a tensor product of Hilbert spaces $\mathcal{H}_{AB} = \mathcal{H}_A \otimes \mathcal{H}_B$, we obtain the reduced state or reduced density operator $\rho_A \in \mathcal{P}(A)$ by taking the partial trace over the complement: $\rho_A = \text{tr}_B[\rho_{AB}]$, and similarly in multipartite situations. Finally, we adopt the standard notation that if μ_n is some sequence of finite measures, we write $\mu_n \Rightarrow \mu$ if μ_n converges weakly (or in distribution) to a finite measure μ , meaning that for any bounded continuous function $f \in C_b(\mathbb{R})$,

$$\int f(x) d\mu_n(x) \rightarrow \int f(x) d\mu(x). \quad (1.6)$$

If μ_n is a sequence of *random* finite measures on \mathbb{R} , we say that the sequence μ_n *converges weakly, in probability*, to a finite measure μ , if, for any bounded continuous function $f \in C_b(\mathbb{R})$, [Eq. \(1.6\)](#) converges in probability, i.e., if for every $\varepsilon > 0$ we have that

$$\lim_{n \rightarrow \infty} \Pr \left(\left| \int f(x) d\mu_n(x) - \int f(x) d\mu(x) \right| \geq \varepsilon \right) = 0.$$

All logarithms are to base 2.

2 Random tensor network states

We first review the random tensor network model, closely following [HNQ⁺16, DQW21]. Let $G = (V, E)$ be a connected undirected graph, and let $V = V_\partial \sqcup V_b$ be a partition of the vertices into a set of boundary vertices V_∂ and bulk vertices V_b . If $A \subseteq V_\partial$, we write $\bar{A} = V_\partial \setminus A$. We assign a bond dimension D_e to each edge, and we will consider families of states with increasing bond dimensions; for example, we may take $D_e = D$ for all edges and let D increase. For each vertex $x \in V$, let $\partial\{x\}$ denote the set of edges $e = (xy) \in E$ connecting x to some $y \in V$. We define Hilbert spaces $\mathcal{H}_{e,x} = \mathbb{C}^{D_e}$ for $e \in \partial\{x\}$, and $\mathcal{H}_x := \bigotimes_{e \in \partial\{x\}} \mathcal{H}_{e,x}$. We call the pair (e, x) a *half-edge*. Moreover, we write $\mathcal{H}_e = \mathcal{H}_{e,x} \otimes \mathcal{H}_{e,y}$ for an edge $e = (xy) \in E$. Let $D_x = \dim(\mathcal{H}_x)$. For a subset $A \subseteq V$, we write $\mathcal{H}_A = \bigotimes_{x \in A} \mathcal{H}_x$, and similarly, for a subset $S \subseteq E$ we write $\mathcal{H}_S = \bigotimes_{e \in S} \mathcal{H}_e$. Similarly, for a set T of half-edges we write $\mathcal{H}_T = \bigotimes_{(e,x) \in T} \mathcal{H}_{e,x}$. At each edge $e = (xy) \in E$, we place a pure state $|\phi_e\rangle \in \mathcal{P}_=(\mathcal{H}_e)$

$$|\phi_e\rangle = \sum_{i=1}^{D_e} \sqrt{\lambda_{e,i}} |ii\rangle \in \mathcal{H}_e = \mathcal{H}_{e,x} \otimes \mathcal{H}_{e,y}, \quad (2.1)$$

that we call a *link state*. Then, $\phi_{e,x} = \phi_{e,y} = \sum_{i=1}^{D_e} \lambda_{e,i} |i\rangle\langle i|$ is the reduced density matrix of the link state on either of the two subsystems. We refer to the vector $\text{spec}(\phi_{e,x}) = \text{spec}(\phi_{e,y})$, which is ordered in non-increasing fashion, as the *entanglement spectrum* of ϕ_e . Let $\phi \in \mathcal{P}_=(V)$ be the full state on edges given by the tensor product of link states

$$|\phi\rangle = \bigotimes_{e \in E} |\phi_e\rangle. \quad (2.2)$$

At every bulk vertex $x \in V_b$, we place a random vector $|\psi_x\rangle \in \mathcal{H}_x$, where the entries of $|\psi_x\rangle$ are independent standard (circularly-symmetric) complex Gaussian random variables: each entry of the tensor can be written as $\frac{1}{\sqrt{2}}(x + iy)$ where x and y are independent real Gaussian random variables of mean 0 and unit variance. We note that, in the model of [HNQ⁺16], the tensors $|\psi_x\rangle$ were not chosen as random Gaussian vectors, but as uniformly random vectors on the unit sphere. However, for our choice of Gaussian $|\psi_x\rangle$, the norm $\| |\psi_x\rangle \|$ is independent of the normalized vector $|\psi_x\rangle / \| |\psi_x\rangle \|$, and $|\psi_x\rangle / \| |\psi_x\rangle \|$ will be a uniformly random vectors on the unit sphere. Therefore, these two models only differ by their normalization. We write $|\psi\rangle = \bigotimes_{x \in V_b} |\psi_x\rangle$. The resulting *random tensor network state* $\rho \in \mathcal{P}(V_\partial)$ is defined by

$$|\rho\rangle = (I_{V_\partial} \otimes \langle \psi |) |\phi\rangle. \quad (2.3)$$

The random tensor network state is obtained by projecting the link states onto random vectors, so that the final state lives in the boundary Hilbert space. We can make this manifest by using the cyclicity of the trace to write the density matrix:

$$\rho = (I_{V_\partial} \otimes \langle \psi |) \phi (I_{V_\partial} \otimes |\psi\rangle) = \text{tr}_{V_b} [(I_{V_\partial} \otimes \psi) \phi]. \quad (2.4)$$

Note that this state need not be normalized, but we chose the standard deviation of the $|\psi_x\rangle$ such that ρ is normalized on average, given that the link state ϕ is normalized:

$$\mathbb{E} \text{tr}[\rho] = \text{tr}[\phi]. \quad (2.5)$$

In Section 2.3.1, we prove the stronger statement that ρ is normalized with high probability for appropriately connected tensor networks and large bond dimension. Note also, that in Eq. (2.1), we have chosen states which have a Schmidt decomposition in a fixed basis (the standard basis). Since we project onto uniformly random tensors, we can choose to do so without loss of generality.

2.1 The replica trick for random tensor networks

We now consider a boundary subset $A \subseteq V_\partial$ and use the *replica trick* to study the Rényi entropies of the reduced density matrix ρ_A . The replica trick for random tensor network models was first studied in [HNQ+16], and it is the key tool we apply throughout this work. Let \mathcal{H} be a Hilbert space. The Rényi entropies of a (normalized) density matrix $\rho \in \mathcal{P}_=(\mathcal{H})$ are defined by

$$H_k(\rho) = \frac{1}{1-k} \log(\text{tr}[\rho^k])$$

for $k \in (0, 1) \cup (1, \infty)$. For $k = 0, 1, \infty$, there are well-defined limits, given by

$$\begin{aligned} H_0(\rho) &:= \log(\text{rank}(\rho)) \\ H_1(\rho) &:= -\text{tr}[\rho \log(\rho)] \\ H_\infty &:= -\log(\|\rho\|_\infty). \end{aligned} \tag{2.6}$$

In particular, we see that $H(\rho) = H_1(\rho)$ is the von Neumann entropy. We will also write $H(A)_\rho := H(\rho_A)$ and $H_k(A)_\rho := H_k(\rho_A)$ for reduced density matrices. If $\rho \in \mathcal{P}_<(\mathcal{H})$ is subnormalized, we let

$$H_k(\rho) = \frac{1}{1-k} \log \frac{\text{tr}[\rho^k]}{\text{tr}[\rho]}. \tag{2.7}$$

Denote by R the representation of S_k on $\mathcal{H}^{\otimes k}$ which permutes the k copies of \mathcal{H} according to the action of S_k . We will write $R_x(\pi)$ when $\mathcal{H} = \mathcal{H}_x$ and $R_A(\pi)$ if $\mathcal{H} = \mathcal{H}_A$ for $A \subseteq V$. We let τ denote the standard k -cycle in S_k , i.e.,

$$\tau = (12 \dots k).$$

The key idea of the replica trick is the observation that the k -th moment of $\rho \in \mathcal{P}(\mathcal{H})$ can be written as

$$\text{tr}[\rho^k] = \text{tr}[R(\tau)\rho^{\otimes k}]. \tag{2.8}$$

Recall the notion of the cycle type of a permutation π : if π can be written as a product of m disjoint cycles of lengths l_1, \dots, l_m , then π has cycle type $C(\pi) = \{l_1, \dots, l_m\}$. Then, for an arbitrary $\pi \in S_k$,

$$\text{tr}[R(\pi)\rho^{\otimes k}] = \prod_{l \in C(\pi)} \text{tr}[\rho^l].$$

Note that this is the generalization of the well-known *swap trick* for two copies of a state ρ . The other crucial ingredient is a property of the Gaussian random vectors:

$$\mathbb{E}[\psi_x^{\otimes k}] = \sum_{\pi \in S_k} R_x(\pi). \tag{2.9}$$

Using Eq. (2.4), we may then compute

$$\begin{aligned} \mathbb{E} \text{tr}[\rho_A^k] &= \mathbb{E} \text{tr}[R_A(\tau)\rho_A^{\otimes k}] \\ &= \mathbb{E} \text{tr}[(R_A(\tau) \otimes R_{V_\partial \setminus A}(\text{id})) \rho^{\otimes k}] \\ &= \mathbb{E} \text{tr}[(R_A(\tau) \otimes R_{V_\partial \setminus A}(\text{id})) (I_{V_\partial} \otimes \psi)^{\otimes k} \phi^{\otimes k}] \\ &= \text{tr}[(R_A(\tau) \otimes R_{V \setminus A}(\text{id})) \mathbb{E}[(I_{V_\partial} \otimes \psi)^{\otimes k}] \phi^{\otimes k}]. \end{aligned} \tag{2.10}$$

To further simplify this expression, we define the following set:

$$\mathcal{S}_{A,\sigma} = \left\{ \{\pi_x\}_{x \in V} : \pi_x \in S_k, \text{ where } \pi_x = \sigma \text{ for } x \in A \text{ and } \pi_x = \text{id for } x \in \bar{A} \right\}, \tag{2.11}$$

for any $\sigma \in S_k$ and $A \subseteq V_\partial$. An element of $\mathcal{S}_{A,\sigma}$ assigns a permutation to each vertex in V subject to a ‘boundary condition.’ Now, using Eq. (2.9), we find that

$$\mathbb{E} \operatorname{tr}[\rho_A^k] = \sum_{\{\pi_x\} \in \mathcal{S}_{A,\tau}} \operatorname{tr} \left[\bigotimes_{x \in V} R_x(\pi_x) \phi^{\otimes k} \right].$$

Finally, we observe that for $e = (xy)$

$$\operatorname{tr} [R(\pi_x) \otimes R(\pi_y) \phi^{\otimes k}] = \prod_{l \in C(\pi_x^{-1}\pi_y)} \operatorname{tr} [\phi_{e,x}^l],$$

where we recall that $\phi_{e,x}$ is the reduced density matrix of the link state on edge $e = (xy)$. Thus, we conclude that

$$\mathbb{E} \operatorname{tr}[\rho_A^k] = \sum_{\{\pi_x\} \in \mathcal{S}_{A,\tau}} \prod_{e=(xy) \in E} \prod_{l \in C(\pi_x^{-1}\pi_y)} \operatorname{tr} [\phi_{e,x}^l]. \quad (2.12)$$

We can interpret the expectation as the partition function of a classical spin model

$$\mathbb{E} \operatorname{tr}[\rho_A^k] = \sum_{\{\pi_x\} \in \mathcal{S}_{A,\tau}} 2^{-\sum_{e=(xy) \in E} J_e(\pi_x, \pi_y)}, \quad (2.13)$$

where the site variables in the spin model are permutations $\pi_x \in S_k$, the interaction at the edges between sites is given by

$$J_e(\pi_x, \pi_y) = - \sum_{l \in C(\pi_x^{-1}\pi_y)} \log(\operatorname{tr}[\phi_{e,x}^l]) = \sum_{l \in C(\pi_x^{-1}\pi_y)} (l-1) H_l(\phi_{e,x}),$$

with H_l the l -th Rényi entropy, and the model as boundary conditions such that the permutation must be τ on A and id on \bar{A} . It turns out that J_e is a metric on the symmetric group S_k – see Appendix D. Similarly, we may place an arbitrary permutation π on A instead of τ , which yields (by exactly the same reasoning)

$$\mathbb{E} \operatorname{tr} [R(\pi) \rho^{\otimes k}] = \sum_{\{\pi_x\} \in \mathcal{S}_{A,\pi}} \prod_{e=(xy) \in E} \prod_{l \in C(\pi_x^{-1}\pi_y)} \operatorname{tr} [\phi_{e,x}^l]. \quad (2.14)$$

2.2 Maximally entangled link states and minimal cuts

We will now discuss the special case where all the link states are maximally entangled states of dimension D , which has been studied extensively in [HNQ⁺16]. We will generalize the results we discuss here to a wider class of link states in Section 3. In this case, the entanglement spectra of the link states are flat: for $e \in E$, we have $\lambda_{e,i} = \frac{1}{D}$ for $i = 1, \dots, D$. In particular, for all $l \in \mathbb{N}$ we have $H_l(\phi_e) = \log(D)$ and hence

$$J_e(\pi_x, \pi_y) = \log(D) \sum_{l \in C(\pi_x^{-1}\pi_y)} (l-1).$$

This leads to the so-called *Cayley distance* on S_k :

$$d(\pi_x, \pi_y) = \sum_{l \in C(\pi_x^{-1}\pi_y)} (l-1) = k - |C(\pi_x^{-1}\pi_y)|, \quad (2.15)$$

where $|C(\pi)|$ is the number of cycles in π . Moreover, $d(\pi_x, \pi_y)$ is a metric and equals the minimal number of transpositions needed to transform π_x into π_y . We say that $\pi \in S_k$ is on a geodesic between π_1 and π_2

if $d(\pi_1, \pi) + d(\pi, \pi_2) = d(\pi_1, \pi_2)$ (recall that d is a metric). We can rewrite the spin model in terms of this distance:

$$\mathbb{E} \operatorname{tr}[\rho_A^k] = \sum_{\{\pi_x\} \in \mathcal{S}_{A,\sigma}} 2^{-\log(D)} \sum_{e=(xy) \in E} d(\pi_x, \pi_y). \quad (2.16)$$

The physically inclined reader may observe that the logarithm of the bond dimension has the role of an inverse temperature, and for large D , the dominant contribution to the partition function will be the ground state of the spin model, subject to the relevant boundary conditions.

To describe the dominant contribution to the sum in Eq. (2.16) for large D , we need the *minimal cuts* for A in G . A cut for A is a subset of the vertices $\Gamma_A \subset V$ such that $\Gamma_A \cap V_\partial = A$. Throughout this work, we will denote the set of all cuts for A by $C(A)$. We will use the convention of denoting cuts (i.e. subsets of vertices) by capital Greek letters. Given a cut $\Gamma_A \in C(A)$, we will denote the set of edges crossing the cut, that is, edges connecting a vertex in Γ_A with a vertex in $V \setminus \Gamma_A$, by lowercase Greek letters γ_A (and by an abuse of language, also refer to this set as a ‘cut’). A minimal cut for A is a cut such that the number of edges $|\gamma_A|$ is minimal. We write $m(A) = |\gamma_A|$ for a minimal cut $\gamma_A \in C(A)$. If $\Gamma_A \in C(A)$, we write $\Gamma_A^c = V \setminus \Gamma_A$. Note that Γ_A^c is a cut for $\bar{A} = V_\partial \setminus A$.

In the simplest case, there is a *unique* minimal cut γ_A . For this case, one can show that the dominant configuration is the one in which $\pi_x = \tau$ for $x \in \Gamma_A$ and $\pi_x = \text{id}$ for $x \in V \setminus \Gamma_A$, see [HLW06], or Proposition 3.3. That is, there are two domains in the spin model corresponding to τ and id , and the minimization of the domain wall corresponds to the minimal cut in the graph.

We will also be interested in the case of exactly two non-intersecting minimal cuts $\Gamma_{A,1}$ and $\Gamma_{A,2}$. In this case, we have that $\Gamma_{A,1} \subset \Gamma_{A,2}$, or $\Gamma_{A,1} \supset \Gamma_{A,2}$. After relabeling, we may assume that the first is the case, and define three domains in the graph: $V = V_1 \sqcup V_2 \sqcup V_3$ given by $V_1 = \Gamma_{A,1}$, $V_2 = \Gamma_{A,2} \setminus \Gamma_{A,1}$ and $V_3 = V \setminus \Gamma_{A,2}$. If there are exactly two minimal cuts, then multiple dominant configurations contribute equally to the partition function Eq. (2.16). These dominant configurations can be constructed as follows: for each π on a geodesic between τ and id , set $\pi_x = \tau$ for $x \in V_1$, $\pi_x = \pi$ for $x \in V_2$ and $\pi_x = \text{id}$ for $x \in V_3$. That these are the dominant configurations follows immediately from the fact that $d(\tau, \pi) + d(\pi, \text{id}) \geq d(\tau, \text{id})$, with equality if and only if π is on a geodesic between τ and id .

To understand this degeneracy, we use the following fact [NS06]: *the set of permutations π on a geodesic between τ and id is in a one-to-one correspondence with the set of non-crossing partitions $NC(k)$ of $[k]$* . See Section 3.1 for a definition and properties of $NC(k)$. Thus, the degeneracy for the the k -th moment is $|NC(k)| = C_k$ where

$$C_k = \frac{1}{k+1} \binom{k}{2k}$$

is the k -th Catalan number. These are the moments of the *Marchenko-Pastur distribution* $MP(t)$

$$\begin{aligned} MP(t) &= \max(1-t, 0) \delta_0 + \nu_t \\ d\nu_t(x) &= \frac{\sqrt{4t - (x-1-t)^2}}{2\pi x} \mathbb{1}_{(x-1-t)^2 \leq 4t} dx. \end{aligned} \quad (2.17)$$

This allows one to show the folklore result (which we prove and extend to more general link states in Theorem 3.4) that upon an appropriate rescaling, the empirical distribution of the spectrum of ρ_A converges to a Marchenko-Pastur distribution. This is in line with the case of a single random tensor, which precisely yields a Wishart matrix (see Section 3.1.1 for a brief introduction to these objects). In the first case, where there is a unique minimal cut, the entanglement spectrum of ρ_A is flat, while, as we have seen, in the second case, the degeneracy gives rise to a nontrivial spectrum in the right scaling limit.

2.3 The replica trick for general background states

In Eq. (2.13), we computed the result of the replica trick for the k -th moment for a random tensor network state. We will also consider the more general setting where the link state is replaced by some arbitrary




Figure 3: The structure of a (purified) random tensor network with a general background state.

state ϕ_V . In this setting, there need not be a graph structure, and the Hilbert space at each vertex $x \in V$ can be some arbitrary Hilbert space, rather than a tensor product of Hilbert spaces labelled by half-edges. We will refer to ϕ_V as a “background state” instead of a “link state” (as the interpretation of links along the edges does not necessarily make sense in this situation). That is, where before we had a link state

$$|\phi\rangle = \bigotimes_{e \in E} |\phi_e\rangle,$$

we will now consider some *arbitrary* possibly mixed and subnormalized $\phi_V \in \mathcal{P}_{\leq}(V)$ in the tensor network construction. We can generalize Eq. (2.4) to also apply for general background states to obtain a state $\rho \in \mathcal{P}(V_{\partial})$ given by

$$\rho = \text{tr}_{V_b}[(I_{V_{\partial}} \otimes \psi) \phi_V] \quad (2.18)$$

where $|\psi\rangle$ is a tensor product of random states at the bulk vertices. If ϕ is pure, then so is ρ . If ϕ is not pure, we can consider a purification $\phi_{VR} \in \mathcal{P}_{\leq}(VR)$ and consider R as an additional boundary system; this leads to a random tensor network state $\rho_{V_{\partial}R}$ which is a purification of $\rho_{V_{\partial}}$. This set-up is illustrated in Figure 3. While formally very similar, the resulting state is no longer a PEPS tensor network state in general.

There are multiple reasons to also allow general background states. The first reason is of a technical nature: they are useful for estimates based on *smooth entropies*, which we discuss in Section 4. In this application, the full state on edges is still pure, but is no longer a tensor product of link states along the edges. A second motivation for considering general background states is that they can be used as a toy model for holographic systems where there is “bulk entropy” present. Finally, these states are closely related to protocols for the quantum information processing task of split transfer [DH10]. We comment on this connection in Appendix B.

Even for a general background state, a version of the replica trick still applies. Consider a boundary subsystem $A \subseteq V_{\partial}$ with corresponding boundary state ρ_A . Then, the computation in Eq. (2.10) is still valid, and we find

$$\mathbb{E} \text{tr}[\rho_A^k] = \sum_{\{\pi_x\} \in \mathcal{S}_{A,\tau}} \text{tr}_V \left[\bigotimes_{x \in V} R_x(\pi_x) \phi_V^{\otimes k} \right] \quad (2.19)$$

where $\tau = (12 \dots k)$. However, Eq. (2.19) no longer has the interpretation of a spin model with local interactions.

For general background states, we will only need the replica trick for $k = 2$. Since S_2 has only two elements, each configuration of permutations is completely characterized by the domain $\Delta_A = \{x \in V \text{ such that } \pi_x = \tau\}$.

Because of the boundary conditions in $\mathcal{S}_{A,\tau}$, the collection of these sets coincides with $C(A)$, and hence

$$\mathbb{E} \operatorname{tr}[\rho_A^2] = \sum_{\Delta_A \in C(A)} \operatorname{tr}[\phi_{\Delta_A}^2] = \sum_{\Delta_A \in C(A)} \operatorname{tr}[\phi] 2^{-H_2(\Delta_A)_\phi}. \quad (2.20)$$

Another useful fact is that by Eq. (2.9),

$$\mathbb{E} \rho_{V_\partial} = \phi_{V_\partial}. \quad (2.21)$$

We remark that if one only uses the $k = 2$ replica trick, one could also use tensors which are drawn from a *projective 2-design*, a distribution which produces tensors with the same first and second moments as uniformly random tensors of unit norm [KR05, GAE07]. An example of a projective 2-design is the set of uniformly random stabilizer states. For tensors $|\psi_x\rangle$ drawn from a projective 2-design of dimension D_x , it holds that

$$\mathbb{E} \psi_x^{\otimes 2} = \frac{1}{D_x(D_x - 1)} I + \frac{1}{D_x(D_x - 1)} R_x(\tau),$$

and hence

$$\mathbb{E} \operatorname{tr}[\rho_A^2] = \frac{D_x}{D_x + 1} \sum_{\Delta_A \in C(A)} \operatorname{tr}[\phi_{\Delta_A}^2],$$

which is close to Eq. (2.20) for large D_x . Thus, it is not hard to see that all random tensor network results which only use the $k = 2$ replica trick are also valid for states with tensors drawn from projective 2-designs. This was already observed in [HNQ⁺16] and random tensor networks with random stabilizer tensors were further studied in [NW20]. The results of Section 4 only use the $k = 2$ replica trick, and thus will extend to states with tensors drawn from projective 2-designs. This will not be true for the results in Section 3, which requires usage of the replica trick for all $k \in \mathbb{N}$.

2.3.1 Normalization of random tensor network states

One immediate consequence of the replica trick for $k = 2$ is that the random tensor network state ρ will be approximately normalized with high probability, so long as a mild condition on the background state is satisfied: the bulk needs to be connected, with sufficiently entangled edges. Let

$$\eta = \max_{\Delta \subseteq V_b, \Delta \neq \emptyset} \operatorname{tr}[\phi_\Delta^2] = \max_{\Delta \subseteq V_b, \Delta \neq \emptyset} \operatorname{tr}[\phi] 2^{-H_2(\Delta)_\phi}. \quad (2.22)$$

If the state has enough correlations along each cut (or more precisely, if $H_2(\Delta)_\phi$ is large for each Δ), then η is small. Concretely, if we consider a random tensor network state with maximally entangled link states of bond dimension D , we will have $\eta \leq \frac{1}{D}$. We then have

Lemma 2.1. *For any background state $\phi \in \mathcal{P}_\leq(V)$, with associated $\rho \in \mathcal{P}(V_\partial)$ as in Eq. (2.18), it holds that for any $\varepsilon > 0$*

$$\Pr(|\operatorname{tr}[\rho] - \operatorname{tr}[\phi]| \geq \varepsilon) \leq 2^{|V_b|} \frac{\eta}{\varepsilon^2}$$

where η is defined in Eq. (2.22).

Proof. This follows from a special case of Eq. (2.20). In this case, the empty cut contributes $\operatorname{tr}[\phi]^2$, so we find

$$\operatorname{Var}(\operatorname{tr}[\rho]) = \mathbb{E}|\operatorname{tr}[\rho] - \operatorname{tr}[\phi]|^2 = \mathbb{E}|\operatorname{tr}[\rho]^2 - \operatorname{tr}[\phi]^2| \leq 2^{|V_b|} \max_{\Delta \subseteq V_b, \Delta \neq \emptyset} \operatorname{tr}[\phi_\Delta^2],$$

where we have used the normalization of ρ in expectation $\mathbb{E} \operatorname{tr}[\rho] = \operatorname{tr}[\phi]$, as in Eq. (2.5). The result follows by an application of Chebyshev's inequality. \square

We can improve this result by taking advantage of the fact that our random projectors are random Gaussian vectors, allowing us to use Gaussian concentration of measure rather than the Chebyshev's inequality. For instance, using a concentration bound for Gaussian polynomials (see [AS17], Corollary 5.49) one can show that for any $\varepsilon \geq (\sqrt{2}e)^{2V_b}\eta$:

$$\Pr(|\text{tr}[\rho] - \text{tr}[\phi]| \geq \varepsilon) \leq \exp\left(-\frac{|V_b|}{2e}\varepsilon^{\frac{1}{|V_b|}}\eta^{-\frac{1}{|V_b|}}\right),$$

where η is defined as in Eq. (2.22). We will not need this refinement.

3 Link states with bounded spectral variation

In this section, we study random tensor network states with link states that have *bounded spectral variation*, meaning that there is an effective bond dimension D such that the Schmidt coefficients of the link state are of the order $\frac{1}{D}$.

We start by providing background material on random matrix theory and *free probability*, which is a key tool in the study of products of random matrices. In Section 3.2, we will precisely define the notion of bounded spectral variation and generalize the results in Section 2.2 for random tensor network states with maximally entangled link states to this wider class of link states. This leads to the main result of this section, Theorem 3.4, which shows that the asymptotic entanglement spectrum can be expressed in terms of a free product of distributions. We will see that the results are similar to the quantum gravity set-up described in Appendix A.3. Finally, in Section 3.3, we investigate the entanglement negativity for random tensor network states with link states of bounded spectral variation.

3.1 Random matrices, free probability and non-crossing partitions

3.1.1 Random matrix theory and Wishart matrices

We start by reviewing relevant concepts from probability and random matrix theory that are relevant for our analyses. This material can be found in any introduction to random matrix theory, e.g. [AGZ10, BS10, PB20].

A fundamental question in random matrix theory is as follows: given a family of $n \times n$ matrices with entries selected according to some distribution, what is the asymptotic distribution of the eigenvalues as $n \rightarrow \infty$? This question has been extensively studied, and in many cases has an elegant and concise answer. We discuss a basic example which is closely related to our purposes: *Wishart matrices*. Consider an $n \times m$ matrix X whose entries are drawn i.i.d. from a Gaussian distribution with mean zero and unit variance. The sample covariance matrix of X is the $n \times n$ matrix defined as

$$Y_{n,m} = \frac{1}{m}XX^T. \quad (3.1)$$

Such random matrices are called (*real*) *Wishart matrices*, and can be thought of as a sample second moment matrix (where one has m realizations of an n -dimensional random variable). One can also consider *complex* Wishart matrices: in this case the entries of the $n \times m$ matrix X are complex i.i.d. standard (circularly symmetric) complex Gaussian random variables. We then let $Y_{n,m} = \frac{1}{m}XX^\dagger$. We would like to understand the spectrum of $Y_{n,m}$, and to that end, we consider the empirical distribution of the eigenvalues. This empirical distribution is itself random, depending on the particular realization of $Y_{n,m}$. To characterize the convergence, we recall that if $\{\mu_n\}_{n \in \mathbb{N}}$ is a sequence of random finite measures on \mathbb{R} , we say that the sequence μ_n *converges weakly, in probability*, to a finite measure μ , if, for any bounded continuous function $f \in C_b(\mathbb{R})$, it holds that for every $\varepsilon > 0$

$$\lim_{n \rightarrow \infty} \Pr\left(\left|\int f(x)d\mu_n(x) - \int f(x)d\mu(x)\right| \geq \varepsilon\right) = 0.$$

The asymptotic distribution of the eigenvalues of Wishart matrices is known to obey the *Marchenko-Pastur law* (see, for instance, Theorem 3.6 and Theorem 3.7 in [BS10]):

Theorem 3.1. Consider (real or complex) Wishart matrices $Y_{n,m}$ and let

$$\mu_{n,m} = \frac{1}{n} \sum_{\lambda \in \text{spec}(Y_{n,m})} \delta_\lambda$$

be the empirical distribution of its eigenvalue spectrum. Suppose that the ratio of dimensions n/m converges to a constant $t > 0$ as $n \rightarrow \infty$. Then $\mu_{n,m}$ converges weakly, in probability, to the Marchenko-Pastur distribution $\text{MP}(t)$ with parameter $t > 0$, as defined in Eq. (2.17).

Generalizations to this result are possible. For example, one still has convergence if the entries of X are chosen according to non-Gaussian distributions with mean zero and unit variance. Also, one can prove weak convergence, almost surely (rather than just in probability); see [BS10].

If $Y_{n,m} = \frac{1}{m} X X^\dagger$ is a complex Wishart matrix, X can also be interpreted as a uniformly random pure quantum state on $\mathbb{C}^n \otimes \mathbb{C}^m$, and $Y_{n,m}$, up to normalization, as the reduced density matrix on \mathbb{C}^n [HLW06]. Note that $\frac{1}{n} Y_{n,m}$ is normalized in expectation in the sense that $\mathbb{E} \frac{1}{n} Y_{n,m} = 1$. So, complex Wishart matrices can be used as a model for the reduced state of a random bipartite quantum state, and this allows one to quantify the ‘typical entanglement’ of a random state. Equivalently, in the tensor network setting, $\frac{1}{\sqrt{n}} X$ can be thought of as a random tensor network state with a single bulk vertex, two boundary vertices, and maximally entangled link states. We can then interpret $\frac{1}{n} Y_{n,m}$ as the reduced density matrix on one of the boundary vertices. We will provide a generalization of Theorem 3.1 for the entanglement spectrum of random tensor network states in Theorem 3.4.

3.1.2 Free probability

The topic of probability distributions in random matrix theory is closely related to *free probability* and, in particular, to the notion of the *free product*. We provide a brief introduction here; the material in this section is very standard, and we only review a few relevant aspects. For an extensive treatment, see, for instance, Chapter 5 in [AGZ10] or the books [NS06, MS17, PB20]. As we will see later, the free product will allow us to concisely formulate replica trick results involving multiple minimal cuts.

A *non-commutative probability space* is a pair (\mathcal{A}, ω) , where \mathcal{A} is a C^* -algebra and ω is a state on \mathcal{A} . An element $a \in \mathcal{A}$ is called a non-commutative random variable. The key example to have in mind is the space of $n \times n$ random matrices, where the matrix entries are distributed according to some probability distribution, and $\omega(a) = \mathbb{E} \frac{1}{n} \text{tr}[a]$ defines a tracial state. If $a \in \mathcal{A}$, the *distribution* (or *law*) μ_a of a is defined as a map on polynomials, which evaluates on a polynomial p as $\mu_a(p) = \omega(p(a))$. If a is self-adjoint, it has real spectrum and we can extend the domain of μ_a to all bounded continuous functions $f \in C_b(\mathbb{R})$, using the functional calculus to define $f(a)$ and letting $\mu_a(f) = \omega(f(a))$. In this case we can identify μ_a with a distribution such that, for $f \in C_b(\mathbb{R})$, we have $\mu_a(f) = \int f(x) d\mu_a(x)$. In particular, if a is an $n \times n$ self-adjoint random matrix, then $\mu_a(f) = \frac{1}{n} \mathbb{E} \sum_{\lambda \in \text{spec}(a)} f(\lambda)$, and we may identify μ_a with the empirical measure of the eigenvalues of a . If \mathcal{A} is a commutative algebra, these notions reduce to the usual notions of probability theory, where ω is the expectation.

We call a set of n non-commutative random variables $\{a_i\}$ on a non-commutative probability space (\mathcal{A}, ω) *freely independent* or just *free* if, for any set of $k \geq 2$ polynomials $\{p_j\}$, the variables satisfy

$$\omega(p_1(a_{i_1}) \dots p_k(a_{i_k})) = 0$$

whenever $\omega(p_m(a_{i_m})) = 0$ for all $1 \leq m \leq k$ and no two adjacent indices i_m and i_{m+1} for $1 \leq m \leq k-1$ are equal. One can see that two freely independent variables a_1, a_2 satisfy:

$$0 = \omega((a_1 - \omega(a_1))(a_2 - \omega(a_2))) = \omega(a_1 a_2) - \omega(a_1) \omega(a_2), \quad (3.2)$$

which, in the commutative case with random variables x_1, x_2 , is the classical bivariate independence condition $\mathbb{E}[x_1 x_2] = \mathbb{E}[x_1] \mathbb{E}[x_2]$. While the definition of free independence is stronger than independence (commuting independent random variables are generically not free), the role of free independence is analogous

to the role of classical independence for commuting random variables: it allows one to, in principle, compute the joint mixed moments of the variables.

We will be interested in the *multiplicative free convolution* or *free product* (there also exists an additive convolution or just *free convolution*) of distributions. Suppose a, b are non-commutative self-adjoint free random variables on (\mathcal{A}, ω) with distributions μ_a and μ_b . Then we denote the distribution of ab by $\mu_{ab} = \mu_a \boxtimes \mu_b$. Note that, generally, ab need not be self-adjoint. However, if ω is tracial (as in the random matrix case) and a is positive, the distribution of ab coincides with that of $\sqrt{ab}\sqrt{a}$ which is self-adjoint, and we can identify μ_{ab} with a distribution on \mathbb{R} . If μ_a and μ_b are compactly supported distributions, then so is $\mu_a \boxtimes \mu_b$.

As a concrete example of the freeness and the free product, let X_n and Y_n be two families of random $n \times n$ positive diagonal matrices with uniformly bounded norm, such that their spectrum converges weakly to probability distributions μ and ν respectively. Let U_n be a family of Haar random unitary $n \times n$ matrices. Then as n goes to infinity, X_n and $Y'_n = U_n Y_n U_n^\dagger$ will be freely independent (so they are *asymptotically free*), and we would like to study their product. The product of positive matrices need not be self-adjoint, so we consider $Z_n = \sqrt{X_n} Y'_n \sqrt{X_n}$ which is a positive matrix. One may then show that the distribution of the spectrum of Z_n weakly converges in probability to $\mu \boxtimes \nu$. See Corollary 5.4.11 in [AGZ10] for a precise statement and proof.

The free product may be analyzed using generating functions: given a (non-commutative) random variable a with distribution μ_a , let

$$m_{a,k} = \int x^k d\mu_a(x)$$

be the k -th moment of μ_a . Then the *moment generating function* is the formal power series

$$M_{\mu_a}(z) = \sum_{k=1}^{\infty} m_{a,k} z^k. \quad (3.3)$$

We define the *S-transform* to be the formal power series

$$S_{\mu_a}(z) = \frac{1+z}{z} M_a^{-1}(z),$$

where $M_a^{-1}(z)$ is the power series corresponding to the formal inverse of $M_\mu(z)$ under composition, which is well-defined as long as $m_{a,1} \neq 0$. For compactly supported distributions, the moment generating function, and hence the S-transform, uniquely determines the distribution.

If a and b are non-commutative self-adjoint free random variables, then

$$S_{\mu_a \boxtimes \mu_b}(z) = S_{\mu_{ab}}(z) = S_{\mu_a}(z) S_{\mu_b}(z). \quad (3.4)$$

This also provides a completely combinatorial interpretation of the free product, without reference to the associated non-commutative probability spaces. That is, given compactly supported distributions μ and ν , we can define $\mu \boxtimes \nu$ by Eq. (3.4): it is the compactly supported distribution with moments prescribed by $S_{\mu_a \boxtimes \mu_b}(z)$, and hence, $M_{\mu_a \boxtimes \mu_b}(z)$. The free product is commutative and associative.

As an example, we compute the S-transform of the Marchenko-Pastur distribution $\mu \sim \text{MP}(1)$. The distribution function is given by

$$d\mu(x) = \frac{1}{2\pi} \sqrt{4x^{-1} - 1} dx.$$

The moments can be computed directly:

$$m_k = \sum_{i=0}^{k-1} \frac{1}{i+1} \binom{k}{i} \binom{k-1}{i} \quad (3.5)$$

After some work, one can show that the moments above lead to a closed-form moment generating function

$$M(z) = \frac{2z - 1 - \sqrt{1 - 4z}}{2z}.$$

One may then invert the expression and obtain the S-transform

$$S(z) = \frac{1}{1+z}.$$

Similarly, for the Marchenko-Pastur distribution $MP(t)$ with parameter t , which has distribution as given in Eq. (2.17), we find that

$$S(z) = \frac{1}{t+z}.$$

See, for instance, [BBCC11].

3.1.3 Non-crossing partitions

Given $k \in \mathbb{N}$, let $NC(k)$ denote the set of *non-crossing partitions* of $[k]$. A non-crossing partition of $[k]$ is a partition $[k] = X_1 \sqcup \dots \sqcup X_m$ which is such that, if $i < j \in X_\alpha$, then there are no $k, l \in X_\beta$ for $\beta \neq \alpha$ with $k < i < l < j$ or $i < k < j < l$. To any non-crossing partition, we associate a permutation $\pi \in S_k$ by mapping each subset $\{i_1, \dots, i_l\}$ to the cycle (i_1, \dots, i_l) with $i_1 < \dots < i_l$. In a slight abuse of notation, we will write $\pi \in NC(k)$. For any $\pi \in S_k$, and for a sequence of numbers f_k for $k = 1, 2, \dots$, we write

$$f_\pi = \prod_{l \in C(\pi)} f_l \tag{3.6}$$

where $C(\pi)$ is the cycle type of π . We will need the following result, which is a straightforward consequence of the combinatorics of the S-transform.

Theorem 3.2. *Consider compactly supported probability distributions μ, ν, η . Suppose that the moments of η are given by*

$$m_k^\eta = \sum_{\pi \in NC(k)} m_\pi^\mu m_{\pi^{-1}\tau_k}^\nu$$

where $\tau_k = (12 \dots k)$ is the full cycle. Then

$$\eta = \text{MP}(1) \boxtimes \mu \boxtimes \nu.$$

Proof. We let \mathcal{F} be the transformation that sends a formal power series $f(z)$ to the power series $\frac{1}{z}f^{-1}(z)$. This is such that for some distribution μ , the S-transform is given by $S_\mu(z) = (1+z)\mathcal{F}(M_\mu)(z)$. Moreover, given power series $f(z) = \sum_k f_k z^k$ and $g(z) = \sum_k g_k z^k$, define a convolution operation \otimes by

$$(f \otimes g)(z) = \sum_k \left(\sum_{\pi \in NC(k)} f_\pi g_{\pi^{-1}\tau_k} \right) z^k$$

where τ_k is the full cycle in S_k . Then Theorem 18.14 in [NS06] states that for any two f and g with $f_1 \neq 0$ and $g_1 \neq 0$, it holds that

$$\mathcal{F}(f \otimes g)(z) = \mathcal{F}(f)(z)\mathcal{F}(g)(z).$$

Then the S-transform of η can be written:

$$S_\eta(z) = (1+z)\mathcal{F}(M_\eta)(z) = (1+z)\mathcal{F}(M_\mu)(z)\mathcal{F}(M_\nu)(z) = \frac{1}{1+z}S_\mu(z)S_\nu(z).$$

This implies the desired result, as the S-transform of $\text{MP}(1)$ is given by $\frac{1}{1+z}$, and the S-transform uniquely determines a compactly supported distribution. \square

We remark briefly that free independence can equivalently be formulated in terms of the vanishing of *free cumulants*, which are themselves defined in terms of sums over non-crossing partitions. We refer the interested reader to any of the previously cited references for a more in-depth discussion on the role of non-crossing partitions in free probability. For our purposes, the fact that non-crossing partitions are intimately related to free independence will allow us to later phrase random tensor network results in terms of free probability.

3.2 Entanglement spectrum of random tensor network states as a free product

We now return to studying random tensor network states. Consider a family of states in $\mathcal{P}_=(V)$ composed of the tensor product of link states $|\phi_e\rangle$ along the edges $e \in E$ as in Eq. (2.1), and assume that along each edge, the bond dimensions scale with a parameter D , so $D_e = \Theta(D)$. Our key assumption is that the link states have *bounded spectral variation* – by this we mean that the empirical distribution of the rescaled entanglement spectrum of the link states

$$\mu_e^{(D)} := \sum_{i=1}^{D_e} \frac{1}{D_e} \delta_{D_e \lambda_{e,i}} \quad (3.7)$$

has all moments converging to the moments $m_{e,k}$ of a compactly supported probability distribution μ_e , as D goes to infinity. We assume that the link states are normalized, so $m_{e,1} = 1$. This condition implies that, up to a vanishing fraction as $D \rightarrow \infty$, the elements of the entanglement spectrum of the link state are of order D^{-1} .

For a minimal cut γ_A , let $\mu_{\gamma_A}^{(D)}$ be the distribution for the spectrum of the tensor product of the link states in γ_A :

$$\mu_{\gamma_A}^{(D)} = \bigotimes_{e \in \gamma_A} \mu_e^{(D)} = \frac{1}{D_{\gamma_A}} \sum_{\{i_e\}} \delta_{D_{\gamma_A} \prod_{e \in \gamma_A} \lambda_{e,i_e}}$$

where $i_e = 1, \dots, D_e$ and $D_{\gamma_A} = \prod_{e \in \gamma_A} D_e$. We define the tensor product of distributions as follows: if X_1 and X_2 are independent real valued random variables with distributions μ_{X_1} and μ_{X_2} , then $\mu_{X_1} \otimes \mu_{X_2}$ is defined as the distribution of the product $X_1 X_2$. The distribution $\mu_{\gamma_A}^{(D)}$ has k -th moment given by $m_{\gamma_A,k}^{(D)} = \prod_{e \in \gamma_A} m_{e,k}^{(D)}$, and we can see that $m_{\gamma_A,k}^{(D)}$ converges to $m_{\gamma_A,k}$, the moments of the distribution

$$\mu_{\gamma_A} := \bigotimes_{e \in \gamma_A} \mu_e.$$

Let $\text{spec}(\rho_A) = \{\lambda_{A,i}\}$ (recall that $\text{spec}(\rho_A)$ is ordered in non-increasing order). Let γ_A be a cut for A . By a standard argument, the number of nonzero eigenvalues of ρ_A (that is, $\text{rank}(\rho_A)$) is upper bounded by D_{γ_A} . If γ_A is the unique minimal cut, then we define

$$\mu_A^{(D)} := \frac{1}{D_{\gamma_A}} \sum_{i=1}^{D_{\gamma_A}} \delta_{D_{\gamma_A} \lambda_{A,i}}. \quad (3.8)$$

If there are multiple minimal cuts, it is ambiguous which γ_A , and hence, which D_{γ_A} , we should pick; we choose the cut for which D_{γ_A} is minimal in Eq. (3.8), and we will denote this minimal cut by $\gamma_{A,1}$. The moments of $\mu_A^{(D)}$ are given by

$$m_{A,k}^{(D)} := \int z^k d\mu_A(z) = D_{\gamma_A}^{k-1} \sum_{i=1}^{D_{\gamma_A}} (\lambda_{A,i})^k.$$

Note that the distribution $\mu_A^{(D)}$ is random, and correspondingly, the moments $m_{A,k}^{(D)}$ are random variables. In contrast, the moments $m_{e,k}^{(D)}$ and $m_{\gamma_A,k}^{(D)}$ are numbers depending only on the bond dimension.

The theorem we want to prove will follow straightforwardly from a key intermediate result: as D goes to infinity, all the moments of the boundary distribution $\mu_A^{(D)}$ converge to the moments of μ_{γ_A} . We use the notation in Eq. (3.6) to write expressions like

$$m_{\gamma_A, \pi} = \prod_{l \in C(\pi)} m_{\gamma_A, l}$$

for a permutation $\pi \in S_k$. We will then apply the *method of moments* to show that convergence of moments implies convergence in distribution. As a remark on notation, in the error bounds in both the current section and Section 4, when we use \mathcal{O} -notation, the constants may depend on the graph underlying the tensor network (typically our bounds scale as $2^{|V_b|}$, where V_b is the set of bulk vertices).

Proposition 3.3. *If there exists a unique minimal cut γ_A for A , then*

$$\lim_{D \rightarrow \infty} \mathbb{E} m_{A, k}^{(D)} = m_{\gamma_A, k}. \quad (3.9)$$

If there exist exactly two minimal cuts $\gamma_{A,1}$ and $\gamma_{A,2}$, which do not intersect (so $\gamma_{A,1} \cap \gamma_{A,2} = \emptyset$) and for which $\frac{D_{\gamma_{A,1}}}{D_{\gamma_{A,2}}}$ converges to a constant $t \leq 1$, then

$$\lim_{D \rightarrow \infty} \mathbb{E} m_{A, k}^{(D)} = \sum_{\pi \in NC(k)} t^{d(\pi, \text{id})} m_{\gamma_{A,1}, \tau^{-1}\pi} m_{\gamma_{A,2}, \pi}. \quad (3.10)$$

Moreover, in both cases the variance goes to zero as D goes to infinity: for every k

$$\mathbb{E} \left[\left(m_{A, k}^{(D)} - \mathbb{E} \left[m_{A, k}^{(D)} \right] \right)^2 \right] = \mathcal{O} \left(\frac{1}{D} \right). \quad (3.11)$$

Proof. We first provide a sketch of the proof. It proceeds via the following steps:

1. Write the expectation of the moments of $\mu_A^{(D)}$ as the partition function for a classical spin model, as in Section 2.1.
2. Show that the contributions from terms of the form given in the statement of the proposition dominate the partition function by carefully tracking the powers of D , and showing that all other contributions are suppressed polynomially in D .
3. Show that the variance of the moments vanishes in the limit $D \rightarrow \infty$ by direct computation.

We begin with Step 1. First, we observe that the k -th moment of $\mu_A^{(D)}$ is given by $m_{A, k}^{(D)} = D_{\gamma_A}^{k-1} \text{tr}[\rho_A^k]$. Consider the expression in Eq. (2.14) for the replica trick with permutation π on A :

$$Z_{k, \pi} := \mathbb{E} \text{tr} [R_A(\pi) \rho_A^{\otimes k}] = \sum_{\{\pi_x\} \in \mathcal{S}_{A, \pi}} \prod_{e=(xy) \in E} \prod_{l \in C(\pi_x^{-1} \pi_y)} \text{tr} [\phi_{e, x}^l]. \quad (3.12)$$

Recall that the set $\mathcal{S}_{A, \pi}$, as defined in Eq. (2.11), consists of assignments of permutations to each $x \in V$, subject to $\pi_x = \pi$ for $x \in A$ and $\pi_x = \text{id}$ for $x \in \bar{A}$. As in Eq. (2.12), if $\pi = \tau$, then we indeed have $Z_{k, \tau} = \mathbb{E} \text{tr}[\rho_A^k]$, so

$$\mathbb{E} m_{A, k}^{(D)} = D_{\gamma_A}^{k-1} Z_{k, \tau}. \quad (3.13)$$

On the other hand, if we let $k = 2n$ and $\pi = \tilde{\tau} = (12 \dots n)(n+1 \ n+2 \dots 2n)$, then $Z_{k, \pi} = \mathbb{E} \left[\text{tr}[\rho_A^n]^2 \right]$, and hence

$$\mathbb{E} \left[\left(m_{A, n}^{(D)} \right)^2 \right] = D_{\gamma_A}^{2n-2} Z_{k, \tilde{\tau}}. \quad (3.14)$$

Recall that $m_{e,l}^{(D)} = D_e^{l-1} \text{tr}[\phi_{e,x}^l]$, and write

$$Z_{k,\pi} = \sum_{\{\pi_x\} \in \mathcal{S}_{A,\pi}} Z_k(\{\pi_x\})$$

where

$$\begin{aligned} Z_k(\{\pi_x\}) &:= \prod_{e=(xy) \in E} \prod_{l \in C(\pi_x^{-1}\pi_y)} \text{tr}[\phi_{e,x}^l] \\ &= \prod_{e=(xy) \in E} D_e^{|C(\pi_x^{-1}\pi_y)|-k} \prod_{l \in C(\pi_x^{-1}\pi_y)} m_{e,l}^{(D)} \\ &= \prod_{e=(xy) \in E} D_e^{-d(\pi_x,\pi_y)} m_{e,\pi_x^{-1}\pi_y}^{(D)}. \end{aligned} \quad (3.15)$$

This accomplishes Step 1: we have recast the problem of computing moments into a question of computing a partition function for a classical spin model with fixed boundary conditions.

For Step 2, we want to show that the dominant contribution(s) to $Z_{k,\pi}$ as D goes to infinity are those given in the statement of the proposition. This will simply be a matter of checking powers of D , and using the triangle inequality property of the Cayley distance. If Γ_A is the unique minimal cut, then we let $\pi_x^{\min} = \pi$ for $x \in \Gamma_A$ and $\pi_x^{\min} = \text{id}$ for $x \in V \setminus \Gamma_A$, and we have

$$Z_k(\{\pi_x^{\min}\}) = D_{\gamma_A}^{-d(\pi,\text{id})} \prod_{e \in \gamma_A} m_{e,\pi}^{(D)} = D_{\gamma_A}^{-d(\pi,\text{id})} m_{\gamma_A,\pi}^{(D)}. \quad (3.16)$$

If there are exactly two minimal cuts $\Gamma_{A,1} \subset \Gamma_{A,2}$, we let $V = V_1 \sqcup V_2 \sqcup V_3$, with $V_1 = \Gamma_{A,1}$, $V_2 = \Gamma_{A,2} \cap \Gamma_{A,1}^c$ and $V_3 = \Gamma_{A,2}^c$. Now consider the permutations $\sigma \in S_k$ that are on a geodesic between π and id (recall this implies $d(\pi,\sigma) + d(\sigma,\text{id}) = d(\pi,\text{id})$), and consider the configuration given by $\pi_x^\sigma = \pi$ for $x \in V_1$, $\pi_x^\sigma = \sigma$ for $x \in V_2$, and $\pi_x^\sigma = \text{id}$ for $x \in V_3$. By hypothesis, $\gamma_{A,1}$ and $\gamma_{A,2}$ do not intersect, and hence, the edges in each cut are distinct. Then this configuration has weight

$$\begin{aligned} Z_k(\{\pi_x^\sigma\}) &= \prod_{e_1 \in \gamma_{A,1}} D_{e_1}^{-d(\pi,\sigma)} \prod_{l_1 \in C(\pi^{-1}\sigma)} m_{e_1,l_1}^{(D)} \prod_{e_2 \in \gamma_{A,2}} D_{e_2}^{-d(\sigma,\text{id})} \prod_{l_2 \in C(\sigma)} m_{e_2,l_2}^{(D)} \\ &= D_{\gamma_{A,1}}^{-d(\pi,\text{id})} \left(\frac{D_{\gamma_{A,1}}}{D_{\gamma_{A,2}}} \right)^{d(\sigma,\text{id})} \prod_{e_1 \in \gamma_{A,1}} m_{e_1,\pi^{-1}\sigma}^{(D)} \prod_{e_2 \in \gamma_{A,2}} m_{e_2,\sigma}^{(D)} \\ &= D_{\gamma_{A,1}}^{-d(\pi,\text{id})} \left(\frac{D_{\gamma_{A,1}}}{D_{\gamma_{A,2}}} \right)^{d(\sigma,\text{id})} m_{\gamma_{A,1},\pi^{-1}\sigma}^{(D)} m_{\gamma_{A,2},\sigma}^{(D)}, \end{aligned} \quad (3.17)$$

where $D_{\gamma_{A,1}}/D_{\gamma_{A,2}}$ converges to t , by assumption. Now, to show that these configurations yield the dominant contributions, we will need to use that $D_e = \Theta(D)$, so let us write $\frac{D}{D_e} = C_e^{(D)} = \Theta(1)$. Then for general configurations labeled by π_x , we may rewrite Eq. (3.15) as

$$Z_k(\{\pi_x\}) = \prod_{e=(xy) \in E} D^{-d(\pi_x,\pi_y)} (C_e^{(D)})^{d(\pi_x,\pi_y)} m_{e,\pi_x^{-1}\pi_y}^{(D)}.$$

The configurations we claimed to be dominant satisfy $Z_k(\{\pi\}) = \Theta(D^{-m(A)d(\pi,\text{id})})$, where we recall that $m(A)$ is the size of a minimal cut for A . Now we will show that all other configurations satisfy $Z_k(\{\pi\}) = \mathcal{O}(D^{-m(A)d(\pi,\text{id})-1})$. To this end, consider some arbitrary configuration $\{\pi_x\} \in \mathcal{S}_{A,\pi}$. Let P be a maximal set of edge-disjoint paths in G from A to \bar{A} . It is a well-known fact that such a set has size $m(A)$, by the max-flow min-cut theorem. Let

$$C_k^{(D)} := \left(\max_{e \in E, l=1,\dots,k} (C_e^{(D)})^{l-1} \right) \left(\max_{e \in E, \pi \in S_k} m_{e,\pi}^{(D)} \right).$$

Then we may bound

$$Z_k(\{\pi_x\}) \leq (C_k^{(D)})^{|E|} \prod_{e=(xy) \in E} D^{-d(\pi_x, \pi_y)} \leq (C_k^{(D)})^{|E|} \prod_{p \in P} D^{-\sum_{e=(xy) \in p} d(\pi_x, \pi_y)}. \quad (3.18)$$

The first inequality is clear from the definition of $C_k^{(D)}$, and in the second inequality, we simply restrict to a subset of the edges we sum over. Note that $C_k^{(D)} = \mathcal{O}(1)$. Then, by the triangle inequality for the Cayley distance d , it holds that

$$\sum_{e=(xy) \in p} d(\pi_x, \pi_y) \geq d(\pi, \text{id})$$

with equality if and only if the only edges (xy) for which $\pi_x \neq \pi_y$ are on a path in P , and each of the paths is a geodesic. Then we conclude

$$Z_k(\{\pi_x\}) \leq C_k^{(D)} \prod_{p \in P} D^{-d(\pi, \text{id})} = C_k^{(D)} D^{-m(A)d(\pi, \text{id})},$$

and we see that the weight of every configuration can be bounded by the product of a $\mathcal{O}(1)$ number and a polynomial in D .

Now, as promised, we show that if $\{\pi_x\}$ is not one of the minimal configurations described above, we actually have

$$Z_k(\{\pi_x\}) = \mathcal{O}(D^{-m(A)d(\pi, \text{id})-1}). \quad (3.19)$$

To see this, we rewrite the triangle inequality for the Cayley distance as:

$$\prod_{e=(xy) \in E} D^{-d(\pi_x, \pi_y)} \leq \prod_{p \in P} D^{-\sum_{e=(xy) \in p} d(\pi_x, \pi_y)} \leq D^{-m(A)d(\pi, \text{id})} \quad (3.20)$$

with equality if and only if the π_x are on a geodesic path in P . We now show that this is satisfied only for the configurations we claimed to be minimal. Assume that $\{\pi_x\} \in \mathcal{S}_{A, \pi}$ is such that the inequalities in Eq. (3.20) are equalities and let

$$\Delta_n = \{x \in V \text{ such that } d(\pi_x, \pi) \leq n\}.$$

Then $\Delta_n \in C(A)$ for $0 \leq n < d(\pi, \text{id})$, and we denote by δ_n the associated set of edges crossing the cut. Each edge $(xy) \in \delta_n$ must be such that $\pi_x \neq \pi_y$, so it must be on a path in P , and because the permutations are geodesics along the paths, they must be on different paths. Hence $|\delta_n| \leq |P| = m(A)$, implying each Δ_n is a minimal cut. This immediately implies the claim if there is a unique minimal cut, since we must have $\Delta_{d(\pi, \text{id})-1} = \Delta_0 = \Gamma_A$. If there are exactly two minimal cuts, then we must have $\pi_x = \pi$ for $x \in V_1$, $\pi_x = \text{id}$ for $x \in V_3$, and there must be some l such that for all $x \in V_2$ we have $d(\pi, \pi_x) = l$ and $d(\pi_x, \text{id}) = d(\pi, \text{id}) - l$. Then in order to have equality in Eq. (3.20), we must have that for all $x \in V_2$, π_x equals some fixed permutation σ , because the assumption of having exactly two cuts implies that V_2 is connected, and we must have $d(\pi_x, \pi_y) = 0$ for all $(xy) \in E$ with $x, y \in V_2$. This proves Eq. (3.19).

In conclusion, if there is a unique minimal cut, then by Eq. (3.16) and Eq. (3.19), we find

$$Z_{k, \pi} = D_{\gamma_A}^{-d(\pi, \text{id})} m_{\gamma_A, \pi}^{(D)} + \mathcal{O}(D^{-m(A)d(\pi, \text{id})-1}), \quad (3.21)$$

and if there are exactly two (non-intersecting) cuts, then by Eq. (3.17) and Eq. (3.19), we find

$$Z_{k, \pi} = \sum_{\sigma, d(\pi, \sigma) + d(\sigma, \text{id}) = d(\pi, \text{id})} D_{\gamma_A, 1}^{-d(\pi, \text{id})} \left(\frac{D_{\gamma_A, 1}}{D_{\gamma_A, 2}} \right)^{d(\sigma, \text{id})} m_{\gamma_A, 1, \pi^{-1}\sigma}^{(D)} m_{\gamma_A, 2, \sigma}^{(D)} + \mathcal{O}(D^{-m(A)d(\pi, \text{id})-1}). \quad (3.22)$$

Finally, we set $\pi = \tau$ for the full cycle τ and we use Eq. (3.13). For a unique minimal cut γ_A , by Eq. (3.21)

$$\begin{aligned}\mathbb{E}m_{\gamma_A,k} &= D_{\gamma_A}^{k-1} Z_{k,\tau} \\ &= m_{\gamma_A,\pi}^{(D)} + \mathcal{O}(D_{\gamma_A}^{k-1} D^{-m(A)(k-1)-1}) \\ &= m_{\gamma_A,k}^{(D)} + \mathcal{O}\left(\frac{1}{D}\right).\end{aligned}\tag{3.23}$$

using $d(\tau, \text{id}) = k - 1$ and $D_{\gamma_A} = \Theta(D^{m(A)})$. This proves Eq. (3.9) as $m_{\gamma_A,k}^{(D)}$ converges to $m_{\gamma_A,k}$.

For two non-intersecting minimal cuts, we saw that the dominant contribution is due to configurations $\{\pi_x^\sigma\}$ for σ on a geodesic between τ and id . Then applying the observation that σ is on such a geodesic if and only if σ is a non-crossing partition similarly yields that by Eq. (3.22)

$$\begin{aligned}\mathbb{E}m_{\gamma_A,k} &= D_{\gamma_A}^{k-1} Z_{k,\tau} \\ &= D_{\gamma_A}^{k-1} \sum_{\sigma \in NC(k)} \left(\frac{D_{\gamma_{A,1}}}{D_{\gamma_{A,2}}}\right)^{d(\sigma,\text{id})} m_{\gamma_{A,1},\tau^{-1}\sigma}^{(D)} m_{\gamma_{A,2},\sigma}^{(D)} + \mathcal{O}(D_{\gamma_A}^{k-1} D^{-m(A)(k-1)-1}) \\ &= \sum_{\sigma \in NC(k)} \left(\frac{D_{\gamma_{A,1}}}{D_{\gamma_{A,2}}}\right)^{d(\sigma,\text{id})} m_{\gamma_{A,1},\tau^{-1}\sigma}^{(D)} m_{\gamma_{A,2},\sigma}^{(D)} + \mathcal{O}\left(\frac{1}{D}\right).\end{aligned}\tag{3.24}$$

Since $m_{\gamma_{A,1},\tau^{-1}\sigma}^{(D)} \rightarrow m_{\gamma_{A,1},\tau^{-1}\sigma}$, $m_{\gamma_{A,2},\sigma}^{(D)} \rightarrow m_{\gamma_{A,2},\sigma}$ and $D_{\gamma_{A,1}}/D_{\gamma_{A,2}} \rightarrow t$, this proves Eq. (3.10).

This accomplishes Step 2: we have shown that the configurations $\{\pi_x^{\min}\}$ (in case of a unique minimal cut for A) and $\{\pi_x^\sigma\}$ (in case there are exactly two non-intersecting minimal cuts for A) dominate in the computation of the expectation of the k -th moment of μ_A in terms of powers of D .

We complete the proof by showing that the variance of $m_{A,k}$ vanishes as $D \rightarrow \infty$. We use the observation in Eq. (3.14), applying the analysis of $Z_{2k,\pi}$ to the case where $\pi = \tilde{\tau} = (12 \dots k)(k+1k+2 \dots 2k)$. If there is a unique minimal cut, then using Eq. (3.21) and the fact that $d(\tilde{\tau}, \text{id}) = 2k - 2$, we find

$$\mathbb{E}\left[\left(m_{A,k}^{(D)}\right)^2\right] = D_{\gamma_A}^{2k-2} Z_{2k,\tilde{\tau}} = \prod_{l \in C(\tilde{\tau})} m_{\gamma_A,l}^{(D)} + \mathcal{O}(D_{\gamma_A}^{2k-2} D^{-m(A)(2k-2)-1}) = \left(m_{\gamma_A,k}^{(D)}\right)^2 + \mathcal{O}\left(\frac{1}{D}\right).$$

By Eq. (3.23) we know that $(\mathbb{E}m_{A,k}^{(D)})^2 = \left(m_{\gamma_A,k}^{(D)} + \mathcal{O}\left(\frac{1}{D}\right)\right)^2$, and we conclude that the variance obeys

$$\mathbb{E}\left[\left(m_{A,k}^{(D)} - \mathbb{E}\left[m_{A,k}^{(D)}\right]\right)^2\right] = \mathbb{E}\left[\left(m_{A,k}^{(D)}\right)^2\right] - \left(\mathbb{E}m_{A,k}^{(D)}\right)^2 = \mathcal{O}\left(\frac{1}{D}\right).$$

For the case with exactly two minimal cuts, a similar argument holds. Here, the key observation is that σ is on a geodesic between $\tilde{\tau}$ and id if and only if $\sigma = \sigma_1 \sigma_2$ where σ_1 is on a geodesic between $(12 \dots k)$ and id and σ_2 is on a geodesic between $(k+1k+2 \dots 2k)$ and id . Using Eq. (3.22), this implies

$$\begin{aligned}\mathbb{E}\left(m_{A,k}^{(D)}\right)^2 &= D_{\gamma_A}^{2k-2} Z_{2k,\tilde{\tau}} \\ &= \sum_{\sigma_1, \sigma_2 \in NC(k)} \left(\frac{D_{\gamma_{A,1}}}{D_{\gamma_{A,2}}}\right)^{d(\sigma_1,\text{id})+d(\sigma_2,\text{id})} m_{\gamma_{A,1},\tau^{-1}\sigma_1}^{(D)} m_{\gamma_{A,1},\tau^{-1}\sigma_2}^{(D)} m_{\gamma_{A,2},\sigma_1}^{(D)} m_{\gamma_{A,2},\sigma_2}^{(D)} + \mathcal{O}\left(\frac{1}{D}\right) \\ &= \left(\sum_{\sigma \in NC(k)} \left(\frac{D_{\gamma_{A,1}}}{D_{\gamma_{A,2}}}\right)^{d(\sigma,\text{id})} m_{\gamma_{A,1},\tau^{-1}\sigma}^{(D)} m_{\gamma_{A,2},\sigma}^{(D)}\right)^2 + \mathcal{O}\left(\frac{1}{D}\right).\end{aligned}$$

By Eq. (3.24), we see that this coincides with $(\mathbb{E}m_{A,k}^{(D)})^2$, up to $\mathcal{O}\left(\frac{1}{D}\right)$, and hence Eq. (3.11) holds. \square

We now have the ingredients to prove that the entanglement spectrum of random tensor networks with link states with bounded spectral variation can be written in a simple fashion. We will use the *method of*

moments to translate the above result on convergence of moments to convergence in distribution. The basic statement is that, given certain conditions on the distributions in question, if the moments of a sequence of distribution μ_n converge to those of μ , then $\mu_n \Rightarrow \mu$ – see for instance Theorem 30.8 in [Bil08].

The method of moments is valid, so long as a distribution μ is completely determined by its moments. This occurs if, for all k , the k -th moment $m_{\mu,k}$ is bounded as

$$m_{\mu,k} \leq AB^k k! \quad (3.25)$$

for constants A, B independent of k . If the distributions have compact support, as in Proposition 3.3, then this condition is satisfied.²

Now that we have established the convergence of moments in Proposition 3.3, we have our main result of the (conditional) convergence in distribution. As in Proposition 3.3 we consider a family of random tensor network states with link states with bounded spectral variation with increasing D , as defined in the beginning of this section.

Theorem 3.4. *If there exists a unique minimal cut γ_A for A , then $\mu_A^{(D)} \Rightarrow \mu_{\gamma_A}$, in probability, as $D \rightarrow \infty$. If there exist exactly two minimal cuts $\gamma_{A,1}$ and $\gamma_{A,2}$, which do not intersect and for which $\lim_{D \rightarrow \infty} \frac{D_{\gamma_{A,1}}}{D_{\gamma_{A,2}}} = t \leq 1$, then $\mu_A^{(D)} \Rightarrow \text{MP}(1) \boxtimes \mu_{\gamma_{A,1}} \boxtimes \mu_{\gamma_{A,2}}(t)$, in probability, where $\mu_{\gamma_{A,2}}(t) = (1-t)\delta_0 + t\mu_{\gamma_{A,2}}$*

Proof. It is straightforward to see that the k -th moment of $\mu_{\gamma_{A,2}}(t)$ is given by $t^{k-1}m_{\gamma_{A,2},k}$, and then the result follows immediately from Proposition 3.3, Theorem 3.2, and the method of moments. Because we assumed that for any minimal cut γ_A for A , the limiting distributions μ_{γ_A} are compactly supported, they are uniquely determined by their moments. Hence, the method of moments is valid, and the convergence of moments implies convergence in distribution. \square

Remark 3.5. In Theorem 3.4, we assumed that the two cuts were non-intersecting. What happens if there are still only exactly two minimal cuts, but $\gamma_{A,1} \cap \gamma_{A,2}$ is nonempty? This extension is straightforward. Let $\gamma_A^{(a)} := \gamma_{A,1} \cap \gamma_{A,2}$ and let $\gamma_{A,i}^{(b)} = \gamma_{A,i} \setminus \gamma_A^{(a)}$ for $i = 1, 2$. In line with previous notation, let $\mu_{\gamma_A^{(a)}}$ and $\mu_{\gamma_{A,i}^{(b)}}$ denote the corresponding limiting distributions of the entanglement spectra along these sets, with moments $m_{\gamma_A^{(a)},k}$ and $m_{\gamma_{A,i}^{(b)},k}$. The only step in the proof of Proposition 3.3 where we used that the cuts were non-intersecting is when we computed the value of $Z_k(\{\pi_x\})$ for the optimal configuration. If the cuts do intersect, and we consider the configuration with $\pi_x = \tau$ for $x \in V_1$ with τ the complete cycle, $\pi_x = \sigma$ for $x \in V_2$ and $\sigma \in \text{NC}(k)$, and $\pi_x = \text{id}$ for $x \in V_3$, then a quick calculation shows

$$Z_k(\{\pi_x\}) \rightarrow D_{\gamma_{A,1}}^{-d(\pi, \text{id})} (D_{\gamma_{A,1}}/D_{\gamma_{A,2}})^{d(\sigma, \text{id})} \prod_{e \in \gamma_A^{(a)}} m_{e,k} \prod_{e_1 \in \gamma_{A,1}^{(b)}} m_{e_1, \pi^{-1}\sigma} \prod_{e_2 \in \gamma_{A,2}^{(b)}} m_{e_2, \sigma}.$$

Apart from this modification, the proof of Proposition 3.3 is still valid, leading to

$$Z_{k,\tau} = m_{\gamma_A^{(a)},k} \sum_{\sigma \in \text{NC}(k)} t^{d(\sigma, \text{id})} m_{\gamma_{A,1}^{(b)}, \tau^{-1}\sigma} m_{\gamma_{A,2}^{(b)}, \sigma}.$$

If, in Theorem 3.4, we do not assume that the cuts are non-intersecting, then the partition function above leads to a limiting distribution given by

$$\mu_{\gamma_A^{(a)}} \otimes \left(\text{MP}(1) \boxtimes \mu_{\gamma_{A,1}^{(b)}} \boxtimes \mu_{\gamma_{A,2}^{(b)}}(t) \right).$$

²A basic example of a distribution which does not have compact support, but is nevertheless uniquely determined by its moments is a standard Gaussian distribution. On the other hand, a standard example of distributions that are not determined by their moments are the densities on $\mathbb{R}_{\geq 0}$ with $d\mu_1(x) = \sqrt{2\pi}x^{-1}e^{-(\log x)^2/2}dx$ and $d\mu_2(x) = (1 + \sin(2\pi \log x))d\mu_1(x)$, for which it can be verified that the n -th moments of both distributions are equal to $e^{-n^2/2}$, while the distributions are clearly not identical.

3.3 Nontrivial link states and entanglement negativity

As another application of the theory of free probability, we will compute the entanglement negativity spectrum for random tensor network states with link states with bounded spectra. In [DQW21], it was shown how to compute the entanglement negativity spectrum for a random tensor network state with maximally entangled link states using a replica trick. Using the methods from the previous subsection, we can analyze the negativity for entangled link states with bounded spectral variation. We remark that similar computations have recently been performed in [DMW21] in the context of replica wormholes, and our assumption on the link states is a generalization of the “pairwise connected regime” in [DMW21]. Another work investigating nontrivial entanglement negativity spectra in random tensor networks is [KFNR21], where they focus on the effect of having multiple minimal cuts in the network. As our analysis will be a straightforward combination of the arguments in [DQW21] and Section 3.2, we will be rather concise; the main message of this section is to show that the language of free probability applies to other random tensor network computations as well.

We first recall how negativity functions as an entanglement measure for mixed states. Let \mathcal{T} be the superoperator which maps an operator X to its transpose X^\top , and \mathcal{I} be the identity superoperator. For $\rho_{AB} \in \mathcal{P}(AB)$,

$$\rho_{AB}^{T_B} := (\mathcal{I}_A \otimes \mathcal{T}_B)(\rho_{AB})$$

is the partial transpose of ρ_{AB} on the B system. The *logarithmic* or *entanglement negativity* is given by

$$E_N(\rho_{AB}) = \log \frac{\|\rho_{AB}^{T_B}\|_1}{\text{tr}[\rho]}.$$

It is a measure for the entanglement of the mixed state ρ_{AB} : if $E_N(\rho_{AB}) > 0$ the state must be entangled. We call $\text{spec}(|\rho_{AB}^{T_B}|)$ the *entanglement negativity spectrum*. In analogy to the Rényi entropies, we can generalize the logarithmic negativity to a one-parameter family of negativities. The k -th Rényi negativity is given by

$$N_k(\rho_{AB}) = \text{tr}[(\rho_{AB}^{T_B})^k].$$

If we let $N_m^{(\text{even})}(\rho_{AB}) = N_{2m}(\rho_{AB})$, then the logarithmic negativity is obtained as an analytic continuation in the Rényi index $m \rightarrow \frac{1}{2}$ of $\log(N_m^{(\text{even})}(\rho_{AB}))$. More precisely, in the expression

$$\log \sum_{\lambda \in \text{spec}(\rho_{AB}^{T_B})} |\lambda|^\alpha,$$


we may take $\alpha \rightarrow \frac{1}{2}$ to obtain $E_N(\rho_{AB}) + \log \text{tr}[\rho]$.

In the context of random tensor networks, we partition the boundary in three regions: $V_\partial = A \sqcup B \sqcup C$, and we would like to compute the Rényi negativities of the reduced state ρ_{AB} . We will then use this to determine the entanglement negativity spectrum, and compute the entanglement negativity. The idea is that the k -th Rényi negativity can be computed using a replica trick, by placing the full cycle $\tau_k = (12 \dots k) \in S_k$ on A and $\tau_k^{-1} = (k k-1 \dots 1)$ on B :

$$N_k(\rho_{AB}) = \text{tr}[\rho_{AB}^{\otimes k} (R_A(\tau) \otimes R_B(\tau^{-1}))] = \text{tr}[\rho_{ABC}^{\otimes k} (R_A(\tau) \otimes R_B(\tau^{-1}) \otimes R_C(\text{id}))].$$

Let us first discuss the case with maximally entangled link states, following [DQW21]. The same arguments as in Section 2.2 show that one can compute the expectation of $N_k(\rho_{AB})$ for a random tensor network state using a spin model, now with boundary conditions of τ_k on A , τ_k^{-1} on B and id on C . We will assume that the minimal cuts Γ_A , Γ_B and Γ_C are unique. Note that the minimal cut for AB is given by $\Gamma_{AB} = \Gamma_C^c$. From the theory of multi-commodity flows, it is known that there exist sets of edge-disjoint paths $P = P_{AB} \cup P_{AC} \cup P_{BC}$, where P_{AB} consists of paths from A to B , and similarly for P_{AC} and P_{BC} , and which are such that

$$|P_{AB}| + |P_{AC}| = |\gamma_A|, \quad |P_{AB}| + |P_{BC}| = |\gamma_B|, \quad |P_{AC}| + |P_{BC}| = |\gamma_C|.$$



(a) Illustration of the case where $\gamma_C = \gamma_A \cup \gamma_B$, where the dominant configuration is given by τ_k on Γ_A , τ_k^{-1} on Γ_B and id on Γ_C .

(b) Illustration of the case where $\gamma_C \neq \gamma_A \cup \gamma_B$, where the dominant configuration is given by τ_k on Γ_A , τ_k^{-1} on Γ_B and id on Γ_C and some non-crossing pairing π_1 on the domain V_1 .

Figure 4: Tensor networks with one and two minimal cuts. The relevant ground state configuration domains are denoted by Γ_A .

This can be used to show (in analogous fashion to the proof of [Proposition 3.3](#)) that, if $k = 2n$ is even, any spin model configuration contributing to $\mathbb{E}N_k(\rho_{AB})$ is of order $\mathcal{O}(D^{-(n-1)(|\gamma_A|+|\gamma_B|)-n|\gamma_C|})$. If $k = 2n + 1$ is odd, any spin model configuration contributing to $\mathbb{E}N_k(\rho_{AB})$ is of order $\mathcal{O}(D^{-n(|\gamma_A|+|\gamma_B|+|\gamma_C|)})$.

In order to determine what happens as $D \rightarrow \infty$, we need to determine the dominant configurations. Let r be the number of connected components of $V \setminus (\Gamma_A \cup \Gamma_B \cup \Gamma_C)$. There are two distinct cases. The first is when the minimal cut for AB (which is the complement of the minimal cut for C) is the union of the minimal cuts for A and B , so $\Gamma_{AB} = \Gamma_A \cup \Gamma_B$ and hence $\gamma_{AB} = \gamma_A \cup \gamma_B$. Then the minimal cuts naturally partition the bulk vertices into three cuts Γ_A , Γ_B and Γ_C , and we have $r = 0$. In this case, the dominant configurations in the spin model are those where the vertices in Γ_A are assigned τ_k , those in Γ_B are assigned τ_k^{-1} and those in Γ_C are assigned id . This is illustrated in [Figure 4a](#).

The second case is when $\Gamma_A \cup \Gamma_B \subsetneq \Gamma_{AB}$ and hence $\gamma_{AB} \neq \gamma_A \cup \gamma_B$. Now, we have again the domains Γ_A , Γ_B and Γ_C , but upon removing these vertices, there may also be connected components V_1, \dots, V_r which are not connected to A , B or C . Here, the minimal configurations are those for which, again, the vertices in Γ_A are assigned τ_k , those in Γ_B are assigned τ_k^{-1} and those in Γ_C are assigned id , and where in each component V_i the vertices are assigned a permutation π_i which is such that it satisfies three conditions: it must be on a geodesic between τ_k and τ_k^{-1} , on a geodesic between τ_k and id and on a geodesic between τ_k^{-1} and id . If $k = 2n$ is even, such permutations are given by *non-crossing pairings*: permutations corresponding to non-crossing partitions in which each cycle has length 2. The set of non-crossing pairings on $2n$ elements is in bijection with the set of non-crossing partitions on n elements, so the number of non-crossing pairings on $2n$ elements is given by $|NC(n)| = C_n$. One way to obtain this correspondence is as follows. If π is a non-crossing pairing, $\tau_{2n}\pi$ will map even numbers to even numbers, and restricting to the even numbers and relabeling $2i \mapsto i$ yields a non-crossing partition $\sigma \in NC(n)$. Moreover, restricting to the odd numbers and relabeling $2i + 1 \mapsto i$ yields the non-crossing partition $\sigma^{-1}\tau \in NC(n)$. This leads to C_n^r dominant contributions to $\mathbb{E}N_{2n}(\rho_{AB})$ of size $D^{-(n-1)(|\gamma_A|+|\gamma_B|)-n|\gamma_C|}$ since we can choose a non-crossing pairing π_i for each component. Such a configuration is illustrated in [Figure 4b](#).

For odd $k = 2n + 1$, we similarly have permutations which correspond to a non-crossing partition, and which have a single fixed point and all other cycles with length 2. This leads to $((2n + 1)C_n)^r$ dominant

contributions to $\mathbb{E}N_{2n}(\rho_{AB})$, of size $D^{-(n-1)(|\gamma_A|+|\gamma_B|)-n|\gamma_C|}$. We also note that $\text{rank}(\rho_{AB}^{T_B}) \leq D^{|\gamma_A|+|\gamma_B|}$. If $\text{spec}(\rho_{AB}^{T_B}) = \{s_i\}$, then we define the measure

$$\mu_{AB}^{(D)} = \frac{1}{D^{|\gamma_A|+|\gamma_B|}} \sum_{i=1}^{D^{|\gamma_A|+|\gamma_B|}} \delta_{D^{\frac{1}{2}(|\gamma_A|+|\gamma_B|+|\gamma_C|)} s_i}. \quad (3.26)$$

This has moments given by

$$m_{AB,k}^{(D)} = \int x^k d\mu_{AB}^{(D)}(x) = D^{(\frac{k}{2}-1)(|\gamma_A|+|\gamma_B|)+\frac{k}{2}|\gamma_C|} N_k(\rho_{AB}).$$

If we take the expectation of the moments, we again need to distinguish the two cases. If $|\gamma_A| + |\gamma_B| = |\gamma_C|$, we see that the powers of D cancel for the dominant configurations, so $m_{AB,k}^{(D)} \rightarrow 1$ for all k . On the other hand, for $|\gamma_A| + |\gamma_B| > |\gamma_C|$, we see that for $D \rightarrow \infty$ with odd k , we have $\mathbb{E}m_{AB,k}^{(D)} \rightarrow 0$. For even k , we recover the degeneracy of the dominant configurations, leading to

$$\lim_{D \rightarrow \infty} m_{AB,k}^{(D)} = \begin{cases} 1 & \text{if } |\gamma_A| + |\gamma_B| = |\gamma_C|, \\ 0 & \text{if } k \text{ odd and } |\gamma_A| + |\gamma_B| > |\gamma_C|, \\ C_{k/2}^r & \text{if } k \text{ even and } |\gamma_A| + |\gamma_B| > |\gamma_C|, \end{cases} \quad (3.27)$$

where r is the number of connected components of $V \setminus (\Gamma_A \cup \Gamma_B \cup \Gamma_C)$. In fact, one can show that, as in [Proposition 3.3](#), the variance goes to zero as well, and hence the method of moments allows one to conclude that $\mu_{AB}^{(D)}$ converges weakly, in probability, to

$$\mu_{AB} = \begin{cases} \sigma^{\otimes r} & \text{if } r > 0, \\ \frac{1}{2}\delta_1 + \frac{1}{2}\delta_{-1} & \text{if } r = 0 \text{ and } |\gamma_A| + |\gamma_B| > |\gamma_C|, \\ \delta_1 & \text{if } r = 0 \text{ and } |\gamma_A| + |\gamma_B| = |\gamma_C|, \end{cases} \quad (3.28)$$

where σ is the semi-circle distribution with density

$$d\sigma(x) = \frac{1}{2\pi} \sqrt{4-x^2} \mathbb{1}_{|x| \leq 2} dx$$

Alternatively, one may study the empirical distribution of the *squared* entanglement negativity spectrum

$$\nu_{AB}^{(D)} = \frac{1}{D^{|\gamma_A|+|\gamma_B|}} \sum_i \delta_{D^{|\gamma_A|+|\gamma_B|+|\gamma_C|} s_i^2}. \quad (3.29)$$

This distribution has k -th moment given by $m_{AB,2k}^{(D)}$, and in comparison with the limiting moments in [Eq. \(3.27\)](#), one can conclude that $\nu_{AB}^{(D)} \Rightarrow \nu_{AB}$, in probability, where

$$\nu_{AB} = MP(1)^{\otimes r}. \quad (3.30)$$

The logarithmic negativity can be computed using the distribution $\mu_{AB}^{(D)}$ or $\nu_{AB}^{(D)}$ as

$$E_N(\rho_{AB}) = \log \int |\lambda| d\mu_{AB}^{(D)}(\lambda) + \frac{\log D}{2} (|\gamma_A| + |\gamma_B| - |\gamma_C|) - \log \text{tr}[\rho] \quad (3.31)$$

$$= \log \int \sqrt{\lambda} d\nu_{AB}^{(D)}(\lambda) + \frac{\log D}{2} (|\gamma_A| + |\gamma_B| - |\gamma_C|) - \log \text{tr}[\rho]. \quad (3.32)$$

The convergence of $\nu_{AB}^{(D)}$ to ν_{AB} implies³ that $E_N(\rho_{AB}) - \frac{\log D}{2} (|\gamma_A| + |\gamma_B| - |\gamma_C|)$ converges in probability to

$$\log \int \sqrt{\lambda} d\nu_{AB}(\lambda) = r \log \frac{8}{3\pi}.$$

³The function $f(\lambda) = \sqrt{\lambda}$ is not in $C_b(\mathbb{R})$, but the method of moments actually shows a stronger convergence, allowing test functions to have polynomial growth.

See Appendix D of [DQW21] for details and proofs.

A straightforward combination of the arguments in Section 3.2 and [DQW21] shows that the same configurations are the dominant contributions for link states with bounded spectral variation as in Section 3.2. To determine the limiting distribution in this case, we can generalize Eq. (3.28) in the same fashion as in Section 3.2. We assume the minimal cuts Γ_A , Γ_B and Γ_C are unique. We also assume that $\gamma_A \cap \gamma_B = \emptyset$, and in the case where $\gamma_C = \gamma_{AB} \neq \gamma_A \cup \gamma_B$ (so $|\gamma_A| + |\gamma_B| > |\gamma_C|$), all pairwise intersections between γ_A , γ_B and γ_C are empty. This excludes the case where $|\gamma_A| + |\gamma_B| > |\gamma_C|$, but $r = 0$. We let $\gamma_{A,i}$ and $\gamma_{B,i}$ denote the components of γ_A and γ_B which are connected to V_i , and we let $\mu_{\gamma_{A,i}}^{(D)}$ and $\mu_{\gamma_{B,i}}^{(D)}$ denote the distribution of the spectrum along these sets, with associated k -th moments $m_{\gamma_{A,i},k}^{(D)}$, $m_{\gamma_{B,i},k}^{(D)}$, which we assume to converge to the moments $m_{\gamma_{A,i},k}$, $m_{\gamma_{B,i},k}$ of compactly supported distributions $\mu_{\gamma_{A,i}}$ and $\mu_{\gamma_{B,i}}$. For convenience, we assume $D_e = D$ for all edges $e \in E$.

We can now compute the dominant contributions to $\mathbb{E}N_k(\rho_{AB})$. If $\gamma_C = \gamma_A \cup \gamma_B$, then there is a unique dominant configuration, which contributes $D^{-(k-1)|\gamma_C|} m_{\gamma_C,k}^{(D)}$. If $|\gamma_A| + |\gamma_B| > |\gamma_C|$ and $k = 2n$ is even, consider the configuration which assigns π_i to V_i , where each π_i is a non-crossing pairing. For each edge $e \in \gamma_C$, we have $m_{e,\pi_i}^{(D)} = (m_{e,2}^{(D)})^n$, so this configuration contributes

$$D^{-(n-1)(|\gamma_A|+|\gamma_B|)-n|\gamma_C|} \left(m_{\gamma_C,2}^{(D)}\right)^n \prod_{i=1}^r \left(m_{\gamma_{A,i},\tau_{2n}^{-1}\pi_i}^{(D)} m_{\gamma_{B,i},\tau_{2n}\pi_i}^{(D)}\right).$$

Recalling the construction of the equivalence between $NC(n)$ and non-crossing pairings on $2n$ elements, we see that

$$m_{\gamma_{B,i},\tau_{2n}\pi_i}^{(D)} = m_{\gamma_{B,i},\sigma_i}^{(D)} m_{\gamma_{B,i},\sigma_i^{-1}\tau_n}^{(D)}$$

for some unique $\sigma_i \in NC(n)$. Similarly, one may verify

$$m_{\gamma_{A,i},\tau_{2n}^{-1}\pi_i}^{(D)} = m_{\gamma_{A,i},\sigma_i}^{(D)} m_{\gamma_{A,i},\sigma_i^{-1}\tau_n}^{(D)}.$$

This implies that the contribution of all dominant configurations is given by

$$\left(m_{\gamma_C,2}^{(D)}\right)^n \prod_{i=1}^r \left(\sum_{\sigma \in NC(n)} m_{\gamma_{A,i},\sigma}^{(D)} m_{\gamma_{B,i},\sigma}^{(D)} m_{\gamma_{A,i},\sigma^{-1}\tau_n}^{(D)} m_{\gamma_{B,i},\sigma^{-1}\tau_n}^{(D)}\right)$$

As in the maximally entangled case, upon rescaling, the odd moments vanish as $D \rightarrow \infty$. In conclusion, the resulting asymptotic moments are given by

$$\lim_{D \rightarrow \infty} \mathbb{E}m_{AB,k}^{(D)} = \begin{cases} m_{\gamma_C,k} & \text{if } |\gamma_A| + |\gamma_B| = |\gamma_C|, \\ 0 & \text{if } k \text{ odd and } |\gamma_A| + |\gamma_B| > |\gamma_C|, \\ m_k & \text{if } k \text{ even and } |\gamma_A| + |\gamma_B| > |\gamma_C| \end{cases} \quad (3.33)$$

with

$$m_{2n} = m_{\gamma_C,2}^n \prod_{i=1}^r \left(\sum_{\sigma \in NC(n)} m_{\gamma_{A,i},\sigma} m_{\gamma_{B,i},\sigma} m_{\gamma_{A,i},\sigma^{-1}\tau_n} m_{\gamma_{B,i},\sigma^{-1}\tau_n}\right).$$

As before, one can also show, in similar fashion to the proof of Proposition 3.3, that the variance of the moments goes to zero as $D \rightarrow \infty$. For the case $|\gamma_A| + |\gamma_B| > |\gamma_C|$, we consider $\nu_{AB}^{(D)}$ similar to Eq. (3.29), but with an additional rescaling by $m_{\gamma_C,2}$:

$$\nu_{AB}^{(D)} = \frac{1}{D^{|\gamma_A|+|\gamma_B|}} \sum_i \delta_{D^{|\gamma_A|+|\gamma_B|+|\gamma_C|} m_{\gamma_C,2}^{-1} s_i^2}.$$

This has moments, which compute $N_k^{(\text{even})}(\rho_{AB})$, converging to

$$\lim_{D \rightarrow \infty} \mathbb{E} \int x^k d\nu_{AB}^{(D)}(x) = \prod_{i=1}^r \left(\sum_{\sigma \in NC(n)} m_{\gamma_{A,i},\sigma} m_{\gamma_{B,i},\sigma} m_{\gamma_{A,i},\sigma^{-1}\tau_n} m_{\gamma_{B,i},\sigma^{-1}\tau_n} \right).$$

Thus, by the method of moments and [Theorem 3.2](#), it holds that $\nu_{AB}^{(D)}$ converges weakly, in probability, to

$$\nu_{AB} = \begin{cases} \bigotimes_{i=1}^r \nu_i & \text{if } r > 0, \\ \mu_{\gamma_C} & \text{if } r = 0, \end{cases} \quad (3.34)$$

where ν_i is given by

$$\nu_i = (\mu_{\gamma_{A,i}} \otimes \mu_{\gamma_{B,i}})^{\boxtimes 2} \boxtimes \text{MP}(1).$$

This reduces to [Eq. \(3.30\)](#) if the link states are maximally entangled. We can use this to compute the logarithmic negativity, as we did previously. For $r > 0$,

$$E_N(\rho_{AB}) = \log \int \sqrt{\lambda} d\nu_{AB}^{(D)}(\lambda) + \frac{\log D}{2} (|\gamma_A| + |\gamma_B| - |\gamma_C|) + \frac{1}{2} \log m_{\gamma_{C,2}} - \log \text{tr}[\rho],$$

from which we find that $E_N(\rho_{AB}) - \frac{\log D}{2} (|\gamma_A| + |\gamma_B| - |\gamma_C|)$ converges in probability to

$$\log \int \sqrt{\lambda} d\nu_{AB}(\lambda) + \frac{1}{2} \log m_{\gamma_{C,2}}.$$

For the case $|\gamma_A| + |\gamma_B| = |\gamma_C|$, it is more elegant to use the limiting distribution of $\mu_{AB}^{(D)}$, as defined in [Eq. \(3.26\)](#). By the method of moments and [Eq. \(3.33\)](#), $\mu_{AB}^{(D)} \Rightarrow \mu_{\gamma_C}$, in probability. We may then compute the entanglement negativity as

$$E_N(\rho_{AB}) = \log \int |\lambda| d\mu_{AB}^{(D)}(\lambda) + \frac{\log D}{2} (|\gamma_A| + |\gamma_B| - |\gamma_C|) - \log \text{tr}[\rho],$$

and hence $E_N(\rho_{AB}) - \frac{\log D}{2} (|\gamma_A| + |\gamma_B| - |\gamma_C|)$ converges in probability to $\log \int |\lambda| d\mu_{AB}(\lambda)$.

4 Link states with unbounded spectral variation

We will now consider a different regime, where the link states have unbounded spectral variation. Our methods in this section are distinct from the previous one, and the two sections can be considered separately.

4.1 One-shot entropies

We begin by introducing one of our main tools for studying entanglement spectra in random tensor network states: *one-shot entropies*. In quantum information theory, the rates of certain important protocols, such as compression or state merging can be expressed as entropic quantities. One-shot entropies are the appropriate analogs for settings where one would like to analyze a task for a single or finite number of copies of the relevant state. Asymptotic rates in terms of ordinary von Neumann entropies are then recovered in the limit of infinitely many independent copies. For an extensive introduction to this point of view, see [\[Tom15\]](#); here we provide the basic definitions and introduce the relevant concepts.

A random tensor network built from link states that are maximally entangled (or more generally have bounded spectral variation) can be analyzed using asymptotic tools. Indeed, if we have a maximally entangled state of large dimension $D = 2^n$, then this is equal to the n -th tensor power of a qubit maximally entangled state, so we are effectively in an asymptotic situation. However, if we allow for link states with unbounded

spectral variation or even completely general background states, as in [Section 2.3](#), then it is more natural to use tools from one-shot quantum information theory.

We take the Rényi entropies as a starting point, which we defined in [Eq. \(2.7\)](#) for subnormalized states. Let \mathcal{H} be some Hilbert space and for $\rho \in \mathcal{P}_{\leq}(\mathcal{H})$ we define the (unconditional) *min-entropy* and the *max-entropy* by

$$\begin{aligned} H_{\min}(\rho) &= -\log \|\rho\|_{\infty} \\ H_{\max}(\rho) &= \log (\operatorname{tr}[\sqrt{\rho}]^2) \end{aligned}$$

which coincide with the Rényi entropies $H_{\infty}(\rho)$ and $H_{\frac{1}{2}}(\rho)$ for $\rho \in \mathcal{P}_{=}(\mathcal{H})$. As usual, if ρ_A is the reduced density matrix on a system A , we write $H_{\min}(A)_{\rho} = H_{\min}(\rho_A)$ and $H_{\max}(A)_{\rho} = H_{\max}(\rho_A)$.

Often, when applied to study quantum information processing tasks, it is useful to allow a small error. This leads to the introduction of smooth entropies. To define these we use a distance measure known as the *purified distance*, which is given for $\rho, \sigma \in \mathcal{P}_{\leq}(\mathcal{H})$ by

$$P(\rho, \sigma) = \sqrt{1 - F_{*}(\rho, \sigma)^2}$$

where $F_{*}(\rho, \sigma)$ is the *generalized fidelity* between ρ and σ , which is defined by

$$F_{*}(\rho, \sigma) = F(\rho, \sigma) + \sqrt{(1 - \operatorname{tr}[\rho])(1 - \operatorname{tr}[\sigma])}$$

in terms of the ordinary *fidelity* $F(\rho, \sigma) = \|\sqrt{\rho}\sqrt{\sigma}\|_1$. We define the *smooth min- and max-entropies* of $\rho \in \mathcal{P}_{\leq}(\mathcal{H})$ as

$$\begin{aligned} H_{\min}^{\varepsilon}(\rho) &= \sup_{\rho^{\varepsilon} \in \mathcal{P}_{\leq}(\mathcal{H}), P(\rho^{\varepsilon}, \rho) \leq \varepsilon} H_{\min}(\rho^{\varepsilon}) \\ H_{\max}^{\varepsilon}(\rho) &= \sup_{\rho^{\varepsilon} \in \mathcal{P}_{\leq}(\mathcal{H}), P(\rho^{\varepsilon}, \rho) \leq \varepsilon} H_{\max}(\rho^{\varepsilon}). \end{aligned}$$

The smooth entropies are such that one recovers the usual von Neumann entropies in the limit of many independent copies. Indeed, the following *asymptotic equipartition property* holds:

$$\lim_{n \rightarrow \infty} \frac{1}{n} H_{\min}^{\varepsilon}(\rho^{\otimes n}) = H(\rho) = \lim_{n \rightarrow \infty} \frac{1}{n} H_{\max}^{\varepsilon}(\rho^{\otimes n})$$

for any $0 < \varepsilon < 1$. Variations on this definition are possible. For instance, one can choose a different distance measure, which will yield different entropies. However, for the usual choices, the differences go to zero as ε goes to zero, so the particular choice is often immaterial. For instance, consider the *trace distance* between $\rho, \sigma \in \mathcal{P}_{\leq}(\mathcal{H})$, which is defined by

$$T(\rho, \sigma) = \frac{1}{2} \|\rho - \sigma\|_1 + \frac{1}{2} |\operatorname{tr}[\rho - \sigma]|,$$

where the last term, which is absent in usual definitions of the trace distance, accounts for subnormalized states. It is easy to see that $T(\rho, \sigma) \leq \|\rho - \sigma\|_1 \leq 2T(\rho, \sigma)$. The Fuchs-van de Graaff inequalities relate the trace distance and purified distance:

$$T(\rho, \sigma) \leq P(\rho, \sigma) \leq \sqrt{2T(\rho, \sigma)} \tag{4.1}$$

for $\rho, \sigma \in \mathcal{P}_{\leq}(\mathcal{H})$.

There are also conditional versions of the Rényi entropies. Consider a bipartite quantum state $\rho_{AB} \in \mathcal{P}_{=}(AB)$. For the von Neumann entropy, the conditional entropy can simply be defined as an entropy difference, namely $H(A|B)_{\rho} = H(AB)_{\rho} - H(B)_{\rho}$. However, it turns out that this is not a good definition in the Rényi case. There are various ways to define a Rényi conditional entropy $H_k(A|B)$; we use a version based on the so-called sandwiched Rényi relative entropy. For $k = 2$, this gives a *quantum conditional collision*

entropy, which will be useful for defining minimal cuts and which is defined as follows. For $\rho_{AB} \in \mathcal{P}_{\leq}(AB)$, let

$$H_2(A|B)_{\rho|\rho} := -\log \operatorname{tr} \left[\left((I \otimes \rho_B)^{-\frac{1}{4}} \rho_{AB} (I \otimes \rho_B)^{-\frac{1}{4}} \right)^2 \right] + \log \operatorname{tr}[\rho]. \quad (4.2)$$

Finally, there are also conditional versions of the min- and max-entropy. For $\rho_{AB} \in \mathcal{P}_{\leq}(AB)$ and $\sigma_B \in \mathcal{P}_{\leq}(B)$, we define

$$\begin{aligned} H_{\min}(A|B)_{\rho|\sigma} &= -\inf\{\lambda : \rho_{AB} \leq 2^\lambda I_A \otimes \sigma_B\} \\ H_{\max}(A|B)_{\rho|\sigma} &= \log \|\sqrt{\rho_{AB}} \sqrt{I \otimes \sigma_B}\|_1^2 \end{aligned}$$

and we let

$$\begin{aligned} H_{\min}(A|B)_\rho &= \sup_{\sigma \in \mathcal{P}_{\leq}(B)} H_{\min}(A|B)_{\rho|\sigma} \\ H_{\max}(A|B)_\rho &= \sup_{\sigma \in \mathcal{P}_{\leq}(B)} H_{\max}(A|B)_{\rho|\sigma}. \end{aligned}$$

We can also define their smoothed versions

$$\begin{aligned} H_{\min}^\varepsilon(A|B)_\rho &= \sup_{\rho^\varepsilon \in \mathcal{P}_{\leq}(AB), P(\rho^\varepsilon, \rho) \leq \varepsilon} H_{\min}(A|B)_{\rho^\varepsilon} \\ H_{\max}^\varepsilon(A|B)_\rho &= \sup_{\rho^\varepsilon \in \mathcal{P}_{\leq}(AB), P(\rho^\varepsilon, \rho) \leq \varepsilon} H_{\max}(A|B)_{\rho^\varepsilon}. \end{aligned}$$

There is a duality between (smooth) max- and min-entropies. If $\rho \in \mathcal{P}_{\leq}(ABC)$ is a *pure* state, it holds that

$$H_{\min}^\varepsilon(A|B)_\rho = -H_{\max}^\varepsilon(A|C)_\rho. \quad (4.3)$$

We will use the fact that for a normalized state $\rho_{AB} \in \mathcal{P}_=(AB)$ (Corollary 5.10 in [Tom15])

$$H_{\min}(A|B)_\rho \leq H_2(A|B)_{\rho|\rho}. \quad (4.4)$$

A final important property of conditional smooth entropies are the *data processing inequalities*. Let Φ and Ψ be completely positive and trace-preserving (CPTP) maps, mapping systems A to A' and B to B' respectively, and let $\sigma = (\Phi \otimes \Psi)(\rho)$. If Φ is also subunital, and $0 \leq \varepsilon \leq \sqrt{\operatorname{tr} \rho}$, then Theorem 6.2 of [Tom15] states

$$H_{\min}^\varepsilon(A'|B')_\sigma \geq H_{\min}^\varepsilon(A|B)_\rho, \quad \text{and} \quad H_{\max}^\varepsilon(A'|B')_\sigma \geq H_{\max}^\varepsilon(A|B)_\rho.$$

In fact, for the smooth min-entropy the data processing inequality is also valid if Φ is only trace-nonincreasing rather than trace-preserving, see [Tom12].

4.2 Recovery isometries

Recall that we study random tensor network states with *link states*, pure states placed on each edge whose tensor product forms the full state on edges $\phi = \bigotimes_{e \in E} \phi_e \in \mathcal{P}_=(V)$ for some graph $G = (V, E)$, and in Section 2.3, we considered more general *background states* $\phi_V \in \mathcal{P}_{\leq}(V)$, where we no longer have a tensor product structure along the edges of some graph, and applying the replica trick does not yield a local spin model for the moments of the tensor network state. This situation is of independent interest, but will also be useful as an intermediate step when applying bounds based on one-shot entropies to link states. In Section 3, we studied link states for which the entanglement spectrum of the edge states ϕ_e had bounded variation, and we used the replica trick to compute the moments of the spectrum of ρ_A for a boundary subsystem A . For general background states we saw that the replica trick for $k = 2$ extends as in Eq. (2.20). What are the minimal cuts in this setting? Based on Eq. (2.20) a first guess would be that $\Gamma_A \in C(A)$ would be a

minimal cut (i.e. correspond to the dominant term in the replica trick) if for all other cuts $\Delta_A \in C(A)$ we would have $H_2(\Gamma_A)_\phi \ll H_2(\Delta_A)_\phi$. If the state is a link state, this corresponds to adding weights to the edges of the graph corresponding to the Rényi-2 entropies along the edges, and computing a weighted minimal cut. Indeed, this would yield an accurate approximation of $\text{tr}[\rho_A^2]$ and hence of $H_2(\rho_A)$. However, if the spectrum of ρ_A is not close to a flat spectrum, this does not imply that $\text{spec}_+(\rho_A)$ is close to $\text{spec}_+(\phi_{\Gamma_A})$. We would like to show that for link states with unbounded spectral variation, and an appropriate minimal cut condition for $\Gamma_A \in C(A)$, it is still true that $\text{spec}_+(\rho_A)$ is close to $\text{spec}_+(\phi_{\Gamma_A})$.

We will adapt the $k = 2$ replica trick for general background states to get a bound on the difference in trace norm between $\text{spec}_+(\rho_A)$ and $\text{spec}_+(\phi_{\Gamma_A})$ in terms of *conditional* Rényi-2 entropies,⁴ as defined in Eq. (4.2). In Section 4.3, we will use this to formulate a condition for cut minimality in terms of smooth entropies for link states.

The main result of this subsection is a tensor network version of *one-shot decoupling*. Let $\phi_V \in \mathcal{P}_\leq(V)$. We allow ϕ_V to be a general state, which need not be pure and also need not be a product state along the edges of some graph. Let R be a purifying system and $\phi_{VR} \in \mathcal{P}_\leq(VR)$ be a purification of ϕ_V . Then we can construct the random tensor network state $\rho_{V_\partial R}$ where the boundary systems are given by $V_\partial \cup R$, which is a purification of the random tensor network state ρ_{V_∂} as in Eq. (2.18) by

$$\phi_{V_\partial R} = \text{tr}_{V_b}[(I_{V_\partial R} \otimes \psi)\phi] \quad (4.5)$$

where ψ is a tensor product of random tensors. We briefly recall our notation for boundary subsystems and cuts: for a boundary subsystem $A \subseteq V_\partial$, we denote its boundary complement by $\bar{A} = V_\partial \setminus A$, and for a cut $\Gamma_A \in C(A)$, we let $\Gamma_A^c = V \setminus \Gamma_A$, which is a cut for \bar{A} . The purifying system R can be thought of as an additional boundary system in the tensor network construction.

In Theorem 4.4, we will assume that we have a cut $\Gamma_A \in C(A)$ which is such that for all cuts $\Delta_A \in C(A)$ for which $\Delta_A \subsetneq \Gamma_A$ we have $H_2(\Gamma_A \setminus \Delta_A | \Gamma_A^c R)_{\phi|\phi} \gg 1$, and similarly for all cuts $\Delta_A \in C(A)$ for which $\Gamma_A \subsetneq \Delta_A$ we have $H_2(\Delta_A \setminus \Gamma_A | \Gamma_A R)_{\phi|\phi} \gg 1$. We show that this condition implies that with high probability there exist isometries $V_A : \mathcal{H}_A \rightarrow \mathcal{H}_{\Gamma_A}$ and $V_{\bar{A}} : \mathcal{H}_{\bar{A}} \rightarrow \mathcal{H}_{\Gamma_A^c}$ such that

$$(V_A \otimes V_{\bar{A}} \otimes I_R) |\rho\rangle \approx |\phi\rangle. \quad (4.6)$$

The approximation accuracy will be measured in trace norm. In particular, this implies that $\text{spec}_+(\rho_A) \approx \text{spec}_+(\phi_{\Gamma_A})$. If the state ϕ is a tensor product of link states, $\text{spec}_+(\phi_{\Gamma_A})$ is precisely the entanglement spectrum along the cut γ_A . The isometries V_A and $V_{\bar{A}}$ are *recovery isometries*, which allow us to ‘recover’ Γ_A from the A system, and similarly we can recover Γ_A^c from \bar{A} .

The result is closely related to quantum error correction. One way to interpret this is as follows: consider a subspace \mathcal{H}_S of \mathcal{H}_V and let R be a reference system of dimension $\dim(\mathcal{H}_S)$, and ϕ_{VR} a maximally entangled state between S and R . Then Eq. (4.6) can be interpreted as saying that if we encode the subspace S by projecting onto random tensors, the information in Γ_A is protected, after encoding, against an erasure error on \bar{A} . This idea is also discussed in [PYHP15] for perfect tensor network models, and in [HNQ⁺16] for random tensor networks with maximally entangled link states. In holography, the notion of local recovery isometries and their error correction interpretation goes under the name of *entanglement wedge reconstruction* or *subregion-subregion duality*. See [AR19, AP22] for a detailed discussion of entanglement wedge reconstruction in holographic systems with bulk entropy, relating to one-shot entropies. We provide more details in Appendix B.

Our approach to showing Eq. (4.6) is that we start by projecting only on the random tensors in Γ_A , and not on the random tensors in Γ_A^c . This yields a random tensor network state σ on $A\Gamma_A^c R$.

We then show that, by a version of one-shot decoupling, the reduced state on $\sigma_{\Gamma_A^c R}$ has not changed much from $\phi_{\Gamma_A^c R}$. By Uhlmann’s theorem, this implies that there exists an isometry V_A such that $(V_A \otimes I_{\Gamma_A^c R}) |\sigma\rangle \approx |\phi\rangle$. Combining this with a similar result for Γ_A^c we obtain Eq. (4.6), as will be made precise in Theorem 4.4.

In our construction of σ , we can relabel the vertices in the graph, and think of the vertices in $\Gamma_A \setminus A$ as the bulk vertices V_b , the boundary subsystem A as the complete boundary V_∂ , and relabel all other subsystems as

⁴Note that while $H(A|B)_\phi = H(AB)_\phi - H(B)_\phi$, in general $H_2(A|B)_{\phi|\phi} \neq H_2(AB)_\phi - H_2(B)_\phi$.

the reference system R . Then we prove the following result, which is closely related to the one-shot decoupling results in [DBWR14].

Proposition 4.1. *Consider a random tensor network state $\rho_{V_\partial R}$ as in Eq. (4.5) with a (purified) background state $\phi_{VR} \in \mathcal{P}_<(VR)$. Let $A = V_\partial$ and let $\Gamma_A = V$ and suppose that for any cut $\Delta_A \in C(A)$ other than Γ_A*

$$H_2(\Gamma_A \setminus \Delta_A | R)_{\phi|\phi} \geq K$$

then

$$\mathbb{E} \|\rho_R - \phi_R\|_1 \leq 2^{\frac{|V_b|}{2}} \sqrt{\text{tr}[\phi]} 2^{-\frac{1}{2}K}.$$

Note that, since $\Gamma_A = V_b \cup A$, the sets $\Gamma_A \setminus \Delta_A$ for $\Delta_A \in C(A) \setminus \{\Gamma_A\}$ are exactly the non-empty subsets of V_b . The formulation in terms of $\Delta_A \in C(A)$ will be natural when we apply this result in Theorem 4.4.

Proof. We closely follow the strategy in [DBWR14, DH10]. We first note a basic fact (Lemma 3.7 in [DBWR14]): for any operator X and ω a subnormalized density matrix, it holds that

$$\|X\|_1 \leq \|\omega^{-\frac{1}{4}} X \omega^{-\frac{1}{4}}\|_2. \quad (4.7)$$

The proof is an application of the Cauchy-Schwarz inequality. We use Eq. (4.7) with $\omega = \phi_R$ and Jensen's inequality to see that

$$\mathbb{E} \|\rho_R - \phi_R\|_1 \leq \sqrt{\mathbb{E} \text{tr}[(\tilde{\rho}_R - \tilde{\phi}_R)^2]}$$

where $\tilde{\rho}_{V_\partial R} = (I \otimes \phi_R)^{-\frac{1}{4}} \rho_{V_\partial R} (I \otimes \phi_R)^{-\frac{1}{4}}$ and $\tilde{\phi}_{VR} = (I \otimes \phi_R)^{-\frac{1}{4}} \phi_{VR} (I \otimes \phi_R)^{-\frac{1}{4}}$. Now $\mathbb{E} \tilde{\rho}_R = \tilde{\phi}_R$ by Eq. (2.21), and the replica trick in Eq. (2.20) yields

$$\begin{aligned} \mathbb{E} \text{tr}[(\tilde{\rho}_R - \tilde{\phi}_R)^2] &= \mathbb{E} \text{tr}[\tilde{\rho}_R^2] - \text{tr}[\tilde{\phi}_R^2] \\ &= \sum_{\Delta_A \in C(A), \Delta_A \not\subseteq \Gamma_A} \text{tr}[\tilde{\phi}_{(\Gamma_A \setminus \Delta_A)R}^2] \\ &= \sum_{\Delta_A \in C(A), \Delta_A \not\subseteq \Gamma_A} \text{tr}[\phi] 2^{-H_2(\Gamma_A \setminus \Delta_A | R)_{\phi|\phi}} \end{aligned}$$

using the definition of $\tilde{\phi}$ and Eq. (4.2) and hence

$$(\mathbb{E} \|\rho_R - \phi_R\|_1)^2 \leq 2^{|V_b|} \text{tr}[\phi] 2^{-K}.$$

□

Suppose that in the set-up of Proposition 4.1, we would have equality $\rho_R = \phi_R$. Then, by Uhlmann's theorem, their purifications ρ_{AR} and $\phi_{\Gamma_A R}$ are related by an isometry V_A from A to Γ_A . The following lemma is useful to extend to the case where the reduced states are close in trace distance.

Lemma 4.2. *Suppose $\rho_{AB} \in \mathcal{P}(AB)$ and $\sigma_{AC} \in \mathcal{P}_<(AC)$ are pure states on Hilbert spaces $\mathcal{H}_A \otimes \mathcal{H}_B$ and $\mathcal{H}_A \otimes \mathcal{H}_C$ respectively. Then*

$$\min_V \|(I_A \otimes V) \rho_{AB} (I_A \otimes V^\dagger) - \sigma_{AC}\|_1 \leq 2 \sqrt{2 \|\rho_A - \sigma_A\|_1 + 2 \|\rho_A - \sigma_A\|_1^2}.$$

where the minimum is over all isometries $V : \mathcal{H}_B \rightarrow \mathcal{H}_C$.

Proof. Uhlmann's theorem states that if $\rho_{AB} \in \mathcal{P}(AB)$ and $\sigma_{AC} \in \mathcal{P}(AC)$ are pure quantum states with $\dim(\mathcal{H}_B) \leq \dim(\mathcal{H}_C)$, then there exists an isometry $V : \mathcal{H}_B \rightarrow \mathcal{H}_C$ such that

$$P(\rho_A, \sigma_A) = P((I_A \otimes V)\rho_{AB}(I_A \otimes V^\dagger), \sigma_{AC}) = P(\rho_A, \sigma_A)$$

and, in particular, the isometry is the solution to an optimization problem:

$$P(\rho_A, \sigma_A) = \min_V P((I_A \otimes V)\rho_{AB}(I_A \otimes V^\dagger), \sigma_{AC}).$$

Moreover, if both ρ and σ are subnormalized, by Eq. (4.1), we can bound

$$\begin{aligned} \min_V \|(I_A \otimes V)\rho_{AB}(I_A \otimes V^\dagger) - \sigma_{AC}\|_1 &\leq \min_V 2P((I_A \otimes V)\rho_{AB}(I_A \otimes V^\dagger), \sigma_{AC}) \\ &= 2P(\rho_A, \sigma_A) \\ &\leq 2\sqrt{2\|\rho_A - \sigma_A\|_1}. \end{aligned}$$

From this it follows that if σ is subnormalized and ρ has $\text{tr}[\rho] > 1$,

$$\min_V \|(I_A \otimes V)\rho_{AB}(I_A \otimes V^\dagger) - \sigma_{AC}\|_1 \leq 2\sqrt{2\text{tr}[\rho]\|\rho_A - \sigma_A\|_1}.$$

Since $\text{tr}[\rho] \leq \text{tr}[\sigma] + \|\rho - \sigma\|_1$ and $\text{tr}[\sigma] \leq 1$ we conclude that

$$\min_V \|(I_A \otimes V)\rho_{AB}(I_A \otimes V^\dagger) - \sigma_{AC}\|_1 \leq 2\sqrt{2\|\rho_A - \sigma_A\|_1 + 2\|\rho_A - \sigma_A\|_1^2}.$$

for arbitrary ρ and subnormalized σ . □

Finally, we will need a basic lemma relating tensor network states with differing background states:

Lemma 4.3. *Suppose we consider random tensor network states $\rho_{V_\partial R}$ and $\tilde{\rho}_{V_\partial R}$ with (purified) background states $\phi_{VR}, \tilde{\phi}_{VR} \in \mathcal{P}_\leq(VR)$ and projecting onto the same random tensors. Then*

$$\mathbb{E}\|\rho_{V_\partial R} - \tilde{\rho}_{V_\partial R}\|_1 \leq \|\phi_{VR} - \tilde{\phi}_{VR}\|_1.$$

Proof. Let $\phi - \tilde{\phi} = \Delta_+ - \Delta_-$ where both Δ_+ and Δ_- are positive semidefinite and are such that $\|\phi - \tilde{\phi}\|_1 = \text{tr}[\Delta_+ + \Delta_-]$. Then we can also consider the random tensor network states σ_+ and σ_- which take Δ_+ and Δ_- as background states, and by the linearity of Eq. (4.5) in the background state we have $\rho - \tilde{\rho} = \sigma_+ - \sigma_-$. By Eq. (2.21), $\mathbb{E}\sigma_\pm = \Delta_\pm$. We then estimate

$$\mathbb{E}\|\rho - \tilde{\rho}\|_1 = \mathbb{E}\|\sigma_+ - \sigma_-\|_1 \leq \mathbb{E}(\|\sigma_+\|_1 + \|\sigma_-\|_1) = \mathbb{E}\text{tr}[\sigma_+ + \sigma_-] = \text{tr}[\Delta_+ + \Delta_-] = \|\phi - \tilde{\phi}\|_1.$$

where we have used that σ_+ and σ_- are positive semidefinite and hence $\|\sigma_\pm\|_1 = \text{tr}[\sigma_\pm]$. □

With all our tools assembled, we are ready to prove the main result of this subsection. We again let $\phi_{VR} \in \mathcal{P}_\leq(VR)$ be a background state with R a purifying system, and we let $\rho_{V_\partial R}$ be the associated random tensor network state as constructed in Eq. (4.5). Let Γ_A be an arbitrary cut for the boundary region A . In Theorem 4.4, we provide a criterion to determine whether Γ_A is a minimal cut in terms of conditional entropies. Informally speaking, the following result shows that if Γ_A is a minimal cut in this sense, we can recover the system Γ_A from the boundary subsystem A , while we can recover Γ_A^c from the boundary subsystem \bar{A} . For general ϕ , Theorem 4.4 is closely related to the task of *split transfer*, see Appendix B for a discussion. The following result closely follows Proposition 18 of [DH10].

Theorem 4.4 (Recovery isometries). *Let $\phi_{VR} \in \mathcal{P}_\leq(VR)$ and let $\rho_{V_\partial R}$ be the associated random tensor network state as in Eq. (4.5). Let $\Gamma_A \in C(A)$ and suppose that*

$$H_2(\Gamma_A \setminus \Delta_A | \Gamma_A^c R)_{\phi|\phi} \geq K_1 \tag{4.8}$$

for all cuts $\Delta_A \in C(A)$ such that $\Delta_A \subsetneq \Gamma_A$ and

$$H_2(\Delta_A \setminus \Gamma_A | \Gamma_A R)_{\phi|\phi} \geq K_2 \quad (4.9)$$

for all cuts $\Delta_A \in C(A)$ such that $\Gamma_A \subsetneq \Delta_A$. Then

$$\mathbb{E} \min_{V_A, V_{\bar{A}}} \|(V_A \otimes V_{\bar{A}} \otimes I_R) \rho_{V_{\partial R}}(V_A^\dagger \otimes V_{\bar{A}}^\dagger \otimes I_R) - \phi_{VR}\|_1 = \mathcal{O}(\text{tr}[\phi]^{\frac{1}{4}} (2^{-\frac{1}{4}K_1} + 2^{-\frac{1}{4}K_2})). \quad (4.10)$$

where the minimum is over isometries $V_A : \mathcal{H}_A \rightarrow \mathcal{H}_{\Gamma_A}$ and $V_{\bar{A}} : \mathcal{H}_{\bar{A}} \rightarrow \mathcal{H}_{\Gamma_A^c}$.

Proof. Let $\sigma_{A\Gamma_A^c R}$ be the state where we have contracted along the tensors in Γ_A but not along those in Γ_A^c , and similarly let $\tau_{\bar{A}\Gamma_A R}$ be the state where we have contracted along the tensors in Γ_A^c but not along those in Γ_A . We first use [Proposition 4.1](#) to show that $\sigma_{A\Gamma_A^c R} \approx \phi_{\Gamma_A^c R}$ and $\tau_{\bar{A}\Gamma_A R} \approx \phi_{\Gamma_A R}$. Indeed, for σ we simply apply [Proposition 4.1](#) with $V_b \cap \Gamma_A$ as the set of bulk vertices, A as the set of boundary vertices and $\Gamma_A^c R$ as the reference system. This gives

$$\mathbb{E} \|\sigma_{A\Gamma_A^c R} - \phi_{\Gamma_A^c R}\|_1 = \mathcal{O}(\sqrt{\text{tr}[\phi]} 2^{-\frac{1}{2}K_1}).$$

A similar application of [Proposition 4.1](#), with $V_b \cap \Gamma_A^c$ as the set of bulk vertices, \bar{A} as the set of boundary vertices and $\Gamma_A R$ as the reference system, shows

$$\mathbb{E} \|\tau_{\bar{A}\Gamma_A R} - \phi_{\Gamma_A R}\|_1 = \mathcal{O}(\sqrt{\text{tr}[\phi]} 2^{-\frac{1}{2}K_2}).$$

We note that for any isometries $V_A : \mathcal{H}_A \rightarrow \mathcal{H}_{\Gamma_A}$ and $V_{\bar{A}} : \mathcal{H}_{\bar{A}} \rightarrow \mathcal{H}_{\Gamma_A^c}$

$$\begin{aligned} \|(V_A \otimes V_{\bar{A}} \otimes I_R) \rho(V_A^\dagger \otimes V_{\bar{A}}^\dagger \otimes I_R) - \phi_{VR}\|_1 &\leq \|(I_{\Gamma_A} \otimes V_{\bar{A}} \otimes I_R) \tau(I_{\Gamma_A} \otimes V_{\bar{A}}^\dagger \otimes I_R) - \phi_{VR}\|_1 \\ &\quad + \|\tau - (V_A \otimes I_{\bar{A}R}) \rho(V_A^\dagger \otimes I_{\bar{A}R})\|_1, \end{aligned}$$

where we have applied the triangle inequality after adding and subtracting $(I_{\Gamma_A} \otimes V_{\bar{A}} \otimes I_R) \tau(I_{\Gamma_A} \otimes V_{\bar{A}}^\dagger \otimes I_R)$, and then using the invariance of the trace norm under isometries in the second term. We use this to estimate

$$\begin{aligned} \mathbb{E} \min_{V_A, V_{\bar{A}}} \|(V_A \otimes V_{\bar{A}} \otimes I_R) \rho(V_A^\dagger \otimes V_{\bar{A}}^\dagger \otimes I_R) - \phi_{VR}\|_1 \\ \leq \mathbb{E} (\min_{V_A} \|(I_{\Gamma_A} \otimes V_{\bar{A}} \otimes I_R) \tau(I_{\Gamma_A} \otimes V_{\bar{A}}^\dagger \otimes I_R) - \phi_{VR}\|_1 + \min_{V_{\bar{A}}} \|\tau - (V_A \otimes I_{\bar{A}R}) \rho(V_A^\dagger \otimes I_{\bar{A}R})\|_1), \end{aligned} \quad (4.11)$$

where the minimum is over isometries $V_A : \mathcal{H}_A \rightarrow \mathcal{H}_{\Gamma_A}$ and $V_{\bar{A}} : \mathcal{H}_{\bar{A}} \rightarrow \mathcal{H}_{\Gamma_A^c}$. For the first term of [Eq. \(4.11\)](#), we apply [Lemma 4.2](#) to get

$$\begin{aligned} \min_{V_{\bar{A}}} \|(I_{\Gamma_A} \otimes V_{\bar{A}} \otimes I_R) \tau(I_{\Gamma_A} \otimes V_{\bar{A}}^\dagger \otimes I_R) - \phi_{VR}\|_1 &\leq 2\sqrt{2\|\tau_{\Gamma_A R} - \phi_{\Gamma_A R}\|_1 + \|\tau_{\Gamma_A R} - \phi_{\Gamma_A R}\|_1^2} \\ &\leq 2\sqrt{2} \left(\sqrt{\|\tau_{\Gamma_A R} - \phi_{\Gamma_A R}\|_1} + \|\tau_{\Gamma_A R} - \phi_{\Gamma_A R}\|_1 \right) \end{aligned}$$

and by Jensen's inequality

$$\begin{aligned} \mathbb{E} \min_{V_{\bar{A}}} \|(I_{\Gamma_A} \otimes V_{\bar{A}} \otimes I_R) \tau(I_{\Gamma_A} \otimes V_{\bar{A}}^\dagger \otimes I_R) - \phi_{VR}\|_1 \\ \leq 2\sqrt{2} \left(\sqrt{\mathbb{E} \|\tau_{\Gamma_A R} - \phi_{\Gamma_A R}\|_1} + \mathbb{E} \|\tau_{\Gamma_A R} - \phi_{\Gamma_A R}\|_1 \right) \\ = \mathcal{O}(\text{tr}[\phi]^{\frac{1}{4}} 2^{-\frac{1}{4}K_2}) \end{aligned} \quad (4.12)$$

For the second term of [Eq. \(4.11\)](#), we can think of τ and $(V_A \otimes I_{\bar{A}R}) \rho(V_A^\dagger \otimes I_{\bar{A}R})$ as the random tensor network states with ϕ and $(V_A \otimes I_{\Gamma_A^c R}) \sigma(V_A^\dagger \otimes I_{\Gamma_A^c R})$ as the full state on edges, applying random tensors in Γ_A^c . Then, denoting by $\mathbb{E}_{\Gamma_A^c}$ the expectation value over all random tensors in Γ_A^c , by [Lemma 4.3](#)

$$\mathbb{E}_{\Gamma_A^c} \|\tau - (V_A \otimes I_{\bar{A}R}) \rho(V_A^\dagger \otimes I_{\bar{A}R})\|_1 = \mathcal{O}(\|\phi_{VR} - (V_A \otimes I_{\bar{A}R}) \sigma(V_A^\dagger \otimes I_{\bar{A}R})\|_1).$$

We thus estimate

$$\mathbb{E} \min_{V_A} \|\tau - (V_A \otimes I_{\bar{A}R}) \rho(V_A^\dagger \otimes I_{\bar{A}R})\|_1 = \mathcal{O}(\mathbb{E}_{\Gamma_A} \min_{V_A} \|\phi_{VR} - (V_A \otimes I_{\bar{A}R}) \sigma(V_A^\dagger \otimes I_{\bar{A}R})\|_1)$$

for which we may argue exactly as in [Eq. \(4.12\)](#) and using [Lemma 4.2](#) that

$$\mathbb{E}_{\Gamma_A} \min_{V_A} \|\phi_{VR} - (V_A \otimes I_{\Gamma_A^c R}) \sigma(V_A^\dagger \otimes I_{\Gamma_A^c R})\|_1 = \mathcal{O}(\text{tr}[\phi]^{\frac{1}{4}} 2^{-\frac{1}{4}K_1}).$$

We conclude that

$$\mathbb{E} \min_{V_A, V_{\bar{A}}} \|(V_A \otimes V_{\bar{A}} \otimes I_R) \rho(V_A^\dagger \otimes V_{\bar{A}}^\dagger \otimes I_R) - \phi_{VR}\|_1 = \mathcal{O}(\text{tr}[\phi]^{\frac{1}{4}} (2^{-\frac{1}{4}K_1} + 2^{-\frac{1}{4}K_2})).$$

□

We hence find that the closeness of the boundary and background state can be bounded via conditional Rényi-2 entropies of cuts. In particular, for large K_1, K_2 , the recovery isometries can recover states to good accuracy, and we find that $\mathbb{E} \|\text{spec}_+(\rho_A) - \text{spec}_+(\phi_{\Gamma_A})\|$ is small. However, this result is not yet completely satisfying. The conditional Rényi-2 entropy is not a ‘robust’ quantity, in the sense that a small deformation of ϕ can drastically change the values of the conditional Rényi-2 entropies in [Eq. \(4.8\)](#) and [Eq. \(4.9\)](#). For this reason, we would like a condition with *smoothed* entropies. We first note that one can actually show that for the condition in [Eq. \(4.8\)](#), we can bound

$$H_2(\Gamma_A \setminus \Delta_A | \Gamma_A^c R)_{\phi|\phi} \geq H_{\min}(\Gamma_A \setminus \Delta_A | \Gamma_A^c R)_{\phi} \quad (4.13)$$

and similarly for [Eq. \(4.9\)](#),

$$H_2(\Delta_A \setminus \Gamma_A | \Gamma_A R)_{\phi|\phi} \geq H_{\min}(\Delta_A \setminus \Gamma_A | \Gamma_A R)_{\phi} \quad (4.14)$$

To make the condition ‘robust’, we would like to replace these by smoothed entropies and express a condition in terms of $H_{\min}^{\varepsilon}(\Gamma_A \setminus \Delta_A | \Gamma_A^c R)_{\phi}$ and $H_{\min}^{\varepsilon}(\Delta_A \setminus \Gamma_A | \Gamma_A R)_{\phi}$. This will require *simultaneous smoothing*: finding a state $\phi_{VR}^{\varepsilon} \in \mathcal{P}_{\leq}(VR)$ which is close to ϕ , such that $H_{\min}(\Gamma_A \setminus \Delta_A | \Gamma_A^c R)_{\phi^{\varepsilon}} \geq H_{\min}^{\varepsilon}(\Gamma_A \setminus \Delta_A | \Gamma_A^c R)_{\phi}$ and $H_{\min}(\Delta_A \setminus \Gamma_A | \Gamma_A R)_{\phi^{\varepsilon}} \geq H_{\min}^{\varepsilon}(\Delta_A \setminus \Gamma_A | \Gamma_A R)_{\phi}$ for all relevant cuts Δ_A . If we have a general background state, it is not known how this can be done [[DF13](#), [Dut11](#)]. However, if the background state is actually a tensor product of link states, we can perform the simultaneous smoothing.

4.3 One minimal cut

The primary result in this subsection is [Theorem 4.7](#), which states that if the background state is actually a tensor product of link states as in [Eq. \(2.2\)](#), then the spectrum of the boundary state is well-approximated by the spectrum of the minimal cut link state in expectation, where the approximation accuracy is controlled by *smooth* one-shot entropies. It is a straightforward application of [Theorem 4.4](#).

For a cut $\Gamma_A \in C(A)$ we define

$$\begin{aligned} \mathcal{C}_1(\Gamma_A) &= \{\Delta_A \in C(A) : \Delta_A \subsetneq \Gamma_A\} \\ \mathcal{C}_2(\Gamma_A) &= \{\Delta_A \in C(A) : \Gamma_A \subsetneq \Delta_A\}. \end{aligned}$$

The key result we need is the following lemma, which we prove in [Appendix C](#), which shows that if we have a link state, we can perform the desired joint smoothing.

Lemma 4.5. *Let $\phi \in \mathcal{P}_{=}(V)$ be a link state, $A \subseteq V_{\partial}$ a boundary subsystem and $\Gamma_A \in C(A)$ a cut for A . Then there exists a pure state $\phi^{\varepsilon} \in \mathcal{P}_{\leq}(V)$ which is such that*

$$P(\phi, \phi^{\varepsilon}) \leq 2 \left(\sqrt{|\mathcal{C}_1(\Gamma_A)|} + \sqrt{|\mathcal{C}_2(\Gamma_A)|} \right) \sqrt{\varepsilon}$$

and it holds that for any $\Delta_A \in \mathcal{C}_1(\Gamma_A)$

$$H_{\min}(\Gamma_A \setminus \Delta_A | \Gamma_A^c)_{\phi^\varepsilon} \geq H_{\min}^\varepsilon(\Gamma_A \setminus \Delta_A | \Gamma_A^c)_\phi$$

and for any $\Delta_A \in \mathcal{C}_2(\Gamma_A)$

$$H_{\min}(\Delta_A \setminus \Gamma_A | \Gamma_A)_{\phi^\varepsilon} \geq H_{\min}^\varepsilon(\Delta_A \setminus \Gamma_A | \Gamma_A)_\phi.$$

We now define the notion of a minimal cut for an arbitrary link state.

Definition 4.6 (Generalized minimal cut). A cut Γ_A is an (ε, K) -minimal cut if for all $\Delta_A \in \mathcal{C}_1(\Gamma_A)$

$$H_{\min}^\varepsilon(\Gamma_A \setminus \Delta_A | \Gamma_A^c)_\phi \geq K$$

and for all $\Delta_A \in \mathcal{C}_2(\Gamma_A)$

$$H_{\min}^\varepsilon(\Delta_A \setminus \Gamma_A | \Gamma_A)_\phi \geq K.$$

This definition is consistent with the smooth entropy conditions for minimal surfaces in holography from [AP20]. The following is now a straightforward consequence of Theorem 4.4 and Lemma 4.5. It justifies our notion of a generalized minimal cut, as it controls the degree to which the spectrum of the corresponding cut link state ϕ_{Γ_A} is close to the boundary state ρ_A .

Theorem 4.7. Consider a random tensor network state ρ constructed with $\phi \in \mathcal{P}_=(V)$ a tensor product of link states as in Eq. (2.2). Let A be a boundary region of the network, ρ_A the corresponding boundary state, and Γ_A an (ε, K) -minimal cut. Then the spectra of ρ_A and the state ϕ_{Γ_A} on A are related as:

$$\mathbb{E} \|\text{spec}_+(\rho_A) - \text{spec}_+(\phi_{\Gamma_A})\|_1 = \mathcal{O}(2^{-\frac{1}{4}K} + \sqrt{\varepsilon}).$$

Proof. Let ϕ^ε be a state as constructed in Lemma 4.5, and let ρ^ε be the random tensor network state using this background state. Then, by Theorem 4.4

$$\begin{aligned} \mathbb{E} \|\text{spec}_+(\rho_A^\varepsilon) - \text{spec}_+(\phi_{\Gamma_A}^\varepsilon)\|_1 &\leq \mathbb{E} \min_{V_A, V_{\bar{A}}} \|(V_A \otimes V_{\bar{A}} \otimes I_R) \rho^\varepsilon (V_A^\dagger \otimes V_{\bar{A}}^\dagger \otimes I_R) - \phi^\varepsilon\|_1 \\ &= \mathcal{O}(\text{tr}[\phi]^{-\frac{1}{4}} (2^{-\frac{1}{4}K_1} + 2^{-\frac{1}{4}K_2})) \end{aligned}$$

where K_1 is the minimal value over $\Delta_A \in \mathcal{C}_1(\Gamma_A)$ of

$$H_2(\Gamma_A \setminus \Delta_A | \Gamma_A^c)_{\phi^\varepsilon | \phi^\varepsilon} \geq H_{\min}(\Gamma_A \setminus \Delta_A | \Gamma_A^c)_{\phi^\varepsilon} - \log \text{tr}[\phi^\varepsilon] \geq H_{\min}^\varepsilon(\Gamma_A \setminus \Delta_A | \Gamma_A^c)_\phi - \log \text{tr}[\phi^\varepsilon]$$

using Eq. (4.4) and accounting for the normalization. Similarly K_2 is the minimal value over $\Delta_A \in \mathcal{C}_2(\Gamma_A)$ of

$$H_2(\Delta_A \setminus \Gamma_A | \Gamma_A)_{\phi^\varepsilon | \phi^\varepsilon} \geq H_{\min}(\Delta_A \setminus \Gamma_A | \Gamma_A)_{\phi^\varepsilon} - \log \text{tr}[\phi^\varepsilon] \geq H_{\min}^\varepsilon(\Delta_A \setminus \Gamma_A | \Gamma_A)_\phi - \log \text{tr}[\phi^\varepsilon]$$

and hence $K_1 \geq K$ and $K_2 \geq K$, so

$$\mathbb{E} \|\text{spec}_+(\rho_A^\varepsilon) - \text{spec}_+(\tilde{\phi}_{\Gamma_A}^\varepsilon)\|_1 = \mathcal{O}(2^{-\frac{1}{4}K}).$$

Moreover, $\|\phi - \phi^\varepsilon\|_1 \leq 2T(\phi, \phi^\varepsilon) \leq 2P(\phi, \phi^\varepsilon) = \mathcal{O}(\sqrt{\varepsilon})$ by Eq. (4.1) and hence by Lemma 4.3

$$\mathbb{E} \|\rho - \rho^\varepsilon\|_1 = \mathcal{O}(\sqrt{\varepsilon}).$$

We conclude that

$$\begin{aligned} \mathbb{E} \|\text{spec}_+(\rho_A) - \text{spec}_+(\phi_{\Gamma_A})\|_1 &\leq \mathbb{E} \|\text{spec}_+(\rho_A^\varepsilon) - \text{spec}_+(\phi_{\Gamma_A}^\varepsilon)\|_1 + \mathbb{E} \|\rho_A - \rho_A^\varepsilon\|_1 + \|\phi_{\Gamma_A} - \phi^\varepsilon \Gamma_A\|_1 \\ &= \mathcal{O}(2^{-\frac{1}{4}K} + \sqrt{\varepsilon}). \end{aligned}$$

□

4.4 Two competing minimal cuts

We now consider the case of two minimal cuts for A , where the link states do not have bounded spectral variation as in [Section 3](#). We cannot directly apply [Theorem 4.7](#) if there are two competing minimal cuts (note that we will need to define what this means exactly), nor can we assume that the empirical measure, or even a rescaling of it, will converge. Instead, we will see that the spectrum can be approximated in trace distance in a way that allows one to compute entropies, as well as showing convergence of certain measures depending on the spectrum of the reduced state. In particular, under these regularity conditions, we will prove the main result of this subsection: the distribution defined by the boundary state converges to the measure defined by the pushforward along the min-function acting on the distributions of the two competing minimal cuts. The intuition behind this is that the state on the edges can be approximated by a superposition of two states which both do have a unique minimal cut. As a corollary of this result, we will then show how to approximate the von Neumann entropy of the boundary state in such a situation.

We begin by introducing the relevant family of probability distributions for our purposes. Given a probability distribution vector $p \in \mathbb{R}^d$ of d outcomes and $f(d) > 0$ and $g(d) \in \mathbb{R}$, consider the random variable which takes value $f(d)[- \log(p_i) - g(d)]$ with probability p_i . It has distribution

$$\nu_p = \sum_{i=1}^d p_i \delta_{f(d)[- \log(p_i) - g(d)]}. \quad (4.15)$$

Typically, we will have families of probability distributions with increasing d , let $f(d) = (\log d)^{-1}$ and choose $g(d)$ such that it corresponds to the entropy of p . We will also write $\nu_p = \nu_{\text{spec}(\rho)}$ for a quantum state ρ . We may also consider [Eq. \(4.15\)](#) for the case where $p \in \mathbb{R}^d$ is positive but normalized, in which case ν_p is a finite measure. To motivate the study of the distribution in [Eq. \(4.15\)](#), we note that it is closely related to the central limit theorem. Let $d = d_0^n$ and $\rho = \rho_0^{\otimes n}$ for some state ρ_0 on \mathbb{C}^{d_0} . Then

$$\nu_\rho^{(n)} = \sum_{\lambda \in \text{spec}(\rho_0^{\otimes n})} \lambda \delta_{\frac{1}{\sqrt{n}}[- \log(\lambda) - nH(\rho_0)]}$$

converges in distribution to a normal distribution by the central limit theorem. We study this particular measure because the empirical measures we investigated in [Section 3](#) may not have good convergence properties if, for example, the full state on edges consists of many copies of a single non-maximally entangled link state. Moreover, the measure in [Eq. \(4.15\)](#) clearly captures information about the entropy of the probability distribution, and turns out to capture information about second order asymptotics of information processing tasks [[Hay08](#), [TH13](#)]. In fact, the example of many independent copies of a single link state is not the only relevant situation. In [[CHLS15](#), [BPSW19](#)], it was argued that the entanglement spectrum of conformal field theories with large central charge (which are the motivating example to study random tensor networks) have similar behavior.

Before diving into more technical details, we can build some intuition for how distributions of the form [Eq. \(4.15\)](#) will behave in the two-cut setting. Consider the situation with a *single* random tensor. Let us take a single bulk vertex x and two boundary vertices a and b , with edges (xa) and (xb) . There is a single random tensor at x , and we take link states $|\phi_1\rangle^{\otimes n}$ along (xa) and $|\phi_2\rangle^{\otimes n}$ along (xb) . We denote

$$|\phi_j\rangle = \sum_i \sqrt{\lambda_{i,j}} |ii\rangle$$

for $j = 1, 2$ and we let $h_j := H(\{\lambda_{i,j}\})$ be the entanglement entropies along the two minimal link states. There are two separating cuts for the boundary subsystem $\{a\}$: if $h_1 < h_2$ then (xa) is the minimal cut, and if $h_1 > h_2$, the minimal cut is given by (xb) . In these cases, for large n , the entanglement spectrum of $\rho_{\{a\}}$ can be approximated by the entanglement spectrum along the minimal cut. What happens at the ‘phase transition’ where $h_1 = h_2 = h$? The intuition is that we can split $|\phi_1\rangle^{\otimes n} \otimes |\phi_2\rangle^{\otimes n}$ into a superposition of the two states, one for which the minimal surface is at (xa) and one for which the minimal surface is at (xb) . We

find that in this case, if we let $\sigma_j^2 = \sum_i \lambda_{i,j} (\log(\frac{1}{\lambda_{i,j}}) - h)^2$, then for any bounded continuous function f , the quantity

$$\sum_{\lambda \in \text{spec}(\rho_{\{a\}})} \lambda f\left(\frac{-\log(\lambda) - nh}{\sqrt{n}}\right)$$

converges to

$$\frac{1}{2\pi\sigma_1\sigma_2} \int \int e^{-\frac{(x_1-h)^2}{2\sigma_1^2} - \frac{(x_2-h)^2}{2\sigma_2^2}} f(\min(x_1, x_2)) dx_1 dx_2$$

in probability as n goes to infinity. Our goal in this section will be to prove a general version of this result for full random tensor networks.

In [Section 3](#), we used the method of moments (if all moments of a distribution converge, and the distribution is uniquely determined by its moments, then we have weak convergence). However, convergence of moments is a stronger condition than weak convergence, and it requires computation of all moments. In this section, we will use that for distributions of the form [Eq. \(4.15\)](#), convergence in distribution follows from convergence in trace norm:

Lemma 4.8. *Consider a sequence of increasing integers $\{d_n\}_{n \in \mathbb{N}}$ and for each n , positive vectors $p^{(n)}, q^{(n)} \in \mathbb{R}^{d_n}$. Let $\sigma, h : \mathbb{N} \rightarrow \mathbb{R}$ be such that $\sigma(d) > 0$ for $d \in \mathbb{N}$ and $\lim_{d \rightarrow \infty} \sigma(d)^{-1} = 0$. Let*

$$\nu_p^{(n)} := \sum_{i=1}^{d_n} p_i^{(n)} \delta_{\frac{1}{\sigma(d_n)}[-\log(p_i^{(n)}) - h(d_n)]}$$

and similarly let

$$\nu_q^{(n)} = \sum_{i=1}^{d_n} q_i^{(n)} \delta_{\frac{1}{\sigma(d_n)}[-\log(q_i^{(n)}) - h(d_n)]}.$$

1. Suppose $\|p^{(n)} - q^{(n)}\|_1$ goes to zero as n goes to infinity, and $\nu_p^{(n)} \Rightarrow \nu$ for some probability distribution ν . Then $\nu_q^{(n)} \Rightarrow \nu$.
2. Suppose the vectors $q^{(n)}$ and $p^{(n)}$ are random, $\mathbb{E}\|p^{(n)} - q^{(n)}\|_1$ goes to zero as n goes to infinity, and $\nu_p^{(n)} \Rightarrow \nu$, in probability. Then $\nu_q^{(n)}$ converges weakly, in probability, to ν .

Proof. We need to show that

$$\left| \int f(x) d\nu_q^{(n)}(x) - \int f(x) d\nu(x) \right| \rightarrow 0$$

for any $f \in C_b(\mathbb{R})$. In fact, it suffices to show this for f uniformly continuous (see for instance [Theorem C.10](#) in [\[AGZ10\]](#)). Let $\varepsilon > 0$, then if we assume f to be uniformly continuous, there exists $\delta > 0$ such that for any x, y for which $|x - y| \leq \delta$, it holds that $|f(x) - f(y)| \leq \varepsilon$. We use a triangle inequality to bound

$$\begin{aligned} \left| \int f(x) d\nu_q^{(n)}(x) - \int f(x) d\nu(x) \right| &\leq \left| \int f(x) d\nu_q^{(n)}(x) - \int f(x) d\nu_p^{(n)}(x) \right| \\ &\quad + \left| \int f(x) d\nu_p^{(n)}(x) - \int f(x) d\nu(x) \right|. \end{aligned} \tag{4.16}$$

Since $\nu_p^{(n)} \Rightarrow \nu$, the second term on the right hand side vanishes as n goes to infinity. If we write $g_n(x) = \frac{1}{\sigma(d_n)} (\log \frac{1}{x} - h(d_n))$, then the first term on the right hand side of [Eq. \(4.16\)](#) is given by

$$\left| \sum_{i=1}^{d_n} q_i^{(n)} f(g_n(q_i^{(n)})) - p_i^{(n)} f(g_n(p_i^{(n)})) \right|.$$

This can be bounded by

$$\sum_{i=1}^{d_n} |q_i^{(n)} f(g_n(q_i^{(n)})) - p_i^{(n)} f(g_n(p_i^{(n)}))| \leq \sum_{i=1}^{d_n} |q_i^{(n)} - p_i^{(n)}| |f(g_n(q_i^{(n)}))| + \sum_{i=1}^{d_n} p_i^{(n)} |f(g_n(q_i^{(n)})) - f(g_n(p_i^{(n)}))|$$

In this expression, the first term is bounded by

$$\sum_{i=1}^{d_n} |q_i^{(n)} - p_i^{(n)}| |f(g_n(q_i^{(n)}))| \leq C \|q^{(n)} - p^{(n)}\|_1 \quad (4.17)$$

where C is a uniform upper bound for f (using $f \in C_b(\mathbb{R})$). For the second term, we partition the sum over $[d_n]$ into three sets

$$\begin{aligned} I_1 &= \{i \in [d_n] \text{ such that } 2^{-\delta\sigma(d_n)} \leq \frac{p_i^{(n)}}{q_i^{(n)}} \leq 2^{\delta\sigma(d_n)}\} \\ I_2 &= \{i \in [d_n] \text{ such that } \frac{p_i^{(n)}}{q_i^{(n)}} > 2^{\delta\sigma(d_n)}\} \\ I_3 &= \{i \in [d_n] \text{ such that } \frac{p_i^{(n)}}{q_i^{(n)}} < 2^{-\delta\sigma(d_n)}\}. \end{aligned}$$

The idea is that for $i \in I_1$, p_i and q_i are sufficiently close that we may use the continuity of f , whereas I_2 and I_3 cannot have too much weight (using that $p^{(n)}$ and $q^{(n)}$ are close in trace distance). Let us now make this precise. For the sum over the elements in I_1 , we have

$$\sum_{i \in I_1} p_i^{(n)} |f(g_n(q_i^{(n)})) - f(g_n(p_i^{(n)}))| \leq \sum_{i \in I_1} p_i^{(n)} \varepsilon \leq \|p^{(n)}\|_1 \varepsilon, \quad (4.18)$$

using the uniform continuity of f and the fact that for $i \in I_1$, we have $g(q_i^{(n)}) - g(p_i^{(n)}) = \frac{1}{\sigma(d_n)} \log\left(\frac{p_i^{(n)}}{q_i^{(n)}}\right)$, implying $|g(q_i^{(n)}) - g(p_i^{(n)})| \leq \delta$ by the definition of I_1 . Next, we observe that

$$\|p^{(n)} - q^{(n)}\|_1 \geq \sum_{i \in I_2} |p_i^{(n)} - q_i^{(n)}| = \sum_{i \in I_2} p_i^{(n)} \left(1 - \frac{q_i^{(n)}}{p_i^{(n)}}\right) \geq \sum_{i \in I_2} p_i^{(n)} (1 - 2^{-\delta\sigma(d_n)}),$$

and hence for $\sigma(d_n) \geq \delta^{-1}$, we have

$$\sum_{i \in I_2} p_i^{(n)} \leq 2 \|p^{(n)} - q^{(n)}\|_1. \quad (4.19)$$

In analogous fashion we see that

$$\|p^{(n)} - q^{(n)}\|_1 \geq \sum_{i \in I_3, p_i^{(n)} > 0} |p_i^{(n)} - q_i^{(n)}| = \sum_{i \in I_3, p_i^{(n)} > 0} p_i^{(n)} \left(\frac{q_i^{(n)}}{p_i^{(n)}} - 1\right) \geq \sum_{i \in I_3} p_i^{(n)} (2^{\delta\sigma(d_n)} - 1),$$

so for $\sigma(d_n) \geq \delta^{-1}$ we have

$$\sum_{i \in I_3} p_i^{(n)} \leq \|p^{(n)} - q^{(n)}\|_1. \quad (4.20)$$

In conclusion, for $\sigma(d_n) \geq \delta^{-1}$, collecting Eq. (4.18), Eq. (4.19), Eq. (4.20) and using that f is uniformly bounded by C , we can bound

$$\sum_{i=1}^{d_n} p_i^{(n)} |f(g_n(q_i^{(n)})) - f(g_n(p_i^{(n)}))| \leq \varepsilon \|p^{(n)}\|_1 + 3C \|p^{(n)} - q^{(n)}\|_1.$$

Together with Eq. (4.17), this implies that for $\sigma(d_n) \geq \delta$, we obtain the bound:

$$\left| \int f(x) d\nu_q^{(n)}(x) - \int f(x) d\nu_p^{(n)}(x) \right| \leq 4C \|q^{(n)} - p^{(n)}\|_1 + \varepsilon \|p^{(n)}\|_1. \quad (4.21)$$

Since $\nu_p^{(n)} \Rightarrow \nu$, $\|p^{(n)}\|_1 \rightarrow 1$, and since $\varepsilon > 0$ was arbitrary, we conclude that as n goes to infinity

$$\left| \int f(x) d\nu_q^{(n)}(x) - \int f(x) d\nu(x) \right| \rightarrow 0$$

proving 1.

To prove 2, we note that by Markov's inequality, it suffices to show that

$$\mathbb{E} \left| \int f(x) d\nu_q^{(n)}(x) - \int f(x) d\nu(x) \right| \rightarrow 0, \quad (4.22)$$

where $f \in C_b(\mathbb{R})$ is a uniformly continuous function. By Eq. (4.16), it suffices to show

$$\mathbb{E} \left| \int f(x) d\nu_q^{(n)}(x) - \int f(x) d\nu_p^{(n)}(x) \right| \rightarrow 0.$$

Let $\varepsilon > 0$, and let δ such that if $|x - y| \leq \delta$, $|f(x) - f(y)| \leq \varepsilon$. Then by Eq. (4.21), for $\sigma(d_n) \geq \delta^{-1}$,

$$\mathbb{E} \left| \int f(x) d\nu_q^{(n)}(x) - \int f(x) d\nu_p^{(n)}(x) \right| \leq 4C \mathbb{E} \|q^{(n)} - p^{(n)}\|_1 + \varepsilon \mathbb{E} \|p^{(n)}\|_1.$$

Since $\varepsilon > 0$ was arbitrary, $\mathbb{E} \|q^{(n)} - p^{(n)}\|_1$ goes to zero, and $\mathbb{E} \|p^{(n)}\|_1$ goes to one, we conclude that Eq. (4.22) holds, proving 2. \square

The value of this result is clear: so long as we restrict ourselves to measures of the form Eq. (4.15), then we can prove convergence results by proving convergence of trace norms.

We will also need a basic lemma to help estimate overlaps of states.

Lemma 4.9. *Suppose we have two bipartite pure states $\psi_{AB}, \phi_{AB} \in \mathcal{P}(AB)$. Then*

$$\|\mathrm{tr}_B[|\psi\rangle\langle\phi|]\|_1 = F(\psi_B, \phi_B).$$

Proof. For any operator M_A on \mathcal{H}_A it holds that

$$\|M_A\|_1 = \sup_{U_A} |\mathrm{tr}[U_A M_A]|$$

where the supremum is over unitary operators U_A on \mathcal{H}_A , so

$$\|\mathrm{tr}_B[|\psi\rangle\langle\phi|]\|_1 = \sup_{U_A} |\mathrm{tr}[U_A \mathrm{tr}_B[|\psi\rangle\langle\phi|]]| = \sup_{U_A} |\mathrm{tr}[U_A \otimes I_B |\psi\rangle\langle\phi|]| = \sup_{U_A} |\langle\phi| U_A \otimes I_B |\psi\rangle|$$

which equals $F(\psi_B, \phi_B)$ by Uhlmann's theorem. \square

With these tools in hand, let us now return to the random tensor network setting. Let $\Gamma_{A,1}$ and $\Gamma_{A,2}$ be two cuts with associated sets of edges $\gamma_{A,1}$ and $\gamma_{A,2}$. We will assume these two cuts are non-intersecting: $\gamma_{A,1} \cap \gamma_{A,2} = \emptyset$, and, without loss of generality $\Gamma_{A,1} \subset \Gamma_{A,2}$. We will assume that the full state on edges is a tensor product of link states along the edges, and that the associated central limit measure for the spectrum along the cuts $\gamma_{A,1}$ and $\gamma_{A,2}$ converges to a continuous probability distribution. The most obvious application is where the link states on each edge ϕ_e are many copies of some single state $\phi_{e,0}$, so $\phi_e = (\phi_{e,0})^{\otimes n}$, in which case the spectrum is subject to a central limit theorem and the assumptions of [Theorem 4.12](#) are satisfied.

We will now formalize what it means for these two cuts to be competing minimal cuts. We have fixed $\Gamma_{A,1}$ and $\Gamma_{A,2}$ with $\Gamma_{A,1} \subsetneq \Gamma_{A,2}$. Then we consider three sets of cuts: $\mathcal{C}_1(\Gamma_{A,1}, \Gamma_{A,2})$ is the collection of cuts strictly contained in $\Gamma_{A,1}$, $\mathcal{C}_2(\Gamma_{A,1}, \Gamma_{A,2})$ is the collection of cuts which strictly contain $\Gamma_{A,2}$, and $\mathcal{C}_3(\Gamma_{A,1}, \Gamma_{A,2})$ is the collection of cuts which are ‘in between’ $\Gamma_{A,1}$ and $\Gamma_{A,2}$. That is,

$$\begin{aligned}\mathcal{C}_1(\Gamma_{A,1}, \Gamma_{A,2}) &= \{\Delta_A \in C(A) : \Delta_A \subsetneq \Gamma_{A,1}\} \\ \mathcal{C}_2(\Gamma_{A,1}, \Gamma_{A,2}) &= \{\Delta_A \in C(A) : \Gamma_{A,2} \subsetneq \Delta_A\} \\ \mathcal{C}_3(\Gamma_{A,1}, \Gamma_{A,2}) &= \{\Delta_A \in C(A) : \Gamma_{A,1} \subsetneq \Delta_A \subsetneq \Gamma_{A,2}\}\end{aligned}$$

and we let $\mathcal{C} = \mathcal{C}_1(\Gamma_{A,1}, \Gamma_{A,2}) \cup \mathcal{C}_2(\Gamma_{A,1}, \Gamma_{A,2}) \cup \mathcal{C}_3(\Gamma_{A,1}, \Gamma_{A,2})$. As in the single cut case, we would like to say that the surfaces $\Gamma_{A,1}$ and $\Gamma_{A,2}$ are minimal cuts if an appropriate set of conditional entropies are sufficiently large for all $\Delta_A \in \mathcal{C}$. We formalize this as follows:

Definition 4.10. A pair of cuts $\Gamma_{A,1}$ and $\Gamma_{A,2}$ for A is a *pair of (ε, K) -minimal cuts* if

$$H_{\min}^\varepsilon(\Gamma_{A,1} \setminus \Delta_A | \Gamma_{A,1}^c) \geq K$$

for all cuts $\Delta_A \in \mathcal{C}_1(\Gamma_{A,1}, \Gamma_{A,2})$

$$H_{\min}^\varepsilon(\Delta_A \setminus \Gamma_{A,2} | \Gamma_{A,2}) \geq K$$

for all cuts $\Delta_A \in \mathcal{C}_2(\Gamma_{A,1}, \Gamma_{A,2})$ and

$$H_{\min}^\varepsilon(\Delta_A \setminus \Gamma_{A,1} | \Gamma_{A,1}) \geq K$$

$$H_{\min}^\varepsilon(\Gamma_{A,2} \setminus \Delta_A | \Gamma_{A,2}^c) \geq K$$

for all cuts $\Delta_A \in \mathcal{C}_3(\Gamma_{A,1}, \Gamma_{A,2})$.

We will need a joint smoothing result similar to [Lemma 4.5](#) for the particular case where $\Gamma_{A,1} = A$ and $\Gamma_{A,2}^c = \bar{A}$. To this end we consider a graph $G = (V, E)$ with a set of boundary vertices V_∂ , and a boundary subsystem $A \subseteq V_\partial$. We denote by $\gamma_{A,1}$ the edges incident to A , and $\gamma_{A,2}$ the edges incident to \bar{A} , and assume these sets do not intersect. We let $E_b = E \setminus (\gamma_{A,1} \cup \gamma_{A,2})$. For a cut $\Delta_A \in C(A)$ we let Y^{Δ_A} be the set of half-edges

$$Y^{\Delta_A} = \{(e, x) : e = (xy), x \in \Delta_A^c, y \in A\}.$$

Lemma 4.11. Let $\phi \in \mathcal{P}_{\leq}(V)$ be a pure background state which is of the form $\phi = \phi_{\gamma_{A,1}} \otimes \phi_{\gamma_{A,2}} \otimes \phi_{E_b}$, where

$$\phi_{E_b} = \bigotimes_{e \in E_b} \phi_e$$

is a product state and the $\phi_{\gamma_{A,i}}$ have a Schmidt decomposition in the standard basis along the half-edges. Then there exists a state ϕ^ε which is such that

$$P(\phi, \phi^\varepsilon) \leq 2\sqrt{2^{V_b}\varepsilon}$$

and for all cuts $\Delta_A \in C(A)$ it holds that

$$H_{\min}(\Delta_A \setminus A | A)_{\phi^\varepsilon} \geq H_{\min}^\varepsilon(\Delta_A \setminus A | AY^{\Delta_A})_\phi.$$

To state the main result of this section, recall that given a measurable function $f : \mathcal{X} \rightarrow \mathcal{Y}$ between measure spaces, the *pushforward* of the function f on a measure μ on \mathcal{X} is defined by $(f_*(\mu))(A) = \mu(f^{-1}(A))$ for any measurable set $A \subseteq \mathcal{Y}$. We apply this to the function $\min : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ in the result:

Theorem 4.12. *Consider a family of random tensor network states on a graph G with pure state on edges ϕ , indexed by an increasing sequence of positive integers n . We assume that $\Gamma_{A,1}, \Gamma_{A,2} \in C(A)$ are a pair of nonintersecting $(\varepsilon(n), K(n))$ -minimal cuts for all n where $\varepsilon(n) = \mathcal{O}(n^{-\gamma})$ for $\gamma > 4$ and $K(n) = \Omega(\log(n)^2)$ as n goes to infinity. We let $H : \mathbb{N} \rightarrow \mathbb{R}$, and we assume that*

$$\nu_{\phi_{\Gamma_{A,i}}} = \sum_{\lambda \in \text{spec}(\phi_{\Gamma_{A,i}})} \lambda \delta_{\frac{1}{\sqrt{n}}[\log(\frac{1}{\lambda}) - H(n)]}$$

for $i = 1, 2$, is such that $\nu_{\phi_{\Gamma_{A,i}}} \Rightarrow \nu_i$, where ν_i is a probability distribution with a continuous cumulative distribution function. Then $\nu_{\rho_A} \Rightarrow \min_*(\nu_1, \nu_2)$ in probability.

Proof. Proving this result will require several intermediate results. We provide a very high-level, enumerated sketch of our proof here, involving the following steps:

1. Study a reduced problem on a subnetwork; this subnetwork is such that the minimal cuts $\gamma_{A,i}$ are incident to the boundary.
2. Approximate the background state with superpositions of maximally entangled states by binning eigenvalues, and construct approximate tensor network states with the approximate background states.
3. Prove that the spectrum of the boundary state converges to the spectrum of the approximate tensor network state, which in turn converges to the spectrum of the approximate background states. We write the background state as a superposition of two states, one of which has $\gamma_{A,1}$ as its minimal cut, whereas the other state has $\gamma_{A,2}$ as its minimal cut. We show that the resulting states are approximately orthogonal.
4. Show that the distribution of the approximate background states converges to the min-distribution, and hence conclude that the spectrum of the boundary state converges to the min-distribution.

Figure 5 provides a more detailed visual sketch of the intuition behind our proof strategy. Lemma 4.8 will be a key tool, as it implies that it will suffice to show convergence in trace norm.

We assume without loss of generality that $\Gamma_{A,1} \subset \Gamma_{A,2}$. We now define the subnetwork that we will analyze in our proof. Let $G' = (V', E')$ be the induced graph on $V' = V'_b \cup V'_\partial$, where $V'_b = \Gamma_{A,2} \setminus \Gamma_{A,1}$, and $V'_\partial = B \cup \bar{B}$, with B the set of vertices in $\Gamma_{A,1}$ which are incident to V'_b and \bar{B} the set of vertices in $V \setminus \Gamma_{A,2}$ which are incident to V'_b . The subgraph G' is depicted in Figure 5b.


We also define the random tensor network state τ as the state obtained by applying random tensors only on bulk vertices in the complement of V'_b . Then for this state, by a slight variation on Theorem 4.7, it holds that

$$\mathbb{E} \min_{V_A, V_{\bar{A}}} \|(V_A \otimes V_{\bar{A}} \otimes I)\tau(V_A^\dagger \otimes V_{\bar{A}}^\dagger \otimes I) - \phi\|_1 = \mathcal{O}(2^{-K(n)} + \sqrt{\varepsilon(n)}),$$


where the minimum is over isometries $V_A : \mathcal{H}_A \rightarrow \mathcal{H}_{\Gamma_{A,1}}$ and $V_{\bar{A}} : \mathcal{H}_{\bar{A}} \rightarrow \mathcal{H}_{\Gamma_{A,2}^c}$. Let $\tilde{\tau} = (V_A \otimes V_{\bar{A}} \otimes I)\tau(V_A^\dagger \otimes V_{\bar{A}}^\dagger \otimes I)$ where the V_A and $V_{\bar{A}}$ are the isometries that realize the minimum.

Now, consider two random tensor network states on G' obtained by applying the same random tensors on V'_b with background states $\tilde{\tau}$ and ϕ respectively. The state where we take $\tilde{\tau}$ as the background state yields $(V_A \otimes V_{\bar{A}} \otimes I)\rho(V_A^\dagger \otimes V_{\bar{A}}^\dagger \otimes I)$. Denote the state where we take ϕ as the background state by σ . Then by Lemma 4.3:


$$\mathbb{E}\|(V_A \otimes V_{\bar{A}})\rho(V_A^\dagger \otimes V_{\bar{A}}^\dagger) - \sigma\|_1 = \mathcal{O}(2^{-K(n)} + \sqrt{\varepsilon(n)}).$$





(a) Consider a tensor network with a boundary region A with two competing minimal cuts.




(b) We show that the spectrum is almost preserved when removing the boundary regions; that is, the spectrum of ρ_A is approximated by that of σ_B .




(c) The distribution of the logarithms of the eigenvalues has width of order \sqrt{n} . We approximate the spectrum of the state along the two minimal cuts by a superposition of maximally entangled states and discard the tails of the spectrum.




(d) We then discard the part of the state where the maximally entangled states along the two cuts are of almost equal dimension to get the approximate state on edges $|\bar{\phi}\rangle$.



(e) We decompose the background state as a superposition over $|\phi^{(j,a)}\rangle$ and $|\phi^{(j,b)}\rangle$. We denote the corresponding tensor network states by $\sigma^{(j,a)}$ and $\sigma^{(j,b)}$.



(f) For the state $|\phi^{(j,a)}\rangle$, the minimal cut is $\gamma_{A,1}$, and along $\gamma_{A,2}$ we have a maximally entangled state, which gives $\sigma_B^{(j,a)} \approx \phi_B^{(j,a)}$. Similarly, for $|\phi^{(j,b)}\rangle$, the minimal cut is $\gamma_{A,2}$ and we get $\sigma_B^{(j,b)} \approx \phi_B^{(j,b)}$.



(g) All the $\sigma_B^{(j,a)}$ have the same eigenbasis, so we can add the spectra. We can do the same for the $\sigma_B^{(j,b)}$, yielding $\sigma_B^{(a)}$ and $\sigma_B^{(b)}$, which has the same spectrum as σ_B . Finally we argue that the $\sigma_B^{(a)}$ and $\sigma_B^{(b)}$ are nearly orthogonal, so we can combine the spectra, as in Eq. (4.37). We complete the proof by showing that the resulting spectrum converges to the pushforward measure of the min function applied to the spectra of the two cuts.

Figure 5: Illustration of the proof of Theorem 4.12.

If we let E_1 denote the set of edges (xy) for which both $x, y \in \Gamma_{A,1}$, and E_2 the set of edges (xy) for which both $x, y \in \Gamma_{A,2}^c$, then $\sigma_{\Gamma_{A,1}} = \phi_{E_1} \otimes \sigma_B$ and $\sigma_{\Gamma_{A,2}} = \phi_{E_2} \otimes \sigma_{\bar{B}}$. This shows that

$$\mathbb{E} \|\text{spec}_+(\rho_A) - \text{spec}_+(\sigma_B)\|_1 = \mathcal{O}(2^{-K(n)} + \sqrt{\varepsilon(n)}) \quad (4.23)$$

and $\text{spec}_+(\sigma_B) = \text{spec}_+(\sigma_{\bar{B}})$, because σ is pure, which follows because its background state ϕ is pure. We will continue to study σ on the reduced graph G' , as in Figure 5b, and at the end of the proof, we will see that Eq. (4.23) will be sufficient to prove the desired properties of $\text{spec}(\rho_A)$.

We have accomplished Step 1 by reducing the full network to a subnetwork, and we now try to construct an approximation to σ . We will do so by coarse-graining the spectrum of the background states along the two minimal cuts: slicing the tails of the distribution, binning the remaining eigenvalues, throwing away the

smallest bins of the binned distribution, and then approximating the states as a superposition of maximally entangled states defined on the bins. Consider the background states along the two minimal cuts in the Schmidt basis:

$$|\phi_{\gamma_{A,i}}\rangle = \sum_J \sqrt{\lambda_{i,J}} |JJ\rangle. \quad (4.24)$$

Here, the $|JJ\rangle$ are a tensor product basis along the half-edges (which we again may take to be the standard basis), so that

$$|JJ\rangle = \bigotimes_{e=(xy)\in\gamma_{A,i}} |i_{e,x}\rangle \otimes |i_{e,y}\rangle,$$

where the $|i_{e,x}\rangle$ and $|i_{e,y}\rangle$ are a basis for $\mathcal{H}_{e,x}$ and $\mathcal{H}_{e,y}$. First, we truncate the allowed range of eigenvalues by slicing off the tails of the distribution, as in the left side of [Figure 5c](#). Let

$$I_n = [2^{-H(n)-C\sqrt{n}\log(n)}, 2^{-H(n)+C\sqrt{n}\log(n)}]$$

for some constant C , and let

$$|\phi_{\gamma_{A,i}}^{(1)}\rangle = \sum_{\lambda_{i,J}\in I_n} \sqrt{\lambda_{i,J}} |JJ\rangle.$$

Note that the entanglement spectrum of $\phi_{\gamma_{A,i}}$ is given by the spectrum of $\phi_{\Gamma_{A,i}}$. This implies that

$$\sum_{\lambda_{i,J}\notin I_n} \lambda_{i,J} = \nu_{\phi_{\Gamma_{A,i}}}((-\infty, -C\log(n)) \cup (C\log(n), \infty)) \rightarrow 0$$

and hence

$$\|\phi_{\gamma_{A,i}} - \phi_{\gamma_{A,i}}^{(1)}\|_1 \rightarrow 0.$$

Next, for $\lambda_{i,J} \in I_n$, we define new $\tilde{\lambda}_{i,J}$ according to $\log(\tilde{\lambda}_{i,J}) = \frac{1}{n^\alpha} \lfloor n^\alpha \log(\lambda_{i,J}) \rfloor$, effectively binning the values of $\log(\lambda_{i,J})$ into intervals of size $n^{-\alpha}$, as in the right side of [Figure 5c](#). We will choose $\alpha > 0$ small (but other choices will be useful for [Corollary 4.14](#)), to be precise, we choose $\alpha < \frac{1}{8}(\gamma - 4)$. We can now define the background state

$$|\phi_{\gamma_{A,i}}^{(2)}\rangle = \sum_{\lambda_{i,J}\in I_n} \sqrt{\tilde{\lambda}_{i,J}} |JJ\rangle.$$

Then by [Lemma C.4](#)

$$\|\phi_{\gamma_{A,i}}^{(1)} - \phi_{\gamma_{A,i}}^{(2)}\|_1 \leq 2 \sqrt{\sum_J |\lambda_{i,J} - \tilde{\lambda}_{i,J}|},$$

which we may bound using the fact that $2^{-\frac{1}{n^\alpha}} \leq \frac{\tilde{\lambda}_{i,J}}{\lambda_{i,J}} \leq 1$, so

$$\sum_J |\lambda_{i,J} - \tilde{\lambda}_{i,J}| = \sum_J \lambda_{i,J} \left(1 - \frac{\tilde{\lambda}_{i,J}}{\lambda_{i,J}}\right) \leq \max_J \left(1 - \frac{\tilde{\lambda}_{i,J}}{\lambda_{i,J}}\right) \leq 1 - 2^{-\frac{1}{n^\alpha}} = \mathcal{O}\left(\frac{1}{n^\alpha}\right).$$

Thus, $\|\phi_{\gamma_{A,i}}^{(1)} - \phi_{\gamma_{A,i}}^{(2)}\|_1 = \mathcal{O}(n^{-\frac{1}{2}\alpha})$. Since the interval I_n has length $\mathcal{O}(\sqrt{n}\log(n))$ and the distinct values $\tilde{\lambda}_{i,J}$ are $n^{-\alpha}$ apart, the $\tilde{\lambda}_{i,J}$ take $\mathcal{O}(n^{\alpha+\frac{1}{2}}\log(n))$ different values. Denote these values by $p_{i,j}$, and let $d_{i,j}$ be their multiplicities. Setting

$$q_{i,j} = d_{i,j} p_{i,j},$$

we can rewrite the state $|\phi_{\gamma_{A,i}}^{(2)}\rangle$ using the collected eigenvalues:

$$|\phi_{\gamma_{A,i}}^{(2)}\rangle = \sum_j \sqrt{q_{i,j}} |\Phi_{i,j}^+\rangle,$$

where the $|\Phi_{i,j}^+\rangle$ are maximally entangled states of dimension $d_{i,j}$ which are orthogonal, i.e. $\langle \Phi_{i,j}^+ | \Phi_{i,k}^+ \rangle = \delta_{j,k}$. Note that $0 \leq q_{i,j} \leq 1$ and $\sum_j q_{i,j} \leq 1$. Now, we discard any bins that are too small: let $\beta > \alpha + \frac{1}{2}$ and consider the state

$$|\phi_{\gamma_{A,i}}^{(3)}\rangle = \sum_{q_{i,j} > n^{-\beta}} \sqrt{q_{i,j}} |\Phi_{i,j}^+\rangle \quad (4.25)$$

where we restricted the sum to terms for which $q_{i,j}$ is sufficiently large (and hence $p_{i,j}$ will be sufficiently close to $d_{i,j}^{-1}$). Then, since the number of terms is $\mathcal{O}(n^{\alpha + \frac{1}{2}} \log(n))$ and using [Lemma C.4](#), we have

$$\|\phi_{\gamma_{A,i}}^{(2)} - \phi_{\gamma_{A,i}}^{(3)}\|_1 = \mathcal{O}\left(n^{\frac{1}{2}\alpha + \frac{1}{4} - \frac{1}{2}\beta} \sqrt{\log(n)}\right)$$

To summarize, $\phi_{\gamma_{A,i}}^{(3)}$ is the state obtained from the original background state $\phi_{\gamma_{A,i}}$ by 1) removing the tails of the spectrum, 2) binning the eigenvalues, and finally 3) dropping any of the bins that are too small. The first approximation incurs an error $\|\phi_{\gamma_{A,i}} - \phi_{\gamma_{A,i}}^{(1)}\|_1$, which converges to zero, and the second and third approximations incur errors of order $\mathcal{O}(n^{-\frac{1}{2}\alpha})$ and $\mathcal{O}\left(n^{\frac{1}{2}\alpha + \frac{1}{4} - \frac{1}{2}\beta} \sqrt{\log(n)}\right)$ respectively.

Now, let $\tilde{\phi}$ be the background state for which we have replaced $\phi_{\gamma_{A,i}}$ by $\phi_{\gamma_{A,i}}^{(3)}$ for $i = 1, 2$. Recall that σ is the state on the random tensor network state constructed with background state ϕ on the subgraph G' , as in [Figure 5b](#). We then define the approximation $\tilde{\sigma}$ to be the tensor network state on G' which uses $\tilde{\phi}$ as its background state instead of ϕ . With these background states, we see

$$\|\phi - \tilde{\phi}\|_1 = \mathcal{O}\left(\|\phi_{\gamma_{A,i}} - \phi_{\gamma_{A,i}}^{(1)}\|_1 + n^{-\frac{1}{2}\alpha} + n^{\frac{1}{2}\alpha + \frac{1}{4} - \frac{1}{2}\beta} \sqrt{\log(n)}\right) \rightarrow 0. \quad (4.26)$$

We then apply [Lemma 4.3](#) to find:

$$\mathbb{E}\|\sigma - \tilde{\sigma}\|_1 = \mathcal{O}(\|\phi - \tilde{\phi}\|_1) \rightarrow 0. \quad (4.27)$$

We now make one final approximation to σ , in which we discard the parts of the background state where the maximally entangled states along each cut are close in dimension, as in [Figure 5d](#). Consider the state

$$|\bar{\phi}_{\gamma_{A,1}, \gamma_{A,2}}\rangle = \sum_{2^{n^{1/4}} p_{2,k} \leq p_{2,j} \text{ or } 2^{-n^{1/4}} p_{2,k} \geq p_{2,j}} \sqrt{q_{1,k} q_{2,j}} |\Phi_{1,k}\rangle \otimes |\Phi_{2,j}\rangle,$$

where the sum is still only over those j and k for which $q_{1,k} > n^{-\beta}$ and $q_{2,j} > n^{-\beta}$. If we let

$$D_n = \{(x, y) \in \mathbb{R}^2 : x - n^{-\frac{1}{4}} - n^{-\frac{1}{2}} \leq y \leq x + n^{-\frac{1}{4}} + n^{-\frac{1}{2}}, |x| \leq \log(n) + 1\},$$

then as n increases, the measure of D_n converges to zero and since the measures $\nu_{\phi_{\gamma_{A,i}}} \Rightarrow \nu_i$ and ν_i has continuous cumulative distribution function

$$\sum_{2^{-n^{1/4}} p_{1,j} \leq p_{2,k} \leq 2^{n^{1/4}} p_{1,j}} q_{1,j} q_{2,k} \leq (\nu_{\phi_{\Gamma_{A,1}}} \times \nu_{\phi_{\Gamma_{A,2}}})(D_n) \rightarrow 0. \quad (4.28)$$

Hence, if we let $\bar{\phi}$ denote the background state with $\tilde{\phi}_{\gamma_{A,1}} \otimes \tilde{\phi}_{\gamma_{A,2}}$ replaced by $\bar{\phi}_{\gamma_{A,1}, \gamma_{A,2}}$, we get that $\|\bar{\phi} - \tilde{\phi}\|_1 \rightarrow 0$. By [Lemma 4.3](#), if we denote by $\bar{\sigma}$ the state we obtain on G' by using $\bar{\phi}$ rather than $\tilde{\phi}$ as the background state, we get

$$\mathbb{E}\|\bar{\sigma} - \sigma\|_1 \rightarrow 0. \quad (4.29)$$

We pause here to note that we have accomplished Step 2: we have constructed an approximation $\bar{\sigma}$ to σ by approximating the background state along the cuts as superpositions of maximally entangled states. The utility in doing so is that working with this approximated tensor network state allows us to reduce to calculations where we restrict to a maximally entangled state along one of the two surfaces. Our next major step is to then show that the spectrum of $\bar{\sigma}$ can be used to approximate the spectrum of ρ , which will then allow us to analyze the convergence of the corresponding distribution. As a first intermediate step, we will show that $\bar{\sigma}$ can be approximated as a superposition of a state with minimal cut $\gamma_{A,1}$ and a state with minimal cut $\gamma_{A,2}$. This will allow us to more easily reason about how the background states are related to the spectrum of $\bar{\sigma}$, in turn, σ , and in turn, ρ_A .

Let us write

$$\begin{aligned} |\phi_{\gamma_{A,1},\gamma_{A,2}}^{(j,a)}\rangle &= \sum_{k \text{ s.t. } p_{1,k} \geq p_{2,j} 2^{n/4}} \sqrt{q_{1,k} q_{2,j}} |\Phi_{1,k}^+\rangle \otimes |\Phi_{2,j}^+\rangle \\ |\phi_{\gamma_{A,1},\gamma_{A,2}}^{(j,b)}\rangle &= \sum_{k \text{ s.t. } p_{2,k} \geq p_{1,j} 2^{n/4}} \sqrt{q_{1,j} q_{2,k}} |\Phi_{1,j}^+\rangle \otimes |\Phi_{2,k}^+\rangle, \end{aligned}$$

allowing us to write the background state $\bar{\phi}_{\gamma_{A,1},\gamma_{A,2}}$ as a different superposition, as depicted in [Figure 5e](#):

$$|\bar{\phi}_{\gamma_{A,1},\gamma_{A,2}}\rangle = \sum_j |\phi_{\gamma_{A,1},\gamma_{A,2}}^{(j,a)}\rangle + \sum_j |\phi_{\gamma_{A,1},\gamma_{A,2}}^{(j,b)}\rangle.$$

Let $|\phi^{(j,a)}\rangle$ and $|\phi^{(j,b)}\rangle$ denote the background states on G' where we have replaced $|\bar{\phi}_{\gamma_{A,1},\gamma_{A,2}}\rangle$ by $|\phi_{\gamma_{A,1},\gamma_{A,2}}^{(j,a)}\rangle$ and $|\phi_{\gamma_{A,1},\gamma_{A,2}}^{(j,b)}\rangle$ respectively, and let

$$|\phi^a\rangle = \sum_j |\phi^{(j,a)}\rangle, \quad |\phi^b\rangle = \sum_j |\phi^{(j,b)}\rangle.$$

Denote by $|\sigma^{(j,a)}\rangle, |\sigma^{(j,b)}\rangle$ the random tensor network states on G' with background states $|\phi^{(j,a)}\rangle$ and $|\phi^{(j,b)}\rangle$, respectively. Similarly, denote by $|\sigma^{(a)}\rangle$ and $|\sigma^{(b)}\rangle$ the random tensor network states on G' with background states $|\phi^{(a)}\rangle$ and $|\phi^{(b)}\rangle$, respectively.

We start with the following bound on the rank of $\sigma_B^{(j,a)}$:

$$\begin{aligned} \text{rank}(\sigma_B^{(j,a)}) &\leq \text{rank}(\phi_B^{(j,a)}) \\ &\leq \sum_{p_{1,k} \geq p_{2,j} 2^{n/4}} d_{1,k} \\ &\leq \sum_{p_{1,k} \geq p_{2,j} 2^{n/4}} \frac{q_{1,k}}{p_{1,k}} \\ &\leq \sum_k \frac{q_{1,k} 2^{-n/4}}{p_{2,j}} \\ &\leq \frac{2^{-n/4}}{p_{2,j}} \end{aligned} \tag{4.30}$$

By the same reasoning we may bound the rank of $\sigma_B^{(j,b)}$ as

$$\text{rank}(\sigma_B^{(j,b)}) \leq \frac{2^{-n/4}}{p_{1,j}}. \tag{4.31}$$

Now, we will argue that the state $\sigma^{(j,a)}$ has minimal cut $\gamma_{A,1}$, and $\sigma^{(j,b)}$ has minimal cut $\gamma_{A,2}$, as in [Figure 5f](#). Intuitively, it is sensible that for $\sigma^{(j,a)}$ the unique minimal cut is along $\gamma_{A,1}$, as for $\phi^{(j,a)}$, we have

a fixed maximally entangled state along $\gamma_{A,2}$, and a superposition of maximally entangled states of lower dimension along $\gamma_{A,1}$. Similarly, for $\sigma^{(j,b)}$ the minimal cut is along $\gamma_{A,2}$. To confirm this intuition, we will now show that $\sigma_B^{(j,a)} \approx \phi_B^{(j,a)}$ and $\sigma_{\bar{B}}^{(j,b)} \approx \phi_{\bar{B}}^{(j,b)}$.

We show this for $\sigma_B^{(j,a)}$. In this case, the ‘minimal’ cut is simply B . Let Δ_B be a cut for B unequal to B or to $V'_b \cup B$. We denote by Δ_A the associated minimal cut for A on the original graph G given by $\Delta_A = \Delta_B \cup \Gamma_{A,1}$. We let

$$Y^{\Delta_B} = \{(e, x) : e = (xy), x \in \Delta_B^c, y \in B\}.$$

Note that as ϕ is a product state we have $H_{\min}^\varepsilon(\Delta_B \setminus B|B)_\phi = H_{\min}^\varepsilon(\Delta_B \setminus B|BY^{\Delta_B})_\phi$. We can obtain $\phi^{(j,a)}$ from ϕ by acting with subunitary CP maps on B and \bar{B} and therefore (by two applications of data processing)

$$H_{\min}^\varepsilon(\Delta_B \setminus B|BY^{\Delta_B})_{\phi^{(j,a)}} \geq H_{\min}^\varepsilon(\Delta_B \setminus B|BY^{\Delta_B})_\phi = H_{\min}^\varepsilon(\Delta_B \setminus B|B)_\phi \geq H_{\min}^\varepsilon(\Delta_A \setminus \Gamma_A|\Gamma_A)_\phi$$

We now consider $\Delta_B = V'_b \cup B$ (in which case Y^{Δ_B} is empty, since we assume $\gamma_{A,1} \cap \gamma_{A,2} = \emptyset$). Then

$$H_{\min}^\varepsilon(\Delta_B \setminus B|B)_{\phi^{(j,a)}} \geq H_{\min}(\Delta_B \setminus B|B)_{\phi^{(j,a)}} = H_{\min}(\bar{B})_{\phi^{(j,a)}} - H_{\max}(B)_{\phi^{(j,a)}}$$

using the product structure of $\phi^{(j,a)}$. Now, we compute

$$H_{\min}(\bar{B})_{\phi^{(j,a)}} = -\log(\|\phi_{\bar{B}}^{(j,a)}\|_\infty) \geq -\log(p_{2,j}),$$

where we recall that $p_{2,j}$ is a binned eigenvalue. By Eq. (4.30), we can bound

$$H_{\max}(B)_{\phi^{(j,a)}} \leq \log(\text{rank}(\phi_B^{(j,a)})) \leq \log\left(\frac{2^{-n^{1/4}}}{p_{2,j}}\right) = -\log(p_{2,j}) - n^{\frac{1}{4}}.$$

In conclusion,

$$H_{\min}^\varepsilon(\Delta_B \setminus B|B)_{\phi^{(j,a)}} \geq n^{\frac{1}{4}}.$$

By Lemma 4.11 we can find $\phi^{(\varepsilon,j,a)}$ such that $P(\phi^{(j,a)}, \phi^{(\varepsilon,j,a)}) = \mathcal{O}(\sqrt{\varepsilon})$ and for all cuts $\Delta_B \neq B$ we have

$$H_{\min}(\Delta_B \setminus B|B)_{\phi^{(\varepsilon,j,a)}} \geq H_{\min}^\varepsilon(\Delta_B \setminus B|BY^{\Delta_B})_{\phi^{(j,a)}}$$

and hence for $\Delta_B \neq V'_b \cup B$

$$H_{\min}(\Delta_B \setminus B|B)_{\phi^{(\varepsilon,j,a)}} \geq H_{\min}^\varepsilon(\Delta_A \setminus \Gamma_A|\Gamma_A)_\phi$$

and for $\Delta_B = V'_b \cup B$

$$H_{\min}(\Delta_B \setminus B|B)_{\phi^{(\varepsilon,j,a)}} \geq n^{\frac{1}{4}}.$$

Therefore, Proposition 4.1 allows us to conclude that if we let $\sigma^{(\varepsilon,j,a)}$ denote the random tensor network state with background state $\phi^{(\varepsilon,j,a)}$, then

$$\mathbb{E}\|\sigma_B^{(\varepsilon,j,a)} - \phi_B^{(\varepsilon,j,a)}\|_1 = \mathcal{O}\left(2^{-\frac{1}{2}K(n)} + 2^{-\frac{1}{2}n^{\frac{1}{4}}}\right),$$

and by Lemma 4.3 and the fact that $T(\phi^{(j,a)}, \phi^{(\varepsilon,j,a)}) \leq P(\phi^{(j,a)}, \phi^{(\varepsilon,j,a)})$ this implies

$$\begin{aligned} \mathbb{E}\|\sigma_B^{(j,a)} - \phi_B^{(j,a)}\|_1 &\leq \mathbb{E}\|\sigma_B^{(\varepsilon,j,a)} - \phi_B^{(\varepsilon,j,a)}\|_1 + \mathbb{E}\|\sigma_B^{(\varepsilon,j,a)} - \sigma_B^{(j,a)}\|_1 + \|\phi_B^{(\varepsilon,j,a)} - \phi_B^{(j,a)}\|_1 \\ &= \mathcal{O}\left(2^{-\frac{1}{2}K(n)} + 2^{-\frac{1}{2}n^{\frac{1}{4}}} + \sqrt{\varepsilon(n)}\right). \end{aligned}$$

It follows that

$$\mathbb{E}\|\sigma_B^{(a)} - \phi_B^{(a)}\|_1 \leq \sum_j \mathbb{E}\|\sigma_B^{(j,a)} - \phi_B^{(j,a)}\|_1 = \mathcal{O}\left(n^{\alpha+\frac{1}{2}} \log(n)(2^{-\frac{1}{2}K(n)} + 2^{-\frac{1}{2}n^{\frac{1}{4}}} + \sqrt{\varepsilon(n)})\right) \rightarrow 0 \quad (4.32)$$

using that the number of terms is $\mathcal{O}(n^{\alpha+\frac{1}{2}} \log(n))$. This expression goes to zero since we assume $K(n) = \Omega(\log(n)^2)$ and $\varepsilon(n) = \mathcal{O}(n^{-\gamma})$. A completely symmetric argument shows that

$$\mathbb{E}\|\sigma_B^{(j,b)} - \phi_B^{(j,b)}\|_1 = \mathcal{O}\left(2^{-\frac{1}{2}K(n)} + 2^{-\frac{1}{2}n^{\frac{1}{4}}} + \sqrt{\varepsilon(n)}\right) \quad (4.33)$$

and hence

$$\mathbb{E}\|\sigma_B^{(b)} - \phi_B^{(b)}\|_1 = \mathcal{O}(n^{\alpha+\frac{1}{2}} \log(n)(2^{-\frac{1}{2}K(n)} + 2^{-\frac{1}{2}n^{\frac{1}{4}}} + \varepsilon(n))) \rightarrow 0. \quad (4.34)$$

We have hence shown that $\sigma_B^{(j,a)} \approx \phi_B^{(j,a)}$ and $\sigma_B^{(j,b)} \approx \phi_B^{(j,b)}$ in expectation.

We now show that we can approximate the state $\bar{\sigma}_B$ on G' by the sum of these states. We first claim that we can sum over the j index to obtain the $\sigma_B^{(a)}$ and $\sigma_B^{(b)}$. To see this, we introduce some notation: given a self-adjoint matrix X , let us write $\text{supp}(X)$ for the image, or support, of X . In other words, $\text{supp}(X)$ is the space spanned by the eigenvectors of X with nonzero eigenvalue. We let $\mathcal{H}_{B,j}$ be the support of the reduced state of $|\Phi_{1,j}^+\rangle$ on the B system, and we similarly let $\mathcal{H}_{\bar{B},j}$ be the support of the reduced state of $|\Phi_{2,j}^+\rangle$ on the \bar{B} system. Then for $j \neq k$, the subspace $\mathcal{H}_{B,j}$ is orthogonal to $\mathcal{H}_{B,k}$ and similarly $\mathcal{H}_{\bar{B},j}$ is orthogonal to $\mathcal{H}_{\bar{B},k}$. By construction, it is clear that $\text{supp}(\sigma_B^{(j,a)}) \subseteq \mathcal{H}_{B,j}$ and $\text{supp}(\sigma_B^{(j,b)}) \subseteq \mathcal{H}_{B,j}$. This orthogonality for indices $j \neq k$ then makes it clear that we can sum the reduced states, as in [Figure 5g](#):

$$\sigma_B^{(a)} = \sum_j \sigma_B^{(j,a)}, \quad \sigma_B^{(b)} = \sum_j \sigma_B^{(j,b)},$$

and by the purity of $\sigma_B^{(b)}$, we have that $\sigma_B^{(b)}$ has the same spectrum as $\sigma_{\bar{B}}^{(b)}$. We remind the reader that the (a) and (b) indices indicate that the minimal cut for the state is given by $\gamma_{A,1}$ and $\gamma_{A,2}$, respectively. Naturally, this also holds for the summed states, by orthogonality of the summands.

We now claim that $\bar{\sigma}_B \approx \sigma_B^{(a)} + \sigma_B^{(b)}$, i.e. the spectrum of the approximate state on G' , as in [Eq. \(4.29\)](#), is well-approximated by the sum of two states with differing minimal cuts. We estimate their difference by

$$\begin{aligned} \mathbb{E}\|\sigma_B^{(a)} + \sigma_B^{(b)} - \bar{\sigma}_B\|_1 &= \mathbb{E}\left\|\sum_{j,k} \text{tr}_{\bar{B}}[\sigma^{(j,a)} \langle \sigma^{(k,b)} | + | \sigma^{(k,b)} \rangle \langle \sigma^{(j,a)} |]\right\|_1 \\ &\leq 2 \sum_{j,k} \mathbb{E}\|\text{tr}_{\bar{B}}[\sigma^{(j,a)} \langle \sigma^{(k,b)} |]\|_1 \\ &= 2 \sum_{j,k} \mathbb{E}F(\bar{\sigma}_B^{(j,a)}, \bar{\sigma}_B^{(k,b)}) \end{aligned}$$

by [Lemma 4.9](#). Now, if ρ, σ_1 and σ_2 are positive operators, then by [Lemma B.9](#) in [\[FR15\]](#), we have

$$|F(\rho, \sigma_1) - F(\rho, \sigma_2)| \leq \sqrt{\|\sigma_1 - \sigma_2\|_1 \text{tr}[\rho]}.$$

So, we find that

$$\begin{aligned} \mathbb{E}F(\sigma_B^{(j,a)}, \sigma_B^{(k,b)}) &\leq \mathbb{E}|F(\sigma_B^{(j,a)}, \sigma_B^{(k,b)}) - F(\sigma_B^{(j,a)}, \phi_B^{(k,b)})| + \mathbb{E}F(\sigma_B^{(j,a)}, \phi_B^{(k,b)}) \\ &\leq \mathbb{E}\sqrt{\text{tr}[\sigma_B^{(j,a)}] \|\sigma_B^{(k,b)} - \phi_B^{(k,b)}\|_1} + \mathbb{E}F(\sigma_B^{(j,a)}, \phi_B^{(k,b)}) \\ &\leq \sqrt{\mathbb{E}\text{tr}[\sigma_B^{(j,a)}]} \sqrt{\mathbb{E}\|\sigma_B^{(k,b)} - \phi_B^{(k,b)}\|_1} + \mathbb{E}F(\sigma_B^{(j,a)}, \phi_B^{(k,b)}), \end{aligned}$$

where we used Cauchy-Schwarz in the last step. For the first term, we may use Eq. (4.33) and $\mathbb{E} \operatorname{tr}[\bar{\sigma}_B^{(j,a)}] \leq 1$ to see

$$\sqrt{\mathbb{E} \operatorname{tr}[\sigma_B^{(j,a)}]} \sqrt{\mathbb{E} \|\sigma_B^{(k,b)} - \phi_B^{(k,b)}\|_1} = \mathcal{O}(2^{-\frac{1}{4}K(n)} + 2^{-\frac{1}{4}n^{\frac{1}{4}}} + (\varepsilon(n))^{\frac{1}{4}}).$$

For the second term we use a basic estimate on the fidelity: if ρ and σ are positive operators, then

$$F(\rho, \sigma) = \|\sqrt{\rho}\sqrt{\sigma}\|_1 \leq \|\sqrt{\rho}\|_1 \|\sqrt{\sigma}\|_\infty \leq \sqrt{\operatorname{rank}(\rho)} \|\sqrt{\rho}\|_2 \|\sqrt{\sigma}\|_\infty = \sqrt{\operatorname{rank}(\rho) \operatorname{tr}[\rho] \|\sigma\|_\infty},$$

which follows by Hölder's inequality and the standard relation between Schatten 1 and 2-norms. Then the second term can be estimated as follows: write $P_{\bar{B},j}$ for the projection onto $\mathcal{H}_{\bar{B},j}$, then

$$\begin{aligned} F(\sigma_B^{(j,a)}, \phi_B^{(k,b)}) &= F(\sigma_B^{(j,a)}, P_{\bar{B},j} \phi_B^{(k,b)} P_{\bar{B},j}) \\ &\leq \sqrt{\operatorname{tr}[\sigma_B^{(j,a)}] \operatorname{rank}(\sigma_B^{(j,a)}) \|P_{\bar{B},j} \phi_B^{(k,b)} P_{\bar{B},j}\|_\infty} \\ &\leq \sqrt{\operatorname{tr}[\sigma_B^{(j,a)}] \frac{2^{-n^{1/4}}}{p_{2,j}} p_{2,j}} \\ &\leq \sqrt{\operatorname{tr}[\sigma_B^{(j,a)}] 2^{-n^{1/4}}} \end{aligned}$$

using Eq. (4.30) and $\|P_{\bar{B},j} \phi_B^{(k,b)} P_{\bar{B},j}\|_\infty \leq p_{2,j}$. Thus

$$\mathbb{E} F(\sigma_B^{(j,a)}, \phi_B^{(k,b)}) \leq \sqrt{\mathbb{E} \operatorname{tr}[\sigma_B^{(j,a)}] 2^{-n^{1/4}}} \leq 2^{-\frac{1}{2}n^{1/4}}$$

and we may estimate

$$\mathbb{E} F(\sigma_B^{(j,a)}, \sigma_B^{(k,b)}) = \mathcal{O}(2^{-\frac{1}{4}K(n)} + 2^{-\frac{1}{4}n^{\frac{1}{4}}} + (\varepsilon(n))^{\frac{1}{4}}).$$

and therefore

$$\mathbb{E} \|\sigma_B^{(a)} + \sigma_B^{(b)} - \bar{\sigma}_B\|_1 \leq 2 \sum_{j,k} \mathbb{E} F(\bar{\sigma}_B^{(j,a)}, \bar{\sigma}_B^{(k,b)}) = \mathcal{O}(n^{2\alpha+1} \log(n)^2 (2^{-\frac{1}{4}K(n)} + 2^{-\frac{1}{4}n^{\frac{1}{4}}} + (\varepsilon(n))^{\frac{1}{4}})) \rightarrow 0 \quad (4.35)$$

using that the number of terms is $\mathcal{O}(n^{2\alpha+1} \log(n)^2)$, $K(n) = \Omega(\log(n)^2)$ and $\varepsilon(n) = \mathcal{O}(n^{-\gamma})$, and our choice of α is such that $\mathcal{O}(n^{2\alpha+1} \log(n)^2 n^{-\frac{1}{4}\gamma})$ goes to zero.

To summarize, we have shown that we can approximate the random tensor network state $\bar{\sigma}_B$ on G' by $\sigma_B^{(a)} + \sigma_B^{(b)}$, and we can approximate $\sigma_B^{(a)}$ by $\phi_B^{(a)}$. Moreover, we can approximate $\sigma_B^{(b)}$ by $\phi_B^{(b)}$, so the spectrum of $\sigma_B^{(b)}$ can be approximated by the spectrum of $\phi_B^{(b)}$. Recall that $\sigma^{(a)}$ and $\sigma^{(b)}$ are the random tensor network states on G' which take $\phi^{(a)}$ and $\phi^{(b)}$ as background states, respectively. Furthermore, recall that our larger goal is to show that the spectra of the reduced background states $\phi_B^{(a)}$ and $\phi_B^{(b)}$ are, in some sense, close to the spectrum of the approximate random tensor network state σ_B on the subgraph G' . This is a two-step process:

1. Show that $\phi_B^{(a)}$ and $\sigma_B^{(a)}$ can be slightly deformed to $\phi_B^{(a,\perp)}$ and $\sigma_B^{(a,\perp)}$, states with support orthogonal to $\operatorname{supp}(\sigma_B^{(j,b)})$ for all j .
2. Use the slightly deformed states to show that $\operatorname{spec}_+(\phi_B^{(a)} \oplus \phi_B^{(b)})$ is close to $\operatorname{spec}_+(\sigma_B^{(a)} + \sigma_B^{(b)})$, and follow the chain of approximations: $\sigma_B^{(a)} + \sigma_B^{(b)} \approx \bar{\sigma}_B \approx \sigma_B$ along with $\operatorname{spec}_+(\sigma_B) \approx \operatorname{spec}_+(\rho_A)$ to conclude that $\operatorname{spec}_+(\phi_B^{(a)} \oplus \phi_B^{(b)}) \approx \operatorname{spec}_+(\rho_A)$

Now, consider the state $\sigma_B^{(j,b)}$. It has support contained in $\mathcal{H}_{B,j}$ and by Eq. (4.31), it has $\text{rank}(\sigma_B^{(j,b)}) \leq 2^{-n^{1/4}} p_{1,j}^{-1}$. Let Q_j be the projection onto $\text{supp}(\sigma_B^{(j,b)})$, so $\text{supp}(Q_j) \subseteq \mathcal{H}_{B,j}$ and $\text{rank}(Q_j) \leq 2^{-n^{1/4}} p_{1,j}^{-1}$. We find that

$$\|Q_j \phi_B^{(a)}\|_1 \leq \text{rank}(Q_j) \|P_{B,j} \phi_B^{(a)} P_{B,j}\|_\infty \leq \frac{2^{-n^{1/4}}}{p_{1,j}} p_{1,j} = 2^{-n^{1/4}}.$$

If we let $Q = \sum_j Q_j$, then we see that $(I - Q)\phi^{(a)}(I - Q)$ is a small deformation of $\phi_B^{(a)}$:

$$\|\phi_B^{(a)} - (I - Q)\phi^{(a)}(I - Q)\|_1 \leq 2\|Q\phi^{(a)}\|_1 \leq 2 \sum_j \|Q_j \phi_B^{(a)}\|_1 = \mathcal{O}(n^{9/2} \log(n) 2^{-n^{1/4}}) \rightarrow 0. \quad (4.36)$$

With this property in hand, we can show that $\sigma_B^{(a)}$ can similarly be deformed. Let $\sigma_B^{(a,\perp)} = (I - Q)\sigma^{(a)}(I - Q)$. By construction, $\sigma_B^{(a,\perp)}$ has support orthogonal to that of $\sigma_B^{(b)}$, and hence, $\text{spec}_+(\sigma_B^{(a,\perp)} + \sigma_B^{(b)}) = \text{spec}_+(\sigma_B^{(a,\perp)} \oplus \sigma_B^{(b)})$. We observe that $\sigma_B^{(a,\perp)}$ is a small deformation of $\sigma_B^{(a)}$:

$$\begin{aligned} \mathbb{E}\|\sigma_B^{(a)} - \sigma_B^{(a,\perp)}\|_1 &\leq \mathbb{E}\|\sigma_B^{(a)} - \phi_B^{(a)}\|_1 \\ &\quad + \mathbb{E}\|(I - Q)\sigma_B^{(a)}(I - Q) - (I - Q)\phi_B^{(a)}(I - Q)\|_1 \\ &\quad + \|\phi_B^{(a)} - (I - Q)\phi^{(a)}(I - Q)\|_1 \\ &\leq 2\mathbb{E}\|\sigma_B^{(a)} - \phi_B^{(a)}\|_1 + \|\phi_B^{(a)} - (I - Q)\phi^{(a)}(I - Q)\|_1 \end{aligned} \quad (4.37)$$

which goes to zero, using Eq. (4.32) for the first term, and Eq. (4.36) for the second term. By construction, $(I - Q)\phi^{(a)}(I - Q)$ has support orthogonal to $\text{supp}(\sigma_B^{(b)})$. This allows us to bound

$$\begin{aligned} \mathbb{E}\|\text{spec}_+(\phi_B^{(a)} \oplus \phi_B^{(a)}) - \text{spec}_+(\sigma_B^{(a)} + \sigma_B^{(b)})\|_1 \\ &\leq \mathbb{E}\|\sigma_B^{(a)} - \sigma_B^{(a,\perp)}\|_1 + \mathbb{E}\|\text{spec}_+(\phi_B^{(a)} \oplus \phi_B^{(a)}) - \text{spec}_+(\sigma_B^{(a,\perp)} \oplus \sigma_B^{(b)})\|_1 \\ &= \mathbb{E}\|\sigma_B^{(a)} - \sigma_B^{(a,\perp)}\|_1 + \mathbb{E}\|\text{spec}_+(\phi_B^{(a)}) - \text{spec}_+(\sigma_B^{(a,\perp)}) + \text{spec}_+(\phi_B^{(a)}) - \text{spec}_+(\sigma_B^{(b)})\|_1 \\ &\leq \mathbb{E}\|\sigma_B^{(a)} - \sigma_B^{(a,\perp)}\|_1 + \mathbb{E}\|\phi_B^{(a)} - \sigma_B^{(a,\perp)}\|_1 + \mathbb{E}\|\text{spec}_+(\phi_B^{(a)}) - \text{spec}_+(\sigma_B^{(b)})\|_1, \end{aligned}$$

where we have used the bound on the difference of spectra by the trace distance of the corresponding states Eq. (1.5). Each term on the RHS can be shown to converge to zero. The first term goes to zero by Eq. (4.37). Similarly, the second term can be bounded by

$$\mathbb{E}\|\phi_B^{(a)} - \sigma_B^{(a,\perp)}\|_1 \leq \mathbb{E}\|\phi_B^{(a)} - \sigma_B^{(a)}\|_1 + \mathbb{E}\|\sigma_B^{(a)} - \sigma_B^{(a,\perp)}\|_1$$

which goes to zero by Eq. (4.37) and Eq. (4.32). The third term can be estimated by observing that $\text{spec}_+(\sigma_B^{(b)}) = \text{spec}_+(\sigma_B^{(b)})$ and

$$\mathbb{E}\|\text{spec}_+(\phi_B^{(a)}) - \text{spec}_+(\sigma_B^{(b)})\|_1 \leq \mathbb{E}\|\phi_B^{(a)} - \sigma_B^{(b)}\|_1$$

which goes to zero by Eq. (4.34). We conclude that $\mathbb{E}\|\text{spec}_+(\phi_B^{(a)} \oplus \phi_B^{(a)}) - \text{spec}_+(\sigma_B^{(a)} + \sigma_B^{(b)})\|_1 \rightarrow 0$.

We now follow a chain of approximations to get the desired closeness between $\text{spec}_+(\phi_B^{(a)} \oplus \phi_B^{(a)})$ and $\text{spec}_+(\rho_A)$. Using Eq. (4.29) and Eq. (4.35), we can see that

$$\mathbb{E}\|\sigma_B - (\sigma_B^{(a)} + \sigma_B^{(b)})\|_1 \leq \mathbb{E}\|\sigma_B - \bar{\sigma}_B\|_1 + \mathbb{E}\|\sigma_B^{(a)} + \sigma_B^{(b)} - \bar{\sigma}_B\|_1 \rightarrow 0,$$

so $\mathbb{E}\|\sigma_B - (\sigma_B^{(a)} + \sigma_B^{(b)})\|_1$ will converge to 0. This then implies

$$\mathbb{E}\|\text{spec}_+(\sigma_B) - \text{spec}_+(\phi_B^{(a)} \oplus \phi_B^{(a)})\|_1 \leq \mathbb{E}\|\sigma_B - (\sigma_B^{(a)} + \sigma_B^{(b)})\|_1$$

$$+ \mathbb{E} \|\text{spec}_+(\phi_B^{(a)} \oplus \phi_B^{(b)}) - \text{spec}_+(\sigma_B^{(a)} + \sigma_B^{(b)})\|_1,$$

and as we have just shown, both terms on the RHS converge to zero. Together with Eq. (4.23), this yields the desired relationship to the spectrum of ρ :

$$\begin{aligned} \mathbb{E} \|\text{spec}_+(\phi_B^{(a)} \oplus \phi_B^{(b)}) - \text{spec}_+(\rho_A)\|_1 &\leq \mathbb{E} \|\text{spec}_+(\phi_B^{(a)} \oplus \phi_B^{(b)}) - \text{spec}_+(\sigma_B)\|_1 \\ &+ \mathbb{E} \|\text{spec}_+(\rho_A) - \text{spec}_+(\sigma_B)\|_1 \rightarrow 0. \end{aligned} \quad (4.38)$$

We pause here again to note that we have accomplished Step 3: approximating the spectrum of ρ_A by the spectra of the (approximate) background states. The final step will then be to consider the convergence properties of the distributions on the background states, which will then translate to convergence for the distribution on ρ_A .

Explicitly, we want to relate the above result to $\min_*(\nu_1, \nu_2)$. First, we observe that by Lemma 4.8 and Eq. (4.26), $\nu_{\tilde{\phi}_B} \Rightarrow \nu_1$ and $\nu_{\tilde{\phi}_{\bar{B}}} \Rightarrow \nu_2$, and since \min is a continuous function, convergence holds for the pushforward measure:

$$\min_*(\nu_{\tilde{\phi}_B}, \nu_{\tilde{\phi}_{\bar{B}}}) \Rightarrow \min_*(\nu_1, \nu_2). \quad (4.39)$$

We compute

$$\begin{aligned} \min_*(\nu_{\tilde{\phi}_B}, \nu_{\tilde{\phi}_{\bar{B}}}) &= \sum_{j,k} q_{1,k} q_{2,j} \delta_{\frac{1}{\sqrt{n}} [\min(\log(\frac{1}{p_{1,k}}), \log(\frac{1}{p_{2,j}})) - H(n)]} \\ &= \sum_j \left(\sum_{p_{1,k} \geq 2^{n^{1/4}} p_{2,j}} q_{1,k} q_{2,j} \delta_{\frac{1}{\sqrt{n}} [\log(\frac{1}{p_{1,k}}) - H(n)]} \right. \\ &\quad \left. + \sum_{p_{1,k} \leq 2^{-n^{1/4}} p_{2,j}} q_{1,k} q_{2,j} \delta_{\frac{1}{\sqrt{n}} [\log(\frac{1}{p_{2,j}}) - H(n)]} \right) \\ &\quad + \sum_{2^{-n^{1/4}} p_{2,j} \leq p_{1,k} \leq 2^{n^{1/4}} p_{2,j}} q_{1,k} q_{2,j} \delta_{\frac{1}{\sqrt{n}} [\min(\log(\frac{1}{p_{1,k}}), \log(\frac{1}{p_{2,j}})) - H(n)]} \\ &= \sum_k q_{1,k} q_{2,>k} \delta_{\frac{1}{\sqrt{n}} [\log(\frac{1}{p_{1,k}}) - H(n)]} + \sum_j q_{1,>j} q_{2,j} \delta_{\frac{1}{\sqrt{n}} [\log(\frac{1}{p_{2,j}}) - H(n)]} + \nu_0 \end{aligned} \quad (4.40)$$

where we have written

$$\begin{aligned} q_{1,>j} &= \sum_{p_{1,k} \geq 2^{n^{1/4}} p_{2,j}} q_{1,k} \\ q_{2,>k} &= \sum_{p_{1,k} \leq 2^{-n^{1/4}} p_{2,j}} q_{2,j} \end{aligned}$$

and

$$\nu_0 = \sum_{2^{-n^{1/4}} p_{2,j} \leq p_{1,k} \leq 2^{n^{1/4}} p_{2,j}} q_{1,k} q_{2,j} \delta_{\frac{1}{\sqrt{n}} [\min(\log(\frac{1}{p_{1,k}}), \log(\frac{1}{p_{2,j}})) - H(n)]}.$$

Then by Eq. (4.28), we see that $\nu_0 \Rightarrow 0$. Using Eq. (4.39), we conclude

$$\min_*(\nu_{\tilde{\phi}_B}, \nu_{\tilde{\phi}_{\bar{B}}}) - \nu_0 \Rightarrow \min_*(\nu_1, \nu_2). \quad (4.41)$$

On the other hand, by construction, we have

$$\nu_{\text{spec}(\phi_B^{(a)} \oplus \phi_B^{(b)})} = \nu_{\phi_B^{(a)}} + \nu_{\phi_B^{(b)}}.$$

We can explicitly write down

$$\nu_{\phi_B^{(a)}} + \nu_{\phi_B^{(b)}} = \sum_k q_{1,k} q_{2,>k} \delta_{\frac{1}{\sqrt{n}}[\log(\frac{1}{p_{1,k} q_{2,>k}}) - H(n)]} + \sum_j q_{1,>k} q_{2,j} \delta_{\frac{1}{\sqrt{n}}[\log(\frac{1}{p_{2,j} q_{1,>k}}) - H(n)]}.$$

Since the $q_{1,k}$ and $q_{2,j}$ are at least $n^{-\beta}$ (if the corresponding term is nonzero), $|\log(q_{1,>j})| \leq \beta \log(n)$ and $|\log(q_{2,>k})| \leq \beta \log(n)$. With these bounds in mind, we can compare to the last line of [Eq. \(4.40\)](#). More precisely, if we let $\nu_{\phi}^t = \min_*(\nu_{\phi_B}, \nu_{\phi_{\bar{B}}}) - \nu_0$, then for uniformly continuous $f \in C_b(\mathbb{R})$, we have

$$\lim_{n \rightarrow \infty} \left| \int f(x) d\nu_{\phi}^t(x) - \int f(x) d\nu_{\phi_B^{(a)}}(x) - \int f(x) d\nu_{\phi_B^{(b)}}(x) \right| = 0$$

We conclude that

$$\nu_{\phi_B^{(a)}} + \nu_{\phi_B^{(b)}} \Rightarrow \min_*(\nu_1, \nu_2). \quad (4.42)$$

Now, we can finally put all of our ingredients together. Recall the statement of convergence in probability implied by the convergence of spectra, in expectation, as in the second part of [Lemma 4.8](#). Then using [Eq. \(4.38\)](#) as the vectors $p^{(n)}$ and $q^{(n)}$ in the statement of [Lemma 4.8](#) and the convergence of distributions in [Eq. \(4.42\)](#), we conclude

$$\nu_{\rho_A} \Rightarrow \min_*(\nu_1, \nu_2) \quad (4.43)$$

□

4.5 Computing entropies with two minimal cuts

Ideally, we would like to use [Theorem 4.12](#) to compute von Neumann entropies of random tensor network states. However, [Theorem 4.12](#) alone is too weak to allow us to directly compute entropies up to $o(\sqrt{n})$ corrections, as weak convergence of the spectrum does not directly imply convergence of the mean. However, we kept track of various approximation errors in the proof of [Theorem 4.12](#), and we will use these to show that with slightly stronger assumptions, these errors allow to compute the entropy up to $\mathcal{O}(\log(n))$ corrections.

To begin, we first bound the difference in the entropy of a sum of density matrices and the sum of the entropies of the individual density matrices:

Lemma 4.13. *Suppose $\{p_i\}_{i=1}^m$ is a subnormalized distribution and $\{\rho_i \in \mathcal{P}(\mathcal{H})\}_{i=1}^m$ is a collection of (unnormalized) density matrices and let $\rho = \sum_i p_i \rho_i \in \mathcal{P}(\mathcal{H})$. Suppose that $C^{-1} \leq \text{tr}[\rho_i] \leq C$, and $C^{-1} \leq \text{tr}[\rho] \leq C$, then*

$$|H(\rho) - \sum_{i=1}^m p_i H(\rho_i)| \leq C(\log(m) + 2\log(C)).$$

Proof. If the ρ_i are normalized and $\sum_i p_i = 1$, then by the Holevo bound we have

$$\sum_i p_i H(\rho_i) \leq H(\sum_i p_i \rho_i) \leq \sum_i p_i H(\rho_i) + H(\{p_i\}) \leq \sum_i p_i H(\rho_i) + \log(m).$$

Now let $q_i = \text{tr}[\rho_i]$, $P = \sum_i q_i p_i$ and let $\sigma_i = \frac{\rho_i}{q_i}$, $r_i = \frac{p_i q_i}{P}$. Then $H(\sum_i p_i \rho_i) = H(P \sum_i r_i \sigma_i) = PH(\sum_i r_i \sigma_i) - P \log(P)$. On the other hand

$$P \sum_i r_i H(\sigma_i) = \sum_i p_i H(\rho_i) + \sum_i p_i \log(q_i)$$

so it follows that

$$|H(\sum_i p_i \rho_i) - \sum_i p_i H(\rho_i)| \leq P \log(m) + |P \log(P)| + \max_i |\log(q_i)| \leq C(\log(m) + 2\log(C)).$$

□

The idea is that for a random tensor network state, we will split up the background state as a superposition of states which are maximally entangled along the two minimal cuts and then use [Lemma 4.13](#). This approach formalizes an argument sketched in [\[AP20\]](#).

To state our result, we introduce a new function: for vectors of positive numbers $p \in \mathbb{R}^{d_1}$, $q \in \mathbb{R}^{d_2}$, we let

$$H^*(p, q) := \sum_{i,j} p_i q_j \min\left(\log\left(\frac{1}{p_i}\right), \log\left(\frac{1}{q_j}\right)\right).$$

A key tool we will need is the continuity of the entropy: if $\rho, \sigma \in \mathcal{P}_{\leq}(\mathcal{H})$ are quantum states on a Hilbert space \mathcal{H} of dimension d with $T(\rho, \sigma) \leq \frac{1}{e}$, then the Fannes-Audenaert inequality states that

$$|H(\rho) - H(\sigma)| \leq T(\rho, \sigma) \log d - T(\rho, \sigma) \log(T(\rho, \sigma)). \quad (4.44)$$

Let η be the function defined by $\eta(x) = x \log x$. Then, if we write $\text{spec}(\rho) = \{p_j\}_{j=1}^d$ and $\text{spec}(\sigma) = \{q_j\}_{j=1}^d$ for [Eq. \(4.44\)](#) one can actually show

$$|H(\rho) - H(\sigma)| \leq \sum_{j=1}^d |\eta(p_j) - \eta(q_j)| \leq T(\rho, \sigma) \log d - T(\rho, \sigma) \log(T(\rho, \sigma)).$$

We will use this to show continuity of H^* as well. Consider $\rho_i, \sigma_i \in \mathcal{P}_{\leq}(\mathcal{H}_i)$ with $\dim(\mathcal{H}_i) \leq d$ for $i = 1, 2$. We let $\text{spec}(\rho_i) = \{p_{i,j}\}_{j=1}^{d_i}$ and $\text{spec}(\sigma_i) = \{q_{i,j}\}_{j=1}^{d_i}$. For real numbers x_i, y_i we have

$$|\min(x_1, y_1) - \min(x_2, y_2)| \leq |x_1 - x_2| + |y_1 - y_2|.$$

Then we see that

$$\begin{aligned} & |H^*(\text{spec}(\rho_1), \text{spec}(\rho_2)) - H^*(\text{spec}(\sigma_1), \text{spec}(\sigma_2))| \\ & \leq \sum_{j,k} |p_{1,j} p_{2,k} \min(\log p_{1,j}, \log p_{2,k}) - q_{1,j} q_{2,k} \min(\log q_{1,j}, \log q_{2,k})| \\ & \leq \sum_{j,k} |p_{1,j} p_{2,k} \log p_{1,j} - q_{1,j} q_{2,k} \log q_{1,j}| + |p_{1,j} p_{2,k} \log p_{2,k} - q_{1,j} q_{2,k} \log q_{2,k}|. \end{aligned}$$

We may estimate the first term by

$$\begin{aligned} \sum_{j,k} |p_{1,j} p_{2,k} \log p_{1,j} - q_{1,j} q_{2,k} \log q_{1,j}| & \leq \sum_{j,k} p_{2,k} |p_{1,j} \log p_{1,j} - q_{1,j} \log q_{1,j}| + \sum_{j,k} |p_{2,k} - q_{2,k}| q_{1,j} \log q_{1,j} \\ & \leq \sum_j |\eta(p_{1,j}) - \eta(q_{1,j})| + \|\rho_2 - \sigma_2\|_1 H(\sigma_1) \\ & \leq T(\rho_1, \sigma_1) \log d - T(\rho_1, \sigma_1) \log(T(\rho_1, \sigma_1)) + 2T(\rho_2, \sigma_2) \log d \end{aligned}$$

The second term may be estimated in similar fashion. We conclude that if $T(\rho_1, \sigma_1) \leq \varepsilon \leq e^{-1}$ and $T(\rho_2, \sigma_2) \leq \varepsilon \leq e^{-1}$

$$|H^*(\text{spec}(\rho_1), \text{spec}(\rho_2)) - H^*(\text{spec}(\sigma_1), \text{spec}(\sigma_2))| \leq 6\varepsilon \log d - 2\varepsilon \log(\varepsilon). \quad (4.45)$$

We will now show that we can approximate the entropy as would be expected from [Theorem 4.12](#), if we make some additional assumptions (which in particular are satisfied if the state along each edge is a copy of n states, $\phi_e = \phi_{e,0}^{\otimes n}$).

Corollary 4.14. *Let ρ be a random tensor network state satisfying the same assumptions as in [Theorem 4.12](#), and assume additionally that the $\nu_{\phi_{\gamma_{A,i}}}$ have uniformly exponentially decaying tail probabilities and $\varepsilon(n) = \mathcal{O}(n^{-\gamma})$ for $\gamma > 10$. Assume that for each edge $e = (xy)$, the bond dimension is $D_e = 2^{\mathcal{O}(n)}$ and $\text{tr}[\phi_{e,x}^2] \leq 2^{-\Omega(n)}$. Then with high probability*

$$|H(\rho_A) - H^*(\text{spec}(\phi_{\gamma_{A,1}}), \text{spec}(\phi_{\gamma_{A,2}}))| = \mathcal{O}(\log(n)).$$

Proof. The basic proof strategy will be that of [Theorem 4.12](#): we study a slightly reduced problem on the approximated tensor network states $\tilde{\sigma}$ with approximate background states $\tilde{\phi}$, work out the entropies for $\tilde{\sigma}$ and $\tilde{\phi}$, and then argue that the closeness of the resulting entropies will continue to hold for the original tensor network state and background state, up to errors that we carefully keep track of.

First of all, we note that we can reduce to the tensor network state σ on the reduced graph G' , with error as in [Eq. \(4.23\)](#); in particular

$$\mathbb{E}\|\text{spec}_+(\rho_A) - \text{spec}_+(\sigma_B)\|_1 = \mathcal{O}(n^{-\frac{\gamma}{2}}). \quad (4.46)$$

Next we adapt the part of the proof of [Theorem 4.12](#) where we modify the state along the minimal cuts. In the proof of [Theorem 4.12](#), we first observed that if the (regularized) spectrum along each cut $\nu_{\phi_{\Gamma_{A,i}}}$ has uniformly exponentially decaying tail probabilities, then for sufficiently large C , we can slice off the tails with vanishing probability mass:

$$\sum_{\lambda_{i,j} \notin I_n} \lambda_{i,j} = \nu_{\phi_{\Gamma_{A,i}}}((-\infty, -C \log(n)) \cup (C \log(n), \infty)) = \mathcal{O}\left(\frac{1}{n^4}\right).$$

We also binned the eigenvalues of ϕ in the reduced spectrum:

$$q_{i,j} = d_{i,j} p_{i,j},$$

where $p_{i,j}$ are the bins and $d_{i,j}$ is the multiplicity of each bin. We then removed bins that were too small, leading to a state $\tilde{\phi}$. In the notation of the proof of [Theorem 4.12](#) we choose α and β such that $\alpha > 2$ and $\frac{5}{2} + \alpha < \beta \leq \frac{1}{2}(\gamma - 1)$. By [Eq. \(4.26\)](#) this yields an error

$$\|\tilde{\phi} - \phi\|_1 = o(n^{-1}). \quad (4.47)$$

Note that in the proof of [Theorem 4.12](#), we performed one more approximation of removing the “middle” or “diagonal” part of the spectrum and obtained a state $\bar{\phi}$; we will not need to do this here. Now, we recall that the binning of eigenvalues allows us to write $\tilde{\phi}$ as a superposition over maximally-entangled states along each cut:

$$|\tilde{\phi}_{\gamma_{A,1}, \gamma_{A,2}}\rangle = \sum_{j,k} \sqrt{q_{1,j} q_{2,k}} |\Phi_{1,j}^+\rangle \otimes |\Phi_{2,k}^+\rangle,$$

and use this to decompose $\tilde{\phi}$ as

$$|\tilde{\phi}\rangle = \sum_{j,k} \sqrt{q_{1,j} q_{2,k}} |\psi^{(j,k)}\rangle,$$

where in $|\psi^{(j,k)}\rangle$, we have replaced $|\tilde{\phi}_{\gamma_{A,1}, \gamma_{A,2}}\rangle$ with the maximally entangled state $|\Phi_{1,j}^+\rangle \otimes |\Phi_{2,k}^+\rangle$. The states $|\psi^{(j,k)}\rangle$ are normalized background states on the graph G' . We let

$$|\phi^{(j,k)}\rangle = \sqrt{q_{1,j} q_{2,k}} |\psi^{(j,k)}\rangle$$

which are subnormalized states. At this point, the key idea of the argument is straightforward. We will consider the random tensor network states which have (a smoothed version) of the background states $|\psi^{(j,k)}\rangle$. These will have entropy close to $\min(\log(d_{1,j}), \log(d_{2,k}))$. Then we will use [Lemma 4.13](#) to argue that the entropy of ρ_A is approximated up to $\mathcal{O}(\log(n))$ terms by the convex combination $\sum_{j,k} q_{1,j} q_{2,k} \min(\log(d_{1,j}), \log(d_{2,k}))$, which we can relate to the desired result. To make this easy argument precise, we will need to take care of smoothing the background state appropriately and ensure that the relevant states are close to normalized.

We will now argue that we can smoothen the states $\phi^{(j,k)}$. We may assume without loss of generality that the states $\phi^{(j,k)}$ have nonnegative coefficients in the standard basis $|I\rangle = \prod_{e \in E'} |i_e i_e\rangle$ where $I = \{i_e\}_{e \in E'}$ runs over all possible basis elements over each edge, so we can write

$$|\phi^{(j,k)}\rangle = \sum_I \sqrt{\lambda_I^{(j,k)}} |I\rangle.$$

By the same argument as in [Theorem 4.12](#) we find a state

$$\phi^{(j,k,\varepsilon)} = \sum_I \sqrt{\lambda_I^{(j,k,\varepsilon)}}$$

which is such that for any $\Delta_B \in C(B)$ not equal to B or $V'_b \cup B$ and $\Delta_A = \Gamma_{A,1} \cup \Delta_B$

$$H_{\min}(\Delta_B \setminus B|B)_{\phi^{(j,k,\varepsilon)}} \geq H_{\min}^\varepsilon(\Delta_A \setminus \Gamma_A|\Gamma_A)_\phi,$$

while for $\Delta_B = B$ and $\Delta_B = B \cup V'_b$ we have

$$\begin{aligned} H_{\min}(B)_{\phi^{(j,k,\varepsilon)}} &\geq H_{\min}(B)_{\phi^{(j,k)}} = \log \frac{1}{q_{2,k} p_{1,j}} \\ H_{\min}(\bar{B})_{\phi^{(j,k,\varepsilon)}} &\geq H_{\min}(\bar{B})_{\phi^{(j,k)}} = \log \frac{1}{q_{1,j} p_{2,k}}. \end{aligned}$$

Moreover, $\phi^{(j,k,\varepsilon)}$ is close to $\phi^{(j,k)}$ in the sense that

$$\|\phi^{(j,k,\varepsilon)} - \phi^{(j,k)}\|_1 = \mathcal{O}\left(\sqrt{\sum_I |\lambda^{(j,k,\varepsilon)} - \lambda^{(j,k)}|}\right) = \mathcal{O}(\sqrt{\varepsilon}).$$

By the remark after [Lemma 4.11](#) we may assume that $\lambda^{(j,k,\varepsilon)} \leq \lambda^{(j,k)}$.

By a chain rule (e.g. Theorem 5.13 in [\[Tom15\]](#), proven in [\[Dup15\]](#)) for any $\Delta_B \in C(B)$ it holds that

$$H_2(\Delta_B)_{\phi^{(j,k,\varepsilon)}} \geq H_{\min}(\Delta_B \setminus B|B)_{\phi^{(j,k,\varepsilon)}} + H_{\min}(B)_{\phi^{(j,k,\varepsilon)}}.$$

We now let

$$|\phi^\varepsilon\rangle = \sum_{j,k} |\phi^{(j,k,\varepsilon)}\rangle.$$

Then, by [Lemma C.4](#)

$$\|\tilde{\phi} - \phi^\varepsilon\|_1 \leq \sqrt{2 \sum_{j,k} \sum_I |\lambda_I^{(j,k)} - \lambda_I^{(j,k,\varepsilon)}|} = \mathcal{O}(\sqrt{n^{2\alpha} \log(n)^2 \varepsilon(n)}) = \mathcal{O}(n^{\alpha - \frac{1}{2}\gamma} \log(n)) = \mathcal{O}(n^{-3}). \quad (4.48)$$

Therefore, if we let σ_B^ε be the random tensor network state with background state ϕ^ε

$$\mathbb{E}\|\sigma - \sigma^\varepsilon\|_1 \leq \|\phi - \tilde{\phi}\|_1 + \|\tilde{\phi} - \phi^\varepsilon\|_1 = o\left(\frac{1}{n}\right). \quad (4.49)$$

Finally, let

$$|\phi^{(j,k,\varepsilon)}\rangle = \sqrt{q_{1,j} q_{2,k}} |\psi^{(j,k,\varepsilon)}\rangle$$

By [Eq. \(C.2\)](#) in the remark after the proof of [Lemma 4.11](#) we have

$$\begin{aligned} |\mathrm{tr}[\psi^{(j,k,\varepsilon)}] - 1| &= \frac{1}{q_{1,j} q_{2,k}} |\mathrm{tr}[\phi^{(j,k,\varepsilon)}] - \mathrm{tr}[\phi^{(j,k)}]| \\ &= \mathcal{O}\left(\frac{\varepsilon}{q_{1,j} q_{2,k}}\right) = \mathcal{O}(n^{2\beta - \gamma}) = \mathcal{O}(n^{-1}). \end{aligned}$$

using that $q_{1,j}, q_{2,k} > n^{-\beta}$ and $\beta \leq \frac{1}{2}(\gamma - 1)$. We denote the random tensor network states with background states $\psi^{(j,k,\varepsilon)}$ by $\sigma^{(j,k)}$, and the random tensor network states with background states $\sum_j \sqrt{q_{1,j}} \psi^{(j,k,\varepsilon)}$ by $\sigma^{(k)}$.

We introduce the event N , which entails that ρ, σ and $\sigma^{(j,k)}$ for all j, k are close to normalized, that is

$$|\mathrm{tr}[\rho] - 1| \leq \frac{1}{n^2} \quad \text{and} \quad |\mathrm{tr}[\sigma] - 1| \leq \frac{1}{n^2} \quad \text{and} \quad |\mathrm{tr}[\sigma^{(j,k)}] - \mathrm{tr}[\psi^{(j,k,\varepsilon)}]| \leq \frac{1}{n^2}.$$

By assumption, for each edge $e = (xy)$ we have $\mathrm{tr}[\phi_{e,x}^2] \leq 2^{-\Omega(n)}$. Moreover, using that $q_{1,j}, q_{2,k} \geq n^{-\beta}$

$$\mathrm{tr}[(\phi_{\Delta}^{(j,k,\varepsilon)})^2] \leq q_{1,j}^{-2} q_{2,k}^{-2} \mathrm{tr}[\phi_{\Delta}^2] = \mathcal{O}(\mathrm{poly}(n)2^{-\Omega(n)})$$

and therefore the quantity η in [Lemma 2.1](#) is $\mathcal{O}(2^{-\Omega(n)})$ for ρ, σ and $\sigma^{(j,k)}$. So, by [Lemma 2.1](#) and the union bound the event N has probability

$$\begin{aligned} p_N &\geq 1 - \left(\Pr(|\mathrm{tr}[\rho] - 1| \geq \frac{1}{n^2}) + \Pr(|\mathrm{tr}[\sigma] - 1| \geq \frac{1}{n^2}) + \sum_{j,k} \Pr(|\mathrm{tr}[\sigma^{(j,k)}] - \mathrm{tr}[\psi^{(j,k,\varepsilon)}]| \geq \frac{1}{n^2}) \right) \\ &= 1 - \mathcal{O}(\mathrm{poly}(n)2^{-\Omega(n)}) = 1 - \mathcal{O}(2^{-\Omega(n)}). \end{aligned}$$

We denote by \mathbb{E}_N the expectation value over the random tensors conditioned on this event.

We now use [Lemma 4.13](#) to approximate the entropy of σ_B^ε conditioned on N :

$$|H(\sigma_B^\varepsilon) - \sum_k q_{2,k} H(\sigma_B^{(k)})| = \mathcal{O}(\log(n)).$$

Because $\sigma^{(k)}$ is pure, we have $H(\sigma_B^{(k)}) = H(\sigma_B^{(k)})$. We apply [Lemma 4.13](#) again, this time to the decomposition of $\sigma_B^{(k)}$ to see:

$$|H(\sigma_B^{(k)}) - \sum_j q_{1,j} H(\sigma_B^{(j,k)})| = \mathcal{O}(\log(n)).$$

Thus,

$$|H(\sigma_B^\varepsilon) - \sum_{j,k} q_{1,j} q_{2,k} H(\sigma_B^{(j,k)})| = \mathcal{O}(\log(n)).$$

Since $\mathrm{rank}(\sigma_B^{(j,k)}) \leq \min\{d_{1,j}, d_{2,k}\}$ (and taking into account the normalization of $\sigma^{(j,k)}$)

$$H(\sigma_B^{(j,k)}) \leq \mathrm{tr}[\sigma^{(j,k)}] \min(\log d_{1,j}, \log d_{2,k}) - \mathrm{tr}[\sigma^{(j,k)}] \log \mathrm{tr}[\sigma^{(j,k)}].$$

Conditioned on N and using $|\mathrm{tr}[\psi^{(j,k,\varepsilon)}] - 1| = \mathcal{O}(n^{-1})$ this implies

$$H(\sigma_B^{(j,k)}) \leq \min(\log d_{1,j}, \log d_{2,k}) + \mathcal{O}(1).$$

For a lower bound we use that

$$\begin{aligned} \mathbb{E}_N H(\sigma_B^{(j,k)}) &\geq \mathbb{E}_N \mathrm{tr}[\sigma^{(j,k)}] \left(-\log \mathrm{tr}[(\sigma_B^{(j,k)})^2] + \log \mathrm{tr}[\sigma_B^{(j,k)}] \right) \\ &\geq (\mathrm{tr}[\psi^{(j,k,\varepsilon)}] - \frac{1}{n^2}) \left(-\log \mathbb{E}_N \mathrm{tr}[(\sigma_B^{(j,k)})^2] + \log(\mathrm{tr}[\psi^{(j,k,\varepsilon)}] - \frac{1}{n^2}) \right) \\ &\geq -\log \mathbb{E}_N \mathrm{tr}[(\sigma_B^{(j,k)})^2] - \mathcal{O}(1) \end{aligned}$$

where in the first and second inequality we have used Jensen's inequality, and in the second and third inequality we have used $\mathrm{tr}[\sigma_B^{(j,k)}] \geq \mathrm{tr}[\psi^{(j,k,\varepsilon)}] - \frac{1}{n^2} \geq 1 - \mathcal{O}(\frac{1}{n})$. We use the replica trick to estimate

$$\mathbb{E} \mathrm{tr}[(\sigma_B^{(j,k)})^2] = \sum_{\Delta_B \in C(B)} \mathrm{tr}[(\psi_{\Delta_B}^{(j,k,\varepsilon)})^2]$$

In this expression, we have contributions from $\Delta_B = B$ and $\Delta_B = B \cup V'_b$, which yield contributions

$$\begin{aligned}\mathrm{tr}[(\psi_B^{(j,k,\varepsilon)})^2] &\leq d_{1,j}^{-1} \\ \mathrm{tr}[(\psi_{\bar{B}}^{(j,k,\varepsilon)})^2] &\leq d_{2,k}^{-1}.\end{aligned}$$

For any other cut Δ_B , we have a contribution at most

$$\begin{aligned}\mathrm{tr}[(\psi_{\Delta_B}^{(j,k,\varepsilon)})^2] &= q_{1,j}^{-2} q_{2,k}^{-2} \mathrm{tr}[\phi^{(j,k,\varepsilon)}] 2^{-H_{\min}(B)_{\phi^{(j,k,\varepsilon)}} - H_{\min}^{\varepsilon}(\Delta_A \setminus \Gamma_A | \Gamma_A)_{\phi}} \\ &\leq q_{1,j}^{-1} q_{2,k}^{-1} 2^{-H_{\min}(B)_{\phi^{(j,k,\varepsilon)}}} \leq d_{1,j}^{-1}\end{aligned}$$

using that $\mathrm{tr}[\phi^{(j,k,\varepsilon)}] \leq q_{1,j} q_{2,k}$ and $2^{-H_{\min}(B)_{\phi^{(j,k,\varepsilon)}}} \leq q_{2,k} p_{1,j} = q_{1,j} q_{2,k} d_{1,j}^{-1}$. Therefore

$$\begin{aligned}\mathbb{E}_N \mathrm{tr}[(\sigma_B^{(j,k,\varepsilon)})^2] &\leq p_N^{-1} \mathbb{E} \mathrm{tr}[(\sigma_B^{(j,k,\varepsilon)})^2] \\ &= d_{1,j}^{-1} + d_{2,k}^{-1} + \mathcal{O}(d_{1,j}^{-1}).\end{aligned}$$

so

$$\begin{aligned}-\log \mathbb{E}_N \mathrm{tr}[(\sigma_B^{(j,k,\varepsilon)})^2] &\geq -\log \left(\max(d_{1,j}^{-1}, d_{2,k}^{-1}) (2 + \mathcal{O}(1)) \right) \\ &\geq \min(\log d_{1,j}, \log d_{2,k}) - \mathcal{O}(1).\end{aligned}$$

We find that

$$\mathbb{E}_N |H(\sigma_B^{(j,k)}) - \min(\log d_{1,j}, \log d_{2,k})| = \mathcal{O}(1)$$

and hence

$$\mathbb{E}_N \left| \sum_{j,k} q_{1,j} q_{2,k} H(\sigma_B^{(j,k)}) - \sum_{j,k} q_{1,j} q_{2,k} \min(\log d_{1,j}, \log d_{2,k}) \right| = \mathcal{O}(1). \quad (4.50)$$

We collect the various estimates we have found:

$$\begin{aligned}\mathbb{E}_N \|\mathrm{spec}_+(\rho_A) - \mathrm{spec}_+(\sigma_B)\|_1 &\leq p_N^{-1} \mathbb{E} \|\mathrm{spec}_+(\rho_A) - \mathrm{spec}_+(\sigma_B)\|_1 = o\left(\frac{1}{n}\right) \\ \mathbb{E}_N \|\sigma_B - \sigma_B^{\varepsilon}\|_1 &\leq p_N^{-1} \mathbb{E} \|\sigma_B - \sigma_B^{\varepsilon}\|_1 = o\left(\frac{1}{n}\right) \\ \mathbb{E}_N \left| \sum_{j,k} q_{1,j} q_{2,k} H(\sigma_B^{(j,k)}) - \sum_{j,k} q_{1,j} q_{2,k} \min(\log d_{1,j}, \log d_{2,k}) \right| &= \mathcal{O}(1).\end{aligned}$$

Let M be the event that N holds and moreover

$$\begin{aligned}\|\mathrm{spec}_+(\rho_A) - \mathrm{spec}_+(\sigma_B)\|_1 &\leq \frac{1}{n} \\ \|\sigma_B - \sigma_B^{\varepsilon}\|_1 &\leq \frac{1}{n} \\ \left| \sum_{j,k} q_{1,j} q_{2,k} H(\sigma_B^{(j,k)}) - \sum_{j,k} q_{1,j} q_{2,k} \min(\log d_{1,j}, \log d_{2,k}) \right| &\leq \log(n).\end{aligned}$$

By Markov's inequality and the union bound, the probability that M holds goes to one as n goes to infinity. Moreover, if M holds, it is easy to verify that by the Fannes-Audenaert inequality [Eq. \(4.44\)](#) and the fact that ρ , σ and σ^{ε} are close to normalized

$$|H\left(\frac{\rho_A}{\mathrm{tr}[\rho]}\right) - H(\sigma_B^{\varepsilon})| = \mathcal{O}(1).$$

Also, if M holds, by Eq. (4.50)

$$\begin{aligned} |H(\sigma_B^\varepsilon) - \sum_{j,k} q_{1,j} q_{2,k} \min(\log d_{1,j}, \log d_{2,k})| &= \left| \sum_{j,k} q_{1,j} q_{2,k} H(\sigma_B^{(j,k)}) - \sum_{j,k} q_{1,j} q_{2,k} \min(\log d_{1,j}, \log d_{2,k}) \right| \\ &\quad + \mathcal{O}(\log(n)) \\ &= \mathcal{O}(\log(n)) \end{aligned}$$

so we conclude that

$$\left| H\left(\frac{\rho_A}{\text{tr}[\rho]}\right) - \sum_{j,k} q_{1,j} q_{2,k} \min(\log d_{1,j}, \log d_{2,k}) \right| = \mathcal{O}(\log(n)).$$

Since $q_{i,j} = d_{i,j} p_{i,j} \geq n^{-\beta}$ from the binning procedure, we have the simple observation

$$\frac{1}{n^\beta p_{i,j}} \leq \frac{q_{i,j}}{p_{i,j}} = d_{i,j} \leq \frac{1}{p_{i,j}},$$

and hence $|\min(\log d_{1,j}, \log d_{2,k}) - \log(\min(\frac{1}{p_{1,k}}, \frac{1}{p_{2,j}}))| = \mathcal{O}(\log(n))$. We conclude that

$$\left| H\left(\frac{\rho_A}{\text{tr}[\rho]}\right) - \sum_{j,k} q_{1,k} q_{2,j} \log\left(\min\left(\frac{1}{p_{1,k}}, \frac{1}{p_{2,j}}\right)\right) \right| = \mathcal{O}(\log(n)).$$

Finally, we need to relate the result back to the original background state. In the above approximation to $H(\tilde{\sigma}_B)$, we see that

$$\sum_{j,k} q_{1,k} q_{2,j} \log\left(\min\left(\frac{1}{p_{1,k}}, \frac{1}{p_{2,j}}\right)\right) = H^*(\text{spec}(\tilde{\phi}_{\gamma_{A,1}}), \text{spec}(\tilde{\phi}_{\gamma_{A,2}})).$$

Then by Eq. (4.47) and Eq. (4.45), this will converge to the appropriate quantity on the non-approximated background state:

$$|H^*(\text{spec}(\tilde{\phi}_{\gamma_{A,1}}), \text{spec}(\tilde{\phi}_{\gamma_{A,2}})) - H^*(\text{spec}(\phi_{\gamma_{A,1}}), \text{spec}(\phi_{\gamma_{A,2}}))| \rightarrow 0,$$

proving the desired result. □

Acknowledgments

NC is supported in part by the Department of Energy via the GeoFlow consortium (QuantISED Award DE-SC0019380). CL acknowledges support from the projects ESQuisses (ANR-20-CE47-0014-01), STARS (ANR-20-CE40-0008), Qtraj (ANR-20-CE40-0024-01) and Random Tensors (ANR-11-LABX-0040) of the French National Research Agency (ANR). GP is supported by the UC Berkeley Physics Department, the Simons Foundation through the ‘‘It from Qubit’’ program, the Department of Energy via the GeoFlow consortium (QuantISED Award DE-SC0019380), and AFOSR award FA9550-22-1-0098. He also acknowledges support from an IBM Einstein Fellowship at the Institute for Advanced Study. MW acknowledges support by the NWO through grant OCENW.KLEIN.267, by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy - EXC 2092 CASA - 390781972, by the BMBF through project Quantum Methods and Benchmarks for Resource Allocation (QuBRA), and by European Research Council (ERC). Funded by the European Union. Views and opinions expressed are however those of the authors only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them.

A Euclidean gravity path integrals and entanglement spectra

In this appendix, we give a heuristic description of certain Euclidean gravity path integrals in holography whose descriptions are in close analogy with random tensor network models. This section serves as a motivation for the random tensor network models we study, but is not needed to understand the random tensor network results. In [Appendix A.1](#), we review the replica trick in quantum field theory and the role of Euclidean path integrals. We review an application of such tools in [Appendix A.2](#), we discuss the problem of studying the entanglement entropy near a phase transition between two minimal surfaces, which agrees with results in [Section 4](#). Then, in [Appendix A.3](#), we discuss a simplified model of quantum gravity which is in very close correspondence to a random tensor network model with link states with bounded spectral variation, as in [Section 3](#).

A.1 The replica trick and Euclidean path integrals

In order to compute entropies in quantum field theory, one often uses a version of the *replica trick* to compute the Rényi entropies, which can then be analytically continued to deduce von Neumann entropies. We consider a pure quantum field theory state $|\rho\rangle$ on a space M , which is prepared by a Euclidean path integral on $M \times (-\infty, 0]$. Correspondingly, $\langle\rho|$ is prepared by the time-reflected path integral on $M \times [0, \infty)$. Let A be a subregion of M . The reduced density matrix on A is given by taking $|\rho\rangle$ and $\langle\rho|$, then integrating over the field configurations on the complement of A (the equivalent of the partial trace for field theories), as shown in [Figure 6a](#). Analogous to [Eq. \(2.8\)](#), we may now compute $\text{tr}[\rho_A^k]$ by taking k copies of this path integral, and gluing the boundaries at the A system cyclically, then integrating over the field configurations at each boundary. This operation is manifestly invariant under cyclic permutations, a symmetry called *replica symmetry*. This is illustrated in [Figure 6b](#). We conclude that $\text{tr}[\rho_A^k]$ is computed by a path integral $Z_{A,k}$ on a manifold $M_{A,k}$, allowing us to compute $H_k(\rho_A)$. The space $M_{A,k}$ is a k -fold cover of $M \times \mathbb{R}$, branching at the boundary ∂A of the subregion A .

Of course, the path integral $Z_{A,k}$ is formally infinite. One ‘normalizes’ the path integral by normalizing by Z_1 which can be thought of as $\text{tr}[\rho]$. We have the expression for the k -th Rényi entropy:

$$H_k(\rho_A) = \log \frac{Z_{A,k}}{Z_1^k}. \quad (\text{A.1})$$

Note that this is a slightly different normalization convention (the denominator is Z_1^k instead of Z_1) to match the standard convention in the quantum gravity literature. To get a finite result, one has to impose a UV cut-off of size ε , an aspect we will ignore in our discussion. By employing such a cutoff, we will pretend that the relevant Hilbert spaces are finite-dimensional Hilbert spaces that factorize with respect to spatial decompositions of M .

What happens if the quantum field theory is a *holographic CFT*? In this case, we have a correspondence between the path integral of the CFT on the one hand, and a bulk quantum gravity path integral on the other hand, which, for large effective central charge, we may approximate by its semiclassical saddle point geometry. This can be used to derive the *Ryu-Takayanagi formula* [\[LM13\]](#), by finding a bulk manifold $B_{A,k}$ which has $M_{A,k}$ as its boundary. In other words, one obtains the gravitational dual by setting $M_{A,k}$ as a boundary condition, then allowing the system to evolve according to the gravitational theory.

The saddle point solution will be such that the bulk copies are glued along a cyclic permutation in an area adjacent to A and along the identity permutation in an area adjacent to \bar{A} , the boundary complement of A . The boundary between these two regions is a surface $\gamma_{A,k}$ anchored at the boundary at ∂A . After orbifolding with respect to the replica symmetry, this leads to a bulk spacetime which has M as its boundary and a ‘cosmic brane’ at the surface $\gamma_{A,k}$. The orbifold procedure gives rise to a conical deficit, corresponding to the fixed points under the action of the replica symmetry. Comparing the action of $B_{A,k}$ and B_1 , we find that in the computation of [Eq. \(A.1\)](#), only the conical deficit at $\gamma_{A,k}$ does not cancel, yielding

$$H_k(\rho_A) \approx \frac{|\gamma_{A,k}|}{4G_N}. \quad (\text{A.2})$$




Figure 6: The path integral replica trick to compute $\text{tr}[\rho_A^k]$.

where $|\gamma_{A,k}|$ is the area of the brane in the saddle-point solution. This can be analytically continued to non-integer values, and in particular, continuation to $k = 1$ yields the RT formula:

$$H(\rho_A) \approx \frac{|\gamma_A|}{4G_N}. \quad (\text{A.3})$$

where γ_A is the surface of minimal area out of all bulk surface homologous to A and anchored at ∂A .

The spectrum of ρ_A can be recovered from the Rényi entropies. In Eq. (A.2), we see that $|\gamma_{A,k}|$ depends on k , but is otherwise fixed as we let G_N go to zero in the classical limit. It was argued in [BPSW19] that this behavior implies a spectrum that is flat at leading order:

$$\begin{aligned} H_{\min}^{\varepsilon}(\rho_A) &= H(\rho_A) - \mathcal{O}(G_N^{-\frac{1}{2}}) = H(\rho_A) - \mathcal{O}(\sqrt{H(\rho_A)}) \\ H_{\max}^{\varepsilon}(\rho_A) &= H(\rho_A) + \mathcal{O}(G_N^{-\frac{1}{2}}) = H(\rho_A) + \mathcal{O}(\sqrt{H(\rho_A)}) \end{aligned}$$

This is precisely the link state regime we investigated in Section 4.4, and shows that the large c limit in a holographic CFT is similar to considering the many-copy limit in quantum information theory.

A.1.1 Fixed area states

A useful variation on the derivation of the RT formula is to consider *fixed-area states* [DHM19]. We consider the ‘area operator’ $\hat{\gamma}_A$ for a subsystem A , which measures the area of a minimal surface. The operator $\hat{\gamma}_A$ actually has fluctuations when we consider a state with a semiclassical gravity dual, and we may write

$$|\rho\rangle = \int d\alpha |\psi_{\alpha}\rangle$$

where ψ_{α} is an eigenvector of \hat{A} with eigenvalue α . The state $|\psi_{\alpha}\rangle$ is a fixed-area state: while it is not itself a physical state, they form a basis with which to construct physical states. These fixed-area states can be thought of as prepared by a bulk path integral where we have restricted to bulk geometries for which $|\gamma_A| = \alpha$. The same derivation as above now leads to

$$H_k(\rho_A) \approx \frac{\alpha}{4G_N},$$

since the area of the minimal surface is fixed to be α . Thus, for fixed-area states, all Rényi entropies are (to good approximation) equal, which implies that the state has flat entanglement spectrum. This corresponds to a random tensor network state with maximally entangled link states.

A.2 States at a minimal surface phase transition

Consider, again, a holographic CFT state ρ_A on a boundary subregion A . In the case where there is a unique minimal surface, the previous subsection showed how the replica trick leads to the RT formula.

We would now like to investigate the entanglement spectrum of ρ_A when there are two RT surfaces for A that have area of the same order of magnitude. Let us denote the two competing minimal surfaces in the bulk by $\gamma_{A,1}$ and $\gamma_{A,2}$. In [MWW20], it was shown how the entanglement entropy should behave at this phase transition between the two minimal surfaces. A similar computation was performed in [AP20] for the setting with two competing minimal surfaces and bulk matter. We will briefly sketch their argument, referring the interested reader to [AP20] for more details.

A.2.1 Fixed area states with two minimal surfaces

We begin by considering bipartite fixed-area states $\rho_{A\bar{A}}$ prepared by a Euclidean path integral, in which we have fixed the size of the two competing surfaces $\gamma_{A,1}$ and $\gamma_{A,2}$. In this case, saddle points of the path integral have to satisfy the equations of motion everywhere except at the surfaces $\gamma_{A,i}$, where there could be conical singularities. The two surfaces divide the bulk into 3 regions: a_1 , a_2 and a_3 .

The saddle points are states with smooth geometries in the regions where the copies of regions a_i are glued to each other – the k copies of region a_1 are glued cyclically, while the copies of region a_3 are glued along the identity permutation. On the middle region a_2 , we are free to glue along an arbitrary permutation π , giving solutions that can break the replica symmetry. However, it is often the case that the dominant solutions of the path integral are those that respect the replica symmetry, and we assume this to be the case for all solutions we consider.

Let us denote a saddle-point solution of the path integral with permutation π by $B_{A,\pi}$, and let us write ϕ_i for the conical singularity angle at $\gamma_{A,i}$. It turns out that these saddle points lead to an action of the form

$$I(B_{A,\pi}) = kI_{\text{away}}(B_{A,\pi}) + (k\phi_1 - 2\pi|C(\pi)|)\frac{|\gamma_{A,1}|}{8\pi G_N} + (k\phi_2 - 2\pi|C(\tau^{-1}\pi)|)\frac{|\gamma_{A,2}|}{8\pi G_N},$$

where I_{away} is the action away from the surfaces, $|\gamma_{A,i}|$ is the area of the surface $\gamma_{A,i}$, and we recall that $|C(\pi)|$ is the number of cycles of π , and τ is the full cycle $(12\dots k)$. In particular, for $k = 1$, we have

$$I(B_{A,\pi}) = I_{\text{away}}(B_{A,\pi}) + (\phi_2 - 2\pi)\frac{|\gamma_{A,1}|}{8\pi G_N} + (\phi_2 - 2\pi)\frac{|\gamma_{A,2}|}{8\pi G_N},$$

so when we look at the normalized path integral, and sum over all permutations

$$\frac{Z_{A,k}}{(Z_{A,1})^k} \approx \sum_{\pi \in S_k} e^{(|C(\pi)|-k)\frac{|\gamma_{A,1}|}{4G_N} + (|C(\tau^{-1}\pi)|-k)\frac{|\gamma_{A,2}|}{4G_N}} = \sum_{\pi \in S_k} e^{d(\text{id},\pi)\frac{|\gamma_{A,1}|}{4G_N} + d(\pi,\tau)\frac{|\gamma_{A,2}|}{4G_N}}.$$

Note that the areas $\gamma_{A,i}$ are of the same order of magnitude, and are divergent. As a result, only the permutations for which $d(\text{id},\pi) + d(\pi,\tau)$ is minimal will contribute. In other words, only the configurations where $\pi \in NC(k)$ contribute, as all other permutations are suppressed by at least a factor of the area of $\gamma_{A,i}$ in the action. We conclude that

$$\frac{Z_{A,k}}{(Z_{A,1})^k} \approx \sum_{\pi \in NC(k)} e^{d(\text{id},\pi)\frac{|\gamma_{A,1}|}{4G_N} + d(\pi,\tau)\frac{|\gamma_{A,2}|}{4G_N}}. \quad (\text{A.4})$$

This computation is in one-to-one correspondence with the computation of the moments for a subsystem of a single random tensor, as observed in [PSSY19]. It also corresponds more generally to a random tensor

network computation with two minimal cuts and maximally entangled link states, as is clear from the computations in [Section 2](#) and [Section 3](#). This is in agreement with the claim that random tensor network states are a model for fixed area states. One can also add bulk matter in this path integral computation, which will again be in correspondence to a similar computation in a random tensor network [\[AP20\]](#) with a background state, as in [Section 2.3](#). From the moment computation in [Eq. \(A.4\)](#) and applying the results for the entanglement of a single random tensor, we observe that for two fixed surfaces of exactly equal size, the (appropriately scaled) entanglement spectrum is a Marchenko-Pastur distribution, giving an $\mathcal{O}(1)$ correction to the entanglement entropy, agreeing with the gravitational replica trick computation in [\[MWW20\]](#).

A.2.2 General states at the minimal surface phase transition

We now relax the fixed-area restriction, and study similar calculations performed in [\[DHM19\]](#), [\[MWW20\]](#), and [\[AP20\]](#). Denote by $Z_{A,k}(\alpha_1, \alpha_2)$ the path integral where we have fixed the areas of $\gamma_{A,i}$ to be α_i . Then, following [Section 2.3](#) in [\[DHM19\]](#), the full path integral is given by

$$Z_{A,k} = \int d\alpha_1 d\alpha_2 Z_k(\alpha_1, \alpha_2).$$

Again, we consider the semiclassical limit, so we take our saddle-point approximation of $Z_k(\alpha_1, \alpha_2)$ in [Eq. \(A.4\)](#), and we also take a saddle-point approximation for the integral over α_1 and α_2 . This saddle point will be at the values for α_i where the deficit angles are $\phi_i = \frac{2\pi}{n}$ (since then the saddle point geometry is smooth), which leads to

$$\frac{Z_{A,k}}{(Z_{A,1})^k} \approx \sum_{\pi \in NC(k)} e^{(|C(\pi)|-k)\frac{|\gamma_{A,1}^{(k)}|}{4G_N} + (|C(\tau^{-1}\pi)|-k)\frac{|\gamma_{A,2}^{(k)}|}{4G_N}} \quad (\text{A.5})$$

where $\gamma_{A,i}^{(k)}$ are now minimal surfaces, with a dependence on k . Analytic continuation to $k = 1$ yields the usual surface prescription. In particular, if there are two surfaces that are of almost equal area, the contribution of the larger term is exponentially suppressed for any $\mathcal{O}(1)$ or larger difference in areas.

To zoom in on the region where the two surfaces are nearly equal, we write the state as a superposition of fixed area states, as in [Section 3](#) of [\[MWW20\]](#). We discretize the area size α_1 and α_2 over $\text{poly}(\frac{1}{G_N})$ values and approximate the full holographic pure state with boundary regions A and $B = A^c$ as a (finite) sum

$$|\psi\rangle_{AB} = \sum_{\alpha_1, \alpha_2} \sqrt{p(\alpha_1, \alpha_2)} |\psi_{\alpha_1, \alpha_2}\rangle$$

where $|\psi_{\alpha_1, \alpha_2}\rangle$ is the state where the areas are fixed as $|\gamma_{A,i}| = \alpha_i$, and p is a probability distribution over the possible areas. Then a straightforward calculation of the reduced density matrix ρ_A yields a state of the form:

$$\begin{aligned} \rho_A &= \sum_{\alpha_1, \alpha_2} p(\alpha_1, \alpha_2) \rho_{A, \alpha_1, \alpha_2} + \sum_{\alpha_1 \neq \alpha'_1, \alpha_2 \neq \alpha'_2} \sqrt{p(\alpha_1, \alpha_2) p(\alpha'_1, \alpha'_2)} \text{tr}_B (|\psi_{\alpha_1, \alpha_2}\rangle \langle \psi_{\alpha'_1, \alpha'_2}|), \\ &= \sum_{\alpha_1, \alpha_2} p(\alpha_1, \alpha_2) \rho_{A, \alpha_1, \alpha_2} + OD_A, \end{aligned}$$

where OD_A are the off-diagonal elements of ρ_A . One can argue that the states $\rho_{A, \alpha_1, \alpha_2}$ are all mutually orthogonal by entanglement wedge reconstruction – the area operator can be reconstructed on A , and hence, each $\rho_{A, \alpha_1, \alpha_2}$ is perfectly distinguishable from each other by measuring the area operator. Then the entropy of the diagonal part of the state is easily computed as

$$H\left(\sum_{\alpha_1, \alpha_2} p(\alpha_1, \alpha_2) \rho_{A, \alpha_1, \alpha_2}\right) = \sum_{\alpha_1, \alpha_2} p(\alpha_1, \alpha_2) H(\rho_{A, \alpha_1, \alpha_2}) - \sum_{\alpha_1, \alpha_2} p(\alpha_1, \alpha_2) \log p(\alpha_1, \alpha_2). \quad (\text{A.6})$$

The second term is the so-called entropy of mixing, and it is a standard argument that this term is suppressed relative to the first term [MWW20] as $\mathcal{O}(\ln G_N)$ or smaller. The entropies appearing in the first term can be computed using the methods in the previous subsection, for which one finds that $H(\rho_{A,\alpha_1,\alpha_2}) = \frac{\min\{\alpha_1,\alpha_2\}}{4G_N}$.

Returning to the off-diagonal terms OD_A , [MWW20] argued that such terms should be subleading in the analytic continuation due to the relevant surfaces breaking replica symmetry. At the same time, [AP20] argued that such terms should be subleading due to reasons similar to those for the orthogonality of the diagonal elements: complementary entanglement wedge reconstruction implies one may reconstruct the bulk area operator on B , and hence, such states are perfectly distinguishable on B . Therefore, the partial trace over B vanishes for $\alpha_1 \neq \alpha'_1, \alpha_2 \neq \alpha'_2$.

One reaches the conclusion:

$$H(\rho_A) = \sum_{\alpha_1,\alpha_2} p(\alpha_1,\alpha_2) \frac{\min\{\alpha_1,\alpha_2\}}{4G_N} + \mathcal{O}(\ln G_N), \quad (\text{A.7})$$

In this computation the $\mathcal{O}(1)$ corrections due to the Marchenko-Pastur distribution along each pair of minimal cuts of equal size (or equivalently, the degeneracy in the contributions to the saddle point approximation) is irrelevant, as the entropy of mixing already leads to $\mathcal{O}(\ln G_N)$ deviations.

Our results in Section 4 can be seen as a rigorous version of the above result for random tensor networks.

A.3 Replica wormholes and JT gravity

One of the most basic holographic models of quantum gravity is *JT gravity*, a 1 + 1-dimensional model of gravity; see [Sár17] for a review. JT gravity is also a useful model for the near-horizon dynamics of extremal black holes in any dimension. In this case, the dual theory should be 0 + 1-dimensional. In other words, it should be regular quantum mechanics rather than a quantum field theory. Indeed, in [SSS19], it was shown that JT gravity theory is dual to a random matrix model, where the Hamiltonian is a random self-adjoint matrix according to some distribution, providing another strong connection between quantum gravity and random matrix theory. It also appears that such gravitational systems may be dual to an ensemble of boundary theories [BW20], rather than a single one. Whether this is fundamental, a special feature of 1+1-dimensional models, or due to averaging over microscopic features of the gravity theory, is a line of active research [SSSY21].

We now sketch a variation on a calculation in [PSSY19], providing a proof-of-principle that the free probability techniques used in Section 3 provide an elegant framework to understand such results. We refer the interested reader to [PSSY19] for more in-depth motivation and detailed computations.

We consider JT gravity with an end of the world (EOW) brane containing a large number n of internal states. This model has action

$$I = I_{\text{JT}} + \mu \int_{\text{brane}} ds,$$


where the action of a manifold M with metric g , induced boundary metric h , (trace of) extrinsic curvature K , and dilaton ϕ is given by

$$I_{\text{JT}}[M, g] = -\frac{S_0}{2\pi} \left[\frac{1}{2} \int_M \sqrt{g} R + \int_{\partial M} \sqrt{h} K \right] - \left[\frac{1}{2} \int_M \sqrt{g} \phi (R + 2) + \int_{\partial M} \sqrt{h} \phi K \right].$$

The details of this action are not very important for us; we just note that we will take the S_0 parameter to be large, and that this suppresses contributions where the manifold M has genus $\gamma > 0$ in the Euclidean path integral.

Such systems are of interest when studying a simple version of an evaporating black hole. Let

$$|\psi\rangle = \frac{1}{\sqrt{n}} \sum_{i=1}^n |\psi_{B,i}\rangle |i_{\text{Rad}}\rangle$$



(a) Path integral which prepares the state $|\psi_i\rangle$. (b) Replica trick for JT gravity. On the left side two diagrams that show up when computing matrix elements of $\psi_{\text{Rad}}^{\otimes 2}$, on the right a diagram contributing a factor $Z_2^2 Z_1 m_2^2 m_1$ to $\text{tr}[\psi_{\text{Rad}}^5]$.

Figure 7: Path integrals and the replica trick for JT gravity. See [PSSY19] for a detailed explanation of the diagrammatic notation.

where $|\psi_{B,i}\rangle$ is the state of the black hole with the EOW brane in state i , and $|i_{\text{Rad}}\rangle$ is a reference state, which can be thought of as the radiation system. Notice that the entanglement spectrum of this state is flat. We generalize this to

$$|\psi\rangle = \sum_{i=1}^n \sqrt{p_i} |\psi_{B,i}\rangle |i_{\text{Rad}}\rangle,$$

where the entanglement between the black hole and the radiation has some nontrivial spectrum, which we will assume to be close to uniform, so that $\frac{p_i}{n}$ is bounded by a constant for all i . In other words, we assume this distribution satisfies the bounded spectral variation assumption from Section 3. We let

$$m_k = \sum_{i=1}^n n^{k-1} p_i^k$$

be the (appropriately scaled) moments of the entanglement spectrum of the EOW brane. Moreover, we write the path integral on a disc geometry with k boundary components and k EOW branes as $e^{-S_0} Z_k$. Then following the arguments of [PSSY19], one can compute the k -th moment of the radiation system for large n and large e^{S_0} (large n enforces a planar limit with only non-crossing partitions, while large e^{S_0} ensures that only genus $\gamma = 0$ geometries contribute), as illustrated in Figure 7b. The contributions of path integral configurations connecting different replicas are called *replica wormholes*. This diagrammatic computation shows that

$$\text{tr}[\psi_{\text{Rad}}^k] = \sum_{\pi \in \text{NC}(k)} m_\pi \frac{Z_{\pi^{-1}\tau}}{Z_1^k} n^{-d(\pi, \text{id})} e^{-S_0 d(\pi, \tau)}. \quad (\text{A.8})$$

In this expression, recall that $Z_\sigma = \prod_{l \in C(\sigma)} Z_l$, where $C(\sigma)$ is the cycle type of σ , and $l \in C(\sigma)$ are the lengths of the cycles of σ . This expression implies that if $n \gg e^{S_0}$, the dominant contribution in Eq. (A.8) is given by $\pi = \tau$. On the other hand, if $n \ll e^{S_0}$, the dominant contribution in Eq. (A.8) is given by $\pi = \text{id}$. This corresponds to the situation where there is a unique minimal surface (more precisely, a unique *quantum extremal surface*). We are interested in the regime at the phase transition, which is analogous to the Page time of an evaporating black hole, so we assume $ne^{-S_0} \rightarrow 1$. The coefficients m_σ correspond to the weight of the σ configuration, as determined by the number and length of the cycles in σ , and the probability distribution of eigenstates p_i . In the case of the flat entanglement spectrum, this number equals the number of closed loops

between the connected components. This will also be the case for the non-trivial entanglement spectrum, but each loop will have a different weight that depend on the p_i 's.

The m_k are the (scaled) moments of a probability distribution. While the explicit expression itself is not important for our purposes, the Z_l can be written as the l -th moments of a probability distribution [PSSY19]. Hence, Eq. (A.8) is a product of moments, summed over all non-crossing partitions of length k . As a result, $\text{tr}[\psi_{\text{Rad}}^k]$ can be calculated in the planar limit very simply by way of free probability theory.

More precisely, we may define moment-generating functions for EOW, JT, and Rad:

$$\begin{aligned} M_{\text{JT}}(z) &= \sum_{k=1}^{\infty} \frac{Z_k}{Z_1^k} z^k, \\ M_{\text{EOW}}(z) &= \sum_{k=1}^{\infty} m_k z^k, \\ M_{\text{Rad}} &= \sum_{k=1}^{\infty} e^{S_0(k-1)} \text{tr}[\psi_{\text{Rad}}^k] z^k. \end{aligned}$$

Given a moment generating function $M(z)$, which is a formal power series, recall that its S-transform is given by

$$S(z) = \frac{1+z}{z} M^{-1}(z).$$

We use this to define the S-transforms S_{JT} , S_{EOW} and S_{Rad} . From Theorem 3.2, we see that the relation between the moments in Eq. (A.8) implies that these are related as

$$S_{\text{Rad}}(z) = \frac{1}{1+z} S_{\text{JT}}(z) S_{\text{EOW}}(z). \quad (\text{A.9})$$

This means that, at the phase transition where $n \approx e^{S_0}$, the spectrum of ψ_{Rad} can be described as a free product of the spectra of the end of the world brane, the JT gravity spectrum and a Marchenko-Pastur distribution.

A.3.1 A recursion relation for the resolvent

Given a moment generating function $M(z)$, we may also define the *resolvent function* $R(z)$ by

$$R(z) = \frac{1}{z} \left(1 + M \left(\frac{1}{z} \right) \right).$$

To relate to previous results, we consider the case where the entanglement with the radiation is maximally entangled. In this case, $S_{\text{EOW}}(z) = 1$ and $S_{\text{Rad}}(z) = \frac{1}{1+z} S_{\text{JT}}(z)$. By definition of the S-transform and setting $z \rightarrow M_{\text{Rad}}(z)$, this implies

$$\frac{1}{1 + M_{\text{Rad}}(z)} S_{\text{JT}}(M_{\text{Rad}}(z)) = \frac{1 + M_{\text{Rad}}(z)}{M_{\text{Rad}}(z)} z,$$

which we may rewrite as (again using the definition of the S-transform):

$$M_{\text{Rad}}(z) = M_{\text{JT}}[z(1 + M_{\text{Rad}}(z))].$$

In terms of the resolvent, this becomes

$$R(z) = \frac{1}{z} + \frac{1}{z} M_R \left(\frac{1}{z} \right)$$

$$\begin{aligned}
&= \frac{1}{z} + \frac{1}{z} M_{\text{JT}}(R(z)) \\
&= \frac{1}{z} + \sum_{k=1}^{\infty} \frac{Z_k}{Z_1^k} \frac{R(z)^k}{z},
\end{aligned}$$

which is a recursion relation previously derived in [PSSY19] by a diagrammatic argument. More generally, we can interpret Eq. (A.9) as a (complicated) recursion relation that directly generalizes the above recursion relation.

B Random tensor network states and split transfer protocols

Our results involving general background states are closely related to the quantum-information-theoretic task of *split transfer* introduced in [DH10], which can be understood as a variant of quantum state merging. The standard setup is as follows: two parties, Alice and Bob, share a state $\phi_{ABC_1\dots C_m}$, with Alice controlling A , Bob controlling B , and the systems C_1, \dots, C_m being m “helpers”. Let R be a reference system and $\phi_{ABRC_1\dots C_m}$ a purification of ϕ . Initially, the state is shared not only by Alice and Bob, but also with all the helper systems C_i . The goal of split transfer is to try to redistribute the state to Alice, Bob, and R , using local quantum operations and classical communication (LOCC) between Alice, Bob and the helper systems, and possibly with the assistance of additional maximally entangled states.

A split transfer protocol consists of

1. A partitioning of the set of the helper systems: $T_A \sqcup T_B = \{C_1, \dots, C_m\}$.
2. For each $C_i \in T_A$, a number $K_{A,i}$ of shared maximally entangled qubits between Alice and C_i , and for each $C_i \in T_B$, a number $K_{B,i}$ of shared maximally entangled qubits between Bob and C_i .
3. An LOCC operation between Alice, Bob and the helper systems, such that after applying the protocol, Alice and Bob share a state $\psi_{ABRC_1\dots C_m}$, which is close to $\phi_{ABRC_1\dots C_m}$, and is such that Alice possesses systems A and T_A , while Bob controls B and T_B . Moreover, after applying the protocol they may be in possession of a number $L_{A,i}$ or $L_{B,i}$ of (approximately) maximally entangled qubits between C_i , and respectively A or B .

In this case, we say that the split transfer protocol has *entanglement costs* $K_{A,i} - L_{A,i}$ for all $C_i \in T_A$ and $K_{B,i} - L_{B,i}$ for all $C_i \in T_B$. A precise definition can be found as Definition 14 in [DH10].

Intuitively, the helper systems need to transfer their correlations with R to Alice and Bob, but without touching R . For instance, a simple protocol would be that the helpers simply teleport their full system to either Alice or Bob, consuming EPR pairs, leading to large entanglement costs. We can construct a potentially much more efficient protocol by way of random measurements, as detailed in Proposition 16 of [DH10]. Roughly speaking, such a protocol functions because random measurements have the effect of *decoupling* the helper systems from R . The helpers perform simultaneous random measurements on their systems, and send the results of their measurements to Alice and Bob. Then, Alice and Bob can use their share of the global state and their portions of the maximally-entangled states to perform a decoding operation conditioned on the results of the random measurements. The state they receive will be a purification of $\psi_{ABC_1\dots C_m}$, which will then be equivalent to the original $\psi_{ABRC_1\dots C_m}$ up to local isometries. The way we set up the split transfer protocol above was in a one-shot fashion: we get a single copy of ϕ and need to determine the optimal entanglement cost for the protocol.

One can also consider asymptotic variants of split transfer, where one has many copies available and aims to achieve an optimal transfer rate. An example application is the *entanglement of assistance*. Suppose that Alice, Bob and the helper systems C_i get many copies of a pure state ϕ . At what rate can they distill maximally entangled pairs between Alice and Bob, if Alice and Bob are allowed to perform LOCC operations with all the helper systems? In this case, the answer is that the rate is given by

$$\min_{T_A} S(AT_A)_\phi,$$

that is, by minimizing the entanglement entropy over all bipartitions. This rate is reminiscent of the importance of minimal cuts in a random tensor network, and the connection was explored in [HNQ⁺16].

To see how the task of split transfer relates to random tensor networks, we consider three boundary regions, A and B , under the control of Alice and Bob, and the purifying system R . Each of the bulk vertices on the network correspond to a ‘helper’ party. We would like to know whether there exists a protocol in which the assisting parties are allowed to perform local operations and classical communication (LOCC) such that the state ϕ_{VR} is redistributed into a state ρ_{ABR} held by Alice and Bob that can be transformed by local isometries, acting only on A and B , to a state close to ϕ_{VR} . The protocol given in [DH10] consists of simultaneous random measurements by each of the helpers. This precisely corresponds to the random projections performed in constructing the random tensor network state with this background state!

In this light, we can interpret [Theorem 4.4](#) as a result on split transfer. Let us assume that $\phi \in \mathcal{P}_=(ABRC_1 \dots C_n)$, denote the associated random tensor network state by ρ_{ABR} , and choose a partitioning $T_A \sqcup T_B$ of the assisting (bulk) parties. Since $H_2(A|B)_{\phi|\phi} \geq H_{\min}(A|B)_{\phi}$, [Theorem 4.4](#) directly yields

Theorem. *Suppose that*

$$H_{\min}(S_A|BRT_B)_{\phi} \geq K_1 \tag{B.1}$$

for all non-empty subsets $S_A \subseteq T_A$ and

$$H_{\min}(S_B|ART_A)_{\phi} \geq K_2 \tag{B.2}$$

for all non-empty subsets $S_B \subseteq T_B$. Then

$$\mathbb{E} \min_{V_A, V_B} \|(V_A \otimes V_B \otimes I_R) \rho(V_A^\dagger \otimes V_B^\dagger \otimes I_R) - \phi_{ABRC_1 \dots C_n}\|_1 = \mathcal{O}((2^{-\frac{1}{4}K_1} + 2^{-\frac{1}{4}K_2})). \tag{B.3}$$

where the minimum is over isometries $V_A : \mathcal{H}_A \rightarrow \mathcal{H}_{AT_A}$ and $V_B : \mathcal{H}_B \rightarrow \mathcal{H}_{BT_B}$.

This result shows that if K_1 and K_2 are sufficiently large, then after measurement in a random basis, the state possessed by Alice and Bob can, with high probability, be used to approximately reconstruct ϕ by acting with local isometries on the systems of Alice and Bob.

Suppose we fix values for K_1 and K_2 . If the conditions in [Eq. \(B.1\)](#) and [Eq. \(B.1\)](#) are not satisfied for the initial state ϕ , we can use another state where we have added an appropriate number of maximally entangled Bell pairs between the assisting parties, increasing the entanglement cost of the protocol.⁵ An interesting open question in this context is whether one can generalize this result using *smooth* entropies in [Eq. \(B.1\)](#) and [Eq. \(B.1\)](#). As alluded to in [Section 4](#), the problem is that for a general state ϕ , one would need to perform simultaneous smoothing for all the relevant subsystems, which remains an open problem.

There is an alternative approach, in which it is straightforwardly possible to use smooth entropies [DH10]. In this approach, one merges each party in T_A one by one, and similarly for T_B . That is, we choose some ordering $T_A = \{1, \dots, m\} = [m]$ and we apply a sequence of state merging protocols where we merge the state in m steps, where a single step merges $A \cup T_A \setminus [i-1]$ into $A \cup T_A \setminus [i]$. In this case it is not hard to see that, if we allow some error the entanglement cost is determined by the smooth conditional entropies $H_{\min}^\epsilon(\{i+1\}|BR[i]T_B)_{\phi}$. We perform a similar protocol for B and the assisting T_B systems.

B.1 Split transfer and recovery in holography

Split transfer is closely related to *subregion-subregion duality*, or *entanglement wedge reconstruction*, in holography. Consider an asymptotically AdS, stationary semiclassical geometry which is dual to a boundary CFT state ρ . We fix a time-reversal invariant spatial slice and partition the boundary into A and \bar{A} . Let γ_A be the minimal surface for A , and recall that $\partial\gamma_A = \partial A$. The region enclosed by A and γ_A is the *entanglement*

⁵In fact, the protocol in [DH10] is slightly more general than what we describe; rather than measuring a random state one could also measure a random projection of rank greater than 1. This can be used to obtain EPR pairs between the helpers and Alice and Bob to get nonzero $L_{A,i}$ and $L_{B,i}$.

wedge of A , which we denote by Γ_A . The claim of entanglement wedge reconstruction is that if we act with a low-energy local bulk operator in the entanglement wedge of A , we can reconstruct the action of this operator as a corresponding operator acting on the boundary system A .

One way to make this more precise is by considering a *code subspace* of bulk states S , which can be thought of as a set of states obtained by acting with low-energy operators on a fixed semiclassical space-time. The action of the operators is small, in the sense that they do not create a significant backreaction that changes the geometry. We then introduce a reference system R of the same dimension as S . In this framework, the AdS/CFT correspondence describes an encoding of the bulk into the boundary, taking a bulk state $\phi_{\Gamma_A \Gamma_A^c R}$ to a boundary state $\rho_{A \bar{A} R}$. In this set-up, the claim of entanglement wedge reconstruction is that we can act with an isometry on A to recover $\phi_{\Gamma_A R}$ (and Γ_A is actually the maximal such region). This situation corresponds to quantum state merging – there is a single decoder A . However, in AdS/CFT, one usually requires the stronger condition of *complementary entanglement wedge reconstruction*, in which the entanglement wedge for \bar{A} is also the complement of Γ_A , so that $\phi_{\Gamma_A^c R}$ is recoverable from \bar{A} . This stronger requirement with two decoders is closely related to split transfer. These ideas and the precise relation to quantum information theory remain an active area of research, e.g. [HNQ⁺16, Har17, AP20, AP22].

In particular, [BPSW19, AP20, AP22] have argued that one-shot quantum information is the correct framework to understand entanglement wedge reconstruction, as holography is fundamentally a one-shot setting – we are provided with a single copy of a gravitational or CFT state, rather than asymptotically-many copies.

The limit of large effective central charge, and hence small G_N , does reproduce certain aspects of the many-copy limit, as we reviewed in Appendix A.1. However, the distinction between the one-shot and asymptotic regimes becomes apparent in the presence of large bulk entropy, where the bulk entropy, and hence R , is large. Such situations arise when studying, for example, the black hole information paradox. In this case, where we assume we have a bulk state ϕ , the minimal surface prescription in Eq. (A.3) is replaced by *quantum extremal surface* prescription:

$$H(\rho_A) = \min \text{ext}_{\gamma_A} \left\{ \frac{|\gamma_A|}{4G_N} + H(\Gamma_A)_\phi \right\},$$

where we minimize over extremal surfaces γ_A , and we minimize the joint contribution of the area of γ_A and the bulk entropy contained in the associated entanglement wedge Γ_A . This formula has a natural tensor network interpretation: consider a random tensor network state with background state $\phi = \phi_{V^{(b)}R} \otimes \phi_{V^{(l)}}$, where $\phi_{V^{(b)}R}$ is a general background state (accounting for bulk entropy) and $\phi_{V^{(l)}}$ is a tensor product of link states on a graph $G = (V, E)$. Let us take maximally entangled link states with dimension D . Then, for some cut Γ_A with edge set γ_A , we have

$$H(\phi_{\Gamma_A}) = \log(D)|\gamma_A| + H(\phi_{\Gamma_A}^{(b)}),$$

and we may hope that minimization over this quantity along the cuts gives a good approximation to the entropy. Whether such a prescription is valid depends on the structure of the background state ϕ . A proposal put forth in [AP20] is that the surface γ_A with entanglement wedge Γ_A gives the *max-entanglement wedge*, if Γ_A is the largest region such that, for any other surface δ_A homologous to A , with Δ_A the region enclosed by A and δ_A , and where Δ_A is contained in Γ_A , it holds that

$$H_{\min}^\varepsilon(\Gamma_A \setminus \Delta_A | \Gamma_A^c R) \gg \frac{|\gamma_A| - |\delta_A|}{4G_N}.$$

In this case, Γ_A should be the largest region which can be (approximately) reconstructed from A . Again, one can think of a random tensor network where the background state is a tensor product of a bulk state and a maximally entangled link state of dimension D with $\log(D) = \Theta(G_N^{-1})$ along the discretization of the space. Then this condition is (apart from the simultaneous smoothing problem) equivalent to Eq. (4.13). Enforcing complementary reconstruction, in which Γ_A is the max-entanglement wedge for A , and its complement Γ_A^c is the max-entanglement wedge for \bar{A} , allows us to interpret the holographic encoding of the bulk state into the boundary as a version of one-shot split transfer. See [AP20] for a detailed discussion of this proposal for holographic systems.

C Joint smoothing of link states

In this section, we provide a proof of [Lemma 4.5](#). In our application of one-shot entropy estimates in [Section 4.3](#) and [Section 4.4](#), we needed to jointly smooth over different subsystems for the background state ϕ . For general background states, joint smoothing is an open problem [[DF13](#), [Dut11](#)]; however, if the background state is a product of link states as in [Eq. \(2.2\)](#), we can straightforwardly do so. In order to prove [Lemma 4.5](#), we first need some intermediate lemmas. These lemmas involve quantum states which can be expanded in some preferential basis with positive coefficients. Therefore, for [Lemmas C.1 to C.3](#) and [C.5](#), we assume that each quantum system has a preferential basis, which we assume without loss of generality to be the standard basis.

Lemma C.1. *Suppose that $\phi \in \mathcal{P}_\leq(XST)$ can be written as*

$$\phi_{XST} = \sum_i |i\rangle\langle i| \otimes \phi_{ST,i}$$

where each $\phi_{ST,i}$ is a pure state such that $|\phi_{ST,i}\rangle$ can be written with positive coefficients in the standard basis. If

$$\tilde{\phi}_{XST} = \sum_i |i\rangle\langle i| \otimes \tilde{\phi}_{ST,i}$$

is such that each $\tilde{\phi}_{ST,i}$ is a pure state satisfying $|\tilde{\phi}_{ST,i}\rangle \leq |\phi_{ST,i}\rangle$ elementwise, then

$$H_{\max}(XS|T)_{\tilde{\phi}} \leq H_{\max}(XS|T)_{\phi}.$$

Proof. We start by arguing that for the max-entropy

$$H_{\max}(XS|T)_{\tilde{\phi}} = \max_{\sigma_T \in \mathcal{P}_\leq(T)} \log F(\tilde{\phi}_{XST}, I_{XS} \otimes \sigma_T)^2 \quad (\text{C.1})$$

we may choose σ_T in the optimization problem with nonnegative matrix elements in the standard basis. Indeed, suppose that σ_T realizes the maximum in [Eq. \(C.1\)](#). Then we can write a spectral decomposition $\sigma_T = \sum_j p_j |e_j\rangle\langle e_j|$. Now let $|\tilde{e}_j\rangle = \sum_k |e_j|k\rangle |k\rangle$. Then $\tilde{\sigma}_T = \sum_j p_j |\tilde{e}_j\rangle\langle \tilde{e}_j| \in \mathcal{P}_\leq(T)$ has nonnegative matrix elements in the standard basis. Moreover,

$$\begin{aligned} F(\tilde{\phi}_{XST}, I_{XS} \otimes \sigma_T) &= \sum_i F(\tilde{\phi}_{ST,i}, I_S \otimes \sigma_T) \\ &\leq \sum_{i,j} p_j |\langle \tilde{\phi}_{ST,i} | I_S \otimes |\psi_j\rangle\langle \psi_j| | \tilde{\phi}_{ST,i}\rangle| \\ &\leq \sum_{i,j} p_j \langle \tilde{\phi}_{ST,i} | I_S \otimes |\tilde{\psi}_j\rangle\langle \tilde{\psi}_j| | \tilde{\phi}_{ST,i}\rangle \\ &= F(\tilde{\phi}_{XST}, I_{XS} \otimes \tilde{\sigma}_T). \end{aligned}$$

Let σ_T be optimal for [Eq. \(C.1\)](#) with nonnegative matrix elements. It is clear that if $|\tilde{\phi}_{ST,i}\rangle \leq |\phi_{ST,i}\rangle$, then

$$\begin{aligned} F(\tilde{\phi}_{XST}, I_{XS} \otimes \sigma_T) &= \sum_i \langle \tilde{\phi}_{ST,i} | I_S \otimes \sigma_T | \tilde{\phi}_{ST,i}\rangle \\ &\leq \sum_i \langle \phi_{ST,i} | I_S \otimes \sigma_T | \phi_{ST,i}\rangle \\ &= F(\phi_{XST}, I_{XS} \otimes \sigma_T) \end{aligned}$$

and hence

$$H_{\max}(XS|T)_{\tilde{\phi}} = \log F(\tilde{\phi}_{XST}, I_{XS} \otimes \sigma_T)^2 \leq \log F(\phi_{XST}, I_{XS} \otimes \sigma_T)^2 \leq H_{\max}(XS|T)_{\phi}.$$

□

Lemma C.2. Suppose that $\phi \in \mathcal{P}_{\leq}(XYST)$ can be written as

$$\phi_{XYST} = \sum_i |i\rangle\langle i| \otimes |j\rangle\langle j| \otimes \phi_{ST,ij}$$

where each $\phi_{ST,ij}$ is a pure state such that $|\phi_{ST,ij}\rangle$ can be written with positive coefficients in the standard basis. If

$$\tilde{\phi}_{XYST} = \sum_i |i\rangle\langle i| \otimes |j\rangle\langle j| \otimes \tilde{\phi}_{ST,ij}$$

is such that each $\tilde{\phi}_{ST,ij}$ is a pure state satisfying $|\tilde{\phi}_{ST,ij}\rangle \leq |\phi_{ST,ij}\rangle$ elementwise, then

$$H_{\max}(XS|YT)_{\tilde{\phi}} \leq H_{\max}(XS|YT)_{\phi}.$$

Proof. If we have a state $\rho \in \mathcal{P}_{\leq}(XYST)$ which is of the form

$$\sum_j |j\rangle\langle j| \otimes \rho_j$$

where each $\rho_j \in \mathcal{P}_{\leq}(XST)$, then (Proposition 4.6 in [Tom15])

$$H_{\max}(XS|YT) = \log \left(\sum_j 2^{H_{\max}(XS|T)_{\rho_j}} \right).$$

Together with Lemma C.1 this implies the result. \square

The background states we consider are tensor products of link states, and they can be expressed with positive coefficients in a product basis along the half-edges. The following lemma will help us to show that when we perform smoothing we can retain some of this structure.

Lemma C.3. Suppose that $\phi \in \mathcal{P}_{\leq}(XST)$ can be written as

$$\phi_{XST} = \sum_i |i\rangle\langle i| \otimes \phi_{ST,i}$$

where each $\phi_{ST,i}$ is pure and has a Schmidt decomposition

$$|\phi_{ST,i}\rangle = \sum_j \sqrt{\lambda_{i,j}} |jj\rangle$$

in the standard basis. Then

$$H_{\min}^{\varepsilon}(XS|T)_{\phi} = H_{\min}(XS|T)_{\tilde{\phi}}$$

for a state $\tilde{\phi} \in \mathcal{P}_{\leq}(XST)$ which is such that $P(\phi, \tilde{\phi}) \leq \varepsilon$ and

$$\tilde{\phi}_{XST} = \sum_i |i\rangle\langle i| \otimes \tilde{\phi}_{ST,i}$$

with

$$|\tilde{\phi}_{ST,i}\rangle = \sum_j \sqrt{\tilde{\lambda}_{i,j}} |jj\rangle.$$

for some $0 \leq \tilde{\lambda}_{i,j} \leq 1$.

Proof. Let Y be a copy of X and let R be an additional reference system of sufficiently large dimension. Then we let

$$|\phi_{XYSTR}\rangle = \sum_{i,j} \sqrt{\lambda_{i,j}} |ii\rangle \otimes |jj\rangle \otimes |0\rangle$$

be a purification of ϕ_{XST} . Now, by duality of smooth entropies

$$H_{\min}^{\varepsilon}(XS|T)_{\phi} = -H_{\max}^{\varepsilon}(XS|Y)_{\phi}$$

By Lemma 6.13 of [Tom15] we can find a state

$$\bar{\phi}_{SXY} = \sum_j |j\rangle\langle j| \otimes \bar{\phi}_{XY}$$

such that $P(\phi, \bar{\phi}) \leq \varepsilon$ and $H_{\max}^{\varepsilon}(XS|Y)_{\phi} = H_{\max}(XS|Y)_{\bar{\phi}}$. Let

$$|\bar{\phi}_{STXYR}\rangle = \sum_j |jj\rangle \otimes |\bar{\phi}_{XYR,j}\rangle$$

be a purification such that $F(\phi_{XYS}, \bar{\phi}_{XYS}) = |\langle \phi | \bar{\phi} \rangle|$. We define the projector $\Pi_{XY} = \sum_j |jj\rangle\langle jj|$. Then for any $\sigma_Y \in \mathcal{P}_{\leq}(Y)$ we note that $(I_S \otimes \Pi_{XY})I_{XS} \otimes \sigma_Y (I_S \otimes \Pi_{XY}) \leq I_{XS} \otimes \sigma'_Y$ where $\sigma'_Y = \sum_j \langle j | \sigma_Y | j \rangle |j\rangle\langle j|$ and hence

$$F((I_S \otimes \Pi_{XY})\bar{\phi}(I_S \otimes \Pi_{XY}), I_{XS} \otimes \sigma_Y) = F(\bar{\phi}, (I_S \otimes \Pi_{XY})I_{XS} \otimes \sigma_Y (I_S \otimes \Pi_{XY})) \leq F(\bar{\phi}, I_{XS} \otimes \sigma'_Y).$$

If we let $|\phi'_{XYSTR}\rangle = (\Pi_{XY} \otimes I_{STR})|\bar{\phi}_{XYSTR}\rangle$ this implies that

$$H_{\max}(XS|Y)_{\phi'} \leq H_{\max}(XS|Y)_{\bar{\phi}}.$$

Moreover it is easy to see that $|\langle \phi' | \phi \rangle| = |\langle \bar{\phi} | \phi \rangle|$. Again using duality we find that

$$H_{\min}^{\varepsilon}(XS|T)_{\phi} \leq H_{\min}(XS|TR)_{\phi'}.$$

Finally, let

$$|\tilde{\phi}\rangle = (I_{XYST} \otimes \langle 0|) |\phi'\rangle$$

then by data processing (Theorem 6.19 of [Tom15]) it holds that

$$H_{\min}(XS|TR)_{\phi'} \leq H_{\min}(XS|T)_{\tilde{\phi}}.$$

By construction $|\langle \tilde{\phi} | \phi \rangle| = |\langle \phi' | \phi \rangle|$ and hence $P(\tilde{\phi}, \phi) \leq \varepsilon$. Finally, by construction $\tilde{\phi}$ is of the desired form

$$|\tilde{\phi}_{STXY}\rangle = \sum_{i,j} \sqrt{\tilde{\lambda}_{i,j}} |ii\rangle \otimes |jj\rangle.$$

□

The following lemmas will be used in our joint smoothing construction to bound the purified distance of the smoothed state.

Lemma C.4. *Suppose $\phi, \psi \in \mathcal{P}_{\leq}(\mathcal{H})$ are pure states, and suppose*

$$|\phi\rangle = \sum_i \sqrt{\lambda_i} |i\rangle \quad |\psi\rangle = \sum_i \sqrt{\mu_i} |i\rangle$$

for some $\lambda, \mu_i \geq 0$. Then

$$T(\phi, \psi) \leq P(\phi, \psi) \leq \sqrt{2 \sum_i |\lambda_i - \mu_i|}.$$

Proof. Let

$$\rho = \Delta(\phi), \quad \sigma = \Delta(\psi)$$

where Δ is the completely dephasing channel, so

$$\rho = \sum_i \lambda_i |i\rangle\langle i| \quad \sigma = \sum_i \mu_i |i\rangle\langle i|.$$

Then since ρ and σ are diagonal in the same basis

$$P(\phi, \psi) = P(\rho, \sigma).$$

We then estimate the trace distance by

$$T(\rho, \sigma) \leq \|\rho - \sigma\|_1 = \sum_i |\lambda_i - \mu_i|$$

and we apply the Fuchs-Van de Graaff inequalities, Eq. (4.1), to estimate

$$T(\phi, \psi) \leq P(\phi, \psi) = P(\rho, \sigma) \leq \sqrt{2T(\rho, \sigma)} \leq \sqrt{2 \sum_i |\lambda_i - \mu_i|}.$$

□

Lemma C.5. *Suppose $\phi, \phi_j \in \mathcal{P}_{\leq}(\mathcal{H})$ are pure states for $j \in [n]$, and suppose*

$$|\phi\rangle = \sum_i \sqrt{\lambda_i} |i\rangle \quad |\phi_j\rangle = \sum_i \sqrt{\lambda_{j,i}} |i\rangle$$

for some $\lambda, \lambda_{i,j} \geq 0$ and suppose $P(\phi, \phi_j) \leq \varepsilon$ for all j . Then if we let

$$|\tilde{\phi}\rangle = \sum_i \min_j \sqrt{\lambda_{j,i}} |i\rangle$$

it holds that $P(\phi, \tilde{\phi}) \leq 2\sqrt{n\varepsilon}$.

Proof. Let

$$\rho = \sum_i \lambda_i |i\rangle\langle i| \quad \rho_j = \sum_i \lambda_{j,i} |i\rangle\langle i|.$$

Then, $P(\rho, \rho_j) = P(\phi, \phi_j)$ and using the Fuchs-Van de Graaff inequalities

$$\sum_i |\lambda_i - \min_j \lambda_{j,i}| \leq \sum_j \sum_i |\lambda_i - \lambda_{j,i}| = \sum_j \|\rho - \rho_j\|_1 \leq 2 \sum_j T(\rho, \rho_j) \leq 2 \sum_j P(\phi, \phi_j) \leq 2n\varepsilon.$$

The result now follows from Lemma C.4. □

Finally, we prove our main joint smoothing result. We again consider the setting of a tensor product of link states on a graph $G = (V, E)$, where we consider some boundary subsystem $A \subseteq V_{\partial}$. Recall that for a cut $\Gamma_A \in C(A)$ we define

$$\begin{aligned} \mathcal{C}_1(\Gamma_A) &= \{\Delta_A \in C(A) : \Delta_A \subsetneq \Gamma_A\} \\ \mathcal{C}_2(\Gamma_A) &= \{\Delta_A \in C(A) : \Gamma_A \subsetneq \Delta_A\}. \end{aligned}$$

Lemma 4.5. *Let $\phi \in \mathcal{P}_=(V)$ be a link state, $A \subseteq V_\partial$ a boundary subsystem and $\Gamma_A \in C(A)$ a cut for A . Then there exists a pure state $\phi^\varepsilon \in \mathcal{P}_\leq(V)$ which is such that*

$$P(\phi, \phi^\varepsilon) \leq 2 \left(\sqrt{|\mathcal{C}_1(\Gamma_A)|} + \sqrt{|\mathcal{C}_2(\Gamma_A)|} \right) \sqrt{\varepsilon}$$

and it holds that for any $\Delta_A \in \mathcal{C}_1(\Gamma_A)$

$$H_{\min}(\Gamma_A \setminus \Delta_A | \Gamma_A^c)_{\phi^\varepsilon} \geq H_{\min}^\varepsilon(\Gamma_A \setminus \Delta_A | \Gamma_A^c)_\phi$$

and for any $\Delta_A \in \mathcal{C}_2(\Gamma_A)$

$$H_{\min}(\Delta_A \setminus \Gamma_A | \Gamma_A)_{\phi^\varepsilon} \geq H_{\min}^\varepsilon(\Delta_A \setminus \Gamma_A | \Gamma_A)_\phi.$$

Proof. We may assume without loss of generality that $\phi = \bigotimes_{e \in E} \phi_e$ is such that each ϕ_e has Schmidt decomposition in the standard basis,

$$|\phi_e\rangle = \sum_{i=1}^{D_e} \sqrt{\lambda_{e,i}} |ii\rangle$$

This means we may write

$$|\phi\rangle = \sum_I \sqrt{\lambda_I} |I\rangle$$

where I runs over all possible basis elements along each edge $I = \{i_e\}_{e \in E}$ and

$$\lambda_I = \prod_{e \in E} \lambda_{e,i_e} \quad |I\rangle = \bigotimes_{e \in E} |i_e i_e\rangle.$$

We also let

$$\begin{aligned} E_1 &= \{e = (xy) \in E, x, y \in \Gamma_A\} \\ E_2 &= \{e = (xy) \in E, x, y \in \Gamma_A^c\}. \end{aligned}$$

so $E = E_1 \sqcup E_2 \sqcup \gamma_A$.

Consider a cut $\Delta_A \in \mathcal{C}_1(\Gamma_A)$. Let

$$\begin{aligned} X &= \{(e, x) : e = (xy), x \in \Gamma_A \setminus \Delta_A, y \in \Delta_A\} \\ S &= \{(e, x) : e = (xy), x \in \Gamma_A \setminus \Delta_A, y \in \Delta_A^c\} \\ T &= \{(e, x) : e = (xy), x \in \Gamma_A^c, y \in \Delta_A^c\} \end{aligned}$$

Then we see that $H_{\min}^\varepsilon(\Gamma_A \setminus \Delta_A | \Gamma_A^c)_\phi = H_{\min}^\varepsilon(XS|T)_\phi$ and we can write (as ϕ is a product state)

$$\phi_{XST} = \phi_X \otimes \phi_{ST}.$$

Here ϕ_{ST} is pure, whereas ϕ_X is diagonal in the standard basis along the half-edges in X . From [Lemma C.3](#) (applied with $\phi_{ST,i} = \phi_{ST}$ for each i) we find that we obtain a state $\phi^{\Delta_A, \varepsilon}$ which is such that $P(\phi^{\Delta_A, \varepsilon}, \phi) \leq \varepsilon$ and

$$H_{\min}(\Gamma_A \setminus \Delta_A | \Gamma_A^c)_{\phi^{\Delta_A, \varepsilon}} \geq H_{\min}^\varepsilon(\Gamma_A \setminus \Delta_A | \Gamma_A^c)_\phi$$

and which can be written as

$$|\phi^{\Delta_A, \varepsilon}\rangle = \sum_I \sqrt{\lambda_I^{\Delta_A, \varepsilon}} |I\rangle.$$

for some coefficients $\lambda_I^{\Delta A, \varepsilon}$. Moreover, we can write

$$|\phi^{\Delta A, \varepsilon}\rangle = \sum_i |ii\rangle \otimes |\phi_{E_1, i}^{\Delta A, \varepsilon}\rangle \otimes |\phi_{E_2}\rangle$$

where i runs over all possible basis elements along the cut γ_A , $i = \{i_e\}_{e \in \gamma_A}$ and $|i\rangle = \bigotimes_{e \in \gamma_A} |i_e\rangle$, and the $\phi_{E_1, i}^{\Delta A, \varepsilon}$ are pure states with positive coefficients in the standard basis. We now let

$$\begin{aligned} |\phi_1^\varepsilon\rangle &= \sum_I \sqrt{\lambda_I^\varepsilon} |I\rangle \\ \lambda_I^\varepsilon &= \min_{\Delta_A \in \mathcal{C}_1(\Gamma_A)} \lambda_I^{\Delta A, \varepsilon}. \end{aligned}$$

By construction, this state can be written as

$$|\phi_1^\varepsilon\rangle = \sum_i |ii\rangle \otimes |\phi_{E_1, i}^\varepsilon\rangle \otimes |\phi_{E_2}\rangle$$

By [Lemma C.5](#), it holds that

$$P(\phi, \phi_1^\varepsilon) \leq 2\sqrt{|\mathcal{C}_1(\Gamma_A)|\varepsilon}.$$

An analogous construction can be used to construct $\phi^{\Delta A, \varepsilon}$ for $\Delta_A \in \mathcal{C}_2(\Gamma_A)$, and taking the minimum over all such cuts, we get a state ϕ_2^ε of the form

$$|\phi_2^\varepsilon\rangle = \sum_i |ii\rangle \otimes |\phi_{E_1}\rangle \otimes |\phi_{E_2, i}^\varepsilon\rangle$$

which satisfies

$$P(\phi, \phi_2^\varepsilon) \leq 2\sqrt{|\mathcal{C}_2(\Gamma_A)|\varepsilon}.$$

We now define

$$|\phi^\varepsilon\rangle = \sum_i |ii\rangle \otimes |\phi_{E_1, i}^\varepsilon\rangle \otimes |\phi_{E_2, i}^\varepsilon\rangle.$$

We will now show that ϕ^ε has the desired properties. First of all, since ϕ is normalized, $F_*(\phi, \phi^\varepsilon) = F(\phi, \phi^\varepsilon)$ and $F(\phi, \phi^\varepsilon) = F(\phi_1^\varepsilon, \phi_2^\varepsilon)$ so

$$P(\phi, \phi^\varepsilon) = P(\phi_1^\varepsilon, \phi_2^\varepsilon) \leq P(\phi, \phi_1^\varepsilon) + P(\phi, \phi_2^\varepsilon) \leq 2 \left(\sqrt{|\mathcal{C}_1(\Gamma_A)|} + \sqrt{|\mathcal{C}_2(\Gamma_A)|} \right) \sqrt{\varepsilon}$$

using the fact that the purified distance is a metric.

Next, consider $\Delta_A \in \mathcal{C}_1(A)$. We note that by duality $H_{\min}(\Gamma_A \setminus \Delta_A | \Gamma_A^c)_{\phi^\varepsilon} = -H_{\max}(\Gamma_A \setminus \Delta_A | \Delta_A)_{\phi^\varepsilon}$. We define the following subsystems of half-edges,

$$\begin{aligned} X' &= \{(e, x) : e = (xy), x \in \Gamma_A \setminus \Delta_A, y \in \Gamma_A^c\} \\ Y' &= \{(e, x) : e = (xy), x \in \Delta_A, y \in \Gamma_A^c\} \\ S' &= \{(e, x) : e = (xy), x \in \Gamma_A \setminus \Delta_A, y \in \Gamma_A\} \\ T' &= \{(e, x) : e = (xy), x \in \Delta_A, y \in \Gamma_A\}. \end{aligned}$$

so $H_{\max}(\Gamma_A \setminus \Delta_A | \Delta_A)_{\phi^\varepsilon} = H_{\max}(X'S' | Y'T')_{\phi^\varepsilon}$. Next, we observe that by construction, ϕ^ε is of the form

$$\phi_{X'Y'S'T'}^\varepsilon = \sum_{i,j} |i\rangle\langle i| \otimes |j\rangle\langle j| \otimes \phi_{S'T', ij}^\varepsilon$$

where i runs over all possible basis elements along $\gamma_A \setminus \delta_A$, $i = \{i_e\}_{e \in \gamma_A \setminus \delta_A}$ and $|i\rangle = \bigotimes_{e \in \gamma_A \setminus \delta_A} |i_e\rangle$, which forms a basis for \mathcal{H}_X ; and j runs over all possible basis elements along $\gamma_A \cap \delta_A$, $j = \{j_e\}_{e \in \gamma_A \cap \delta_A}$ and $|j\rangle = \bigotimes_{e \in \gamma_A \cap \delta_A} |j_e\rangle$, which forms a basis for \mathcal{H}_Y . Each $\phi_{S'T',ij}^\varepsilon$ is a pure state. Similarly,

$$\phi_{X'Y'S'T'}^{\Delta_A, \varepsilon} = \sum_{i,j} |i\rangle\langle i| \otimes |j\rangle\langle j| \otimes \phi_{S'T',ij}^{\Delta_A, \varepsilon}$$

where the $\phi_{S'T',ij}^{\Delta_A, \varepsilon}$ are pure states. Moreover, by construction $|\phi_{S'T',ij}^\varepsilon\rangle \leq |\phi_{S'T',ij}^{\Delta_A, \varepsilon}\rangle$ elementwise in the standard basis. Therefore, by [Lemma C.2](#) it holds that

$$H_{\max}(X'S'|Y'T')_{\phi^\varepsilon} \leq H_{\max}(X'S'|Y'T')_{\phi^{\Delta_A, \varepsilon}}.$$

Finally, by duality $H_{\min}(\Gamma_A \setminus \Delta_A | \Gamma_A^c)_{\phi^{\Delta_A, \varepsilon}} = -H_{\max}(X'S'|Y'T')_{\phi^{\Delta_A, \varepsilon}}$ and we conclude

$$H_{\min}(\Gamma_A \setminus \Delta_A | \Gamma_A^c)_{\phi^\varepsilon} \geq H_{\min}(\Gamma_A \setminus \Delta_A | \Gamma_A^c)_{\phi^{\Delta_A, \varepsilon}} \geq H_{\min}^\varepsilon(\Gamma_A \setminus \Delta_A | \Gamma_A^c)_\phi.$$

A similar argument is valid for $\Delta_A \in \mathcal{C}_2(\Gamma_A)$, showing that

$$H_{\min}(\Delta_A \setminus \Gamma_A | \Gamma_A)_{\phi^\varepsilon} \geq H_{\min}(\Delta_A \setminus \Gamma_A | \Gamma_A)_{\phi^{\Delta_A, \varepsilon}} \geq H_{\min}^\varepsilon(\Delta_A \setminus \Gamma_A | \Gamma_A)_\phi.$$

□

In the proof of [Theorem 4.12](#) we need a similar smoothing lemma for a slightly different situation as stated in [Lemma 4.11](#).

Recall the set-up: we consider a graph $G = (V, E)$ and we have a set of boundary vertices V_∂ , and $A \subseteq V_\partial$. We denote by $\gamma_{A,1}$ the edges incident to A , and $\gamma_{A,2}$ the edges incident to \bar{A} , and we assume these sets do not intersect. We let $E_b = E \setminus (\gamma_{A,1} \cup \gamma_{A,2})$. For a cut $\Delta_A \in C(A)$ we let Y^{Δ_A} be the set of half-edges

$$Y^{\Delta_A} = \{(e, x) : e = (xy), x \in \Delta_A^c, y \in A\}.$$

Lemma 4.11. *Let $\phi \in \mathcal{P}_{\leq}(V)$ be a pure background state which is of the form $\phi = \phi_{\gamma_{A,1}} \otimes \phi_{\gamma_{A,2}} \otimes \phi_{E_b}$, where*

$$\phi_{E_b} = \bigotimes_{e \in E_b} \phi_e$$

is a product state and the $\phi_{\gamma_{A,i}}$ have a Schmidt decomposition in the standard basis along the half-edges. Then there exists a state ϕ^ε which is such that

$$P(\phi, \phi^\varepsilon) \leq 2\sqrt{2^{V_b}\varepsilon}$$

and for all cuts $\Delta_A \in C(A)$ it holds that

$$H_{\min}(\Delta_A \setminus A | A)_{\phi^\varepsilon} \geq H_{\min}^\varepsilon(\Delta_A \setminus A | AY^{\Delta_A})_\phi.$$

Proof. The argument is much the same as the proof of [Lemma 4.5](#). We again assume without loss of generality that each ϕ_e has Schmidt decomposition in the standard basis, and we may write

$$|\phi\rangle = \sum_I \sqrt{\lambda_I} |I\rangle$$

as in the proof of [Lemma 4.5](#). Consider a cut $\Delta_A \in C(A)$. Then we let

$$\begin{aligned} X &= \{(e, x) : e = (xy), x \in \Delta_A, y \in \Delta_A^c\} \\ S &= \{(e, x) : e = (xy), x \in \Delta_A, y \in \Delta_A \setminus A\} \\ T &= \{(e, x) : e = (xy), x \in A, y \notin A\} \cup Y^{\Delta_A} \end{aligned}$$

so $H_{\min}^{\varepsilon}(\Delta_A \setminus A|AY^{\Delta_A})_{\phi} = H_{\min}^{\varepsilon}(XS|T)_{\phi}$. Moreover, by the structure of the state, $\phi_{XST} = \phi_X \otimes \phi_{ST}$, where ϕ_{ST} is pure and ϕ_X is diagonal in the standard basis. So, from [Lemma C.3](#) we obtain a state

$$|\phi^{\Delta_A, \varepsilon}\rangle = \sum_I \sqrt{\lambda_I^{\Delta_A, \varepsilon}} |I\rangle$$

for some real coefficients $\lambda_I^{\Delta_A, \varepsilon} \geq 0$ which is such that $P(\phi, \phi^{\Delta_A, \varepsilon}) \leq \varepsilon$ and $H_{\min}(XS|T)_{\phi^{\varepsilon}} \geq H_{\min}^{\varepsilon}(XS|T)_{\phi}$. As before, we define

$$|\phi^{\varepsilon}\rangle = \sum_I \sqrt{\lambda_I^{\varepsilon}} |I\rangle$$

$$\lambda_I^{\varepsilon} = \min_{\Delta_A \in C(A)} \lambda_I^{\Delta_A, \varepsilon}.$$

By [Lemma C.5](#) $P(\phi, \phi^{\varepsilon}) \leq 2\sqrt{|C(A)|\varepsilon} = 2\sqrt{2^{|V_b|}\varepsilon}$. If we define the following subsystems

$$X' = \{(e, x) : e = (xy), x \in \Delta_A, yA\}$$

$$Y' = \{(e, x) : e = (xy), x \in \Delta_A^c, yA\}$$

$$S' = \{(e, x) : e = (xy), x \in \Delta_A, y \notin A\}$$

$$T' = \{(e, x) : e = (xy), x \in \Delta_A^c, y \notin A\}$$

then $\phi_{X'Y'S'T'}^{\varepsilon}$ can be written as

$$\phi_{X'Y'S'T'}^{\varepsilon} = \sum_{i,j} |i\rangle\langle i| \otimes |j\rangle\langle j| \otimes \phi_{S'T', ij}^{\varepsilon}$$

where i runs over all possible basis elements along $\gamma_{A,1} \setminus \delta_A$, $i = \{i_e\}_{e \in \gamma_{A,1} \setminus \delta_A}$ and $|i\rangle = \bigotimes_{e \in \gamma_{A,1} \setminus \delta_A} |i_e\rangle$, which forms a basis for $\mathcal{H}_{X'}$; and j runs over all possible basis elements along $\gamma_{A,1} \cap \delta_A$, $j = \{j_e\}_{e \in \gamma_{A,1} \cap \delta_A}$ and $|j\rangle = \bigotimes_{e \in \gamma_{A,1} \cap \delta_A} |j_e\rangle$, which forms a basis for $\mathcal{H}_{Y'}$. The $\phi_{S'T', ij}^{\varepsilon}$ are pure states. We can therefore apply [Lemma C.1](#), and

$$H_{\max}(X'S'|T'Y')_{\phi^{\varepsilon}} \leq H_{\max}(X'S'|T'Y')_{\phi^{\Delta_A, \varepsilon}}$$

and hence, by applying duality and data processing

$$\begin{aligned} H_{\min}(\Delta_A \setminus A|A)_{\phi^{\varepsilon}} &= -H_{\max}(\Delta_A \setminus A|T'Y')_{\phi^{\varepsilon}} = -H_{\max}(X'S'|T'Y')_{\phi^{\varepsilon}} \\ &\geq H_{\max}(X'S'|T'Y')_{\phi^{\Delta_A, \varepsilon}} = H_{\min}(\Delta_A \setminus A|A)_{\phi^{\Delta_A, \varepsilon}} \\ &\geq H_{\min}(\Delta_A \setminus A|AY^{\Delta_A})_{\phi^{\Delta_A, \varepsilon}} \geq H_{\min}^{\varepsilon}(\Delta_A \setminus A|AY^{\Delta_A})_{\phi}. \end{aligned}$$

□

Remark C.6. We may make two observations on [Lemma 4.5](#) and [Lemma 4.11](#). In both cases, if we construct the coefficients as $\min \lambda_I^{\varepsilon}, \lambda_I$ we find that we may assume the resulting state has coefficients in the basis $|I\rangle$ which are upper bounded by λ_I (affecting only the constant factor in the upper bound for $P(\phi, \phi^{\varepsilon})$). Secondly, we note that

$$|\text{tr}[\phi] - \text{tr}[\phi^{\varepsilon}]| \leq \sum_I |\lambda_I - \lambda_I^{\varepsilon}| = \mathcal{O}(\varepsilon) \quad (\text{C.2})$$

rather than the naive $\mathcal{O}(\sqrt{\varepsilon})$ estimate.

D A family of metrics on the symmetric group

In analogy to the Cayley distance on the symmetric group S_k , we consider the following function $d_\rho: S_k \times S_k \rightarrow \mathbb{R}_{\geq 0}$ given some arbitrary density matrix $\rho \in \mathcal{P}_=(\mathcal{H})$:

$$d_\rho(\pi_1, \pi_2) = \sum_{l \in C(\pi_1^{-1}\pi_2)} (l-1)H_l(\rho) = - \sum_{l \in C(\pi_1^{-1}\pi_2)} \log \text{tr}[\rho^l].$$

As we saw in [Section 2.1](#), this function is closely related to the replica trick for random tensor networks with nontrivial link states. In the case where ρ is a maximally mixed state, this function is precisely proportional to the Cayley distance ([2.15](#)). In this appendix we show that d_ρ is a metric for *any* non-pure quantum state ρ . (If it is pure, then d_ρ vanished identically.)

We say that two permutations $\alpha, \beta \in S_k$ are *disjoint* if any point not fixed by α is fixed by β and vice versa.

Lemma D.1. *The function d_ρ defines a metric on S_k for any state ρ which is not pure. Moreover, if ρ does not have flat spectrum the following holds for all $\pi_1, \pi_2, \pi_3 \in S_k$:*

$$d_\rho(\pi_1, \pi_2) + d_\rho(\pi_2, \pi_3) = d_\rho(\pi_1, \pi_3)$$

if and only if $\pi_1^{-1}\pi_2$ and $\pi_2^{-1}\pi_3$ are disjoint permutations.

Proof. The fact that $d_\rho(\pi_1, \pi_2) = 0$ if and only if $\pi_1 = \pi_2$ follows from the assumption that ρ is not pure. The symmetry $d_\rho(\pi_1, \pi_2) = d_\rho(\pi_2, \pi_1)$ is clear from $C(\pi_1^{-1}\pi_2) = C(\pi_2^{-1}\pi_1)$. Thus, the only nontrivial property we have to show in order for d_ρ to be a metric is the triangle inequality. We let $d = \dim(\mathcal{H})$ and we write $\text{spec}(\rho) = \{\lambda_i\}_{i=1}^d$. Then, by letting $\alpha = \pi_1^{-1}\pi_2$ and $\beta = \pi_2^{-1}\pi_3$ the triangle inequality

$$d_\rho(\pi_1, \pi_2) + d_\rho(\pi_2, \pi_3) \geq d_\rho(\pi_1, \pi_3)$$

is equivalent to

$$\prod_{l \in C(\alpha)} \left(\sum_{i=1}^d \lambda_i^l \right) \prod_{m \in C(\beta)} \left(\sum_{i=1}^d \lambda_i^m \right) \leq \prod_{n \in C(\alpha\beta)} \left(\sum_{i=1}^d \lambda_i^n \right). \quad (\text{D.1})$$

Moreover, we need to show that if the spectrum is not flat, we have equality if and only if α and β are disjoint permutations. It suffices to show this for the case where β is a cycle, as we can write β as a product of disjoint cycles in general and iteratively apply the result for the case where β is a cycle. We write $\beta = (i_1 \dots i_m)$ and $\beta_q = (i_1 \dots i_q)$ for $q \leq m$ and we let $\alpha_q = \alpha\beta_q$ and $\alpha_0 = \alpha$.

We can then find a unique sequence of numbers $q_0 = 0 < q_1 < q_2 < \dots < m$ such that, if one compares α_{q_j} and $\alpha_{q_{j+1}}$, then either $q_{j+1} - q_j + 1$ cycles have merged into a single cycle, or a single cycle has split into $q_{j+1} - q_j + 1$ cycles. Moreover, these two operations are alternating in the sense that if α_{q_j} to $\alpha_{q_{j+1}}$ is a merge then $\alpha_{q_{j+1}}$ to $\alpha_{q_{j+2}}$ is a split and vice versa. Indeed, $\beta_{q+1} = \beta_q(i_q i_{q+1})$, so $\alpha_{q+1} = \alpha_q(i_q i_{q+1})$. If i_q and i_{q+1} are in different cycles in α_q , then applying $(i_q i_{q+1})$ merges these two cycles, and if i_q and i_{q+1} are in the same cycle in α_q , then applying $(i_q i_{q+1})$ splits this cycle into two cycles. Let

$$J = \{j : \alpha_{q_j} \text{ to } \alpha_{q_{j+1}} \text{ is a merge}\}$$

and for $j \in J$ let $m_j = q_{j+1} - q_j + 1$, then $\sum_{j \in J} m_j \leq m$. Now, it is clear that for any collection n_1, \dots, n_r of numbers with $\sum_{j=1}^r n_j \leq n$ it holds that

$$\prod_{j=1}^r \left(\sum_{i=1}^d \lambda_i^{n_j} \right) \geq \sum_{i=1}^d \lambda_i^n \quad (\text{D.2})$$

where the inequality is strict unless $r = 1$ and $n_1 = n$ (since we assume that ρ is not pure). In particular (recall that we assumed β to be a cycle of length m) we may estimate the left hand side of Eq. (D.1) by

$$\prod_{l \in C(\alpha)} \left(\sum_{i=1}^d \lambda_i^l \right) \left(\sum_{i=1}^d \lambda_i^m \right) \leq \prod_{l \in C(\alpha)} \left(\sum_{i=1}^d \lambda_i^l \right) \prod_{j \in J} \left(\sum_{i=1}^d \lambda_i^{m_j} \right). \quad (\text{D.3})$$

with equality if and only if $\{m_j\}_{j \in J} = \{m\}$ (so β only merges cycles). We will next argue that for $j \notin J$

$$\prod_{l \in C(\alpha_{q_j})} \left(\sum_{i=1}^d \lambda_i^l \right) \leq \prod_{l \in C(\alpha_{q_{j+1}})} \left(\sum_{i=1}^d \lambda_i^l \right). \quad (\text{D.4})$$

and for $j \in J$

$$\prod_{l \in C(\alpha_{q_j})} \left(\sum_{i=1}^d \lambda_i^l \right) \left(\sum_{i=1}^d \lambda_i^{m_j} \right) \leq \prod_{l \in C(\alpha_{q_{j+1}})} \left(\sum_{i=1}^d \lambda_i^l \right) \quad (\text{D.5})$$

with equality if and only if the spectrum is flat or all the cycles that are merged are 1-cycles. Then, combining Eq. (D.3), Eq. (D.4) and Eq. (D.5) we may conclude that Eq. (D.1) holds, with equality if and only if β is a disjoint cycle from α .

Eq. (D.4) follows immediately from Eq. (D.2), so it remains to show Eq. (D.5). To this end we will apply Jensen's inequality. In Eq. (D.5) let us assume that s cycles of lengths l_1, \dots, l_s are merged into a cycle of length l , so $l = \sum_{p=1}^s l_p$. We let $l_{s+1} = s$. Let $f_p : [d] \rightarrow \mathbb{R}_{\geq 0}$ be the function defined by $i \mapsto \lambda_i^{l_p - 1}$, for $p \in [s+1]$. Moreover, let

$$\eta_p = \frac{l_p - 1}{l - 1} \quad \text{for } p \in [s].$$

We consider expectation values over the probability measure on $[d]$ where i has probability λ_i . Since $\eta_p \leq 1$ for all p , by Jensen's inequality for $\eta_p \neq 0$

$$\left(\mathbb{E} f_p^{\frac{1}{\eta_p}} \right)^{\eta_p} \geq \mathbb{E} f_p \quad (\text{D.6})$$

with equality if and only if the spectrum is flat or $\eta_p = 1$. Note that for $\eta_p = 0$ (equivalently $l_p = 1$), $\mathbb{E} f_p = 1$. Using that

$$\sum_{p=1}^{s+1} \eta_p = \sum_{p=1}^s \frac{l_p - 1}{l - 1} + \frac{s - 1}{l - 1} = \frac{l - s + s - 1}{l - 1} = 1$$

and for $\eta_p \neq 0$

$$\mathbb{E} f_p^{\frac{1}{\eta_p}} = \sum_{i=1}^d \lambda_i^l$$

we find that

$$\prod_{p=1}^s \left(\sum_{i=1}^d \lambda_i^{l_p} \right) \sum_{i=1}^d \lambda_i^s = \prod_{p=1}^{s+1} \mathbb{E} f_p \leq \prod_{p, \eta_p \neq 0} \left(\mathbb{E} f_p^{\frac{1}{\eta_p}} \right)^{\eta_p} = \sum_{i=1}^d \lambda_i^l.$$

Here we have equality if and only if the spectrum is flat or for all $p \in [s+1]$ it holds that $\eta_p = 1$ or $\eta_p = 0$. Since $\eta_p < 1$ for $p \in [s]$ and $l_{s+1} > 1$ this only happens if $l_p = 1$ for $p \in [s]$ and $s = l_{s+1} = l$ (in other words, if all the merged cycles are 1-cycles). Applying this with $s = m_j$ for each $j \in J$ proves Eq. (D.5). \square

References

- [AEMM19] Ahmed Almheiri, Netta Engelhardt, Donald Marolf, and Henry Maxfield. The entropy of bulk quantum fields and the entanglement wedge of an evaporating black hole. *Journal of High Energy Physics*, 2019(12):1–47, 2019.
- [AFLR21] Chris Akers, Thomas Faulkner, Simon Lin, and Pratik Rath. Reflected entropy in random tensor networks. *arXiv preprint arXiv:2112.09122*, 2021.
- [AGZ10] Greg W Anderson, Alice Guionnet, and Ofer Zeitouni. *An Introduction to Random Matrices*. Cambridge University Press, 2010.
- [AHM⁺20] Ahmed Almheiri, Thomas Hartman, Juan Maldacena, Edgar Shaghoulian, and Amirhossein Tajdini. Replica wormholes and the entropy of Hawking radiation. *Journal of High Energy Physics*, 2020(5):1–42, 2020.
- [AN12] Guillaume Aubrun and Ion Nechita. Realignment random states. *Journal of Mathematical Physics*, 53(10):102210, 2012.
- [AP20] Chris Akers and Geoff Penington. Leading order corrections to the quantum extremal surface prescription. *arXiv preprint arXiv:2008.03319*, 2020.
- [AP22] Chris Akers and Geoff Penington. Quantum minimal surfaces from quantum error correction, 2022.
- [AR19] Chris Akers and Pratik Rath. Holographic Rényi entropy from quantum error correction. *Journal of High Energy Physics*, 2019(5):1–24, 2019.
- [AS17] Guillaume Aubrun and Stanisław J Szarek. *Alice and Bob meet Banach*, volume 223. American Mathematical Society, 2017.
- [ASY12] Guillaume Aubrun, Stanisław J Szarek, and Deping Ye. Phase transitions for random states and a semicircle law for the partial transpose. *Physical Review A*, 85(3):030302, 2012.
- [Aub12] Guillaume Aubrun. Partial transposition of random states and non-centered semicircular distributions. *Random Matrices: Theory and Applications*, 1(02):1250001, 2012.
- [BCCC11] Teodor Banica, Serban Teodor Belinschi, Mireille Capitaine, and Benoit Collins. Free Bessel laws. *Canadian Journal of Mathematics*, 63(1):3–37, 2011.
- [Bil08] Patrick Billingsley. *Probability and Measure*. John Wiley & Sons, 2008.
- [BPSW19] Ning Bao, Geoffrey Penington, Jonathan Sorce, and Aron C Wall. Beyond toy models: distilling tensor networks in full AdS/CFT. *Journal of High Energy Physics*, 2019(11):1–63, 2019.
- [BS10] Zhidong Bai and Jack W Silverstein. *Spectral Analysis of Large Dimensional Random Matrices*, volume 20. Springer, 2010.
- [BW20] Raphael Bousso and Elizabeth Wildenhain. Gravity/ensemble duality. *Physical Review D*, 102(6):066005, 2020.
- [CDKW14] Matthias Christandl, Brent Doran, Stavros Kousidis, and Michael Walter. Eigenvalue distributions of reduced density matrices. *Communications in Mathematical Physics*, 332(1):1–52, 2014.
- [CGGPG13] Benoît Collins, Carlos E González-Guillén, and David Pérez-García. Matrix product states, random matrix theory and the principle of maximum entropy. *Communications in Mathematical Physics*, 320(3):663–677, 2013.

- [CHLS15] Bartłomiej Czech, Patrick Hayden, Nima Lashkari, and Brian Swingle. The information theoretic interpretation of the length of a curve. *Journal of High Energy Physics*, 2015(6):1–40, 2015.
- [CN16] Benoit Collins and Ion Nechita. Random matrix techniques in quantum information theory. *Journal of Mathematical Physics*, 57(1):015215, 2016.
- [CNŻ10] Benoît Collins, Ion Nechita, and Karol Życzkowski. Random graph states, maximal flow and Fuss–Catalan distributions. *Journal of Physics A: Mathematical and Theoretical*, 43(27):275–303, 2010.
- [CNŻ13] Benoît Collins, Ion Nechita, and Karol Życzkowski. Area law for random graph states. *Journal of Physics A: Mathematical and Theoretical*, 46(30):305302, 2013.
- [CPGSV21] J Ignacio Cirac, David Perez-Garcia, Norbert Schuch, and Frank Verstraete. Matrix product states and projected entangled pair states: Concepts, symmetries, theorems. *Reviews of Modern Physics*, 93(4):045003, 2021.
- [DBWR14] Frédéric Dupuis, Mario Berta, Jürg Wullschleger, and Renato Renner. One-shot decoupling. *Communications in Mathematical Physics*, 328(1):251–284, 2014.
- [DF13] Lukas Drescher and Omar Fawzi. On simultaneous min-entropy smoothing. In *2013 IEEE International Symposium on Information Theory*, pages 161–165. IEEE, 2013.
- [DH10] Nicolas Dutil and Patrick Hayden. One-shot multiparty state merging. *arXiv preprint arXiv:1011.1974*, 2010.
- [DHM19] Xi Dong, Daniel Harlow, and Donald Marolf. Flat entanglement spectra in fixed-area states of quantum gravity. *Journal of High Energy Physics*, 2019(10):1–25, 2019.
- [DMW21] Xi Dong, Sean McBride, and Wayne W Weng. Replica wormholes and holographic entanglement negativity. *arXiv preprint arXiv:2110.11947*, 2021.
- [DQW21] Xi Dong, Xiao-Liang Qi, and Michael Walter. Holographic entanglement negativity and replica symmetry breaking. *arXiv preprint arXiv:2101.11029*, 2021.
- [Dup15] Frédéric Dupuis. Chain rules for quantum Rényi entropies. *Journal of Mathematical Physics*, 56(2):022203, 2015.
- [Dut11] Nicolas Dutil. Multiparty quantum protocols for assisted entanglement distillation. *arXiv preprint arXiv:1105.4657*, 2011.
- [EW15] Netta Engelhardt and Aron C Wall. Quantum extremal surfaces: holographic entanglement entropy beyond the classical regime. *Journal of High Energy Physics*, 2015(1):1–27, 2015.
- [FLM13] Thomas Faulkner, Aitor Lewkowycz, and Juan Maldacena. Quantum corrections to holographic entanglement entropy. *Journal of High Energy Physics*, 2013(11):1–18, 2013.
- [FR15] Omar Fawzi and Renato Renner. Quantum conditional mutual information and approximate Markov chains. *Communications in Mathematical Physics*, 340(2):575–611, 2015.
- [GAE07] David Gross, Koenraad Audenaert, and Jens Eisert. Evenly distributed unitaries: On the structure of unitary designs. *Journal of Mathematical Physics*, 48(5):052104, 2007.
- [Har17] Daniel Harlow. The Ryu-Takayanagi formula from quantum error correction. *Communications in Mathematical Physics*, 354(3):865–912, 2017.
- [Has17] Matthew B Hastings. The asymptotics of quantum max-flow min-cut. *Communications in Mathematical Physics*, 351(1):387–418, 2017.

- [Hay08] Masahito Hayashi. Second-order asymptotics in fixed-length source coding and intrinsic randomness. *IEEE Transactions on Information Theory*, 54(10):4619–4637, 2008.
- [HLW06] Patrick Hayden, Debbie W Leung, and Andreas Winter. Aspects of generic entanglement. *Communications in Mathematical Physics*, 265(1):95–117, 2006.
- [HNQ⁺16] Patrick Hayden, Sepehr Nezami, Xiao-Liang Qi, Nathaniel Thomas, Michael Walter, and Zhao Yang. Holographic duality from random tensor networks. *Journal of High Energy Physics*, 2016(11):1–56, 2016.
- [HRT07] Veronika E Hubeny, Mukund Rangamani, and Tadashi Takayanagi. A covariant holographic entanglement entropy proposal. *Journal of High Energy Physics*, 2007(07):062, 2007.
- [KFNR21] Jonah Kudler-Flam, Vladimir Narovlansky, and Shinsei Ryu. Negativity spectra in random tensor networks and holography. *arXiv preprint arXiv:2109.02649*, 2021.
- [KR05] Andreas Klappenecker and Martin Rotteler. Mutually unbiased bases are complex projective 2-designs. In *Proceedings. International Symposium on Information Theory, 2005. ISIT 2005.*, pages 1740–1744. IEEE, 2005.
- [LC21] Ryan Levy and Bryan K Clark. Entanglement entropy transitions with random tensor networks. *arXiv preprint arXiv:2108.02225*, 2021.
- [LM13] Aitor Lewkowycz and Juan Maldacena. Generalized gravitational entropy. *Journal of High Energy Physics*, 2013(8):1–29, 2013.
- [LPG21] Cécilia Lancien and David Pérez-García. Correlation length in random MPS and PEPS. *Annales Henri Poincaré*, pages 1–82, 2021.
- [LPWV20] Javier Lopez-Piqueres, Brayden Ware, and Romain Vasseur. Mean-field entanglement transitions in random tree tensor networks. *Physical Review B*, 102(6):064202, 2020.
- [LVFL21] Yaodong Li, Romain Vasseur, Matthew Fisher, and Andreas WW Ludwig. Statistical mechanics model for Clifford random tensor networks and monitored quantum circuits. *arXiv preprint arXiv:2110.02988*, 2021.
- [Mal99] Juan Maldacena. The large- N limit of superconformal field theories and supergravity. *International Journal of Theoretical Physics*, 38(4):1113–1133, 1999.
- [MB21] Erica Morgan and Fernando GSL Brandão. A classical model correspondence for G -symmetric random tensor networks. *Journal of Physics Communications*, 2021.
- [MM20] Donald Marolf and Henry Maxfield. Transcending the ensemble: baby universes, spacetime wormholes, and the order and disorder of black hole information. *Journal of High Energy Physics*, 2020(8):1–72, 2020.
- [MS17] James A Mingo and Roland Speicher. *Free Probability and Random Matrices*, volume 35. Springer, 2017.
- [MVS21] Raimel Medina, Romain Vasseur, and Maksym Serbyn. Entanglement transitions from restricted Boltzmann machines. *Physical Review B*, 104(10):104205, 2021.
- [MWW20] Donald Marolf, Shannon Wang, and Zhencheng Wang. Probing phase transitions of holographic entanglement entropy with fixed area states. *Journal of High Energy Physics*, 2020(12):1–41, 2020.

- [NRSR21] Adam Nahum, Sthitadhi Roy, Brian Skinner, and Jonathan Ruhman. Measurement and entanglement phase transitions in all-to-all quantum circuits, on quantum trees, and in Landau-Ginsburg theory. *PRX Quantum*, 2(1):010352, 2021.
- [NS06] Alexandru Nica and Roland Speicher. *Lectures on the combinatorics of free probability*, volume 13. Cambridge University Press, 2006.
- [NW20] Sepehr Nezami and Michael Walter. Multipartite entanglement in stabilizer tensor networks. *Physical Review Letters*, 125(24):241602, 2020.
- [PB20] Marc Potters and Jean-Philippe Bouchaud. *A First Course in Random Matrix Theory: For Physicists, Engineers and Data Scientists*. Cambridge University Press, 2020.
- [Pen20] Geoffrey Penington. Entanglement wedge reconstruction and the information paradox. *Journal of High Energy Physics*, 2020(9):1–84, 2020.
- [PSSY19] Geoff Penington, Stephen H Shenker, Douglas Stanford, and Zhenbin Yang. Replica wormholes and the black hole interior. *arXiv preprint arXiv:1911.11977*, 2019.
- [PYHP15] Fernando Pastawski, Beni Yoshida, Daniel Harlow, and John Preskill. Holographic quantum error-correcting codes: Toy models for the bulk/boundary correspondence. *Journal of High Energy Physics*, 2015(6):1–55, 2015.
- [QSY21] Xiao-Liang Qi, Zhou Shangnan, and Zhenbin Yang. Holevo information and ensemble theory of gravity. *arXiv preprint arXiv:2111.05355*, 2021.
- [QY18] Xiao-Liang Qi and Zhao Yang. Space-time random tensor networks and holographic duality. *arXiv preprint arXiv:1801.05289*, 2018.
- [QYY17] Xiao-Liang Qi, Zhao Yang, and Yi-Zhuang You. Holographic coherent states from random tensor networks. *Journal of High Energy Physics*, 2017(8):1–29, 2017.
- [RT06a] Shinsei Ryu and Tadashi Takayanagi. Aspects of holographic entanglement entropy. *Journal of High Energy Physics*, 2006(08):045, 2006.
- [RT06b] Shinsei Ryu and Tadashi Takayanagi. Holographic derivation of entanglement entropy from the anti-de Sitter space/conformal field theory correspondence. *Physical Review Letters*, 96(18):181602, 2006.
- [Sár17] Gábor Sárosi. AdS₂ holography and the SYK model. *arXiv preprint arXiv:1711.08482*, 2017.
- [SSS19] Phil Saad, Stephen H Shenker, and Douglas Stanford. JT gravity as a matrix integral. *arXiv preprint arXiv:1903.11115*, 2019.
- [SSSY21] Phil Saad, Stephen H Shenker, Douglas Stanford, and Shunyu Yao. Wormholes without averaging. *arXiv preprint arXiv:2103.16754*, 2021.
- [Swi12a] Brian Swingle. Constructing holographic spacetimes using entanglement renormalization. *arXiv preprint arXiv:1209.3304*, 2012.
- [Swi12b] Brian Swingle. Entanglement renormalization and holography. *Physical Review D*, 86(6):065007, 2012.
- [TH13] Marco Tomamichel and Masahito Hayashi. A hierarchy of information quantities for finite block length analysis of quantum tasks. *IEEE Transactions on Information Theory*, 59(11):7693–7710, 2013.

- [Tom12] Marco Tomamichel. *A framework for non-asymptotic quantum information theory*. PhD thesis, ETH Zurich, 2012.
- [Tom15] Marco Tomamichel. *Quantum Information Processing with Finite Resources: Mathematical Foundations*, volume 5. Springer, 2015.
- [VC04] Frank Verstraete and J Ignacio Cirac. Valence-bond states for quantum computation. *Physical Review A*, 70(6):060302, 2004.
- [VMC08] Frank Verstraete, Valentin Murg, and J Ignacio Cirac. Matrix product states, projected entangled pair states, and variational renormalization group methods for quantum spin systems. *Advances in Physics*, 57(2):143–224, 2008.
- [VPYL19] Romain Vasseur, Andrew C Potter, Yi-Zhuang You, and Andreas WW Ludwig. Entanglement transitions from holographic random tensor networks. *Physical Review B*, 100(13):134203, 2019.
- [Wan22] Jinzhao Wang. *ON THE INFORMATION-THEORETIC ASPECTS OF BLACK HOLES*. PhD thesis, ETH Zurich, 2022.
- [WW21] Michael Walter and Freek Witteveen. Hypergraph min-cuts from quantum entropies. *Journal of Mathematical Physics*, 62(9):092203, 2021.
- [YHQ16] Zhao Yang, Patrick Hayden, and Xiao-Liang Qi. Bidirectional holographic codes and sub-AdS locality. *Journal of High Energy Physics*, 2016(1):1–24, 2016.
- [YLFC21] Zhi-Cheng Yang, Yaodong Li, Matthew Fisher, and Xiao Chen. Entanglement phase transitions in random stabilizer tensor networks. *arXiv preprint arXiv:2107.12376*, 2021.
- [YYQ18] Yi-Zhuang You, Zhao Yang, and Xiao-Liang Qi. Machine learning spatial geometry from entanglement features. *Physical Review B*, 97(4):045153, 2018.