Appearance of an Infalling Star in Black Holes with Multiple Photon Spheres

Yiqian Chen^a,^{*} Guangzhou Guo^a,[†] Peng Wang^a,[‡] Houwen Wu^{a,b},[§] and Haitang Yang^a¶

^aCenter for Theoretical Physics, College of Physics, Sichuan University, Chengdu, 610064, China and ^bDepartment of Applied Mathematics and Theoretical Physics, University of Cambridge, Wilberforce Road, Cambridge, CB3 0WA, UK

Photon spheres play a pivotal role in the imaging of luminous objects near black holes. In this paper, we examine observational appearances of a star freely falling in hairy black holes, which can possess one or two photon spheres outside the event horizon. When there exists a single photon sphere, the total luminosity measured by distant observers decreases exponentially with time at late times. Due to successive arrivals of photons orbiting around the photon sphere different times, a specific observer would see a series of light flashes with decreasing intensity, which share a similar frequency content. Whereas in the case with two photon spheres, photons temporarily trapped between the photon spheres can cause a peak of the total luminosity, which is followed by a slow exponential decay, at late times. In addition, these photons lead to one more series of light flashes seen by the specific observer.

^{*} chenyiqian@stu.scu.edu.cn

[†] guangzhouguo@stu.scu.edu.cn

[‡] pengw@scu.edu.cn

[§] hw598@damtp.cam.ac.uk

[¶] hyanga@scu.edu.cn

CONTENTS

I.	Introduction	2
II.	Set up	4
III.	Numerical Results	6
	A. Single-peak Potential	7
	B. Double-peak Potential	12
IV.	Conclusions	18
	Acknowledgments	19
	References	19

I. INTRODUCTION

The recent announcements of the images of the supermassive black holes M87^{*} [1–8] and Sgr A^{*} [9–14] by the Event Horizon Telescope (EHT) collaboration open a new window to test general relativity in the strong field regime. These images display two main features, a central brightness depression, dubbed "shadow", and a bright ring, which is closely relevant to a class of circular and unstable photon orbits. For static spherically symmetric black holes, these circular and unstable photon orbits would form photon spheres outside the event horizon. Since the shadow and the bright ring are originated from light deflections by the strong gravitational field near the photon spheres [15–22], black hole images can encode valuable information of the geometry in the vicinity of the photon spheres. In particular, black hole images have been widely studied in the context of different theories of gravity, e.g., nonlinear electrodynamics [23–29], the Gauss-Bonnet theory [30–33], fuzzball [34], the Chern-Simons type theory [35, 36], f(R) gravity [37–39], string inspired black holes [40–43], wormholes [44–50], bosonic stars [51, 52] and other theories [53–65]. Furthermore, the EHT observations can also be applied to impose constraints on the cosmological parameters [66–71] and the size of extra dimensions [72–74], test the equivalence principle [75–77], and probe some fundamental physics issues including dark matter [78–83] and dark energy [84–88].

Apart from stationary observational appearances of black holes, photon spheres also play an important role in dynamic observations of luminous objects around black holes. A particularly interesting scenario is luminous matter freely falling onto black holes, which occurs frequently and was reported near the Cyg X-1 black hole [89] and the Sgr A* source [90, 91]. Recently, the authors of [92] numerically simulated the appearance of a point-like source, which emits photons or gravitons as it falls into a Schwarzschild black hole. Note that there exists a single photon sphere outside the event horizon in a Schwarzschild black hole. It showed that, owing to radiations temporarily trapped just outside the photon sphere, late-time radiations are dominated by blueshifted photons or gravitons with a near-critical impact parameter, and the total luminosity decreases exponentially as $e^{-\lambda t}$, where λ is the Lyapunov exponent of circular null geodesics at the photon sphere.

On the other hand, a novel class of Einstein-Maxwell-scalar (EMS) models have recently been constructed to understand the formation of hairy black holes [93–97]. In such models, the scalar field non-minimally couples to the electromagnetic field, which can trigger a tachyonic instability to form spontaneously scalarized hairy black holes from Reissner-Nordström (RN) black holes. Properties of the hairy black holes have been extensively studied in the literature, e.g., different non-minimal coupling functions [98–100], massive and self-interacting scalar fields [101, 102], horizonless reflecting stars [103], stability analysis [104–108], higher dimensional scalar-tensor models [109], quasinormal modes [110, 111], two U(1) fields [112], quasi-topological electromagnetism [113], topology and spacetime structure influences [114], and in dS/AdS spacetime [96, 115–117].

Intriguingly, the scalarized hairy black holes have been shown to possess two photon spheres outside the event horizon in certain parameter regions [118, 119]. The existence of an extra photon sphere can significantly affect optical appearances of surrounding accretion disks (e.g., producing bright rings of different radius in the black hole images [118] and noticeably increasing the flux of the observed images [119]) and luminous celestial spheres (e.g., tripling higher-order images of the celestial spheres [120]). Moreover, the effective potential for a scalar perturbation in the hairy black holes with two photon spheres was found to exhibit a double-peak structure, leading to long-lived quasinormal modes [121] and echo signals [122]. It is worth noting that the existence of two photon spheres outside the event horizon has also been reported for dyonic black holes with a quasi-topological electromagnetic term [123], black holes in massive gravity [124, 125] and wormholes in the black-bounce spacetime [126–128].

In this paper, we aim to study how an extra photon sphere affects observational appearances of a freely-falling star in the context of the hairy black holes. The rest of the paper is organized as follows. In Section II, we briefly review hairy black hole solutions in the EMS model and introduce the observational settings. Numerical results are presented in Section III. Section IV is devoted to our main conclusions. We set $16\pi G = c = 1$ throughout the paper.

II. SET UP

We consider a 4-dimensional EMS model with the action [93]

$$S = \int d^4x \sqrt{-g} \left[\mathcal{R} - 2\partial_\mu \phi \partial^\mu \phi - e^{\alpha \phi^2} F_{\mu\nu} F^{\mu\nu} \right], \tag{1}$$

where \mathcal{R} is the Ricci scalar, and $F_{\mu\nu} = \partial_{\mu}A_{\nu} - \partial_{\nu}A_{\mu}$ is the electromagnetic field strength tensor. In this EMS model, the scalar field ϕ is non-minimally coupled to the electromagnetic field A_{μ} with the coupling function $e^{\alpha\phi^2}$. Restricting to static and spherically symmetric black hole solutions, one can have the generic ansatz

$$ds^{2} = -N(r)e^{-2\delta(r)}dt^{2} + \frac{1}{N(r)}dr^{2} + r^{2}\left(d\theta^{2} + \sin^{2}\theta d\varphi^{2}\right),$$

$$A_{\mu}dx^{\mu} = V(r)dt \text{ and } \phi = \phi(r).$$
(2)

For this ansatz, the equations of motion are

$$N'(r) = \frac{1 - N(r)}{r} - \frac{Q^2}{r^3 e^{\alpha \phi^2(r)}} - rN(r) \left[\phi'(r)\right]^2,$$

$$\left[r^2 N(r)\phi'(r)\right]' = -\frac{\alpha Q^2 \phi(r)}{r^2 e^{\alpha \phi^2(r)}} - r^3 N(r) \left[\phi'(r)\right]^3,$$

$$\delta'(r) = -r \left[\phi'(r)\right]^2,$$

$$V'(r) = \frac{Q}{r^2 e^{\alpha \phi^2(r)}} e^{-\delta(r)},$$

(3)

where the integration constant Q can be interpreted as the electric charge of the black hole, and primes stand for derivative with respect to r. In addition, the ADM mass M can be obtained via $M = \lim_{r \to \infty} r \left[1 - N(r)\right]/2$. Moreover, we impose proper boundary conditions at the event horizon r_h and spatial infinity,

$$N(r_h) = 0, \ \delta(r_h) = \delta_0, \ \phi(r_h) = \phi_0, \ V(r_h) = 0,$$

$$N(\infty) = 1, \ \delta(\infty) = 0, \ \phi(\infty) = 0, \ V(\infty) = \Phi,$$
 (4)

where δ_0 and ϕ_0 can be used to characterize black hole solutions, and Φ is the electrostatic potential. Specifically, $\phi_0 = \delta_0 = 0$ correspond to the scalar-free solutions with $\phi = 0$, i.e., RN black holes. When non-zero values of ϕ_0 and δ_0 are admitted, hairy black hole solutions with a non-trivial scalar field ϕ are obtained by a shooting method built in the *NDSolve* function of *Wolfram* (*Mathematica*.

In this paper, we study a point-like star freely falling along the radial direction at $\theta = \frac{\pi}{2}$ and $\varphi = 0$. The equations governing geodesics on the equatorial plane can be derived from the Lagrangian

$$\mathcal{L} = \frac{1}{2} \left[-N(r)e^{-2\delta(r)}\dot{t}^2 + \frac{1}{N(r)}\dot{r}^2 + r^2\dot{\varphi}^2 \right],\tag{5}$$

where dots stand for derivative with respect to some affine parameter λ . Note that $\mathcal{L} = -1/2$ for time-like geodesics if λ is the proper time. Since t and φ do not explicitly appear in the Lagrangian, one can characterize geodesics by their energy E and angular momentum L, namely

$$p_t \equiv -N(r)e^{-2\delta(r)}\dot{t} = -E, \quad p_{\varphi} \equiv r^2\dot{\varphi} = L.$$
(6)

For simplicity, we specialize to a radially freely falling star starting at infinity with a zero initial velocity, whose four-velocity is

$$v_e^{\mu}(r) = \left(\frac{e^{2\delta(r)}}{N(r)}, -\sqrt{e^{2\delta(r)} - N(r)}, 0, 0\right).$$
(7)

Suppose that the star emits photons when it falls radially onto a black hole. Due to spherical symmetry, we can confine ourselves to emissions on the equatorial plane. The Lagrangian (5) can also be used to describe null geodesics on the equatorial plane. Moreover, the constancy of the Lagrangian $\mathcal{L} = 0$ rewrites the radial component of the null geodesic equations as

$$\frac{e^{-2\delta(r)}}{L^2}\dot{r}^2 = \frac{1}{b^2} - V_{\text{eff}}(r),\tag{8}$$

where $b \equiv L/E$ is the impact parameter, and $V_{\text{eff}} \equiv e^{-2\delta(r)}N(r)r^{-2}$ is the effective potential. Unstable circular null geodesics at radius r_{ph} , which constitute a photon sphere of radius r_{ph} , are determined by

$$V_{\rm eff}(r_{\rm ph}) = \frac{1}{b_{\rm ph}^2}, \quad V_{\rm eff}'(r) = 0, \quad V_{\rm eff}''(r_{\rm ph}) < 0, \tag{9}$$

where $b_{\rm ph}$ is the corresponding impact parameter. In short, a local maximum of the effective potential corresponds to a photon sphere. Remarkably, it showed that, when the black hole charge is large enough, $V_{\rm eff}(r)$ of the hairy black hole solutions (2) can have two local maxima, leading to a double-peak structure [118, 119]. Consequently, the hairy black holes with the double-peak effective potential possess two photon spheres outside the event horizon. Indeed, the effective potential of the hairy black hole with $\alpha = 0.9$ and Q = 1.064M is plotted in FIG. 6, which exhibits a double-peak structure.

To trace light rays from the star to a far-away observers, one need to supply initial conditions for Eqs. (6) and (8). For a photon of four-momentum p_{μ} , the momentum measured in the rest frame of the star at $r = r_e$ is

$$p^{\hat{t}} = -\left[N(r_e)e^{-2\delta(r_e)}\right]^{-1} p_t + \sqrt{e^{2\delta(r_e)} - N(r_e)}p_r,$$

$$p^{\hat{r}} = -\left[N(r_e)e^{-2\delta(r_e)}\right]^{-1} \sqrt{1 - N(r_e)e^{-2\delta(r_e)}}p_t + \sqrt{e^{2\delta(r_e)}}p_r,$$

$$p^{\hat{\theta}} = 0, \ p^{\hat{\varphi}} = \frac{p_{\varphi}}{r_e}.$$
(10)

The emission angle β is defined as

$$\cos\beta = \frac{p^{\hat{r}}}{p^{\hat{t}}},\tag{11}$$

which is the angle between the propagation direction of the photon and the radial direction in the rest frame of the star. With Eqs. (10) and (11), one has the normalized frequency of the photon measured by the far-away observer,

$$\frac{\omega_o}{\omega_e} \equiv \frac{p^t}{p^{\hat{t}}} = 1 - \cos\beta \sqrt{1 - N(r_e)e^{-2\delta(r_e)}},\tag{12}$$

where ω_e and ω_o are rest-frame and observed photon frequencies, respectively. Furthermore, the luminosity is defined as $L_i \equiv dE_i/d\tau_i$, where E_i is the total energy of the photon, τ_i is the proper time, and i = e and o denote quantities corresponding to the emitter and observer, respectively. Similar to the normalized frequency, we define the normalized luminosity as

$$\frac{L_o}{L_e} = \frac{dE_o/d\tau_o}{dE_e/d\tau_e} \approx \frac{\omega_o dn_o}{\omega_e dn_e} \left(\frac{dt_o}{d\tau_e}\right)^{-1},$$

where n_o and n_e are the observed and emitted photon numbers, respectively, and we replaced $d\tau_o$ by dt_o since they are almost the same for the far-away observer.

III. NUMERICAL RESULTS

In this section, we numerically study observational appearances of a star falling radially towards hairy black holes with a single or two photon spheres outside the event horizon. During the free fall of the star, photons are emitted isotropically in its rest frame. Specifically, we assume that the star starts emitting photons at t = 0 and $r_e = 31.155M$, and emits 3200 photons, which are uniformly distributed in the emission angle β , every proper time interval $\delta \tau = 0.002M$. In addition, observers collect photons on a celestial sphere at the radius $r_o = 100M$. To calculate observed luminosities, the collected photons are grouped into packets of 50 (i.e., $dn_o = 50$) according to their arrival time. Without loss of generality, we set M = 1 in the rest of this section.



FIG. 1. The black line denotes the effective potential of null geodesics in the hairy black hole with $\alpha = 0.9$ and Q = 1.054. There is only one potential maximum at $r_{\rm ph} = 1.610$, corresponding to the photon sphere with the critical impact parameter $b_{\rm ph} = 3.673$. Colored regions are presented in the *r*-*b* parameter space, for which a photon with the impact parameter *b* is emitted at *r*. A far-away observer can not receive photons emitted in the gray region. Photons emitted in the pink and purple regions have impact parameters close to $b_{\rm ph}$, hence can be temporarily trapped around the photon sphere. In particular, after photons in the pink (purple) region are emitted outward inside (inward outside) the photon sphere, they would circle around the photon sphere more than once before reaching an observer.

A. Single-peak Potential

We first consider the hairy black hole with $\alpha = 0.9$ and Q = 1.054, which has a single-peak potential as shown in FIG. 1. The photon sphere is located at $r_{\rm ph} = 1.610$, and the corresponding critical impact parameter is $b_{\rm ph} = 3.673$. Ingoing photons with $b < b_{\rm ph}$ can overcome the effective potential barrier and get captured by the black hole. Therefore for $b < b_{\rm ph}$, a distant observer can only receive outward-emitted photons. On the other hand, inward-emitted photons with $b > b_{\rm ph}$ can be observed since they bounce outward from the potential barrier. Interestingly, strong gravitational lensing near the photon sphere causes extreme bending of light rays with a nearcritical impact parameter, which makes them temporarily trapped around the photon sphere and take a long time to reach the observer. As a result, light rays with a near-critical impact parameter play a crucial role in the observational appearance of the infalling star at late times. To illustrate photons with a near-critical impact parameter, we classify photons into four categories according to their b,

- b < 3.564. Yellow region in FIG. 1 and yellow dots in FIGs. 4 and 5.
- $3.564 \le b < b_{\rm ph}$. Pink region in FIG. 1 and pink dots in FIGs. 4 and 5. The left panel of



FIG. 2. Photon trajectories in the hairy black hole with a single potential. The blue dashed circles denote the photon sphere. Left: The trajectory of a photon with b = 3.564 emitted at $r_e = 1.2$. The light ray has the angular coordinate change $\Delta \varphi = 2\pi$ when reaching a celestial sphere at $r_o = 100$. Right: The trajectory of a photon with b = 3.900 emitted at $r_e = 5$. The light ray has the angular coordinate change $\Delta \varphi = 2\pi$ when reaching the celestial sphere.

FIG. 2 depicts a light ray of b = 3.564, which corresponds to an outward-emitted photon that is emitted at $r_e = 1.2$ inside the photon sphere and reaches the celestial sphere after the change of angular coordinate $\Delta \varphi = 2\pi$. It is worth emphasizing that near-critical photons of $b < b_{\rm ph}$ that are emitted outward inside the photon sphere would linger near the photon sphere for some time.

- $b_{\rm ph} < b \leq 3.900$. Purple region in FIG. 1 and purple dots in FIGs. 4 and 5. The right panel of FIG. 2 shows the trajectory of an inward-emitted photon with b = 3.900, which is emitted at $r_e = 5$ outside the photon sphere and reaches the celestial sphere after the change of angular coordinate $\Delta \varphi = 2\pi$. Near-critical photons of $b > b_{\rm ph}$ that are emitted inward can circle around the photon sphere more than once before being observed, and contribute significantly to late-time observations. In this case, the photon sphere acts as a reflecting wall to scatter photons, which indicates that the near-critical photons can undergo a blueshift.
- b > 3.900. Green region in FIG. 1 and green dots in FIGs. 4 and 5.

Considering photons emitted with an impact parameter very close to $b_{\rm ph}$ in the pink and purple regions of FIG. 1, their normalized frequencies measured by observers on the celestial sphere are plotted against the position of the star in FIG. 3. Particularly, the purple line represents photons emitted inward in the purple region, and the pink line denotes photons emitted outward inside the photon sphere in the pink region. The normalized frequency of a photon is determined by the



FIG. 3. The normalized frequency ω_o/ω_e as a function of the position of the star r_e for photons of impact parameter very close to $b_{\rm ph}$, which are emitted outward and inward in the pink and purple regions of FIG. 1, respectively. For inward-emitted photons, the Doppler effect competes with the gravitational redshift to determine the observed frequency, leading to a blueshift peak $\omega_o/\omega_e = 1.378$ at $r_e = 8.970$.

gravitational redshift and the Doppler effect, which are controlled by the position and the velocity of the photon when it is emitted, respectively. Due to light-bending near the photon sphere, the Doppler effect can increase the observed frequency of inward-emitted photons. At large r_e , the gravitational redshift is weak, and hence the photons in purple line are blueshifted. The blueshift reaches the maximum $\omega_o/\omega_e = 1.378$ at $r_e = 8.970$ and becomes 1 at $r_e = 3.673$. For $r_e < 3.673$, as a result of strong gravitational redshift near the black hole, the observed frequency is redshifted and approaches zero as r_e goes to r_h .

In the left panel of FIG. 4, we display the normalized frequency distribution of photons, which are emitted from the infalling star and collected by the observers distributed on the celestial sphere. At early times, the received photons are dominated by those emitted in the green region of FIG. 1. Later, photons emitted inward in the near-critical purple region start to reach the observers after circling around the photon sphere, and come to dominate the high-frequency observation. On the other hand, photons emitted outward in other color regions are found to be severely redshifted since both the Doppler effect and the gravitational redshift decrease the observed frequency. The corresponding normalized total luminosity are presented in the right panel of FIG. 4, where a dot corresponds to a packet of 50 photons, and the color of the dot is that having most photons in the packet. The luminosity gradually increases until reaching a peak at early times, and are dominated by photons emitted in the green region roughly before $t_o = 215$, which is in agreement with the frequency observation. At late times, the luminosity is mostly determined by photons with a near-critical impact parameter emitted in the purple region. Interestingly, we find that the



FIG. 4. The normalized frequency distribution and total luminosity of photons emitted from a freelyfalling star in the hairy black hole with a single-peak potential. The star radially falls along $\phi = 0$ on the the equatorial plane from spatial infinity at rest and starts emitting isotropically in its rest frame at $r_e = 31.155$. The emitted photons are collected by observers on a celestial sphere at $r_o = 100$. Left: The observers receive photons with a wide range of frequencies. In the early stage, photons emitted in the green region of FIG. 1 dominate the frequency observation. The late-time high-frequency observation is determined by photons emitted in the purple region, which undergo extreme bending near the photon sphere. Right: The luminosity is calculated by grouping received photons into packets of 50. At late times, the luminosity is mainly controlled by photons emitted with a near-critical impact parameter in the purple region. The luminosity decays exponentially as $e^{-\lambda t} \simeq e^{-0.140t}$, where λ is the Lyapunov exponent of circular null geodesics at the photon sphere. Note that the imaginary part of the associated lowest-lying quasinormal modes is $-\lambda/2$.

luminosity decays exponentially as $e^{-\lambda t} \simeq e^{-0.140t}$, where λ is the Lyapunov exponent of circular null geodesics at the photon sphere. Indeed, it was argued that the late-time decay of the luminosity \mathcal{L} is determined by the imaginary part of quasinormal frequencies ω , i.e., $\mathcal{L} \propto e^{2 \operatorname{Im} \omega t}$, in the eikonal limit [92]. For the hairy black hole with a single-peak potential, the imaginary part of the lowestlying quasinormal frequencies is $-\lambda/2$ in the eikonal limit [121], which gives $\mathcal{L} \propto e^{-\lambda t}$. Note that the finite number of photons in a packet results in the non-smoothness of the luminosity in FIG. 4. The luminosity can become smoother by including more photons in a packet, which, nevertheless, reduces the accuracy of our numerical results with fixed number of total emitted photons.

For a specific observer located at $\varphi = 0$ and $\theta = \pi/2$ on the celestial sphere, the angular coordinate change $\Delta \varphi$ of light rays connecting the star with the observer is

$$\Delta \varphi = 2n\pi,\tag{13}$$

where $n = 0, 1, 2 \cdots$ is the number of orbits that the light rays complete around the black hole.



FIG. 5. The normalized frequency and luminosity of the infalling star measured by a far-away observer at $r_o = 100$, $\theta = \pi/2$ and $\phi = 0$ in the hairy black hole with a single-peak potential. The colored dots denote photons emitted in the regions with the same color in FIG. 1. **Left:** The received photons form several frequency lines indexed by the number n of photon orbits around the black hole. From left to right, the frequency lines correspond to n = 0, 1, 2 and 3, respectively. In particular, photons with a impact parameter close to $b_{\rm ph}$ give rise to the self-similar $n \ge 1$ lines, which display blueshifts caused by the Doppler effect and the strong gravitational lensing. The time delay between the $n \ge 1$ lines is roughly the period of circular null geodesics at the photon sphere, $\Delta T \simeq 2\pi b_{\rm ph} \simeq 23$. **Right:** At early times, the luminosity is dominated by outward-emitted photons with a small impact parameter and gradually decreases. Subsequently, blueshifted n = 1 photons start to reach the observer and then become the most dominant contribution, which results in the growth of the luminosity. Later, the luminosity drops sharply with the decrease in the frequency of the n = 1 photons. Consequently, a flash is observed around $t_o \simeq 215$. When the n = 2 photons play out, a similar steady rise followed by a steep fall is also observed, leading to a much fainter flash.

To obtain the observational appearance of the star seen by the observer, we focus on the collected photons with $\cos \varphi > 0.99$. We present the frequency observation in the left panel of FIG. 5, which shows a discrete spectrum separated by the received time. The leftmost line is formed by photons with n = 0, which radially propagate to the observer. As expected, the observed frequency of the n = 0 photons decreases with the received time because of the gravitational redshift. The rest three lines correspond to light rays with n = 1, 2 and 3 from left to right, and have a similar shape since the emission angle is almost same for the $n \ge 1$ light rays. As discussed earlier, the $n \ge 1$ photons can be blueshifted due to the Doppler effect and the strong gravitational lensing. With a fixed n, the observed frequencies increase slowly, reach a maximum and then decrease rapidly. In addition, the time delay between the adjacent lines roughly equals to the time it takes to orbit the photon sphere, i.e., $\Delta T \simeq 2\pi b_{\rm ph} \simeq 23^1$. Furthermore, although the received n = 1 photons

¹ Eqn. (6) leads to $dt/d\phi|_{r_{\rm ph}} = b^{-1}V_{\rm eff}^{-1}(r_{\rm ph}) = b_{\rm ph}$, which gives $\Delta T \simeq 2\pi b_{\rm ph}$.

mainly come from the purple and pink regions of FIG. 1, those emitted in the green and yellow regions are detected at early and late times, respectively. Note that the angular coordinate changes of light rays with a given impact parameter vary with the emitted position r_e . When r_e is small (large) enough, photons emitted in the yellow (green) regions can circle around the black hole once before being received. For $n \ge 2$, the observer only receive photons emitted in the purple and pink regions, which can circle around the black hole more than once due to the near-critical impact parameter.

The left panel of FIG. 5 shows the observed normalized luminosity as a function of the observed time, which exhibits a steady decline at early times owing to the gravitational redshift. Around $t_o \simeq 200$, blueshifted photons with n = 1 start to play a dominant role, leading to a gradual rise of luminosity until reaching a peak. Afterwards, a sharp drop of luminosity is observed due to strong redshifts of photons emitted at small r_e . Later, photons with n = 2 start to dominate the contribution to the luminosity. The luminosity of the n = 2 photons is about 40 times smaller than that of the n = 1 photons since the n = 2 photons are much less than the n = 1 photons. Despite different magnitudes, the luminosities of the photons with n = 1 and 2 have a similar shape. In short, successive receptions of photons orbiting around the photon sphere bring about a series of observed flashes labelled by n. Note that our numerical simulation does not have enough emitted photons to produce the luminosity of the n > 2 photons.

B. Double-peak Potential

The hairy black hole with $\alpha = 0.9$ and Q = 1.064 has a double-peak effective potential as shown in FIG. 6. The inner peak corresponds to the inner photon sphere with the critical impact parameter $b_{\rm in} = 3.455$ at $r_{\rm in} = 0.403$, and the outer peak to the outer photon sphere with $b_{\rm out} = 3.584$ at $r_{\rm out} = 1.463$. Similarly, emitted photons are sorted into seven categories by their impact parameter $b_{\rm in}$

- b < 2.884. Yellow region in FIG. 6 and yellow dots in FIGs. 9 and 10.
- 2.884 ≤ b < b_{in}. Pink region in FIG. 6 and pink dots in FIGs. 9 and 10. In this category, photons emitted outward inside the inner photon sphere can circle around the inner photon sphere before reaching a distant observer. For example, a light ray with b = 2.884, which has Δφ = 2π, is displayed in the upper-left panel of FIG. 7.
- $b_{\rm in} < b \le 3.458$. Brown region in FIG. 6 and brown dots in FIGs. 9 and 10. In this category,



FIG. 6. The effective potential of null geodesics in the hairy black hole with $\alpha = 0.9$ and Q = 1.064. The potential has two peaks at $r_{\rm in} = 0.403$ and $r_{\rm out} = 1.463$, corresponding to the inner photon sphere with $b_{\rm in} = 3.455$ and the outer one with $b_{\rm out} = 3.584$, respectively. The right panel highlights the region near the inner peak. When photons are emitted outward (inward) at $r < r_{\rm in}$ ($r > r_{\rm out}$) in the pink (purple) region, they usually orbit the black hole with $\Delta \varphi \ge 2\pi$. Photons emitted in the brown, blue and orange regions have $b_{\rm in} < b < b_{\rm out}$ and can be temporarily trapped in the region between the inner and outer photon spheres by orbiting the black hole with $\Delta \varphi \ge 6\pi$. Hence, these photons play a key role in late-time observations.

photons emitted inward would circle around the inner photon sphere roughly with $\Delta \varphi \ge 10\pi$ before escaping to the celestial sphere. For example, a light ray with b = 3.458, which has $\Delta \varphi = 10\pi$, is displayed in the upper-right panel of FIG. 7.

- 3.458 < b ≤ 3.547. Blue region in FIG. 6 and blue dots in FIGs. 9 and 10. In this category, photons can be temporarily trapped between the inner and outer photon spheres by circling around the black hole approximately between three and five times. For example, a light ray with b = 3.547, which has Δφ = 10π, is displayed in the lower-left panel of FIG. 7.
- $3.547 < b < b_{out}$. Orange region in FIG. 6 and orange dots in FIGs. 9 and 10. In this category, if photons are emitted inward outside the outer photon sphere, they would linger for some time around the outer photon sphere by orbiting the black hole approximately with $\Delta \varphi \geq 10\pi$.
- b_{out} < b ≤ 3.847. Purple region in FIG. 6 and purple dots in FIGs. 9 and 10. In this category, photons emitted inward outside the outer photon sphere usually circle around the outer photon sphere more than once. For example, a light ray with b = 3.847, which has Δφ = 2π, is displayed in the lower-right panel of FIG. 7.
- b > 3.847. Green region in FIG. 6 and green dots in FIGs. 9 and 10.



FIG. 7. Photon trajectories in the hairy black hole with a double-peak potential. The blue dashed circles denote the photon spheres. **Upper-Left**: A photon is emitted at $r_e = 0.35$ with b = 2.884, and the light ray has $\Delta \varphi = 2\pi$. **Upper-Right**: A photon is emitted at $r_e = 5$ with b = 3.458, and the light ray has $\Delta \varphi = 10\pi$. Lower-Left: A photon is emitted at $r_e = 5$ with b = 3.547, and the light ray has $\Delta \varphi = 10\pi$. Lower-Left: A photon is emitted at $r_e = 5$ with b = 3.547, and the light ray has $\Delta \varphi = 10\pi$.

In FIG. 8, we plot the normalized frequency ω_o/ω_e as a function of the emitted position r_e for near-critical photons. Specifically, we focus on inward-emitted photons with b very close to $b_{\rm in}$ at $r_e > r_{\rm in}$ in the brown region, outward-emitted photons with b very close to $b_{\rm in}$ at $r_e < r_{\rm in}$ in the pink region, inward-emitted photons with b very close to $b_{\rm out}$ at $r_e > r_{\rm out}$ in the purple region and photons emitted with b very close to $b_{\rm out}$ at $r_e < r_{\rm out}$ in the orange region. The colors of the lines in FIG. 8 match those of the corresponding emitted regions in FIG. 6. Moreover, photons with b very close to $b_{\rm in}$ and $b_{\rm out}$ are denoted by solid and dashed lines, respectively. Similar to the single-peak case, strong gravitational lensing around the inner and outer photon spheres can cause blueshifts of the near-critical photons when r_e is large enough. In particular, the normalized frequency of photons with b very close to $b_{\rm in}$ ($b_{\rm out}$) reaches the maximum $\omega_o/\omega_e = 1.387$ at $r_e = 8.512$ ($r_e = 8.784$) and becomes the unit at $r_e = 3.455$ ($r_e = 3.584$). In addition, the inset



FIG. 8. The normalized frequency ω_o/ω_e as a function of the emitted position r_e for photons, whose impact parameter b is very close to b_{in} (solid lines) or b_{out} (dashed lines), in the hairy black hole with a double-peak potential. For large r_e , the inward-emitted and near-critical photons are blueshifted since the Doppler effect dominates over the gravitational redshift. The photons emitted inward and outward between the two photon spheres can both reach a distant observer, and denote the upper and lower branches of the dashed line in the inset, respectively.

shows that the dashed line is divided into two branches for $r_{\rm in} < r_e < r_{\rm out}$, and the upper and lower branches correspond to inward-emitted and outward-emitted photons emitted between the two photon spheres, respectively.

The normalized frequency distribution of photons received by the observers distributed on the celestial sphere is presented in the left panel of FIG. 9. Roughly for $t_o < 250$, the frequency distribution of the received photons bears a resemblance to the single-peak case. At early stage, most of the received photons are emitted in the green region. Afterwards, photons that are emitted inward in the purple region come to dominate blueshifted photons received by the observers, and photons that are emitted outward in other colored regions suffer from a severe redshift. Remarkably, photons emitted inward in the blue region start reaching the celestial sphere around $t_o \simeq 240$ after they orbit the black hole several times between the inner and outer photon spheres. Subsequently, the observers receive photons emitted inward in the orange and brown regions, which would linger between the inner and outer photon spheres for a longer time. As indicated previously, the high-frequency and low-frequency photons come from photons emitted at large and small r_e , respectively. When r_e is large enough, the received photons are blueshifted.

The normalized total luminosity of the infalling star is displayed in the right panel of FIG. 9, where colored dots correspond to packets of 50 photons. Before $t_o \simeq 250$, the luminosity behaves similarly to the single-peak case, i.e., a gradual rise followed by an exponentially decaying tail



FIG. 9. The normalized frequency distribution (Left) and total luminosity (Right) of the infalling star measured by observers on a celestial sphere at $r_o = 100$ in the hairy black hole with a double-peak potential. Before $t_o \simeq 250$, the frequency and luminosity observations are quite similar to the single-peak case as the outer photon sphere plays the role of the photon sphere in the single-peak case. For $200 \lesssim t_o \lesssim 250$, the luminosity scales as a decay $\mathcal{L} \propto e^{-0.114t}$, in agreement with the Lyapunov exponent $\lambda_{\text{out}} = 0.11420$ at the outer photon sphere. Interestingly, the inward-emitted photons with $b_{\text{in}} < b < b_{\text{out}}$ (i.e., those emitted in the brown, blue and orange regions) usually linger for a long time between the inner and outer photon spheres before being received and kick in after $t_o \simeq 250$. For $250 \lesssim t_o \lesssim 300$, a large number of photons with a wide range of frequencies are detected by the observers, thereby leading to an increase of the observed luminosity and a luminosity peak at $t_o \simeq 300$. After $t_o \gtrsim 300$, less and less photons leak out the outer photon sphere, and the luminosity decays exponentially as $\mathcal{L} \propto e^{-0.058t}$, which may be associated with sub-long-lived quasinormal modes residing at the outer photon sphere.

as $\mathcal{L} \propto e^{-0.114t}$. The tail is determined by photons that are emitted in the purple region and linger outside the outer photon sphere. As expected, the Lyapunov exponent $\lambda_{\text{out}} = 0.11420$ at the outer photon sphere controls the decay rate of the tail. It should be emphasized that the Lyapunov exponent at the inner photon sphere is $\lambda_{\text{in}} = 0.14399$, and hence the contribution to the luminosity from photons temporarily trapped near the inner photon sphere is suppressed. After $t_o \simeq 250$, photons temporarily trapped between the inner and outer photon spheres start to play a dominant role. In particular, photons emitted in the blue region give rise to a noticeable increase of the luminosity and a sharp peak. Moreover, the late-time behavior is dominated by photons emitted in the orange region and can be described by an exponential decay, $\mathcal{L} \propto e^{-0.058t}$. In [121], we found that the photons emitted in the orange region are associated with a class of sub-longlived quasinormal modes, thereby leading to a more slowly decaying tail than $e^{-0.114t}$. However unlike the single-peak case, the imaginary parts of the sub-long-lived modes depend on the angular quantum number, and hence the decay rate of the tail can not be universally determined.



FIG. 10. The normalized frequency and luminosity of the infalling star measured by a far-away observer at $r_o = 100, \theta = \pi/2$ and $\phi = 0$ in the hairy black hole with a double-peak potential. Left: The frequency lines correspond to the orbit number $n = 0, 1, 2, \cdots$ from left to right, and the $n \ge 1$ frequency lines are separated by $\Delta T \simeq 2\pi b_{\rm in} \simeq 2\pi b_{\rm out} \simeq 22$. At early times, the $1 \le n \le 4$ frequency lines are primarily determined by photons lingering outside the outer photon sphere. After $t_o \simeq 250$, photons emitted from the star successively reach the observer after they orbit around the black hole between the inner and outer photon spheres, causing the n > 4 frequency lines. Right: Before the photons temporarily trapped between the photon spheres arrive, the observer sees a luminosity drop followed by two flashes, which is quite similar to the single-peak case. After $t_o \simeq 250$, the received n = 5 photons lead to a notable flash. Subsequently, the received n = 6 and 7 photons give two more but much less luminous flashes.

In the left panel of FIG. 10, we exhibit the normalized frequency of photons received by an observer located at $\varphi = 0$ and $\theta = \pi/2$ on the celestial sphere. After photons are emitted from the star, they would orbit around the black hole different times before being captured, which results in a discrete set of the observed frequencies indexed by the number of orbits n. The orbit number n of the frequency lines increases from left to right, and the leftmost yellow line corresponds to n = 0. As previously stated, the $n \ge 1$ frequency lines are similar and separated by the time delay $\Delta T \simeq 2\pi b_{\rm in} \simeq 2\pi b_{\rm out} \simeq 22$. For $1 \le n \le 3$, the frequency lines are mainly contributed by photons that are emitted in the purple region and circle around the outer photon sphere n times. When n = 4, photons emitted in the purple region are rarely seen, and some observed low-frequency photons are from emission between the photon spheres in the blue and orange regions. Inward-emitted photons in the blue, orange and brown regions can circle around the black hole 5 times between the inner and outer photon spheres and come back to start arriving after $t_o \simeq 70 \times 2 + 22 \times 5 \simeq 250$, which forms the n = 5 frequency line. Later, photons orbiting around the black hole more than 5 times successively reach the observer and give rise to the n > 5 frequency lines.

The normalized luminosity of the star measured by the observer is displayed in the right panel of FIG. 10. Before $t_o \simeq 250$, the luminosity observation resembles that in the single-peak case, i.e., two flashes following the decrease of the luminosity. Owing to the leaking of photons temporarily trapped between the photon spheres, a noticeable flash appears around $t_o \simeq 300$, and two much fainter flashes are seen later on. These three flashes are caused by the received photons orbiting around the black hole 5, 6 and 7 times, respectively. For n > 7, we do not have sufficient photons to compute the luminosity.

IV. CONCLUSIONS

In this paper, we investigated observational appearances of a point-like freely-falling star, which emits photons isotropically in its rest frame, in hairy black holes, which possess one or two photon spheres outside the event horizon. In particular, we considered the frequency and luminosity of received photons measured by all observers on a celestial sphere and a specific observer. Interestingly, it was found that the existence of an extra photon sphere can significantly change the late-time observations, which may provide a new tool to detect black holes with multiple photon spheres.

For hairy black holes with a single-peak potential, emitted photons can be temporarily trapped just outside the photon sphere and subsequently reemitted, resulting in several interesting observational signatures. First, photons with a near-critical impact parameter can be blueshifted since the photon sphere acts as a reflecting wall (see FIGs. 3 and 4). Second, at late times, the total luminosity measured by observers on the celestial sphere decreases exponentially with time, which is controlled by the Lyapunov exponent of circular null geodesics at the photon sphere (see FIG. 4). Last but not least, photons orbiting around the photon sphere different times produce a cascade of flashes observed by the specific observer. The luminosity of the flashes decreases sharply with the orbit number, and the frequency contents of the flashes are similar (see FIG. 5). It is worth emphasizing that the observational appearances in the hairy black holes with a single-peak potential are similar to those in Schwarzschild black holes reported in [92].

When an extra photon sphere appears outside the event horizon, we found that light rays can get trapped between the two photon spheres longer than just outside the photon spheres, which appreciably impacts late-time observations. In fact, due to arrivals of photons trapped between the photon spheres, the total luminosity rises to a peak, which is succeeded by a slow exponential decay (see FIG. 9). This slow decay may be related to sub-long-lived quasinormal modes living near the outer photon sphere, as found in [121]. The specific observer would see two cascades of flashes, which contain a similar frequency content. While the earlier cascade is mainly determined by photons orbiting outside the outer photon sphere, photons trapped between the photon spheres primarily give rise to the later one (see FIG. 10). In summary, black holes with multiple photon spheres will lead to distinctive optical appearances of an infalling star. In the future studies, it will be of great interest if our analysis can be generalized to more astrophysically realistic models.

ACKNOWLEDGMENTS

We are grateful to Qingyu Gan and Xin Jiang for useful discussions and valuable comments. This work is supported in part by NSFC (Grant No. 12105191, 11947225 and 11875196). Houwen Wu is supported by the International Visiting Program for Excellent Young Scholars of Sichuan University.

- Kazunori Akiyama et al. First M87 Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole. Astrophys. J. Lett., 875:L1, 2019. arXiv:1906.11238, doi:10.3847/2041-8213/ ab0ec7. I
- [2] Kazunori Akiyama et al. First M87 Event Horizon Telescope Results. II. Array and Instrumentation. Astrophys. J. Lett., 875(1):L2, 2019. arXiv:1906.11239, doi:10.3847/2041-8213/ab0c96.
- [3] Kazunori Akiyama et al. First M87 Event Horizon Telescope Results. III. Data Processing and Calibration. Astrophys. J. Lett., 875(1):L3, 2019. arXiv:1906.11240, doi:10.3847/2041-8213/ ab0c57.
- [4] Kazunori Akiyama et al. First M87 Event Horizon Telescope Results. IV. Imaging the Central Supermassive Black Hole. Astrophys. J. Lett., 875(1):L4, 2019. arXiv:1906.11241, doi: 10.3847/2041-8213/ab0e85.
- [5] Kazunori Akiyama et al. First M87 Event Horizon Telescope Results. V. Physical Origin of the Asymmetric Ring. Astrophys. J. Lett., 875(1):L5, 2019. arXiv:1906.11242, doi:10.3847/2041-8213/ab0f43.
- [6] Kazunori Akiyama et al. First M87 Event Horizon Telescope Results. VI. The Shadow and Mass of the Central Black Hole. Astrophys. J. Lett., 875(1):L6, 2019. arXiv:1906.11243, doi:10.3847/ 2041-8213/ab1141.
- [7] Kazunori Akiyama et al. First M87 Event Horizon Telescope Results. VII. Polarization of the Ring. Astrophys. J. Lett., 910(1):L12, 2021. arXiv:2105.01169, doi:10.3847/2041-8213/abe71d.
- [8] Kazunori Akiyama et al. First M87 Event Horizon Telescope Results. VIII. Magnetic Field Structure

near The Event Horizon. Astrophys. J. Lett., 910(1):L13, 2021. arXiv:2105.01173, doi:10.3847/2041-8213/abe4de. I

- [9] Kazunori Akiyama et al. First Sagittarius A* Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole in the Center of the Milky Way. Astrophys. J. Lett., 930(2):L12, 2022. doi:10.3847/2041-8213/ac6674. I
- [10] Kazunori Akiyama et al. First Sagittarius A* Event Horizon Telescope Results. II. EHT and Multiwavelength Observations, Data Processing, and Calibration. Astrophys. J. Lett., 930(2):L13, 2022. doi:10.3847/2041-8213/ac6675.
- [11] Kazunori Akiyama et al. First Sagittarius A* Event Horizon Telescope Results. III. Imaging of the Galactic Center Supermassive Black Hole. Astrophys. J. Lett., 930(2):L14, 2022. doi:10.3847/ 2041-8213/ac6429.
- [12] Kazunori Akiyama et al. First Sagittarius A* Event Horizon Telescope Results. IV. Variability, Morphology, and Black Hole Mass. Astrophys. J. Lett., 930(2):L15, 2022. doi:10.3847/2041-8213/ ac6736.
- [13] Kazunori Akiyama et al. First Sagittarius A* Event Horizon Telescope Results. V. Testing Astrophysical Models of the Galactic Center Black Hole. Astrophys. J. Lett., 930(2):L16, 2022. doi:10.3847/2041-8213/ac6672.
- [14] Kazunori Akiyama et al. First Sagittarius A* Event Horizon Telescope Results. VI. Testing the Black Hole Metric. Astrophys. J. Lett., 930(2):L17, 2022. doi:10.3847/2041-8213/ac6756. I
- [15] J. L. Synge. The Escape of Photons from Gravitationally Intense Stars. Mon. Not. Roy. Astron. Soc., 131(3):463-466, 1966. doi:10.1093/mnras/131.3.463. I
- [16] James M. Bardeen, William H. Press, and Saul A Teukolsky. Rotating black holes: Locally nonrotating frames, energy extraction, and scalar synchrotron radiation. Astrophys. J., 178:347, 1972. doi: 10.1086/151796.
- [17] J. M. Bardeen. Timelike and null geodesics in the Kerr metric. In Les Houches Summer School of Theoretical Physics: Black Holes, 1973.
- [18] K. S. Virbhadra and George F. R. Ellis. Schwarzschild black hole lensing. *Phys. Rev. D*, 62:084003, 2000. arXiv:astro-ph/9904193, doi:10.1103/PhysRevD.62.084003.
- [19] Clarissa-Marie Claudel, K. S. Virbhadra, and G. F. R. Ellis. The Geometry of photon surfaces. J. Math. Phys., 42:818-838, 2001. arXiv:gr-qc/0005050, doi:10.1063/1.1308507.
- [20] K. S. Virbhadra. Relativistic images of Schwarzschild black hole lensing. Phys. Rev. D, 79:083004, 2009. arXiv:0810.2109, doi:10.1103/PhysRevD.79.083004.
- [21] Valerio Bozza. Gravitational Lensing by Black Holes. Gen. Rel. Grav., 42:2269-2300, 2010. arXiv: 0911.2187, doi:10.1007/s10714-010-0988-2.
- [22] K. S. Virbhadra. Distortions of images of Schwarzschild lensing. 4 2022. arXiv:2204.01879. I
- [23] Farruh Atamurotov, Sushant G. Ghosh, and Bobomurat Ahmedov. Horizon structure of rotating Einstein-Born-Infeld black holes and shadow. *Eur. Phys. J. C*, 76(5):273, 2016. arXiv:1506.03690,

doi:10.1140/epjc/s10052-016-4122-9. I

- [24] Zdenek Stuchlik and Jan Schee. Shadow of the regular Bardeen black holes and comparison of the motion of photons and neutrinos. *Eur. Phys. J. C*, 79(1):44, 2019. doi:10.1140/epjc/ s10052-019-6543-8.
- [25] Tian-Chi Ma, He-Xu Zhang, Peng-Zhang He, Hao-Ran Zhang, Yuan Chen, and Jian-Bo Deng. Shadow cast by a rotating and nonlinear magnetic-charged black hole in perfect fluid dark matter. *Mod. Phys. Lett. A*, 36(17):2150112, 2021. arXiv:2010.00151, doi:10.1142/S0217732321501121.
- [26] Zezhou Hu, Zhen Zhong, Peng-Cheng Li, Minyong Guo, and Bin Chen. QED effect on a black hole shadow. Phys. Rev. D, 103(4):044057, 2021. arXiv:2012.07022, doi:10.1103/PhysRevD.103. 044057.
- [27] S. I. Kruglov. The shadow of M87* black hole within rational nonlinear electrodynamics. Mod. Phys. Lett. A, 35(35):2050291, 2020. arXiv:2009.07657, doi:10.1142/S0217732320502910.
- [28] Zhen Zhong, Zezhou Hu, Haopeng Yan, Minyong Guo, and Bin Chen. QED effects on Kerr black hole shadows immersed in uniform magnetic fields. *Phys. Rev. D*, 104(10):104028, 2021. arXiv: 2108.06140, doi:10.1103/PhysRevD.104.104028.
- [29] Aoyun He, Jun Tao, Peng Wang, Yadong Xue, and Lingkai Zhang. Effects of Born-Infeld electrodynamics on black hole shadows. 5 2022. arXiv:2205.12779. I
- [30] Liang Ma and H. Lu. Bounds on photon spheres and shadows of charged black holes in Einstein-Gauss-Bonnet-Maxwell gravity. *Phys. Lett. B*, 807:135535, 2020. arXiv:1912.05569, doi:10.1016/j.physletb.2020.135535. I
- [31] Shao-Wen Wei and Yu-Xiao Liu. Testing the nature of Gauss-Bonnet gravity by four-dimensional rotating black hole shadow. *Eur. Phys. J. Plus*, 136(4):436, 2021. arXiv:2003.07769, doi:10.1140/ epjp/s13360-021-01398-9.
- [32] Xiao-Xiong Zeng, Hai-Qing Zhang, and Hongbao Zhang. Shadows and photon spheres with spherical accretions in the four-dimensional Gauss-Bonnet black hole. *Eur. Phys. J. C*, 80(9):872, 2020. arXiv: 2004.12074, doi:10.1140/epjc/s10052-020-08449-y.
- [33] Minyong Guo and Peng-Cheng Li. Innermost stable circular orbit and shadow of the 4D Einstein-Gauss-Bonnet black hole. Eur. Phys. J. C, 80(6):588, 2020. arXiv:2003.02523, doi: 10.1140/epjc/s10052-020-8164-7. I
- [34] Fabio Bacchini, Daniel R. Mayerson, Bart Ripperda, Jordy Davelaar, Héctor Olivares, Thomas Hertog, and Bert Vercnocke. Fuzzball Shadows: Emergent Horizons from Microstructure. *Phys. Rev. Lett.*, 127(17):171601, 2021. arXiv:2103.12075, doi:10.1103/PhysRevLett.127.171601. I
- [35] Dimitry Ayzenberg and Nicolas Yunes. Black Hole Shadow as a Test of General Relativity: Quadratic Gravity. Class. Quant. Grav., 35(23):235002, 2018. arXiv:1807.08422, doi:10.1088/1361-6382/aae87b. I
- [36] Leonardo Amarilla, Ernesto F. Eiroa, and Gaston Giribet. Null geodesics and shadow of a rotating black hole in extended Chern-Simons modified gravity. *Phys. Rev. D*, 81:124045, 2010. arXiv:

1005.0607, doi:10.1103/PhysRevD.81.124045. I

- [37] Sara Dastan, Reza Saffari, and Saheb Soroushfar. Shadow of a Charged Rotating Black Hole in f(R)Gravity. 6 2016. arXiv:1606.06994. I
- [38] Andrea Addazi, Salvatore Capozziello, and Sergei Odintsov. Chaotic solutions and black hole shadow in f(R) gravity. *Phys. Lett. B*, 816:136257, 2021. arXiv:2103.16856, doi:10.1016/j.physletb. 2021.136257.
- [39] Guo-Ping Li and Ke-Jian He. Observational appearances of a f(R) global monopole black hole illuminated by various accretions. *Eur. Phys. J. C*, 81(11):1018, 2021. doi:10.1140/epjc/ s10052-021-09817-y. I
- [40] Leonardo Amarilla and Ernesto F. Eiroa. Shadow of a rotating braneworld black hole. *Phys. Rev. D*, 85:064019, 2012. arXiv:1112.6349, doi:10.1103/PhysRevD.85.064019. I
- [41] Minyong Guo, Shupeng Song, and Haopeng Yan. Observational signature of a near-extremal Kerr-Sen black hole in the heterotic string theory. *Phys. Rev. D*, 101(2):024055, 2020. arXiv:1911.04796, doi:10.1103/PhysRevD.101.024055.
- [42] Tao Zhu, Qiang Wu, Mubasher Jamil, and Kimet Jusufi. Shadows and deflection angle of charged and slowly rotating black holes in Einstein-Æther theory. *Phys. Rev. D*, 100(4):044055, 2019. arXiv: 1906.05673, doi:10.1103/PhysRevD.100.044055.
- [43] Rahul Kumar, Sushant G. Ghosh, and Anzhong Wang. Gravitational deflection of light and shadow cast by rotating Kalb-Ramond black holes. *Phys. Rev. D*, 101(10):104001, 2020. arXiv:2001.00460, doi:10.1103/PhysRevD.101.104001. I
- [44] Rajibul Shaikh, Pritam Banerjee, Suvankar Paul, and Tapobrata Sarkar. A novel gravitational lensing feature by wormholes. *Phys. Lett. B*, 789:270-275, 2019. [Erratum: Phys.Lett.B 791, 422-423 (2019)].
 arXiv:1811.08245, doi:10.1016/j.physletb.2018.12.030. I
- [45] Maciek Wielgus, Jiri Horak, Frederic Vincent, and Marek Abramowicz. Reflection-asymmetric wormholes and their double shadows. *Phys. Rev. D*, 102(8):084044, 2020. arXiv:2008.10130, doi:10.1103/PhysRevD.102.084044.
- [46] Merce Guerrero, Gonzalo J. Olmo, Diego Rubiera-Garcia, and Diego Sáez-Chillón Gómez. Shadows and optical appearance of black bounces illuminated by a thin accretion disk. JCAP, 08:036, 2021. arXiv:2105.15073, doi:10.1088/1475-7516/2021/08/036.
- [47] Jun Peng, Minyong Guo, and Xing-Hui Feng. Observational signature and additional photon rings of an asymmetric thin-shell wormhole. *Phys. Rev. D*, 104(12):124010, 2021. arXiv:2102.05488, doi:10.1103/PhysRevD.104.124010.
- [48] Cosimo Bambi and Dejan Stojkovic. Astrophysical Wormholes. Universe, 7(5):136, 2021. arXiv: 2105.00881, doi:10.3390/universe7050136.
- [49] Gonzalo J. Olmo, Diego Rubiera-Garcia, and Diego Sáez-Chillón Gómez. New light rings from multiple critical curves as observational signatures of black hole mimickers. *Phys. Lett. B*, 829:137045, 2022. arXiv:2110.10002, doi:10.1016/j.physletb.2022.137045.

- [50] Merce Guerrero, Gonzalo J. Olmo, Diego Rubiera-Garcia, and Diego Gómez Sáez-Chillón. Light ring images of double photon spheres in black hole and wormhole spacetimes. *Phys. Rev. D*, 105(8):084057, 2022. arXiv:2202.03809, doi:10.1103/PhysRevD.105.084057. I
- [51] Pedro V. P. Cunha, José A. Font, Carlos Herdeiro, Eugen Radu, Nicolas Sanchis-Gual, and Miguel Zilhão. Lensing and dynamics of ultracompact bosonic stars. *Phys. Rev. D*, 96(10):104040, 2017. arXiv:1709.06118, doi:10.1103/PhysRevD.96.104040. I
- [52] Hector Olivares, Ziri Younsi, Christian M. Fromm, Mariafelicia De Laurentis, Oliver Porth, Yosuke Mizuno, Heino Falcke, Michael Kramer, and Luciano Rezzolla. How to tell an accreting boson star from a black hole. Mon. Not. Roy. Astron. Soc., 497(1):521-535, 2020. arXiv:1809.08682, doi: 10.1093/mnras/staa1878. I
- [53] Shao-Wen Wei and Yu-Xiao Liu. Observing the shadow of Einstein-Maxwell-Dilaton-Axion black hole. JCAP, 11:063, 2013. arXiv:1311.4251, doi:10.1088/1475-7516/2013/11/063. I
- [54] Mingzhi Wang, Songbai Chen, and Jiliang Jing. Shadow casted by a Konoplya-Zhidenko rotating non-Kerr black hole. JCAP, 10:051, 2017. arXiv:1707.09451, doi:10.1088/1475-7516/2017/10/051.
- [55] Yosuke Mizuno, Ziri Younsi, Christian M. Fromm, Oliver Porth, Mariafelicia De Laurentis, Hector Olivares, Heino Falcke, Michael Kramer, and Luciano Rezzolla. The Current Ability to Test Theories of Gravity with Black Hole Shadows. *Nature Astron.*, 2(7):585–590, 2018. arXiv:1804.05812, doi: 10.1038/s41550-018-0449-5.
- [56] Ming Zhang and Jie Jiang. Shadows of accelerating black holes. Phys. Rev. D, 103(2):025005, 2021. arXiv:2010.12194, doi:10.1103/PhysRevD.103.025005.
- [57] Xiao-Xiong Zeng, Ke-Jian He, and Guo-Ping Li. Effect of dark matter on shadows and rings of a brane-world black hole illuminated by various accretions. 11 2021. arXiv:2111.05090.
- [58] Yang Guo and Yan-Gang Miao. Charged black-bounce spacetimes: Photon rings, shadows and observational appearances. 12 2021. arXiv:2112.01747.
- [59] Yuan Meng, Xiao-Mei Kuang, and Zi-Yu Tang. Photon regions, shadow observables and constraints from M87* of a charged rotating black hole. 4 2022. arXiv:2204.00897.
- [60] Chen-Kai Qiao and Ming Li. A Geometric Approach on Circular Photon Orbits and Black Hole Shadow. 4 2022. arXiv:2204.07297.
- [61] Igor Bogush, Dmitri Gal'tsov, Galin Gyulchev, Kirill Kobialko, Petya Nedkova, and Tsvetan Vetsov. Photon surfaces, shadows and accretion disks in gravity with minimally coupled scalar field. 5 2022. arXiv:2205.01919.
- [62] Sen Guo, Guan-Ru Li, and En-Wei Liang. Observable characteristics of the charged black hole surrounded by thin disk accretion in Rastall gravity. *Class. Quant. Grav.*, 39(13):135004, 2022. arXiv:2205.11241, doi:10.1088/1361-6382/ac6fa8.
- [63] Hui-Min Wang, Zi-Chao Lin, and Shao-Wen Wei. Optical appearance of Einstein-Æther black hole surrounded by thin disk. 5 2022. arXiv:2205.13174.
- [64] Zelin Zhang, Songbai Chen, and Jiliang Jing. Image of Bonnor black dihole with a thin accretion disk

and its polarization information. 5 2022. arXiv: 2205.13696.

- [65] Yehui Hou, Peng Liu, Minyong Guo, Haopeng Yan, and Bin Chen. Multi-level images around Kerr-Newman black holes. 3 2022. arXiv:2203.02755. I
- [66] Volker Perlick, Oleg Yu. Tsupko, and Gennady S. Bisnovatyi-Kogan. Black hole shadow in an expanding universe with a cosmological constant. *Phys. Rev. D*, 97(10):104062, 2018. arXiv:1804.04898, doi:10.1103/PhysRevD.97.104062. I
- [67] Oleg Yu. Tsupko, Zuhui Fan, and Gennady S. Bisnovatyi-Kogan. Black hole shadow as a standard ruler in cosmology. *Class. Quant. Grav.*, 37(6):065016, 2020. arXiv:1905.10509, doi:10.1088/ 1361-6382/ab6f7d.
- [68] Sunny Vagnozzi, Cosimo Bambi, and Luca Visinelli. Concerns regarding the use of black hole shadows as standard rulers. Class. Quant. Grav., 37(8):087001, 2020. arXiv:2001.02986, doi:10.1088/ 1361-6382/ab7965.
- [69] Peng-Cheng Li, Minyong Guo, and Bin Chen. Shadow of a Spinning Black Hole in an Expanding Universe. *Phys. Rev. D*, 101(8):084041, 2020. arXiv:2001.04231, doi:10.1103/PhysRevD.101. 084041.
- [70] Rittick Roy and Sayan Chakrabarti. Study on black hole shadows in asymptotically de Sitter spacetimes. Phys. Rev. D, 102(2):024059, 2020. arXiv:2003.14107, doi:10.1103/PhysRevD.102.024059.
- [71] Abhishek Chowdhuri and Arpan Bhattacharyya. Shadow analysis for rotating black holes in the presence of plasma for an expanding universe. 12 2020. arXiv:2012.12914. I
- [72] Muhammed Amir, Balendra Pratap Singh, and Sushant G. Ghosh. Shadows of rotating fivedimensional charged EMCS black holes. Eur. Phys. J. C, 78(5):399, 2018. arXiv:1707.09521, doi:10.1140/epjc/s10052-018-5872-3. I
- [73] Sunny Vagnozzi and Luca Visinelli. Hunting for extra dimensions in the shadow of M87*. *Phys. Rev.* D, 100(2):024020, 2019. arXiv:1905.12421, doi:10.1103/PhysRevD.100.024020.
- [74] Indrani Banerjee, Sumanta Chakraborty, and Soumitra SenGupta. Silhouette of M87*: A New Window to Peek into the World of Hidden Dimensions. *Phys. Rev. D*, 101(4):041301, 2020.
 arXiv:1909.09385, doi:10.1103/PhysRevD.101.041301. I
- [75] Sheng-Feng Yan, Chunlong Li, Lingqin Xue, Xin Ren, Yi-Fu Cai, Damien A. Easson, Ye-Fei Yuan, and Hongsheng Zhao. Testing the equivalence principle via the shadow of black holes. *Phys. Rev. Res.*, 2(2):023164, 2020. arXiv:1912.12629, doi:10.1103/PhysRevResearch.2.023164. I
- [76] Samuel E. Gralla. Can the EHT M87 results be used to test general relativity? Phys. Rev. D, 103(2):024023, 2021. arXiv:2010.08557, doi:10.1103/PhysRevD.103.024023.
- [77] Chunlong Li, Hongsheng Zhao, and Yi-Fu Cai. New test on the Einstein equivalence principle through the photon ring of black holes. *Phys. Rev. D*, 104(6):064027, 2021. arXiv:2102.10888, doi:10.1103/ PhysRevD.104.064027. I
- [78] Yifan Chen, Jing Shu, Xiao Xue, Qiang Yuan, and Yue Zhao. Probing Axions with Event Horizon Telescope Polarimetric Measurements. *Phys. Rev. Lett.*, 124(6):061102, 2020. arXiv:1905.02213,

doi:10.1103/PhysRevLett.124.061102. I

- [79] R. A. Konoplya. Shadow of a black hole surrounded by dark matter. *Phys. Lett. B*, 795:1-6, 2019. arXiv:1905.00064, doi:10.1016/j.physletb.2019.05.043.
- [80] Kimet Jusufi, Mubasher Jamil, Paolo Salucci, Tao Zhu, and Sumarna Haroon. Black hole surrounded by a dark matter halo in the m87 galactic center and its identification with shadow images. *Physical Review D*, 100(4), Aug 2019. URL: http://dx.doi.org/10.1103/PhysRevD.100.044012, doi:10. 1103/physrevd.100.044012.
- [81] Hooman Davoudiasl and Peter B. Denton. Ultralight boson dark matter and event horizon telescope observations of m87*. *Physical Review Letters*, 123(2), Jul 2019. URL: http://dx.doi.org/10.1103/ PhysRevLett.123.021102, doi:10.1103/physrevlett.123.021102.
- [82] Rittick Roy and Urjit A. Yajnik. Evolution of black hole shadow in the presence of ultralight bosons. *Phys. Lett. B*, 803:135284, 2020. arXiv:1906.03190, doi:10.1016/j.physletb.2020.135284.
- [83] K. Saurabh and Kimet Jusufi. Imprints of dark matter on black hole shadows using spherical accretions.
 Eur. Phys. J. C, 81(6):490, 2021. arXiv:2009.10599, doi:10.1140/epjc/s10052-021-09280-9. I
- [84] Ahmadjon Abdujabbarov, Bobir Toshmatov, Zdeněk Stuchlík, and Bobomurat Ahmedov. Shadow of the rotating black hole with quintessential energy in the presence of plasma. Int. J. Mod. Phys. D, 26(06):1750051, 2016. arXiv:1512.05206, doi:10.1142/S0218271817500511. I
- [85] Xiao-Xiong Zeng and Hai-Qing Zhang. Influence of quintessence dark energy on the shadow of black hole. Eur. Phys. J. C, 80(11):1058, 2020. arXiv:2007.06333, doi:10.1140/epjc/ s10052-020-08656-7.
- [86] Xin Qin, Songbai Chen, and Jiliang Jing. Image of a regular phantom compact object and its luminosity under spherical accretions. *Class. Quant. Grav.*, 38(11):115008, 2021. arXiv:2011.04310, doi:10.1088/1361-6382/abf712.
- [87] Yuan Chen, He-Xu Zhang, Tian-Chi Ma, and Jian-Bo Deng. Optical properties of a nonlinear magnetic charged rotating black hole surrounded by quintessence with a cosmological constant. 9 2020. arXiv: 2009.03778, doi:10.1142/S0217732321500760.
- [88] Aoyun He, Jun Tao, Yadong Xue, and Lingkai Zhang. Shadow and photon sphere of black hole in clouds of strings and quintessence *. Chin. Phys. C, 46(6):065102, 2022. arXiv:2109.13807, doi:10.1088/1674-1137/ac56cf. I
- [89] Joseph F. Dolan and Daria C Auerswald. A Search for Dying Pulse Trains in Cyg X-1 using RXTE. 4 2011. arXiv:1104.3164. I
- [90] M. Bauböck et al. Modeling the orbital motion of Sgr A*'s near-infrared flares. Astron. Astrophys., 635:A143, 2020. arXiv:2002.08374, doi:10.1051/0004-6361/201937233. I
- [91] R. Abuter et al. The flux distribution of Sgr A*. Astron. Astrophys., 638:A2, 2020. arXiv: 2004.07185, doi:10.1051/0004-6361/202037717. I
- [92] Vitor Cardoso, Francisco Duque, and Arianna Foschi. Light ring and the appearance of matter accreted by black holes. *Phys. Rev. D*, 103(10):104044, 2021. arXiv:2102.07784, doi:10.1103/PhysRevD.

103.104044. I, III A, IV

- [93] Carlos A.R. Herdeiro, Eugen Radu, Nicolas Sanchis-Gual, and José A. Font. Spontaneous Scalarization of Charged Black Holes. *Phys. Rev. Lett.*, 121(10):101102, 2018. arXiv:1806.05190, doi:10.1103/PhysRevLett.121.101102. I, II
- [94] R. A. Konoplya and A. Zhidenko. Analytical representation for metrics of scalarized Einstein-Maxwell black holes and their shadows. *Phys. Rev. D*, 100(4):044015, 2019. arXiv:1907.05551, doi:10.1103/ PhysRevD.100.044015.
- [95] Peng Wang, Houwen Wu, and Haitang Yang. Scalarized Einstein-Born-Infeld black holes. Phys. Rev. D, 103(10):104012, 2021. arXiv:2012.01066, doi:10.1103/PhysRevD.103.104012.
- [96] Guangzhou Guo, Peng Wang, Houwen Wu, and Haitang Yang. Scalarized Einstein-Maxwell-scalar black holes in anti-de Sitter spacetime. Eur. Phys. J. C, 81(10):864, 2021. arXiv:2102.04015, doi:10.1140/epjc/s10052-021-09614-7. I
- [97] Guangzhou Guo, Peng Wang, Houwen Wu, and Haitang Yang. Thermodynamics and phase structure of an Einstein-Maxwell-scalar model in extended phase space. *Phys. Rev. D*, 105(6):064069, 2022. arXiv:2107.04467, doi:10.1103/PhysRevD.105.064069. I
- [98] Pedro G. S. Fernandes, Carlos A. R. Herdeiro, Alexandre M. Pombo, Eugen Radu, and Nicolas Sanchis-Gual. Spontaneous Scalarisation of Charged Black Holes: Coupling Dependence and Dynamical Features. *Class. Quant. Grav.*, 36(13):134002, 2019. [Erratum: Class.Quant.Grav. 37, 049501 (2020)]. arXiv:1902.05079, doi:10.1088/1361-6382/ab23a1. I
- [99] Pedro G.S. Fernandes, Carlos A.R. Herdeiro, Alexandre M. Pombo, Eugen Radu, and Nicolas Sanchis-Gual. Charged black holes with axionic-type couplings: Classes of solutions and dynamical scalarization. Phys. Rev. D, 100(8):084045, 2019. arXiv:1908.00037, doi:10.1103/PhysRevD.100.084045.
- [100] Jose Luis Blázquez-Salcedo, Carlos A.R. Herdeiro, Jutta Kunz, Alexandre M. Pombo, and Eugen Radu. Einstein-Maxwell-scalar black holes: the hot, the cold and the bald. *Phys. Lett. B*, 806:135493, 2020. arXiv:2002.00963, doi:10.1016/j.physletb.2020.135493. I
- [101] De-Cheng Zou and Yun Soo Myung. Scalarized charged black holes with scalar mass term. *Phys. Rev.* D, 100(12):124055, 2019. arXiv:1909.11859, doi:10.1103/PhysRevD.100.124055. I
- [102] Pedro G.S. Fernandes. Einstein-Maxwell-scalar black holes with massive and self-interacting scalar hair. Phys. Dark Univ., 30:100716, 2020. arXiv:2003.01045, doi:10.1016/j.dark.2020.100716. I
- [103] Yan Peng. Scalarization of horizonless reflecting stars: neutral scalar fields non-minimally coupled to Maxwell fields. *Phys. Lett. B*, 804:135372, 2020. arXiv:1912.11989, doi:10.1016/j.physletb. 2020.135372. I
- [104] Yun Soo Myung and De-Cheng Zou. Instability of Reissner-Nordström black hole in Einstein-Maxwell-scalar theory. Eur. Phys. J. C, 79(3):273, 2019. arXiv:1808.02609, doi:10.1140/epjc/ s10052-019-6792-6. I
- [105] Yun Soo Myung and De-Cheng Zou. Stability of scalarized charged black holes in the Einstein-Maxwell-Scalar theory. Eur. Phys. J. C, 79(8):641, 2019. arXiv:1904.09864, doi:10.1140/

epjc/s10052-019-7176-7.

- [106] De-Cheng Zou and Yun Soo Myung. Radial perturbations of the scalarized black holes in Einstein-Maxwell-conformally coupled scalar theory. *Phys. Rev. D*, 102(6):064011, 2020. arXiv:2005.06677, doi:10.1103/PhysRevD.102.064011.
- [107] Yun Soo Myung and De-Cheng Zou. Onset of rotating scalarized black holes in Einstein-Chern-Simons-Scalar theory. *Phys. Lett. B*, 814:136081, 2021. arXiv:2012.02375, doi:10.1016/j.physletb.2021.136081.
- [108] Zhan-Feng Mai and Run-Qiu Yang. Stability analysis of a charged black hole with a nonlinear complex scalar field. *Phys. Rev. D*, 104(4):044008, 2021. arXiv:2101.00026, doi:10.1103/PhysRevD.104.044008. I
- [109] Dumitru Astefanesei, Carlos Herdeiro, João Oliveira, and Eugen Radu. Higher dimensional black hole scalarization. JHEP, 09:186, 2020. arXiv:2007.04153, doi:10.1007/JHEP09(2020)186. I
- [110] Yun Soo Myung and De-Cheng Zou. Quasinormal modes of scalarized black holes in the Einstein-Maxwell-Scalar theory. *Phys. Lett. B*, 790:400-407, 2019. arXiv:1812.03604, doi:10.1016/ j.physletb.2019.01.046. I
- [111] Jose Luis Blázquez-Salcedo, Carlos A.R. Herdeiro, Sarah Kahlen, Jutta Kunz, Alexandre M. Pombo, and Eugen Radu. Quasinormal modes of hot, cold and bald Einstein-Maxwell-scalar black holes. 8 2020. arXiv:2008.11744. I
- [112] Yun Soo Myung and De-Cheng Zou. Scalarized charged black holes in the Einstein-Maxwell-Scalar theory with two U(1) fields. *Phys. Lett. B*, 811:135905, 2020. arXiv:2009.05193, doi:10.1016/j. physletb.2020.135905. I
- [113] Yun Soo Myung and De-Cheng Zou. Scalarized black holes in the Einstein-Maxwell-scalar theory with a quasitopological term. Phys. Rev. D, 103(2):024010, 2021. arXiv:2011.09665, doi:10.1103/ PhysRevD.103.024010. I
- [114] Hong Guo, Xiao-Mei Kuang, Eleftherios Papantonopoulos, and Bin Wang. Topology and spacetime structure influences on black hole scalarization. 12 2020. arXiv:2012.11844. I
- [115] Yves Brihaye, Betti Hartmann, Nathália Pio Aprile, and Jon Urrestilla. Scalarization of asymptotically anti-de Sitter black holes with applications to holographic phase transitions. *Phys. Rev. D*, 101(12):124016, 2020. arXiv:1911.01950, doi:10.1103/PhysRevD.101.124016. I
- [116] Yves Brihaye, Carlos Herdeiro, and Eugen Radu. Black Hole Spontaneous Scalarisation with a Positive Cosmological Constant. *Phys. Lett. B*, 802:135269, 2020. arXiv:1910.05286, doi: 10.1016/j.physletb.2020.135269.
- [117] Cheng-Yong Zhang, Peng Liu, Yunqi Liu, Chao Niu, and Bin Wang. Dynamical charged black hole spontaneous scalarization in Anti-de Sitter spacetimes. 3 2021. arXiv:2103.13599. I
- [118] Qingyu Gan, Peng Wang, Houwen Wu, and Haitang Yang. Photon spheres and spherical accretion image of a hairy black hole. *Phys. Rev. D*, 104(2):024003, 2021. arXiv:2104.08703, doi:10.1103/ PhysRevD.104.024003. I, II

- [119] Qingyu Gan, Peng Wang, Houwen Wu, and Haitang Yang. Photon ring and observational appearance of a hairy black hole. *Phys. Rev. D*, 104(4):044049, 2021. arXiv:2105.11770, doi: 10.1103/PhysRevD.104.044049. I, II
- [120] Guangzhou Guo, Xin Jiang, Peng Wang, and Houwen Wu. Gravitational Lensing by Black Holes with Multiple Photon Spheres. 4 2022. arXiv:2204.13948. I
- [121] Guangzhou Guo, Peng Wang, Houwen Wu, and Haitang Yang. Quasinormal modes of black holes with multiple photon spheres. JHEP, 06:060, 2022. arXiv:2112.14133, doi:10.1007/JHEP06(2022)060.
 I, III A, III B, IV
- [122] Guangzhou Guo, Peng Wang, Houwen Wu, and Haitang Yang. Echoes from hairy black holes. JHEP, 06:073, 2022. arXiv:2204.00982, doi:10.1007/JHEP06(2022)073. I
- [123] Hai-Shan Liu, Zhan-Feng Mai, Yue-Zhou Li, and H. Lü. Quasi-topological Electromagnetism: Dark Energy, Dyonic Black Holes, Stable Photon Spheres and Hidden Electromagnetic Duality. Sci. China Phys. Mech. Astron., 63:240411, 2020. arXiv:1907.10876, doi:10.1007/s11433-019-1446-1. I
- [124] Claudia de Rham, Gregory Gabadadze, and Andrew J. Tolley. Resummation of Massive Gravity. Phys. Rev. Lett., 106:231101, 2011. arXiv:1011.1232, doi:10.1103/PhysRevLett.106.231101. I
- [125] Ruifeng Dong and Dejan Stojkovic. Gravitational wave echoes from black holes in massive gravity. *Phys. Rev. D*, 103(2):024058, 2021. arXiv:2011.04032, doi:10.1103/PhysRevD.103.024058. I
- [126] Naoki Tsukamoto. Gravitational lensing by two photon spheres in a black-bounce spacetime in strong deflection limits. 5 2021. arXiv:2105.14336. I
- [127] Naoki Tsukamoto. Linearization stability of reflection-asymmetric thin-shell wormholes with double shadows. *Phys. Rev. D*, 103(6):064031, 2021. arXiv:2101.07060, doi:10.1103/PhysRevD.103.064031.
- [128] Naoki Tsukamoto. Retrolensing by two photon spheres of a black-bounce spacetime. *Phys. Rev. D*, 105(8):084036, 2022. arXiv:2202.09641, doi:10.1103/PhysRevD.105.084036. I