

The cosmological vector modes from a monochromatic primordial power spectrum

Zhe Chang^{a,b} Xukun Zhang,^{a,b} Jing-Zhi Zhou,^{a,b,1}

^aInstitute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China

^bUniversity of Chinese Academy of Sciences, Beijing 100049, China

E-mail: changz@ihep.ac.cn, zhangxukun@ihep.ac.cn, zhoujingzhi@ihep.ac.cn

Abstract. The cosmological background of higher order vector modes can be generated by the first order scalar perturbations. We investigate the second order and the third order vector modes systematically. The explicit expressions of two point functions $\langle V^{(n),\lambda} V^{(n),\lambda'} \rangle$ ($n = 2, 3$) and power spectra corresponded are presented. In the case of a monochromatic primordial power spectrum, the second order vector modes do not exist. However, the third order vector modes can be generated by a monochromatic primordial power spectrum. And it is found that the third order vector modes sourced by the second order scalar perturbations dominate the two point function $\langle V^{(3),\lambda} V^{(3),\lambda'} \rangle$ and power spectrum corresponded.

¹Corresponding author.

Contents

1	Introduction	1
2	Second order vector perturbations	2
2.1	Second order kernel function	2
2.2	Second order power spectrum	4
3	Third order vector perturbations	5
3.1	Third order kernel function	5
3.2	Third order power spectrum	9
4	Conclusions and discussions	13
A	Decomposition Operators	13
B	Higher order perturbations of Einstein equation	14
B.1	Second order perturbation equation	14
B.2	Third order perturbation equation	15
C	The correlation function	16
C.1	Four-point correlation function	16
C.2	Six-point correlation function	17
D	The power spectra of vector modes	17
D.1	The power spectrum of second order vector mode	17
D.2	The power spectrum of third order vector mode	18
E	The expressions of $\mathbb{P}^{ij}(\eta, \mathbf{k})$	19

1 Introduction

In the inflation theory, the cosmological perturbations are originated from the quantum fluctuations during inflation. The information about the early Universe is encoded in these perturbations. The cosmological perturbations can be decomposed as scalar, vector, and tensor perturbations based on symmetry of Friedmann-Robertson-Walker (FRW) spacetime. According to the observations of the cosmic microwave background (CMB) and large-scale structure, the primordial power spectra are well constricted on large scales ($\gtrsim 1$ Mpc) [1, 2]. It indicates a nearly scale-invariant primordial power spectrum of scalar perturbations with amplitude $\sim 2 \times 10^{-9}$, and tensor perturbations with tensor-to-scalar ratio $r < 0.06$. However, contrary to the scalar modes and the tensor modes, the vector modes decay as $1/a^2$ after they leave the Hubble horizon during inflation, and the scalar perturbations can not generate vector modes at linear order [3]. Therefore, the vector perturbations are always neglected at first order.

On small scales ($\lesssim 1$ Mpc), the constraints of first order primordial scalar perturbations are much weaker than the constraints on large scales [4]. The amplitude of the small-scale primordial scalar perturbations can be large enough to generate the primordial black holes,

and it has close relations with the scalar induced gravitational waves [5–19]. Similarly, the cosmological background of higher order vector modes can be generated by the first order scalar perturbations [20–25]. These higher order vector modes will affect many cosmological observations, such as CMB polarization [21, 26], redshift-space distortions [27], and weak lensing [28–30].

In the study of second order scalar induced gravitational waves, it is convenient to consider a monochromatic primordial power spectrum $P_\phi = Ak_*\delta(k - k_*)$. However, it is not the case for the second order vector modes. As mentioned in Ref. [20], the second order vector modes can not be generated by a monochromatic primordial power spectrum. More precisely, if we calculate the power spectrum of second order vector modes $\mathcal{P}_V^{(2)}$ in terms of a monochromatic primordial power spectrum, we will obtain $\mathcal{P}_V^{(2)} = 0$. This property of second order vector modes is very different from the second order scalar perturbations and the second order gravitational waves.

In this paper, we investigate the second order and the third order vector modes systematically. We conclude that the vector modes can be generated by a monochromatic primordial power spectrum at third order. In this case, the first order and the second order vector modes do not exist. The third order vector mode is the first non-trivial order. The explicit expressions of two point functions $\langle V^{(n),\lambda}V^{(n),\lambda'} \rangle (n = 2, 3)$ and power spectra corresponded are presented. It shows that the third order vector modes sourced by the second order scalar perturbations dominate the two point functions $\langle V^{(3),\lambda}V^{(3),\lambda'} \rangle$ and power spectrum corresponded of third order vector modes.

This paper is organized as follows. In Sec. 2, we study the kernel function and the power spectrum of the second order vector modes. We show that the second order vector modes can not be generated by a monochromatic primordial power spectrum. In Sec. 3, we investigate the kernel functions and the power spectrum of the third order vector modes. The explicit expressions of the power spectrum of the third order vector modes are presented. The conclusions and discussions are summarized in Sec. 4.

2 Second order vector perturbations

In flat Friedmann-Robertson-Walker (FRW) spacetime, the background metric is given by

$$g_{\mu\nu}^{(0)}dx^\mu dx^\nu = a^2(\eta) (-d\eta^2 + \delta_{ij}dx^i dx^j) , \quad (2.1)$$

where η is the conformal time. The second order metric perturbations in Newtonian gauge is

$$ds^2 = -a^2 \left[(1 + 2\phi^{(1)}) d\eta^2 + V_i^{(2)} d\eta dx^i + (1 - 2\psi^{(1)}) \delta_{ij} dx^i dx^j \right] , \quad (2.2)$$

where $\phi^{(1)}$ and $\psi^{(1)}$ are the first order scalar perturbations, $V_i^{(2)}$ is the second order vector perturbation. In this section, we consider second order vector mode $V_i^{(2)}$ generated by the first order scalar perturbations. So that we neglect the first order vector perturbation $V_i^{(1)}$, the second order scalar perturbations $\phi^{(2)}$ and $\psi^{(2)}$, and the tensor perturbations $h_{ij}^{(n)}$.

2.1 Second order kernel function

We consider second order vector mode $V_i^{(2)}$ generated by the first order scalar perturbations. The equation of motion of second order vector perturbation is

$$V_l^{(2)'}(\eta, \mathbf{x}) + 2\mathcal{H}V_l^{(2)}(\eta, \mathbf{x}) = -4\Delta^{-1}\mathcal{T}_l^r \partial^s \mathcal{S}_{rs}^{(2)}(\eta, \mathbf{x}) , \quad (2.3)$$

where the source term is given by

$$\begin{aligned} \Delta^{-1}\mathcal{T}_l^r\partial^s\mathcal{S}_{rs}^{(2)}(\eta, \mathbf{x}) = & \Delta^{-1}\mathcal{T}_l^r\partial^s\left(\partial_r\phi^{(1)}\partial_s\phi^{(1)} - \frac{1}{\mathcal{H}}\left(\partial_r\phi^{(1)'}\partial_s\phi^{(1)} + \partial_r\phi^{(1)}\partial_s\phi^{(1)'}\right)\right. \\ & \left.+ 4\phi^{(1)}\partial_r\partial_s\phi^{(1)} - \frac{1}{\mathcal{H}^2}\partial_r\phi^{(1)'}\partial_s\phi^{(1)'}\right). \end{aligned} \quad (2.4)$$

The details of Eq. (2.3) are given in Appendix B.1. After making use of the Fourier transformation, we obtain the equation of motion of second order vector perturbation in momentum space

$$V^{\lambda,(2)'}(\eta, \mathbf{k}) + 2\mathcal{H}V^{\lambda,(2)}(\eta, \mathbf{k}) = 4\mathcal{S}_V^{\lambda,(2)}(\eta, \mathbf{k}), \quad (2.5)$$

where $V^{\lambda,(2)}(\eta, \mathbf{k}) = e^{\lambda,l}(\mathbf{k})V_l^{(2)}(\eta, \mathbf{k})$, $e^{\lambda,l}(\mathbf{k})$ is the polarization vectors with respect to \mathbf{k} , which satisfies $\sum_\lambda e^{\lambda,m}(\mathbf{k})e^{\lambda,r}(\mathbf{k}) + \frac{k^m k^r}{k^2} = \delta^{mr}$, $e^{\lambda,l}(\mathbf{k})e_l^{\lambda'}(\mathbf{k}) = \delta^{\lambda\lambda'}$, and $e^{\lambda,l}(\mathbf{k})k_l = 0$. The source term can be written as

$$\begin{aligned} \mathcal{S}_V^{\lambda,(2)}(\eta, \mathbf{k}) = & i\frac{p^s e^{\lambda,r}(\mathbf{k})}{k^2}\mathcal{S}_{rs}(\eta, \mathbf{k}) \\ = & \int \frac{d^3p}{(2\pi)^{3/2}} i\frac{k^s e^{\lambda,r}(\mathbf{k})}{k^2} p_r p_s \left(3\phi^{(1)}(\mathbf{k}-\mathbf{p})\phi^{(1)}(\mathbf{p}) + \frac{2}{\mathcal{H}}\phi^{(1)'}(\mathbf{k}-\mathbf{p})\phi^{(1)}(\mathbf{p}) \right. \\ & \left. + \frac{1}{\mathcal{H}^2}\phi^{(1)'}(\mathbf{k}-\mathbf{p})\phi^{(1)'}(\mathbf{p}) \right). \end{aligned} \quad (2.6)$$

We rewrite the first order scalar perturbations in the form of

$$\psi(\eta, \mathbf{p}) = \phi(\eta, \mathbf{p}) = \Phi_{\mathbf{p}}T_\phi(p\eta), \quad (2.7)$$

where $\Phi_{\mathbf{p}}$ is initial value originated from primordial curvature perturbation, and the transfer function $T_\phi(y) = \frac{9}{y^2}\left(\frac{\sqrt{3}}{y}\sin\left(\frac{y}{\sqrt{3}}\right) - \cos\left(\frac{y}{\sqrt{3}}\right)\right)$. Therefore, the formal expression of the second order vector mode can be written as

$$V^{\lambda,(2)}(\eta, \mathbf{k}) = \int \frac{d^3p}{(2\pi)^{3/2}} i\frac{k^s e^{\lambda,r}(\mathbf{k})}{k^2} p_r p_s I_V^{(2)}(|k-p|, p, \eta) \Phi_{\mathbf{k}-\mathbf{p}}\Phi_{\mathbf{p}}. \quad (2.8)$$

We define $k \equiv |\mathbf{k}|$, $p \equiv |\mathbf{p}| = vk$, $|\mathbf{k}-\mathbf{p}| = uk$, and $x = k\eta$. Then, the corresponding second order kernel function is

$$I_V^{(2)}(u, v, x) = \frac{4}{k} \int_0^x f_V^{(2)}(u, v, \bar{x}) \frac{\bar{x}^2}{x^2} d\bar{x}, \quad (2.9)$$

where

$$f_V(u, v, x) = 3T_\phi(ux)T_\phi(vx) + 2ux\frac{d}{d(ux)}T_\phi(ux)T_\phi(vx) + uvx^2\frac{d}{d(ux)}T_\phi(ux)\frac{d}{d(vx)}T_\phi(vx). \quad (2.10)$$

We present the second order kernel function $I_V^{(2)}(u=1, v=1, x)$ as a function of $x = k\eta$ in Fig. 1.




Figure 1. The square of the second order vector kernel function. Here we have set $u = v = 1$.

2.2 Second order power spectrum

The power spectrum of second order vector mode $\mathcal{P}_V(\eta, k)$ is defined as

$$\delta_{\lambda\lambda'} \langle V^{\lambda,(2)}(\eta, \mathbf{k}) V^{\lambda',(2)}(\eta, \mathbf{k}') \rangle = 2\delta(\mathbf{k} + \mathbf{k}') \frac{2\pi^2}{k^3} \mathcal{P}_V^{(2)}(\eta, k). \quad (2.11)$$

Substituting Eq. (2.8) into Eq. (2.11), we obtain

$$\begin{aligned} 2\delta(\mathbf{k} + \mathbf{k}') \frac{2\pi^2}{k^3} \mathcal{P}_V^{(2)}(\eta, k) &= -\delta_{\lambda\lambda'} \int \frac{d^3p d^3p'}{(2\pi)^3} \frac{k^s e^{\lambda,r}(\mathbf{k})}{k^2} \frac{k'^l e^{\lambda',m}(\mathbf{k}')}{k'^2} p_r p_s p'_l p'_m \\ &\times I_V^{(2)}(|k-p|, p, \eta) I_V^{(2)}(|k'-p'|, p', \eta) \langle \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{p}} \Phi_{\mathbf{k}'-\mathbf{p}'} \Phi_{\mathbf{p}'} \rangle, \end{aligned} \quad (2.12)$$

where the minus sign comes from the $(i)^2$. The integral and measure $\int \frac{d^3p d^3p'}{(2\pi)^3}$ can be simplified as the integrals of u and v with the explicit upper and lower limits in terms of a given coordinate system. And it is convenient to set $\mathbf{k}/|\mathbf{k}| = (0, 0, 1)$ in the calculations of integral and measure. The polynomial of momentums $\frac{k^s e^{\lambda,r}(\mathbf{k})}{k^2} \frac{k'^l e^{\lambda',m}(\mathbf{k}')}{k'^2} p_r p_s p'_l p'_m$ comes from the decomposition operator of vector mode $\Delta^{-1} \mathcal{T}_l^r \partial^s$ in momentum space and the polarization vectors $e^{\lambda,l}(\mathbf{k})$. It can be simplified as a polynomial of u , v , and k . The product of second order kernel functions $I_V^{(2)}(|\mathbf{k}-\mathbf{p}|, p, \eta) I_V^{(2)}(|\mathbf{k}'-\mathbf{p}'|, p', \eta)$ have been studied in Sec. 2.1. The four point function $\langle \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{p}} \Phi_{\mathbf{k}'-\mathbf{p}'} \Phi_{\mathbf{p}'} \rangle$ can be simplified in terms of Wick's theorem in Appendix C.1. It is noticed that the left hand side of Eq. (2.12) and the last line of Eq. (C.1) both contain the three dimensional delta function $\delta(\mathbf{k} + \mathbf{k}')$. We can integral over \mathbf{k}' to obtain the expression of $\mathcal{P}_V^{(2)}(\eta, k)$. The Wick's expansion in Eq. (C.1) also contains the summation of two delta functions ($\delta(\mathbf{p} + \mathbf{p}') + \delta(\mathbf{k} - \mathbf{p} + \mathbf{p}')$). We can integral over \mathbf{p}' to obtain the substitutions $\mathbf{p}' \rightarrow -\mathbf{p}$ and $\mathbf{p}' \rightarrow \mathbf{p} - \mathbf{k}$. The explicit expression of the second order power spectrum $\mathcal{P}_V^{(2)}(\eta, k)$ is

$$\mathcal{P}_V^{(2)} = \frac{1}{4} \int_0^\infty dv \int_{|1-v|}^{1+v} du I_V^{(2)}(u, v, x) \sum_{a=1}^2 \mathbb{P}_a(u, v) I_{V,a}^{(2)}(u, v, x) k^2 \mathcal{P}_\Phi(ku) \mathcal{P}_\Phi(kv), \quad (2.13)$$

where

$$\begin{aligned}\mathbb{P}_1(u, v) &= \frac{(1 + v^2 - u^2) \left(4v^2 - (1 + v^2 - u^2)^2\right)}{16u^2v^2}, \\ \mathbb{P}_2(u, v) &= \frac{\left((1 + v^2 - u^2)^2 - 2(1 + v^2 - u^2)\right) \left(4v^2 - (1 + v^2 - u^2)^2\right)}{16v^2u^2}.\end{aligned}\tag{2.14}$$

The summation of index a in Eq. (2.13) comes from the two terms in Wick's expansions of four point function. The calculation details are given in Appendix D.1. One can use Eq. (2.13) to calculate the power spectrum of second order vector mode for a given primordial power spectrum $\mathcal{P}_\Phi(k)$.

Here, we consider a monochromatic primordial power spectrum $\mathcal{P}_\Phi(k) = Ak_*\delta(k - k_*)$. Then Eq. (2.13) can be simplified as

$$\mathcal{P}_V^{(2)} = \frac{A^2}{2\tilde{k}^2} \int_0^\infty dv \int_{|1-v|}^{1+v} du \left(kI_V^{(2)}(u, v, x)\right)^2 \left(\delta\left(u - \frac{1}{\tilde{k}}\right)\delta\left(v - \frac{1}{\tilde{k}}\right)\right) \sum_{a=1}^2 \mathbb{P}_a(u, v).\tag{2.15}$$

Integrating over u and v , we obtain the substitutions $u = v = \frac{k_*}{k} = \frac{1}{\tilde{k}}$. It is not difficult to find that the summation of the polynomials is equal to zero, namely $\sum_{a=1}^2 \mathbb{P}_a\left(\frac{1}{\tilde{k}}, \frac{1}{\tilde{k}}\right) = 0$. So that we conclude that the second order vector mode $V^{\lambda,(2)}$ can not be generated by a monochromatic primordial power spectrum [20].

3 Third order vector perturbations

In this section, we consider the third order vector perturbations. The third order metric perturbations in Newtonian gauge take the form of

$$\begin{aligned}ds^2 &= -a^2 \left[\left(1 + 2\phi^{(1)} + \phi^{(2)}\right) d\eta^2 + \left(V_i^{(2)} + \frac{1}{3}V_i^{(3)}\right) d\eta dx^i \right. \\ &\quad \left. + \left(\left(1 - 2\psi^{(1)} - \psi^{(2)}\right) \delta_{ij} + \frac{1}{2}h_{ij}^{(2)}\right) dx^i dx^j \right],\end{aligned}\tag{3.1}$$

where $\phi^{(n)}$ and $\psi^{(n)}$ ($n = 1, 2$) are the n -order scalar perturbations, $V_i^{(n)}$ ($n = 2, 3$) are the n -order vector perturbation, and $h_{ij}^{(2)}$ is the second order tensor perturbations.

3.1 Third order kernel function

We study the equation of motion and the kernel functions of the third order vector mode $V_i^{(3)}$ in this section. The equation of motion of the third order vector mode is rewritten as follows,

$$V_i^{(3)'}(\eta, \mathbf{x}) + 2\mathcal{H}V_i^{(3)}(\eta, \mathbf{x}) = -12\Delta^{-1}\mathcal{T}_i^l\partial^m S_{lm}^{(3)}(\eta, \mathbf{x}),\tag{3.2}$$

where the expression of traceless operator is $\mathcal{T}_i^l = \delta_i^l - \partial^l\Delta^{-1}\partial_i$. The details of the decomposed operators are shown in Appendix A. The explicit details of Eq. (3.2) are given in Appendix B.2. The source term $S_{lm}^{(3)}(\eta, \mathbf{x})$ can be divided into four parts,

$$S_{lm}^{(3)}(\eta, \mathbf{x}) = S_{lm,1}^{(3)}(\eta, \mathbf{x}) + S_{lm,2}^{(3)}(\eta, \mathbf{x}) + S_{lm,3}^{(3)}(\eta, \mathbf{x}) + S_{lm,4}^{(3)}(\eta, \mathbf{x}),\tag{3.3}$$

where $S_{lm,1}^{(3)}(\eta, \mathbf{x})$ is composed of the first order scalar perturbation $\phi^{(1)}$,

$$\begin{aligned}
S_{lm,1}^{(3)}(\eta, \mathbf{x}) &= 12\phi^{(1)}\partial_l\phi^{(1)}\partial_m\phi^{(1)} - \frac{4}{\mathcal{H}}\phi^{(1)'}\partial_l\phi^{(1)}\partial_m\phi^{(1)} + \frac{2}{3\mathcal{H}^2}\Delta\phi^{(1)}\partial_l\phi^{(1)}\partial_m\phi^{(1)} \\
&+ \frac{2}{3\mathcal{H}^4}\Delta\phi^{(1)}\partial_l\phi^{(1)'}\partial_m\phi^{(1)'} - \frac{3}{\mathcal{H}^2}\phi^{(1)'}\partial_l\phi^{(1)'}\partial_m\phi^{(1)} - \frac{3}{\mathcal{H}^2}\phi^{(1)'}\partial_m\phi^{(1)'}\partial_l\phi^{(1)} \\
&+ \frac{2}{3\mathcal{H}^3}\Delta\phi^{(1)}\partial_l\phi^{(1)'}\partial_m\phi^{(1)} + \frac{2}{3\mathcal{H}^3}\Delta\phi^{(1)}\partial_m\phi^{(1)'}\partial_l\phi^{(1)} \\
&- \frac{2}{\mathcal{H}^3}\phi^{(1)'}\partial_l\phi^{(1)'}\partial_m\phi^{(1)'} - \frac{4}{\mathcal{H}^2}\phi^{(1)}\partial_l\phi^{(1)'}\partial_m\phi^{(1)'} .
\end{aligned} \tag{3.4}$$

The source term $S_{lm,2}^{(3)}(\eta, \mathbf{x})$ is composed of the first order scalar perturbation $\phi^{(1)}$ and the second order tensor perturbation $h_{lm}^{(2)}$,

$$\begin{aligned}
S_{lm,2}^{(3)}(\eta, \mathbf{x}) &= -\frac{1}{2}\phi^{(1)}\left(h_{lm}^{(2)''} + 2\mathcal{H}h_{lm}^{(2)'} - \Delta h_{lm}^{(2)}\right) - \phi^{(1)}\Delta h_{lm}^{(2)} - \phi^{(1)'}\mathcal{H}h_{lm}^{(2)} - \frac{1}{3}\Delta\phi^{(1)}h_{lm}^{(2)} \\
&- \partial^b\phi^{(1)}\partial_b h_{lm}^{(2)} .
\end{aligned} \tag{3.5}$$

The source term $S_{lm,3}^{(3)}(\eta, \mathbf{x})$ is composed of the first order scalar perturbation $\phi^{(1)}$ and the second order vector perturbation $V_l^{(2)}$,

$$\begin{aligned}
S_{lm,3}^{(3)}(\eta, \mathbf{x}) &= \phi^{(1)}\partial_l\left(V_m^{(2)'} + 2\mathcal{H}V_m^{(2)}\right) + \phi^{(1)}\partial_m\left(V_l^{(2)'} + 2\mathcal{H}V_l^{(2)}\right) + \phi^{(1)'}\left(\partial_l V_m^{(2)} + \partial_m V_l^{(2)}\right) \\
&- \frac{\phi^{(1)}}{8\mathcal{H}}\left(\partial_m\Delta V_l^{(2)} + \partial_l\Delta V_m^{(2)}\right) - \frac{\phi^{(1)'}}{8\mathcal{H}^2}\left(\partial_m\Delta V_l^{(2)} + \partial_l\Delta V_m^{(2)}\right) .
\end{aligned} \tag{3.6}$$

And the source term $S_{lm,4}^{(3)}(\eta, \mathbf{x})$ is composed of the first order scalar perturbation $\phi^{(1)}$ and the second order scalar perturbations $\phi^{(2)}$ and $\psi^{(2)}$,

$$\begin{aligned}
S_{lm,4}^{(3)}(\eta, \mathbf{x}) &= \frac{1}{\mathcal{H}}\left(\phi^{(1)}\partial_l\partial_m\psi^{(2)'}\right) + \frac{1}{\mathcal{H}}\left(\phi^{(1)'}\partial_l\partial_m\phi^{(2)}\right) + \frac{1}{\mathcal{H}^2}\left(\phi^{(1)'}\partial_l\partial_m\psi^{(2)'}\right) \\
&+ 3\left(\phi^{(1)}\partial_l\partial_m\phi^{(2)}\right) .
\end{aligned} \tag{3.7}$$

The second order perturbations $h_{lm}^{(2)}$, $V_l^{(2)}$ and $\phi^{(2)}$ and $\psi^{(2)}$ are induced by the first order scalar perturbations $\psi^{(1)}(=\phi^{(1)})$. At the next iteration, the first order scalar, the second order scalar, vector and tensor perturbations all induce the third order vector perturbation. The kernel functions of the second order perturbations have been studied in previous work [5]. It is similar to the calculations of the second order vector mode. In order to solve the equations of motion of third order vector mode, we rewrite Eq. (3.2) in momentum space as

$$V^{\lambda,(3)'}(\eta, \mathbf{p}) + 2\mathcal{H}V^{\lambda,(3)}(\eta, \mathbf{p}) = 12\sum_{j=1}^4 \mathcal{S}_j^{\lambda,(3)}(\eta, \mathbf{k}) , \tag{3.8}$$

where $V^{\lambda,(3)}(\eta, \mathbf{k}) = e^{\lambda,l}(\mathbf{k})V_l^{(3)}(\eta, \mathbf{k})$, $e^{\lambda,l}(\mathbf{k})$ is the polarization vectors with respect to \mathbf{k} , which satisfies $\sum_{\lambda} e^{\lambda,m}(\mathbf{k})e^{\lambda,r}(\mathbf{k}) + \frac{k^m k^r}{k^2} = \delta^{mr}$, $e^{\lambda,l}(\mathbf{k})e_l^{\lambda'}(\mathbf{k}) = \delta^{\lambda\lambda'}$, and $e^{\lambda,l}(\mathbf{k})k_l = 0$. Unlike the second order tensor perturbation, the third order vector mode have four types of source terms $\mathcal{S}_i^{\lambda,(3)}(\eta, \mathbf{k})$, ($i = 1, 2, 3, 4$) (Eqs. (3.4)–(3.7)). We rewrite the source terms

$S_i^{\lambda,(3)}(\eta, \mathbf{k})$ in terms of initial value of the first order scalar perturbations $\Phi_{\mathbf{p}}$ in momentum space, i.e.,

$$S_1^{\lambda,(3)}(\eta, \mathbf{k}) = \int \frac{d^3p}{(2\pi)^{3/2}} \int \frac{d^3q}{(2\pi)^{3/2}} i \frac{k^l e^{\lambda,m}(\mathbf{k})}{k^2} (p_l - q_l) q_m \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{p}-\mathbf{q}} \Phi_{\mathbf{q}} \\ \times f_1^{(3)}(|\mathbf{k}-\mathbf{p}|, |\mathbf{p}-\mathbf{q}|, \mathbf{q}, \eta) , \quad (3.9)$$

$$S_2^{\lambda,(3)}(\eta, \mathbf{k}) = \int \frac{d^3p}{(2\pi)^{3/2}} \int \frac{d^3q}{(2\pi)^{3/2}} i \frac{k^l e^{\lambda,m}(\mathbf{k})}{k^2} \Lambda_{lm}^{rs}(\mathbf{p}) q_r q_s \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{p}-\mathbf{q}} \Phi_{\mathbf{q}} \\ \times f_2^{(3)}(|\mathbf{k}-\mathbf{p}|, |\mathbf{p}-\mathbf{q}|, \mathbf{q}, \eta) , \quad (3.10)$$

$$S_3^{\lambda,(3)}(\eta, \mathbf{k}) = \int \frac{d^3p}{(2\pi)^{3/2}} \int \frac{d^3q}{(2\pi)^{3/2}} i \frac{k^l e^{\lambda,m}(\mathbf{k})}{k^2} (\mathcal{T}_m^r(\mathbf{p}) p_l + \mathcal{T}_l^r(\mathbf{p}) p_m) \frac{p^s}{p^2} q_r q_s \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{p}-\mathbf{q}} \Phi_{\mathbf{q}} \\ \times f_3^{(3)}(|\mathbf{k}-\mathbf{p}|, |\mathbf{p}-\mathbf{q}|, \mathbf{q}, \eta) , \quad (3.11)$$

$$S_4^{\lambda,(3)}(\eta, \mathbf{k}) = \int \frac{d^3p}{(2\pi)^{3/2}} \int \frac{d^3q}{(2\pi)^{3/2}} i \frac{k^l e^{\lambda,m}(\mathbf{k})}{k^2} p_l p_m \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{p}-\mathbf{q}} \Phi_{\mathbf{q}} \\ \times f_4^{(3)}(|\mathbf{k}-\mathbf{p}|, |\mathbf{p}-\mathbf{q}|, \mathbf{q}, \eta) , \quad (3.12)$$

and the expression of $f_i^{(3)}(|\mathbf{k}-\mathbf{p}|, |\mathbf{p}-\mathbf{q}|, \mathbf{q}, \eta)$ are given by,

$$f_1^{(3)}(u, \bar{u}, \bar{v}, x, y) = 12T_\phi(ux)T_\phi(\bar{u}y)T_\phi(\bar{v}y) - 4ux \frac{d}{d(ux)} T_\phi(ux)T_\phi(\bar{u}y)T_\phi(\bar{v}y) \\ - \frac{2u^2x^2}{3} T_\phi(ux)T_\phi(\bar{u}y)T_\phi(\bar{v}y) - 6\bar{u}uxy \frac{d}{d(ux)} T_\phi(ux) \frac{d}{d(\bar{u}y)} T_\phi(\bar{u}y)T_\phi(\bar{v}y) \\ - \frac{4u^2\bar{u}x^2y}{3} T_\phi(ux) \frac{d}{d(\bar{u}y)} T_\phi(\bar{u}y)T_\phi(\bar{v}y) - 4\bar{u}\bar{v}y^2 T_\phi(ux) \frac{d}{d(\bar{u}y)} T_\phi(\bar{u}y) \frac{d}{d(\bar{v}y)} T_\phi(\bar{v}y) \\ - 2\bar{u}\bar{v}uy^2x \frac{d}{d(ux)} T_\phi(ux) \frac{d}{d(\bar{u}y)} T_\phi(\bar{u}y) \frac{d}{d(\bar{v}y)} T_\phi(\bar{v}y) \\ - \frac{2u^2\bar{u}\bar{v}x^2y^2}{3} T_\phi(ux) \frac{d}{d(\bar{u}y)} T_\phi(\bar{u}y) \frac{d}{d(\bar{v}y)} T_\phi(\bar{v}y) , \quad (3.13)$$

$$f_2^{(3)}(u, \bar{u}, \bar{v}, x, y) = -6T_\phi(ux)T_\phi(\bar{u}y)T_\phi(\bar{v}y) - 4\bar{u}y T_\phi(ux) \frac{d}{d(\bar{u}y)} T_\phi(\bar{u}y)T_\phi(\bar{v}y) \\ - T_\phi(ux)p^2 I_h^{(2)}(\bar{u}, \bar{v}, y) - 2\bar{u}\bar{v}y^2 T_\phi(ux) \frac{d}{d(\bar{u}y)} T_\phi(\bar{u}y) \frac{d}{d(\bar{v}y)} T_\phi(\bar{v}y) \\ + \frac{u}{v^2x} \frac{d}{d(ux)} T_\phi(ux)p^2 I_h^{(2)}(\bar{u}, \bar{v}, y) - \frac{u^2}{3v^2} T_\phi(ux)p^2 I_h^{(2)}(\bar{u}, \bar{v}, y) \\ - \frac{1-u^2-v^2}{2v^2} T_\phi(ux)p^2 I_h^{(2)}(\bar{u}, \bar{v}, y) , \quad (3.14)$$

$$\begin{aligned}
f_3^{(3)}(u, \bar{u}, \bar{v}, x, y) &= \frac{u}{v} \frac{d}{d(ux)} T_\phi(ux) p I_V^{(2)}(\bar{u}, \bar{v}, y) + \frac{y}{8} T_\phi(ux) p I_V^{(2)}(\bar{u}, \bar{v}, y) \\
&+ \frac{xuy}{8} \frac{d}{d(ux)} T_\phi(ux) p I_V^{(2)}(\bar{u}, \bar{v}, y) - 4T_\phi(ux) T_\phi(\bar{u}y) T_\phi(\bar{v}y) \\
&+ 8\bar{u}y T_\phi(ux) \frac{d}{d(\bar{u}y)} T_\phi(\bar{u}y) T_\phi(\bar{v}y) + 16T_\phi(ux) T_\phi(\bar{u}y) T_\phi(\bar{v}y) \\
&+ 4\bar{u}\bar{v}y^2 T_\phi(ux) \frac{d}{d(\bar{u}y)} T_\phi(\bar{u}y) \frac{d}{d(\bar{v}y)} T_\phi(\bar{v}y) ,
\end{aligned} \tag{3.15}$$

$$\begin{aligned}
f_4^{(3)}(u, \bar{u}, \bar{v}, \eta) &= y \left(T_\phi(ux) \frac{\partial}{\partial y} I_\psi^{(2)}(\bar{u}, \bar{v}, y) \right) + uxy \left(\frac{d}{d(ux)} T_\phi(ux) \frac{\partial}{\partial y} I_\psi^{(2)}(\bar{u}, \bar{v}, y) \right) \\
&+ ux \left(\frac{d}{d(ux)} T_\phi(ux) (I_\psi^{(2)}(\bar{u}, \bar{v}, y) + f_\phi^{(2)}(\bar{u}, \bar{v}, y)) \right) \\
&+ 3 \left(T_\phi(ux) (I_\psi^{(2)}(\bar{u}, \bar{v}, y) + f_\phi^{(2)}(\bar{u}, \bar{v}, y)) \right) ,
\end{aligned} \tag{3.16}$$

where we have set $|\mathbf{k} - \mathbf{p}| = uk$, $p = vk$, $|\mathbf{k} - \mathbf{q}| = wk$, $|\mathbf{p} - \mathbf{q}| = \bar{u}p$, $q = \bar{v}p$, $x = k\eta$, and $y = p\eta$. As shown in Eqs. (3.14)–(3.16), we need to calculate the kernel function of the second order scalar, vector, tensor perturbations induced by the first order scalar perturbations.

Corresponding to four kinds of the source terms in Eqs. (3.9)–(3.12), we obtain the formal expression of third order vector mode

$$V^{\lambda,(3)}(\eta, \mathbf{k}) = V_1^{\lambda,(3)}(\eta, \mathbf{k}) + V_2^{\lambda,(3)}(\eta, \mathbf{k}) + V_3^{\lambda,(3)}(\eta, \mathbf{k}) + V_4^{\lambda,(3)}(\eta, \mathbf{k}) , \tag{3.17}$$

where $V_i^{\lambda,(3)}$, ($i = 1, 2, 3, 4$) are defined as

$$\begin{aligned}
V_1^{\lambda,(3)}(\eta, \mathbf{k}) &= \int \frac{d^3p}{(2\pi)^{3/2}} \int \frac{d^3q}{(2\pi)^{3/2}} i \frac{k^l e^{\lambda,m}(\mathbf{k})}{k^2} (p_l - q_l) q_m \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{p}-\mathbf{q}} \Phi_{\mathbf{q}} \\
&\times I_1^{(3)}(|\mathbf{k} - \mathbf{p}|, |\mathbf{p} - \mathbf{q}|, \mathbf{q}, \eta) ,
\end{aligned} \tag{3.18}$$

$$\begin{aligned}
V_2^{\lambda,(3)}(\eta, \mathbf{k}) &= \int \frac{d^3p}{(2\pi)^{3/2}} \int \frac{d^3q}{(2\pi)^{3/2}} i \frac{k^l e^{\lambda,m}(\mathbf{k})}{k^2} \Lambda_{lm}^{rs}(\mathbf{p}) q_r q_s \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{p}-\mathbf{q}} \Phi_{\mathbf{q}} \\
&\times I_2^{(3)}(|\mathbf{k} - \mathbf{p}|, |\mathbf{p} - \mathbf{q}|, \mathbf{q}, \eta) ,
\end{aligned} \tag{3.19}$$

$$\begin{aligned}
V_3^{\lambda,(3)}(\eta, \mathbf{k}) &= \int \frac{d^3p}{(2\pi)^{3/2}} \int \frac{d^3q}{(2\pi)^{3/2}} i \frac{k^l e^{\lambda,m}(\mathbf{k})}{k^2} (\mathcal{T}_m^s(\mathbf{p}) p_l + \mathcal{T}_l^s(\mathbf{p}) p_m) \frac{p^s}{p^2} q_r q_s \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{p}-\mathbf{q}} \Phi_{\mathbf{q}} \\
&\times I_3^{(3)}(|\mathbf{k} - \mathbf{p}|, |\mathbf{p} - \mathbf{q}|, \mathbf{q}, \eta) ,
\end{aligned} \tag{3.20}$$

$$\begin{aligned}
V_4^{\lambda,(3)}(\eta, \mathbf{k}) &= \int \frac{d^3p}{(2\pi)^{3/2}} \int \frac{d^3q}{(2\pi)^{3/2}} i \frac{k^l e^{\lambda,m}(\mathbf{k})}{k^2} p_l p_m \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{p}-\mathbf{q}} \Phi_{\mathbf{q}} \\
&\times I_4^{(3)}(|\mathbf{k} - \mathbf{p}|, |\mathbf{p} - \mathbf{q}|, \mathbf{q}, \eta) .
\end{aligned} \tag{3.21}$$

Substituting Eqs. (3.9)–(3.12) and (3.18)–(3.21) into Eq. (3.8), we obtain the equations of motion of kernel functions $I_i^{(3)}(u, \bar{u}, \bar{v}, x)$ in the form of

$$I_i^{(3)'}(u, \bar{u}, \bar{v}, x) + 2\mathcal{H}I_i^{(3)}(u, \bar{u}, \bar{v}, x) = 12f_i^{(3)}(u, \bar{u}, \bar{v}, x) , \quad (i = 1, 2, 3, 4) . \tag{3.22}$$




Figure 2. The squares of the kernel functions of the second order and four third order vector perturbations. Here we have set $u = v = \bar{u} = \bar{v} = 1$.

The kernel functions can be expressed as

$$I_i^{(3)}(u, \bar{u}, \bar{v}, x) = \frac{12}{k} \int_0^x d\bar{x} \left(\frac{\bar{x}^2}{x^2} f_i^{(3)}(u, \bar{u}, \bar{v}, \bar{x}) \right), \quad (i = 1, 2, 3, 4). \quad (3.23)$$

It is similar to the second order kernel function. We present $(I_i(u = 1, v = 1, \bar{u} = 1, \bar{v} = 1, x))^2$, $(i, j = 1, 2, 3, 4)$ as function of $x = k\eta$ in Fig. 2. We notice that $I_4^{(3)}$ has the largest amplitudes for large x .

3.2 Third order power spectrum

In this section, we will study the power spectrum of the third order vector mode. An explicit expression of the power spectrum $\mathcal{P}_V^{(3)}(\eta, k)$ is presented. Substituting Eq. (3.17) into the definition of the power spectrum, we obtain

$$\delta_{\lambda\lambda'} \langle V^{\lambda, (3)}(\eta, \mathbf{k}) V^{\lambda', (3)}(\eta, \mathbf{k}') \rangle = \sum_{i,j=1}^4 \langle V_i^{\lambda, (3)}(\eta, \mathbf{k}) V_j^{\lambda', (3)}(\eta, \mathbf{k}') \rangle = 2\delta(\mathbf{k} + \mathbf{k}') \frac{2\pi^2}{k^3} \sum_{i,j=1}^4 \mathcal{P}_V^{ij}(\eta, k). \quad (3.24)$$

The power spectrum of third order vector mode $\mathcal{P}_V^{(3)}(\eta, \mathbf{k})$ can be written as the sum of sixteen terms,

$$\mathcal{P}_V^{(3)}(\eta, \mathbf{k}) = \sum_{i,j=1}^4 \mathcal{P}_V^{ij}(\eta, \mathbf{k}). \quad (3.25)$$

The summation of i and j is originated from the four kinds of source terms of the third order vector mode $V_i^{(3)}$. Based on the formal expressions of $V_i^{\lambda}(\eta, \mathbf{k})$ in Eqs. (3.18)–(3.21), we obtain the formal expression of power spectra $\mathcal{P}_V^{ij}(\eta, \mathbf{k})$ in the form of

$$\begin{aligned} \mathcal{P}_V^{ij}(\eta, k) &= \frac{k^3 (2\pi^2)^2}{2} \int \frac{d^3 p \, d^3 q \, d^3 p' \, d^3 q'}{(2\pi)^6} \mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, \mathbf{p}', \mathbf{q}, \mathbf{q}') \mathcal{C}(\mathbf{k}, \mathbf{k}', \mathbf{p}, \mathbf{p}', \mathbf{q}, \mathbf{q}') \\ &\quad \times I_i^{(3)}(|\mathbf{k} - \mathbf{p}|, |\mathbf{p} - \mathbf{q}|, p, q, \eta) I_j^{(3)}(|\mathbf{k}' - \mathbf{p}'|, |\mathbf{p}' - \mathbf{q}'|, p', q', \eta), \end{aligned} \quad (3.26)$$

where $\mathcal{C}(\mathbf{k}, \mathbf{k}', \mathbf{p}, \mathbf{p}', \mathbf{q}, \mathbf{q}')$ is derived from the six-point correlation function, and its explicit expression is shown in Eq. (C.2). The integral and measure $\int \frac{d^3 p d^3 q d^3 p' d^3 q'}{(2\pi)^6}$ should be simplified as the integrals of u, v, \bar{u}, \bar{v} , and w with the explicit upper and lower limits in terms of a given coordinate system. Here the calculations are similar to the calculations in the power spectrum of second order vector mode, and it will be more complicated. The polynomial of momentums $\mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, \mathbf{p}', \mathbf{q}, \mathbf{q}')$ is originated from the decomposition operators, the polarization vectors, and the polarization tensors. We will simplify these polynomial of momentums as a polynomial of u, v, \bar{u}, \bar{v} , and w . The product of third order kernel functions $I_i^{(3)}(|\mathbf{k} - \mathbf{p}|, |\mathbf{p} - \mathbf{q}|, p, q, \eta) I_j^{(3)}(|\mathbf{k}' - \mathbf{p}'|, |\mathbf{p}' - \mathbf{q}'|, p', q', \eta)$ have been studied in Sec. 3.1. The Wick's expressions of six point function $\mathcal{C}(\mathbf{k}, \mathbf{k}', \mathbf{p}, \mathbf{p}', \mathbf{q}, \mathbf{q}')$ are shown in Appendix C.2. There are six non-trivial terms in the Wick's expressions, they both contain the three dimensional delta functions of \mathbf{p}' and \mathbf{q}' . For example, the first term in Eq. (C.2) contains $\delta(\mathbf{p} + \mathbf{p}') \delta(\mathbf{q} + \mathbf{q}')$. When we integral over \mathbf{p}' and \mathbf{q}' , we will obtain the substitutions $\mathbf{p}' \rightarrow -\mathbf{p}$ and $\mathbf{q}' \rightarrow -\mathbf{q}$. These substitutions will change the variables \mathbf{p}' and \mathbf{q}' in the polynomial $\mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, \mathbf{p}', \mathbf{q}, \mathbf{q}')$ and the second kernel functions $I_j^{(3)}(|\mathbf{k}' - \mathbf{p}'|, |\mathbf{p}' - \mathbf{q}'|, p', q', \eta)$. The explicit expression of the power spectra \mathcal{P}_V^{ij} in Eq. (3.26) can be written as,

$$\begin{aligned} \mathcal{P}_V^{ij} &= \frac{1}{2\pi} \int_0^\infty dv \int_0^\infty d\bar{v} \int_{|1-\bar{v}v|}^{1+\bar{v}v} dw \int_{|1-v|}^{1+v} du \int_{\bar{u}_-}^{\bar{u}_+} d\bar{u} \left\{ \frac{w}{u^2 \bar{u}^2 v \bar{v}^2 \sqrt{Y(1-X^2)}} I_i^{(3)}(u, v, \bar{u}, \bar{v}, x) \right. \\ &\quad \left. \times \sum_{a=1}^6 \mathbb{P}_a^{ij}(k, u, v, \bar{u}, \bar{v}, w, x) I_j^{(3),a}(u, v, \bar{u}, \bar{v}, w, x) P_\Phi(ku) P_\Phi(k\bar{u}v) P_\Phi(k\bar{v}v) \right\}, \end{aligned} \quad (3.27)$$

where

$$\begin{aligned} \bar{u}_\pm &= \left(1 + \bar{v}^2 - \frac{(1+v^2-u^2)(1+(\bar{v}v)^2-w^2)}{2v^2} \right. \\ &\quad \left. \pm 2\bar{v} \sqrt{\left(1 - \left(\frac{1+v^2-u^2}{2v} \right)^2 \right) \left(1 - \left(\frac{1+(\bar{v}v)^2-w^2}{2\bar{v}v} \right)^2 \right)} \right)^{\frac{1}{2}}, \end{aligned} \quad (3.28)$$

$$\begin{aligned} X &= (-1 + u^2 + v^2 - 2\bar{u}^2 v^2 + v^2 \bar{v}^2 + u^2 v^2 \bar{v}^2 - v^4 \bar{v}^2 + w^2 - u^2 w^2 + v^2 w^2) \\ &\quad \times (1 - 2u^2 + u^4 - 2v^2 - 2u^2 v^2 + v^4) (1 - 2v^2 \bar{v}^2 + v^4 \bar{v}^4 - 2w^2 - 2v^2 \bar{v}^2 w^2 + w^4)^{-\frac{1}{2}}, \end{aligned} \quad (3.29)$$

$$Y = (1 - 2u^2 + u^4 - 2v^2 - 2u^2 v^2 + v^4) (1 - 2v^2 \bar{v}^2 + v^4 \bar{v}^4 - 2w^2 - 2v^2 \bar{v}^2 w^2 + w^4). \quad (3.30)$$

Some details of Eq.(3.27) are given in Appendix D.2. The summation of index a in Eq. (3.27) comes from the six terms in Wick's expansions. More precisely, the polynomials \mathbb{P}_a^{ij} and the

second kernel functions $I_j^{(3),a}(|\mathbf{k}' - \mathbf{p}'|, |\mathbf{p}' - \mathbf{q}'|, p', q', \eta)$ take the form of

$$\mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, \mathbf{p}', \mathbf{q}, \mathbf{q}') \rightarrow \mathbb{P}_a^{ij} = \left(\mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}), \mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, \mathbf{q} - \mathbf{p}), \mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{q}) \right. \\ \left. , \mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{q} - \mathbf{p}), \mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{k} - \mathbf{q}, \mathbf{q}, \mathbf{p} - \mathbf{k}) \right. \\ \left. , \mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{k} + \mathbf{p}) \right), \quad (3.31)$$

$$I_j^{(3)}(|\mathbf{k}' - \mathbf{p}'|, |\mathbf{p}' - \mathbf{q}'|, p', q', \eta) \rightarrow I_j^{(3),a} = \left(I_j^{(3)}(|\mathbf{k} - \mathbf{p}|, |\mathbf{p} - \mathbf{q}|, p, q, \eta), I_j^{(3)}(|\mathbf{k} - \mathbf{p}|, |\mathbf{p} - \mathbf{q}|, p, |\mathbf{p} - \mathbf{q}|, \eta) \right. \\ \left. , I_j^{(3)}(|\mathbf{k} - \mathbf{p}|, |\mathbf{p} - \mathbf{q}|, |\mathbf{p} - \mathbf{q} - \mathbf{k}|, q, \eta), I_j^{(3)}(|\mathbf{k} - \mathbf{p}|, |\mathbf{p} - \mathbf{q}|, |\mathbf{q} - \mathbf{k}|, |\mathbf{p} - \mathbf{q}|, \eta) \right. \\ \left. , I_j^{(3)}(|\mathbf{k} - \mathbf{p}|, |\mathbf{p} - \mathbf{q}|, |\mathbf{p} - \mathbf{q} - \mathbf{k}|, |\mathbf{p} - \mathbf{k}|, \eta) \right. \\ \left. , I_j^{(3)}(|\mathbf{k} - \mathbf{p}|, |\mathbf{p} - \mathbf{q}|, |\mathbf{q} - \mathbf{k}|, |\mathbf{p} - \mathbf{k}|, \eta) \right). \quad (3.32)$$

In order to obtain a specific result of $\mathcal{P}_V^{(3)}(\eta, \mathbf{k})$, we consider a monochromatic primordial power spectrum,

$$P_\Phi(k) = Ak_*\delta(k - k_*). \quad (3.33)$$

In this case, the power spectrum \mathcal{P}_V^{ij} reduce to

$$\mathcal{P}_V^{ij}(x, \tilde{k}) = \frac{\mathcal{A}^3 \tilde{k}^3}{2\pi} \Theta(3 - \tilde{k}) \int_{|1 - \frac{1}{\tilde{k}}|}^{\min\{\frac{2}{\tilde{k}}, 1 + \frac{1}{\tilde{k}}\}} dv \int_{w_-}^{w_+} dw \left(\frac{vw}{\sqrt{Y(1 - X^2)}} \right. \\ \left. \times I_i^{(3)}(u, v, \bar{u}, \bar{v}, x) \sum_{a=1}^3 \mathbb{P}_a^{ij} I_j^{(3),a}(u, v, \bar{u}, \bar{v}, w, x) \right)_{u=\frac{1}{\tilde{k}}, \bar{u}=\bar{v}=\frac{1}{v\tilde{k}}}, \quad (3.34)$$

where we have defined $\tilde{k} \equiv \frac{k}{k_*}$ and

$$w_\pm = \left(\frac{1}{2} + \frac{3}{2\tilde{k}^2} - \frac{1}{2}v^2 \pm \frac{1}{2v} \sqrt{\left(v^2 - \frac{4}{\tilde{k}^2} \right) \left(v^2 - \left(1 - \frac{1}{\tilde{k}} \right)^2 \right) \left(v^2 - \left(1 + \frac{1}{\tilde{k}} \right)^2 \right)} \right)^{\frac{1}{2}}. \quad (3.35)$$

In the case of monochromatic primordial power spectrum $P_\Phi(k) = Ak_*\delta(k - k_*)$, the summation of index a reduce to three terms

$$\mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, \mathbf{p}', \mathbf{q}, \mathbf{q}') \rightarrow \mathbb{P}_a^{ij} = \left(\mathbb{P}_1^{ij}, \mathbb{P}_2^{ij}, \mathbb{P}_3^{ij} \right), \quad (3.36)$$

where the explicit expressions of \mathbb{P}_a^{ij} , ($a = 1, 2, 3$) in Eq. (3.36) are shown in the Appendix E. The corresponding kernel functions can be expressed as

$$I_j^{(3)}(|\mathbf{k}' - \mathbf{p}'|, |\mathbf{p}' - \mathbf{q}'|, p', q', \eta) \rightarrow I_j^{(3),a}(|\mathbf{k}' - \mathbf{p}'|, |\mathbf{p}' - \mathbf{q}'|, p', q', \eta) \\ = 2 \left(I_j^{(3),1}(k_*, k_*, p, k_*, \eta), I_j^{(3),2}(k_*, k_*, |\mathbf{k} - \mathbf{q}|, k_*, \eta), I_j^{(3),3}(k_*, k_*, |\mathbf{p} - \mathbf{q} - \mathbf{k}|, k_*, \eta) \right). \quad (3.37)$$




Figure 3. The power spectra of the third order vector perturbations. The black dot-dashed curve denotes the total power spectrum $\mathcal{P}_V^{(3)} \equiv \sum_{i,j=1}^4 \mathcal{P}_V^{ij}$. The diagonal terms \mathcal{P}_V^{11} (blue solid curve), \mathcal{P}_V^{22} (orange dashed curve) and \mathcal{P}_V^{44} (green dot-dashed curve) are shown. It shows that $\mathcal{P}_V^{(3)} \simeq \mathcal{P}_V^{44}$.

Namely,

$$\begin{aligned}
I_j^{(3),1}(u', v', \bar{u}', \bar{v}', x) &= I_j^{(3)}\left(\frac{1}{\tilde{k}}, v, \frac{1}{v\tilde{k}}, \frac{1}{v\tilde{k}}, x\right), \\
I_j^{(3),2}(u', v', \bar{u}', \bar{v}', x) &= I_j^{(3)}\left(\frac{1}{\tilde{k}}, w, \frac{1}{w\tilde{k}}, \frac{1}{w\tilde{k}}, x\right), \\
I_j^{(3),3}(u', v', \bar{u}', \bar{v}', x) &= I_j^{(3)}\left(\frac{1}{\tilde{k}}, \left(1 - v^2 - w^2 + 3/\tilde{k}^2\right)^{\frac{1}{2}}, \frac{1}{\tilde{k}} \left(1 - v^2 - w^2 + 3/\tilde{k}^2\right)^{-\frac{1}{2}}, \right. \\
&\quad \left. \frac{1}{\tilde{k}} \left(1 - v^2 - w^2 + 3/\tilde{k}^2\right)^{-\frac{1}{2}}, x\right)
\end{aligned} \tag{3.38}$$

where we have defined

$$|\mathbf{k}' - \mathbf{p}'| = u'k', \quad |\mathbf{p}' - \mathbf{q}'| = \bar{u}'p' = \bar{u}'v'k', \quad q' = \bar{v}'p' = \bar{v}'v'k'. \tag{3.39}$$

In the end of this section, we calculate the third order power spectrum $\mathcal{P}_V^{(3)}$, i.e.,

$$\mathcal{P}_V^{(3)} = \sum_{i,j=1}^4 \mathcal{P}_V^{ij}(x, \tilde{k}). \tag{3.40}$$

We plot the third order power spectra in Fig. 3. It shows that the third order gravitational waves sourced by the second order scalar perturbations dominate the energy density spectrum, i.e., $\mathcal{P}_V^{(3)} \simeq \mathcal{P}_V^{44}$. This is consistent with the kernel function shown in Fig. 2, where $I_4^{(3)}$ have the largest amplitude for large x and contributes most to the total power spectrum.

4 Conclusions and discussions

In this paper, we studied the second order and third order vector modes induced by first order primordial scalar perturbations. The explicit expressions of the power spectra were presented. We conclude that the second order vector modes can not be generated by a monochromatic primordial power spectrum. Moreover, we calculated the kernel functions and the power spectra of the third order vector modes for a monochromatic primordial power spectrum. It shows that the power spectra of the third order vector modes are not zero, the third order vector mode is the first non-trivial order.

The third order vector modes have four kinds of source terms. More, precisely, the first order scalar perturbation induces the second order scalar, vector and tensor perturbations. Then, the first order scalar, the second order scalar, vector and tensor perturbations all induce the third order vector modes. The third order vector modes sourced by the second order scalar perturbations dominate the two point function $\langle V^{(3),\lambda} V^{(3),\lambda'} \rangle$ and power spectrum corresponded. And the third order vector modes sourced by the second order vector perturbations are equal to zero, this result is consistent with the conclusion of the second order vector modes.

It would be helpful to compare the third order vector modes with the third order scalar induced gravitational waves for a monochromatic primordial power spectrum [5]. First, they both have four kinds of source terms, and their total power spectra are both dominated by the source term of the second order scalar perturbations. Second, the source term of the second order vector perturbation does not affect their total power spectra. Third, the third order power spectrum of gravitational waves is composed of $\langle h^{(3),\lambda} h^{(3),\lambda'} \rangle$ and $\langle h^{(2),\lambda} h^{(4),\lambda'} \rangle$. Therefore, if we want to calculate the third order power spectrum of gravitational waves completely, we need to study the fourth order scalar induced gravitational waves $h^{(4),\lambda}$. Similarly, it is necessary to consider the correlation function $\langle V^{(2),\lambda} V^{(4),\lambda'} \rangle$ for a general primordial power spectra.

Acknowledgments

We thank Dr. Q.H. Zhu for useful discussions. This work has been funded by the National Nature Science Foundation of China under grant No. 12075249 and 11690022, and the Key Research Program of the Chinese Academy of Sciences under Grant No. XDPB15

A Decomposition Operators

We review the decomposition operators on FRW spacetime. The spatial tensor field S_{ij} on FRW spacetime could be decomposed as scalar, vector and tensor modes

$$S_{ij} = S_{ij}^{(H)} + 2\delta_{ij} S^{(\Psi)} + 2\partial_i \partial_j S^{(E)} + \partial_j S_i^{(C)} + \partial_i S_j^{(C)}. \quad (\text{A.1})$$

We define the decomposed operators as follow

$$S_{ij}^{(H)} \equiv \Lambda_{ij}^{kl} S_{kl} = \left(\mathcal{T}_i^k \mathcal{T}_j^l - \frac{1}{2} \mathcal{T}_{ij} \mathcal{T}^{kl} \right) S_{kl}, \quad S^{(\Psi)} \equiv \frac{1}{4} \mathcal{T}^{kl} S_{kl}. \quad (\text{A.2})$$

$$S^{(E)} \equiv \frac{1}{2} \Delta^{-1} \left(\partial^k \Delta^{-1} \partial^l - \frac{1}{2} \mathcal{T}^{kl} \right) S_{lk}, \quad S_i^{(C)} \equiv \Delta^{-1} \partial^l \mathcal{T}_i^k S_{lk}. \quad (\text{A.3})$$

The transverse operators is defined by

$$\mathcal{T}_j^i \equiv \delta_i^j - \partial^i \Delta^{-1} \partial_j . \quad (\text{A.4})$$

These decomposed operators can only use in FRW spacetime. For arbitrary spacetime, we can't distinguish between $S_i^{(C)}$ and $S^{(E)}$ [31].

B Higher order perturbations of Einstein equation

In this appendix, we use the `xPand` package to study the higher order perturbations of Einstein equation on FRW spacetime [32]. The `xPand` package can help us to obtain and simplify the perturbation equations.

B.1 Second order perturbation equation

In this appendix, we study the second order perturbation equation and derive the equation of motion of second order vector mode $V_i^{(2)}$. The space-space part of the second order perturbed equation is presented as follows

$$\begin{aligned} & \frac{1}{4} h_{ij}^{(2)''} + \frac{1}{2} \mathcal{H} h_{ij}^{(2)'} - \frac{1}{2} \mathcal{H}^2 h_{ij}^{(2)} - \mathcal{H}' h_{ij}^{(2)} - \frac{1}{6} \kappa a^2 h_{ij}^{(2)} \rho^{(0)} - \frac{4}{3} \kappa a^2 u_i^{(1)} u_j^{(1)} \rho^{(0)} - \frac{1}{4} \Delta h_{ij}^{(2)} \\ & - \frac{1}{2} \mathcal{H} \partial_i V_j^{(2)} - \frac{1}{4} \partial_i V_j^{(2)'} - \frac{1}{2} \mathcal{H} \partial_j V_i^{(2)} - \frac{1}{4} \partial_j V_i^{(2)'} + \partial_i \phi^{(1)} \partial_j \phi^{(1)} - \partial_i \psi^{(1)} \partial_j \phi^{(1)} - \partial_j \psi^{(1)} \partial_i \phi^{(1)} \\ & + 3 \partial_i \psi^{(1)} \partial_j \psi^{(1)} + 2 \phi^{(1)} \partial_i \partial_j \phi^{(1)} + 2 \psi^{(1)} \partial_i \partial_j \psi^{(1)} - \frac{1}{2} \partial_i \partial_j \phi^{(2)} + \frac{1}{2} \partial_i \partial_j \psi^{(2)} = 0 , \end{aligned} \quad (\text{B.1})$$

where $u_i^{(1)}$ is the first order transverse part of three dimensional velocity in energy-momentum tensor, $\rho^{(0)}$ is the zero order density in energy-momentum tensor, \mathcal{H}' is the derivative of \mathcal{H} with respect to η . The time-time component and space-time component of lower order perturbation equations can help us to express $u_i^{(1)}$, $\rho^{(0)}$, and \mathcal{H}' in terms of \mathcal{H} , and the first order perturbations. Namely,

$$\mathcal{H}' = -\mathcal{H}^2 , \quad \rho^{(0)} = \frac{3\mathcal{H}^2}{\kappa a^2} , \quad (\text{B.2})$$

$$\rho^{(1)} = \frac{-6\mathcal{H} (\mathcal{H} \phi^{(1)} + \psi^{(1)'}) + 2\Delta \psi^{(1)}}{\kappa a^2} , \quad u_i^{(1)} = -\frac{(\mathcal{H} \partial_i \phi^{(1)} + \partial_i \psi^{(1)'})}{2\mathcal{H}^2} , \quad (\text{B.3})$$

Substituting Eqs. (B.2)–(B.3) into Eq. (B.1), we obtain

$$\begin{aligned} & \frac{1}{4} h_{ij}^{(2)''} + \frac{1}{2} \mathcal{H} h_{ij}^{(2)'} - \frac{1}{4} \Delta h_{ij}^{(2)} - \frac{1}{2} \mathcal{H} \partial_i V_j^{(2)} - \frac{1}{4} \partial_i V_j^{(2)'} - \frac{1}{2} \mathcal{H} \partial_j V_i^{(2)} - \frac{1}{4} \partial_j V_i^{(2)'} - \frac{1}{2} \partial_i \partial_j \phi^{(2)} \\ & + \frac{1}{2} \partial_i \partial_j \psi^{(2)} + \partial_i \phi^{(1)} \partial_j \phi^{(1)} - \partial_i \psi^{(1)} \partial_j \phi^{(1)} - \partial_j \psi^{(1)} \partial_i \phi^{(1)} - \frac{\partial_i \psi^{(1)'} \partial_j \phi^{(1)}}{\mathcal{H}} - \frac{\partial_i \phi^{(1)} \partial_j \psi^{(1)'}}{\mathcal{H}} \\ & - \frac{\partial_i \psi^{(1)'} \partial_j \psi^{(1)'}}{\mathcal{H}^2} + 3 \partial_i \psi^{(1)} \partial_j \psi^{(1)} + 2 \phi^{(1)} \partial_i \partial_j \phi^{(1)} + 2 \psi^{(1)} \partial_i \partial_j \psi^{(1)} = 0 . \end{aligned} \quad (\text{B.4})$$

Using the decomposition operator $\Delta^{-1} \partial^l \mathcal{T}_i^k$ in Eq. (A.3) to extract the equation of motion of second order vector perturbation, we obtain

$$V_l^{(2)'}(\eta, \mathbf{x}) + 2\mathcal{H} V_l^{(2)}(\eta, \mathbf{x}) = -4\Delta^{-1} \mathcal{T}_l^r \partial^s \mathcal{S}_{rs}^{(2)}(\eta, \mathbf{x}) , \quad (\text{B.5})$$

where the source term is given by

$$\begin{aligned} \Delta^{-1} \mathcal{T}_l^r \partial^s \mathcal{S}_{rs}^{(2)}(\eta, \mathbf{x}) = & \Delta^{-1} \mathcal{T}_l^r \partial^s \left(\partial_r \phi^{(1)} \partial_s \phi^{(1)} - \frac{1}{\mathcal{H}} \left(\partial_r \phi^{(1)'} \partial_s \phi^{(1)} + \partial_r \phi^{(1)} \partial_s \phi^{(1)'} \right) \right. \\ & \left. + 4\phi^{(1)} \partial_r \partial_s \phi^{(1)} - \frac{1}{\mathcal{H}^2} \partial_r \phi^{(1)'} \partial_s \phi^{(1)'} \right). \end{aligned} \quad (\text{B.6})$$

Here, we have set $\psi^{(1)} = \phi^{(1)}$ from the equation of motion of first order scalar modes.

B.2 Third order perturbation equation

In this appendix, we investigate the third order perturbation equation and obtain the equation of motion of third order vector mode $V_i^{(3)}$. The space-space part of the third order perturbed equation is presented as follows,

$$\begin{aligned} & -\frac{1}{6} \mathcal{H} \partial_i V_j^{(3)} - \frac{1}{6} \mathcal{H} \partial_j V_i^{(3)} - \frac{1}{12} \partial_i V_j^{(3)'} - \frac{1}{12} \partial_j V_i^{(3)'} + \frac{1}{6} \partial_i \partial_j \psi^{(3)} - \frac{1}{6} \partial_i \partial_j \phi^{(3)} \\ & + \frac{1}{12} h_{ij}^{(3)''} + \frac{1}{6} h_{ij}^{(3)'} \mathcal{H} - \frac{1}{6} h_{ij}^{(3)} \mathcal{H}^2 - \frac{1}{3} h_{ij}^{(3)} \mathcal{H}' - \frac{1}{18} \kappa a^2 h_{ij}^{(3)} \rho^{(0)} - \frac{1}{12} \Delta h_{ij}^{(3)} - \frac{2}{3} \kappa a^2 V_j^{(2)} u_i^{(1)} \rho^{(0)} \\ & - \frac{2}{3} \kappa a^2 V_i^{(2)} u_j^{(1)} \rho^{(0)} - \frac{2}{3} \kappa a^2 u_i^{(1)} u_j^{(2)} \rho^{(0)} - \frac{2}{3} \kappa a^2 u_j^{(1)} u_i^{(2)} \rho^{(0)} - \frac{1}{6} \kappa a^2 h_{ij}^{(2)} \rho^{(1)} - \frac{4}{3} \kappa a^2 u_i^{(1)} u_j^{(1)} \rho^{(1)} \\ & - h_{ij}^{(2)'} \mathcal{H} \phi^{(1)} + h_{ij}^{(2)} \mathcal{H}^2 \phi^{(1)} + 2h_{ij}^{(2)} \mathcal{H}' \phi^{(1)} + \frac{16}{3} \kappa a^2 u_i^{(1)} u_j^{(1)} \rho^{(0)} \phi^{(1)} + 4h_{ij}^{(2)} \mathcal{H} \phi^{(1)'} + \frac{3}{2} h_{ij}^{(2)} \phi^{(1)''} \\ & - \frac{1}{2} h_{ij}^{(2)} \Delta \phi^{(1)} + \frac{1}{2} h_i^{b(2)} \partial_b \partial_j \phi^{(1)} + \frac{1}{2} h_j^{b(2)} \partial_b \partial_i \phi^{(1)} - \partial_b h_{ij}^{(2)} \partial^b \phi^{(1)} + \mathcal{H} \phi^{(1)} \partial_i V_j^{(2)} + \frac{1}{2} \phi^{(1)'} \partial_i V_j^{(2)} \\ & + \frac{1}{2} \partial^b \phi^{(1)} \partial_i h_{jb}^{(2)} - \frac{1}{2} V_j^{(2)'} \partial_i \phi^{(1)} - V_j^{(2)} \mathcal{H} \partial_i \phi^{(1)} - \frac{1}{2} V_j^{(2)} \partial_i \phi^{(1)'} + \mathcal{H} \phi^{(1)} \partial_j V_i^{(2)} + \frac{1}{2} \phi^{(1)'} \partial_j V_i^{(2)} \\ & + \frac{1}{2} \phi^{(1)} \partial_j V_i^{(2)} + \frac{1}{2} \partial^b \phi^{(1)} \partial_j h_{ib}^{(2)} - \frac{1}{2} V_i^{(2)'} \partial_j \phi^{(1)} - V_i^{(2)} \mathcal{H} \partial_j \phi^{(1)} - \frac{1}{2} V_i^{(2)} \partial_j \phi^{(1)'} + 8\phi^{(1)} \partial_i \phi^{(1)} \partial_j \phi^{(1)} \\ & + \partial_i \psi^{(2)} \partial_j \phi^{(1)} + \partial_i \phi^{(1)} \partial_j \psi^{(2)} + \phi^{(2)} \partial_i \partial_j \phi^{(1)} + \phi^{(1)} \partial_i \partial_j \phi^{(2)} + \psi^{(2)} \partial_i \partial_j \phi^{(1)} + \phi^{(1)} \partial_i \partial_j \psi^{(2)} \\ & - \frac{1}{2} \phi^{(1)} \Delta h_{ij}^{(2)} - \frac{1}{2} h_{ij}^{(2)''} \phi^{(1)} + \frac{1}{2} \phi^{(1)} \partial_i V_j^{(2)'} = 0, \end{aligned} \quad (\text{B.7})$$

where $u_i^{(1)}$ and $u_i^{(2)}$ are first and second order transverse part of three dimensional velocity in energy-momentum tensor. We have set $\psi^{(1)} = \phi^{(1)}$ from the equation of motion of first order scalar perturbations. We express $\rho^{(0)}$, $\rho^{(1)}$, $u^{(1)}$, $u^{(2)}$, and \mathcal{H}' in Eq. (B.7) in terms of \mathcal{H} , the first order perturbations, and the second order perturbations,

$$\mathcal{H}' = -\mathcal{H}^2, \quad \rho^{(0)} = \frac{3\mathcal{H}^2}{\kappa a^2}, \quad (\text{B.8})$$

$$\rho^{(1)} = \frac{-6\mathcal{H} \left(\mathcal{H} \phi^{(1)} + \psi^{(1)'} \right) + 2\Delta \psi^{(1)}}{\kappa a^2}, \quad u_i^{(1)} = -\frac{\left(\mathcal{H} \partial_i \phi^{(1)} + \partial_i \psi^{(1)'} \right)}{2\mathcal{H}^2}, \quad (\text{B.9})$$

$$\begin{aligned} \rho^{(2)} = & \frac{1}{4\kappa a^2 \mathcal{H}} \left[2\mathcal{H} \left(12\mathcal{H}^3 \left(4(\phi^{(1)})^2 - \phi^{(2)} \right) - 12\mathcal{H}^2 \psi^{(2)'} - 8\partial_b \phi^{(1)'} \partial^b \phi^{(1)} \right. \right. \\ & \left. \left. + \mathcal{H} \left(12(\phi^{(1)'})^2 + 32\phi^{(1)} \Delta \phi^{(1)} + 4\Delta \psi^{(2)} + 8\partial_b \phi^{(1)} \partial^b \phi^{(1)} \right) \right) - 8\partial_b \phi^{(1)'} \partial^b \phi^{(1)'} \right], \end{aligned} \quad (\text{B.10})$$

$$\begin{aligned}
u_i^{(2)} = & -V_i^{(2)} + \frac{1}{32\mathcal{H}^4} \left[\frac{64}{3} \Delta\phi^{(1)} \partial_i \phi^{(1)'} - \frac{64}{3} \mathcal{H} \left(-\Delta\phi^{(1)} \partial_i \phi^{(1)} + 3\phi^{(1)'} \partial_i \phi^{(1)'} \right) \right. \\
& + 3\mathcal{H}^2 \left(\frac{4}{3} \Delta V_i^{(2)} - 32\phi^{(1)'} \partial_i \phi^{(1)} - \frac{160}{3} \phi^{(1)} \partial_i \phi^{(1)'} - \frac{16}{3} \partial_i \psi^{(2)'} \right) \\
& \left. + 12\mathcal{H}^3 \left(-\frac{8}{3} \phi^{(1)} \partial_i \phi^{(1)} - \frac{4}{3} \partial_i \phi^{(2)} \right) \right]. \tag{B.11}
\end{aligned}$$

Substituting Eqs. (B.8)–(B.11) into Eq. (B.7), we obtain

$$\begin{aligned}
& + \frac{2}{3\mathcal{H}^2} \Delta\phi^{(1)} \partial_l \phi^{(1)} \partial_m \phi^{(1)} + \frac{2}{3\mathcal{H}^4} \Delta\phi^{(1)} \partial_l \phi^{(1)'} \partial_m \phi^{(1)'} - \frac{3}{\mathcal{H}^2} \phi^{(1)'} \partial_l \phi^{(1)'} \partial_m \phi^{(1)} - \frac{3}{\mathcal{H}^2} \phi^{(1)'} \partial_m \phi^{(1)'} \partial_l \phi^{(1)} \\
& + \frac{2}{3\mathcal{H}^3} \Delta\phi^{(1)} \partial_l \phi^{(1)'} \partial_m \phi^{(1)} + \frac{2}{3\mathcal{H}^3} \Delta\phi^{(1)} \partial_m \phi^{(1)'} \partial_l \phi^{(1)} - \frac{2}{\mathcal{H}^3} \phi^{(1)'} \partial_l \phi^{(1)'} \partial_m \phi^{(1)'} - \frac{4}{\mathcal{H}^2} \phi^{(1)} \partial_l \phi^{(1)'} \partial_m \phi^{(1)'} \\
& - \frac{1}{6} \mathcal{H} \partial_l V_m^{(3)} - \frac{1}{6} \mathcal{H} \partial_m V_l^{(3)} - \frac{1}{12} \partial_l V_m^{(3)'} - \frac{1}{12} \partial_m V_l^{(3)'} + \frac{1}{6} \partial_l \partial_m \psi^{(3)} - \frac{1}{6} \partial_l \partial_m \phi^{(3)} \\
& + h_{lm}^{(3)''}(\eta, \mathbf{x}) + 2\mathcal{H} h_{ilm}^{(3)'}(\eta, \mathbf{x}) - \Delta h_{lm}^{(3)}(\eta, \mathbf{x}) + 12\phi^{(1)} \partial_l \phi^{(1)} \partial_m \phi^{(1)} - \frac{4}{\mathcal{H}} \phi^{(1)'} \partial_l \phi^{(1)} \partial_m \phi^{(1)} \\
& - \frac{1}{2} \phi^{(1)} \left(h_{lm}^{(2)''} + 2\mathcal{H} h_{lm}^{(2)'} - \Delta h_{lm}^{(2)} \right) - \phi^{(1)} \Delta h_{lm}^{(2)} - \phi^{(1)'} \mathcal{H} h_{lm}^{(2)} - \frac{1}{3} \Delta\phi^{(1)} h_{lm}^{(2)} - \partial^b \phi^{(1)} \partial_b h_{lm}^{(2)} \\
& + \phi^{(1)} \partial_l \left(V_m^{(2)'} + 2\mathcal{H} V_m^{(2)} \right) + \phi^{(1)} \partial_m \left(V_l^{(2)'} + 2\mathcal{H} V_l^{(2)} \right) + \phi^{(1)'} \left(\partial_l V_m^{(2)} + \partial_m V_l^{(2)} \right) \\
& - \frac{\phi^{(1)}}{8\mathcal{H}} \left(\partial_m \Delta V_l^{(2)} + \partial_l \Delta V_m^{(2)} \right) - \frac{\phi^{(1)'}}{8\mathcal{H}^2} \left(\partial_m \Delta V_l^{(2)'} + \partial_l \Delta V_m^{(2)'} \right) + \frac{1}{\mathcal{H}} \left(\phi^{(1)} \partial_l \partial_m \psi^{(2)'} \right) \\
& + \frac{1}{\mathcal{H}} \left(\phi^{(1)'} \partial_l \partial_m \phi^{(2)} \right) + \frac{1}{\mathcal{H}^2} \left(\phi^{(1)'} \partial_l \partial_m \psi^{(2)'} \right) + 3 \left(\phi^{(1)} \partial_l \partial_m \phi^{(2)} \right) = 0. \tag{B.12}
\end{aligned}$$

Using the decomposition operator $\Delta^{-1} \partial^l \mathcal{T}_i^k$ in Eq. (A.3), we can extract the equation of motion of third vector perturbation.

C The correlation function

In the appendix, we simplify the four-point correlation function and the six-point correlation function in the power spectra of vector mode.

C.1 Four-point correlation function

The four-point function can be simplified in terms of Wick's theorem.

$$\begin{aligned}
\langle \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{p}} \Phi_{\mathbf{k}'-\mathbf{p}'} \Phi_{\mathbf{p}'} \rangle & = \langle \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{p}} \rangle \langle \Phi_{\mathbf{k}'-\mathbf{p}'} \Phi_{\mathbf{p}'} \rangle + \langle \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{k}'-\mathbf{p}'} \rangle \langle \Phi_{\mathbf{p}} \Phi_{\mathbf{p}'} \rangle + \langle \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{p}'} \rangle \langle \Phi_{\mathbf{p}} \Phi_{\mathbf{k}'-\mathbf{p}'} \rangle \\
& = \frac{(2\pi^2)^2}{|k-p|^3 |k'-q|^3} \delta(\mathbf{k}) \delta(\mathbf{k}') \mathcal{P}_\Phi(k-p) \mathcal{P}_\Phi(k'-p') \\
& + \frac{(2\pi^2)^2}{|k-p|^3 p^3} \delta(\mathbf{k}-\mathbf{p}+\mathbf{k}'-\mathbf{p}') \delta(\mathbf{p}+\mathbf{p}') \mathcal{P}_\Phi(k-p) \mathcal{P}_\Phi(p) \\
& + \frac{(2\pi^2)^2}{|k-p|^3 p^3} \delta(\mathbf{k}-\mathbf{p}+\mathbf{p}') \delta(\mathbf{p}+\mathbf{k}'-\mathbf{p}') \mathcal{P}_\Phi(k-p) \mathcal{P}_\Phi(p) \\
& = \frac{(2\pi^2)^2}{|k-p|^3 p^3} \delta(\mathbf{k}+\mathbf{k}') \left(\delta(\mathbf{p}+\mathbf{p}') + \delta(\mathbf{k}-\mathbf{p}+\mathbf{p}') \right) \mathcal{P}_\Phi(k-p) \mathcal{P}_\Phi(p), \tag{C.1}
\end{aligned}$$

where in the last line we have neglected the first term which corresponds to the bubble diagram $\int d^3p \mathcal{P}_\Phi(p)$.

C.2 Six-point correlation function

In this appendix, we use Wick's theorem to simplify the six-point correlation function in Sec. 3.2. We define $\mathcal{C}(\mathbf{k}, \mathbf{k}', \mathbf{p}, \mathbf{p}', \mathbf{q}, \mathbf{q}')$ as

$$\begin{aligned}
\mathcal{C}(\mathbf{k}, \mathbf{k}', \mathbf{p}, \mathbf{p}', \mathbf{q}, \mathbf{q}') &= \frac{1}{(2\pi^2)^3} \int d^3\mathbf{k}' \langle \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{p}-\mathbf{q}} \Phi_{\mathbf{q}} \Phi_{\mathbf{k}'-\mathbf{p}'} \Phi_{\mathbf{p}'-\mathbf{q}'} \Phi_{\mathbf{q}'} \rangle \\
&= \left[\left(\frac{1}{(k-p)^3(p-q)^3q^3} \delta(\mathbf{p} + \mathbf{p}') \delta(\mathbf{q} + \mathbf{q}') P_\Phi(k-p) P_\Phi(p-q) P_\Phi(q) \right) \right. \\
&+ \left(\frac{1}{(k-p)^3(p-q)^3q^3} \delta(\mathbf{p} + \mathbf{p}') \delta(\mathbf{q} + \mathbf{p}' - \mathbf{q}') P_\Phi(k-p) P_\Phi(p-q) P_\Phi(q) \right) \\
&+ \left(\frac{1}{(k-p)^3(p-q)^3q^3} \delta(\mathbf{p} - \mathbf{q} - \mathbf{k} - \mathbf{p}') \delta(\mathbf{q} + \mathbf{q}') P_\Phi(k-p) P_\Phi(p-q) P_\Phi(q) \right) \\
&+ \left(\frac{1}{(k-p)^3(p-q)^3q^3} \delta(\mathbf{p} - \mathbf{q} + \mathbf{q}') \delta(\mathbf{q} - \mathbf{k} - \mathbf{p}') P_\Phi(k-p) P_\Phi(p-q) P_\Phi(q) \right) \\
&+ \left(\frac{1}{(k-p)^3(p-q)^3q^3} \delta(\mathbf{p} - \mathbf{k} - \mathbf{q}') \delta(\mathbf{q} + \mathbf{p}' - \mathbf{q}') P_\Phi(k-p) P_\Phi(p-q) P_\Phi(q) \right) \\
&\left. + \left(\frac{1}{(k-p)^3(p-q)^3q^3} \delta(-\mathbf{k} + \mathbf{p} - \mathbf{q}') \delta(\mathbf{q} - \mathbf{k} - \mathbf{p}') P_\Phi(k-p) P_\Phi(p-q) P_\Phi(q) \right) \right]. \tag{C.2}
\end{aligned}$$

Here, we have neglected the bubble diagram in the Wick's expansion.

D The power spectra of vector modes

In this appendix, we calculate the power spectra of second order and third order vector modes. The explicit expressions of the power spectra will be given in this appendix.

D.1 The power spectrum of second order vector mode

Substituting Eq. (C.1) into Eq. (2.12) and integrating over \mathbf{k}' and \mathbf{p}' , we obtain

$$P_V^{(2)} = \sum_{a=1}^2 \mathcal{P}_a^{(2)} = \mathcal{P}_1^{(2)} + \mathcal{P}_2^{(2)}, \tag{D.1}$$

where the summation of a comes from the Wick's expression in the last line of Eq. (C.1). More precisely, substituting the first term in the last line of Eq. (C.1) into Eq. (2.12) and

integrating over \mathbf{k}' and \mathbf{p}' , we obtain

$$\begin{aligned}
\mathcal{P}_1^{(2)} &= k^3 \pi^2 \int \frac{d^3 p}{(2\pi)^3} \frac{1}{|k-p|^3 p^3} \delta_{\lambda\lambda'} \frac{k^s e^{\lambda,r}(\mathbf{k})}{k^2} \frac{k^l e^{\lambda',m}(\mathbf{k})}{k^2} p_r p_s p_l p_m \left(I_V^{(2)}(|k-p|, p, \eta) \right)^2 \mathcal{P}_\Phi(k-p) \mathcal{P}_\Phi(p) \\
&= k^3 \pi^2 \int \frac{d^3 p}{(2\pi)^3} \frac{1}{|k-p|^3 p^3} \frac{k^s k^l (\delta^{mr} - \frac{k^m k^r}{k^2})}{k^4} p_r p_s p_l p_m \left(I_V^{(2)}(|k-p|, p, \eta) \right)^2 \mathcal{P}_\Phi(k-p) \mathcal{P}_\Phi(p) \\
&= \pi^2 k^3 \int \frac{p^2 dp \sin \theta d\theta d\phi}{(2\pi)^3} \left\{ \frac{1}{|k-p|^3 p^3} \left(\frac{(k \cdot p)^2 (p^2 - \frac{(k \cdot p)^2}{k^2})}{k^4} \right) \left(I_V^{(2)}(|k-p|, p, \eta) \right)^2 \mathcal{P}_\Phi(k-p) \mathcal{P}_\Phi(p) \right\} \\
&= -\frac{1}{4} \int dp d \cos \theta \left\{ \frac{k^3}{|k-p|^3 p} \left(\frac{(k \cdot p)^2 (p^2 - \frac{(k \cdot p)^2}{k^2})}{k^4} \right) \left(I_V^{(2)}(|k-p|, p, \eta) \right)^2 \mathcal{P}_\Phi(k-p) \mathcal{P}_\Phi(p) \right\} \\
&= \frac{1}{4} \int_0^\infty dv \int_{|1-v|}^{1+v} du \left(\frac{(1+v^2-u^2) (4v^2 - (1+v^2-u^2)^2)}{16u^2 v^2} \right) \left(k I_V^{(2)}(|k-p|, p, \eta) \right)^2 \mathcal{P}_\Phi(ku) \mathcal{P}_\Phi(kv), \tag{D.2}
\end{aligned}$$

where in the second line we have used the property of the polarization vectors $\sum_\lambda e^{\lambda,m}(\mathbf{k}) e^{\lambda,r}(\mathbf{k}) + \frac{k^m k^r}{k^2} = \delta^{mr}$. In the last line, we have used the relations $|\mathbf{k} - \mathbf{p}| = uk$, $|\mathbf{p}| = vk$, and $\cos \theta = \frac{1+v^2-u^2}{2v}$. Substituting the second term in the last line of Eq. (C.1) into Eq. (2.12) and integrating over \mathbf{k}' and \mathbf{p}' , we obtain

$$\begin{aligned}
\mathcal{P}_2^{(2)} &= k^3 \pi^2 \int \frac{d^3 p}{(2\pi)^3} \frac{1}{|k-p|^3 p^3} \delta_{\lambda\lambda'} \frac{k^s e^{\lambda,r}(\mathbf{k})}{k^2} \frac{k^l e^{\lambda',m}(\mathbf{k})}{k^2} p_r p_s (p_l - k_l) (p_m - k_m) \\
&\quad \times I_V^{(2)}(|k-p|, p, \eta) I_V^{(2)}(p, |k-p|, \eta) \mathcal{P}_\Phi(k-p) \mathcal{P}_\Phi(p) \\
&= k^3 \pi^2 \int \frac{d^3 p}{(2\pi)^3} \frac{1}{|k-p|^3 p^3} \frac{(k \cdot p) (k \cdot p - k^2) (p^2 - \frac{(k \cdot p)^2}{k^2})}{k^4} \\
&\quad \times I_V^{(2)}(|k-p|, p, \eta) I_V^{(2)}(p, |k-p|, \eta) \mathcal{P}_\Phi(k-p) \mathcal{P}_\Phi(p) \\
&= \frac{1}{4} \int_0^\infty dv \int_{|1-v|}^{1+v} du \left(\frac{\left((1+v^2-u^2)^2 - 2(1+v^2-u^2) \right) (4v^2 - (1+v^2-u^2)^2)}{16v^2 u^2} \right) \\
&\quad \times k^2 I_V^{(2)}(|k-p|, p, \eta) I_V^{(2)}(p, |k-p|, \eta) \mathcal{P}_\Phi(ku) \mathcal{P}_\Phi(kv). \tag{D.3}
\end{aligned}$$

D.2 The power spectrum of third order vector mode

The calculations of the power spectrum of third order vector mode are similar to the second order power spectrum. The integral and measure in Eq. (3.26) are much more difficult to calculate, compared with the second order power spectrum. As we mentioned in Sec. 3.2, the variables \mathbf{p}' and \mathbf{q}' can be integrated in terms of the three dimensional delta functions in Eq. (C.2). Then, the integral and measure in Eq. (3.26) in spherical coordinate system can be written as

$$\int d^3 p \int d^3 q = \int p^2 dp d(\cos \theta_p) d\phi_p \int q^2 dq d(\cos \theta_q) d\phi_q. \tag{D.4}$$

We set $\mathbf{k}/|\mathbf{k}|=(0, 0, 1)$, then $\cos \theta_p$, $\cos \theta_q$, $\cos \phi_p$, and ϕ_q in Eq. (D.4) take the form of

$$\begin{aligned}
\cos \phi_p &= X = \frac{(\mathbf{p} - \frac{1}{k^2} \mathbf{k}(\mathbf{k} \cdot \mathbf{p})) \cdot (\mathbf{q} - \frac{1}{k^2} \mathbf{k}(\mathbf{k} \cdot \mathbf{q}))}{|\mathbf{p} - \frac{1}{k^2} \mathbf{k}(\mathbf{k} \cdot \mathbf{p})| |\mathbf{q} - \frac{1}{k^2} \mathbf{k}(\mathbf{k} \cdot \mathbf{q})|} \\
&= \frac{\mathbf{p} \cdot \mathbf{q} - \frac{1}{k^2} (\mathbf{k} \cdot \mathbf{p})(\mathbf{k} \cdot \mathbf{q})}{\sqrt{(p^2 - \frac{1}{k^2} (\mathbf{k} \cdot \mathbf{p})^2) (q^2 - \frac{1}{k^2} (\mathbf{k} \cdot \mathbf{q})^2)}} \\
&= \frac{\frac{k^2 v^2 (1 + \bar{v}^2 - \bar{u}^2)}{2} - \frac{1}{k^2} \left(\frac{k^2 (1 + v^2 - u^2)}{2} \right) \left(\frac{k^2 (1 + \bar{v}^2 v^2 - w^2)}{2} \right)}{\sqrt{\left(v^2 k^2 - \frac{1}{k^2} \left(\frac{k^2 (1 + v^2 - u^2)}{2} \right)^2 \right) \left((\bar{v} v k)^2 - \frac{1}{k^2} \left(\frac{k^2 (1 + \bar{v}^2 v^2 - w^2)}{2} \right)^2 \right)}} \\
&= \frac{-1 + u^2 + v^2 - 2\bar{u}^2 v^2 + v^2 \bar{v}^2 + u^2 v^2 \bar{v}^2 - v^4 \bar{v}^2 + w^2 - u^2 w^2 + v^2 w^2}{\sqrt{(1 - 2u^2 + u^4 - 2v^2 - 2u^2 v^2 + v^4) (1 - 2v^2 \bar{v}^2 + v^4 \bar{v}^4 - 2w^2 - 2v^2 \bar{v}^2 w^2 + w^4)}} , \tag{D.5}
\end{aligned}$$

$$\cos \theta_p = \frac{|\mathbf{k} - \mathbf{p}|^2 - k^2 - p^2}{-2pk} = \frac{1 + v^2 - u^2}{2v} , \tag{D.6}$$

$$\cos \theta_q = \frac{|\mathbf{k} - \mathbf{q}|^2 - k^2 - q^2}{-2qk} = \frac{1 + (\bar{v}v)^2 - w^2}{2\bar{v}v} , \tag{D.7}$$

$$\int d\phi_q = 2\pi , \tag{D.8}$$

where the variables u , v , \bar{u} , \bar{v} , and w are defined in Sec. 3.1. Substituting Eq. (D.5)–Eq. (D.8) and Eq. (C.2) into Eq. (3.26), we can obtain Eq. (3.27).

E The expressions of $\mathbb{P}^{ij}(\eta, \mathbf{k})$

Explicit expressions of \mathbb{P}_a^{ij} in Eq. (3.36) are shown as follows,

$$\begin{aligned}
\mathbb{P}_1^{11} &= \mathbb{P}^{11}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{11}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, \mathbf{q} - \mathbf{p}) \\
&= -\frac{k^2}{8\tilde{k}^8} (ks^2 (v^2 + w^2) - 2) (ks^6 w^2 (w^2 - 1) (v^2 + w^2 - 1) - ks^4 (v^2 (2w^2 + 1) + 4w^4 - 2w^2) \\
&\quad + ks^2 (v^2 + 5w^2 + 2) - 2) , \tag{E.1}
\end{aligned}$$

$$\begin{aligned}
\mathbb{P}_2^{11} &= \mathbb{P}^{11}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{q} - \mathbf{p}) + \mathbb{P}^{11}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{k} + \mathbf{p}) \\
&= -\frac{k^2 (ks^2 (w^2 + 1) - 1) (ks^2 (v^2 + w^2) - 2) (ks^4 v^2 (v^2 + w^2 - 1) - ks^2 (3v^2 + w^2 - 2) + 2)}{8ks^8} , \tag{E.2}
\end{aligned}$$

$$\begin{aligned}
\mathbb{P}_3^{11} &= \mathbb{P}^{11}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{11}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{p} - \mathbf{k}) \\
&= \frac{k^2 (ks^2 (v^2 + w^2) - 2) (ks^6 v^2 w^2 (w^2 - 1) - ks^4 (v^2 (2w^2 + 1) + w^4 - 2w^2 - 1) + ks^2 (v^2 + 2w^2) - 1)}{8ks^8} , \tag{E.3}
\end{aligned}$$

$$\begin{aligned}
\mathbb{P}_1^{22} &= \mathbb{P}^{22}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{22}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, \mathbf{q} - \mathbf{p}) \\
&= \frac{k^2}{512ks^{12}v^4} \left(-4ks^4v^2 (ks^2v^2 - 4) (ks^6v^2 (3v^4 + 2v^2 (8w^2 - 5) + 16w^4 - 16w^2 + 3) \right. \\
&\quad + ks^4 (-18v^4 + v^2 (34 - 48w^2) + 4) + ks^2 (27v^2 - 8) + 4) - (ks^6v^2 (v^4 + v^2 (8w^2 - 6) + 8w^4 - 8w^2 + 1) \\
&\quad \left. + ks^4 (-6v^4 + v^2 (22 - 24w^2) + 4) + ks^2 (9v^2 - 8) + 4 \right)^2, \tag{E.4}
\end{aligned}$$

$$\begin{aligned}
\mathbb{P}_2^{22} &= \mathbb{P}^{22}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{q} - \mathbf{p}) + \mathbb{P}^{22}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{k} + \mathbf{p}) \\
&= -\frac{k^2}{512ks^{12}v^2w^2} \left(ks^{12}v^2w^2 (8v^8 + 8v^6 (9w^2 - 1) + v^4 (129w^4 - 62w^2 - 7) + 2v^2 (36w^6 - 31w^4 - 7w^2 + 4) \right. \\
&\quad + 8w^8 - 8w^6 - 7w^4 + 8w^2 - 1) + ks^{10} (-72v^8w^2 + v^6 (-438w^4 + 38w^2 + 4) + v^4 (-438w^6 + 170w^4 \\
&\quad + 74w^2) + v^2 (-72w^8 + 38w^6 + 74w^4 + 2w^2 - 4) + 4w^2 (w^4 - 1)) + ks^8 (v^6 (225w^2 - 8) \\
&\quad + 12v^4 (63w^4 - 11w^2 - 2) + v^2 (225w^6 - 132w^4 - 144w^2 + 8) - 8 (w^6 + 3w^4 - w^2 + 2)) \\
&\quad + ks^6 (4v^6 + v^4 (48 - 206w^2) + v^2 (-206w^4 + 254w^2 + 32) + 4 (w^6 + 12w^4 + 8w^2 + 8)) \\
&\quad \left. - 3ks^4 (8v^4 + v^2 (37w^2 + 24) + 8w^2 (w^2 + 3)) + 4ks^2 (9v^2 + 9w^2 - 8) + 16 \right), \tag{E.5}
\end{aligned}$$

$$\begin{aligned}
\mathbb{P}_3^{22} &= \mathbb{P}^{22}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{22}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{p} - \mathbf{k}) \\
&= -\frac{k^2}{512ks^{10}v^2 (ks^2 (v^2 + w^2 - 1) - 3)} \left(ks^{12}v^2 (v^{10} + 3v^8 (w^2 - 1) + v^6 (-37w^4 + 36w^2 + 1) \right. \\
&\quad - v^4 (w^2 - 1)^2 (79w^2 - 3) + v^2 (-32w^8 + 98w^6 - 109w^4 + 44w^2 - 2) + w^2 (8w^8 - 32w^6 + 41w^4 - 19w^2 \\
&\quad + 2)) + ks^{10} (-9v^{10} + 20v^8 + v^6 (267w^4 - 266w^2 - 1) + v^4 (234w^6 - 488w^4 + 254w^2 + 26) + v^2 (-48w^8 \\
&\quad + 134w^6 - 113w^4 + 66w^2 - 36) + 4w^2 (w^4 - 3w^2 + 2)) + ks^8 (27v^8 - 3v^6 (27w^2 + 13) + v^4 (-459w^4 \\
&\quad + 552w^2 - 66) + v^2 (81w^6 - 45w^4 - 66w^2 - 48) - 8w^6 + 12w^4 + 8w^2 + 16) + ks^6 (-27v^6 \\
&\quad + 10v^4 (11w^2 + 3) + v^2 (-79w^4 - 170w^2 + 168) + 4 (w^6 + 3w^4 - 10w^2 - 8)) \\
&\quad \left. + 12ks^4 (v^4 + 2v^2 (7w^2 - 2) - w^2 (w^2 - 2)) - 4ks^2 (9v^2 - 8) - 16 \right), \tag{E.6}
\end{aligned}$$

$$\mathbb{P}_1^{33} = \mathbb{P}^{33}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{33}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, \mathbf{q} - \mathbf{p}) = 0, \tag{E.7}$$

$$\mathbb{P}_2^{33} = \mathbb{P}^{33}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{q} - \mathbf{p}) + \mathbb{P}^{33}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{k} + \mathbf{p}) = 0, \tag{E.8}$$

$$\mathbb{P}_3^{33} = \mathbb{P}^{33}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{33}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{p} - \mathbf{k}) = 0, \tag{E.9}$$

$$\begin{aligned}
\mathbb{P}_1^{44} &= \mathbb{P}^{44}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{44}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, \mathbf{q} - \mathbf{p}) \\
&= -\frac{k^2 \left(ks^4 (v^2 - 1)^2 - 2ks^2 (v^2 + 1) + 1 \right) (ks^2 (v^2 + 1) - 1)^2}{8ks^8}, \tag{E.10}
\end{aligned}$$

$$\begin{aligned}
\mathbb{P}_2^{44} &= \mathbb{P}^{44}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{q} - \mathbf{p}) + \mathbb{P}^{44}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{k} + \mathbf{p}) \\
&= -\frac{k^2 (\text{ks}^2 (v^2 + 1) - 1) (\text{ks}^2 (w^2 + 1) - 1) (\text{ks}^4 (v^2 (w^2 + 1) + w^2 - 1) - \text{ks}^2 (v^2 + w^2) + 1)}{8\text{ks}^8},
\end{aligned} \tag{E.11}$$

$$\begin{aligned}
\mathbb{P}_3^{44} &= \mathbb{P}^{44}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{44}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{p} - \mathbf{k}) \\
&= -\frac{k^2 (\text{ks}^2 (v^2 + 1) - 1) (\text{ks}^2 (v^2 + w^2 - 2) - 2) (\text{ks}^4 (v^4 + v^2 (w^2 - 1) + w^2) - \text{ks}^2 (3v^2 + w^2 + 2) + 2)}{8\text{ks}^8},
\end{aligned} \tag{E.12}$$

$$\begin{aligned}
\mathbb{P}_1^{12} &= \mathbb{P}^{12}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{12}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, \mathbf{q} - \mathbf{p}) \\
&= \frac{k^2}{64\text{ks}^{10}v^2} (\text{ks}^2 (v^2 + w^2) - 2) (\text{ks}^8 v^2 (v^4 (w^2 + 1) + v^2 (8w^4 - 6w^2) + 8w^6 - 16w^4 + 9w^2 - 1) \\
&\quad - \text{ks}^6 (v^6 + 2v^4 (7w^2 + 3) + v^2 (32w^4 - 22w^2 - 1) - 4w^2 + 4) + \text{ks}^4 (6v^4 + v^2 (33w^2 + 9) - 8w^2 + 4) \\
&\quad + \text{ks}^2 (-9v^2 + 4w^2 + 4) - 4) ,
\end{aligned} \tag{E.13}$$

$$\begin{aligned}
\mathbb{P}_2^{12} &= \mathbb{P}^{12}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{q} - \mathbf{p}) + \mathbb{P}^{12}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{k} + \mathbf{p}) \\
&= \frac{k^2}{64\text{ks}^{10}w^2} (\text{ks}^2 (w^2 + 1) - 1) (\text{ks}^2 (v^2 + w^2) - 2) \left(\text{ks}^6 w^2 (8v^4 + 8v^2 (w^2 - 1) + (w^2 - 1)^2) \right. \\
&\quad \left. + \text{ks}^4 ((6 - 24v^2) w^2 - 6w^4 + 4) + \text{ks}^2 (9w^2 - 8) + 4) ,
\end{aligned} \tag{E.14}$$

$$\begin{aligned}
\mathbb{P}_3^{12} &= \mathbb{P}^{12}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{12}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{p} - \mathbf{k}) \\
&= \frac{k^2}{64\text{ks}^8 (\text{ks}^2 (v^2 + w^2 - 1) - 3)} (\text{ks}^2 (v^2 + w^2) - 2) (\text{ks}^8 (v^6 (w^2 + 1) + v^4 (-5w^4 + 6w^2 - 3) \\
&\quad + v^2 (-5w^6 + 9w^4 - 6w^2 + 2) + w^2 (w^2 - 2) (w^2 - 1)^2) + \text{ks}^6 (-v^6 + 2v^4 w^2 + v^2 (23w^4 - 24w^2 + 4) \\
&\quad - 4w^6 + 8w^4 - 8w^2 + 4) + \text{ks}^4 (3v^4 - 6v^2 (3w^2 + 1) + 3w^4 + 14w^2 - 4) - 4\text{ks}^2 (w^2 + 1) + 4) ,
\end{aligned} \tag{E.15}$$

$$\begin{aligned}
\mathbb{P}_1^{21} &= \mathbb{P}^{21}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{21}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, \mathbf{q} - \mathbf{p}) \\
&= \frac{k^2}{64\text{ks}^{10}v^2} (\text{ks}^{10} v^2 w^2 (v^6 + v^4 (9w^2 - 7) + v^2 (16w^4 - 22w^2 + 7) + 8w^6 - 16w^4 + 9w^2 - 1) \\
&\quad - \text{ks}^8 (v^8 + 17v^6 w^2 + v^4 (62w^4 - 38w^2 - 3) + v^2 (48w^6 - 54w^4 + 5w^2 + 6) - 4w^2 (w^2 - 1)) \\
&\quad + \text{ks}^6 (8v^6 + v^4 (67w^2 + 2) + v^2 (97w^4 - 51w^2 - 6) - 4 (2w^4 + w^2 - 2)) - \text{ks}^4 (21v^4 + v^2 (71w^2 + 2) \\
&\quad - 4 (w^4 + 5w^2 - 2)) + 2\text{ks}^2 (7v^2 - 6w^2 - 4) + 8) ,
\end{aligned} \tag{E.16}$$

$$\begin{aligned}
\mathbb{P}_2^{21} &= \mathbb{P}^{21}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{q} - \mathbf{p}) + \mathbb{P}^{21}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{k} + \mathbf{p}) \\
&= \frac{k^2}{64\text{ks}^{10}v^2} (\text{ks}^{10}v^4 (v^6 + v^4 (9w^2 - 1)) + v^2 (16w^4 - 8w^2 - 1) + 8w^6 - 8w^4 - w^2 + 1) + \text{ks}^8 v^2 (-9v^6 \\
&\quad + v^4 (4 - 55w^2) + v^2 (-56w^4 + 22w^2 + 13) - 8w^6 + 8w^4 + 11w^2 - 4) + \text{ks}^6 (29v^6 + v^4 (103w^2 - 11) \\
&\quad + v^2 (40w^4 - 22w^2 - 26) - 4w^2) + \text{ks}^4 (-35v^4 + v^2 (24 - 53w^2) + 8 (w^2 + 1)) \\
&\quad + 2\text{ks}^2 (3v^2 - 2 (w^2 + 4)) + 8) , \tag{E.17}
\end{aligned}$$

$$\begin{aligned}
\mathbb{P}_3^{21} &= \mathbb{P}^{21}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{21}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{p} - \mathbf{k}) \\
&= -\frac{k^2}{64\text{ks}^{10}v^2} (\text{ks}^{10}v^4 w^2 (v^4 + 8v^2 (w^2 - 1) + 8w^4 - 16w^2 + 7) + \text{ks}^8 v^2 (-v^6 + v^4 (1 - 15w^2) \\
&\quad + v^2 (-40w^4 + 38w^2 + 3) - 8w^6 + 16w^4 + 3w^2 - 7) + \text{ks}^6 (7v^6 + v^4 (47w^2 - 4) + v^2 (32w^4 - 38w^2 - 5) \\
&\quad - 4w^2 + 4) + \text{ks}^4 (-15v^4 + v^2 (7 - 29w^2) + 8w^2 - 4) + \text{ks}^2 (5v^2 - 4 (w^2 + 1)) + 4) , \tag{E.18}
\end{aligned}$$

$$\mathbb{P}_1^{13} = \mathbb{P}^{13}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{13}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, \mathbf{q} - \mathbf{p}) = 0 , \tag{E.19}$$

$$\mathbb{P}_2^{13} = \mathbb{P}^{13}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{q} - \mathbf{p}) + \mathbb{P}^{13}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{k} + \mathbf{p}) = 0 , \tag{E.20}$$

$$\mathbb{P}_3^{13} = \mathbb{P}^{13}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{13}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{p} - \mathbf{k}) = 0 , \tag{E.21}$$

$$\begin{aligned}
\mathbb{P}_1^{31} &= \mathbb{P}^{31}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{31}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, \mathbf{q} - \mathbf{p}) \\
&= -\frac{k^2 (\text{ks}^2 (v^2 + 1) - 1) (\text{ks}^2 (v^2 + 2w^2 - 1) - 3) (\text{ks}^4 w^2 (v^2 + w^2 - 1) - \text{ks}^2 (v^2 + 3w^2) + 2)}{16\text{ks}^8} , \tag{E.22}
\end{aligned}$$

$$\begin{aligned}
\mathbb{P}_2^{31} &= \mathbb{P}^{31}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{q} - \mathbf{p}) + \mathbb{P}^{31}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{k} + \mathbf{p}) \\
&= -\frac{k^2}{16\text{ks}^8} (\text{ks}^8 v^2 (v^6 + v^4 (3w^2 - 1) + v^2 (2w^4 - 2w^2 + 1) + w^2 - 1) - \text{ks}^6 (7v^6 + v^4 (13w^2 - 2) \\
&\quad + v^2 (4w^4 + 1) + 2w^4 - w^2 - 2) + \text{ks}^4 (17v^4 + v^2 (17w^2 + 3) + 2 (w^4 + 3w^2 - 1)) \\
&\quad - \text{ks}^2 (17v^2 + 7w^2 + 6) + 6) , \tag{E.23}
\end{aligned}$$

$$\begin{aligned}
\mathbb{P}_3^{31} &= \mathbb{P}^{31}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{31}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{p} - \mathbf{k}) \\
&= \frac{k^2}{16\text{ks}^8} (\text{ks}^8 v^2 w^2 (v^4 + 2v^2 (w^2 - 1) - 1) + \text{ks}^6 (-v^6 + v^4 (1 - 7w^2) + v^2 (-4w^4 + 4w^2 + 1) - 2w^4 \\
&\quad + 3w^2 + 1) + \text{ks}^4 (5v^4 + v^2 (11w^2 - 2) + 2w^4 + 2w^2 - 3) - \text{ks}^2 (7v^2 + 5w^2 + 1) + 3) , \tag{E.24}
\end{aligned}$$

$$\begin{aligned}
\mathbb{P}_1^{14} &= \mathbb{P}^{14}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{14}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, \mathbf{q} - \mathbf{p}) \\
&= \frac{k^2 (\text{ks}^2 (v^2 + 1) - 1) (\text{ks}^2 (v^2 + w^2) - 2) (\text{ks}^4 (v^2 (w^2 + 1) + w^2 - 1) - \text{ks}^2 (v^2 + w^2) + 1)}{8\text{ks}^8},
\end{aligned} \tag{E.25}$$

$$\begin{aligned}
\mathbb{P}_2^{14} &= \mathbb{P}^{14}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{q} - \mathbf{p}) + \mathbb{P}^{14}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{k} + \mathbf{p}) \\
&= \frac{k^2 (\text{ks}^4 (w^2 - 1)^2 - 2\text{ks}^2 (w^2 + 1) + 1) (\text{ks}^2 (w^2 + 1) - 1) (\text{ks}^2 (v^2 + w^2) - 2)}{8\text{ks}^8},
\end{aligned} \tag{E.26}$$

$$\begin{aligned}
\mathbb{P}_3^{14} &= \mathbb{P}^{14}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{14}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{p} - \mathbf{k}) \\
&= \frac{k^2}{8\text{ks}^8} (\text{ks}^2 (v^2 + w^2 - 2) - 2) (\text{ks}^2 (v^2 + w^2) - 2) (\text{ks}^4 (v^2 (w^2 + 1) + w^2 (w^2 - 1)) \\
&\quad - \text{ks}^2 (v^2 + 3w^2 + 2) + 2),
\end{aligned} \tag{E.27}$$

$$\begin{aligned}
\mathbb{P}_1^{41} &= \mathbb{P}^{41}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{41}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, \mathbf{q} - \mathbf{p}) \\
&= \frac{k^2 (\text{ks}^2 (v^2 + 1) - 1)^2 (\text{ks}^4 w^2 (v^2 + w^2 - 1) - \text{ks}^2 (v^2 + 3w^2 - 2) + 2)}{8\text{ks}^8},
\end{aligned} \tag{E.28}$$

$$\begin{aligned}
\mathbb{P}_2^{41} &= \mathbb{P}^{41}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{q} - \mathbf{p}) + \mathbb{P}^{41}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{k} + \mathbf{p}) \\
&= \frac{k^2}{8\text{ks}^8} (\text{ks}^2 (v^2 + 1) - 1) (\text{ks}^6 v^2 (v^2 - 1) (v^2 + w^2 - 1) - \text{ks}^4 (4v^4 + 2v^2 (w^2 - 1) + w^2) \\
&\quad + \text{ks}^2 (5v^2 + w^2 + 2) - 2),
\end{aligned} \tag{E.29}$$

$$\begin{aligned}
\mathbb{P}_3^{41} &= \mathbb{P}^{41}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{41}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{p} - \mathbf{k}) \\
&= -\frac{k^2 (\text{ks}^2 (v^2 + 1) - 1) (\text{ks}^6 v^2 (v^2 - 1) w^2 - \text{ks}^4 (v^4 + 2v^2 (w^2 - 1) + w^2 - 1) + \text{ks}^2 (2v^2 + w^2) - 1)}{8\text{ks}^8},
\end{aligned} \tag{E.30}$$

$$\mathbb{P}_1^{23} = -\mathbb{P}^{23}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{23}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, \mathbf{q} - \mathbf{p}) = 0, \tag{E.31}$$

$$\mathbb{P}_2^{23} = \mathbb{P}^{23}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{q} - \mathbf{p}) + \mathbb{P}^{23}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{k} + \mathbf{p}) = 0, \tag{E.32}$$

$$\mathbb{P}_3^{23} = \mathbb{P}^{23}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{23}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{p} - \mathbf{k}) = 0, \tag{E.33}$$

$$\begin{aligned}
\mathbb{P}_1^{32} &= \mathbb{P}^{32}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{32}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, \mathbf{q} - \mathbf{p}) \\
&= \frac{k^2}{128ks^{10}v^2} (ks^2(v^2 + 1) - 1) (ks^2(v^2 + 2w^2 - 1) - 3) (ks^6v^2(v^4 + 8v^2w^2 + 8w^4 - 8w^2 + 1) \\
&\quad - 2ks^4(3v^4 + v^2(12w^2 + 1) - 2) + ks^2(9v^2 - 8) + 4) ,
\end{aligned} \tag{E.34}$$

$$\begin{aligned}
\mathbb{P}_2^{32} &= \mathbb{P}^{32}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{q} - \mathbf{p}) + \mathbb{P}^{32}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{k} + \mathbf{p}) \\
&= \frac{k^2}{128ks^{10}w^2} (ks^{10}w^2(8v^8 + 8v^6(3w^2 - 1) + v^4(17w^4 - 14w^2 + 9) + 2v^2(w^6 + w^4 + 4w^2 - 4) + 2w^6 \\
&\quad - w^4 - 2w^2 + 1) - 2ks^8(28v^6w^2 + v^4(51w^4 - 11w^2 - 2) + v^2(16w^4 + w^2 + 6)w^2 + w^8 + 7w^6 - 2w^4 \\
&\quad - 3w^2 - 2) + ks^6(v^4(129w^2 - 8) + 2v^2(57w^4 - 4w^2 - 8) + 15w^6 + 34w^4 - 12w^2 - 8) \\
&\quad + 2ks^4(2v^4 + v^2(16 - 50w^2) - 18w^4 - 7w^2 + 8) + ks^2(-16v^2 + 19w^2 - 24) + 12) ,
\end{aligned} \tag{E.35}$$

$$\begin{aligned}
\mathbb{P}_3^{32} &= \mathbb{P}^{32}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{32}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{p} - \mathbf{k}) \\
&= \frac{k^2}{128ks^8(ks^2(v^2 + w^2 - 1) - 3)} (ks^{10}(v^{10} - 3v^8(w^2 + 1) + v^6(-15w^4 + 12w^2 + 3) - v^4(9w^6 - 19w^4 \\
&\quad + w^2 + 3) + v^2(2w^8 - 2w^6 + 5w^4 - 8w^2 + 2) + w^2(2w^6 - 7w^4 + 7w^2 - 2)) - ks^8(7v^8 - 2v^6(15w^2 + 7) \\
&\quad + v^4(-63w^4 + 62w^2 + 11) + 2v^2(7w^4 + 9w^2 - 7) + 2w^8 + 2w^6 - 13w^4 + 6w^2 - 4) + ks^6(15v^6 \\
&\quad - v^4(81w^2 + 5) + v^2(-39w^4 + 76w^2 - 2) + 9w^6 - 11w^4 + 10w^2 - 16) - ks^4(13v^4 + v^2(30 - 46w^2) \\
&\quad + 9w^4 + 10w^2 - 8) + 8ks^2(2v^2 + w^2 + 2) - 12) ,
\end{aligned} \tag{E.36}$$

$$\begin{aligned}
\mathbb{P}_1^{24} &= \mathbb{P}^{24}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{24}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, \mathbf{q} - \mathbf{p}) \\
&= -\frac{k^2}{64ks^{10}v^2} (ks^2(v^2 + 1) - 1)^2 (ks^6v^2(v^4 + v^2(8w^2 - 2) + 8w^4 - 8w^2 + 1) \\
&\quad + ks^4(-6v^4 + v^2(6 - 24w^2) + 4) + ks^2(9v^2 - 8) + 4) ,
\end{aligned} \tag{E.37}$$

$$\begin{aligned}
\mathbb{P}_2^{24} &= \mathbb{P}^{24}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{q} - \mathbf{p}) + \mathbb{P}^{24}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{k} + \mathbf{p}) \\
&= -\frac{k^2}{64ks^{10}v^2} (ks^2(w^2 + 1) - 1) (ks^8v^2(v^4(w^2 + 1) + v^2(8w^4 - 6w^2) + 8w^6 - 16w^4 + 9w^2 - 1) \\
&\quad - ks^6(v^6 + 2v^4(7w^2 + 3) + v^2(32w^4 - 22w^2 - 1) - 4w^2 + 4) \\
&\quad + ks^4(6v^4 + v^2(33w^2 + 9) - 8w^2 + 4) + ks^2(-9v^2 + 4w^2 + 4) - 4) ,
\end{aligned} \tag{E.38}$$

$$\begin{aligned}
\mathbb{P}_3^{24} &= \mathbb{P}^{24}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{24}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{p} - \mathbf{k}) \\
&= -\frac{k^2}{64ks^{10}v^2} (\text{ks}^2 (v^2 + w^2 - 2) - 2) (\text{ks}^8 v^2 (v^6 + 9v^4 w^2 + v^2 (16w^4 - 6w^2 - 1) + 8w^6 - 8w^4 + w^2) \\
&\quad - 2\text{ks}^6 (4v^6 + v^4 (23w^2 + 2) + v^2 (20w^4 - 3w^2 - 5) - 2w^2) + \text{ks}^4 (21v^4 + v^2 (57w^2 + 4) - 8(w^2 + 1)) \\
&\quad + \text{ks}^2 (4(w^2 + 4) - 14v^2) - 8) , \\
\end{aligned} \tag{E.39}$$

$$\begin{aligned}
\mathbb{P}_1^{42} &= \mathbb{P}^{42}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{42}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, \mathbf{q} - \mathbf{p}) \\
&= -\frac{k^2}{64ks^{10}v^2} (\text{ks}^2 (v^2 + 1) - 1)^2 (\text{ks}^6 v^2 (v^4 + v^2 (8w^2 - 2) + 8w^4 - 8w^2 + 1) \\
&\quad + \text{ks}^4 (-6v^4 + v^2 (6 - 24w^2) + 4) + \text{ks}^2 (9v^2 - 8) + 4) , \\
\end{aligned} \tag{E.40}$$

$$\begin{aligned}
\mathbb{P}_2^{42} &= \mathbb{P}^{42}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{q} - \mathbf{p}) + \mathbb{P}^{42}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{k} + \mathbf{p}) \\
&= -\frac{k^2}{64ks^{10}w^2} (\text{ks}^2 (v^2 + 1) - 1) \left(\text{ks}^8 w^2 (8v^6 + 8v^4 (w^2 - 2) + v^2 (w^2 - 3)^2 + w^4 - 1) \right. \\
&\quad \left. - \text{ks}^6 (32v^4 w^2 + 2v^2 (7w^4 - 11w^2 - 2) + w^6 + 6w^4 - w^2 + 4) + \text{ks}^4 (v^2 (33w^2 - 8) + 6w^4 + 9w^2 + 4) \right. \\
&\quad \left. + \text{ks}^2 (4v^2 - 9w^2 + 4) - 4 \right) , \\
\end{aligned} \tag{E.41}$$

$$\begin{aligned}
\mathbb{P}_3^{42} &= \mathbb{P}^{42}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{42}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{p} - \mathbf{k}) \\
&= -\frac{k^2}{64ks^8 (\text{ks}^2 (v^2 + w^2 - 1) - 3)} (\text{ks}^2 (v^2 + 1) - 1) (\text{ks}^8 (v^8 - v^6 (5w^2 + 4) + v^4 (-5w^4 + 9w^2 + 5) \\
&\quad + v^2 (w^6 + 6w^4 - 6w^2 - 2) + w^2 (w^4 - 3w^2 + 2)) + \text{ks}^6 (-4v^6 + v^4 (23w^2 + 8) + 2v^2 (w^4 - 12w^2 - 4) \\
&\quad - w^6 + 4w^2 + 4) + \text{ks}^4 (3v^4 - 2v^2 (9w^2 - 7) + 3w^4 - 6w^2 - 4) - 4\text{ks}^2 (v^2 + 1) + 4) , \\
\end{aligned} \tag{E.42}$$

$$\begin{aligned}
\mathbb{P}_1^{34} &= \mathbb{P}^{34}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{34}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, \mathbf{q} - \mathbf{p}) \\
&= \frac{k^2 (\text{ks}^2 (v^2 + 1) - 1) (\text{ks}^4 (v^4 + 1) - 2\text{ks}^2 (v^2 + 1) + 1) (\text{ks}^2 (v^2 + 2w^2 - 1) - 3)}{16ks^8} , \\
\end{aligned} \tag{E.43}$$

$$\begin{aligned}
\mathbb{P}_2^{34} &= \mathbb{P}^{34}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{q} - \mathbf{p}) + \mathbb{P}^{34}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{k} + \mathbf{p}) \\
&= \frac{k^2}{16ks^8} (\text{ks}^2 (w^2 + 1) - 1) (\text{ks}^6 (v^4 (w^2 + 1) + 2v^2 w^4 + 2w^4 - 3w^2 + 1) \\
&\quad - \text{ks}^4 (v^4 + v^2 (6w^2 + 4) + 2w^4 + 2w^2 + 1) + \text{ks}^2 (4v^2 + 5w^2 + 3) - 3) , \\
\end{aligned} \tag{E.44}$$

$$\begin{aligned}
\mathbb{P}_3^{34} &= \mathbb{P}^{34}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{34}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{p} - \mathbf{k}) \\
&= \frac{k^2}{16ks^8} (\text{ks}^2 (v^2 + w^2 - 2) - 2) (\text{ks}^6 (v^6 + 3v^4 w^2 + 2v^2 w^4 + v^2 + 2w^4 - w^2) \\
&\quad - 2\text{ks}^4 (3v^4 + v^2 (5w^2 + 2) + w^4 + 3w^2 + 1) + \text{ks}^2 (11v^2 + 7w^2 + 8) - 6) , \\
\end{aligned} \tag{E.45}$$

$$\mathbb{P}_1^{43} = \mathbb{P}^{43}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{43}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, \mathbf{q} - \mathbf{p}) = 0, \quad (\text{E.46})$$

$$\mathbb{P}_2^{43} = \mathbb{P}^{43}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{q} - \mathbf{p}) + \mathbb{P}^{43}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{k} + \mathbf{p}) = 0, \quad (\text{E.47})$$

$$\mathbb{P}_3^{43} = \mathbb{P}^{43}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{43}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{p} - \mathbf{k}) = 0. \quad (\text{E.48})$$

References

- [1] Elcio Abdalla et al. Cosmology intertwined: A review of the particle physics, astrophysics, and cosmology associated with the cosmological tensions and anomalies. *JHEAp*, 34:49–211, 2022.
- [2] N. Aghanim et al. Planck 2018 results. VI. Cosmological parameters. *Astron. Astrophys.*, 641:A6, 2020. [Erratum: *Astron. Astrophys.* 652, C4 (2021)].
- [3] Bruce A. Bassett, Shinji Tsujikawa, and David Wands. Inflation dynamics and reheating. *Rev. Mod. Phys.*, 78:537–589, 2006.
- [4] Torsten Bringmann, Pat Scott, and Yashar Akrami. Improved constraints on the primordial power spectrum at small scales from ultracompact minihalos. *Phys. Rev. D*, 85:125027, 2012.
- [5] Jing-Zhi Zhou, Xukun Zhang, Qing-Hua Zhu, and Zhe Chang. The third order scalar induced gravitational waves. *JCAP*, 05(05):013, 2022.
- [6] Kishore N. Ananda, Chris Clarkson, and David Wands. The Cosmological gravitational wave background from primordial density perturbations. *Phys. Rev. D*, 75:123518, 2007.
- [7] Kazunori Kohri and Takahiro Terada. Semianalytic calculation of gravitational wave spectrum nonlinearly induced from primordial curvature perturbations. *Phys. Rev. D*, 97(12):123532, 2018.
- [8] Ioannis Dalianis and Chris Kouvaris. Gravitational waves from density perturbations in an early matter domination era. *JCAP*, 07:046, 2021.
- [9] Daniel Baumann, Paul J. Steinhardt, Keitaro Takahashi, and Kiyotomo Ichiki. Gravitational Wave Spectrum Induced by Primordial Scalar Perturbations. *Phys. Rev. D*, 76:084019, 2007.
- [10] Keisuke Inomata and Tomohiro Nakama. Gravitational waves induced by scalar perturbations as probes of the small-scale primordial spectrum. *Phys. Rev. D*, 99(4):043511, 2019.
- [11] Christian T. Byrnes, Philippa S. Cole, and Subodh P. Patil. Steepest growth of the power spectrum and primordial black holes. *JCAP*, 06:028, 2019.
- [12] Rong-gen Cai, Shi Pi, and Misao Sasaki. Gravitational Waves Induced by non-Gaussian Scalar Perturbations. *Phys. Rev. Lett.*, 122(20):201101, 2019.
- [13] Chen Yuan, Zu-Cheng Chen, and Qing-Guo Huang. Probing primordial–black-hole dark matter with scalar induced gravitational waves. *Phys. Rev. D*, 100(8):081301, 2019.
- [14] Keisuke Inomata. Analytic solutions of scalar perturbations induced by scalar perturbations. *JCAP*, 03:013, 2021.
- [15] Misao Sasaki, Teruaki Suyama, Takahiro Tanaka, and Shuichiro Yokoyama. Primordial black holes—perspectives in gravitational wave astronomy. *Class. Quant. Grav.*, 35(6):063001, 2018.
- [16] Laila Alabidi, Kazunori Kohri, Misao Sasaki, and Yuuiti Sendouda. Observable Spectra of Induced Gravitational Waves from Inflation. *JCAP*, 09:017, 2012.
- [17] Guillem Domènech. Scalar Induced Gravitational Waves Review. *Universe*, 7(11):398, 2021.

- [18] Shyam Balaji, Joseph Silk, and Yi-Peng Wu. Induced gravitational waves from the cosmic coincidence. *JCAP*, 06(06):008, 2022.
- [19] Alba Romero-Rodriguez, Mario Martinez, Oriol Pujolàs, Mairi Sakellariadou, and Ville Vaskonen. Search for a Scalar Induced Stochastic Gravitational Wave Background in the Third LIGO-Virgo Observing Run. *Phys. Rev. Lett.*, 128(5):051301, 2022.
- [20] Teresa Hui-Ching Lu, Kishore Ananda, and Chris Clarkson. Vector modes generated by primordial density fluctuations. *Phys. Rev. D*, 77:043523, 2008.
- [21] Silvia Mollerach, Diego Harari, and Sabino Matarrese. CMB polarization from secondary vector and tensor modes. *Phys. Rev. D*, 69:063002, 2004.
- [22] Adam J. Christopherson, Karim A. Malik, and David R. Matravers. Vorticity generation at second order in cosmological perturbation theory. *Phys. Rev. D*, 79:123523, 2009.
- [23] Adam J. Christopherson, Karim A. Malik, and David R. Matravers. Estimating the amount of vorticity generated by cosmological perturbations in the early universe. *Phys. Rev. D*, 83:123512, 2011.
- [24] Teresa Hui-Ching Lu, Kishore Ananda, Christopher Clarkson, and Roy Maartens. The cosmological background of vector modes. *JCAP*, 02:023, 2009.
- [25] Shohei Saga. *The Vector Mode in the Second-order Cosmological Perturbation Theory*. PhD thesis, Nagoya U. (main), 2017.
- [26] Marc Kamionkowski, Arthur Kosowsky, and Albert Stebbins. A Probe of primordial gravity waves and vorticity. *Phys. Rev. Lett.*, 78:2058–2061, 1997.
- [27] Robert E. Smith, Ravi K. Sheth, and Roman Scoccimarro. An analytic model for the bispectrum of galaxies in redshift space. *Phys. Rev. D*, 78:023523, 2008.
- [28] Ruth Durrer. Light deflection in perturbed Friedmann universes. *Phys. Rev. Lett.*, 72:3301–3304, 1994.
- [29] Daisuke Yamauchi, Toshiya Namikawa, and Atsushi Taruya. Weak lensing generated by vector perturbations and detectability of cosmic strings. *JCAP*, 10:030, 2012.
- [30] Shohei Saga, Daisuke Yamauchi, and Kiyotomo Ichiki. Weak lensing induced by second-order vector mode. *Phys. Rev. D*, 92(6):063533, 2015.
- [31] Kouji Nakamura. Construction of gauge-invariant variables of linear metric perturbations on an arbitrary background spacetime. *PTEP*, 2013:043E02, 2013.
- [32] Cyril Pitrou, Xavier Roy, and Obinna Umeh. xPand: An algorithm for perturbing homogeneous cosmologies. *Class. Quant. Grav.*, 30:165002, 2013.