

Lectures on Introduction to Quantum Field Theory

Ghanashyam Date^{1,*}

¹*Chennai Mathematical Institute*

H1, SIPCOT IT Park, Siruseri, Kelambakkam 603103, INDIA.

Abstract

These are lecture notes of the QFT-I course I gave in an online mode at Chennai Mathematical Institute. The course focussed on the free relativistic quantum fields, their interactions in the perturbative scattering framework, standard computations of QED processes, radiative corrections at 1-loop with renormalization and an introduction to the toolbox of path integrals.

arXiv:2207.02243v1 [hep-th] 5 Jul 2022

*Electronic address: gdate@cmi.ac.in, ghanashyamdate@gmail.com

Contents

1. What is Quantum Field Theory?	6
A. Relativistic Covariance	6
B. Compatibility with quantum conditions	7
C. Manifest Covariance	9
2. The Poincare group and its representation: mass and spin/helicity	12
A. Poincare group, Lie algebra and Casimir invariants	12
B. Representations of the proper, orthochronous Poincare group:	15
C. Little groups	17
D. The Discrete Subgroups and their actions	20
3. Representations suitable for manifest covariance: “field representations”	24
A. Induced representation of Poincare from inducing representation of Lorentz	25
B. Irreducibility and field equations	27
C. Clifford Algebras and Spinor representations	28
4. Unitary representations on the space of solutions: anti-particles	32
5. Parity, time reversal and charge conjugation	39
A. Representations of the Clifford algebra and relations among them	42
B. Dirac-Majorana-Charge Conjugates	43
6. Relativistic Actions: classical fields	47
A. Variational Principle, Symmetries of the Action and Noether’s theorem	51
B. Conserved Poincare charges for scalar field solutions	53
7. Fourier decompositions of fields: collection of harmonic oscillators	57
A. Maxwell Field	59
B. Dirac Field	62
C. Interaction with source: Green’s functions	66
8. Covariance of quantum fields and relativistic causality	70
A. Poincare Covariance of Quantum fields	70

B. Space Inversion, Time Reversal, Charge Conjugation of Dirac Field	71
C. CPT theorem for Dirac Field	74
D. Relativistic Causality and the Spin-Statistics Theorem	75
9. States of Free Quantum Fields: Particles, coherence and coherent states	81
A. Particle/anti-particle wave packets	81
B. Correlation Functions and Coherence of states	82
C. Coherent States	85
1. An aside: Harmonic Oscillator States	85
2. Correlation functions	86
D. Evolution into a coherent state	90
10. Quantum fields in Scattering Phenomena	92
A. General scattering framework	94
B. Scattering with quantum fields: Heisenberg Picture	98
C. Kallen-Lehmann Representation	102
D. The S -matrix and its properties	105
E. Lehmann-Symanzik-Zimmermann Reduction of S -matrix	107
1. LSZ reduction for Klein-Gordon field	107
2. LSZ reduction for Dirac field	109
3. LSZ reduction for Maxwell field	112
11. Covariant Perturbation Theory	114
A. Normal ordering and Wick's theorem	118
B. Differential Cross-section for $2 \rightarrow n$ process	129
1. The Special case of $2 \rightarrow 2$ processes	131
12. Diagrammatic recipe for S-matrix elements	134
13. Elementary processes in Yukawa and QED: NR limit	137
14. Basic QED processes	142
A. Electron-muon processes:	143
B. Compton Scattering:	147

C. Electron-Positron Annihilation:	152
15. Numerical Estimates of Cross-sections and Applications	155
16. Radiative corrections in QED	159
A. The fermion propagator: self-energy	159
B. The photon propagator: photon self-energy	161
1. The Ward identity claim: $q^\mu \Pi_{\mu\nu}(q) = 0$:	162
C. The Vertex function: Form factors	164
D. Electric Charge and Anomalous Magnetic Moment	166
17. Radiative corrections at 1-loop: Divergences	169
A. Isolation of Divergence: Fermion Self Energy	171
B. Isolation of Divergence: Photon Self-Energy	173
C. Isolation of Divergence: Vertex function	176
D. Bremsstrahlung Cross-section to $o(\alpha)$	178
18. Treatment of Divergences:	183
A. Treatment of the IR divergences	185
B. Treatment of the UV divergences	187
1. 1-Loop Renormalization: Charge Screening and Lamb shift	189
C. The Method of Counter terms	192
19. Renormalized perturbation series	194
A. Necessary Conditions for UV divergence: Power Counting	194
B. An Example: The $(\Phi^3)_6$ theory	198
1. At 2-Loops	201
C. Renormalization with massless particles	202
20. Path Integrals in Quantum Mechanics	207
A. The Ab Initio Path Integral	207
B. Derivation From Transition Amplitude	210
C. Functional Derivative	214
D. Ground State-to-ground state Amplitude: $Z[J]$	216

E. Explicit evaluation of a path integral	217
F. Alternative Expression for $Z[J]$	221
21. Path Integrals in Quantum Field Theory	225
A. The 1-point function: Renormalization \leftrightarrow normal ordering	231
B. Path Integrals and Statistical Mechanics	233
22. Path Integrals as Generating Functionals	234
A. The Generating Functionals: $Z[J]$, $W[J]$, $\Gamma[\Phi]$	234
B. The Renormalization Group Equation	239
C. The Background Field Method	243
23. Closing Remarks	246
References	248

1. WHAT IS QUANTUM FIELD THEORY?

There is of course no short answer to this question. There are different facets of quantum fields. An aspect that is used extensively in condensed matter physics is the ‘QFT as a framework for computing processes in interacting many body systems’. Here, each of the many bodies typically has its internal energy states eg atoms/molecules at various lattice sites, localized states in periodic potentials etc. The interactions take place by making transitions among these internal states. Since the internal states are discrete, the interactions also involve discrete exchanges of energy/momentum/spin etc which is described by a *quantum field* eg., $[\Psi(\vec{r}), \Psi^\dagger(\vec{r}')] \sim \delta^3(\vec{r} - \vec{r}')$. With large number of bodies involved, a quantum field capable of indefinite number of transitions is a well suited framework.

There is another aspect which makes quantized fields *essential* when we want to describe interactions of even few bodies in a relativistically covariant manner. Special relativity imposes two crucial modifications: (a) the notion of causality is modified by the finite upper speed limit and (b) equivalence of mass and energy demands that the rest masses be also included in the energy conservation. The latter one allows annihilation and creation of ‘particles’ at the expense of energy. The framework of non-relativistic quantum mechanics is not capable of accommodating these possibilities. Why is it so?

For this, we need to be sharper about what is meant by “relativistic covariance”.

A. Relativistic Covariance

Special relativity posits that the space-time that is appropriate arena for describing physical processes is the Minkowski space-time - \mathbb{R}^4 with the flat metric. This allows the space-time to be described in terms of coordinates $x^\mu \leftrightarrow (t, x^i)$ and metric $\eta^{\mu\nu} = \text{diag}(-1, 1, 1, 1) = \eta_{\mu\nu}$. All inertial observers, use such a space-time and relate their descriptions of the experiments by using the coordinate transformations: $x'^\mu = \Lambda^\mu_\nu x^\nu + a^\mu$. The Λ^μ_ν denote the Lorentz transformations. These transformations, termed Poincare transformations, preserve the space-time metric. Under composition of transformations, these form the Poincare group.

In general, *covariance of a ‘structure’ with respect to some group* means that there is a homomorphism of that group, onto the invariance group of the structure. Here, invariance group of the structure means the group of transformations of the structure which preserves

the structure. For instance, if the ‘structure’ is a vector space, then its invariance group must be the group of linear transformations. For the quantum framework, the structure is a projective Hilbert space (Hilbert space modulo non-zero scaling) and the basic observable quantities are the transition probabilities: $|\langle\psi|\phi\rangle|^2/(\|\psi\|\|\phi\|)$. It is a theorem due to Wigner that the transformations of physical states (elements of the projective Hilbert space) preserving the transition probabilities can be represented by linear or anti-linear transformation of the Hilbert space, $\psi' = A\psi$, satisfying $\langle\psi'|\phi'\rangle = \langle\psi|\phi\rangle$ or $\langle\psi'|\phi'\rangle = \langle\phi|\psi\rangle$. The former defines A to be a *unitary* operator while the latter defines an *anti-unitary* operator. The anti-unitary operator is necessarily anti-linear as well. The only anti-unitary operator one encounters is the *time reversal* operator (more on this later). *Thus covariance requires the Hilbert space of the quantum system to carry a unitary representation of the Poincare group.*

If $g \in G \rightarrow R(g)$ denotes a group action which satisfies the composition rule: $R(g_1 \cdot g_2) = R(g_1) \cdot R(g_2)$, $R(g)$ constitutes a representation of the group G . On a Hilbert space, we denote a group action as $|\psi'\rangle := g \cdot |\psi\rangle := U(g)|\psi\rangle$. It follows that $g_1 \cdot (g_2 \cdot |\psi\rangle) = U(g_1)(U(g_2)|\psi\rangle) = U(g_1 \cdot g_2)|\psi\rangle = (g_1 \cdot g_2) \cdot |\psi\rangle$. This action also induces a natural action on the operators via $\langle\psi|g \cdot A|\psi\rangle := \langle g^{-1} \cdot \psi|A|g^{-1} \cdot \psi\rangle \leftrightarrow A' := g \cdot A := U(g)AU(g)^\dagger$. Check that (a) this is indeed a homomorphism: $(g_1 \cdot g_2) \cdot A = g_1 \cdot (g_2 \cdot A)$ and (b) $\langle\psi'|A'|\phi'\rangle = \langle\psi|A|\phi\rangle$. All algebraic relations among the operators are preserved under this group action. Incidentally, this is exactly how one defines the action of diffeomorphisms on functions on a manifold: $[g \cdot f](p) := f(g^{-1} \cdot p)$ (this is not the pull back definition). All this essentially says that covariance is implemented through a representation of the group. *Which representation?*

B. Compatibility with quantum conditions

Any *particular* quantum system is however distinguished by what Dirac called *quantum conditions* eg. $[q^i, p_j] = i\hbar\delta^i_j$. It is on the Hilbert space carrying a representation of the quantum conditions that the covariance group must be represented. To appreciate what this means, let us take the specific examples of the Galilean group and the Poincare group and try to seek unitary representations on the Hilbert space of a single particle with the above quantum conditions with, $i, j = 1, 2, 3$. Taking the generators of the infinitesimal transformations to be represented by self-adjoint operators, the Lie algebras of the two

groups have the following commutation relations:

$$\text{Galilean transformations} : x'^i = R^i_j x^j + v^i t + a^i \quad , \quad t' = t + b \quad , \quad R^i_m R^j_n \delta^{mn} = \delta^{ij};$$

Generators : P_0, P_i, C_i, M_{ij} ; with non-vanishing commutators:

$$[M_{ij}, M_{kl}] = i \left(\delta_{ik} M_{jl} - (i \leftrightarrow j) - (k \leftrightarrow l) + (i, k \leftrightarrow j, l) \right);$$

$$[M_{ij}, P_k] = i \left(\delta_{ik} P_j - \delta_{jk} P_i \right) \quad ; \quad [M_{ij}, C_k] = i \left(\delta_{ik} C_j - \delta_{jk} C_i \right);$$

$$[C_i, P_j] = i M \delta_{ij} \quad , \quad [C_i, P_0] = i P_i .$$

$$\text{Poincare transformations} : x'^\mu := \Lambda^\mu_\nu x^\nu + a^\mu \quad , \quad \Lambda^\mu_\alpha \Lambda^\nu_\beta \eta^{\alpha\beta} = \eta^{\mu\nu} , \eta = \text{diag}(-1, 1, 1, 1);$$

Generators : P_0, P_i, K_i, M_{ij} ; with non-vanishing commutators:

$$[M_{ij}, M_{kl}] = i \left(\delta_{ik} M_{jl} - (i \leftrightarrow j) - (k \leftrightarrow l) + (i, k \leftrightarrow j, l) \right);$$

$$[M_{ij}, P_k] = i \left(\delta_{ik} P_j - \delta_{jk} P_i \right) \quad ; \quad [M_{ij}, K_k] = i \left(\delta_{ik} K_j - \delta_{jk} K_i \right);$$

$$[K_i, P_j] = i \delta_{ij} P_0 \quad , \quad [K_i, P_0] = i P_i \quad \text{and,}$$

$$[K_i, K_j] = i M_{ij} .$$

Both have the same number of generators, 10, and almost the same Lie algebra except that the Galilean algebra has a ‘central extension’ denoted by the (mass) parameter, M and the ‘Galilean boost’ generators C_i commute among themselves unlike the ‘Lorentz boost’ generators K_i .

These can be expressed in terms of the basic operators q^i, p_j as follows¹.

Galilean , Poincare

$$M_{ij} := q_i p_j - q_j p_i \quad , \quad P_i := p_i \quad , \quad M_{ij} := q_i p_j - q_j p_i \quad , \quad P_i := p_i \quad ;$$

$$C_i := M q_i \quad , \quad P_0 := \frac{\vec{p} \cdot \vec{p}}{2M} \quad , \quad K_i := q_i P_0 \quad , \quad P_0 := \sqrt{\vec{p} \cdot \vec{p} + M^2} .$$

The parameter M is naturally identified with the mass of our particle system. Notice that only the boost generators and the P_0 generators are different in the two groups.

Ex: Verify that the above definitions of the generators, satisfy the respective algebras, up to factors of \hbar .

¹ Thomas Jordan, 0810.4637 and 1st reference from it. The reference 1 shows that if q^i is taken to represent position and transform under Galilean group accordingly, then the Galilean brackets suffice to fix the remaining generators, including P up to a constant. This also shows that there are no functions on the phase space which are invariant under the Galilean group apart from constant functions.

Ex: Under infinitesimal group action, the infinitesimal change in any operator is given by $\delta_{\vec{\epsilon}}A := i[A, \vec{\epsilon} \cdot \vec{T}]$, where $\vec{\epsilon}$ denotes the infinitesimal parameters and \vec{T} denotes the generators. Show that the infinitesimal changes, $\delta_{\vec{\epsilon}}q^i, \delta_{\vec{\epsilon}}p_i$ are at the most *linear* in q^i, p_i for the Galilean case but not so for the Poincare case for the K_i, P_0 .

In particular, the Poincare group action is *non-linear* on the basic operators. This is problematic for the following reason. At the classical level, the observables are sufficiently smooth functions on the phase space and the set of all observables can be given the structure of an *algebra*: i.e. a vector space with a multiplication (product of two functions) defined. A similar feature holds in the quantum case. Observables are self-adjoint operators. The set of observables too can be given the structure of an algebra (product of two operators). In both cases, the algebras can be thought of as being generated by the basic observables. The action of a group on the basic variables induces a corresponding action on all operators. There are domain issues in the quantum case, but much more importantly, the multiplication is non-commutative which leads to the ordering issues.

C. Manifest Covariance

When the basic observables transform non-linearly, it is much harder to deduce the transformation properties of the other operators. This is compounded further by the non-commutativity in the quantum case. When the transformations of basic variables are linear, the non-commutativity issue exists but is usually easier to manage. With this in mind we now put the additional requirement that the basic observables transform *linearly* i.e. in effect tensorially (modulo inhomogeneous action of translations). This is sometimes called the requirement of *manifest covariance*.

Invoking manifest covariance discards Poincare group action on the phase space of a particle. The action of the Galilean group is however permissible.

This could of course be generalized to N -particles, $i, j = 1, \dots, 3N$. We see that non-relativistic quantum theory of N -particles is not capable of implementing manifest Poincare covariance. and we can expect potential hazards in incorporating the relativistic causality and the creation/annihilation processes.

Could we not extend the quantum conditions to a relativistically covariant form?

For instance, *define* a quantum system by postulating quantum conditions as $[q^\mu, p_\nu] =$

$i\hbar\delta_{\nu}^{\mu}$. Then the Poincare generators can be taken to be $M_{\mu\nu} := q_{\mu}p_{\nu} - q_{\nu}p_{\mu}$ and $P_{\mu} := p_{\mu}$. It is trivial to check the Poincare Lie algebra. These also show that the basic observables are tensors of ranks as indicated by their index positions. This could be a model for a relativistic particle. However, now there is another problem in *quantum* theory.

From the basic commutators, we can define the creation/annihilation operators: $a^{\mu} := \frac{1}{\sqrt{2\hbar}}(q^{\mu} + i\eta^{\mu\nu}p_{\nu})$ and its adjoint $(a^{\mu})^{\dagger}$, satisfying the commutation relations $[a^{\mu}, (a^{\nu})^{\dagger}] = \eta^{\mu\nu}$. As usual, the states are labeled by the eigenvalues of the number operators and the vacuum state is defined by $a^{\mu}|0\rangle = 0 \forall \mu$. This is a Lorentz covariant form of defining the vacuum state. It follows immediately that $\langle 0|[a^0, (a^0)^{\dagger}]|0\rangle = \|(a^0)^{\dagger}|0\rangle\|^2 = \eta^{00} = -1$.

Additionally, if we were to interpret x^0 as a “time” coordinate so that p_0 is interpreted as energy, then *the energy eigenvalues must necessarily be unbounded below*. The Stone-von Neumann theorem implies that if the canonical commutation relations arise from the infinitesimal form of the Weyl relations, then the spectra of both the conjugate operators consists of all real numbers.

Thus, *the manifestly Lorentz covariant quantum conditions cannot be realized on a Hilbert space!*

Let us turn attention to systems with infinitely many degrees of freedom. The best known example is classical electrodynamics. This is also relativistically covariant (in fact, led to special relativity!). Its consequence, the electromagnetic waves, surprisingly show particulate behavior. While classically, an accelerated charge radiates waves at the frequency of its mechanical motion, in atomic systems the electrons only radiate at frequencies related by energy differences between levels. When photographic films are seen at low intensity light, the ‘light marks’ are localized. In interference experiments, the interference pattern builds up grain by grain. The photo-electric effect also shows a threshold behavior. All these are very particulate manifestations. The observation of intensity correlations, the $(g-2)$ measurement and the Lamb shift all point to going beyond classical electrodynamics. The validation by these high precision measurements support the quantum electrodynamical theory - a quantum field theory. The most recent example is of course the prediction of the Higgs particle and its discovery. These are positive arguments in favor of quantum field theoretic framework being well suited for a quantum theory in relativistic regimes.

What *is* a QFT framework? Let us keep in mind the familiar classical electromagnetic field. As an example, consider a reflective cavity in which some classical electromagnetic

field exist. This field satisfies the source free, first order in time, Maxwell equations which imply that the electric and magnetic fields satisfy a wave equation. Let us put some boundary conditions (the cavity is closed, say). We can write the general solution of the wave equation as a linear combination of an infinite set of *mode functions*, which are by themselves convenient solutions satisfying the boundary condition. Typically, this selects a set of frequencies/wavelength. The expansion coefficients encode the relative weightage of the mode functions. One can express the energy and the momentum contained in the cavity field, in terms of the expansion coefficients with the disappearance of the mode functions - their memory remains only in the frequencies/wavelengths. Classically, the expansion coefficients are complex numbers without any particular restriction (except that the total energy/momentum should be finite!). Schematically, we quantize the electromagnetic field by putting hats on the fields or equivalently on the *expansion coefficients*. The quantization procedure requires imposition of quantum conditions which make the expansion coefficients similar to creation/annihilation operators - *and we have the quanta of the cavity field*. Any exchange of energy/momentum within the cavity or with any outside environment is done in terms of these quanta, typically referred to as *photons*. The many body facet mentioned at the beginning does essentially this process in reverse.

It is useful to recall two equivalent ways of thinking about quantum harmonic oscillator. We may think of it as a single particle performing motion which is somehow “discretized” or as a collection of “quanta”, each carrying an energy $\hbar\omega$ and the ‘particle motion’ being understood as consequences of emission/absorption of these quanta. It is the latter view which is more convenient in the context of a field.

In nutshell then, *A quantum field is a means to hold or supply an arbitrary collection of quanta and all interactions are transactions of energy/momentum/charges etc in terms of these quanta.*

This view is certainly borne out in perturbative analysis but by no means the most general one.

2. THE POINCARÉ GROUP AND ITS REPRESENTATION: MASS AND SPIN/HELICITY

A. Poincaré group, Lie algebra and Casimir invariants

The Poincaré group: The Poincaré group or inhomogeneous Lorentz group is defined by the transformations of space-time coordinates,

$$x'^{\mu} = \Lambda^{\mu}_{\nu} x^{\nu} + a^{\mu}, \quad \Lambda^{\mu}_{\alpha} \Lambda^{\nu}_{\beta} \eta^{\alpha\beta} = \eta^{\mu\nu}, \quad \eta^{\mu\nu} = \text{diag}(-1, 1, 1, 1) = \eta_{\mu\nu}, \quad a^{\mu} \in \mathbb{R}^4. \quad (2.1)$$

The defining conditions on Λ^{μ}_{ν} imply that,

$$\eta_{\mu\nu} \Lambda^{\mu}_{\alpha} \Lambda^{\nu}_{\beta} = \eta_{\alpha\beta} = \Lambda_{\nu\alpha} \Lambda^{\nu}_{\beta} \Rightarrow \delta^{\alpha}_{\beta} = \Lambda_{\nu}^{\alpha} \Lambda^{\nu}_{\beta} \quad (2.2)$$

$$\therefore \Lambda_{\nu}^{\alpha} = (\Lambda^{-1})^{\alpha}_{\nu} \quad \leftrightarrow \quad (\Lambda^{-1})^{\alpha}_{\nu} = \Lambda^{\alpha}_{\nu} \quad \text{Note the index positions} \quad (2.3)$$

It is convenient to view Λ^{μ}_{ν} as matrices. It is then important to adopt a consistent convention regarding the index positions. Viewed as matrices, in the last equation, Λ_{ν}^{α} is the *left inverse* of Λ while Λ^{α}_{ν} is the *right inverse* of Λ .

The *infinitesimal Transformations* are defined by taking $\Lambda^{\mu}_{\nu} = \delta^{\mu}_{\nu} + \omega^{\mu}_{\nu}$ and $a^{\mu} = \epsilon^{\mu}$. The defining equation for Lorentz transformations then imply $\omega^{\mu\nu} = -\omega^{\nu\mu}$, $\omega^{\mu\nu} := \omega^{\mu}_{\alpha} \eta^{\alpha\nu}$. This also explains the parameter counting of 6 for the Lorentz and 4 for the translations.

Thus, the Poincaré group is a 10 parameter continuous group (actually a Lie group) with the 6 parameters in Λ constituting the *Lorentz* subgroup and the 4 parameters a^{μ} constituting the *translations* subgroup. We denote a generic element of the group as (Λ, a) .

Clearly $\det \Lambda = \pm 1$. The transformations with positive determinant are called *proper Lorentz* transformations while those with negative one are called *improper*. Furthermore, $-(\Lambda^0_0)^2 + \sum_{i=1}^3 (\Lambda^0_i)^2 = -1 \Rightarrow \Lambda^0_0 = \pm \sqrt{1 + \sum_i (\Lambda^0_i)^2}$. Those with positive Λ^0_0 are called *orthochronous* Lorentz transformations. The identity transformation, being both proper and orthochronous, the subgroup of proper, orthochronous transformations is the continuous subgroup connected to the identity. The two improper transformations with $\Lambda = \text{diag}(1, -1, -1, -1)$ and $\Lambda = \text{diag}(-1, 1, 1, 1)$ are called *space inversion* and *time reversal* transformations. It can be shown that any Lorentz transformation can be obtained from a proper, orthochronous transformation followed by space inversion and/or time reversal.

Group composition: The composition of two Poincaré transformations is defined as:

$(\Lambda_2, a_2) \cdot (\Lambda_1, a_1) = (\Lambda_2\Lambda_1, \Lambda_2a_1 + a_2)$. It follows that $(\mathbb{1}, 0)$ is the identity and $(\Lambda, a)^{-1} = (\Lambda^{-1}, -\Lambda^{-1}a)$.

Check: $(\Lambda, a) \cdot (\mathbb{1}, 0) \cdot (\Lambda, a)^{-1} = (\mathbb{1}, \Lambda b)$ i.e. $gHg^{-1} \in H \forall g$. Such a subgroup H is called an *invariant or normal subgroup*. Thus the translations subgroup is a normal subgroup of the Poincare group and the *Poincare group is said to be a semi-direct product of the Lorentz and the translations groups*.

Check: Any group element can be *uniquely* written as:

$$(\Lambda, a) = (\Lambda, 0) \cdot (\mathbb{1}, \Lambda^{-1}a) = (\mathbb{1}, a) \cdot (\Lambda, 0)$$

. While it is possible to proceed with the abstract identification of the Lie algebra via the vector fields generating the infinitesimal transformation, we can simplify by directly considering a *unitary representation* of the Poincare group. That is, a set of unitary operators $U(g) : \mathcal{H} \rightarrow \mathcal{H}$ such that (i) $U^\dagger(g)U(g) = \mathbb{1} = U(g)U(g)^\dagger$ and (ii) $U(g_2g_1) = U(g_2)U(g_1)$, $\forall g$'s $\in G$. To allow for the possibility of infinite dimensional representations, we use operators on Hilbert space. We also restrict to *irreducible representations* i.e. the representation space has no proper subspace which is invariant under the $U(g)$ operators. It is a result (from Schur's lemma) that if $U(g)$ is a unitary, irreducible representation, then the only bounded operator that commutes with all $U(g)$ operators is a multiple of the identity operator. Its converse also holds, namely, if the only operator that commutes with all the $U(g)$'s is a multiple of the identity operator, then the unitary representation is irreducible. A simple application of this result is that unitary, irreducible representations of abelian groups are one dimensional. Hence the translation subgroup has only 1-dimensional irreducible representations.

We define the infinitesimal generators as,

$$U(\Lambda, a) = U(\delta^\mu_\alpha + \omega^\mu_\alpha, \epsilon^\mu) := \mathbb{1} + \frac{i}{2}\omega_{\mu\nu}M^{\mu\nu} - i\epsilon_\mu P^\mu + o(\omega^2, \epsilon^2), \quad M, P \text{ are self-adjoint.} \quad (2.4)$$

The commutation relations among the generators are obtained from using the homomorphism property, $U(\Lambda, a)^{-1}U(\lambda, b)U(\Lambda, a) = U[(\Lambda, a)^{-1} \cdot (\lambda, b) \cdot (\Lambda, a)]$ and taking $\lambda = \mathbb{1} + \omega, b =$

ϵ . Using the definition of the generators, it follows,

$$\omega_{\mu\nu}U(\Lambda, a)^{-1}M^{\mu\nu}U(\Lambda, a) = (\Lambda^{-1}\omega\Lambda)_{\alpha\beta}M^{\alpha\beta} - 2(\Lambda^{-1}\omega a)_\alpha P^\alpha \quad (2.5)$$

$$\epsilon_\mu U(\Lambda, a)^{-1}P^\mu U(\Lambda, a) = (\Lambda^{-1}\epsilon)_\alpha P^\alpha . \text{ Using}$$

$$(\Lambda^{-1}\epsilon)_\alpha = \eta_{\alpha\sigma}(\Lambda^{-1})^\sigma{}_\mu \epsilon^\mu = (\Lambda^{-1})_\alpha{}^\mu \epsilon_\mu = \Lambda^\mu{}_\alpha \epsilon^\alpha, \quad (2.6)$$

$$(\Lambda^{-1}\omega\Lambda)_{\alpha\beta} = \omega_{\mu\nu}\Lambda^\mu{}_\alpha\Lambda^\nu{}_\beta \text{ and } (\Lambda^{-1}\omega a)_\alpha = \omega_{\mu\nu}\Lambda^\mu{}_\alpha a^\nu ;$$

$$U(\Lambda, a)^{-1}M^{\mu\nu}U(\Lambda, a) = \Lambda^\mu{}_\alpha\Lambda^\nu{}_\beta M^{\alpha\beta} - (\Lambda^\mu{}_\alpha a^\nu - \Lambda^\nu{}_\alpha a^\mu)P^\alpha, \quad (2.7)$$

$$U(\Lambda, a)^{-1}P^\mu U(\Lambda, a) = \Lambda^\mu{}_\nu P^\nu$$

The last two equations show that under the ‘adjoint’ action of the group, the generators transform as Lorentz tensors. To deduce the commutation relations, take Λ, a to be infinitesimal i.e. $U(\Lambda, a)^{\pm 1} = \mathbb{1} \pm \frac{i}{2}\omega_{\alpha\beta}M^{\alpha\beta} \mp i\epsilon_\alpha P^\alpha$. Substitution and reading off coefficients gives,

$$[M^{\mu\nu}, M^{\alpha\beta}] = i(\eta^{\mu\alpha}M^{\nu\beta} - (\mu \leftrightarrow \nu) - (\alpha \leftrightarrow \beta) + (\mu, \alpha \leftrightarrow \nu, \beta)) \quad (2.8)$$

$$[M^{\mu\nu}, P^\alpha] = i(\eta^{\mu\alpha}P^\nu - \eta^{\nu\alpha}P^\mu) \quad (2.9)$$

$$[P^\mu, P^\nu] = 0 . \quad (2.10)$$

It is convenient to introduce the notation, $K^i := M^{i0}$ and $J_i := \frac{1}{2}\epsilon_{ijk}M^{jk} \leftrightarrow M^{ij} = \epsilon^{ijk}J_k$ and also $H := P^0$. Then the Poincare commutators take the form (non-zero commutators only),

$$[J_i, J_j] = i\epsilon_{ij}{}^k J_k , [J_i, K_j] = i\epsilon_{ij}{}^l K_l \quad (2.11)$$

$$[K_i, K_j] = -i\epsilon_{ij}{}^l J_l , [J_i, P_j] = i\epsilon_{ij}{}^k P_k \quad (2.12)$$

$$[K_i, P_j] = -iH\delta_{ij} , [K_i, H] = iP_i \quad (2.13)$$

Define the *Pauli-Lubanski vector*, $W_\mu := \frac{1}{2}\epsilon_{\mu\nu\alpha\beta}M^{\nu\alpha}P^\beta$, $\epsilon_{0123} := 1$. It follows that $[P^\mu, W^\nu] = 0$, $[M^{\mu\nu}, W^\lambda] = i(\eta^{\mu\alpha}W^\nu - \eta^{\nu\alpha}W^\mu)$ and $[W_\mu, W_\nu] = -i\epsilon_{\mu\nu\alpha\beta}W^\alpha P^\beta$. These relations imply that $P^2 := \eta_{\mu\nu}P^\mu P^\nu$, $W^2 := \eta^{\mu\nu}W_\mu W_\nu$ commute with all the generators of the Poincare group and hence also with all the group elements of the proper, orthochronous Poincare group. These are the two independent *Casimir invariants* of the Poincare group and must be multiples of identity operator in irreducible representations. These multiples serve to label the unitary, irreducible representations.

B. Representations of the proper, orthochronous Poincare group:

Each of its representations induces a representation of its Lie algebra 2.11. Since the P^μ 's commute, their simultaneous eigenvectors can be taken as a group of labels for a basis.

$$P^\mu |k^\mu, \xi\rangle = k^\mu |k^\mu, \xi\rangle \Rightarrow P^2 |k^\mu, \xi\rangle = k^2 |k^\mu, \xi\rangle, \quad k^2 = \eta_{\mu\nu} k^\mu k^\nu. \quad (2.14)$$

Here, ξ collectively labels the degenerate eigenvectors and k^2 is the value of the Casimir P^2 . Thus, first level of classification is done by the value of the Lorentz invariant k^2 . There are four classes that arise naturally:

(i) $m^2 := -k^2 > 0$ (massive), (ii) $k^2 = 0, k^\mu \neq 0$ (massless), (iii) $k^2 > 0$ (tachyonic), and (iv) $k^\mu = 0$ (vacuum). The vacuum representation is just a single vector. The tachyonic representations have not showed up in experiments yet.

- Since P^μ transforms as a Lorentz vector, so does k^μ and for $k^2 \leq 0$, the $sgn(k^0)$ distinguishes two further subclasses of the massive and the massless representations.

We restrict to the massive and the massless representations with $k^0 > 0$ as being physically relevant.

- Under a Lorentz transformation, $(\Lambda, 0)$, k^μ goes to another point on the hyperboloid $k^2 = \text{constant}$. How does the collective label ξ transform? We would like to deduce the action of $U(\Lambda, a)$ on the vectors $|k^\mu, \xi\rangle$. Since $U(\mathbb{1}, a) = \exp(-ia^\mu P_\mu)$, we already know that $U(\mathbb{1}, a)|k^\mu, \xi\rangle = \exp(-ia^\mu k_\mu)|k^\mu, \xi\rangle$ and we focus on the Lorentz transformations $U(\Lambda, 0) =: U(\Lambda)$.

The Lorentz transformation relation (2.7) imply,

$$\begin{aligned} P^\mu (U(\Lambda)|k, \xi\rangle) &= \Lambda^\mu{}_\nu U(\Lambda) P^\nu |k, \xi\rangle = (\Lambda^\mu{}_\nu k^\nu) (U(\Lambda)|k, \xi\rangle) \Rightarrow \\ U(\Lambda)|k, \xi\rangle &= \sum_{\xi'} C_{\xi, \xi'} |\Lambda k, \xi'\rangle. \end{aligned} \quad (2.15)$$

We want to find the coefficients $C_{\xi, \xi'}$ corresponding to irreducible representations of the Lorentz group. In principle, these coefficients could (and do) depend on k label, however, their number (the degree of degeneracy) must be the same over a given hyperboloid. This is essentially a continuity argument - the label k changes continuously while the degree of degeneracy is integer valued.

Let \hat{k} be some convenient, fixed vector on the $k^2 \leq 0$ hyperboloid and let $L(k)$ be some arbitrarily chosen Lorentz transformation such that $k^\mu = L(k)^\mu{}_\nu \hat{k}^\nu$. The Lorentz

transformation L depends explicitly on k (and also implicitly on \hat{k}) and also is not unique. For instance, for $\hat{k} = (M, \vec{0})$, any k can be obtained by a combination of a rotation and a boost transformation. Having made a choice of $L(k)$, define,

$$|k, \xi\rangle := N(k)U(L(k))|\hat{k}, \xi\rangle \quad (2.16)$$

Since this is a definition, the same label ξ appears on both sides. It follows,

$$U(\Lambda)|k, \xi\rangle = N(k)U(\Lambda L(k))|\hat{k}, \xi\rangle = N(k) [U(L(\Lambda k)) \cdot U((L(\Lambda k))^{-1})] U(\Lambda) \cdot U(L(k))|\hat{k}, \xi\rangle .$$

The last three factors of U combine to give $U(L^{-1}(\Lambda k)) \cdot \Lambda \cdot L(k)$, which acting on \hat{k} takes $\hat{k} \rightarrow k \rightarrow \Lambda k \rightarrow \hat{k}$ since $(\Lambda k)^\mu = [L(\Lambda k)]^\mu_\nu \hat{k}^\nu$. Denote: $W(\Lambda, k) := L^{-1}(\Lambda k) \cdot \Lambda \cdot L(k)$. It follows that $W(\Lambda, k)\hat{k} = \hat{k} \forall \Lambda$.

The Lorentz transformations, $W(\Lambda, k)$ leaving the \hat{k} invariant, form a subgroup of the Lorentz transformations. It is known as the *little group of \hat{k}* or *stability subgroup of \hat{k}* . Defining a representation $D(W)$ by, $U(W(\Lambda, k))|\hat{k}, \xi\rangle := \sum_{\xi'} D_{\xi'\xi}(W)|\hat{k}, \xi'\rangle$, we get,

$$\begin{aligned} U(\Lambda)|k, \xi\rangle &= N(k)U(L(\Lambda k)) \sum_{\xi'} D_{\xi'\xi}(W)|\hat{k}, \xi'\rangle \\ &= N(k) \sum_{\xi'} D_{\xi'\xi}(W) [U(L(\Lambda k))|\hat{k}, \xi'\rangle] \\ \therefore U(\Lambda)|k, \xi\rangle &= \frac{N(k)}{N(\Lambda k)} \sum_{\xi'} D_{\xi'\xi}(W)|\Lambda k, \xi'\rangle \end{aligned} \quad (2.17)$$

In the last equation, we have used the definition (2.16). The irreducible representations of the little group can thus be used to characterize the irreducible representations of the Lorentz group. We are not done yet, the normalization factors $N(k)$ need to be determined.

Determination of $N(k)$:

For different \hat{k}, \hat{k}' , the corresponding little groups are different. The unitarity of the Poincare (and hence of the Lorentz group) representation implies the representations of the little groups be also unitary. We can *choose* $\langle \hat{k}', \xi' | \hat{k}, \xi \rangle = \delta_{\xi, \xi'} \delta^3(\hat{k} - \hat{k}')$ and we would like this to be preserved under Lorentz transformations i.e. with the hatted k 's being replaced by their Lorentz transforms. This is possible only for a specific choice of $N(k)$ which we determine now.

Note that this requirement associates the ortho-normalization with the hyperboloids determined by $\hat{k}^2, (\hat{k}')^2$. Since a delta function is defined in the context of an integration, we are

implicitly envisaging an integration over the hyperboloids and *not* over \mathbb{R}^4 . The integration is to be Lorentz invariant. This can be inferred as follows.

$$\begin{aligned} \int_{\text{massshell}} \theta(k^0) f(k) &:= \int_{\mathbb{R}^4} d^4 k \theta(k^0) \delta(k^2 + m^2) f(k) = \int d^3 k dk^0 \theta(k^0) \delta(-(k^0)^2 + \vec{k} \cdot \vec{k} + m^2) f(k) \\ &= \int d^3 k \frac{f(\sqrt{\vec{k}^2 + m^2}, \vec{k})}{2\sqrt{\vec{k}^2 + m^2}} \quad \because \delta(f(x)) = \sum_i \frac{\delta(x - x_i)}{|f'(x_i)|}. \end{aligned} \quad (2.18)$$

Although there are two roots from the delta function of the mass shell, the $\theta(k^0)$ picks out only one term with $\omega_k := k^0 := +\sqrt{\vec{k} \cdot \vec{k} + m^2}$. The $\frac{d^3 k}{2\omega_k}$ is the *Lorentz invariant volume element (measure)* on the mass shell. Since we have,

$$f(\vec{k}) = \int_{\mathbb{R}^3} d^3 k' f(\vec{k}') \delta^3(\vec{k} - \vec{k}') = \int_{\mathbb{R}^3} \frac{d^3 k'}{2\omega_k} f(\vec{k}') \left[2\omega_k \delta^3(\vec{k} - \vec{k}') \right], \quad (2.19)$$

we identify the *invariant delta function* on the mass shell as:

$$\delta_{inv}^3(\vec{k} - \vec{k}') := 2\omega_k \delta^3(\vec{k} - \vec{k}') = 2\sqrt{\vec{k}^2 + m^2} \delta^3(\vec{k} - \vec{k}').$$

Let $k = L(k)\hat{k}$ and define $k' := L(k)\hat{k}'$ with the same $L(k)$.

$$\langle k', \xi' | k, \xi \rangle = N^*(k') N(k) \langle \hat{k}', \xi' | U^\dagger(L(k)) U(L(k)) | \hat{k}, \xi \rangle = |N(k)|^2 \delta_{\xi', \xi} \delta^3(\hat{k}' - \hat{k}).$$

We have used unitarity of $U(L(k))$. But we also know that $\delta^3(\vec{k}' - \vec{k})$ and $\delta(\hat{k}' - \hat{k})$ are related by the same Lorentz transformation. Therefore using invariant delta function we have,

$$\omega_k \delta^3(\vec{k}' - \vec{k}) = \omega_{\hat{k}} \delta^3(\hat{k}' - \hat{k}) \quad (2.20)$$

$$\therefore \langle k', \xi' | k, \xi \rangle = |N(k)|^2 \delta_{\xi', \xi} \left[\frac{k^0}{\hat{k}^0} \right] \delta^3(\vec{k}' - \vec{k}) \quad \text{and the choice,} \quad (2.21)$$

$$|N(k)| := \sqrt{\frac{\hat{k}^0}{k^0}} \Rightarrow \quad (2.22)$$

$$\langle k', \xi' | k, \xi \rangle = \delta_{\xi', \xi} \delta^3(\vec{k}' - \vec{k}) \quad \text{and,} \quad (2.23)$$

$$U(\Lambda) | k, \xi \rangle = \sqrt{\frac{(\Lambda k)^0}{k^0}} \sum_{\xi'} D_{\xi' \xi}(W(\Lambda, k)) | \Lambda k, \xi' \rangle \quad (2.24)$$

C. Little groups

It remains to determine the little groups for the various cases.

The vacuum representation ($\mathbf{k}^\mu = \mathbf{0}$) :

The little group is all of the Lorentz group. However the representation is the trivial one: $U(\Lambda) = \mathbb{1}, \forall \Lambda$.

Massive representation ($\mathbf{k}^2 = -\mathbf{m}^2$) : Choose $\hat{k}^\mu = (M, \vec{0})$. The little group is defined by $\Lambda^\mu_\nu \hat{k}^\nu = \hat{k}^\mu \Rightarrow \Lambda^0_0 = 1, \Lambda^i_0 = 0$. The defining condition on $\Lambda, \eta^{00} = \Lambda^0_0 \Lambda^0_0 \eta^{00} + (\Lambda^0_i)^2 \delta^{ii}$, gives $\Lambda^i_0 = 0$. Hence Λ is block diagonal, with $\delta^{ij} = \Lambda^i_k \Lambda^j_l \delta^{kl}$ and the little group is $SO(3)$, the group of rotations. Its unitary representations are all finite dimensional and labeled by its Casimir, $J^2 = j(j+1), j \in \mathbb{N}/2$. The half integer ones come from the double cover $SU(2)$. The $D_{\xi'\xi}$ are the usual $(2s+1)$ dimensional representations.

The representations of this class are thus labeled by ‘mass’, m and ‘spin’ s .

Massless representation ($\mathbf{k}^2 = 0$) :

Choose $\hat{k}^\mu = (k, 0, 0, k), k > 0$. It is more convenient to determine the generators of the little group. These are some linear combinations of the Lorentz generators $M^{\alpha\beta}$. Consider the commutator,

$$\begin{aligned} [\epsilon_{\alpha\beta} M^{\alpha\beta}, P^\lambda] &= i(\epsilon^\lambda_\beta P^\beta - \epsilon_\alpha^\lambda P^\alpha) \Rightarrow \\ [\epsilon_{\alpha\beta} M^{\alpha\beta}, P^\lambda] |\hat{k}, \xi\rangle &= i(\epsilon^\lambda_\beta \hat{k}^\beta - \epsilon_\alpha^\lambda \hat{k}^\alpha) |\hat{k}, \xi\rangle = 2i\epsilon^\lambda_\alpha \hat{k}^\alpha. \end{aligned}$$

The l.h.s. of the above equation vanishes because, the generators of the little group, $\epsilon \cdot M$ leave the eigenstate of \hat{k} invariant. The r.h.s. Then gives the conditions on ϵ 's. Explicitly, $0 = \epsilon^\lambda_\alpha \hat{k}^\alpha = (\epsilon^\lambda_0 + \epsilon^\lambda_3)k \Rightarrow \epsilon_{\alpha 0} + \epsilon_{\alpha 3} = 0$. This in turn gives $\epsilon_{03} = 0, \epsilon_{01} = \epsilon_{13}, \epsilon_{02} = \epsilon_{23}$. Hence,

$$\begin{aligned} \frac{1}{2} \epsilon_{\alpha\beta} M^{\alpha\beta} |_{\text{little}} &= \epsilon_{12} M^{12} + \epsilon_{13} (M^{13} - M^{01}) + \epsilon_{23} (M^{23} - M^{02}) \\ &:= \epsilon_1 (J_1 - K_2) + \epsilon_2 (-J_2 - K_2) + \epsilon_3 J_3 \end{aligned} \quad (2.25)$$

It is conventional to denote: $A := J_2 + K_1, B := -J_1 + K_2$ and write the little algebra of $\hat{k} = k(1, 0, 0, 1)$ as,

$$[A, B] = 0, [J_3, A] = iB, [J_3, B] = -iA. \quad (2.26)$$

This is the Euclidean group in 2 dimensions: two translations and one rotation. Its representations are not as familiar. To study these, define $R(\theta) := \exp i\theta J_3$. It is easy to check that $R(\theta) A R^{-1}(\theta) = A \cos(\theta) - B \sin(\theta)$ and $R(\theta) B R^{-1}(\theta) = A \sin(\theta) + B \cos(\theta)$.

Consider a representation D in which $A|\hat{k}, a, b\rangle = a|\hat{k}, a, b\rangle$ and $B|\hat{k}, a, b\rangle = b|\hat{k}, a, b\rangle$. Define $|\hat{k}, a, b, \theta\rangle := R(\theta)|\hat{k}, a, b\rangle$. It follows that,

$$A|\hat{k}, a, b, \theta\rangle = (a \cos \theta - b \sin \theta) |\hat{k}, a, b, \theta\rangle, \quad B|\hat{k}, a, b, \theta\rangle = (a \sin \theta + b \cos \theta) |\hat{k}, a, b, \theta\rangle.$$

Hence, if a, b are non-zero, then for each θ , the states $|\hat{k}, a, b, \theta\rangle$ are simultaneous eigenstates of A and B . Such a representation has the continuous parameter θ and no such additional parameter is seen in observations. To avoid such a parameter, we *restrict the representation of the Little group to those for which $a = b = 0$ or A, B vanish in the representation*. The little group then effectively reduces to just $U(1)$ generated by J_3 . Its irreducible representations are labeled by *half integers*. We denote these representations as, $J_3|\hat{k}, \sigma\rangle = \sigma|\hat{k}, \sigma\rangle, \sigma \in \frac{1}{2}\mathbb{Z}$. The label σ is called *helicity*. The one dimensional representation is denoted as $D_{\sigma'\sigma}$. The representations with a, b nonzero, are called “continuous spin representations”.

In summary:

$$|k, \sigma\rangle := \sqrt{\frac{\hat{k}^0}{k^0}} U(L(k)) |\hat{k}, \sigma\rangle \quad (2.27)$$

For $m > 0$:

$$U(\Lambda)|k, \sigma\rangle = \sqrt{\frac{(\Lambda k)^0}{k^0}} \sum_{\sigma'} D_{\sigma\sigma'}^j [W(\Lambda, k)] |\Lambda k, \sigma'\rangle \quad \text{where,}$$

$$W(\Lambda, k) := L^{-1}(\Lambda k) \cdot \Lambda \cdot L(k) \quad , \quad L(k)\hat{k} := k \quad \sigma \text{ is the } j_3 \text{ eigenvalue} \quad (2.28)$$

For $m = 0$:

$$U(\Lambda)|k, \sigma\rangle = \sqrt{\frac{(\Lambda k)^0}{k^0}} \exp\{i\sigma\theta(\Lambda, k)\} |\Lambda k, \sigma\rangle \quad \text{where, } \sigma \text{ is the helicity label,}$$

$$W(\Lambda, k) := L^{-1}(\Lambda k) \cdot \Lambda \cdot L(k) := S(\alpha(\Lambda, k), \beta(\Lambda, k)) R(\theta(\Lambda, k)) \quad (2.29)$$

Let us evaluate the second Casimir invariant of the Poincare group, W^2 . For this we evaluate $W_\mu|\hat{k}, \sigma\rangle$.

(i) $-\hat{\mathbf{k}}^2 = \mathbf{m}^2 \neq \mathbf{0}$: For $\hat{k} = (m, 0, 0, 0)$,

$W_\mu|\hat{k}, \sigma\rangle = \frac{1}{2}\epsilon_{\mu\nu\alpha 0} M^{\nu\alpha} m|\hat{k}, \sigma\rangle$. This vanishes for $\mu = 0$ and for $\mu = i$, we get $W_i|\hat{k}, \sigma\rangle = -\frac{1}{2}\epsilon_{0ijk} M^{jk} m|\hat{k}, \sigma\rangle = -mJ_i|\hat{k}, \sigma\rangle$. Therefore,

$$W^2|\hat{k}, \sigma\rangle = m^2 j(j+1)|\hat{k}, \sigma\rangle.$$

(ii) $-\hat{\mathbf{k}}^2 = \mathbf{0}$: For $\hat{k} = k(1, 0, 0, 1)$,

$W_\mu|\hat{k}, \sigma\rangle = \frac{1}{2}(\epsilon_{\mu\nu\alpha 0} + \epsilon_{\mu\nu\alpha 3} M^{\nu\alpha}) k|\hat{k}, \sigma\rangle$. This implies,

$$W_0|\hat{k}, \sigma\rangle = k\sigma|\hat{k}, \sigma\rangle \quad , \quad W_1|\hat{k}, \sigma\rangle = k(-J_1 + K_2)|\hat{k}, \sigma\rangle = 0 \quad (2.30)$$

$$W_3|\hat{k}, \sigma\rangle = -k\sigma|\hat{k}, \sigma\rangle \quad , \quad W_2|\hat{k}, \sigma\rangle = k(-J_2 - K_1)|\hat{k}, \sigma\rangle = 0. \quad (2.31)$$

Hence $W^2|\hat{k}, \sigma\rangle = [-(k\sigma)^2 + (-k\sigma)^2]|\hat{k}, \sigma\rangle = 0$, for our choice of representations.

Note that the action of the translations subgroup on the Poincare representation is given by, $U(\mathbb{1}, a)|k, \sigma\rangle = \exp(-ia^\mu k_\mu)|k, \sigma\rangle$ while that of the Lorentz group is given in the box. Since any element of the Poincare group can be uniquely written as a product of a Lorentz transformation times a translation, we have specified the full action. Since this action is determined by the choice of the unitary, irreducible representation of the little group, the Poincare representation is said to be *induced* by a representation of the little group.

These representations are taken to identify elementary quanta.

D. The Discrete Subgroups and their actions

Recall that we have two improper Lorentz transformations, the space inversion $\mathcal{P} : \Lambda^\mu_\nu = \text{diag}(1, -1, -1, -1)$ and the time reversal $\mathcal{T} : \Lambda^\mu_\nu = \text{diag}(-1, 1, 1, 1)$. All Lorentz transformations can be generated from the proper, orthochronous transformations combined with either or both of these. Clearly these transformations are order 2 i.e. $\mathcal{P}^{-1} = \mathcal{P}$ and $\mathcal{T}^{-1} = \mathcal{T}$. Let us assume that these have an action on the physical state space preserving probabilities. Then by Wigner's theorem, these are represented either as linear and unitary or anti-linear and anti-unitary operators on the Hilbert space. *Let us use the same symbols to denote the corresponding operators which should be clear from the context.*

The homomorphism property,

$$U(\Lambda, a)^{-1}U(\lambda, b)U(\Lambda, a) = U((\Lambda, a)^{-1} \cdot (\lambda, b) \cdot (\lambda, a)) \quad (2.32)$$

gives us the action of space inversion and time reversal on the proper, orthochronous transformations. For $a = 0$ and $\Lambda = \mathcal{P}$ or \mathcal{T} leads to:

$$\mathcal{P}^{-1}U(\lambda, b)\mathcal{P} = U(\mathcal{P}^{-1}\lambda\mathcal{P}, \mathcal{P}^{-1}b) \quad , \quad \mathcal{T}^{-1}U(\lambda, b)\mathcal{T} = U(\mathcal{T}^{-1}\lambda\mathcal{T}, \mathcal{T}^{-1}b). \quad (2.33)$$

The same homomorphism (2.32), together with the infinitesimal form, $U(\mathbb{1} + \omega, \epsilon) = \mathbb{1} + \frac{i}{2}\omega_{\alpha\beta}M^{\alpha\beta} - i\epsilon_\alpha P^\alpha$ for $(U(\lambda, b))$, gives us the relations:

$$U^{-1}(\Lambda)M^{\mu\nu}U(\Lambda) = \Lambda^\mu_\alpha \Lambda^\nu_\beta M^{\alpha\beta} \quad , \quad U^{-1}(\Lambda)P^\mu U(\Lambda) = \Lambda^\mu_\nu P^\nu.$$

Following the same steps as before but *not* canceling the factors of i to allow for anti-linearity, we get,

$$\mathcal{P}^{-1}(iP^0)\mathcal{P} = iP^0 \quad , \quad \mathcal{P}^{-1}(iP^i)\mathcal{P} = -iP^i \quad (2.34)$$

$$\mathcal{T}^{-1}(iP^0)\mathcal{T} = -iP^0 \quad , \quad \mathcal{T}^{-1}(iP^i)\mathcal{T} = iP^i \quad (2.35)$$

Let if possible \mathcal{P} be anti-linear. Then it *anti-commutes* with P^0 which represents the energy. This in turn means that for every positive energy state, there is a negative energy state. This conflicts with the energy being bounded below. Identical implication follows if \mathcal{T} is *linear*! Hence, *the space inversion operator must be linear and unitary while the time reversal operator must be anti-linear and anti-unitary.*

With this understood, the action of these discrete operators on the Poincare generators is given by,

$$\mathcal{P}^{-1}P^0\mathcal{P} = P^0, \quad \mathcal{P}^{-1}P^i\mathcal{P} = -P^i, \quad \mathcal{P}^{-1}J_i\mathcal{P} = J_i, \quad \mathcal{P}^{-1}K_i\mathcal{P} = -K_i \quad (2.36)$$

$$\mathcal{T}^{-1}P^0\mathcal{T} = P^0, \quad \mathcal{T}^{-1}P^i\mathcal{T} = -P^i, \quad \mathcal{T}^{-1}J_i\mathcal{T} = -J_i, \quad \mathcal{T}^{-1}K_i\mathcal{T} = K_i \quad (2.37)$$

From these defining actions, we can determine how these operators act on the unitary, irreducible representations of the Poincare group.

It is easy to see that under space inversion and time reversal, the two Casimir invariants are invariant: $\mathcal{P}^{-1}P^2\mathcal{P} = P^2$ and $\mathcal{P}^{-1}W^2\mathcal{P} = W^2$. This is obvious from the action of these operators on the generators which transform as their tensor indices indicate. The Casimir invariants are Lorentz *scalars* (not pseudo-scalars) and hence invariant under inversions and time reversal. Thus both parity and time reversal will not mix different irreducible representations. To evaluate their actions on individual vectors, we note that the irreducible representations are characterized in terms of \hat{k} vector and an eigenvalue(s) of J_3 generator.

Action of \mathcal{P} :

$m > 0$:

Here, $\hat{k} = (m, 0, 0, 0)$ and $\sigma = -s, -s+1, \dots, s-1, s$, s being the spin. Since \mathcal{P} commutes with P^0 , anti-commutes with P^i and commutes with J_3 , we see that $\mathcal{P}|\hat{k}, \sigma\rangle$ also has the same eigenvalues of the P^μ, J_3 . Hence, $\boxed{\mathcal{P}|\hat{k}, \sigma\rangle = \eta_\sigma|\hat{k}, \sigma\rangle}$, with η_σ being a phase (since the vectors are non-degenerate). Next, from the usual angular momentum algebra, we know

$$(J_1 \pm iJ_2)|\hat{k}, \sigma\rangle = \sqrt{(s \mp \sigma)(s \pm \sigma + 1)}|\hat{k}, \sigma \pm 1\rangle \Rightarrow$$

$$(J_1 \pm iJ_2)(\mathcal{P}|\hat{k}, \sigma\rangle) = \sqrt{(s \mp \sigma)(s \pm \sigma + 1)}(\mathcal{P}|\hat{k}, \sigma \pm 1\rangle) \Rightarrow \eta_\sigma = \eta_{\sigma \pm 1} .$$

Thus η_σ is independent of σ and equal ± 1 since $\mathcal{P}^2 = \mathbb{1}$. The phase η is called the *intrinsic parity* of the representation. What about action on the states $|k, \sigma\rangle$? Recall that these states are obtained as (2.27),

$$|k, \sigma\rangle = \sqrt{\frac{\hat{k}^0}{k^0}}U(L(k))|\hat{k}, \sigma\rangle, \quad k := L(k)\hat{k} \quad \text{for a chosen } L(k).$$

Acting by the matrix \mathcal{P} on the defining equation for the Lorentz transformation $L(k)$, gives $(\mathcal{P}k) = (\mathcal{P}L(k)\mathcal{P}^{-1})(\mathcal{P}\hat{k})$ which implies that $(\mathcal{P}L(k)\mathcal{P}^{-1}) = L(\mathcal{P}k)$. The homomorphism property, (2.32), then gives $\mathcal{P}^{-1}U(L(k))\mathcal{P} = U(L(\mathcal{P}k))$. Acting on $|k, \sigma\rangle$ gives,

$$\mathcal{P}|k, \sigma\rangle = \sqrt{\frac{\hat{k}^0}{k^0}}U(L(\mathcal{P}k))\eta|\hat{k}, \sigma\rangle = \eta|\mathcal{P}k, \sigma\rangle. \quad (2.38)$$

Thus \mathcal{P} acts on all vectors of the representation with the same η and thus, η is a property of the irreducible representation.

$m = 0$:

Now $\hat{k} = (k, 0, 0, k)$ and the helicity σ is the eigenvalues of the J_3 , the angular momentum direction along the momentum direction. Since \mathcal{P} changes the sign of the momentum but not of the angular momentum, a state $|\hat{k}, \sigma\rangle \propto |-\hat{k}, -\sigma\rangle$. Unlike the massive case, the direction of the reference momentum \hat{k} is changed and this makes it inconvenient to infer the action of parity on a general state. For this reason, it is useful to choose a rotation R_2 about, say, the 2-axis to bring back $-\hat{k} \rightarrow \hat{k}$. Hence consider a new operator $\mathcal{P}' := U_2\mathcal{P}$, with $U_2 := e^{-i\pi J_2}$. But U_2 also rotates J_3 (just as \hat{k}) and hence the helicity is unchanged by the U_2 operation. Hence, $\mathcal{P}'|\hat{k}, \sigma\rangle = \eta_\sigma|\hat{k}, -\sigma\rangle$. Here η_σ is some other phase factor. What about the action of \mathcal{P} on a general state $|p, \sigma\rangle$ defined in the equation (2.27)? Note that we are using $p := (|p|, \vec{p})$ instead of k to denote the general vector.

Let $R(p, \hat{k})$ be the rotation that rotates the reference \hat{k} in the direction of the general vector $p = (|\vec{p}|, \vec{p})$. Let $B(p, \hat{k})$ denote the boost along the 3-direction which changes the reference magnitude k to $|\vec{p}|$. We can thus go from \hat{k} to p by first using the boost B followed by the rotation $R(p, \hat{k}) : |p, \sigma\rangle \sim U(R(p, \hat{k})B(p, \hat{k}))|\hat{k}, \sigma\rangle$. Noting that \mathcal{P}' commutes with boosts along the 3-axis, we have,

$$\begin{aligned} U(B(p, \hat{k})) &= \mathcal{P}'U(B(p, \hat{k}))(\mathcal{P}')^{-1} = U_2\mathcal{P}B\mathcal{P}^{-1}U_2^{-1} \Rightarrow \mathcal{P}U(B(p, \hat{k})) = U_2^{-1}U(B)U_2\mathcal{P} \\ \therefore \mathcal{P}|p, \sigma\rangle &\sim U(R(p, \hat{k}))\mathcal{P}U(B(p, \hat{k}))|\hat{k}, \sigma\rangle = U(R(p, \hat{k})R_2^{-1}B) (\eta_\sigma|\hat{k}, -\sigma\rangle) \end{aligned}$$

Now, although $R(p, \hat{k})R_2^{-1}$ is a rotation that takes the 3-axis along $-\vec{p}$, its unitary representative is not quite $U(R(-\vec{p}, \hat{k}))$. It introduces additional phases. I refer you to the equation (2.6.21) in [1]. The net result is that, $\mathcal{P}|p, \sigma\rangle = \eta_\sigma \exp\{\mp i\pi\sigma\}|\mathcal{P}(p), -\sigma\rangle$. The additional phase $\mp\pi\sigma$ correlates with the sign of the 2-component of the momentum \vec{p} .

Action of \mathcal{T} :

$m > 0$:

Now \mathcal{T} commutes with P^0 , anti commutes with P^i and J_i .

$$\therefore P^i \mathcal{T} |\hat{k}, \sigma\rangle = 0, \quad P^0 \mathcal{T} |\hat{k}, \sigma\rangle = m |\hat{k}, \sigma\rangle, \quad J_3 |\hat{k}, \sigma\rangle = -\sigma |\hat{k}, \sigma\rangle.$$

The last one implies that $\mathcal{T} |\hat{k}, \sigma\rangle = \zeta_\sigma |\hat{k}, -\sigma\rangle$, where ζ is a phase factor. From the angular momentum algebra we get,

$$\begin{aligned} (J_1 \pm iJ_2) |\hat{k}, \sigma\rangle &= \sqrt{(s \mp \sigma)(s \pm \sigma + 1)} |\hat{k}, \sigma \pm 1\rangle \Rightarrow \\ (-J_1 \pm iJ_2) (\mathcal{T} |\hat{k}, \sigma\rangle) &= \sqrt{(s \mp \sigma)(s \pm \sigma + 1)} (\mathcal{T} |\hat{k}, \sigma \pm 1\rangle) \\ &= \sqrt{(s \mp \sigma)(s \pm \sigma + 1)} \zeta_{\sigma \pm 1} |\hat{k}, -\sigma \mp 1\rangle \Rightarrow -\zeta_\sigma = \zeta_{\sigma \pm 1}. \end{aligned}$$

$\zeta_{\sigma+1} = \zeta_{\sigma-1}$. Choosing the phase for say $\zeta := \zeta_{\sigma=j}$, the phase for $j-1, j-2, \dots$ alternate between $\pm\zeta$. Thus we can write the phase as $\zeta_\sigma = \zeta(-1)^{j-\sigma}$, ζ is an arbitrarily chosen phase. The phase ζ can actually be absorbed away by putting $|\hat{k}, \sigma\rangle' := \sqrt{\zeta} |\hat{k}, \sigma\rangle$. The action of \mathcal{T} on the new vector gives,

$$\mathcal{T} |\hat{k}, \sigma\rangle' = \sqrt{\zeta^*} \mathcal{T} |\hat{k}, \sigma\rangle = |\zeta| (-1)^{j-\sigma} |\hat{k}, -\sigma\rangle.$$

Note: A similar manipulation with the η phase will retain the phase in the redefined vectors and thus the intrinsic parity cannot be absorbed away. The anti-linearity of \mathcal{T} is responsible for this feature. We may retain the irrelevant phase ζ .

The action of time reversal operator on the general vectors $|k, \sigma\rangle$ proceeds in exactly the same manner as above leading to, $\mathcal{T} |k, \sigma\rangle = \zeta (-1)^{j-\sigma} |k, -\sigma\rangle$. It follows immediately that $\mathcal{T}^2 |k, \sigma\rangle = (-1)^{2j} |k, \sigma\rangle$ which distinguishes different spins.

$m = 0$:

The analysis here proceeds in an exactly same manner as that for the parity and we just note the final result as: $\boxed{\mathcal{T} |p, \sigma\rangle = \xi_\sigma \sigma \exp\{\mp i\pi\sigma\} |\mathcal{T}(p), \sigma\rangle}$. The additional phase $\mp\pi\sigma$ correlates with the sign of the 2-component of the momentum \vec{p} .

This completes the basic definitions related to the Poincare group and its unitary, irreducible representations.

These representations specify the attributes that we may assign to elementary quanta. However, these are *not* suitable for manifest covariance. Firstly, the label σ transforms covariantly only under the little group and not the full Lorentz group. Secondly, the label k is restricted to the positive hyperboloid and not the full \mathbb{R}^4 . This makes it inconvenient to take Fourier transform to link them with space-time coordinates. For this we need to construct representations on vector valued functions on the space-time. This is done next.

3. REPRESENTATIONS SUITABLE FOR MANIFEST COVARIANCE: “FIELD REPRESENTATIONS”

As noted earlier, the abstractly classified unitary, irreducible representations of the Poincare group are not suitable for manifest Lorentz covariance. For this, we begin by defining vector space of suitably chosen function and define an action (homomorphism) of the Poincare group. These will be reducible in general and irreducibility will be imposed by partial differential equation (“field equations”). To make the action unitary, requires an inner product to be defined. This will further show a ‘doubling of representations’ leading to the particle/anti-particle interpretation.

Let $\Psi^A(x)$ be a finite dimensional column vector of a suitable class of complex/real valued functions of the space-time coordinates x^μ . The defining action of the Poincare group is: $x \rightarrow gx := x_{(\Lambda,a)} = \Lambda x + a$. It induces a corresponding action on the functions which we denote as: $\Psi \rightarrow (g\Psi) := \Psi_g$,

$$\Psi_g(x) = D(h(g))\Psi(g^{-1}x) \leftrightarrow \Psi'_{(\Lambda,a)}(x) = D^A_B(h(\lambda, a))\Psi^B((\Lambda, a)^{-1}x)$$

Here, D is a finite dimensional representation of the *Lorentz group*, $h(\Lambda, a)$ is a map from the Poincare group to the Lorentz group. The above action is required to be a homomorphism, i.e. $(g\Psi)(x) = D(h(g))\Psi(g^{-1}x)$ such that

$$(g'(g\Psi))(x) = (g'g\Psi)(x) \quad \forall g, g' \in \text{Poincare and } \forall x. \quad (3.1)$$

$$\begin{aligned} l.h.s. &= D(h(g'))(g\Psi)(g'^{-1}x) = D(h(g'))D(h(g))\Psi(g^{-1}g'^{-1}x) \\ &= D(h(g')h(g))\Psi((g'g)^{-1}x). \end{aligned} \quad (3.2)$$

$$r.h.s. = D(h(g'g))\Psi(g'g)^{-1}x \quad \therefore h(g)'s \text{ must satisfy,} \quad (3.3)$$

$$D(h(g'))D(h(g)) = D(h(g'g)) \quad \forall g, g' \in \text{Poincare. Equivalently,} \quad (3.4)$$

$$\begin{aligned} D(h(g')h(g)) &= D(h(g'g)) \quad \text{which implies,} \\ h(g')h(g) &= h(g'g), \text{ or } h(g) \text{ must be a homomorphism.} \end{aligned} \quad (3.5)$$

Since our primary focus is on the Lorentz group, and $h(g) = h((\mathbb{1}, a) \cdot (\Lambda, 0)) = h((\mathbb{1}, a))h((\Lambda, 0))$, we may *choose* $h((\mathbb{1}, a)) = \mathbb{1}$. This reduces the homomorphism from Poincare into Lorentz, to a homomorphism from Lorentz to Lorentz. We can and do take, this homomorphism to the identity homomorphism. With this we write,

$$\Psi_{(\Lambda,a)}(x) := D(\Lambda)\Psi(\Lambda^{-1}(x - a)). \quad (3.6)$$

A. Induced representation of Poincare from inducing representation of Lorentz

We have thus got an *induced representation* of the Poincare group from an *inducing representation* $D(\Lambda)$ of the Lorentz group and we take D to be an irreducible representation. There is no requirement of unitarity as $\Psi^A(x)$ is *not* a quantum mechanical wave function. Additionally, it is a fact that the Lorentz group has *no non-trivial finite dimensional, unitary representations*. Our task now reduces to finding out all possible, irreducible, finite dimensional representations of the Lorentz group.

To appreciate irreducibility, consider the infinitesimal action: $(\Lambda, a) = (\mathbb{1} + \omega, \epsilon)$, $D(\Lambda) = D(\mathbb{1} + \frac{i}{2}\omega_{\alpha\beta}T^{\alpha\beta})$ where, T represent the Lorentz generators in the D representation. Then,

$$\Psi_{\mathbb{1}+\omega,\epsilon}(x) = (\mathbb{1} + \frac{i}{2}\omega \cdot T)(\Psi(x - \omega x - \epsilon)) = \Psi(x) + \frac{i}{2}\omega \cdot T\Psi(x) - (\omega^\alpha_\beta x^\beta + \epsilon^\alpha)\partial_\alpha\Psi|_x \quad (3.7)$$

This defines the Poincare generators acting on the $\Psi(x)$. Explicitly,

$$M^{\alpha\beta}\Psi := (T^{\alpha\beta})^A_B\Psi^B(x) + i(\eta^{\gamma\alpha}x^\beta - \eta^{\gamma\beta}x^\alpha)\partial_\gamma\Psi^A(x) \quad (3.8)$$

$$P_\mu\Psi := -i\partial_\mu\Psi^A(x) \quad (3.9)$$

The two Casimir invariants, P^2, W^2 acting on Ψ should evaluate to some constants to satisfy the necessary condition for irreducibility of the Poincare representation. The P^2 Casimir is easy to evaluate and gives,

$$(P^2\Psi)^A = \eta^{\mu\nu}(-i)^2\partial_\mu\partial_\nu\Psi^A = -\square\Psi^A \quad \square := -\partial_0^2 + \nabla^2. \quad (3.10)$$

For the second Casimir we have,

$$\begin{aligned} (W_\mu\Psi)^A &= \frac{1}{2}\epsilon_{\mu\nu\alpha\beta}(M^{\nu\alpha}P^\beta\Psi)^A \\ &= \frac{1}{2}\epsilon_{\mu\nu\alpha\beta} \left[(T^{\nu\alpha})^A_B + i\delta^A_B(\eta^{\gamma\nu}x^\alpha - \eta^{\gamma\alpha}x^\nu)\partial_\gamma \right] \left[-i\delta^B_C\partial^\beta \right] \Psi^C \\ &= -\frac{i}{2}\epsilon_{\mu\nu\alpha\beta}(T^{\nu\alpha})^A_B\partial^\beta\Psi^B + \epsilon_{\mu\nu\alpha\beta}x^\alpha\partial^\nu\partial^\beta\Psi^A = -\frac{i}{2}\epsilon_{\mu\nu\alpha\beta}(T^{\nu\alpha})^A_B\partial^\beta\Psi^B + \mathcal{O}(3.11) \end{aligned}$$

Note that the 2-derivative term has canceled. Then,

$$\begin{aligned} (W^2\Psi)^A &= \frac{1}{2}\epsilon^{\mu\nu'\alpha'\beta'}(M_{\nu'\alpha'}P_{\beta'})^A_B(W_\mu\Psi)^B \\ &= \frac{1}{2}\epsilon^\mu_{\nu'\alpha'\beta'} \left[(T^{\nu'\alpha'})^A_B + i\delta^A_B(\eta^{\gamma\nu'}x^{\alpha'} - \eta^{\gamma\alpha'}x^{\nu'})\partial_\gamma \right] \left[-i\delta^B_C\partial^{\beta'} \right] (W_\mu\Psi)^C \\ &= -\frac{i}{2}\epsilon^\mu_{\nu'\alpha'\beta'}(T^{\nu'\alpha'})^A_B\partial^{\beta'}(W_\mu\Psi)^B + \epsilon^\mu_{\nu'\alpha'\beta'}x^{\alpha'}\partial^{\nu'}\partial^{\beta'}(W_\mu\Psi)^A \\ &= -\frac{1}{4}\epsilon^\mu_{\nu'\alpha'\beta'}\epsilon_{\mu\nu\alpha\beta}(T^{\nu'\alpha'}T^{\nu\alpha})^A_B\partial^{\beta'}\partial^\beta\Psi^B \quad (3.12) \end{aligned}$$

Notice that $W^2\Psi = 0$ for non-one dimensional D representation.

As illustrations consider two examples: (i) D is the trivial representation of the Lorentz group and (ii) D is the defining representation of the Lorentz group.

$D(\Lambda) = 1$: Now $W^2\Psi = 0$ while $P^2\Psi = -\square\Psi = -m^2\Psi$. Thus, irreducibility requires that Ψ must satisfy $(\square - m^2)\Psi(x) = 0$ and we recognize this as the Klein-Gordon equation.

The Ψ transforms as: $\Psi_{(\Lambda,a)}(x) = \Psi(\Lambda^{-1}(x - a))$.

$D(\Lambda)$ is the defining representation: This means $\Lambda^\mu{}_\nu = \delta^\mu{}_\nu + \omega^\mu{}_\nu := \delta^\mu{}_\nu + \frac{i}{2}(\omega_{\alpha\beta}T^{\alpha\beta})^\mu{}_\nu$ and hence

$$(T^{\alpha\beta})^\mu{}_\nu := -i(\eta^{\alpha\mu}\delta^\beta{}_\nu - \eta^{\beta\mu}\delta^\alpha{}_\nu)$$

This gives,

$$(T^{\nu'\rho'}T^{\nu\alpha})^\rho{}_\sigma = - \left[\eta^{\nu'\rho}\eta^{\nu\alpha'}\delta^\alpha{}_\sigma - \eta^{\alpha'\rho}\eta^{\nu\nu'}\delta^\alpha{}_\sigma - \eta^{\nu'\rho}\eta^{\alpha\alpha'}\delta^\nu{}_\sigma + \eta^{\alpha'\rho}\eta^{\alpha\nu'}\delta^\nu{}_\sigma \right]$$

This is manifestly antisymmetric in $(\nu'\alpha')$ and $(\nu\alpha)$. Contraction with the epsilons give equal contributions and cancel the factor of 4, leading to $(W^2\Psi)^\rho = \epsilon^{\mu\rho\nu}{}_{\beta'}\epsilon_{\mu\nu\sigma\beta}\partial^{\beta\beta'}\Psi^\sigma$.

Recall the general identity:

$$\epsilon^{a_1\dots a_j a_{j+1}\dots a_n}\epsilon_{a_1\dots a_j b_{j+1}\dots b_n} = (-1)^s (n-j)! j! \delta^{[a_{j+1}} \dots \delta^{a_n]}{}_{b_{j+1}} \dots b_n, \quad (3.13)$$

where s is the index of the metric i.e. number of negative eigenvalues of the metric. In our case the metric is the Minkowski metric and $s = 1$. This gives,

$$\epsilon^{\mu\rho\nu}{}_{\beta'}\epsilon_{\mu\nu\sigma\beta} = -\epsilon^{\mu\nu\rho}{}_{\beta'}\epsilon_{\mu\nu\sigma\beta} = -(-1)2!2!\frac{1}{2}(\delta^\rho{}_\sigma\eta_{\beta'\beta} - \eta_{\beta'\sigma}\delta^\rho{}_\beta),$$

leading to,

$$(W^2\Psi)^\rho = 2(\delta^\rho{}_\sigma\eta^{\beta'\beta} - \eta^{\beta\rho}\delta^\beta{}_\sigma)\partial^2_{\beta'\beta}\Psi^\sigma = 2[\square\Psi^\rho - \partial^\rho(\partial_\sigma\Psi^\sigma)], \text{ and} \quad (3.14)$$

$$(P^2\Psi)^\rho = -\square\Psi^\rho \quad (3.15)$$

Note: We have specified the action of the Poincare group on Ψ^A . How does $\partial_\mu\Psi^A$ transform?

We can think of the derivative as a new quantity with two indices. The index A will transform by $(T^{\alpha\beta})^A{}_B$ as before while the index μ will transform by the *defining* representation of the Lorentz group, $(T^{\alpha\beta})^\mu{}_\nu = +i(\eta^{\alpha\nu}\delta^\beta{}_\mu - \eta^{\beta\nu}\delta^\alpha{}_\mu)$. In the above calculation of the second W^μ , this was missed. However it may be checked that this extra contribution vanishes in the W^2 calculation.

B. Irreducibility and field equations

The Poincare representation will be irreducible provided the Casimir invariants have fixed values. Let $P^2\Psi^\rho = -m^2\Psi^\rho$ and $W^2\Psi^\rho = C\Psi^\rho$. For non-zero mass, with spin j , we know that W^2 equals $m^2 j(j+1)$ and hence $C = m^2 j(j+1)$. Thus Ψ^ρ satisfies,

$$\square\psi^\rho = m^2\Psi^\rho \quad , \quad 2(m^2\Psi^\rho - \partial^\rho\partial\cdot\Psi) = m^2 j(j+1)\Psi^\rho$$

We have two cases to consider. The defining vector representation of the Lorentz group, splits as $1 \oplus 0$ under the rotation subgroup. Hence $j = 1$ or 0 . For $j = 1$, taking the divergence of the second equation implies $\partial\cdot\Psi = 0$. For $j = 0$, this argument gives an identity. However, vanishing of W^2 gives $\partial_\rho\partial\cdot\Psi = m^2\psi_\rho$. Differentiating by ∂_σ and anti-symmetrising in the indices gives, $\partial_\rho\Psi_\sigma = \partial_\sigma\Psi_\rho \Rightarrow \Psi_\rho = \partial_\rho\Phi$ for some scalar Φ which is defined up to a constant. This scalar satisfies $(\square - m^2)\Phi = \text{constant}$ which can be taken to be zero. Thus,

if the defining representation is to give the massive, spin one irreducible representation, Ψ^ρ must satisfy: $(\square - m^2)\Psi^\rho = 0 = \partial\cdot\Psi$. If it is to give the massive, spin zero irreducible representation, then $\Psi_\rho = \partial_\rho\Phi$ with $(\square - m^2)\Phi = 0$.

For $m = 0$ and our restriction on the representations of the little group, the Casimir conditions give, $\square\Psi_\mu = 0$ and $\partial_\mu\partial_\nu\Psi^\nu = 0$. Introduce a new field, $A_\mu := \Psi_\mu + \partial_\mu\Lambda$. It follows that,

$$\square A_\mu = 0 + \partial_\mu\square\Lambda \quad , \quad \partial_\mu\partial^\nu A_\nu = 0 + \partial_\mu\square\Lambda \quad \Rightarrow \quad \square A_\mu - \partial_\mu\partial^\nu A_\nu = 0 = \partial^\nu(\partial_\nu A_\mu - \partial_\mu A_\nu)$$

We recognize the last equality as the source free Maxwell equation, *including its gauge invariance*: $A_\mu \rightarrow A_\mu + \partial_\mu\Lambda$!

These examples show that the induced representation Ψ^A of the *Poincare group* defined through the inducing representation of the *Lorentz* subgroup, is in general *reducible*. Requiring that the Poincare Casimir invariants, evaluated on Ψ gives the values corresponding to the unitary, irreducible representation of the Poincare group, imposes *differential conditions* on Ψ eg the Klein-Gordon equation and the subsidiary conditions. Usually, these equations are proposed as the *field equations*. Here we see them arising as irreducibility conditions.

Let us return to the classification of the finite dimensional, representations of the Lorentz group.

We already know that the defining representation acts as: $v'^{\mu} = \Lambda^{\mu}_{\nu} v^{\nu} \leftrightarrow v'_{\mu} = \Lambda_{\mu}^{\nu} v_{\nu} = (\Lambda^{-1})^{\nu}_{\mu} v_{\nu}$. From these we can trivially construct other irreducible representations by taking tensor products and subtracting suitable ‘traces’. For instance, a rank-2 tensor $V^{\mu\nu}$ transforms as $V'^{\mu\nu} = \Lambda^{\mu}_{\alpha} \Lambda^{\nu}_{\beta} V^{\alpha\beta}$. This is a reducible representation though. Why? Because its symmetrised and anti-symmetrised parts transform among themselves and thus form invariant subspaces. Furthermore, tensors of the form $\eta^{\mu\nu} C$ also transforms into the same form. Hence the symmetric combination splits further into traceless and trace part. Thus we get the space \mathcal{V} of the second rank tensors decomposes as:

$$\mathcal{V} = \mathcal{V}_{anti-sym} \oplus \mathcal{V}_{sym,traceless} \oplus \mathcal{V}_{trace}, \quad \text{each being an irreducible representation.}$$

Analogous construction can be carried out for higher rank tensors. These however do *not* exhaust all the finite dimensional irreducible representations. There are the spinor representations which are missed. It is a result that *all finite dimensional representations of pseudo-orthogonal groups, $SO(p, q), p + q \geq 2$ can be constructed from the irreducible representations of the corresponding Clifford algebra.*

C. Clifford Algebras and Spinor representations

Clifford algebras are algebras generated by elements, $\{\mathbb{1}, \gamma^{\mu}, \mu = 0, \dots, (p + q - 1)\}$ satisfying $\gamma^{\mu} \gamma^{\nu} + \gamma^{\nu} \gamma^{\mu} = 2\bar{\eta}^{\mu\nu} \mathbb{1}, \bar{\eta}^{\mu\nu} = \text{diag}(-1, \dots, -1, +1, \dots, +1)$. There are p negative and q positive signs. We have put a bar on the metric since we will relate it to the η defining the Lorentz group for which $p = 1$ and $q = 3$. Note that for $\mu \neq \nu$ the gamma’s anti-commute, $(\gamma^0)^2 = \bar{\eta}^{00}$ and $(\gamma^i)^2 = \bar{\eta}^{ii}$. Since the gamma’s anti-commute, their bilinear satisfy commutation relations.

Let $\Sigma^{\mu\nu} := a(\gamma^{\mu} \gamma^{\nu} - \gamma^{\nu} \gamma^{\mu})$. We will choose the proportionality constant a suitably. Consider,

$$[\Sigma^{\mu\nu}, \Sigma^{\alpha\beta}] = a^2 \{ [\gamma^{\mu} \gamma^{\nu}, \gamma^{\alpha} \gamma^{\beta}] - (\mu \leftrightarrow \nu) - (\alpha \leftrightarrow \beta) + (\mu, \alpha \leftrightarrow \nu, \beta) \} \quad (3.16)$$

$$\begin{aligned} \gamma^{\mu} \gamma^{\nu} \gamma^{\alpha} \gamma^{\beta} &= 2\bar{\eta}^{\nu\alpha} \gamma^{\mu} \gamma^{\beta} - 2\bar{\eta}^{\mu\alpha} \gamma^{\nu} \gamma^{\beta} + 2\bar{\eta}^{\nu\beta} \gamma^{\alpha} \gamma^{\mu} - 2\bar{\eta}^{\mu\beta} \gamma^{\alpha} \gamma^{\nu} \\ \therefore \frac{1}{2} [\gamma^{\mu} \gamma^{\nu}, \gamma^{\alpha} \gamma^{\beta}] &= [-\bar{\eta}^{\mu\alpha} \gamma^{\nu} \gamma^{\beta} + \bar{\eta}^{\nu\alpha} \gamma^{\mu} \gamma^{\beta} - \bar{\eta}^{\mu\beta} \gamma^{\alpha} \gamma^{\nu} + \bar{\eta}^{\nu\beta} \gamma^{\alpha} \gamma^{\mu}] \\ \therefore [\Sigma^{\mu\nu}, \Sigma^{\alpha\beta}] &= (4a) \{ -\bar{\eta}^{\mu\alpha} \Sigma^{\nu\beta} + (\mu \leftrightarrow \nu) + (\alpha \leftrightarrow \beta) - (\mu, \alpha \leftrightarrow \nu, \beta) \} \quad (3.17) \end{aligned}$$

This has the same form as the algebra satisfied by the Lorentz generators: $[M^{\mu\nu}, M^{\alpha\beta}] = i(\eta^{\mu\alpha} M^{\nu\beta} - \dots)$, the M ’s were defined through $U(\Lambda = 1 + \omega) = \mathbb{1} + \frac{i}{2} \omega_{\mu\nu} M^{\mu\nu}$. Thus we

need $-4a\bar{\eta}^{\mu\nu} = i\eta^{\mu\nu}$. There are two ways to satisfy this and we *choose our conventions as:* $\bar{\eta} = -\eta$ and $a = \frac{i}{4}$. Thus,

$$\boxed{\gamma^\mu\gamma^\nu + \gamma^\nu\gamma^\mu = -2\eta^{\mu\nu} \quad , \quad \Sigma^{\mu\nu} := \frac{i}{4}(\gamma^\mu\gamma^\nu - \gamma^\nu\gamma^\mu) = \frac{i}{4}[\gamma^\mu, \gamma^\nu].} \quad (3.18)$$

The Σ 's satisfy the Lorenz Lie algebra. Furthermore, the definitions imply that

$$\boxed{[\Sigma^{\mu\nu}, \gamma^\lambda] = i(\eta^{\mu\lambda}\gamma^\nu - \eta^{\nu\lambda}\gamma^\mu)} \text{ i.e. the } \gamma\text{'s transform as Lorentz vectors.}$$

It is useful to define $\boxed{\gamma_5 := +i\gamma^0\gamma^1\gamma^2\gamma^3}$. Just from the Clifford algebra and definitions it follows that $\gamma_5^2 = +\mathbb{1}$, $\gamma_5\gamma^\mu = -\gamma^\mu\gamma_5$, and $[\gamma_5, \Sigma^{\mu\nu}] = 0$.

Claim: *It is always possible to choose the γ 's to be finite dimensional unitary matrices.*

This is based on the following facts: (a) The set of elements $G := \{\pm\mathbb{1}, \pm\gamma^\mu, \pm\gamma^\mu\gamma^\nu, \pm\gamma^\mu\gamma^\nu\gamma^\lambda, \dots, \pm\gamma_5\}$ with all indices distinct in the products, forms a finite group of 32 elements (in our 4 dimensional case); (b) All representations of finite groups can be made unitary; (c) representation theory of finite groups applied to the group G gives that there is exactly one, non-trivial, unitary, irreducible representation of G and hence of the Clifford algebra and that its dimension is $2^{\lfloor 4/2 \rfloor} = 4$. The results also extends to other dimensions.

Unitarity of the γ 's and their squares being $\pm\mathbb{1}$, imply further that $(\gamma^0)^\dagger = \gamma^0$ and $(\gamma^i)^\dagger = -\gamma^i$. This in turn gives $(\Sigma^{0i})^\dagger = -\Sigma^{0i}$, $(\Sigma^{ij})^\dagger = \Sigma^{ij}$ and $\gamma_5^\dagger = \gamma_5$.

Hence, the representation D of the Lorenz group provided by Σ 's is (i) *non-unitary* and (ii) *reducible*, since γ_5 is Hermitian and commutes with the generators.

Since γ 's are finite dimensional, their traces are defined and all of them including γ_5 are traceless. In particular, this implies that γ_5 has two eigenvalues equal to +1 and other two equal to -1. Hence $\frac{1\pm\gamma_5}{2}$ are projection matrices and reduce the 4 dimensional representation of the Lorenz group into two irreducible representations of dimensions 2.

By convention, $\Psi_L := \frac{1-\gamma_5}{2}\Psi$, $\leftrightarrow \gamma_5\Psi_L = -\Psi_L$ is called a *left handed Weyl spinor* while $\Psi_R := \frac{1+\gamma_5}{2}\Psi$, $\leftrightarrow \gamma_5\Psi_R = +\Psi_R$ is called a *right handed Weyl spinor*.

The 4-component Ψ^A 's are called *Dirac spinors* while the 2-component projections, $\Psi_\pm := \frac{1\pm\gamma_5}{2}\Psi$ are called *Weyl spinors*. Choosing D to be generated by the $\Sigma_\pm := \frac{1\pm\gamma_5}{2}\Sigma$ give irreducible representations of the *Lorentz group* and in turn an irreducible representation of the Poincare group .

Returning to Poincare representation, consider the combination $\gamma^\mu P_\mu$, apparently a

Lorentz scalar. Indeed, recalling that $[M^{\mu\nu}, P_\lambda] = +i(\delta_\lambda^\mu P^\nu - \delta_\lambda^\nu P^\mu)$, it follows that,

$$[T^{\mu\nu}, \gamma^\lambda P_\lambda] = [\Sigma^{\mu\nu}, \gamma^\lambda] P_\lambda + \gamma^\lambda [M^{\mu\nu}, P_\lambda] = 0 .$$

We have used T to denote a general representation, M to denote the defining representation and Σ to denote the reducible spinor representation of the Lorentz group. We use the conventional abbreviation $\not{a} := \gamma^\mu a_\mu$ for all 4-vectors a_μ .

Consider the Poincare Casimir P^2 . Irreducibility forces $P^2\Psi = -m^2\Psi$. Let $m \neq 0$. The combinations, $m\mathbb{1} \pm \not{p}$ commutes with the Lorentz generators and also satisfy, $(m \pm \not{p})^2 = m^2 + (\not{p})^2 \pm 2m\not{p} = 2(m \pm \not{p})$. Therefore, $\pi_\pm := \frac{m \pm \not{p}}{2m}$ is a projection matrix operator (since P is an operator) that also commutes with the Lorentz generators. It trivially commutes with the Poincare generators as well. Hence the Poincare representation induced by the Dirac spinor is *reducible* in yet another way, the irreducible subspaces being provided by $(\pm m + \not{p})\Psi = 0 = (-i\gamma^\mu \partial_\mu \pm m)\Psi = 0$. This is just the Dirac equation! Note that the projection property holds *only for non-zero mass*.

Note: The projectors $\frac{1 \pm \gamma_5}{2}$ reduce the D representation of the Lorentz group and each of these induces an irreducible representation of the Poincare group. By contrast, $\frac{m \pm \not{p}}{2m}$, do *not* reduce the D representation, but nevertheless reduces the Poincare representations. Are either of these Poincare representations further reduced by the other projector? The answer is ‘No’ because γ_5 anti-commutes with \not{p} and thus exchanges the two projectors.

Note: Consider the action of \not{p} on the left(right) handed Weyl spinors. It follows immediately that $\gamma_5(\not{p})\Psi_{L/R} = \pm \not{p}\Psi_{R/L}$. Therefore, if we restrict to any one irreducible Weyl representation, then \not{p} must annihilate it. That is *Weyl spinors satisfy the massless Dirac equation - also known as the Weyl equation*. Conversely, if we have a Dirac spinor Ψ that satisfies the Weyl equation, then its decomposition into its Weyl spinors also satisfy the Weyl equation individually: $\not{p}(\Psi_L + \Psi_R) = 0 \Rightarrow \not{p}\Psi_{L/R} = 0$.

Exercises:

1. Form the anti-symmetrized products of the γ -matrices, satisfying the Clifford algebra

$$\{\gamma^\mu, \gamma^\nu\} = -2\eta^{\mu\nu},$$

$$\Gamma^\mu := \gamma^\mu$$

$$\Gamma^{\mu\nu} := \gamma^\mu \gamma^\nu - \gamma^\nu \gamma^\mu$$

$$\Gamma^{\mu\nu\lambda} := \gamma^\mu \gamma^\nu \gamma^\lambda + \gamma^\nu \gamma^\lambda \gamma^\mu + \gamma^\lambda \gamma^\mu \gamma^\nu - \gamma^\nu \gamma^\mu \gamma^\lambda - \gamma^\mu \gamma^\lambda \gamma^\nu - \gamma^\lambda \gamma^\nu \gamma^\mu$$

$\Gamma^{\mu\nu\alpha\beta} := \gamma^{[\mu}\gamma^\nu\gamma^\alpha\gamma^{\beta]}$ with no numerical overall factors. Since the indices take 4 values only, we cannot have any more antisymmetric products. Together with $\mathbb{1}$, this set comprises of 16 matrices.

Show that products of any number of γ 's can be expressed in terms of the these 16 matrices together with products of η 's. Conclude that these 16 matrices constitute a basis for arbitrary products of γ 's. Since they have different Lorentz transformation properties, they are all independent. The vector space of $k \times k$ matrices has dimension k^2 , hence the minimum matrix order for the γ 's is 4 and they are necessarily irreducible representation of the Clifford algebra. Hence our γ 's are 4×4 . Analogous arguments hold for Clifford algebra of n dimensions (indices taking n values).

2. For spinors satisfying the Dirac equation with $M \neq 0$, evaluate the Pauli-Lubanski scalar, $W_\mu W^\mu \Psi$.
3. The Lorentz J_i, K_i satisfy the algebras $[J_i, J_j] = i\epsilon_{ij}^l J_l$; $[K_i, K_j] = -i\epsilon_{ij}^l K_l$; $[J_i, K_j] = i\epsilon_{ij}^l J_l$. Define: $A_i := \frac{1}{2}(J_i + iK_i), B_i := \frac{1}{2}(J_i - iK_i)$. Check that the $[A_i, B_j] = i\epsilon_{ij}^l A_l$, $[B_i, B_j] = i\epsilon_{ij}^l B_l$, $[A_i, B_j] = 0$. This the $M^{\mu\nu}$ Lie algebra is equivalent to the direct sum of two, mutually commuting $SU(2)$ algebras. They have the Casimir invariants \vec{A}^2, \vec{B}^2 . Evaluate these for the spinor representation provided by $\Sigma^{\mu\nu}$'s.

In summary:

$$\Psi_{\Lambda,a}(x) := D(\Lambda)\Psi(\Lambda^{-1}x - \Lambda^{-1}a) \text{ (Reducible) Poincare action} \quad (3.19)$$

Require Poincare Casimir invariants to take constant values

$$(\square - m^2)\Phi = 0 \quad : \quad \text{Klein-Gordon equation (scalar)} \quad (3.20)$$

$$(\square - m^2)v^\mu = 0, \partial \cdot v = 0 \quad : \quad \text{Proca equation (massive vector)} \quad (3.21)$$

$$\square A^\mu = 0, \partial \cdot A = 0 \leftrightarrow \partial_\mu F^{\mu\nu} = 0, F_{\mu\nu} := \partial_\mu A_\nu - \partial_\nu A_\mu \text{ Maxwell equation) } \quad (3.22)$$

$$(-i\cancel{\partial} + m)\Psi = 0 \quad : \quad \text{Dirac equation, } m = 0 \text{ is Weyl equation} \quad (3.23)$$

4. UNITARY REPRESENTATIONS ON THE SPACE OF SOLUTIONS: ANTI-PARTICLES

At this stage, we have two sets of irreducible representations of the Poincare group - the ‘‘Particle’’ representations $\{|k, \sigma\rangle\}$, and the ‘‘field’’ representations $\Psi^A(x)$ satisfying the appropriate field equations (or irreducibility conditions). The former are unitary while the latter have no such notion defined for them. We fill this gap now.

The space of vector valued functions, Ψ^A can be made a complex vector space easily enough, but we need to define an inner product to define the notion of unitary. Since the irreducible representations are solutions of the field equations and the equations are linear, we consider the complex vector space of the *solutions of the field equations* and look for an inner product.

Consider the **massive scalar field** first: $(\partial^\mu \partial_\mu - m^2)\Phi(x) = 0$, $\Phi_{\Lambda, a}(x) = \Phi(\Lambda^{-1}(x - a))$. Let u, v be two solutions of the Klein-Gordon equation. Then,

$$u^*(\square - m^2)v - v(\square - m^2)u^* = \partial_\mu (u^* \partial^\mu v - v \partial^\mu u^*) = 0.$$

Hence, $J^\mu := \lambda (u^* \partial^\mu v - v \partial^\mu u^*)$ is a conserved current. This has an immediate implication. Consider a 4 dimensional region bounded by two hypersurfaces of constant value of the time coordinate. More generally, these are two Cauchy surfaces. Restricting to solutions which vanish sufficiently rapidly as one approaches asymptotic infinity along the space-like directions (eg $|\vec{x}| \rightarrow \infty$),

$$\int_{region} d^4x \partial_\mu J^\mu = 0 = \int_{\Sigma_2 \cup \Sigma_1} d^3x J^0 \Rightarrow \int_{\Sigma_2} d^3x J^0 = \int_{\Sigma_1} d^3x J^0.$$

Here we have used that both the Cauchy surfaces have their normals directed ‘outward’ (future pointing for the later one and past pointing for the earlier one). This suggests that we define a candidate inner product between two solutions as,

$$(v, u) := \lambda \int_{\Sigma} d^3x J^0(v, u), \quad J^0(v, u) := v^* \partial^0 u - u \partial^0 v^*, \quad \Sigma \text{ a Cauchy surface.} \quad (4.1)$$

The conserved current implies that the inner product is independent of the Cauchy surface.

Note: On a Cauchy surface, the solution and its time derivative form an initial data and the inner product is really defined on these data. However the Σ -independence of the inner product allows us to think of this as an inner product on the space of solutions.

Is (v, u) really an inner product? It is (i) linear in u and anti-linear in v ; (ii) $(v, u)^* = (u, v)$ provided $\lambda^* = -\lambda$ and (iii) $(u, u) \geq 0$ with equality for $u = 0$?

To check the third property, let us write a solution of the field equation in the form $u(t, \vec{x}) = e^{\pm i\omega t} u_\omega(\vec{x})$, $\omega > 0$. Let us *conventionally* call $e^{-i\omega t}$ as a *positive frequency solution*. Then, $(u, u) = \lambda(-2i\omega) \int_\Sigma d^3x |u_\omega(\vec{x})|^2$. The inner product then satisfies the crucial third property provided we choose $\lambda = +i|\lambda| := i$. The absolute value of λ has been taken to be one as it only affects the normalization of the solutions.

Thus, with the convention adopted, the (v, u) with $\lambda = i$ is indeed an inner product on the *subspace of positive frequency solution*. *Equally well*, the choice $\lambda = -i$ defines an inner product on the *subspace of negative frequency solutions*.

Consider a family of solution, the plane wave solution, $u_{\vec{k}}(x) := A_{\vec{k}} e^{ik \cdot x}$, $k \cdot x := -k^0 t + \vec{k} \cdot \vec{x}$, $k^0 := \sqrt{\vec{k}^2 + m^2} =: \omega_{\vec{k}}$. Substitution gives,

$$(u_{\vec{k}'}, u_{\vec{k}}) = i A_{\vec{k}'}^* A_{\vec{k}} \int_{\Sigma_t} d^3x (-ik^0 - ik'^0) e^{i(k-k')x} \quad (4.2)$$

$$= +A_{\vec{k}'}^* A_{\vec{k}} (k^0 + k'^0) e^{-i(k^0 - k'^0)t} \left((2\pi)^3 \delta^3(\vec{k} - \vec{k}') \right) \quad (4.3)$$

$$= [(2\pi)^3 |A_{\vec{k}}|^2] \left[(2k^0) \delta^3(\vec{k} - \vec{k}') \right] \quad (4.4)$$

Since k^0 depends only on the magnitude of \vec{k} , the delta function forces the frequencies to be equal. We recognize the second square bracket as the Lorentz invariant delta function and *choose* the normalization constant to $A_{\vec{k}} := (2\pi)^{-3/2}$ so that,

$$u_{\vec{k}}(x) := \frac{1}{(2\pi)^{3/2}} e^{ik \cdot x} \quad , \quad k \cdot x := -\sqrt{\vec{k}^2 + m^2} t + \vec{k} \cdot \vec{x} \quad , \quad \vec{k} \in \mathbb{R}^3 \quad (4.5)$$

$$(u_{\vec{k}'}, u_{\vec{k}}) = \delta_{inv}^3(\vec{k} - \vec{k}') = (2k^0) \delta^3(\vec{k} - \vec{k}') \quad . \quad (4.6)$$

The above family of solutions formally form an orthonormal set in the space of *positive frequency solutions*. Technically, these do *not* belong to the space of solutions which have to die off suitably at spatial infinity. This is understood as usual by either using ‘box normalization’ or forming wave-packets. Manipulations done using the above do not lead to any inconsistency.

Now we are ready to check if the Poincare action on the above inner product space is unitary. We will check this by showing that under the Poincare action, the orthonormality is preserved.

Consider,

$$[u_{\vec{k}}(x)]_{\Lambda,a} = u_{\vec{k}}(\Lambda^{-1}(x-a)) = \frac{1}{(2\pi)^{3/2}} e^{ik \cdot \Lambda^{-1}(x-a)} \quad (4.7)$$

$$= \frac{1}{(2\pi)^{3/2}} e^{i(\Lambda k) \cdot (x-a)} \quad \because k_\mu (\Lambda^{-1})^\mu{}_\nu x^\nu = (\Lambda_\nu{}^\mu k_\mu) x^\nu = (\Lambda k) \cdot x$$

$$\therefore [u_{\vec{k}}(x)]_{\Lambda,a} = e^{-i(\Lambda k) \cdot a} u_{\Lambda k}(x) \quad \text{and,} \quad (4.8)$$

$$\begin{aligned} ([u_{\vec{k}'}]_{\Lambda,a}, [u_{\vec{k}}]_{\Lambda,a}) &= e^{i(\Lambda(k-\vec{k}')) \cdot a} ([u_{\Lambda \vec{k}'}], [u_{\Lambda \vec{k}}]) = e^{i(\Lambda(k-\vec{k}')) \cdot a} \delta_{inv}^3(\Lambda(k-\vec{k}')) \\ &= ([u_{\vec{k}'}], [u_{\vec{k}}]) \quad \text{Hence, unitarity!} \end{aligned} \quad (4.9)$$

Note: The orthochronous Lorentz group preserves the sign of the frequency, $k^0 > 0 \Rightarrow (\Lambda k)^0 > 0$, and hence maps positive (negative) frequency solutions into positive (negative) frequency solutions. Thus we see that *the space of solutions of the irreducibility condition (field equations) itself decomposes into two Lorentz invariant subspaces of positive/negative frequency solutions*. There is no contradiction with the Casimir being constant - we just happen to have *two* unitary representations which have the *same* values of the Poincare (orthochronous, proper) Casimir invariants. These in fact represent ‘particle’ and ‘anti-particle’ representations respectively.

The plane wave orthonormal basis, $\{u_{\vec{k}}(x)\}$ is in one-to-one and onto correspondence with the ‘particle basis’ $\{|k\rangle\}$.

The same features are exhibited by the other solution spaces, it remains to identify a conserved current on the space of solutions, define an inner product, obtain an orthonormal set and show unitarity. Since all field equations imply that the fields always satisfy the Klein-Gordon equation, we will always have the positive/negative frequency subspaces and the particle/anti-particle identification.

Consider the **Proca equation with divergence condition**: $(\square - m^2)v^\mu = 0 = \partial \cdot v$. Let V denote the space of solutions of these equations. As before, let v^μ and u^μ be two solutions. Then it follows that $J^\mu := \lambda(v^{*\nu} \partial^\mu u_\nu - u^\nu \partial^\mu v_\nu^*)$ is conserved exactly as before. The divergence condition just selects a subspace of the space of solution of the Klein-Gordon equation. The inner product is defined as (with the same convention of positive frequency solutions),

$$(v, u) := i \int_{\Sigma_t} d^3x [v^{*\nu} \partial^\mu u_\nu - u^\nu \partial^\mu v_\nu^*]$$

Consider the family of solutions,

$$u_{\vec{k}}^\mu(x) := \frac{1}{(2\pi)^{3/2}} \varepsilon^\mu(k) e^{ik \cdot x} , \quad k_\mu \varepsilon^\mu(k) = 0 , \quad k \cdot x := -k^0 t + \vec{k} \cdot \vec{x} \quad (4.10)$$

$$\begin{aligned} (u_{\vec{k}'}^\mu, u_{\vec{k}}^\mu) &= \frac{1}{(2\pi)^3} (k^0 + k'^0) [\varepsilon^{*\mu}(k') (\varepsilon_\mu(k))] e^{-i(k^0 - k'^0)t} \left[(2\pi)^3 \delta^3(\vec{k} - \vec{k}') \right] \\ &= [\varepsilon^{*\mu}(k') (\varepsilon_\mu(k))] \delta_{inv}^3(\vec{k}' - \vec{k}) , \quad \text{here } k^0 := \sqrt{\vec{k}^2 + m^2} . \end{aligned} \quad (4.11)$$

The polarizations $\varepsilon(k)$ satisfying the transversality condition, $k \cdot \varepsilon(k) = 0$ selects 3 independent vectors for $m \neq 0$ and 2 independent vectors for $m = 0$ thanks to the equivalence of A_μ and $A_\mu + \partial_\mu \Lambda$. To distinguish the different polarization vectors, we introduce an additional label, a taking 3 and 2 values respectively for the massive and massless cases. We choose them to satisfy the orthonormality relations, $\boxed{\varepsilon_\mu^*(k, a) \varepsilon^\mu(k, b) = \delta_{ab}}$ and of course $\varepsilon(k, a) \cdot k = 0 \forall a = 1, 2, 3$. For massless case, a is usually denoted by λ and takes only two values. The plane wave family of solutions so defined, constitute an orthonormal set.

Under the Poincare action,

$$\begin{aligned} \left[u_{\vec{k}, a}^\mu(x) \right]_{\Lambda, b} &= \Lambda^\mu{}_\nu u_{\vec{k}, a}^\nu(\Lambda^{-1}(x - b)) = \frac{1}{(2\pi)^{3/2}} \Lambda^\mu{}_\nu \varepsilon^\nu(\vec{k}, a) e^{ik \Lambda^{-1}(x-b)} \\ &= e^{-i(\Lambda k) \cdot b} (\Lambda^\mu{}_\nu \varepsilon^\nu(k, a)) e^{i(\Lambda k) \cdot x} \quad \text{But,} \\ (\Lambda k)_\mu (\Lambda^\mu{}_\nu \varepsilon^\nu(k, a)) &= k_\mu \varepsilon^\mu(k, a) = 0 \quad \therefore \text{ r.h.s.} = e^{i(\Lambda k) \cdot b} \varepsilon^\mu(\Lambda k, a) e^{i(\Lambda k) \cdot x} \\ \left[u_{\vec{k}, a}^\mu(x) \right]_{\Lambda, b} &= e^{i(\Lambda k) \cdot b} u_{\Lambda \vec{k}, a}^\mu(x) . \end{aligned} \quad (4.12)$$

Noting further that $(\Lambda \varepsilon(k', a))^* \cdot (\Lambda \varepsilon(k, b)) = \varepsilon^*(\Lambda k', a) \cdot \varepsilon(\Lambda k, b)$, unitarity of the Poincare action now follows in exactly the same manner as for the scalar. As in the case of the scalar, here too we have two unitary, irreducible representations of the (orthochronous, proper) Poincare group corresponding to the ‘particle’ and ‘anti-particle’ tags. For the zero mass case, the only difference is that the polarizations are restricted to the two spatially transverse directions.

The case of the spinors is more interesting. We follow the same strategy of looking for a conserved current and a corresponding inner product, but we will do this for the Dirac equation which is a first order differential equation. We have:

$$\Psi_{\Lambda, a}^A(x) = D_B^A(\Lambda) \Psi^B(\Lambda^{-1}(x - a)) , \quad (-i\gamma^\mu \partial_\mu \pm m) \Psi = 0 .$$

Note that there are *two* equations.

To look for a conserved current, we need to consider the (matrix) adjoint of the Dirac equation. The adjoint will involve γ^\dagger which are inconvenient for manipulations. However, it follows readily that $\boxed{(\gamma^\mu)^\dagger = \gamma^0 \gamma^\mu \gamma^0}$. This suggests that we use the *Dirac conjugate* defined as: $\bar{\Psi} := \Psi^\dagger \gamma^0$. It follows that

$$(-i\gamma^\mu \partial_\mu \pm m)\Psi = 0 \quad \leftrightarrow \quad i\partial_\mu \bar{\Psi} \gamma^\mu \pm m\bar{\Psi} = 0 .$$

Using these equations for two solutions, Ψ_1, Ψ_2 and their Dirac conjugates, it follows that $\boxed{J^\mu(\Psi_2, \Psi_1) := \lambda(\bar{\Psi}_2 \gamma^\mu \Psi_1)}$ is the conserved current for both $\pm m$ equations. With the choice $\lambda = +1$, the inner product on the solutions of Dirac equations is defined as:

$$(\Psi_2, \Psi_1) := \int_{\Sigma_t} d^3x \bar{\Psi}_2 \gamma^0 \Psi_1 \quad \leftrightarrow \quad (\Psi, \Psi) = \int_{\Sigma_t} d^3x \Psi^\dagger \Psi \geq 0 .$$

Caution: For γ_0 in the above equation, $(\Psi, \Psi) \leq 0$.

To construct a family of orthonormal solutions, consider the ‘plain wave ansatz’, $\Phi^A(\vec{k}, \sigma, x) := \frac{1}{(2\pi)^{3/2}} \varphi^A(k, \sigma) e^{ik \cdot x}$. The Dirac equations then require $(\not{k} \pm m)\varphi(k, \sigma) = 0$ or $\not{k}\varphi = \mp m\varphi$. Multiplying by \not{k} and using $\not{k}\not{k} = -k^2$ leads to $k^2 + m^2 = 0 \leftrightarrow k^0 := \pm \sqrt{\vec{k}^2 + m^2}$. We fix k^0 to be positive and call a solution with $e^{ik \cdot x}$ as having positive frequency and a solution with $e^{-ik \cdot x}$ as having negative frequency. The *two* Dirac equations can now be taken as a single Dirac equation with $+m$, admitting positive and negative frequency solutions. Explicitly, we refine the ansatz as $\Phi_+^A(\vec{k}, \sigma, x) := \frac{1}{(2\pi)^{3/2}} u^A(k, \sigma) e^{ik \cdot x}$ and $\Phi_-^A(\vec{k}, \sigma, x) := \frac{1}{(2\pi)^{3/2}} v^A(k, \sigma) e^{-ik \cdot x}$. Both satisfying $(-i\not{\partial} + m)\Phi_\pm = 0$. This implies that $(\not{k} + m)u(k, \sigma) = 0 = (\not{k} - m)v(k, \sigma)$. The u, v spinors get their k -dependence through these defining equations. These spinors are eigen-spinors of \not{k} with eigenvalues $\pm m$. Since the trace of the γ 's vanishes, \not{k} is traceless too and hence each eigenvalue is doubly degenerate i.e. the ‘ σ ’ label in the eigen-spinors takes *two* values. These will be linked to the helicities later on.

Substitution of these solutions in the inner product leads to,

$$\begin{aligned} (\Phi(k', \sigma',), \Phi(k, \sigma)) &= \int_{\Sigma_t} d^3x \frac{1}{(2\pi)^3} u^\dagger(k', \sigma') u(k, \sigma) e^{-i(k' - k) \cdot x} \\ &= u^\dagger(k', \sigma') u(k, \sigma) e^{-i(k^0 - k'^0)t} \delta^3(\vec{k} - \vec{k}') \end{aligned} \quad (4.13)$$

$$\therefore (\Phi(k', \sigma',), \Phi(k, \sigma)) = \left[\frac{u^\dagger(k', \sigma') u(k, \sigma)}{2k^0} \right] \delta_{inv}^3(\vec{k} - \vec{k}') := \delta_{\sigma', \sigma} \delta_{inv}^3(\vec{k} - \vec{k}') . \quad (4.14)$$

We have chosen a normalization of the u -spinors. Identical orthonormality relations follow for the v -spinors.

Under Lorentz transformations, the solutions transform as,

$$[\Phi(k, \sigma)]_{\Lambda}(x) = [D(\Lambda)u(k, \sigma)] \frac{e^{i(\Lambda k) \cdot x}}{(2\pi)^{3/2}} \stackrel{?}{=} \Phi(\Lambda k, \sigma)(x)$$

The last equality will hold if $D(\Lambda)u(k, \sigma) = u(\Lambda k, \sigma)$ i.e. if $((\Lambda k) + m) [D(\Lambda)u(k, \sigma)] = 0$.

We already have $[\Sigma^{\mu\nu}, \gamma^{\lambda}] = i(\eta^{\mu\lambda}\gamma^{\nu} - \eta^{\nu\lambda}\gamma^{\mu})$. This implies (may be checked by taking the infinitesimal form of $D(\Lambda)$),

$$D^{-1}(\Lambda)\gamma^{\mu}D(\Lambda) = \Lambda^{\mu}_{\nu}\gamma^{\nu} \Rightarrow \gamma^{\mu}D(\Lambda) = \Lambda^{\mu}_{\nu}D(\Lambda)\gamma^{\nu} .$$

It follows,

$$[(\Lambda k) + m] (D(\Lambda)u(k, \sigma)) = [(\Lambda k)_{\mu}\gamma^{\mu}D(\Lambda) + D(\Lambda)m] u(k, \sigma) \quad (4.15)$$

$$= [(\Lambda k)_{\mu} \Lambda^{\mu}_{\nu}D(\Lambda)\gamma^{\nu} + D(\Lambda)m] u(k, \sigma)$$

$$= D(\Lambda) [\Lambda_{\mu}^{\alpha}k_{\alpha}\Lambda^{\mu}_{\nu}\gamma^{\nu} + m] u(k, \sigma) \quad \text{but, } \Lambda_{\mu}^{\alpha}\Lambda^{\mu}_{\nu} = \delta^{\alpha}_{\nu}$$

$$\therefore [(\Lambda k) + m] (D(\Lambda)u(k, \sigma)) = D(\Lambda) [k_{\nu}\gamma^{\nu} + m] u(k, \sigma) = 0. \quad (\text{Covariance Result})(4.16)$$

Thus, indeed we have $\boxed{D(\Lambda)u(k, \sigma) = u(\Lambda k, \sigma)}$ and hence,

$$[\Phi(k, \sigma)]_{\Lambda, a}(x) = e^{-i(\Lambda k) \cdot a} \Phi(\Lambda k, \sigma)(x) .$$

Under Lorentz transformations then the orthonormality relation transforms as,

$$\begin{aligned} (\Phi(k', \sigma')_{\Lambda, a}, \Phi(k, \sigma)_{\Lambda, a}) &= e^{i\Lambda(k' - k) \cdot a} (\Phi(\Lambda k', \sigma'), \Phi(\Lambda k, \sigma)) \\ &= \left[\frac{u^{\dagger}(\Lambda k', \sigma')u(\Lambda k, \sigma)}{2(\Lambda k)^0} \right] \delta_{in\nu}^3(\vec{k}' - \vec{k}) = \delta_{\sigma', \sigma} \delta_{in\nu}^3(\vec{k}' - \vec{k}) \end{aligned} \quad (4.17)$$

In the last equality we have used the normalization of the u spinors. This proves the unitarity of the Poincare representation on the positive frequency solutions. As before, Lorentz transformations do not mix the subspaces of the positive/negative frequency solutions.

The normalization definition does not look Lorentz invariant, but it is. Both $u^{\dagger}u = \bar{u}\gamma^0u$ and k^0 transform the same way. A more convenient form will be displayed later on.

Note: The covariance result allows us to define the u, v spinors from their definition in the rest frame (for massive case) i, $\hat{k} = (m, \vec{0})$. In this frame, the spinors satisfy the equations: $-m\gamma^0u(\hat{k}, \sigma) = -mu(\hat{k}, \sigma)$, and $-m\gamma^0v(\hat{k}, \sigma) = +mv(\hat{k}, \sigma)$. Thus these spinors at \hat{k} are eigen-spinors of γ^0 with eigenvalues ± 1 respectively. γ^0 being Hermitian, these spinors are orthogonal: $u^{\dagger}(\hat{k}, \sigma)v(\hat{k}, \sigma) = 0$. As noted before, these eigenvalues are doubly degenerate

and σ labels these. For the massless case, $\hat{k} = (k, 0, 0, k)$ and u, v are eigen-spinors of J_3 generator of the Little group.

Note: Incidentally, covariance result and the identification of $u(\hat{k}, \sigma), v(\hat{k}, \sigma)$ as eigen-spinors of γ^0 also leads to a *completeness relation* as follows. Given that $\gamma^0 u(\hat{k}, \sigma) = u(\hat{k}, \sigma)$, $\gamma^0 v(\hat{k}, \sigma) = -v(\hat{k}, \sigma)$ and each eigen-space being two dimensional, \Rightarrow completeness relations ('spectral representation' - $A = \sum_n \lambda_n |n\rangle \langle n|$) takes the form

$$\begin{aligned} \sum_{\sigma} u(\hat{k}, \sigma) u^{\dagger}(\hat{k}, \sigma) &= \frac{\mathbb{1} + \gamma^0}{2} \quad , \quad \sum_{\sigma} v(\hat{k}, \sigma) v^{\dagger}(\hat{k}, \sigma) = \frac{\mathbb{1} - \gamma^0}{2} \\ \sum_{\sigma} u(\hat{k}, \sigma) \bar{u}(\hat{k}, \sigma) &= \frac{\gamma^0 + \mathbb{1}}{2} = \frac{-\hat{k} + m}{2m} \quad , \quad \sum_{\sigma} v(\hat{k}, \sigma) \bar{v}(\hat{k}, \sigma) = \frac{\gamma^0 - \mathbb{1}}{2} = -\frac{\hat{k} + m}{2m} \\ \sum_{\sigma} u(\vec{k}, \sigma) \bar{u}(\vec{k}, \sigma) &= \frac{-\vec{k} + m}{2m} \quad , \quad \sum_{\sigma} v(\vec{k}, \sigma) \bar{v}(\vec{k}, \sigma) = -\frac{\vec{k} + m}{2m} \end{aligned}$$

In the second equation, we used $\gamma^0 = -\frac{\hat{k}}{m}$ and in the last equation we multiplied by $D(L)$ on the left and $D^{-1}(L)$ on the right and used the covariance result. Here L is a boost which takes $\hat{k} \rightarrow \vec{k}$. Notice that multiplying by $u(\vec{k}, \sigma')$ and $v(\vec{k}, \sigma')$ respectively, and using the equations satisfied by the spinors, gives $\boxed{\bar{u}(\vec{k}, \sigma) u(\vec{k}, \sigma') = \delta_{\sigma, \sigma'} = -\bar{v}(\vec{k}, \sigma) v(\vec{k}, \sigma')}$. These are consistent with the previously chosen normalizations: $u^{\dagger}(\vec{k}, \sigma) u(\vec{k}, \sigma') = 2k^0 \delta_{\sigma, \sigma'} = v^{\dagger}(\vec{k}, \sigma) v(\vec{k}, \sigma')$.

Thus, for all the field equations we have found unitary, irreducible representations of the Poincare group. We have also discovered that the *field representations automatically also include 'anti-particles'*. This is really a consequence of the requirement of *manifest Poincare covariance* which necessitates the field representations.

Note: From the expressions above, it should be clear that the ket vectors $|k, \sigma\rangle$ are in one-to-one correspondence with the "Plane wave solutions" with positive *and* negative frequencies, displayed above.

We need to study the behavior of these representations under the space inversion, time reversal and the new possibility of 'charge conjugation' (particle-particle exchange).

5. PARITY, TIME REVERSAL AND CHARGE CONJUGATION

For the field representations above, we focussed on the subgroup of the Poincare group, connected to the identity. The remaining elements of the Poincare group - improper and non-orthochronous are generated by the two transformations of space inversion and time reversal. Their actions on the generators is given in eq.(2.36). These relations continue to hold in *any* representation, in particular also on the field representations. Looking at the general form of the Poincare action on the $\Psi^A(x)$'s, $\Psi_{\Lambda,a}^A(x) = D_B^A(\Lambda)\Psi^B(\Lambda^{-1}(x-a))$, we see that the action on the space-time point is common to *all* $D(\Lambda)$ representations and we may focus on the $D(\Lambda)$ part of it. Recalling that the time reversal operation is anti-linear and anti-unitary, we have to be careful about taking complex conjugates. Thus we define the actions as,

$$\Psi_{\mathcal{P}}^A(x) := D_B^A(\mathcal{P})\Psi^B(t, -\vec{x}) \quad , \quad D(\mathcal{P}) =: \Pi \quad (5.1)$$

$$\Psi_{\mathcal{T}}^A(x) := D_B^A(\mathcal{T})(\Psi^B(-t, \vec{x}))^* \quad , \quad D(\mathcal{T}) =: \tau, \quad (5.2)$$

It is customary to denote the anti-unitary, anti-linear operator \mathcal{T} as τK where τ is a unitary operator and K takes complex conjugate of numbers on its right. $\mathcal{T}^2 = \mathbb{1}$ implies $\tau\tau^* = \mathbb{1}$.

Consider the translation generators. These act as differential operators on the Ψ 's: $P_\mu\Psi^A(x) = -i\partial_\mu\Psi^A$. Consider the \mathcal{P} and \mathcal{T} actions on P_μ as given in eqn. (2.36). For instance,

$$\begin{aligned} (\mathcal{T}^{-1}P_0\mathcal{T})\Psi^A(t, \vec{x}) &= \mathcal{T}^{-1}\{-i\partial_0(\tau_B^A\Psi^B(-t, \vec{x})^*)\} = \mathcal{T}^{-1}\{+i\partial_{-t}(\tau_B^A\Psi^B(-t, \vec{x})^*)\} \\ &= -i\partial_t(\tau\tau^*)^A_B\Psi^B(t, \vec{x}) = P_0\Psi^A(t, \vec{x}) . \end{aligned} \quad (5.3)$$

The time has been reversed twice and the complex conjugation has been effected twice. For the spatial components, there is no 't' reversal and the \mathcal{T} introduces a sign. Space inversion is likewise straight forward. We now focus on the Lorentz generators only.

The scalar case has no non-trivial matrices and the vector case is simpler and is left as an exercise. The spinner case is non-trivial. We have the Lorentz generators:

$$\Sigma^{\mu\nu} = \frac{i}{4} [\gamma^\mu, \gamma^\nu] \leftrightarrow J_i = \frac{\epsilon_{ijk}}{2} \Sigma^{jk} \quad , \quad K^i = \Sigma^{i0} .$$

The Π and τ matrices are also 4×4 matrices. We note a result.

Result: Any 4×4 matrix can be written as a linear combination of the 16 Γ matrices, $\{\Gamma\} = \{\mathbb{1}, \gamma^\mu, \gamma^{[\mu}\gamma^{\nu]}, \gamma^{[\mu}\gamma^\nu\gamma^\lambda], \gamma_5\}$, the square brackets denoting anti-symmetrization ($16 = 1 + 4 + 6 + 4 + 1$).

The result follows by showing that the 16 matrices are linearly independent.

We now deduce the Π and τ matrices from the commutation relations (see eq.(2.36)):

$$\Sigma_{jk}\Pi = \Pi\Sigma_{jk}, \quad \Sigma_{0i}\Pi = -\Pi\Sigma_{0i}, \quad \Sigma_{jk}\tau = -\tau(\Sigma_{jk})^*, \quad \Sigma_{0i}\tau = \tau(\Sigma_{0i})^* .$$

The determination of Π is straight forward. Commutation with Σ_{ij} implies that Π is a linear combination of $\mathbb{1}, \gamma_5, \gamma_0$. Anti-commutation with $\Sigma_{0i} = \frac{i}{2}\gamma_0\gamma_i$ implies that only γ_0 survives and $\Pi = c\gamma^0$. Now $\Pi^2 = \mathbb{1}$ implies $c = \pm 1$. Hence, $\boxed{(\Psi_{\mathcal{P}})^A(x) = \pm(\gamma^0)^A_B\Psi^B(t, -\vec{x})}$. It follows immediately that *left(right) handed Weyl spinor becomes right(left) handed Weyl spinor under space inversion*.

The determination of τ involves complex conjugation. From the unitarity of the γ matrices and the defining Clifford relations, the hermiticity properties are determined: $\gamma_0^\dagger = \gamma_0, \gamma_i^\dagger = -\gamma_i$. The transpose/complex conjugation properties however depend on the explicit choice of the γ 's. The τ matrix thus depends on the explicit representation of the γ 's. There are three commonly employed representations:

$$\text{Dirac-Pauli} : \gamma^0 := \begin{pmatrix} \mathbb{1} & 0 \\ 0 & -\mathbb{1} \end{pmatrix}, \quad \gamma^i := \begin{pmatrix} 0 & \sigma^i \\ -\sigma^i & 0 \end{pmatrix}, \quad \gamma_5 := \begin{pmatrix} 0 & \mathbb{1} \\ \mathbb{1} & 0 \end{pmatrix}; \quad (5.4)$$

$$\text{Weyl} : \gamma^0 := \begin{pmatrix} 0 & \mathbb{1} \\ \mathbb{1} & 0 \end{pmatrix}, \quad \gamma^i := \begin{pmatrix} 0 & \sigma^i \\ -\sigma^i & 0 \end{pmatrix}, \quad \gamma_5 := \begin{pmatrix} -\mathbb{1} & 0 \\ 0 & \mathbb{1} \end{pmatrix}; \quad (5.5)$$

$$\begin{aligned} \text{Majorana} : \gamma^0 &:= \begin{pmatrix} 0 & \sigma_2 \\ \sigma_2 & 0 \end{pmatrix}, \quad \gamma^1 := \begin{pmatrix} i\sigma_3 & 0 \\ 0 & i\sigma_3 \end{pmatrix}, \quad \gamma^2 := \begin{pmatrix} 0 & -\sigma_2 \\ \sigma_2 & 0 \end{pmatrix}; \quad (5.6) \\ &: \gamma^3 := \begin{pmatrix} -i\sigma_1 & 0 \\ 0 & -i\sigma_1 \end{pmatrix}, \quad \gamma_5 := \begin{pmatrix} \sigma_2 & 0 \\ 0 & -\sigma_2 \end{pmatrix}, \quad \text{All purely imaginary.} \quad (5.7) \end{aligned}$$

In the Dirac-Pauli and Weyl representations, only γ_2 is imaginary. The τ matrix thus satisfies:

$$\begin{aligned} \Sigma_{12}\tau &= -\tau\Sigma_{12}, \quad \Sigma_{23}\tau = -\tau\Sigma_{23}, \quad \Sigma_{31}\tau = \tau\Sigma_{31}, \\ \Sigma_{01}\tau &= -\tau\Sigma_{01}, \quad \Sigma_{02}\tau = \tau\Sigma_{02}, \quad \Sigma_{03}\tau = -\tau\Sigma_{03}, \end{aligned}$$

The last relation in the first line suggests $\tau = \lambda\gamma^0\gamma^2$. It also checks with all other relations. The λ is determined as follows.

The operator $D(\mathcal{T})$ is anti-unitary. Hence,

$$(D(\mathcal{T})D(\mathcal{T})\Psi_1, D(\mathcal{T})D(\mathcal{T})\Psi_2) = (D(\mathcal{T})\Psi_2, D(\mathcal{T})\Psi_1) = (\Psi_1, \Psi_2) \Rightarrow D^2(\mathcal{T}) \text{ is unitary.}$$

Next, $D^2(\mathcal{T}) = (\tau K)(\tau K) = \tau\tau^* = |\lambda|^2(\gamma^0\gamma^2\gamma^0(-\gamma^2)) = -|\lambda|^2\mathbb{1}$. The unitarity then implies that $|\lambda| = 1$ i.e. λ is a phase and $D^2(\mathcal{T}) = -\mathbb{1}$ in the spinor representation.

In the Majorana representation, all γ 's are imaginary. Hence, $\Sigma_{\mu\nu}^* = -\Sigma_{\mu\nu}$. Hence Σ_{jk} commute with τ and anti-commute with Σ_{0i} . This is exactly as for the space inversion and we deduce that $D(\mathcal{T}) = \lambda\gamma^0K$. Once again $D^2(\mathcal{T})$ is unitary and equal to $-\mathbb{1}$ and λ is a phase.

For vector representation, the 'A' index will be a tensorial index and it is left as an exercise to work out the D matrices for space inversion and time reversal.

The existence of the anti-particle representations suggest one more discrete transformation of order 2, called *Charge Conjugation*.

Recall that the anti-particle representation is the subspace of negative frequency solutions. These subspaces are spanned by plane wave solutions and involve the operation of complex conjugation which takes $e^{ik\cdot x} \rightarrow e^{-ik\cdot x}$. For the non-trivial $D(\Lambda)$ representations, complex conjugation also takes the complex conjugate of the $D(\Lambda)$ matrices. However, a complex conjugation of a solution need not be a solution again (especially for spinor as we will see) and hence the charge conjugation must also involve additional transformations over and above complex conjugation. For the Klein-Gordon, Proca and Maxwell equations, complex conjugate of a solution is also a solution as the differential operators are real. For the Dirac equation we need to do further work.

For instance, let Ψ be a positive frequency solution of the Dirac equation, $(-i\gamma^\mu\partial_\mu + m)\Psi = 0$. Taking complex conjugate of the equation, we get $(+i(\gamma^\mu)^*\partial_\mu + m)\Psi^* = 0$. If we could find an invertible matrix B such that $(\gamma^\mu)^* = -B\gamma^\mu B^{-1}$, then $\Psi^c := (B^{-1}\Psi^*)$ satisfies the same Dirac equation and of course is a negative frequency solution. Ψ^c is called the *charge conjugate*² of the Ψ . Apparently, B depends on the choice of explicit γ matrices.

² The terminology comes when coupling to external electromagnetic field is considered by the minimal substitution $\partial_\mu \rightarrow \partial_\mu - ieA_\mu$. Under complex conjugation, $e \rightarrow -e$. So if Ψ is thought of as charge 'e' solution then Ψ^c is a charge '-e' solution.

As a preparation for subsequent development, we have a subsection on properties of γ matrices.

A. Representations of the Clifford algebra and relations among them

Quite generally, for any group, given a (matrix) representation $R(G)$, we have three other representations, namely, $R^*(G)$, $(R^T)^{-1}(G)$ and $(R^\dagger)^{-1}(G)$. From the basic relation $R(g_1)R(g_2) = R(g_1.g_2)$, taking complex conjugate, transpose inverse and adjoint inverse immediately verifies the assertion. If in addition, the representation $R(G)$ is *unitary* (always true for finite groups and compact Lie groups), then $(R^T)^{-1}(G) = R^*(G)$ and $(R^\dagger)^{-1}(G) = R(G)$. $R(G)$ and $R^*(G)$ are then the only independent representations. These are either equivalent, $R^*(g) = SR(g)S^{-1}$ or inequivalent. It turns out that for unitary, irreducible representations, if R and R^* are equivalent, then S is either symmetric or anti-symmetric. The following terminology ensues.

For unitary, irreducible representations:

$$\begin{aligned} R(G) \text{ is complex} & \quad \text{if } R^*(G) \approx R(G) \\ R(G) \text{ is pseudo-real} & \quad \text{if } R^*(G) = SR^*(G)S^{-1} \quad , \quad S^T = -S \\ R(G) \text{ is real} & \quad \text{if } R^*(G) = SR^*(G)S^{-1} \quad , \quad S^T = S \end{aligned}$$

This is relevant for representations of internal symmetry groups as well as the Clifford group of the 32 elements. The $D(\Lambda)$ representations are finite dimensional but not unitary since Lorentz group does not have unitary finite dimensional representations.

For the Clifford group in 4 dimensions (and in even number of dimensions) there is only one (up-to unitary equivalence) non-trivial representation which is 4 dimensional (or $2^{N/2}$ dimensional for N -even number of dimensions). Since $\pm\gamma^*$, $\pm\gamma^T$ and $\pm\gamma^\dagger$ all satisfy the same Clifford algebra and the representation is unique, \exists matrices B, C, D such that,

$$(\gamma^\mu)^* = -B\gamma^\mu B^{-1} \quad , \quad (\gamma^\mu)^T = -C\gamma^\mu C^{-1} \quad , \quad (\gamma^\mu)^\dagger = +D\gamma^\mu D^{-1} \quad .$$

The choice of signs above is conventional. Notice that replacing any of the B, C, D matrices by multiplying by γ_5 on the right, reverses the signs.

It follows immediately that,

$$-(\Sigma^{\mu\nu})^* = B\Sigma^{\mu\nu}B^{-1} \quad , \quad -(\Sigma^{\mu\nu})^T = C\Sigma^{\mu\nu}C^{-1} \quad , \quad (\Sigma^{\mu\nu})^\dagger = D\Sigma^{\mu\nu}D^{-1}$$

Given a representation $D(\Lambda)$ we have the $(D^\dagger)^{-1}(\Lambda)$, $(D^T)^{-1}(\Lambda)$ and $D^*(\Lambda)$ representations. The infinitesimal forms are $(\mathbb{1} + \frac{i}{2}\omega \cdot \Sigma)$, $(\mathbb{1} - \frac{i}{2}\omega \cdot \Sigma)^\dagger$, $(\mathbb{1} - \frac{i}{2}\omega \cdot \Sigma^T)$, $(\mathbb{1} - \frac{i}{2}\omega \cdot \Sigma^*)$ which imply that the corresponding generators are $\Sigma^{\mu\nu}$, $(\Sigma^{\mu\nu})^\dagger$, $-(\Sigma^{\mu\nu})^T$, $-(\Sigma^{\mu\nu})^*$. The B, C, D matrices precisely relate these generators.

These relations can be exponentiated to get corresponding relations among the $D(\Lambda)$, $D^\dagger(\Lambda)$, $D^T(\Lambda)$, $D^*(\Lambda)$, namely,

$$D^\dagger(\Lambda) = DD^{-1}(\Lambda)D^{-1} \quad , \quad D^T(\Lambda) = CD^{-1}(\Lambda)C^{-1} \quad , \quad D^*(\Lambda) = BD(\Lambda)B^{-1} \quad .$$

These can be checked easily by using the series form of the exponentials derived from $D(\Lambda) = \sum_{k=0}^{\infty} \frac{(i\omega \cdot \Sigma)^k}{k!}$.

B. Dirac-Majorana-Charge Conjugates

This allows us to construct Lorentz invariant bi-linears from the spinors. For instance, define the *Majorana conjugate* $\tilde{\Psi} := \Psi^T C$. Then,

$$(\tilde{\Psi})_\Lambda := (\Psi_\Lambda^T)C = \Psi^T(D^T(\Lambda))C = \Psi^T CD^{-1}(\Lambda) = \tilde{\Psi}D^{-1}(\Lambda) \quad .$$

Thus the Majorana conjugate, like the Dirac conjugate $\bar{\psi} := \Psi^\dagger D$, transforms by $D^{-1}(\Lambda)$. Recall that the charge conjugate, Ψ^c transforms by $D(\Lambda)$. *Consequently, $\bar{\Psi}\Psi$, $\bar{\Psi}\Psi^c$, $\tilde{\Psi}\Psi$, $\tilde{\Psi}\Psi^c$ are all Lorentz scalars.*

Recalling that $D^{-1}(\Lambda)\gamma^\mu D(\Lambda) = \Lambda^\mu{}_\nu \gamma^\nu$, it is easy to see that,

$\bar{\Psi}\Psi$ is a Scalar

$\bar{\Psi}\gamma^\mu\Psi$ is a Vector

$\bar{\Psi}\gamma^\mu\gamma^\nu\Psi$ is a Tensor of rank 2

$\bar{\Psi}\gamma^\mu\gamma_5\Psi$ is a Axial (or pseudo) vector

$\bar{\Psi}\gamma_5\Psi$ is a Pseudo-scalar

Combine these with the definitions,

Dirac Spinor : $\Psi = \Psi$, $(\Psi)_\Lambda = D(\Lambda)\Psi$;

Dirac Conjugate : $\bar{\Psi} = \Psi^\dagger D$, $(\bar{\Psi})_\Lambda = \bar{\Psi}D^{-1}(\Lambda)$;

Charge Conjugate : $\Psi^c = B^{-1}\Psi^*$, $(\Psi^c)_\Lambda = D(\Lambda)\Psi^c$;

Majorana Conjugate : $\tilde{\Psi} = \Psi^T C$, $(\tilde{\Psi})_\Lambda = \tilde{\Psi}D^{-1}(\Lambda)$.

Thus, the same Lorentz transformation properties hold with $\Psi \rightarrow \Psi^c$ and/or $\bar{\Psi} \rightarrow \tilde{\Psi}$. The last two need a little explanation.

We defined $\gamma_5 := i\gamma^0\gamma^1\gamma^2\gamma^3 = \frac{i}{4!}\varepsilon_{\mu\nu\alpha\beta}\gamma^\mu\gamma^\nu\gamma^\alpha\gamma^\beta$, where ε is the completely anti-symmetric symbol, with $\varepsilon_{0123} := 1$. This symbol is also used to define the determinant of a matrix, eg, $\varepsilon_{\mu'\nu'\alpha'\beta'}\Lambda_{\mu'}^{\mu'}\Lambda_{\nu'}^{\nu'}\Lambda_{\alpha'}^{\alpha'}\Lambda_{\beta'}^{\beta'} = \det(\Lambda)\varepsilon_{\mu\nu\alpha\beta}$. It follows immediately that $D^{-1}(\Lambda)\gamma_5 D(\Lambda) = \det(\Lambda)\gamma_5$.

Consider now an axial vector combination,

$$[\bar{\Psi}\gamma^\mu\gamma_5\Psi]_\Lambda = \bar{\Psi}D^{-1}(\Lambda)\gamma^\mu\gamma_5 D(\Lambda)\Psi = \Lambda^\mu_\nu\bar{\Psi}\gamma^\nu D^{-1}\gamma_5 D(\Lambda)\Psi = \det(\Lambda)\Lambda^\mu_\nu\bar{\Psi}\gamma^\nu\gamma_5\Psi.$$

Note that for proper Lorentz transformations, the axial vector and the pseudo-scalar transform as vector and scalar respectively.

Quantities that transform as tensors but with an extra factor of determinant of the transformation are called *pseudo-tensors*. There are also tensor looking quantities that transform with additional factor of $|\det|^w$ and these are called *tensor densities of weight w*.

Let us return to determine the matrices B, C, D .

Since the γ 's are unitary, we can always choose any of the B, C, D matrices to be unitary. Quite generally, if $R' = SRS^{-1}$ for two irreducible, equivalent unitary representations, then $R'^\dagger R' = \mathbb{1} \Rightarrow R^\dagger S^\dagger S = S^\dagger S R \Rightarrow S^\dagger S = \alpha\mathbb{1}$. Positivity of $S^\dagger S$ gives $\alpha > 0$. Hence $S' := S/\sqrt{\alpha}$ is a unitary matrix. This proves the claim.

Next, $\gamma^\mu = (\gamma^{\mu*})^* = -B^*\gamma^{\mu*}(B^{-1})^* = +B^*B\gamma^m(B^*B)^{-1} \Rightarrow B^*B = \lambda\mathbb{1}$ with λ being real since B^*B is. The determinant of B being a phase (unitarity of B) fixes $\lambda := \epsilon_B := \pm 1$ and $B^*B = \epsilon_B\mathbb{1}$. Similar manipulation with $\gamma^\mu = (\gamma^{\mu T})^T$ leads to $C^{-1}C^T = \lambda\mathbb{1}$. Unitarity gives $C^{-1}C^T = C^\dagger C^T = (C^T)^*C^T$, hence λ is real and determinant gives $\lambda^4 = 1$. Hence, $C^T = \epsilon_C C, \epsilon_C = \pm 1$. For $\gamma = (\gamma^\dagger)^\dagger$ we get $D^\dagger = \lambda D$ for some phase λ . Since D is determined to within a phase, we can define $D' := e^{i\alpha}D$ and choose α so that $\lambda = e^{-2i\alpha}$ to get $(D')^\dagger = D'$. Thus, without loss of generality, we can always choose $D^\dagger = D$. Now, knowing the hermiticity properties of the γ 's, we get $D = \gamma^0$, independent of any choice of explicit γ matrices.

The ϵ_B, ϵ_C are correlated since $\gamma^T = (\gamma^*)^\dagger = (\gamma^\dagger)^*$. $\gamma^T = (\gamma^*)^\dagger$ leads to,

$$-C\gamma C^{-1} = -(B^\dagger)^{-1}D\gamma D^{-1}B^\dagger \Rightarrow C = \lambda B D.$$

$\gamma^T = (\gamma^\dagger)^*$ leads to,

$$-C\gamma C^{-1} = -(D^*)B\gamma B^{-1}(D^{-1})^* \Rightarrow C = \lambda' D^* B.$$

Eliminating C gives $BD = \frac{\lambda'}{\lambda} D^* B$ or $D^* = \frac{\lambda'}{\lambda} B D B^{-1}$. However, $D = \gamma^0$ can always be achieved as shown above. Therefore, $\lambda' = -\lambda$ and $C = \lambda B D = -\lambda D^* B$. Next,

$$C^T = \lambda D^T B^T = \lambda (D^\dagger)^* (B^\dagger)^* = \lambda D^* (B^*)^{-1} = \lambda D^* (\epsilon_B B) = -\epsilon_B C \Rightarrow \boxed{\epsilon_C = -\epsilon_B.}$$

Note that this is independent of explicit γ matrices and also independent of the phase λ ! We still need to determine ϵ_B , say.

Claim: The ϵ_B does not depend on the choice of γ matrices.

This is easily proved. Let γ' and γ be two distinct choices of explicit γ -matrices. Both being unitary are related by some unitary matrix, S as $\gamma = S\gamma'S^{-1}$. Then substitution in $\gamma^* = -B\gamma B^{-1}$ gives, $(\gamma')^* = -B'\gamma'(B')^{-1}$ with $B' := (S^*)^{-1}BS$. It follows that $(B')^*B' = (S^{-1}B^*S^*)((S^*)^{-1}BS) = \epsilon_B \mathbb{1}$, proving the claim. It therefore suffices to choose an explicit representation and evaluate ϵ_B .

Referring to say the Dirac-Pauli representation, see (5.4), only γ^2 is pure imaginary. Hence, $\gamma^{0,1,3}B = -B\gamma^{0,1,3}$ and $\gamma^2B = +B\gamma^2$. Furthermore, $\gamma_5^* = \gamma_5 \Rightarrow \gamma_5 B = -B\gamma_5$ as well. Therefore B must be made up of odd number of γ 's. By inspection, $B = \alpha\gamma^2$ satisfies the conditions. Unitarity of B restricts α to be a phase and $B^*B = |\alpha|^2(-\gamma^2)(\gamma^2) = \mathbb{1} \Rightarrow \epsilon_B = +1$ and $\epsilon_C = -1$. Also $C = \lambda B D = \lambda\alpha\gamma^2\gamma^0 = (-\lambda\alpha)\gamma^0\gamma^2$. *The phases λ, α are arbitrary and convention dependent. We choose $\lambda = 1$ and $\alpha = i$ so that $\boxed{B = i\gamma^2, C = i\gamma^2\gamma^0}$.*

This also verifies for the Weyl representation. For Majorana representation, all γ^μ 's are imaginary and thus *commute* with B . Hence B must be a phase multiple of $\mathbb{1}$. This is also consistent with commutation with γ_5 . Clearly $\epsilon_B = +1$ and $C = (\lambda\alpha)\gamma^0$.

This completes the discussion of representations and relations among them.

It is customary to introduce a *charge conjugation operator*, $C := B^{-1}K$, where K instructs to take the complex conjugate of the numbers on its right. This operator is anti-linear and since B is unitary, it is also anti-unitary operator. It follows,

$$C\Sigma^{\mu\nu} = B^{-1}(\Sigma^{\mu\nu})^*K = B^{-1}(-B\Sigma^{\mu\nu}B^{-1})K = -\Sigma^{\mu\nu}C \Rightarrow CD(\Lambda) = D(\Lambda)C.$$

Thus charge conjugation acts invariantly on Lorentz representations. *Provided $C^2 = \mathbb{1}$* , we have $(\frac{1\pm C}{2})^2 = \frac{1\pm C}{2}$ and the Lorentz representation can be reduced further. The corresponding subspaces satisfy $C\psi = \pm\Psi$ and these spinors are called *Majorana spinors*. Thus,

Majorana spinors are Dirac spinors which satisfy $C\Psi = \pm\Psi \leftrightarrow B^{-1}\psi^* = \pm\Psi$. So is $C^2 = \mathbb{1}$? We have $C^2 = B^{-1}KB^{-1}K = (B^*B)^{-1}K^2 = \epsilon_B\mathbb{1} = \mathbb{1}$. Therefore, we *do* have Majorana spinors (in 4 dimensions).

Do we have Weyl-Majorana spinors? Well, $C\gamma_5 = B^{-1}\gamma_5^*K = -\gamma_5B^{-1}K = -\gamma_5C$ and we *cannot have* Weyl-Majorana spinors in 4 dimensions.

A similar analysis can be carried out for spinors in any dimensions and with any metric signature. This may be seen for instance in [2].

6. RELATIVISTIC ACTIONS: CLASSICAL FIELDS

So far we focused on the representation theory of the Poincare group. The abstract, algebraic approach revealed the attributes of these representations, namely mass and spin/helicity. In order to have a framework which is manifestly Poincare covariant, we constructed and analyzed representations on vector valued (complex in general) function, Ψ^A - henceforth these will be generically referred to as *fields*. The irreducibility condition emerged as “field equations”. These equations are all homogeneous, linear and with at the most two derivatives. They all admit plane waves and their linear combinations as solutions, but no other ‘phenomenon’ involving (say) different types of waves modifying their propagation etc. Intuitively, there are *no interactions*.

We would like to have a framework which is not only Poincare covariant but also involves “interaction” or non-trivial “dynamics”. The well tested and successful strategy is to have an *action formulation*. Let us quickly note several advantages of having an action formulation:

1. The equations of motion are retrieved invoking a variational principle as the Euler-Lagrange equations of motion;
2. “Interactions” can be understood naturally as leading to non-linear and/or coupled equations which can be easily introduced as more than quadratic order terms in the action;
3. Covariance of equations of motion (or dynamics) can be easily incorporated by requiring appropriate invariance of the action;
4. The Noether’s theorem gives a recipe for obtaining quantities conserved by equations of motion (and hence during interactions as well);
5. It leads to a canonical framework of symplectic structure (“Poisson brackets”) and a Hamiltonian evolution. This provides a systematic method of identifying “degrees of freedom” (eg Dirac’s theory of constrained systems).

A canonical structure is already inherent in the quantum framework: the imaginary part of the inner product provides the symplectic structure while the Schrodinger equation gives a Hamiltonian evolution;

6. Path integral quantization - very well suited for gauge theories especially the non-abelian ones - has action as the central quantity.

Without further ado, let us proceed with an action formulation. Our first aim is to obtain the Klein-Gordon, the Dirac and the Proca/Maxwell equations. Because the equations are local, partial differential equations, the action must be an integral over the Minkowski space-time, of a Lagrangian density built out of the Poincare covariant fields and their derivatives i.e. it must of the form $S = \int d^4x \sqrt{|\det(\eta)|} \mathcal{L}$ with \mathcal{L} is a Lorentz scalar built out of the fields. It is called the *Lagrangian density*. Here, η denotes the Minkowski metric and is necessary to absorb the Jacobian of Lorentz transforms of the space-time coordinates. The absolute value of the determinant happens to be 1 and is suppressed throughout. A variational principle has the form,

$$\delta S[\Phi] := S[\Phi + \delta\Phi] - S[\Phi] = \int d^4x \delta\mathcal{L} := \int d^4x \delta\Phi \text{ " } \frac{\delta\mathcal{L}}{\delta\Phi} \text{ " } .$$

Extremization of the action under arbitrary variation $\delta\Phi$ leads to the Euler-Lagrange equations: $\frac{\delta\mathcal{L}}{\delta\Phi} = 0$. Since the equation we want to derive are linear, it suffices to have \mathcal{L} to be quadratic in the fields. Furthermore, the equations have no more than two derivatives which can be obtained by restricting to first derivatives in the Lagrangian.

Let us begin with the scalar field,

$$\begin{aligned} \phi_{\Lambda,a}(x) &= \phi(\Lambda^{-1}(x-a)) \quad , \quad \Lambda_\mu^\alpha \partial_\alpha \phi(\Lambda^{-1}(x-a)) \quad \Rightarrow \\ [\partial^\mu \phi \partial_\mu \phi]_{\Lambda,a} &= \eta^{\mu\nu} [\partial_\mu \phi \partial_\nu \phi]_{\Lambda,a} = \eta^{\mu\nu} \Lambda_\mu^\alpha \Lambda_\nu^\beta [\partial_\alpha \phi \partial_\beta \phi] (\Lambda^{-1}(x-a)) \\ &= \eta^{\alpha\beta} \partial_\alpha \phi \partial_\beta \phi (\Lambda^{-1}(x-a)) = [\partial^\alpha \phi \partial_\alpha \phi] (\Lambda^{-1}(x-a)) . \end{aligned}$$

When integrated over the space-time, we can change the dummy variable $x \rightarrow \Lambda x + a$ which restores the argument of the ϕ 's without changing the integration measure. Hence $\int d^4x \partial^\mu \phi \partial_\mu \phi$ is clearly Poincare invariant. Since all fields have the same shift of the space-time argument, we will suppress the translation part and focus on the Lorentz. This works as long as there are no externally prescribed fields/functions, which break translation in-

variance. Thus, let

$$S[\phi] := \int_M d^4x \left[-\frac{1}{2} \partial^\mu \phi \partial_\mu \phi - \frac{1}{2} m^2 \phi^2 \right] =: \int_M d^4x \mathcal{L}(\phi, \partial\phi) \quad (6.1)$$

$$\begin{aligned} \delta S[\phi] &= \int_M d^4x \left[-\partial_\mu \phi \partial^\mu \delta\phi - m^2 \phi \delta\phi \right] \\ &= \int_M d^4x \left[-\partial_\mu (\partial^\mu \phi \delta\phi) + (\square\phi - m^2\phi) \delta\phi \right] \\ &= \int_M d^4x \delta\phi \left[(\square - m^2)\phi \right] - \int_{\partial M} d^3x n_\mu (\partial^\mu \phi \delta\phi) \end{aligned} \quad (6.2)$$

The integration is over all of Minkowski space-time which has no boundary and hence the second term should be zero. In practice, such space-times extending to infinite coordinates are handled/defined putting the system in a large box and taking a limit. The fields must satisfy suitable boundary conditions so that the boundary contribution again vanishes. Alternatively requiring the field to vanish fast enough in the asymptotic regions, makes the boundary term vanish. We can always choose the variational principle to set $\delta\phi = 0$ at the boundary/asymptotic regions. While the boundary contribution are important in some context, we will restrict ourselves to boundary contribution being zero.

Demanding that the action be stationary for arbitrary variations of the field, we get the equation of motion $(\square - m^2)\phi = 0$. Incidentally, we will get the same equation of motion from another Lagrangian $\mathcal{L}' = \mathcal{L} + \partial_\mu \Lambda^\mu$. Thus, several Lagrangians can give the same equations of motion (although the Hamiltonian formulation will vary with the Lagrangians.)

Consider now a vector field. For the massive case, the equations we want are $(\square - m^2)v^\mu = 0 = \partial_\mu v^\mu$.

Consider $\mathcal{L} = aF^{\mu\nu}F_{\mu\nu} + bm^2v^\mu v_\mu$, where $F_{\mu\nu} := \partial_\mu v_\nu - \partial_\nu v_\mu$ and a, b are non-zero constants to be determined.

$$\begin{aligned} \delta\mathcal{L} &= 2aF^{\mu\nu}(\partial_\mu \delta v_\nu - \partial_\nu \delta v_\mu) + 2bm^2v^\mu \delta v_\mu = 4aF^{\mu\nu}\partial_\mu \delta v_\nu + 2bm^2v^\nu \delta v_\nu \\ &= \partial_\mu (4aF^{\mu\nu}\delta v_\nu) - 4a(\partial_\mu F^{\mu\nu})\delta v_\nu + 2bm^2v^\nu \delta v_\nu \Rightarrow -4a\partial_\mu F^{\mu\nu} + 2bm^2v^\nu = 0. \\ \therefore 0 &= \left(\square v^\nu - \frac{b}{2a}m^2v^\nu \right) - \partial^\nu \partial \cdot v \quad \text{taking divergence and choosing } b = 2a \text{ gives} \\ 0 &= (\square - m^2)v^\nu \text{ and } \partial \cdot v = 0. \end{aligned} \quad (6.3)$$

We got $b = 2a$ and we choose $b = -\frac{1}{2}$ similar to the scalar field case. The reason for this choice will be clear little later.

Note: For $m = 0$, we denote the vector field by A^μ . Now we cannot conclude $\partial \cdot A = 0$, and the equation we get is: $\partial_\mu F^{\mu\nu} = 0$ which is just the Maxwell equation. It has the well

known *gauge invariance*: $A_\mu \rightarrow A_\mu + \partial_\mu \Lambda$ which allows us to take one of the components of A_μ to be zero.

Exercise: Another natural choice of the Lagrangian is: $\mathcal{L}' = a\partial^\mu v^\nu \partial_\mu v_\nu + bm^2 v_\mu v^\mu + c(\partial_\mu v^\mu)^2$. Repeat the steps and deduce the choices for the constants so as to get the Proca equations. Check that $\mathcal{L}, \mathcal{L}'$ differ by a divergence.

Lastly, let us consider the action for getting the Dirac equation. We have already noted the Lorentz transformations of $\Psi, \bar{\Psi}, \gamma^\mu, \partial_\mu$. From these it follows that we can have the following Lorentz invariant terms, quadratic in the fields and with a single derivative: $\bar{\Psi}\gamma^\alpha\partial_\alpha\Psi, \partial^\alpha\bar{\Psi}\partial_\alpha\Psi, \bar{\Psi}\Psi$ and the same terms with a γ_5 inserted between the spinors. The terms with the γ_5 are all pseudo-scalars and may be dropped if parity (space-inversion plus a rotation) is required to be a symmetry. We assume so for the present. The candidate Lagrangian is then,

$$\mathcal{L} := a\bar{\Psi}\gamma^\alpha\partial_\alpha\Psi + b\partial^\alpha\bar{\Psi}\partial_\alpha\Psi + cm\bar{\Psi}\Psi \text{ and } \delta_{\bar{\Psi}}\mathcal{L} = \delta\bar{\Psi}[a\not{\partial}\Psi - b\Box\Psi + cm\Psi]. \quad (6.4)$$

For the choice $a = -i, b = 0$ and $c = 1$, we get the Dirac equation. If we varied ψ , then for the same choice, we will get the conjugate Dirac equation: $i\partial_\alpha\bar{\Psi}\gamma^\alpha + m\bar{\Psi} = 0$.

In summary, we have $S = \int_M d^4x \sqrt{|det\eta|} \mathcal{L}$ with,

$$\mathcal{L}_{scalar} = -\frac{1}{2}\partial^\mu\phi\partial_\mu\phi - \frac{1}{2}m^2\phi^2 ; \quad (6.5)$$

$$\mathcal{L}_{vector} = -\frac{1}{4}F^{\mu\nu}F_{\mu\nu} - \frac{1}{2}m^2v^\mu v_\mu ; \quad (6.6)$$

$$\mathcal{L}_{spinor} = -i\bar{\Psi}\gamma^\mu\partial_\mu\Psi + m\bar{\Psi}\Psi . \quad (6.7)$$

$\delta S = 0$ leads to the field equations implementing the irreducibility condition. For massless vector, the action is invariant under the gauge transformation: $A_\mu \rightarrow A_\mu + \partial_\mu \Lambda$. The actions are also invariant under parity.

Note: We have taken the scalar and vector fields to be real. We could have complex scalar field (say) with two field equations: $(\Box - m^2)\phi = 0 = (\Box - m^2)\phi^*$. The complex field may be considered as two real fields, $\phi = \phi_1 + i\phi_2$ and two terms may be included in the action. Alternatively, the complex scalar field equations may be derived from $\mathcal{L} = -\partial^\mu\phi^*\partial_\mu\phi - m^2\phi^*\phi$.

Notice that all actions are *real* (for spinors it is convenient to take matrix hermitian conjugate). We will always take the actions to be real so that $e^{\frac{i}{\hbar}S}$ will be a phase. Only when dissipation of energy/momentum/angular momentum is to be incorporated we need to take the action to be complex. In this course, we will not do so.

A. Variational Principle, Symmetries of the Action and Noether's theorem

Let us denote a generic field by X with all indices suppressed. Let an action be expressed as, $S[X] := \int_M d^4x \mathcal{L}(X, \partial X)$. Let $\delta X := \epsilon \chi(X(x))$ be an arbitrary, infinitesimal variation of the field $X(x)$. Then,

$$\begin{aligned} \delta S &= \int_M d^4x \left(\frac{\delta \mathcal{L}}{\delta X} \delta X + \frac{\delta \mathcal{L}}{\delta X_{,\mu}} \delta X_{,\mu} \right) = \int_M d^4x \left(\frac{\delta \mathcal{L}}{\delta X} \delta X + \frac{\delta \mathcal{L}}{\delta X_{,\mu}} \partial_{\mu} \delta X \right) \\ &= \int_M d^4x \left(\frac{\delta \mathcal{L}}{\delta X} - \partial_{\mu} \frac{\delta \mathcal{L}}{\delta X_{,\mu}} \right) \delta X + \int_M d^4x \partial_{\mu} \left(\frac{\delta \mathcal{L}}{\delta X_{,\mu}} \delta X \right) \end{aligned}$$

Here, $X_{,\mu} = \partial_{\mu} X$. The $\frac{\delta \mathcal{L}}{\delta X}$ is really like partial derivative, but since both \mathcal{L} and X depend on the coordinate, these should be mentioned too. We have kept these implicit and used the δ as a reminder. We follow this customary practice. The last term is a divergence and can be expressed as a boundary integral: $\int_{\partial M} d^3y n_{\mu} \frac{\delta \mathcal{L}}{\delta X_{,\mu}} \delta X$. This is typically dropped/vanishes for various reasons.

Note: For an action principle to be well defined (this includes the specification of the class of variations), it is necessary that the boundary contribution must vanish. If in some cases it does not vanish, additional ‘surface action’ needs to be added to cancel the total boundary contribution. This is typically encountered in gravitational actions on manifolds with boundaries.

To summarize: *a variational principle asserts that $\delta S[X]$ vanishes for arbitrary variation δX around X iff $\frac{\delta \mathcal{L}}{\delta X} - \partial_{\mu} \frac{\delta \mathcal{L}}{\delta X_{,\mu}} = 0$ i.e. iff $X(x)$ is a solution of the equation of motion.*

We now consider a different situation. We consider special, restricted variations such that $\delta S[X] = 0$ at *all fields* $X(x)$. Such variations are called *infinitesimal symmetries of the action*. This is translated in terms of the Lagrangian density as,

$$\delta S[X] = \int_M d^4x \delta \mathcal{L} = 0 \quad \text{if either } \delta \mathcal{L} = 0 \quad \text{or} \quad \delta \mathcal{L} = \partial_{\mu} \delta \Lambda^{\mu} ,$$

where Λ^{μ} is some 4-vector which vanishes/falls off on the boundary. Now we have the Noether's theorem:

Noether's Theorem: For every infinitesimal symmetry of the action, \exists a conserved current, conserved on every solution of the equation of motion.

Proof: We already have the infinitesimal variation of the \mathcal{L} which must equal a divergence i.e. for a symmetry variation,

$$\left(\frac{\delta\mathcal{L}}{\delta X} - \partial_\mu \frac{\delta\mathcal{L}}{\delta X_{,\mu}}\right) \delta X + \partial_\mu \left(\frac{\delta\mathcal{L}}{\delta X_{,\mu}} \delta X\right) = \delta\mathcal{L} = \partial_\mu \Lambda^\mu \Rightarrow,$$

$$\left(\frac{\delta\mathcal{L}}{\delta X} - \partial_\mu \frac{\delta\mathcal{L}}{\delta X_{,\mu}}\right) \delta X + \partial_\mu \left(\frac{\delta\mathcal{L}}{\delta X_{,\mu}} \delta X - \partial_\mu \Lambda^\mu\right) = 0.$$

Thus, if X satisfies the equation of motion, the first term vanishes and we have a conserved current, $\delta J^\mu := \frac{\delta\mathcal{L}}{\delta X_{,\mu}} \delta X - \partial_\mu \Lambda^\mu$, $\partial_\mu \delta J^\mu = 0$

This is neat prescription to discover conserved currents and their corresponding conserved charges: $\delta Q := \int_{\partial M} d^3\sigma n^\mu \delta J_\mu$. Discovering conserved currents by inspection of the equation of motion is easy only in the simplest of cases.

Let us see an example, particularly a symmetry variation induced by infinitesimal Poincare transformations for which $\delta S = 0$ by construction. Consider a real scalar field action for simplicity.

Under an infinitesimal Poincare transformation,

$$\begin{aligned} \delta\phi_{\Lambda,a}(x) &:= \phi(\Lambda^{-1}(x-a)) - \phi(x) = \phi((\delta^\mu_\nu - \omega^\mu_\nu)(x - \epsilon)^\nu) - \phi(x) \\ &= (-\omega^\mu_\nu x^\nu - \epsilon^\mu) \partial_\mu \phi := -\xi^\mu \partial_\mu \phi(x), \quad \xi^\mu := \omega^\mu_\nu x^\nu + \epsilon^\mu. \end{aligned}$$

This leads to

$$\begin{aligned} \delta\mathcal{L} &= -\partial^\mu \phi \partial_\mu (-\xi \cdot \partial\phi) - m^2 \phi (-\xi \cdot \partial\phi) = \partial^\mu \phi \partial_\mu (\xi^\nu \partial_\nu \phi) + m^2 \phi \xi^\nu \partial_\nu \phi \\ &= (\partial_\mu \xi^\nu) \partial^\mu \phi \partial_\nu \phi + \xi^\nu (\partial^\mu \phi \partial_{\nu\mu}^2 \phi + m^2 \phi \partial_\nu \phi) \\ &= \frac{1}{2} (\partial_\mu \xi_\nu + \partial_\nu \xi_\mu) \partial^\mu \phi \partial^\nu \phi + \xi^\nu \left(\frac{1}{2} \frac{\delta\mathcal{L}}{\delta\phi_{,\mu}} \partial_\nu \phi_{,\mu} + \frac{1}{2} m^2 \frac{\delta\mathcal{L}}{\delta\phi} \partial_\nu \phi \right) \end{aligned}$$

Explicitly, $\partial_\mu \xi_\nu = \partial_\mu (\omega_{\nu\alpha} x^\alpha + \epsilon^\alpha) = \omega_{\nu\mu} + 0$ which is antisymmetric in $\mu \leftrightarrow \nu$ and hence the first term vanishes. The second term is just $\xi^\nu \partial_\nu \mathcal{L}$.

$$\therefore \delta\mathcal{L} = \xi^\mu \partial_\mu \mathcal{L} = \partial_\mu (\xi^\mu \mathcal{L}) - (\partial \cdot \xi) \mathcal{L} = \partial_\mu \delta\Lambda^\mu + 0, \quad \text{or} \quad \boxed{\delta\mathcal{L} = \partial_\mu \delta\Lambda^\mu, \quad \delta\Lambda^\mu = \xi^\mu \mathcal{L}}.$$

Note: We ensured that \mathcal{L} is a Lorentz scalar, but it is *not* a translation scalar. Action gets its Poincare invariance thanks to the $\int d^4x$. Since all fields under translations shift their

coordinates by a , in all cases we will have $\delta\mathcal{L} = \xi \cdot \partial\mathcal{L}$ provided ξ_μ satisfies $\partial_\mu\xi_\nu + \partial_\nu\xi_\mu = 0$ i.e. ξ^μ is a ‘‘Killing vector’’ of the Minkowski metric.

By Noether’s theorem, we get

$$\delta J^\mu = \frac{\delta\mathcal{L}}{\delta\phi_{,\mu}}\delta\phi - \delta\Lambda^\mu = \partial^\mu\phi(-\xi^\nu\partial_\nu\phi) - \xi^\mu\mathcal{L} = -\xi_\nu(\partial^\mu\phi\partial^\nu\phi + \eta^{\mu\nu}\mathcal{L}) =: -\xi_\nu T^{\mu\nu} .$$

Conservation of the Noether current gives $(\partial_\mu\xi_\nu)T^{\mu\nu} + \xi_\nu\partial_\mu T^{\mu\nu} = 0$. Provided $T^{\mu\nu}$ is symmetric (as it is here), the first term vanishes and independence of ξ^μ , we get $\partial_\mu T^{\mu\nu} = 0$. The tensor, $\boxed{T^{\mu\nu} := \partial^\mu\phi\partial^\nu\phi + \eta^{\mu\nu}\mathcal{L}}$ is called the *canonical stress tensor* for the scalar field.

Thus, for each Killing vector ξ^μ , we have $J^\mu := T^{\mu\nu}\xi_\nu$ which is conserved on the solutions of the field equation. We define the corresponding charges as, $Q^\mu(\xi) := \int_{\Sigma_t} d^3x T^{0\nu}\xi_\nu$ which are independent of the hypersurfaces Σ_t .

Here are practice exercises:

(1) For the Proca Lagrangian, show that $\delta\mathcal{L} = \xi^\mu\partial_\mu\mathcal{L}$ for $\xi^\mu = \omega^\mu_\nu x^\nu + \epsilon^\mu$, ω, ϵ being infinitesimal, and obtain the canonical stress tensor.

(2) For $m = 0$, show that this stress tensor is not gauge invariant and one needs to ‘‘improve’’ it to get a symmetric, gauge invariant and conserved stress tensor. Guess it.

(3) For the scalar and the Maxwell field, obtain the conserved charges corresponding to the 10 Killing vectors. The charges corresponding to the translations are the energy and momentum while those corresponding to the ω_{ij} are the angular momentum components.

B. Conserved Poincare charges for scalar field solutions

Let us see an explicit example of Noether charges for a general solution of the Klein-Gordon equation. We noted above that the canonical stress tensor for the scalar field is given by,

$$\begin{aligned} T_{\mu\nu} &= \partial_\mu\phi\partial_\nu\phi + \eta_{\mu\nu}\mathcal{L} \quad , \quad \mathcal{L} := -\frac{1}{2}\partial^\mu\phi\partial_\mu\phi - \frac{1}{2}m^2\phi^2 \\ \partial_\mu T^{\mu\nu} &= 0 \quad , \quad \forall \quad (\square - m^2)\phi = 0 . \end{aligned}$$

$$\text{Define: } M^{\mu\nu\lambda} := T^{\mu\nu}x^\lambda - T^{\mu\lambda}x^\nu \Rightarrow \partial_\mu M^{\mu\nu\lambda} = T^{\lambda\nu} - T^{\nu\lambda}$$

$$\therefore \partial_\mu M^{\mu\nu\lambda} = 0 \text{ on solutions} \quad \text{iff} \quad \partial_\mu T^{\mu\nu} = 0 \text{ and } T^{\mu\nu} = T^{\nu\mu}$$

These 6 conserved currents ($M^{\mu\nu\lambda}$ are antisymmetric in the last two indices) together with the conserved stress tensor give the 10 Poincare charges:

$$P^\mu := \int_{\Sigma_t} d^3x T^{0\mu} \quad , \quad M^{\mu\nu} := \int_{\Sigma_t} d^3x M^{0\mu\nu} .$$

We had noted the plane wave solutions of the Klein-Gordon equation, labeled by $\vec{k} \in \mathbb{R}^3$ namely,

$$u_{\vec{k}}(x) = \frac{1}{\sqrt{2\omega_{\vec{k}}(2\pi)^3}} e^{ik \cdot x} \quad , \quad k \cdot x := -\omega_{\vec{k}}t + \vec{k} \cdot \vec{x} \quad , \quad \omega_{\vec{k}} := \sqrt{\vec{k}^2 + m^2}$$

and their complex conjugates. The general solution is thus expressed as $\phi(x) = \int d^3k [a(\vec{k})u_{\vec{k}}(x) + a^*(\vec{k})u_{\vec{k}}^*(x)]$ with $a(\vec{k})$ being complex numbers with a suitable dependence on \vec{k} so that the field has the appropriate boundary behavior. The general solution is manifestly real.

To compute the energy-momentum and angular momentum charges, we have integration of expressions quadratic in the solutions. So we note: $\partial_\mu u_{\vec{k}}(x) = ik_\mu u_{\vec{k}}(x)$, $\partial_\mu u_{\vec{k}}^*(x) = -ik_\mu u_{\vec{k}}^*(x)$, $k^0 = \omega_{\vec{k}}$ and the orthogonality relations:

$$\begin{aligned} \int d^3x u_{\vec{k}}^*(x) u_{\vec{k}'}(x) &= \frac{e^{i(\omega_{\vec{k}} - \omega_{\vec{k}'})t}}{\sqrt{2\omega_{\vec{k}}}\sqrt{2\omega_{\vec{k}'}}} = \frac{\delta^3(\vec{k} - \vec{k}')}{2\omega_{\vec{k}}} = \delta_{inv}^3(\vec{k} - \vec{k}') \\ \int d^3x u_{\vec{k}}(x) u_{\vec{k}'}(x) &= e^{-2i\omega_{\vec{k}}t} \delta_{inv}^3(\vec{k} + \vec{k}') \quad , \quad \int d^3x u_{\vec{k}}^*(x) u_{\vec{k}'}^*(x) = e^{2i\omega_{\vec{k}}t} \delta_{inv}^3(\vec{k} + \vec{k}') . \end{aligned}$$

We need,

$$P^\mu = \int d^3x \partial^0 \phi \partial^\mu \phi + \eta^{0\mu} \left(-\frac{1}{2} \partial^\alpha \phi \partial_\alpha \phi - \frac{1}{2} m^2 \phi^2 \right) \quad (6.8)$$

$$M^{\mu\nu} = \int d^3x \{ (\partial^0 \phi \partial^\mu \phi + \eta^{0\mu} \mathcal{L}) x^\nu - (\partial^0 \phi \partial^\nu \phi + \eta^{0\nu} \mathcal{L}) x^\mu \} \quad (6.9)$$

And we have,

$$\begin{aligned} \mathcal{L} = -\frac{1}{2} \int d^3k \int d^3k' \{ & (a(\vec{k})a(\vec{k}')u_{\vec{k}}u_{\vec{k}'} + a^*(\vec{k})a^*(\vec{k}')u_{\vec{k}}^*u_{\vec{k}'}^*) (-\vec{k} \cdot \vec{k}' + m^2) \\ & (a(\vec{k})a^*(\vec{k}')u_{\vec{k}}u_{\vec{k}'}^* + a^*(\vec{k})a(\vec{k}')u_{\vec{k}}^*u_{\vec{k}'}) (+\vec{k} \cdot \vec{k}' + m^2) \} \quad (6.10) \end{aligned}$$

$$\begin{aligned} \partial^0 \phi \partial^\mu \phi = \int d^3k \int d^3k' \{ & a(\vec{k})a(\vec{k}')u_{\vec{k}}u_{\vec{k}'} + a^*(\vec{k})a^*(\vec{k}')u_{\vec{k}}^*u_{\vec{k}'}^* \\ & - a(\vec{k})a^*(\vec{k}')u_{\vec{k}}u_{\vec{k}'}^* - a^*(\vec{k})a(\vec{k}')u_{\vec{k}}^*u_{\vec{k}'} \} (-\vec{k}^0 \vec{k}'^\mu) \quad (6.11) \end{aligned}$$

Substituting and carrying out the $\int d^3x$ using the orthogonality relations gives,

$$\begin{aligned}
P^i &= \int d^3k \int d^3k' \left[\frac{k^0 k^i}{2\omega_{\vec{k}}} \left\{ a(\vec{k})a(-\vec{k})e^{-2i\omega_{\vec{k}}t} + a^*(\vec{k})a^*(-\vec{k})e^{2i\omega_{\vec{k}}t} \right\} \delta^3(\vec{k} + \vec{k}') \right. \\
&\quad \left. \frac{k^0 k^i}{2\omega_{\vec{k}}} \left\{ a(\vec{k})a^*(\vec{k}) + a^*(\vec{k})a(\vec{k}) \right\} \delta^3(\vec{k} - \vec{k}') \right] \\
&= \frac{1}{2} \int d^3k k^i \left\{ a(\vec{k})a(-\vec{k})e^{-2i\omega_{\vec{k}}t} + a^*(\vec{k})a^*(-\vec{k})e^{2i\omega_{\vec{k}}t} + a(\vec{k})a^*(\vec{k}) + a^*(\vec{k})a(\vec{k}) \right\} \quad (6.12)
\end{aligned}$$

The first two, time decedent terms in the braces are symmetric under $\vec{k} \leftrightarrow -\vec{k}$ while the last two time independent term go just change their argument. Hence, under symmetric integration, the time dependent terms vanish and we are left with,

$$P^i = \int d^3k k^i \frac{a(\vec{k})a^*(\vec{k}) - a(-\vec{k})a^*(-\vec{k})}{2} \quad (6.13)$$

Since the a 's are ordinary complex numbers the factor of 2 cancels and a new factor arises from explicit anti-symmetrization w.r.t. \vec{k} . The claimed conserved quantity is manifestly independent of t .

The calculation of P^0 proceeds much the same way. We have the t -dependent term proportional to $\delta^3(\vec{k} + \vec{k}')$ while the t -independent term is proportional to $\delta^3(\vec{k} - \vec{k}')$. We get,

$$\begin{aligned}
P^0 &= \int \frac{d^3k}{2\omega_{\vec{k}}} \left[-\omega_{\vec{k}}^2 \left\{ a(\vec{k})a(-\vec{k})e^{-2i\omega_{\vec{k}}t} + a^*(\vec{k})a^*(-\vec{k})e^{2i\omega_{\vec{k}}t} \right\} + \omega_{\vec{k}}^2 \left\{ 2a(\vec{k})a^*(\vec{k}) \right\} \right. \\
&\quad + \frac{1}{2} \left\{ a(\vec{k})a(-\vec{k})e^{-2i\omega_{\vec{k}}t} + a^*(\vec{k})a^*(-\vec{k})e^{2i\omega_{\vec{k}}t} \right\} \left(\omega_{\vec{k}}^2 + \vec{k}^2 + m^2 \right) \\
&\quad \left. + \frac{1}{2} \left\{ 2a(\vec{k})a^*(\vec{k}) \right\} \left(-\omega_{\vec{k}}^2 + \vec{k}^2 + m^2 \right) \right] \quad (6.14)
\end{aligned}$$

The t -dependent terms cancel while the t -independent terms give,

$$P^0 = \int d^3k \omega_{\vec{k}} a(\vec{k})a^*(\vec{k}) \quad (6.15)$$

The calculation of $M^{\mu\nu}$ is similar. The new feature is the explicit x^λ in the integral. In the M^{ij} we have $\int T^{0i}x^j - T^{0j}x^i$. We trade x^i for ∂_{k_i} by noting that

$$\partial_{k_i} u_{\vec{k}}(x) = u_{\vec{k}}(x) \left\{ -it \frac{k^j}{\omega_{\vec{k}}} + ix^j \right\} \quad \text{or} \quad x^j u_{\vec{k}} = -i\partial_{k_j} u + \frac{tk^j}{\omega_{\vec{k}}} u_{\vec{k}} .$$

As before the spatial integration will produce δ functions and also derivatives of δ functions from the ∂_{k_i} . While doing the k' -integration, we need to flip the derivative as per the rules

of δ function. As before, the $e^{\pm 2i\omega_{\vec{k}}t}$ will turn out to be symmetric under $\vec{k} \leftrightarrow -\vec{k}$ and will not contribute. Filling in the algebra leads to the final results,

$$M^{ij} = i \int d^3k \left[a(\vec{k}) (k^i \partial_{k_j} - k^j \partial_{k_i}) a_{\vec{k}}^* - a^*(\vec{k}) (k^i \partial_{k_j} - k^j \partial_{k_i}) a_{\vec{k}} \right] \quad (6.16)$$

$$M^{0i} = -\frac{i}{2} \int d^3k \omega_{\vec{k}} \{ a_{\vec{k}} \partial_{k_i} a_{\vec{k}}^* - a_{\vec{k}}^* \partial_{k_i} a_{\vec{k}} \} . \quad (6.17)$$

All are manifestly independent of time.

Apart from verifying that the conserved quantities are indeed time independent, the expressions for energy and momentum show that these quantities are a sum (integral) of energy-momentum of individual solutions. Thus each of these plane waves, at least heuristically be thought of as carrying an energy $\omega_{\vec{k}}$ and momentum \vec{k} . Note that there is no \hbar yet so these are mathematically defined conserved quantities much like the energy-momentum fluxes of electromagnetic fields identified via the Poynting theorem. This is helpful for interpreting the solutions of the field equations as physical entities carrying energy-momentum. To strengthen this further, we would like to see if the fields and the action can be cast in the form of a “dynamical system”.

7. FOURIER DECOMPOSITIONS OF FIELDS: COLLECTION OF HARMONIC OSCILLATORS

In the previous section, we wrote the general solution of the Klein-Gordon equation. Now we want to see if the action takes the ‘form of a dynamical system’. The meaning will be clear shortly. Consider again the scalar field. Let Σ_t be a constant t surface. On Σ_t , the set of functions, $\{\varphi_{\vec{k}} = \frac{1}{(2\pi)^{3/2}} e^{i\vec{k}\cdot\vec{x}}, \vec{k} \in \mathbb{R}^3\}$ forms a complete, orthonormal set of functions, $\int_{\Sigma_t} d^3x \varphi_{\vec{k}}^*(\vec{x}) \varphi_{\vec{k}'}(\vec{x}) = \delta^3(\vec{k} - \vec{k}')$. Let us *Fourier decompose* the field as,

$$\phi(t, \vec{x}) := \int_{\mathbb{R}^3} d^3k f_{\vec{k}}(t) \varphi_{\vec{k}}(\vec{x}). \quad (7.1)$$

The reality of the field, $\phi^*(x) = \phi(x) \Rightarrow f_{\vec{k}}^*(t) = f_{-\vec{k}}(t)$. The expansion coefficient functions are determined by the equations of motion. It is trivial to check that $(\square - m^2)\phi(x) = 0 \Rightarrow \int d^3k (-\ddot{f}_{\vec{k}} - \vec{k}^2 f_{\vec{k}} - m^2 f_{\vec{k}}) \varphi_{\vec{k}}(x) = 0$. By independence of the basis functions, we get $\ddot{f}_{\vec{k}} + \omega_{\vec{k}}^2 f_{\vec{k}} = 0 \forall \vec{k} \in \mathbb{R}^3, \omega_{\vec{k}}^2 := \vec{k}^2 + m^2$. Its general solution is $f_{\vec{k}}(t) = a_{\vec{k}} e^{-i\omega_{\vec{k}} t} + b_{\vec{k}} e^{i\omega_{\vec{k}} t}$. The reality condition $f_{\vec{k}}^* = f_{-\vec{k}}$ gives $a_{\vec{k}}^* = b_{-\vec{k}}$ or $b_{\vec{k}} = a_{-\vec{k}}^*$. Substitution gives,

$$\begin{aligned} \phi_{soln}(x) &= \int_{\mathbb{R}^3} d^3k (a_{\vec{k}} e^{-i\omega_{\vec{k}} t} + b_{\vec{k}} e^{i\omega_{\vec{k}} t}) \frac{e^{i\vec{k}\cdot\vec{x}}}{\sqrt{2\omega_{\vec{k}}(2\pi)^3}} \\ &= \int \frac{d^3k}{\sqrt{2\omega_{\vec{k}}(2\pi)^3}} \left(a_{\vec{k}} e^{-i\omega_{\vec{k}} t + i\vec{k}\cdot\vec{x}} + b_{-\vec{k}} e^{i\omega_{\vec{k}} t - i\vec{k}\cdot\vec{x}} \right) \\ \therefore \phi_{soln}(x) &= \int d^3k (a_{\vec{k}} u_{\vec{k}}(x) + a_{\vec{k}}^* u_{\vec{k}}^*(x)) \quad , \quad u_{\vec{k}} := \frac{e^{ik\cdot x}}{\sqrt{2\omega_{\vec{k}}(2\pi)^3}} \end{aligned} \quad (7.2)$$

This is exactly the same form we had for the general solution. We have adjusted the normalization factor for future convenience. The last equation will be referred to as a *mode decomposition*, with $u_{\vec{k}}(x)$ ’s denoting the *mode functions*.

Let us rewrite the Fourier decomposition in a manifestly real form:

$$\phi(t, \vec{x}) = \int_{\mathbb{R}^3/2} d^3k [f_{\vec{k}}(t) \varphi_{\vec{k}}(\vec{x}) + f_{\vec{k}}^*(t) \varphi_{\vec{k}}^*(\vec{x})]$$

We have separated the $-\vec{k}$ tagged terms and used the reality condition. The integration is

over *positive half* of \mathbb{R}^3 . Substituting the Fourier decomposition into the Lagrangian, we get,

$$\begin{aligned}
L &= \int_{\Sigma_t} d^3x \mathcal{L} = \int_{\Sigma_t} d^3x \left[-\frac{1}{2} \left(-\dot{\phi}^2 + (\nabla\phi)^2 \right) - \frac{m^2}{2} \phi^2 \right] \\
\dot{\phi}(t, \vec{x}) &= \int_{\mathbb{R}^3/2} d^3k \left[\dot{f}_{\vec{k}}(t) \varphi_{\vec{k}}(\vec{x}) + \dot{f}_{\vec{k}}^*(t) \varphi_{\vec{k}}^*(\vec{x}) \right] \\
\partial_j \phi(t, \vec{x}) &= \int_{\mathbb{R}^3/2} d^3k (ik_j) \left[f_{\vec{k}}(t) \varphi_{\vec{k}}(\vec{x}) - f_{\vec{k}}^*(t) \varphi_{\vec{k}}^*(\vec{x}) \right] \quad \text{using ortho-normality, we get} \\
L &= \int_{\mathbb{R}^3/2} d^3k \left[\dot{f}_{\vec{k}} \dot{f}_{\vec{k}}^* - \omega_{\vec{k}}^2 f_{\vec{k}} f_{\vec{k}}^* \right] \quad , \quad \omega_{\vec{k}}^2 = \vec{k}^2 + m^2 \quad . \quad (7.3)
\end{aligned}$$

Let $f_{\vec{k}}(t) := \frac{1}{\sqrt{2}}(q_{\vec{k}}(t) + iq'_{\vec{k}}(t))$, where q, q' are real and \vec{k} are in the positive half of \mathbb{R}^3 . Substitution in the Lagrangian gives,

$$L = \int_{\mathbb{R}^3/2} d^3k \frac{1}{2} \left[(\dot{q}_{\vec{k}}^2 - \omega_{\vec{k}}^2 q_{\vec{k}}^2) + (\dot{q}'_{\vec{k}}^2 - \omega_{\vec{k}}^2 q'_{\vec{k}}^2) \right] .$$

Denoting $q'_{\vec{k}} =: q_{-\vec{k}}$, we can express the Lagrangian as,

$$L = \int_{\mathbb{R}^3} d^3k \frac{1}{2} \left[(\dot{q}_{\vec{k}}^2 - \omega_{\vec{k}}^2 q_{\vec{k}}^2) \right] . \quad (7.4)$$

\vec{k} now range over full \mathbb{R}^3 .

The Lagrangian is manifestly expressed as a sum (integral) of Lagrangians for harmonic oscillators $q_{\vec{k}}$ each with frequency $\omega_{\vec{k}}$ for $\vec{k} \in \mathbb{R}^3$. One can easily pass to the Hamiltonian form by defining $p_{\vec{k}} := \frac{\partial L}{\partial \dot{q}_{\vec{k}}} = \dot{q}_{\vec{k}}$. We get,

$$H = \int_{\mathbb{R}^3} d^3k \left[\frac{1}{2} (p_{\vec{k}}^2 + \omega_{\vec{k}}^2 q_{\vec{k}}^2) \right] .$$

To complete the canonical form, introduce the usual basic Poisson brackets: $\{q_{\vec{k}}, q_{\vec{k}'}\} = 0 = \{p_{\vec{k}}, p_{\vec{k}'}\}$ and $\{q_{\vec{k}}, p_{\vec{k}'}\} = \delta^3(\vec{k} - \vec{k}')$, $\vec{k} \in \mathbb{R}^3$.

Introduce the field, $\pi(t, \vec{x}) := \dot{\phi}(t, \vec{x})$ and *define* a Poisson brackets among the fields ϕ, π fields from the Poisson brackets among the $q_{\vec{k}}$ and $p_{\vec{k}}$ using the Fourier decompositions. Noting that $f_{\vec{k}} = \frac{1}{\sqrt{2}}(q_{\vec{k}} + iq_{-\vec{k}})$ and $\dot{f}_{\vec{k}} = \frac{1}{\sqrt{2}}(p_{\vec{k}} + ip_{-\vec{k}})$, $\vec{k} \in \mathbb{R}^3/2$. It follows,

$$\begin{aligned}
\{\phi(t, \vec{x}), \pi(t, \vec{y})\} &= \int_{\mathbb{R}^3/2} d^3k \int_{\mathbb{R}^3/2} d^3k' \{f_{\vec{k}} \varphi_{\vec{k}}(\vec{x}) + f_{\vec{k}}^* \varphi_{\vec{k}}^*(\vec{x}), \dot{f}_{\vec{k}'} \varphi_{\vec{k}'}(\vec{y}) + \dot{f}_{\vec{k}'}^* \varphi_{\vec{k}'}^*(\vec{y})\} \\
&= \int_{\mathbb{R}^3/2} d^3k \int_{\mathbb{R}^3/2} d^3k' \\
&\quad \left[\varphi_{\vec{k}}(\vec{x}) \varphi_{\vec{k}'}(\vec{y}) \{f_{\vec{k}}, f_{\vec{k}'}\} + \varphi_{\vec{k}}(\vec{x}) \varphi_{\vec{k}'}^*(\vec{y}) \{f_{\vec{k}}, f_{\vec{k}'}^*\} \right. \\
&\quad \left. + \varphi_{\vec{k}}^*(\vec{x}) \varphi_{\vec{k}'}(\vec{y}) \{f_{\vec{k}}^*, f_{\vec{k}'}\} + \varphi_{\vec{k}}^*(\vec{x}) \varphi_{\vec{k}'}^*(\vec{y}) \{f_{\vec{k}}^*, f_{\vec{k}'}^*\} \right] \\
&= \int_{\mathbb{R}^3/2} d^3k \left[\varphi_{\vec{k}}(\vec{x}) \varphi_{\vec{k}}^*(\vec{y}) + \varphi_{\vec{k}}^*(\vec{x}) \varphi_{\vec{k}}(\vec{y}) \right] = \int_{\mathbb{R}^3} d^3k \varphi_{\vec{k}}(\vec{x}) \varphi_{\vec{k}}^*(\vec{y}) \quad (7.5)
\end{aligned}$$

$$\therefore \{\phi(t, \vec{x}), \pi(t, \vec{y})\} = \delta^3(\vec{x} - \vec{y}) \quad (7.6)$$

The remaining Poisson brackets $\{\phi, \phi\}, \{\pi, \pi\}$, defined similarly, are zero.

Note: The same t is taken for the fields since the q 's and p 's are also at the same time.

Starting from the Lagrangian, using the Fourier decomposition, we saw explicitly that the Lagrangian can be expressed as a sum of Lagrangians of harmonic oscillators. Furthermore, we could define Poisson brackets among the fields *from* the same oscillator system. It is apparent now that *a field satisfying the Klein-Gordon equation as irreducibility condition, can be given a canonical formulation wherein it appears as a system of infinitely many, uncoupled harmonic oscillators.*

Note: The integration domains appearing above can be confusing. For real valued fields, using a *manifestly real form* of a Fourier decomposition, the $\vec{k} \in \mathbb{R}^3/2$. Once the *solutions for the time dependent coefficients* of the Fourier decomposition are substituted back in the Fourier decomposition, we get the *mode decomposition* of the field, with $\vec{k} \in \mathbb{R}$. For complex valued fields, in both Fourier and mode decompositions, $\vec{k} \in \mathbb{R}^3$.

We now repeat the exercise for the Maxwell field and the Dirac field. We will use the same orthonormal set of $\varphi_{\vec{k}}(\vec{x})$ in developing the Fourier decomposition. The steps being very similar, we will be brief and suppress the integration range of the \vec{k} .

A. Maxwell Field

Knowing the plane wave solutions having two polarizations, we write the Fourier decomposition of the vector field $A_\mu(t, \vec{x})$ as,

$$A_\mu(t, \vec{x}) = \int d^3k \varepsilon_\mu(\vec{k}, a) f_{\vec{k}, a}(t) \varphi_{\vec{k}}(\vec{x}) \quad , \quad \varphi_{\vec{k}}(\vec{x}) = \frac{e^{i\vec{k}\cdot\vec{x}}}{(2\pi)^{3/2}} .$$

The ε_μ is a polarization vector which depends on \vec{k} while the label a enumerates the numbers of polarization vectors. A priori, there are *four* independent polarization vectors (a tetrad) and a takes 4 values. One of the Maxwell equations fixes the 0^{th} component of all polarization vectors in terms of the spatial components. One of the second set of equations is trivially true for the *longitudinal* polarization, $\vec{\varepsilon}(\vec{k}) \propto \vec{k}$ and we are left with only two independent equations for the $f_{\vec{k}, a}$ corresponding to the transverse polarizations. We could have kept the $f(t, a)$ inside the polarization vector, but this is more convenient. We will suppress the sum-over- a till the final expression.

This leads to,

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu = \int d^3k [\varepsilon_\nu \partial_\mu (f\varphi) - \varepsilon_\mu \partial_\nu (f\varphi)].$$

$$\therefore F_{0i} = \int d^3k \varphi_{\vec{k}}(\vec{x}) \left\{ \varepsilon_i \dot{f}_{\vec{k}} - ik_i \varepsilon_0 f_{\vec{k}} \right\}, \quad F_{ij} = \int d^3k f_{\vec{k}} \varphi_{\vec{k}}(\vec{x}) \{ i(k_i \varepsilon_j - k_j \varepsilon_i) \}$$

The equations of motion, using independence of $\varphi_{\vec{k}}(\vec{x})$ give,

$$\begin{aligned} \partial^i F_{i0} = 0 & : \varepsilon_0(\vec{k}, a) f_{\vec{k}}(t) = -\frac{i\vec{k} \cdot \vec{\varepsilon}}{k^2} \dot{f}_{\vec{k}}(t) \Rightarrow \varepsilon_0 \dot{f}_{\vec{k}} = -\frac{i\vec{k} \cdot \vec{\varepsilon}}{k^2} \ddot{f}_{\vec{k}}(t) \\ \partial^0 F_{0i} + \partial^j F_{ji} = 0 & : \varepsilon_i \left(-\ddot{f}_{\vec{k}} - k^2 f_{\vec{k}} \right) + ik_i (\varepsilon_0 \dot{f}_{\vec{k}}) + (\vec{k} \cdot \vec{\varepsilon}) k_i f_{\vec{k}} = 0 \\ \therefore 0 & = \ddot{f}_{\vec{k}} \left\{ -\varepsilon_i + \frac{k_i k^j \varepsilon_j}{k^2} \right\} + k^2 f_{\vec{k}} \left\{ -\varepsilon_i + \frac{k_i k^j \varepsilon_j}{k^2} \right\} \\ \text{or,} \quad 0 & = \left[\left(-\delta_i^j + \frac{k_i k^j}{k^2} \right) \varepsilon_j \right] \left[\ddot{f}_{\vec{k}} + k^2 f_{\vec{k}} \right]. \end{aligned}$$

The prefactor is projector, projecting a 3-vector onto the plane perpendicular to \vec{k} . Thus,

$\boxed{\underline{\varepsilon}_i(\vec{k}, a) := \left(\delta_i^j - \frac{k_i k^j}{k^2} \right) \varepsilon_j(\vec{k}, a)}$ define *the transverse polarizations* and there are two independent ones. Since the prefactor is non-zero, the Maxwell equations imply that $f_{\vec{k}}$ satisfy the same harmonic oscillator equation as before, with $\omega_k^2 = k^2$ and the mass is zero. It is important to remember that transverse polarization, $\underline{\varepsilon}_i(\vec{k}, a)$, is defined by \vec{k} and hence acquire the \vec{k} dependence as well as the label a . Hence the $f_{\vec{k},a}$ satisfy the oscillator equations only for the transverse polarization. We may not display it always to avoid clutter.

The next task is to compute the Lagrangian, $\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} = \frac{1}{2} (F_{0i})^2 - \frac{1}{4} (F_{ij})^2$.

$$F_{0i}^2 = \int d^3k \int d^3k' \left\{ \varepsilon_i \dot{f}_{\vec{k}} - ik_i \varepsilon_0 \right\} \left\{ \varepsilon'_i \dot{f}_{\vec{k}'} - ik'_i \varepsilon'_0 f_{\vec{k}} \right\} \varphi_{\vec{k}}(\vec{x}) \varphi_{\vec{k}'}(\vec{x})$$

Eliminating $\varepsilon_0 f_{\vec{k}} = -i \frac{\vec{k} \cdot \vec{\varepsilon}}{k^2} \dot{f}_{\vec{k}}$ and likewise $\varepsilon'_0 f_{\vec{k}'}$, simplifies the product of the $\{ \dots \}$ to³,

$$\{ \dots \} \{ \dots \} = \dot{f}_{\vec{k}} \dot{f}_{\vec{k}'} \left[\vec{\varepsilon} \cdot \vec{\varepsilon}' + \vec{k} \cdot \vec{k}' \frac{\vec{k} \cdot \vec{\varepsilon}}{k^2} \frac{\vec{\varepsilon}' \cdot \vec{k}'}{k'^2} - (\vec{\varepsilon} \cdot \vec{k}') \frac{\vec{\varepsilon}' \cdot \vec{k}'}{(k')^2} - (\vec{\varepsilon}' \cdot \vec{k}) \frac{\vec{\varepsilon} \cdot \vec{k}}{(k)^2} \right]$$

³ This is equivalent to using one of the Maxwell equations, the Gauss law equation which is a constraint.

Integration over \vec{x} gives $\delta^3(\vec{k} + \vec{k}')$ which cancels two term and we are left with,

$$\begin{aligned} \frac{1}{2} \int_{\Sigma_t} d^3x (F_{0i})^2 &= \frac{1}{2} \int d^3k \left[\vec{\varepsilon}(\vec{k}) \cdot \vec{\varepsilon}'(-\vec{k}) - \frac{(\vec{\varepsilon}(\vec{k}) \cdot \vec{k})(\vec{\varepsilon}'(-\vec{k}) \cdot \vec{k})}{\vec{k}^2} \right] \dot{f}_{\vec{k}} \dot{f}_{-\vec{k}} \\ \text{But } [\dots] &= \varepsilon_i(\vec{k}) \left(\delta^{ij} - \frac{k^i k^j}{\vec{k}^2} \right) \varepsilon_j(-\vec{k}) = \underline{\underline{\varepsilon}}(\vec{k}) \cdot \underline{\underline{\varepsilon}}(-\vec{k}) \\ \therefore \frac{1}{2} \int_{\Sigma_t} d^3x (F_{0i})^2 &= \int d^3k \frac{1}{2} \underline{\underline{\varepsilon}}(\vec{k}) \cdot \underline{\underline{\varepsilon}}(-\vec{k}) \dot{f}_{\vec{k}} \dot{f}_{-\vec{k}} \quad \text{and similarly,} \\ -\frac{1}{4} \int d^3x (F_{ij})^2 &= \int d^3k -\frac{\vec{k}^2}{2} \underline{\underline{\varepsilon}}(\vec{k}) \cdot \underline{\underline{\varepsilon}}(-\vec{k}) \dot{f}_{\vec{k}} \dot{f}_{-\vec{k}} \end{aligned}$$

The dot product of the transverse polarization includes the sum over the polarizations. The longitudinal polarization has dropped out explicitly. Making it explicit, we have,

$$L_{Maxwell} = \int_{\mathbb{R}^3/2} d^3k \sum_{a,b=1}^2 \left[\underline{\underline{\varepsilon}}(\vec{k}, a) \cdot \underline{\underline{\varepsilon}}(-\vec{k}, b) \right] \left[\dot{f}_{\vec{k},a} \dot{f}_{-\vec{k},b} - \omega_k^2 f_{\vec{k},a} f_{-\vec{k},b} \right] \quad (7.7)$$

We may now choose the transverse polarizations such that $\underline{\underline{\varepsilon}}(\vec{k}, a) \cdot \underline{\underline{\varepsilon}}(-\vec{k}, b) = \delta_{a,b}$ (note the $\pm\vec{k}$ for both polarizations). Using this, the Maxwell action becomes,

$$S[A] = \int dt \int_{\mathbb{R}^3/2} d^3k \sum_{a=1,2} \left[\dot{f}_{\vec{k},a} \dot{f}_{-\vec{k},a} - \omega_k^2 f_{\vec{k},a} f_{-\vec{k},a} \right] \quad (7.8)$$

$$= \int dt \int_{\mathbb{R}^3} d^3k \frac{1}{2} \sum_{a=1,2} \left[\dot{q}_{\vec{k},a}^2 - \omega_k^2 q_{\vec{k},a}^2 \right] \quad \text{for,} \quad (7.9)$$

$$A_\mu(t, \vec{x}) = \int_{\mathbb{R}^3/2} d^3k \sum_a \varepsilon_\mu(\vec{k}, a) f_{\vec{k},a}(t) \varphi_{\vec{k}}(\vec{x}) + \varepsilon_\mu^*(\vec{k}, a) f_{\vec{k},a}^*(t) \varphi_{\vec{k}}^*(\vec{x}) \quad \text{with,} \quad (7.10)$$

$$\varepsilon_0(\vec{k}, a) f_{\vec{k},a}(t) = -i \frac{\vec{k} \cdot \vec{\varepsilon}(\vec{k}, a)}{\vec{k}^2} \dot{f}_{\vec{k},a} \quad , \quad \underline{\underline{\varepsilon}}_i(\vec{k}, a) := \left(\delta_i^j - \frac{k_i k^j}{\vec{k}^2} \right) \varepsilon_j(\vec{k}, a) \quad (7.11)$$

$$\delta_{a,b} = \underline{\underline{\varepsilon}}(\vec{k}, a) \cdot \underline{\underline{\varepsilon}}^*(\vec{k}, b) \quad (\text{Normalization}). \quad (7.12)$$

As in the case of the scalar, we have gone through the introducing the ‘real $q_{\vec{k}a}$ ’ degrees of freedom and expressed the Maxwell action too is a sum of *twice as many* harmonic oscillators. The canonical form goes through as well.

Note: If we were to begin with the Maxwell action and attempt to get its Hamiltonian formulation, we would obtain the Gauss Law equation of motion as a constraint equation. By using the Fourier decomposition, we have explicitly solved this constraint and eliminated the ε_0 polarization. Once this is done, we get the transverse projector which eliminates the longitudinal polarization. This gives the action involving only the physical degrees of freedom.

B. Dirac Field

Lastly, let us consider the Dirac action. Here the Lagrangian is first order in the derivatives and we do not expect a harmonic oscillator form. We will also find it clearer to write the Fourier decomposition using the positive half in \vec{k} space. We take the Fourier decomposition of the spinor as,

$$\psi(t, \vec{x}) = \int_{\mathbb{R}^3/2} d^3k \left[\sum_{\sigma} u(\vec{k}, \sigma) f_{\vec{k},\sigma}(t) \right] \varphi_{\vec{k}}(\vec{x}) + \left[\sum_{\sigma} v(\vec{k}, \sigma) g_{\vec{k},\sigma}(t) \right] \varphi_{\vec{k}}^*(\vec{x})$$

Substitution in the Dirac equation, keeping summation over σ implicit, gives,

$$0 = -i\gamma^{\mu} \partial_{\mu} \psi + m\psi = \int d^3k \left[\left\{ -i\gamma^0 u_{\vec{k}} \dot{f}_{\vec{k}} + (\vec{k} \cdot \vec{\gamma} + m) u_{\vec{k}} f_{\vec{k}} \right\} \varphi_{\vec{k}} \right. \\ \left. + \left\{ -i\gamma^0 v_{\vec{k}} \dot{g}_{\vec{k}} + (-\vec{k} \cdot \vec{\gamma} + m) v_{\vec{k}} g_{\vec{k}} \right\} \varphi_{\vec{k}}^* \right]$$

Linear independence of the basis functions gives two equations:

$$-i\gamma^0 u_{\vec{k}} \dot{f}_{\vec{k}} + (\vec{k} \cdot \vec{\gamma} + m) u_{\vec{k}} f_{\vec{k}} = 0 = -i\gamma^0 v_{\vec{k}} \dot{g}_{\vec{k}} + (-\vec{k} \cdot \vec{\gamma} + m) v_{\vec{k}} g_{\vec{k}} .$$

Differentiating w.r.t. time once more, multiplying by $-i\gamma^0$ on the left and using the above equations again leads to,

$$- \sum_{\sigma} u_{\vec{k},\sigma} \left\{ \ddot{f}_{\vec{k},\sigma} + \omega_{\vec{k}}^2 f_{\vec{k},\sigma} \right\} = 0 = - \sum_{\sigma} v_{\vec{k},\sigma} \left\{ \ddot{g}_{\vec{k},\sigma} + \omega_{\vec{k}}^2 g_{\vec{k},\sigma} \right\} , \quad \omega_{\vec{k}}^2 := \vec{k}^2 + m^2 .$$

Since $u_{\vec{k},\sigma}, v_{\vec{k},\sigma}$ are presumed linearly independent, each of the f and the g functions satisfy the same, familiar harmonic oscillator equation, with solutions $e^{\pm i\omega_{\vec{k}} t}$. Of course one of the solutions for each of f, g is spurious since the original equations are first order. Substituting $f_{\vec{k},\sigma} \sim e^{-i\omega_{\vec{k}} t}$ and $g_{\vec{k},\sigma} \sim e^{+i\omega_{\vec{k}} t}$ in the first order equations *requires* the u, v spinors to satisfy: $\boxed{(\vec{k} + m)u_{\vec{k},\sigma} = 0 = (\vec{k} - m)v_{\vec{k},\sigma}}$. Choice of the other signs does not give a Lorentz covariant equation for the spinors. Hence we restrict to: $\boxed{\dot{f}_{\vec{k},\sigma} = -i\omega_{\vec{k}} f_{\vec{k},\sigma} , \dot{g}_{\vec{k},\sigma} = +i\omega_{\vec{k}} g_{\vec{k},\sigma}}$. These equations satisfied by the spinors are analogous to the transversality conditions we got on the polarization tensors. These are consequences of the Dirac equation i.e. hold for *solutions*.

To compute the Lagrangian, we need the decomposition of the Dirac conjugate,

$$\bar{\psi}(t, \vec{x}) = \int_{\mathbb{R}^3/2} d^3k \left[\left\{ \sum_{\sigma} \bar{u}(\vec{k}, \sigma) f_{\vec{k},\sigma}^*(t) \right\} \varphi_{\vec{k}}^*(\vec{x}) + \left\{ \sum_{\sigma} \bar{v}(\vec{k}, \sigma) g_{\vec{k},\sigma}^*(t) \right\} \varphi_{\vec{k}}(\vec{x}) \right]$$

We will now choose the u, v spinors to be solutions of the equations: $(\not{k} + m)u(\vec{k}, \sigma) = 0 = (\not{k} - m)v(\vec{k}, \sigma)$ and express the Dirac action in terms of the f, g functions alone.

$$\begin{aligned}
-i\gamma^\mu \partial_\mu \psi &= \int_{\mathbb{R}^3/2} d^3k \left[\left\{ -i\gamma^0 u(\vec{k}, \sigma) \dot{f}_{\vec{k}, \sigma} \varphi_{\vec{k}}(\vec{x}) + -i\gamma^0 v(\vec{k}, \sigma) \dot{g}_{\vec{k}, \sigma} \varphi_{\vec{k}}^*(\vec{x}) \right\} \right. \\
&\quad \left. + \left\{ k_i \gamma^i u(\vec{k}, \sigma) f_{\vec{k}, \sigma} \varphi_{\vec{k}}(\vec{x}) - k_i \gamma^i v(\vec{k}, \sigma) g_{\vec{k}, \sigma} \varphi_{\vec{k}}^*(\vec{x}) \right\} \right] \\
\therefore (-i\gamma^\mu \partial_\mu + m)\psi &= \int d^3k \left[\left\{ -i\gamma^0 u(\vec{k}, \sigma) \dot{f}_{\vec{k}, \sigma} + k_i \gamma^i u(\vec{k}, \sigma) f_{\vec{k}, \sigma} \right. \right. \\
&\quad \left. \left. + m u(\vec{k}, \sigma) f_{\vec{k}, \sigma} \right\} \varphi_{\vec{k}}(\vec{x}) \right. \\
&\quad \left. + \left\{ -i\gamma^0 v(\vec{k}, \sigma) \dot{g}_{\vec{k}, \sigma} - k_i \gamma^i v(\vec{k}, \sigma) g_{\vec{k}, \sigma} \right. \right. \\
&\quad \left. \left. + m v(\vec{k}, \sigma) g_{\vec{k}, \sigma} \right\} \varphi_{\vec{k}}^*(\vec{x}) \right] \\
\therefore L &= \int_{\Sigma_t} d^3x \bar{\psi} (-i\gamma^\mu \partial_\mu + m)\psi = \int_{\mathbb{R}^3/2} d^3k' \int_{\mathbb{R}^3/2} d^3k \int_{\Sigma_t} d^3x \\
&\quad \left[\left\{ \bar{u}(\vec{k}', \sigma) f_{\vec{k}', \sigma'}^*(t) \right\} \varphi_{\vec{k}'}^*(\vec{x}) + \left\{ \bar{v}(\vec{k}', \sigma') g_{\vec{k}', \sigma'}^*(t) \right\} \varphi_{\vec{k}'}(\vec{x}) \right] \times \\
&\quad \left[\left\{ -i\gamma^0 u(\vec{k}, \sigma) \dot{f}_{\vec{k}, \sigma} + k_i \gamma^i u(\vec{k}, \sigma) f_{\vec{k}, \sigma} + m u(\vec{k}, \sigma) f_{\vec{k}, \sigma} \right\} \varphi_{\vec{k}}(\vec{x}) \right. \\
&\quad \left. + \left\{ -i\gamma^0 v(\vec{k}, \sigma) \dot{g}_{\vec{k}, \sigma} - k_i \gamma^i v(\vec{k}, \sigma) g_{\vec{k}, \sigma} + m v(\vec{k}, \sigma) g_{\vec{k}, \sigma} \right\} \varphi_{\vec{k}}^*(\vec{x}) \right]
\end{aligned}$$

Thanks to the integration domain of the \vec{k}, \vec{k}' integrations, only $\varphi_{\vec{k}'}^* \varphi_{\vec{k}}$ terms contribute $\delta^3(\vec{k} - \vec{k}')$. Hence only \bar{u} pairs with u and \bar{v} pairs with v . Using the equations satisfied by the spinors, we eliminate $(k_i \gamma^i + m)u = +\omega_{\vec{k}} \gamma^0 u$ and $(-k_i \gamma^i + m)v = -\omega_{\vec{k}} \gamma^0 v$. This leads to,

$$\begin{aligned}
L_{Dirac} &= \int_{\mathbb{R}^3/2} d^3k \left\{ \sum_{\sigma, \sigma'} \left[\bar{u}(\vec{k}, \sigma') \gamma^0 u(\vec{k}, \sigma) \right] \left[f_{\vec{k}, \sigma'}^* (-i \dot{f}_{\vec{k}, \sigma} + \omega_{\vec{k}} f_{\vec{k}, \sigma}) \right] \right. \\
&\quad \left. + \sum_{\sigma, \sigma'} \left[\bar{v}(\vec{k}, \sigma') \gamma^0 v(\vec{k}, \sigma) \right] \left[g_{\vec{k}, \sigma'}^* (-i \dot{g}_{\vec{k}, \sigma} - \omega_{\vec{k}} g_{\vec{k}, \sigma}) \right] \right\} \quad (7.13)
\end{aligned}$$

Choosing the normalization of the spinors so that $\bar{u}(\vec{k}, \sigma') \gamma^0 u(\vec{k}, \sigma) = \delta_{\sigma, \sigma'} = \pm \bar{v}(\vec{k}, \sigma') \gamma^0 v(\vec{k}, \sigma)$ gives the Dirac action as,

$$S[\bar{\psi}, \psi] = \int dt \int_{\mathbb{R}^3/2} d^3k (-i) \sum_{\sigma} \left[f_{\vec{k}, \sigma}^* (\dot{f}_{\vec{k}, \sigma} + i\omega_{\vec{k}} f_{\vec{k}, \sigma}) \pm g_{\vec{k}, \sigma}^* (\dot{g}_{\vec{k}, \sigma} - i\omega_{\vec{k}} g_{\vec{k}, \sigma}) \right] \quad (7.14)$$

We can group the f, g 's together to get a uniform Lagrangian with $\vec{k} \in \mathbb{R}^3$. Denote $f_{-\vec{k}, \sigma} := g_{\vec{k}, \sigma}^* \forall \vec{k} \in \mathbb{R}^3/2$. Do a partial integration of the $g_{\vec{k}, \sigma}^* \dot{g}_{\vec{k}, \sigma} = f_{-\vec{k}, \sigma} \dot{f}_{-\vec{k}, \sigma}^*$. This makes the relative signs the same in both the terms. Choosing the '-' sign for the normalization condition for

the v -spinors, allows us to combine the two terms and extend the integration domain to full \mathbb{R}^3 . Thus, we get,

$$S_{Dirac}[\bar{\psi}, \psi] = \int dt \int_{\mathbb{R}^3} d^3k \sum_{\sigma} \left[f_{\vec{k},\sigma}^* (-i\dot{f}_{\vec{k},\sigma} + \omega_{\vec{k}} f_{\vec{k},\sigma}) \right] \quad \text{for} \quad (7.15)$$

$$\psi(t, \vec{x}) = \int_{\mathbb{R}^3/2} d^3k \sum_{\sigma} \left[u(\vec{k}, \sigma) f_{\vec{k},\sigma}(t) \varphi_{\vec{k}}(\vec{x}) + v(\vec{k}, \sigma) f_{-\vec{k},\sigma}^*(t) \varphi_{\vec{k}}^*(\vec{x}) \right] \quad (7.16)$$

$$0 = (\not{k} + m)u(\vec{k}, \sigma) = (-\not{k} + m)v(\vec{k}, \sigma) \quad \text{normalized as,} \quad (7.17)$$

$$\delta_{\sigma',\sigma} = \bar{u}(\vec{k}, \sigma') \gamma^0 u(\vec{k}, \sigma) = -\bar{v}(\vec{k}, \sigma') \gamma^0 v(\vec{k}, \sigma). \quad (7.18)$$

Note that the action is real again.

An aside: It is an interesting exercise to express the action in terms of the real q type variables as we did for the scalar. Since the action is manifestly a sum over variables with uncoupled labels σ, \vec{k} , we focus on just one of the terms. Thus consider the manifestly real Lagrangian, $L(f, f^*) := \frac{1}{2} \{-i f^* \dot{f} + i \dot{f}^* f\} + \omega f^* f$. Introduce the real variables x, y as: $f := x + iy, \dot{f} = \dot{x} + i\dot{y}$. The Lagrangian then takes the form,

$$L(x, y, \dot{x}, \dot{y}) = x\dot{y} - y\dot{x} + \omega(x^2 + y^2) \Rightarrow p_x = -y, p_y = +x$$

Notice that the equations defining the conjugate momenta *cannot* be inverted to solve for the velocities - *we have a constrained system* [3]. This is not the place to discuss it in detail, the reference gives the necessary background. I will just list the steps.

$$\phi := p_x + y, \quad \chi := p_y - x \quad (\text{primary constraints});$$

$$H_{canonical} := p_x \dot{x} + p_y \dot{y} - L = -\omega(x^2 + y^2);$$

$$H_{total} = -\omega(x^2 + y^2) + \lambda\phi + \mu\chi \quad \text{Preservation of primary constraints} \Rightarrow$$

$$H_{total} = -\omega(xp_y - yp_x) \quad \because \lambda = \omega y, \mu = -\omega x;$$

$$\text{Dirac brackets} : \{x, p_x\}_* = \{y, p_y\}_* = \frac{1}{2}, \{x, y\}_* = \{p_x, p_x\}_* = -\frac{1}{2}. \quad \text{Use:}$$

$$H_{total} = -\omega(x^2 + y^2) \quad \text{and Dirac brackets for equations of motion:}$$

$$\dot{x} = \omega y, \quad \dot{y} = -\omega x; \quad \text{finally drop } p_x, p_y \text{ and set } x := \frac{p}{\sqrt{2}}, y := \frac{q}{\sqrt{2}},$$

$$H_{total} = -\frac{1}{2}\omega(p^2 + q^2), \quad \{q, p\} = 1, \quad \dot{p} = \omega q, \quad \dot{q} = -\omega p.$$

Re-introducing the (\vec{k}, σ) labels shows that the spinor field Hamiltonian too is a sum of harmonic oscillators for each of the $\vec{k} \in \mathbb{R}^3$ and $\sigma = \pm$ labels.

Puzzle: Why is the Hamiltonian of a wrong sign? Is it merely a choice of the overall sign in the Lagrangian density (will not change the equation of motion)? What is the rationale for a choice of sign? We will not pursue this classical formulation further here.

Remarks: The scalar and the vector fields that we discussed have been real. This was reflected in the Fourier decomposition with complex conjugate coefficients. What if we have a complex field? Well, we can always write the Fourier decomposition as (see the spinorial case),

$$\phi(t, \vec{x}) = \int_{\mathbb{R}^3/2} d^3k [f_{\vec{k}}(t)\varphi_{\vec{k}}(\vec{x}) + g_{\vec{k}}^*(t)\varphi_{\vec{k}}(\vec{x})]$$

For real scalar field, the reality condition simply identifies $g_{\vec{k}}(t) = f_{-\vec{k}}(t)$. For a complex field, there is no such identification. We will then have *two* harmonic oscillators for each $\vec{k} \in \mathbb{R}^3$. For the spinorial case, we had a constrained system and hence half the number of oscillators.

Remark: Consider the time-Fourier transform of a scalar field: $\phi(\omega, \vec{x}) := \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dt e^{-i\omega t} \phi(t, \vec{x})$. The Fourier transform of its complex conjugate is given by: $\phi_c(\omega) := \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dt e^{+i\omega t} \phi^*(t, \vec{x})$. Clearly, $\phi_c(\omega) = \phi^*(-\omega)$. Therefore, a positive frequency $\phi(t, \vec{x})$ has $\phi(\omega, \vec{x}) = 0$, for $\omega < 0$. This immediately also give that $\phi^*(t, \vec{x})$ is a negative frequency field, i.e. $\phi_c(\omega, \vec{x}) = 0$ for $\omega > 0$. This gets reflected in the *mode decomposition*.

While the Fourier decomposition displays the “degrees of freedom” of the field, the mode decomposition will turn out to be convenient for passage to quantum fields. To appreciate this, consider a single oscillator (a single \vec{k}). The degrees of freedom view gives: $\ddot{q} + \omega^2 q^2 = 0, p = \dot{q}, H = \frac{1}{2}(p^2 + q^2), \{q, p\} = 1$. The mode decomposition gives (a, a^* are constants),

$$\begin{aligned} q(t) &= \underline{a}e^{-i\omega t} + \underline{a}^*e^{i\omega t} &\Rightarrow & p(t) = -i\omega (\underline{a}e^{-i\omega t} - \underline{a}^*e^{i\omega t}) \text{ inverting gives,} \\ \underline{a} &= \frac{e^{i\omega t}}{2} \left(q + i\frac{p}{\omega} \right) &, & \underline{a}^* = \frac{e^{-i\omega t}}{2} \left(q - i\frac{p}{\omega} \right) \\ \therefore \{\underline{a}, \underline{a}^*\} &= -\frac{i}{2\omega} \text{ Define: } a := \sqrt{2\omega}\underline{a}, a^* := \sqrt{2\omega}\underline{a}^* &\Rightarrow & \\ \{a, a^*\} &= -i &, & q(t) = \left(\frac{e^{-i\omega t}}{\sqrt{2\omega}} \right) a + \left(\frac{e^{i\omega t}}{\sqrt{2\omega}} \right) a^* \end{aligned}$$

Notice the normalization factor of $\sqrt{2\omega}$. This is correlated with the normalization of the Poisson bracket of a, a^* (and of course has nothing to do with the Lorentz covariance!). As per the usual canonical quantization procedure, $\{, \} \rightarrow -i/(\hbar = 1) \times [,]$. This would lead to $[a, a^\dagger] = 1$.

Keeping this in mind, we summarize the mode decompositions for the fields:

$$\phi(x) = \int_{\mathbb{R}^3} \frac{d^3k}{\sqrt{2\omega(2\pi)^3}} [a_{\vec{k}} e^{ik \cdot x} + b_{\vec{k}}^* e^{-ik \cdot x}] \quad k \cdot x := -\omega_{\vec{k}} t + \vec{k} \cdot \vec{x} \quad (7.19)$$

If $\phi^* = \phi$ then $b_{\vec{k}} = a_{\vec{k}}$.

$$A_\mu(x) = \int_{\mathbb{R}^3} \frac{d^3k}{\sqrt{2\omega(2\pi)^3}} \sum_{\lambda=1,2} [a_{\vec{k},\lambda} \varepsilon_\mu(\vec{k}, \lambda) e^{ik \cdot x} + a_{\vec{k},\lambda}^* \varepsilon_\mu^*(\vec{k}, \lambda) e^{-ik \cdot x}] \quad (7.20)$$

$$\varepsilon_0(\vec{k}, \lambda) := -\frac{\vec{k} \cdot \vec{\varepsilon}(\vec{k}, \lambda)}{\omega_{\vec{k}}} \quad , \quad \omega_{\vec{k}} = |\vec{k}|$$

$$\varepsilon_i(\vec{k}, \lambda) := \left(\delta_i^j - \frac{k_i k^j}{k^2} \right) \varepsilon_j(\vec{k}, \lambda) \quad , \quad \vec{\varepsilon}(\vec{k}, \lambda) \cdot \vec{\varepsilon}^*(\vec{k}, \lambda') = \delta_{\lambda,\lambda'}$$

$$\psi(x) = \int_{\mathbb{R}^3} \frac{d^3k}{\sqrt{2\omega(2\pi)^3}} \sum_{\sigma=\pm} [b_{\vec{k},\sigma} u(\vec{k}, \sigma) e^{ik \cdot x} + d_{\vec{k},\sigma}^* v(\vec{k}, \sigma) e^{-ik \cdot x}] \quad (7.21)$$

$$(\not{k} + m)u(\vec{k}, \sigma) = 0 = (-\not{k} + m)v(\vec{k}, \sigma)$$

$$\bar{u}(\vec{k}, \sigma)u(\vec{k}, \sigma') = \delta_{\sigma,\sigma'} = -\bar{v}(\vec{k}, \sigma)v(\vec{k}, \sigma')$$

$$\sum_{\sigma} u(\vec{k}, \sigma)\bar{u}(\vec{k}, \sigma) = \frac{-\not{k} + m}{2m} \quad , \quad \sum_{\sigma} v(\vec{k}, \sigma)\bar{v}(\vec{k}, \sigma) = -\frac{\not{k} + m}{2m}$$

For a real scalar field, we note,

$$\{a_{\vec{k}}, a_{\vec{k}'}^*\} = -i\delta^3(\vec{k} - \vec{k}') \quad \leftrightarrow \quad \{\phi(t, \vec{x}), \pi(t, \vec{x}')\} = \delta^3(\vec{x} - \vec{x}') \quad (7.22)$$

NOTE: So far whatever we have done has no quantum in it. We have obtained a classical theory of non-interacting fields.

C. Interaction with source: Green's functions

A simplest kind of ‘interaction’ we are familiar with, eg. from classical electrodynamics, is interaction with a “source”. So let us consider a real scalar field interacting with a source $J(x)$ which is a real function. The interaction is described by the equation⁴:

$$\begin{aligned} (\square - m^2)\phi(x) &= -J(x) \quad \leftrightarrow \quad (\square_x - m^2)G(x, x') = -\delta^4(x - x') \quad \text{Green's Function} \\ \Rightarrow \phi(x) &= + \int_{\text{Source}} d^4x' G(x, x') J(x') \end{aligned}$$

⁴ Follows from the Lagrangian $\mathcal{L} = \mathcal{L}_{\text{scalar}} + J(x)\phi(x)$ with $\mathcal{L}_{\text{scalar}}$ given in (6.5).

Poincare invariance of the defining equation for Green's function implies that it is a Lorentz invariant function of $(x - x')$. It can be represented as,

$$G(x - x') = \int_{\mathbb{R}^4} \frac{d^4k}{(2\pi)^4} \frac{e^{ik \cdot (x - x')}}{k^2 + m^2}, \quad k^2 = -(k^0)^2 + \vec{k}^2. \quad (7.23)$$

The denominator vanishes for $k^2 = -m^2$ and thus we need a *definition* of the integral and hence of the Green's function.

The usual method is to define $\int d^4k = \int d^3k \int dk^0$ and interpret the k^0 integral as a contour integral along a suitably chosen contour. Each choice is supposed to reflect a "boundary condition" for the wave operator. In the complex k^0 plane, there are poles at $k^0 = \pm \omega_{\vec{k}} := \sqrt{\vec{k}^2 + m^2}$. The integral is sought to be defined by shifting the poles off the real axis. This can be done in 4 ways: (i) shift both poles in the Lower Half Plane (LHP), $k^0 \rightarrow k^0 + i\epsilon$, (ii) shift both poles in the UHP, $k^0 \rightarrow k^0 - i\epsilon$, (iii) shift the pole at $+\omega_{\vec{k}}$ in the LHP and shift the pole at $-\omega_{\vec{k}}$ in the UHP, $\omega_{\vec{k}}^2 \rightarrow \omega_{\vec{k}}^2 - i\epsilon$ and (iv) reverse of the (iii). To see the consequences of any of these choices, let us consider the special case of a source localized at the space-time origin: $J(t, x) = \delta^4(x)$. The solution takes the form,

$$\begin{aligned} \phi(x) &= \int dt' \int d^3x' \int \frac{d^4k}{(2\pi)^4} \frac{e^{ik \cdot x}}{k^2 + m^2} e^{-i(-k^0 t' + \vec{k} \cdot \vec{x}')} \delta(t') \delta^3(\vec{x}') \\ &= \int \frac{d^3k}{(2\pi)^3} e^{i\vec{k} \cdot \vec{x}} \left[\int \frac{dk^0}{2\pi} \frac{e^{-ik^0 t}}{(k^0)^2 - \omega_{\vec{k}}^2} \right] \quad \text{and,} \\ [\dots] &= \frac{1}{2\omega_{\vec{k}}} \int \frac{dk^0}{2\pi} \left(\frac{1}{k_0 - \omega_{\vec{k}}} - \frac{1}{k_0 + \omega_{\vec{k}}} \right) e^{-ik^0 t} \end{aligned} \quad (7.24)$$

Without the $i\epsilon$'s, both sides of the above equations are real. Now we introduce the $i\epsilon$'s. The contour integrals will have the segment of a semicircle at infinity whose contribution should vanish. On the semicircle at large R , $k^0 = Re^{i\theta}$, the integrand behaves as $R^{-2} e^{R \sin(\theta)t}$ and the measure provides factor of R . This suffices for convergence for $t = 0$. However, for a non-zero t , we can get an exponential divergence unless θ is restricted appropriately. This dictates how the contour should be closed in UHP or LHP. Since $\phi(x)$ is a function of the Lorentz invariant $x^2 = -t^2 + \vec{x}^2$, we can consider the two cases as: (a) $x^2 \leq 0$ (sign of t is invariant) and we may evaluate the integrals for $\vec{x} = 0$; (b) $x^2 > 0$ and we can evaluate the integrals for $t = 0$.

(i) *Both poles to LHP*: To pickup the residues, we should close the contour in the LHP. For space-like interval, taking $t = 0$, we see that the residues at the two poles cancel each

other and the solution vanishes. For time-like intervals, $t \neq 0$ but we can take $\vec{x} = 0$. To have a vanishing contribution from the semicircle at infinity, we have to restrict $t > 0$. The evaluation of the integral is trivial and gives,

$$[\dots]_{Retarded} = -\frac{1}{\omega_{\vec{k}}} \theta(t) \sin(\omega_{\vec{k}} t) .$$

(ii) *Both poles to UHP*: To pick-up the residues, the contour should be closed in the UHP and for time-like interval, we need to restrict to $t < 0$. The evaluation gives,

$$[\dots]_{Advanced} = \frac{1}{\omega_{\vec{k}}} \theta(-t) \sin(\omega_{\vec{k}} t) .$$

As before, for space-like interval, the solution vanishes.

(iii) *Positive pole to LHP and negative to UHP*: Now closing the contour in either of UHP/LHP will always pick up a contribution. For time-like interval, for LHP closure, we have to take $t > 0$ and we will pick up the contribution from the first term. Likewise, for the UHP closure, we have to take $t < 0$ and we will pick up the contribution from the second term. Evaluation gives, for time-like or light-like separation

$$[\dots]_{Feynman} = -i \frac{1}{2\omega_{\vec{k}}} [\theta(t)e^{-i\omega_{\vec{k}}t} + \theta(-t)e^{+i\omega_{\vec{k}}t}] = -i \frac{1}{2\omega_{\vec{k}}} e^{-i\omega_{\vec{k}}|t|} [\theta(t) + \theta(-t)] .$$

For space-like interval, taking $t = 0$, we get the *non-zero* answer, $[\dots]_{Feynman} = -i \frac{1}{2\omega_{\vec{k}}}$ for closure in either LHP or UHP.

The case (iv) is similar to (iii) and is obtained in an obvious manner. The solution $\phi(x)$ is obtained by the spatial Fourier transform as given above (7.24).

Remarks: There are two properties of the Retarded and the Advanced choices that stand out. In both cases, the solution $\phi(x)$ (a) is real for a real source function and (b) it reflects the expected *causal* behavior - vanishing outside the future/past light cones. By contrast, the Feynman choice gives a non-real $\phi(x)$ even for a real source function and the solution is non-vanishing outside the light cones! The same holds for the choice (iv). *What do we make of this?*

The retarded and advanced choices support the interpretation that the field can be regarded as an *observable* responding to a source in a causally consistent manner. This is what we would have in a classical field theory. The Feynman choice however disallows such an interpretation - *the particular solution $\phi(x)$ cannot be interpreted as causally consistent response to a source.*

Both the causal and the Feynman Green's functions appear naturally when $\phi(x)$ is promoted to be an operator field i.e. in *Quantum Field Theory*. In the scattering theory, section 10, the causal Green's function are used in articulating the *asymptotic conditions* while the Feynman Green's function (or Feynman propagator) appears naturally in the Lehmann-Symanzik-Zimmermann (LSZ) reduction of the scattering matrix.

For completeness, we note the full Green's functions⁵:

$$G_{ret}(x) = +\frac{\theta(t)}{2\pi}\delta(x^2) - \theta(t)\theta(-x^2)\frac{mJ_1(m\sqrt{-x^2})}{4\pi\sqrt{-x^2}} \quad (7.25)$$

$$G_{adv}(x) = +\frac{\theta(-t)}{2\pi}\delta(x^2) - \theta(-t)\theta(-x^2)\frac{mJ_1(m\sqrt{-x^2})}{4\pi\sqrt{-x^2}} \quad (7.26)$$

$$G_{Feynman}(x^2 < 0) = +\frac{\delta(x^2)}{4\pi} - \frac{m}{8\pi\sqrt{-x^2}}H_1^{(2)}(m\sqrt{-x^2})$$

$$G_{Feynman}(x^2 > 0) = +\frac{i}{4\pi^2}\frac{m}{\sqrt{x^2}}K_1(m\sqrt{x^2}) \quad (7.27)$$

⁵ The overall signs are convention dependent. It is easiest to match it for the massless case. For detail see: [\[4, 5\]](#)

8. COVARIANCE OF QUANTUM FIELDS AND RELATIVISTIC CAUSALITY

So far we have studied representations of the Poincare group and seen emergence of a “field dynamical system” - an infinite collection of harmonic oscillators. In our attempt to incorporate interactions with classical sources, we encountered the Green’s functions. We met the expected and familiar retarded and advanced Green’s functions, but also encountered the mathematical possibility of the Feynman alternative which does not gel with a classical field theory interpretation. Taking it as a hint of an opportunity, we attempt a ‘quantum’ interpretation. Here, the field as a collection of harmonic oscillators gives a strong clue: *use the mode decomposition and promote the expansion coefficients to operators*. We will then have a collection of creation and annihilation operators and can tag the states of the system by the number of quanta. We already saw (7.22) in the simpler case of a real scalar field that postulating Poisson brackets for the coefficients and using the completeness of mode functions, we can deduce the Poisson brackets for the fields themselves. The same method works for operators as well. That is, not only do we promote the fields to operators but we also postulate the necessary commutation relations.

A. Poincare Covariance of Quantum fields

The first feature, fields as operators, immediately modifies the implementation of covariance: a covariant quantum field requires specific transformations for the coefficient functions and conversely. Transformation laws of the mode functions do not suffice.

To see this, consider a linear combination of classical fields, $\psi^A(x) := \sum_n c_n \psi_n^A(x)$. A group action on ψ^A is deduced from that on the individual ψ_n^A : $g \cdot \psi_n^A(x) = D_B^A(g) \psi_n^B(g^{-1}x)$,

$$g \cdot \psi^A(x) := \sum_n c_n g \cdot \psi_n^A(x) = \sum_n D_B^A(g) \psi_n^B(g^{-1}x) = D_B^A(g) \psi^B(g^{-1}x).$$

This of course presumes quite naturally that the coefficients are *invariant* under the group action. When a similar linear combination is promoted to an operator by making the c_n as operators, the above procedure breaks down.

$$U(g) \hat{\psi}^A(x) U^\dagger(g) := \sum_n [U(g) \hat{c}_n U(g)^\dagger] \psi_n^A(x) \stackrel{?}{=} D_B^A(g^{-1}) \hat{\psi}^B(gx).$$

Evidently, if we assume the \hat{c}_n to be invariant, then the field cannot be covariant. So the \hat{c}_n should transform in a manner that shifts the group action on to the $\psi_n^A(x)$ and then

the \sum_n to work out correctly so as to produce the expected right hand side⁶. Taking now $c_n \rightarrow \hat{a}, \hat{a}^\dagger, \sum_n \rightarrow \int \frac{d^3k}{\dots}$ and using the mode expansion form, we observe that *if* we postulate the transformation laws for the \hat{a}, \hat{a}^\dagger as,

$$U(g)\hat{a}(\vec{k}, \sigma)U^\dagger(g) := \hat{a}(\Lambda\vec{k}, "D(\Lambda)"\sigma) \quad , \quad U(g)\hat{a}^\dagger(\vec{k}, \sigma)U^\dagger(g) := \hat{a}^\dagger(\Lambda\vec{k}, "D(\Lambda)"\sigma) \quad ,$$

then we can shift the changed labels from the operators to the mode function labels using the change of integration (and summation) variables. Provided the integration measure is invariant, the right hand side takes the expected form and we have the *covariance of the quantum field*. Note that this essentially fixes the form of the mode decomposition! In equations (for a Dirac field say),

$$\begin{aligned} [U(g)\hat{\psi}U^\dagger(g)](x) &:= \int \frac{d^3k}{\sqrt{2\omega_{\vec{k}}(2\pi)^3}} \sum_{\sigma} \left[(U\hat{b}_{\vec{k},\sigma}U^\dagger)u(\vec{k}, \sigma)e^{ik \cdot x} + (U\hat{d}_{\vec{k},\sigma}^\dagger U^\dagger)v(\vec{k}, \sigma)e^{-ik \cdot x} \right] \\ &= \int \frac{d^3k}{\sqrt{2\omega_{\vec{k}}(2\pi)^3}} \sum_{\sigma} \left[\hat{b}_{\Lambda\vec{k},\sigma_\Lambda} u(\vec{k}, \sigma)e^{ik \cdot x} + \hat{d}_{\Lambda\vec{k},\sigma_\Lambda}^\dagger v(\vec{k}, \sigma)e^{-ik \cdot x} \right] \\ &= \int \frac{d^3(\Lambda^{-1}k)}{\sqrt{2\omega_{(\Lambda^{-1}k)}(2\pi)^3}} \sum_{\sigma_{\Lambda^{-1}}} \left[\hat{b}_{\vec{k},\sigma} u((\Lambda^{-1}k), \sigma_{\Lambda^{-1}})e^{i(\Lambda^{-1}k) \cdot x} \right. \\ &\quad \left. + \hat{d}_{\vec{k},\sigma}^\dagger v((\Lambda^{-1}k), \sigma_{\Lambda^{-1}})e^{-i(\Lambda^{-1}k) \cdot x} \right] \\ &= \int \frac{d^3k}{\sqrt{2\omega_{\vec{k}}(2\pi)^3}} \sum_{\sigma} \left[\hat{b}_{\vec{k},\sigma} \left\{ D(\Lambda^{-1})u(\vec{k}, \sigma) \right\} e^{ik \cdot (\Lambda x)} \right. \\ &\quad \left. + \hat{d}_{\vec{k},\sigma}^\dagger \left\{ D(\Lambda^{-1})v(\vec{k}, \sigma) \right\} e^{-ik \cdot (\Lambda x)} \right] \\ &= D(\Lambda^{-1})\hat{\psi}(\Lambda x) \end{aligned}$$

Note: If we postulate a Poincare invariant state $|0\rangle$ and define $|\vec{k}, \sigma\rangle := b^\dagger(\vec{k}, \sigma)|0\rangle$, then these states transform precisely as per the particle representation. Thus the postulated group action on the b, d operators are precisely as needed for building up the particle representations and we will use it shortly.

B. Space Inversion, Time Reversal, Charge Conjugation of Dirac Field

Let us quickly see how the discrete symmetries act on quantum fields. Again, we will consider a Dirac field for illustration and now suppress the ‘hats’ on the operators.

⁶ The homomorphism dictates the expected form on the r.h.s.

Space Inversion:

We want $\Psi_{\mathcal{P}}(t, \vec{x}) := \mathcal{P}\Psi\mathcal{P}^\dagger = D(\mathcal{P})\Psi(t, -\vec{x})$. From the defining relations for \mathcal{P} , namely, $\mathcal{P}P_i\mathcal{P}^\dagger = -P_i$ and $\mathcal{P}J_i\mathcal{P}^\dagger = J_i$, we postulate,

$$\mathcal{P}b^\dagger(\vec{k}, \sigma)\mathcal{P}^\dagger := \eta b^\dagger(-\vec{k}, \sigma) \quad , \quad \mathcal{P}d^\dagger(\vec{k}, \sigma)\mathcal{P}^\dagger := \eta d^\dagger(-\vec{k}, \sigma) \quad , \quad \eta^2 = \pm 1 .$$

As noted while discussing the discrete symmetry actions on the particle representations, the phase is σ independent.

Substituting the mode expansion and using these postulated actions gives,

$$\mathcal{P}\Psi^\dagger(\vec{k}, \sigma)\mathcal{P}^\dagger = \int [d^3k]_{inv} \sum_{\sigma} \left\{ \eta^* b(\vec{k}, \sigma) u_{-\vec{k}, \sigma} e^{ik \cdot (\mathcal{P}x)} + \eta d^\dagger(\vec{k}, \sigma) v_{-\vec{k}, \sigma} e^{-ik \cdot (\mathcal{P}x)} \right\}$$

Claim: The u, v spinors satisfy, $u(-\vec{k}, \sigma) = \gamma^0 u(\vec{k}, \sigma)$ and $v(-\vec{k}, \sigma) = -\gamma^0 v(\vec{k}, \sigma)$.

This follows by noting that

$$\begin{aligned} 0 &= (\not{k} + m)u(\vec{k}, \sigma) = (-\omega\gamma^0 + \vec{k} \cdot \vec{\gamma} + m)u(\vec{k}, \sigma) \quad , \quad u(\vec{k}, \sigma) = \gamma^2 u(\vec{k}, \sigma) \\ \therefore 0 &= (-\omega\gamma^0\gamma^0 + \vec{k} \cdot \vec{\gamma}\gamma^0 + m\gamma^0)(\gamma^0 u(\vec{k}, \sigma)) \\ &= (-\omega\gamma^0 + (-\vec{k} \cdot \vec{\gamma} + m)(\gamma^0 u(\vec{k}, \sigma)) \quad \Rightarrow \quad \gamma^0 u(\vec{k}, \sigma) \propto u(-\vec{k}, \sigma) \end{aligned}$$

Similarly, $\gamma^0 v(\vec{k}, \sigma) \propto v(-\vec{k}, \sigma)$. The proportionality factors are fixed by noting that for $\vec{k} = \hat{k}$, the u, v spinors are eigen-spinors of γ^0 with eigenvalues ± 1 . This implies,

$$\mathcal{P}b^\dagger(\vec{k}, \sigma)\mathcal{P}^\dagger = \gamma^0 \int [d^3k]_{inv} \sum_{\sigma} \left\{ \eta^* b(\vec{k}, \sigma) u_{\vec{k}, \sigma} e^{ik \cdot (\mathcal{P}x)} - \eta d^\dagger(\vec{k}, \sigma) v_{\vec{k}, \sigma} e^{-ik \cdot (\mathcal{P}x)} \right\}$$

Clearly η should be imaginary for the right hand side to have the field $\psi(\mathcal{P}x)$. We choose $\eta = -i$ and deduce that, $\boxed{\mathcal{P}\Psi(x)\mathcal{P}^\dagger = i\gamma^0\Psi(\mathcal{P}x)}$, *i.e.*, $D(\mathcal{P}) = i\gamma^0$ and $\eta^2 = -1$.

Time reversal:

We want $\Psi_{\mathcal{T}}(t, \vec{x}) := \mathcal{T}\Psi\mathcal{T}^\dagger = D(\mathcal{T})\Psi(-t, \vec{x})$. The defining relations where both P_i and J_i change sign, we postulate,

$$\mathcal{T}b^\dagger(\vec{k}, \sigma)\mathcal{T}^\dagger = \xi_\sigma b^\dagger(-\vec{k}, -\sigma) \quad , \quad \mathcal{T}d^\dagger(\vec{k}, \sigma)\mathcal{T}^\dagger = \xi_\sigma d^\dagger(-\vec{k}, -\sigma) .$$

As noted while discussing the discrete symmetry actions on the particle representations, the phase is σ dependent.

Substituting the mode decomposition and noting the \mathcal{T} operator is anti-unitary, we get,

$$\begin{aligned} [\mathcal{T}\Psi\mathcal{T}^\dagger](x) &= \int [d^3k]_{inv} \sum_{\sigma} \left\{ \xi_{\sigma}^* b(-\vec{k}, -\sigma) u^*(\vec{k}, \sigma) e^{-ik \cdot x} + \xi_{\sigma} d^\dagger(-\vec{k}, -\sigma) v^*(\vec{k}, \sigma) e^{ik \cdot x} \right\} \\ &= \int [d^3k]_{inv} \sum_{\sigma} \left\{ \xi_{-\sigma}^* b(\vec{k}, \sigma) u^*(-\vec{k}, -\sigma) e^{-ik \cdot (\mathcal{T}x)} + \xi_{-\sigma} d^\dagger(\vec{k}, \sigma) v^*(-\vec{k}, -\sigma) e^{ik \cdot (\mathcal{T}x)} \right\} \end{aligned}$$

Now we need to use the relations,

$$u^*(-\vec{k}, -\sigma) = -\sigma C \gamma_5 u(\vec{k}, \sigma) \quad , \quad v^*(-\vec{k}, -\sigma) = -\sigma C \gamma_5 v(\vec{k}, \sigma) \quad .$$

Once again these are proved from the defining equations for the spinors.

Clearly, if we choose $\xi_{\sigma} = \sigma (= \pm 1)$, then we get $\Psi(\mathcal{T}x)$ on the right hand side and we get,

$$\boxed{\mathcal{T}\Psi(x)\mathcal{T}^\dagger = C\gamma_5\Psi(\mathcal{T}x), i.e.D(\mathcal{T}) = C\gamma_5 \quad , \quad C\gamma^\mu C^\dagger = -(\gamma^\mu)^T} .$$

Charge conjugation:

Lastly, consider the charge conjugation action. This is defined through the particle-anti-particle exchange and we postulate,

$$Cb_{\vec{k},\sigma}^\dagger C^\dagger = \xi d_{\vec{k},\sigma}^\dagger \quad , \quad Cd_{\vec{k},\sigma}^\dagger C^\dagger = \xi b_{\vec{k},\sigma}^\dagger \quad . \quad (8.1)$$

Substitution of mode expansion gives,

$$[C\Psi C^\dagger](x) = \int [d^3k]_{inv} \sum_{\sigma} \left\{ \xi^* d_{\vec{k},\sigma} u(\vec{k}, \sigma) e^{ik \cdot x} + \xi b_{\vec{k},\sigma}^\dagger v(\vec{k}, \sigma) e^{-ik \cdot x} \right\}$$

Now we need to use: $v(\vec{k}, \sigma) = C^\dagger \bar{u}^T(\vec{k}, \sigma)$ and recall that the C matrix is given by $C = i\gamma^2\gamma^0$, $C^T = -C$, $C^\dagger C = \mathbb{1}$ so that $v = -C\bar{u}^T$ and $u = -C\bar{v}^T$. This gives,

$$\begin{aligned} [C\Psi C^\dagger](x) &= \int [d^3k]_{inv} \sum_{\sigma} \left\{ \xi^* d_{\vec{k},\sigma} (-C\bar{v}^T(\vec{k}, \sigma)) e^{ik \cdot x} + \xi b_{\vec{k},\sigma}^\dagger (-C\bar{u}^T(\vec{k}, \sigma)) e^{-ik \cdot x} \right\} \quad \text{and,} \\ \bar{\Psi}^T(x) &= \int [d^3k]_{inv} \sum_{\sigma} \left\{ b_{\vec{k},\sigma}^\dagger (\bar{u}^T(\vec{k}, \sigma)) e^{-ik \cdot x} + d_{\vec{k},\sigma} (\bar{v}^T(\vec{k}, \sigma)) e^{ik \cdot x} \right\} \end{aligned}$$

Hence, if we choose $\xi^* = \xi = -1$, then we get, $\boxed{C\Psi(x)C^\dagger = +C(\bar{\Psi})^T(x)}$. Note that C operator is unitary!

From these it is easy to get the transformations of Dirac conjugates. Here is a summary:

$$\mathcal{P}\Psi\mathcal{P}^\dagger(x) = i\gamma^0\Psi(\mathcal{P}x) \quad , \quad \mathcal{P}\bar{\Psi}\mathcal{P}^\dagger(x) = -i\bar{\Psi}(\mathcal{P}x)\gamma^0 ; \quad (8.2)$$

$$\mathcal{T}\Psi\mathcal{T}^\dagger(x) = C\gamma_5\Psi(\mathcal{T}x) \quad , \quad \mathcal{T}\bar{\Psi}\mathcal{T}^\dagger(x) = \bar{\Psi}(\mathcal{T}x)\gamma_5C^\dagger ; \quad (8.3)$$

$$C\Psi C^\dagger(x) = C\bar{\Psi}^T(x) \quad , \quad C\bar{\Psi}C^\dagger(x) = \Psi^T(x)C ; \quad (8.4)$$

$$C\gamma^\mu C^\dagger = -(\gamma^\mu)^T \quad , \quad C^\dagger C = \mathbb{1} \quad , \quad C^T = -C \quad , \quad C := i\gamma^2\gamma^0 \quad (8.5)$$

We are now ready to verify one of the important general theorem of relativistic quantum field theory, *the CPT theorem*. We verify it for the spinorial quantum field.

C. CPT theorem for Dirac Field

We want to consider the combined action of the discrete transformations. Let us choose one ordering, say, \mathcal{CPT} . Then,

$$\begin{aligned} (\mathcal{CPT})\Psi(x)(\mathcal{CPT})^\dagger &= \mathcal{C}\mathcal{P}(C\gamma_5\Psi(\mathcal{T}x))\mathcal{P}^\dagger\mathcal{T}^\dagger = C\gamma_5\mathcal{C}(i\gamma^0\Psi(\mathcal{PT}x))C^\dagger \\ &= iC\gamma_5\gamma^0C\bar{\Psi}^T(-x) \quad \because \mathcal{PT}x = -x \\ (\mathcal{CPT})\bar{\Psi}(x)(\mathcal{CPT})^\dagger &= \mathcal{C}\mathcal{P}(\bar{\Psi}(\mathcal{T}x)\gamma_5C^\dagger)\mathcal{P}^\dagger\mathcal{T}^\dagger = \mathcal{C}(\bar{\Psi}(\mathcal{PT}x))(-i\gamma^0)\gamma_5C^\dagger C^\dagger \\ &= i\Psi^T(-x)C\gamma^0\gamma_5C \quad \because C^\dagger = -C \end{aligned}$$

Next, consider the CPT transform of a general bilinear of the form $\bar{\Psi}(x)A\Psi(x)$. Using $(\mathcal{CPT})^\dagger(\mathcal{CPT}) = \mathbf{1}$, the CPT transform will be the CPT transform of the Dirac conjugate, that of the matrix A and that of Ψ . The CPT transform of A is just A^* since only the complex conjugate acts on A .

$$\begin{aligned} &[(\mathcal{CPT})\bar{\Psi}(x)(\mathcal{CPT})^\dagger] [(\mathcal{CPT})A(\mathcal{CPT})^\dagger] [(\mathcal{CPT})\Psi(x)(\mathcal{CPT})^\dagger] \\ &= [i\Psi^T(-x)C\gamma^0\gamma_5C] [A^*] [iC\gamma_5\gamma^0C\bar{\Psi}^T(-x)] \\ &= i^2 [\bar{\Psi}(-x)(C\gamma_5\gamma^0C)^T A^\dagger (C\gamma^0\gamma_5C)^T \Psi(-x)]^T \times (-1) \\ &= +\bar{\Psi}(-x) \{ (C\gamma_5\gamma^0C)^T A^\dagger (C\gamma^0\gamma_5C)^T \} \Psi(-x) \end{aligned} \quad (8.6)$$

The explicit (-1) in the third line is due to the *fermionic* nature of the spinors which we will see shortly in the discussion of the spin-statistics theorem.

The factors of the gamma matrices simplify as,

$$C\gamma_5\gamma^0C = +\gamma_5C\gamma^0C = -\gamma_5C\gamma^0C^\dagger = +\gamma_5(\gamma^0)^T = \gamma_5\gamma^0 \quad \text{and} \quad C\gamma^0\gamma_5C = \gamma^0\gamma_5 .$$

$$(\mathcal{CPT}) [\bar{\Psi}(x)A\Psi(x)] (\mathcal{CPT})^\dagger = \bar{\Psi}(-x) \{(\gamma^0\gamma_5)A^\dagger(\gamma_5\gamma^0)\} \Psi(-x) \quad (8.7)$$

It remains to evaluate the middle braces for $A = \mathbb{1}, \gamma^\mu, \gamma^\mu\gamma^\nu, \gamma^\mu\gamma_5$ and γ_5 . These are obtained easily as,

$$\begin{aligned} \gamma^0\gamma_5\mathbb{1}\gamma_5\gamma^0 &= \mathbb{1} \\ \gamma^0\gamma_5\gamma^\mu\gamma_5\gamma^0 &= -\gamma^\mu \\ \gamma^0\gamma_5\Sigma^{\mu,\nu}\gamma_5\gamma^0 &= +\Sigma^{\mu\nu} \\ \gamma^0\gamma_5\gamma^\mu\gamma_5\gamma_5\gamma^0 &= -\gamma^\mu \\ \gamma^0\gamma_5(i\gamma_5)\gamma_5\gamma^0 &= \gamma_5 \end{aligned}$$

The Lagrangian density is made up of the bi-linears and also involves derivatives. For a $\partial_\mu\Psi(x)$, there is an extra minus sign due to $\mathcal{PT}x = -x$. Thus we see that *every tensor index, on γ or on ∂ , gives a minus sign under CPT transform*. For a Hermitian Lagrangian (coefficients are appropriately real or pure imaginary) which is a Lorentz scalar (so even rank tensor), CPT contributes no extra sign and only changes the space-time argument i.e.

$$\boxed{(\mathcal{CPT})\mathcal{L}(x)(\mathcal{CPT})^\dagger = \mathcal{L}(-x)}. \text{ Thus,}$$

a Hermitian action, invariant under proper and orthochronous Lorentz transformations, is invariant under the discrete CPT transformations.

We have explicitly verified it for the quantum, Dirac field. Verification for scalar and Maxwell field is left as an exercise.

Note: Had we not used the extra minus sign attributed to the fermionic nature of the fields, we would have got the CPT transform of the Lagrangian to be minus of $\mathcal{L}(-x)$ and got CPT non-invariance of the action! This minus sign is tightly connected with the spin-statistics theorem which we discuss next.

D. Relativistic Causality and the Spin-Statistics Theorem

Now we come to the second feature of quantum fields - the commutation relations. Here we get another surprise. The Pauli exclusion principle follows from the requirement of relativistic causality!

The Fourier decomposition suggested field as a collection of harmonic oscillators. A natural quantization procedure is to postulate $[a_{\vec{k}}, a_{\vec{k}'}^\dagger] = \delta^3(\vec{k} - \vec{k}')$. This has a potential

problem for $\vec{k} = \vec{k}'$. The usual way this is interpreted is to imagine the field being confined to a large box satisfying periodic or Dirichlet boundary conditions. This discretizes the momentum labels, $\vec{k} \rightarrow \vec{k}_n \sim 2\pi\vec{n}/L$ and also replaces the delta function by Kronecker delta. Another way is to choose a suitable, countable orthonormal set of functions $\{\varphi_n(\vec{x})\}$ such that $\nabla^2\varphi_n = -\omega_n^2\varphi_n$. Completeness of the φ_n 's allows translating $[a_m, a_n^\dagger]$ to $[\Phi(t, \vec{x}), \Pi(t, \vec{y})] = \delta^3(x - y)$. We will implicitly assume such a procedure and proceed formally using the delta function.

From the postulated commutation relations we have, for each \vec{k} ,

$$\begin{aligned} N_{\vec{k}} &:= a_{\vec{k}}^\dagger a_{\vec{k}} \ , \ [N_{\vec{k}}, a_{\vec{k}}] = -a_{\vec{k}} \ , \ [N_{\vec{k}}, a_{\vec{k}}^\dagger] = a_{\vec{k}}^\dagger \ , \\ N_{\vec{k}}|n_{\vec{k}}\rangle &= n_{\vec{k}}|n_{\vec{k}}\rangle \ , \ n_{\vec{k}} = 0, 1, \dots; \ \langle m_{\vec{k}}|n_{\vec{k}}\rangle = \delta_{m_{\vec{k}}, n_{\vec{k}}} \\ |n_{\vec{k}}\rangle &:= \frac{(a_{\vec{k}}^\dagger)^{n_{\vec{k}}}}{\sqrt{(n_{\vec{k}})!}}|0_{\vec{k}}\rangle \ , \ a_{\vec{k}}|n_{\vec{k}}\rangle = \sqrt{n_{\vec{k}}}|n_{\vec{k}} - 1\rangle \ , \ a_{\vec{k}}^\dagger|n_{\vec{k}}\rangle = \sqrt{n_{\vec{k}} + 1}|n_{\vec{k}} + 1\rangle \end{aligned}$$

The state label $n_{\vec{k}}$ is interpreted as denoting the number of quanta, created by $a_{\vec{k}}^\dagger$ and destroyed by $a_{\vec{k}}$.

Note: In this case of a scalar field, all quanta have the same mass parameter, but different $k_{\vec{k}}$ label. For a vector field, we have the polarization label λ and for the spinor field we have the spin projection label σ . All quanta of a given mass m and spin/helicity are identical.

Now we add a postulate of *relativistic causality* [1] also referred to as *micro-causality*, namely, *all pairs of fields and their adjoints, commute or anti-commute for space-like separation*.

This requirement comes about as follow. In a Lorentz invariant theory, all observables would be Lorentz tensors of various ranks. Spinors being double valued quantities, they themselves are *not observables*. Spinorial fields must come in even powers (single valued) in an observable. As per general principles of quantum theory, namely, simultaneously measurable observables must commute and the notion of relativistic causality that simultaneously measurable observables must be space-like separated, suggest that for a space-time dependent operators to be observables, they must commute for space-like separation. Operators built out of spinorial fields being at least quadratic in the spinorial fields, allows possible anti-commutation rules for spinor fields⁷.

⁷ Weinberg bases his argument in favor of relativistic causality by invoking Lorentz invariance of the scattering matrix.

All our mode decompositions have annihilation operators as coefficients of the positive frequency modes and the creation operators as coefficients of the negative frequency modes. Let us denote these parts as $\Psi_+(x)$ and $\Psi_-(x)$ respectively. Consider a general linear combination of these fields: $\Psi(x) := \alpha\Psi_+(x) + \beta\Psi_-(x)$ and let $\Psi^\dagger(x)$ denote its adjoint. Let all observables be built out of $\Psi(x)$ and $\Psi^\dagger(x)$. Let $[A, B]_\pm := AB \pm BA$, the anti-commutator and commutator respectively. The requirement of micro-causality is then stated as: $\boxed{[\Psi(x), \Psi(y)]_\pm = 0 = [\Psi(x), \Psi^\dagger(y)]_\pm \text{ for } (x - y)^2 > 0}$. This requirement is quite restrictive, as seen below.

Let us begin with a real scalar field. We have,

$$\phi_+(x) := \int \frac{d^3k}{\sqrt{2\omega_{\vec{k}}}(2\pi)^3} a_{\vec{k}} e^{ik \cdot x}, \quad \phi_-(x) := \int \frac{d^3k}{\sqrt{2\omega_{\vec{k}}}(2\pi)^3} a_{\vec{k}}^\dagger e^{-ik \cdot x} = \phi_+^\dagger(x).$$

$$\therefore [\phi_+(x), \phi_-(y)]_\pm = \int \frac{d^3k}{\sqrt{2\omega_{\vec{k}}}(2\pi)^3} \int \frac{d^3k'}{\sqrt{2\omega_{\vec{k}'}(2\pi)^3}} e^{ik \cdot x - ik' \cdot y} [a_{\vec{k}}, a_{\vec{k}'}^\dagger]_\pm$$

For the scalar field, we use the commutator given above. Let us assume the same form for the anti-commutator i.e. we assume, $[a_{\vec{k}}, a_{\vec{k}'}^\dagger]_\pm = \delta^3(\vec{k} - \vec{k}')$. This gives,

$$[\phi_+(x), \phi_-(y)]_\pm = \int \frac{d^3k}{2\omega_{\vec{k}}(2\pi)^3} e^{ik \cdot (x-y)} =: \Delta_+(x-y) \text{ is Poincare invariant. For}$$

space-like separation, choose $(x-y)^2 = (\vec{x} - \vec{y})^2 =: r^2$,

$$\text{Thus } \Delta_+(x^2) = \frac{1}{(2\pi)^3} \int_0^\infty dk \frac{k^2}{2\sqrt{k^2 + m^2}} \int_{-1}^1 d\cos(\theta) \int_0^{2\pi} d\varphi e^{ikr\cos\theta}$$

$$= \frac{1}{4\pi^2} \int_0^\infty dk \frac{k^2}{\sqrt{k^2 + m^2}} \frac{\sin(kr)}{kr} \quad \text{put } k = m\alpha,$$

$$\Delta_+(x^2) = \frac{1}{4\pi^2} \frac{m}{r} \int_0^\infty d\alpha \frac{\alpha}{\sqrt{\alpha^2 + 1}} \sin(mr\alpha)$$

$$= -\frac{1}{4\pi^2} \frac{1}{r} \frac{\partial}{\partial r} \int_0^\infty d\alpha \frac{\alpha}{\sqrt{\alpha^2 + 1}} \cos(mr\alpha)$$

Thus, $\boxed{\Delta_+(x^2 > 0) = K_1(m\sqrt{x^2})}$ (Modified Bessel function of the second kind), and it is *non-zero for space-like separation*.

Define $\phi(x) := \alpha\phi_+(x) + \beta\phi_-(x)$, $\phi^\dagger(y) = \alpha^*\phi_-(y) + \beta^*\phi_+(y)$. It follows,

$$\begin{aligned} [\phi(x), \phi(y)]_\pm &= \alpha\beta [\phi_+(x), \phi_-(y)]_\pm + \beta\alpha [\phi_-(x), \phi_+(y)]_\pm \\ &= \alpha\beta (\Delta_+(x-y) \pm \Delta_+(y-x)) \\ &= \alpha\beta(1 \pm 1)\Delta_+(x-y) \quad \text{since } \Delta_+ \text{ is symmetric in } x, y. \\ [\phi(x), \phi^\dagger(y)]_\pm &= |\alpha|^2 [\phi_+(x), \phi_-(y)]_\pm + |\beta|^2 [\phi_-(x), \phi_+(y)]_\pm \\ &= (|\alpha|^2 \pm |\beta|^2)\Delta_+(x-y) \end{aligned}$$

To satisfy the requirement of causality, both the brackets must vanish i.e. in both equations we *must* choose the *minus* sign i.e. a *commutator* as well as the condition $|\alpha| = |\beta|$. This is quite a strong restriction on both the sign as well as the coefficients α, β .

Let us consider the spinor field now. Let,

$$\begin{aligned} \psi_+(x) &:= \int \frac{d^3k}{\sqrt{2\omega_{\vec{k}}}(2\pi)^3} \sum_{\sigma} (u(\vec{k}, \sigma) e^{ik \cdot x}) b_{\vec{k}, \sigma}, & \longleftrightarrow \\ (\psi_+)^\dagger(x) &:= \int \frac{d^3k}{\sqrt{2\omega_{\vec{k}}}(2\pi)^3} \sum_{\sigma} (u^*(\vec{k}, \sigma) e^{-ik \cdot x}) b_{\vec{k}, \sigma}^\dagger; \\ \psi_-(x) &:= \int \frac{d^3k}{\sqrt{2\omega_{\vec{k}}}(2\pi)^3} \sum_{\sigma} (v(\vec{k}, \sigma) e^{-ik \cdot x}) d_{\vec{k}, \sigma}^\dagger, & \longleftrightarrow \\ (\psi_-)^\dagger(x) &:= \int \frac{d^3k}{\sqrt{2\omega_{\vec{k}}}(2\pi)^3} \sum_{\sigma} (v^*(\vec{k}, \sigma) e^{+ik \cdot x}) d_{\vec{k}, \sigma}. \end{aligned}$$

The Dirac index is suppressed and the dagger refers to the adjoint of the operators and not of the spinors.

Let $\psi(x) := \mu\psi_+(x) + \nu\psi_-(x)$ and $\psi^\dagger(y) := \mu^*\psi_+^\dagger(y) + \nu^*\psi_-^\dagger(y)$. Then,

$$\begin{aligned} [\psi_\alpha(x), \psi_\beta^\dagger(y)]_\pm &= \int \frac{d^3k}{2\omega_{\vec{k}}(2\pi)^3} \sum_{\sigma} \left\{ |\mu|^2 u_\alpha(\vec{k}, \sigma) u_\beta^*(\vec{k}, \sigma) e^{ik \cdot (x-y)} \right. \\ &\quad \left. \pm |\nu|^2 v_\alpha(\vec{k}, \sigma) v_\beta^*(\vec{k}, \sigma) e^{-ik \cdot (x-y)} \right\} \end{aligned}$$

The sums over σ are given by,

$$\begin{aligned} \sum_{\sigma} u_\alpha(\vec{k}, \sigma) u_\beta^*(\vec{k}, \sigma) &= \sum_{\sigma} [u(\vec{k}, \sigma) \bar{u}(\vec{k}, \sigma) \gamma^0]_{\alpha\beta} = \frac{-\not{k} + m}{2m} \gamma^0 \Rightarrow \\ \sum_{\sigma} u_\alpha(\vec{k}, \sigma) u_\beta^*(\vec{k}, \sigma) e^{ik \cdot (x-y)} &= (2m)^{-1} (i\not{\partial} + m) \gamma^0 e^{ik \cdot (x-y)} \quad \text{and likewise,} \\ \sum_{\sigma} v_\alpha(\vec{k}, \sigma) v_\beta^*(\vec{k}, \sigma) &= \sum_{\sigma} [v(\vec{k}, \sigma) \bar{v}(\vec{k}, \sigma) \gamma^0]_{\alpha\beta} = \frac{-\not{k} - m}{2m} \gamma^0 \Rightarrow \\ \sum_{\sigma} v_\alpha(\vec{k}, \sigma) v_\beta^*(\vec{k}, \sigma) e^{-ik \cdot (x-y)} &= -(2m)^{-1} (i\not{\partial} + m) \gamma^0 e^{-ik \cdot (x-y)}; \end{aligned}$$

and the integration over \vec{k} just gives the $\Delta_+((x-y)^2)$ function computed above. Hence,

$$\left[\psi_\alpha(x), \psi_\beta^\dagger(y) \right]_{\pm} = (|\mu|^2 \mp |\nu|^2) \frac{i\not{\partial} + m}{2m} \gamma^0 \Delta_+((x-y)^2)$$

Once again, the right hand side vanishes for space-like separation provided the upper signs are chosen *and* $|\mu| = |\nu|$. Thus, the for spinors we have to choose anti-commutation relations and the weights of the positive and negative frequency solutions must be the same up to a phase factor. We may just take $\mu = \nu = 1$ which gives back our previous mode decomposition.

Note: While writing the mode decomposition, we did not bother about the normalizations of the coefficients, b, d^* etc. Now, in choosing the brackets, $[b_{\vec{k},\sigma}, b_{\vec{k}',\sigma'}^\dagger]_+ = \delta_{\sigma,\sigma'} \delta^3(\vec{k} - \vec{k}')$ and ditto for the d 's, we have fixed these normalizations. We could still have arbitrary multiple in each term. The requirement of causality restricts this freedom to the above.

The two examples considered above, generalize to other representations as well and what we have is the celebrated *spin-statistics theorem*:

A relativistic quantum field theory satisfies the requirement of causality provided integer spin/helicity field quantum conditions use commutators (bosons) while the half integer spin/helicity field quantum conditions use anti-commutators (fermions).

Note: While discussing quantum statistics in statistical mechanics, we invoke the additional attribute of *indistinguishability* among identical particles and require the Bose-Einstein/Fermi-Dirac statistics. Identical particles are defined by having identical intrinsic attributes such as mass, spin, color, flavor etc. The (in)distinguishability arises from a spatial localization and subsequent ability to tag them through, say, a scattering process. This is lost when their wave functions overlap, the ‘quantum’ becomes operative and quantum statistics becomes essential. In the above discussion, no such additional property emerged. We already have the ‘quantum’ operative so indistinguishability is also operative. It is the demand of relativistic causality that forced the spin-statistics correlation.

With the anti-commutators around, $[b_{\vec{k},\sigma}, b_{\vec{k}',\sigma'}^\dagger]_+ = \delta_{\sigma,\sigma'} \delta^3(\vec{k} - \vec{k}')$, $[b_{\vec{k},\sigma}, b_{\vec{k}',\sigma'}]_+ = 0$, we define the number operators for each (\vec{k}, σ) as before: $N := b^\dagger b$ It follows that $[N, b]_- = -b$, $[N, b^\dagger]_- = +b^\dagger$. Eigenvalues of N continue to be integers but thanks to $(b^\dagger)^2 = 0$, they take only two values, 0, 1. The corresponding eigenstates are: $|0\rangle$ and $|1\rangle$. This is the incorporation of the Pauli exclusion principle.

Consider states of two modes (scalar for simplicity), say \vec{k}_1, \vec{k}_2 . A general basis state

would be $|\vec{k}_1, \dots, \vec{k}_1, \vec{k}_2, \dots, \vec{k}_2\rangle \sim (a_{\vec{k}_1}^\dagger)^m (a_{\vec{k}_2}^\dagger)^n |0, 0\rangle$.

If the a 's anti-commute, then $m, n \leq 1$ and we have 4 possible states: $|0, 0\rangle, |\vec{k}_1, 0\rangle, |0, \vec{k}_2\rangle, |\vec{k}_1, \vec{k}_2\rangle$. Thanks to anti-commutation, $|\vec{k}_2, \vec{k}_1\rangle = -|\vec{k}_1, \vec{k}_2\rangle$. For bosonic operators, there is no minus sign under exchange. Generalizing from this, it is clear that (anti-)commutation relations among the creation operators automatically ensure complete (anti-)symmetry under permutations of the labels.

To understand the extra minus sign in the third line of eq. (8.6), momentarily write the $\bar{\Psi}$ as $\bar{\Psi}_1$ and Ψ as Ψ_2 . Then in the third line of the equation, we see an explicit exchange of Ψ_1 and Ψ_2 . The fermionic commutation relation then gives that extra minus sign. As noted before, this is also important for the CPT theorem.

To summarize: Quantum fields are operators *tagged by space-time points* and obeying certain commutation relations. We may think of them as a collection of creation-annihilation operators for each mode (solution of the equations of motion). A general state of a quantum field is a linear combination of basis states which are multi-quanta states. Each quantum with momentum label \vec{k} and a spin/helicity label σ carries energy-momentum-angular momentum as given by the Poincare charges. As per quantum field theory, all processes involve exchanges of quanta among different quantum fields.

9. STATES OF FREE QUANTUM FIELDS: PARTICLES, COHERENCE AND COHERENT STATES

While we argued that particle dynamics with position-momentum is not compatible with relativity, we observe particles in a variety of experiments eg as tracks in bubble chambers, photographic emulsions, stacks of particle counters etc and also use the particle view in designing accelerator beams. So the more sophisticated relativistic framework should also have a suitable description of “particles”. Since quantum fields are operators, the description must be in terms of *states of a quantum field*. In the section (7), we saw that from a dynamical view point, a (free) field is an infinite collection of non-interacting harmonic oscillators. Furthermore, its quantization follows from the quantization of oscillators and A quantum field emerges as a linear combination of creation-annihilation operators of its modes. Its states are built by the action of the creation operators on vacuum state of each mode. We have to identify “particles” in this huge space of states of a quantum field.

A. Particle/anti-particle wave packets

Let $|0\rangle$ denote the unique vacuum state annihilated by *all* annihilation operators $a_{\vec{k}}$. We can generate “1-particle states” by taking a linear combination of the form $\sum_{\vec{k}} \alpha(\vec{k}) a_{\vec{k}}^\dagger |0\rangle$, “2-particle states” by taking a linear combination of the form $\sum_{\vec{k}, \vec{l}} \alpha(\vec{k}, \vec{l}) a_{\vec{k}}^\dagger a_{\vec{l}}^\dagger |0\rangle$ and so on. The totality of all such states forms the state space of a free quantum field. For the n -particle states, the (anti-)commutators of the creation operators automatically take care of (anti-)symmetrization of the states. Notice that 1-particle state does *not* mean a *single* $a_{\vec{k}}^\dagger |0\rangle$, but a linear combination of several creation operators. These allow us to form *wave packets*.

As an explicit example, let us construct a wave packet of an anti-fermion with some *momentum and spin distribution*. We have the Dirac field,

$$\Psi(x) = \int [d^3k] \left(b_{\vec{k}, \sigma} u(\vec{k}, \sigma) e^{ik \cdot x} + d_{\vec{k}, \sigma}^\dagger v(\vec{k}, \sigma) e^{-ik \cdot x} \right) \quad , \text{ where,}$$

$$[d^3k] := \frac{d^3k}{\sqrt{2\omega_{\vec{k}}(2\pi)^3}} \quad , \quad \text{and} \quad (\not{k} + m)u = 0 = (-\not{k} + m)v.$$

Acting on the vacuum, it will create a *1-anti-fermion* state, the linear combination from the

$d_{\vec{k},\sigma}^\dagger$ creation operators. Let $f(\vec{k}', \sigma')$ be a suitable complex valued function. Define

$$\langle f| := \int d^3k' \sum_{\sigma'} f(\vec{k}', \sigma') \langle 0|d_{\vec{k}',\sigma'} \quad , \quad \int d^3k \sum_{\sigma} |f(\vec{k}, \sigma)|^2 = 1 . \quad (9.1)$$

$$\begin{aligned} (\chi)_c(x) &:= \langle f|\Psi(x)|0\rangle \\ &= \int [d^3k] \sum_{\sigma} f(\vec{k}, \sigma)v(\vec{k}, \sigma)e^{-ik \cdot x} \quad \because \quad \langle 0|d_{\vec{k}',\sigma'}d_{\vec{k},\sigma}^\dagger|0\rangle = \delta_{\sigma,\sigma'}\delta^3(\vec{k}' - \vec{k}) \end{aligned} \quad (9.2)$$

The so constructed $(\chi)_c(x)$ is a spinor-valued function of x which satisfies the Dirac equation. The $v(\vec{k}, \sigma)$ spinor signifies that it is an anti-fermion wave packet and the suffix c on χ is a reminder. How do we get a fermion wave packet? Well, in place of $\Psi(x)$ operator, we use its charge conjugate, $C\bar{\Psi}^T(x)$ (see 8.2), and in place of the $d_{\vec{k},\sigma}^\dagger$ in $\langle f|$ we use $b_{\vec{k},\sigma}$. The corresponding fermion wave packet takes the form,

$$\langle g| := \int d^3k' \sum_{\sigma'} g(\vec{k}', \sigma') \langle 0|b_{\vec{k}',\sigma'} \quad , \quad \int d^3k \sum_{\sigma} |g(\vec{k}, \sigma)|^2 = 1 . \quad (9.3)$$

$$\begin{aligned} \chi(x) &:= \langle g|C\bar{\Psi}^T(x)|0\rangle \quad \text{and} \quad \langle 0|b_{\vec{k}',\sigma'}b_{\vec{k},\sigma}^\dagger|0\rangle = \delta_{\sigma,\sigma'}\delta^3(\vec{k}' - \vec{k}) \Rightarrow \\ &= \int [d^3k] \sum_{\sigma} g(\vec{k}, \sigma)u^c(\vec{k}, \sigma)e^{-ik \cdot x} \end{aligned} \quad (9.4)$$

where, $u^c(\vec{k}, \sigma) = (i\gamma^2 u^*(\vec{k}, \sigma))$ is the charge conjugate of the $u(\vec{k}, \sigma)$ spinor.

The generalization for wave packets of other fields should be obvious. All these wave packets are positive or negative frequency solutions of the corresponding field equations. We already have discussed the inner products on these spaces and we may view these as the usual probability amplitudes. Of course we do not have the analogue of position operator unless we restrict to a non-relativistic approximation/limit. Suffice it to say that one can use Dirac equation to analyze beams of relativistic fermions, as is done for instance in [8].

B. Correlation Functions and Coherence of states

There are of course more general states which are not labeled by the number of quanta. Even for a single oscillator, states with a given occupation number is only a class of states. We can have finite linear combination or infinite linear combinations of these numbers states, can have squeezed states, coherent states etc. They are distinguished by their properties and utility. A some what more general characterization of quantum states is in terms of *correlation functions*. Let us see an example of this.

Consider a real scalar quantum field. We have its mode decomposition and had also defined the $\phi_{\pm}(x)$ fields consisting of the positive/negative frequency parts. Consider the measurement of the operators $\phi_{-}(x)\phi_{+}(y)$. Its average value is *some state* which is kept implicit, is denoted as

$$\begin{aligned} G(x, x') &:= \langle \phi_{-}(x)\phi_{+}(y) \rangle \quad (\text{a correlation function}) \\ &= \int \frac{d^3k}{\sqrt{2\omega_{\vec{k}}}(2\pi)^3} \int \frac{d^3k'}{\sqrt{2\omega_{\vec{k}'}}(2\pi)^3} e^{-i\vec{k}\cdot x + i\vec{k}'\cdot x'} \langle a_{\vec{k}}^{\dagger} a_{\vec{k}'} \rangle \end{aligned}$$

where, $\langle a_{\vec{k}}^{\dagger} a_{\vec{k}'} \rangle := \text{Tr}(\rho a_{\vec{k}}^{\dagger} a_{\vec{k}'})$ and thus the measured averaged value depends on the density matrix or “state”, ρ . The usual Schrodinger equation for the ket vectors becomes the equation $id_t\rho = [H, \rho]$ whose solution is: $\rho(t) = e^{-itH}\rho_0 e^{itH}$ for a time independent Hamiltonian. These are the only Hamiltonians considered below. For example, the Hamiltonian of our scalar field is given by the quantum version of (6.15).

A time-independent density matrix ρ is said to represent a *stationary* state. Thus for a stationary state, $[\rho, H] = 0$ which implies, $\rho = e^{-itH}\rho e^{itH} \forall t$.

$$\begin{aligned} \text{Tr} \left[\rho a_{\vec{k}}^{\dagger} a_{\vec{k}'} \right] &= \text{Tr} \left[e^{-itH} \rho e^{itH} a_{\vec{k}}^{\dagger} a_{\vec{k}'} \right] = \text{Tr} \left[\rho e^{itH} a_{\vec{k}}^{\dagger} a_{\vec{k}'} e^{-itH} \right] \\ &= \text{Tr} \left[\rho (e^{itH} a_{\vec{k}}^{\dagger} e^{-itH}) (e^{itH} a_{\vec{k}'} e^{-itH}) \right] \\ &= \text{Tr} \left[\rho a_{\vec{k}}^{\dagger} a_{\vec{k}'} \right] e^{it(\omega_{\vec{k}} - \omega_{\vec{k}'})} \quad \because H \sim \int [d^3k''] \omega_{\vec{k}''} N_{\vec{k}''} \end{aligned}$$

The left hand side is time independent, so $\text{Tr} \left[\rho a_{\vec{k}}^{\dagger} a_{\vec{k}'} \right] = 0$ unless $\omega_{\vec{k}} = \omega_{\vec{k}'}$ i.e. unless $|\vec{k}| = |\vec{k}'|$. For the same frequency, the trace need not vanish and we have a matrix in the labels \vec{k}, \vec{k}' . This matrix is hermitian (ρ is hermitian) and can be diagonalised. Hence, without loss of generality, $\boxed{\text{for a stationary state, } \text{Tr} \left[\rho a_{\vec{k}}^{\dagger} a_{\vec{k}'} \right] =: \bar{n}_{\vec{k}} \delta_{\vec{k}, \vec{k}'}}$, where $\bar{n}_{\vec{k}}$ denotes the average number of quanta in the \vec{k} mode. This leads to,

$$G(x, x') = \int \frac{d^3k}{2\omega_{\vec{k}}(2\pi)^3} \bar{n}_{\vec{k}} e^{i\omega_{\vec{k}}(t-t') - i\vec{k}\cdot(\vec{x}-\vec{x}')} \quad (\text{In a stationary state}).$$

Clearly, a *stationary state* leads to translational invariance of the above correlation function and this correlation function contains the information of the average number of quanta in each mode.

Assume now that the average number of quanta is vanishingly small outside a small neighbourhood of some direction \hat{k}_0 (collimation) and also outside a small neighbourhood

of $\omega_{\vec{k}_0}$ (monochromaticity). We may then approximate the integral as ,

$$G(x, x') \approx \frac{N_0}{2\omega_{\vec{k}_0}(2\pi)^3} e^{i\omega_{\vec{k}_0}(t-t') - i\vec{k}_0 \cdot (\vec{x} - \vec{x}')} , \quad N_0 \approx \int_{ngbd(\vec{k}_0)} \frac{d^3k}{2\omega_{\vec{k}}(2\pi)^3} \bar{n}_{\vec{k}}$$

It is apparent that the correlation function has a factorised form: $G(x, x') \approx \varphi^*(x)\varphi(x')$, $\varphi(x) := \sqrt{\frac{N_0}{2\omega_{\vec{k}_0}(2\pi)^3}} e^{ik_0 \cdot x}$ which is a plane wave. A state ρ with such a form of the correlation function is said to have a *first order coherence*⁸.

Notice that the factorised form results when the integral can be *approximated*. This in turn is valid for $|\vec{x} - \vec{x}'| \approx (2\pi)/|\vec{k}_0|$ and $|t - t'| \approx (2\pi)/\omega_{\vec{k}_0}$. Furthermore, the imperfections in the monochromaticity and collimation affect the approximation and also limit the validity of the first order coherence. This introduces the ideas of *coherence time* and *coherence length* as time and space intervals over which the factorization property or coherence property holds. Thus a stationary state with first order coherence, with finite coherence length and time, may be interpreted as a “beam” along the direction \hat{k}_0 with intensity proportional to $N_0 \omega_{\vec{k}_0}$.

Let us continue further and assume perfect collimation along say the x-axis. The beam may have a distribution of frequencies though. The phase⁹ then becomes $i\omega_{\vec{k}}(t - t' - x + x')$ and $G(x, x') =: G(\tau)$, $\tau := (t - t' - x + x')$. Since we have a single direction, we can denote the dependence on \vec{k} by the frequency ω . Suppose further that $\bar{n}_{\vec{k}} =: \bar{n}_\omega$ has a Gaussian dependence on ω : $\bar{n}_\omega = A \exp[-\frac{(\omega - \omega_0)^2}{2\sigma^2}]$. Now the integral can be done exactly and get,

$$G(\tau) = A \int_0^\infty \frac{d\omega}{\omega(2\pi)} e^{[-\frac{(\omega - \omega_0)^2}{2\sigma^2}]} e^{i\omega\tau} = G(0) e^{i\omega_0\tau} e^{-\frac{\sigma^2\tau^2}{2}} .$$

Thus, a highly collimated beam with a frequency range of σ around ω_0 , has the correlation function $G(x, x')$ decaying with $\tau = (t - t' - x + x') \sim \sigma^{-1}$. For example, if we have say a laser beam with $\omega_0 \sim 10^{14}$ Hz and $\sigma \sim 10^6$ Hz, then $\tau \sim 10^{-6}$ seconds or about 10^2 meters. Since the beam is presumed to be a stationary state, the τ does *not* vary with the time, t (or t'). However, beyond the coherence length, the beam will show frequency dispersion. Note that first order coherence suffices for the notions of coherence length and time.

⁸ k^{th} order coherence is defined in terms of the factorization property of $G^n(x_1, \dots, x_n, y_1, \dots, y_n) := \langle \phi_-(x_1), \dots, \phi_+(y_n) \rangle = E^*(x_1) \dots E^*(x_n) E(y_1) \dots E(y_n)$, where $E(x)$ is independent of n for all $1 \leq n \leq k$ [7]. The coherent states, eigenstates of the annihilation operators, familiar from the harmonic oscillator have infinite order coherence.

⁹ The formula is given for a massless field. More generally, we will have $\tau(\omega) = t - t' - \sqrt{1 - m^2/\omega^2}(x - x')$. For very high frequency/energy, the mass can be neglected.

Remark: In this example, we have stipulated the properties of stationarity and 1st order coherence to restrict the implicit state of the quantum field. These stipulations do not single out a particular state, but permits a subset of states ρ . All such states suffice to describe a beam with stable properties over the coherence interval.

C. Coherent States

There is a very important class of states of quantum fields, especially for electromagnetic field, namely the class of coherent states. These are familiar from the quantum harmonic oscillator. Let us recall their construction briefly.

1. An aside: Harmonic Oscillator States

With the usual creation-annihilation operators, $[a, a^\dagger] = 1$ we take the oscillator variables as:

$$\begin{aligned} q(t) &:= ae^{-i\omega t} + a^\dagger e^{i\omega t} := q_+(t) + q_-(t) , \\ p(t) &:= -\frac{i}{2}ae^{-i\omega t} + \frac{i}{2}a^\dagger e^{-\omega t} \leftrightarrow [q(t), p(t)] - i . \end{aligned}$$

Define:

(i) *Displacement operators:* For each $\alpha \in \mathbb{C}$,

$$\begin{aligned} D(\alpha) &:= e^{\alpha a^\dagger - \alpha^* a} = e^{\alpha a^\dagger} e^{-\alpha^* a} e^{-\frac{|\alpha|^2}{2}} \quad \text{Using the BCH formula,} \\ \therefore D^\dagger(\alpha) &= e^{-\frac{|\alpha|^2}{2}} e^{-\alpha a^\dagger} e^{\alpha^* a} = D(-\alpha) \quad \text{and} \quad D^\dagger(\alpha)D(\alpha) = \mathbb{1} \\ D^\dagger(\alpha) a D(\alpha) &= a + \alpha \mathbb{1} \quad , \quad D^\dagger(\alpha) a^\dagger D(\alpha) = a^\dagger + \alpha^* \mathbb{1} \quad , \quad \text{and} \\ D(\alpha + \beta) &= D(\alpha) D(\beta) e^{-\frac{i}{2}(\alpha\beta^* - \alpha^*\beta)} . \end{aligned}$$

(ii) *Squeezing operators:* for each complex number $\epsilon := re^{i\phi}$,

$$\begin{aligned} S(\epsilon) &:= e^{\frac{1}{2}(\epsilon^* a^2 - \epsilon (a^\dagger)^2)} , \quad S^\dagger(\epsilon) = S(-\epsilon) \quad , \quad S^\dagger(\epsilon)S(\epsilon) = \mathbb{1} \\ S^\dagger(\epsilon) a S(\epsilon) &= a \operatorname{ch}(r) - a^\dagger e^{-2i\phi} \operatorname{sh}(r) \quad , \quad S^\dagger(\epsilon) a^\dagger S(\epsilon) = a^\dagger \operatorname{ch}(r) - a e^{2i\phi} \operatorname{sh}(r) \end{aligned}$$

These operators define a two parameter family of states, *squeezed coherent states*, as

$$|\alpha, \epsilon\rangle := D(\alpha)S(\epsilon)|0\rangle \quad ; \quad \epsilon = 0 \leftrightarrow (\text{coherent states}), \quad \alpha = 0 \leftrightarrow (\text{squeezed states}).$$

For this family of states, it is straightforward to see,

$$\begin{aligned}\langle \alpha, \epsilon | a^2 | \alpha, \epsilon \rangle &= -e^{-2i\phi} \text{ch}(r) \text{sh}(r) + \alpha^2, & \boxed{\langle \alpha, \epsilon | a | \alpha, \epsilon \rangle = \alpha, \forall \epsilon} \\ \langle \alpha, \epsilon | (a^\dagger)^2 | \alpha, \epsilon \rangle &= -e^{2i\phi} \text{ch}(r) \text{sh}(r) + (\alpha^*)^2, & \langle \alpha, \epsilon | a^\dagger | \alpha, \epsilon \rangle = \alpha^* \\ \langle \alpha, \epsilon | N | \alpha, \epsilon \rangle &= |\alpha|^2 + \text{sh}^2(r) \\ a | \alpha, \epsilon \rangle &= -\text{sh}(r) e^{-2i\phi} D(\alpha) S(\epsilon) a^\dagger | 0 \rangle + \alpha | \alpha, \epsilon \rangle.\end{aligned}$$

The last equation shows that only for the squeezing parameter, $r = 0$, $|\alpha, 0\rangle$ is an eigenstate of the annihilation operator which is the usual definition of coherent states.

The Heisenberg uncertainty relation takes the form,

$$\begin{aligned}A &:= 1 + 2\langle N \rangle - 2|\langle a \rangle|^2 = 1 + 2\text{sh}^2(r) \\ B &:= (\Delta a)^2 e^{-2i\omega t} + (\Delta a^\dagger)^2 e^{2i\omega t} = -2\text{ch}(r) \text{sh}(r) \cos(\omega t + \phi) \\ (\Delta q)^2(t) &= A + B, \quad 4(\Delta p)^2(t) = A - B\end{aligned}$$

Clearly, for the coherent states $r = 0 \Rightarrow A = 1, B = 0$ and $(\Delta q)^2(\Delta p)^2 = \frac{1}{4}$ which saturates the uncertainty bound. The uncertainties are also time independent. Some of the basic properties of the coherent states are:

$$|\alpha\rangle = |\alpha, 0\rangle = D(\alpha)|0\rangle \Rightarrow \langle \alpha | \alpha \rangle = 1 \quad (9.5)$$

$$|\alpha\rangle = e^{-\frac{|\alpha|^2}{2}} \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} |n\rangle \quad (9.6)$$

$$\langle \beta | \alpha \rangle = e^{-\frac{|\alpha|^2 + |\beta|^2}{2} + \alpha\beta^*} \quad (9.7)$$

$$\mathbb{1} = \frac{1}{\pi} \int d^2\alpha |\alpha\rangle \langle \alpha| = \sum_{n \geq 0} |n\rangle \langle n|. \quad (9.8)$$

2. Correlation functions

This subsection is based on [6, 7].

We begin with the definition:

$$G^n(t_1, \dots, t_n; t_{n+1} \dots t_{2n}) := \langle q_-(t_1) \dots q_-(t_n) q_+(t_{n+1}) \dots q_+(t_{2n}) \rangle \quad (9.9)$$

$$= [e^{i\omega(t_1 + \dots t_n - t_{n+1} \dots - t_{2n})}] \text{Tr} [\rho (a^\dagger)^n a^n] \quad (9.10)$$

Since the \hat{q}_- and \hat{q}_+ commute among themselves, the ordering of (t_1, \dots, t_n) and $(t_{n+1} \dots t_{2n})$ is unimportant. Note that for (and only for) $\rho = |0\rangle \langle 0|$, the correlation functions vanish identically. In the following, this density matrix is *excluded*.

We invoke the general result: $\boxed{\forall \hat{A}, \text{ and } \forall \rho, \text{Tr}[\rho \hat{A}^\dagger \hat{A}] \geq 0.}$

Taking $A := \sum_i \alpha_i \hat{q}_+(t_i)$,

$$\langle A^\dagger A \rangle = \sum_{i,j} \alpha_i^* G^1(t_i, t_j) \alpha_j \geq 0 \quad \forall \alpha_i s .$$

Hence, the matrix $G^1(t_i, t_j)$ is non-negative. So its determinant is non-negative. Thus,

$$G^1(t_i, t_i) G^1(t_j, t_j) - G^1(t_i, t_j) G^1(t_j, t_i) \geq 0 .$$

From the definition of G^1 , it follows that $G^1(t_j, t_i) = G^1(t_i, t_j)^*$. Therefore $G^1(t, t)$ is real and in fact non-negative. The non-negative determinant condition becomes $\boxed{|G^1(t_i, t_j)|^2 \leq G^1(t_i, t_i) G^1(t_j, t_j)}$. This leads to the definition of the *normalized correlation functions*,

$$g^n(t_1, \dots, t_{2n}) := \frac{G^n(t_1, \dots, t_{2n})}{\sqrt{G^1(t_1, t_1) \dots G^1(t_{2n}, t_{2n})}} , \quad (9.11)$$

The determinant condition may be expressed as, $|g^1(t_i, t_j)| \leq 1$. Using the explicit definition,

$$g^1(t_i, t_j) = \frac{e^{i\omega(t_i - t_j)} \text{Tr}(\rho a^\dagger a)}{\sqrt{(\text{Tr}(\rho a^\dagger a) \text{Tr}(\rho a^\dagger a))}} = e^{i\omega(t_i - t_j)} \quad \therefore \quad |g^1(t_i, t_j)| = 1.$$

k^{th} order Coherence :

A state ρ is said to have *k^{th} -order coherence*, if $|g^n(t_1, \dots, t_{2n})| = 1 \quad \forall 1 \leq n \leq k$.

Remark: This is a property of a state, defined with respect to a specific kind of correlation function. Thus we could have different notions of coherence, if we adopt different correlation functions. The choice of this specific class of correlation functions has its roots in the measurements in optics which involves electromagnetic field. The correlation functions are also defined for classical fields (with the averages defined as ensemble averages over repeated observations) and have very similar properties as the quantum counterparts. The distinctions begin to show up when the intensities of the light beams is decreased to almost “single photon” level.

Remark: In light of the result in the preceding line, *every* state has 1st order coherence. It follows immediately that for k^{th} order coherence, $|G^n(t_1, \dots, t_{2n})|^2 = \prod_i G^1(t_i, t_i)$.

Result: *A state has a k^{th} order coherence iff there exist a function $E(t)$, independent of n , such that*

$$G^n(t_1, \dots, t_{2n}) = E^*(t_1) \dots E^*(t_n) E(t_{n+1}) \dots E(t_{2n}) \quad \forall 1 \leq n \leq k.$$

The proof is easy. First prove it for $G^1(t_1, t_2)$. From this the result for G^n follows. This factorization property is another characterization of k^{th} order coherence.

Among the pure states, we have states which are finite linear combinations of number states and those with infinite linear combinations. Let $|\psi\rangle$ be a state with finite linear combinations of the number eigenstates. Let K be the maximum eigenvalue of N appearing in the linear combination. Then $a^n|\psi\rangle = 0 \forall n > K$. Since, for pure states, $|g^n| = \frac{\langle\psi|(a^\dagger)^n a^n|\psi\rangle}{(\langle\psi|a^\dagger a|\psi\rangle)^n}$, it follows that $|g^n| = 0 \forall n > K$. Hence, *finite linear combinations are necessarily partially coherent i.e. have maximum order of coherence to be K . A fully coherent state must have infinite linear combinations.*

But not all infinite linear combinations have full coherence! Here are two example.

Coherent states $|\alpha, 0\rangle$: We have $|g^n| = \frac{(\alpha^*)^n \alpha^n}{(|\alpha|^2)^n} = 1 \forall n \geq 1$ and coherent states are fully coherent. These are the *only* non-vacuum states which have infinite order coherence. This may be checked by considering an arbitrary, infinitesimal perturbation of a coherent state, $|\Psi\rangle := (1 - \epsilon/2)|\alpha\rangle + \epsilon/2|\phi\rangle, |\phi\rangle \neq |0\rangle$ and computing $\langle\Psi|(a^\dagger)^n a^n|\Psi\rangle - (\langle\Psi|a^\dagger a|\Psi\rangle)^n$ to first order in ϵ .

Squeezed vacuum $|0, \epsilon\rangle$: Now,

$$|g^2| = \frac{\langle 0|S^\dagger(\epsilon)(a^\dagger)^2 a^2 S(\epsilon)|0\rangle}{\sqrt{(\langle 0|S^\dagger(\epsilon)a^\dagger a S(\epsilon)|0\rangle)^n}}$$

This evaluates to,

$$\begin{aligned} \text{Numerator:} &= \langle 0|S^\dagger a^\dagger S S^\dagger a^\dagger S S^\dagger a S S^\dagger a S|0\rangle \\ &= \langle 0|(\text{ch}(r)a^\dagger - \text{sh}(r)e^{2i\phi}a)(\text{ch}(r)a^\dagger - \text{sh}(r)e^{2i\phi}a) \\ &\quad (\text{ch}(r)a - \text{sh}(r)e^{-2i\phi}a^\dagger)(\text{ch}(r)a - \text{sh}(r)e^{-2i\phi}a^\dagger)|0\rangle \\ &= (\text{sh}(r))^2 \langle 0|a(\text{ch}(r)a^\dagger - \text{sh}(r)e^{2i\phi}a)(\text{ch}(r)a - \text{sh}(r)e^{-2i\phi}a^\dagger)a^\dagger|0\rangle \\ &= (\text{sh}(r))^2 [(\text{ch}(r))^2 + 2(\text{sh}(r))^2] \end{aligned}$$

$$\text{Denominator:} = \langle 0|S^\dagger N S|0\rangle = (\text{sh}(r))^2 \implies \boxed{|g^2| = 1 + 3(\text{sh}(r))^2 > 1 \forall |0, \epsilon\rangle.}$$

Thus while full coherence requires infinite linear combinations, they do not guarantee even a second order coherence!

Points to note as a summary:

- Quantum framework has vastly more number of states compared to the classical framework. The observed measurements are always averages with uncertainties. There are

many states which have the same average of a given set of observable. While uncertainties serve to distinguish among these, alternative attribute are provided by the correlation functions and notion of coherence. There is at least one class of states, the coherent states, which have full coherence, have time independent uncertainties (for oscillator dynamics) and have the smallest uncertainties compatible with the Heisenberg relation. This class of states is large enough to provide (over-complete) basis for the Hilbert space.

- We used a specific kind of correlation function and defined coherence of states w.r.t. these. These correlators have equal number of creation and annihilation operators which are ordered with annihilation operators to the right (the so-called *normal order*). This is prejudiced on typical detectors working by *absorbing* quanta. For opposite types of detectors, anti-normal order would be appropriate. The correlators with unequal number of a, a^\dagger operators, *are used* in measurement of phase information.

All these correlation functions are premised on processes being understood as emission and absorption of quanta.

- Generalizing, the coherent states of fields are essentially tensor product of single mode coherent states. We may thus denote a coherent state of a scalar field as $|\{\alpha_{\vec{k}}\}\rangle$ with $\alpha_{\vec{k}} := \langle \alpha_{\vec{k}} | a_{\vec{k}} | \alpha_{\vec{k}} \rangle$. Then,

$$\varphi(x) := \langle \{\alpha_{\vec{k}}\} | \hat{\phi}(x) | \{\alpha_{\vec{k}}\} \rangle = \int [d^3k] (\alpha_{\vec{k}} e^{ik \cdot x} + \alpha_{\vec{k}}^* e^{-ik \cdot x}).$$

Thus every coherent state of the quantum field gives a classical solution determined by the complex parameters $\alpha_{\vec{k}}$'s. Conversely, we may interpret a classical (free) field as the expectation value of the quantum field in the corresponding *coherent* state. It will faithfully mimic the usual classical description eg descriptions of wave solutions etc. This is how one understands the usual classical electromagnetic waves.

- There is also a generalization of coherent states for fermionic fields. This uses Grassmann variables, their algebra and calculus. The constructions are very analogous. I refer you to [7]

D. Evolution into a coherent state

Since we see a large set of classical field solutions, a question arises as to why does a quantum system, go into a coherent state? Under what conditions are these states manifested?

Under a free evolution (linear equation), a state which is say a finite linear combination of the number states will continue to maintain the finite linear combination and not evolve into a coherent state, interactions are essential. It suffices to have an interaction with a *classical* source i.e. a c-number source. This is demonstrated most easily by using interaction picture for a single harmonic oscillator.

Recall:

Let $|\psi\rangle_S$ denote a state vector in the Schrodinger picture and A_S denote a generic operator without explicit time dependence and let the Hamiltonian be also time independent. The *Schrodinger picture* is defined by: $id_t|\psi(t)\rangle_S = H|\psi(t)\rangle$, $d_t A_S = 0$. Define a new ‘picture I’ with an arbitrary unitary operator $V(t)$, $V(0) = \mathbb{1}$ as,

$$\begin{aligned} |\psi(t)\rangle_I &:= V(t)|\psi(t)\rangle_S \quad , \quad A_I(t) := V^\dagger(t)A_S V(t) \quad \Rightarrow \\ id_t|\psi(t)\rangle_I &= id_t V^\dagger(t)|\psi(t)\rangle_S + V^\dagger(t)H_S|\psi(t)\rangle_S = V^\dagger(t)[(-id_t + H_S)V(t)|\psi(t)\rangle_I \\ \therefore id_t|\psi(t)\rangle_I &= [H_I - V^\dagger(t)id_t V(t)]|\psi(t)\rangle_I \quad , \quad H_I(t) := V^\dagger(t)H_S V(t) \quad . \quad \text{Similarly,} \\ id_t A_I(t) &= [A_I(t), V^\dagger(t)id_t V(t)] \quad . \end{aligned}$$

For the choice $d_t V(t) = 0$, we get back the Schrodinger picture. For the choice $id_t V(t) = V(t)H_I(t) \leftrightarrow id_t V(t) = H_S V(t)$, we get $d_t|\psi(t)\rangle_I = 0$, $id_t A_I(t) = [A_I(t), H_I(t)]$ and we have the *Heisenberg picture*.

Let us split the Schrodinger Hamiltonian as $H_S := H_0 + H'$ where H_0 is the “free” Hamiltonian and H' is the “interaction” Hamiltonian. Now we choose, $id_t V_I(t) = H_0 V_I(t)$. This results in the equations,

$$id_t|\psi(t)\rangle_I = (H_I - (H_0)_I)|\psi(t)\rangle_I := H'_I|\psi(t)\rangle \quad , \quad id_t A_I(t) = [A_I, (H_0)_I] \quad (9.12)$$

This is the “Dirac” or the “interaction” picture wherein the states evolve by interaction Hamiltonian while the operators evolve by the free Hamiltonian. Notice that $(H_0)_I(t) = H_0$.

Consider now a quantum harmonic oscillator interacting with a “classical source” i.e. $H = (\frac{1}{2}\hat{p}^2 + \frac{1}{2}\omega^2\hat{q}^2) + J(t)\hat{q}(t)$, where $J(t)$ is an externally prescribed real function of time.

We express the Hamiltonian as,

$$H = \omega(a^\dagger a + \frac{1}{2}) + J(t)(a + a^\dagger) \quad := \quad H_0 + H' .$$

In the interaction picture,

$$\begin{aligned} id_t a_I(t) &= [a_I(t), H_0] = \omega [a_I, a_I^\dagger a_I] = \omega a_I \\ \therefore a_I(t) &= e^{-i\omega t} a_I(0) \quad , \quad a_I^\dagger(t) = e^{+i\omega t} a_I^\dagger(0) \\ \therefore H'_I &= J(t)e^{i\omega t} a_I(0) + J(t)e^{-i\omega t} a_I^\dagger(0) . \end{aligned}$$

The state vector evolves by $H'_I(t)$ which gives, $|\psi(t)\rangle = \exp\left(-i \int_0^t dt' H'_I(t')\right) |\psi(0)\rangle_I$. The exponential simplifies as,

$$\begin{aligned} e^{-i \int_0^t dt' H'_I(t')} &= e^{-i \int_0^t dt' [J(t')e^{i\omega t'} a + e^{-i\omega t'} a^\dagger]} \quad a := a_I(0) \quad , \quad a^\dagger := a_I^\dagger(0) \\ &= e^{\alpha(t)a^\dagger - \alpha^*(t)a} \quad , \quad \alpha(t) := -i \int_0^t dt' J(t') e^{-i\omega t'} . \text{ Evidently,} \\ |\psi(t)\rangle_I &= D(\alpha(t)) |\psi(0)\rangle_I . \end{aligned}$$

We see immediately that if $|\psi(0)\rangle_I = |0\rangle$, then at each t , $|\psi(t)\rangle_I$ is the coherent state $|\alpha(t)\rangle$.

Thus a quantum oscillator interacting with a classical source (linearly coupled) will evolve into a coherent state from its ground state.

The generalization to a quantum field is immediate. A quantum field in its ground state (vacuum state), upon interacting (linear coupling) with a classical source, will evolve into a coherent state.

Remark: You may try out different coupling to a classical source eg $J(t)q^2$, to see what happens. This is indeed what happens with a quantum field in a FRW expanding universe. The result is *squeezed states* of the field modes, in exactly the same manner. If the source is also an operator, then in general we do not expect the state to evolve into a coherent state.

10. QUANTUM FIELDS IN SCATTERING PHENOMENA

So far we have introduced and studied quantum fields in the “non-interacting” context in the sense, the equations of motion that they satisfied were linear. We alluded to interactions of the fields with sources/detectors, involving emission/absorption of quanta. A more detailed version of this is typically treated in interactions of quantum fields with atoms/molecules which are non-relativistic system making transitions among their bound states. There is another class of interactions which are manifested in *scattering* phenomena. These can be studied in high energy collisions and are a *direct tool to probe the basic microscopic interactions*.

We first discuss typical scattering arrangement, identify the relevant measurable physical quantities, set up a theoretical framework and see how quantum fields gel with the framework.

A typical scattering experiment involves shooting particles at some target to be studied and observing the deflected or emerging set of particles. Instead of fixed target we could also have colliding beams. It is usually the case that,

(i) we understand the projectile particles reasonably well and able to control their properties such as energy, momentum, spin, charge, shooting angles etc fairly accurately. It is presumed (a necessary condition) that the presence of target and/or other particles do not affect these properties significantly at least as long as they are well separated (spatially and/or temporally);

(ii) after a large but finite time, we can observe the emergent particles whose properties can be similarly ascertained and they too are presumed well understood;

(iii) the correlations among the incoming and outgoing particles, though determined by the intervening target region, can be *measured* independent of the target.

If we wish to view the combined system of projectile, target and emergent particles as a *single* dynamical system, then the dynamical system must have three separate evolutions at least approximately. These are: (a) evolution in the distant past ($t \rightarrow -\infty$); (b) evolution in the distant future ($t \rightarrow +\infty$) and (c) evolution during the intermediate time of interaction. For instance, in a classical scattering, we will have a trajectory beginning and ending in a region of the phase space, marked as “asymptotic region”. In these regions, the exact trajectory may be well approximated by a different, usually “free” Hamiltonian. In a

quantum setting, we may replace a trajectory in phase space by one in the state space. It should be noted right away that *not all trajectories are of the scattering type - classically or quantum mechanically*.

What are the typically observed quantities?

In practice, it is not a single particle that is sent in but rather a well collimated beam directed at the target region. After individual particles emerge, there is a spread in their direction as the initial conditions are not identical. Typically, these are detected at different locations by detectors which have finite aperture and therefore the state of the detected particles is also known approximately. Given the aperture of a detector and its distance to the target, we know the solid angle subtended at the scattering region. Thus, the basic measurements are: (i) $I(\hat{n})d\Omega :=$ number of particles or events detected per second along the direction \hat{n} . This may be measured for each species of particle separately or just the total; (ii) the prepared beam has a flux $J :=$ number of incident particles per second per unit transverse area of the beam. Both these numbers are known directly. They are obviously proportional and define: $I(\hat{n})d\Omega := d\sigma(\hat{n}, \dots)J$. The $d\sigma(\hat{n}, \dots)$ is called the *differential cross-section* and \dots refer to its dependence on other properties such as energy etc. These reflect the averaged attributes of the beam - average energy, momentum, polarization etc. The detection can be made finer with particle detection in coincidence, select specific polarization, charge etc. Depending upon the attributes tagged *and included* in $I(\hat{n})d\Omega$ we get *exclusive/inclusive cross-sections*. If one does not care even about the angle, we have the *total cross-section*. These are typically used in determining decay rates and in applications in statistical mechanics as input from the microscopic dynamics.

In summary, the only information that is used about the beam is the flux while finer details of the final state are used in defining various differential cross-sections. Thus any cross-section is defined as the ratio above and has dimensions of *area*. Experimentalist give these measured numbers while a theorist has to choose a favorite model and compute these numbers. For this, we turn to the theoretical formulation of scattering processes in the quantum framework. Here are some basic observations.

A. General scattering framework

Let \mathcal{H} be the Hilbert space of the combined system (projectile-target-scattered/emerged). Let H_0, H'_0 and H be the three Hamiltonian operators governing the time evolution in the remote past, distant future and all through. For simplicity, we take $H'_0 = H_0$, though they could be different. The evolution generated by H_0 is said to be “free”. All are assumed to have no explicit time dependence. All are assumed to be self-adjoint.

Let their corresponding eigenvalue equations be, $H_0\phi_n = \epsilon_n\phi_n$ and $H\psi_\alpha = \epsilon_\alpha\psi_\alpha$. We have the most general solutions expressed as,

$$\begin{aligned}\psi(t) &= \sum_{\alpha} c_{\alpha} e^{-i\epsilon_{\alpha}t} \psi_{\alpha}(0) \quad , \quad \phi(t) = \sum_n a_n e^{-i\epsilon_n t} \phi_n(0) \\ \psi_{\alpha}(0) &= \sum_n b_{\alpha,n} \phi_n(0) \quad \because \{ \psi_{\alpha}(0) \} , \{ \phi_n(0) \} , \text{ are complete.} \\ \therefore \psi(t) &= \sum_{\alpha,n} c_{\alpha} b_{\alpha,n} e^{-i\epsilon_{\alpha}t} \phi_n(0) \quad , \quad \text{all coefficients are time independent.}\end{aligned}$$

We are interested in those solutions $\psi(t)$ which resembles some free solution as $t \rightarrow \pm\infty$. Thus, as $t \rightarrow -\infty$,

$$\begin{aligned}\psi(t) - \phi(t) \rightarrow 0 &\Rightarrow \sum_n \left\{ \sum_{\alpha} c_{\alpha} b_{\alpha,n} e^{-i\epsilon_{\alpha}t + i\epsilon_n t} - a_n \right\} \phi_n(0) \rightarrow 0 . \\ \therefore a_n &\rightarrow \sum_{\alpha} c_{\alpha} b_{\alpha,n} e^{i(\epsilon_n - \epsilon_{\alpha})t} \quad \forall n . \quad t\text{-independence of coefficients implies,} \\ c_{\alpha} b_{\alpha,n} \text{ (no sum)} &= 0 \quad \text{whenever } \epsilon_{\alpha} \neq \epsilon_n .\end{aligned}$$

So, if H_0, H do not have any eigenvalue in common, then for every $c_{\alpha_*} \neq 0, b_{\alpha_*,n} = 0 \forall n$. Hence, *no* solution $\psi_{\alpha}(0)$ is approximable by a free evolution. For a sufficiently general solution $\phi(t)$ to approximate *some* exact solution $\psi(t)$, spectrum of H_0 should be a subset of the spectrum of H . Conversely, every solution with $c_{\alpha} = 0$ for every $\epsilon_{\alpha} \notin \text{Spec}(H_0)$, can be asymptotically approximated, by some free evolution. Such solutions are called *scattering solutions*.

Note: Usually, the spectrum of H_0 is continuous and bounded below. Its eigenstates span the Hilbert space. The scattering states of H however span only a subspace of \mathcal{H} . To span the full Hilbert space we need to include the bound states of H . We have been somewhat informal in the above argument, here are the sharper definitions.

Definition: A solution $|\psi(t)\rangle$ is said to be *incoming (outgoing)* if \exists a solution $|\phi_{in}(t)\rangle$ ($|\phi_{out}(t)\rangle$) such that

$$\lim_{t \rightarrow -\infty} \|\psi(t) - \phi_{in}(t)\| = 0 \quad \left(\lim_{t \rightarrow +\infty} \|\psi(t) - \phi_{out}(t)\| \right).$$

Note: If the prepared beam or the detected scattered particles are to be associated with a state of the full system, then it is necessary that *at the most only one* $|\psi(t)\rangle$ *can be asymptote to a given* $|\phi_{in}(t)\rangle$ *or a given* $|\phi_{out}(t)\rangle$. Establishing such a property for a model of scattering is called the basic existence and uniqueness problem.

The solutions can be tagged by states at any particular instance for example, “ $|\psi(t=0)\rangle$ ”. We identify $\mathcal{H}_{in} := \{\psi \in \mathcal{H} / \psi(t) \text{ is incoming}\}$ and $\mathcal{H}_{out} := \{\psi \in \mathcal{H} / \psi(t) \text{ is outgoing}\}$.

Potentially it is possible to have (i) $\psi \notin \mathcal{H}_{in}, \psi \notin \mathcal{H}_{out}$ (no scattering); (ii) $\psi \in \mathcal{H}_{in}$, but $\psi \notin \mathcal{H}_{out}$ (‘capture’ process); (iii) $\psi \notin \mathcal{H}_{in}$, but $\psi \in \mathcal{H}_{out}$ (‘decay’ process) and (iv) $\psi \in \mathcal{H}_{in} \cap \mathcal{H}_{out}$ (scattering).

Definition: A scattering system is *weakly asymptotically complete* if $\mathcal{H}_{in} = \mathcal{H}_{out}$. It may so happen that the subspace of bound states, \mathcal{H}_{bound} is such that $\mathcal{H} = \mathcal{H}_{bound} \oplus \mathcal{H}_{in} \oplus \mathcal{H}_{out}$. Then the system is said to be *asymptotically complete*.

These distinctions have a bearing on the definition of S -matrix and its unitarity.

Let $U(t), U_0(t)$ denote the unitary evolution operators corresponding to H_0, H respectively. Let $|\psi(t)\rangle = U(t-t')|\psi(t')\rangle$ be an incoming solution. Then $\exists \phi_{in}(t) = U_0(t-t')\phi_{in}(t')$ (t' arbitrarily chosen instant) such that,

$$\text{As } t \rightarrow -\infty \quad , \quad \|U(t-t')\psi(t') - U_0(t-t')\phi_{in}(t')\| \rightarrow 0$$

$$\therefore \quad \|\psi(t') - U^\dagger(t-t')U_0(t-t')\phi_{in}(t')\| \rightarrow 0$$

$$\therefore \lim_{t \rightarrow -\infty} \psi(t) = \lim_{t \rightarrow -\infty} U^\dagger(t-t')U_0(t-t')\phi_{in}(t') .$$

$$\text{Define: } \Omega_+ := \lim_{t \rightarrow -\infty} U^\dagger(t-t')U_0(t-t') = \lim_{t \rightarrow -\infty} U^\dagger(t)U_0(t) \quad \text{Using } U^\dagger(t) = U(-t),$$

$$\text{we have, } \boxed{\Omega_+ = \lim_{t \rightarrow -\infty} U(-t)U_0(t) \text{ and } |\psi(t')\rangle = \Omega_+|\phi_{in}(t')\rangle \forall t' .}$$

Likewise, for an asymptotically outgoing solution, we have,

$$\boxed{\Omega_- = \lim_{t \rightarrow +\infty} U(-t)U_0(t) \text{ and } |\psi(t')\rangle = \Omega_-|\phi_{out}(t')\rangle \forall t' .}$$

The Ω_\pm are called the *Moller operators*. They map free solutions to asymptotic solutions. The assumption of existence and uniqueness of scattering solutions, guarantees the existence

of these operators. Their adjoints are defined as, $\Omega_+^\dagger := \lim_{t \rightarrow -\infty} U_0(-t)U(t)$, $\Omega_-^\dagger := \lim_{t \rightarrow +\infty} U_0(-t)U(t)$. The adjoints give $\phi_{in}(t') = \Omega_+^\dagger \psi(t')$, $\phi_{out}(t') = \Omega_-^\dagger \psi(t')$ $\forall t'$.

It follows that $\psi(t) = \Omega_\pm \Omega_\pm^\dagger \psi(t)$, $\Omega_+^\dagger \Omega_+ \phi_{in}(t) = \phi_{in}(t)$ and $\Omega_-^\dagger \Omega_- \phi_{out}(t) = \phi_{out}(t)$. Since the $\phi_{in/out}$ span the full Hilbert space, the last two equalities imply that $\Omega_\pm^\dagger \Omega_\pm = \mathbb{1}$. The same is not true for the $\psi(t)$'s and hence $\Omega_\pm \Omega_\pm^\dagger \neq \mathbb{1}$. Thus the Moller operators are isometry operators but not unitary operators. For the special case of asymptotic completeness, we can write $\Omega_\pm \Omega_\pm^\dagger = \mathbb{1} - P_{bound}$ where P_{bound} is the projection operator on \mathcal{H}_{bound} . From the Schrodinger equations satisfied by the $\psi(t)$ and $\phi(t)$, it is easy to see that

$$\boxed{H\Omega_\pm = \Omega_\pm H_0 \text{ and } H_0\Omega_\pm^\dagger = \Omega_\pm^\dagger H .}$$

Consider now a solution which is both asymptotically incoming and outgoing. Then we have, $\psi(t) = \Omega_+ \phi_{in}(t)$ and $\phi_{out}(t) = \Omega_-^\dagger \psi(t)$. Therefore,

$$\phi_{out}(t) = \Omega_-^\dagger \Omega_+ \phi_{in}(t) , \quad \phi_{in}(t) = \Omega_+^\dagger \Omega_- \phi_{out}(t) \text{ and we define,} \quad (10.1)$$

$$S := \Omega_-^\dagger \Omega_+ \text{ on } [Range(\Omega_+)] \cap [Domain(\Omega_-)] \neq \emptyset . \quad (10.2)$$

Note that the range of Ω_+ are states in \mathcal{H}_{in} while the domain of Ω_-^\dagger are states in \mathcal{H}_{out} . The *scattering operator*, S , is thus well defined when weak asymptotic completeness holds. Similarly, we define $S^\dagger := \Omega_+^\dagger \Omega_-$ which is also well defined when weak asymptotic completeness holds. If in addition asymptotic completeness holds, then

$$S^\dagger S = \Omega_+^\dagger \Omega_- \Omega_-^\dagger \Omega_+ = \Omega_+^\dagger (\mathbb{1} - P_{bound}) \Omega_+ = \mathbb{1} - \mathbb{0} \text{ likewise,} \quad (10.3)$$

$$S S^\dagger = \Omega_-^\dagger \Omega_+ \Omega_+^\dagger \Omega_- = \Omega_-^\dagger (\mathbb{1} - P_{bound}) \Omega_- = \mathbb{1} - \mathbb{0} . \quad (10.4)$$

Thus the scattering operator is unitary when asymptotic completeness holds. This is regardless of the Moller operators being non-unitary. *The scattering operator S maps from free solutions $\phi_{in}(t)$ to free solutions ϕ_{out} .* Its matrix elements between the basis of free solutions define the *S -matrix*. Explicitly, let $\phi_{in}^a(t)$ and $\phi_{out}^b(t)$ denote incoming and out going free solutions. Then,

$$\boxed{S_{ba} := \langle \phi_{out}^b(t) | S | \phi_{in}^a(t) \rangle =: \langle \phi_{out}^b(t) | \tilde{\phi}_{out}^a(t) \rangle .} \quad (10.5)$$

The $\tilde{\phi}_{out}^a(t)$ state is a state that evolved from a specific incoming state while $\phi_{out}^b(t)$ is an arbitrary free state and S_{ba} is the inner product between the two.

Hence, S_{ba} gives the probability amplitude for a free state $|\phi_{in}(t)\rangle$ to evolve into an outgoing state $|\phi_{out}^b(t)\rangle$ - in short a transition amplitude.

The S operator defined in terms of the Moller operators is manifestly time independent. Its matrix elements however seem to have a time dependence through those of the $\phi_{in/out}(t)$. This apparent time dependence is actually absent and can be seen as follows.

Recall the boxed relations among $\Omega_{\pm}, \Omega_{\pm}^{\dagger}, H$ and H_0 . From these we get,

$$\begin{aligned} SH_0 &= \Omega_-^{\dagger} \Omega_+ H_0 = \Omega_-^{\dagger} H \Omega_+ = H_0 \Omega_-^{\dagger} \Omega_+ = H_0 S. \\ id_t S_{ba} &= id_t (\langle \phi_{out}^b(t) | S | \phi_{in}^a(t) \rangle) \\ &= -\langle \phi_{out}^b(t) | H_0 S | \phi_{in}^a(t) \rangle + 0 + \langle \phi_{out}^b(t) | i S H_0 | \phi_{in}^a(t) \rangle \\ &= \langle \phi_{out}^b(t) | [S, H_0] | \phi_{in}^a(t) \rangle = 0. \end{aligned}$$

In summary:

- A generic scattering system has \mathcal{H} and the three Hamiltonians H, H_0, H'_0 specified;
- There exist asymptotic subspaces of $\mathcal{H}, \mathcal{H}_{in}, \mathcal{H}_{out}$ elements of which give full solutions approaching some free solutions as $t \rightarrow \pm\infty$;
- Provided the system admits scattering states, there exist the Moller operators, Ω_{\pm} on \mathcal{H} with ranges in $\mathcal{H}_{in/out}$ respectively;
- If weak asymptotic completeness holds, \exists the scattering operators $S : \mathcal{H}_{in} \rightarrow \mathcal{H}_{out}$ and its adjoint S^{\dagger} . The scattering operator S is unitary provided asymptotic completeness holds.
- Once S_{ba} are known, the cross-section can be computed, since the number of scattered particles is proportional to $|S_{ba}|^2$, while the flux of the incoming particles is given by the probability current corresponding to $\phi_{in}(t)$.
- Much of the rigorous analysis goes in addressing the existence and uniqueness properties of the proposed scattering system. Further details may be seen in [9, 10].

In practice, rarely a scattering system is specified in sufficient details to address these issues, but they are at the very basic definitional level.

Returning to interacting quantum fields, we have the candidate free states - the states of free quantum fields with known free Hamiltonian H_0 . We may specify the interacting system by additional Lorentz invariant terms in the Lagrangian or specifying the Hamiltonian H .

However, we do not know what the Hilbert space of interacting quantum fields is let alone if it admits scattering states. But we do know what properties we would like to have for a scattering interpretation. The task now is to make suitable assumptions and build a computational recipe to define and compute the S -matrix elements.

B. Scattering with quantum fields: Heisenberg Picture

The above discussion has been in terms of the states evolving in time i.e. in Schrodinger picture. For interacting quantum fields we do not know the state space (except for the free fields). We do have Poincare covariance as a guide and this puts the field operators and their Poincare transforms at the center stage. We are thus led to use the Heisenberg picture and formulate the recipe in terms of the evolving operators. As states do not evolve in Heisenberg picture, we postulate certain properties for the states, postulate Poincare covariance on the operator evolution and make specific additional assumption of “asymptotic condition” to incorporate the scattering feature. A reference for this material is [11].

1. We choose the basis states of the interacting quantum fields to be labeled by mutually commuting, conserved Noether charges whose existence is guaranteed by the presumed symmetries. Among these symmetries is included the Poincare group (proper, orthochronous with/without discrete symmetries). The Poincare symmetry immediately gives eigenvalues of P^μ as a set of labels. Our first assumption is regarding the spectrum of P^μ .
2.
 - There exists a unique vacuum, eigenstate $|0\rangle$ with $P^\mu|0\rangle = 0$;
 - Eigenvalues, p^μ , of P^μ lie in the forward light cone i.e. $p^2 \leq 0, p^0 > 0$;
 - \exists stable, *single particle states* with masses m_i : $P^\mu|p^i\rangle = p_i^\mu|p^i\rangle$, $p_i^2 = -m_i^2$.
 - The eigenvalues 0 and m_i^2 of P^2 are discrete in the sense of being isolated values. The “multi-particle” states have their total momentum time-like but the corresponding $-p^2$ can take any value above their total rest mass.

Note: If there are “internal symmetries” (i.e. non-space-time symmetries), then it is conceivable that the zero eigenvalue of P^μ is degenerate, and carries some representation of the internal symmetry group (usually compact). Every one of these states




FIG. 1: Spectrum for a single, massive scalar field. The two particle states form a continuum above and including $-(p_1 + p_2)^2 = 4m^2$ hyperboloid.

must still be singlets of the Poincare group. But now, these vacua themselves carry some quantum numbers and quantum fields acting on these will create single particle states labeled by tensor product of representations of the internal symmetry group. This is not done and one stipulates the vacuum to be a singlet under *all symmetries*. Note: We have taken the stable particles to have a non-zero mass. This is a statement about the spectrum of P^2 in the full, exact interacting quantum field theory and not at some approximation of it. This would seem to exclude the massless representations of the Poincare group for interacting quantum fields. In practice i.e. in perturbation theory we will use massless photons and see the accompanying infrared problems.

3.
 - An interacting field operator, denoted schematically by $\Phi(x)$ satisfies an operator equation of the form, $(\square - m_0^2)\Phi(x) = J(x)$ where $J(x)$ is an operator built out of Φ and possibly other field operators. We re-write the field equation as $(\square - m^2)\Phi(x) = J(x) - (m^2 - m_0^2)\Phi(x) =: \tilde{J}(x)$
 - The field obeys the equal time (anti-)commutation relations:

$$[\Phi(t, \vec{x}), \Phi(t, \vec{y})] = 0 = [\Pi(t, \vec{x}), \Pi(t, \vec{y})] \quad , \quad [\Phi(t, \vec{x}), \Pi(t, \vec{y})] = i\delta^3(\vec{x} - \vec{y}) \quad .$$

If $J(x)$ has no derivatives of fields, the momentum field is $\Pi(t, \vec{x}) = \dot{\Phi}(t, \vec{x})$. This stipulates the quantized nature of the field while the equation of motion stipulates

the nature of interaction.

- \exists another set of fields, quantized $\Phi_{in}(x), \Phi_{out}(x)$, build out of the interacting field $\Phi(x)$, and satisfying the same set of equal time (anti-)commutation relations but obeying free field equations: $(\square - m^2)\Phi_{in/out} = 0$ where m is the “physical” mass. The in and out fields thus have the same mode decomposition introducing the creation/annihilation operators which create/destroy stable particles with physical mass m . These fields also transform the same way as the field $\Phi(x)$ under the symmetries, in particular, $\Phi_{in}(x + a) = e^{ia \cdot P}\Phi_{in}(x)e^{-ia \cdot P}$.

4. Asymptotic conditions: The exact interacting quantum field $\Phi(x)$ is linked to the in and out fields by the asymptotic conditions,

$$\boxed{\Phi(x)|_{t \rightarrow -\infty} \longrightarrow \sqrt{Z}\Phi_{in}(x) \quad , \quad \Phi(x)|_{t \rightarrow \infty} \longrightarrow \sqrt{Z}\Phi_{out}(x)}$$

Here Z is a possible normalization constant which we will show to be necessarily different from 1.

Note: The asymptotic conditions are *not* operator equations but are *weak* equations i.e. hold for arbitrary matrix elements in the basis of normalized states.

Below we elaborate the conditions and note a number of properties.

Asymptotic conditions are consistent with Poincare covariance: A formal solution of the equation satisfied by $\Phi(x)$ is given by,

$$\Phi(x) = \int d^4y G_{ret}(x - y, m)\tilde{J}(y) + \sqrt{Z}\Phi_{in}(x). \quad (10.6)$$

As $t \rightarrow -\infty$, the first term vanishes (property of retarded Green function) recovering the asymptotic condition. This is a heuristic argument since $J(y)$ it built out of $\Phi(x)$ and some iterative procedure is implicit. The consistency with covariance is seen as,

$$\begin{aligned} \sqrt{Z}\Phi_{in}(x - a) &= \Phi(x - a) - \int d^4y G_{ret}(x - a - y, m)\tilde{J}(y) \\ &= e^{-ia \cdot P}\Phi(x)e^{ia \cdot P} - \int d^4y G_{ret}(x - y, m)\tilde{J}(y - a) \\ &= e^{-ia \cdot P} \left[\Phi(x) - \int d^4y G_{ret}(x - y, m)\tilde{J}(y) \right] e^{ia \cdot P} = \sqrt{Z}e^{-ia \cdot P}\Phi_{in}(x)e^{ia \cdot P} . \end{aligned}$$

Note: Only the G_{ret} is used in the formal solution and not $G_{Feynman}$.

$\Phi_{in}(x)$ creates states with physical mass m : The infinitesimal form of the transformation law for $\Phi_{in}(x)$ implies, $[P^\mu, \Phi_{in}(x)] = -i\partial^\mu\Phi_{in}(x)$ and we also have the equation of motion $(\square - m^2)\Phi_{in} = 0$.

For any physical state $|k, \alpha\rangle$, $P^\mu|k, \alpha\rangle = k^\mu|k, \alpha\rangle$. Therefore,

$$\begin{aligned} -i\partial^\mu\langle k, \alpha|\Phi_{in}(x)|0\rangle &= \langle k, \alpha|[P^\mu, \Phi_{in}(x)]|k, \alpha\rangle = k^\mu\langle k, \alpha|\Phi_{in}(x)|0\rangle \\ \therefore -\square\langle k, \alpha|\Phi_{in}(x)|0\rangle &= k^2\langle k, \alpha|\Phi_{in}(x)|0\rangle \\ \therefore (-\square + m^2)\langle k, \alpha|\Phi_{in}(x)|0\rangle &= 0 = (k^2 + m^2)\langle k, \alpha|\Phi_{in}(x)|0\rangle \Rightarrow k^2 = -m^2. \end{aligned}$$

Thus, $\Phi_{in}(x)$ behaves exactly like a free field producing physical states from the vacuum and we can be taken to have the usual mode decomposition,

$$\begin{aligned} \Phi_{in}(x) &= \int \frac{d^3k}{\sqrt{2\omega_{\vec{k}}}(2\pi)^3} \left[a_{in}(\vec{k})e^{ik\cdot x} + a_{in}^\dagger(\vec{k})e^{-ik\cdot x} \right] \\ \partial_0\Phi_{in}(x) &= \int \frac{d^3k}{\sqrt{2\omega_{\vec{k}}}(2\pi)^3} \left[-i\omega_{\vec{k}}a_{in}(\vec{k})e^{ik\cdot x} + i\omega_{\vec{k}}a_{in}^\dagger(\vec{k})e^{-ik\cdot x} \right] \Rightarrow \\ a_{in}^\dagger(\vec{k}) &= \frac{-i}{\sqrt{2\omega_{\vec{k}}}(2\pi)^3} \int d^3x e^{ik\cdot x} \overleftrightarrow{\partial}_0 \Phi_{in}(x) \quad , \quad f \overleftrightarrow{\partial} g := f\partial g - \partial(f)g. \quad (10.7) \\ \text{And, } a_{in}(\vec{k}) &= \frac{i}{\sqrt{2\omega_{\vec{k}}}(2\pi)^3} \int d^3x e^{-ik\cdot x} \overleftrightarrow{\partial}_0 \Phi_{in}(x) \end{aligned}$$

Note: In writing the inversion formula, we have done the spatial integration over a $t = \text{constant}$ hypersurface Σ_t . The choice of this hypersurface does not matter ($a_{in}^\dagger(\vec{k})$ is independent of t). *This is kept implicit in all the inversion formulae below.*

It follows immediately, $\langle 0|\Phi_{in}(x)|0\rangle = 0$ and,

$$\begin{aligned} \langle k, \alpha|\Phi_{in}(x)|0\rangle &= \langle k, \alpha|e^{-ix\cdot P}\Phi_{in}(0)e^{ix\cdot P}|0\rangle = e^{-ik\cdot x}\langle k, \alpha|\Phi_{in}(0)|0\rangle \\ \langle k, \alpha|\Phi_{in}(0)|0\rangle &= \frac{1}{\sqrt{2\omega_{\vec{k}}}(2\pi)^2} \Rightarrow \boxed{\langle k, \alpha|\Phi_{in}(x)|0\rangle = \frac{e^{-ik\cdot x}}{\sqrt{2\omega_{\vec{k}}}(2\pi)^3}}. \end{aligned}$$

Suffice it to say that corresponding expressions exist for ϕ_{out} . In particular the heuristic expression take the form,

$$\Phi(x) = \sqrt{Z}\Phi_{out}(x) + \int d^4y G_{adv}(x-y, m^2)\tilde{J}(y).$$

How far can we get with Lorentz covariance and the assumptions on the spectrum?

C. Kallen-Lehmann Representation

Consider the vacuum expectation values of the commutator of interacting field, $i\Delta'(x, x') := \langle 0 | [\Phi(x), \Phi(x')] | 0 \rangle$. We had evaluated this for the free fields while considering the spin-statistics, $\sim \Delta_+(x - x')$. We evaluate it as,

$$i\Delta(x, x') = \sum_n \langle 0 | \Phi(x) | n \rangle \langle n | \Phi(x') | 0 \rangle - (x \leftrightarrow x'), \quad (10.8)$$

$$= \sum_n \langle 0 | \Phi(0) e^{ix \cdot P} | n \rangle \langle n | e^{-ix' \cdot P} \Phi(0) | 0 \rangle - (x \leftrightarrow x'), \quad (10.9)$$

$$= \sum_n |\langle 0 | \Phi(0) | n \rangle|^2 \left(e^{i(x-x') \cdot k_n} - e^{-i(x-x') \cdot k_n} \right) \quad \text{a function of } (x - x'). \quad (10.10)$$

The states $|n\rangle$ contain the vacuum, the single particle states on the isolated hyperboloid $k_n^2 = -m^2$ as well as the multi-particles continuum. To collect terms with the same total k_n^μ , insert $1 = \int d^4q \delta^4(q - k_n)$ and write,

$$\begin{aligned} \Delta'(x - x') &= -\frac{i}{(2\pi)^3} \int d^4q \left[(2\pi)^3 \sum_n \delta^4(q - k_n) |\langle 0 | \Phi(0) | n \rangle|^2 \left(e^{i(x-x') \cdot k_n} - e^{-i(x-x') \cdot k_n} \right) \right] \\ &:= -\frac{i}{(2\pi)^3} \int d^4q \left(e^{i(x-x') \cdot q} - e^{-i(x-x') \cdot q} \right) \rho(q) \quad \text{where,} \\ \rho(q) &:= \left[(2\pi)^3 \sum_n \delta^4(q - k_n) |\langle 0 | \Phi(0) | n \rangle|^2 \right] \quad (\text{the spectral density}). \end{aligned}$$

Claim: The spectral density is a scalar function of q^2 .

Proof: Under a Lorentz transformation, $U(\Lambda)$, we have $U(\Lambda)|0\rangle = |0\rangle$ and $U(\Lambda)\Phi(0)U^{-1}(\Lambda) = \Phi(\Lambda 0) = \Phi(0)$. Hence,

$$\rho(q) = (2\pi)^3 \sum_n \delta^4(q - k_n) |\langle 0 | \Phi(0) U(\Lambda) | n \rangle|^2.$$

Next,

$$\begin{aligned} \delta^4(k) &\sim \int d^4x e^{ik \cdot x} = \int d^4(\Lambda x) e^{ik \cdot (\Lambda x)} = \int d^4x e^{i(\Lambda^{-1}k) \cdot x} \sim \delta^4(\Lambda^{-1}k). \\ \therefore \rho(q) &= (2\pi)^3 \sum_n \delta^4(\Lambda^{-1}(q - k_n)) |\langle 0 | \Phi(0) U^{-1}(\Lambda) | n \rangle|^2. \end{aligned}$$

Let $U(\Lambda)|n\rangle = |m\rangle$. Then $U^{-1}(\Lambda)P^\mu U(\Lambda) = \Lambda^\mu_\nu P^\nu$. Acting on $|n\rangle$ implies, $P^\mu |m\rangle = k_m^\mu |m\rangle$ and we already had $P^\mu |n\rangle = k_n^\mu |n\rangle$. This in turn implies, $k_m^\mu U^{-1}(\Lambda) |m\rangle = \Lambda^\mu_\nu k_n^\nu |n\rangle$ or, $k_m^\mu = \Lambda^\mu_\nu k_n^\nu$. Lowering the Lorentz index gives, $(k_m)_\mu = (k_n)_\nu (\Lambda^{-1})^\nu_\mu$. Replacing the

sum over n by that over m gives,

$$\rho(q) = (2\pi)^3 \sum_m \delta^4(k_m - \Lambda^{-1}q) |\langle 0|\Phi(0)|m\rangle|^2 = \rho(\Lambda^{-1}q) .$$

Thus, $\rho(q)$ can only depend on q through q^2 .

Since the physical spectrum has $q^2 < 0$ with $q^0 > 0$, we write $\rho(q) := \theta(q^0)\rho(q^2)$, $\rho(q^2) = 0$ for $q^2 > 0$. Substituting in Δ' gives,

$$\begin{aligned} \Delta'(x-x') &= \frac{-i}{(2\pi)^3} \int d^4q \rho(q^2) \theta(q^0) \left(e^{iq \cdot (x-x')} - e^{-iq \cdot (x-x')} \right) \\ &= \int_0^\infty d\sigma^2 \rho(\sigma^2) \left[\frac{-i}{(2\pi)^3} \int d^4q \delta(q^2 - \sigma^2) \epsilon(q^0) e^{iq \cdot (x-x')} \right] . \\ \therefore \Delta'(x-x') &= \int_0^\infty d\sigma^2 \rho(\sigma^2) \Delta(x-x', \sigma^2) . \end{aligned} \quad (10.11)$$

Here we have introduced the function $\epsilon(x) = \theta(x) - \theta(-x)$ to combine the two exponentials and also recognized that the the square bracket is just $-i[\Phi_{in}(x), \Phi_{in}(x')]$ for $m^2 = \sigma^2$. The last equation is known as the *Kallen-Lehmann representation* or the *spectral representation* for the commutator function $\Delta'(x-x')$ of the exact interacting quantum fields. This has followed from Poincare covariance (separation of the x -dependence), the assumptions regarding the spectrum of the interacting system ($\rho(q^2 > 0) = 0$) and normalizations used for Φ_{in} in identifying the $\Delta(x-x', \sigma^2)$.

We can separate the contribution of the 1-particle states at $\sigma^2 = m^2$. We obtain the $\langle 0|\Phi(x)|k\rangle$ using the asymptotic condition,

$$\begin{aligned} \langle 0|\Phi(x)|k\rangle &= \sqrt{Z} \langle 0|\Phi_{in}(x)|k\rangle + \int d^4y G_{ret}(x-y, m) \langle 0|\tilde{J}(y)|k\rangle , \text{ but} \\ \langle 0|\tilde{J}(y)|k\rangle &= \langle 0|(\square - m^2)\Phi(x)|k\rangle = (\square - m^2) (\langle 0|\Phi(0)|k\rangle e^{ik \cdot x}) \\ &= (-k^2 - m^2) (\langle 0|\Phi(0)|k\rangle e^{ik \cdot x}) = 0. \text{ The first term then gives,} \\ \langle 0|\Phi(x)|k\rangle &= \sqrt{Z} \langle 0|\Phi_{in}(x)|k\rangle = \sqrt{Z} \frac{e^{ik \cdot x}}{\sqrt{2\omega_{\vec{k}}(2\pi)^3}} \Rightarrow |\langle 0|\Phi(x)|k\rangle|^2 = \frac{Z}{2\omega_{\vec{k}}(2\pi)^3} . \\ \therefore \rho(q^2)|_{1\text{-particle}} &= (2\pi)^3 \int d^3k \delta^4(k-q) \frac{Z}{2\omega_{\vec{k}}(2\pi)^3} = Z\delta(q^2 - m^2)\theta(q^0) \end{aligned} \quad (10.12)$$

$$\therefore \boxed{\Delta'(x-x') = Z\Delta(x-x', m^2) + \int_{m_1^2}^\infty d\sigma^2 \rho(\sigma^2) \Delta(x-x', \sigma^2)} . \quad (10.13)$$

The lower limit on the integral is the smallest invariant mass of the multi-particle continuum and $m_1^2 > m^2$ is assumed.

We can now derive a bound on Z . Observe that

$$\lim_{t' \rightarrow t} (i\partial_t \Delta'(x - x')) = \langle 0 | [\dot{\Phi}(x), \Phi(x')] | 0 \rangle = -i\delta^3(x - x') = (i\partial_t \Delta(x - x', m^2)) .$$

We are using the assumption that there are no derivative terms in $J(x)$ or equivalently, $\Pi(x) = \dot{\Phi}(x)$ and noting that the in/out fields satisfy the same commutation relations.

Taking $-i\partial_t$ on eq.(10.13) and taking the limit $t' \rightarrow t$, we deduce

$$\begin{aligned} -i\delta^3(\vec{x} - \vec{x}') &= Z(-i\delta^3(\vec{x} - \vec{x}')) + \int_{m_1^2}^{\infty} d\sigma^2 \rho(\sigma^2) [-i\delta^3(\vec{x} - \vec{x}', \sigma)] \\ \text{Or, } 1 &= Z + \int_{m_1^2}^{\infty} d\sigma^2 \rho(\sigma^2) \implies \boxed{0 \leq Z < 1.} \end{aligned} \quad (10.14)$$

The inequalities on Z can be understood as follows. Since asymptotic condition is needed to link $\Phi(x)$ with $\Phi_{in/out}(x)$, $Z \neq 0$. To understand the upper limit, notice that the free fields can only produce 1-particles states since it is linear in creation/annihilation operators. The interacting field has no such restriction and can produce multi-particle states as well. Hence the probability for it to produce 1-particle states is *less* than that for the free fields i.e. $|\langle 0 | \Phi(x) | k \rangle|^2 < |\langle 0 | \Phi_{in/out}(x) | k \rangle|^2$. This results in $Z < 1$.

In principle the Z factors for the in and out fields could have been different in the asymptotic conditions. However, following identical steps as above would lead us to the same relation (10.13) implying equality of the Z 's.

An important point to note is that $Z < 1$ has a physical reason, quite independent of any infinities and their renormalization that we will encounter later on.

Remark: In the Kallen-Lehmann spectral representation we considered the vacuum expectation value of the commutator of the fully interacting quantum field presumed to satisfy the usual canonical commutation relations. Because of this, the interacting quantum field gets identified with the so called ‘‘bare’’ quantum field in the renormalized perturbation theory. We could instead use a ‘‘renormalized’’ quantum field, $\Phi_R(x) := \Phi(x)/\sqrt{Z}$ and correspondingly define, $\Delta'_R(x - x') := Z^{-1}\Delta'(x - x')$. It follows that $\rho_R(q^2) = Z^{-1}\rho(q^2)$ and $[\Phi_R(x), \dot{\Phi}_R(x')] = iZ^{-1}\delta^3(x - x')$. In terms of the renormalized quantities, there is no factor of Z in equation (10.13). The manipulations leading to the bound on Z will now give,

$$\lim_{t' \rightarrow t} (i\partial_t \Delta'_R(x - x')) = \langle 0 | [\dot{\Phi}_R(x), \Phi_R(x')] | 0 \rangle = -iZ^{-1}\delta^3(x - x') = Z^{-1}(i\partial_t \Delta(x - x', m^2)) ,$$

and the eq.(10.13) will lead to $1 = Z [1 + \int_0^{\infty} d\sigma^2 \rho_R(\sigma^2)]$ implying again that $Z < 1$. These bounds are true provided the spectral integral is *finite*. In perturbation theory, the two point

functions (commutator or the Feynman propagator) of the unrenormalized fields are Ultra-Violet divergent and the bound cannot be inferred (See section 10.7 of [1]). The Kallen-Lehmann representation eq.(10.11) itself does *not* depend on the canonical commutation relations for the interacting field and can indeed be derived for composite fields as well.

D. The S -matrix and its properties

Having the $\Phi_{in/out}(x)$ and their corresponding mode expansions at our disposal, we can now define the in/out states as, $|k\rangle_{in} := a_{in}^\dagger(\vec{k})|0\rangle$ and $|k\rangle_{out} := a_{out}^\dagger(\vec{k})|0\rangle$. Similarly, multi-particle states can be defined. We can form wave packets for normalizable states and take limit of infinite sharpness at the end. We bypass these steps and work with the momentum eigenstates. Note that these states are *time independent*. While we can certainly define the general in/out states and take their spans, we need to (and do) make a further assumption that these spans generate the full Hilbert space of the interacting quantum fields¹⁰. The sets of in and out states just constitute two orthonormal bases, conveniently transforming by representations of the Poincare group. This guarantees existence of a unitary operator connecting the two orthonormal bases. This is our scattering operator in this Heisenberg picture formulation. Taking the basis elements to be generated by *monomials of $a_{in}^\dagger(\vec{k}, \sigma, \dots)$ and $a_{out}^\dagger(\vec{k}, \sigma, \dots)$* , the S -matrix is defined as,

$$S_{\beta\alpha} := \langle \beta \text{ out} | \alpha \text{ in} \rangle =: \langle \beta \text{ in} | S | \alpha \text{ in} \rangle \quad (10.15)$$

Note that α, β label *basis states* which are created by *monomials* in $a_{in}^\dagger, a_{out}^\dagger$. The definition of S operator shows that it preserves the labels: $\langle \beta \text{ out} | =: \langle \beta \text{ in} | S$. We stipulate some conditions on the S -matrix.

Stability of the vacuum and 1-particle states: We expect vacuum state to suffer no scattering spontaneously except may be acquiring a phase (uniqueness of vacuum allows a phase since rays are defined unto phases). We choose the phase to be 1, and stipulate $\boxed{S_{00} = 1}$. Likewise presumed stability of single particle states also disallows any spontaneous change (nothing to scatter against). Thus we stipulate that $S_{k',k} = \delta^3(\vec{k}' - \vec{k})$. Scattering takes place with multi-particle states.

¹⁰ This automatically precludes bound states in the interacting system and amounts to postulating asymptotic completeness property mentioned earlier.

Claim: $\Phi_{out}(x) = S^{-1}\Phi_{in}(x)S$.

Proof: Consider, $\langle \beta out | \Phi_{out}(x) | \alpha in \rangle$ and evaluate in two ways.

$$\langle \beta out | \Phi_{out}(x) | \alpha in \rangle = (\langle \beta in | S) \Phi_{out}(x) | \alpha in \rangle.$$

Next, $\langle \beta' out | := \langle \beta out | \Phi_{out}(x)$ is a linear combination of the out-basis elements. Each of this can be expressed as corresponding in-basis element times S . And the in-basis element is also obtained by action of $\Phi_{in}(x)$. In equation,

$$\begin{aligned} \langle \beta' out | &= \sum_{\gamma} C_{\gamma} \langle \gamma out | = \sum_{\gamma} C_{\gamma} (\langle \gamma in | S) = \\ & \left(\sum_{\gamma} C_{\gamma} \langle \gamma in | \right) S =: \langle \beta' in | S = (\langle \beta in | \Phi_{in}(x)) S. \end{aligned}$$

Taking inner product of the two expressions with $|\alpha in \rangle$ gives,

$$\langle \beta out | \Phi_{out}(x) | \alpha in \rangle = \langle \beta in | S \Phi_{out}(x) | \alpha in \rangle = \langle \beta in | \Phi_{in}(x) S | \alpha in \rangle \quad \forall \alpha, \beta.$$

Completeness of the in-basis implies, $\boxed{\Phi_{out}(x) = S^{-1}\Phi_{in}(x)S}$ proving the claim.

Claim: *Covariance of the in/out fields gives invariance of the scattering operator.*

Proof: We have,

$$\begin{aligned} U(\Lambda, a) \Phi_{in}(x) U^{-1}(\Lambda, a) &= \Phi_{in}(\Lambda x + a) \\ U(\Lambda, a) \Phi_{out}(x) U^{-1}(\Lambda, a) &= \Phi_{out}(\Lambda x + a) \quad \text{and} \quad \Phi_{out}(x) = S^{-1} \Phi_{in}(x) S \\ \therefore U(S^{-1} \Phi_{in}(x) S) U^{-1} &= S^{-1} \Phi_{in}(\Lambda x + a) S = S^{-1} U \Phi_{in}(x) U^{-1} S \quad (10.16) \\ \therefore U^{-1} S U S^{-1} \Phi_{in}(x) &= \Phi_{in}(x) U^{-1} S U S^{-1} \Rightarrow \end{aligned}$$

$$U^{-1} S U S^{-1} = \mathbb{1} \text{ or } \boxed{S U(\Lambda, a) = U(\Lambda, a) S} . \quad (10.17)$$

Question: Are the Schrodinger picture definitions of the scattering matrix element (10.5) and the Heisenberg picture scattering matrix element defined using $\Phi_{out} = S^{-1}\Phi_{in}S$, related? If so, how?

We now relate the S-matrix elements to vacuum expectation values of time ordered products of the interacting fields - the Lehmann-Symanzik-Zimmermann (LSZ) reduction formulae. This is followed by covariant perturbation series leading to Feynman rules. We will first detail these steps for the notationally simpler case of a scalar field and then summarize the corresponding steps for the Dirac and the Maxwell fields.

E. Lehmann-Symanzik-Zimmermann Reduction of S -matrix

We have defined the S -matrix elements using the in and out states which form the basis elements created by monomials in $a_{in}^\dagger(\vec{k}, \dots)$ and $a_{out}^\dagger(\vec{k}, \dots)$ acting on the unique vacuum. The in and out fields are not independent or uncorrelated, they are linked through the interacting field $\Phi(x)$ via the asymptotic condition. The LSZ reduction of the S -matrix elements expresses them in terms of the interacting field explicitly. For simplicity, let us first consider the case of an interacting Hermitian scalar field.

1. LSZ reduction for Klein-Gordon field

Consider a matrix element of the form $\langle \beta \text{ out} | \alpha, k \text{ in} \rangle$ where we have separated the k label in the in-state. The definition gives,

$$\langle \beta \text{ out} | \alpha, k \text{ in} \rangle = \langle \beta \text{ out} | a_{in}^\dagger(\vec{k}) | \alpha \text{ in} \rangle = \langle \beta \text{ out} | \left(a_{in}^\dagger(\vec{k}) - a_{out}^\dagger(\vec{k}) + a_{out}^\dagger(\vec{k}) \right) | \alpha \text{ in} \rangle .$$

The $a_{out}^\dagger(\vec{k})$ acting on the out-state will remove a particle with label \vec{k} if it is present in the label set β , or will annihilate the out-state. For simplicity, let us assume that there is no such state in the β label. Then we have just added and subtracted a 0. For the first two term in the brackets, use the inversion formula (10.7).

$$\begin{aligned} \langle \beta \text{ out} | \alpha, k \text{ in} \rangle &= \langle \beta \text{ out} | \left[\frac{-i}{\sqrt{2\omega_{\vec{k}}}(2\pi)^3} \int d^3x e^{ik \cdot x} \overleftrightarrow{\partial}_0 \Phi_{in}(x) \right. \\ &\quad \left. + \frac{+i}{\sqrt{2\omega_{\vec{k}}}(2\pi)^3} \int d^3x e^{ik \cdot x} \overleftrightarrow{\partial}_0 \Phi_{out}(x) \right] | \alpha \text{ in} \rangle . \\ [\dots] &= \left(\lim_{t \rightarrow +\infty} - \lim_{t \rightarrow -\infty} \right) \left\{ \frac{+i}{\sqrt{2\omega_{\vec{k}}}(2\pi)^3} \int d^3x \frac{e^{ik \cdot x} \overleftrightarrow{\partial}_0 \Phi(x)}{\sqrt{Z}} \right\} \\ \therefore \sqrt{Z} [\dots] &= \int d^4x \partial_0 \left\{ e^{ik \cdot x} \overleftrightarrow{\partial}_0 \Phi(x) \right\} \\ &= \int d^4x \left\{ e^{ik \cdot x} \partial_0^2 \Phi(x) - \partial_0^2 (e^{ik \cdot x}) \Phi(x) \right\} , \quad (-\partial_0^2 e^{ik \cdot x} = (-\nabla^2 + m^2) e^{ik \cdot x}) \\ &= \int d^4x e^{ik \cdot x} \left\{ \partial_0^2 \Phi(x) + (-\nabla^2 + m^2) \Phi(x) \right\} , \quad (\nabla^2 \text{ flipped onto } \Phi(x)). \\ \therefore \langle \beta \text{ out} | \alpha, k \text{ in} \rangle &= \frac{-i}{\sqrt{Z}} \int \frac{d^4x e^{ik \cdot x}}{\sqrt{2\omega_{\vec{k}}}(2\pi)^3} (\square - m^2) \langle \beta \text{ out} | \Phi(x) | \alpha \text{ in} \rangle . \end{aligned}$$

Now let us separate a particle with label k' from the out-state and write $\langle \beta \text{ out} | =$

$\langle \gamma, k' \text{ out} |$ the inner product on the right hand side of the above equation. Consider,

$$\begin{aligned} \langle \gamma, k' \text{ out} | \Phi(x) | \alpha \text{ in} \rangle &= \langle \gamma \text{ out} | a_{\text{out}}(\vec{k}') \Phi(x) | \alpha \text{ in} \rangle \\ &= \langle \gamma \text{ out} | (a_{\text{out}}(\vec{k}') \Phi(x) - \Phi(x) a_{\text{in}}(\vec{k}') + \Phi(x) a_{\text{in}}(\vec{k}')) | \alpha \text{ in} \rangle \end{aligned}$$

As before, for simplicity, assume α -label does not contain k' and thus drop the third term in the brackets. Notice that we have added and subtracted the zero term with $a_{\text{in}}(\vec{k}')$ to the right of $\Phi(x)$ while $a_{\text{out}}(\vec{k}')$ is naturally to the left of $\Phi(x)$. We need to maintain this order as we do not have the commutation relations between the in/out field and the interacting field. As before, using the inversion formula (10.7) and using the asymptotic condition, we get

$$\begin{aligned} \langle \gamma, k' \text{ out} | \Phi(x) | \alpha \text{ in} \rangle &= \langle \gamma \text{ out} | \left[\frac{i}{\sqrt{2\omega_{\vec{k}'}(2\pi)^3}} \int d^3x' e^{-ik' \cdot x'} \left\{ \overleftrightarrow{\partial}_0 \Phi(x')_{\text{out}} \Phi(x) \right. \right. \\ &\quad \left. \left. - \Phi(x) \overleftrightarrow{\partial}_0 \Phi(x')_{\text{in}} \right\} \right] | \alpha \text{ in} \rangle \\ \therefore \sqrt{Z} [\dots] &= \frac{i}{\sqrt{2\omega_{\vec{k}'}(2\pi)^3}} \int d^3x' \left\{ \lim_{t' \rightarrow \infty} (e^{-ik' \cdot x'} \overleftrightarrow{\partial}_0 \Phi(x')) \Phi(x) \right. \\ &\quad \left. - \lim_{t' \rightarrow -\infty} \Phi(x) (e^{-ik' \cdot x'} \overleftrightarrow{\partial}_0 \Phi(x')) \right\} \end{aligned}$$

It is convenient to define a time ordering instruction. Define a *time ordered product* as:

$$\boxed{T \{A(t_1, \vec{x}) B(t_2, \vec{y})\} := \theta(t_1 - t_2) A(t_1, \vec{x}) B(t_2, \vec{y}) + \theta(t_2 - t_1) B(t_2, \vec{y}) A(t_1, \vec{x})}.$$

Notice that the two limits already have the time ordering incorporated. Hence we can combine the terms and write,

$$\begin{aligned} \sqrt{Z} [\dots] &= \frac{i}{\sqrt{2\omega_{\vec{k}'}(2\pi)^3}} \int d^3x' \left(\lim_{t' \rightarrow \infty} - \lim_{t' \rightarrow -\infty} \right) e^{-ik' \cdot x'} \overleftrightarrow{\partial}_0 T \{ \Phi(x') \Phi(x) \} \\ &= \frac{i}{\sqrt{2\omega_{\vec{k}'}(2\pi)^3}} \int d^4x' \partial'_0 \left(e^{-ik' \cdot x'} \overleftrightarrow{\partial}_0 T \{ \Phi(x') \Phi(x) \} \right) \end{aligned}$$

Proceeding exactly as before we get,

$$\begin{aligned} \langle \gamma, k' \text{ out} | \alpha, k \text{ in} \rangle &= \left(\frac{-i}{\sqrt{Z}} \right)^2 \int d^4x' \int d^4x \frac{e^{-ik' \cdot x'}}{\sqrt{2\omega_{\vec{k}'}(2\pi)^3}} \frac{e^{ik \cdot x}}{\sqrt{2\omega_{\vec{k}}(2\pi)^3}} \\ &\quad (\square_{x'} - m^2)(\square_x - m^2) \langle \gamma, k' \text{ out} | T \{ \Phi(x') \Phi(x) \} | \alpha, k \text{ in} \rangle \quad (10.18) \end{aligned}$$

The generalization is immediate and we state the master formula for the S -matrix element

as,

$$\begin{aligned}
\langle k_1, \dots, k_n \text{ out} | k_1, \dots, k_m \text{ in} \rangle &= \left(\frac{-i}{\sqrt{Z}} \right)^{m+n} \int d^4 x'_1 \dots d^4 x'_n \int d^4 x_1 \dots d^4 x_m \\
&\prod_j^n \frac{e^{-ik'_j \cdot x'_j}}{\sqrt{2\omega_{\vec{k}'_j} (2\pi)^3}} \prod_j^m \frac{e^{+ik_j \cdot x_j}}{\sqrt{2\omega_{\vec{k}_j} (2\pi)^3}} (\square_{x'_1} - m^2) \\
&\dots (\square_{x'_m} - m^2) (\square_{x_1} - m^2) \dots (\square_{x_n} - m^2) \\
&\langle 0 | T \{ \Phi(x'_1) \dots \Phi(x'_n) \Phi(x_1) \dots \Phi(x_n) \} | 0 \rangle \quad (10.19)
\end{aligned}$$

The vacuum expectation value of the time ordered product of n quantum fields (interacting or free) is called an n -point function/Green function/ n -point correlation function.

To get the S -matrix elements, the n -point function is operated by the free field equation expression, multiplied by the in/out mode functions and integrated over all the space-time points. Note that the time ordering arises because the definition of S -matrix requires all the creation operators, $a_{in}^\dagger(\vec{k})$, have to be on the right while the annihilation operators, $a_{out}(\vec{k})$ have to be on the left. This will also naturally lead to the Feynman Green's function (free, 2-point function).

Let us note the steps that were followed to get the S -matrix elements in terms of the n -point functions.

- Mode decomposition of the free field and the inversion formulae;
- Equation of motion for the interacting field with a 'source' term on the right hand side;
- Equal time(anti-)commutation relations for both the interacting and the in/out fields;
- Asymptotic conditions;
- Reduction process;

2. LSZ reduction for Dirac field

We begin by recalling the mode decomposition.

$$\begin{aligned}
\Psi(x) &= \int \frac{d^3 k}{\sqrt{2\omega_{\vec{k}} (\pi)^3}} \left[b(\vec{k}, \sigma) u(\vec{k}, \sigma) e^{ik \cdot x} + d^\dagger(\vec{k}, \sigma) v(\vec{k}, \sigma) e^{-ik \cdot x} \right] \\
\Psi^\dagger(x) &= \int \frac{d^3 k}{\sqrt{2\omega_{\vec{k}} (\pi)^3}} \left[b^\dagger(\vec{k}, \sigma) u^\dagger(\vec{k}, \sigma) e^{-ik \cdot x} + d(\vec{k}, \sigma) v^\dagger(\vec{k}, \sigma) e^{ik \cdot x} \right]
\end{aligned}$$

We have used the Ψ^\dagger field instead of $\bar{\Psi}$ because the inversion formulae take a more convenient form.

We had defined the u, v spinors as,

$$(k+m)u(\vec{k}, \sigma) = 0, \quad \bar{u}(\vec{k}, \sigma)u(\vec{k}, \sigma') = \delta_{\sigma, \sigma'} = -\bar{v}(\vec{k}, \sigma)v(\vec{k}, \sigma'), \quad (-k+m)v(\vec{k}, \sigma) = 0$$

$$\sum_{\sigma} u(\vec{k}, \sigma)\bar{u}(\vec{k}, \sigma) = \frac{-k+m}{2m}, \quad \sum_{\sigma} v(\vec{k}, \sigma)\bar{v}(\vec{k}, \sigma) = -\frac{k+m}{2m}$$

We need to express these in terms of the adjoints rather than in terms of the Dirac adjoints. This alternative form can be derived as follows. We have,

$$\bar{u}(\vec{k}, \sigma)\gamma^\mu u(\vec{k}, \sigma') = \Lambda_{\nu}^{\mu} \bar{u}(\hat{k}, \sigma)\gamma^\mu u(\hat{k}, \sigma'), \quad k^\mu = \Lambda_{\nu}^{\mu} \hat{k}^\nu, \quad \hat{k}^\mu = (m, \vec{0}). \quad \text{For } \mu = 0,$$

$$u^\dagger(\vec{k}, \sigma)u(\vec{k}, \sigma') = \Lambda_0^0 u^\dagger(\hat{k}, \sigma)\gamma^0 \gamma^0 u(\hat{k}, \sigma') = \Lambda_0^0 u^\dagger(\hat{k}, \sigma)\gamma^0 u(\hat{k}, \sigma') \quad \because \gamma^0 u(\hat{k}, \sigma') = u(\hat{k}, \sigma')$$

$$= \Lambda_0^0 u^\dagger(\hat{k}, \sigma)\gamma^0 u(\hat{k}, \sigma') + \Lambda_0^i u^\dagger(\hat{k}, \sigma)\gamma^i u(\hat{k}, \sigma')$$

$$\therefore u^\dagger(\vec{k}, \sigma)u(\vec{k}, \sigma') = \Lambda_0^0 \bar{u}(\hat{k}, \sigma)u(\hat{k}, \sigma') + 0 = \Lambda_0^0 \delta_{\sigma, \sigma'} + 0$$

The second term in the last equation is zero because, for \hat{k} spinors $\gamma^0 u(\hat{k}, \sigma) = u(\hat{k}, \sigma)$. This implies,

$$u^\dagger \gamma^i u = u^\dagger \gamma^0 \gamma^i u = -u^\dagger \gamma^i \gamma^0 u = -u^\dagger \gamma^i u.$$

Similarly, $v^\dagger(\vec{k}, \sigma)v(\vec{k}, \sigma') = -\Lambda_0^0 \bar{v}(\hat{k}, \sigma)v(\hat{k}, \sigma') = +\Lambda_0^0 \delta_{\sigma, \sigma'}$. But Λ_0^0 is defined through, $k^\mu = \Lambda_{\nu}^{\mu} \hat{k}^\nu$. For $\mu = 0$ this gives, $\Lambda_0^0 = \frac{\omega_{\vec{k}}}{m}$. Our orthonormality relations then take the form,

$$\boxed{u^\dagger(\vec{k}, \sigma)u(\vec{k}, \sigma') = \frac{\omega_{\vec{k}}}{m} \delta_{\sigma, \sigma'} = v^\dagger(\vec{k}, \sigma)v(\vec{k}, \sigma'), \quad u^\dagger v = 0 = v^\dagger u.}$$

We can use these relations to derive the inversion formulae. For instance, multiplying the mode decomposition for $\Psi(x)$ by $u^\dagger(\vec{k}, \sigma)e^{-ik \cdot x}$ and integrating over d^3x yields,

$$b(\vec{k}, \sigma) = \frac{2m}{\sqrt{2\omega_{\vec{k}}(2\pi)^3}} \int d^3x u^\dagger(\vec{k}, \sigma)e^{-ik \cdot x} \Psi(x)$$

It is convenient to define

$$U(\vec{k}, \sigma) := 2m \frac{u(\vec{k}, \sigma)}{\sqrt{2\omega_{\vec{k}}(2\pi)^3}}, \quad V(\vec{k}, \sigma) := 2m \frac{v(\vec{k}, \sigma)}{\sqrt{2\omega_{\vec{k}}(2\pi)^3}}$$

The inversion formulae are,

$$b(\vec{k}, \sigma) = \int d^3x e^{-ik \cdot x} U^\dagger(\vec{k}, \sigma) \Psi(x) \quad (10.20)$$

$$d^\dagger(\vec{k}, \sigma) = \int d^3x e^{+ik \cdot x} V^\dagger(\vec{k}, \sigma) \Psi(x) \quad (10.21)$$

$$b^\dagger(\vec{k}, \sigma) = \int d^3x e^{+ik \cdot x} \Psi^\dagger(x) U(\vec{k}, \sigma) \quad (10.22)$$

$$d(\vec{k}, \sigma) = \int d^3x e^{-ik \cdot x} \Psi^\dagger(x) V(\vec{k}, \sigma) \quad (10.23)$$

In the reduction process, we will have for instance,

$$b_{in}^\dagger(\vec{k}, \sigma) - b_{out}^\dagger(\vec{k}, \sigma) = \int d^3x e^{ik \cdot x} \left(\Psi^\dagger(x)_{in} - \Psi^\dagger(x)_{out} \right) U(\vec{k}, \sigma)$$

$$\text{and using asymptotic condition} \longrightarrow \frac{1}{\sqrt{Z}} \int d^4x \partial_0 \left(e^{ik \cdot x} \Psi^\dagger(x) \right) U(\vec{k}, \sigma)$$

Next,

$$\partial_0 (e^{ik \cdot x} \Psi^\dagger(x)) U(\vec{k}, \sigma) = e^{ik \cdot x} (\partial_0 \Psi^\dagger + ik_0 \Psi^\dagger) U;$$

$$ik_0 \Psi^\dagger U = i\bar{\Psi} (k_0 \gamma^0) U = i\bar{\Psi} (\not{k} - k_i \gamma^i) U = i\bar{\Psi} (-mU) + \bar{\Psi} \gamma^i (-ik_i U)$$

$$\therefore e^{ik \cdot x} ik_0 \Psi^\dagger U = e^{ik \cdot x} (-i\bar{\Psi} mU) + \bar{\Psi} \gamma^i (-\partial_i e^{ik \cdot x}) U = e^{ik \cdot x} (-im\bar{\Psi} + \partial_i \bar{\Psi} \gamma^i) U$$

$$\therefore \partial_0 (e^{ik \cdot x} \Psi^\dagger) U(\vec{k}, \sigma) = e^{ik \cdot x} \{ \partial_0 \bar{\Psi} \gamma^0 + \partial_i \bar{\Psi} \gamma^i - im\bar{\Psi} \} U = (-ie^{ik \cdot x}) \left\{ i\bar{\Psi} \overleftarrow{\not{\partial}} + m\bar{\Psi} \right\} U$$

Hence, instead of $(-\square_x + m^2)$ acting on $\Phi(x)$, we will have $(-2mi)(i\overleftarrow{\not{\partial}} + m)$ acting on $\bar{\Psi}(x)$ and $(-2mi)(-i\overrightarrow{\not{\partial}} + m)$ acting on $\Psi(x)$. The wavefunction factors will be:

$$\text{Particle in in-state: } U(\vec{k}, \sigma) e^{ik \cdot x}, \text{ Particle in out-state: } \bar{U}(\vec{k}, \sigma) e^{-ik \cdot x} \quad (10.24)$$

$$\text{anti-Particle in in-state: } \bar{V}(\vec{k}, \sigma) e^{ik \cdot x}, \text{ anti-Particle in out-state: } V(\vec{k}, \sigma) e^{-ik \cdot x} \quad (10.25)$$

For multi-particle case we will again have time ordering instruction. For fermions, it is defined with an relative minus sign,

$$\boxed{T \{ \Psi_\alpha(x) \Psi_\beta(y) \} := \theta(x^0 - y^0) \Psi_\alpha(x) \Psi_\beta(y) - \theta(y^0 - x^0) \Psi_\beta(y) \Psi_\alpha(x)}$$

and $T \{ \Psi_\alpha(x) \Psi_\beta(y) \} = -T \{ \Psi_\beta(y) \Psi_\alpha(x) \}$. Thus for fermions, changing the ordering inside the time ordering generates a minus sign for each exchange. We summarize the general

matrix element for m fermions going into n fermions as,

$$\begin{aligned}
{}_{out}\langle \dots \vec{k}'_j \sigma'_{j'} \dots | \dots \vec{k}_j \sigma_j \dots \rangle_{in} &= \left(\frac{-i}{\sqrt{Z_\Psi}} \right)^{m+n} \prod_{j,j'} \int d^4 x'_j d^4 x_j U(\vec{k}_j, \sigma_j) \bar{U}(\vec{k}'_{j'}, \sigma'_{j'}) \\
&\quad e^{-i \sum_{j'} k'_{j'} \cdot x'_{j'}} (-i \overleftarrow{\partial}'_1 + m) \dots (-i \overleftarrow{\partial}'_n + m) \\
&\quad \langle 0 | T \{ \Psi(x'_1) \dots \Psi(x'_n) \bar{\Psi}(x_1) \dots \bar{\Psi}(x_m) \} | 0 \rangle \\
&\quad (-i \overleftarrow{\partial}'_1 + m) \dots (-i \overleftarrow{\partial}'_m + m) e^{+i \sum_j k_j \cdot x_j} \quad (10.26)
\end{aligned}$$

For anti-fermions in the in/out states, the spinors U, \bar{U} are changed to V, \bar{V} as appropriate.

3. LSZ reduction for Maxwell field

This is a bosonic field so there is no additional minus sign. Like the Dirac field, it has two polarizations and thanks to the zero mass, these are transverse polarizations. We begin by recalling the mode decomposition.

$$\begin{aligned}
A_\mu(x) &= \int \frac{d^3 k}{\sqrt{2\omega(2\pi)^3}} \sum_{\lambda=1,2} \left[a_{\vec{k},\lambda} \varepsilon_\mu(\vec{k}, \lambda) e^{ik \cdot x} + a_{\vec{k},\lambda}^* \varepsilon_\mu^*(\vec{k}, \lambda) e^{-ik \cdot x} \right] \\
\varepsilon_0(\vec{k}, \lambda) &:= -\frac{\vec{k} \cdot \vec{\varepsilon}(\vec{k}, \lambda)}{\omega_{\vec{k}}} \quad , \quad \varepsilon_i(\vec{k}, \lambda) := \left(\delta_i^j - \frac{k_i k^j}{k^2} \right) \varepsilon_j(\vec{k}, \lambda) \quad , \quad \omega_{\vec{k}} = |\vec{k}| \\
\vec{\varepsilon}(\vec{k}, \lambda) \cdot \vec{\varepsilon}^*(\vec{k}, \lambda') &= \delta_{\lambda,\lambda'} \quad , \quad \sum_\lambda \varepsilon_i(\vec{k}, \lambda) \varepsilon_j^*(\vec{k}, \lambda) = \left(\delta_i^j - \frac{k_i k^j}{k^2} \right)
\end{aligned}$$

This gives the inversion formulae as,

$$a(\vec{k}, \lambda) = i \varepsilon_i(\vec{k}, \lambda) \int d^3 x e^{-ik \cdot x} \overleftrightarrow{\partial}_0^i A^i(x) \quad , \quad a^\dagger(\vec{k}, \lambda) = -i \varepsilon_i^*(\vec{k}, \lambda) \int d^3 x e^{+ik \cdot x} \overleftrightarrow{\partial}_0^i A^i(x)$$

In the transverse/radiation/Coulomb gauge, $\varepsilon_\mu(\vec{k}, \lambda) = \varepsilon_\mu(\vec{k}, \lambda)$, $\vec{k} \cdot \vec{\varepsilon}(\vec{k}, \lambda) = 0$ and the equations of motion are $\square A_i(x) = 0$. The commutation relations are:

$$[A_i(t, \vec{x}), \pi_j(t, \vec{y})] = i \left(\delta_{ij} - \frac{\partial_i \partial_j}{\nabla^2} \right) := i \int \frac{d^3 k}{(2\pi)^3} e^{ik \cdot (x-y)} \left(\delta_{ij} - \frac{k_i k_j}{k^2} \right)$$

The reduction formula goes similar to the scalar. We will have $(-\square_x + m^2) \rightarrow -\square_x$ and the wave function factors would be $\mathcal{E}_\mu(\vec{k}, \lambda) := \frac{\varepsilon(\vec{k}, \lambda)}{\sqrt{2\omega_{\vec{k}}(2\pi)^3}}$. Then the general S -matrix

element for $m \rightarrow n$ particles takes the form,

$$\begin{aligned}
{}_{out}\langle \dots \vec{k}'_{j'} \lambda'_{j'} \dots | \dots \vec{k}_j \lambda \dots \rangle_{in} &= \left(\frac{-i}{\sqrt{Z_A}} \right)^{m+n} \prod_{j,j'} \int d^4 x'_{j'} d^4 x_j \mathcal{E}(\vec{k}_j, \lambda) \mathcal{E}^*(\vec{k}'_{j'}, \lambda'_{j'}) \\
&\quad e^{-i \sum_{j'} k'_{j'} \cdot x'_{j'}} (-\square'_1) \dots (-\square'_n) \\
&\quad \langle 0 | T \{ A_{\mu'_1}(x'_1) \dots A_{\mu'_n}(x'_n) A_{\mu_1}(x_1) \dots A_{\mu_m}(x_m) \} | 0 \rangle \\
&\quad (-\overleftarrow{\square}_1) \dots (-\overleftarrow{\square}_m) e^{+i \sum_j k_j \cdot x_j} \tag{10.27}
\end{aligned}$$

Having related the S -matrix elements to the n -point functions of the interacting fields, our next task is to try see if these can be expressed in terms of the free fields in some systematic, well defined way. This is achieved in the so-called covariant perturbation series.

11. COVARIANT PERTURBATION THEORY

In formulating the scattering theory for interacting fields, we postulated the in/out fields satisfying the free field equations. More importantly, we also postulated them to satisfy the same equal time (anti-)commutation relations. With some assumptions of existence, these suffice to develop a perturbation scheme to compute the n -point functions.

Since the interacting and the free (in/out) fields satisfy the same basic quantum conditions, it is plausible that they are related by some *unitary transformation*. For systems with finitely many degrees of freedom, it is theorem that guarantees existence of canonical transformations at the classical level and unitary transformations at the quantum level (thanks to the Stone-von-Neumann theorem). For field theories, there is no such guarantee and we need to postulate the required existence. Their utility gives a post-facto justification for the assumptions. Let,

$$\boxed{\Phi(t, \vec{x}) = U^{-1}(t)\Phi_{in}(t, \vec{x})U(t) \quad , \quad \Pi(t, \vec{x}) = U^{-1}(t)\Pi_{in}(t, \vec{x})U(t) \quad .}$$

The fields satisfy the equations of motion in the form,

$$\partial_t \Phi_{in}(x) = i[H_{in}(\Phi_{in}, \Pi_{in}), \Phi_{in}(x)] \quad , \quad \partial_t \Pi_{in}(x) = i[H_{in}(\Phi_{in}, \Pi_{in}), \Pi_{in}(x)] \quad (11.1)$$

$$\partial_t \Phi(x) = i[H(\Phi, \Pi), \Phi(x)] \quad , \quad \partial_t \Pi(x) = i[H(\Phi, \Pi), \Pi(x)] \quad (11.2)$$

These enable us to derive an equation for $U(t)$.

$$\begin{aligned} \partial_t \Phi_{in} &= \partial_t [U(t)\Phi U^{-1}(t)] = \partial_t U \cdot U^{-1} \Phi_{in} + U(\partial_t \Phi)U^{-1} + \Phi_{in} \cdot U \partial_t U^{-1} \\ &= [\partial_t U \cdot U^{-1}, \Phi_{in}] + iU[H(\Phi, \Pi), \Phi]U^{-1} \quad \text{But,} \end{aligned}$$

$$U[H(\Phi, \Pi), \Phi]U^{-1} = [UH(\Phi, \Pi)U^{-1}, U\Phi U^{-1}] = [H(\Phi_{in}, \Pi_{in}), \Phi_{in}]$$

$$\therefore \partial_t \Phi_{in} = [\partial_t U \cdot U^{-1} + i\{H(\Phi_{in}, \Pi_{in}) - H_{in}(\Phi_{in}, \Pi_{in}) + H_{in}(\Phi_{in}, \Pi_{in})\}, \Phi_{in}]$$

$$\partial_t \Phi_{in} = \partial_t \Phi_{in} + [\partial_t U \cdot U^{-1} + iH_I(\Phi_{in}, \Pi_{in}), \Phi_{in}] \quad \text{Similarly,}$$

$$\partial_t \Pi_{in} = \partial_t \Pi_{in} + [\partial_t U \cdot U^{-1} + iH_I(\Phi_{in}, \Pi_{in}), \Pi_{in}] \quad .$$

Since $\partial_t U \cdot U^{-1} + iH_I(\Phi_{in}, \Pi_{in})$ commutes with both Φ_{in}, Π_{in} , it must be a multiple of identity, say $E_0 \mathbb{1}$. This gives the equation determining the $U(t)$ as,

$$\boxed{i\partial_t U(t) = H'_I(\Phi_{in}, \Pi_{in})U(t)} \quad , \quad \boxed{H'_I(\Phi_{in}, \Pi_{in}) := H(\Phi_{in}, \Pi_{in}) - H_{in}(\Phi_{in}, \Pi_{in}) + E_0(t)} \quad (11.3)$$

To solve the equation, it is convenient to define the combination $U(t, t') := U(t)U^{-1}(t')$. It follows that this combination too satisfies the same first order equation with the initial condition, $U(t, t) = \mathbb{1}$. The integral form of this equation is:

$$U(t, t') = \mathbb{1} - i \int_{t'}^t dt'' H'_I(t'') U(t'', t') .$$

This is solved by iteration.

Define: $U_0(t, t') = \mathbb{1}$, and for $n \geq 1$, $U_n(t, t') := \mathbb{1} - i \int_{t'}^t dt_n H'(t_n) U_{n-1}(t_n, t')$. Then $U(t, t') = \sum_n U_n(t, t')$ is the formal solution of the integral equation.

It is trivial to verify the formal solution. It is formal because no conditions are imposed for convergence of the series. The first few terms are,

$$\begin{aligned} U_0(t, t') &= \mathbb{1} \\ U_1(t, t') &= \mathbb{1} - i \int_{t'}^t dt_1 H'(t_1) \mathbb{1} \\ U_2(t, t') &= \mathbb{1} - \int_{t'}^t dt_2 H'(t_2) U_1(t_2, t') \\ &= \mathbb{1} + (-i) \int_{t'}^t dt_2 H'(t_2) + (-i)^2 \int_{t'}^t dt_2 \int_{t'}^{t_2} dt_1 H'(t_2) H'(t_1) \\ U_3(t, t') &= \mathbb{1} + (-i) \int_{t'}^t dt_3 H'(t_3) \left\{ \mathbb{1} + (-i) \int_{t'}^t dt_2 H'(t_2) \right. \\ &\quad \left. + (-i)^2 \int_{t'}^t dt_2 \int_{t'}^{t_2} dt_1 H'(t_2) H'(t_1) \right\} \\ &= \mathbb{1} + (-i) \int_{t'}^t dt_3 H'(t_3) + (-i)^2 \int_{t'}^t dt_3 \int_{t'}^{t_3} dt_2 H'(t_3) H'(t_2) \\ &\quad + (-i)^3 \int_{t'}^t dt_3 \int_{t'}^{t_3} dt_2 \int_{t'}^{t_2} dt_1 H'(t_3) H'(t_2) H'(t_1) \\ \therefore U_n(t, t') &= \mathbb{1} + \sum_{k=1}^n (-i)^k \int_{t'}^t dt_k \int_{t'}^{t_k} dt_{k-1} \cdots \int_{t'}^{t_2} dt_1 H'(t_k) H'(t_{k-1}) \cdots H'(t_1) \quad (11.4) \end{aligned}$$

Consider the term $\int_{t'}^t dt_n \int_{t'}^{t_n} dt_{n-1}$. By interchanging the order of integration we can write

it as ($H'_n := H'(t_n)$),

$$\begin{aligned}
\int_{t'}^t dt_n \int_{t'}^{t_n} dt_{n-1} H'_n H'_{n-1} &= \int_{t'}^t dt_{n-1} \int_{t_{n-1}}^t dt_n H'_n H'_{n-1} = \int_{t'}^t dt_n \int_{t_n}^t dt_{n-1} H'_{n-1} H'_n \\
\therefore LHS &= \frac{1}{2}(LHS + RHS) = \frac{1}{2} \left[\int_{t'}^t dt_n \int_{t'}^{t_n} dt_{n-1} H'_n H'_{n-1} + \int_{t'}^t dt_n \int_{t_n}^t dt_{n-1} H'_{n-1} H'_n \right] \\
&= \frac{1}{2} \left[\int_{t'}^t dt_n \int_{t'}^{t_n} dt_{n-1} T \{ H'_n H'_{n-1} \} \right. \\
&\qquad\qquad\qquad \left. + \int_{t'}^t dt_n \int_{t_n}^t dt_{n-1} T \{ H'_{n-1} H'_n \} \right] \\
&= \frac{1}{2} \int_{t'}^t dt_n \int_{t'}^t dt_{n-1} T \{ H'_n H'_{n-1} \}
\end{aligned}$$

Similarly, we can symmetrize the higher order terms to replace each product by their time ordered products and then combine the integrals to have the same limits of integrations. The n^{th} order term then takes the form: $\frac{1}{n!} \int_{t'}^t dt_n \int_{t'}^{t_n} dt_{n-1} \cdots \int_{t'}^{t_1} dt_1 T \{ H'(t_n) H'(t_{n-1}) \cdots H'(t_1) \}$.

We write the iterated formal solution of the (11.3) as,

$$U(t, t') := T \left(\exp \left\{ -i \int_{t'}^t dt' H'_I(\Phi_{in}(t'), \Pi_{in}(t')) \right\} \right) \quad (11.5)$$

$$= \mathbb{1} + \sum_{n=1}^{\infty} \frac{(-i)^n}{n!} \int_{t'}^t dt_1 \cdots \int_{t'}^{t_n} dt_n T \{ H'_I(t_1) \cdots H'_I(t_n) \} \quad (11.6)$$

The solution is thus obtained entirely in terms of the ‘in’ fields. Note that the solution gives $U(t, t')$ and *not* $U(t)$. While the equations satisfied by $U(t)$ and $U(t, t')$ are the same, there is no “initial” condition provided for $U(t)$ and in a sense it is ill-defined. There is no canonical way to deduce/define $U(t)$ from $U(t, t')$. Fortunately, $U(t, t')$ suffice.

Note: From the definition it follows the useful composition property that $U(t, t') = U(t, t'')U(t'', t')$, regardless of any inequalities for t'' . This may also be verified directly.

Let us see an implication of the existence of $U(t)$. Consider an n -point function for a scalar, $G(x_1, \dots, x_n) = \langle 0 | T \{ \Phi(x_1) \cdots \Phi(x_n) \} | 0 \rangle$. Insert in this $\Phi(x) = U^{-1}(t)\Phi_{in}(x)U(t)$ which gives,

$$\begin{aligned}
\Phi(x_1)\Phi(x_2)\cdots\Phi(x_n) &= U^{-1}(t_1)\Phi_{in}(x_1)U(t_1) \cdot U^{-1}(t_2)\Phi_{in}(x_2)U(t_2) \cdots U^{-1}(t_n)\Phi_{in}(x_n)U(t_n) \\
&= U^{-1}(t) [U(t, t_1)\Phi_{in}(x_1)U(t_1, t_2)\Phi_{in}(x_2) \cdots \\
&\qquad\qquad\qquad U(t_{n-1}, t_n)\Phi_{in}(x_n)U(t_n, -t)] U(-t) \quad , \quad U(t, t') := U(t)U^{-1}(t')
\end{aligned}$$

Observe that in the limit $t \rightarrow \infty$, the factor $U^{-1}(t)$ goes to the extreme left and $U(t)$ to the extreme right. All other space-time arguments being finite, we can take these factors

outside the time ordering symbol. The product of the fields $\Phi(x)$'s is already under time ordering which then allows us to arrange all the factors within $T\{\dots\}$ to be conveniently grouped. In particular, all the $U(t_i, t_j)$ can be combined using their composition property noted above.

With the limit of $t \rightarrow \infty$ implicit, we can write the n -point function as,

$$G(x_1, \dots, x_n) = \langle 0|U^{-1}(t)T \left\{ \Phi_{in}(x_1) \dots \Phi_{in}(x_n) \exp \left(-i \int_{-t}^t dt' H_I'(t') \right) \right\} U(t)|0\rangle . \quad (11.7)$$

Next, we evaluate the $U(-t)|0\rangle$ in the limit $t \rightarrow \infty$ and its adjoint.

Claim: $U(-t)|0\rangle \rightarrow \lambda_-|0\rangle$ as $t \rightarrow \infty$.

Proof: Consider an in-state containing at least one particle of momentum $k, |\alpha, k\rangle_{in}$.

Then,

$$\begin{aligned} {}_{in}\langle \alpha, k|U(-t)|0\rangle &= {}_{in}\langle \alpha|a_{in}(\vec{k})U(-t)|0\rangle \\ &= i \int_{\Sigma_{t'}} \frac{d^3x}{\sqrt{2\omega_{\vec{k}}(2\pi)^3}} e^{-ik \cdot x} \overleftrightarrow{\partial}_0 {}_{in}\langle \alpha|\Phi_{in}(-t', \vec{x})U(-t)|0\rangle \end{aligned}$$

We have used the inversion formula for a_{in} by choosing an arbitrary $\Sigma_{t'}$ hyper-surface and a_{in} is of course independent of this.

In the integrand above, substitute $\Phi_{in}(t', \vec{x}) = U(t')\Phi(t', \vec{x})U^{-1}(t')$ and evaluate the time derivative.

$$\begin{aligned} \text{Integrand} &= e^{-ik \cdot x} \langle \alpha| \left[\dot{U}(-t')U^{-1}(-t')\Phi_{in}(-t', \vec{x})U(-t) + \Phi_{in}(-t', \vec{x})U(-t')\partial_{t'}U^{-1}(t')U(-t) \right. \\ &\quad \left. + U(-t')\partial'_0\Phi(-t')U^{-1}(-t')U(-t) - U(-t')\Phi(-t')U^{-1}(-t')U(-t)(\partial'_0(-ik \cdot x)) \right] |0\rangle \end{aligned}$$

Now take choose $t' = t$. Then, the first two terms on the r.h.s. combine as,

$$[\dot{U}(-t)U^{-1}(-t), \Phi_{in}(-t, \vec{x})]U(-t) = -i[H_I'(\Phi_{in}, \Pi_{in}), \Phi_{in}(-t, \vec{x})]U(-t) = 0 .$$

In the usual theories, the interaction Hamiltonian has no Π dependence and hence at equal time, commutes with the field giving 0.

In the last two terms, the $U^{-1}(-t')U(-t)$ factor becomes $\mathbb{1}$ for $t' = t$. These terms combine to give, ${}_{in}\langle \alpha|U(-t) \left(e^{-ik \cdot x} \overleftrightarrow{\partial}_0 \Phi(-t) \right) |0\rangle$. Taking the limit $t \rightarrow \infty$ and invoking the asymptotic condition, converts $\Phi(x)$ into $\sqrt{Z}\Phi_{in}(x)$. The spatial integration converts the expression into $\sqrt{Z}{}_{in}\langle \alpha|U(-t)a_{in}(\vec{k})|0\rangle = 0$.

Thus we conclude that ${}_{in}\langle \alpha, k|U(-t)|0\rangle \rightarrow 0$ as $t \rightarrow \infty$ for *all* states containing at least one particle. $U(-t)|0\rangle$ must therefore be proportional to $|0\rangle$ itself thereby proving the claim.

By similar reasoning, we obtain $U(t)|0\rangle \rightarrow \lambda_+|0\rangle$ as $t \rightarrow \infty$.

Corollary:

$$\lambda_+^* \lambda_- = \left[\langle 0|T \exp \left(-i \int_{-\infty}^{\infty} dt' H_I(t') \right) |0\rangle \right]^{-1} \quad (11.8)$$

The proof is simple,

$$\begin{aligned} \lambda_+^* \lambda_- &= \lim_{t \rightarrow \infty} \langle 0|U^{-1}(t)|0\rangle \langle 0|U(-t)|0\rangle \\ &\approx \langle 0|U^{-1}(t)|U(-t)|0\rangle \quad \text{since intermediate states do not contribute by above result} \\ &\approx \langle 0|U(-t)U^{-1}(t)|0\rangle = \langle 0|U(-t, t)|0\rangle \\ &\approx \langle 0|(U(t, -t))^{-1}|0\rangle = (\langle 0|U(t, -t)|0\rangle)^{-1}. \end{aligned}$$

The final expression follows by substituting the formal solution in the limit $t \rightarrow \infty$.

Using the equations (11.7, 11.8), we express the n -point function as,

$$G(x_1, \dots, x_n) = \frac{\langle 0|T \left\{ \Phi_{in}(x_1) \dots \Phi_{in}(x_n) \exp -i \int_{-\infty}^{\infty} dt' H_I(t') \right\} |0\rangle}{\langle 0|T \left\{ \exp -i \int_{-\infty}^{\infty} dt' H_I(t') \right\} |0\rangle} \quad (11.9)$$

Notice that the multiple of identity $E_0(t)\mathbb{1}$ has been canceled between the numerator and the denominator before taking the limit $t \rightarrow \infty$.

Suffice it to say that analogous expressions exist for other fields as well. However we will not write them explicitly.

The reduction formulae give S -matrix elements in terms of the n -point functions and the n -point functions have a perturbative expansion in terms of the free fields alone. These two together are the master formulae for computations.

A further simplification is provided by the *Wick's Theorem*.

A. Normal ordering and Wick's theorem

Recall that the mode expansion of the *free* quantum fields can be grouped as $\Phi(x) = \Phi_+(x) + \Phi_-(x)$ with $\Phi_{\pm}(x)$ denoting the sum over positive/negative frequency modes. The positive frequency part is made up of annihilation operators while the negative frequency part is made up of creation operators alone. For a Hermitian quantum field, the negative frequency part is the Hermitian adjoint of the positive frequency part. A product of fields at different points will mix these parts and the idea is to bring all annihilation operators to

the right and the creation operators to the left. The vacuum expectation value of so ordered groups will of course be zero. In this process, several commutators of fields at different space-time points are generated, but *they all are c-numbers*. This leads to a simplification.

We define *normal ordered products* of two *free* fields as:

$$\text{For bosons} \quad : \Phi_{-}(x)\Phi_{+}(y) : = \Phi_{-}(x)\Phi_{+}(y) \quad , \quad : a^{\dagger}a : := a^{\dagger}a \quad (11.10)$$

$$: \Phi_{+}(x)\Phi_{-}(y) : = \Phi_{-}(y)\Phi_{+}(x) \quad , \quad : aa^{\dagger} : := a^{\dagger}a \quad (11.11)$$

$$\text{For fermions} \quad : \Psi_{-}(x)\Psi_{+}(y) : = \Psi_{-}(x)\Psi_{+}(y) \quad , \quad : b^{\dagger}b : := b^{\dagger}b \quad (11.12)$$

$$: \Psi_{+}(x)\Psi_{-}(y) : = -\Psi_{-}(y)\Psi_{+}(x) \quad , \quad : bb^{\dagger} : := -b^{\dagger}b \quad (11.13)$$

For product of two or more positive (or negative) frequency fields, the normal ordering does not change order. It is immediately obvious that $\langle 0 | : Operator : | 0 \rangle = 0$.

Note that the normal ordering are meaningful only for products of free fields (in/out). Hence the in/out suffixes are suppressed for these.

To get a general form for several bosonic as well as fermionic free fields, notice that for a quantum field $Q(x) = Q_{+}(x) + Q_{-}(x)$ a product of n fields generates terms with the k number of Q_{+} 's interspersed with $(n - k)$ number of Q_{-} 's, with the order of the space-time points maintained, with $k = 0, \dots, n$. Under normal ordering each of these terms will shift the Q_{+} 's to the right and Q_{-} 's to the left, generating a permutation of the space-time points, say p . If Q is a fermionic field, the term with permutation p will get a factor of $\sigma_p = \text{sgn}(p)$ while a bosonic field will have $\sigma_p = 1$. Note that for any given k , there will be several terms eg Q_1Q_2, Q_3Q_7, \dots etc. Each will generate its own permutation under normal ordering. With this understood, we can write the general expression for normal ordered product of n -fields as,

$$: Q(x_1) \dots Q(x_n) : := \sum_{A,B} \sigma_p \prod_{i \in A} Q_{-}(x_{p(i)}) \prod_{j \in B} Q_{+}(x_{p(j)}) .$$

The A, B are two groups of space-time points corresponding to the $k, (n - k)$ mentioned above and the sum refers to various possible groupings of the space-time points within each class.

Exercise: Verify for $n = 4$.

Consider now the relation between time ordered products and normal ordered products. For a single field, both orderings are trivial and their vacuum expectation value vanishes: $\langle 0 | T \{ \phi_{in}(x) \} | 0 \rangle = 0 = \langle 0 | : \phi(x) : | 0 \rangle$.

For time ordered product of two fields, for any particular instances of time they appear in a particular order which can be explicitly put in the normal ordered form, eg. $a(\vec{k})a^\dagger(\vec{k}') = \mathbb{1} + a^\dagger(\vec{k}')a(\vec{k}) =: a(\vec{k})a^\dagger(\vec{k}') + \mathbb{1}$. The last term is a c-number (multiple of the identity operator). Similarly, $\Phi(x)\Phi(y) =: \Phi(x)\Phi(y) + c\text{-number}$. And the c-number is trivial to evaluate by just taking the vev (vacuum expectation value): c-number = $\langle 0|\Phi(x)\Phi(y)|0\rangle$. Noting that $T\{\Phi(x)\Phi(y)\} = \theta(x^0 - y^0)\Phi(x)\Phi(y) + \theta(y^0 - x^0)\Phi(y)\Phi(x)$, we get, $T\{\Phi(x)\Phi(y)\} =: T\{\Phi(x)\Phi(y)\} + \langle 0|T\{\Phi(x)\Phi(y)\}|0\rangle$. Notice that normal ordering and time ordering commute: $:T\{A(x)B(y)\} := T\{ :A(x)B(y) : \}$, for all non-trivial $A(x), B(y)$ operators i.e. for operators which are polynomials in the basic field operators of degree greater than zero. The generalization of this to product of arbitrary number of field is the Wick's theorem. We state it explicitly for a scalar field.

Wick's Theorem:

$$\begin{aligned}
T\{\Phi(x_1) \dots \Phi(x_n)\} &= :T\{\Phi(x_1) \dots \Phi(x_n)\} : \\
&+ [\langle 0|T\{\Phi(x_1)\Phi(x_2)\}|0\rangle :T\{\Phi(x_3) \dots \Phi(x_n)\} : + \text{permutations}] \\
&+ [\langle 0|T\{\Phi(x_1)\Phi(x_2)\}|0\rangle \langle 0|T\{\Phi(x_3)\Phi(x_4)\}|0\rangle \times \\
&\quad :T\{\Phi(x_5) \dots \Phi(x_n)\} : + \text{permutations}] \\
&\quad \vdots \\
&+ \left\{ \begin{array}{l} [\langle 0|T\{\Phi_1\Phi_2\}|0\rangle \dots \langle 0|T\{\Phi_{n-1}\Phi_n\}|0\rangle + \text{permutations}] \\ \hspace{15em} (n \text{ even}) \\ [\langle 0|T\{\Phi_1\Phi_2\}|0\rangle \dots \langle 0|T\{\Phi_{n-2}\Phi_{n-1}\}|0\rangle \Phi_n \\ \hspace{15em} + \text{permutations}] (n \text{ odd}) \end{array} \right.
\end{aligned}$$

In the last line we have used the abbreviation $\Phi_i := \Phi(x_i)$ and we will use the same when convenient.

Proof:

The proof is by induction and we have already seen the validity for $n = 1, 2$. Assume the theorem is true for $n = n$. Include an extra field $\Phi(x_{n+1})$ and choose t_{n+1} to be the earliest

instant so that it stays on the extreme right. Then,

$$\begin{aligned}
T\{\Phi_1 \dots \Phi_{n+1}\} &= T\{\Phi_1 \dots \Phi_n\}\Phi_{n+1} \\
&= \left[:T\{\Phi_1 \dots \Phi_n\}: + \sum_{perm} \langle 0|T\{\Phi_1\Phi_2\}|0\rangle :T\{\dots\}: + \dots \right] \Phi_{n+1}, \\
:\Phi_1 \dots \Phi_n : \Phi_{n+1} &= \left\{ \sum_{A,B} \sigma_p \prod_{i \in A} \Phi_i^- \prod_{j \in B} \Phi_j^+ \right\} (\Phi_{n+1}^+ + \Phi_{n+1}^-) \\
&= \sum_{A,B} \sigma_p \prod_{i \in A} \Phi_i^- \prod_{j \in B} \Phi_j^+ \Phi_{n+1}^+ + \\
&\quad \sum_{A,B} \sigma_p \prod_{i \in A} \Phi_i^- \left\{ \Phi_{n+1}^- \prod_{j \in B} \Phi_j^+ + \left[\prod_{j \in B} \Phi_j^+, \Phi_{n+1}^- \right] \right\}
\end{aligned}$$

The first two terms in the last equation have the requisite normal ordered form for $n + 1$ fields. The third term has several commutators, $[\Phi_j^+, \Phi_{n+1}^-]$ which again are c-numbers and can be evaluated by taking vev. That is,

$$\begin{aligned}
[\Phi_j^+, \Phi_{n+1}^-] &= \langle 0|(\Phi_j^+ \Phi_{n+1}^- - \Phi_{n+1}^- \Phi_j^+)|0\rangle = \langle 0|\Phi_j^+ \Phi_{n+1}^-|0\rangle + 0 \\
&= \langle 0|\Phi_j \Phi_{n+1}|0\rangle = \langle 0|T\{\Phi_j \Phi_{n+1}\}|0\rangle, \text{ since } t_{n+1} \text{ is the earliest.}
\end{aligned}$$

$$\therefore :\Phi_1 \dots \Phi_n : \Phi_{n+1} = :\Phi_1 \dots \Phi_n \Phi_{n+1}: + \sum_j \sigma_p :\Phi_1 \dots \Phi_{j-1} \Phi_{j+1} \dots \Phi_n : \langle 0|T\{\Phi_j \Phi_{n+1}\}|0\rangle$$

Thus, inserting the time ordering, the first term, $:T\{\Phi_1 \dots \Phi_n\} : \Phi_{n+1}$ gets expressed as $:T\{\Phi_1 \dots \Phi_n \Phi_{n+1}\} : + \{\text{terms which have a structure similar to the second term}\}$ and so on. Hence the induction hypothesis allows the expression to hold for $n + 1$ and the theorem is proved by induction.

Exercise: Verify the theorem for $n = 3, 4$.

Observe that $G(x_1, \dots, x_n) = \langle 0|T\{\Phi(x_1) \dots \Phi(x_n)\}|0\rangle$ vanishes for odd n and equals $\sum_{perm} \sigma_p \langle 0|T\{\Phi(x_1)\Phi(x_2)\}|0\rangle \dots \langle 0|T\{\Phi(x_{n-1})\Phi(x_n)\}|0\rangle$ for even n .

Thus, *2n-point function of free fields is the sum of products of n, 2-point functions of free fields. We denote:* $\langle 0|T\{\Phi(x)\Phi(y)\}|0\rangle =: i\Delta_F(x-y) =: (\text{Feynman propagator})$.

We have the S -matrix elements in terms of the n -point functions of interacting quantum fields, expressed in terms of vev of time ordered products of free fields and thanks to Wick's theorem these are expressed in terms of 2-point functions of free fields. In short, the S -matrix elements are obtained from a bunch of 2-point functions of free fields, together with some integrations and operations of $(\square - m^2)$. It remains to have a convenient algorithm for computations.

A few remarks are in order. The normal ordering is introduced here as a simplification step. It works because the postulated *unique* Poincare invariant vacuum state is annihilated by all (free field) annihilation operators. This also *requires* the Poincare generators of the free fields to be normal ordered¹¹. The Poincare generators of the interacting fields also need to annihilate the vacuum. Thanks to the assumed unitary operator linking the free and the interacting fields, these generators too are expressed in terms of the free field creation-annihilation operator. These expressions, in particular the Hamiltonian, must also be normal ordered for Poincare invariance. We will make this explicit for the interaction Hamiltonian $H_I(\Phi_{in}, \Pi_{in})$.

In the quantum optics context, the normal ordering was used since the detectors worked by absorbing a quantum from the field and in principle, for a detector working by emission of a quantum would require anti-normal ordering. Such an option is not available with manifest Poincare invariance *with a unique invariant vacuum state*.

An explicit example of evaluation of an n -point function will be useful. Consider $G_2(x_1, x_2)$ computed to second order in H_I . Consider the numerator first. *All fields are 'in' fields and the 'in' subscript is suppressed below.*

$$\begin{aligned} G_2(x_1, x_2)|_{Nr} &= \langle 0|T\{\Phi(x_1)\Phi(x_2)\left(\mathbb{1} - i\int_{-\infty}^{\infty} dt' H_I(t') \right. \\ &\quad \left. + \frac{(-i)^2}{2!}\int_{-\infty}^{\infty} dt_1 dt_2 T\{H_I(t_1)H_I(t_2)\}\right)\}|0\rangle \\ &= \langle 0|T\{\phi(x_1)\Phi(x_2)\}|0\rangle - i\int_{-\infty}^{\infty} dt_1 \langle 0|T\{\Phi(x_1)\Phi(x_2)H_I(t_1)\}|0\rangle \\ &\quad + \frac{(-i)^2}{2}\int_{-\infty}^{\infty} dt_1 dt_2 \langle 0|T\{\Phi(x_1)\Phi(x_2)H_I(t_1)H_I(t_2)\}|0\rangle \end{aligned}$$

Now let us take a specific interaction Hamiltonian, simplest non-trivial being $H_I(t) := g \int d^3y : \Phi(t, \vec{y})^3 :$. As noted above, for the case of Poincare invariant vacuum, the Poincare generators and hence in particular the Hamiltonian must be normal ordered. The time integrations combine with these spatial integration to give a space time integral. Thus, we

¹¹ Note that while abelian symmetries could allow a non-zero multiple of identity for invariance (i.e. $P^\mu|0\rangle = \alpha^\mu|0\rangle$), non-abelian symmetry generators have to annihilate the vacuum. Hence the Lorentz generators must annihilate the vacuum and then the commutator of $M^{\mu\nu}$ with P^λ shows that the $\alpha^\mu = 0$ must also hold. Poincare group being non-abelian must have its generators being normal ordered for the vacuum to be invariant.

get,

$$\begin{aligned}
G_2(x_1, x_2)|_{Nr} &= \langle 0|T\{\phi(x_1)\Phi(x_2)\}|0\rangle - ig \int d^4y \langle 0|T\{\Phi(x_1)\Phi(x_2) : \Phi^3(y) : \}|0\rangle \\
&\quad + \frac{(-i)^2}{2} g^2 \int d^4y_1 \int d^4y_2 \langle 0|T\{\Phi(x_1)\Phi(x_2) : \Phi^3(y_1) :: \Phi^3(y_2) : \}|0\rangle \\
&= i\Delta_F(x_1 - x_2) - \frac{g^2}{2} \int d^4y_1 \int d^4y_2 \langle 0|T\{\Phi(x_1)\Phi(x_2) : \Phi^3(y_1) :: \Phi^3(y_2) : \}|0\rangle
\end{aligned}$$

The first term is just the Feynman propagator and the $o(g)$ term vanishes since it is a 5-point function. The non-trivial term is the $o(g^2)$ term. This is not in the form of the Wick's theorem, in that there are normal ordered factors. We can replace the normal ordered product in terms of unordered product minus the product of two point functions i.e. using the Wick's theorem in reverse form. For instance, consider (for three distinct points)

$$\begin{aligned}
:\Phi(y_1)\Phi(y'_1)\Phi(y''_1): &= \Phi(y_1)\Phi(y'_1)\Phi(y''_1) - \langle 0|\Phi(y_1)\Phi(y'_1)|0\rangle\Phi(y''_1) \\
&\quad - \langle 0|\Phi(y'_1)\Phi(y''_1)|0\rangle\Phi(y_1) - \langle 0|\Phi(y_1)\Phi(y''_1)|0\rangle\Phi(y'_1)
\end{aligned}$$

And likewise for the second factor of $:\Phi(y_2)^3:$. If we now substitute these in the $o(g^2)$ term and eventually take the coincidence limit of the primed arguments, we see that the two point functions of coincident points cancel out. The net result result is that in writing the permutation terms, we omit the Feynman propagators of coincident points and this holds because we use the normal ordered interaction Hamiltonian. If we did not normal order the interaction Hamiltonian then these terms, which are formally divergent, will remain and will have to be handled differently. This is what will happen in the functional integral method which will be discussed later. Keeping this in mind, the $o(g^2)$ term under vev becomes,

$$\begin{aligned}
&\langle 0|T\{\Phi(x_1)\Phi(x_2) : \Phi^3(y_1) :: \Phi^3(y_2) : \}|0\rangle \\
&= i\Delta_F(x_1 - x_2).i\Delta_F(y_1 - y_2).i\Delta_F(y_1 - y_2).i\Delta_F(y_1 - y_2) + \text{permutations} \\
&= (i^4) [\Delta_F(x_1 - x_2)\Delta_F(y_1 - y_2)^3 + \Delta_F(x_1 - y_1)\Delta_F(x_2 - y_2)\Delta_F(y_1 - y_2)^2 \\
&\quad + \Delta_F(x_1 - y_2)\Delta_F(x_2 - y_1)\Delta_F(y_1 - y_2)^2]
\end{aligned}$$

There are no other 'pairings' of space-time points as they would involve Feynman propagators with coincident arguments.

The denominator of the n -point function has the same structure as the numerator except

for the $\Phi(x_1)\Phi(x_2)$ fields i.e.

$$\begin{aligned}
G_2(x_1, x_2)|_{Dr} &= \langle 0|0\rangle - ig \int d^4y \langle 0|T\{\Phi^3(y)\}|0\rangle \\
&\quad + \frac{(-i)^2}{2} g^2 \int d^4y_1 \int d^4y_2 \langle 0|T\{\Phi^3(y_1) :: \Phi^3(y_2)\}|0\rangle \\
&= 1 - \frac{g^2}{2} \int d^4y_1 \int d^4y_2 \langle 0|T\{\Phi^3(y_1) :: \Phi^3(y_2)\}|0\rangle \\
&= 1 + \frac{(-ig)^2}{2} (CF) \int d^4y_1 \int d^4y_2 (i\Delta_F(y_1 - y_2))^3
\end{aligned}$$

Here, CF is the *combinatorial factor* - number of distinct ways of effecting the same pairings - and equals $3!$ in the example above.

The generalization is fairly obvious. For definiteness consider the interaction Hamiltonian *density* to be $H_I(\Phi(y)) := \frac{g}{k!} : \Phi^k(y) :$. Then, it is apparent that the numerator of a general n -point function is a sum of the m^{th} order contributions, $m = 0, 1, \dots$. The m^{th} order contribution has the form:

- $\frac{(-ig/k!)^m}{m!} \int d^4y_1 \dots d^4y_m \langle 0|T\{\Phi(x_1) \dots \Phi(x_n) : \Phi^k(y_1) : \dots : \Phi^k(y_m)\}|0\rangle$;
- The Wick's theorem gives a sum of terms, each of which is a product of a total of $(n + km)/2$ Feynman propagators, $i\Delta_F(x_j - y_k)$, whose arguments give the 'pairing' of space-time points from the set: x_1, \dots, x_n, k -copies of y_1, \dots, k -copies of y_m . The normal ordered form of interaction Hamiltonian ensures that the paired points are necessarily *distinct*. The x_1, \dots, x_n points are assumed to be distinct. A 'pairing' is also called a 'Wick contraction'. The sum of terms is generated from any particular one by permutations. If $n + km$ is odd, the contribution vanishes;

- There would also be combinatorial factor counting distinct ways of getting the same set of pairing (or Feynman propagators). The denominator has similar contributions except the x_1, \dots, x_n points.

There is a convenient diagrammatic representation to keep track of the generated terms as well as associating specific factors and integrals with them. The diagrams are the *Feynman Diagrams* while the prescription to construct associated integral is given by the *Feynman rules*.

Procedure to generate Feynman diagrams for Green's function:

- For each distinct x_1, \dots, x_n , draw a *vertex* with a single edge. Call these vertices as *external vertices*;

- For normal ordered monomial of order k , draw a vertex y_j with k edges sticking out.





Call these as the *internal vertices*;

- Wick contract or pair, by joining two edges from two distinct vertices, internal or external. If the pair has one or both vertices to be external vertices, the line is called an *external line* while a pairing with both vertices being internal, is called an *internal line*. This constructs a Feynman diagram with vertices and internal/external lines.

Procedure to associate integral with a Feynman diagram:

- With each contraction, associate a Feynman propagator, $i\Delta_F(z' - z'')$.
- With each internal vertex, associate a factor of $(-i) \times$ coefficient of the monomial of the fields in $H_I(\Phi(y))$ and an integral over its space-time location, $\int d^4y$.
- Associate a numerical factor as a ratio. In the numerator, count the number of *distinct* ways of generating the same diagram (or the same contractions). In the denominator, put $m!$ if m is the total number of internal vertices.

For the above example of $G_2(x_1, x_2)$ to second order for $H_I = g : \Phi(y)^3$:, here is a summary:

$o(g)$		Has one loose end and vev is zero.
$o(g^2)$		
(a)		$(-ig)^2 \left(\frac{3.3.2.1}{2!}\right) \int d^4 y_1 d^4 y_2 (i\Delta_F(x_1 - y_1)) \times (i\Delta_F(x_2 - y_2))(i\Delta_F(y_1 - y_2))^2$
(b)		$(-ig)^2 \left(\frac{3.3.2.1}{2!}\right) \int d^4 y_1 d^4 y_2 (i\Delta_F(x_1 - y_2)) \times (i\Delta_F(x_2 - y_1))(i\Delta_F(y_1 - y_2))^2$
(c)		$(-ig)^2 \left(\frac{1.3.2.1}{2!}\right) \int d^4 y_1 d^4 y_2 (i\Delta_F(x_1 - x_2))(i\Delta_F(y_1 - y_2))^3$

The diagrams (a) and (b) are examples of *connected diagrams* while (c) is an example of a *topologically disconnected diagram*. Here (dis)connection refers to (dis)connection with *external vertices*. A diagram with no external vertices is called a *vacuum bubble*. The diagram in (c) has a disconnected piece which is a vacuum bubble. The denominator diagrams are all vacuum bubbles.

It is actually possible to separate the vacuum bubble components of the diagrams in the numerator and cancel them against those in the denominator. The proof goes as follows.

In an n -point function, at the k^{th} order, the numerator has the form,

$$Nr = \sum_{k=0}^{\infty} \frac{(-i)^k}{k!} \int d^4 y_1 \dots d^4 y_k \langle 0 | T \{ \Phi(x_1) \dots \Phi(x_n) : H_I(y_1) : \dots : H_I(y_k) : \} | 0 \rangle$$

Group the contractions into two groups: (i) those that have l of the internal vertices with a contraction with *at least one* of the external vertices and (ii) the remaining $(k - l)$ internal vertices which have no segment connecting an external vertex. The above vev then splits

as,

$$\begin{aligned} \langle 0|T\{\Phi(x_1)\dots\Phi(x_n):H_I(y_1):\dots:H_I(y_k):\}|0\rangle = \\ \langle 0|T\{\Phi(x_1)\dots\Phi(x_n):H_I(y_1):\dots:H_I(y_l):\}|0\rangle \times \\ \langle 0|T\{H_I(y_1):\dots:H_I(y_{k-l})\}|0\rangle \end{aligned}$$

And this split can happen in kC_l ways. Hence,

$$\begin{aligned} Nr = \sum_{k=0}^{\infty} \frac{(-i)^k}{k!} \sum_{l=0}^k \frac{k!}{l!(k-l)!} \int d^4y_1\dots d^4y_l \\ \langle 0|T\{\Phi(x_1)\dots\Phi(x_n):H_I(y_1):\dots:H_I(y_l):\}|0\rangle \times \\ \int d^4y_{l+1}\dots d^4y_k \langle 0|T\{H_I(y_{l+1})\dots:H_I(y_k)\}|0\rangle \end{aligned}$$

Putting $(-i)^k = (-i)^l(-i)^{(k-l)}$ and interchanging the order of summation we can write,

$$\begin{aligned} Nr = \left[\sum_{l=0}^{\infty} \frac{(-i)^l}{l!} \int d^4y_1\dots d^4y_l \langle 0|T\{\Phi(x_1)\dots\Phi(x_n):H_I(y_1):\dots:H_I(y_l):\}|0\rangle \right] \\ \times \left[\sum_{k=l}^{\infty} \frac{(-i)^{(k-l)}}{(k-l)!} \int d^4y_{l+1}\dots d^4y_k \langle 0|T\{H_I(y_{l+1})\dots:H_I(y_k)\}|0\rangle \right] \end{aligned}$$

$k \rightarrow k+l$ in the second square bracket, shows it as $\langle 0|T\{\exp -i \int_{-\infty}^{\infty} dt H(t)\}|0\rangle$ which is just the denominator!

The first factor is called the *connected Green's function* and denoted as G_n^c . It consists of only the diagrams connected to external vertices. The diagrams may be topologically disconnected. *From now on we focus on the connected Green's functions and drop the denominator.*

Having gotten Feynman diagrams and Feynman integrals, we further simplify the expression for the S -matrix elements.

To go from Green's functions to S -matrix elements (a) we operate on the Green's function by the equation of motion differential operator for each external vertex; (b) insert the wave function factors for each external vertex; (c) divide by \sqrt{Z} for each external vertex and (d) integrate over the external vertices.

We have only connected diagrams. Among these are topologically disconnected diagrams too. There is a special subclass of topologically disconnected diagrams in which two external vertices are Wick contracted. These represent an incoming particle which goes out *without any scattering*. We can separate such un-scattered processes from the S -matrix (which

we will do little later) and focus on processes wherein every incoming particle necessarily scatters i.e. *every external vertex is necessarily connected to an internal vertex*.

Consider an external vertex x connected with an internal vertex y . Associated with the external vertex is a wavefunction factor $\frac{e^{ik \cdot x}$ (or $e^{-ik' \cdot x'}$), integration over x (or x'), Feynman propagator $\Delta_F(x - y)$ and $(\square_x - m^2)$ acting on the propagator. Since the propagator is a Green's function, $(\square_x - m^2)\Delta_F(x - y) = -\delta^4(x - y)$. Integration over x then removes the delta function and $e^{ik \cdot x} \rightarrow e^{ik \cdot y}$. For the external vertex of an out-going particle, $e^{-ik' \cdot x'} \rightarrow e^{-ik' \cdot y}$. Thus the integration over external vertices is trivially carried out and the propagators involving an external vertex is removed - an “external leg is said to be amputated”.

Now use the Fourier representation of the remaining propagators, $\Delta_F(y_i - y_j) = \int \frac{d^4l}{(2\pi)^4} \frac{e^{il \cdot (y_i - y_j)}}{l^2 + m^2 - i\epsilon}$. Each internal vertex y_i is connected to possibly several other internal vertices as well as possibly several external vertices. For internal vertex, the Fourier transform supplies a factor of $e^{il \cdot y_i}$ and each external vertex provides a factor of $e^{k \cdot y_i}$. All of these combine and the $\int d^4y_i$ then gives a $(2\pi)^4 \delta^4(\sum k - \sum k' + \sum l)$. Thus, the integration over internal vertices result in a momentum conserving delta function. The space-time integrations are done but we are left with integration over the Fourier momenta which are associated with internal lines. The delta functions trivialize many momentum integrations leaving an overall delta function involving only the external momenta and enforcing the momentum conservation. We also left with some unconstrained *loop momenta*.

To be explicit, let n_E, n_I denote the number of external and internal lines and let n_v denote the number internal vertices. We have then $n_E + n_I$ number of momenta and n_v number of conservation equations. Thus $n_E + n_I - n_v + 1$ is the number of undetermined momenta or the loop momenta. The +1 in the counting signifies the loss of one conservation equation due to the left over delta function enforcing conservation of external momenta.

There are no dependence on space-time points, no space-time integrations left and the scattering matrix element is given by integration over a bunch of loop momenta and a product of momentum space propagators forming the integrand and of course the various factors of i, π and numerical combinatorial factors.

$$S_{fi} \sim \delta^4(\sum k - \sum k') \prod_j \int d^4l_j \prod_{n_I} \left(\frac{1}{p_i^2(k, l) + m^2 - i\epsilon} \right)$$

We will elaborate the numerical factors in explicit examples. The structure of the S -matrix elements (non-trivial scattering) should be clear from the above discussion.

To separate out the trivial scattering it is customary to introduce the so-called T -matrix as,

$$S := \mathbb{1} + iT \quad , \quad \langle k'_{j'} | T | k_j \rangle := (2\pi)^4 \delta^4(\Sigma_{j'} k'_{j'} - \Sigma_j k_j) \mathcal{M}(k_j \rightarrow k'_{j'})$$

Here the k, k' denote the on-shell momenta of the incoming and outgoing particles. The delta function enforcing momentum conservation is a consequence of translation invariance and (M) is called the *invariant matrix element*. This is what is computed in practice. The S_{fi} given above is really the iT_{fi} .

In practice, computation of the T -matrix elements is only part of what is needed to compare with experiments which measure *cross-sections*. To compute cross-sections, recall that the S -matrix elements are the transition probability amplitudes whose non-trivial contributions are identified as $\langle f | i \rangle = (2\pi)^4 \delta^4(k'_{total} - k_{total}) (i\mathcal{M}_{i \rightarrow f})$. The cross-section is the ratio of the number of outgoing particles per second to the incident flux. The numerator is proportional to the transition probability rate while the denominator is determined by the initial state.

We have given the S -matrix elements in the plane wave basis which is strictly incorrect. One should use wave packets for representing the asymptotic states. Alternatively, a commonly used practice is to put the system in a finite space-time box and take the limit of infinite box at the end. This is simpler to implement in practice and suffices for most purposes. We will use this [12] and refer to [13] for the wave packet treatment.

B. Differential Cross-section for $2 \rightarrow n$ process

Imagine the scattering experiment to be enclosed in a large spatial box of volume $V = L^3$, with periodic boundary conditions imposed on the mode functions. Let the duration of the experiment be a large time interval T . The probability of transition from an initial state $|i\rangle$ to a final state $|f\rangle$ is given by,

$$Prob_{i \rightarrow f} = \frac{|\langle f | i \rangle|^2}{\langle i | i \rangle \langle f | f \rangle} = \frac{[(2\pi)^4 \delta^4(\Sigma k'_{j'} - \Sigma k_j)] [(2\pi)^4 \delta^4(0)] |i\mathcal{M}_{i \rightarrow f}|^2}{\langle i | i \rangle \langle f | f \rangle}$$

The square of the momentum conservation delta function is written with one factor as $\delta(0)$ which is to be understood as: $(2\pi)^4 \delta^4(0) = \int d^4x e^{i0 \cdot x} = \int d^4x := VT$. The initial and

final states are normalized using $\langle k|k \rangle := \lim_{k' \rightarrow k} \delta^3(\vec{k}' - \vec{k}) = \delta^3(0) := \frac{V}{(2\pi)^3}$. Typically, the initial state consists of *two particles*. So let us specialize to this case of $2 \rightarrow n$ processes. Then, $\langle i|i \rangle \langle f|f \rangle = [\frac{V}{(2\pi)^3}]^{(n+2)}$. Therefore,

$$\frac{Prob_{i \rightarrow f}}{T} = \frac{[(2\pi)^4 \delta^4(\Sigma k'_{j'} - \Sigma k_j)] |i\mathcal{M}_{i \rightarrow f}|^2 V}{[V/(2\pi)^3]^{n+2}}$$

Real detectors have finite aperture and hence the detected particle's momentum is anywhere within a small window around the central value. The corresponding probabilities are thus to be added. An estimate for such a window follows from the box normalization we have taken. The momenta are given by $\vec{k}_j = \frac{2\pi}{L} \vec{n}_j$. Summing over the momenta within a window is same as summing over the integers within a window and for large volume, $\Sigma \vec{n}_j \approx \frac{V}{(2\pi)^3} \int d^3 k$, for each final state particle. This leads to the total probability per unit time for the transition,

$$\begin{aligned} \frac{Prob_{2 \rightarrow n}}{T} d\Gamma_n &= \frac{[(2\pi)^4 \delta^4(\Sigma k'_{j'} - \Sigma k_j)] |i\mathcal{M}_{2 \rightarrow n}|^2 V}{[V/(2\pi)^3]^{n+2}} \prod_{j=1}^n \left[\frac{d^3 k_j}{(2\pi)^3} V \right] \\ &= \left\{ \frac{[(2\pi)^4 \delta^4(\Sigma k'_{j'} - k_1 - k_2)] |i\mathcal{M}_{2 \rightarrow n} \sqrt{\prod_{j=1}^{n+2} [2\omega_{\vec{k}_j} (2\pi)^3]}|^2}{V(2\omega_{\vec{k}_1})(2\omega_{\vec{k}_2})} \right\} \prod_{j=1}^n \left[\frac{d^3 k_j}{2\omega_{\vec{k}_j} (2\pi)^3} \right] \end{aligned}$$

In going from the first to the second line, we have divided and multiplied by the product of $(2\omega_{\vec{k}_j})$ for all the $(n+2)$ particles. The last product is the $d\Gamma_n$ which is the *Lorentz invariant phase space volume*. The square root factor will get absorbed in the invariant matrix element and will get rid of similar factors coming from the wave functions of the incoming and outgoing particles.

We also need the incident flux. For the initial state of two articles, consider the *laboratory frame* where the particle '2' is at rest. The particle '1' has a speed $|\vec{k}_1|/E_1$ and gives the number density of one particle per unit volume, $1/V$. Hence, the incident number flux is $|\vec{k}_1|/(E_1 V)$. Dividing by the flux, the *exclusive, differential cross-section* for $2 \rightarrow n$ process is given by,

$$\begin{aligned}
d\sigma_{lab} &= \frac{(2\pi)^4 \delta^4(k_1 + k_2 - \sum_j k_j) |i\tilde{\mathcal{M}}_{2 \rightarrow n}|^2}{4m_2 |(\vec{k}_1)_{lab}|} d\Gamma_n && \text{where,} \\
\tilde{\mathcal{M}}_{2 \rightarrow n} &:= \mathcal{M}_{2 \rightarrow n} \sqrt{\prod_{j=1}^{n+2} [2\omega_{\vec{k}_j} (2\pi)^3]} && \text{and,} \\
d\Gamma_n &:= \prod_{j=1}^n \frac{d^3 k_j}{2\omega_{\vec{k}_j} (2\pi)^3}
\end{aligned}$$

We have used $\omega_{\vec{k}_2} = m_2$ and $\omega_{\vec{k}_1} = E_1$. Notice that all factors of the volume and the duration T have disappeared. The explicit square root factors will also cancel out.

Many basic calculations involve scattering processes with two particles going into two particles. We have already used the initial state of two particles to obtain the incident flux. We will now also specialize to $n = 2$ for out going particles and write the phase space integrals more explicitly.

1. The Special case of $2 \rightarrow 2$ processes

Let the initial momenta be denoted by k_1, k_2 and the final momenta be denoted by k'_1, k'_2 . Lorentz invariance implies that the scattering amplitude will have Lorentz indices (tensorial or spinorial) carried by the wavefunctions and the invariant amplitude will be a function of Lorentz invariants. We have three independent momenta thanks to the overall conservation due to translation invariance and we can form 6 Lorentz scalars of the form $p_i \cdot p_j$. Of these three are masses and the remaining three are conveniently defined as ‘‘center of mass energy’’ and two types of ‘‘squared momentum transferred’’. These are known as the *Mandelstam variables* and are defined as $(k_1 + k_2 = k'_1 + k'_2)$,

$$s := -(k_1 + k_2)^2 = -(k'_1 + k'_2)^2 \quad (\text{Centre of Mass Energy}) \quad (11.14)$$

$$t := -(k'_1 - k_1)^2 = -(k'_2 - k_2)^2 \quad (\text{Squared Momentum transfer}) \quad (11.15)$$

$$u := -(k'_2 - k_1)^2 = -(k_2 - k'_1)^2 \quad (\text{Squared Momentum transfer}) \quad (11.16)$$

The definitions imply the *Mandelstam identity*: $s + u + t = m_{k_1}^2 + m_{k_2}^2 + m_{k'_1}^2 + m_{k'_2}^2$.

Laboratory and Centre of Mass Frames:

The lab frame, is defined by regarding particle 2, say, at rest while particle 1 is incident

on it. Thus,

$$k_1^{lab} = (\sqrt{\vec{k}_1^2 + m_1^2}, \vec{k}_1^{lab}), \quad k_2^{lab} = (m_2, \vec{0})$$

The lab frame is convenient for expressing the incident flux.

The center of mass frame is defined by $\vec{k}_{total} := \vec{k}_1 + \vec{k}_2 = 0 = \vec{k}'_1 + \vec{k}'_2$. Thus,

$$k_1^{cm} = (\sqrt{\vec{k}_{cm}^2 + m_1^2}, \vec{k}_{cm}), \quad k_2^{cm} = (\sqrt{\vec{k}_{cm}^2 + m_2^2}, -\vec{k}_{cm})$$

This frame is more convenient for defining scattering angle. The common momentum direction singles out say, the z-axis. Orienting the frame accordingly, we take

$$\begin{aligned} k_1 &= (E_1, k_{cm}\hat{z}), \quad k_2 = (E_2, -k_{cm}\hat{z}), \\ k'_1 &= (E'_1, \vec{k}'_1), \quad k'_2 = (E'_2, -\vec{k}'_1), \quad \hat{k}'_1 \cdot \hat{z} =: \cos(\Theta_{cm}). \end{aligned} \quad (11.17)$$

The Mandelstam invariant s can be used to relate the lab frame momentum $|k_1|^{lab}$ and the center of mass momentum $|k|_{cm}$. The invariant $s := -(k_1 + k_2)^2 = -(-m_1^2 - m_2^2 - 2E_1E_2 + 2\vec{k}_1 \cdot \vec{k}_2)$ has two equivalent expressions:

$$\begin{aligned} s|_{lab} &= m_1^2 + m_2^2 + 2m_2\sqrt{m_1^2 + \vec{k}_1^2} \\ s|_{cm} &= m_1^2 + m_2^2 + 2\sqrt{m_1^2 + \vec{k}_{cm}^2}\sqrt{m_2^2 + \vec{k}_{cm}^2} + 2\vec{k}_{cm}^2. \end{aligned}$$

These can be solved for the momenta and give manifestly Lorentz invariant expressions,

$$|k_1|_{lab} = \frac{1}{2m_2}\sqrt{s^2 - 2s(m_1^2 + m_2^2) + (m_1^2 - m_2^2)^2} \quad (11.18)$$

$$|k|_{cm} = \frac{1}{2\sqrt{s}}\sqrt{s^2 - 2s(m_1^2 + m_2^2) + (m_1^2 - m_2^2)^2} \quad (11.19)$$

Phase space volume in center of mass frame: We have the Lorentz invariant definition,

$$d\Gamma_2 = (2\pi)^4 \delta^4(k'_1 + k'_2 - k_1 - k_2) \frac{d^3k'_1}{2\omega_{k'_1}(2\pi)^3} \frac{d^3k'_2}{2\omega_{k'_2}(2\pi)^3}.$$

In the CM frame we simplify it using: $\delta^4 \rightarrow \delta(E'_1 + E'_2 - \sqrt{s})\delta^3(k'_1 + k'_2)$ since $\vec{k}_1 + \vec{k}_2 = 0$, $s = (E_1 + E_2)^2$. We can remove the momentum delta function by integrating over \vec{k}'_2 . Denoting $d^3k'_1 := d^3k'_{cm} = dk'_{cm}(k'_{cm})^2 d\Theta_{cm} \sin^2(\Theta_{cm}) d\phi$, we write,

$$\int d^3k'_2 d\Gamma_2 = \frac{1}{(2\pi)^2} \delta(E'_1 + E'_2 - \sqrt{s}) \frac{1}{4E'_1 E'_2} dk'_{cm} (k'_{cm})^2 d\Theta_{cm} \sin^2(\Theta_{cm}) d\phi$$

Since $E'_1 + E'_2 = \sqrt{(m'_1)^2 + (k'_{cm})^2} + \sqrt{(m'_2)^2 + (k'_{cm})^2}$ is a function of k'_{cm} , we simplify the delta function using $\delta(f(x)) = \sum_i \frac{\delta(x-x_i)}{f'(x_i)}$, $f(x_i) = 0$. We have

$$f'(k'_{cm}) = \frac{k'_{cm}}{E'_1} + \frac{k'_{cm}}{E'_2} = k'_{cm} \frac{E'_1 + E'_2}{E'_1 E'_2} = \frac{k'_{cm} \sqrt{s}}{E'_1 E'_2},$$

$$f(k'_{cm}) = 0 \Rightarrow k'_{cm} = \frac{\sqrt{s^2 - 2s((m'_1)^2 + (m'_2)^2) + ((m'_1)^2 - (m'_2)^2)^2}}{2\sqrt{s}}$$

Doing the dk'_{cm} integration using the delta function gives,

$$\int d(k'_{cm})^2 \delta(E'_1 + E'_2 - \sqrt{s}) = \frac{(k'_{cm})^2 E'_1 E'_2}{k'_{cm} \sqrt{s}} \text{ which gives } d\Gamma_2 = \frac{1}{16\pi^2} d^2\Omega_{cm} \frac{k'_{cm}}{\sqrt{s}}. \quad (11.20)$$

Thus, for the special case of a process with 2 particles going to 2 particles, one gets the *differential cross-section* as,

$$\frac{d\sigma}{d\Omega_{cm}} = \frac{1}{64\pi^2} \frac{1}{s} \frac{k'_{cm}}{k_{cm}} |\mathcal{M}|^2 \quad \text{where we have used } 2m_2 |k_1|^{lab} = 2\sqrt{s} |k|_{cm}, \text{ and} \quad (11.21)$$

$$\frac{k'_{cm}}{k_{cm}} = \sqrt{\frac{s^2 - 2s((m'_1)^2 + (m'_2)^2) + ((m'_1)^2 - (m'_2)^2)^2}{s^2 - 2s(m_1^2 + m_2^2) + (m_1^2 - m_2^2)^2}}. \quad (11.22)$$

The invariant amplitude has dependence on the scattering angle Θ_{cm} and the total cross-section is obtained by integrating over the center of mass solid angle.

12. DIAGRAMMATIC RECIPE FOR S-MATRIX ELEMENTS

We now specify explicit interacting fields by giving an interaction Lagrangian and state the Feynman rules to complete the diagrammatic recipe for the T – *matrix* elements. We will state the rules for the Φ^4 theory, the Yukawa theory and the Quantum electrodynamics (QED). In the next section we will compute specific processes. There is a good deal of conventions of normalization etc and they have to be kept track of carefully. The free (quadratic) part of the Lagrangian density sets the normalization of the propagators while the interaction terms (beyond quadratic in fields) contribute to the numerical factors. Here are the terms in the *Lagrangian densities*.

$$\text{Free Scalar} : -\frac{1}{2}\partial^\mu\Phi\partial_\mu\Phi - \frac{1}{2}m^2\Phi^2(x) ; \quad (12.1)$$

$$\text{Free Spinor} : -i\bar{\Psi}\not{\partial}\Psi + m\bar{\Psi}\Psi ; \quad (12.2)$$

$$\text{Massless Vector} : -\frac{1}{4}F^{\mu\nu}F_{\mu\nu} , \quad F_{\mu\nu} := \partial_\mu A_\nu - \partial_\nu A_\mu ; \quad (12.3)$$

$$\text{Scalar self coupling} : -\frac{g}{3!}\Phi^3(x) - \frac{\lambda}{4!}\Phi(x)^4 ; \quad (12.4)$$

$$\text{Yukawa coupling} : -g\Phi(x)\bar{\Psi}(x)\Psi(x) ; \quad (12.5)$$

$$\text{QED coupling} : -ieA_\mu(x)\bar{\Psi}(x)\gamma^\mu\Psi(x) . \quad (12.6)$$

$$\text{In the interaction} \quad \textit{Hamiltonian}, \text{ all coupling terms will change signs.} \quad (12.7)$$

The propagators are the Feynman Green's functions for the free equations of motion. Consider the massless vector field as this has a new feature.

The free Lagrangian (Maxwell) can be expressed as $\frac{1}{2}A_\mu(\eta^{\mu\nu}\square - \partial^\mu\partial^\nu)A_\nu$ + divergence terms. The equation of motion is $(\eta^{\mu\nu}\square - \partial^\mu\partial^\nu)A_\nu(x) = 0$. Equivalently, in Fourier space equation takes the form $-(\eta^{\mu\nu}k^2 - k^\mu k^\nu)\tilde{A}_\nu(k) = 0$. However, the differential operator is not invertible and hence does not admit a Green's function! The non-invertibility follows because every A_ν of the form $\partial_\nu f(x)$, solves the equation. That is, $\partial_\nu f$ is a non-trivial eigenvector of the differential operator, with zero eigenvalue. For perturbation theory though we need a propagator for which extra terms are added to the action to break its gauge invariance - invariance under $\delta A_\mu(x) = \partial_\mu\Lambda(x)$. This can be done in several ways and each choice corresponds to a gauge.

A common and convenient choice is to add $-\frac{1}{2}(\partial_\mu A^\mu)^2$ term to the action. Up to a divergence term, this is just $+\frac{1}{2}A_\mu\partial^\mu\partial^\nu A_\nu$ and precisely cancels the term in the equation

of motion operator, making it $\eta^{\mu\nu}\square$ which is invertible. The added term is manifestly Lorentz invariant and the corresponding gauge is called the *Lorentz gauge*. For our purposes, this gauge will suffice. The propagator is an inverse of the differential operator since $(\eta^{\mu\nu}\square_x)(D_F)_{\nu\lambda}(x-y) = \delta^\mu_\lambda \delta^4(x-y)$.

The QED coupling arises from the minimal substitution rule: $\vec{P} \rightarrow \vec{P} + e\vec{A} \leftrightarrow -i\partial_\mu \rightarrow -i\partial_\mu + eA_\mu \leftrightarrow \partial_\mu \rightarrow \partial_\mu + ieA_\mu$. Substitution in the free Dirac action gives a $-ieA_\mu$ as the QED coupling.

Let us gather the various factors for the scalar field in the T -matrix elements.

- Each in/out particle gives a wave function factor of $[2\omega_{\vec{k}}(2\pi)^3]^{-1/2}e^{\pm ik\cdot x}$, an $\int d^4x$ and $(\square_x - m^2)$ acting on the n -point function. For a spinor field, the amputation operator changes to $(-i\not{\partial} + m)$ and we have additionally the u, v, \bar{u}, \bar{v} spinors. For a vector field, the amputation operator changes and we have the polarization vectors in addition. Other factors remain the same.
- Each order in H_I , gives $(-i) \int d^4y$ from the T -ordered exponential and a $(m!)^{-1}$ for the m^{th} order, from the exponential;
- Each Wick contraction gives $(i\Delta_F(z-z')) := \int \frac{d^4l}{(2\pi)^4} \frac{ie^{il\cdot(z-z')}}{l^2+m^2-i\epsilon}$;
- Each amputation, action of $(\square - m^2)$ gives a $(-\delta^4(x-y))$ while each integration over x or y gives $(2\pi)^4$ times a momentum conserving δ^4 . The momentum integrations do not produce any factors.
- Let E, I, V denote the number of external lines (= number of external vertices), number of internal lines and number of internal vertices (= order of H_I) respectively. Then, the factors of $[2\omega_{\vec{k}}(2\pi)^3]^{-1/2}$, precisely cancel the explicit factor we found in the amplitude $\tilde{M}_{2 \rightarrow n}$. This is due to the normalization choices which cancel out in $\frac{|\langle f|i \rangle|^2}{\langle i|i \rangle \langle f|f \rangle}$ and can now be dropped from both \tilde{M} and from the T -matrix element.

Factors of i : $(-i)^V (i)^I = (-1)^V i^{V+I}$;

Factors of 2π : $(2\pi)^{-4I+4E+4V-4}$. The last -4 is because it has been taken out in the definition of the \mathcal{M} due to the overall momentum conservation.

Vertex factors, including the i in the QED vertex are to be taken case-by-case. There is the $(m!)^{-1}$ and a combinatorial factor that will come for each diagram. These too

are taken case-by-case.

With these, we now state the Feynman rules for Feynman diagrams.

External Line:				
scalar		1		1
fermion		$u(p, \sigma)$		$\bar{u}(p, \sigma)$
anti-fermion		$\bar{v}(p, \sigma)$		$v(p, \sigma)$
photon		$\varepsilon_\mu^*(p, \lambda)$		$\varepsilon_\mu(p, \lambda)$

Internal Line:		
scalar		$\frac{-i}{k^2+m^2-i\epsilon}$
fermion		$\frac{(-i)(-\not{k}+m)}{k^2+m^2-i\epsilon}$
photon (Lorentz gauge)	$\mu \text{ ~~~~~ } \nu$	$\frac{-i\eta^{\mu\nu}}{k^2-i\epsilon}$

Vertex:		
Φ^3		$i\frac{g}{3!}$
Φ^4		$i\frac{\lambda}{4!}$
Yukawa		ig
QED		$ie\gamma^\mu$

13. ELEMENTARY PROCESSES IN YUKAWA AND QED: NR LIMIT

We begin with scattering of a fermion off another fermion, interacting via the Yukawa coupling and compute the T -matrix element to the leading order. We will take the non-relativistic limit and identify an *equivalent potential*. We will consider anti-fermion scattering and fermion-anti-fermion scattering as well. We will then compare the qed coupling and briefly the gravitational coupling. This will lead to appreciate the dependence the attractive/repulsive nature of the interaction on spin of the exchanged particle.

Consider a general process depicted below together with its ‘expansion’.

$$\begin{array}{c}
 \begin{array}{ccc}
 p & & p' \\
 \swarrow & & \searrow \\
 \text{[Shaded Circle]} & & \\
 \nwarrow & & \nearrow \\
 k & & k'
 \end{array}
 \quad = \quad
 \begin{array}{ccc}
 p & \longrightarrow & p' \\
 k & \longrightarrow & k'
 \end{array}
 \quad + \quad
 \begin{array}{ccc}
 p & & p' \\
 \swarrow & & \searrow \\
 & \text{---} & \\
 \nwarrow & & \nearrow \\
 k & & k'
 \end{array}
 \quad + \quad
 \begin{array}{ccc}
 p & & p' \\
 \swarrow & & \searrow \\
 & \text{---} & \\
 \nwarrow & & \nearrow \\
 k & & k'
 \end{array}
 \quad + \dots \quad (13.1)
 \end{array}$$

The first term ($o(g^0)$) denotes ‘no scattering’. There are two contributions at $o(g^2)$ corresponding to exchange of two out-going fermions. The overall momentum conservation delta function is the same, enforcing $p + k = p' + k'$. The Feynman rules give the expression as,

$$\begin{aligned}
 i\mathcal{M} = (ig)^2 \left\{ \bar{u}(p')u(p) \frac{-i}{(p' - p)^2 + m_\varphi^2 - i\epsilon} \bar{u}(k')u(k) \right. \\
 \left. - \bar{u}(k')u(p) \frac{-i}{(k' - p)^2 + m_\varphi^2 - i\epsilon} \bar{u}(p')u(k) \right\} \quad (13.2)
 \end{aligned}$$

The *relative* minus sign between the two terms is due to the T-ordering definition for the fermions. The overall sign of the amplitude is determined by the convention adopted for ordering the initial/final state labels for the fermions. See [12, 13]. Notice how the fermion arrows are followed.

Note: If we had anti-fermion scattering, then all the fermion arrows will be reversed and their momenta will be denoted as minus the previous momenta¹². Apart from the reversal of fermion arrows, the $u(p), \bar{u}(p')$ spinors go to $v(p'), \bar{v}(p)$ spinors.

If it is a fermion-anti-fermion scattering, then the second exchange diagram will be absent.

¹² Thus, we may adopt a convention that the diagram displays the fermion arrow and the momentum is also in the same direction. For a fermion the momentum is p and for anti-fermion it is $-p$.

A specific scattering arrangement may permit the initial and the final state fermions to be *distinguishable*. Then only one of the two scattering diagrams will contribute.

Note: The reduction formula for fermions gave a factor of $2m$ for the wavefunctions since we had normalized the spinors as $\bar{u}(p, \sigma)u(p, \sigma') = \delta_{\sigma, \sigma'}$. With the Feynman rules we have adopted in the table the wavefunctions have only the spinors. *This is equivalent to using the normalization:* $\bar{u}(p, \sigma)u(p, \sigma') = 2m\delta_{\sigma, \sigma'} = -\bar{v}(p, \sigma)v(p, \sigma')$, m is of course the fermion mass.

With this noted, the amplitude becomes,

$$i\mathcal{M} = (ig)^2(-i)(2m)^2 \left\{ \frac{\delta_{\sigma_p, \sigma_{p'}} \delta_{\sigma_k, \sigma_{k'}}}{(p' - p)^2 + m_\varphi^2 - i\epsilon} - \frac{\delta_{\sigma_p, \sigma_{k'}} \delta_{\sigma_{p'}, \sigma_k}}{(k' - p)^2 + m_\varphi^2 - i\epsilon} \right\} \quad (13.3)$$

Noting that the scalar field momentum is just the momentum transfer in both the diagrams, we denote it by q . And $q^2 = -(q^0)^2 + |\vec{q}|^2 = |\vec{q}|^2$ in both cases. We thus write the amplitude more conveniently as,

$$i\mathcal{M} = \frac{+ig^2 4m^2}{|\vec{q}|^2 + m_\varphi^2 - i\epsilon} \left\{ \delta_{\sigma_p, \sigma_{p'}} \delta_{\sigma_k, \sigma_{k'}} - \delta_{\sigma_p, \sigma_{k'}} \delta_{\sigma_{p'}, \sigma_k} \right\}.$$

As noted above, only one of the two terms will contribute if the fermions are distinguishable.

In the non-relativistic scattering theory, we have $\langle \vec{k}' | S | \vec{k} \rangle := \langle \vec{k}' | \vec{k} \rangle - 2\pi i \delta(E_{k'} - E_k) \langle \vec{k}' | T | \vec{k} \rangle$ with,

$$\langle \vec{k}' | T | \vec{k} \rangle = \langle \vec{k}' | H' | \vec{k} \rangle = \langle \vec{k}' | V(\vec{q}) | \vec{k} \rangle = V(|\vec{q} = \vec{k}' - \vec{k}|).$$

We can read-off the potential as,

$$V(|q|) = -\frac{4m^2 g^2}{|\vec{q}|^2 + m_\varphi^2} (\delta\delta - \delta\delta) := -\frac{g_{eff}^2}{q^2 + m_\varphi^2}. \quad (13.4)$$

The inverse Fourier transform gives,

$$\begin{aligned} V(\vec{x}) &= -g_{eff}^2 \int \frac{d^3q}{(2\pi)^3} \frac{e^{i\vec{q}\cdot\vec{x}}}{q^2 + m_\varphi^2} = -\frac{g_{eff}^2}{(2\pi)^2} \int_0^\infty dq \frac{q^2}{q^2 + m_\varphi^2} \int_{-1}^1 d(\cos(\theta)) e^{iqr\cos(\theta)} \\ &= -\frac{g_{eff}^2}{4\pi^2(ir)} \int_{-\infty}^\infty dq \frac{qe^{iqr}}{q^2 + m_\varphi^2} \quad \text{contour integrate to pick up } q = im_\varphi, \\ \therefore V(r) &= -\frac{g_{eff}^2}{4\pi} \frac{e^{-m_\varphi r}}{r} \quad \text{(Yukawa Potential)} \end{aligned} \quad (13.5)$$

This is a simple illustration how an effective potential can be inferred from the underlying relativistically specified interaction. Several remarks are in order.

Remark: The most important point is the sign of the potential which renders it *attractive*. But *is the sign unambiguous?*

We have already noted that the overall sign of the amplitude is determined by the convention adopted for ordering of the fermion labels in the in/out states while the relative sign is due to the Pauli principle. When the fermions are distinguishable, either of the two diagrams may be chosen, there is no preference. But the diagrams contribute opposite signs!

The convention arises as follows [13]. We have taken $|p, k\rangle \sim b_p^\dagger b_k^\dagger |0\rangle$ (sequence of creation operators follows label order in the in-state) and $\langle p', k'| \sim \langle 0| b_{k'} b_{p'}$ which is opposite to that of the in-state, but consistent with the Hermitian conjugation. Hence,

$$\begin{aligned} \langle p', k'| P, k\rangle &\sim \langle 0| b_{k'} b_{p'} b_p^\dagger b_k^\dagger |0\rangle = \langle 0| b_{k'} b_k^\dagger |0\rangle \delta^3(p' - p) - \langle 0| b_{k'} b_p^\dagger b_p b_k^\dagger |0\rangle \\ &\approx \delta^3(k' - k) \delta^3(p' - p) - \delta^3(p' - k) \delta^3(k' - p) \end{aligned}$$

Actually there are $\bar{\Psi}\Psi$ fields in between, but being a bilinear it does not change the relative sign. This explains the overall sign.

In a non-relativistic comparison, $|q| \ll m_\varphi$, the small momentum transfer means large spatial separation and hence fermions are distinguishable. If so, there is no Pauli principle or anti-commutation and no relative sign either! So either of the two diagrams will give the same answer.

Ideally, we should consider one of the fermions to be very massive to mimic a source of potential (means $\vec{q} \rightarrow \vec{0}$ limit) and match all the factors for a precise comparison.

The effective coupling contains the Kronecker deltas enforcing conservation of the spin projections. Hence, the Yukawa potential preserves spin projection. For matching with non-relativistic normalization we have to take $2m_{fermion} \delta_{\sigma_p, \sigma_{p'}} \rightarrow 1$. This is needed for inferring the *strength* of the Yukawa potential.

Remark: Suppose we consider fermion-anti-fermion scattering. Then we will have one of the Kronecker deltas to have a minus sign. However, in this case, $b^\dagger \rightarrow d^\dagger$ and doing the contractions with Ψ 's from the Yukawa coupling, we pick up another minus sign. Hence the overall sign does not change. Hence, the Yukawa potential between fermion-anti-fermion is also *attractive*. The same holds for anti-fermion scattering.

Thus, the inferred Yukawa potential is always attractive. Its underlying QFT description is an exchange of a (virtual) massive scalar particle of mass m_φ . Yukawa proposed this potential as a model for binding of nucleons and working with the properties of nucleons,

estimated the scalar mass to be about 200 MeV which is close to the mass of the pion. Of course pions come in three varieties, π^\pm, π^0 while we have taken only a single scalar.

Consider fermion scattering in QED, by replacing the Yukawa coupling by the QED coupling. The relevant diagrams are:



The corresponding expression is,

$$i\mathcal{M} = (ie)^2 \left\{ \bar{u}(p')\gamma^\mu u(p) \frac{-i\eta_{\mu\nu}}{(p' - p)^2 - i\epsilon} \bar{u}(k')\gamma^\nu u(k) - \bar{u}(k')\gamma^\mu u(p) \frac{-i\eta_{\mu\nu}}{(k' - p)^2 - i\epsilon} \bar{u}(p')\gamma^\nu u(k) \right\}, \quad (13.6)$$

with the same relative minus sign.

As before, consider the case of non-relativistic limit with distinguishable fermions and consider only the first diagram. The denominator simplifies the same way: $(p' - p)^2 - i\epsilon \rightarrow |\vec{q}|^2 - i\epsilon$. In the numerator we have $[\bar{u}(p', \sigma')\gamma^\mu u(p, \sigma)][\bar{u}(k', \rho')\gamma_\nu u(k, \rho)]$.

Claim: In the NR limit, $\bar{u}(p', \sigma')\gamma^\mu u(p, \sigma) \rightarrow \bar{u}\gamma^0 u$.

Proof: The spinor u satisfies, $(\not{p} + m)u = 0$. Consider say the Dirac representation of the gamma matrices. Then in terms of the two component notation we have,

$$\begin{pmatrix} -\omega_{\vec{p}} + m & \vec{p} \cdot \vec{\sigma} \\ -\vec{p} \cdot \vec{\sigma} & \omega_{\vec{p}} + m \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = 0 \Rightarrow -\vec{p} \cdot \vec{\sigma} u_1 = (\omega_{\vec{p}} + m)u_2, \vec{p} \cdot \vec{\sigma} u_2 = (-\omega_{\vec{p}} + m)u_1$$

The NR limit $\omega_{\vec{p}} \approx m + \frac{\vec{p}^2}{2m}$ then implies $u_2 \sim o(|p|/m)u_1$. Also, in the two component notation, we have

$$\begin{aligned} \bar{u}'\gamma^i u &= ((u'_1)^\dagger, (u'_2)^\dagger) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 0 & \sigma^i \\ -\sigma^i & 0 \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = (u'_1)^\dagger \sigma^i u_2 + (u'_2)^\dagger \sigma^i u_1 \sim o(p/m); \\ \bar{u}'\gamma^0 u &= ((u'_1)^\dagger, (u'_2)^\dagger) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = (u'_1)^\dagger u_1 + (u'_2)^\dagger u_2 \sim o(1) + o(|p|/m). \end{aligned}$$

This proves the claim.

Hence the numerator approximates to $[\bar{u}(\vec{p}', \sigma')\gamma^0 u(\vec{p}, \sigma)][\bar{u}(\vec{k}', \rho')\gamma^0 u(\vec{k}, \rho)]$ and we get,

$$i\mathcal{M} \approx \frac{ie^2\eta_{00}[u^\dagger(\vec{p}'\sigma')u(\vec{p}, \sigma)][u^\dagger(\vec{k}', \rho')u(\vec{k}, \rho)]}{q^2 - i\epsilon}. \quad (13.7)$$

Since the square brackets are positive ($p' \approx p, k' \approx k$ for small momentum transfer) and the $\eta_{00} < 0$, relative to the Yukawa matrix element we have an opposite sign and hence, the effective potential from the QED coupling will be *repulsive* for fermion scattering. The form of the potential itself can be obtained from the Yukawa one by taking $m_\varphi \rightarrow 0$ and as expected, we get the *Coulomb potential*, $V(r) = \frac{(e')^2}{4\pi} =: \frac{\alpha}{4\pi}$. We have absorbed the normalization factors in e' . By redefining the original e , we can take $e' \rightarrow e$, the measured electric charge (in the natural units). The constant $\alpha := \frac{e^2}{4\pi} \approx \frac{1}{137}$ is called the *fine structure constant*.

If we consider fermion-anti-fermion scattering via the QED coupling, the $u \rightarrow v$ does *not* introduce a negative sign since it is $u^\dagger u$ and not $\bar{u}u$. The $b^\dagger \rightarrow d^\dagger$ introduces a minus sign as before and hence the overall sign changes. Hence, for fermion-anti-fermion, the potential is *attractive*. For anti-fermion scattering of course the potential is repulsive.

Note: It seems the sign depends on the η_{00} which is convention dependent. However, the signs in the propagator will also change accordingly and the sign of the potential will be metric signature independent.



Note: For a tensorial interaction like gravity, helicity 2, we will have two factors of η_{00} and the overall sign will *not* change. The *Newtonian potential* will be *attractive* and will be so for fermion-anti-fermion scattering as well. This argument is to be taken as heuristic since we need to be explicit about the gravitational coupling to infer the numerator factors of the spinors which are sensitive to $u \rightarrow v$ changes.

These examples indicate that exchange of quanta can be interpreted as giving an equivalent potential (and hence force) in the non-relativistic limit (where the concept of potential is meaningful). The qualitative properties of attractive/repulsive are already encoded in the Feynman rules. The comparison with effective potential also allows the coupling parameters to be determined experimentally.

14. BASIC QED PROCESSES

The quantum electrodynamics has basic scattering of two charged particles, the scattering of light by a charged particle, particle-anti-particle annihilation and pair creation. These are analyzed as: (i) electron-muon scattering, (ii) electron-positron scattering as well as production of muon-anti-muon, (iii) electron-photon scattering (Compton scattering) and (iv) electron-positron annihilation into two photons. We will evaluate these in the leading e^2 approximation. We will also compute the cross-sections for comparison with experiments. These cross-sections will be for the *un*-polarized particles in the initial state, for which we will *average the cross section over the initial spins/polarizations*. We will also not detect the final particles spin/polarizations and hence *sum the cross-section over the final spins/polarizations*.

Here is the list of the processes with their diagrams and the corresponding invariant amplitudes.

$e^-(p)\mu^-(k) \rightarrow e^-(p')\mu^-(k')$	(for μ replaced by e it is <i>Moller Scattering</i>)
	$i\mathcal{M} = [\bar{u}(p')(ie\gamma^\mu)u(p)] \left[\frac{-i\eta_{\mu\nu}}{q^2 - i\epsilon} \right] [\bar{u}(k')(ie\gamma^\mu)u(k)]$
$e^+(p')e^-(p) \rightarrow \mu^+(k')\mu^-(k)$	(for μ replaced by e it is <i>Bhabha Scattering</i>)
	$i\mathcal{M} = [\bar{v}(p')(ie\gamma^\mu)u(p)] \left[\frac{-i\eta_{\mu\nu}}{q^2 - i\epsilon} \right] [\bar{u}(k')(ie\gamma^\mu)v(k)]$

In the next two processes, there are two diagrams contributing, obtained by exchanging the initial and final state photons in the Compton scattering and the two final state photons in the annihilation process. Although the diagrams may not look ‘crossed’, they are!

$e^-(p)\gamma(k) \rightarrow e^-(p')\gamma(k')$	(Compton Scattering)
	$i\mathcal{M} = \varepsilon_\nu(\vec{k}', \lambda') \bar{u}(p') (ie\gamma^\nu) \left[\frac{-i(-(\not{p} + \not{k}) + m)}{(p+k)^2 + m_e^2 - i\epsilon} \right] \times (ie\gamma^\mu) u(p) \varepsilon_\mu^*(\vec{k}, \lambda)$
	$i\mathcal{M} = \varepsilon_\nu^*(\vec{k}, \lambda) \bar{u}(p') (ie\gamma^\nu) \left[\frac{-i(-(\not{p}' - \not{k}) + m)}{(p-k')^2 + m_e^2 - i\epsilon} \right] \times (ie\gamma^\mu) u(p) \varepsilon_\mu(\vec{k}', \lambda')$
$e^+(p')e^-(p) \rightarrow \gamma(k)\gamma(k')$	(Annihilation process)
	$i\mathcal{M} = \varepsilon_\nu(\vec{k}', \lambda') \bar{v}(p') (ie\gamma^\nu) \left[\frac{-i(-(\not{p} - \not{k}) + m)}{(p-k)^2 + m_e^2 - i\epsilon} \right] \times (ie\gamma^\mu) u(p) \varepsilon_\mu(\vec{k}, \lambda)$
	$i\mathcal{M} = \varepsilon_\nu(\vec{k}, \lambda) \bar{v}(p') (ie\gamma^\nu) \left[\frac{-i(-(\not{k} - \not{p}') + m)}{(p-k')^2 + m_e^2 - i\epsilon} \right] \times (ie\gamma^\mu) u(p) \varepsilon_\mu(\vec{k}', \lambda')$

Since we will be computing un-polarized (average over initial spin/polarizations), inclusive (sum over final spin/polarizations) cross-sections, we have to evaluate $\frac{1}{2} \sum_{\sigma, \sigma'} |\mathcal{M}|^2$ for the fermion scattering and likewise for the other cases. For the Compton scattering and pair annihilation where two diagrams contribute, we need to add them *before* taking the mod-square.

A. Electron-muon processes:

Consider the electron-muon scattering. We have,

$$(i\mathcal{M})(-i\mathcal{M}^*) = \frac{e^4}{(q^2 - i\epsilon)(q^2 + i\epsilon)} \{ (\bar{u}(p')\gamma^\mu u(p)) (u^\dagger(p)(\gamma^\nu)^\dagger (\gamma^0)^\dagger u(p')) \} \\ \times \{ (\bar{u}(k')\gamma_\mu u(k)) (u^\dagger(k)(\gamma_\nu)^\dagger (\gamma^0)^\dagger u(k')) \}$$

The spins, $\sigma, \sigma', \rho, \rho'$ are implicit in the above expression. Summing over all of these replaces $u\bar{u}$ using the completeness relation. Each of the braces can be expressed as traces over the

gamma matrices. With the new normalizations used, the completeness relations take the form,

$$\boxed{\sum_{\sigma} u(p, \sigma) \bar{u}(p, \sigma) = -\not{p} + m, \quad \sum_{\sigma} v(p, \sigma) \bar{v}(p, \sigma) = -\not{p} - m.} \quad (14.1)$$

Thus the electron and muon spin sums become,

$$\{\dots\}_{p,p'} = Tr[\gamma^{\mu}(-\not{p} + m_e)\gamma^{\nu}(-\not{p}' + m_e)] \quad , \quad \{\dots\}_{k,k'} = Tr[\gamma_{\mu}(-\not{k} + m_{\mu})\gamma_{\nu}(-\not{k}' + m_{\mu})] .$$

This leads to,

$$\begin{aligned} \frac{1}{4} \sum_{\sigma, \sigma', \rho, \rho'} |\mathcal{M}|_{e\mu \rightarrow e\mu}^2 &= \frac{1}{4} \frac{e^4}{(q^2 - i\epsilon)(q^2 + i\epsilon)} Tr[\gamma^{\mu}(-\not{p} + m_e)\gamma^{\nu}(-\not{p}' + m_e)] \\ &\quad \times Tr[\gamma_{\mu}(-\not{k} + m_{\mu})\gamma_{\nu}(-\not{k}' + m_{\mu})] \end{aligned} \quad (14.2)$$

For the $e^+e^- \rightarrow \mu^+\mu^-$ we will have,

$$\begin{aligned} (i\mathcal{M})(-i\mathcal{M}^*) &= \frac{e^4}{(q^2 - i\epsilon)(q^2 + i\epsilon)} \{(\bar{v}(p')\gamma^{\mu}u(p))(u^{\dagger}(p)(\gamma^{\nu})^{\dagger}(\gamma^0)^{\dagger}v(p'))\} \\ &\quad \times \{(\bar{u}(k')\gamma_{\mu}v(k))(v^{\dagger}(k)(\gamma_{\nu})^{\dagger}(\gamma^0)^{\dagger}u(k'))\} \end{aligned}$$

This is the same as before except the p', k' spinors are v spinors. Using the corresponding completeness relation we get,

$$\begin{aligned} \frac{1}{4} \sum_{\sigma, \sigma', \rho, \rho'} |\mathcal{M}|_{e^+e^- \rightarrow \mu^+\mu^-}^2 &= \frac{1}{4} \frac{e^4}{(q^2 - i\epsilon)(q^2 + i\epsilon)} Tr[\gamma^{\mu}(-\not{p} + m_e)\gamma^{\nu}(-\not{p}' - m_e)] \\ &\quad \times Tr[\gamma_{\mu}(-\not{k} + m_{\mu})\gamma_{\nu}(-\not{k}' - m_{\mu})] \end{aligned} \quad (14.3)$$

The Compton scattering and annihilation processes have two diagrams each and each of these has a *single* Dirac trace. But each diagram also has photon polarization completeness relations. Incidentally, for Moller and Bhabha processes, there are ‘crossed diagrams’ too. We will do these later.

To proceed with the two processes above, we need to traces of 2 and 4 gamma matrices. Here are the relevant formulae.

$$Tr\mathbf{1} = 4; \quad (14.4)$$

$$Tr\gamma^{\mu}\gamma^{\nu} = -4\eta^{\mu\nu}; \quad (14.5)$$

$$Tr\gamma^{\mu}\gamma^{\nu}\gamma^{\alpha}\gamma^{\beta} = 4(\eta^{\mu\nu}\eta^{\alpha\beta} - \eta^{\mu\alpha}\eta^{\nu\beta} + \eta^{\mu\beta}\eta^{\nu\alpha}) \quad (14.6)$$

Using these, we get

$$\begin{aligned}
Tr[\gamma^\mu(-\not{p} + m_e)\gamma^\nu(-\not{p}' + m_e)] &= p_\alpha p'_\beta Tr[\gamma^\mu\gamma^\alpha\gamma^\nu\gamma^\beta] + m_e^2 Tr[\gamma^\mu\gamma^\nu] \\
&= 4(p^\mu p'^\nu - \eta^{\mu\nu} p \cdot p' + p^\nu p'^\mu) - 4m_e^2 \eta^{\mu\nu} \quad \text{and,} \\
Tr[\gamma_\mu(-\not{k} + m_\mu)\gamma_\nu(-\not{k}' + m_\mu)] &= k_\alpha k'_\beta Tr[\gamma_\mu\gamma^\alpha\gamma_\nu\gamma^\beta] + m_\mu^2 Tr[\gamma_\mu\gamma_\nu] \\
&= 4(k_\mu k'_\nu - \eta_{\mu\nu} k \cdot k' + k_\nu k'_\mu) - 4m_\mu^2 \eta_{\mu\nu}.
\end{aligned}$$

Dotting the traces gives,

$$\begin{aligned}
Tr[\dots]Tr[\dots] &= 32[(p \cdot k)(p' \cdot k') + (p \cdot k')(p' \cdot k) + m_\mu^2 p \cdot p' + m_e^2 k \cdot k' + 2m_e^2 m_\mu^2] \\
\therefore \frac{1}{4} \sum_{spins} |\mathcal{M}|_{e\mu \rightarrow e\mu}^2 &= \frac{8e^4}{q^4} [(p \cdot k)(p' \cdot k') + (p \cdot k')(p' \cdot k) + m_\mu^2 p \cdot p' + m_e^2 k \cdot k' + 2m_e^2 m_\mu^2] \quad [14.7]
\end{aligned}$$

where $q = p' - p$.

For the $e^+e^- \rightarrow \mu^+\mu^-$ process, the only difference is m_e^2, m_μ^2 changing signs. Thus we have,

$$\frac{1}{4} \sum_{spins} |\mathcal{M}|_{e^+e^- \rightarrow \mu^+\mu^-}^2 = \frac{8e^4}{q^4} [(p \cdot k)(p' \cdot k') + (p \cdot k')(p' \cdot k) - m_\mu^2 p \cdot p' - m_e^2 k \cdot k' + 2m_e^2 m_\mu^2] \quad (14.8)$$

where $q = p + p'$.

To relate to cross-section, we can use the center of mass frame. For the electron-muon scattering, we may choose

$$p^\mu = (E_e, p\hat{z}), \quad k^\mu = (E_\mu, -p\hat{z}), \quad p'^\mu = (E'_e, \vec{p}'), \quad k'^\mu = (E'_\mu, -\vec{p}'), \quad \cos(\Theta_{cm}) := \hat{z} \cdot \hat{p}'.$$

Notice that $E_e + E_\mu = \sqrt{p^2 + m_e^2} + \sqrt{p^2 + m_\mu^2} = E'_e + E'_\mu = \sqrt{(p')^2 + m_e^2} + \sqrt{(p')^2 + m_\mu^2}$, implies that $|p'| = |p|$. The magnitude of the center of mass momentum and the scattering angle are the only two independent parameters given the electron and muon masses.

For $e^+e^- \rightarrow \mu^+\mu^-$, a good deal of convenience ensues. The equality of magnitudes of momenta and masses being identical in both initial and final states, implies that the energies of individual particles in initial and final state are equal and total energy conservation implies these energies are equal too. The initial and final momenta magnitudes are then simply related by the masses. Thus we may choose,

$$p^\mu = (E, p\hat{z}), \quad (p')^\mu = (E, -p\hat{z}), \quad k^\mu = (E, \vec{k}), \quad (k')^\mu = (E, -\vec{k}), \quad \cos(\Theta_{cm}) := \hat{z} \cdot \hat{k}.$$

Again only the center of mass energy and the scattering angle are the only independent parameters given the masses.

It remains to simplify the expression for the cross-sections. We will give it for the muon production process and leave the $e\mu$ scattering process as an exercise. We will also neglect the electron mass compared to the muon mass.

The dot products of momenta, in the center of mass take the form, $q^2 = (p + p')^2 = -(2E)^2 = -4(k^2 + m_e^2) \approx -4k^2$. Next, $p \cdot p' = -E^2 - p^2 \approx -2E^2$ and

$$p \cdot k = p' \cdot k' = -E^2 + Ekc\cos(\Theta_{cm}) \quad , \quad p \cdot k' = p' \cdot k = -E^2 - Ekc\cos(\Theta_{cm}) \quad .$$

Substituting in the invariant amplitude gives ($m_e = 0$ set),

$$\begin{aligned} \frac{1}{4} \sum_{spins} |\mathcal{M}|_{e^+e^- \rightarrow \mu^+\mu^-}^2 &= \frac{8e^4}{(-4E^2)^2} \left\{ (E^2 + Ekc\cos(\Theta_{cm}))^2 + (E^2 - Ekc\cos(\Theta_{cm}))^2 - m_\mu^2(-2E^2) \right\} \\ &= e^4 \left\{ 1 + \frac{k^2 \cos^2(\Theta_{cm})}{E^2} + \frac{m_\mu^2}{E^2} \right\} \\ &= e^4 \left[\left(1 + \frac{m_\mu^2}{E^2}\right) + \left(1 - \frac{m_\mu^2}{E^2}\right) \cos^2(\Theta_{cm}) \right] \end{aligned} \quad (14.9)$$

The differential cross-section is given by (11.21),

$$\begin{aligned} \frac{d\sigma}{d\Omega_{cm}} &= \frac{1}{64\pi^2} \frac{1}{s} \frac{k'_{cm}}{k_{cm}} \left(\frac{1}{4} \sum_{spins} |\mathcal{M}|^2 \right) \quad \text{where, } s = 4E^2, m_e = 0, m'_1 = m'_2 = m_\mu \Rightarrow \\ &= \frac{1}{64\pi^2} \frac{1}{s} \sqrt{\frac{s^2 - 4sm_\mu^2}{s}} \left[\left(1 + \frac{m_\mu^2}{E^2}\right) + \left(1 - \frac{m_\mu^2}{E^2}\right) \cos^2(\Theta_{cm}) \right] \\ &= \left(\frac{e^2}{4\pi}\right)^2 \frac{1}{4 \cdot 4E^2} \sqrt{1 - \frac{m_\mu^2}{E^2}} \left[\left(1 + \frac{m_\mu^2}{E^2}\right) + \left(1 - \frac{m_\mu^2}{E^2}\right) \cos^2(\Theta_{cm}) \right] \end{aligned} \quad (14.10)$$

Putting $\alpha := \frac{e^2}{4\pi}$, the fine structure constant and $E_{cm} = 2E$, we write the *unpolarized*, $e^+e^- \rightarrow \mu^+\mu^-$ cross-section as,

$$\frac{d\sigma}{d\Omega_{cm}}(E, \Theta_{cm}) = \frac{\alpha^2}{4E_{cm}^2} \sqrt{1 - \frac{m_\mu^2}{E^2}} \left[\left(1 + \frac{m_\mu^2}{E^2}\right) + \left(1 - \frac{m_\mu^2}{E^2}\right) \cos^2(\Theta_{cm}) \right] \quad (14.11)$$

$$\sigma_{total} := \int d\Omega \frac{d\sigma}{d\Omega} = \frac{\pi\alpha^2}{3E_{cm}^2} \sqrt{1 - \frac{m_\mu^2}{E^2}} \left(1 + \frac{m_\mu^2}{2E^2}\right) \quad (14.12)$$

Remarks: The square root factor shows that $E \geq m_\mu$ must hold for the muon production to proceed. In the ultra-relativistic limit, $E \gg m_\mu$, the differential cross-section has the characteristic $(1 + \cos^2(\Theta))$ angular dependence.

Since the cross-section is a measured quantity and the center of mass energy is under our control, by comparing the pair production cross-sections for two different final states, eg μ, τ , and taking the ratio, we can obtain bounds on the masses of the heavier leptons.

Exercises: Obtain the differential and the total cross-section for $e\mu \rightarrow e\mu$ process. Also for the Moller and the Bhabha processes.

B. Compton Scattering:

Recall the total amplitude from the two diagrams (we take the photon polarizations to be real),

$$i\mathcal{M} = \varepsilon_\nu(\vec{k}', \lambda') \bar{u}(p') (ie\gamma^\nu) \left[\frac{-i(-(\not{p}' + \not{k}) + m)}{(p+k)^2 + m_e^2 - i\epsilon} \right] (ie\gamma^\mu) u(p) \varepsilon_\mu(\vec{k}, \lambda) \\ + \varepsilon_\nu(\vec{k}, \lambda) \bar{u}(p') (ie\gamma^\nu) \left[\frac{-i(-(\not{p}' - \not{k}) + m)}{(p'-k)^2 + m_e^2 - i\epsilon} \right] (ie\gamma^\mu) u(p) \varepsilon_\mu(\vec{k}', \lambda') \quad (14.13)$$

$$\mathcal{M} = e^2 \varepsilon_\mu(\vec{k}, \lambda) \varepsilon_\nu(\vec{k}', \lambda') \bar{u}(p') \left[\frac{\gamma^\nu(-(\not{p}' + \not{k}) + m)\gamma^\mu}{(p+k)^2 + m_e^2 - i\epsilon} + \frac{\gamma^\mu(-(\not{p}' - \not{k}) + m)\gamma^\nu}{(p'-k)^2 + m_e^2 - i\epsilon} \right] u(p) \quad (14.14)$$

In the second (crossed diagram) terms, we have used $p-k' = p'-k$ which is more convenient.

Simplification: Since the momenta are on shell, $p^2 = -m^2, k^2 = 0 = (k')^2$, we get

$$(p+k)^2 + m^2 = p^2 + m^2 + k^2 + 2p \cdot k = 2p \cdot k \quad , \quad (p'-k)^2 + m^2 = -2p' \cdot k;$$

$$(-\not{p}' + m)\gamma^\mu u(p) = (2p^\mu + \gamma^\mu(\not{p}' + m))u(p) = 2p^\mu u(p) \quad ,$$

$$\bar{u}(p')\gamma^\mu(-\not{p}' + m) = \bar{u}(p')(2(p')^\mu + (\not{p}' + m))\gamma^\mu = 2(p')^\mu \bar{u}(p') \quad .$$

$$\mathcal{M} = e^2 \varepsilon_\mu(\vec{k}, \lambda) \varepsilon_\nu(\vec{k}', \lambda') \bar{u}(p') \left[\frac{\gamma^\nu(2p^\mu - \not{k}\gamma^\mu)}{2p \cdot k - i\epsilon} + \frac{(2(p')^\mu + \gamma^\mu \not{k})\gamma^\nu}{-2p' \cdot k - i\epsilon} \right] u(p) \quad (14.15)$$

$$\mathcal{M}^\dagger = e^2 \varepsilon_\alpha(\vec{k}, \lambda) \varepsilon_\beta(\vec{k}', \lambda') \bar{u}(p) \left[\frac{(2p^\alpha - \gamma^\alpha \not{k})\gamma^\beta}{2p \cdot k + i\epsilon} + \frac{\gamma^\beta(2(p')^\alpha + \not{k}\gamma^\alpha)}{-2p' \cdot k + i\epsilon} \right] u(p') \quad (14.16)$$

In the second equation, we have used

$$[\bar{u}(p')\Gamma u(p)]^* = \bar{u}(p)(\gamma^0)\Gamma^\dagger\gamma^0 u(p') = \bar{u}(p)\Gamma_{reversed}u(p').$$

Here $\Gamma_{reversed}$ has the order of the gamma matrices reversed due to the \dagger operation.

Summing and averaging over the spins and polarizations the $|\mathcal{M}|^2$, we get,

$$\frac{1}{4} \sum_{\sigma, \sigma', \lambda, \lambda'} |\mathcal{M}|^2 = \frac{e^4}{4} \left[Tr \left\{ \left(\frac{\gamma^\nu(2p^\mu - \not{k}\gamma^\mu)}{2p \cdot k - i\epsilon} + \frac{(2(p')^\mu + \gamma^\mu \not{k})\gamma^\nu}{-2p' \cdot k - i\epsilon} \right) (-\not{p}' + m) \right. \right. \\ \left. \left. \left(\frac{(2p^\alpha - \gamma^\alpha \not{k})\gamma^\beta}{2p \cdot k + i\epsilon} + \frac{\gamma^\beta(2(p')^\alpha + \not{k}\gamma^\alpha)}{-2p' \cdot k + i\epsilon} \right) (-\not{p}' + m) \right\} \times \right. \\ \left. \left\{ \sum_\lambda \varepsilon_\mu(\vec{k}, \lambda) \varepsilon_\alpha(\vec{k}, \lambda) \right\} \left\{ \sum_{\lambda'} \varepsilon_\nu(\vec{k}', \lambda') \varepsilon_\beta(\vec{k}', \lambda') \right\} \right] \quad (14.17)$$

We have used the completeness relation (14.1) to convert the spin sum. We have a trace over strings of γ matrices, but now we also have sums over the photon polarizations. An explicit expression can be obtained by introducing an explicit set of 4 orthonormal vectors.

Consider the polarization sum. Given a $k, k^2 = 0$ we can write $k^\mu = (|\vec{k}|, \vec{k})$. Introduce another vector $\tilde{k} := C(|\vec{k}|, -\vec{k})$ so that $\tilde{k}^2 = 0$. Fix C by demanding $k \cdot \tilde{k} = -2 \leftrightarrow C = 1/|\vec{k}|^2$. We have two transverse directions and we take polarizations $\varepsilon(\vec{k}, \lambda)$ along these two directions and mutually orthonormalized. These are space-like vectors and have no time component. These 4 vectors, $k, \tilde{k}, \varepsilon(\vec{k}, 1), \varepsilon(\vec{k}, 2)$ are independent and the completeness relation takes the form,

$$-\frac{k^\mu \tilde{k}^\nu + k^\nu \tilde{k}^\mu}{2} + \varepsilon^\mu(\vec{k}, 1)\varepsilon^\nu(\vec{k}, 1) + \varepsilon^\mu(\vec{k}, 2)\varepsilon^\nu(\vec{k}, 2) = \eta^{\mu\nu}$$

Thus we get the sum over photon polarizations as,

$$\sum_{\lambda=1,2} \varepsilon_\mu(\vec{k}, \lambda)\varepsilon_\nu(\vec{k}, \lambda) = \eta_{\mu\nu} + \frac{k_\mu \tilde{k}_\nu + k_\nu \tilde{k}_\mu}{2} \quad (14.18)$$

Claim: The terms containing k, \tilde{k} in the polarization sum, do not contribute to the $\sum |\mathcal{M}|^2$.

Proof: Observe that we can always express $\mathcal{M} = \varepsilon_\mu(\vec{k}, \lambda)M^\mu$, by just taking out the ε_μ . Then $\sum_\lambda |\mathcal{M}|^2 = M^\mu (M^*)^\nu \{\eta_{\mu\nu} + \frac{1}{2}(k_\mu \tilde{k}_\nu + k_\nu \tilde{k}_\mu)\}$. The k terms contracting with M can be seen by replacing $\varepsilon_\mu \rightarrow k_\mu$. It is more convenient to write the Fermion propagators using, $\frac{-\not{q}+m}{q^2+m^2-i\epsilon} = \frac{1}{\not{q}+m}$ since $(-\not{q}+m)(\not{q}+m) = q^2+m^2$. Then,

$$\begin{aligned} M|_{\varepsilon_\mu=k_\mu} &= e^2 \bar{u}(p') \left[\not{\varepsilon}(k') \frac{1}{\not{p} + \not{k} + m} \{\not{k} + \not{p} + m - \not{p} - m\} \right. \\ &\quad \left. + \{\not{k} - \not{p}' - m + \not{p}' + m\} \frac{1}{\not{p}' - \not{k} + m} \not{\varepsilon}(k') \right] u(p) \\ &= e^2 \left[\bar{u}(p') \not{\varepsilon}(k') u(p) - \bar{u}(p') \not{\varepsilon}(k') \frac{1}{\not{p} + \not{k} + m} (\not{p} + m) \right] u(p) \\ &\quad \left[\bar{u}(p') (-\not{\varepsilon}(k')) u(p) + \bar{u}(p') (\not{p}' + m) \frac{1}{\not{p}' - \not{k} + m} \not{\varepsilon}(k') u(p) \right] = 0! \end{aligned}$$

We have used $p - k' = p' - k$ in the second term in the first equation. In the last equation, the first and the third terms cancel while the second and the fourth terms vanish by equation of motion. This proves the claim. Hence, *effectively, each of the polarization sums give only the $\eta_{\mu\nu}$.*

Note: The *total scattering amplitude* vanishes when the photon transverse polarization is replaced by a longitudinal one, is a general result known as a ‘Ward identity’. This is a

consequence of gauge invariance. While we do not discuss the general proof here, it suffices to note that (a) we used the Dirac equation: $(\not{p} + m)u(p) = 0 = \bar{u}(p')(\not{p}' + m)$; (b) the QED coupling has the form $\sim A_\mu J^\mu$ with $J^\mu \sim \bar{\Psi}\gamma^\mu\Psi \sim \bar{u}(p')\gamma^\mu u(p)$; (c) $\delta A_\mu = \partial_\mu\Lambda$ is a gauge transformation ($\leftrightarrow \varepsilon_\mu(k) \rightarrow \varepsilon_\mu(k) + k_\mu\lambda(k)$) and gauge invariance of the interaction implies $\partial_\mu J^\mu = 0$. Thus, the vanishing of the amplitude is related to gauge invariance.

The averaged $|\mathcal{M}|^2$ in equation (14.17) becomes,

$$\begin{aligned}
\sum_{\sigma,\sigma',\lambda,\lambda'} |\mathcal{M}|^2 &= e^4 \left[Tr \left\{ \left(\frac{\gamma^\nu(2p^\mu - \not{k}\gamma^\mu)}{2p \cdot k - i\epsilon} + \frac{(2(p')^\mu + \gamma^\mu \not{k})\gamma^\nu}{-2p' \cdot k - i\epsilon} \right) (-\not{p} + m) \right. \right. \\
&\quad \left. \left. \left(\frac{(2p_\mu - \gamma_\mu \not{k})\gamma_\nu}{2p \cdot k + i\epsilon} + \frac{\gamma_\nu(2(p')_\mu + \not{k}\gamma_\mu)}{-2p' \cdot k + i\epsilon} \right) (-\not{p}' + m) \right\} \right] \quad (14.19) \\
&= e^4 \left[Tr \left\{ \frac{(2p^\mu\gamma^\nu - \gamma^\nu \not{k}\gamma^\mu)\not{p}(2p_\mu\gamma_\nu - \gamma_\mu \not{k}\gamma_\nu)\not{p}'}{(2p \cdot k)^2} \right\} \textcircled{1} + \right. \\
&\quad Tr \left\{ \frac{(2(p')^\mu\gamma^\nu + \gamma^\mu \not{k}\gamma^\nu)\not{p}(2p'_\mu\gamma_\nu + \gamma_\nu \not{k}\gamma_\mu)\not{p}'}{(-2p' \cdot k)^2} \right\} \textcircled{2} + \\
&\quad Tr \left\{ \frac{(2p^\mu\gamma^\nu - \gamma^\nu \not{k}\gamma^\mu)\not{p}(2p'_\mu\gamma_\nu + \gamma_\nu \not{k}\gamma_\mu)\not{p}'}{-4p \cdot k p' \cdot k} \right\} \textcircled{3} + \\
&\quad \left. Tr \left\{ \frac{(2(p')^\mu\gamma^\nu + \gamma^\mu \not{k}\gamma^\nu)\not{p}(2p_\mu\gamma_\nu - \gamma_\mu \not{k}\gamma_\nu)\not{p}'}{-4p \cdot k p' \cdot k} \right\} \textcircled{4} \right] \\
&+ m^2 e^4 \left[Tr \left\{ \frac{(2p^\mu\gamma^\nu - \gamma^\nu \not{k}\gamma^\mu)(2p_\mu\gamma_\nu - \gamma_\mu \not{k}\gamma_\nu)}{(2p \cdot k)^2} \right\} \textcircled{5} + \right. \\
&\quad Tr \left\{ \frac{(2(p')^\mu\gamma^\nu + \gamma^\mu \not{k}\gamma^\nu)(2p'_\mu\gamma_\nu + \gamma_\nu \not{k}\gamma_\mu)}{(-2p' \cdot k)^2} \right\} \textcircled{6} + \\
&\quad Tr \left\{ \frac{(2p^\mu\gamma^\nu - \gamma^\nu \not{k}\gamma^\mu)(2p'_\mu\gamma_\nu + \gamma_\nu \not{k}\gamma_\mu)}{-4p \cdot k p' \cdot k} \right\} \textcircled{7} + \\
&\quad \left. Tr \left\{ \frac{(2(p')^\mu\gamma^\nu + \gamma^\mu \not{k}\gamma^\nu)(2p_\mu\gamma_\nu - \gamma_\mu \not{k}\gamma_\nu)}{-4p \cdot k p' \cdot k} \right\} \textcircled{8} \right] \quad (14.20)
\end{aligned}$$

The terms linear on m have odd number of γ 's and hence vanish under trace.

It may be checked easily that the $\textcircled{2}$ is obtained from $\textcircled{1}$ by $p \leftrightarrow -p'$ and likewise for $5 \leftrightarrow 6$. Noting the identity $Tr(\gamma_1\gamma_2 \cdots \gamma_n) = Tr(\gamma_n\gamma_{n-1} \cdots \gamma_2\gamma_1)$, it follows that $3 = 4$ and $7 = 8$. The first group of 4 terms involves trace of a maximum of 8 gamma matrices while the last 4 terms involve a maximum of 6 gamma's. These are simplified using various identities among the gamma matrices.

Consider (1). We have the traces,

$$Tr \{ 4p^2 \gamma^\nu \not{p} \gamma_\nu \not{p}' - 2\gamma^\nu \not{p} \not{p}' \gamma_\nu \not{p}' - 2\gamma^\nu \not{k} \not{p}' \gamma_\nu \not{p}' + \gamma^\nu \not{k} \gamma^\mu \not{p}' \gamma_\mu \not{k} \gamma_\nu \not{p}' \}$$

Use: $\gamma^\nu \not{p} \gamma_\nu = +2\not{p}$, $\not{p} \not{p} = -p^2$, $\not{k} \not{p} + \not{p} \not{k} = -2p \cdot k$, $Tr \not{p} \not{p}' = -4p \cdot p'$ and the cyclic property of the trace to get the above trace as,

$$\begin{aligned} Tr\{\dots\}_{(1)} &= (4p^2)(2)(-4p \cdot p') - 2p^2(2)(-4k \cdot p') \times 2 + (2)(2)Tr(\not{k} \not{p}' \not{k} \not{p}') \\ &= 32m^2(p \cdot p' - k \cdot p') + 32(k \cdot p)(k \cdot p') \end{aligned} \quad (14.21)$$

We have used $p^2 = -m^2$ and $k^2 = 0$.

Similarly we get the other traces (without the denominator factors) as,

$$\begin{aligned} Tr\{\dots\}_{(3)} &= 4p \cdot p' Tr(\gamma^\nu \not{p}' \gamma_\nu \not{p}) - 2Tr(\not{k} \not{p}' \not{p}' \gamma_\nu \not{p}' \gamma^\nu) + 2Tr(\gamma^\nu \not{p}' \gamma_\nu \not{k} \not{p}' \not{p}') \\ &\quad - Tr(\gamma^\nu \not{k} \gamma^\mu \not{p}' \gamma_\nu \not{k} \gamma_\mu \not{p}') \end{aligned} \quad (14.22)$$

$$= -32(p \cdot p')^2 + (k \cdot p - k \cdot p')(16m^2 + 32p \cdot p') + 0 \quad (14.23)$$

$$\begin{aligned} Tr\{\dots\}_{(5)} &= 4p^2 Tr(\gamma^\nu \gamma_\nu) - 2Tr(\not{k} \not{p}'(-4\mathbb{1})) + 8Tr(\not{p}' \not{k}) + 16Tr(\not{k} \not{k}) \\ &= 64m^2 - 64k \cdot p \end{aligned} \quad (14.24)$$

$$\begin{aligned} Tr\{\dots\}_{(7)} &= -64p \cdot p' + 32(k \cdot p - k \cdot p') - Tr(\gamma^\nu \not{k} \gamma^\mu \gamma_\nu \not{k} \gamma_\mu) \\ &= -64p \cdot p' + 32(k \cdot p - k \cdot p') + 0 \end{aligned} \quad (14.25)$$

To simplify the expression further, we need to use: $p \cdot p' = -m^2 + k \cdot p - k \cdot p'$ which follows from $0 = (k')^2 = (p + k - p')^2$. For comparison with [13], it is useful to note: $k \cdot p = k' \cdot p'$, $p \cdot k' = p' \cdot k$.

The final expression is,

$$\frac{1}{4} \sum_{spin/Pol} |\mathcal{M}|^2 = 2e^4 \left[\left(\frac{p \cdot k'}{p \cdot k} + \frac{p \cdot k}{p \cdot k'} \right) + 2m^2 \left(\frac{1}{p \cdot k'} - \frac{1}{p \cdot k} \right) + m^4 \left(\frac{1}{p \cdot k} - \frac{1}{p \cdot k'} \right)^2 \right] \quad (14.26)$$

Exercise: Check the algebra!

The Compton cross-section is usually presented in the lab frame with the electron initially at rest, i.e. $p^\mu = (m, \vec{0})$, $k^\mu = (\omega, \omega \hat{z})$, $(p')^\mu = (E', \vec{p}')$, $(k')^\mu = (\omega', \omega' \sin \theta, 0, \omega' \cos(\theta))$. The final electron and photon momenta define a plane which is taken to be the $z-x$ plane with the final photon making an angle θ to the z -axis. These choices give: $p \cdot k' = -m\omega'$, $p \cdot k = -m\omega$

and,

$$\sum_{spin/Pol} |\mathcal{M}|^2 = 2e^4 \left[\frac{\omega'}{\omega} + \frac{\omega}{\omega'} + 2m \left(-\frac{1}{\omega'} + \frac{1}{\omega} \right) + m^2 \left(-\frac{1}{\omega} + \frac{1}{\omega'} \right)^2 \right] \quad (14.27)$$

Next, $(p')^2 = (p + k - k')^2$ implies,

$$-m^2 = -m^2 + 2p \cdot (k - k') + (k - k')^2 \Rightarrow 0 = m(\omega - \omega') - \omega\omega'(1 - \cos(\theta)),$$

leading to the Compton formula for shift in the photon wavelength with the scattering angle:

$$\text{Compton Formula: } \boxed{\frac{1}{\omega'} - \frac{1}{\omega} = \frac{1 - \cos\theta}{m}} \leftrightarrow \omega' = \frac{\omega}{1 + \frac{\omega}{m}(1 - \cos\theta)}. \quad (14.28)$$

To get the differential cross-section, we simplify the phase space integral as,

$$\int \frac{d^3k'}{2\omega'(2\pi)^3} \int \frac{d^3p'}{2E'(2\pi)^3} (2\pi)^4 \delta^4(p; +k' - p - k) = \int \frac{d\omega'(\omega')^2 d\Omega}{2\omega'(2\pi)^3} \frac{1}{2E'(2\pi)^3} (2\pi)^4 \delta(\omega' + E' - m - \omega)$$

We have used up a δ^3 to get $\vec{p}' = \vec{k}' - \vec{k} - \vec{p} = (\omega' \sin\theta, 0, \omega' \cos\theta - \omega)$ which gives $(E')^2 - (\omega')^2 + \omega^2 - 2\omega'\omega \cos\theta + m^2$. Hence

$$\int d\omega' \frac{\omega'}{4E'(\omega')} \delta(\omega' + E'(\omega') - m - \omega) = \frac{\omega'_*}{4E'(\omega'_*)} \frac{1}{|1 + \frac{dE'}{d\omega'_*}|}, \quad \frac{dE'}{d\omega'} = \frac{\omega' - \omega \cos\theta}{E'}.$$

Inserting in the phase space integral,

$$\begin{aligned} \int_{\Gamma_2} &= \int \frac{d\cos\theta}{2\pi} \frac{\omega'_*}{4E'_*} \frac{1}{|1 + \frac{\omega' - \omega \cos\theta}{E'}|_*}, \quad \omega'_* + E'(\omega'_*) = m + \omega \\ &= \frac{1}{8\pi} \int d\cos\theta \frac{\omega'^2}{m\omega}. \quad \text{We have,} \end{aligned} \quad (14.29)$$

$$\begin{aligned} \frac{d\sigma}{d\cos\theta} &= \frac{1}{4m\omega} \frac{1}{8\pi} \frac{(\omega')^2}{m\omega} [\sum |\mathcal{M}|^2] \\ &= \frac{e^4}{16\pi} \frac{1}{m^2} \left(\frac{\omega'}{\omega} \right)^2 \left[\frac{\omega'}{\omega} + \frac{\omega}{\omega'} + \left\{ 2m \frac{-1 + \cos\theta}{m} + m^2 \frac{(1 - \cos\theta)^2}{m^2} \right\} \right] \end{aligned}$$

This give the differential cross-section for the Compton scattering, known as *Klein-Nishina formula*,

$$\boxed{\left(\frac{d\sigma}{d\cos\theta_{lab}} \right) = \frac{\pi\alpha^2}{m^2} \left(\frac{\omega'}{\omega} \right)^2 \left[\frac{\omega'}{\omega} + \frac{\omega}{\omega'} - \sin^2\theta_{lab} \right], \text{ with } \frac{\omega'}{\omega} = \frac{1}{1 + \frac{\omega}{m}(1 - \cos\theta)}. \quad (14.30)}$$

Remarks:

(a) There has been no approximation at the α^2 level computation. It can now be used for various limits. At $\theta = 0$, forward scattering, there is no frequency change and the cross-section equaling $\frac{\pi\alpha^2}{m}$ is independent of the photon energy. For a *massive charged particle*, one defines its *Compton length* to be $\lambda_c := h/(mc) = 2\pi/m$ in the natural units, $\hbar = 1 = c$. The forward scattering cross-section is then $\alpha^2(\pi\lambda^2)$, where $\lambda := \lambda/(2\pi)$.

(b) There are two scales in the problems: m, ω . This gives two natural limits: (i) $\omega \rightarrow 0 \leftrightarrow \omega/m \ll 1$ and (ii) $\omega \rightarrow \infty \leftrightarrow \omega/m \gg 1$.

For a generic θ , consider $\omega \rightarrow 0$. Then $\omega/\omega' \rightarrow 1 + (\omega/m)(1 - \cos\theta)$ and

$$\left. \frac{d\sigma}{d\cos\theta} \right|_{\omega/m \ll 1} = \frac{\pi\alpha^2}{m^2} \left[1 + \cos^2\theta - 4\sin^2(\theta/2) \frac{\omega}{m} + \dots \right], \quad \sigma_{tot} = \frac{8\pi\alpha^2}{3m^2} + \dots \quad (14.31)$$

The leading term is the *Thomson cross-section* for classical radiation scattering off free electrons.

In the opposite limit, $m/\omega \ll 1$, we have

$$\omega'/\omega = \frac{m}{\omega} \frac{1}{1 - \cos\theta} \left(1 - \frac{m}{\omega} \frac{1}{1 - \cos\theta} + \dots \right) := \epsilon(1 - \epsilon \dots), \quad \epsilon(\theta) := \frac{m}{\omega} \frac{1}{1 - \cos\theta}.$$

In this limit,

$$\frac{d\sigma}{d\cos\theta} \approx \frac{\pi\alpha^2}{m^2} \epsilon(\theta) = \frac{\pi\alpha^2}{m^2} \frac{m}{\omega(1 - \cos\theta)}, \quad (14.32)$$

$$\sigma_{tot}(\delta) := \int_{-1}^{1-\delta} d\cos\theta \frac{d\sigma}{d\cos\theta} \approx \frac{\pi\alpha^2}{m\omega} [-\ln(\delta/2)] \rightarrow \infty, \quad \text{as } \theta \rightarrow 0. \quad (14.33)$$

For the $\epsilon(\theta) \ll 1$ condition to hold so that the expansion is meaningful, $\theta \gg \sqrt{2m/\omega}$ must hold. At $\theta = 0$ we know the exact answer which is finite.

C. Electron-Positron Annihilation:

Having evaluated the Compton scattering amplitude, this is actually simple to evaluate. From the table giving the amplitudes (notice the use of $p' - k$ and $k - p'$ momenta in the crossed diagrams), we can see that the annihilation diagrams expressions are obtained by making the substitutions: $\boxed{p \rightarrow p, k \rightarrow -k, p' \rightarrow -p', k' \rightarrow k' \text{ and } \bar{u} \rightarrow \bar{v}}$. These substitutions preserve the momentum conservations appropriate for the processes: $p + k = p' + k' \rightarrow p - k = -p' + k'$. The photon polarizations are insensitive to the ‘sign’ of momentum and will give $\eta_{\mu\nu}$ as before. One of the spin sums however changes from

$[-\not{p}' + m] \rightarrow [-(-\not{p}' + m)] = [\not{p} - m]$ which generates an *overall minus sign in the summed, squared amplitude*. This is actually an example of what is known as the *crossing symmetry* of the S -matrix: *A scattering amplitude for a particle with momentum k in the initial state is the same as the amplitude with the initial particle moved to the final state anti-particle with momentum $-k$.*

Without any explicit calculation we can write down the summed squared amplitude for the annihilation process as,

$$\sum_{spin/Pol} |\mathcal{M}|^2 = (-)2e^4 \left[- \left(\frac{p \cdot k'}{p \cdot k} + \frac{p \cdot k}{p \cdot k'} \right) + 2m^2 \left(\frac{1}{p \cdot k'} + \frac{1}{p \cdot k} \right) + m^4 \left(\frac{1}{p \cdot k} + \frac{1}{p \cdot k'} \right)^2 \right] \quad (14.34)$$

It is more convenient to express the cross-section in the center of mass frame:

$$p^\mu = (E, p\hat{z}) \quad , \quad (p')^\mu = (E, -p\hat{z}) \quad , \quad E^2 = p^2 + m^2 \quad , \quad s = -(p + p')^2 = 4E^2$$

$$k^\mu = (E, E\sin\Theta, 0, E\cos\Theta) \quad , \quad (k')^\mu = (E, -E\sin\Theta, 0, -E\sin\Theta)$$

This gives,

$$p \cdot k = -E^2 + pE\cos\Theta = -E(E - p\cos\Theta) \quad ,$$

$$p \cdot k' = -E^2 - pE\cos\Theta = -E(E + p\cos\Theta) \quad ,$$

$$\frac{|k'|_{cm}}{|k|_{cm}} = \sqrt{\frac{s^2 - 0 + 0^2}{s^2 - 2s(2m^2) + 0}} = \frac{1}{1 - m^2/E^2} = \frac{E}{\sqrt{E^2 - m^2}} = \frac{E}{p}$$

The differential cross-section takes the form,

$$\frac{d\sigma}{d\Omega_{cm}} = \frac{1}{64\pi^2} \frac{1}{4E^2} \frac{E}{|p|} \left(\sum_{spin/Pol} |\mathcal{M}|^2 \right) \quad \text{with}$$

$$\sum_{spin/Pol} |\mathcal{M}|^2 = (-2e^4) \left[- \left(\frac{E + p\cos\Theta}{E - p\cos\Theta} + \frac{E - p\cos\Theta}{E + p\cos\Theta} \right) \right] \quad (14.35)$$

$$+ \frac{2m^2}{-E} \left(\frac{1}{E + p\cos\Theta} + \frac{1}{E - p\cos\Theta} \right) \quad (14.36)$$

$$+ \frac{m^4}{E^2} \left(\frac{1}{E + p\cos\Theta} + \frac{1}{E - p\cos\Theta} \right)^2 \right]$$

$$= 4e^4 \left[\frac{E^2 + p^2\cos^2\Theta}{E^2 - p^2\cos^2\Theta} + \frac{2m^2}{E^2 - p^2\cos^2\Theta} - \frac{2m^4}{(E^2 - p^2\cos^2\Theta)^2} \right] \quad (14.37)$$

The differential cross-section for unpolarized, e^+e^- annihilation process is then given by,

$$\boxed{\frac{d\sigma}{d\Omega_{cm}} = \frac{\alpha^2 E}{s |p|} \left[\frac{E^2 + p^2 \cos^2 \Theta + 2m^2}{m^2 + p^2 \sin^2 \Theta} - \frac{2m^4}{(m^2 + p^2 \sin^2 \Theta)^2} \right]}, \quad s = 4E^2. \quad (14.38)$$

In the high energy limit, $E \gg m$, this reduces to,

$$\frac{d\sigma}{d\Omega_{cm}} \rightarrow \frac{\alpha^2}{s} \left(\frac{1 + \cos^2 \Theta}{\sin^2 \Theta} \right) \quad (14.39)$$

To obtain the total cross-section, we need to integrate over the final state. Since the two photons are identical, Θ needs to be integrated between $0, \pi/2$. The Bose factor of 2 (which we did not include in the amplitude) cancels the factor of 2 from the angular integration.

We get,

$$\sigma_{tot} = \frac{\alpha^2}{s} (2\pi) \int_0^1 dx \frac{1+x^2}{1-x^2} \quad (14.40)$$

The integral is clearly divergent from the ‘forward’ direction ($\Theta = 0$). This is artificial since we set $m = 0$ in the integrand. Split the integration as $\int_0^{1-\Delta} + \int_{1-\Delta}^1$. The first term is a finite number and this contribution falls off as s^{-1} . For the second term, we go back to the exact differential cross-section, approximate the integrand for $1 - \Delta \leq \cos(\Theta) \leq 1$. Then, the factors of E all cancel and second term $\sim \alpha^2(1 - \Delta)m^{-2}$ where, $\Delta \ll m^2/p^2$ so that the denominator can be approximated. Thus, the total cross-section does *not* diverge but is bounded by m^{-2} . This is an example of bounds satisfied by total cross-sections at very high energies.

15. NUMERICAL ESTIMATES OF CROSS-SECTIONS AND APPLICATIONS

We have obtained examples of cross-sections for some of the basic processes in quantum electrodynamics. We also saw how they relate to non-relativistic versions of the processes eg potential scattering and inferred the modification due to relativistic effects as well as estimate ‘strengths’ of interactions. Before we study the higher order corrections, it is useful to have a sense of order of magnitudes of cross-sections and how these numbers are used in applications. We will focus on the QED processes.

What is the value of the fine structure constant?

This is related to the electric charge, the Planck constant due to quantum framework and the speed of light due to special relativity. We first obtain the expression in terms of “engineering units” and then use the measured values to compute the fine structure constant.

Observe that the Feynman rules are derived from the Lagrangian density which has properly normalized kinetic (quadratic) terms, $-\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + i\bar{\Psi}\gamma^\mu\partial_\mu\Psi - m\bar{\Psi}\Psi$ and the interaction term introduces ‘ e ’ as $e\bar{\Psi}\gamma^\mu\Psi A_\mu$ and thus is dimensionless. This definition gives the equation of motion, $\partial_\mu F^{\mu\nu} = -e\bar{\Psi}\gamma^\nu\Psi \leftrightarrow \partial_i E^i = e\Psi^\dagger\Psi$, using the identifications: $F^{0i} := E^i, \partial_\mu F^{\mu\nu} = -J^\nu, J^\mu = (\rho, J^i)$. Furthermore, the Hamiltonian and hence the energy density is $\vec{E}^2/2$.

The MKSA (SI) units has the Gauss law in the form: $\vec{\nabla} \cdot \vec{E}_{SI} = \rho_{SI}/\epsilon_0$. The field energy density (as inferred from the Poynting theorem) is $\epsilon_0\vec{E}_{SI}^2/2$. Comparing the energy densities, we introduce the identification, $E := \sqrt{\epsilon_0}E_{SI}$. Then $\rho = \vec{\nabla} \cdot \vec{E} = \sqrt{\epsilon_0}\vec{\nabla} \cdot \vec{E}_{SI} := \sqrt{\epsilon_0}\frac{\rho_{SI}}{\epsilon_0}$. This leads to the identification, $e := \frac{e_{SI}}{\sqrt{\epsilon_0}}$. Thus, we identify the variables used in the action with those used in engineering variables by comparing the equations of motion *and* the energy densities.

In the natural units, $\hbar = 1 = c$ used in writing the action, the fine structure constant was defined as $\alpha := e^2/(4\pi)$ which is dimensionless as noted above. Substituting e in terms of e_{SI} and introducing \hbar, c , we write $\alpha := \frac{e_{SI}^2}{4\pi\epsilon_0}\hbar^a c^b$ and determine the powers a, b by requiring α to be dimensionless. For this, we need the dimensions of e_{SI} and ϵ_0 ! Now we also use the SI expression of the Lorentz force, $F = e_{SI}E_{SI}$. Hence, *dimensionally* (Force)²/(energy density) $\sim e_{SI}^2/\epsilon_0 \sim M^1L^3T^{-2}$. This and α being dimensionless gives $a = b = -1$ and $\alpha = \frac{e_{SI}^2}{4\pi\epsilon_0\hbar c} \simeq 1/137$, using the values: $(4\pi\epsilon_0)^{-1} = c^2 10^{-7}, e_{SI} = 1.6 \times 10^{-19}$ Coulomb, $\hbar \simeq 1.05 \times 10^{-34}$ Joules.sec and $c \simeq 3 \times 10^8$ m/sec.

An Aside: Another set of units are commonly used in gravitational physics, the so called *geometrized units*. These are defined by setting $G = c = 1$. The choice of $c = 1$ gives: 1 sec = 3×10^8 meters. Newton's constant in SI units is, $G \approx 6.67 \times 10^{-11} kg^{-1} meter^3 sec^{-2}$. Hence setting $G = 1$ along with $c = 1$, gives $1 kg = 7.4 \times 10^{-28}$ meters.

It is conventional in gravitational physics to use the Gauss law in the form $\vec{\nabla} \cdot \vec{E} = 4\pi\rho$ and the electric field energy density as $E^2/(8\pi)$. This gives the identifications: $E_{geom} = \sqrt{4\pi\epsilon_0}E_{SI}$, $q_{geom} = q_{SI}/\sqrt{4\pi\epsilon_0}$. As before, the Lorentz force equation gives the dimensions as: $[q_{SI}^2\epsilon_0] = [4\pi q_{geom}^2] = M^1 L^3 T^{-2} = L^2$.


Setting $q_{geom} = \frac{q_{SI}}{\sqrt{4\pi\epsilon_0}} G^\alpha c^\beta$ must have dimensions of L , we infer $\alpha = 1/2$ and $\beta = -2$ and $q_{geom} = \frac{q_{SI}}{\sqrt{4\pi\epsilon_0}} G^{1/2} c^{-2} = q_{SI} \sqrt{c^2 \times 10^{-7}} G^{1/2} c^{-2} \approx (8.6 \times 10^{-18}) q_{SI}$ meters.

Looking at the expressions for the cross-sections and noting that in the natural units energy \sim length $^{-1}$, we see that the cross-sections are of the form α/E^2 . This gets more and more accurate at ultra relativistic energies where masses can be neglected. Since the cross-section has dimensions of area while energies are typically given in MeV/GeV/TeV etc, it is useful to have a conversion between energy units and length units. As already notes, $c = 1 \Rightarrow 1 \text{ second} = 3 \times 10^8 \text{ meters}$. $\hbar = 1 \Rightarrow 1 \text{ Joule} = 10^{34} sec^{-1} \simeq 3.3 \times 10^{25} meter^{-1}$. Next, $100 MeV = 1.6 \times 10^{-19+8} J \simeq 5 \times 10^{14} meter^{-1}$. (This is also expressed as $200 MeV \cdot fermi = 1$). Clearly, a cross-section at 100 MeV (electron mass being 0.5 Mev) is approximately equal to $\sigma_{100Mev} \simeq \alpha^2/E^2 \simeq 2 \times 10^{-33} m^2$. The conventional unit for these cross-sections is $1 \text{ barn} := 100 \text{ fermi}^2 = 10^{-28} m^2$. Typical numbers encountered in high energy processes (around 1 GeV) dominated by strong interactions $\sim 10^{-30} m^2$, electromagnetic interactions $\sim 10^{-36} m^2$ and weak interactions $\sim 10^{-42} m^2$.

So, scattering experiments will provide us with such numbers. How are they *useful*?

In many applications, say passage of a collimated set of particles through some medium, we have the estimate of the cross-section of scattering of individual particles comprising the 'beam' and the medium. Intuitively, a cross-section may be viewed as a target disk of area σ which is bombarded by 'marbles' which get reflected from the disk. They are reflected if they hit the disk or pass by if not. Thus, σ gives the likelihood of a scattering interaction. Suppose we have multiple targets (medium) and multiple beam particles. The probability of interaction is clearly proportional to the ratio of the effective area of the target particles and

the area exposed to the beam Thus, probability of scattering = (number of target particles



exposed o the beam) $\times\sigma$ /Total area presented by the medium). Let n be the number of target particles exposed per unit area perpendicular to the beam. Then the number of particles exposed = $n \times$ total area and the probability of scattering = $n \cdot \sigma$.

Let ρ be the number density of the target particles i.e. number per unit volume. Let d be the thickness of the medium. Then the areal density $n = \rho \cdot d$. The probability of scattering is $1n\sigma = \rho \cdot d \cdot \sigma$ for $d = \hat{d} = (\rho \cdot \sigma)^{-1}$. Let $\langle v \rangle$ be the average speed of the beam particles. Then the average *reaction time* := $\tau_r := \hat{d}/\langle v \rangle = 1/(\rho\langle v \rangle\sigma)$.

Thus, if we have a confined plasma of say electrons and we bombard it with a beam of positrons and look for production of μ^\pm pairs, then we should expect to wait for a time $\tau = 1/(\rho\langle v \rangle\sigma_{e^+e^- \rightarrow \mu^+\mu^-})$. This type of use of microscopic cross-sections is required in discussing bulk processes eg in nuclear physics or astrophysics and are useful in estimating thermalization times.

Sometimes, we aren't interested in tracking individual processes, any interaction may suffice. In such cases we need to sum over *all possible final states*, $\sigma_{tot} = \sum_f \sigma_{i \rightarrow f}$. The $\sigma_{i \rightarrow f} \sim |\mathcal{M}_{i \rightarrow f}|^2$. We already have expressions using phase space integration for fixed number of final state particles. Now we have to sum over the number of particles and the labels,

$$\sigma_{tot} = \frac{1}{4E_k E_p} v_{lab} \sum_{f,\alpha} \int \prod_{i=1}^{N_f} \frac{d^3 k'_i}{2\omega_{k'_i} (2\pi)^3} (2\pi)^4 \delta^4(\Sigma k' - P_{in}) |\mathcal{M}|^2 .$$

This can be related to the forward scattering amplitude via the ‘‘Optical Theorem’’.

We have $S^\dagger S = \mathbb{1} \Rightarrow \sum_f \langle i | S^\dagger | f \rangle \langle S | i \rangle = 1 = \sum_f |S_{fi}|^2$. Substituting for S_{fi} we get,

$$|S_{fi}|^2 = \delta_{fi} \delta_{fi} + (2\pi)^4 \delta^4(P_f - P_i) \delta_{fi} [i(\mathcal{M}_{fi} - \mathcal{M}_{fi}^*)] + (2\pi)^8 [\delta^4(P_f - P_i)]^2 |\mathcal{M}_{fi}|^2 .$$

Summing over f and using unitarity gives,

$$(2\pi)^4 \sum_f \delta^4(P_f - P_i) \delta_{fi} [i(\mathcal{M}_{fi} - \mathcal{M}_{fi}^*)] = -(2\pi)^8 \sum_f \delta^4(P_f - P_i) \delta^4(P_f - P_i) |\mathcal{M}_{fi}|^2$$

Or,

$$\boxed{2Im(\mathcal{M}_{ii}) = (2\pi)^4 \sum_f \delta^4(P_f - P - i) |\mathcal{M}_{fi}|^2.} \quad (15.1)$$

Substituting the rhs in the expression for σ_{tot} gives,

$$\sigma_{tot} = \frac{2Im(\mathcal{M}_{ii})}{4k\sqrt{s}} \quad , \quad \text{where, } k = k_{CM} \text{ and } s = -(\sum P_i)^2 \quad .$$

The \mathcal{M}_{ii} is the forward scattering, $i \rightarrow i$, *amplitude* and σ_{tot} is the inclusive cross-section for $i \rightarrow$ any f processes. This could also be used to estimate the total cross-section.

Remark: Let us note that we defined the scattering matrix in the basis of number states or Fock basis. Further, we chose the idealization of plane waves for both in and out states. This is convenient but loses the information about *spatial* locations of the scattering particles. This also loses any notion of ‘‘impact parameter’’ - how far the projectile particle is from the target particle or equivalently the ‘orbital angular momentum’ of the projectile about the target particle. The information about the transverse position is of course contained in the parameters of the outgoing particles. This should be kept in mind while interpreting some of the divergences of cross-sections.

16. RADIATIVE CORRECTIONS IN QED

Consider now corrections due to higher powers of interaction Hamiltonian, also called *radiative corrections*. Having seen the correspondence between the correction terms and Feynman diagrams, we can list the corrections directly in terms of the diagrams. A Feynman diagram is made up of lines (free propagators) and vertices. In QED, there are two types of propagators and one type of vertex. So consider corrections to two point functions and the three point function.

A. The fermion propagator: self-energy

$$\begin{aligned}
 \text{Diagram with shaded blob '1PI'} &= \text{Free propagator} + \text{Propagator with photon loop} + \text{Propagator with photon loop and fermion loop} + \text{Propagator with photon loop and fermion loop and another photon loop} + \dots \\
 &= \text{Free propagator} + \text{Propagator with shaded '1PI' blob} + \text{Propagator with two shaded '1PI' blobs} + \dots
 \end{aligned}
 \tag{16.1}$$

Here is a graphical expansion of the exact two point function of the fermions. In the second line, the diagrams have been grouped into a series of terms involving *1-particle irreducible (1PI)* blobs, connected by free propagators. The 1PI blob represents the sum of all diagrams with two external legs and which cannot be disconnected by removing a single fermion line. The 2nd and the 3rd diagrams in the first line are 1PI (their extreme propagators being the external lines) while the last diagram is 1-particle reducible. The second line follows from first one by inspection.

Let $iS'_F(p)$ denote the lhs - the full 2-point function or full propagator - and $iS_F(p)$ denote the free propagator. Let the 1PI blob be denoted by $i\Sigma(p)$. Then the series is:

$$\begin{aligned}
 -iS'_F(p) &= -iS_F(p) + -iS_F \cdot i\Sigma(p) \cdot -iS_F + -iS_F \cdot i\Sigma \cdot -iS_F \cdot i\Sigma \cdot -iS_F + \dots \\
 &= -iS_F + -iS_F \cdot i\Sigma(-iS_F - -iS_F \cdot i\Sigma \cdot -iS_F \dots) \\
 \therefore -iS'_F &= -iS_F + -iS_F \cdot i\Sigma \cdot -iS'_F \\
 \therefore S_F &= (1 - S_F \Sigma) S'_F \leftrightarrow \mathbb{1} = S_F^{-1}(-\Sigma) S'_F \quad \text{and} \quad S_F^{-1} = \not{p} + m, \\
 \therefore & \boxed{S'_F(p) = \frac{\mathbb{1}}{\not{p} + m - \Sigma(p)}, \quad \Sigma(p) \text{ is called the fermion self energy.}}
 \end{aligned}$$

The $S'_F(p), \Sigma(p)$ are matrices with the Dirac spinor indices though we may not always be explicit about it. What can we say about its singularities?

We can appeal to Lorentz covariance and decompose $\Sigma(p) = a(p^2)\not{p} + b(p^2)\mathbb{1} + c(p^2)\gamma_5$. With parity conserving interactions such as qed, we can take $c(p^2) = 0$. Furthermore, *to the leading order*, $a(p^2) = 0 = b(p^2)$. Thus the denominator of the exact propagator is,

$$[S'_F(p)]^{-1} = (1 - a(p^2))\not{p} + (m - b(p^2)) = (1 - a(p^2)) \left[\not{p} + \frac{m - b(p^2)}{1 - a(p^2)} \right] \Rightarrow$$

$$S'_F(p) = \frac{1}{1 - a(p^2)} \left[\frac{-\not{p} + \frac{m - b(p^2)}{1 - a(p^2)}}{p^2 + \left(\frac{m - b(p^2)}{1 - a(p^2)} \right)^2} \right]$$

Clearly, the denominator vanishes for $p^2 = -m_{ph}^2, m_{ph} := \frac{m - b(-m_{ph}^2)}{1 - a(-m_{ph}^2)}$. Since the numerator has the form $-\not{p} + m_{ph}$, we can regard the pole to be defined by $\not{p} = -m_{ph}$. Since a, b are functions of $p^2 = -\not{p}\not{p}$, we may regard the self energy as a function of \not{p} and define the pole by the condition: $\boxed{\not{p} + m - \Sigma(\not{p})|_{\not{p} = -m_{ph}} = 0}$. Clearly, $m_{ph} \neq m$.

The exact propagator can be expanded about its pole as,

$$\not{p} + m - \Sigma(\not{p}) = 0 + (\not{p} + m_{ph}) \left[1 - \frac{d\Sigma(\not{p})}{d\not{p}} \Big|_{\not{p} = -m_{ph}} \right] + o((\not{p} + m_{ph})^2)$$

$$\therefore S'_F(p) \simeq \frac{1}{\not{p} + m_{ph}} \left[\frac{1}{1 - \frac{d\Sigma(\not{p})}{d\not{p}} \Big|_{\not{p} = -m_{ph}}} \right] := \frac{Z_2}{\not{p} + m_{ph}}$$

The shift of the position of the pole in the exact propagator from that of the free propagator, $\boxed{\delta m := m_{ph} - m = -\Sigma(\not{p} = -m_{ph})}$ is important in an S -matrix element which has the spinors on the external lines. These spinors satisfy the equation of motion with *physical mass* ($= m_{ph}$) and *not* the mass parameter in the free propagator inferred from the Lagrangian. In particular this means that in the LSZ formula for S -matrix element, the amputation of external fermion legs is to be done with $(-i\not{\partial} + m_{ph})$. This in turn means that in the Green's functions with external fermion legs, the external lines must include the self-energy corrections i.e. should have S'_F propagator instead of the free propagator S_F (and of course no more self energy corrections on the external lines). We will return to this point later again while discussing renormalized perturbation series.

We also see that the residue at the pole in the exact propagator is not 1 and we identify it with the Z_2 which we know from the Kallen-Lehmann representation to be *less than 1*.

B. The photon propagator: photon self-energy

$$\text{All} = \text{free} + \text{1-loop} + \text{2-loop} + \text{2-loop} \quad (16.3)$$

$$\text{All} = \text{free} + \text{1PI} + \text{2PI} + \dots \quad (16.4)$$

Here is a graphical representation of the exact photon propagator. As before, we group together the 1-PI diagrams connected by free propagators. Let the exact propagator be denoted by $-iD'_{\mu\nu}(q)$ and the free propagator by $-iD_{\mu\nu}(q)$. Let the 1PI blob be denoted by $i\Pi_{\mu\nu}(q)$. In the Lorentz gauge, the free propagator is: $D_{\mu\nu}(q) = \eta_{\mu\nu}/(q^2 - i\epsilon)$. While the series can be formally summed as before, it is more convenient to separate the tensor indices in the $\Pi_{\mu\nu}(q)$ tensor.

Lorentz covariance implies, $\Pi_{\mu\nu}(q) = \eta_{\mu\nu}A(q^2) + q_\mu q_\nu B(q^2)$. The $\Pi_{\mu\nu}$ tensor may be thought of as $\gamma \rightarrow \gamma$ process, although $q^2 \neq 0$. We now appeal to a *Ward identity*: $q^\mu \Pi_{\mu\nu}(q) = 0$. This is done separately below. Presently, it implies $A(q^2) = -q^2 B(q^2)$ and we define: $\boxed{\Pi_{\mu\nu}(q) := [\eta_{\mu\nu}q^2 - q_\mu q_\nu]\Pi(q^2)}$. With this notation, the exact 2-point function takes the form,

$$\begin{aligned} -iD'_{\mu\nu}(q) &= -i\frac{\eta_{\mu\nu}}{q^2 - i\epsilon} + -i\frac{\eta_{\mu\alpha}}{q^2 - i\epsilon} \cdot [i(q^2\eta^{\alpha\beta} - q^\alpha q^\beta)\Pi(q^2)] \cdot -i\frac{\eta_{\beta\nu}}{q^2 - i\epsilon} + \\ &\quad - iD_{\mu\alpha} \cdot [i(\dots)^{\alpha\beta}] \cdot -iD_{\beta\rho} \cdot [i(\dots)^{\rho\sigma}] \cdot -iD_{\sigma\nu} + \dots \end{aligned}$$

Define,

$$i(q^2\eta^{\alpha\beta} - q^\alpha q^\beta)(-i\eta_{\beta\nu}q^{-2}) = \left(\delta^\alpha_\nu - \frac{q^\alpha q_\nu}{q^2} \right) =: \Delta^\alpha_\nu \Rightarrow \Delta^\alpha_\nu \Delta^\nu_\beta = \Delta^\alpha_\beta.$$

Then the exact propagator series takes the form,


$$\begin{aligned} -iD'_{\mu\nu}(q) &= -i\frac{\eta_{\mu\nu}}{q^2 - i\epsilon} + -i\frac{\eta_{\mu\alpha}}{q^2 - i\epsilon} \Delta^\alpha_\nu \Pi(q^2) + -i\frac{\eta_{\mu\alpha}}{q^2 - i\epsilon} \Delta^\alpha_\beta \Delta^\beta_\nu \Pi(q^2) + \dots \\ &= -i\frac{\eta_{\mu\nu}}{q^2 - i\epsilon} + -i\frac{\eta_{\mu\alpha}}{q^2 - i\epsilon} \Delta^\alpha_\nu (\Pi + \Pi^2 + \Pi^3 + \dots) \\ &= \frac{-i}{q^2 - i\epsilon} \left[\left\{ \eta_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} + \frac{q_\mu q_\nu}{q^2} \right\} + \left(\eta_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \right) (\Pi + \Pi^2 + \Pi^3 \dots) \right] \\ \therefore D'_{\mu\nu}(q^2) &= \frac{1}{q^2(1 - \Pi(q^2))} \left(\eta_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \right) + \frac{1}{q^2} \frac{q_\mu q_\nu}{q^2}. \end{aligned}$$

In any S -matrix calculation, the D' propagator will land on a fermion line and thanks to the Ward identity, the terms proportional to q^μ, q^ν will vanish. Hence, for S -matrix element

computations, we can take $\boxed{D'_{\mu\nu}(q) = \frac{\eta_{\mu\nu}}{(q^2 - i\epsilon)(1 - \Pi(q^2))}}$. The $\Pi(q^2)$ is called the photon self energy. To the leading order, the self energy vanishes and hence in perturbation theory, it can never equal 1 and cause another pole at some other q^2 . As long as it is regular at $q^2 = 0$, the exact propagator *continues to have a simple pole at $q^2 = 0$* , just like the free propagator. Hence, the photon wavefunction, $\epsilon_\mu(\vec{k})$ continues to be transverse and longitudinal polarizations will decouple.

Unlike the fermion self energy which shifts the mass, the photon self energy does *not* shift the photon mass from zero thanks to the Ward identity. However, like the fermion self energy, the residue at the pole does shift away from 1, $\boxed{(Z_3)^{-1} := 1 - \Pi(q^2 = 0)}$.

1. *The Ward identity claim: $q^\mu \Pi_{\mu\nu}(q) = 0$:*



Proof: Recall that $\Pi_{\mu\nu}$ is defined without external propagators i.e. it is an amputated 2-point function. The momentum q however, need not be on-shell and hence it does *not* correspond to an S -matrix element (even if we disregard the polarization factors). Consider diagrams contributing to the photon self energy with some fixed number of vertices (or order in the coupling). The set of these diagrams will be generated by vertex injecting momentum q^μ at various points on the available fermion lines. The fermion lines will necessarily be loops as there are no external fermions contributing to the photon self energy. Consider the subset of diagrams wherein the q^μ vertex is on some particular loop which contributes a factor of the form: (the fermion arrow points from $i = 1$ to $i = n$ and so do the fermion momenta)

$$Tr [S_F(p_n + q)\gamma^{\mu_n} \dots S_F(p_{i+1} + q)\gamma^{\mu_{i+1}} \{S_F(p_i + q)\not{q}S_F(p_i)\} \gamma^{\mu_i} \dots S_F(p_1)\gamma^{\mu_1}]$$

The q^μ vertex is inserted between the i^{th} and $(i + 1)^{th}$ vertices and adds the propagator $S_F(p_i + q)$. All momenta from p_{i+1} to p_n are shifted by q . We have also contracted with q^μ .

The expression between the braces simplifies as ¹³,

$$\frac{1}{\not{p}_i + \not{q} + m} \not{q} \frac{1}{\not{p}_i + m} = \frac{1}{\not{p}_i + m} - \frac{1}{\not{p}_i + \not{q} + m} = S_F(p_i) - S_F(p_i + q) .$$

The trace thus becomes,

$$\begin{aligned} & Tr [S_F(p_n + q)\gamma^{\mu_n} \dots S_F(p_{i+1} + q)\gamma^{\mu_{i+1}} \{S_F(p_i)\} \gamma^{\mu_i} \dots S_F(p_1)\gamma^{\mu_1}] \\ & - Tr [S_F(p_n + q)\gamma^{\mu_n} \dots S_F(p_{i+1} + q)\gamma^{\mu_{i+1}} \{S_F(p_i + q)\} \gamma^{\mu_i} \dots S_F(p_1)\gamma^{\mu_1}] \end{aligned}$$


Note that the momenta after i (and $i - 1$) have q added to them. Summing over the diagrams where the q^μ vertex is on this fermion loop i.e from $i = 1, \dots, n$, there is a pairwise cancellation. The first term at i cancels the second term at $i + 1$ and so on. We are left with the second term at $i = 1$ and the first term at $i = n$, i.e.

$$\sum_{i=1}^n Tr [\dots] = Tr [S_F(p_n)\gamma^{\mu_n} \dots S_F(p_1)\gamma^{\mu_1}] - Tr [S_F(p_n + q)\gamma^{\mu_n} \dots S_F(p_1 + q)\gamma^{\mu_1}]$$

Because we have a fermion loop, there is an integration over say p_1 . *Provided the integral is finite, we can shift the integration variable and absorb away the q .* The integrals of the two traces, then cancel out and the claim is proved.

Remark: We have not yet introduced loop diagrams. The above fermion loop, before the q insertion, has n vertices and n momenta entering at those vertices. The conservation of momenta, taking out the overall conservation, thus leaves one momentum undetermined and this is to be integrated.

Remark: As an extension of the result, consider an open fermion line with incoming momentum p_0 , outgoing momentum p_n and insertions on it.



¹³ This identity holds as long as both the fermion propagators have the same mass, m_{ph} or m . This is relevant while dealing with external fermion lines.

For the insertion between the $i, i + 1$ vertices, we will have a string of factors as,

$$S_F(p_n + q)\gamma^{\mu_n}S_F(p_n + q) \dots \gamma^{\mu_{i+1}} \{S_F(p_i + q)\not{q}S_F(p_i)\} \gamma^{\mu_i} \dots S_F(p_1)\gamma^{\mu_1}S_F(p_0)$$

Replacing $S_F(p_i + q)\not{q}S_F(p_i) = S_F(p_i) - S_F(p_i + q)$ and summing over the insertions from $i = 0, \dots, n$, we get

$$\begin{aligned} \sum_{i=0}^n [\dots] &= S_F(p_n)\gamma^{\mu_n}S_F(p_{n-1}) \dots S_F(p_1)\gamma^{\mu_1}S(p_0) \\ &\quad - S_F(p_n + q)\gamma^{\mu_n}S_F(p_{n-1} + q) \dots S_F(p_1 + q)\gamma^{\mu_1}S(p_0 + q) \end{aligned}$$

To get an S -matrix contribution, we have to multiply by $\bar{u}(p_n + q)[S_F(p_n + q)]^{-1}$ on the left and by $[S_F(p_0)]^{-1}u(p_0)$ on the right and take the limits $p_0^2 \rightarrow -m^2, (p_n + q)^2 \rightarrow -m^2$. Now we see that each of the two terms has only one of the poles which can cancel the inverse propagator. In the on-shell limit then, each of the terms vanishes, thereby proving that *for S -matrix elements with external fermion lines, sum over insertion of photon lines followed by contraction with the photon momentum gives zero.*

Note: There is one subtlety here. For external lines, we should be using the exact propagator as per the LSZ rules. The $S\not{q}S$ identity requires both propagators to have the same mass, the physical mass. This would require all internal fermion lines to use the exact propagator (and then do not consider the fermion self energy correction). For fermion loop, this issue does not arise. Notice however that $S'_F(p)^{-1} = S_F(p)^{-1} - \Sigma(p)$ and $\sigma(p)$ is always of order α and higher. Thus, in using free inverse propagator instead of exact inverse propagator, we are dropping a higher order contribution which we can do in a perturbation theory.

The conclusion is then that $q_\mu \mathcal{M}^\mu(q, \dots) = 0$ which is a statement of Ward identity.

C. The Vertex function: Form factors

The expectation is that the electron-photon coupling will undergo a change due to higher order corrections. Unlike the 2-point functions above which were considered for arbitrary momentum, we consider the 3-point function with the fermion momenta “on-shell” (and this means the q^2 is necessarily space-like (prove this)) and hence an internal line. When viewed as part of an S -matrix element, the external fermion legs are amputated with *physical mass* inverse propagator and replaced by the $\bar{u}(p'), u(p)$ spinors respectively. Here is a graphical view of the possible corrections.

$$\mu \xrightarrow{q} \text{Vertex}(p, p') = \text{Tree} + \text{Self-energy on } p + \text{Self-energy on } p' + \text{Photon self-energy} \quad (16.5)$$

$$ie\Gamma^\mu(p, p') = ie\gamma^\mu + \mu \xrightarrow{q} \text{1PI} + \text{1PI} \text{ photon} + \text{1PI} \text{ photon}^2 + \dots \quad (16.6)$$

Notice that some corrections on the fermion lines and the photon line are just self-energy corrections. The remaining corrections have photon lines necessarily connecting the two fermion lines and these diagrams are already 1PI. The last diagram in the second equation is 1PR.

We can simplify the form of the exact vertex function, $\Gamma^\mu(p, p')$ by appealing to Lorentz covariance.

$$\Gamma^\mu(p, p') = \gamma^\mu A(q^2) + (p^\mu + p'^\mu)B(q^2) + (p'^\mu - p^\mu)C(q^2) .$$

With two independent momenta, we have 3 Lorentz scalars, $p^2, (p')^2, p \cdot p'$. The first two equal $-m_{ph}^2$ while the third is traded for q^2 for convenience.

Claim: $\bar{u}(p')q_\mu\Gamma^\mu u(p) = 0$.

Proof: This is the Ward identity argument given above.

The above form for Γ^μ then implies,

$$\begin{aligned} 0 &= \bar{u}(p')\not{q}u(p)A(q^2) + (-m_{ph}^2 + m_{ph}^2)\bar{u}(p')B(q^2)iu(p) + q^2\bar{u}(p')C(q^2)u(p) \\ &= 0 + 0 + q^2\bar{u}(p')C(q^2)u(p) \Rightarrow \bar{u}(p')C(q^2)u(p) = 0 . \end{aligned}$$

The A, B terms can be rewritten using ‘‘Gordon identity’’,

$$\boxed{\bar{u}(p')\gamma^\mu u(p) = \frac{1}{2m_{ph}}\bar{u}(p') [(p + p')^\mu - 2i\sigma^{\mu\nu}q_\nu] u(p)}$$

This can be established as follows ($m_{ph} \rightarrow m$ for convenience):

We have, $u(p) = -\frac{1}{m}\not{p}u(p)$, $\bar{u}(p') = -\frac{1}{m}\bar{u}(p')\not{p}'$. Therefore

$$\begin{aligned}
\bar{u}(p')\gamma^\mu u(p) &= -\frac{1}{2m}(\gamma^\mu \not{p} + \not{p}'\gamma^\mu) \quad \text{But, } \gamma^\mu \not{p} = p_\nu \gamma^\mu \gamma^\nu = p_\nu \left(-\eta^{\mu\nu} + \frac{1}{2}[\gamma^\mu, \gamma^\nu] \right) \\
\therefore \gamma^\mu \not{p} &= -p^\mu - 2i\Sigma^{\mu\nu} p_\nu \quad , \quad \not{p}'\gamma^\mu = -(p')^\mu - 2i\Sigma^{\nu\mu} p'_\nu \quad , \quad \Sigma^{\mu\nu} := \frac{i}{4}[\gamma^\mu, \gamma^\nu] ; \\
\therefore \bar{u}(p')\gamma^\mu u(p) &= -\frac{1}{2m}\bar{u}(p')[-p^\mu - (p')^\mu - 4i\Sigma^{\mu\nu}(p_\nu - p'_\nu)]u(p) \\
&= \frac{1}{2m}\bar{u}(p')[p^\mu + (p')^\mu - 2i\sigma^{\mu\nu}q_\nu]u(p) \quad , \quad \sigma^{\mu\nu} := 2\Sigma^{\mu\nu} \quad , \quad q := p' - p .
\end{aligned}$$

Eliminating the $(p + p')^\mu$ from the $B(q^2)$ term, and identifying $A + 2m_{ph}B =: F_1(q^2)$ and $4\Sigma^{\mu\nu}B(q^2) =: \frac{\sigma^{\mu\nu}}{2m_{ph}}F_2(q^2)$, we take the general form of $\Gamma^\mu(p, p')$ as,

$$\boxed{\Gamma^\mu(p, p') = \gamma^\mu F_1(q^2) + i\frac{\sigma^{\mu\nu}q_\nu}{2m_{ph}}F_2(q^2) \quad , \quad q := p' - p .}$$

The $F_i(q^2)$ are called *form factors*. To the lowest order, $F_1(q^2) = 1, F_2(q^2) = 0$. These get corrected at higher orders.

D. Electric Charge and Anomalous Magnetic Moment

The form factor decomposition helps us identify the *electric charge* and the *anomalous magnetic moment* of electron (and other charged fermions). This is seen as follows.

Consider the scattering of an electron off a heavy charged particle. We imagine the scattering to be effected by exchanging a single photon between the heavy charge and the electron. Schematically the amplitude can be written as,

$$i\mathcal{M} = ie^2 [\bar{u}(p')\Gamma^\mu(p', p)u(p)] \frac{1}{q^2} [\bar{u}(k')\gamma_\mu u(k)]$$

The first factor is from electron, second is the single photon propagator and the third factor is from the heavy charge. The Γ vertex has only 1PI diagrams and excludes the photon self energy on the propagator. This represents the modification of the electron response due to the 1PI corrections, the photon self energy would indicate modification in the electromagnetic field felt by the electron and hence is also referred to as *vacuum polarization*. This is ignored in this calculation.

In the limit of infinitely heavy fermion, the second and third factors are replaced by an external classical potential $A_\mu(q)$ and the amplitude is defined as,

$$i\mathcal{M} (2\pi) \delta((p' - p)^0) = -ie\bar{u}(p')\Gamma^\mu u(p)\tilde{A}_\mu^{cl}(p' - p)$$

We consider two cases - Coulomb potential and magnetic field, and consider non-relativistic limit of the scattering.

Coulomb potential: $A_\mu^{cl}(x) = (\phi(x), \vec{0})$. Coulomb field is static and we also assume it to be slowly varying spatially. This means that we take $\tilde{A}_\mu^{cl}(q) = 2\pi\delta(q^0)(\tilde{\phi}(q), \vec{0})$ with the $\tilde{\phi}(q)$ having support near $q \simeq 0$. We can thus take the limit $q \rightarrow 0$ of the first factor. This in turn means that only the F_1 term contributes. In the non-relativistic limit, $\bar{u}(p')\gamma^\mu u(p) \rightarrow \bar{u}(p')\gamma^0 u(p) = 2mu^\dagger(p)u(p)$ and the amplitude takes the form, $i\mathcal{M} \rightarrow -ieF_1(0)\tilde{\phi}(q)(2mu^\dagger(p)u(p))$. Comparing with the Born approximation for scattering off a potential leads to the identification, $V(x) = eF_1(0)\phi(\vec{x})$. Thus, electric charge = $eF(0)$. Since $F_1(0) = 1$ at the leading order, the radiative corrections to $F_1(q^2)$ should vanish as $q^2 \rightarrow 0$.

Magnetic field: $A_\mu^{cl} = (0, \vec{A}_i^{cl}(x))$. Again this is taken to be time independent and spatially slowly varying. Thus we consider the limit $q \rightarrow 0$. Earlier, we dropped the F_2 term, because F_1 term was non-zero. Now however $\bar{u}(p')\gamma^i u(p) \simeq o(p/m)$ in the non-relativistic limit which is comparable to the F_2 term.

Recall from the NR limit of the Dirac spinors, $u^T := (u_1^T, u_2^T)$, $(\not{p} + m)u(p) = 0 \Rightarrow u_2(p) \simeq -\frac{\vec{p}\cdot\vec{\sigma}}{2m}u_1$ and we have the normalization $u^\dagger u(p) = 2m$. We also had,

$$\bar{u}(p')\gamma^i u(p) \simeq u_1^\dagger(p')\sigma^i u_2(p) + u_2^\dagger(p')\sigma^i u_1(p) \simeq -\frac{1}{2m}u_1^\dagger(p')\{\sigma^i \vec{p} \cdot \vec{\sigma} + \vec{p}' \cdot \vec{\sigma} \sigma^i\}u_1(p).$$

Using $\sigma^i \sigma^j = \delta^{ij} + i\epsilon_k^{ij}\sigma^k$, we get,

$$\bar{u}(p')\vec{\gamma} \cdot \vec{A}^{cl} u(p) = -\frac{1}{2m}u_1^\dagger(p') \left(\vec{p} \cdot \vec{A}^{cl} + \vec{p}' \cdot \vec{A}^{cl} - iA_i^{cl} \epsilon_k^{ij} q_j \sigma^k \right) u_1(p)$$

Noting that $q^0 = 0$, the F_2 term has $\sigma^{ij}q_j$. Going over to the two component spinors, we get

$$\frac{i}{2m}\bar{u}(p')\sigma^{ij}u(p)q_j \approx \frac{i}{2m}\epsilon_k^{ij}[u_1^\dagger(p')\sigma^k u_1(p)]q_j$$

The contribution of the u_2 spinors is negligible in the NR limit.

Combining the terms we get in the NR limit,

$$\bar{u}(p')\Gamma^i u(p)\tilde{A}_i^{cl}(q) \approx u_1^\dagger(p') \left[\frac{(\vec{p} + \vec{p}') \cdot \vec{A}^{cl}}{2m} F_1(0) - \frac{i}{2m} \epsilon_k^{ij} q_j A_i^{cl} \sigma^k (F_1(0) + F_2(0)) \right] u_1(p)$$

In the non-relativistic Hamiltonian, $(p - eA)^2/(2m)$ we have the $e(p \cdot + A \cdot p)$ terms which are recovered. The remaining terms however are new. Focusing only on those, we write the

amplitude as,

$$i\mathcal{M} = -ie u_1^\dagger(p') \left[-\frac{1}{2m} (F_1(0) + F_2(0)) \sigma^k B_k \right] u_1(p) \quad , \quad B_k := -i \epsilon_k^{ij} q_i \tilde{A}_j^{cl}(q) .$$

This amplitude is interpreted as the Born approximation to a potential scattering with potential, $V(x) = -\langle \vec{\mu} \rangle \cdot \vec{B}(\vec{x})$, with the effective magnetic moment,

$$\langle \vec{\mu} \rangle = \frac{e}{m_{ph}} [F_1(0) + F_2(0)] u_1^\dagger(p) \frac{\vec{\sigma}}{2} u_1(p) := g \frac{e}{2m_{ph}} \vec{S} \quad , \quad g := 2[F_1(0) + F_2(0)] = 2[1 + F_2(0)] .$$




g is called the *Lande's g-factor*, its value being 2 is prediction of Dirac equation while $F_2(0) \neq 0$ is a prediction of QED. It is called the *anomalous magnetic moment*.

Without actually calculating the higher order corrections, we noted what they mean for 2-point functions (self-energies). We used Lorentz covariance to get a form factor decomposition and interpreted them by appealing to the NR limit. One of the immediate prediction was the anomalous magnetic moment.

We also note that to the leading order (see diagrams), we have $Z_2^{-1} = 1, Z_3^{-1} = 1, F_1(q^2) = 1, F_2(q^2) = 0$. This is the reason that at the tree level calculations that we did, did not have any factors of Z 's. This will change when we do explicit evaluations of the leading corrections in the next section.

17. RADIATIVE CORRECTIONS AT 1-LOOP: DIVERGENCES

We now calculate the leading radiative corrections to the self energies and the QED vertex function i.e. the 1-loop contributions to $\Sigma(p)$, $\Pi(q^2)$, $F_1(q^2)$, $F_2(q^2)$. Here are the Feynman diagrams and the corresponding invariant amplitudes.

	Fermion self energy
	$i\Sigma(p) = (ie)^2 \int \frac{d^4k}{(2\pi)^4} \gamma^\mu \frac{-i(-\not{k} + m)}{k^2 + m^2 - i\epsilon} \gamma^\nu \frac{-i\eta_{\mu\nu}}{(p-k)^2 + \mu^2 - i\epsilon}$
	Photon self energy (vacuum polarization)
	$i\Pi^{\mu\nu}(q) = (ie)^2(-1) \int \frac{d^4k}{(2\pi)^4} \text{Tr} \left[\gamma^\mu \frac{-i}{\not{k} + m} \gamma^\nu \frac{-i}{\not{k} + \not{q} + m} \right]$
	QED Vertex function, $\delta\Gamma^\mu := \Gamma^\mu - \gamma^\mu$
	$\delta\Gamma^\mu(p, p') = (ie)^2 \int \frac{d^4k}{(2\pi)^4} \frac{-i\eta_{\alpha\beta}}{(k)^2 + \mu^2 - i\epsilon} \times$ $\left[\gamma^\alpha \frac{-i(-(\not{p}' - \not{k}) + m)}{(p' - k)^2 + m^2 - i\epsilon} \gamma^\mu \frac{-i(-(\not{p} - \not{k}) + m)}{(p - k)^2 + m^2 - i\epsilon} \gamma^\beta \right]$

In the photon self energy expression, the (-1) is due to the fermion loop. The photon propagator is added a small mass μ^2 in anticipation. In the vertex correction, the expression is understood to be sandwiched between $\bar{u}(p')$ and $u(p)$.

There are some new points to be noted.

- There is a momentum integration over k , unrelated to the external momenta. This k can be space-like, time-like or light-like. In any frame, we may consider the region where $k^\mu \rightarrow \infty$, then: (i) the electron self-energy $\sim \int d^4k k^\alpha/k^4 \sim \int dk$ which is naively *linearly divergent*; (ii) the photon self-energy $\sim \int d^4k/k^2 \sim \int k dk$ which is naively *quadratically divergent* and (iii) the vertex correction $\sim d^4k k^2/k^6 \sim \int dk/k$ which is naively *logarithmically divergent*.

- There are also other regions of integration space which give divergent contributions, but these will be visible after the integrand is put in a convenient form.

- The $i\epsilon$ in the denominator is quite significant now. It implies that the 4-dimensional integration must be defined with the k^0 integration being done first and the spatial integrations done subsequently.

- The naively divergent integrals need to be *regularised* i.e. a prescription for the integration must be supplied which will manifest the divergence in an explicit form. The divergent contribution so obtained, must be *subtracted* to obtain finite answers. This is the process of *renormalization*.

- The integrand has a numerator which has tensor/spinor indices while the denominator is a product of scalar factors. One combines the denominator factors into a single factor form using the so called *Feynman parameters*. This is followed by the momentum integration with a regularization and a finite part is identified. The integration over the Feynman parameters is done last to get the answer. We explain these steps now.

The Feynman/Schwinger trick:

$$\text{Claim: } \frac{1}{A_1 \dots A_n} = (n-1)! \int_0^\infty dx_1 \dots dx_n \frac{\delta(\sum_i x_i - 1)}{(\sum_i x_i A_i)^n}.$$

Proof: Observe that $A^{-1} = \int_0^\infty d\alpha e^{-\alpha A}$, $Re(A) > 0$. Therefore,

$$\begin{aligned} \frac{1}{A_1 \dots A_n} &= \int_0^\infty d\alpha_1 \dots d\alpha_n e^{-\sum_i \alpha_i A_i} \cdot 1, \quad 1 = \int_0^\infty dt \delta(t - \sum_i \alpha_i) \quad \text{and} \quad \alpha_i \rightarrow tx_i, \\ &= \int_0^\infty dt \int_0^\infty dx_1 \dots dx_n t^n e^{-t \sum_i x_i A_i} \frac{1}{t} \delta(\sum_i x_i - 1) \\ &= \int_0^1 dx_1 \dots dx_n \delta(\sum_i x_i - 1) \int_0^\infty dt t^{n-1} e^{-t(\sum_i x_i A_i)} \\ &= (n-1)! \int_0^\infty dx_1 \dots dx_n \frac{\delta(\sum_i x_i - 1)}{(\sum_i x_i A_i)^n} \quad \because \int_0^\infty dt t^{n-1} e^{-t\Lambda} = \Lambda^{-n} \Gamma(n). \end{aligned}$$

Note: Since each of the denominator factors are of the form $A_i = (p_i - k)^2 + m_i^2 - i\epsilon$, we see that

$$\begin{aligned} \sum_i x_i A_i &= k^2 - 2k \cdot (\sum_i x_i p_i) + \sum_i x_i (p_i^2 + m_i^2) - i\epsilon \\ &= (k - \sum_i x_i p_i)^2 + M^2 - i\epsilon, \quad M^2(x_i, p_i, m_i) := \sum_i x_i (p_i^2 + m_i^2) - (\sum_i x_i p_i)^2. \end{aligned}$$

The obvious step is to shift the integration variable, $k \rightarrow k + \sum_i x_i p_i$ which simplifies the denominator - it has the same form of a single propagator with the same $-i\epsilon$.

Let us begin with the fermion self energy.

A. Isolation of Divergence: Fermion Self Energy

$$i\Sigma(p) = (ie)^2 \int \frac{d^4k}{(2\pi)^4} \gamma^\mu \frac{-i(-\not{k} + m)}{k^2 + m^2 - i\epsilon} \gamma^\nu \frac{-i\eta_{\mu\nu}}{(p-k)^2 + \mu^2 - i\epsilon}$$

The Feynman trick gives,

$$\frac{1}{(k^2 + m^2 - i\epsilon)((p-k)^2 + \mu^2 - i\epsilon)} = \int_0^1 dx \frac{1}{[(k-xp)^2 + M^2 - i\epsilon]^2},$$

with, $M^2 := x(1-x)p^2 + (1-x^2)m^2 + x\mu^2$. Shifting $k \rightarrow k + xp$, and using $\gamma^\mu \gamma_\mu = -4\mathbb{1}$, $\gamma^\mu \not{k} \gamma_\mu = 2\not{k}$, gives

$$i\Sigma = e^2 \int_0^1 \int \frac{d^4k}{(2\pi)^4} \frac{-4m - 2(\not{k} + x)\not{p}}{[k^2 + M^2(x, p) - i\epsilon]^2}$$

Recalling that the k^0 integration is to be done first and that the poles are at $k^0 = \pm\sqrt{\vec{k}^2 + M^2} \mp i\epsilon$, we rotate the contour anti-clockwise without crossing any singularity. This is equivalent to putting $k^0 = i\underline{k}^0$ and $k^2 \rightarrow (\underline{k}^0)^2 + \vec{k}^2 = \text{Euclidean vector norm}$. This sends $\int d^4k \rightarrow i \int d^4\underline{k}$. For Euclidean integrals $\int d^4\underline{k} \frac{\underline{k}^\alpha \gamma_\alpha}{(\underline{k}^2 + M^2)^2} = 0$, since the integrand is odd under $k \rightarrow -k$. We are left with,

$$i\Sigma = -ie^2 \int_0^1 dx \int \frac{d^4\underline{k}}{(2\pi)^4} \frac{4m + 2x\not{p}}{(\underline{k}^2 + M^2)^2}, \quad M^2 := x(1-x)p^2 + (1-x^2)m^2 + x\mu^2$$

The k -integral is logarithmically divergent and needs to be regulated. There are several ways of doing this. One is the *Pauli-Villars* regularization which subtracts from the photon propagator, another identical piece with arbitrarily large mass squared, Λ^2 , i.e.

$$\frac{1}{(p-k)^2 + \mu^2} \rightarrow \frac{1}{(p-k)^2 + \mu^2} - \frac{1}{(p-k)^2 + \Lambda^2}.$$

For large \underline{k} , both terms go as \underline{k}^{-2} and cancel each other leaving a finite answer. Thus $i\Sigma \rightarrow i\Sigma_{reg} := i\Sigma(\Lambda = \infty) - i\Sigma(\Lambda)$. The same combining of denominators will produce identical terms with $M^2(\mu^2) \rightarrow M^2(\Lambda^2)$. The momentum integration becomes,

$$\int \frac{d^4\underline{k}}{(2\pi)^4} \left[\frac{1}{(\underline{k}^2 + M_\mu^2)^2} - \frac{1}{(\underline{k}^2 + M_\Lambda^2)^2} \right] = \int \frac{d\Omega_3}{(2\pi)^4} \int_0^\infty dk |k|^3 [\dots]$$

The angular integration gives: $\Omega_{n-1} = \frac{2\pi^{n/2}}{\Gamma(n/2)} = 2\pi^2$ for $n = 4$ and the integral becomes,

$$\begin{aligned} \int \frac{d^4 \underline{k}}{(2\pi)^2} [\dots - \dots] &= \frac{1}{8\pi^2} \int_0^\infty \frac{dy}{2} \left[\frac{y}{(y + M_\mu^2)^2} - \frac{y}{(y + M_\Lambda^2)^2} \right], \quad k^2 =: y \text{ substituted.} \\ &= \frac{1}{16\pi^2} \int_0^\infty dy \left[\frac{1}{y + M_\mu^2} - \frac{1}{y + M_\Lambda^2} - \frac{M_\mu^2}{(y + M_\mu^2)^2} + \frac{M_\Lambda^2}{(y + M_\Lambda^2)^2} \right] \\ &= \frac{1}{16\pi^2} \left[\ln \left(\frac{y + M_\mu^2}{y + M_\Lambda^2} \right) \Big|_0^\infty + \left(\frac{M_\mu^2}{y + M_\mu^2} - \frac{M_\Lambda^2}{y + M_\Lambda^2} \right) \Big|_0^\infty \right] \\ \therefore \int \frac{d^4 \underline{k}}{(2\pi)^2} [\dots - \dots] &= \frac{1}{16\pi^2} \ln \left(\frac{M_\Lambda^2}{M_\mu^2} \right). \end{aligned}$$

In the limit $\Lambda \rightarrow \infty, M_\Lambda^2 \rightarrow x\Lambda^2$ and we get,

$$\boxed{i\Sigma(p) = -i \frac{e^2}{8\pi^2} \int_0^1 dx (2m + x\not{p}) \ln \left[\frac{x\Lambda^2}{x(1-x)p^2 + (1-x)m^2 + x\mu^2} \right]} \quad (17.1)$$

Recall that the mass shift as well as the Z_2 are obtained from the self energy as, $\delta m := m_{ph} - m = -\Sigma(\not{p} = -m_{ph}), Z_2^{-1} = 1 - \frac{\Sigma(\not{p})}{\not{p}} \Big|_{\not{p} = -m_{ph}}$. Since the self energy already has an explicit factor of e^2 , in the integrand, we can use $\not{p} = -m_{ph} \approx -m$. The derivative of Σ is obtained as,

$$\begin{aligned} \frac{d\Sigma}{d\not{p}} \Big|_{\not{p} = -m_{ph}} &= -\frac{\alpha}{2\pi} \int_0^1 dx \left[x \ln \left(\frac{x\Lambda^2}{-x(1-x)m^2 + (1-x)m^2 + x\mu^2} \right) + \right. \\ &\quad \left. (2m - x \cdot m) \frac{-1}{-x(1-x)m^2 + (1-x)m^2 + x\mu^2} (-2x(1-x)(-m)) \right] \end{aligned}$$

Simplification leads to,

$$\boxed{\frac{d\Sigma}{d\not{p}} \Big|_{\not{p} = -m_{ph}} = \frac{\alpha}{2\pi} \int_0^1 dx \left[-x \ln \left(\frac{x\Lambda^2}{(1-x)^2 m^2 + x\mu^2} \right) + 2(2-x) \frac{x(1-x)m^2}{(1-x)^2 m^2 + x\mu^2} \right]} \quad (17.2)$$

$$\boxed{\delta m = -\Sigma(\not{p} = -m_{ph}) = \frac{\alpha}{2\pi} \int_0^1 dx (2-x) \ln \left(\frac{x\Lambda^2}{(1-x)^2 m^2 + x\mu^2} \right)} \quad (17.3)$$

The integrands are well behaved at both end points so the integrals are finite and the leading contribution is given by the Λ^2 dependent part alone and for dimensional reasons we divide by the fermion mass. Thus, the *divergent contributions* as $\Lambda \rightarrow \infty$ take the form,

$$\begin{aligned} \delta m &\rightsquigarrow \frac{\alpha}{2\pi} \int_0^1 dx (2-x) \ln(\Lambda^2/m^2) = \frac{3\alpha}{4\pi} \ln(\Lambda^2/m^2) \\ Z_2^{-1} &\rightsquigarrow 1 - \frac{\alpha}{2\pi} \int_0^1 (-x \ln(\Lambda^2/m^2)) = 1 + \frac{\alpha}{4\pi} \ln(\Lambda^2/m^2) \Rightarrow Z_2 \approx 1 - \frac{\alpha}{4\pi} \ln(\Lambda^2/m^2) \end{aligned}$$

The logarithmic divergence is thus manifested in terms of $\ln(\Lambda^2/m^2)$. There are of course finite pieces, but until we take care of the divergent parts the finite parts are irrelevant.

B. Isolation of Divergence: Photon Self-Energy

$$i\Pi^{\mu\nu}(q) = -e^2 \int \frac{d^4k}{(2\pi)^4} \frac{\text{Tr}[\gamma^\mu(-\not{k} + m)\gamma^\nu(-\not{k} - \not{q} + m)]}{(k^2 + m^2 - i\epsilon)((k+q)^2 + m^2 - i\epsilon)}$$

Using the trace formulae for the Dirac matrices (14.4), we get

$$\begin{aligned} \text{Numerator:} &= \text{Tr}[\gamma^\mu \not{k} \gamma^\nu (\not{k} + \not{q}) + m^2 \gamma^\mu \gamma^\nu] \\ &= 4[k^\mu(k+q)^\nu - \eta^{\mu\nu} k \cdot (k+q) + (k+q)^\mu k^\nu - m^2 \eta^{\mu\nu}] \\ &= 4[2k^\mu k^\nu - \eta^{\mu\nu}(k^2 + m^2) + (k^\mu k^\nu + k^\nu q^\mu - \eta^{\mu\nu} k \cdot q)] \\ \text{Denominator:} &= \frac{1}{k^2 + m^2 - i\epsilon} \frac{1}{(k+q)^2 + m^2 - i\epsilon} \\ &= \int_0^1 dx \frac{1}{[(k+xq)^2 + x(1-x)q^2 + m^2]^2} \quad \text{shift } k \rightarrow k - xq, \end{aligned}$$

The shift generates additional terms linear in k in the numerator. Under integration, these terms drop out and we are left with,

$$i\Pi^{\mu\nu}(q) = (-4e^2) \int \frac{d^4k}{(2\pi)^4} \frac{\{2k^\mu k^\nu + 2x^2 q^\mu q^\nu - \eta^{\mu\nu}(k^2 + x^2 q^2 + m^2) - x(2q^\mu q^\nu - \eta^{\mu\nu} q^2)\}}{(k^2 + M^2 - i\epsilon)^2}$$

where, $M^2 = x(1-x)q^2 + m^2$. Doing Wick rotation as before,

$$i\Pi^{\mu\nu}(q) = (-4e^2)i \int \frac{d^4\mathbf{k}}{(2\pi)^4} \frac{\{2\mathbf{k}^\mu \mathbf{k}^\nu - 2x(1-x)q^\mu q^\nu + x(1-x)\eta^{\mu\nu} q^2 - \eta^{\mu\nu}(\mathbf{k}^2 + m^2)\}}{(\mathbf{k}^2 + M^2 - i\epsilon)^2}$$

The tensorial integral,

$$\int \frac{d^4\mathbf{k}}{(2\pi)^4} \frac{\mathbf{k}^\mu \mathbf{k}^\nu}{(\mathbf{k}^2 + M^2)^2} = A\delta^{\mu\nu} \quad , \quad \text{with } A = \frac{1}{4} \int \frac{d^4\mathbf{k}}{(2\pi)^4} \frac{\mathbf{k}^2}{(\mathbf{k}^2 + M^2)^2}$$

There are several problems to be faced now. First, the integral is quadratically divergent and shift in the momentum variable is unjustified. Second, we have now $\delta^{\mu\nu}$ and $\eta^{\mu\nu}$ in the numerator, confusing Lorentz covariance. Third, naively we see a quadratically divergent piece with coefficient $\eta^{\mu\nu}$ but only logarithmically divergent one with coefficient $q^\mu q^\nu$ casting doubts on possibility of gauge invariance. We could try the Pauli-Villars regulation, but we will introduce a different one, the *dimensional regularization*. The Pauli-Villars may be seen in [13]. The dimensional regularization has the main virtue of keeping the Ward identity satisfied, even for non-abelian gauge theories.

The idea is to think of the perturbation theory to be a member of a class of similar theories formulated in general n dimensions. One chooses a value of n where the integrals are well

defined ($n < 4$) and analytically continues $n \rightarrow 4$. This isolates the original divergence in a particular form (poles) providing the needed regularization. Since the mass/length dimensions also depend on space-time dimensions we need the coupling constants to be dimensionfull. To maintain space-time covariance, we need to regard the external momenta and the γ matrices etc also to be n -dimensional. We set $\epsilon := 4 - n$ and consider the limit $\epsilon \rightarrow 0$. Since the integrals are over Euclidean momenta due to the Wick rotation, *the external momenta and the metric tensor need to continue back to Minkowski signature at the end of the calculations.*

In this regularization scheme, the following rules are adopted.

$$\int \frac{d^n k}{(2\pi)^n} f(|k|^2) = \int \frac{d\Omega_{n-1}}{(2\pi)^n} \int dk |k|^{n-1} f(|k|^2) , \quad \boxed{\int d\Omega_{n-1} = \frac{2\pi^{n/2}}{\Gamma(n/2)}} \quad (17.4)$$

$$\therefore \int \frac{d^n k}{(2\pi)^n} f(|k|^2) = \left[\frac{2\pi^{n/2}}{\Gamma(n/2)} \right] \frac{1}{(2\pi)^n} \int dk k^{n-1} f(k^2) \quad (17.5)$$

$$\begin{aligned} \int_0^\infty dk \frac{k^{n-1}}{(k^2 + M^2)^\alpha} &= \frac{1}{2} \int_0^\infty dx \frac{x^{\frac{n}{2}-1}}{(x + M^2)^\alpha} , \quad \text{put } y = \frac{M^2}{x + M^2} \\ &= \frac{M^{n-2\alpha}}{2} \int_0^1 dy y^{\alpha-1-\frac{n}{2}} (1-y)^{\frac{n}{2}-1} = \frac{M^{n-2\alpha}}{2} \beta(\alpha - n/2, n/2) \end{aligned}$$

$$\boxed{\therefore \int_0^\infty dk \frac{k^{n-1}}{(k^2 + M^2)^\alpha} = \frac{M^{n-2\alpha}}{2} \frac{\Gamma(\alpha - n/2)\Gamma(n/2)}{\Gamma(\alpha)}} \quad (17.6)$$

and

$$\boxed{\int \frac{d^n k}{(2\pi)^n} \frac{1}{(k^2 + M^2)^\alpha} = \frac{1}{(4\pi)^{n/2}} \frac{\Gamma(\alpha - n/2)}{\Gamma(\alpha)} (M^2)^{n/2-\alpha}} \quad (17.7)$$

The Γ function has isolated poles at $\alpha - n/2 = 0, -1, -2, \dots$ i.e. at $n = 2(\alpha + m), m = 0, 1, \dots$. The Gamma function has the expansion $\Gamma(\epsilon/2) = 2/\epsilon - \gamma + o(\epsilon)$ where, $\gamma = 0.57721 \dots$ is the *Euler-Mascheroni Constant*.

Since finite parts can be affected we need to keep ϵ in various place too eg $\{\gamma^\mu, \gamma^\nu\} = -2\delta^{\mu\nu}\mathbb{1}, Tr(\mathbb{1}) = n = 4 - \epsilon, \gamma^\mu \gamma^\nu \gamma_\mu = (2 - \epsilon)\gamma^\nu$. These terms will give contributions from the $1/\epsilon$ terms. The coupling constant dimensions work as follows. $[\Psi] = (n - 1)/2, [A_\mu] = (n - 2)/2 \Rightarrow [e] = n - (n - 1) - (n/2 - 1) = \epsilon/2$. Therefore we write $e \rightarrow e\mu_0^{\epsilon/2}$ where μ_0 is an arbitrary mass scale which should disappear from physical quantities. The dimensions are in mass units. The tensorial integral will now follow from $k^\mu k^\nu \rightarrow \frac{1}{n}\delta^{\mu\nu}k^2$. We use these in evaluating the $\Pi^{\mu\nu}(q)$.

$$\begin{aligned}
\Pi^{\mu\nu}(\underline{q}) &= (-ne^2\mu_0^\epsilon) \int_0^1 dx \left[\{-2x(1-x)\underline{q}^\mu\underline{q}^\nu + \delta^{\mu\nu}(x(1-x)\underline{q}^2 - m^2)\} \times \right. \\
&\quad \left. \int \frac{d^n \underline{k}}{(2\pi)^n} \frac{1}{(\underline{k}^2 + M^2)^2} + \delta^{\mu\nu}(2/n - 1) \int \frac{d^n \underline{k}}{(2\pi)^n} \frac{\underline{k}^2 + M^2 - M^2}{(\underline{k}^2 + M^2)^2} \right], \text{ where,} \\
&\quad \boxed{M^2 = x(1-x)q^2 + m^2} \\
&= (-ne^2\mu_0^\epsilon) \int_0^1 dx \left[\delta^{\mu\nu} \left(\frac{2}{n} - 1 \right) \underbrace{\int \frac{d^n \underline{k}}{(2\pi)^n} \frac{1}{\underline{k}^2 + M^2}}_{\textcircled{1}} + \underbrace{\int \frac{d^n \underline{k}}{(2\pi)^n} \frac{\underline{k}^2}{(\underline{k}^2 + M^2)^2}}_{\textcircled{2}} \times \right. \\
&\quad \left. \left\{ -2x(1-x)\underline{q}^\mu\underline{q}^\nu + \delta^{\mu\nu} \left(-m^2 - x(1-x)\underline{q}^2 - \left(\frac{2}{n} - 1 \right) M^2 \right) \right\} \right]
\end{aligned}$$

$$\textcircled{1} = \frac{1}{(4\pi)^{n/2}} \frac{\Gamma(1 - n/2)}{\Gamma(1)} (M^2)^{n/2-1} = \frac{M^2}{(4\pi)^2} \left(\frac{M^2}{4\pi} \right)^{-\epsilon/2} \frac{\Gamma(\epsilon/2)}{(\epsilon/2 - 1)}$$

$$\textcircled{2} = \frac{1}{(4\pi)^{n/2}} \frac{\Gamma(2 - n/2)}{\Gamma(1)} (M^2)^{n/2-2} = \frac{1}{(4\pi)^2} \left(\frac{M^2}{4\pi} \right)^{-\epsilon/2} \Gamma(\epsilon/2)$$

$$\therefore \textcircled{1} = \left(\frac{M^2}{\epsilon/2 - 1} \right) \textcircled{2}, \quad \frac{\epsilon}{2} - 1 = 1 - \frac{n}{2} = \frac{n}{2} \left(\frac{2}{n} - 1 \right)$$

$$\begin{aligned}
\Pi^{\mu\nu}(q) &= (-ne^2\mu_0^\epsilon) \int_0^1 dx \left[\delta^{\mu\nu} \left(\frac{2}{n} - 1 \right) \frac{M^2}{\frac{n}{2} \left(\frac{2}{n} - 1 \right)} \right. \\
&\quad \left. + \delta^{\mu\nu} \left(-m^2 - \left(\frac{2}{n} - 1 \right) M^2 + x(1-x)q^2 \right) - 2x(1-x)q^\mu q^\nu \right] \times \textcircled{2}
\end{aligned}$$

$$\text{But, } [\dots] = \delta^{\mu\nu} \left\{ \frac{2}{n} \cancel{M^2} - m^2 - \underbrace{\left(\frac{2}{n} \cancel{M^2} - M^2 \right)}_{2x(1-x)q^2} + x(1-x)q^2 \right\} - 2x(1-x)q^\mu q^\nu$$

$$= 2x(1-x)(q^2\delta^{\mu\nu} - q^\mu q^\nu) \Rightarrow \boxed{\Pi^{\mu\nu}(\underline{q}) = (q^2\delta^{\mu\nu} - q^\mu q^\nu)\Pi(q^2) \text{ with,}}$$

$$\begin{aligned}
\Pi(q^2) &:= -ne^2\mu_0^\epsilon \int_0^1 dx \frac{1}{(4\pi)^2} \left(\frac{M^2}{4\pi} \right)^{-\epsilon/2} \Gamma(\epsilon/2)(2x)(1-x) \\
&= -\frac{2(4-\epsilon)e^2}{(4\pi)^2} \int_0^1 dx x(1-x) \left\{ \frac{2}{\epsilon} - \gamma \right\} \left\{ \ln \left(\frac{4\pi\mu_0^2}{M^2} \right) \frac{\epsilon}{2} + 1 \right\}
\end{aligned}$$

Using $a^{\epsilon/2} = 1 + \frac{\epsilon}{2}\ln(a) + o(\epsilon^2)$, we write,

$$\boxed{\Pi(q^2) = -\frac{2\alpha}{\pi} \int_0^1 dx x(1-x) \left(\frac{2}{\epsilon} - \gamma \right) \left(1 + \frac{\epsilon}{2}\ln \left(\frac{4\pi\mu_0^2}{M^2} \right) \left(1 - \frac{\epsilon}{4} \right) \right)} \quad (17.8)$$

Notice that for q^2 , $M^2 = m^2$ and

$$\boxed{\Pi(0) = -\frac{2\alpha}{\pi} \frac{2}{\epsilon} \int_0^1 dx(x-x^2) + \text{constant} = -\frac{2\alpha}{3\pi} \frac{1}{\epsilon}} \Rightarrow \boxed{Z_3 = 1 - \frac{2\alpha}{3\pi\epsilon}} \quad (17.9)$$

We see the divergence in $\Pi(0)$, Z_3 and there is of course no mass shift. We return to the interpretation of $\Pi(q^2)$ later.

Remark: The ϵ^{-1} pole in dimensional regularization corresponds to logarithmic divergence in Pauli-Villars. this is indicated by comparing the regulated integral $\int[(k^2 + M^2)^{-2} - (k^2\Lambda^2)^{-2}]$ in Pauli-Villars, which goes over to $\frac{1}{4\pi^2} \ln(x\Lambda^2/M^2)$. In the dimensional regularization, our integral (2) goes as $\frac{1}{4\pi^2} \frac{2}{\epsilon}$. Hence, $\epsilon^{-1} \leftrightarrow \ln(\Lambda/M)$. Thanks to the dimensional regularization maintaining the Ward identity, the $q^2\delta^{\mu\nu} - q^\mu q^\nu$ got pulled out and the superficially quadratically divergent integral go converted to a logarithmically divergent one.

C. Isolation of Divergence: Vertex function

We compute $\delta\Gamma^\mu = \Gamma^\mu - \gamma^\mu$, with $p^2 = (p')^2 = -m^2$ and sandwiching by $\bar{u}(p')$, $u(p)$ is implicit.

$$\delta\Gamma^\mu(p, p') = -ie^2 \int \frac{d^4k}{(2\pi)^4} \frac{\eta_{\alpha\beta}}{k^2 + \mu^2 - i\epsilon} \left[\gamma^\alpha \frac{-(\not{p}' - \not{k}) + m}{(p' - k)^2 + m^2 - i\epsilon} \gamma^\mu \frac{-(\not{p} - \not{k}) + m}{(p - k)^2 + m^2 - i\epsilon} \gamma^\beta \right]$$

We have three denominators and the Feynman trick leads to,

$$\begin{aligned} & \frac{1}{k^2 + \mu^2 - i\epsilon} \frac{1}{(p - k)^2 + m^2 - i\epsilon} \frac{1}{(p' - k)^2 + m^2 - i\epsilon} \\ &= \int_0^1 dx \int_0^1 dy \int_0^1 dz \frac{2}{[(k - xp - yp')^2 + M^2 - i\epsilon]^3} \quad \text{where,} \\ & M^2 := x(p^2 + m^2) + y((p')^2 + m^2) + z\mu^2 - (xp + yp')^2 \\ & \therefore M^2 = 0 + 0 + z\mu^2 + m^2(x + y)^2 + q^2xy \end{aligned}$$

Since $q^2 > 0$ (space-like) for this diagram, $\boxed{M^2 = (x + y)^2 m^2 + xyq^2 + z\mu^2}$ is manifestly positive. As usual shifting the momentum $k \rightarrow k + xp + xp'$, doing the Wick rotation and dropping the terms linear in k in the numerator, we get,

$$\delta\Gamma^\mu(p, p') = e^2 \int_0^1 dx \int_0^1 dy \int_0^1 dz \delta(1 - x - y - z) \int \frac{d^4k}{(2\pi)^4} \frac{2 \times Nr}{[k^2 + M^2]^3}$$

$$\begin{aligned} Nr &= \gamma^\alpha \{(\not{p}' - \not{k} - x\not{p} - y\not{p}')\gamma^\mu(\not{p}' - \not{k} - x\not{p} - y\not{p}')\} \gamma_\alpha + m^2 \gamma^\alpha \gamma^\mu \gamma_\alpha \\ &\quad - m \gamma^\alpha \{\gamma^\mu(\not{p} - x\not{p} - y\not{p}') + (\not{p}' - x\not{p} - y\not{p}')\gamma^\mu\} \gamma_\alpha + \text{linear in } k \end{aligned}$$

Now we need to use the identities,

$$\gamma^\alpha \gamma^\mu \gamma_\alpha = -2\eta^{\alpha\mu} \gamma_\alpha - \gamma^\mu(-4) = 2\gamma^\mu \quad (17.10)$$

$$\gamma^\alpha \gamma^\rho \gamma^\sigma \gamma_\alpha = 4\eta^{\rho\sigma} \quad (17.11)$$

$$\gamma^\alpha \gamma^\rho \gamma^\mu \gamma^\sigma \gamma_\alpha = 2\gamma^\sigma \gamma^\mu \gamma^\rho \quad (17.12)$$

$$\therefore \gamma^\alpha \not{k} \gamma^\mu \not{k} \gamma_\alpha = 2\not{k} \gamma^\mu \not{k} = -4k^\mu \not{k} + 2\gamma^\mu k^2 \quad \because \not{k} \not{k} = -k^2 \quad (17.13)$$

$$\gamma^\alpha \{\dots\} \gamma^\mu \{\dots\} \gamma_\alpha = 2\{(1-x)\not{p} - y\not{p}'\} \gamma^\mu \{(1-y)\not{p}' - x\not{p}\} \quad (17.14)$$

$$\gamma^\alpha \{\gamma^\mu(\dots) + (\dots)\gamma^\mu\} \gamma_\alpha = 4\{(1-x)p^\mu - y(p')^\mu + (1-y)(p')^\mu - xp^\mu\} \quad (17.15)$$

Using these, the numerator takes the form,

$$\begin{aligned} \bar{u}(p')(Nr)u(p) = & \bar{u}(p') \left([-4k^\mu \not{k} + 2k^2 \gamma^\mu] \textcircled{1} + 2[\{(1-x)\not{p} - y\not{p}'\} \gamma^\mu \{(1-y)\not{p}' - x\not{p}\}] \textcircled{2} \right. \\ & \left. + 2[m^2 \gamma^\mu] \textcircled{3} - 4m[(1-2x)p^\mu + (1-2y)(p')^\mu] \textcircled{4} \right) u(p) \end{aligned}$$

Observe that $\not{p}u(p) = -mu(p)$, $\bar{u}\not{p}' = -m\bar{u}p'$. In $\textcircled{1}$, $\int d^4k k^\mu \not{k} \propto \gamma^\mu$ (after Pauli-Villars).

We simplify $\textcircled{2}$ as,

$$\begin{aligned} \frac{1}{2}\bar{u}(p')\textcircled{2}u(p) = & \bar{u}(p')[(1-x)(1-y)\not{p}\gamma^\mu\not{p}' + xym^2\gamma^\mu + x(1-x)m\not{p}\gamma^\mu + y(1-y)m\gamma^\mu\not{p}']u(p) \\ \not{p}\gamma^\mu \rightarrow & -2p^\mu - (-m)\gamma^\mu, \quad \gamma^\mu\not{p}' \rightarrow -2(p')^\mu + m\gamma^\mu \quad \text{between the spinors.} \end{aligned}$$

Thus we see that, between the spinors, all the terms in the numerator can be put in the form of $(p+p')^\mu[\dots] + \gamma^\mu[\dots]$ and therefore the integral can be arranged as contributing to the form factors F_1, F_2 . The intermediate steps, including the Pauli-Villars regularization, are left as an exercise. The result is (from $\delta\Gamma^\mu$):

$$F_1(q^2) - 1 = \frac{\alpha}{2\pi} \int_0^1 dx dy dz \delta(1-x-y-z) \left[\ln\left(\frac{z\Lambda^2}{M^2}\right) + \frac{1}{M^2} \left((1-x)(1-y)q^2 + (1-4z+z^2)m^2 \right) \right] \quad (17.16)$$

$$F_2(q^2) = \frac{\alpha}{2\pi} \int_0^1 dx dy dz \delta(1-x-y-z) \left[\frac{2m^2 z(1-z)}{M^2} \right], \quad \text{with} \quad (17.17)$$

$$\boxed{M^2 := q^2 xy + m^2(1-z)^2 + z\mu^2}$$

Note:

(i) In the δF_1 , there is a UV divergence ($\Lambda \rightarrow \infty$) coming from the large k region, even for $q^2 = 0$. This violates the $\delta F_1(0) = 0$ condition and thus changes the electric charge. The condition can be naturally restored by defining $\boxed{\delta F_{1,ren}(q^2) := \delta F_1(q^2) - \delta F_1(0)}$. This is a *renormalization prescription*.

(ii) The δF_1 also has an ‘‘infrared divergence’’ ($\mu^2 \rightarrow 0$) at $q^2 = 0$ coming from the $M^{-2}(0) = m^{-2}(1-z)^{-2}$ from the $z \rightarrow 1$ region.

(iii) The $F_2(q^2)$ has no divergences. Its value at $q^2 = 0$ gives the anomalous magnetic moment and can be evaluated explicitly as,

$$F_2(0) = \frac{\alpha}{2\pi} \int_0^1 dx dy dz \delta(1-x-y-z) \frac{2z}{1-z} = \frac{\alpha}{\pi} \int_0^1 dz \int_0^{1-z} \frac{z}{1-z} = \frac{\alpha}{2\pi}$$

$$\Rightarrow \boxed{\frac{g-2}{2} = \frac{\alpha}{2\pi}} \quad \text{One of the triumphs of QED!} \quad (17.18)$$


The *renormalized* $F_1(q^2)$ is obtained as,

$$F_{1,ren}(q^2) = 1 + \frac{\alpha}{2\pi} \int_0^1 dx dy dz \delta(1-x-y-z) \left[\ln \left(\frac{m^2(1-z)^2}{m^2(1-z)^2 + q^2 xy} \right) + \frac{m^2(1-4z+z^2) + q^2(1-x)(1-y)}{m^2(1-z)^2 + q^2 xy + \mu^2 z} - \frac{m^2(1-4z+z^2)}{m^2(1-z)^2 + \mu^2 z} \right] \quad (17.19)$$

We have to face the divergences now. All three radiative corrections have UV divergence while the vertex function has IR divergence as well. For dealing with the IR divergence, we need to look at another process called ‘‘Bremsstrahlung’’ (breaking radiation in German).

D. Bremsstrahlung Cross-section to $o(\alpha)$

We know from the classical electrodynamics that accelerated charge radiates. A quantum mechanical view of such a radiation is depicted in the diagrams below.



The diagrams depict a scattering process which causes the top electron to ‘accelerate’ and ‘emit’ a (on-shell) photon. This is the QFT view of ‘radiation from an accelerated charge’. The bottom charged line represents some heavy fermion/boson/source, while the connecting wavy line denotes mediation of the interaction. The first diagram suggests photon emission *before* the kick while the second diagram suggests photon emission *after* the kick. As per

the rules of the relativistic QFT, both contributions are to be added and mod-squared to get to the cross-section. To the leading order - single photon emission - the cross-section is order α . This cross-section also has a an IR divergence as $k \rightarrow 0$. An on-shell photon with arbitrarily small momentum is called a ‘soft photon’. To isolate the divergence, focus on the following diagrams:



We are computing the amplitude for the process $e \rightarrow e + \gamma$. Let $\mathcal{M}_0(k', l')$ denote the part of the amplitude of electron interacting with the external source, it is depicted as the blob. Here k', l' denote the out-going and in-coming momenta respectively. The full amplitude is then given by (polarizations $\varepsilon(\vec{k}, \lambda)$ taken to be real),

$$i\mathcal{M}(p', p, k) = (ie)\bar{u}(p') \left[\mathcal{M}_0(p', p-k) \frac{-i(-\not{p}' - \not{k}) + m}{(p-k)^2 + m^2 - \epsilon} \gamma^\mu \varepsilon_\mu \right. \\ \left. \varepsilon_\mu \gamma^\mu \frac{-i(-\not{p}' + \not{k}) + m}{(p'+k)^2 + m^2 - i\epsilon} \mathcal{M}_0(p'+k, p) \right] u(p)$$

To compare with the classical radiation formula, consider the limit wherein the external source interaction approximates as, $\mathcal{M}_0(p', p-k) \approx \mathcal{M}_0(p'+k, p) \approx \mathcal{M}_0(p', p)$. Next, since the photon is soft, we neglect the \not{k} in the fermion numerators. The denominators simplify as, $(p-k)^2 + m^2 = -m^2 - 2p \cdot k + 0 = m^2 = -2p \cdot k$ and likewise, $(p'+k)^2 + m^2 = +2p' \cdot k$. Furthermore, $(-\not{p}' + m)\gamma^\mu u(p) \rightarrow 2p^\mu u(p)$, and $\bar{u}(p')\gamma^\mu(-\not{p}' + m) \rightarrow \bar{u}(p')(p')^\mu$. Thus the amplitude takes the form,

$$i\mathcal{M}(p', p, k) = e \underbrace{[\bar{u}(p')\mathcal{M}_0(p', p)u(p)]}_{\text{elastic scattering amplitude}} \times \underbrace{\left(-\frac{\varepsilon \cdot p}{p \cdot k} + \frac{\varepsilon \cdot p'}{p' \cdot k} \right)}_{\text{extra } k \text{ dependent factor}}$$

The total, unpolarized cross-section is obtained as,

$$d\sigma(p \rightarrow p' + k) = d\sigma(p \rightarrow p') \int \frac{d^3k}{2|k|(2\pi)^3} \sum_{\lambda=1,2} e^2 \left| \frac{\varepsilon \cdot p'}{p' \cdot k} - \frac{\varepsilon \cdot p}{p \cdot k} \right|^2 \quad (17.20)$$

The integration is over soft photons i.e. $|\vec{k}| < k_{max}$. Notice that The integral is dimensionless.

The integrand is the probability density of radiating a photon of momentum k within d^3k and any transverse polarization, accompanying the electron scattering with $p \rightarrow p'$

(“acceleration kick”, $p' \neq p + k$) while *the integral is the total probability of emission of a soft photon with energy up to k_{max} .*

Multiplying the integrand by the photon energy, $|\vec{k}|$ gives the *expectation value of the energy carried away by soft photons up to energy k_{max} ,*

$$\mathcal{E}_{soft}(k_{max}) := \langle E \rangle = \int \frac{d^3k}{(2\pi)^3} \sum_{\lambda} \frac{e^2}{2} \left| \vec{\varepsilon} \cdot \left(\frac{\vec{p}'}{p' \cdot k} - \frac{\vec{p}}{p \cdot k} \right) \right|^2$$

The polarization sum in the cross-section can be done as follows. Noting that $k \cdot \left(\frac{p'}{p' \cdot k} - \frac{p}{p \cdot k} \right) = 0$, we can drop the $k\vec{k}$ terms in the completeness relation for the polarizations, effectively taking $\sum_{\lambda} \varepsilon_{\mu} \varepsilon_{\nu} = \eta_{\mu\nu}$. Hence,

$$\sum_{\lambda} \left[\left(\frac{p'}{p' \cdot k} - \frac{p}{p \cdot k} \right)^{\mu} \varepsilon_{\mu} \right] \left[\left(\frac{p'}{p' \cdot k} - \frac{p}{p \cdot k} \right)^{\nu} \varepsilon_{\nu} \right] = \left(\frac{p'}{p' \cdot k} - \frac{p}{p \cdot k} \right) \cdot \left(\frac{p'}{p' \cdot k} - \frac{p}{p \cdot k} \right),$$

and the expectation value of energy carried by soft photons is given by,

$$\boxed{\mathcal{E}_{soft}(k_{max}) = \int \frac{d^3k}{(2\pi)^3} \sum_{\lambda} \frac{e^2}{2} \left[-\frac{m^2}{(p' \cdot k)^2} - \frac{m^2}{(p \cdot k)^2} - 2 \frac{p \cdot p'}{(p \cdot k)(p' \cdot k)} \right]} \quad (17.21)$$

Note that the integral represents average radiated energy inferred from the quantum mechanical probability distribution for emission of a single soft photon with $|\vec{k}| < k_{max}$.

An Aside: Incidentally, integrand in the r.h.s. is also the *classical* formula for energy carried away by the k^{th} Fourier mode of the electromagnetic field. To see this, consider a classical trajectory $z^{\mu}(\tau) = p^{\mu}/m$ for $\tau < 0$ and equals $(p')^{\mu}/m$ for $\tau > 0$. The corresponding current is,

$$J^{\mu}(x) = \int_0^{\infty} d\tau \frac{(p')^{\mu}}{m} \delta^4(x - z(\tau)) + \int_{-\infty}^0 d\tau \frac{p^{\mu}}{m} \delta^4(x - z(\tau)) \Rightarrow$$

$$A^{\mu}(\vec{k}) = -\frac{e}{|\vec{k}|} \left(\frac{(p')^{\mu}}{p' \cdot k} - \frac{p^{\mu}}{p \cdot k} \right) \quad (\text{using the Retarded Green function.})$$

The energy of the above radiation field is the precisely as given by the integrand in the r.h.s. of eq.(17.21) [13]. The integral now represents *the total energy carried by Fourier modes with momenta $|\vec{k}| < k_{max}$.*

The integral, $\mathcal{E}_{soft}(k_{max})$, thus has two different interpretations.

Returning to the evaluation of the integral in eq.(17.21), let us choose a frame in which the initial and final energies of the electron are the same: $E' = E \Rightarrow (p')^0 = p^0$ and we write $p = E(1, \vec{v})$, $p' = E(1, \vec{v}')$, $k = (k := |\vec{k}|, \vec{k})$. This is equivalent to $p^2 = -m^2$ and

$m^2/E^2 = 1 - \vec{v}^2$. With this choice, $p \cdot k = Ek(-1 + \hat{k} \cdot \vec{v})$, $p' \cdot k = Ek(-1 + \hat{k} \cdot \vec{v}')$ and $p \cdot p' = E^2(-1 + \vec{v} \cdot \vec{v}')$. The k^2 from the d^3k cancels and the integrated emitted energy takes the form,

$$\begin{aligned} \mathcal{E}_{soft}(k_{max}) &= -\frac{e^2}{2} \int dk \int \frac{d\Omega}{8\pi^3} \left[\frac{m_0^2}{E^2} \left(\frac{1}{(1 - \hat{k} \cdot \vec{v})^2} + \frac{1}{(1 - \hat{k} \cdot \vec{v}')^2} \right) - \frac{2(1 - \vec{v} \cdot \vec{v}')}{(1 - \hat{k} \cdot \vec{v})(1 - \hat{k} \cdot \vec{v}')} \right] \\ I(\vec{v}, \vec{v}') &:= \int \frac{d\Omega_{\hat{k}}}{4\pi} \left[\frac{2(1 - \vec{v} \cdot \vec{v}')}{(1 - \hat{k} \cdot \vec{v})(1 - \hat{k} \cdot \vec{v}')} - \frac{m^2}{E^2} \left(\frac{1}{(1 - \hat{k} \cdot \vec{v})^2} + \frac{1}{(1 - \hat{k} \cdot \vec{v}')^2} \right) \right] \Rightarrow \\ \mathcal{E}_{soft}(k_{max}) &= \frac{e^2}{4\pi^2} \int dk I(\vec{v}, \vec{v}') \approx \frac{\alpha}{\pi} k_{max} I(\vec{v}, \vec{v}') \end{aligned}$$

The integral is actually divergent at the upper end. It has been cut off at k_{max} which is provided by the *inverse of the duration over which the electron receives the kick*. This is given by $k_{max} \sim |\vec{p} - \vec{p}'| := |\vec{q}|$. This is a physical cutoff.

The angular integration receives the maximum contribution from when \hat{k} is parallel to \vec{v} or \vec{v}' depending upon if it is the initial or the final state bremsstrahlung. Given the directions $v\hat{v}$, $v'\hat{v}'$, the \hat{k} varies between these two directions (smaller angle). Thus, to pick up contribution when \hat{k} is parallel to \vec{v} , we define $\cos\theta$ by $\hat{k} \cdot \vec{v} = v\cos\theta$ which varies between $|\vec{v}'|\cos\theta = \vec{v} \cdot \vec{v}'$ and $\cos\theta = 1$. Similarly for contribution from nearly parallel to \vec{v}' , we define $\cos\theta$ by $\hat{k} \cdot \vec{v}' = |\vec{v}'|\cos\theta$ which varies from $|\vec{v}'|\cos\theta = \vec{v} \cdot \vec{v}'$ to $\cos\theta = 1$.

Additionally, in the extreme relativistic limit where we can neglect the m^2/E^2 terms, we can approximate $(1 - \hat{k} \cdot \vec{v})(1 - \hat{k} \cdot \vec{v}') \approx (1 - |\vec{v}'|\cos\theta)(1 - \vec{v} \cdot \vec{v}')$ or $\approx (1 - |\vec{v}'|\cos\theta)(1 - \vec{v} \cdot \vec{v}')$. The angular integral then approximates as,

$$\begin{aligned} I(\vec{v}, \vec{v}') &\approx \int_{|\vec{v}'|\cos\theta=\vec{v}\cdot\vec{v}'}^1 d\cos\theta \frac{1 - \vec{v} \cdot \vec{v}'}{(1 - v\cos\theta)(1 - \vec{v} \cdot \vec{v}')} + \int_{|\vec{v}'|\cos\theta=\vec{v}\cdot\vec{v}'}^1 d\cos\theta \frac{1 - \vec{v} \cdot \vec{v}'}{(1 - v'\cos\theta)(1 - \vec{v} \cdot \vec{v}')} \\ &\simeq \ln\left(\frac{1 - \vec{v} \cdot \vec{v}'}{1 - |\vec{v}'|}\right) + \ln\left(\frac{1 - \vec{v} \cdot \vec{v}'}{1 - |\vec{v}'|}\right) \approx \ln\left(\frac{(p \cdot p')^2}{E^2(E - p)(E - p')}\right) \\ &\simeq 2\ln\frac{q^2}{m^2} \quad \text{where } q := p' - p \end{aligned}$$

We have used: $(1 - \vec{v} \cdot \vec{v}') = -p \cdot p'/E^2$, $(1 - |\vec{v}'|) = (E - |\vec{p}'|)/E$, $(E - |\vec{p}'|)(E - |\vec{p}|) \approx (E - |\vec{p}'|)^2$ and $E(E - |\vec{p}'|) \approx E^2 - \vec{p}^2 = m^2$ in getting to the last equation.

Thus the radiated energy in the soft modes is given by,

$$\mathcal{E}_{soft}(k_{max}) = \frac{\alpha}{\pi} \int_0^{k_{max}} dk I(\vec{v}, \vec{v}') \xrightarrow{E \gg m} \frac{2\alpha}{\pi} \int_0^{k_{max}} dk \ln(q^2/m^2). \quad (17.22)$$

Interpreting the above energy classically, what would be the total number of photons emitted, N_γ ? Notice that $I(\vec{v}, \vec{v}')dk$ is the contribution to the radiated energy from the

Fourier modes with energy, $|\vec{k}|$. The equivalent number of photons would be $\frac{I(\vec{v}, \vec{v}') dk}{|\vec{k}|}$ and the total number N_γ is obtained by integrating over the energies:

$$N_\gamma = \frac{\alpha}{\pi} \int_0^{k_{max}} \frac{dk}{k} I(\vec{v}, \vec{v}') \approx \frac{2\alpha}{\pi} \int_0^{k_{max}} \frac{dk}{k} \ln(q^2/m^2)$$

This is clearly divergent from the lower limit.

But this is the same expression for the total quantum mechanical probability for emission of a soft photon up to energy k_{max} and thus is divergent.

This is the IR divergence of the QED cross-section for the bremsstrahlung process.

If the photon is given a small mass μ , then the $\int dk/k$ will give $\ln(|q| \simeq k_{max})/\mu$ while $I(\vec{v}, \vec{v}')$ gives the $\ln(q^2/m^2)$. Hence,

$$\boxed{d\sigma(p \rightarrow p' + k) = d\sigma(p \rightarrow p') \cdot \frac{\alpha}{\pi} \underbrace{[\ln(q^2/\mu^2) \cdot \ln(q^2/m^2)]}_{\text{Sudakov double log}}} \quad (17.23)$$

18. TREATMENT OF DIVERGENCES:

At the first non-trivial attempt at computing radiative corrections, we encounter divergences of the UV type (from large loop momentum) and of IR type (from small loop momentum in massless propagators). How do we understand the physical origin of these? How do we adjust the computational procedure so as to make *unambiguous predictions* to be confronted with observations? Let us recapitulate what we have got.

- Quite generally, using only Poincare covariance and assumption about the possible spectrum of a theory with mass gap, the Kallen-Lehmann representation gave us,

$$\begin{aligned}\Delta'_F(p) &= Z\Delta_0(p, m_{ph}) + \int_{m_{th}^2}^{\infty} d\sigma^2 \rho(\sigma^2) \Delta_0(p, \sigma) \quad , \quad 0 < Z < 1 \\ \Delta_0(p, \sigma) &:= \frac{-i}{p^2 + \sigma^2 - i\epsilon} \quad (\text{Free propagator of mass } \sigma) \\ S'_F(p) &= Z_2 S_0(p, m_{ph}) + \int_{m_{th}^2}^{\infty} d\sigma^2 \frac{-i\{\rho_1(\sigma^2)\not{p} + \rho_2(\sigma^2)\}}{p^2 + \sigma^2 - i\epsilon} \\ S_0(p, \sigma) &:= -i \frac{(-\not{p} + \sigma)}{p^2 + \sigma^2 - i\epsilon} \quad (\text{Free propagator of mass } \sigma) \\ (D'_F)_{\mu\nu}(q) &= \left(\eta_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \right) \left\{ \frac{Z_3}{q^2 - i\epsilon} + \int_0^{\infty} d\sigma^2 \frac{\Pi(\sigma^2)}{q^2 + \sigma^2 - i\epsilon} \right\}\end{aligned}$$

The Z 's are the field (or wavefunction) renormalization constants. They are determined as the residues at the (isolated) pole at the *physical mass*, m_{ph} . This is without any perturbation series.

In perturbation series though we obtained (for fermion and photon),

$$S'_F(p) = \frac{1}{\not{p} + m - \Sigma(\not{p})} \quad \& \quad D'_F(q) = \left(\eta_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \right) \left[\frac{1}{q^2} \frac{1}{1 - \Pi(q^2)} \right]$$

This is consistent with the general expectation that the physical masses are determined by the poles in the exact propagator while the corresponding residues determine the Z 's. Thus,

$$\delta m := m_{ph} - m = -\Sigma(\not{p} = -m_{ph}) \quad , \quad Z_2^{-1} = 1 - \frac{d\Sigma(\not{p})}{d\not{p}} \Big|_{\not{p} = -m_{ph}} \quad , \quad Z_3^{-1} = 1 - \Pi(0).$$

and of course the photon physical mass is zero.

- The vertex function for on shell fermion masses takes the form,

$$\gamma^\mu(p, p') := \gamma^\mu F_1(q^2) + i\sigma^{\mu\nu} q_\nu F_2(q^2) \quad , \quad \sigma_{\mu\nu} := \frac{i}{2} [\gamma_\mu, \gamma_\nu] \quad , \quad \not{p} = -m_{ph} = \not{p}' \quad .$$

This decomposition is understood to be sandwiched between $\bar{u}(p')$ and $u(p)$. Since $m_{ph} = m + \delta m = m + o(\alpha)$, for the first order corrections, we can take $m_{ph} = m$. Here are the expressions we obtained:

$$(I) \quad \Sigma(p) = \frac{\alpha}{2\pi} \int_0^1 dx (2m + xp) \ln \frac{x\Lambda^2}{x(1-x)p^2 + (1-x)m^2 + x\mu^2 - i\epsilon} \quad (18.1)$$

$$\delta m = \frac{\alpha}{2\pi} \int_0^1 dx (2-x) \ln \frac{x\Lambda^2}{(1-x)^2 m^2 + x\mu^2}, \quad (18.2)$$

$$Z_2 = 1 + \frac{\alpha}{4\pi} \ln(\Lambda^2/m^2) \quad (18.3)$$

$$(II) \quad \Pi(q^2) = -\frac{2\alpha}{\pi} \int_0^1 dx x(1-x) \left\{ \frac{2}{\epsilon} - \gamma - \ln \left(\frac{m^2 + q^2 x(1-x)}{\mu_0^2} \right) \right\} \quad (18.4)$$

$$= -\frac{2\alpha}{3\pi} \frac{1}{\epsilon} + o(\epsilon^0) \Rightarrow Z_3 = 1 - \frac{2\alpha}{3\pi} \frac{1}{\epsilon} \quad (18.5)$$

$$(III) \quad F_1(q^2) = 1 + \frac{\alpha}{2\pi} \int dx dy dz \delta(1-x-y-z) \left\{ \ln \left(\frac{z\Lambda^2}{M^2} \right) \right. \\ \left. - \frac{-(1-x)(1-y)q^2 + (1-4z+z^2)m^2}{M^2} \right\} \quad (18.6)$$

$$F_2(q^2) = \frac{\alpha}{2\pi} \int dx dy dz \delta(1-x-y-z) \left\{ \frac{2m^2 z(1-z)}{M^2} \right\}, \quad (18.7)$$

$$M^2 := q^2 xy + m^2(1-z)^2 + \mu^2 z \quad (18.8)$$

$$\therefore F_1(0) = 1 + \frac{\alpha}{2\pi} \int dx dy dz \delta(1-x-y-z) \left\{ \ln \left(\frac{z\Lambda^2}{(1-z)^2 m^2} \right) \right. \\ \left. + \frac{1-4z+z^2}{(1-z)^2} \right\} \quad (18.9)$$

$$F_2(0) = \frac{\alpha}{2\pi} \int dx dy dz \delta(1-x-y-z) \left(\frac{z}{1-z} \right) = \frac{\alpha}{2\pi} =: \frac{g-2}{2} \quad (18.10)$$

$$(IV) \quad d\sigma_{p \rightarrow p'+k} = d\sigma_{p \rightarrow p'} \int \frac{d^3 k}{2k(2\pi)^2} \sum_{\lambda} e^2 \left(\frac{\vec{\epsilon} \cdot \vec{p}'}{\vec{p} \cdot \vec{k}} - \frac{\vec{\epsilon} \cdot \vec{p}}{\vec{p}' \cdot \vec{k}} \right)^2 \quad (18.11)$$

$$= d\sigma_{p \rightarrow p'} \left[\frac{e^2}{4\pi^2} \int \frac{dk}{k} I(\vec{v}, \vec{v}') \right] \quad \text{where,} \quad (18.12)$$

$$I(\vec{v}, \vec{v}') = \int \frac{d\Omega}{4\pi} \left\{ \frac{2(1 - \vec{v} \cdot \vec{v}')}{(1 - \hat{k} \cdot \vec{v})(1 - \hat{k} \cdot \vec{v}')} - \frac{m^2/E^2}{(1 - \hat{k} \cdot \vec{v})^2} - \frac{m^2/E^2}{(1 - \hat{k} \cdot \vec{v}')^2} \right\} \quad (18.13)$$

$$\text{where } p^\mu = (E, E\vec{v}), (p')^\mu = (E, E\vec{v}'), k^\mu = (|\vec{k}|, \vec{k}) \text{ is used.} \quad (18.14)$$

For $|\vec{v}| \approx |\vec{v}'| \approx 1$, $I(\vec{v}, \vec{v}') \approx 2\ln(\bar{q}^2/m^2)$, $q := p' - p \Rightarrow$

$$d\sigma_{p \rightarrow p'+k} = d\sigma_{p \rightarrow p'} \left[\frac{\alpha}{\pi} \ln(\bar{q}^2/\mu^2) \ln(\bar{q}^2/m^2) \right] \quad (18.15)$$

A. Treatment of the IR divergences

Let us consider the IR problem first. Since F_1 has both the UV and the IR divergence, we will separate these by using the UV renormalized $F_{1,ren}$ which was defined through, $\delta F_{1,ren}(q^2) := \delta F_1(q^2) - \delta F_1(0)$. This guarantees that the renormalized F_1 satisfies the condition $F_1(0) = 1$. Using the expressions above, we see that,

$$\delta F_{1,ren}(q^2) = \frac{\alpha}{2\pi} \int dx dy dz \delta(1-x-y-z) \left\{ \frac{-(1-x)(1-y)q^2 + (1-4z+z^2)m^2}{M^2(q^2)} - \frac{(1-4z+z^2)m^2}{(1-z)^2m^2 + z\mu^2} \right\}$$

The IR divergence comes from $z \rightarrow 1 \leftrightarrow x \sim y \sim 0 \leftrightarrow M(q^2, x, y)$ would vanish but for the photon mass μ^2 . To isolate the divergence, it suffice to take $x = y = 0$, $z = 1$ in the numerator. Doing the x integration using the delta function gives $x = 1 - y - z > 0 \Rightarrow y < 1 - z$ and leads to,

$$\delta F_{1,ren} \approx \frac{\alpha}{2\pi} \int_0^1 dz \int_0^{1-z} dy \left\{ \frac{-2m^2 - q^2}{(1-z)^2m^2 + y(1-y-z)q^2 + \mu^2} \frac{2m^2}{(1-z)^2m^2 + \mu^2} \right\}$$

Substituting $y := (1-z)u$, $v := 1-z$, the leading contribution in the limit $\mu \rightarrow 0$ is expressed as [13],

$$\boxed{F_{1,ren}(q^2) \approx 1 - \frac{\alpha}{2\pi} f_{IR}(q^2) \ln \left(\frac{m^2 \text{ or } q^2}{\mu^2} \right)} \quad \text{where,} \quad (18.16)$$

$$\boxed{f_{IR}(q^2) = \int_0^1 du \left[\frac{m^2 + q^2/2}{m^2 + u(1-u)q^2} \right] - 1} \quad (18.17)$$

Since F_1 is the coefficient of the γ^μ term, we can replace $e \rightarrow eF_{1,ren}$ in the electron scattering off a *classical potential*. The cross-section is then given by,

$$\left(\frac{d\sigma}{d\sigma} \right) \simeq \left(\frac{d\sigma}{d\sigma} \right)_{tree} \left[1 - \frac{\alpha}{\pi} f_{IR}(q^2) \ln \left(\frac{m^2 \text{ or } q^2}{\mu^2} \right) \right].$$

Notice that $e \rightarrow eF_1 \Rightarrow e^2 \rightarrow e^2 F_1^2 \Rightarrow \alpha/(2\pi) \rightarrow \alpha/(\pi)$.

Not only is this divergent as $\mu \rightarrow 0$, for non-zero μ it is actually *negative* implying negative cross-section! In the limit of $q^2 \rightarrow \infty$ (q^2 is space-like and hence positive), $f_{IR}(q^2) \rightarrow \ln(q^2/m^2) \Rightarrow F_{1,ren}(q^2) \simeq 1 - \frac{\alpha}{2\pi} \ln(q^2/m^2) \ln(q^2/\mu^2)$ and hence,

$$\boxed{d\sigma(p \rightarrow p') \simeq d\sigma_{tree}(p \rightarrow p') \left[1 - \frac{\alpha}{\pi} \ln(q^2/m^2) \ln(q^2/\mu^2) \right] , \quad q^2 \rightarrow \infty , \quad \mu^2 \rightarrow 0.} \quad (18.18)$$

The Bremsstrahlung cross-section on the other hand is,

$$\boxed{d\sigma(p \rightarrow p' + k) = d\sigma(p \rightarrow p') \left[\frac{\alpha}{\pi} \ln \frac{q^2}{m^2} \ln \frac{q^2}{\mu^2} \right] = d\sigma(p \rightarrow p')_{tree} \left[\frac{\alpha}{\pi} \ln \frac{q^2}{m^2} \ln \frac{q^2}{\mu^2} \right].} \quad (18.19)$$

Both the cross-sections above are IR divergent and both suffer from ambiguity from contamination of soft photons.

We already noted that detectors are unable to distinguish a charged particle accompanied by soft photons below detector sensitivity. What is the appropriate theoretically computed quantity which reflects this limitation of the detection process?

When an experimenter reports a detection of a scattered electron, he/she is actually giving an estimate of the probability that *an electron, $e(p')$ is detected and a photon is not detected*. This probability is the probability for a process with no emitted photon plus the probability that there are accompanying soft photons with energies below the detection threshold, ε_{th} . In equation,

$$\boxed{(d\sigma)_{measured} = d\sigma (p \rightarrow p') + d\sigma (p \rightarrow p' + k ; |\vec{k}| < \varepsilon_{th}) .} \quad (18.20)$$

But this is precisely the sum of the two cross-sections given in eqns.(18.18,18.19). We see that the leading contribution as $q^2 \rightarrow \infty, \mu^2 \rightarrow 0$ is exactly canceled out in the observed cross-section!

Note: For a general q^2 , the sum of the two cross-sections is given by [13],

$$(d\sigma)_{measured} = d\sigma_{tree}(p \rightarrow p') \left\{ 1 - \frac{\alpha}{\pi} f_{IR}(q^2) \ln \left(\frac{q^2 \text{ or } m^2}{\mu^2} \right) + \frac{\alpha}{2\pi} I(\vec{v}, \vec{v}') \ln(\varepsilon_{th}/\mu^2) \right\}$$

It turns out that without the limit $q^2 \rightarrow \infty$, it still holds that $I(\vec{v}, \vec{v}') \rightarrow 2f_{IR}$ [13]. For general q^2 , the coefficient of f_{IR} is $\ln(q^2)$ or $\ln(m^2)$. For large q^2 we can of course drop m^2 . Furthermore, experimentally it is easier to track the behavior of the cross-section as a function of q^2 , so taking $q^2 \gg m^2$, we write the unambiguous and measurable prediction as,

$$\boxed{\left(\frac{d\sigma}{d\Omega} \right)_{measured} = \left(\frac{d\sigma}{d\Omega} \right)_{tree} (p \rightarrow p') \left[1 - \frac{\alpha}{\pi} \ln \frac{q^2}{m^2} \ln \frac{q^2}{\varepsilon_{th}^2} + o(\alpha^2) \right]_{q^2 \gg m^2}} \quad (18.21)$$

Appreciate that we had parametrised the IR divergence in terms of the photon mass μ^2 ; we used the renormalized $F_{1,ren}(q^2)$ to separate the IR from the UV and finally, identified the quantity which is actually reported by experiments which is the sum of the two cross-sections.

This has been a demonstration at the 1-loop. It is non-trivial result of a great deal of work, that the basic mechanism of cancellation works to all orders in α . There are other types of divergences analogous to the IR divergences, the so called “mass singularities” which do not cancel but can be factorised in a convenient form and then eliminated from the observed cross-sections. See the book by G. Sterman [14].

UV divergences is a different ball game and needs a different procedure.

B. Treatment of the UV divergences

All the three radiative corrections in, Σ, Π and F_1 have the UV divergences parametrised in terms of the Pauli-Villars cut-off Λ or the dimensional regularization ϵ^{-1} pole. Note that the bremsstrahlung process does not have a UV divergence as there is physical cut-off for the energy of the “soft photons”. We need to pay attention to the Z factors now.

Recall that the S -matrix element definition, via the LSZ reduction procedure, had a factor of $\frac{1}{\sqrt{Z}}$ for each external line. There was also an amputation of the external line, effected by the equation of motion operator eg $\square - m^2$ acting on the Feynman propagator Δ_F for the external line. The \sqrt{Z} entered from the asymptotic condition relating the interacting field to the in/out fields. The in/out fields satisfy their respective field equations with *physical masses* and also have the normalization factors with ω_k which also contain the physical masses. In the momentum space, these factors associated with external lines are of the form $\frac{1}{\sqrt{Z}}(p^2 + m^2)\Delta'_F(p)$ where the Δ_F contains the self-energy giving it the form $\Delta'_F(p) = (p^2 + m^2 - \Pi(p^2))^{-1}$ (for scalars).

In perturbative computation of the self-energy however, the m is the mass parameter in the \mathcal{L} which is not the physical mass. In fact from the Kallen-Lehmann representation, we know that the physical mass is determined from $p^2 + m^2 - \Pi(p^2)|_{p^2 = -m_{ph}^2} = 0 \leftrightarrow -m_{ph}^2 + m^2 = \Pi(-m_{ph}^2)$. Expanding $\Pi(p^2)$ about $-m_{ph}^2$ gives us,

$$\begin{aligned} \Pi(p^2) &= \pi(-m_{ph}^2) + (p^2 + m_{ph}^2)\Pi'(-m_{ph}^2) + \dots \Rightarrow \\ (\Delta'_F)^{-1}(p) &= p^2 + m^2 - \Pi(-m_{ph}^2) - (p^2 + M_{ph}^2)\Pi'(-m_{ph}^2) + \dots \\ &= (p^2 + m_{ph}^2) \{1 - \Pi'(-m_{ph}^2) + \dots\} = (p^2 + m_{ph}^2) \{Z_\Phi^{-1} + \dots\} \\ \therefore \frac{1}{\sqrt{Z_\Phi}}(p^2 + m_{ph}^2)\Delta'_F(p) &= \sqrt{Z_\Phi} \{1 + o((p^2 + m_{ph}^2)^2)\} \end{aligned}$$

Since we evaluate the S -matrix elements on shell, all the higher order terms vanish. The

net result is that:

In an S -matrix element, for each external line introduce a factor of $\sqrt{Z_{field}} \times$ appropriately normalized wavefunction and now do not include self-energy corrections on the external lines.

Consider the electron scattering off external potential. The corresponding invariant amplitude is given by $e(\sqrt{Z_2})^2 \bar{u}(p') \Gamma^\mu(p', p) u(p)$ with $\Gamma^\mu = \gamma^\mu F_1 + (\dots) F_2$. The F_2 has no divergences while Z_2, F_1 both are divergent. Thus we write, $Z_2 = 1 + \delta Z_2, F_1 = 1 + \delta F_1$ with the δ 's representing the $o(\alpha)$ divergent corrections. Thus we write,

$$\begin{aligned} \mathcal{M}_{e \rightarrow e}(p', p) &= e(1 + \delta Z_2) \left\{ \bar{u}(p') \gamma^\mu u(p) (1 + \delta F_1(q^2)) + \bar{u}(p') \frac{i\sigma^{\mu\nu} q_\nu}{2m_{ph}} u(p) \delta F_2(q^2) \right\} \\ &= e [\bar{u}(p') \gamma^\mu u(p)] \{1 + \delta F_1 + \delta Z_2\} + \bar{u}(p') \frac{i\sigma^{\mu\nu} q_\nu}{2m_{ph}} u(p) \delta F_2(q^2) \quad \text{But,} \\ \delta Z_2 &= \frac{\alpha}{2\pi} \int_0^1 dz \left[-z \ln \frac{z\Lambda^2}{M^2} + 2(2-z) \frac{z(1-z)m^2}{M^2} \right], \quad M^2 = (1-z)^2 m^2 + z\mu^2 \\ \delta F_1(0) &= \frac{\alpha}{2\pi} \int_0^1 dz (1-z) \left[\ln \frac{z\Lambda^2}{M^2} + (1+z^2-4z) \frac{m^2}{M^2} \right] \\ \therefore \delta F_1(0) + \delta Z_2 &= \frac{\alpha}{2\pi} \int_0^1 dz \left[(1-2z) \ln \frac{z\Lambda^2}{M^2} + \frac{m^2}{M^2} \times \right. \\ &\quad \left. \{ (4-2z)(1-z)z + (1-z)(1+z^2-4z) \} \right] \end{aligned}$$

Integrate the first term by parts,

$$\begin{aligned} \int_0^1 dz (1-2z) \ln \frac{z\Lambda^2}{M^2} &= (z-2z^2) \ln \frac{z\Lambda^2}{M^2} \Big|_0^1 - \int_0^1 dz (z-z^2) \left\{ \frac{1}{z} - \frac{2(1-z)(-1)m^2 + \mu^2}{M^2} \right\} \\ &= - \int_0^1 dz \frac{1-z}{M^2} \{ M^2 + 2z(1-z)m^2 - z\mu^2 \} \\ &= - \int_0^1 dz \frac{1-z}{M^2} \{ m^2(1-z)(1+z) \} = - \int_0^1 dz \frac{m^2(1-z)^2(1+z)}{m^2 \left((1-z)^2 + z \frac{\mu^2}{m^2} \right)} \\ \therefore \delta F_1(0) + \delta Z_2 &= \frac{\alpha}{2\pi} \int_0^1 dz \left[- \frac{m^2(1-z)(1-z^2)}{M^2} + \frac{m^2}{M^2} (1-z) \{ 4z - 2z^2 + 1 + z^2 - 4z \} \right] \\ &= \frac{\alpha}{2\pi} \int_0^1 dz \frac{m^2}{M^2} (1-z)(1-z^2) \{ -1 + 1 \} = 0 \quad (!) \end{aligned}$$

Thus not only has the $\ln \Lambda^2$ divergence canceled, we have got $\delta Z_2 = -\delta F_1(0)$ and thus the entire correction takes the form,

$$\mathcal{M}_{e \rightarrow e}(p', p) = e \bar{u}(p') \gamma^\mu u(p) \left\{ 1 + \underbrace{\delta F_1(q^2) - \delta F_1(0)}_{\delta F_{1,ren}} \right\} + i \bar{u}(p') \frac{i\sigma^{\mu\nu} q_\nu}{2m_{ph}} u(p) \delta F_2(q^2)$$

We recover the *ad hoc* prescription of using $\delta F_{1,ren}$ introduced earlier while discussing the IR divergences. Note that the subtracted term was crucial for the IR divergence cancellation. It was of course physically expected since $\delta F_1(q^2 = 0) = 0$ should actually hold to all orders. Note that the UV divergence in the vertex function, $\delta F_1(q^2)$, is canceled against the divergence in δZ_2 coming from electron self-energy.

Now we are left with the divergence in Z_3 from the $\Pi(q^2 = 0)$ for the photon self-energy. We still have the electron mass-shift δm which is divergent.

1. 1-Loop Renormalization: Charge Screening and Lamb shift

Consider charged particle scattering by exchanging a photon, necessarily off-shell (space-like in fact). Replace the free propagator $(D_F)_{\mu\nu}(q^2)$ by the exact propagator which includes the photon self-energy, $(D'_F)_{\mu\nu}(q^2)$. The exact propagator has additional terms $q_\mu q_\nu$. But thanks to the Ward identity, these do not contribute to S -matrix elements and effectively, $(D'_F)_{\mu\nu}(q^2) = -i\eta_{\mu\nu}[q^2(1 - \Pi(q^2))]^{-1}$. This has a pole at $q^2 = 0$ with residue $Z_3 = [1 - \Pi(0)]^{-1}$. The replacement thus gives a factor of Z_3 . The photon internal line connects two vertices contributing e^2 . Thus, if we *identify* $\boxed{e^2 Z_3 =: e_{ph}^2}$, then all divergences coming from the photon self-energy can be neatly absorbed in the *physical, measured charge* which is finite. For the mass parameter too, we identified $m_{ph}^2 = m^2 + \delta m$, which absorbs the divergence in δm into the Lagrangian parameter m . The identification $e_{ph} := e\sqrt{Z_3}$ is called *charge renormalization* while $m_{ph}^2 := m^2 + \delta m$ is called *mass renormalization*. For external photon line, we will just have $\sqrt{Z_3}$ factor and no self-energy corrections and of course no mass shift.

Thus the exact photon propagator may be replaced by the free propagator by simultaneously replacing α by α_{eff} defined below.

$$\alpha(D'_F)_{\mu\nu}(q^2) = (D_F)_{\mu\nu}(q^2) \frac{\alpha}{1 - \Pi(q^2)} = \frac{\eta_{\mu\nu}}{q^2 - i\epsilon} \left[\frac{\alpha_{ph}(1 - \Pi(0))}{1 - \Pi(q^2)} \right] := \alpha_{eff}(q^2) \left[\frac{\eta_{\mu\nu}}{q^2 - i\epsilon} \right]$$

with, $\boxed{\alpha_{eff}(q^2) = \frac{\alpha_{ph}}{1 - (\Pi(q^2) - \Pi(0))}}$

To appreciate the implication of the above procedure, recall the discussion of the effective potential inferred from the tree level scattering for both the Yukawa and the QED coupling. Let us use the same formula (for QED), but include the self-energy correction for the photon

propagator. The inferred potential is,

$$V(\vec{x}) = \int \frac{d^3q}{(2\pi)^3} e^{i\vec{q}\cdot\vec{x}} \frac{-e_{ph}^2}{\vec{q}^2(1 - \Pi(q^2) + \Pi(0))} \quad (18.22)$$

$$\Pi(q^2) = -\frac{2\alpha}{\pi} \int_0^1 dx x(1-x) \left\{ \frac{2}{\epsilon} - \gamma - \ln \left(\frac{m^2 + q^2 x(1-x)}{\mu_0^2} \right) \right\} \quad (18.23)$$

$$\Pi(q^2) - \Pi(0) = -\frac{2\alpha}{\pi} \int_0^1 dx x(1-x) \left\{ \ln \left(\frac{m^2}{m^2 + q^2 x(1-x)} \right) \right\} \quad (18.24)$$

$$\xrightarrow{q^2 \ll m^2} \frac{2\alpha}{\pi} \int_0^1 dx x^2(1-x)^2 \frac{q^2}{m^2} = \frac{\alpha}{15\pi} \frac{q^2}{m^2} \quad (\text{NR limit}) \quad (18.25)$$

$$\therefore V(\vec{x}) \approx \int \frac{d^3q}{(2\pi)^3} e^{i\vec{q}\cdot\vec{x}} \frac{-e_{ph}^2}{\vec{q}^2(1 - \frac{\alpha}{15\pi} \frac{q^2}{m^2})} \quad (18.26)$$

$$\approx -\frac{\alpha_{ph}}{r} \underbrace{\frac{4\alpha_{ph}^2}{15m^2}}_{\text{perturbation}} \delta^3(\vec{x}) \quad (18.27)$$

The perturbed Coulomb potential induces a shift in the energy levels of the Coulomb potential, say in Hydrogen atom. The first order perturbation theory gives the shift as,

$$\Delta E = -\frac{4\alpha_{ph}^2}{15m^2} \frac{\alpha_{ph}^3 m^3}{8\pi} = -\frac{\alpha_{ph}^5 m}{30\pi} \simeq -1.123 \times 10^{-7} \text{eV (Lamb Shift)} \quad (18.28)$$

This is the contribution of the photon self-energy to the famous Lamb shift. The photon self-energy $\leftrightarrow \alpha_{eff}$ is thus an observable and observed effect. More exact calculation may be seen in [13].

We can also consider the ultra-relativistic regime $q^2 \gg m^2$. Now

$$\begin{aligned} \Pi(q^2) - \Pi(0) &= -\frac{2\alpha}{\pi} \int_0^1 dx x(1-x) \left\{ \ln \left(\frac{m^2}{q^2} \right) - \ln \left(x(1-x) + \frac{m^2}{q^2} \right) \right\} \\ &\approx -\frac{2\alpha}{\pi} \left[\frac{1}{6} \ln(m^2/q^2) - \int_0^1 dx x(1-x) \ln \{ x(1-x) + o(m^2/q^2) \} \right] \\ &\simeq \frac{\alpha}{3\pi} \left[\ln(q^2/m^2) - 5/3 + o(m^2/q^2) \right] \\ &\therefore \alpha_{eff} = \frac{\alpha_{ph}}{1 - \frac{\alpha}{3\pi} \ln \left(\frac{q^2}{m^2 e^{5/3}} \right)} \xrightarrow{q^2 \gg m^2} -\frac{3\pi}{\ln(q^2 e^{-5/3}/m^2)} \end{aligned} \quad (18.29)$$

Note that the denominator is less than 1 and hence effective coupling is *larger* than the physical coupling. Thus, the effective coupling gets stronger at shorter distances. This is interpreted as saying that the photon self-energy correction polarizes the space between say the nucleus and the electron, shielding the nuclear charge. As the shielding cloud is

penetrated, higher nuclear charge is seen and hence α_{eff} is larger. For this reason, the photon self-energy $\Pi(q^2)$ is also called the *vacuum polarization*.

To summarize:

1. The UV divergences in the self energies and the vertex correction are absorbed away by introducing the physical mass for electron, physical charge for the electron and electron wavefunction renormalization constant to cancel the divergence in $F_1(q^2)$;
2. The IR divergences in the amplitudes, imply IR divergence in the cross-section. For the $e - e$ scattering, the IR divergence is canceled against the bremsstrahlung process, once the measured quantity is correctly identified taking into account the finite detector resolution;
3. The hiding of divergences in α_{ph} and m_{ph}^2 imply that $\alpha_{eff}, \delta m^2$ acquire q^2 dependence - they “run” with q^2 .

This seems satisfactory within the context of the $o(\alpha)$ corrections and the magic of the Ward identity. But it raises more questions.

- Are the divergences generic? What are the mathematical and physical reasons for their existence?
- Can we sometimes/always take care of the divergences and make unambiguous predictions?
- What kind of predictions can be made?
- What are the effective/running parameters?
- Are the divergences an artifact of perturbation theory? Etc, etc

Addressing these questions is the genesis of the renormalization theory.

As a matter of strategy, we will stay within perturbative framework and try to make sense of the divergences. After all, despite divergences, we could obtain prediction from QED at 1-loop which have been well tested!

We begin with a comment that the UV divergences persist at higher loops as well and have to be faced. Can we always absorb them away in physical masses, couplings and the field renormalizations in S -matrix elements/cross-sections automatically and make finite, unambiguous predictions? Indeed it is so for QED and for the class of (super-)renormalizable theories. The proof needs a somewhat modified procedure, introducing additional diagrams,

Note that we have only rewritten the bare Lagrangian density split into two sets of terms. Since we have not stipulated any conditions, the split is completely arbitrary. Two sets of conditions, called *renormalization conditions*, are now provided: (i) The exact propagator is given by $\frac{-i}{p^2+m^2-i\epsilon}$ + terms regular at $p^2 = -m^2$. That is, the renormalized mass should be identified with the physical mass *and* the residue should be 1;

(ii) The exact 4-point, 1PI function should equal $-i\lambda$ at $s = 4m^2, t = u = 0$. s, t, u are the Mandelstam variables.

The definition of λ could be changed, but essentially it is the value of the exact 4-point scattering amplitude which is measurable.

Diagrams generated by these vertices will again have divergences which must be regulated. The counter terms coefficients, $\delta_Z, \delta_m, \delta_\lambda$ are to be so chosen as to ensure that the renormalization conditions are satisfied, order-by-order. Since the conditions are finite and cut-off independent (regularization independent), all divergences are spent in defining the counter term coefficients. By construction, we have generated finite quantities to all orders!

Note: There are different ways of absorbing the divergences in the counter term coefficients, especially when massless particles are involved. The different methods of absorbing the UV divergences is generically referred to as a *subtraction scheme*. This will be illustrated below.

The perturbation series generated using the bare form of the Lagrangian is called “bare perturbation series” while that generated using the renormalized quantities together with the renormalization conditions, is called “renormalized perturbation series”.

Does such a simple splitting procedure always generate UV finite (cut-off independent) quantities in terms of *finitely many* renormalization conditions? The answer is YES for a class of theories called “renormalizable theories”. There are non-renormalizable theories (= Lagrangians) for which this procedure fails.

19. RENORMALIZED PERTURBATION SERIES

Let a theory be specified by a (bare) Lagrangian density, \mathcal{L} as,

$$\mathcal{L}(\Phi_0, m_0, g_{0,k}) = -\frac{1}{2}\partial_\mu\Phi_0\partial^\mu\Phi_0 - \frac{1}{2}m_0^2\Phi_0^2 - \sum_{k\geq 3} \frac{g_{0,k}}{k!}\Phi_0^k.$$

Introduce renormalized quantities, Φ, m^2, g_k and *scaling parameters*, Z_ϕ, Z_m, Z_{g_k} through the definitions: $\Phi_0 =: \sqrt{Z_\phi}\Phi$, $m_0^2 Z_\phi = Z_m m^2$, $g_{0,k} Z_\phi^{k/2} =: Z_{g_k} g_k$. Writing the Z 's in $1 + \delta$ form, we recast the Lagrangian density as,

$$\begin{aligned} \mathcal{L}(\Phi, m, g_k; \delta_\Phi, \delta_m, \delta_k) = & -\frac{1}{2}\partial_\mu\Phi\partial^\mu\Phi - \frac{1}{2}m^2\Phi^2 - \sum_{k\geq 3} \frac{g_k}{k!}\Phi^k \\ & -\frac{1}{2}\delta_\Phi\partial_\mu\Phi\partial^\mu\Phi - \frac{1}{2}\delta_m m^2\Phi^2 - \sum_{k\geq 3} \delta_k \frac{g_k}{k!}\Phi^k \quad \leftarrow \text{(Counter terms)} \end{aligned}$$

We *choose* the renormalization conditions to be:

(i) Exact propagator = $\frac{-i}{p^2+m^2-i\epsilon}$, which is equivalent to the two conditions: (a) $\Pi(p^2 = -m^2) = 0$, and (b) $\Pi'(p^2 = -m^2) = 0$;

(ii) $g_k = k$ -point amputated function, also called *k-point vertex function*, at some chosen values of its momentum arguments.

Note: Due to the conditions (i), this scheme is called *on-shell renormalization*.

Calculations of the vertex functions will be functions of the external momenta, $m, g_k, \delta_\Phi, \delta_m, \delta_k$ and the regularization parameter - Λ for a momentum cut-off, $\epsilon = 4 - n$ for the dimensional regularization. The renormalization conditions will serve to define the counter term coefficients in terms of the regularization parameter and eliminate them from the vertex functions. We will be left with vertex function having dependence on the momenta, m and the couplings g_k . This is what we seek to understand.

A. Necessary Conditions for UV divergence: Power Counting

We begin by finding *necessary conditions* for occurrence of UV divergence. These are obtained by estimating the Feynman integrals in the region where *all* loop momenta become large. Feynman integrands are rational functions of momenta and in the regime of large momenta, simply give a power of the large momenta.

Consider an arbitrary, connected and topologically connected diagram made up of E -external lines, I -internal lines, n_k - vertices of k^{th} order and L -loops. The loops arise

because we have I number of momenta with $V := \sum_{k \geq 3} n_k$ number of vertices enforcing momentum conservation (and -1 since an overall momentum conservation) condition which leaves some momenta undetermined. All internal momenta are linear functions of some external momenta, p_i and some loop momenta, k_l . There are several integration regions in the $d \times L$ dimensional space (d is the space-time dimension). These regions correspond to various subsets of internal lines vanishing as $q(p_i, k_l)^{-2}$. It suffices to consider region wherein *all* loop momenta diverge which means that all internal momenta also vanish. If we take the loop momenta to diverge as $k_l = \Lambda \hat{k}_l, \Lambda \rightarrow \infty$, then we get the *superficial degree of divergence*, $D = dL - 2I$.

For vertex functions, there are no external lines and $2I = \sum_{k \geq 3} k n_k - E$. The Green's functions will have $+E$. As noted before, the number of loop momenta is given by $L = I - V + 1$. Using these the superficial degree of divergence is given by,

$$D = d - \frac{d-2}{2}E + \sum_{k \geq 3} n_k \left\{ k \left(\frac{d-2}{2} \right) - d \right\}$$

A necessary condition for divergence is that $D \geq 0$.





For $D < 0$, there is no UV divergence, $D = 0$ is (possibly) logarithmically divergent, $D = 1$ is (possibly) linearly divergent, $D = 2$ is (possibly) quadratically divergent. We have taken $k \geq 3$ to have some interaction, we take at least *one* $n_k \neq 0$ for a non-trivial diagram and of course $d \geq 2, E \geq 1$.

As an example, consider $d = 4$. Then $D = 4 - E + \sum_{k \geq 3} n_k (k - 4) \geq 0$ for divergence. That is $4 \geq E + n_3 - \sum_{k \geq 5} n_k (k - 4)$. If we have $n_{k \geq 5} = 0$, i.e. only Φ^3, Φ^4 terms, then the condition for divergence is independent of the number of Φ^4 vertices while with increasing Φ^3 vertices, the E must decrease correspondingly.

- For $n_3 = 0$, we must have $E \leq 4$ i.e. the 2-point and the 4-point vertex functions are superficially divergent, quadratically and logarithmically respectively. For pure Φ^4 theory, the vertex functions with odd number of external lines vanish identically. Thus, in 4 dimensional Φ^4 theory, the self-energy and the 4-point vertex function are divergent to *all orders*. Since we do have $\delta_\Phi, \delta_m, \delta_4$ coefficients, we can satisfy the renormalization conditions to all orders. This theory is *renormalizable*.

- For pure Φ^3 theory, $n_4 \geq 0, D = 4 - E - n_3$. For $n_3 = 1$, we can have only tree level, 3-point function. This gives $D = 0$. But of course there is no loop integration and hence

no divergence (hence superficial degree of divergence is not a sufficient condition). Let us also require $L \geq 1$.

For $n_3 = 2$, we can have 2-point function diagrams  ,  with $E = 2, L = 1, D = 0$ which are logarithmically divergent and also  with $E = 4, D = -1, L = 0$ which are (trivially) “convergent”. For $n_3 = 3, E = 2$ we can have the diagram  which is divergent even though $D < 0$. But this is because of the 1PR nature of the diagram. To exclude this triviality, we stipulate that contributing diagrams should all be 1PI. Then, for all $n_3 \geq 4$, all E -point vertex functions are convergent. This theory thus has divergences (eg 2-point function), but only up to a *finite order*. Beyond that, there are no divergences. Such a theory is called *super-renormalizable*.

- For a pure $\Phi^k, k \geq 5, D = 4 - E + (k - 4)n_k$. With increasing n_k , the number of loops also increase and D keeps increasing. Equivalently, more and more E -point functions turn divergent and we will need infinitely many counter terms to absorb the divergences, in the *same* E -point function. Such a theory is *non-renormalizable*.

- For sake of variety, take $d = 3$, so that $D = 3 - E/2 + n_k(k/2 - 3)$. For $k = 6, D$ is independent of n_k . The 2, 4, 6 point vertex functions need counter terms to all orders and Φ^6 theory is renormalizable.


- Take $d = 6, \Rightarrow D = 6 - 2E + n_k(2k - 6) = 2[(3 - E) + n_k(k - 3)]$. For $k = 3, D$ is independent of n_3 . The 1, 2, 3 point functions are divergent to all orders and the theory is renormalizable.

Remarks:

- The formula for the superficial degree of divergence can be generalised to include fermions and photons. The fermion internal line contributes -1 (instead of -2) to the power counting. The fermion number conservation must also be paid attention to by restricting the types of vertices. If there are derivative couplings, the numerator contributes positive powers to D .

- As noted above, $D \geq 0$ indicates the possibility of a UV divergence. The actual diagram may be less divergent due to symmetries eg $\Pi(q^2)$ in QED is only log divergent though $D = 2$ suggests quadratic divergence. It is also possible the coefficient of the indicated divergence is

actually zero! An example of this is the photon 3–point vertex function in QED, at 1-loop.



Here $D = 4 - 3 = 1$. However due to *Furry's theorem*, in QED, any photon amplitude with odd number of external lines is zero. Hence the diagram is identically zero. However, what is true is that *if $D < 0$ for a diagram and each of its sub-diagram, then the diagram has no UV divergence* (Dyson-Weinberg theorem).

The superficial degree of divergence, with the caveats mentioned above, suggest a classification of theories as:

- (i) Super-renormalizable: Only a finitely many diagrams have $D \geq 0$. The D has a dependence on the number of vertices such that it decreases with increase in n_k ;
- (ii) Renormalizable: Only a finite number of E –point vertex functions have $D \geq 0$. For these functions though, D is non-negative at all orders;
- (iii) Non-renormalizable: All vertex functions are superficially divergent at sufficiently high orders. The D increases with increase in n_k .

A given theory, may be in any one of the classes in different dimensions.

Dimensional analysis gives another convenient criterion for renormalizability. This goes as follows.

Dimensional Analysis:

In d dimensions $[\Phi] = \frac{d-2}{2}$, $[\Phi^k] = k\frac{d-2}{2}$ $\therefore [g_k] = d - k\frac{d-2}{2} = d(1 - \frac{k}{2}) + k$. An E –point vertex function can come from a term in the Lagrangian as $g_E\Phi^E$ with $[g_E] = d(1 - \frac{E}{2}) + E$. If a diagram contributing to such as vertex function has a momentum cut-off Λ and the number of vertices of order k is n_k , then the divergent part is proportional to $g_k^{n_k}\Lambda^D$. For $k = E$ and $n_E = 1$, there is no loop integration and dimension comes only from the coupling g_E . Hence $[g_E] = [g_k^{n_k}\Lambda^D]$, i.e.

$$d(1 - E/2) + E = n_k \left\{ d - k \frac{d-2}{2} \right\} + D \Rightarrow$$

$$\text{or, } D = d - \frac{d-2}{2}E - \underbrace{\left\{ d - \frac{d-2}{2}k \right\}}_{[g_k]} n_k$$

Thus, D is independent of n_k if $[g_k] = 0$ and we have finitely many vertex functions potentially divergent. If $[g_k] > 0$, then only finitely many diagram can be divergent. If $[g_k] < 0$, then every vertex function has divergence at some order. Thus, the dependence on the space-time dimension, d can be hidden in the dimensions of the couplings and

$g_k \Phi^k$ is renormalizable	if $[g_k] = 0$;
$g_k \Phi^k$ is super-renormalizable	if $[g_k] > 0$;
$g_k \Phi^k$ is non-renormalizable	if $[g_k] < 0$.


Having identified simple criteria for presence of UV divergences, we see now how the counter terms are used for renormalization.

B. An Example: The $(\Phi^3)_6$ theory

As specific example, we will consider the $(\Phi^3)_6$. Here are the Feynman rules for the renormalized perturbation series.

$\frac{p}{\text{---}}$	$\frac{-i}{p^2+m^2-i\epsilon}$,		$+ig$,
$\frac{p}{\text{---} \otimes \text{---}}$	$-i(\delta_\Phi p^2 + \delta_m m^2)$,		$+ig\delta_3$.

$D = 6 - 2E \geq 0 \Rightarrow E = 2$ (quadratically divergent) and $E = 3$ (logarithmically divergent) are the only vertex functions with UV divergence. Consider the 1-loop divergences first.



$$i\Pi(p^2) = \left[\left(\frac{1}{2} \right) (ig)^2 \int \frac{d^d k}{(2\pi)^d} \left(\frac{-i}{(p+k)^2 + m^2 - i\epsilon} \frac{-i}{k^2 + m^2 - i\epsilon} \right) \right] \textcircled{1}$$

$$+ [1 \cdot (-i)(\delta_\Phi p^2 + \delta_m m^2) + \dots] \textcircled{2}$$

The $\frac{1}{2}$ comes from: $\frac{6 \times 2 \times 3}{2!(3!)^2} = \frac{1}{2}$ and

The 1 comes from: $\frac{2 \times 1}{1!2!} = 1$

δ_Φ, δ_m to be determined from $\Pi(-m^2) = 0 = \pi'(-m^2)$.

The first term, after using the Feynman parameters and Wick rotation and equations (17.4, 17.6, 17.7) gives,

$$\begin{aligned}
\textcircled{1} &= \frac{g^2}{2} I(p^2), \quad I(p^2) := i \int_0^1 dx \int \frac{d^d k}{(2\pi)^d} \frac{1}{(\underline{k}^2 + M^2)^2}, \quad \boxed{M^2 := x(1-x)\underline{p}^2 + m^2} \\
I(p^2) &= \frac{\Gamma(-1 + \epsilon/2)}{(4\pi)^3} \int_0^1 dx M^2 (4\pi/M^2)^{\epsilon/2}, \quad \epsilon := 6 - d, \quad \text{put } g \rightarrow g(\bar{\mu})^{\epsilon/2} \alpha := \frac{g^2}{(4\pi)^3} \\
\textcircled{2} &= -i(\delta_\Phi p^2 + \delta_m m^2) \\
\therefore \Pi(p^2) &= -\frac{\alpha}{2} \left[\left(1 + \frac{2}{\epsilon}\right) \left(\frac{p^2}{6} + m^2\right) + \int_0^1 dx M^2 \ln \left(\frac{4\pi \bar{\mu}^2}{e^\gamma M^2} \right) \right] - \delta_\Phi p^2 - \delta_m m^2 + o(\alpha^2).
\end{aligned}$$

Putting $\boxed{\mu := \sqrt{4\pi} e^{-\gamma} \bar{\mu}}$, the logarithm becomes $\ln(\mu^2/M^2)$. We continue the Euclidean momenta back to Minkowski momenta, $\underline{p}^2 \rightarrow p^2$. Club the first group of terms with the counter terms. This gives the self energy as, [12]

$$\begin{aligned}
\Pi(p^2) &= \frac{\alpha}{2} \int_0^1 dx M^2 \ln(M^2/\mu^2) \\
&\quad - \left\{ \frac{\alpha}{6} \left(\frac{1}{\epsilon} + \frac{1}{2} \right) + \delta_\Phi \right\} p^2 - \left\{ \alpha \left(\frac{1}{\epsilon} + \frac{1}{2} \right) + \delta_m \right\} m^2 + o(\alpha^2) \quad (19.1)
\end{aligned}$$

Now, in the first term, use $\ln(M^2/\mu^2) = \ln(M^2/m^2) + \ln(m^2/\mu^2)$ and absorb the contribution of the second term into the counter term coefficients along with the UV divergences (the ϵ^{-1} pole) by *choosing*,

$$\delta_\Phi := -\frac{\alpha}{6} \left(\frac{1}{\epsilon} + \frac{1}{2} + C_\Phi \right) + o(\alpha^2), \quad \delta_m := -\alpha \left(\frac{1}{\epsilon} + \frac{1}{2} + C_m \right) + o(\alpha^2). \quad (19.2)$$

Note that we have only absorbed the pole together with some, μ -dependent *finite parts* (the $\ln(m^2/\mu^2)$ terms) and subsumed these in the undetermined, finite constants C_Φ, C_m . These constants are determined by the renormalization conditions. The self-energy then takes the form,

$$\Pi(p^2) = \frac{\alpha}{2} \int_0^1 dx M^2 \ln(M^2/m^2) + \frac{\alpha}{6} C_\Phi p^2 + \alpha C_m m^2 + o(\alpha^2). \quad (19.3)$$

The C 's are now determined by imposing $\Pi(-m^2) = 0 = \Pi'(-m^2)$. This leads to,

$$C_m - \frac{C_\Phi}{6} = \int_0^1 dx \frac{M_0^2}{m^2} \ln(M_0^2/m^2), \quad \frac{M_0^2}{m^2} = 1 - x + x^2, \quad (19.4)$$

$$-\frac{C_\Phi}{6} = \frac{\alpha}{2} \int_0^1 dx x(1-x) \{ \ln(M_0^2/m^2) + 1 \}. \quad (19.5)$$

Notice that the C 's so determined, are independent of the arbitrary parameter μ that entered in the dimensional regularization scheme. We have expressed the 2-point vertex function

explicitly free of UV divergence *and* the arbitrary μ parameter. In $\Pi(p^2)$ we had two terms with the ϵ^{-1} pole, one with coefficient p^2 and one with coefficient m^2 . The two counter terms δ_Φ, δ_m sufficed to absorb these.

Consider now the 3-point vertex function, $\Gamma(p_1, p_2, p_3)$:

$$i\Gamma_3 : \quad \begin{array}{c} \downarrow p_1 \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ \uparrow p_2 \quad \uparrow p_3 \end{array} := \begin{array}{c} \downarrow p_1 \\ \text{---} \\ \text{---} \\ \uparrow p_2 \quad \uparrow p_3 \end{array} + \begin{array}{c} \downarrow p_1 \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ \uparrow p_2 \quad \uparrow p_3 \end{array} + \begin{array}{c} \downarrow p_1 \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ \uparrow p_2 \quad \uparrow p_3 \end{array} + \dots$$

At 1-loop we have,

$$i\delta\Gamma_3 : \quad \begin{array}{c} \downarrow p_1 \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ \uparrow p_2 \quad \uparrow p_3 \end{array} + \begin{array}{c} \downarrow p_1 \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ \uparrow p_2 \quad \uparrow p_3 \end{array} + \begin{array}{c} \downarrow p_1 \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ \uparrow p_2 \quad \uparrow p_3 \end{array} + \begin{array}{c} \downarrow p_1 \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ \uparrow p_2 \quad \uparrow p_3 \end{array} + \dots$$

$$= [ig\delta_3] \textcircled{1} + (ig)^3 (-i)^3 \times \int \frac{d^d k}{(2\pi)^d} \left[\frac{1}{k^2 + m^2 - i\epsilon} \frac{1}{(p_2 + k)^2 + m^2 - i\epsilon} \frac{1}{(k - p_1)^2 + m^2 - i\epsilon} \right] \textcircled{2}$$

In the second term, Feynman parameterization, shifting momentum and Wick rotation gives,

$$[\dots] = 2 \int dx \int dy \int dz \frac{\delta(1-x-y-z)}{(k^2 + M^2)^3}, \quad \boxed{M^2 := zp_1^2 + zp_2^2 + xyp_3^2 + m^2}.$$

As before replacing $g \rightarrow g\bar{\mu}^{\epsilon/2}$, $\epsilon = 6 - d$ and using the (17.7) gives,

$$\int \frac{d^d k}{(2\pi)^d} \frac{1}{(k^2 + M^2)^3} = \frac{\Gamma(3 - d/2)}{2(4\pi)^{d/2}} M^{-(3-d/2)}.$$

Hence, putting $\alpha := g^2/(4\pi)^3$, $\mu^2 := 4\pi e^{-\gamma} \bar{\mu}^2$

$$\begin{aligned} \frac{\delta\Gamma(p_1, p_2, p_3)}{g} &= \delta_g + \frac{\alpha}{2} \left[\frac{2}{\epsilon} + 2 \int dx dy dz \delta(1-x-y-z) \left(\frac{4\pi\bar{\mu}^2}{M^2} \right)^{\epsilon/2} \right]; \\ &= \left\{ \frac{\alpha}{\epsilon} + \delta_g \right\} \\ &\quad - \alpha \int dx dy dz \delta(1-x-y-z) \ln(M^2/m^2) + o(\alpha^2) \end{aligned} \quad (19.6)$$

$$\text{Choosing } \boxed{\delta_g := -\frac{\alpha}{\epsilon} - \alpha C_g + o(\alpha^2)} \Rightarrow \quad (19.7)$$

$$\Gamma(p_1, p_2, p_3) = g \left[1 - \alpha \int dx dy dz \delta(1-x-y-z) \ln(M^2/m^2) - \alpha C_g + o(\alpha^2) \right] \quad (19.8)$$

As before, we have obtained a completely finite and μ -independent vertex function with one undetermined constant C_g . This is to be fixed by a suitable renormalization condition.

What condition do we choose?

In QED we had a natural choice $\Gamma^\mu(q^2 \rightarrow 0) = e_{ph}$. Here however, there is no natural choice. *Any* cross-section will involve both g and C_g and the value of a cross-section gives one condition. A *convenient* choice is $\Gamma_3(0,0,0) = g \leftrightarrow C_g = 0$.


Thus, at the 1-loop level, we see how the counter terms absorb the divergence and how the renormalization conditions determine the constants C 's. Note that we could have added any finite constant to the pole in defining the counter terms. The renormalization conditions then would give different expressions for the constants.

1. At 2-Loops

A natural question is: How does this work at higher orders? In our $(\Phi^3)_6$ theory, only the 2-point and 3-point vertex functions are divergent. At 2-loops the contributing diagrams are:

		I
2-point		II
		III
3-point		IV

The counter terms have the same form as the original bare terms. When they are clubbed to-

gether into the ‘interaction’ Lagrangian or Hamiltonian, one would expect that from $H_{int}^k/k!$ terms, we would get the counter vertices also appearing k times. For example, at $o(g^2)$, we would have  vertices. In vertex functions, the 1PR diagrams from the self-energy counter terms are omitted. Secondly, the coefficients of the counter terms are explicitly instructed to be adjusted to enforce the renormalization conditions at any given order in g . Hence, the counter vertices are *not* on par with the elementary vertices. The order of perturbation is determined by the number of elementary vertices and not by counter vertices.

In the first of the group of diagrams, we have a subdiagram that is divergent (only $|k_1| \rightarrow \infty$, k_2 remains finite). This divergence is absorbed by the 1-loop counter vertices - the second and the third diagrams of group I. To absorb the divergence when both k_1, k_2 loop momenta become large, we need the *new 2-loop counter term* as shown in III.

The group II exhibit the so called “overlapping divergences”: the vertical line’s momentum diverges when either of the loop momenta diverge. There are two subdiagrams which are divergent and to absorb these, we need the δ_3 1-loop counter term. The group III counter terms are needed to absorb the divergence coming from both momenta becoming large.

At higher loops, the procedure is thus recursive. The counter vertices themselves have an expansion in g^2 with coefficients absorbing divergences from the subdiagrams (lower loop orders). The renormalization conditions determine the highest loop order coefficients.

This explains how the divergences arise and how they are absorbed systematically via the counter terms and the renormalization conditions.

We have taken for illustration the on-shell renormalization condition. This method fails when the physical masses vanish and the procedure needs to be adopted suitably. This impacts what exactly is subtracted - (divergent part) or (divergent part + a finite piece)? This leads to different *schemes of renormalization*. Let us see how the problem arises and how different renormalization conditions can be chosen. We continue with the $(\Phi^3)_6$ theory.

C. Renormalization with massless particles

We have the self energy in equation (19.1). Taking derivative w.r.t. p^2 gives,

$$\Pi'(p^2) = - \left[\delta_\Phi + \frac{\alpha}{6} \left(\frac{1}{\epsilon} + \frac{1}{2} \right) \right] + \frac{\alpha}{2} \int_0^1 dx x(1-x) \{ \ln(M^2/\mu^2) + 1 \} + o(\alpha^2) .$$

As $m \rightarrow 0$, $M^2 \rightarrow x(1-x)p^2$. Then $\Pi(0) = 0$ identically while $\Pi'(0)$ is ill-defined (depends on how $p^2, m^2 \rightarrow 0$ is taken). Thus the previous renormalization conditions *cannot* be used and the counter term coefficients would remain undetermined! We noted while discussing the Kallen-Lehmann representation, that the physical spectrum was assumed to have a mass gap i.e. single particle pole being separated from the multi-particle branch point. Whenever this is violated, the above *on-shell renormalization procedure fails*.

All is not lost though. We can try different renormalization conditions.

There are two commonly used *subtraction schemes* within dimensional regularization - the so called *Minimal Subtraction scheme (MS)* and the *Modified Subtraction Scheme (\overline{MS})*. They are defined as:

$\boxed{MS \leftrightarrow \text{define the counter term constants by absorbing only the } \epsilon \text{ pole(s);}}$ while the \overline{MS} scheme uses $\mu^2 := 4\pi e^{-\gamma} \bar{\mu}^2$ in the MS scheme.

With these definitions, we give the finite self-energy in the three schemes:

$$\begin{aligned} \Pi_{on-shell}(p^2) &= -\frac{\alpha}{12}(p^2 + m^2) + \frac{\alpha}{2} \int_0^1 dx M^2 \ln(M^2/M_0^2) + o(\alpha^2), \\ \text{where, } M^2 &= x(1-x)p^2 + m^2 \text{ and } M_0^2 := M^2(p^2 = -m^2) \end{aligned} \quad (19.9)$$

$$\Pi_{MS}(p^2) = -\frac{\alpha}{12}(p^2 + m^2) + \frac{\alpha}{2} \int_0^1 dx M^2 \ln(M^2/\bar{\mu}^2) + o(\alpha^2) \quad (19.10)$$

$$\Pi_{\overline{MS}}(p^2) = -\frac{\alpha}{12}(p^2 + m^2) + \frac{\alpha}{2} \int_0^1 dx M^2 \ln(M^2/\mu^2) + o(\alpha^2) \quad (19.11)$$

The $\Pi_{on-shell}$ is ill-defined as $m \rightarrow 0$ as noted above. For non-zero m though, it is unambiguous, free of the arbitrary scale μ and m is also the physical mass.

By contrast, $\Pi_{\overline{MS}}$ is well defined as $m \rightarrow 0$, but it depends on the mathematical artifact parameter μ ! Now the m parameter *cannot* be the pole of the propagator and hence is *not* the physical mass. The physical mass is determined from $[p^2 + m^2 - \Pi(p^2)]_{p^2=-m_{ph}^2} = 0 \leftrightarrow m_{ph}^2 = m^2 - \Pi_{\overline{MS}}(-m_{ph}^2)$. Since Π is order α , we can write, to order α , $\boxed{m_{ph}^2 = m^2 - \Pi_{\overline{MS}}(-m^2)}$. Substitution gives,

$$\begin{aligned} m_{ph}^2 &= m^2 + \frac{\alpha}{12}m^2(1-6) + \frac{\alpha}{2} \int_0^1 dx m^2(1-x+x^2) \ln((1-x+x^2)m^2/\mu^2) \\ \text{or } \boxed{m_{ph}^2} &= m^2 \left[1 + \frac{5}{12}\alpha \left\{ \ln \frac{\mu^2}{m^2} + C' \right\} + o(\alpha^2) \right], \quad C' = \frac{34 - 3\pi\sqrt{3}}{15} \approx 1.18 \end{aligned} \quad (19.12)$$

Observe that the physical mass explicitly depends on μ which it should not. Hence $\boxed{\frac{d}{d\mu^2} m_{ph}^2(m^2, \alpha, \mu) = 0}$ must hold. Clearly this would be possible if α and/or m also have μ dependence. The independence of the physical mass then relates the μ dependences of m, α .

The residue, R , at the physical pole is as before, $R^{-1} = \frac{\Delta'_E}{dp^2}|_{p^2=-m_{ph}^2} = 1 - \Pi'_{\overline{MS}}(-m_{ph}^2) = 1 - \Pi'_{\overline{MS}}(-m^2) + o(\alpha^2)$. This evaluates to,

$$R^{-1} = 1 + \frac{\alpha}{12} \{ \ln(\mu^2/m^2) + C'' \} , \quad C'' = \frac{17 - 3\pi\sqrt{3}}{\sqrt{3}} \approx 0.23 . \quad (19.13)$$


The residue gives the wave function renormalization constant $Z_\Phi = R$.

Finally, in the vertex function, we choose $\delta_g = -\alpha/\epsilon$ in the \overline{MS} scheme. This leads to,

$$\Gamma_{3,\overline{MS}}(p_1, p_2, p_3) = g \left[1 - \alpha \int dx dy dz \delta(-x - y - z) \ln(M^2/\mu^2) + o(\alpha^2) \right] , \quad \text{where,} \\ M^2 = xyp_1^2 + yzp_2^2 + zxp_3^2 + m^2 . \quad (19.14)$$

To see a possible μ -dependence of the coupling α , we need to identify a *physical quantity* which has explicit μ dependence. As discussed by Srednicky [12], we consider the two particle going to two particle process which is related to the 4-point vertex function which is UV finite. In the limit $m \rightarrow 0$, this process too suffers from IR divergence which is handled as in QED - carefully considering what is observed and including the soft particles contribution. From [12], we borrow the formulae below.

The amplitude for the process, $p_1 + p_2 \rightarrow p_3 + p_4$ is depicted below to order α .



The contribution of (a) in the high energy limit, is given by,

$$\mathcal{M} = \mathcal{M}_0 \left[1 - \frac{11}{12} \alpha (\ln(s/m^2) + o(m^0)) + o(\alpha^2) \right] \quad \text{with,} \\ \mathcal{M}_0 = -g^2 \left(\frac{1}{s} + \frac{1}{t} + \frac{1}{u} \right) , \quad s := -(p_1 + p_2)^2, \quad t = -(p'_1 - p_1)^2, \quad u := -(p'_2 - p_1)^2 .$$

The $o(m^0)$ term is free of $lm(m \rightarrow 0)$ singularity. The amplitude \mathcal{M} is divergent as $m \rightarrow 0$ (see equation (26.1) of Srednicky [12].)

In the limit $m \rightarrow 0$, the above process is experimentally indistinguishable from the one in which there are on-shell, *collinear* particles associated with any/all external lines¹⁴. This is

¹⁴ For mass less particles, $(\sum_i p_i)^2 = \sum_{i \neq j} p_i \cdot p_j = \sum_{ij} |p_i||p_j|(-1 + \cos(\theta_{ij}))$ which vanishes if all particles are collinear, $\theta_{ij} = 0$.

shown in (b) above, with the dotted lines denoting the collinear particles. We must integrate the squared amplitude over momenta collinear within detector's angular resolution.

For observable cross-section with collinear emission included (from all 4 external lines), we have

$$\begin{aligned}
|\mathcal{M}|_{obs}^2 &= |\mathcal{M}|^2 \left[1 + \frac{4\alpha}{12} \left(\ln \left(\frac{\Delta^2 \vec{k}^2}{m^2} \right) + C \right) + \dots \right], \quad C := 4 - 3\sqrt{3}\pi, \quad \vec{k}^2 = s/4, \\
&= |\mathcal{M}_0|^2 \left[1 - \frac{11}{6}\alpha \ln(s/m^2) \right] \times \left[1 + \frac{\alpha}{3} (\ln(\Delta^2 s/(4m^2)) + C) + \dots \right] \\
&= |\mathcal{M}_0|^2 \left[1 - \alpha \left\{ \frac{3}{2}\ln(s/m^2) + \frac{1}{3}\ln(1/\Delta^2) + o(m^0) \right\} + o(\alpha^2) \right]. \mathbf{26.15} \text{ of [12]}
\end{aligned}$$

The Δ is related to the detector angular resolution. In going to the second equation, we have included collinear emission from all 4 lines. There are no UV divergences here. The above formulae implicitly assume on-shell renormalization scheme where we cannot take the $m \rightarrow 0$ limit.

The same computation of the amplitude can be done in the \overline{MS} scheme. This changes the amplitude \mathcal{M} by changing the $\ln(s/m^2) \rightarrow \ln(s/\mu^2)$. The correction due to inclusion of collinear emission continues to have $\ln(\Delta^2 s/(4m^2))$. Additionally, the residue at the physical pole is not 1 but R and hence the amplitude is multiplied by R^2 . Including these changes, the observed squared amplitude is given by,

$$\begin{aligned}
|\mathcal{M}|_{obs}^2 &= |\mathcal{M}_0|^2 \left[1 - \frac{11}{6}\alpha \ln(s/\mu^2) \right] \left[1 + \frac{\alpha}{3}\ln(\Delta^2 s/m^2) \right] \left[1 - \frac{\alpha}{3}\ln(\mu^2/m^2) \right] \\
&= |\mathcal{M}_0|^2 \left[1 - \alpha \left\{ \frac{11}{6}\ln(s/\mu^2) - \frac{1}{3}\ln(\Delta^2 s/m^2) + \frac{1}{3}\ln(\mu^2/m^2) \right\} + \dots \right] \\
\therefore |\mathcal{M}|_{obs}^2 &= |\mathcal{M}_0|^2 \left[1 - \alpha \left\{ \frac{3}{2}\ln(s/\mu^2) + \frac{1}{3}\ln(1/\Delta^2) + o(m^0) \right\} + o(\alpha^2) \right] \quad (19.15)
\end{aligned}$$

There is no dependence on m now and the limit $m \rightarrow 0$ can be taken. However, there is explicit μ dependence and the observed cross-section *cannot* depend on the arbitrary μ . Hence $\boxed{\frac{d}{d\mu}|\mathcal{M}|_{obs}^2 = 0}$ must hold. Writing

$$\ln|\mathcal{M}|_{obs}^2 := \ln|\mathcal{M}_0|^2 + [\alpha \frac{3}{2}\ln(\mu^2) + \alpha C_2] := C_1 + 2\ln\alpha + 3\alpha\ln(\mu) + 3\alpha C_2$$

where C_1, C_2 are functions of s, t, u but independent of α and μ , the condition of

μ -independence gives,

$$0 = \left(\frac{2}{\alpha} + 3(C_2 + \ln(\mu)) \right) \frac{d\alpha}{d\ln(\mu)} + 3\alpha \Rightarrow \quad (19.16)$$

$$\boxed{\beta(\alpha) := \frac{d\alpha}{d\ln(\mu)}} = -\frac{3}{2}\alpha^2 + o(\alpha^3). \quad \text{We also recall,} \quad (19.17)$$

$$0 = \frac{dm_{ph}^2}{d\ln(\mu)} \Rightarrow \quad (19.18)$$

$$\boxed{\gamma_m(\alpha) := \frac{d\ln(m)}{d\ln(\mu)}} = -\frac{\alpha}{12} + o(\alpha^2) \quad (19.19)$$

The γ_m is called the *anomalous mass dimension* of the mass parameter and $\beta(\alpha)$ is the famous ‘‘beta’’ function of the theory. We have just obtained these functions at 1-loop, for the $(\Phi^3)_6$ theory and these govern how the renormalized parameter must change with the ‘‘renormalization scale’’, μ , in order that the physical quantities are independent of the renormalization scale. If we identify $\mu = s$, then we have the running of the renormalized parameters with the center-of-mass energy scale.

The differential equations can be trivially solved giving the running as, $\mu = \hat{\mu}e^t$,

$$\boxed{\alpha(t) = \frac{\hat{\alpha}}{1 + \frac{3}{2}\hat{\alpha}t}, \quad m(t) = \hat{m} \left(1 + \frac{3}{2}\hat{\alpha}t \right)^{-5/18}} \quad (19.20)$$

As $\mu \sim \sqrt{s} \gg \hat{\mu}$, the coupling *decreases*. Such a theory is said to be *asymptotically free*.

To summarize: We have seen how the introduction of counter terms makes it possible to absorb the UV divergences into unobservable coefficients such as $\delta_\Phi, \delta_m, \delta_g$. This process of absorption or ‘subtraction’ has an inherent ambiguity: exactly what is subtracted. This is implicitly determined by imposition of a set of renormalization conditions (eg the on-shell renormalization) or explicitly (eg the MS/\overline{MS} scheme). When the pole (in p^2) in the self-energy is not a simple pole, which happens with mass less particle ($m_{ph} = 0$), alternative subtraction schemes are needed. Such schemes, typically imply renormalization scale dependence in the renormalized parameters such as masses and couplings. This is governed by the beta function(s) and the anomalous dimension(s). For the sub-class of asymptotically free theories, this helps in improving perturbative predictions at high energies.

A more general discussion of the renormalization group and its application is given in subsection (22 B).

20. PATH INTEGRALS IN QUANTUM MECHANICS

We consider an alternative strategy to view quantum dynamics.

Recall that in a quantum framework, we have a (projective) Hilbert space (or more generally a density operator on a Hilbert space) that encodes the kinematics while a family of unitary operators encodes the dynamics. Observable quantities are provided by operators whose expectation values and uncertainties in a given state (or density operator), provide numbers to be matched against experiments. To track time evolution of observable quantities, the central quantity of interest is the *transition probability amplitude, for a state $|\Psi\rangle$ at time t to make a transition at $t' > t$ to another state $|\Psi'\rangle$* . The state $|\Psi\rangle$ evolves to $\exp(-i(t' - t)H)|\Psi\rangle$ by Schrodinger equation (H is taken time independent for simplicity). Its inner product with $|\Psi'\rangle$ gives the probability amplitude. This is denoted as:

$$\text{Transition probability amplitude} := \langle \Psi' | \left(e^{-i(t'-t)H} |\Psi\rangle \right) = \langle \Psi' | e^{-i(t'-t)H} | \Psi \rangle$$

Assuming the quantum system to be describing a particle with a configuration space $\{\vec{q}\}$, we can use the position representation completeness relation and express the amplitude as,

$$\begin{aligned} \langle \Psi' | e^{-i(t'-t)H} | \Psi \rangle &= \int d\vec{q}' \int d\vec{q} \langle \Psi' | \vec{q}' \rangle \langle \vec{q}' | e^{-i(t'-t)H} | \vec{q} \rangle \langle \vec{q} | \Psi \rangle \\ &:= \int d\vec{q}' \int d\vec{q} \Psi^*(\vec{q}') K(\vec{q}', t'; \vec{q}, t) \Psi(\vec{q}) \end{aligned}$$

The idea is to focus on the kernel K and get a convenient representation for it.

Note: We have already presumed the usual framework of quantum theory. It is possible to develop the ideas ab initio taking the Kernel as a central quantity with a proposed form and arrive at the usual quantum framework. The book of Feynman and Hibbs follows this line. We will first sketch the heuristic approach and then relate it to the usual quantum framework.

A. The Ab Initio Path Integral

Consider a classical system with a configuration space Q which, for convenience of notation we take to be \mathbb{R} . Let q_1, q_2 be two points of it. The particle is supposed to make a transition from q_1 at t_1 to q_2 at t_2 . Such a transition takes place classically as well and we use it in the Lagrangian framework and deduce that the transition takes place along a curve

$q(t)$ which extremises the action. In a departure from the classical dynamics, it is proposed that there is certain *probability for the transition* $(q_1, t_1) \rightarrow (q_2, t_2)$. This is to be computed by squaring the total probability amplitude obtained by summing over the probability amplitudes for every path connecting the (q_1, t_1) and (q_2, t_2) . The amplitude for each path is given as a certain phase. Thus, heuristically, $K(q_2, t_2; q_1, t_1) \sim \sum_{paths} e^{i\varphi[q(t)]}$

Questions: What do we choose for the phase? How do we restrict the paths? How is the ‘sum’ to be performed?

- We expect to recover classical dynamics in the limit $\hbar \rightarrow 0$. So we expect the phase must be such as the amplitude is dominated by a single path corresponding to the classical transition. the obvious choice is $\varphi[q(t)] \sim S[q(t)]/\hbar!$

- First guess about the paths would be smooth paths, but this turns out to be not enough. Since the action involves derivatives of $q(t)$, we expect at least piecewise differentiability. But even this may not suffice - we can define derivatives as differences. It seems, continuity suffices. Without worrying about this too much, let us proceed by *discretizing* the time interval. This will also lead us to answer the third question.

Let $T := t_2 - t_1 := N\epsilon$ where N is a large number eventually to be taken to infinity. For definiteness, let the action be

$$S = \int_0^T dt \left[\frac{m}{2} \dot{q}^2 - V(q) \right] \rightarrow \sum_{k=0}^{N-1} \left\{ \frac{m}{2} \frac{(q_{k+1} - q_k)^2}{\epsilon} - \epsilon V \left(\frac{q_{k+1} + q_k}{2} \right) \right\}, \quad q_k := q(t_k)$$

For uniformity of notation, denote $q_0 =: q_{initial}, q_N =: q_{final}$. We have used the natural discretization of the action. The space of paths is now described by the $N - 1$ q_k 's varying independently over \mathbb{R} . The integration measure may be taken as the product measure $\prod_{k=1}^{N-1} \frac{dq_k}{C(\epsilon)}$, where the constant $C(\epsilon)$ is left free and will be chosen shortly. Since N is arbitrary, we define a *path integral* as:

$$K(q_{in}, q_f; T) := \lim_{\epsilon \rightarrow 0} \frac{1}{C(\epsilon)} \prod_{k=0}^{N-1} \int_{-\infty}^{\infty} \frac{dq_k}{C(\epsilon)} \exp \frac{i}{\hbar} \left\{ \frac{m}{2} \frac{(q_{k+1} - q_k)^2}{\epsilon} - \epsilon V \left(\frac{q_{k+1} + q_k}{2} \right) \right\} \quad (20.1)$$

We can rewrite it by separating the $k = (N - 1)^{th}$ integral as,

$$K(q_{in}, q_f; T) = \int_{-\infty}^{\infty} \frac{dq'}{C(\epsilon)} \left[\exp \frac{i}{\hbar} \left\{ \frac{m}{2} \frac{(q_f - q')^2}{\epsilon} - \epsilon V \left(\frac{q_f + q'}{2} \right) \right\} \cdot K(q_{in}, q'; T - \epsilon) \right]$$

As $\epsilon \rightarrow 0$, the rapid oscillations of the $\frac{1}{\epsilon}$ term imply that dominant contribution to the integral comes from q' very close to q_f . Therefore Taylor expanding the potential and the $K(q_{in}, q'; T - \epsilon)$ about q_f , we get

$$K(q_{in}, q_f; T) \approx \int_{-\infty}^{\infty} \frac{dq'}{C(\epsilon)} e^{\frac{i}{\hbar} \frac{m}{2\epsilon} (q_f - q')^2} \cdot \left\{ 1 + \frac{-i}{\hbar} \epsilon V(q_f) + \left(\frac{-i}{\hbar} \right)^2 \frac{\epsilon^2}{2} \frac{V_f^2}{2} \dots \right\} \\ \times \left\{ 1 + (q' - q_f) \partial_{q'} + \frac{(q' - q_f)^2}{2!} \partial_{q'}^2 + \dots \right\} K(q_{in}, q'; T - \epsilon) \Big|_{q'=q_f}$$

Notice that the first braces are independent of q' ($V(q)$ is assumed to be reasonably well behaved). The second braces is a power series in $(q' - q_f)$.

$$K(q_{in}, q_f; T) \approx e^{\frac{-i}{\hbar} \epsilon V(q_f)} \int_{-\infty}^{\infty} \frac{dx}{C(\epsilon)} e^{\frac{i}{\hbar} \frac{m}{2\epsilon} x^2} \cdot \left\{ 1 + x \partial_{q_f} + \frac{x^2}{2!} \partial_{q_f}^2 + \dots \right\} K(q_{in}, q_f; T - \epsilon)$$

In effect, we have a series of Gaussian integrals of the form: $\int_{-\infty}^{\infty} dx x^m e^{\frac{i}{\hbar} \frac{m}{2\epsilon} x^2}$. The Gaussian integrals are all well known,

$$\int_{-\infty}^{\infty} dx e^{-ax^2} = \sqrt{\frac{\pi}{a}}, \quad \int_{-\infty}^{\infty} dx x^{2k} e^{-ax^2} = \frac{1}{a^{k+1/2}} \Gamma(k + 1/2), \quad \int_{-\infty}^{\infty} dx x^{2k+1} e^{-ax^2} = 0.$$

For us, $a = \frac{im}{2\hbar\epsilon}$. For this to be defined, we have to define the usual Gaussian integral for $a > 0$ and continue analytically in the complex a -plane with real part of a being positive. Hence, we must put $m \rightarrow m + i\epsilon$ to provide the convergence factor. Thus,

$$K(q_{in}, q_f; T) \approx \sqrt{\frac{\pi}{-im/(2\hbar\epsilon)}} \frac{1}{C(\epsilon)} \times \left\{ 1 - i \frac{\epsilon}{\hbar} V(q_f) + \frac{i\epsilon\hbar}{2m} \partial_{q_f}^2 + \dots \right\} K(q_{in}, q_f; T - \epsilon).$$

The terms in the braces are regular as $\epsilon \rightarrow 0$ and hence the right hand side can be arranged to have a regular limit as $\epsilon \rightarrow 0$ by choosing $C(\epsilon) = \sqrt{\frac{2\pi\epsilon\hbar}{-im}}$. Additionally, we also get,

$$K(q_{in}, q_f; T) - K(q_{in}, q_f; T - \epsilon) = \left(\frac{i\epsilon\hbar}{2m} \partial_{q_f}^2 - \frac{i\epsilon}{\hbar} V(q_f) \right) K(q_{in}, q_f; T) \quad (20.2)$$

$$\text{or} \quad i\hbar \partial_T K(q_{in}, q_f; T) = \left(-\frac{\hbar^2}{2m} \partial_{q_f}^2 + V \right) K(q_{in}, q_f; T) \quad (20.3)$$

Thus, the K defined above, *satisfies the time dependent Schrodinger equation!*

What about the initial condition? To see the limit $T \rightarrow 0$, take $N = 1(T = \epsilon)$. Then there is no $\int dq$. Only the $k = 0$ term in the exponent survives and we get,

$$K(q_{in}, q_f; \epsilon \rightarrow 0) = \lim_{\epsilon \rightarrow 0} \frac{1}{C(\epsilon)} e^{\frac{i}{\hbar} \frac{m}{2} (q_f - q_{in})^2 - \epsilon V((q_f + q_{in})/2)}, \quad C(\epsilon) = \sqrt{\frac{2\pi\epsilon\hbar}{-im}}.$$

The right hand side is just the representation of $\delta(q_f - q_{in})$. The usual quantum framework definition of the transition amplitude satisfies the time dependent Schrodinger equation with the same initial condition! So we have a strong hint that the $K(q_{in}, q_f; T)$ defined may indeed be identified with $\langle q_f | e^{-iTH} | q_{in} \rangle$ We can see this directly as well.

B. Derivation From Transition Amplitude

Divide the interval $[0, T]$ in to N intervals of size ϵ each, $N\epsilon = T$. Since the Hamiltonian is assumed to be time independent, we can write,

$$e^{-\frac{i}{\hbar}TH} = e^{-\frac{i}{\hbar}\sum_{k=0}^{N-1} H(t_{k+1}-t_k)} = \prod_{k=0}^{N-1} e^{-\frac{i}{\hbar}(t_{k+1}-t_k)H} \approx \prod_{k=0}^{N-1} \left(1 - \frac{i\epsilon}{\hbar}H + \dots \right)_k .$$

Insert the completeness relation $\mathbb{1} = \int d\vec{q}_k |\vec{q}_k\rangle \langle \vec{q}_k|$ between each factor so that

$$\begin{aligned} \langle \vec{q}_f | e^{-\frac{iT}{\hbar}H} | \vec{q}_{in} \rangle &= \int d\vec{q}_1 \dots d\vec{q}_{N-1} \prod_{k=0}^{N-1} \langle \vec{q}_{k+1} | e^{-\frac{i\epsilon H}{\hbar}} | \vec{q}_k \rangle \quad \vec{q}_0 =: \vec{q}_{in}, \quad \vec{q}_N =: \vec{q}_f \\ e^{-\frac{i\epsilon H}{\hbar}} &\approx 1 - \frac{i\epsilon}{\hbar}H + \dots \quad \text{for small } \epsilon. \end{aligned}$$

To simplify further, we consider types of Hamiltonian operators (we suppress the vector arrows).

(i) $H(\hat{q}, \hat{p}) = f(\hat{p}) + g(\hat{q})$. Then

$$\begin{aligned} \langle q_{k+1} | f(\hat{p}) | q_k \rangle &= \int dp_k \langle q_{k+1} | f(\hat{p}) | p_k \rangle \langle p_k | q_k \rangle \\ &= \int dp_k f(p_k) \langle q_{k+1} | p_k \rangle \langle p_k | q_k \rangle \quad \text{Use: } \langle p_k | q_k \rangle = \frac{e^{-\frac{i}{\hbar}\vec{q}_k \cdot \vec{p}_k}}{\sqrt{2\pi\hbar}} \\ \therefore \langle q_{k+1} | f(\hat{p}) | q_k \rangle &= \int dp_k f(\hat{p}) \frac{e^{\frac{i}{\hbar}\vec{p}_k \cdot (\vec{q}_{k+1} - \vec{q}_k)}}{2\pi\hbar} \quad \text{Likewise,} \\ \langle q_{k+1} | g(\hat{q}) | q_k \rangle &= q(\vec{q}_k) \delta(\vec{q}_{k+1} - \vec{q}_k) = g(\vec{q}_k) \frac{1}{2\pi\hbar} \int dp_k e^{\frac{i}{\hbar}\vec{p}_k \cdot (\vec{q}_{k+1} - \vec{q}_k)} \\ \therefore \langle q_{k+1} | \hat{H}(\hat{q}, \hat{p}) | q_k \rangle &= \int \frac{d\vec{p}_k}{2\pi\hbar} H_{cl} \left(\frac{\vec{q}_{k+1} + \vec{q}_k}{2}, \vec{p}_k \right) e^{\frac{i}{\hbar}\vec{p}_k \cdot (\vec{q}_{k+1} - \vec{q}_k)} \end{aligned}$$

We have used $g(q_k) \rightarrow g(q_{k+1} + q_k)/2$ when multiplied by the $\delta(q_{k+1} - q_k)$.

(ii) \hat{H} contains the \hat{q}, \hat{p} in various ordering such as, $\hat{q}\hat{p}, \hat{p}\hat{q}, \hat{q}^2\hat{p}, \hat{q}\hat{p}\hat{q}, \hat{p}\hat{q}^2, \dots$ etc. As an

example, consider a term $\hat{q}^a \hat{p}_b$ in the quantum Hamiltonian. Then,

$$\begin{aligned}\langle q_{k+1} | \hat{q}^a \hat{p}_b | q_k \rangle &= \int d\vec{p}_k \langle q_{k+1} | \hat{q}^a | p_k \rangle \langle p_k | \hat{p}_b | q_k \rangle = q_{k+1}^a (p_k)_b \int \frac{d\vec{p}_k}{2\pi\hbar} e^{\frac{i}{\hbar} \vec{p}_k \cdot (\vec{q}_{k+1} - \vec{q}_k)} \\ \langle q_{k+1} | \hat{p}_b \hat{q}^a | q_k \rangle &= \int d\vec{p}_k \langle q_{k+1} | \hat{p}_b | p_k \rangle \langle p_k | \hat{q}^a | q_k \rangle = q_k^a (p_k)_b \int \frac{d\vec{p}_k}{2\pi\hbar} e^{\frac{i}{\hbar} \vec{p}_k \cdot (\vec{q}_{k+1} - \vec{q}_k)} \\ \therefore \langle q_{k+1} | \frac{\hat{q}^a \hat{p}_b + \hat{p}_b \hat{q}^a}{2} | q_k \rangle &= \left(\frac{q_{k+1}^a + q_k^a}{2} \right) (p_k)_b \int \frac{d\vec{p}_k}{2\pi\hbar} e^{\frac{i}{\hbar} \vec{p}_k \cdot (\vec{q}_{k+1} - \vec{q}_k)}\end{aligned}$$

Thus for the natural ordering, the classical Hamiltonian has its \vec{q} dependence in the averaged form. Here is another example involving \hat{q}^2, \hat{p}^2 , taken in a particular order, called *Weyl order*: $Weyl(\hat{q}^2, \hat{p}^2) := \frac{1}{4}(\hat{q}^2 \hat{p}^2 + 2\hat{q} \hat{p}^2 \hat{q} + \hat{p}^2 \hat{q}^2)$. It is easily seen that,

$$\begin{aligned}\langle q_{k+1} | Weyl(\hat{q}^2, \hat{p}^2) | q_k \rangle &= \frac{1}{4} \{ q_{k+1}^2 \langle q_{k+1} | p^2 | q_k \rangle + 2q_{k+1} q_k \langle q_{k+1} | p^2 | q_k \rangle + q_k^2 \langle q_{k+1} | p^2 | q_k \rangle \} \\ &= \left(\frac{\vec{q}_{k+1} + \vec{q}_k}{2} \right)^2 \langle q_{k+1} | \hat{p}^2 | q_k \rangle\end{aligned}$$

More generally, for monomials in \hat{q}, \hat{p} , the Weyl order $Weyl(\hat{p}^m, \hat{q}^n)$ equals the fully symmetrized and averaged product. It is a result that

$$\langle q_{k+1} | Weyl(\hat{q}^n, \hat{p}^m) | q_k \rangle = \left(\frac{\vec{q}_{k+1} + \vec{q}_k}{2} \right)^n \langle q_{k+1} | \hat{p}^m | q_k \rangle$$

Hence, modulo Weyl ordering, we do get,

$$\boxed{\langle q_{k+1} | e^{-\frac{i\epsilon}{\hbar} H} | q_k \rangle = \int \frac{dp_k}{2\pi\hbar} e^{-\frac{i}{\hbar} \epsilon H(p_k, \frac{q_{k+1} + q_k}{2})} e^{\frac{i}{\hbar} \vec{p}_k \cdot (\vec{q}_{k+1} - \vec{q}_k)}$$

$$\begin{aligned}\langle q_f | e^{-\frac{iT}{\hbar} H} | q_{in} \rangle &= \int d\vec{q}_1 \dots d\vec{q}_{N-1} \int \frac{d\vec{p}_1}{2\pi\hbar} \dots \frac{d\vec{p}_N}{2\pi\hbar} \exp \frac{i}{\hbar} \left\{ \right. \\ &\quad \left. \epsilon \sum_{k=0}^{N-1} \vec{p}_{k+1} \cdot \frac{(\vec{q}_{k+1} - \vec{q}_k)}{\epsilon} - \epsilon H(p_k, \frac{q_{k+1} + q_k}{2}) \right\} \quad (20.4)\end{aligned}$$

Notice that we have N, p_i integrations, one for each interval and $N - 1, q_i$ integrations, two less than the number of points since q_0, q_N are fixed. The exponent is clearly a discretized form of $\int_0^T dt \vec{p}(t) \cdot \dot{\vec{q}}(t) - H(\vec{p}(t), \vec{q}(t)) = \int_0^T dt L(\vec{q}(t), \dot{\vec{q}}(t))$. The $d\vec{q} \frac{d\vec{p}}{(2\pi\hbar)}$ measure suggests that we have paths in the ‘‘phase space’’. What type of paths?

The mis-match in the number of integration variables makes it harder to view the multiple integrals as a measure on the space of paths in phase space. Introduce an arbitrary momentum \vec{p}_0 so that (\vec{q}_0, \vec{p}_0) denotes a point in the phase space just as $(\vec{q}_k, \vec{p}_k), k = 1, \dots, N$

do. The $\int d\vec{p}_N$ shows that the momentum at the final point is integrated over. Thus, the phase space path may be specified by an initial point (\vec{q}_0, \vec{p}_0) . The arbitrary momentum does not enter anywhere and just serves to anchor the initial point. The final point however is the set of all points (\vec{q}_N, \vec{p}_N) with \vec{p}_N integrated over. Thus, the space of paths is (Γ is a phase space of dimension $2n$):

$$\boxed{\text{Space of paths} =: \mathcal{P}_\Gamma := \left\{ \gamma(t) \in \Gamma / \gamma(0) = (\vec{q}_{in}, \vec{p}_0), \gamma(T) = (\vec{q}_f, \vec{p}), \vec{p} \in \mathbb{R}^n. \right\}}$$

There are arbitrary constants here, the \vec{p}_0 is an arbitrary constant vector. Since the \vec{p}_k, \vec{q}_k are all independent, the paths are continuous but non-differentiable everywhere. Thus we denote:

$$\begin{aligned} \langle q_f | e^{-\frac{iT\hat{H}}{\hbar}} | q_{in} \rangle &:= \lim_{\epsilon \rightarrow 0} \int d\vec{q}_1 \dots d\vec{q}_{N-1} \int \frac{d\vec{p}_1}{2\pi\hbar} \dots \frac{d\vec{p}_N}{2\pi\hbar} \\ &\exp \left[\frac{i}{\hbar} \epsilon \sum_{k=0}^{N-1} \left\{ \vec{p}_{k+1} \cdot \frac{(\vec{q}_{k+1} - \vec{q}_k)}{\epsilon} - H \left(\vec{p}_k, \frac{\vec{q}_{k+1} + \vec{q}_k}{2} \right) \right\} \right] \quad (20.5) \end{aligned}$$

$$= \int_{\mathcal{P}_\Gamma} \mathcal{D}q(t) \mathcal{D}p(t) \exp \left[\frac{i}{\hbar} \int_0^T dt \left(\vec{p}(t) \cdot \dot{\vec{q}}(t) - H(\vec{p}(t), \vec{q}(t)) \right) \right], \quad (20.6)$$

$$\int_{\mathcal{P}_\Gamma} \mathcal{D}q(t) \mathcal{D}p(t) := \lim_{N \rightarrow \infty} \prod_{i=1}^{N-1} \int d\vec{q}_k \prod_{j=1}^N \int \frac{d\vec{p}_j}{2\pi\hbar} \quad (20.7)$$

For the special case of $H(\vec{p}, \vec{q}) = \frac{\vec{p}^2}{2m} + V(\vec{q})$, the momentum dependence is quadratic and the momentum integrals can be done trivially. For instance,

$$\begin{aligned} \int \frac{d\vec{p}_{k+1}}{(2\pi\hbar)^d} e^{\frac{i}{\hbar} \left\{ \vec{p}_{k+1} \cdot (\vec{q}_{k+1} - \vec{q}_k) - \epsilon \left(\frac{\vec{p}_{k+1}^2}{2m} + V \left(\frac{\vec{q}_{k+1} + \vec{q}_k}{2} \right) \right) \right\}} &= \frac{1}{(2\pi\hbar)^d} e^{\frac{i}{\hbar} \frac{m}{2\epsilon} (\vec{q}_{k+1} - \vec{q}_k)^2} \times \\ &\int d^d p_k e^{-\frac{i\epsilon}{2m\hbar} \left(\vec{p} - \frac{m(\vec{q}_{k+1} - \vec{q}_k)}{\epsilon} \right)^2} e^{-\frac{i}{\hbar} \epsilon V} \\ &= \left(\frac{m}{2\pi\hbar(i\epsilon)} \right)^{d/2} e^{\frac{i}{\hbar} \epsilon \left\{ \frac{m}{2} \frac{(\vec{q}_{k+1} - \vec{q}_k)^2}{W p^2} - V \left(\frac{\vec{q}_{k+1} + \vec{q}_k}{2} \right) \right\}} \end{aligned}$$

The square root prefactor is just $C(\epsilon)^{-1}$. Doing all the momentum integrals gives,

$$\langle q_f | e^{-\frac{iT\hat{H}}{\hbar}} | q_{in} \rangle = \lim_{\epsilon \rightarrow 0} \int \frac{d\vec{q}_1}{(2\pi\hbar(i\epsilon/m))^{d/2}} \dots \int \frac{d\vec{q}_{N-1}}{(2\pi\hbar(i\epsilon/m))^{d/2}} e^{\frac{i\epsilon}{\hbar} \left\{ \frac{m}{2} \frac{(\vec{q}_{k+1} - \vec{q}_k)^2}{\epsilon^2} - V \right\}} (20.8)$$

Each $d\vec{q}$ integral gets a factor of $C(\epsilon)^{-d}$ and extra such factor is left over from the extra momentum integration. This was the factor introduced and determined by a regular behavior as $\epsilon \rightarrow 0$.

Thus, the upshot is that we can either take the heuristically motivated definition of K and show that it matched with quantum mechanical definition of transition amplitude *or* directly deduce it from the quantum mechanical definition. The K of the transition amplitude is the central quantity in the path integral approach.

A note: We defined $K(q_{in}, q_f; T)$ and identified it with $\langle q_f | \exp(-i/\hbar)TH | q_{in} \rangle$ and called it the probability amplitude for a particle at q_{in} at t_0 to transit to q_f at time $t_0 + T$. This is sometimes also denoted as $\langle q_f, T | q_{in}, 0 \rangle$ or $\langle q'', t'' | q', t' \rangle$ with $t'' - t' = T$. This notation can be confusing since $|q', t' \rangle$ does *not* denote a solution of the Schrodinger equation. Rather, it denotes the *instantaneous eigenvector of the Heisenberg picture operator* $\hat{Q}(t) := e^{itH/\hbar} Q_{Sch}(0) e^{-itH/\hbar}$. Thus,

$$Q(t)|q, t\rangle = q|q, t\rangle \leftrightarrow Q_{Sch}(e^{-itH/\hbar}|q, t\rangle) = q(e^{-itH/\hbar}|q, t\rangle) \Rightarrow e^{-itH/\hbar}|q, t\rangle = |q\rangle$$

Or, $|q, t\rangle = e^{itH/\hbar}|q\rangle$. If $|q, t\rangle$ were the time evolution of $|q\rangle$, we should have $e^{itH/\hbar}|q\rangle$. Thus we have the consistent notation:

$$\boxed{\langle q_f | e^{-\frac{iTH}{\hbar}} | q_{in} \rangle = \langle q_f | e^{-\frac{i(t''-t')H}{\hbar}} | q_{in} \rangle = \langle q_f | e^{-\frac{it''H}{\hbar}} \cdot e^{\frac{it'H}{\hbar}} | q_{in} \rangle = \langle q_f, t'' | q_{in}, t' \rangle .}$$

Consider now $\langle q_f, T | Q_H(t) | q_{in}, -T \rangle$, for $t \in [-T, T]$.

$$\begin{aligned} \langle q_f, T | Q_H(t) | q_{in}, -T \rangle &= \langle q_f | e^{-iTH/\hbar} e^{itH/\hbar} Q_S e^{-itH/\hbar} e^{-iTH/\hbar} | q_{in} \rangle \\ &= \int dq' dq'' \langle q_f | e^{i(t-T)H/\hbar} | q' \rangle \langle q' | Q_S | q'' \rangle \langle q'' | e^{-i(t+T)H/\hbar} | q_{in} \rangle \\ &= \int dq' dq'' \langle q_f, T | q', t \rangle \langle q' | Q_S | q'' \rangle \langle q'', t | q_{in}, -T \rangle \\ &= \int dq' \langle q_f, T | q', t \rangle q'(t) \langle q', t | q_{in}, -T \rangle \end{aligned}$$

In the last equation, we have used $\langle q' | Q_S | q'' \rangle = q' \delta(q' - q'')$ - the property of the Schrodinger picture operator, carried out the dq'' integration and as a *reminder* inserted the argument t in $q'(t)$. Using the path integral notation, we write the last equation as,

$$\boxed{\langle q_f, T | Q_H(t) | q_{in}, -T \rangle = \int \mathcal{D}q q(t) e^{\frac{i}{\hbar}S[-T, T]} = \int_{\mathbb{R}} dq \int \mathcal{D}q e^{\frac{i}{\hbar}S[t, T]} q(t) \int \mathcal{D}q e^{\frac{i}{\hbar}S[-T, t]} \quad (20.9)}$$

Next, consider the quantity,

$$\begin{aligned} \int \mathcal{D}q q(t_1) q(t_2) e^{\frac{i}{\hbar}S[-T, T]} &:= \int dq_1 \int dq_2 \left[\int \mathcal{D}q e^{\frac{i}{\hbar}S[t_1, T]} \right] q_1(t_1) \times \\ &\quad \left[\int \mathcal{D}q e^{\frac{i}{\hbar}S[t_2, t_1]} \right] q_2(t_2) \left[\int \mathcal{D}q e^{\frac{i}{\hbar}S[-T, t_2]} \right] \quad -T \leq t_2 \leq t_1 \leq T; \end{aligned}$$

For $t_2 \geq t_1$ the factors will switch accordingly. We summarize the formulae as,

$$\langle q_f, t'' | q_{in}, t' \rangle = \int \mathcal{D}q e^{\frac{i}{\hbar} S[t', t'']} =: \langle q_f | e^{-\frac{i}{\hbar} (t'' - t') \hat{H}} | q_{in} \rangle \quad (20.10)$$

$$\begin{aligned} \langle q_f, T | Q_H(t) | q_{in}, -T \rangle &= \int \mathcal{D}q q(t) e^{\frac{i}{\hbar} S[-T, T]} \\ &:= \int dq_t \langle q_f, T | q, t \rangle q_t \langle q, t | q_{in}, -T \rangle \end{aligned} \quad (20.11)$$

$$\begin{aligned} \langle q_f, T | T \{ Q_H(t_1) Q_H(t_2) \} | q_{in}, -T \rangle &= \int \mathcal{D}q q(t_1) q(t_2) e^{\frac{i}{\hbar} S[-T, T]} := \int dq_1 \int dq_2 \\ &\begin{cases} \langle q_f, T | q_1, t_1 \rangle q_1(t_1) \langle q_1, t_1 | q_2, t_2 \rangle q_2(t_2) \langle q_2, t_2 | q_{in}, -T \rangle, \\ \qquad \qquad \qquad t_1 \leq t_2 \\ \langle q_f, T | q_2, t_2 \rangle q_2(t_2) \langle q_2, t_2 | q_1, t_1 \rangle q_1(t_1) \langle q_1, t_1 | q_{in}, -T \rangle, \\ \qquad \qquad \qquad t_2 \leq t_1 \end{cases} \end{aligned} \quad (20.12)$$

C. Functional Derivative

It will be convenient to have the notation of *functional derivative*. Recall from the variational principle of mechanics, the action functional, $S[q(t)] := \int_{t_1}^{t_2} dt L(q(t), \dot{q}(t))$. For any given function $q : [t_1, t_2] \rightarrow q(t) \in \mathbb{R}$, the right hand side computes a number and that number is the action. It is a *function on the space of paths/curves/functions on $[t_1, t_2]$* , and in short is regarded as a *functional of any given $q(t)$* . This is *not* a functional in the sense of being in the dual of vector space - it is not linear in $q(t)$'s. Under a variation of a curve, $q(t) \rightarrow q(t) + \delta q(t)$, we compute $\delta S := S[q + \delta q] - S[q] = \int dt \delta q \left\{ \frac{\partial L}{\partial q} - \frac{d}{dt} \frac{\partial L}{\partial \dot{q}} \right\} + \text{end-point contributions}$. Comparing this with $df(x_1, \dots, x_n) = \sum_{i=1}^n dx^i \partial_i f$, we can identify and denote, *the functional derivative* of the action with respect to the curve as: $\boxed{\frac{\delta S}{\delta q(t)} := \frac{\partial L}{\partial q} - \frac{d}{dt} \frac{\partial L}{\partial \dot{q}}}$. Note that S has no explicit t dependence while its functional derivative does. Thus the idea of a functional derivative is to consider the first order variation of a functional and read off the coefficient of the δ .

We can also notice that any function $f(t)$ we can write, $\delta f(t) = \int dt' \delta(t - t') \delta f(t') \leftrightarrow \frac{\delta f(t)}{\delta f(t')} = \delta(t - t')$. Thus we define the derivative with respect to a function as:

(i)	$\frac{\delta}{\delta f(t)} (\alpha F[f] + \beta G[f]) = \alpha \frac{\delta F}{\delta f(t)} + \beta \frac{\delta G}{\delta f(t)}$	(linearity)
(ii)	$\frac{\delta}{\delta f(t)} F[f] \cdot G[f] = \frac{\delta F}{\delta f(t)} \cdot G[f] + F[f] \cdot \frac{\delta G}{\delta f(t)}$	(Leibnitz rule)
(iii)	$\frac{\delta}{\delta f(t)} F(G[f]) = \frac{\partial F}{\partial G} \frac{\delta G}{\delta f(t)}$	(Chain rule)
(iv)	$\frac{\delta f(t')}{\delta f(t)} := \delta(t' - t)$	

For an arbitrary “source function”, $J(t)$ define the functional,

$$\langle q'', t'' | q', t' \rangle [J] := \int \mathcal{D}q e^{\frac{i}{\hbar} \int_{t'}^{t''} dt \{L(q, \dot{q}) + J(t)q(t)\}} . \quad (20.13)$$

Then it follows that,

$$\begin{aligned} \frac{\delta}{\delta J(t)} \langle q'', t'' | q', t' \rangle [J] &= \int \mathcal{D}q e^{\frac{i}{\hbar} \int_{t'}^{t''} dt \{L(q, \dot{q}) + J(t)q(t)\}} \cdot \frac{i}{\hbar} q(t) \\ \text{Or, } \left(-i\hbar \frac{\delta}{\delta J(t)} \right) \langle q'', t'' | q', t' \rangle [J] &= \int \mathcal{D}q q(t) e^{\frac{i}{\hbar} \int_{t'}^{t''} dt \{L(q, \dot{q}) + J(t)q(t)\}} \\ &= \langle q'', t'' | Q(t) | q', t' \rangle_J . \\ \left(-i\hbar \frac{\delta}{\delta J(t_1)} \right) \left(-i\hbar \frac{\delta}{\delta J(t_2)} \right) \langle q'', t'' | q', t' \rangle [J] &= \int \mathcal{D}q q(t_1) q(t_2) e^{\frac{i}{\hbar} \int_{t'}^{t''} dt \{L(q, \dot{q}) + J(t)q(t)\}} \\ &= \langle q'', t'' | T \{ Q(t_1) Q(t_2) \} | q', t' \rangle_J . \end{aligned}$$

The generalization is obvious. Evaluating the functional derivatives at $J(t) = 0$ gives us,

$$\boxed{\langle q'', t'' | T \{ Q(t_1) \dots Q(t_n) \} | q', t' \rangle = (-i\hbar)^n \frac{\delta^n}{\delta J(t_1) \dots \delta J(t_n)} \langle q'', t'' | q', t' \rangle \Big|_{J=0}} \quad (20.14)$$

Thus, the “correlation functions” (left hand side) are given by the functional derivatives of the transition amplitude in presence of a source function, evaluated at vanishing source function.

To relate it to ground state/vacuum expectation values of time ordered products of Heisenberg picture operators, we take a closer look at the transition amplitude with non-zero source function.

Choose the source function to have a compact support, $J(t) = 0$ for $t < t', t > t''$. Choose $T' < t'$ and $T'' > t''$. Then,

$$\langle Q'', T'' | Q', T' \rangle_J = \int dq'' dq' \langle Q'', T'' | q'', t'' \rangle_{J=0} \langle q'', t'' | q', t' \rangle_J \langle q', t' | Q', T' \rangle_{J=0} .$$

The $J = 0$ amplitudes can be written in the energy representation (energy eigenvalues assumed to be discrete for convenience),

$$\begin{aligned}
\langle q', t' | Q', T' \rangle &= \langle q' | e^{-i(t'-T')H/\hbar} | Q' \rangle = \sum_n \langle q' | \varphi_n \rangle \langle \varphi_n | Q' \rangle e^{-i(t'-T')E_n/\hbar} \\
&= \sum_n \varphi_n(q') \varphi_n^*(Q') e^{-i(t'-T')E_n/\hbar} \\
\therefore e^{-iT'E_0/\hbar} \langle q', t' | Q', T' \rangle &= \sum_n \varphi_n(q') \varphi_n^*(Q') e^{-it'E_n/\hbar} e^{iT'(E_n-E_0)/\hbar} \\
\therefore \lim_{T' \rightarrow i\infty} e^{-iT'E_0/\hbar} \langle q', t' | Q', T' \rangle &= \varphi_0(q') \varphi_0^*(Q') e^{-it'E_0/\hbar} \quad \because E_n \neq E_0 \text{ terms drop out, likewise} \\
\therefore \lim_{T'' \rightarrow -i\infty} e^{iT''E_0/\hbar} \langle Q'', T'' | q'', t'' \rangle &= \varphi_0^*(q'') \varphi_0(Q'') e^{it''E_0/\hbar}
\end{aligned}$$

We have assumed that the lowest energy eigenvalue is non-degenerate, otherwise the \sum_n will reduce to the sum over the degenerate states. Thus we get,

$$\boxed{\lim_{\substack{T' \rightarrow i\infty \\ T'' \rightarrow -i\infty}} \frac{\langle Q'', T'' | Q', T' \rangle_J}{e^{-iE_0(T''-T')} \varphi_0(Q'') \varphi_0^*(Q')} = \int dq' dq'' \varphi_0^*(q'', t'') \varphi_0(q', t') \langle q'', t'' | q', t' \rangle_J} \quad (20.15)$$

D. Ground State-to-ground state Amplitude: $Z[J]$

The right hand side of eq. (20.15) is the ground state-to-ground state transition amplitude, the one that we were looking for. The left hand side tells us how to compute it from $\langle Q'', T'' | Q', T' \rangle$ which is similar to $\langle q'', t'' | q', t' \rangle_J$. The factor in the denominator is independent of J and will not matter. We introduce the definition,

$$\begin{aligned}
Z[J] &:= \int dq' dq'' \varphi_0^*(q'', t'') \langle q'', t'' | q', t' \rangle_J \varphi_0(q', t') \quad (20.16) \\
\frac{\delta^n Z[J]}{\delta J(t_1) \dots \delta J(t_n)} \Big|_{J=0} &= \left(\frac{i}{\hbar} \right)^n \int dq' dq'' \varphi_0^*(q'', t'') \times \\
&\quad \langle q'', t'' | T \{ Q(t_1) \dots Q(t_n) \} | q', t' \rangle \varphi_0(q', t') \quad (20.17)
\end{aligned}$$

The previous result tells us that $Z[J]$ may be computed as,

$$Z[J] \sim \lim_{\substack{T' \rightarrow i\infty \\ T'' \rightarrow -i\infty}} \langle Q'', T'' | Q', T' \rangle_J = \lim_{\substack{T' \rightarrow i\infty \\ T'' \rightarrow -i\infty}} \int \mathcal{D}q \exp \left[\frac{i}{\hbar} \int_{T'}^{T''} dt \{ L(q, \dot{q}) + J(t)q(t) \} \right]$$

We have dropped the unimportant denominator on the left hand side.

We have continued $T' \rightarrow i\infty, T'' \rightarrow -i\infty$, but not the intermediate times, t_i appearing as arguments of the Heisenberg operators. We will do so now and get to the *Euclidean formulation*.

We begin with,

$$\langle Q'', T'' | T \{Q(t_1) \dots Q(t_n)\} | Q', T' \rangle_{J=0} \sim \lim_{\substack{T' \rightarrow i\infty \\ T'' \rightarrow -i\infty}} \int dq_1 \dots dq_n \langle Q'', T'' | q_1, t_1 \rangle q_1 \langle q_1, t_1 | q_2, t_2 \rangle \dots q_n \langle q_n, t_n | Q', T' \rangle, \text{ where,}$$

$$\langle q_i, t_i | q_{i+1}, t_{i+1} \rangle \sim \int \prod_{j=1}^{N-1} \frac{dq_j}{\sqrt{2\pi i \epsilon}} \exp \left[\frac{i}{\hbar} \epsilon \sum_j L \left(\frac{q_i + q_{i+1}}{2}, \frac{q_{i+1} - q_i}{\epsilon} \right) \right], \quad N\epsilon = t_{i+1} - t_i$$

$$\langle q_i, -i\tau_i | q_{i+1}, -i\tau_{i+1} \rangle \rightsquigarrow \int \prod_{j=1}^{N-1} \frac{dq_j}{\sqrt{2\pi \epsilon'}} \exp \left[\frac{\epsilon'}{\hbar} \sum_j L \left(\frac{q_i + q_{i+1}}{2}, \frac{q_{i+1} - q_i}{-\epsilon'} \right) \right], \quad N\epsilon' := \tau_{i+1} - \tau_i$$

The analytic continuation of the $\langle T \{ \dots \} \rangle$ is defined through the $\langle .. | .. \rangle$. Thus,

$$\langle Q'', T'' | T \{Q(t_1) \dots Q(t_n)\} | Q', T' \rangle_{J=0} \Big|_{t_i = -i\tau_i} \sim \lim_{\substack{\tau_{i_n} \rightarrow -\infty \\ \tau_{i_f} \rightarrow \infty}} \int \mathcal{D}q \, q(\tau_1) \dots q(\tau_N) \times \exp \left[\int_{\tau_{i_n}}^{\tau_{i_f}} d\tau L(q, -\frac{dq}{d\tau}) \right]$$

This suggests going over to a *Euclidean formulation*,

$$Z_E[J] := \int \mathcal{D}q \exp \left[\int_{-\infty}^{\infty} d\tau \left\{ L \left(q, i \frac{dq}{d\tau} \right) + J(\tau) q(\tau) \right\} \right] \quad (20.18)$$

and the paths are between some $Q' = \lim_{\tau \rightarrow -\infty} q(\tau)$ and $Q'' = \lim_{\tau \rightarrow \infty} q(\tau)$. The Euclidean Z_E and the Minkowskian Z are related through,

$$\boxed{\frac{1}{Z[J]} \frac{\delta^n Z[J]}{\delta J(t_1) \dots \delta J(t_n)} \Big|_{J=0} = i^n \frac{1}{Z_E[J]} \frac{\delta^n Z_E[J]}{\delta J(\tau_1) \dots \delta J(\tau_n)} \Big|_{J=0}}$$

This will be used in the field theory Green's functions.

E. Explicit evaluation of a path integral

We would like to see the various expressions above explicitly for a 1-dimensional harmonic oscillator with a source function. We have $L = \frac{1}{2}(\dot{q}^2 - \omega^2 q^2) + J(t)q(t)$ and we want to compute $\langle q'', t'' | q', t' \rangle_J$. Discretizing the corresponding action, the definition gives,

$$\langle q'', t'' | q', t' \rangle_J = \lim_{\epsilon \rightarrow 0} \frac{1}{C(\epsilon)} \int_{-\infty}^{\infty} \prod_{k=1}^{N-1} \frac{dq_k}{C(\epsilon)} \exp \left[\frac{i}{\hbar} \left\{ \sum_{k=0}^{N-1} \frac{1}{2} \frac{(q_{k+1} - q_k)^2}{\epsilon} - \frac{\epsilon \omega^2}{2} \left(\frac{q_k + q_{k+1}}{2} \right)^2 + \epsilon J_k \left(\frac{q_k + q_{k+1}}{2} \right) \right\} \right], \quad q_0 := q(t'), q_N := q(t''), J_k := J(t_k)$$

All are Gaussian integrals with coupled variables. It is more convenient to discretize a *different* action.

The equations of motion are: $\ddot{q} + \omega^2 q^2 = J(t)$, $q(t') = q'$, $q(t'') = q''$. Let $q_{cl}(t)$ be a classical solution of this equation satisfying the end point conditions. Let us assume that there is just *one* such solution. Introduce $\eta(t)$ via the definition: $q(t) := q_{cl}(t) + \eta(t)$. Then $\eta(t') = 0 = \eta(t'')$. The action becomes,

$$\begin{aligned} S(t', t'') &= \int_{t'}^{t''} dt \left\{ \frac{1}{2}(\dot{q}_{cl} + \dot{\eta})^2 - \frac{\omega^2}{2}(q_{cl} + \eta)^2 + J(t)(q_{cl} + \eta) \right\} \\ &= \int_{t'}^{t''} dt \left[\left\{ \frac{1}{2}(\dot{q}_{cl}^2 - \omega^2 q_{cl}^2) + J(t)q_{cl} \right\} + \left\{ \frac{1}{2}(\dot{\eta}^2 - \omega^2 \eta^2) \right\} + \{ \dot{\eta}\dot{q}_{cl} - \omega^2 \eta q_{cl} + J\eta \} \right] \end{aligned}$$

Up to a total derivative, the last term is $-\eta(\ddot{q}_{cl} + \omega^2 q_{cl} - J) = 0$. The total derivative term gives $\eta\dot{q}_{cl}|_{t'}^{t''} = 0$ thanks to the end point condition satisfied by $\eta(t)$. Thus,

$S(t', t'') = S_{cl}(t', t'') + \int_{t'}^{t''} \frac{1}{2}(\dot{\eta}^2 - \omega^2 \eta^2)$, where S_{cl} is the action evaluated at the presumed solution q_{cl} . The terms linear in η have vanished thanks to q_{cl} being a solution. We now “quantize” the η variable, i.e. we *define*

$$\langle q'', t'' | q', t' \rangle_J := \left[e^{\frac{i}{\hbar} S_{cl}^J(t', t'')} \right] \left[\lim_{\epsilon \rightarrow 0} \frac{1}{C(\epsilon)} \int_{-\infty}^{\infty} \prod_{k=1}^{N-1} \frac{d\eta_k}{C(\epsilon)} \exp \left\{ \frac{i}{\hbar} \left(\sum_{k=0}^{N-1} \frac{1}{2} \frac{(\eta_{k+1} - \eta_k)^2}{\epsilon} - \frac{\epsilon \omega^2}{2} ((\eta_k + \eta_{k+1})/2)^2 \right) \right\} \right] \quad (20.19)$$

The paths $\eta(t)$ begin and end at $\eta = 0$. The second factor is independent of both J and t', t'' . It depends on $T = t'' - t'$. Our task is to evaluate the first factor and carry out the coupled Gaussian integral.

The exponent in the second factor is of the form,

$$\begin{aligned} \frac{i}{2\hbar}(\dots) &= \sum_{k=0}^{N-1} \frac{1}{\epsilon} (\eta_{k+1} - \eta_k)^2 - \epsilon \omega^2 (\eta_{k+1} + \eta_k)/2)^2 := \sum_{k,l=0}^{N-1} \eta_k A_{kl} \eta_l \quad , \quad \text{where,} \\ A_{lk} &:= \underbrace{2(1/\epsilon - \epsilon \omega/4)}_a \delta_{lk} - \underbrace{(1/\epsilon + \epsilon \omega^2/4)^2}_b \{\delta_{l,k+1} + \delta_{l,k-1}\} \quad , \quad l, k = 1, \dots, N-1. \\ &= \begin{pmatrix} a & -b & 0 & \dots & 0 \\ -b & a & -b & \dots & 0 \\ 0 & -b & a & -b & \dots & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & \dots & -b & a \end{pmatrix} \quad \text{A tri-diagonal, symmetric matrix.} \end{aligned}$$

$$\therefore \langle q'', t'' | q', t' \rangle_J := \left[e^{\frac{i}{\hbar} S_{cl}^J(t', t'')} \right] \left[\lim_{\epsilon \rightarrow 0} \frac{1}{C(\epsilon)} \int_{-\infty}^{\infty} \frac{d^{N-1} \eta}{C(\epsilon)^{N-1}} \exp \left\{ \frac{i}{2\hbar} \sum_{i,j=1}^{N-1} \eta_i A_{ij} \eta_j \right\} \right]$$

For a single variable we have $\int_{-\infty}^{\infty} dx e^{-ax^2} = \sqrt{\pi/a} \Rightarrow \int_{-\infty}^{\infty} dx e^{iax^2} = \sqrt{\pi/(-ia)}$. Its multi-dimensional generalization is

$$\int_{-\infty}^{\infty} d^n x e^{i \sum_{ij} x_i A_{ij} x_j} = \frac{(i\pi)^{n/2}}{\sqrt{\det A}}$$

Using this, our transition amplitude takes the form,

$$\langle q'', t'' | q', t' \rangle_J = \left[e^{\frac{i}{\hbar} S_{cl}^J(t', t'')} \right] \left[\lim_{\epsilon \rightarrow 0} \frac{1}{C(\epsilon)} \frac{1}{(\epsilon)^{N-1}} \frac{(i\pi\hbar)^{(N-1)/2}}{\sqrt{\det A}} \right]$$

Using $C(\epsilon) = \sqrt{2i\pi\epsilon\hbar}$ since we have taken unit mass, $m = 1$, we have,

$$\boxed{\langle q'', t'' | q', t' \rangle_J = \left[e^{\frac{i}{\hbar} S_{cl}^J(t', t'')} \right] \left[\lim_{\epsilon \rightarrow 0} \frac{1}{\sqrt{2\pi i \hbar} (\epsilon)^{N/2}} \frac{1}{\sqrt{\det A(\epsilon)}} \right]} \quad (20.20)$$

Now we need to compute the determinant of the tridiagonal, symmetric matrix. This is usually solved by using a recursion relation. Denote $D_n := \det A_{n \times n}$ where A has the form given above. Clearly, $D_1 = a$ and $D_2 = a^2 - b^2$. By checking for $4 \times 4, 5 \times 5$ matrices, it is easy to see that D_n satisfies the recursion relation,

$$\boxed{D_n = aD_{n-1} - b^2 D_{n-2} \quad , \quad D_0 := 1, D_1 = a \quad ; \quad a = 2 \left(\frac{1}{\epsilon} - \frac{\epsilon \omega^2}{4} \right) \quad , \quad b = \frac{1}{\epsilon} + \frac{\epsilon \omega^2}{4} .}$$

We can pull out a factor of ϵ^{-1} from A and since our A is of order $(N-1)$, we pull out a factor of $(\epsilon)^{-(N-1)/2}$. This replaces the second [...] by $\lim_{\epsilon \rightarrow 0} (2\pi\hbar\sqrt{\epsilon \det A'})^{-1}$. The A' matrix has elements $a' := 2 - \epsilon^2 \omega^2/2$, $b' := 1 + \epsilon^2 \omega^2/4$.

Note: For a given $N, \epsilon = T/N$, hence (suppressing the primes) a, b have an N dependence. The matrix itself is also $(N-1) \times (N-1)$ and our notation D_n as the determinant of $A_{n \times n}$ is valid for $n \leq N-1$. For a given N , the $D_{n \geq N}$ is *not* defined. Hence, the recursion relation is a *difference equation with constant coefficients* [15]. It is important to keep the distinction between fixed N and a variable n . The a, b are functions of N but independent of n .

Such difference equations are solved by the ansatz, $D_n = \lambda^n$. Shifting $n \rightarrow n+2$, we write the difference equation as,

$$\forall 0 \leq n \leq N-3 : D_{n+2} - aD_{n+1} + b^2D_n = 0 \quad , \quad D_0 = 1, D_1 = a.$$

Substitution gives the characteristic equation $\lambda^2 - a\lambda + b^2 = 0$. Its solutions, for $\epsilon \ll 1$, are

$$\begin{aligned} \lambda_{\pm} &\approx \left(1 - \frac{\epsilon^2 \omega^2}{4}\right) \pm \sqrt{\left(1 - \frac{\epsilon^2 \omega^2}{2}\right) - \left(1 + \frac{\epsilon^2 \omega^2}{2}\right)} \approx 1 - \frac{\epsilon^2 \omega^2}{4} \pm i\epsilon \omega \\ \therefore \lambda_{\pm} &\simeq 1 \pm i\epsilon \omega \quad . \Rightarrow D_n = \alpha \lambda_+^n + \beta \lambda_-^n \quad \forall n \in [0, N-1]. \\ \text{Initial conditions give, } &\alpha = \frac{1}{2} \left(1 - \frac{i}{\epsilon \omega}\right) \quad , \quad \beta = \frac{1}{2} \left(1 + \frac{i}{\epsilon \omega}\right) . \end{aligned}$$

The desired determinant is then given by,

$$\begin{aligned} D_{N-1} &= \frac{1}{2} \left(1 - \frac{i}{\epsilon \omega}\right) (1 + i\epsilon \omega)^{N-1} + \frac{1}{2} \left(1 + \frac{i}{\epsilon \omega}\right) (1 - i\epsilon \omega)^{N-1} \quad , \quad \epsilon = T/N \\ &\xrightarrow{N \rightarrow \infty} \frac{1}{2} i \left(1 - \frac{iN}{\omega T}\right) (1 + i\omega T/N)^N + \frac{1}{2} \left(1 + \frac{iN}{\omega T}\right) (1 - i\omega T/N)^N \\ &\rightarrow \left(-\frac{iN}{2\omega T}\right) (e^{i\omega T} - e^{-i\omega T}) = \frac{N}{\omega T} \sin(\omega T) \\ \therefore \epsilon D_N &\rightarrow \frac{\sin(\omega T)}{\omega} . \end{aligned}$$

We finally get,

$$\langle q'', t'' | q', t' \rangle_J = \left[e^{\frac{i}{\hbar} S_{cl}^J(t', t'')} \right] \left[\frac{1}{\sqrt{2\pi i \hbar}} \sqrt{\frac{\omega}{\sin(\omega T)}} \right] . \quad (20.21)$$

Note: All the J dependence is in the first factor only while the “quantum correction” are in the second factor and independent of J .

Note: This method of evaluating the amplitude near a classical solution can also be adopted for more general (non-linear) equations of motion. The S_{cl} always comes out, the term linear in η always vanishes while the $o(\eta^2)$ terms always gives the (determinant) $^{-1/2}$. In the general context, it is termed as a semi-classical approximation which is exact for the oscillator.

The calculation of the first factor requires solving the equation of motion with the source function and then evaluating the action for $q_{cl}(t)$. This is little involved as the equation is inhomogeneous and requires use of Green's function. The Green's function can be obtained by directly solving the differential equation with δ - function source and matching the discontinuity due to the delta function. We just note the final result [Problem 3.11 of Feynman-Hibbs, Abers-Lee].

$$\begin{aligned}
S_{cl}^J(t', t'') &= \frac{\omega}{2\sin(\omega T)} [((q')^2 + (q'')^2)\cos(\omega T) - 2q'q''] \\
&+ \frac{q''}{\sin(\omega T)} \int_{t'}^{t''} dt J(t) \sin(\omega(t - t')) + \frac{q'}{\sin(\omega T)} \int_{t'}^{t''} dt J(t) \sin(\omega(t'' - t)) \\
&- \frac{1}{\omega \sin(\omega T)} \int_{t'}^{t''} d\sigma \int_{t'}^{\sigma} d\tau J(\sigma) J(\tau) \sin(\omega(t'' - \sigma)) \sin(\omega(\tau - t')) \quad (20.22)
\end{aligned}$$

This exercise was done to illustrate the schematics of evaluating the path integral directly using the definition.

Note: If we have a system of two degrees of freedom, the $q(t)$ and the $J(t)$ which are coupled by a linear coupling, $\int_{t'}^{t''} dt J(t) q(t)$, then the $S_{cl}^J(t', t'')$ can be viewed as an effective action contribution after integrating out the $q(t)$ degree of freedom. This has the form $\sim \int dt J(t) \alpha(t) - \int dt J(t) \int^t d\sigma \beta(t, \sigma) J(\sigma)$. This last term is a *non-local term*.

F. Alternative Expression for $Z[J]$

We now consider another method which is closer to what is done in field theory. We begin by modifying the Hamiltonian operator as: $\hat{H} \rightarrow (1 - i\epsilon)\hat{H}$ [12]. Then,

$$|q', t'\rangle = e^{\frac{i}{\hbar} t' \hat{H}} |q'\rangle = \sum_n \varphi_n^*(q') e^{\frac{i}{\hbar} t' E_n} |n\rangle \rightarrow \sum_n \varphi_n^*(q') e^{\frac{i}{\hbar} t' (1-i\epsilon) E_n} |n\rangle$$

Assuming $E_0 = 0$ for convenience and taking the limits, we get

$$\boxed{|q', t'\rangle \xrightarrow{t' \rightarrow -\infty} \varphi_0^*(q') |0\rangle \quad \text{and} \quad \langle q'', t''| \xrightarrow{t'' \rightarrow \infty} \langle 0| \varphi_0(q'')}. \quad (20.23)}$$

Thus, $|q', t'\rangle, \langle q'', t''|$ both go to the (presumed non-degenerate) ground state as $t' \rightarrow -\infty$ and $t'' \rightarrow \infty$, *provided* we make the substitution: $\hat{H} \rightarrow (1 - i\epsilon)\hat{H}$. We will work with the limits.

Consider, for our 1-dimensional oscillator, $H = p^2/2 + \omega^2 q^2/2$,

$$\langle 0|0\rangle_J = \int \mathcal{D}p \mathcal{D}q \exp \left[\frac{i}{\hbar} \int_{-\infty}^{\infty} dt \{p\dot{q} - (1 - i\epsilon)H + J(t)q\} \right]$$

We do momentum integration by using $(1 - i\epsilon)p^2/2 \approx \frac{p^2}{2(1+i\epsilon)}$. This gives the term $\frac{1}{2}(1+i\epsilon)\dot{q}^2$ and we get,

$$\langle 0|0\rangle_J = \int \mathcal{D}q \exp \left[\frac{i}{\hbar} \int_{-\infty}^{\infty} dt \left\{ \frac{1}{2}(1+i\epsilon)\dot{q}^2 - \frac{1}{2}\omega^2(1-i\epsilon)q^2 + J(t)q \right\} \right]. \quad (20.24)$$

Define the Fourier transform,

$$\tilde{q}(\nu) := \int_{-\infty}^{\infty} dt e^{i\nu t} q(t) \leftrightarrow q(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} d\nu e^{-i\nu t} \tilde{q}(\nu)$$

and similarly for $J(t)$. Substitution give,

$$L = \frac{1}{2} \int_{-\infty}^{\infty} \frac{d\nu}{2\pi} \int_{-\infty}^{\infty} \frac{d\nu'}{2\pi} e^{-i(\nu+\nu')t} \left[\left\{ -(1+i\epsilon)\nu\nu' - (1-i\epsilon)\omega^2 \right\} \tilde{q}(\nu)\tilde{q}(\nu') \right. \\ \left. + \frac{\tilde{J}(\nu)\tilde{q}(\nu') + \tilde{J}(\nu')\tilde{q}(\nu)}{2} \right]$$

The expression in the braces simplifies to $\{\dots\} = \nu^2 - \omega^2 + i\epsilon(\nu)$, $\epsilon(\nu) := \epsilon(\nu^2 + \omega^2)$. Define $\tilde{x}(\nu) := \tilde{q}(\nu) + \frac{\tilde{J}(\nu)}{\nu^2 - \omega^2 + i\epsilon(\nu)}$. The the action becomes,

$$S = \frac{1}{2} \int_{-\infty}^{\infty} \frac{d\nu}{2\pi} \left[\tilde{x}(\nu)(\nu^2 - \omega^2 + i\epsilon(\nu))\tilde{x}(-\nu) - \frac{\tilde{J}(\nu)\tilde{J}(-\nu)}{\nu^2 - \omega^2 + i\epsilon(\nu)} \right]$$

The shift, $\tilde{q}(\nu) \rightarrow \tilde{x}(\nu)$ is a *constant shift*, the shift is independent of \tilde{q} . Hence the Jacobian of the transformation will be 1 and $\mathcal{D}\tilde{q}(\nu) = \mathcal{D}\tilde{x}(\nu)$. Taking Fourier transform, the shift takes the form $q(t) \rightarrow x(t) + f(t)$ with $f(t)$ independent of $q(t)$. This is also a constant shift and in the time domain too, and we expect $\mathcal{D}q(t) = \mathcal{D}x(t)$.

$$\therefore \langle 0|0\rangle_J = \exp \left[\frac{i}{2\hbar} \int_{-\infty}^{\infty} \frac{d\nu}{2\pi} \left(\frac{\tilde{J}(\nu)\tilde{J}(-\nu)}{-\nu^2 + \omega^2 - i\epsilon(\nu)} \right) \right] \times \quad (20.25)$$

$$\left[\int \mathcal{D}\tilde{x}(\nu) \exp \frac{i}{2\hbar} \int_{-\infty}^{\infty} \frac{d\nu}{2\pi} \tilde{x}(\nu)(\nu^2 - \omega^2 + i\epsilon(\nu))\tilde{x}(-\nu) \right] \quad (20.26)$$

The second factor is a path integral independent of J and hence equals $\langle 0|0\rangle_{J=0}$. However, without any source interaction, the ground state remains a ground state and hence

$\langle 0|0\rangle_{J=0} = 1!$. This gives finally,

$$\langle 0|0\rangle_J = \exp \frac{i}{2\hbar} \int_{-\infty}^{\infty} \frac{d\nu}{2\pi} \left(\frac{\tilde{J}(\nu)\tilde{J}(-\nu)}{-\nu^2 + \omega^2 - i\epsilon(\nu)} \right) \quad (20.27)$$

We never needed to evaluate the path integral!

The above expression can also be expressed in the time domain as,

$$\langle 0|0\rangle_J = \exp\frac{i}{2\hbar} \int_{-\infty}^{\infty} dt' dt'' J(t'') G(t'' - t') J(t') \quad \text{with,} \quad (20.28)$$

$$G(t'' - t') = \int_{-\infty}^{\infty} \frac{d\nu}{2\pi} \frac{e^{-i\nu(t''-t')}}{-\nu^2 + \omega^2 - i\epsilon(\nu)} = \frac{i}{2\omega} e^{-i\omega|t''-t'|} \text{(by contour integration)} \quad (20.29)$$

Note: The ν dependence in the $\epsilon(\nu)$ does not affect the contour integration.

We have discussed two different methods of computing the transition amplitude in the limit of infinite time separation. Notice that the $\langle 0|0\rangle_J$ above is the same as the $Z[J]$ defined earlier in eq. (20.16) since $\int dq' |q', t'\rangle \varphi_0(q', t') = |0, t'\rangle$ and likewise for $\langle 0, t''|$.

The above frequency domain form is very convenient to obtain the correlation functions as seen below.

Our general formula (20.14) tells us that $\langle 0|T\{Q(t_1) \dots Q(t_n)\}|0\rangle$ is given by the functional derivatives of $Z[J]$ evaluated at $J = 0$. Choosing the above form of $Z[J]$ we see that,

$$\begin{aligned} \langle 0|T\{Q(t_1)Q(t_2)\}|0\rangle &= (-i\hbar)^2 \frac{\delta^2 \langle 0|0\rangle_J}{\delta J(t_1) \delta J(t_2)} \Big|_{J=0} & (20.30) \\ &= (-i\hbar)^2 \frac{\delta}{\delta J(t_1)} \left\{ \frac{i}{2\hbar} \int_{-\infty}^{\infty} dt' G(t_2 - t_1) J(t') \times \right. \\ &\quad \left. \exp\left(\frac{i}{2\hbar} \int \int JGJ\right) \right\} \Big|_{J=0} \\ &= (-i\hbar)^2 (i/\hbar) G(t_2 - t_1) = -i\hbar G(t_2 - t_1). & (20.31) \end{aligned}$$

The derivative of the $e^{\int JGJ}$ does not contribute in the limit of $J = 0$. Taking one more derivative to get the three point function, we have

$$\begin{aligned} \frac{\delta}{\delta J_1} \frac{\delta}{\delta J_2} \left\{ \frac{i}{\hbar} \int dt G(t_3 - t) J(t) \cdot \langle 0|0\rangle_J \right\} &= \\ \frac{\delta}{\delta J_1} \left\{ \frac{i}{\hbar} G(t_3 - t_2) \cdot \langle 0|0\rangle_J + \frac{i}{\hbar} \int dt G(t_3 - t) J(t) \cdot \frac{i}{\hbar} \int dt' G(t_2 - t') J(t') \cdot \langle 0|0\rangle_J \right\} \\ &= 0 + 0 = 0 \quad \because \text{there is always a factor of } J \text{ which kills the term at } J = 0. \end{aligned}$$

For 4 derivatives, we will have,

$$\begin{aligned} \frac{\delta}{\delta J_0} & \left\{ \frac{i}{\hbar} G(t_3 - t_2) \cdot \frac{i}{\hbar} \int dt G(t_1 - t) J(t) \cdot \langle 0|0 \rangle_{J+} \right. \\ & \frac{i}{\hbar} G(t_3 - t_1) \cdot \frac{i}{\hbar} \int dt G(t_2 - t) J(t) \cdot \langle 0|0 \rangle_{J+} \\ & \frac{i}{\hbar} \int dt G(t_3 - t) J(t) \cdot \frac{i}{\hbar} G(t_2 - t_1) \cdot \langle 0|0 \rangle_{J+} \\ & \left. \left(\frac{i}{\hbar} \right)^3 \int dt G(t_3 - t) J(t) \cdot \int dt' G(t_2 - t') J(t') \cdot \int dt'' G(t_1 - t'') J(t'') \cdot \langle 0|0 \rangle_J \right\} \end{aligned}$$

Carrying out the J_0 differentiation and putting $J = 0$, only the first 3 terms contribute since they have a single factor of $J(t)$. For a 4-point function, this is multiplied by $(-i\hbar)^4$ and we get,

$$\boxed{\langle 0|T\{Q(t_0)Q(t_1)Q(t_2)Q(t_3)\}|0\rangle = (-i\hbar)^2 \left\{ G_{01}G_{23} + G_{02}G_{31} + G_{03}G_{12} \right\}, \quad G_{ij} \leftrightarrow G(t_i - t_j).} \quad (20.32)$$

We recognize this as the same pattern seen in the vacuum expectation value calculation using the Wick's theorem. Indeed, the quantum field theory Wick's theorem shows up here just as a result of the functional differentiation.

We now have all the definitions, notations and pattern of computation needed in the field theory generalization.

21. PATH INTEGRALS IN QUANTUM FIELD THEORY

We consider the path integral formulation of a field theory, specifically a scalar field theory. While discussing the field as a dynamical system, we had already noted that the notation $\phi(t, \vec{x})$ refers to infinitely many degrees of freedom labeled by $\vec{x} \in \mathbb{R}^3$ (for us). Thus we could view $\phi(t, \vec{x})$ as $\phi_{\vec{x}}(t)$ and draw analogy with $q_i(t)$. Each of these degrees of freedom can be quantized a la path integral exactly as the single degree of freedom discussed before. In place of $\int \mathcal{D}q(t) \xrightarrow{T=\epsilon N} \int \prod_{k=0}^{N-1} \frac{dq_k}{C(\epsilon)}$, we now have $\int \mathcal{D}\phi(x) \xrightarrow{T=\epsilon N} \int \prod_{\vec{x} \in \Sigma} \prod_{k=0}^{N-1} d\phi_{\vec{x},k}$. We have dropped the $C(\epsilon)$ which will be subsumed in the overall normalization constant. The action for the field theory is as the usual one. Thus, the vacuum-to-vacuum transition amplitude in presence of a source is denoted as,

$$\langle 0|0 \rangle_J := \int \mathcal{D}\phi \exp \frac{i}{\hbar} \int d^4x \left\{ \mathcal{L}(\phi, \partial_\mu \phi) + J(x)\phi(x) \right\} =: Z[J] \quad (21.1)$$

The ‘paths’ implicit in the measure $\mathcal{D}\phi$ are of course paths in the configuration space of the field i.e. the space of all $\phi(\vec{x})$ at any fixed t . A path itself connects $\phi_1(\vec{x}, t_1)$ to $\phi_2(\vec{x}, t_2)$, $\forall \vec{x}$. The paths are *not* paths in the space-time.

For the Lagrangian density \mathcal{L} , we write $\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_1$, with $\mathcal{L}_0 := -\frac{1}{2}\partial^\mu \phi \partial_\mu \phi - \frac{1}{2}m^2 \phi^2$. The m^2 has $-i\epsilon$ implicit, in anticipation. The \mathcal{L}_1 will typically involve interaction terms, polynomials in ϕ such as $g_3\phi^2(x) + g_4\phi^4(x) \dots$ etc. Consider

$$Z_0[J] := \int \mathcal{D}\phi \exp \left\{ \frac{i}{\hbar} \int d^4x (\mathcal{L}_0 + J(x)\phi(x)) \right\}.$$

We go to the momentum space as in the case of the single particle.

Let $\tilde{\phi}(k) := \int d^4x e^{-ik \cdot x} \phi(x) \leftrightarrow \phi(x) = \int \frac{d^4k}{(2\pi)^4} e^{ik \cdot x} \tilde{\phi}(k)$, with $k \cdot x := -k^0 x^0 + \vec{k} \cdot \vec{x}$ and k^0 is the integration variable in d^4k and not k_0 . Substituting in the action and doing the d^4x integration gives,

$$S_0 = \frac{1}{2} \int \frac{d^4k}{(2\pi)^4} \left[-\tilde{\phi}(k)(k^2 + m^2)\tilde{\phi}(-k) + \tilde{J}(k)\tilde{\phi}(-k) + \tilde{J}(-k)\tilde{\phi}(k) \right]$$

Note that S_0 is *not* $\int \mathcal{L}_0$, but includes the source.

Define $\tilde{\phi}(k) := \tilde{\chi}(k) + \frac{\tilde{J}(k)}{k^2 + m^2}$. Then $\mathcal{D}\phi = \mathcal{D}\chi$ since we have a ‘‘constant shift transformation’’. This also holds in the $\phi(x)$ space. Then, as in the case of the single oscillator, we get

$$S_0 = \frac{1}{2} \int \frac{d^4k}{(2\pi)^4} \left[\frac{\tilde{J}(k)\tilde{J}(-k)}{k^2 + m^2} - \tilde{\chi}(k)(k^2 + m^2)\tilde{\chi}(-k) \right].$$

The path integral $\int \mathcal{D}\chi$, just gives a factor independent of \tilde{J} , namely $\langle 0|0\rangle_{J=0} = 1$ and we write, $\hbar = 1$ from now on,

$$\begin{aligned}\langle 0|0\rangle_J &= Z_0[J] = \exp \left[i \int \frac{d^4k}{(2\pi)^4} \left\{ \frac{1}{2} \frac{\tilde{J}(k)\tilde{J}(-k)}{k^2 + m^2 - i\epsilon} \right\} \right] \\ &= \exp \left\{ i \int d^4x \int d^4x' \frac{1}{2} J(x)\Delta(x-x')J(x') \right\} \quad \text{where,} \\ \Delta(x-x') &= \int \frac{d^4k}{(2\pi)^4} \frac{e^{ik\cdot(x-x')}}{k^2 + m^2 - i\epsilon} \quad , \quad (\text{Feynman propagator})\end{aligned}$$

Note that the Feynman propagator arose because of the $-i\epsilon$ included in the mass term in the \mathcal{L} .

As discussed for the oscillator, we get

$$\begin{aligned}\langle 0|T \{ \phi(x_1)\phi(x_2) \} |0\rangle &= (-i) \frac{\delta}{\delta J(x_1)} \cdot (-i) \frac{\delta}{\delta J(x_2)} \cdot Z_0[J] \Big|_{J=0} \\ &= (-i) \frac{\delta}{\delta J(x_1)} \left\{ Z_0[J] \cdot \int \Delta(x' - x_2) J(x') d^4x' \right\} (-i)(i) \Big|_{J=0} \Rightarrow \\ \langle 0|T \{ \phi(x_1)\phi(x_2) \} |0\rangle &= -i\Delta(x_1 - x_2) .\end{aligned}$$

Exactly as for the Oscillator, we get

$$\begin{aligned}\langle 0|T \{ \phi(x_1)\phi(x_2)\phi(x_3)\phi(x_4)0 \} |0\rangle &= \\ &= (-i)^2 \left[\Delta(x_1 - x_2)\delta(x_3 - x_4) + \Delta(x_1 - x_3)\delta(x_2 - x_4) + \Delta(x_1 - x_4)\delta(x_2 - x_3) \right]\end{aligned}$$

which easily generalizes to,

$$\begin{aligned}\langle 0|T \{ \phi(x_1) \dots \phi(x_{2n}) \} |0\rangle &= \\ &= (-i)^n \left[\sum_{\text{pairings}} \Delta(x_1 - x_2)\delta(x_3 - x_4) \dots \Delta(x_{2n-1} - x_{2n}) + \text{permutations} \right]\end{aligned}$$

This is what we had obtained as the *Wick's theorem*.

Note: Wick's theorem expressed the time ordered product of quantum fields in terms of the normal ordered product plus contractions. The normal ordering was for all Poincare generators to ensure invariance of the vacuum. The theorem was also proved for free fields for which we do have Fourier decomposition.

Here too the "Wick's theorem" is again seen for free fields. Here it is no more than the chain rule of differentiation.

Consider now interacting fields, by which we mean $\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_1$. Recall that $\langle 0|Q(t)|0\rangle = \int \mathcal{D}q q(t)e^{iS}$ and we can get the $q(t)$ in the integrand by $\frac{\delta}{\delta J(t)} S_J|_{J=0}$ (remember $S = S_J|_{J=0}$).

Thus, insertion of $q(t)$'s or $\phi(x)$'s in field theory, can be effected by $\frac{\delta}{\delta J(x)}S_J|_{J=0}$. If we expand $e^{i\int\mathcal{L}_1(\phi)} = \sum_{n=0}^{\infty}[i\int\mathcal{L}_1]^n/n!$, then we have integrals of polynomials of the fields which can be expressed as $\frac{\delta}{\delta J}S_J$. The δ_J can be taken outside of the path integration. Thus we write,

$$\begin{aligned} Z[J] &:= \langle 0|0\rangle_J^{int} := \int \mathcal{D}\phi \exp i \int d^4x \left(\mathcal{L}_0(\phi) + \mathcal{L}_1(\phi) + J(x)\phi(x) \right) \\ &\propto e^{i\mathcal{L}(-i\delta_J(x))} \cdot \int \mathcal{D}\phi e^{i\int d^4x \{ \mathcal{L}_0 + J(x)\phi(x) \}} \quad \text{or,} \\ Z[J] &\propto \left[\exp i \int d^4x \mathcal{L}_1 \left(-i \frac{\delta}{\delta J(x)} \right) \right] \cdot Z_0[J], \quad \text{with} \end{aligned} \quad (21.2)$$

$$Z_0[J] := \exp \left(\frac{i}{2} \int d^4x \int d^4y J(x)\Delta(x-y)J(y) \right) \quad (21.3)$$

In the oscillator case, for $Z_0[J]$ we had the $\exp \frac{i}{2} \int \int J\Delta J$ term and a Gaussian path integral which turned out to be equal to $Z_0[J=0]$. By taking the asymptotic time limits, we could assert this to be equal to 1 since it was vacuum-to-vacuum amplitude without any source. In presence of interactions, even with $J=0$, the vacuum may not remain unaffected and we cannot justify $\langle 0|0\rangle_{J=0}^{int} = 1$. Instead, the proportionality constant is determined by *demanding* $Z[J=0] = 1$.

The prescription to compute $Z[J]$ via the $e^{i\mathcal{L}_1(-i\delta_J)}Z_0[J]$, is in effect the perturbative prescription. Let us see how this works in an example $\mathcal{L}_1(\phi) = g\phi^3(x)/3!$. Then $Z[J] \propto e^{\frac{g}{6}\delta_J^3} \cdot Z_0[J]$. Expand the exponential,

$$\begin{aligned} Z[J] = \mathcal{N} \sum_{v=0}^{\infty} \frac{1}{v!} \left[\frac{ig}{3!} \int d^4x \left(-i \frac{\delta}{\delta J(x)} \right)^3 \right]^v \times \\ \sum_{p=0}^{\infty} \frac{1}{p!} \left[\frac{i}{2} \int d^4y \int d^4z J(y)\Delta(y-z)J(z) \right]^p. \end{aligned}$$

This too is a power series in J as is $Z_0[J]$, but with different coefficients. Consider a particular term in this double sum with a fixed v and p .



- Then we have $3v$ derivatives acting on $2p$ J 's, leaving $E = (2p - 3v)J$'s. Clearly, $E \geq 0$ must hold and there are several such term;

- The overall *numerical factor* associated with such a group of terms is:

$$\left(\frac{ig}{3!} \right)^v \frac{(-i)^{3v}}{v!} \frac{(i/2)^p}{p!} = \frac{i^{v-3v+p}}{(3!)^v v! p!} g^v = \left(\frac{g}{3!} \right)^v \frac{1}{v! p!} (i)^{E+v-p}$$

- The *combinatorial factor* resulting from the number of ways the derivative acts is: $3v$ derivatives on $2pJ$'s give $\frac{(2p)!}{(2p-3v)!}$ (This is because the first derivative can act on $2p$ J 's, second on $(2p-1)$, ... $(3v)^{th}$ on $(2p-3v+1)$ J 's.)


We can generate and keep track of the various derivatives by denoting

$\frac{ig}{3!} \delta^3 J^3(x)$	\leftrightarrow	$\frac{ig}{3!}$		Has $\int d^4x$
$\frac{i}{2} J(x) \Delta(x-y) J(y)$	\leftrightarrow	$\frac{i}{2}$		Has $\int d^4y \int d^4z$,


and the operation of evaluating the derivative by joining the free ends from the ‘vertices’ to the free ends of the ‘propagator lines’ - exactly as we represented the Wick contractions.


The number of terms with a give v, p is the number of ways of joining the free ends and generating a *diagram*. Some of the diagrams may have the identical factors associated with them For example, the $3\delta J$'s at a given vertex can be permuted in joining up with the propagator lines. This clearly gives a factor of $3!$. Likewise ends of propagator lines gives a factor of $2!$. The diagram will look the same if the v vertices are themselves permuted (the $\int d^4x$ are dummy variables) and this gives $v!$. Similarly the propagator lines give $p!$. Thus, all the numerical factors, except i, g cancel out.

Compared to the previous counting based on Wick’s theorem, we have a double expansion and lines (pairs of J 's) to be contracted with the edges of the vertices. Secondly, in the Wick’s theorem, thanks to the normal ordering of the interaction terms, we don’t have self contractions at the vertices ($\Delta(x, x)$ propagators). In the present approach, there is no “normal ordering” and we do get such term. Eg.



Thus we have more diagrams and different ways of working out the total combinatorial factor. The above bulleted points have gotten rid of all explicit factors, but could involve an over counting. This is compensated by *dividing* by a factor called the *symmetry factor*. This is a geometrical problem of identifying the groups of diagrams. Here are a couple of examples [12].


 We can exchange the two loops in 2 ways. We can exchange the ends of each loop in 2×2 ways. Therefore the total symmetry factor is $2 \cdot 2 \cdot 2 = 8$.


 The 3 propagators can be permuted in $3 \cdot 2$ ways. The 2 vertices can be exchanged in 2 ways. So the symmetry factor is: $3!2! = 12$.

A general statement and a derivation may be seen in the appendix of Serman's book [14].

In the double expansion, we have terms with no J 's, a single J , $2J$'s ... etc. When $J = 0$ is put, only the terms with no J survive. These are termed as the "vacuum bubbles" and are the contributions to the $Z[J = 0]$.

A general contribution to a *given number* of left over J 's would consist of product of topologically connected diagrams. Let C_γ denote the contribution of a particular topologically connected diagram, γ . In a given diagram, Γ , let γ occur n_γ times. The symmetry factor resulting from permutations within each γ , are included in the contribution C_γ . But now we can also have permutations across different, topologically connected diagrams. Such permutations can leave the product diagram invariant if the different copies of a given Γ are permuted as a whole. Hence, we have an additional symmetry factor of $n_\gamma!$ by which we have to divide. A product diagram with different γ 's thus has a symmetry factor of $\prod_\gamma n_\gamma!$.

Returning to the full $Z[J]$ which we can now represent as a contribution from a sum of diagrams Γ , each of which can have several topologically connected γ 's with n_γ copies. Hence,

$$\begin{aligned}
 Z[J] &\propto \sum_{\{n_\gamma\}} C_\Gamma \sim \sum_{\{n_\gamma\}} \prod_\gamma \frac{(C_\gamma)^{n_\gamma}}{n_\gamma!} := \mathcal{N} \prod_\gamma \sum_{n_\gamma=0}^{\infty} \frac{(C_\gamma)^{n_\gamma}}{n_\gamma!} \\
 &= \mathcal{N} \prod_I \exp[C_\gamma] = \mathcal{N} \exp\left[\sum_\gamma C_\gamma\right].
 \end{aligned}$$

Thus, $Z[J]$ is proportional to the exponential of the sum of contributions of topologically connected diagrams. In this sum, are also the contributions topologically connected vacuum diagrams i.e. $Z[J = 0]$. Imposing the normalization condition, $Z[J = 0] = 1$, gives $\boxed{\mathcal{N} \times \exp(\text{topologically connected vacuum bubbles}) = 1}$. Hence, the normalization is trivially incorporated by simply dropping the contributions of the vacuum diagrams and we are

left with,

$$\boxed{Z[J] = \exp[iW[J]] , \quad iW[J] = \sum (\text{topologically connected diagrams.})} \quad (21.4)$$

The Green's functions that we get from $Z[J]$, now get expressed as,

$$\langle 0|\phi(x)|0\rangle = -i \frac{\delta Z[J]}{\delta J(x)} \Big|_{J=0} = \frac{\delta W[J]}{\delta J(x)} \Big|_{J=0} , \quad \because \frac{1}{W[J=0]} = 1.$$

This gives the contribution of diagrams with a single $J(x)$ and gets removed when $J = 0$ is put.

Similarly other Green's functions can be expressed in terms of the derivatives of the $W[J]$.

As an illustration of the organization of the topologically connected diagrams in $W[J]$, consider the two point function.

$$\begin{aligned} G(x_1, x_2) &:= (-i)^2 \frac{\delta^2}{\delta J_1 \delta J_2} Z[J] \Big|_{J=0} = -i \delta_{J_1} (e^{iW} \delta_{J_2} W) \\ &= e^{iW} (\delta_{J_1} W \cdot \delta_{J_2} W - i \delta_{J_1 J_2}^2 W) \Big|_{J=0} \\ \therefore G(x_1, x_2) &= G_c(x_1) G_c(x_2) - G_c(x_1, x_2) \quad , \quad \text{where,} \\ G_c(x) &:= \delta_{J(x)} W[J] \Big|_{J=0} , \quad G_c(x, y) := i \delta_{J(x)J(y)}^2 W[J] \Big|_{J=0} \end{aligned}$$

$$x_1 \text{ --- } \textcircled{G} \text{ --- } x_2 = \underbrace{x_1 \text{ --- } \textcircled{G_c} \text{ --- } \textcircled{G_c} \text{ --- } x_2}_{\text{Top. disconnected}} - \underbrace{x_1 \text{ --- } \textcircled{G_c} \text{ --- } x_2}_{\text{Top. connected}}$$

The G_c denote, contributions from connected (to external points) and topologically connected diagrams. This continues to the other n -point functions, as proved by the argument leading to eqn. (21.4). To conclude,

$G(x_1, \dots, x_n) :$	Σ of all connected diagrams	$:(-i)^n \frac{\delta^n}{\delta J(x_1) \dots \delta J(x_n)} Z[J] \Big _{J=0}$
$G_c(x_1, \dots, x_n) :$	Σ of topologically connected diagrams	$:(-i)^{n-1} \frac{\delta^n}{\delta J(x_1) \dots \delta J(x_n)} W[J] \Big _{J=0}$
$n = 0$	$:\Sigma$ of <i>all</i> vacuum diagrams	$: Z[J = 0] = 1$
$n = 0$	$:\Sigma$ of all topologically connected vacuum diagrams	$: W[J = 0] = 0$

The computation of the diagrams uses the same Feynman rules that we discussed earlier. The Z and W provide a convenient way of dealing all the diagrams together and serve as *generating functions* for the Greens functions.

A. The 1-point function: Renormalization \leftrightarrow normal ordering

The 1-point function of a scalar field has a distinction. Since this represents the $\langle 0|\Phi(x)|0\rangle$, the Poincare invariance allows this to be a *non-zero constant*, say v . No other Lorentz covariant field can have its vacuum expectation value to be non-zero since there are no Lorentz invariant spinors or vectors¹⁵. However, if we want the quanta exchanged during interactions to have the appropriate quantum numbers, the vacuum should *not have any quantum numbers*. As noted in the discussion of uniqueness of vacuum, a non-unique vacuum would have to carry non-trivial representation labels of the symmetry group which will add to the labels of the exchanged quanta. Consequently, we must have a unique vacuum and then, by the mode expansion of the quantum field, its expectation value must vanish. Does our path integral definition of n -point function satisfy this property?

Consider computation of the 1-point function in the $(\phi)_n^3$ theory. Notice that topologically disconnected diagrams contributing to the 1-point function can only be the vacuum bubbles giving a multiplicative contribution of 1. So we can limit to only topologically connected diagrams. Furthermore, if we have $1PR$ diagrams, their contribution is again of a product form: $(1PI) \times \Delta(x-y) \times (1PI) \times \Delta(z-w) \times (1PI) \dots$. Hence it suffices to consider only the $1PI$ diagrams. Thus we have,

$$\bullet \text{---} \textcircled{\text{shaded}} \text{---} G_1(x) = \bullet \text{---} \textcircled{} + \bullet \text{---} \textcircled{\text{vertical}} + \dots$$

At 1-loop (and higher), the diagrams are non-zero and in fact divergent. We invoke the counter term method to absorb away these divergences. Thus, we add a term in \mathcal{L}_1 , of the form $Y\phi(x) = -iY \frac{\delta}{\delta J(x)}$. This is represented by the diagram $\bullet \text{---} \otimes \text{---}$. With its inclusion, the contributing diagrams are:

$$\begin{aligned} \bullet \text{---} \textcircled{\text{shaded}} \text{---} G_1(x) &= x \bullet \text{---} \textcircled{y} + \bullet \text{---} \otimes Y + \bullet \text{---} \textcircled{\text{vertical}} + \bullet \text{---} \otimes Y + \dots \\ &= \quad (a) \quad (b) \quad (2\text{-loop}) \end{aligned}$$

¹⁵ A second rank, symmetric tensor field *can have* $\langle 0|\hat{h}_{\mu\nu}(x)|0\rangle = \Lambda\eta_{\mu\nu}$. But this typically arises only when gravity is included.

These give, to 1-loop,

$$\langle 0|\phi(x)|0\rangle = \left[\underbrace{iY}_{(b)} + \frac{ig}{2} \cdot \underbrace{-i\Delta(0)}_{(a)} \right] \cdot \int d^n y (i\Delta(x-y)) + o(g^3)$$

For the left hand side to be zero to $o(g)$, we must set $\boxed{Y = i\frac{g}{2}\Delta(0) + o(g^3)}$ with,

$$\begin{aligned} \Delta(0) &= \int \frac{d^n k}{(2\pi)^n} \frac{1}{k^2 + m^2 - i\epsilon} \quad (\text{divergent for } n \geq 2.) \\ &= i \int \frac{d^n k}{(2\pi)^n} \frac{1}{k^2 + m^2 - i\epsilon} = i \frac{1}{(4\pi)^{n/2}} \frac{\Gamma(1 - n/2)}{\Gamma(1)} (m^2)^{n/2-1} \quad \text{for } n = 4 - \epsilon \text{ (say)} \\ &= i \frac{1}{16\pi^2} (-1) \frac{2}{\epsilon} (m^2)^{1-\epsilon/2} . \\ \therefore Y &= \frac{g}{16\pi^2} \frac{m^2}{\epsilon} \left(1 - \frac{\epsilon}{2} \ln(m^2) + \dots \right) \approx \frac{g}{16\pi^2} \frac{m^2}{\epsilon} + o(\epsilon^0) + \dots . \end{aligned}$$

Thus, we fix the counter term Y by demanding that the 1-point function be zero, as the renormalization condition and this can be continued at higher orders. The upshot is that we can ensure $\langle 0|\phi(x)|0\rangle = 0$ by means of a counter term. Hence, *the sum of all connected diagrams with a single external line vanishes.*

Clearly, a tadpole can attach to any other diagram only in a 1PR manner (eg replacing the source end of the tadpole by the other diagram). This is a multiplicative contribution and immediately renders all such diagrams to be zero. Hence *all diagrams with a tadpole attached, when summed to a given order, vanish.* We may thus simply remove the tadpole diagrams and obtain results equivalent to those obtained by normal ordering prescription employed before.

Note: Had the 1-point function been *finite*, we would still need to use the counter term method to set it to zero to ensure uniqueness of the vacuum. In the dimensional regularization, massless tadpole diagrams are zero.

Qn: Normal ordering eliminates *all* self contractions - 1-point function in Φ^3 as well as 2-point function in Φ^4 and likewise if higher order vertices are included. For instance, the 1-loop diagram contributing to the 2-point function in Φ^4 . Now there will be a 1-loop counter term for the 2-point function (absent in the normal ordered version). *Does it affect the finite part?*

Ans: The self contractions at any vertex, in any diagram produces a 1-loop contribution as a multiplicative factor, $\Delta_F(0)$ and this multiplicative factor is independent of any momenta entering/exiting the vertex. Consider such a contribution to the self-energy, $\Pi(p^2)$,

in the Φ^4 theory. Unless a counter vertex is introduced, it is not possible to satisfy the renormalization condition, $\Pi(-m^2) = 0$ at the order λ^1 (here we consider on-shell renormalization for simplicity, so that $m^2 = m_{ph}^2$). So introduce an order λ counter vertex. Then at 1-loop, the self energy equals $\Delta_F(0) + \delta_1$. Both are *independent* of p^2 and the renormalization condition implies that two must add to zero. The self energy thus receives no contribution at 1-loop, exactly as if normal ordering has been invoked. In any other diagram (1PI), if a self contraction appears, then the same counter vertex, δ_1 will again automatically cancel the contribution, effectively enforcing normal ordering. *Again this is independent of UV divergence and is a consequence of the requirement of the renormalization condition on self-energy.*

B. Path Integrals and Statistical Mechanics

We discuss briefly an interesting connection of the path integral representation of the transition amplitude and the (grand) partition function of statistical mechanics [16].

The basic observation is that the transition amplitude, schematically $\langle Q_2, t_2 | Q_1, t_1 \rangle = \int \mathcal{D}q \exp\{\frac{i}{\hbar} \int_{t_1}^{t_2} dt L(q(t), \dot{q}(t))\}$, can be taken as a *postulate* rather than a derivation from matrix element of the quantum unitary evolution operator. We can also separately postulate that the transition amplitude *is* the matrix element $\langle Q_2 | e^{-\frac{i}{\hbar}(t_2-t_1)H} | Q_1 \rangle$. Mathematically, set $t_2 - t_1 =: -i\beta$, set $Q_2 = Q_1 =: Q$ and integrate over Q . On the left hand side, we get $Tr e^{-\beta H}$ which is the usual canonical partition function. On the right hand side, the path integral turns to a path integral over *all closed paths* of an integrand which is exponential of a *Euclidean continuation* ($t \rightarrow -i\tau$) of the action. We thus obtain a path integral representation of the statistical mechanical canonical partition function, in terms of Euclidean continuation of a classical action. This is used in the context of the black hole entropy which may be seen in [16].

22. PATH INTEGRALS AS GENERATING FUNCTIONALS

We have noted the two basic quantities, $Z[J]$ and $W[J] = -i \ln Z[J]$. Their functional derivatives with respect to $J(x)$ give the contributions to the n -point ($n \geq 1$) functions: $(-i)^n \delta^n Z[J]|_{J=0}$ contains the contributions of all the connected diagrams while $(-i)^{n-1} \delta^n W[J]$ contains the contributions of all the connected and topologically connected diagrams. The normalization condition: $Z[J=0] = 1 \leftrightarrow W[J=0] = 0$ omits the contributions of the vacuum bubbles. We had encountered the 1PI diagrams while discussing the self-energies. Their contributions too can be obtained by functional differentiation of another quantity which we obtain below. Note that the n -point functions being obtained by functional derivatives also means that $Z[J]$, $W[J]$ can be viewed as *generating functional for Green's functions*.

A. The Generating Functionals: $Z[J]$, $W[J]$, $\Gamma[\Phi]$

Consider a connected and topologically connected diagram contributing to $\delta^n W[J]|_{J=0}$. It is said to be *1-particle reducible* (1PR), if it can be disconnected by cutting *1 internal line*. A diagram which is not 1PR is *1-particle irreducible* (1PI). Any given diagram can be viewed as a collection of 1PI sub-diagrams connected by one internal line. As noted in the discussion of self energy diagrams, all 1PI diagrams with two external lines when strung together by an internal line (free propagator) lead to the exact propagator. Thus, the *sum of all connected and topologically connected diagrams* contributing to an n -point function can be viewed as new connected and topologically connected diagrams whose “vertices”, γ_k are all 1PI diagrams with $k > 2$ edges, connected by lines representing the *exact propagator*. These new diagrams all have a “tree topology” - there are no loops in such a reorganization of diagrams since any loop would make a 1PI sub-diagram and will already be included in one of the γ_k 's. These vertices can be of any order, unlike elementary vertices dictated by the Lagrangian defining a theory. The set of these new diagrams can be generated from another generating functionals,

$$Z_\gamma[J] := \int \mathcal{D}\varphi \exp \left[i \left(\gamma[\varphi] + i \int d^4x J(x) \varphi(x) \right) \right] := e^{iW_\gamma[J]},$$


$$\gamma[\varphi] = \sum_{n \geq 2} \int dx_1 \cdots dx_n \frac{\gamma_n(x_1, \cdots, x_n)}{n!} \varphi(x_1) \cdots \varphi(x_n).$$

Note: The coefficients of the φ^n terms are not simple numbers and this action is *non-local*. The coefficient of the $n = 2$ term, $\gamma_2(x_1, x_2)$ is the Fourier transform of the inverse of the exact propagator. Note that this establishes that $W_S[J] = W_\gamma[J] \Big|_{tree}$.

However, such a $Z_\gamma[J], W_\gamma[J]$ will generate diagrams with loops as well and we are interested in generating only the tree diagrams. To pick out these alone, introduce a fictitious dimensionless parameter, λ and define

$$Z_{\gamma,\lambda}[J] := \int \mathcal{D}\varphi \exp \left[\frac{i}{\lambda} \left(\gamma[\varphi] + i \int d^4x J(x)\varphi(x) \right) \right] := \exp[iW_{\gamma,\lambda}[J]] .$$

Let us count the powers of λ in a any connected and topologically connected diagram. Noting that the γ_2 term in $\gamma[\varphi]$ is the (Fourier transform of the) inverse of the exact propagator, the scaling by λ^{-1} means that that each propagator gives a factor of λ while each vertex and every J, gives a factor of λ^{-1} . Thus, for any diagram contributing to $W_{\gamma,\lambda}[J]$ we get the factor $(\lambda)^{P-V-E}$, where P, V, E are the number of propagator lines, number of vertices and the number “external” lines (lines connected to a source) respectively. The number of “internal” lines is $P - E$. In the diagram below, we have $P = 6, V = 2, E = 4$ giving λ^0 .



For topologically connected diagrams we also have $L = I - V + 1 = P - E - V + 1$. Hence the factor of λ is $(\lambda)^{L-1}$. We can thus write, $W_{\gamma,\lambda}[J] = \sum_{L=0}^{\infty} \lambda^{L-1} W_{\gamma,L}[J]$ and organize the diagrams by the number of loops. In the formal limit, $\lambda \rightarrow 0$, $W_{\gamma,\lambda \rightarrow 0}[J] \rightarrow \frac{1}{\lambda} W_{\gamma,L=0}$ which is the contribution of the tree diagrams (with exact propagators and exact vertices alone).

On the other hand, we may evaluate $Z_{\gamma,\lambda}[J]$ in the limit $\lambda \rightarrow 0$ by stationary phase approximation. Clearly, the path integral is dominated by the fields, $\varphi(x)$ which satisfy $\frac{\delta\gamma}{\delta\varphi} + J = 0$. Denoting $\varphi_{cl}(x)$ as its solution, we get,

$$Z_{\gamma,\lambda \rightarrow 0}[J] \simeq \exp \frac{i}{\lambda} \left\{ \gamma[\varphi_{cl}] + \int d^4x J(x)\varphi_{cl}(x) \right\} = e^{iW_{\gamma,\lambda \rightarrow 0}[J]} \simeq \exp \frac{i}{\lambda} W_{\gamma,L=0}[J].$$

$$\therefore \gamma[\varphi_{cl}] + \int d^4x J(x)\varphi_{cl}(x) = W_\gamma[J] \Big|_{tree} = W_S[J].$$

The last equation shows that $\gamma[\varphi]$ is just the Legendre transform of $W[J]$ and by construction, it is the generating function of connected, topologically connected, 1PI diagrams with external legs amputated! The last property follows because the vertex functions $\gamma_n(x_1, \dots, x_n)$ explicitly do not have external propagator lines included in them.

Note: We have followed a route of organizing the diagrams as a tree of 1PI diagrams connected by exact propagators and inferred the corresponding generating function $\gamma[\varphi]$ as a Legendre transform of the $W[J]$. The usual approach is to begin with a Legendre transform definition and arrive at its interpretation. This conventional approach goes as follows.

Define, $\boxed{\Phi(x)[J] := \frac{\delta W[J]}{\delta J(x)} = \langle 0 | \hat{\Phi}(x) | 0 \rangle_J}$, J is not set to zero and hence this is not the 1-point function. We have the path integral representation as,

$$\Phi(x)[J] = \frac{-i \delta \ln(Z[J])}{\delta J(x)} = \frac{-i}{Z[J]} \frac{\delta Z[J]}{\delta J(x)} = \frac{\int \mathcal{D}\varphi \varphi(x) e^{iS[J]}}{\int \mathcal{D}\varphi e^{iS[J]}} \quad , \quad \boxed{v := \Phi(x)[J=0]} .$$

Define, $\bar{\Phi}(x) := \Phi(x) - v$. Then $\langle 0 | \hat{\Phi}(x) | 0 \rangle = 0$. This generalizes to,

Claim:

$$\frac{\delta^n}{\delta J(x_1) \dots \delta J(x_n)} W[J] \Big|_{J=0} = i^{n-1} \langle 0 | T \{ \bar{\Phi}(x_1) \dots \bar{\Phi}(x_n) \} | 0 \rangle , \quad \forall n \geq 2 .$$

The proof is by induction starting at $n = 2$. For $n = 2$ we have,

$$\begin{aligned} \frac{\delta^2}{\delta J_1 \delta J_2} W[J] \Big|_{J=0} &= \left[\frac{\delta}{\delta J_2} \left(-\frac{i}{Z[J]} \frac{\delta Z[J]}{\delta J_1} \right) \right]_{J=0} = \left[\frac{i}{Z^2} \frac{\delta Z[J]}{\delta J_2} \frac{\delta Z[J]}{\delta J_1} - i \frac{\delta^2 Z[J]}{\delta J_2 \delta J_1} \right]_{J=0} \\ &= \frac{i}{1} (iv)(iv) - i(i^2) \langle 0 | \Phi(x_1) \Phi(x_2) | 0 \rangle \\ &= i \{ \langle 0 | \Phi(x_1) \Phi(x_2) | 0 \rangle - v^2 \} = \langle 0 | \bar{\Phi}(x_1) \bar{\Phi}(x_2) | 0 \rangle . \end{aligned}$$

Which verifies the claim for $n = 2$. The pattern repeats and the proof follows.

As a corollary, we can write

$$W[J] = \sum_{n \geq 2} \frac{1}{n!} \int dx_1 \dots dx_n J(x_1) \dots J(x_n) \frac{\delta^n W}{\delta J_1 \dots J_n} \Big|_{J=0} \quad (22.1)$$

$$= \sum_{n \geq 2} \frac{(i)^{n-1}}{n!} \int dx_1 \dots dx_n \langle 0 | T \{ \bar{\Phi}(x_1) \dots \bar{\Phi}(x_n) \} | 0 \rangle J(x_1) \dots J(x_n) \quad (22.2)$$

Using $\Phi(x)[J] = \delta_{J(x)} W[J]$, define the Legendre transform of $W[J]$,

$$\boxed{\Gamma[\Phi] := W[J] - \int d^4x J(x) \Phi(x)} \quad , \quad (\text{compare with: } -H(p) = L(\dot{q}) - p\dot{q} \text{)} \quad (22.3)$$

Notice that this is exactly the same as the $\gamma[\varphi]$ defined above.

The $\Gamma[\Phi]$ is defined by expressing the right hand side as a function of Φ , in particular $J = J[\Phi]$ is understood which is obtained by inverting the relation $\Phi[J] = \delta_J W[J]$. It follows as usual for a Legendre transform,

$$\frac{\delta\Gamma[\Phi]}{\delta\Phi(x)} = \int d^4y \frac{\delta W}{\delta J(y)} \frac{\delta J(y)}{\delta\Phi(x)} - \int d^4y \frac{\delta J(y)}{\delta\Phi(x)} \Phi(y) - \int d^4y J(y) \frac{\delta\Phi(y)}{\delta\Phi(x)} = -J(x)$$

In the last equality we have used $\frac{\delta\Phi(y)}{\delta\Phi(x)} = \delta^4(x-y)$. Since $\Phi(x)[J=0] = v$, we also have

$\left. \frac{\delta\Gamma[\Phi]}{\delta\Phi(x)} \right|_{\Phi(x)=v} = 0$. Thus, v is that value which extremises $\Gamma[\Phi(x)]$ and hints at $\Gamma[\Phi]$ being some sort of action whose extremization leads to a solution. This is supported further as follows.

We have $Z[J] = e^{iW[J]} = \int \mathcal{D}\varphi e^{iS[J,\varphi]}$. Let $\varphi_{cl}(x)$ be a classical solution i.e. $\delta S[J, \varphi_{cl}] = 0$. Expand the action around such a solution by setting $\varphi(x) = \varphi_{cl}(x) + \eta(x)$. We get (schematically to avoid clutter),

$$\begin{aligned} S[\varphi] &= S[\varphi_{cl} + \eta] = S[\varphi_{cl}] + \underbrace{\frac{\delta S}{\delta\varphi_{cl}}}_{=0} \cdot \eta + \frac{1}{2} \frac{\delta^2 S}{\delta\varphi^2} \eta^2 + \dots \\ &= S[\varphi_{cl}] + \frac{1}{2} \int d^4x d^4y \frac{\delta^2 S[J, \varphi]}{\delta\varphi_{cl}(x)\delta\varphi_{cl}(y)} \eta(x)\eta(y) \\ \therefore e^{iW[J]} &= e^{iS[J,\varphi_{cl}]} \int \mathcal{D}\eta \exp \left[\frac{i}{2} \int d^4x \int d^4y \left\{ \eta(x) \frac{\delta^2 S}{\delta\varphi_{cl}(x)\delta\varphi_{cl}(y)} \eta(y) \right\} + \dots \right] \end{aligned}$$

The path integral does have an implicit dependence on $J(x)$ through the $\varphi_{cl}(x)$, unless the action itself is quadratic as was the case of the oscillator.

Momentarily let us *just ignore* the path integral altogether. Then, $W[J] \approx W_0[J] := S[J, \varphi_{cl}] = \int d^4x [\mathcal{L}(\varphi_{cl}) + J(x)\varphi_{cl}(x)]$ where the classical solution is to first obtained by solving $\delta S[J, \Phi] = 0$ and then substituted back in the action.

Applying our definition, $\Phi(x) = \delta_{J(x)} W[J] \approx \delta_{J(x)} W_0[J] = \delta_{J(x)} S_{cl}[J]$, $S_{cl} := S[J, \varphi_{cl}(x)[J]]$. Straight forward evaluation gives,

$$\Phi(x) = \int d^4y \frac{\delta S_{cl}}{\delta\varphi_{cl}(y)} \underbrace{\frac{\delta\varphi_{cl}(y)}{\delta J(x)}}_{=0} + \varphi_{cl}(x)$$

the last term coming from $\delta_J \int J\varphi$. Hence, in this approximation, $\Phi(x) = \varphi_{cl}(x)$.

Proceeding with the Legendre transform, we get

$$\begin{aligned}\Gamma_0[\Phi] &= W_0[J] - \int d^4x J(x)\Phi(x) \\ &= \underbrace{S[\varphi_{cl}]}_{\text{no explicit } J} + \cancel{\int d^4x J(x)\varphi_{cl}(x)} - \cancel{\int d^4x J(x)\Phi(x)} = \underbrace{S[\varphi_{cl}]}_{\text{no explicit } J} .\end{aligned}\quad (22.4)$$

Thus, within the approximation, $\Gamma[\Phi]$ is just the classical action ($\varphi_{cl}(x) \rightarrow \Phi(x)$).

But this also shows that when the path integral included, $W[J]$ will be very different and so also the corresponding $\Gamma[\Phi]$. Thus, $\Phi(x)$ is identified as the “quantum corrected solution” while the $\Gamma[\Phi]$ is called a “quantum action” or an “effective action” incorporating quantum corrections.

Consider Taylor expanding the effective action about $\Phi[x] = v$. $\Gamma[v] = W[J=0] - \int 0 \cdot \phi = 0$ since $W[0] = 0$ by our normalization, $Z[0] = 1$. We have also seen that $\delta_v \Gamma = 0$. The remaining terms are a power series in $(\Phi(x) - v) =: \bar{\Phi}(x)$ with the coefficients evaluated at v . This is the same as regarding Γ as a function of $\bar{\Phi}(x)$ and expanding it about $\bar{\Phi}(x) = 0$ i.e.

$$\Gamma[\bar{\Phi}] = \sum_{n \geq 2} \int dx_1 \cdots dx_n \frac{\Gamma(x_1, \cdots, x_n)}{n!} \bar{\Phi}(x_1) \cdots \bar{\Phi}(x_n) .$$

What do these coefficients represent? We already have the answer. We just need to recognize that $\Gamma[\bar{\Phi}] = \gamma[\varphi]!$ The coefficients are the contribution of all connected, topologically connected, 1PI diagrams without external legs. An explicit demonstration of amputation of the external legs may be seen in [17]. The generating functions are summarized in the table below.

In summary

$$Z[J] := \int \mathcal{D}\varphi e^{iS[J,\varphi]} \quad , \quad Z[J=0] = 1 \quad , \quad (\text{normalization}) \quad (22.5)$$

$$G(x_1, \dots, x_n) := \left(-i \frac{\delta}{\delta J(x)} \right)^n Z[J] \Big|_{J=0} = \langle 0|T\{\Phi(x_1) \dots \Phi(x_n)\}|0\rangle \quad (22.6)$$

$$W[J] := -i \ln(Z[J]) \quad , \quad W[J=0] = 0 \quad (22.7)$$

$$G_c(x_1, \dots, x_n) := (-i)^{n-1} \frac{\delta^n W[J]}{\delta J(x_1) \dots \delta J(x_n)} \Big|_{J=0} = \langle 0|T\{\Phi(x_1) \dots \Phi(x_n)\}|0\rangle_c \quad (22.8)$$

$$\Phi(x)[J] := \frac{\delta W[J]}{\delta J(x)} \quad , \quad v := \Phi(x)[J=0] \quad , \quad \bar{\Phi}(x) := \Phi(x) - v \quad (22.9)$$

$$\Gamma[\Phi] := W[J] - \int d^4x J(x)\varphi(x) \quad (22.10)$$

$$J(x) = -\frac{\delta\Gamma[\Phi]}{\delta\Phi(x)} \quad , \quad \frac{\delta\Gamma[\Phi]}{\delta\Phi(x)} \Big|_{\Phi(x)=v} = 0 \quad (22.11)$$

$$Z[J] = \sum_{n \geq 1} \frac{i^{n-1}}{n!} \int d^4x_1 \dots d^4x_n G(x_1, \dots, x_n) J(x_1) \dots J(x_n) \quad (22.12)$$

$$W[J] = \sum_{n \geq 1} \frac{i^{n-1}}{n!} \int d^4x_1 \dots d^4x_n G_c(x_1, \dots, x_n) J(x_1) \dots J(x_n) \quad (22.13)$$

$$\Gamma[\bar{\Phi}] = \sum_{n \geq 2} \frac{1}{n!} \int d^4x_1 \dots d^4x_n \Gamma_n(x_1, \dots, x_n) \bar{\Phi}(x_1) \dots \bar{\Phi}(x_n) \quad (22.14)$$

$$G(x_1, \dots, x_n) \leftrightarrow \text{all connected diagrams} \quad (22.15)$$

$$G_c(x_1, \dots, x_n) \leftrightarrow \text{all connected and topologically diagrams} \quad (22.16)$$

$$\Gamma(x_1, \dots, x_n) \leftrightarrow \text{all connected, topologically connected, 1PI diagrams} \\ \text{with external legs amputated} \quad (22.17)$$

B. The Renormalization Group Equation

Recall that while discussing the renormalized perturbation series, we introduced ϕ_0, m_0, g_0 as the ‘bare’ quantities with $\mathcal{L}(\phi_0, m_0, g_0)$ generating the diagrams containing the UV divergences. We then introduced the renormalized variables ϕ, m, g_k defined through the scaling: $\phi_0 := \sqrt{Z_\phi} \phi$, $m_0^2 Z_\phi := m^2 Z_m$, $g_0 Z_\phi^{k/2} := Z_{g_k} g_k$ which were finite by definition. We also introduced counter terms to take care of the UV divergences. With the Z, G, Γ we have the

formal representation of the totality of all diagrams. Going over to the momentum space labels we have (we take the 1-point function to be zero and a single coupling constant for convenience),

$$\begin{aligned}\Gamma_{0,n}(p_1, \dots, p_n) &:= \frac{\delta\Gamma_0[\phi_0]}{\delta\phi_0(p_1) \dots \delta\phi_0(p_n)} \Big|_{\phi_0=0} \quad , \quad \Gamma_n(p_1, \dots, p_n) := \frac{\delta\Gamma_0[\sqrt{Z_\phi}\phi]}{\delta\phi(p_1) \dots \delta\phi(p_n)} \Big|_{\phi=0} \\ \Rightarrow \Gamma_{0n}(p_i, m_0, g_0) &= Z_\phi^{-n/2} \Gamma_n(p_i, m, g)\end{aligned}\quad (22.18)$$

The renormalized parameters are defined by a set of renormalization conditions, which introduce a scale, say, μ (eg a MS or \overline{MS} scheme). The bare quantities know nothing about this scale and therefore $\mu \frac{d\Gamma_0(p_i, m_0, g_0)}{d\mu} = 0$. Substitution gives,

$$\begin{aligned}0 &= \left(\mu \frac{d}{d\mu} Z_\phi^{-n/2} \right) \Gamma_n + Z_\phi^{-n/2} \mu \frac{d}{d\mu} \Gamma_n \quad \text{and,} \\ \mu \frac{d}{d\mu} \Gamma_n &= \left[\mu \frac{\partial}{\partial \mu} + \mu \frac{dm}{d\mu} \frac{\partial}{\partial m} + \mu \frac{dg}{d\mu} \frac{\partial}{\partial g} \right] \Gamma_n(p_i, m, g)\end{aligned}$$

Thus we have the renormalization group (RG) equation for the renormalized vertex function as,

$$\mu \frac{\partial}{\partial \mu} \Gamma_n(p_i, m(\mu), g(\mu)) \Big|_{m,g} + \beta(g) \frac{\partial}{\partial g} \Gamma_n \Big|_{\mu,m} - \gamma_m(g) m \frac{\partial}{\partial m} \Gamma_n \Big|_{\mu,g} - n\gamma_\phi \Gamma_n = 0, \quad (22.19)$$

where,

$$\beta(g, m) := \mu \frac{dg}{d\mu} \quad , \quad \gamma(g, m) := -\mu \frac{d \ln(m)}{d\mu} \quad , \quad \gamma_\phi(g, m) := \frac{1}{2} \mu \frac{d \ln(Z_\phi)}{d\mu}. \quad (22.20)$$

The first is the *beta function*, the second is called the *mass anomalous dimension* while the last is called the *anomalous dimension*. These are computed typically in a perturbation theory as a power series in the renormalized coupling (see below). Different schemes give different simplifications and there is some *scheme dependence* in the coefficients of these functions.

As mentioned above, we describe a method of computation of the renormalization constants and their dependence on the couplings in the MS scheme [18].

We have the defining equations (22.19, 22.20). These were obtained by noting that the bare vertex functions as a function of the bare parameters and the regulator are independent of μ , for fixed g_0, m_0 and ϵ . When the renormalized parameters are *not* defined in terms of physically measured quantities, how do we compute the μ dependence of these parameters?

For this, we go back to the basic definitions and recall the relation between the bare and the renormalized parameters, in particular, $g_0 := \mu^\epsilon Z_g g$. Note that we have absorbed the Z_ϕ factor into Z_g and have also introduced the μ dependence by taking g to be dimensionless. The $\mu \partial_\mu$ is now evaluated with the bare parameters and the regulator ϵ held fixed i.e.

$$\beta(g) = \mu \partial_\mu g(g_0 \mu^{-\epsilon}, \epsilon) \Big|_{g_0, \epsilon} = \mu \partial_\mu (g_0 \mu^{-\epsilon} Z_g^{-1}) = -\epsilon g - g \mu \partial_\mu \ln(Z_g) .$$

Noting that Z_g is determined as a function of g , $Z_g(g, \epsilon) = 1 + \sum_{k \geq 1} \epsilon^{-k} Z_g^k$, we can write the defining equation for the β function in the form,

$$\beta(g, \epsilon) = -\epsilon g - g \beta(g, \epsilon) \partial_g \ln(Z_g(g, \epsilon)) \leftrightarrow \beta(g, \epsilon) \partial_g (g Z_g(g, \epsilon)) + \epsilon g Z(g, \epsilon) = 0.$$

The $\beta(g, \epsilon)$ must have a smooth limit as $\epsilon \rightarrow 0$ and we may consider $\beta \sim \beta_0 + \beta_1 \epsilon + \beta_2 \epsilon^2 + \dots$. The $\partial_g (g Z_g(g, \epsilon)) \sim 1 + \epsilon^{-1} + \dots$ while $\epsilon g Z_g(g, \epsilon) \sim \epsilon + \epsilon^0 + \epsilon^{-1} + \dots$. Clearly, the coefficients of the positive powers of ϵ greater than 1 must vanish. Hence we take, $\beta(g, \epsilon) := \beta_0(g) + \beta_1 \epsilon$. It follows that $\beta_1 = -g$ and the equation reduces to,

$$0 = (\beta_0 - \epsilon g)(Z_g + g \partial_g Z_g) \quad (22.21)$$

$$0 = \beta_0 \left[1 + \sum_{k \geq 1} \frac{1}{\epsilon^k} \partial_g (g Z_g^k) \right] - \left[g^2 \partial_g Z_g^1 + \sum_{k \geq 1} \frac{g^2 \partial_g Z_g^{k+1}}{\epsilon^k} \right] \quad (22.22)$$

$$\therefore \boxed{\beta_0(g) = g^2 \partial_g Z_g^1, \quad \beta_0 \partial_g (g Z_g^k) = g^2 \partial_g Z_g^{k+1}.} \quad (22.23)$$

The $\beta_0(g)$ is the usual beta function and is obtained from the renormalization constant Z_g . The $Z_g^{k > 2}$ coefficients are also determined recursively.

Note: The above expressions are in the context of the Φ^4 coupling. For Yukawa or the Yang-Mills cubic coupling, the coupling has $g_{Yukawa} = \mu^{\epsilon/2} g_{dimensionless}$ and the above equations will change accordingly.

As a practical application of the renormalization group equation, consider a theory which is massless and remains so perturbatively. Then the anomalous mass dimension term is absent. Putting $t := \ln(\mu/\mu_0)$ for some arbitrary scale μ_0 at which the theory is defined (renormalization conditions are imposed), we write $\mu \frac{\partial}{\partial \mu} = \frac{\partial}{\partial t}$ and the equation takes the form,

$$\left[\frac{\partial}{\partial t} + \beta(g) \frac{\partial}{\partial g} - n \gamma(g) \right] \Gamma_n(g, t, p_i) = 0 \quad \text{with} \quad \beta(g) = \frac{dg(t)}{dt}$$


This immediately gives, $\frac{d\Gamma_n(g(t), t, p_i)}{dt} = 0$. Its solution is obtained as,

$$\boxed{\Gamma_n(t, g_0, p_i) = \Gamma_{n,0}(g(t, g_0), p) \exp \left\{ n \int_0^t dt' \gamma(g(t', g_0)) \right\}} \quad \text{where,} \quad (22.24)$$

$g(t, g_0)$ is the solution: $\frac{dg(t, g_0)}{dt} = \beta(g(t, g_0))$, $g(0, g_0) = g_0$.

Although typically the beta function is a power series in the coupling. Once $g(t)$ is known, the t dependence of *all* vertex functions is determined. A good deal of qualitative behavior can be gleaned from studying the beta function - especially near its zeros. We know it has a zero at $g = 0$ (perturbative calculation), but it may have other zeros. The derivative of $\beta(g)$ near its fixed points, controls how $g(t)$ evolves with t .

Let us assume it to be some given function of the form as shown in the figure.



It has multiple zeros, say $g_0 = 0 < g_1 < g_2 < g_3 \dots$. Let I, II, III, \dots denote the intervals on the g -axis, bounded by the zeros. Clearly, given a g_0 in one of these intervals, it will remain so for all t . If $\beta(g) > 0$ in an interval, then $g(t)$ will evolve to its upper bound and the opposite if $\beta(g) < 0$. A coupling will always flow to a UV-stable (we are considering increasing t) fixed point. In the figure, g_1, g_3 are the stable fixed points while g_0, g_2 are unstable ones.

Let g_0 be in region I so that as $t \rightarrow \infty, g(t, g_0) \rightarrow g_1$. The asymptotic behavior of Γ_n is controlled by the integral of the anomalous dimension. Let us write,

$$\int_0^t dt' \gamma(g(t', g_0)) = \int_0^t dt' [\gamma(g(t', g_0)) - \gamma(g_1)] + \int_0^t dt' \gamma(g_1).$$

In the first term, we may take the upper limit $t \rightarrow \infty$ since the integrand provides exponential suppression. This integral is then some constant, C . In the second term, $\gamma(g_1)$ is independent of t' and thus evaluates to $\gamma(g_1) \cdot t$. Hence,

$$\Gamma_n(t, g_0, p_i) \xrightarrow{t \rightarrow \infty} \left[\Gamma_{n,0}(g_1, p_i) \cdot C \right] \cdot \exp\{\gamma(g_1)t\} \quad , \quad t = \ln(\mu/\mu_0) .$$

μ is some large scale eg some of the invariants $p_i \cdot p_j$. Thus, we get $\Gamma_n(\ln(\mu/\mu_0), g_0, p_i) \xrightarrow{\mu \rightarrow \infty} (\mu/\mu_0)^{\gamma(g_1)} \forall n$. The exponent is the same for all n -point vertex functions and is governed by

a stable fixed point in the vicinity of g_0 . All these vertex functions have the same asymptotic behavior. For further application, I refer you to [19].

C. The Background Field Method

As an illustration of the utility of the formal advantage of the functional methods, we discuss the so called *background field method*, extremely useful for non-abelian gauge theories [20]. For illustration, we consider the simplest case of a scalar field φ .

We have the basic definitions

$$e^{iW[J]} = Z[J] := \int \mathcal{D}\varphi e^{iS[\varphi] + \int J(x)\varphi(x)}, \quad \bar{\varphi}(x) := \frac{W[J]}{J(x)}, \quad \Gamma[\bar{\varphi}] := W[J(\bar{\varphi})] - \int J(x)\bar{\varphi}(x).$$

Introduce a new, arbitrary *background field*, $\chi(x)$ and define,

$$e^{i\tilde{W}[J,\chi]} = \tilde{Z}[J,\chi] := \int \mathcal{D}\varphi e^{iS[\varphi+\chi] + \int J(x)\varphi(x)},$$

$$\tilde{\varphi}(x) := \frac{\tilde{W}[J,\chi]}{J(x)}, \quad \tilde{\Gamma}[\tilde{\varphi},\chi] := W[J(\varphi,\chi)] - \int J(x)\tilde{\varphi}(x).$$

In the new generating functionals defined, shift the integration variable as $\varphi \rightarrow \varphi - \chi$. This is a simple translation and gives the Jacobian to be 1. It is immediate that $\tilde{Z}[J,\chi] = Z[J]e^{-i\int J(x)\chi(x)}$ and $\tilde{W}[J,\chi] = W[J] - \int J(x)\chi(x)$. Clearly, $\tilde{\varphi}(x) = \bar{\varphi}(x) - \chi(x)$ and

$$\tilde{\Gamma}[\tilde{\varphi},\chi] = W[J] - \int J(x)\chi(x) - \int J(x)(\bar{\varphi}(x) - \chi(x)) = \Gamma[\bar{\varphi}] = \Gamma[\tilde{\varphi} + \chi].$$

If we had arranged that $\bar{\varphi} = 0$, then $\Gamma[\chi] = \tilde{\Gamma}[0,\chi]$. The background field χ being arbitrary, we get an alternate method of computing the effective action $\Gamma[\chi]$ using the shifted fields.

The shifted action has the form:

$$S[\varphi + \chi] = S[\chi] + \int \mathcal{L}_1(\chi)\varphi(x) + \int \mathcal{L}_2(\chi)\varphi^2(x) + \dots.$$

Since there is no functional integration over χ , the first term is just the classical action. Since χ is arbitrary, the second term does *not* vanish (it vanishes if χ is an exact solution of the classical equations of motion). The third term gives the propagator for the φ field and the higher order terms give interactions among the φ, χ fields.

Now, the shifted effective action, $\tilde{\Gamma}$ is a generator of 1PI diagrams in presence of $\tilde{\varphi}$ and χ . That is, its derivatives with respect to $\tilde{\varphi}$ will give the 1PI diagrams in presence of χ .

The φ propagator, which comes from the terms in the action which are quadratic in φ , will in general be χ -dependent and the vertices will have factors of $\chi(x)$. When $\tilde{\varphi} = 0$, only the diagrams with no external $\tilde{\varphi}$ lines i.e. only the *vacuum diagrams* will contribute. Thus the *desired effective action can be computed using only vacuum diagrams, albeit with vertex factors having the χ field.*

There are two ways to approach the computation. If we treat the background field exactly, then in obtaining the Feynman rules the propagator of the φ field will have the χ field as well (the terms quadratic in φ in $S[\varphi + \chi]$). Except when χ is space-time independent, this propagator is complicated. And for a general background field, this method is not useful. An alternative is to treat χ also perturbatively i.e. the φ propagator remains exactly same as before and the additional χ -dependent terms are treated as additional couplings (the φ factors are replaced as $\frac{\delta}{\delta J}$). The Feynman rules have the same φ -propagator, there is no propagator for the χ field and the vertices have extra edges denoting the background field. Since there is no χ propagator, these vertices generate *only* external χ lines.

As an explicit example, consider the $(\varphi^3)_6$ theory. The shift by a background field gives,

$$S[\varphi + \chi] = S[\chi] + S[\varphi] + \int d^4x \left[-\partial_\mu \varphi \partial^\mu \chi - m^2 \varphi(x) \chi(x) - \frac{g}{2!} (\varphi(x) \chi^2(x) + \varphi^2(x) \chi(x)) \right]$$

The $S[\chi]$ comes out of the path integral, the $S[\varphi]$ gives the usual φ -propagator and the φ^3 vertex while the last four terms give the additional interaction vertices.

Since in the 1PI diagrams we are interested in, internal lines are only φ -lines and the external lines are only χ lines, the first three of the four additional vertices are irrelevant and only the φ^3 and $\varphi^2 \chi$ vertices remain. It is a simple exercise in power counting to determine the superficially divergent 1PI diagrams with no internal χ -lines and no external φ -lines. Consider a 1PI diagram with $n_{0,3}$ number of φ^3 vertices and $n_{1,2}$ number of $\chi \varphi^2$ vertices. Then the internal φ -lines is given by $2I_\varphi = 3n_{0,3} + 2n_{1,2}$ and the external χ -lines if given by $E_\chi = n_{1,2}$. It is easy to see that the superficial degree of divergence, D , is then given by $D = 6 - 2E_\chi$. Notice that $n_{0,3}$ drops out of the degree of divergence. The renormalizability of the theory is thus manifest and only the 2-point and the 3-point vertex functions are divergent.

How does renormalization work in background field method? Re-scale the fields as $\varphi \rightarrow \sqrt{Z_\varphi} \varphi$, $\chi \rightarrow \sqrt{Z_\chi} \chi$. Then the φ -propagator will get a factor of Z_φ^{-1} . The $n_{0,3}$ vertices give a factor of $(Z_\varphi^{3/2})^{n_{0,3}}$ and the $n_{1,2}$ vertices give a factor of $Z_\varphi^{n_{1,2}} \cdot (Z_\chi^{1/2})^{n_{1,2}}$. The I_φ lines

give a factor of $Z_\varphi^{-I_\varphi}$. Clearly, the factors of Z_φ cancel out since $-I_\varphi + \frac{3}{2}n_{0,3} + \frac{2}{2}n_{1,2} = 0$. We may thus choose not to renormalise the φ fields. The left over factor is $Z_\chi^{n_{1,2}/2} = Z_\chi^{E_\chi/2}$ as expected.

Notice that the diagrams that need to be computed for any vertex function, are exactly the same as without background field except for the vertex factors from the vertices connecting the external lines. The renormalization procedure(s) then proceed as usual eg as discussed in the subsection 19 B.

For scalar fields used in the illustration, there is no particular advantage. But with gauge theories the method allows enormous simplification [20]. This is beyond the scope of this course though.

Returning to our generating functionals Z, W, Γ ; we have seen how the perturbative diagrams for the Green's functions and the S -matrix elements (through the vertex functions) can be subsumed by formal manipulations with the basic path integral. The utility of the formalism goes beyond perturbation theory and computations of cross-sections. When it comes to gauge theories, especially the non-abelian ones with/without spontaneous symmetry breaking, the generating functional provide a convenient tool to establish renormalizability and unitarity. The proof of the Ward identities - constraints or relations enforced by gauge invariance on different n -point functions - is most economical in this framework. The existence of *anomalies* - violation of classical symmetries at the quantum level and their treatment is also much more transparent in these functional methods.

The basic path integral is defined without any pre-supposition of perturbation theory. If the basic definition can be implemented eg via a lattice discretization, we can potentially have a non-perturbative handle through the computation of the vertex functions.

23. CLOSING REMARKS

The lectures were organized around four strands. The main references used for each strand are also cited.

1. Poincare symmetry realization and its consequences [1, 2]:

This involved the particle representations which identified the attributes of the permissible quanta;

Manifest covariance needs the field representation and the irreducibility condition imposed the *linear* field equations; The unitarity analysis revealed anti-particles;

Action formulation showed the free classical fields as a dynamical system of infinitely many independent oscillators; Introduction of interaction with non-dynamical source brought in the various Green's function, the Feynman propagator conflicting with a causally consistent classical interpretation;

Quantization of free fields required their covariance to be formulated differently and led to the CPT theorem; requirement of causality lead to the spin-statistics theorem;

States of Free fields naturally contain particles as the relativistic wave packets and pave the way for recovering the usual non-relativistic limit.

2. Interacting quantum fields [9–11]:

This was discussed within the context of scattering experiments; corresponding scattering theory postulates were discussed leading to the Kallen-Lehmann spectral representation; vanishing of the 1-point function is required for the stability of (unique) vacuum;

LSZ reduction was discussed to obtain the S-matrix elements to vacuum expectation values of time ordered fields; covariant perturbation theory was premised on the interacting fields have the same canonical commutation relation as the in/out fields which are deemed *physical - masses and residues at poles*; S-matrix elements are given entirely in terms of free quantum fields with arbitrarily specified but Lorentz invariant interactions and a diagrammatic recipe follows;

This characterizes interacting quantum fields as facilitating discrete transactions of energy/momentum/spin/charge etc via *virtual* quanta. The off-shell quanta do not

satisfy the equations of motion and hence represent a model of quantum fluctuations.

3. Application to the Yukawa interaction and QED up to 1-loop [12, 13]:

The early successes at tree level were discussed followed by the Radiative corrections; The IR divergences highlighted the care needed in applying the formalism to what is actually measured;

The UV divergences highlighted the unacceptable, unbounded dominance of the quantum fluctuations and questioned the purely theoretical parameters in the Lagrangian ('bare'); Some renormalization procedure needs to be adopted to identify the physically measured parameters;

The generic problem is to be handled recursively using counter terms which were illustrated at the 2-loop level; The renormalization process also led to the renormalization group equation.

4. The path integral formulation [12, 17, 18]:

The classical action has a direct and central role in computing quantum transition amplitudes; The path integral provides generating functionals for various n -point functions and is a powerful tool for both formal studies as well as for richer non-abelian gauge theories with/without spontaneous symmetry breaking (SSB);

When extended to the more general theories - non-abelian gauge theories with/out SSB, it follows that one needs to identify the 'correct fields' and their interactions *before computing the various n -point functions*; The qualitative physics guesses are invoked to propose the choice of *the unique vacuum*. In the gauge theory context, for instance, gauge invariance under large gauge transformations (non-trivial at infinity) reveal the θ -vacua and some choice is made by nature.

-
- [1] Steven Weinberg, *The Quantum Theory of Fields volume I*, Cambridge University Press, 1995.
- [2] C. Wetterich, Massless Spinors in More Than Four Dimensions, *Nucl. Phys. B* **211**, 177 (1983).
- [3] G Date, Lecture on Constrained Systems, [arXiv:1010.2062](https://arxiv.org/abs/1010.2062).
- [4] Hong-Hao Zhang, Kai-Xi Feng, Si-Wei Qiu, An Zhao, Xue-Song Li, A Note on analytic formulas of Feynman propagators in position space, *Chin.Phys.* **C34**, 1576 (2010), [arXiv:0811.1261](https://arxiv.org/abs/0811.1261).
- [5] W. Greiner and J. Reinhardt, *Quantum Electrodynamics*, 4th edition, Springer, 2009, page 68.
- [6] D F Walls and G J Milburn, *Quantum Optics*, second edition, Springer-Verlag, 2008.
- [7] R J Glauber, *Quantum Theory of Optical Coherence: Selected Papers and Lectures*, Wiley-VCH Verlag GmbH & Co. KGaA, 2007.
- [8] R. Jagannathan and S. A. Khan, *Quantum Mechanics of Charged Particle Beam Optics*, CRC Press, 2019.
- [9] M Reed and B Simon, *Methods of Modern Mathematical Physics, Scattering Theory, Vol III*, Academic Press, 1979.
- [10] R G Newton, *Scattering Theory of Waves and Particles*, chapters 6 and 7, Springer-Verlag, NY, 1982.
- [11] J D Bjorken and S D Drell, *Relativistic Quantum Fields*, chapters 16 and 17, McGraw Hill, 1965.
- [12] M Srednicky, *Quantum Field Theory*, Cambridge University Press, 2007.
- [13] M Peskin and D V Schroeder, *Introduction to Quantum Field Theory*, Addison-Wesley, 1995.
- [14] George Stermann, *An Introduction to Quantum Field Theory*, Cambridge University Press, 1993.
- [15] S. Elaydi, *An Introduction to Difference Equations*, Springer, 2005.
- [16] G W Gibbons and S W Hawking, Action integrals and partition functions in quantum gravity, *Phys. Rev. D*, **15**, 2752 (1977).
- [17] E S Abers and B W Lee, Gauge Theories, *Physics Reports*, **9**, 1, (1973).
- [18] D Gross, *Method in Field Theory*, Les Houches, Ed. R Balian and J Zinn-Justin, 1975.
- [19] S Coleman, *Quantum Field Theory: Lectures of Sydney Coleman*, Chapter 50, World Scien-

tific, 2019.

- [20] L F Abbot, Introduction to the background field method, *Acta Physica Polonica*, a**B13**, 33 (1982).