

# A Simple Proof that Major Index and Inversions are Equidistributed

Michael J. Collins  
 Daniel H. Wagner Associates  
 mjcollins10@gmail.com

July 13, 2022

## Abstract

We present a short proof of MacMahon’s classic result that the number of permutations with  $k$  inversions equals the number whose major index (sum of positions at which descents occur) is  $k$ .

## 1 Introduction

Let  $p = p_0 p_1 \cdots p_{n-1}$  be a permutation of  $[n] = \{0, 1, \dots, n-1\}$ . A *descent* of  $p$  is an index  $i$  at which  $p_{i-1} > p_i$ , and an *inversion* of  $p$  is a pair of indices  $i < j$  with  $p_i > p_j$ . Define  $\text{inv}(p)$  to be the number of inversions in  $p$ , and define  $\text{maj}(p)$ , the “major index” of  $p$ , to be the sum of all descent positions (so for instance  $\text{maj}(241350) = 2 + 5 = 7$ ). See ([1, 4]) for other standard definitions and results regarding permutations.

MacMahon ([3]) proved that  $\text{inv}$  and  $\text{maj}$  are *equidistributed*: the number of length- $n$  permutations with  $\text{inv}(p) = k$  equals the number of such permutations with  $\text{maj}(p) = k$ . This common value is denoted  $b(n, k)$ . MacMahon originally proved this by showing that the generating functions coincide, and Foata ([2]) gave a bijective proof; in this note we present a simpler proof.

## 2 Proof of Equidistribution

An *inversion table* of length  $n$  is an  $n$ -tuple of nonnegative integers  $(a_0, a_1, \dots, a_{n-1})$  such that  $a_j \leq j$  for all  $j$ . Clearly there are  $n!$  inversion tables, and each represents a distinct permutation of  $[n]$  as follows: starting with the empty permutation, we repeatedly insert  $j$  so that it will have  $a_j$  items to its right<sup>1</sup>. For instance the inversion table  $(0, 1, 0, 3, 3)$

---

<sup>1</sup>to simplify our presentation we have reversed the usual convention which would have  $a_j \leq n - j$

yields the permutation 31402, building it as

$$\begin{array}{r}
 0 \\
 10 \\
 102 \\
 3102 \\
 34102
 \end{array} \tag{1}$$

The insertion of  $j$  creates  $a_j$  inversions, proving the well-known result that  $b(n, k)$  is the number of inversion tables whose elements sum to  $k$ . To prove equidistribution, we reinterpret  $(a_0, a_1, \dots, a_{n-1})$  as meaning repeated insertion of  $j$  at a position that will increase the major index by  $a_j$ . Finding such a position is always possible. For instance (using boldface to emphasize descents) we have  $\text{maj}(24\mathbf{1}35\mathbf{0}) = 2 + 5 = 7$ , and the possibilities for insertion of 6 are:

$$\begin{array}{r}
 \text{maj}(24\mathbf{1}35\mathbf{0}6) = 7 + 0 = 2 + 5 \\
 \text{maj}(24\mathbf{1}35\mathbf{6}0) = 7 + 1 = 2 + 6 \\
 \text{maj}(24\mathbf{1}36\mathbf{5}0) = 7 + 6 = 2 + 5 + 6 \\
 \text{maj}(24\mathbf{1}6\mathbf{3}50) = 7 + 5 = 2 + 4 + 6 \\
 \text{maj}(246\mathbf{1}350) = 7 + 2 = 3 + 6 \\
 \text{maj}(26\mathbf{4}1350) = 7 + 4 = 2 + 3 + 6 \\
 \text{maj}(6\mathbf{2}41350) = 7 + 3 = 1 + 3 + 6
 \end{array} \tag{2}$$

In general, say the  $\kappa$  inversions of a permutation of  $[j]$  occur at positions  $d_\kappa < d_{\kappa-1} < \dots < d_1$ . Inserting  $j$  at the rightmost position will not change the major index. Insertion at  $d_t$  ( $1 \leq t \leq \kappa$ ) will create no new descents, but the descents at  $d_t$  through  $d_1$  will be shifted to positions  $d_t + 1, \dots, d_1 + 1$ , so  $\text{maj}$  will increase by  $t$ . Finally, consider inserting  $j$  at the  $r^{\text{th}}$  position (from the left) which is *not* a descent: if there are  $r'$  descents to the left of this position, we create a new descent at  $r + r'$  and shift  $\kappa - r'$  old descents to the right, increasing the major index by  $\kappa + r$ . Thus the number of permutations with  $\text{maj}(p) = k$  is again the number of inversion tables with entries summing to  $k$ .

### 3 Symmetric Joint Distribution

Our proof of equidistribution is simpler than Foata's, but the machinery of Foata's proof can be used to prove the stronger result that  $\text{maj}$  and  $\text{inv}$  have a *symmetric joint distribution* ([2, 4]): for any pair of integers  $k, k'$  the number of  $p$  with  $\text{inv}(p) = k, \text{maj}(p) = k'$  equals the number with  $\text{inv}(p) = k', \text{maj}(p) = k$ . We now note that this result can be stated entirely in terms of inversion tables.

To do this we define another way to interpret an inversion table  $(a_0, \dots, a_{n-1})$  as a way to build a permutation, one which makes the relationship between  $\text{inv}$  and  $\text{maj}$  more direct. Now  $a_j$  will mean "put  $j - a_j$  in the rightmost position, and increment all other elements

which are greater than or equal to  $j - a_j$ ". More formally, if  $(a_0 \cdots a_{j-1})$  generates the permutation  $p_0 \cdots p_{j-1}$  then  $(a_0 \cdots a_j)$  generates the permutation  $p'$  with  $p'_j = j - a_j$  and otherwise

$$p'_k = p_k + [p_k > j - a_j]$$

Here we make use of the "Iverson bracket" notation, where  $[S] = 1$  if the statement  $S$  is true, 0 if it is false.

In fact this just yields the inverse of the permutation generated by reading  $(a_0, \cdots a_{n-1})$  as an inversion table. For instance our previous example of  $(0, 1, 0, 3, 3)$  now yields the permutation 32401, building it as

$$\begin{array}{r} 0 \\ 10 \\ 102 \\ 2130 \\ 32401 \end{array} \tag{3}$$

At step  $j$  we create  $a_j$  new inversions; the increments do not change any existing inversions, since a pair  $r < s$  is either unchanged or becomes  $r + 1 < s + 1$  or  $r < s + 1$ . Furthermore we create a descent at position  $j$  if and only if  $a_j > a_{j-1}$  (i.e. if position  $j$  is an *ascent* of  $a$ ), and similarly the increments do not destroy or create any descents. So the resulting permutation  $p$  has

$$\begin{aligned} \text{inv}(p) &= \sum a_j \\ \text{maj}(p) &= \sum_{a_j > a_{j-1}} j \end{aligned} \tag{4}$$

Therefore, since  $\text{inv}$  and  $\text{maj}$  are equidistributed over permutations, the "sum of elements" and "sum of ascent positions" are equidistributed over the set of all inversion tables.

## References

- [1] Miklos Bona. *Combinatorics of Permutations*, volume 1 of *Discrete Mathematics and its Applications*. Chapman and Hall/CRC, 3 edition, 2022.
- [2] Dominique Foata. On the netto inversion number of a sequence. *Proceedings of the American Mathematical Society*, 19(1):236–240, 1968.
- [3] P. MacMahon. Two applications of general theorems in combinatory analysis. *Proceedings of the London Mathematical Society*, 15:314–321, 1916.
- [4] Richard P. Stanley. *Enumerative Combinatorics*, volume 1 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, New York, NY, 2 edition, 2012.