

A SHORT PROOF OF A CONJECTURE OF MATSUSHITA

BENJAMIN BAKKER

ABSTRACT. In this note we build on the arguments of van Geemen and Voisin [18] to prove a conjecture of Matsushita that a Lagrangian fibration of an irreducible hyperkähler manifold is either isotrivial or of maximal variation. We also complete a partial result of Voisin [20] regarding the density of torsion points of sections of Lagrangian fibrations.

Let X be an irreducible compact hyperkähler manifold, that is, a simply-connected compact Kähler manifold X for which $H^0(X, \Omega_X^2) = \mathbb{C}\sigma$ for a nowhere-degenerate holomorphic two-form σ . A Lagrangian fibration of X is a proper morphism $f : X \rightarrow B$ to a normal compact analytic variety B whose generic fiber is smooth, connected, and Lagrangian (see [11] for a recent survey). It follows that every smooth fiber is an abelian variety. We let $B^\circ \subset B$ be a dense Zariski open smooth subset over which the base-change $f^\circ : X^\circ \rightarrow B^\circ$ is smooth. By the period map of f we mean the period map $\varphi : B^\circ \rightarrow S$ to an appropriate moduli space S of polarized abelian varieties associated to the natural variation of (polarized) weight one integral Hodge structures on B° with underlying local system $R^1 f_*^\circ \mathbb{Z}_{X^\circ}$. We say f is *isotrivial* if the period map is trivial (equivalently if $R^1 f_*^\circ \mathbb{Z}_{X^\circ}$ has finite monodromy) and of *maximal variation* if the period map is generically finite.

Our main result is to resolve a conjecture of Matsushita:

Theorem 1. *Let X be an irreducible hyperkähler manifold (or more generally a primitive symplectic variety in the sense of [2]). Then any Lagrangian fibration $f : X \rightarrow B$ is either isotrivial or of maximal variation.*

Both possibilities in Theorem 1 occur for K3 surfaces S —see for example [10, Chapter 11]—and therefore also for their Hilbert schemes $S^{[g]}$ in each (even) dimension. Primitive symplectic varieties are the natural singular analog (as far as deformation theory is concerned) of irreducible hyperkähler manifolds; see below for the definition and the precise meaning of a Lagrangian fibration in this context.

Let $T_0 \subset H^2(X, \mathbb{Q})$ be the rational transcendental lattice, namely, the smallest rational Hodge substructure containing $[\sigma] \in H^{2,0}(X)$. Theorem 1 was proven by van Geemen and Voisin [18, Theorem 5] assuming X is projective, that T_0 has generic (special) Mumford–Tate group (namely $\mathrm{SO}(T_0, q_X)$, where q_X is the Beauville–Bogomolov–Fujiki form), and that $\mathrm{rk} T_0 \geq 5$, by showing that under these conditions any fiber of a Lagrangian fibration that is not of maximal variation must be a factor of the Kuga–Satake variety of T_0 . Their result in particular applies to the generic projective deformation of $f : X \rightarrow B$ assuming $b_2(X) \geq 7$, which includes all known deformation types. The Beauville–Mukai system of a nonprimitive basepoint-free ample divisor on a K3 surface has recently been shown to be of maximal variation in many cases by Dutta–Huybrechts [7] (see Corollary 7 below) by understanding the derivative of the period map.

We will instead prove Theorem 1 by considering the complex variation of Hodge structures on $R^1 f_*^\circ \mathbb{C}_{X^\circ}$. We first recall the basic properties of complex variations. A

complex variation of Hodge structures on a Zariski open subset of a compact analytic variety (see for example [5]) consists of a \mathbb{C} -local system V and a holomorphic (resp. antiholomorphic) descending filtration F^\bullet (resp. \overline{F}^\bullet) such that we have a splitting of the sheaf of C^∞ sections $A^0(V) = \bigoplus_p A^0(V^p)$ where $V^p = F^p \cap \overline{F}^{-p}$ and the flat connection maps $A^0(V^p)$ to $A^{1,0}(V^{p-1}) \oplus A^1(V^p) \oplus A^{0,1}(V^{p+1})$. We refer to the grading V^p as the Hodge grading and we say the level of the variation is the difference $p_{\max} - p_{\min}$ where p_{\max} (resp. p_{\min}) is the maximum (resp. minimum) Hodge degree p for which $V^p \neq 0$. Observe that the level of a tensor product $V \otimes W$ is the sum of the levels of V and W . A polarization of the variation is a flat hermitian form h for which the splitting is orthogonal and $(-1)^p h$ is positive definite on V^p . In this case $\overline{F}^{-p} = (F^{p+1})^\perp$. A variation which admits a polarization is said to be polarizable. We define $\mathbb{C}(-d)$ to be the polarizable complex Hodge structure on $V = \mathbb{C}$ with $V^d = V$.

Recall that the category of polarizable complex variations of Hodge structures is semi-simple. The theorem of the fixed part [17] says that for two polarizable complex variations V, W , the group $\text{Hom}(V, W)$ of morphisms of local systems has a natural complex Hodge structure whose degree p part is exactly the morphisms of complex variations $V \rightarrow W(p)$. We have the following further consequence due to Deligne:

Theorem 2 ([5, 1.13 Proposition]). *Suppose V is a \mathbb{C} -local system underlying a polarizable complex variation of Hodge structures and that we have a splitting of \mathbb{C} -local systems*

$$(1) \quad V = \bigoplus_i N_i \otimes A_i$$

where the N_i are irreducible and pairwise non-isomorphic and the A_i are nonzero complex vector spaces. Then

- (1) Each N_i underlies a polarizable complex variation of Hodge structures, unique up to shifting the Hodge grading.
- (2) Each polarizable complex variation of Hodge structures with underlying local system V arises from (1) by equipping each N_i with its unique (up to shifts) polarizable complex variation of Hodge structures and each A_i with a uniquely determined polarizable complex Hodge structure, namely $A_i = \text{Hom}(N_i, V)$.

In particular, the theorem implies a polarizable complex variation is irreducible if and only if the underlying local system is.

Remark 3. The \mathbb{C} -local system V underlying a polarizable complex variation of Hodge structures on an algebraic space is semi-simple. This is a consequence of an unpublished result of Nori (see [5, §1.12]) and uses that the orthogonal complement of flat subbundle with respect to the hodge metric (which is a harmonic metric) is flat. In our setting, we will only need to apply this to V underlying a polarizable integral variation of Hodge structures. In this case, the semi-simplicity of the underlying \mathbb{Q} -local system is a result of Deligne [6, 4.2.6]. Note that for a perfect field K and a field extension $K \subset L$, a K -local system V is semi-simple if and only if V_L is, as semi-simplicity is equivalent to being generated by simple sub-local systems.

Given an \mathbb{R} -local system V , a polarizable real variation of Hodge structures¹ on V in the usual sense naturally induces a polarizable complex variation on $V_{\mathbb{C}}$. Conversely, a polarizable complex variation on $V_{\mathbb{C}}$ comes from a polarizable real variation on V if complex conjugation flips the Hodge grading, or more precisely if for some (hence any) polarization h the isomorphism of local systems $V_{\mathbb{C}} \rightarrow V_{\mathbb{C}}^{\vee}$ given by $y \mapsto h(-, \overline{y})$ induces

¹Throughout, by a real variation we mean a pure real variation, unless otherwise specified.

an isomorphism of complex variations $V_{\mathbb{C}} \rightarrow V_{\mathbb{C}}^{\vee}(-w)$ for some (uniquely determined) w . Indeed, if this is the case then $V^p \xrightarrow{\cong} (V^{w-p})^{\vee}$ so $\overline{V^p} = V^{w-p}$. Moreover, for even w (resp. odd w) a real polarization is provided by the symmetric (resp. antisymmetric) real form $q(x, y) = h(x, \bar{y}) + h(y, \bar{x})$ (resp. $q(x, y) = i(h(x, \bar{y}) - h(y, \bar{x}))$), since $q(x, \bar{x}) = h(x, x) + h(\bar{x}, \bar{x})$ (resp. $-iq(x, \bar{x}) = h(x, x) - h(\bar{x}, \bar{x})$) is definite of alternating sign on V^p .

The category of polarizable real variations is also semi-simple. Observe that by Theorem 2 any isotypic component W of an \mathbb{R} -local system V underlying a polarizable real variation is canonically a real subvariation, as the same is true over \mathbb{C} and the isomorphism $V_{\mathbb{C}} \rightarrow V_{\mathbb{C}}^{\vee}(-w)$ coming from h restricts to an isomorphism $W_{\mathbb{C}} \rightarrow W_{\mathbb{C}}^{\vee}(-w)$. If V is a single isotypic factor, then $V_{\mathbb{C}}$ either has one self-conjugate irreducible factor N or has two non-isomorphic conjugate irreducible factors N, \bar{N} . Note that $N^{\vee} \cong \bar{N}$ via the polarization. Note also that the level of a polarizable real variation V is at least as large as the level of any of the irreducible factors of $V_{\mathbb{C}}$.

We say that a real or complex variation is *isotrivial* if the Hodge filtration is flat, or equivalently if the irreducible factors of the complexification are level zero². To summarize the above discussion:

Lemma 4. *Let V be an irreducible polarizable real variation of Hodge structures of level one. Then V is either isotrivial, or every irreducible factor of $V_{\mathbb{C}}$ is level one.*

Before turning to the proof of Theorem 1 we recall the definition of a primitive symplectic variety. Let X be a symplectic variety in the sense of Beauville³ [3], that is, a compact Kähler variety with rational singularities and a nowhere degenerate 2-form σ on its regular locus X^{reg} . We say that X is primitive symplectic if $H^1(X, \mathcal{O}_X) = 0$ and $H^0(X^{\text{reg}}, \Omega_{X^{\text{reg}}}^2) = \mathbb{C}\sigma$. As the singularities are rational, for any resolution $\pi : Y \rightarrow X$ the form σ extends to a two-form on Y [12, Corollary 1.7]. Moreover, $\pi^* : H^2(X, \mathbb{Q}) \rightarrow H^2(Y, \mathbb{Q})$ is injective, so the Hodge structure on $H^2(X, \mathbb{Q})$ is pure, and we have an induced isomorphism $\pi^* : H^{2,0}(X) \rightarrow H^{2,0}(Y)$ (see [2] for details). In particular we have a well defined class $[\sigma] \in H^{2,0}(X)$.

By a Lagrangian fibration of a primitive symplectic variety we still mean a proper morphism $f : X \rightarrow B$ to a normal compact analytic variety B whose generic fiber is smooth, connected, and Lagrangian. Each smooth fiber will still be an abelian variety. Moreover, B is in fact Kähler and Moishezon [19] (and in particular an algebraic space) since f is equidimensional as in [11, Lemma 1.17], now using functorial pullback of reflexive forms [12, Theorem 1.11] and the fact that $R\pi_*\omega_Y \cong \omega_X \cong \mathcal{O}_X$ by the rationality of the singularities of X [13, §5.1].

We use the same notation as above: $B^{\circ} \subset B$ is a dense Zariski open smooth subset over which the restriction $f^{\circ} : X^{\circ} \rightarrow B^{\circ}$ is smooth and $\varphi : B^{\circ} \rightarrow S$ is the period map associated to the variation of (polarized) weight one integral Hodge structures on B° with underlying local system $R^1 f_*^{\circ} \mathbb{Z}_{X^{\circ}}$.

Proof of Theorem 1. Let $V_{\mathbb{Z}} := R^1 f_*^{\circ} \mathbb{Z}_{X^{\circ}}$. We start with the following result of Voisin, whose proof we give for convenience (and to extend it slightly).

Lemma 5 ([20, Lemma 5.5]). *$V_{\mathbb{R}}$ is irreducible as a polarizable real variation of Hodge structures.*

Proof. First assume X is smooth. By a result of Matsushita [14, Lemma 2.2] the restriction map $H^2(X, \mathbb{R}) \rightarrow H^2(X_b, \mathbb{R})$ to a generic fiber of f° is rank one and by

²Or equivalently, if the monodromy is unitary (by Theorem 2); since there may not be an integral structure, this does not necessarily mean the monodromy is finite.

³This definition is equivalent to Beauville's original one.

Deligne's global invariant cycles theorem $H^2(X, \mathbb{R}) \rightarrow H^0(B^\circ, R^2 f_* \mathbb{R}_{X^\circ})$ is surjective [6]. If $V_{\mathbb{R}}$ splits as a variation then the polarizations of the factors would yield a larger than one-dimensional space of sections of $R^2 f_* \mathbb{R}_{X^\circ} = \wedge^2 V_{\mathbb{R}}$, which is a contradiction.

Now if X is a primitive symplectic variety, one easily checks using the results of [2] that Matsushita's proof carries through verbatim and that $H^2(X, \mathbb{R}) \rightarrow H^0(B^\circ, R^2 f_* \mathbb{R}_{X^\circ})$ is still surjective, since the cokernel of $\pi^* : H^2(X, \mathbb{R}) \rightarrow H^2(Y, \mathbb{R})$ is generated by exceptional divisors for a log resolution $\pi : Y \rightarrow X$ since X has rational singularities. \square

Suppose now that f is not of maximal variation. Define the real transcendental lattice $T \subset H^2(X, \mathbb{R})$ to be the polarizable real Hodge substructure spanned by $[\sigma]$ and $[\bar{\sigma}]$. We next claim that the polarizable real variation of Hodge structures $V_{\mathbb{R}} \otimes T^\vee$ has a nontrivial subvariation of level at most one after a finite base-change; the argument below is that of [18] at its core, with some mild modifications.

Let $\nu : B^{o'} \rightarrow B^\circ$ be a finite Galois étale cover so that the base-change $V_{\mathbb{Z}}' := \nu^* V_{\mathbb{Z}}$ is pulled back along its period map $\varphi' : B^{o'} \rightarrow S'$, where S' is a level cover of S . Note that up to replacing $B^{o'}$ with a further finite cover, we may assume φ' can be embedded in a proper map $\bar{\varphi}' : \bar{B}^{o'} \rightarrow S'$ [9]. Denote by $Z \subset S'$ the image of $\bar{\varphi}'$, by $\psi : B^{o'} \rightarrow Z$ the resulting map, and by $V_{\mathbb{Z}}''$ the variation on Z so that $V_{\mathbb{Z}}' = \psi^* V_{\mathbb{Z}}''$. The map $\bar{\varphi}'$ and its image Z are in fact algebraic [4, Theorem 3.1]. We shrink Z (and $B^\circ, B^{o'}, X^\circ$) so that it is smooth and so that $R^1 \psi_* \mathbb{R}_{B^{o'}}$ is a local system, naturally underlying a graded polarizable real variation of mixed Hodge structures whose only nonzero Hodge components are $(0, 0), (1, 0), (0, 1), (1, 1)$ (for example using Saito's theory of mixed Hodge modules [15, 16]). Let $f^{o'} : X^{o'} \rightarrow B^{o'}$ be the base-change of f . The natural map $H^2(X, \mathbb{C}) \rightarrow H^0(B^{o'}, R^2 f_* \mathbb{C}_{X^{o'}})$ sends $[\sigma]$ and hence $T_{\mathbb{C}}$ to zero, since the fibers of f are Lagrangian. By the Leray spectral sequence we have a natural morphism $\text{pt}_Z^* T \rightarrow R^1 \psi_* V_{\mathbb{R}}' \cong V_{\mathbb{R}}'' \otimes R^1 \psi_* \mathbb{R}_{B^{o'}}$ in the category of real variations of mixed Hodge structures. This map is nonzero from the following geometric description as in [18].

Through a very general point $b \in B^{o'}$, say above a point $z \in Z$, let F be the positive-dimensional fiber of ψ through b . The restricted family $X_F \rightarrow F$ has trivial monodromy so the following natural diagram commutes

$$\begin{array}{ccc} T_{\mathbb{C}} & \longrightarrow & (V_{\mathbb{C}}'' \otimes R^1 \psi_* \mathbb{C}_{B^{o'}})_z \\ \downarrow & & \downarrow \\ H^2(X_F, \mathbb{C}) & \longrightarrow & H^1(X_b, \mathbb{C}) \otimes H^1(F, \mathbb{C}) \end{array}$$

where the bottom arrow comes from the degeneration of the Leray spectral sequence for $X_F \rightarrow F$. In the projective case we have $X_F \cong X_b \times F$ (possibly after a further base-change) and this map is just the Künneth projection. The image of $[\sigma]$ is nonzero in the bottom right corner since: (i) σ is nonzero when restricted to X_F since $\dim X_F > \frac{1}{2} \dim X$; (ii) $\sigma|_{X_F}$ extends to a smooth compactification since σ extends to a smooth compactification of X° , so $[\sigma] \neq 0 \in H^2(X_F, \mathbb{C})$; (iii) the image of $[\sigma]$ in $H^2(X_b, \mathbb{C}) \otimes H^0(F, \mathbb{C})$ vanishes and $[\sigma]$ is not in the image of $H^0(X_b, \mathbb{C}) \otimes H^2(F, \mathbb{C})$, as it is not pulled back from F .

Thus, there is a nonzero morphism of real variations

$$(2) \quad \text{gr}_{-1}^W \psi^* (R^1 \psi_* \mathbb{R}_{B^{o'}})^\vee \rightarrow V_{\mathbb{R}}' \otimes T^\vee.$$

As the category of polarizable real variations of (pure) Hodge structures is semi-simple, we therefore have a splitting

$$V_{\mathbb{R}}' \otimes T^\vee = U \oplus W$$

of real variations, where $U \neq 0$ is the image of (2). In particular, U has level at most one and weight -1.

Now by Lemma 5 the Galois group of ν acts transitively on the isotypic factors of $V'_\mathbb{R}$. In particular, if f (and therefore $V_\mathbb{R}$) is not isotrivial, no factor of $V'_\mathbb{R}$ (as a variation) is isotrivial, or else its entire isotypic component would be, and so would $V_\mathbb{R}$. But then there can be no nonzero morphism of variations $V'_\mathbb{R} \otimes T^\vee \rightarrow U$. Indeed, by Lemma 4, an irreducible factor N of $V'_\mathbb{C}$ has level one of degrees 0,1, and $N \otimes T^\vee_\mathbb{C}$ can only map nontrivially to an irreducible factor of $U_\mathbb{C}$ of the form $N(1)$, while $\text{Hom}(N \otimes T^\vee_\mathbb{C}, N(1)) = T_\mathbb{C}(1) \cong \mathbb{C}(-1) \oplus \mathbb{C}(1)$ has no degree 0 elements. Thus, f must be isotrivial. \square

Example 6. We revisit the example from [18, §4]. Let $p \geq 5$ be a prime and λ a p th root of unity. Consider a family of abelian varieties $f : X \rightarrow B$ with a cyclic automorphism such that the induced automorphism α of $V_\mathbb{R} = R^1 f_* \mathbb{R}_X$ has λ as an eigenvalue on $V^{1,0}$ but not on $V^{0,1}$. Let α' be the automorphism of T^\vee with eigenvalue λ^{-1} on $(T^\vee)^{-2,0}$ and eigenvalue λ on $(T^\vee)^{0,-2}$. Then $V_\mathbb{R} \otimes T^\vee$ has a level one factor, namely the 1 eigenspace $(V_\mathbb{R} \otimes T^\vee)^1$ of $\alpha \otimes \alpha'$. But the condition on the eigenvalues means the eigenspaces $(V_\mathbb{C})^\lambda$ and $(V_\mathbb{C})^{\lambda^{-1}}$ are level zero, and the real variation $(V_\mathbb{C})^\lambda \oplus (V_\mathbb{C})^{\lambda^{-1}}$ is an isotrivial real factor.

We now discuss a few applications of Theorem 1. Dutta–Huybrechts have proven many cases [7, Theorem 1.1] of the following corollary, and have also shown that Theorem 1 implies the general case:

Corollary 7. *Let H be a basepoint-free ample divisor on a K3 surface S . Then the complete linear system $|H|$ is of maximal variation.*

Proof. The genus 2 case is proven unconditionally in [7, Prop. 5.4], and the genus $g \geq 3$ case in [7, Prop. 5.2] assuming Theorem 1. \square

The next application uses a result of Gao, which is a simple application of the Ax–Schanuel theorem for universal families of abelian varieties [8, Theorem 1.1]. Recall that for a projective family $f : X \rightarrow B$ of g -dimensional abelian varieties⁴ equipped with a section s and letting $\tilde{B} \rightarrow B^{\text{an}}$ be the universal cover, the Betti map $\beta : \tilde{B} \rightarrow H_1(X_b, \mathbb{R})$ is the real analytic map obtained by taking the coordinates of the section s with respect to the flat real-analytic trivialization of f . Observe that $\beta^{-1}(H_1(X_b, \mathbb{Q}))$ is the set of points of \tilde{B} at which s is torsion. If $\varphi : B \rightarrow S$ is the period map of f and $\mathcal{X} \rightarrow S$ the universal family of abelian varieties, then s naturally yields a map $B \rightarrow \mathcal{X}$ lifting φ . We say that s is of maximal variation if $B \rightarrow \mathcal{X}$ is generically finite.

Proposition 8 ([8, Theorem 9.1]). *Let $f : X \rightarrow B$ be a projective family of g -dimensional abelian varieties with $\dim B \geq g$ and whose very general fiber has no nontrivial \mathbb{Q} -factor. Let $s : B \rightarrow X$ be a non-torsion section of f which is of maximal variation. Then the Betti map $\beta : \tilde{B} \rightarrow H_1(X_b, \mathbb{R})$ associated to s is generically submersive.*

Corollary 9. *Let X be a primitive symplectic variety and $f : X \rightarrow B$ a Lagrangian fibration. Let L be a line bundle whose restriction to the smooth fibers is topologically trivial. Then the set of points $b \in B^\circ(\mathbb{C})$ for which $L|_{X_b}$ is torsion is analytically dense in B .*

Corollary 9 was proven by Voisin [20, Theorem 1.3] assuming either f is of maximal variation and $\dim X \leq 8$ or isotrivial with no restriction on the dimension. Our use of Proposition 8 replaces the results of André–Corvaja–Zannier [1] in [20].

⁴Meaning X is an abelian scheme over B , so there is in particular a 0-section of f .

Proof of Corollary 9. By Voisin’s result and Theorem 1 we may assume f is of maximal variation. Note that Voisin’s proof works equally well in the singular case; we leave the details to the reader. Consider the family of abelian varieties $h : \text{Pic}^0(X^\circ/B^\circ) \rightarrow B^\circ$ and the section $s : b \mapsto L|_{X_b}$. Let $\nu : B^{o'} \rightarrow B^\circ$ be a Galois finite base-change for which the \mathbb{Q} -factors of the very general fiber of h are defined over $B^{o'}$. As the Galois group of ν acts transitively on the factors by Lemma 5, the d factors all have the same dimension g' , and the image of the period map of each factor must have dimension $\geq g'$, or else the image of the period map of f would have dimension smaller than $dg' = \dim(X^\circ/B^\circ) = \dim(B^\circ)$. The base-change of the section s is also Galois invariant, so it suffices to prove the density statement for its projection to a single factor $Y^\circ \rightarrow B^{o'}$. Applying Proposition 8, the Betti map $\beta : \tilde{B} \rightarrow H_1(Y_b, \mathbb{R})$ is submersive, so $\beta^{-1}(H_1(Y_b, \mathbb{Q}))$ is analytically dense in \tilde{B} as claimed. \square

Corollary 9 has an interesting interpretation in terms of the Beauville conjecture for irreducible hyperkähler manifolds X , see the discussion in [20, §1.2]. There it is also shown how corollaries 7 and 9 can be used to construct constant cycle curves on K3 surfaces.

Acknowledgements. The author would like to thank C. Voisin for conversations related to Theorem 1 and Corollary 9, D. Maulik for bringing the paper [20] to his attention, and C. Lehn for conversations related to Lagrangian fibrations in the singular case. This work was partially supported by NSF grant DMS-2131688.

REFERENCES

- [1] Y. André, P. Corvaja, and U. Zannier. The Betti map associated to a section of an abelian scheme. *Invent. Math.*, 222(1):161–202, 2020.
- [2] B. Bakker and C. Lehn. The global moduli theory of symplectic varieties. *Journal für die reine und angewandte Mathematik (Crelles Journal)*, 2022.
- [3] A. Beauville. Symplectic singularities. *Invent. Math.*, 139(3):541–549, 2000.
- [4] A. Borel. Some metric properties of arithmetic quotients of symmetric spaces and an extension theorem. *J. Differential Geometry*, 6:543–560, 1972.
- [5] P. Deligne. Un théorème de finitude pour la monodromie. In *Discrete groups in geometry and analysis (New Haven, Conn., 1984)*, volume 67 of *Progr. Math.*, pages 1–19. Birkhäuser Boston, Boston, MA, 1987.
- [6] Pierre Deligne. Théorie de Hodge. II. *Inst. Hautes Études Sci. Publ. Math.*, (40):5–57, 1971.
- [7] Y. Dutta and D. Huybrechts. Maximal variation of curves on K3 surfaces. [arXiv:2105.04787](https://arxiv.org/abs/2105.04787), 2021.
- [8] Z. Gao. Mixed Ax-Schanuel for the universal abelian varieties and some applications. *Compos. Math.*, 156(11):2263–2297, 2020.
- [9] P. A. Griffiths. Periods of integrals on algebraic manifolds. III. Some global differential-geometric properties of the period mapping. *Inst. Hautes Études Sci. Publ. Math.*, (38):125–180, 1970.
- [10] D. Huybrechts. *Lectures on K3 surfaces*, volume 158 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2016.
- [11] D. Huybrechts and M. Mauri. Lagrangian fibrations. *Milan J. Math.*, 2022.
- [12] S. Kebekus and C. Schnell. Extending holomorphic forms from the regular locus of a complex space to a resolution of singularities. *J. Amer. Math. Soc.*, 34(2):315–368, 2021.
- [13] J. Kollár and S. Mori. *Birational geometry of algebraic varieties*, volume 134 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1998.
- [14] D. Matsushita. On deformations of Lagrangian fibrations. In *K3 surfaces and their moduli*, volume 315 of *Progr. Math.*, pages 237–243. Birkhäuser/Springer, [Cham], 2016.
- [15] M. Saito. Modules de Hodge polarisables. *Publ. Res. Inst. Math. Sci.*, 24(6):849–995 (1989), 1988.
- [16] M. Saito. Mixed Hodge modules. *Publ. Res. Inst. Math. Sci.*, 26(2):221–333, 1990.
- [17] W. Schmid. Variation of Hodge structure: the singularities of the period mapping. *Invent. Math.*, 22:211–319, 1973.
- [18] B. van Geemen and C. Voisin. On a conjecture of Matsushita. *Int. Math. Res. Not. IMRN*, (10):3111–3123, 2016.

- [19] J. Varouchas. Sur l'image d'une variété kählérienne compacte. In *Fonctions de plusieurs variables complexes, V (Paris, 1979–1985)*, volume 1188 of *Lecture Notes in Math.*, pages 245–259. Springer, Berlin, 1986.
- [20] C. Voisin. Torsion points of sections of Lagrangian torus fibrations and the Chow ring of hyper-Kähler manifolds. In *Geometry of moduli*, volume 14 of *Abel Symp.*, pages 295–326. Springer, Cham, 2018.

B. BAKKER: DEPT. OF MATHEMATICS, STATISTICS, AND COMPUTER SCIENCE, UNIVERSITY OF ILLINOIS AT CHICAGO, CHICAGO, USA.

Email address: `bakker.uic@gmail.com`