

On the dynamics and integrability of the Ziegler pendulum

Ivan Yu. Polekhin *

*Steklov Mathematical Institute of Russian Academy of Sciences,
Moscow, Russia*
ivanpolekhin@mi-ras.ru

Abstract

We prove that the Ziegler pendulum — a double pendulum with a follower force — can be integrable, provided that the stiffness of the elastic spring located in the pivot point of the pendulum is zero. We show that the integrability of the system follows from the existence of two-parameter families of periodic solutions. We explain the mechanism for the transition from the integrable to the chaotic dynamics and show that this mechanism is different from the typical phenomena observed in conservative Hamiltonian systems. The case when the stiffnesses of the both springs are non-zero is studied numerically. We show that the regular dynamics co-exists with the chaotic dynamics.

Key words: first integrals, non-potential force, system on a torus, non-integrability, periodic solutions.

1 Introduction

The Ziegler pendulum is a well known non-conservative mechanical system with many real-life applications. This system is mostly known for the ‘paradox’ of loss of stability in the presence of friction, which was discovered 70 years ago [1] (see also [2], [3]). More detailed exposition of this and related systems and the history of the problem can be found in [4]. The full bibliography of works on the Ziegler pendulum and other non-conservative systems is too extensive to be presented here and we also refer the reader to book [4]. However, we would like to note that, for the best of our knowledge, the dynamics *in the large* and integrability of the Ziegler pendulum has not been studied before in the literature. At the same time, there are many papers on the dynamics of the classical double pendulum, where it is shown experimentally [5], numerically [6], and analytically [7] by means of the Poincare-Melnikov method that this system is chaotic. First non-trivial results on the non-integrability of systems in non-potential force fields have been obtained in [8]. The problem of integrability of the Ziegler pendulum has been also formulated in [8].

In the paper we study the dynamics of a planar double pendulum in the presence of a follower force of constant magnitude. We assume that this follower force is acting along the rod that is not connected

*Electronic address: ivanpolekhin@mi-ras.ru

to the fixed pivot point. There is no gravity force in the system, yet there are two springs of linear stiffness located in the joints of the double pendulum.

The paper contains both analytical and numerical results. By means of an analytical approach we study the case when the stiffness of the spring located in the pivot point is zero. Thanks to the rotational symmetry, we can consider the reduced system of the third order. We prove that in the reduced system there exists a two-parametric family of periodic solutions. It means that locally there exist two first integrals of the system and one can consider this system as integrable, at least for some initial conditions. We explain the mechanism of loss of periodicity of solutions associated with the change of parameters or initial conditions. We show how the birth of non-periodic solutions leads to the non-integrability of the system.

The case when the stiffnesses of the springs are non-zero is studied numerically. We again obtain the co-existence of two types of solutions in the phase space: regular solutions which lie on two-dimensional surfaces and chaotic trajectories. We show this by means of the Poincare sections and Lyapunov characteristic exponents.

2 Equations of motion

Let us consider a planar double mathematical pendulum. One of its absolutely rigid massless rod of length l_2 can rotate freely in the plane about a fixed point O . Two other rigid massless rods of lengths l_1 and l_3 are connected to the free end A of the first rod OA . Let us denote the ends of these rods by B and C , correspondingly (Fig. 1). We assume that points A , B , and C always remain on a straight line when the pendulum moves. There is a point mass m_2 in point A and point masses m_1 and m_3 in points B and C , correspondingly. There is a force F of constant magnitude acting along segment BC . Two springs of stiffnesses k_1 and k_2 are located in points A and O , correspondingly.




Figure 1: The Ziegler pendulum.

Let φ_2 be a generalized coordinate, the angle between a constant direction in the plane and segment OA and φ_1 be the angle between OA and BC . The kinetic energy of the system can be expressed in these coordinates as follows:

$$T = \frac{m_1}{2} (\dot{\varphi}_2^2 l_2^2 + (\dot{\varphi}_1 + \dot{\varphi}_2)^2 l_1^2 + 2l_1 l_2 \dot{\varphi}_2 (\dot{\varphi}_1 + \dot{\varphi}_2) \cos \varphi_1) + \frac{m_2}{2} \dot{\varphi}_2^2 l_2^2 + \frac{m_3}{2} (\dot{\varphi}_2^2 l_2^2 + (\dot{\varphi}_1 + \dot{\varphi}_2)^2 l_3^2 - 2l_2 l_3 \dot{\varphi}_2 (\dot{\varphi}_1 + \dot{\varphi}_2) \cos \varphi_1). \quad (1)$$

The potential energy:

$$\Pi = \frac{k_1\varphi_1^2}{2} + \frac{k_2\varphi_2^2}{2}. \quad (2)$$

The components of generalized force:

$$Q_1 = 0, \quad Q_2 = -Fl_2 \sin \varphi_1. \quad (3)$$

From (1), (2), and (3) we obtain the Lagrange equations, which can be presented as follows

$$\begin{aligned} A_{11}\ddot{\varphi}_1 + A_{12}\ddot{\varphi}_2 &= r_1, \\ A_{21}\ddot{\varphi}_1 + A_{22}\ddot{\varphi}_2 &= r_2, \end{aligned} \quad (4)$$

where

$$\begin{aligned} A_{11} &= m_1l_1^2 + m_3l_3^2, \\ A_{12} = A_{21} &= m_1l_1^2 + m_3l_3^2 + m_1l_1l_2 \cos \varphi_1 - m_3l_2l_3 \cos \varphi_1, \\ A_{22} &= m_1l_2^2 + m_1l_1^2 + m_2l_2^2 + m_3l_2^2 + m_3l_3^2 + 2m_1l_1l_2 \cos \varphi_1 - 2m_3l_2l_3 \cos \varphi_1, \\ r_1 &= -k_1\varphi_1 - m_1l_1l_2\dot{\varphi}_2^2 \sin \varphi_1 + m_3l_2l_3\dot{\varphi}_2^2 \sin \varphi_1, \\ r_2 &= -Fl_1 \sin \varphi_1 - k_2\varphi_2 + m_1l_1l_2\dot{\varphi}_1(\dot{\varphi}_1 + 2\dot{\varphi}_2) \sin \varphi_1 - m_3l_2l_3\dot{\varphi}_1(\dot{\varphi}_1 + 2\dot{\varphi}_2) \sin \varphi_1. \end{aligned}$$

If $k_2 = 0$ then the right-hand side does not depend on variable φ_2 and the system of the third order for variables φ_1 , $\dot{\varphi}_1 = v_1$, and $\dot{\varphi}_2 = v_2$ can be considered independently

$$\begin{aligned} \dot{\varphi}_1 &= v_1, \\ \dot{v}_1 &= \left(r_1 - \frac{A_{12}}{A_{22}}r_2 \right) \left(A_{11} - \frac{A_{12}A_{21}}{A_{22}} \right)^{-1}, \\ \dot{v}_2 &= \left(r_2 - \frac{A_{21}}{A_{11}}r_1 \right) \left(A_{22} - \frac{A_{12}A_{21}}{A_{11}} \right)^{-1}. \end{aligned} \quad (5)$$

An important property of system (4) is the existence of a smooth invariant measure. Indeed, it is a system with circulatory forces, i.e. the Lagrange equations have the form

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{q}_i} - \frac{\partial L}{\partial q_i} = Q_i(q), \quad i = 1, 2.$$

Since the components of the generalized force depend only on the generalized coordinates and do not depend on the generalized momenta, then in the Hamiltonian coordinates this system has an invariant measure of constant density.

Let us briefly discuss various definitions of integrability which can be applied to systems (4) and (5). First, if system (4) is Hamiltonian (for instance, for $F = 0$), then it is natural to call the system integrable if the Liouville-Arnold theorem holds. Second, if this system is not Hamiltonian, we can consider integrability in the sense of the Jacobi theorem on the last multiplier (see, for instance, [9]). We say that system (4) is integrable if there exist two independent first integrals. Similarly, we say that system (5) is integrable if there exist two first integrals for this system. In this case the integrability follows from the classical theorem on the integrability for general ODE systems: a system in \mathbb{R}^n is integrable if it has $n - 1$ first integrals.

3 Cases $F = 0$ and $m_1 l_1 = m_3 l_3$

Let us consider two cases in which the dynamics of system (5) (for $k_2 = 0$) can be fully understood. First, if there is no follower force, then the system is Hamiltonian and there exist two first integrals: the energy integral $H = T + \Pi$ and the cyclic integral (the integral of angular momentum)

$$K = \frac{\partial T}{\partial \dot{\varphi}_2} = \frac{m_1}{2} (2\dot{\varphi}_2 l_2^2 + 2(\dot{\varphi}_1 + \dot{\varphi}_2) l_1^2 + 2l_1 l_2 \cos \varphi_1 (\dot{\varphi}_1 + 2\dot{\varphi}_2)) + m_2 \dot{\varphi}_2 l_2^2 + \frac{m_3}{2} (2\dot{\varphi}_2 l_2^2 + 2(\dot{\varphi}_1 + \dot{\varphi}_2) l_3^2 - 2l_2 l_3 (\dot{\varphi}_1 + 2\dot{\varphi}_2) \cos \varphi_1). \quad (6)$$

Therefore, the system is integrable in the sense of the Liouville-Arnold theorem, as a general system of ordinary differential equations (system (5) defined in a three-dimensional space has two first integrals), and in the sense of the Jacobi theorem on the last multiplier. The latter follows from the fact that the original system (4) has an invariant measure for any F . The density of this measure is constant if we rewrite our system in the Hamiltonian coordinates.

Another case in which system (5) is simplified and can be integrated in quadratures is when $m_1 l_1 = m_3 l_3$. This equality allows one to consider two separate independent systems:

$$\ddot{\varphi}_1 = \frac{F l_1 \sin \varphi_1 - k_1 \varphi_1}{(m_1 + m_2 + m_3) l_2^2}, \quad (7)$$

$$v_2(t) = v_2(0) - \int_0^t \ddot{\varphi}_1 + \frac{k_1}{m_1 l_1^2 + m_3 l_3^2} \varphi_1 dt.$$

This system is integrable by the Jacobi theorem: the first integral is the energy integral of the first equation, the phase volume is an invariant measure, if we present the system in the Hamiltonian form.

The periodic solutions that intersect the plane $\varphi_1 = 0$ in two points, are of particular interest for our considerations. We show that such solutions exist in the both considered cases (Fig. 2).




Figure 2: Examples of families of periodic solutions intersecting plane $\varphi_1 = 0$ in two points.

Proposition 1. For $F = 0$ and large k_1 there exists a two-parameter family of periodic solutions of system (5) which intersect the plane $\varphi_1 = 0$ transversely.

Proof. Find an expression for $\dot{\varphi}_2$ from (6):

$$\dot{\varphi}_2 = \frac{K - V(\varphi_1)\dot{\varphi}_1}{U(\varphi_1)},$$

where

$$\begin{aligned} V(\varphi_1) &= m_1 l_1^2 + m_3 l_3^2 + m_1 l_1 l_2 \cos \varphi_1 - m_3 l_2 l_3 \cos \varphi_1, \\ U(\varphi_1) &= m_1 l_1^2 + m_1 l_2^2 + m_2 l_2^2 + m_3 l_2^2 + 2m_1 l_1 l_2 \cos \varphi_1 - 2m_3 l_2 l_3 \cos \varphi_1. \end{aligned}$$

Substitute this expression for $\dot{\varphi}_2$ into the energy integral

$$\begin{aligned} H(\varphi_1, \dot{\varphi}_1) &= \frac{1}{2} l_2^2 (m_1 + m_2 + m_3) \left(\frac{K - V\dot{\varphi}_1}{U} \right)^2 + \frac{1}{2} (m_1 l_1^2 + m_3 l_3^2) \left(\frac{K + (U - V)\dot{\varphi}_1}{U} \right)^2 + \\ &\quad \left(\frac{K - V\dot{\varphi}_1}{U} \right) \left(\frac{K + (U - V)\dot{\varphi}_1}{U} \right) \cos \varphi_1 (m_1 l_1 l_2 - m_3 l_2 l_3) + \frac{k_1 \varphi_1^2}{2}. \end{aligned}$$

Let us prove that there is an extremum on the axis $\varphi_1 = 0$. Since $U'(0) = V'(0) = 0$, then for $\varphi_1 = 0$, we have

$$\frac{\partial H}{\partial \varphi_1} = 0.$$

From equation

$$\frac{\partial H}{\partial \dot{\varphi}_1} = 0$$

we obtain

$$\dot{\varphi}_1 = \frac{VKl_2^2(m_1 + m_2 + m_3) + K(V - U)(m_1 l_1^2 + m_3 l_3^2) + (2V - U)K(m_1 l_1 l_2 - m_3 l_2 l_3)}{V^2 l_2^2 (m_1 + m_2 + m_3) + (U - V)^2 (m_1 l_1^2 + m_3 l_3^2) - 2V(U - V)(m_1 l_1 l_2 - m_3 l_2 l_3)}.$$

It can be shown that the denominator of the fraction is strictly positive for all values of parameters. In particular, we obtain that the first minor of the Hessian matrix is positive. Non-diagonal elements of this matrix equal zero and the second diagonal element is positive provided that k_1 is sufficiently large.

Finally, we have that for all values of the energy integral which is close to the critical value of function H at the point of local minimum, the trajectory is periodic and intersects the plane $\varphi_1 = 0$ transversely. Moreover, this result holds for all values of first integral K .

□

Proposition 2. For $k_1 > 0$ system (7) has a two-parameter family of periodic solutions which intersect the plane $\varphi_1 = 0$ transversely.

Proof. The first equation of system (7) is the Lagrange equation with the following potential energy

$$\Pi = \frac{Fl_1 \cos \varphi_1 + \frac{1}{2} k_1 \varphi_1^2}{(m_1 + m_2 + m_3) l_2^2}.$$

If the value of total energy is sufficiently large, then the phase trajectories encircle the point of origin $\varphi_1 = 0$, $\dot{\varphi}_1 = 0$ and these trajectories are symmetric with respect to the coordinate axes. Therefore, $\varphi_1(t)$ is a periodic function and its mean value equals zero. We obtain that $v_2(t)$ is also a periodic function with the same period. □

Proposition 3. For $k_1 = 0$ and $F < 0$ system (7) a two-parameter family of periodic solutions which intersect the plane $\varphi_1 = 0$ transversely.

Proof. For the first equation of the system, point $\varphi_1 = 0$, $\dot{\varphi}_1 = 0$ is a local minimum. \square

Note that the Zielger pendulum can be considered as a Hamiltonian system for some values of the parameters [10]. However, for the case $k_2 = 0$ it follows from the conditions presented in [10] that $m_1l_1 = m_3l_3$ and $k_1 = 0$. Therefore, from (4) we obtain a trivial system, which is less general than (7).

4 Integrability

For the case when $k_2 = 0$ it is natural to call the original system integrable if the reduced system (5) has two first integrals. In the general case, for which it is also allowed that $k_2 \neq 0$, it is also natural to call the system with two first integrals integrable. For any k_2 we have a system defined on $\mathbb{T}^2 \times \mathbb{R}^2$ and this system has an invariant measure. Therefore, this system is integrable if it has two first integrals, similarly to Hamiltonian systems with two degrees of freedom and to the case $k_2 = 0$.

Let us prove that system (5) for $k_2 = 0$ remains integrable for $|F|$ or $|m_1l_1 - m_3l_3|$ sufficiently small. The key step in the proof is given by the following lemma.

Lemma 1. Let functions $\varphi_1(t)$, $v_1(t)$, $v_2(t)$ be a solution of system (5). Then functions $-\varphi_1(-t)$, $v_1(-t)$, $v_2(-t)$ are also a solution.

Proof. The proof follows from the substitution of given functions into the system. Terms A_{11} , A_{12} , A_{22} do not change after substitution $\varphi_1 \mapsto -\varphi_1$. Terms r_1 and r_2 change their signs. At the same time, the left-hand side also changes its sign. \square

From the lemma we obtain the following

Proposition 4. Let $\varphi_1(t)$, $v_1(t)$, $v_2(t)$ be a solution which intersects the plane $\varphi_1 = 0$ in two points. Then this solution is periodic.

Proof. A periodic solution can be obtained from the segment between the two points using the symmetry with respect to the plane $\varphi_1 = 0$. \square

Finally, we obtain the following three results on the existence of families of periodic solutions for system (5).

Proposition 5. There exists K_1 such that for any $k_1 > K_1$ there exists $\varepsilon = \varepsilon(k_1)$ and for all $|F| < \varepsilon$ system (5) has a two-parameter family of periodic solutions.

Proposition 6. Let $k_1 > 0$, then there exists $\varepsilon = \varepsilon(k_1)$, such that for all $|m_1l_1 - m_3l_3| < \varepsilon$ system (5) has a two-parameter family of periodic solutions.

Proposition 7. Let $k_1 = 0$ and $F < 0$, then there exists $\varepsilon = \varepsilon(k_1)$ such that for all $|m_1l_1 - m_3l_3| < \varepsilon$ system (5) has a two-parameter family of periodic solutions.

In all three propositions the proofs are based on the existence of families of solutions which intersect the plane $\varphi_1 = 0$ transversely for $F = 0$ and $m_1l_1 - m_3l_3 = 0$. From Proposition 4 we obtain that these families of solutions exist for small $|F|$ and $|m_1l_1 - m_3l_3|$.

From the existence of a two-parametric family of periodic solutions it follows that locally (in a neighborhood of these solutions) the system is integrable: there are two local first integrals, which can be defined in various ways.




Figure 3: Non-periodic (a) and periodic (b) solutions of the system for $F \neq 0$ and $m_1 l_1 \neq m_3 l_3$. The parameters of the system are the same in the both cases, initial conditions vary.

A natural question of whether or not it is possible to continue these two first integrals globally to the whole three-dimensional space with coordinates φ_1, v_1, v_2 is expected to have a negative answer. It follows from the numerical results which show that in a general case for some initial conditions there exist non-periodic trajectories lying on a two-dimensional surface (Fig. 3). Therefore, there are less than two independent first integrals.

The mechanism for the appearance of such trajectories is as follows. Let for $F = 0$ we have a family of periodic solutions, if $|F|$ is sufficiently small, then all these solutions still intersect the plane $\varphi_1 = 0$ in two points. Therefore, we obtain a family of periodic solutions. When F becomes sufficiently large, some solutions of this family do not intersect the plane $\varphi_1 = 0$ and may become non-periodic. This breaks the integrability (or super-integrability) of the system. In Fig. 4 a family of solutions starting at the same point is shown. The only parameter of system that varies from solution to solution is the magnitude of force F . For small F all solutions intersect the plane $\varphi_1 = 0$, but a non-periodic solution appears when F becomes larger than some critical value. This solution does not intersect the considered plane.

Note that the above conditions for the existence of periodic solutions for system (5) (for $F = 0$) and (7) can be weakened and the families of periodic solutions are much wider. We present simple conditions that can be obtained without cumbersome calculations and detailed estimates.

For the integrability in a four-dimensional space with an invariant measure, the most natural definition of the integrability stems from the Jacobi theorem: the system is integrable if it has two independent first integrals.

For any periodic solution in the reduced space with coordinates φ_1, v_1, v_2 , from the independent equation

$$\dot{\varphi}_2 = v_2,$$




Figure 4: The mechanism for the appearance of a non-periodic solution.

it follows that in the four-dimensional space $\varphi_1, \varphi_2, v_1, v_2$ this solution lies on a two-dimensional surface (the angular variables are considered modulo 2π). If the solution is not periodic in the reduced space φ_1, v_1, v_2 , then its projection from the four-dimensional space into a three-dimensional space with, for instance, coordinates $\varphi_1, \varphi_2, v_1$ can densely fill three-dimensional domains.

In the first case we have two first integrals. In the latter case, there may be only one or no first integrals. As an illustration (Fig. 5) we present the section of projections of two solutions into the coordinate space $\varphi_1, \varphi_2, v_1$ by the plane $\varphi_2 = \pi$: in the first case the solution is periodic in the space with coordinates φ_1, v_1, v_2 , in the second case the solution is not periodic in this space.




Figure 5: Regular (a) and irregular (b) behavior of solutions. The parameters of the system are the same in the both cases, initial conditions vary.

In the conclusion of the section, we consider a sufficient generalization of system (5) which is integrable. Indeed, the key property which is used in the proves of integrability is the symmetry of the original

system. Hence, if we consider perturbations relatively small and satisfying the properties of symmetry. For instance, let us consider the following system (here we assume that $F = 0$)

$$\begin{aligned}\dot{\varphi}_1 &= v_1, \\ \dot{v}_1 &= \left(r_1 - \frac{A_{12}}{A_{22}}r_2\right) \left(A_{11} - \frac{A_{12}A_{21}}{A_{22}}\right)^{-1} + f_1(\varphi_1, v_1, v_2), \\ \dot{v}_2 &= \left(r_2 - \frac{A_{21}}{A_{11}}r_1\right) \left(A_{22} - \frac{A_{12}A_{21}}{A_{11}}\right)^{-1} + f_2(\varphi_1, v_1, v_2),\end{aligned}\tag{8}$$

where $f_i(-\varphi_1, v_1, v_2) = -f_i(\varphi_1, v_1, v_2)$, $i = 1, 2$ are sufficiently small. Then this system is integrable in the above sense. For instance, a family of periodic solutions exists for system (8) if we put $f_1 = -\alpha\varphi_1 \sin(v_2)$, $f_2 = \alpha\varphi_1 \sin(v_1)$, for sufficiently small values of α . At the same time, there may exist asymptotically stable equilibria in this system. Therefore, there is no smooth invariant measure in this system and there cannot exist two first integrals, which are defined globally in the whole phase space.

5 Case $k_2 \neq 0$

In the above considerations we assume that the stiffness of the spring located in the fixed point O equals zero. This allows us to study the reduced system with the three-dimensional phase space $\mathbb{S} \times \mathbb{R}^2$.

When $k_2 \neq 0$ we have to consider the system defined on $\mathbb{T}^2 \times \mathbb{R}^2$, and this system has an invariant measure. Based on the results from KAM theory for the reversible systems (for details, please, see [11]), one can expect the co-existence of regions with regular and chaotic behavior of the solutions.

A detailed numerical study of this case is out of the scope of this paper. We only present some results showing that for the Ziegler pendulum the regular dynamics can co-exists with the irregular behavior of trajectories.

After the change of time variable

$$t \mapsto t \cdot (A_{11}A_{22} - A_{12}A_{21})$$

system (5) can be presented in the following form (here we also the equation for $\dot{\varphi}_2$)

$$\begin{aligned}\dot{\varphi}_1 &= v_1(A_{11}A_{22} - A_{12}A_{21}), \\ \dot{\varphi}_2 &= v_2(A_{11}A_{22} - A_{12}A_{21}), \\ \dot{v}_1 &= A_{22}r_1 - A_{12}r_2, \\ \dot{v}_2 &= A_{11}r_2 - A_{12}r_1,\end{aligned}\tag{9}$$

where we use the same notations as above.

One of the most popular methods of numerical study of chaos is *mLCE* (a simple method based on the estimation of the maximum Lyapunov characteristic exponent). A detailed explanation of this approach can be found in [12]–[14].

For a given solution, the Lyapunov characteristic exponents can be calculated as follows

$$\chi(X(t), x) = \limsup_{t \rightarrow +\infty} \frac{1}{t} \ln \frac{\|X(t)x(0)\|}{\|x(0)\|},$$

where $X(t)$ is the fundamental matrix of the linearized system for system (9). The system is linearized along the considered solution. The evolution of vector $x(t)$ is described by the following linear system

$$\dot{x} = A(t)x.$$

For our system, we have

$$A(t) = \begin{pmatrix} v_1 A_{11} \frac{\partial A_{22}}{\partial \varphi_1} - 2v_1 A_{12} \frac{\partial A_{12}}{\partial \varphi_1} & 0 & A_{11} A_{22} - A_{12} A_{21} & 0 \\ v_2 A_{11} \frac{\partial A_{22}}{\partial \varphi_1} - 2v_2 A_{12} \frac{\partial A_{12}}{\partial \varphi_1} & 0 & 0 & A_{11} A_{22} - A_{12} A_{21} \\ \frac{\partial A_{22}}{\partial \varphi_1} r_1 + A_{22} \frac{\partial r_1}{\partial \varphi_1} - \frac{\partial A_{12}}{\partial \varphi_1} r_2 - A_{12} \frac{\partial r_2}{\partial \varphi_1} & -A_{12} \frac{\partial r_2}{\partial \varphi_2} & A_{22} \frac{\partial r_1}{\partial v_1} - A_{12} \frac{\partial r_2}{\partial v_1} & A_{22} \frac{\partial r_1}{\partial v_1} - A_{12} \frac{\partial r_2}{\partial v_1} \\ \frac{\partial A_{22}}{\partial \varphi_1} r_1 + A_{22} \frac{\partial r_1}{\partial \varphi_1} - \frac{\partial A_{12}}{\partial \varphi_1} r_2 - A_{12} \frac{\partial r_2}{\partial \varphi_1} & A_{11} \frac{\partial r_2}{\partial \varphi_2} & A_{22} \frac{\partial r_1}{\partial v_1} - A_{12} \frac{\partial r_2}{\partial v_1} & A_{22} \frac{\partial r_1}{\partial v_1} - A_{12} \frac{\partial r_2}{\partial v_1} \end{pmatrix}$$

In practice, the following equation is used to obtain the maximum Lyapunov characteristic exponent

$$\chi(X(t)) = \lim_{t \rightarrow +\infty} \frac{1}{t} \ln \frac{\|X(t)x(0)\|}{\|x(0)\|}.$$

It is known that for almost all initial vectors $x(0)$ and for a wide range of matrices $X(t)$ this equation gives the maximum Lyapunov characteristic exponent.

Let us consider the system with the following values of mechanical parameters:

$$\begin{aligned} m_1 &= 1, & l_1 &= 1, & k_1 &= 1, \\ m_2 &= 1, & l_2 &= 1, & k_2 &= 1, \\ m_3 &= 3/2, & l_3 &= 1, & F &= 2, \end{aligned}$$

and two solutions with initial conditions $(\pi, 0, 1/10, 1/10)$ and $(\pi, 0, 20, 20)$ (the order of the coordinates is as follows: $\varphi_1, \varphi_2, v_1, v_2$). Below we present the results of calculation of $mLCE$ for these solutions (Fig. 5). It can be seen that in one case the value of $mLCE$ always remains positive (a property of a chaotic trajectory) and for another solution this value tends to zero.




Figure 6: Zero and positive maximum Lyapunov characteristic exponents.

Regularity and irregularity of solutions can be also seen in the projection of these solutions into the three-dimensional space (φ_1, v_1, v_2) . In the case of regular behavior with zero $mLCE$, the solution lies on a two-dimensional surface. For the trajectory with strictly positive $mLCE$, the closure of the projection of the trajectory covers three-dimensional domains (Fig. 6).

We also present ‘the Poincare sections’ of a family of solutions, from which it is also can be seen that the regular dynamics co-exists with the chaotic dynamics. We consider ten initial conditions $(\pi, 0, 1/10, (n+1)/10)$, $0 \leq n < 10$, $n \in \mathbb{Z}$. For each corresponding solution we consider its projection




Figure 7: Regular (a) and irregular (b) behavior of the solutions. The parameters of the system are the same in the both cases, initial conditions vary.




Figure 8: Regular and chaotic solutions.

into the three-dimensional space (φ_1, v_1, v_2) , and then the section of this projection by the plane $v_1 = 0$.

The obtained sets of points are not the Poincaré section in the usual sense since we consider the section of the projections of solutions. In particular, it is possible that there exist more than one solution which intersects the plane $v_1 = 0$ in a given point. The resulting sections are presented in Fig. 7. It can be seen that for relatively small initial velocity the solutions are regular (lie on a two-dimensional surface) and become chaotic as velocity becomes larger.

6 Conclusion

The dynamics of a planar double pendulum in the presence of a follower force is considered. The system is studied in the case when the governing system of equation can be reduced to a system in a three-dimensional space, and in the general case when the system is considered in a four-dimensional space. The following results are obtained:

1. It is proved that, in the case of zero stiffness of the spring located in the point of suspension, there exist two-parametric families of periodic solutions (for some values of the mechanical parameters).
2. It is shown numerically that the periodic solutions do not fill the whole phase space. In particular, some solutions lie on two-dimensional surfaces and the system does not have two independent first integrals defined globally in the space with coordinates φ_1, v_1, v_2 . Moreover, the system is not integrable in the sense of the Jacobi theorem if we consider the dynamics in the original (not reduced) phase space $\mathbb{T}^2 \times \mathbb{R}^2$.
3. It is shown that in the general case of the Ziegler pendulum when $k_2 \neq 0$ the regular and the chaotic dynamics co-exist in different regions of the phase space.

Let us briefly discuss the above results and outline some possible directions of further research. First, let us note that we make the conclusion on the integrability of the system without explicitly presenting the required set of the first integrals. We only prove that there exist periodic solutions for the system and, therefore, the system can be locally integrable. Moreover, there is a strong evidence that these local first integrals cannot be continued to the whole phase space. It might be interesting to understand this result deeper: to the best of our knowledge there are no other mechanical systems that are integrable in one part of the phase space and not integrable, or integrable in another sense, in the rest of the phase space.

An important property of the circulatory system (5), which cannot be seen in non-integrable Hamiltonian systems, is that there are regions in the phase space of system (5) such that *all* invariant tori are not destroyed when we add a perturbation to the system. For Hamiltonian systems it is typical that resonant tori are destroyed by perturbations. Since resonant tori are dense in the phase space, we can usually conclude that the system becomes non-integrable.

At the same time, the co-existence of the regular and the chaotic behavior of solutions is typical for Hamiltonian systems (see, for instance, [15], [16]). Nevertheless, we have shown that chaotic trajectories in circulatory systems (the Ziegler pendulum for $k_2 = 0$) may appear in a different manner comparing to Hamiltonian systems. The existence of chaotic regions in analytic Hamiltonian systems means that the system is not integrable, yet the existence of *local* set of independent first integrals is still possible if we have irregular behavior of solutions in a circulatory system.

Acknowledgments

The author is grateful to Valery Kozlov for his invaluable advice and useful suggestions.

The work was supported by the Russian Science Foundation (Project No. 19-71-30012).

References

- [1] H. Ziegler, “Die stabilitätskriterien der elastomechanik,” *Ingenieur-Archiv*, vol. 20, no. 1, pp. 49–56, 1952.
- [2] G. Herrmann and R. Bungay, “On the stability of elastic systems subjected to nonconservative forces,” 1964.
- [3] V. Bolotin and N. Zhinzher, “Effects of damping on stability of elastic systems subjected to nonconservative forces,” *International Journal of Solids and Structures*, vol. 5, no. 9, pp. 965–989, 1969.
- [4] O. N. Kirillov, “Nonconservative stability problems of modern physics,” in *Nonconservative Stability Problems of Modern Physics*, de Gruyter, 2013.
- [5] T. Shinbrot, C. Grebogi, J. Wisdom, and J. A. Yorke, “Chaos in a double pendulum,” *American Journal of Physics*, vol. 60, no. 6, pp. 491–499, 1992.
- [6] T. Stachowiak and T. Okada, “A numerical analysis of chaos in the double pendulum,” *Chaos, Solitons & Fractals*, vol. 29, no. 2, pp. 417–422, 2006.
- [7] H. R. Dullin, “Melnikov’s method applied to the double pendulum,” *Zeitschrift für Physik B Condensed Matter*, vol. 93, no. 4, pp. 521–528, 1994.
- [8] V. V. Kozlov, “On the integrability of circulatory systems,” *Regular and Chaotic Dynamics*, vol. 27, no. 1, pp. 11–17, 2022.
- [9] —, “The Euler-Jacobi-Lie integrability theorem,” *Regular and Chaotic Dynamics*, vol. 18, no. 4, pp. 329–343, 2013.
- [10] —, “Integrals of circulatory systems which are quadratic in momenta,” *Regular and Chaotic Dynamics*, vol. 26, no. 6, pp. 647–657, 2021.
- [11] M. B. Sevryuk, *Reversible systems*. Springer, 2006, vol. 1211.
- [12] G. Benettin, L. Galgani, A. Giorgilli, and J.-M. Strelcyn, “Lyapunov characteristic exponents for smooth dynamical systems and for hamiltonian systems; a method for computing all of them. part 1: Theory,” *Meccanica*, vol. 15, no. 1, pp. 9–20, 1980.
- [13] A. J. Lichtenberg and M. A. Leiberman, *Regular and chaotic dynamics*. Springer Science & Business Media, 2013, vol. 38.
- [14] C. Skokos, “The lyapunov characteristic exponents and their computation,” in *Dynamics of Small Solar System Bodies and Exoplanets*, Springer, 2010, pp. 63–135.
- [15] V. I. Arnold, V. V. Kozlov, and A. I. Neishtadt, *Mathematical aspects of classical and celestial mechanics*. Springer Science & Business Media, 2007, vol. 3.
- [16] V. V. Kozlov, *Symmetries, topology and resonances in Hamiltonian mechanics*. Springer Science & Business Media, 2012, vol. 31.