JORDAN CONSTANT FOR CREMONA GROUP OF RANK 2 OVER A FINITE FIELD

ANASTASIA V.VIKULOVA

ABSTRACT. In this paper we find the exact value of the Jordan constant for Cremona group of rank 2 over all finite fields. During the proof we construct a cubic surface over \mathbb{F}_2 with a regular action of the group S₆ which is the maximal automorphism group of cubic surfaces over \mathbb{F}_2 . Moreover, we prove the uniqueness up to isomorphism of such a cubic surface.

1. INTRODUCTION

The Cremona group $\operatorname{Cr}_n(\mathbf{F})$ of rank n is a group of birational automorphisms of the projective space \mathbb{P}^n over \mathbf{F} . Despite the fact that this group appears naturally, its structure is extremely complicated for $n \ge 2$. Moreover, the description of finite subgroups seems very hard to obtain. Even in the first non-trivial case of rank 2 there is a description of conjugacy classes of finite subgroups only over \mathbb{C} (see [7]). Nevertheless, we can understand some properties of finite subgroups of Cremona groups.

Definition 1 ([10, Definition 2.1]). A group G is called *Jordan* if there is a constant J such that any finite subgroup of the group G has a normal abelian subgroup of index at most J. The minimal such constant J is called the *Jordan constant* of the group G and is denoted by J(G).

J.-P. Serre proved in [12, Theorem 5.3] that the rank 2 Cremona group $\operatorname{Cr}_2(\mathbf{F})$ over the field \mathbf{F} of characteristic zero is Jordan. However, it is not true for algebraically closed field \mathbf{F} of characteristic p > 0, because the group $\operatorname{Cr}_2(\mathbf{F})$ contains simple subgroup $\operatorname{PSL}_2(\mathbb{F}_{p^n})$ whose order grows with n. In the article [11] Yu. Prokhorov and C. Shramov proved that the Cremona group of rank 2 over a finite field is Jordan. Moreover, they obtained the following inequalities for Jordan constants.

Theorem 1 ([11, Theorem 1.2]). Let $J(\operatorname{Cr}_2(\mathbb{F}_q))$ be the Jordan constant for the Cremona group $\operatorname{Cr}_2(\mathbb{F}_q)$. Then we have

$J(\operatorname{Cr}_2(\mathbb{F}_q)) = q^3(q^2 - 1)(q^3 - 1),$	<i>if</i> $q \notin \{2, 4, 8\};$
$J(\operatorname{Cr}_2(\mathbb{F}_q)) \leqslant 696729600,$	<i>if</i> $q \in \{2, 4, 8\}$ <i>.</i>

Our goal is to find the exact value of the Jordan constant for the Cremona group $\operatorname{Cr}_2(\mathbb{F}_q)$ for $q \in \{2, 4, 8\}$, where \mathbb{F}_q is the field with q elements. For this it is enough to study the groups of biregular automorphisms of del Pezzo surfaces and conic bundles over \mathbb{F}_q , because every finite subgroup G of $\operatorname{Cr}_2(\mathbb{F}_q)$ is regularized on a G-minimal model of a rational surface. According to [11, Corollary 5.3]

This work is supported by the Russian Science Foundation under grant no.18-11-00121.

and [11, Lemma 6.1], the Jordan constant for groups acting biregularly on conic bundles over \mathbb{P}^1 and del Pezzo surfaces of degrees $4 \leq d \leq 9$ and d = 2 is at most $q^3(q^2-1)(q^3-1)$ which is equal to the order of the automorphism group of \mathbb{P}^2 over \mathbb{F}_q . Therefore, it is enough to study the group of regular automorphisms of del Pezzo surface of degrees 1 and 3 and their normal abelian subgroups.

In this paper we prove the following theorem.

Theorem 2. The Jordan constant $J(\operatorname{Cr}_2(\mathbb{F}_q))$ of the Cremona group $\operatorname{Cr}_2(\mathbb{F}_q)$ is equal to

$$J(\operatorname{Cr}_2(\mathbb{F}_q)) = \begin{cases} 16\,482\,816, & \text{if } q = 8; \\ 60\,480, & \text{if } q = 4; \\ 720, & \text{if } q = 2. \end{cases}$$

Corollary 1. The Jordan constant $J(\operatorname{Cr}_2(\mathbb{F}_q))$ for the Cremona group $\operatorname{Cr}_2(\mathbb{F}_q)$ is equal to

$$J(\operatorname{Cr}_2(\mathbb{F}_q)) = \begin{cases} |\operatorname{PGL}_3(\mathbb{F}_q)|, & \text{if } q \neq 2; \\ |\operatorname{S}_6| > 168 = |\operatorname{PGL}_3(\mathbb{F}_2)|, & \text{if } q = 2. \end{cases}$$

As a complement to the proof of Theorem 2 we prove the following fact about the automorphisms of cubic surface over \mathbb{F}_2 .

Theorem 3. Let S be a smooth cubic surface in \mathbb{P}^3 over \mathbb{F}_2 . Then

$$|\operatorname{Aut}(S)| \leq 720.$$

If the equality holds then we have $\operatorname{Aut}(S) \simeq S_6$. Moreover, the cubic surface with such automorphism group is unique up to isomorphism.

Acknowledgment. The author wish to warmly thank C. A. Shramov for suggesting this problem, a lot of discussion and constant attention to this work. Also the author would like to thank A. S. Trepalin for the interesting conversations about cubic surfaces and Yu. G. Prokhorov for important remarks.

2. Smooth cubic surfaces

It is well-known that the automorphism group of a smooth cubic surface in \mathbb{P}^3 is a subgroup in the Weyl group $W(\mathcal{E}_6)$ (see, for instance, [5, Corollary 8.2.40]). Recall that

$$|W(E_6)| = 2^7 \cdot 3^4 \cdot 5 = 51\,840.$$

According to the classification of the automorphism groups of smooth cubic surfaces over an algebraically closed field (see [6, Table 1]), we get that the largest automorphism group of a cubic surface over the algebraically closed field $\overline{\mathbb{F}}_2$ is $\mathrm{PSU}_4(\mathbb{F}_2)$. This group is the automorphism group of the Fermat cubic $S \subset \mathbb{P}^3$ (see [6, Table 8]) which is defined by the equation

$$x^3 + y^3 + z^3 + t^3 = 0.$$

Recall that $|PSU_4(\mathbb{F}_2)| = 2^6 \cdot 3^4 \cdot 5 = 25\,920.$

However, the situation changes if the base field is \mathbb{F}_2 . We prove that the largest automorphism group of a smooth cubic surface over \mathbb{F}_2 is of order 720.

Theorem 4. Let S be a smooth cubic surface in \mathbb{P}^3 over \mathbb{F}_2 . Then we have

 $|\operatorname{Aut}(S)| \leq 720.$

Moreover, if the order of $\operatorname{Aut}(S)$ is equal to 720, then we have $\operatorname{Aut}(S) \simeq S_6$.

Proof. The group $\operatorname{Aut}(S)$ is contained in the group $\operatorname{PGL}_4(\mathbb{F}_2)$, because the embedding $S \subset \mathbb{P}^3$ is given by the linear system $|-K_S|$ which is invariant under the automorphism group $\operatorname{Aut}(S)$. On the other hand, the group $\operatorname{Aut}(S)$ is contained in the Weyl group $W(\operatorname{E}_6)$. Suppose that there is a cubic surface S such that $720 < |\operatorname{Aut}(S)|$.

The maximal subgroup in the group $PGL_4(\mathbb{F}_2)$ is isomorphic to one of the following groups (see the list of the maximal subgroups of $PGL_4(\mathbb{F}_2)$ in [2, p. 22]):

$$A_7, \ (\mathbb{Z}/2\mathbb{Z})^3 \rtimes PGL_3(\mathbb{F}_2), \ S_6, \ (\mathbb{Z}/2\mathbb{Z})^4 \rtimes (S_3 \times S_3), \ (A_5 \times \mathbb{Z}/3\mathbb{Z}) \rtimes \mathbb{Z}/2\mathbb{Z}.$$

Therefore, we get that $\operatorname{Aut}(S)$ is either isomorphic to $\operatorname{PGL}_4(\mathbb{F}_2)$, or to a subgroup in A_7 , or to a subgroup in $(\mathbb{Z}/2\mathbb{Z})^3 \rtimes \operatorname{PGL}_3(\mathbb{F}_2)$.

The first case is impossible, as $|PGL_4(\mathbb{F}_2)|$ does not divide $|W(E_6)|$. Assume that the second case holds, i.e. the order of the group Aut(S) divides both the order of the group A_7 , and the order of the group $W(E_6)$, that is

$$|\operatorname{Aut}(S)| \leq \operatorname{gcd}(|\operatorname{A}_7|, |W(\operatorname{E}_6)|) = 2^3 \cdot 3^2 \cdot 5 = 360 < 720.$$

Thus, this case is also impossible.

Finally, assume that the third case holds. In other words, we obtain

 $|\operatorname{Aut}(S)| \leq \gcd(|(\mathbb{Z}/2\mathbb{Z})^3 \rtimes \operatorname{PGL}_3(\mathbb{F}_2)|, |W(\operatorname{E}_6)|) = 2^6 \cdot 3 = 192 < 720.$

Therefore, the order of the automorphism group of a smooth cubic surface is at most 720.

Using [2, p. 22], we get that the only subgroup of order 720 in $PGL_4(\mathbb{F}_2)$ is the group S_6 , because it is the maximal subgroup and the other maximal subgroups do not contain subgroups of order 720.

Let us construct the cubic surface with automorphism group S_6 .

Example 1. Let us consider the cubic surface $S \subset \mathbb{P}^3$ over \mathbb{F}_2 which is defined by the equation

(1)
$$x^2t + y^2z + z^2y + t^2x = 0.$$

It is obvious that this equation defines a smooth cubic surface. Moreover, it passes through every point of \mathbb{P}^3 . We show that $\operatorname{Aut}(S) \simeq S_6$.

Let us consider the matrix

$$\Omega = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}.$$

This matrix corresponds to a skew-symmetric bilinear form. Consider the group which fixes this matrix. It consists of the elements $g \in PGL_4(\mathbb{F}_2)$ such that

$$g^T \Omega g = \Omega.$$

This group is isomorphic to $PSp_4(\mathbb{F}_2)$.

Note that the left-hand side of (1) is equal to

$$(x^2, y^2, z^2, t^2)^T \Omega(x, y, z, t).$$

So the group $PSp_4(\mathbb{F}_2)$ is contained in the automorphism group of S. There is a well-known isomorphism $PSp_4(\mathbb{F}_2) \simeq S_6$ (see, for instance, [4, §5]). Thus, we get the inclusion $S_6 \subset Aut(S)$. But by Theorem 4 the group S_6 is the maximal possible automorphism group of a smooth cubic surface over \mathbb{F}_2 . Therefore, we have the isomorphism $Aut(S) \simeq S_6$.

Note that the cubic surface (1) is rational. Indeed, there are two skew lines l_1 and l_2 defined by the equations x = y = 0 and z = t = 0, respectively. This gives us a birational isomorphism $\mathbb{P}^1 \times \mathbb{P}^1 \dashrightarrow S$.

Remark 1. In the article [9] by the computer calculations it was shown that the automorphism group of the cubic surface (1) is of order 720. However, the structure of the automorphism group was not studied.

3. Del Pezzo surfaces of degree 1

In this section we consider smooth del Pezzo surfaces of degree 1.

Proposition 1. Let S be a smooth del Pezzo surface of degree 1 over \mathbb{F}_q . Then the order of the group $\operatorname{Aut}(S)$ satisfies the inequality

$$\operatorname{Aut}(S) \leq 2q^4(q-1)^2(q+1) < q^3(q^3-1)(q^2-1).$$

Proof. A degree 2 regular map

(2)
$$\phi_{|-2K_S|}: S \to \mathbb{P}(1,1,2)$$

defined by the linear system $|-2K_S|$ gives us the following exact sequence:

$$1 \to G \to \operatorname{Aut}(S) \to \operatorname{Aut}(\mathbb{P}(1,1,2)),$$

where $|G| \leq 2$. Let us find the automorphism group Aut($\mathbb{P}(1, 1, 2)$). It is obvious (see [1, §8.3]) that the automorphisms of $\mathbb{P}(1, 1, 2)$ can be written in the following way:

$$[x:y:z] \mapsto [ax+by:cx+dy:ez+f(x,y)],$$

where the matrix

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

is non-degenerate, $e \in \mathbb{F}_q^*$, and f(x, y) is a homogeneous polynomial of degree 2. So we obtain

$$\operatorname{Aut}(\mathbb{P}(1,1,2)) \simeq \frac{\left(\operatorname{GL}_2(\mathbb{F}_q) \times \mathbb{F}_q^*\right) \ltimes \left(\mathbb{F}_q\right)^3}{\mathbb{F}_q^*}.$$

It follows that the order of Aut(S) satisfies the inequality

$$|\operatorname{Aut}(S)| \leq 2 \cdot |\operatorname{Aut}(\mathbb{P}(1,1,2))| = 2q^4(q-1)^2(q+1) < q^3(q^3-1)(q^2-1).$$

Remark 2. It can be shown that the morphism (2) is separable (see, for example, [8, Proposition 1.2]). Therefore, in the proof of Proposition 1 the group G is isomorphic to $\mathbb{Z}/2\mathbb{Z}$.

4. Proof of Theorem 2

In this section we prove Theorem 2. First of all, let us remind a result from the article [11]:

Proposition 2 ([11, Corollary 5.3]). Let S be a smooth del Pezzo surface of degree not equal to 1 or 3 over \mathbb{F}_q . Then the group $\operatorname{Aut}(S)$ contains a normal abelian subgroups of index at most $q^3(q^2-1)(q^3-1)$.

Now we prove our main theorem.

Proof of Theorem 2. As it was mentioned above, any finite subgroup G of the Cremona group $\operatorname{Cr}_2(\mathbb{F}_q)$ is regularized on a G-minimal model of a rational surface. Thus, for all automorphism groups of del Pezzo surfaces and conic bundles over \mathbb{P}^1 it is enough to find the minimal index of a normal abelian subgroup. The maximal among such numbers is the required Jordan constant.

According to Proposition 2, a normal abelian subgroup in the automorphism group of a del Pezzo surface of degree not equal to 1 or 3 over \mathbb{F}_q has index at most $q^3(q^2-1)(q^3-1)$. And according to [11, Lemma 6.1], a normal abelian subgroup in the automorphism group of a conic bundle over \mathbb{P}^1 also has index at most $q^3(q^2-1)(q^3-1)$. By Proposition 1 the automorphism group of a smooth del Pezzo surface of degree 1 over \mathbb{F}_q is of order at most $q^3(q^2-1)(q^3-1)$.

It is enough to find the maximum of all minimal indexes of normal abelian subgroups in every automorphism group for smooth cubic surfaces. If q = 4 or 8 then

$$|W(\mathcal{E}_6)| = 51\,840 < 60\,480 = |\mathrm{PGL}_3(\mathbb{F}_4)| \leqslant |\mathrm{PGL}_3(\mathbb{F}_q)|,$$

therefore, the order of the automorphism group of a smooth cubic surface is less than $q^3(q^2-1)(q^3-1)$. And if q=2 then by Theorem 4 the order of the automorphism group of a smooth cubic surface is at most

$$720 > 168 = q^3(q^2 - 1)(q^3 - 1).$$

All these values, i.e. 720 for \mathbb{F}_2 and $q^3(q^2 - 1)(q^3 - 1)$ for \mathbb{F}_4 and \mathbb{F}_8 are realized as orders of the automorphism groups of some rational surfaces. Indeed, for q = 4 and 8 these values are attained on the automorphism group of \mathbb{P}^2 . This group is isomorphic to $\mathrm{PGL}_3(\mathbb{F}_q)$ and it does not have non-trivial normal abelian subgroups (see [11, Lemma 2.4]). For q = 2 the value of the Jordan constant is attained on the rational cubic surface with the automorphism group isomorphic to S_6 (see Example 1). This group does not have non-trivial normal abelian subgroups either.

Appendix A. Uniqueness of the cubic surface with the automorphism group S_6 over \mathbb{F}_2

In this section we prove Theorem 3.

Lemma 1. Let S be a smooth cubic surface in the projective space \mathbb{P}^3 over \mathbb{F}_2 , such that it passes through all 15 points in \mathbb{P}^3 . Then S is isomorphic to a cubic surface (1) and its automorphism group is S_6 .

Proof. Note that the cubic surface passing through all points in \mathbb{P}^3 can be defined by the following equation:

 $a_1xy(x+y) + a_2xz(x+z) + a_3xt(x+t) + a_4yz(y+z) + a_5yt(y+t) + a_6zt(z+t) = 0,$

where $a_i \in \mathbb{F}_2$. Thus, there are 63 such cubic surface, not necessarily smooth.

Let us consider the action of the group $\operatorname{PGL}_4(\mathbb{F}_2)$ on smooth cubic surfaces in \mathbb{P}^3 over \mathbb{F}_2 , which pass through 15 points. Denote by $\operatorname{Orb}(S)$ the orbit of S under the action of S₆. We have

$$\frac{|\operatorname{PGL}_4(\mathbb{F}_2)|}{|\operatorname{Aut}(S)|} = |\operatorname{Orb}(S)|.$$

Let us find the number of elements in $\operatorname{Orb}(S)$. Consider a singular cubic surface which is a union of three hyperplanes in \mathbb{P}^3 passing through one line. It is obvious that there are 15 points on such a cubic surface. It is not hard to see that there are exactly 35 lines in \mathbb{P}^3 over \mathbb{F}_2 . It follows that there are at least 35 such singular cubic surfaces. Thus, there are at most 28 smooth cubic surfaces passing through 15 points. Therefore, we obtain $|\operatorname{Orb}(S)| \leq 28$. So we have the inequality

$$|\operatorname{Aut}(S)| \ge 720$$

According to Theorem 4, this inequality holds true if and only if $\operatorname{Aut}(S) = S_6$ and $|\operatorname{Orb}(S)| = 28$. In other words, all smooth cubic surfaces passing through 15 points are isomorphic to each other. As the cubic surface of the form (1) passes through 15 points, we get that S is isomorphic to the cubic surface defined by the equation (1).

Lemma 2. Assume that the group S_6 is the automorphism group of a smooth cubic surface S over \mathbb{F}_2 . Then S is isomorphic to a cubic surface defined by the equation (1).

Proof. Note that according to the Chevalley–Warning theorem, there is a point on the cubic surface S. Denote by p one of such points. The action of the group S_6 on S defines its action on \mathbb{P}^3 . Assume that the length of the orbit of the point p is $l \neq 5, 10, 15$. So the stabilizer of every point in the orbit is a subgroup of S_6 of index l. Moreover, the stabilizer contains a subgroup $\mathbb{Z}/5\mathbb{Z}$, because l and 5 are coprime. So the group $\mathbb{Z}/5\mathbb{Z}$ acts non-trivially on the tangent space to \mathbb{P}^3 at the point p (see, for instance, [3, Theorem 3.7]). This means that $\mathbb{Z}/5\mathbb{Z} \subset \mathrm{GL}_3(\mathbb{F}_2)$. But it is impossible, because $|\mathrm{GL}_3(\mathbb{F}_2)| = 168$.

The case l = 5 is impossible as well, because the group S_6 does not contain subgroups of index 5. If l = 10 then there are orbits of the action of S_6 on \mathbb{P}^3 of length at most 5. But as we have just shown, it is also impossible.

Therefore, the only possible case is l = 15. Thus, the cubic surface S with an action of S₆ passes through 15 points. By Lemma 1 such cubic surfaces are isomorphic to the cubic surface of the form (1).

Now we prove Theorem 3.

Proof of Theorem 3. The first part of the theorem follows from Theorem 4. The existence and uniqueness follow from Lemmas 1 and 2.

6

Also we obtain the following corollary from Lemmas 1 and 2.

Corollary 2. The group S_6 acts on a smooth cubic surface S over \mathbb{F}_2 if and only if S passes through all 15 points in \mathbb{P}^3 .

References

- A. Al Amrani, Classes d'idéaux et groupe de Picard des fibrés projectifs tordus, K-Theory, 2 (1989), no. 5, 559–578.
- [2] J. H. Conway, R. T. Curtis, S. Ph. Norton, R. A. Parker, R. A. Wilson, ATLAS of Finite Groups, Oxford University Press, (1985).
- [3] Y. Chen, C. Shramov, Automorphisms of surfaces over fields of positive characteristic, arXiv:2106.15906 [math.AG].
- [4] J. Dieudonné, Les Isomorphismes Exceptionnels Entre Les Groupes Classiques Finis, CJM, 6 (1954), 305–315.
- [5] I. Dolgachev, Classical algebraic geometry. A modern view, Cambridge University Press, Cambridge, 2012
- [6] I. Dolgachev, A. Duncan, Automorphisms of cubic surfaces in positive characteristic, Izv. RAN. Ser. Mat., 83 (2019), no.3, 15–92.
- [7] I. V. Dolgachev, V. A. Iskovskikh, Finite Subgroups of the Plane Cremona Group, In: Y. Tschinkel, Y. Zarhin, (eds) Algebra, Arithmetic, and Geometry. Progress in Mathematics, 269 (2009), Birkhäuser Boston.
- [8] I. Dolgachev, G. Martin, Automorphisms of del Pezzo surfaces of degree 2 in characteristic 2, arXiv:2206.08913 [math.AG].
- [9] F. Karaoğlu, Non-Singular Cubic Surfaces over $\mathbb{F}_{2^k},$ Turk. J. Math., 45 (2021), 2492–2510.
- [10] V. L. Popov, On the Makar-Limanov, Derksen invariants, and finite automorphism groups of algebraic varieties, Affine algebraic geometry: the Russell Festschrift, CRM Proceedings and Lecture Notes, 54, Amer. Math. Soc. (2011), 289–311.
- [11] Yu. Prokhorov, C. Shramov, Jordan property for Cremona group over a finite field, arXiv:2111.13367 [math.AG].
- [12] J.-P. Serre, A Minkowski-style bound for the orders of the finite subgroups of the Cremona group of rank 2 over an arbitrary field, Mosc. Math. J., 9 (2009), no.1, 183–198.

LABORATORY OF ALGEBRAIC GEOMETRY, NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS, 6 USACHEVA STR., MOSCOW, 119048, RUSSIA.

STEKLOV MATHEMATICAL INSTITUTE OF RUSSIAN ACADEMY OF SCIENCES, 8 GUBKIN STR., Moscow, 119991, Russia

Email address: vikulovaav@gmail.com