

MATCHING WITH INCOMPLETE PREFERENCES

Aditya Kuvalekar*

University of Essex

October, 2023

ABSTRACT

I study a two-sided marriage market in which agents have incomplete preferences—i.e., they find some alternatives incomparable. The strong (weak) core consists of matchings wherein no coalition wants to form a new match between themselves, leaving some (all) agents better off without harming anyone. The strong core may be empty, while the weak core can be too large. I propose the concept of the “compromise core”—a nonempty set that sits between the weak and the strong cores. Similarly, I define the men-(women-) optimal core and illustrate its benefit in an application to India’s engineering college admissions system.

Keywords: matching, market design, incomplete preferences

JEL codes: D47, D61, D63.

1. Introduction

In market design, it is typically assumed that agents have complete preferences—they can compare any two options. Several authors have questioned the assumption of completeness in various contexts.¹ Especially in applications such as school choice, incomplete preferences may arise naturally. Consider the Indian Institute of Technology’ (IIT) admission system. Each year, 1.3 million students seek admission for about 34,000 engineering seats across IITs and a few non-IITs. The current procedure involves ranking students based on their performance on two exams ([Baswana et al. \(2019\)](#)). Thereafter, students submit their preferences—a

*Email: a.kuvalekar@essex.ac.uk

I am indebted to Kazuhiro Hara and Elliot Lipnowski for their thoughts and help on this paper. I would like to thank Nageeb Ali, Joyee Deb, Laura Doval, Bhaskar Dutta, Piotr Dworczak, Mikhail Freer, Niccolò Lomys, Ana Mauleon, Marek Pycia, Ariel Rubinstein, Arunava Sen and Alex Teytelboym for comments and suggestions.

¹For example, [Aumann \(1962\)](#), [Bewley \(1986\)](#), [Ok et al. \(2002\)](#).

strict, complete ordering over available choices. A seat is a two-dimensional object consisting of the institute—e.g. IIT Bombay, IIT Delhi—and the discipline of study—e.g. computer science, mathematics, chemical engineering. Suppose Anita prefers IIT Bombay over IIT Delhi (the former being closer to her home), whereas her preferred discipline is computer science, followed by chemical engineering, followed by mathematics. Then, Anita might find computer science at IIT Delhi *incomparable* with chemical engineering at IIT Bombay. Many students face this dilemma each year—making a choice between a more-preferred institute and a more-preferred discipline—and a number of websites offer advice to students on these dimensions.²

When forced to report complete preferences, students must artificially resolve this indecisiveness one way or the other. However, might it be “better” to allow students to express incompleteness? How should we think of stability, and the core, when preferences are incomplete? These questions motivate the current paper.

I study the two-sided marriage market problem as in Gale and Shapley (1962), but with one distinction—agents have *transitive but possibly incomplete* preferences.³ With incompleteness, two natural notions of the core (and stability) immediately come to mind. The weak core consists of matchings wherein no coalition wants to deviate and form a match between themselves that improves the allocation of *all* the agents in the coalition. The strong core consists of matchings wherein no coalition wants to deviate and form a match between themselves that improves the allocation of *at least one* agent in the coalition *without harming anyone* in the coalition.⁴

When agents have incomplete preferences, the strong core may be empty and the weak core may be too large. However, these problems are not the sole motivation for an intermediate solution concept. Consider the example below.

EXAMPLE 1: Let $M = \{m_1, m_2\}$ and $W = \{w_1, w_2\}$. Each w_i likes m_1 better than m_2 . However, women cannot compare remaining single with being matched with either man. Men cannot compare any alternatives. In particular, they cannot compare staying single with being matched. ■

The weak core consists of all possible matchings which we enumerate below, where an ordered pair denotes a match. For example, in μ_1 all the agents are

²For example, <https://www.quora.com/What-branch-and-IIT-should-I-choose>.

³An important question is whether the distinction between indecisiveness and indifference is important. Why can we not just replace indecisiveness with indifference? I discuss this at the end of the introduction.

⁴Notice that, in contrast to the situation with complete preferences, some agents participating in a coalitional deviation here may move to an allocation that is incomparable with their original allocation. See Definition 6 for a formal version.

single, while in μ_2 , m_1 and w_1 are matched to each other, and m_2 and w_2 are single.

$$\begin{aligned}\mu_1 &= \{\}, & \mu_2 &= \{(m_1, w_1)\}, \mu_3 = \{(m_2, w_2)\} \\ \mu_4 &= \{(m_2, w_1)\}, & \mu_5 &= \{(m_1, w_2)\}, \mu_6 = \{(m_1, w_1), (m_2, w_2)\} \\ \mu_7 &= \{(m_2, w_1), (m_1, w_2)\}\end{aligned}$$

The strong core is $\{\mu_1, \mu_2, \mu_5\}$.

I claim that the weak core is too permissive, while the strong core is too restrictive in the above example. Notice that the weak core contains even μ_3 and μ_4 —matchings wherein each woman is assigned to her second-ranked man, while the top-ranked man remains single. This seems somewhat unsatisfactory. The strong core justifiably eliminates those, but, in doing so, it also eliminates μ_6 and μ_7 —matchings wherein all the agents are matched.

Compare μ_6 with μ_2 , for instance. The only difference between the two is that μ_6 matches m_2 with w_2 while μ_2 keeps them single. More importantly, neither m_2 nor w_2 can compare remaining single with being matched to each other. Therefore, as an analyst, we may not want to necessarily assert that μ_6 is better than μ_2 just because it forms more matches. However, by including μ_2 while excluding μ_6 , the strong core implicitly takes a strong stand *against* forming matches in the above example.⁵

In short, even when the strong core is nonempty, it may refine the weak core to an unappealing subset. Can one do “better”? If yes, in what sense? The main contribution of my paper is to answer these questions by proposing a concept of the core: “*the compromise core*”—a nonempty set that sits between the weak and the strong cores.

As the name suggests, it is a compromise that circumvents the emptiness of the strong core while retaining the power to refine the weak core. Moreover, Theorem 1 provides the sense in which the compromise core is a meaningful refinement: it is characterized by three normatively natural axioms. In Example 1, the compromise core selects μ_6 and μ_7 along with the strong core.

The compromise core is predicated on the concept of “strong moves”—moves by coalitions where a subset of agents inside a coalition become strictly better off without harming anyone inside the coalition. Strong moves—also responsible for the potential emptiness of the strong core—motivate an order called “dominance” on the weak core. Loosely speaking, a matching μ dominates μ' if a coalition

⁵This is not to suggest that the strong core takes a stand against forming matches more generally.

can precipitate a move from μ' to μ (can redistribute matches among themselves without disturbing the matches outside), and some members of the coalition find such a move strictly desirable, while no member finds it undesirable.

Unsurprisingly, dominance is sometimes intransitive and sometimes incomplete. If this relation were transitive, we could simply select the maximal elements. Therefore, the question is, how should one refine the weak core endowed with a natural but an intransitive order? Incidentally, an extensive literature in social choice theory constructs reasonable choice rules that circumvent the problem of nonexistence of clear winners due to intransitivities (e.g. majority rule in voting). A particularly appealing choice rule is the uncovered set due to [Miller \(1980\)](#).⁶ The Miller covering induced by my dominance relation, henceforth *covering*, says that a matching μ covers μ' if, (i) μ dominates μ' and (ii) μ dominates any matching that μ' dominates. The compromise core is the nonempty set of uncovered elements.

Let me briefly comment on a natural question raised earlier: Is the distinction between indifference and indecisiveness important? Indeed, if one views choice data as a primitive and preferences as derived objects, then incomparability is indistinguishable from indifference *if we allow for intransitive preferences*.

Let us go back to Anita's example and say that she is choosing among three colleges, C_1, C_2 , and C_3 . She ranks C_2 better than C_3 because it is closer to her home and offers a discipline that she prefers. However, she cannot compare C_1 with either C_2 or C_3 because C_1 is the farthest from her home but offers the discipline she most prefers. Notice that if we replace all the incomparabilities for Anita with indifference, then her preference would be $C_1 \sim C_2, C_1 \sim C_3$ and $C_2 \succ C_3$ —the preference would be intransitive.⁷ However, at least in the context of school choice, there is an important distinction between indifference and incompleteness. [Roth and Sotomayor \(1992\)](#) suggest:

Loosely speaking, the reason is that indifference is in some sense a “knife edge” phenomenon; if an agent is indifferent between two alternatives, a small improvement in one of them would presumably cause him or her to prefer it to the other. (p.35)

Translating this into Anita's environment, if, hypothetically, C_1 were to move closer to her home by a tiny distance, Anita should strictly prefer C_1 over both C_2 and C_3 *if she were indifferent*. It is the incomparability between these colleges that would prevent Anita from having such a strict ranking.

⁶The notions of the uncovered set due to [Miller \(1980\)](#) and [Fishburn \(1977\)](#) are (not) equivalent when the underlying binary relation is complete (incomplete) and strict.

⁷[Luce and Raiffa \(1989\)](#) discuss the possibility of intransitivities occurring when subjects compare inherently incomparable alternatives.

While agnostic about the foundation of incompleteness *vs.* intransitive indifference, I adopt the interpretation of indecisiveness throughout. However, my proposed solution concepts and results remain equally valid when preferences are allowed to be intransitive with a transitive strict part.

Finally, I briefly discuss the IIT admission system. I show that the current process may generate matchings in the weak core that are strictly dominated according to students' preference by another matching in the weak core. Therefore, allowing students to express incomplete preferences may materially improve outcomes.

2. Model

2.1. Preliminaries

Let M and W denote disjoint finite sets, which I refer to as men and women following the literature. Let $A := M \cup W$. A generic agent in M [resp. W] will be denoted by $m[w]$. When the distinction is immaterial, I denote an agent by i . Let $O(i) = W \cup \{i\}$ if $i \in M$, and $O(i) = M \cup \{i\}$ if $i \in W$, denote the *opposite side* for agent i .

Each agent $i \in A$ is endowed with a preference relation, \succsim_i over $O(i)$. \succsim_i is a binary relation that is reflexive, transitive, and antisymmetric, but possibly incomplete.⁸ That is, i may find two agents, $j, k \in O(i)$ incomparable. I denote this by $j \otimes_i k$. Moreover, since \succsim_i is antisymmetric, $j \succsim_i k$ means that i strictly prefers j over k , denoted $j \succ_i k$, whenever $j \neq k$.⁹ I say \succsim'_i is “*more complete*” than \succsim_i , if $j \succsim_i k \implies j \succsim'_i k$ for all $i \in A$. Lastly, $\succsim := (\succsim_i)_{i \in A}$ is called a preference profile.

Let S be a set endowed with a binary relation R , which need not be complete or transitive. Let P be the strict part of R , i.e. xPy if xRy but not yRx . We say that x is a maximal element of S if no $y \in S$ has yPx ; x is a maximum element of S if xPy for all other $y \in S$. The set of maximal elements is denoted by $MAX(S, R)$.

A bijection $\mu : A \rightarrow A$ is called a matching if $\mu(i) \in O(i)$ for all $i \in A$, and $\mu(i) = j \implies \mu(j) = i$. I let \mathcal{M} denote the set of all matchings. Say a matching

⁸A binary relation R on a set X is reflexive if xRx , it is transitive if $xRy, yRz \implies xRz$, and it is antisymmetric if $xRy, yRx \implies x = y$.

⁹While I take preferences as the primitive, one could, alternatively start with a choice behaviour that is consistent with incomplete preferences as a primitive to define stability and core as in [Che et al. \(2019\)](#) for example. [Eliaz and Ok \(2006\)](#) provide a revealed preference foundation of incomplete preferences.

$\mu \in \mathcal{M}$ is *individually rational* if no $i \in A$ has $i \succ_i \mu(i)$. Below, I define some natural notions of stability.

DEFINITION 1: A matching μ is “weakly stable” if it is individually rational and no $(m, w) \in M \times W$ has $w \succ_m \mu(m)$ and $m \succ_w \mu(w)$. If there exists such a (m, w) , then I say that (m, w) is a “strong blocking pair” for μ .

DEFINITION 2: A matching μ is in the “weak core” if $\nexists S \subset A$ and a matching $\mu' \neq \mu$ such that $\mu'(i) \in S$ for all $i \in S$, and $\mu'(i) \succ_i \mu(i)$ for all $i \in S$. If there is such an S , then I say that S strongly blocks μ .

The reason to call the above as a “strong blocking pair” is that we could define a weaker notion—a weak blocking pair—as below.

DEFINITION 3: Given a matching μ , a pair (m, w) is a “weak blocking pair” for μ if one of the following holds.

1. (m, w) is a strong blocking pair.
2. $m \succ_w \mu(w)$ and $w \otimes_m \mu(m)$.
3. $w \succ_m \mu(m)$ and $m \otimes_w \mu(w)$.

In contrast to the strong blocking pair, the notion of a weak blocking pair is somewhat nonstandard. If (m, w) constitute a weak blocking pair for μ , it means the following: by leaving their respective match in μ and matching with each other, (1) neither agent is worse off, and, (2) at least one of them is strictly better off. One may wonder why an agent would participate in a move where (s)he switches between two allocations that (s)he finds incomparable. The reason why such moves maybe reasonable stems from the observation that every incomplete preference relation is an intersection of a set of complete preference relations. Therefore, we can view an agent’s incomplete preference as if it arises from using multiple criteria to evaluate the alternatives, as in Anita’s example from the Introduction. An agent, say m , may find two alternatives incomparable because some of the criteria he uses, do not rank the said two alternatives the same way. In such a case, the reason why m may participate in a move from one alternative to the other is simply because he is not worse off from such a move. In other words, participating in such a move assumes acquiescence rather than hesitancy on the part of the agent in moving between two incomparable allocations.

DEFINITION 4: A matching μ is “strongly stable” if it is individually rational and has no weak blocking pair.

DEFINITION 5: Let $\mu(S) := \{\mu(i) : i \in S\}$. Say that $\mu \succ_S \mu'$ if

- (i) $\mu(S) = \mu'(S)$, and,

(ii) $\mu(i) \not\prec_i \mu'(i)$ for all $i \in S$, and, $\mu(i) \succ_i \mu'(i)$ for at least one $i \in S$.

Notice that, if the preferences were complete, then \succ_S would coincide with the Pareto ranking (for the agents in S), namely, $\mu \succ_S \mu'$ if $\mu(i) \succ_i \mu'(i)$ for all $i \in S$, and $\mu(i) \succ_i \mu'(i)$ for some $i \in S$. On the other hand, with incomplete but strict preferences, matchings undominated according to the Pareto ranking (as defined on the previous line), would coincide with the weak core. Of course, the definition of \succ_S as in Definition 5 is less demanding than the Pareto ranking when the agents have incomplete preferences. Similar to the weak blocking pairs, it captures the idea that some agents in S may be willing to participate in a move from μ' to μ simply because they are not worse off from such a move.

DEFINITION 6: A matching μ is in the “strong core” if $\nexists S \subset A$ and a matching μ' , such that $\mu' \succ_S \mu$. If there is such an S , then I say that S weakly blocks μ .

Proposition 1 below, whose analogues for complete preferences are well-known in the literature, establishes that the weak (strong) core and the set of weakly (strongly) stable matchings coincide. Therefore, I refer to these two concepts interchangeably henceforth.

PROPOSITION 1: A matching μ is weakly [resp. strongly] stable iff it is in the weak [strong] core.

Let \mathcal{W} [resp. \mathcal{S}] denote the weak [strong] core.

A binary relation $\widehat{\succ}$ is a “completion of \succ ” if, for each i , $\widehat{\succ}_i$ is a complete, reflexive, transitive, and antisymmetric binary relation over $O(i)$, such that $j \succ_i k \implies j \widehat{\succ}_i k$.

When the underlying preference is complete, the notions of weak and strong stability coincide. Therefore, when agents have complete preferences, a matching is stable if it is weakly stable. It is easy to characterize the weakly stable matchings in terms of completions of \succ . To this end, let $C(\succ)$ denote the set of completions of \succ .

PROPOSITION 2: A matching μ is weakly stable according to \succ iff it is stable according to some completion $\widehat{\succ}$.

The following corollaries are immediate.

COROLLARY 1: If \succ is more complete than \succ' , then $\mathcal{W}(\succ) \subseteq \mathcal{W}(\succ')$.

COROLLARY 2: The weak core is nonempty.

In contrast to Corollary 2, and as one might expect, the strong core may be empty, as the following example demonstrates.

EXAMPLE 2: Let $M = \{m_1, m_2, m_3\}$, $W = \{w_1, w_2, w_3\}$. All men have identical preferences: $w_1 \otimes_m w_2, w_1 \otimes_m w_3, w_2 \succ_m w_3$ for all $m \in M$. All women have identical preferences, too: $m \otimes_w \tilde{m}$, for all $w \in W$ and different $m, \tilde{m} \in M$. Moreover, all the agents strictly prefer being matched over remaining single. ■

First, it is obvious that $\mathcal{W} = \{\mu \in \mathcal{M} : \mu(i) \neq i \forall i \in A\}$. Consider an arbitrary matching $\mu \in \mathcal{W}$. Letting $m = \mu(w_3)$, notice that (m, w_2) is a weak blocking pair as $w_2 \succ_m w_3$ and $m \otimes_{w_2} \mu(w_2)$.

Analogous to Proposition 2, one might wonder whether the strong core coincides with matchings that are stable with respect to every completion of \succ . This is, however, not true.

PROPOSITION 3: If μ is stable with respect to every completion of \succ , then μ is in the strong core. However, the strong core can be strictly larger than the set of matchings that are stable with respect to every completion.

To see the strictly larger part, consider a market with one man and one woman such that no agent is able to compare remaining single to being matched. The strong core contains all possible matchings. However, no matching is stable with respect to every completion.

3. Compromise Core

While the strong core may be empty as we just saw, Example 1 demonstrates how it can discard intuitively appealing matchings even when it is non-empty. These observations motivate the search for an alternative solution concept like the compromise core. At its heart are coalitional moves. Weakly stable matchings are prone to *strong moves* by coalitions—moves where only a subset of agents need to strictly improve while nobody inside the coalition is worse off. Unsurprisingly, it is possible to cycle through weakly stable matchings when one allows for strong moves. Therefore, I begin the process of refining the weak core by endowing the weak core with the dominance order: a matching μ dominates μ' if there is a coalition that can “enforce” a move from μ' to μ , and the coalition also finds such a move “preferable”—some members of the coalition find this move strictly better off while nobody in the coalition is worse off. Below, I make these ideas precise.

DEFINITION 7: Given a matching μ , a coalition $S \subset A$ is said to be able to enforce a matching μ' over μ if the following conditions hold: (a) $\mu'(i) \notin \{\mu(i), i\} \implies \{i, \mu'(i)\} \subset S$ and (b) $\mu'(i) = i \neq \mu(i) \implies \{i, \mu(i)\} \cap S \neq \emptyset$.

The above definition is due to [Mauleon et al. \(2011\)](#). As they say, “This enforceability condition implies both that any new match in μ' that does not exist

in μ should be between agents in S , and that to destroy an existing match in μ , one of the two agents involved in that match should belong to coalition S .”

Below, I define dominance—an order that forms the basis of refining the weak core.

DEFINITION 8: Consider two matchings $\mu, \mu' \in \mathcal{W}$.¹⁰ Say that μ dominates μ' , denoted by $\mu \succeq \mu'$, if there is $S \subset A$ such that

1. S is able to enforce a matching μ over μ' , and,
2. $\mu \succ_S \mu'$.¹¹

\triangleright denotes the strict part of \succeq . That is, $\mu \triangleright \mu'$ if $\mu \succeq \mu'$ and $\neg(\mu' \succeq \mu)$.

Notice that predicating a refinement on the dominance relation entails considering blockings that involve incomparable matches. That is, when $\mu \triangleright \mu'$, a coalition that enforces a move from μ' to μ will have some agents' allocation in μ and μ' being incomparable. If these were indifferences instead of incomparabilities, then such moves are easy to justify—the agents who strictly prefer their allocation in μ can pay a small “bribe” (not formally in the model) to induce the indifferent agents to accede to the coalitional move. However, if one adopts a (reasonable) perspective that moves involving transitions between incomparable allocations should not be allowed, then it seems that the weak core cannot be refined further in light of Proposition 2. Therefore, to the extent that one views refining the weak core as a meaningful exercise, the dominance relation offers a reasonable alternative towards that goal.

Dominance, while intuitively appealing as an order on \mathcal{W} , can be incomplete or intransitive. Moreover, \succeq need not be antisymmetric either, i.e., there can exist $\mu, \mu' \in \mathcal{W}$ such that $\mu \succeq \mu' \succeq \mu$ (Lemma 1 in the Appendix). Also, one may wonder if \succeq , when restricted to \mathcal{W} , is essentially a form of Pareto dominance. That is, say that μ strongly Pareto dominates μ' if $\mu \succ_{M \cup W} \mu'$. Of course, if μ strongly Pareto dominates μ' then $\mu \succeq \mu'$. The converse is not true.

If \succeq were a nice binary relation (e.g. complete and transitive), the task of refining \mathcal{W} would be straightforward: Choose the \succeq -maximal elements. However, due to intransitivities, a maximal element may not exist. This is neither surprising (given how dominance is defined), nor is the problem unique—especially for the practitioners of social choice theory. Such intransitivities abound in problems dealing with voting rules where the majority rule, while being complete, often fails to be transitive. As Dutta (1988) says, one of the popular pastimes of social choice theorists has been to construct choice rules that handle the difficulty posed

¹⁰The dominance relation can be defined over \mathcal{M} and not just \mathcal{W} . However, since the objective of the exercise is to refine \mathcal{W} , I directly define it over \mathcal{W} .

¹¹Recall that $\mu \succ_S \mu'$ means that no $i \in S$ has $\mu'(i) \succ_i \mu(i)$, and some $i \in S$ has $\mu(i) \succ_i \mu'(i)$.

by the nonexistence of a clear winner.

One such choice rule that seems particularly well-suited for our environment is the “*uncovered set*”, proposed in Fishburn (1977) and Miller (1980). To this end, I introduce the notion of covering below.

DEFINITION 9: Given $\mu, \mu' \in \mathcal{W}$, we say that μ covers μ' , denoted by $\mu \ll \mu'$ if, (i) $\mu \triangleright \mu'$ and, (ii) every $\mu'' \in \mathcal{W}$ with $\mu' \triangleright \mu''$ has $\mu \triangleright \mu''$.¹²

While \triangleright may itself be intransitive or incomplete, \ll is an antisymmetric and transitive (but possibly incomplete) binary relation (Lemma 2 in the Appendix). The compromise core, formally defined below, is the set of \ll -maximal elements or, the uncovered set: matchings that are not covered by any other matching.

DEFINITION 10: The compromise core \mathcal{C} is the set of all the matchings $\mu \in \mathcal{W}$ that are not covered by any other matching $\mu' \in \mathcal{W}$. Formally, $\mathcal{C} := \{\mu \in \mathcal{W} : \nexists \mu' \in \mathcal{W} \text{ s.t. } \mu' \ll \mu\}$.

REMARK 1: A detailed expository example illustrating the compromise core is available in the Appendix A.1.

Several notions of covering are extensively studied in social choice theory.¹³ While they vary in their discriminatory power, I find the covering relation due to Miller (1980) the most appealing. To see why, let us ask why we would want a certain matching admitted by our solution concept? One reason would be that it is undominated. The other would be that it dominates a subset of matchings that makes it compelling. However, if μ covers μ' , then μ' is neither undominated nor does it dominate any matching that μ does not. Therefore, there is arguably no additional advantage of selecting μ' over μ . Consequently, when equipped with the goal of refining \mathcal{W} , if one must include one of μ and μ' , it should be μ . Theorem 1 offers further justification for the compromise core through three simple axioms.

There is another well-studied notion of covering, the Fisher covering (Fishburn (1977)), that is close to Miller-covering, i.e., the notion of covering used in this paper. We say that a matching μ *F-covers* μ' if, $\mu \triangleright \mu'$ and $\mu'' \triangleright \mu \implies \mu'' \triangleright \mu'$. If \triangleright were a strict and a complete order, F-covering and Miller-coverings are equivalent. However, especially with the incompleteness of \triangleright , the Fisher uncovered set seems unappealing. To see this, suppose that $\mathcal{W} = \{\mu_1, \mu_2, \mu_3\}$ for some matching environment wherein, $\mu_1 \triangleright \mu_2$ and $\mu_2 \triangleright \mu_3$ but μ_1 and μ_3 are not \triangleright -comparable (e.g. Example 4 in the Appendix is one such instance). Consider the following two properties that one may consider desirable from a refinement. First, any

¹²Recall that the definition of dominance, \triangleright , is given in Definition 8, while the definition of \succ_S (in the definition of dominance) is given in Definition 5.

¹³See Duggan (2013) for a detailed discussion and comparison of some variants. Some prominent ones are Gillies (1959), Fishburn (1977), Miller (1980), Bordes (1983).

\triangleright -undominated matching must be selected. Second, any \triangleright -dominated matching that does not dominate any other matching, must be excluded. That is, if $\mu \triangleright \mu'$ and there is no μ'' such that $\mu' \triangleright \mu''$, then μ' should be excluded. Putting these two together, the two possible candidate refined sets are: $\{\mu_1\}$ and $\{\mu_1, \mu_2\}$. Notice that, μ_2 Miller-covers μ_3 while μ_1 and μ_2 are Miller-uncovered. On the other hand, μ_1 F-covers μ_2 . Therefore, the compromise core is $\{\mu_1, \mu_2\}$ while the Fisher uncovered set is $\{\mu_1, \mu_3\}$. It is difficult to justify a refinement that includes μ_3 while excluding μ_2 .

I wish to point out two structural differences between \mathcal{C} and the Miller-uncovered set defined on a set endowed with a complete order. First, when the underlying relation is complete, any alternative in the Miller uncovered set beats any other alternative in at most two steps. That is, given any μ in the Miller uncovered set and any $\mu' \in \mathcal{W}$, either $\mu \triangleright \mu'$ or $\exists \mu_1 \in \mathcal{W}$ such that $\mu \triangleright \mu_1 \triangleright \mu'$, if \triangleright were complete. With incompleteness, this is not true. More importantly, whenever the underlying relation is complete, the Miller uncovered set is contained in the top cycle set defined as follows:

$$TC := MAX(\mathcal{W}, \triangleright^T)$$

where $MAX(A, R)$ denotes the maximal elements of set A according to a binary relation R , while R^T denotes the transitive closure of R .¹⁴ However, \mathcal{C} can be strictly larger than the top cycle.

3.1. Relationship to the strong and the weak core

We now see that the strong core is a subset of the compromise core.

PROPOSITION 4: *If $\mu \in \mathcal{S}$, then $\nexists \mu' \in \mathcal{W}$ such that $\mu' \triangleright \mu$. Therefore, $\mathcal{S} \subseteq \mathcal{C}$.*

PROPOSITION 5: *The strong core is a subset of the compromise core, while the compromise core is a nonempty subset of the weak core. Moreover, these inclusions can be strict even when the strong core is nonempty.*

Proof. By Proposition 4, $\mathcal{S} \subseteq \mathcal{C}$. The second inclusion, $\mathcal{C} \subset \mathcal{W}$, is an immediate consequence of the definition of the compromise core. That, \mathcal{C} is nonempty follows from the transitivity of \ll as mentioned before. The second part of the proposition is shown in Example 1. Recall that, in that example, $\mathcal{S} = \{\mu_1, \mu_2, \mu_5\}$, and therefore they belong to \mathcal{C} . Finally, notice that $\mu_2 \triangleright \mu_7$ and $\mu_7 \triangleright \mu_3$, but μ_2 and μ_3 are not \triangleright -comparable. Moreover, there is no other μ such that $\mu \triangleright \mu_7$. Therefore, μ_7 is uncovered. Similarly, μ_6 is uncovered. Therefore, $\mathcal{C} = \{\mu_1, \mu_2, \mu_5, \mu_6, \mu_7\}$. \square

¹⁴Transitive closure of a relation R is defined as $xR^T y$ iff $\exists x_1, x_2, \dots, x_n$ such that $xRx_1Rx_2 \dots x_nRy$.

Notice that the top-cycle in the above example is $\{\mu_1, \mu_2, \mu_5\}$ which is strictly smaller than \mathcal{C} .

In Example 1, the compromise core contains, beyond the strong core, the matchings in which no agent is single. On the other hand, matchings that are not in the compromise core match only one woman to her second-best partner, leaving her best partner unmatched. This highlights an advantage of the compromise core over the weak and the strong cores. The compromise core creates more pairs than does the strong core, to fulfill some individuals by exploiting the incomparability of others. In doing so, the compromise core is more selective than the weak core.

4. Normative Foundations of the Compromise Core

In this section, I show that a set of normatively appealing axioms characterize the compromise core.

To be concrete, consider the problem of designing a map $\Gamma : 2^{\mathcal{W}} \rightarrow 2^{\mathcal{W}}$ such that $\emptyset \neq \Gamma(A) \subset A$ for all $A \subset \mathcal{W}$. That is, given a set of matchings in \mathcal{W} , $\Gamma(\cdot)$ refines them to produce a subset of those matchings. For example, the compromise core, $\mathcal{C}(A)$ —the set of \llcorner -uncovered elements in A —is one such map.

Let us introduce some notation. First, $\mu^\uparrow := \{\mu' \in \mathcal{W} : \mu' \triangleright \mu\}$ and $\mu^\downarrow := \{\mu' \in \mathcal{W} : \mu \triangleright \mu'\}$. Second, for any two matchings, μ, μ' , and a set $T \subset \mathcal{W}$ such that $\mu, \mu' \in T$, let $\mathcal{A}_3(\mu, \mu', T) := \{\{\mu, \mu', \hat{\mu}\} : \hat{\mu} \in T\}$. That is, $\mathcal{A}_3(\mu, \mu', T)$ is a collection of all possible subsets of T of size at most 3 that contain μ and μ' . Notice that by having $\hat{\mu} = \mu$ or μ' , $\{\mu, \mu'\} \in \mathcal{A}_3(\mu, \mu', T)$.

Now, I present three normatively desirable axioms for a refinement. The main result of this section, Theorem 1, shows that these axioms characterize \mathcal{C} .

Include the Maximal Element (IM):

For any $T \subset \mathcal{W}$ and $\mu \in T$, if $\mu^\uparrow \cap T = \emptyset$, then $\mu \in \Gamma(T)$.
 Moreover, if \triangleright is a strict and a complete order on T , then $\Gamma(T) = \{\mu \in T : \mu \triangleright \mu' \forall \mu' \in T\}$.

This axiom requires that, if μ is a maximal element in T , then $\Gamma(T)$ must include μ , and if T is totally ordered by \triangleright , then $\Gamma(T)$ must select only the unique such element.

Exclude the bottom (EB):

For any $T \subset \mathcal{W}$, if $\mu^\downarrow \cap T = \emptyset$ and $\mu^\uparrow \cap T \neq \emptyset$, then $\mu \notin \Gamma(T)$.

This axiom requires a refinement to exclude the “bottom layer” according

to \triangleright . That is, if a matching μ does not dominate any other matching in \mathcal{W} , and is also strictly dominated by some other matching $\mu' \in \mathcal{W}$, then a good refinement should exclude μ according to EB.

Expansion from triples (ET):

If, for some μ, μ', T such that $\mu, \mu' \in T \subset \mathcal{W}$, $\mu' \notin \Gamma(T') \forall T' \in \mathcal{A}_3(\mu, \mu', T)$, then $\mu' \notin \Gamma(T)$.

This axiom is a form of consistency requirement from observing $\Gamma(\cdot)$'s behaviour on sets of size 3 or less. Alternatively, it can also be loosely interpreted as a form of revealed preference. The axiom says the following. Fix a set T that contains μ and μ' . If μ' is never chosen from any subset of T of size 3 or less that includes both μ and μ' , then μ' should not be chosen from T .

While the IM and EB axioms seem relatively natural, the ET axiom perhaps merits some discussion. This axiom above is, essentially, a weakening of WARP. Similar weakenings of WARP have appeared in the literature on incomplete preferences in choice theory, e.g., [Eliaz and Ok \(2006\)](#). More closely, this axiom is similar to the “upward consistency” axiom from [Nishimura \(2018\)](#) that deals with inferring a transitive part of a possibly cyclic choice behavior.

That the above three axioms are mutually independent is straightforward to see. In light of that observation, I present the main result of this section—a characterization of \mathcal{C} using IM, EB and ET.

THEOREM 1: *Any refinement $\Gamma(\cdot) : 2^{\mathcal{W}} \rightarrow 2^{\mathcal{W}}$ that satisfies IM, EB and ET has $\Gamma(\cdot) \subseteq \mathcal{C}(\cdot)$. Conversely, $\mathcal{C}(\cdot)$ satisfies IM, EB and ET. Hence, $\mathcal{C}(\cdot)$ is the most permissive refinement of \mathcal{W} satisfying IM, EB and ET.*

4.1. Relation to other solution concepts

The characterization in [Theorem 1](#) leaves open the possibility that the compromise core can contain matchings μ and μ' such that $\mu \triangleright \mu'$. This may be unsatisfactory if one holds the perspective that a solution concept describes a “system at rest.” That is, we may want a solution concept to not include two matchings wherein one dominates the other. The vNM stable set ([von Neumann and Morgenstern \(1944\)](#)) is one such concept. In our context, one would say that a set S is a vNM stable set if it satisfies the following two properties.

Internal stability: No $\mu, \mu' \in S$ has $\mu \triangleright \mu'$.

External stability: If $\mu \notin S$, then some $\mu' \in S$ has $\mu' \triangleright \mu$.

The above two axioms, while seemingly natural, can easily lead to the situations where no vNM stable set exists. For example, consider [Example 2](#). There, $\mathcal{W} =$

$\{\mu_1, \mu_2, \mu_3\}$ with $\mu_1 \triangleright \mu_2 \triangleright \mu_3 \triangleright \mu_1$. Internal stability demands that any candidate stable set can have at most one element. However, no such set can satisfy external stability.

Another solution concept popularly used in cooperative games is the bargaining set.¹⁵ Loosely speaking, if $\mu' \triangleright \mu$ due to a move enforced by S , then we say that (S, μ') is an objection to μ . However, if there is a set T such that some (but not all) members of S are in T and (T, μ'') is an objection to μ' , then we say that (T, μ'') is a counterobjection against an objection (S, μ') . A matching μ is in the bargaining set if, for every objection there is a counterobjection.

Regardless of whether the bargaining set is always nonempty (which I conjecture it is), or whether a vNM stable set exists, they are unappealing because they can fail a natural requirement of EB: Exclude the bottom. Consider Example 4 given in the Appendix.¹⁶ Its weak core contains three elements, $\{\mu_1, \mu_2, \mu_3\}$ such that $\mu_2 \triangleright \mu_1$, $\mu_3 \triangleright \mu_2$, but μ_3 and μ_1 are not \triangleright -comparable. What would we want as a reasonable solution concept in this example? First, it is uncontentious that it should include μ_3 , an \triangleright -undominated matching. Should it include μ_1 ? For one, μ_1 is \triangleright -dominated by μ_2 , and it does not dominate any other matching. Therefore, if a refinement needs to include μ_1 , we may do better by including μ_2 and excluding μ_1 instead.

Indeed, this is precisely the compromise core, i.e., $\mathcal{C} = \{\mu_2, \mu_3\}$. On the other hand, the only vNM stable set and the bargaining set are $\{\mu_1, \mu_3\}$. By including μ_1 and excluding μ_2 , the vNM stable set and the bargaining set not only include a dominated matching which does not dominate any other matchings, but also exclude its upper contour set.¹⁷

It is a worthwhile goal to refine \mathcal{C} further, given that it is the most permissive solution concept satisfying the three natural normative properties mentioned before Theorem 1. A straightforward way to do that, while achieving internal stability, would be to select the largest internally stable subsets of \mathcal{C} . Of course, any such set may not satisfy external stability that \mathcal{C} does, as Proposition 6 below shows.

PROPOSITION 6: *If $\mu \in \mathcal{W} \setminus \mathcal{C}$, then $\exists \mu' \in \mathcal{C}$ such that $\mu' \triangleright \mu$.*

¹⁵For an example of the bargaining set in the matching environment, see [Atay et al. \(2021\)](#).

¹⁶I have avoided presenting the full example here primarily because I only use its weak core for the present argument.

¹⁷An upper contour set of a matching μ is μ^\uparrow .

5. Men- and Women-Optimal Core

We now turn our attention to another object of interest for applications—“best” stable matchings according to one side of the market. For example, in school choice, it is reasonable to favor students and select a stable matching that they prefer the most. As is standard, I define an order \succ_M on \mathcal{W} to mean, $\mu \succ_M \mu'$ iff there is no $i \in M$ such that $\mu'(i) \succ_i \mu(i)$ and for at least one $i \in M$, $\mu(i) \succ_i \mu'(i)$. Order according to women, \succ_W , is defined analogously.¹⁸

When the preferences are complete, strict and transitive, the set of stable matchings is a lattice under the order \succ_M or \succ_W . Therefore, there is a unique men- (women-) optimal stable matching that is the maximum element of the set of stable matchings according to \succ_M (\succ_W). With weak *but transitive* (and complete) preferences, a maximum element need not exist. A natural remedy is to weaken the definition to *maximal* elements—i.e., say that μ is men-optimal if there is no weakly stable μ' such that $\mu' \succ_M \mu$. As mentioned before, [Erdil and Ergin \(2017\)](#) show that such a matching exists. Unfortunately, when preferences are incomplete, matchings that satisfy even this weaker definition may not exist. The reason for this lack of existence of a maximal element is, unsurprisingly, the intransitivity of \succ_M .

This intransitivity motivates an obvious definition of men- and women-optimal core analogous to the compromise core.

DEFINITION 11: μ *M-covers* μ' , denoted by $\mu \ll^M \mu'$ if $\mu \succ_M \mu'$ and $\mu' \succ_M \mu'' \implies \mu \succ_M \mu''$. The *men-optimal core*, \mathcal{W}^M , is defined as $\text{MAX}(\mathcal{C}, \ll^M)$. *Women optimal core*, \mathcal{W}^W , is defined analogously as the set of \ll^W -uncovered elements.

REMARK 2: Notice that the men-optimal core selects the \ll^M -maximal elements from \mathcal{C} and not \mathcal{W} . One could, alternatively, define the men-optimal core to be the \ll^M -maximal elements of \mathcal{W} . Unsurprisingly, the two notions need not coincide. I take \mathcal{C} as a primitive primarily to ensure that the men-optimal core is a subset of \mathcal{C} .

In applications such as school choice, when students have incomplete preferences, forcing them to report complete preferences could yield outcomes that are Pareto dominated for the students. This issue is not new: Similar problems arise if we replace incompleteness with indifference. However, so long as the weak preferences are transitive one can use the [Erdil and Ergin \(2017\)](#)’s algorithm to obtain a men-optimal stable matching. In contrast, reinterpreting incompleteness

¹⁸This definition is, unfortunately, slightly different from \succ_S defined in Definition 5. In particular, we do not require $\mu(M) = M$ here. Instead of choosing a different notation, I still choose to denote it by \succ_M for ease of exposition.

as indifference may result in intransitivities, rendering this algorithm unusable.

Let us consider the IIT-JEE admissions process mentioned in the Introduction. In the current system, students are ranked according to their performance on an exam. Therefore, each object (seat at one of the IITs) ranks the students the same way. Each student submits a strict ranking over the set of objects—i.e., institute and undergraduate major. Then, a version of the student-proposing deferred acceptance algorithm (DAA henceforth) is executed to achieve a matching.¹⁹ As the following example shows, by forcing agents to report complete preferences and running a student-proposing DAA, we may obtain a strictly suboptimal matching for the students.

EXAMPLE 3: Let $W = \{w_1, w_2, w_3\}$, $M = \{m_1, m_2, m_3\}$. Men rank women as $w_1 \succ_m w_2 \succ_m w_3$. Women's preferences are: $m_1 \otimes_w m_2, m_1 \otimes_w m_3, m_2 \succ_w m_3$ for all $w \in W$. Finally, all the agents strictly prefer being matched over remaining single. ■

It is straightforward to check that $\mathcal{W} = \{\mu_1, \mu_2, \mu_3\}$, where

$$\begin{aligned}\mu_1 &= \{(w_1, m_1), (w_2, m_2), (w_3, m_3)\} \\ \mu_2 &= \{(w_1, m_2), (w_2, m_1), (w_3, m_3)\} \\ \mu_3 &= \{(w_1, m_2), (w_2, m_3), (w_3, m_1)\}\end{aligned}$$

First, notice that $\mathcal{C} = \mathcal{W}$. Moreover, $\mu_1 \succ_W \mu_3$ while μ_2 and μ_3 or μ_1 and μ_2 are not \succ_W comparable. Therefore, the set of women-optimal stable matchings—the set of \lll^W uncovered elements in \mathcal{C} —is $\{\mu_1, \mu_2\}$.

However, notice that forcing women to submit a complete ranking and running a DAA thereafter can generate all three matchings in \mathcal{C} . In particular, if the women submit the following preference: $m_2 \succ_w m_3 \succ_w m_1$ —a completion of their original preference—then the outcome would be μ_3 , a strictly inferior outcome for women compared to μ_1 .

If we had allowed the women to submit incomplete preferences, we could have chosen between μ_1 and μ_2 . Therefore, allowing students to express incomplete preferences may generate welfare gains. These could, in principle, be substantial given the size of the market in applications such as college admissions. For example, the number of students that gained admissions just through the single IIT-JEE seat-allocation mechanism was 13,500 in 2019, and 300 seats (2% of the total capacity) went vacant.²⁰

¹⁹Baswana et al. (2015) provide precise details of the procedure.

²⁰<https://www.hindustantimes.com/mumbai-news/over-300-iit-seats-allotted-in-final-round-of-admissions/story-VTKpYlsDxQ9TRpK0iW48NJ.html>.

6. Related Literature

This paper contributes to the literature on matching with nonstandard preferences. A few papers study matching problems with incomplete preferences from a theoretical point of view. [Manlove \(2002\)](#) studies the lattice structure of the stable matchings for three different notions of stability, two of which are the weak and strong stability considered in this paper. [Irving \(1994\)](#) provides algorithms for weak and strong stable matchings allowing incomplete preferences. There is a recent and burgeoning literature studying matchings with incomplete preferences from a computational point of view.²¹ For example, [Bade \(2016\)](#) studies optimality of matching algorithms when preferences are modeled by choice functions. [Chambers and Yenmez \(2017\)](#) develop a deferred acceptance algorithm for matchings when preferences are given as choice correspondences.²² [Che et al. \(2019\)](#) show an existence theorem of many to one stable matchings for general preferences that allow for incompleteness. In particular, they give a fixed-point characterization of the strong core when one side of the market can have incomplete preferences. In a contemporaneous paper, [Kitahara and Okumura \(2021\)](#) study school choice problems allowing for schools to find some students incomparable. They propose strategy proof mechanisms to obtain weakly stable matchings. In contrast, my focus is to offer a new solution concept—the compromise core. In matching with externalities, an agent’s preference over two outcomes may be incomplete *until an agent knows how the rest of the market is organized*. Several papers have studied such markets in the context of various applications, e.g. [Sasaki and Toda \(1996\)](#), [Dutta and Massó \(1997\)](#), [Echenique and Yenmez \(2007\)](#), [Pycia \(2012\)](#), [Pycia and Yenmez \(2021\)](#). Recently, ? study a classical two-sided matching model when agents may exhibit nonstandard choice behaviour. They provide conditions under which a stable matching exists and also when a strategyproof mechanism exists to achieve a stable matching.

While all of the above papers study properties of existing solution concepts under nonstandard preferences, including questions related to their existence, my focus is on proposing new solution concepts suitable for matching problems when agents have incomplete preferences.

Preference incompleteness has been widely studied in decision theory and behavioural economics (e.g. [Ok et al. \(2002\)](#), [Eliaz and Ok \(2006\)](#)), and its implications have also been studied in strategic environments, e.g., [Bade \(2016\)](#). Incompleteness of preferences is closely related to indifference. Indeed, most of

²¹See for example [Manlove et al. \(2002\)](#), [Irving et al. \(2003\)](#), [Aziz et al. \(2017\)](#) and [Cseh and Juhos \(2018\)](#).

²²Also see [Hatfield and Milgrom \(2005\)](#) and [Aygün and Sönmez \(2013\)](#) for more seminal work on matching where preferences are modeled by choice functions.

the findings in this paper can be viewed as results for matchings with intransitive indifference. As such, I encounter problems arising from indifference. For example, under indifference, stable matchings may not be efficient. Erdil and Ergin (2017) study this issue and propose an algorithm to obtain efficient matchings.

In highlighting the importance of allowing the agents to express indifferences, Erdil and Ergin (2017) say, “...allowing, even encouraging, agents to express indifferences when ranking alternatives not only simplifies preference revelation and market participation, but also improves efficiency”. They argue that indifference has importance beyond theoretical interest.²³ Incompleteness exacerbates these concerns as efficient matchings may not even exist. I have proposed alternative notions of stability that seem to be more suitable for such environments.

Lastly, the definition of the compromise core draws on a rich tradition of uncovered sets in social choice theory. The basic principle therein (adopted in this paper) is to construct a transitive order using the covering relation using the underlying intransitive order. Besides the Miller covering (used in this paper), there are several other notions of covering, e.g Gillies (1959), Fishburn (1977), Bordes (1983), McKelvey (1986).²⁴ Fishburn (1977) has also been found useful in problems of choice theory, e.g. Nishimura (2018). In fact, Dutta and Laslier (1999) offer a more general treatment of the problem of constructing a transitive order on a choice set that goes beyond covering using “comparison functions” that have the uncovered sets as a special case.

7. Conclusion

I study a two-sided matching market in which agents may have incomplete preferences. Incompleteness is pervasive and, especially in the context of matching, it has important consequences for stability. Indeed, choice-theoretically, incompleteness is indistinguishable from intransitive indifference. I prefer not to take a strong stand on which interpretation is more reasonable, as the main contribution in this paper—formulation of the compromise core and the men- and women-optimal core—is equally valid under either interpretation.

The goal of this paper has been largely exploratory, in the sense that I have attempted to study the effects of agents having incomplete preferences on the usual notions of stability. I hope that concepts such as the compromise core offer a good candidate for stability that is more permissive than the strong core and more restrictive than the weak core. At the same time, should one find

²³For example, Abdulkadiroğlu et al. (2009) show empirically that the tie-breaking rule in a school choice mechanism has significant welfare implications.

²⁴These, and many others, are extensively studied in Duggan (2013).

my proposals reasonable, a natural next step is to seek efficient algorithms that produce a matching in the compromise core, or in the men-optimal core. When the strong core is non-empty, [Irving \(1994\)](#) provides such an algorithm. Therefore, it is natural to explore such algorithms when the strong core is empty.

Equally importantly, I have abstracted away from strategic considerations. Again, from a practical standpoint, these are of paramount importance. Therefore, another worthwhile goal would be to explore strategy-proof mechanisms (for the proposing side), such as the DAA, which produce, say, matchings in the men-optimal core.

References

- A. Abdulkadiroğlu, P. A. Pathak, and A. E. Roth. Strategy-proofness versus efficiency in matching with indifference: Redesigning the NYC high school match. *American Economic Review*, 99(5):1954–78, 2009. [18](#)
- A. Atay, A. Mauleon, and V. Vannetelbosch. A bargaining set for roommate problems. *Journal of Mathematical Economics*, 94:102465, 2021. [14](#)
- R. J. Aumann. Utility theory without the completeness axiom. *Econometrica: Journal of the Econometric Society*, pages 445–462, 1962. [1](#)
- O. Aygün and T. Sönmez. Matching with contracts: Comment. *American Economic Review*, 103(5):2050–51, 2013. [17](#)
- H. Aziz, P. Biró, T. Fleiner, S. Gaspers, and R. d. Haan. Stable matching with uncertain pairwise preferences. 2017. [17](#)
- S. Bade. Pareto-optimal matching allocation mechanisms for boundedly rational agents. *Social Choice and Welfare*, 47(3):501–510, 2016. [17](#)
- S. Baswana, P. P. Chakrabarti, V. Kamakoti, Y. Kanoria, A. Kumar, U. Patange, and S. Chandran. Joint seat allocation: An algorithmic perspective, 2015. [16](#)
- S. Baswana, P. P. Chakrabarti, S. Chandran, Y. Kanoria, and U. Patange. Centralized admissions for engineering colleges in india. *Interfaces, INFORMS*, 49(5):338–354, 2019. [1](#)
- T. Bewley. Knightian uncertainty theory: Part i. Technical report, Cowles Foundation Discussion Paper, 1986. [1](#)
- G. Bordes. On the possibility of reasonable consistent majoritarian choice: Some positive results. *Journal of Economic Theory*, 31(1):122–132, 1983. [10](#), [18](#)

- C. P. Chambers and M. B. Yenmez. Choice and matching. *American Economic Journal: Microeconomics*, 9(3):126–47, 2017. 17
- Y.-K. Che, J. Kim, and F. Kojima. Weak monotone comparative statics. *Available at SSRN 3486620*, 2019. 5, 17
- Á. Cseh and A. Juhos. Pairwise preferences in the stable marriage problem. *arXiv preprint arXiv:1810.00392*, 2018. 17
- J. Duggan. Uncovered sets. *Social Choice and Welfare*, 41(3):489–535, 2013. 10, 18
- B. Dutta. Covering sets and a new condorcet choice correspondence. *Journal of Economic Theory*, 44(1):63–80, 1988. 9
- B. Dutta and J.-F. Laslier. Comparison functions and choice correspondences. *Social Choice and Welfare*, 16(4):513–532, 1999. 18
- B. Dutta and J. Massó. Stability of matchings when individuals have preferences over colleagues. *journal of economic theory*, 75(2):464–475, 1997. 17
- F. Echenique and M. B. Yenmez. A solution to matching with preferences over colleagues. *Games and Economic Behavior*, 59(1):46–71, 2007. 17
- K. Eliaz and E. A. Ok. Indifference or indecisiveness? Choice-theoretic foundations of incomplete preferences. *Games and economic behavior*, 56(1):61–86, 2006. 5, 13, 17
- A. Erdil and H. Ergin. Two-sided matching with indifferences. *Journal of Economic Theory*, 171:268–292, 2017. 15, 18
- P. C. Fishburn. Condorcet social choice functions. *SIAM Journal on applied Mathematics*, 33(3):469–489, 1977. 4, 10, 18
- D. Gale and L. S. Shapley. College admissions and the stability of marriage. *The American Mathematical Monthly*, 69(1):9–15, 1962. 2
- D. B. Gillies. Solutions to general non-zero-sum games. *Contributions to the Theory of Games*, 4:47–85, 1959. 10, 18
- J. W. Hatfield and P. R. Milgrom. Matching with contracts. *American Economic Review*, 95(4):913–935, 2005. 17
- R. W. Irving. Stable marriage and indifference. *Discrete Applied Mathematics*, 48(3):261–272, 1994. 17, 19

- R. W. Irving, D. F. Manlove, and S. Scott. Strong stability in the hospitals/residents problem. In *Annual Symposium on Theoretical Aspects of Computer Science*, pages 439–450. Springer, 2003. 17
- M. Kitahara and Y. Okumura. Stable mechanisms in controlled school choice. *Available at SSRN 3806916*, 2021. 17
- R. D. Luce and H. Raiffa. *Games and decisions: Introduction and critical survey*. Courier Corporation, 1989. 4
- D. F. Manlove. The structure of stable marriage with indifference. *Discrete Applied Mathematics*, 122(1-3):167–181, 2002. 17
- D. F. Manlove, R. W. Irving, K. Iwama, S. Miyazaki, and Y. Morita. Hard variants of stable marriage. *Theoretical Computer Science*, 276(1-2):261–279, 2002. 17
- A. Mauleon, V. J. Vannetelbosch, and W. Vergote. von neumann–morgenstern farsightedly stable sets in two-sided matching. *Theoretical Economics*, 6(3):499–521, 2011. 8
- R. D. McKelvey. Covering, dominance, and institution-free properties of social choice. *American Journal of Political Science*, pages 283–314, 1986. 18
- N. R. Miller. A new solution set for tournaments and majority voting: Further graph-theoretical approaches to the theory of voting. *American Journal of Political Science*, pages 68–96, 1980. 4, 10
- H. Nishimura. The transitive core: Inference of welfare from nontransitive preference relations. *Theoretical Economics*, 13(2):579–606, 2018. 13, 18
- E. A. Ok et al. Utility representation of an incomplete preference relation. *Journal of Economic Theory*, 104(2):429–449, 2002. 1, 17
- M. Pycia. Stability and preference alignment in matching and coalition formation. *Econometrica*, 80(1):323–362, 2012. 17
- M. Pycia and M. B. Yenmez. Matching with externalities. *University of Zurich, Department of Economics, Working Paper*, (392), 2021. 17
- A. Roth and M. Sotomayor. Sotomayor,(1990). two-sided matching: A study in game-theoretic modeling and analysis. *Econometric Society Monographs*, 18, 1992. 4
- H. Sasaki and M. Toda. Two-sided matching problems with externalities. *Journal of Economic Theory*, 70(1):93–108, 1996. 17

J. von Neumann and O. Morgenstern. Morgenstern, O. *Theory of games and economic behavior*, 3, 1944. 13

A. Appendix

Proof of Proposition 1. If μ is in the weak core, it is obviously weakly stable. For the opposite, suppose μ is weakly stable but is not in the weak core. Therefore, there is $S \subset A$ and a matching μ' such that $\mu'(i) \in S$ for all $i \in S$ and, $\mu'(i) \succ_i \mu(i) \forall i \in S$. Pick any $i \in S$ and let $j := \mu'(i)$. Then $j \succ_i \mu(i)$ and $i \succ_j \mu(j) \neq i$. Therefore, (i, j) is a strong blocking pair for μ , a contradiction. Therefore, μ is in the weak core.

Similarly, if μ is in the strong core then it is strongly stable. For the reverse, suppose μ is strongly stable but there is $S \subset A$ and a μ' such that $\mu'(i) \in S$ for all $i \in S$ and, either $\mu'(i) \succ_i \mu(i)$ or $\mu'(i) \otimes_i \mu(i)$ for all $i \in S$. Moreover, at least for one $i \in S$, $\mu'(i) \succ_i \mu(i)$. Let i^* be such an agent. But then, $(i^*, \mu'(i^*))$ is a weak blocking pair, a contradiction. Therefore, μ is in the strong core. \square

Proof of Proposition 2. Suppose μ is stable according to some completion $\widehat{\succ}$. Notice that any blocking pair in \succ remains to be a blocking pair according to $\widehat{\succ}$. Therefore, μ is weakly stable according to \succ .

For the converse, suppose μ is weakly stable. Let $C(\succ)$ denote the set of all the completions of \succ . Define,

$$T := \{(m, w) : (m, w) \text{ is a blocking pair for } \mu \text{ for some } \widehat{\succ} \in C(\succ)\}.$$

We now prove a claim towards completing the proof.

CLAIM 1: *If $(m, w) \in T$, then either $w \otimes_m \mu(w)$ or $m \otimes_w \mu(w)$, or both, according to \succ .*

Proof. Suppose not. So, $\exists(m, w) \in T$ such that m finds w and $\mu(m)$ comparable, and w finds m and $\mu(w)$ comparable. Since it is a blocking pair for some $\widehat{\succ}$, it must be the case that $w \widehat{\succ}_m \mu(m)$ and $m \widehat{\succ}_w \mu(w)$. Since w and $\mu(m)$ are comparable according to \succ_m , we have $w \succ_m \mu(m)$. Similarly, $m \succ_w \mu(w)$. Therefore, (m, w) is a strong blocking pair for μ according to \succ , i.e. μ is not weakly stable. A contradiction. \square

Now, for each $i \in T$, define $S(i) := \{j : (i, j) \in T \text{ or } (j, i) \in T \text{ and } j \otimes_i \mu(i)\}$.²⁵ Consider a completion $\widehat{\succ}^*$ of \succ such that, for all $i \in A$, $\mu(i) \widehat{\succ}_i^* j$ for all $j \in S(i)$.²⁶

²⁵The distinction between (i, j) and (j, i) is merely because we have defined a blocking pair as an ordered pair (m, w) . Therefore, whether $i \in M$ or $i \in W$ would matter for notational consistency.

²⁶That such a completion exists is a straightforward consequence of the Szpilrajn extension theorem. We define a new partial order, $\widehat{\succ}_i$ in two steps. First, for any j such that $\mu(i) \otimes_i j$, we set $\mu(i) \widehat{\succ}_i j$. Then, we take its transitive closure to make $\widehat{\succ}_i$ a partial order. Finally, by using the Szpilrajn extension theorem, we extend this partial order to a complete linear order.

We complete the proof by showing that μ is stable according to $\widehat{\succ}^*$. Suppose not. \exists a blocking pair (m, w) according to $\widehat{\succ}^*$. But then, $w \in S(m)$ or $m \in S(w)$ by Claim 1. Suppose, wlog, $w \in S(m)$. But then, by construction, $\mu(w) \widehat{\succ}_w^* m$. Therefore, (m, w) cannot be a blocking pair for μ according to $\widehat{\succ}^*$. A contradiction. Hence, μ is stable according to $\widehat{\succ}^*$. \square

Proof of Proposition 3. Suppose that μ is stable according to every completion of \succ but is not strongly stable. That means, wlog, there is a blocking pair (m, w) , such that $m \succ_w \mu(w)$ and $w \otimes_m \mu(m)$. Consider a completion, $\widehat{\succ}$, of \succ such that $w \succ_m \mu(m)$. Such a completion exists by the Szpilrajn theorem as in the proof of Proposition 2. Therefore, (m, w) is a blocking pair in μ in $\widehat{\succ}$, i.e., μ is not stable in $\widehat{\succ}$. A contradiction. Therefore, μ must be strongly stable. \square

LEMMA 1: \triangleright is a possibly incomplete and intransitive binary relation over \mathcal{M} . Also, it is possible to have $\mu \triangleright \mu' \triangleright \mu$.

Proof. That \triangleright is possibly incomplete is obvious. Consider $M = \{m_1, m_2\}, W = \{w_1, w_2\}$. Suppose that all the agents strictly prefer being matched over remaining single. Moreover, $w_1 \succ_{m_1} w_2, w_2 \succ_{m_2} w_1, m_2 \succ_{w_1} m_1$ and $m_1 \succ_{w_2} m_2$. It is easy to check $\mu = \{(m_1, w_1), (m_2, w_2)\}$ and $\mu' = \{(m_1, w_2), (m_2, w_1)\}$ are in \mathcal{W} and are not ranked according to \triangleright .

The intransitivity of \triangleright is seen in a number of examples throughout the paper, one being Example 1. \square

LEMMA 2: \ll is antisymmetric and transitive.

Proof. For transitivity, suppose that $\mu \ll \mu'$ and $\mu' \ll \mu''$. Consider any matching $\hat{\mu}$ such that $\mu'' \triangleright \hat{\mu}$. Since $\mu' \ll \mu''$, we have $\mu' \triangleright \hat{\mu}$. Since $\mu \ll \mu'$, $\mu \triangleright \hat{\mu}$. Therefore, $\mu \ll \mu''$.

For its antisymmetry, suppose that $\mu \ll \mu' \ll \mu$. Therefore, $\mu' \triangleright \mu$. Since $\mu \ll \mu'$, we have $\mu \triangleright \mu'$. Since \triangleright is not-reflexive, this is not possible. \square

Proof of Proposition 4. Suppose $\mu \in \mathcal{S}$ and there is a μ' such that $\mu' \triangleright \mu$. Therefore, there exists an $S \subset A$ such that S can enforce μ' over μ and $\mu' \succ_S \mu$. Hence, there is $i \in S$ such that $\mu' \succ_i \mu$. Moreover, for the partner of i in μ' , denoted by $j := \mu'(i)$, either $i \succ_j \mu(j)$ or $i \otimes_j \mu(j)$. But then, (i, j) is a weak blocking pair for μ , i.e. $\mu \notin \mathcal{S}$, a contradiction. \square

Proof of Theorem 1. That \mathcal{C} satisfies IM and EB is obvious.

CLAIM 2: $\mathcal{C}(\cdot)$ satisfies ET.

Proof. Suppose that, for some $\mu, \mu' \in A \subset \mathcal{W}$, $\mu' \notin \mathcal{C}(A') \forall A' \in \mathcal{A}_3(\mu, \mu', T)$ but $\mu' \in \mathcal{C}(A)$. Therefore, μ' is uncovered in A . First, suppose that $\mu \triangleright \mu'$. Since μ' is uncovered in A , $\exists \hat{\mu} \in A$ such that $\mu' \triangleright \hat{\mu}$ but not $\mu \triangleright \hat{\mu}$. But then, $\mu' \in \mathcal{C}(\{\mu, \mu', \hat{\mu}\})$. Since $\{\mu, \mu', \hat{\mu}\} \in \mathcal{A}_3(\mu, \mu', T)$, this is a contradiction. Therefore, $\mu \triangleright \mu'$ is not possible. On the other hand, if $\mu' \triangleright \mu$ or if μ and μ' are not \triangleright ranked, then $\mu' \in \mathcal{C}(\{\mu, \mu'\})$: a contradiction as $\{\mu, \mu'\} \in \mathcal{A}_3(\mu, \mu', T)$. \square

For the reverse, suppose Γ satisfies the IM, EB and ET. Let \ll^A denote the covering relation on a set A . That is, for any $\mu, \mu' \in A$, $\mu \ll^A \mu'$ if $\mu \triangleright \mu'$ and $\mu' \triangleright \hat{\mu} \implies \mu \triangleright \hat{\mu}$ for all $\hat{\mu} \in A$.

CLAIM 3: $\mu \ll^A \mu' \implies \mu' \notin \Gamma(A)$.

Proof. Since $\mu \ll^A \mu'$, we have that $\mu \triangleright \mu'$. Therefore, if $A = \{\mu, \mu'\}$, $\mu' \notin \Gamma(A)$ due to IM. Suppose $A \neq \{\mu, \mu'\}$. Consider any $\hat{\mu} \in A$ and let $A' := \{\mu, \mu', \hat{\mu}\}$. We now prove that $\mu' \notin \Gamma(A')$.

First, suppose that $\mu' \triangleright \hat{\mu}$. Notice that, $\mu \ll^A \mu' \implies \mu \triangleright \hat{\mu}$. Therefore, A' is totally ordered, and hence, by IM, $\Gamma(A') = \{\mu\}$.

Alternatively, if $\mu' \triangleright \hat{\mu}$ (and not $\mu' \triangleright \hat{\mu}$) or if μ' and $\hat{\mu}$ are not \triangleright -ranked, then $\mu'^{\uparrow} \cap A' \ni \mu$ and $\mu'^{\downarrow} \cap A' = \emptyset$. Therefore, $\mu' \notin \Gamma(A')$ by EB.

Therefore, $\mu' \notin \Gamma(A')$. However, if $\mu' \notin \Gamma(\{\mu, \mu', \hat{\mu}\})$ for any $\hat{\mu} \in A$, then $\mu' \notin \Gamma(A)$ by ET. \square

That is, if μ' is covered in A , then $\mu' \notin \Gamma(A)$. In other words, $\Gamma(A) \subset \mathcal{C}(A) \forall A \subset \mathcal{W}$. \square

Proof of Proposition 6. Suppose that $\mu \in \mathcal{W} \setminus \mathcal{C}$. Therefore, $\exists \mu_1 \ll \mu$. If $\mu_1 \in \mathcal{C}$ we are done. If not, $\exists \mu_2 \ll \mu_1 \implies \mu_2 \triangleright \mu$ (since $\mu_1 \triangleright \mu$). Since \ll is transitive and strict and \mathcal{W} is finite, $\exists \mu_n$ such that μ_n is not covered by any other matching and $\mu_n \triangleright \mu$. Since μ_n is uncovered, $\mu_n \in \mathcal{C}$ establishing the proposition. \square

EXAMPLE 4: Let $M = \{m_1, m_2, m_3\}$ and $W = \{w_1, w_2, w_3\}$. Men's preferences are the following: $w_2 \succ_{m_1} w_1 \succ_{m_1} w_3$, $w_2 \succ_{m_2} w_3, w_1 \otimes_{m_2} w_2, w_3$, $w_1 \succ_{m_3} w_3, w_2 \otimes_{m_3} w_1, w_3$. All the men strictly prefer being matched over remaining single. Women's preferences are the following: $m_1 \succ_{w_1} m_3$ and $m_2 \otimes_{w_1} m_1, m_3$. Also, w_1 strictly prefers being matched over remaining single. For w_2 , being matched to m_1 or m_2 is strictly preferred to remaining single. But, being single is strictly preferred over being matched to m_3 . She cannot compare m_1 and m_2 . Similarly, w_3 strictly prefers being matched to m_2 or m_3 over remaining single, and prefers remaining single over being matched to m_1 . She cannot compare m_2 and m_3 . \blacksquare

CLAIM: $\mathcal{W} = \{\mu_1, \mu_2, \mu_3\}$ where $\mu_1 = \{(m_1, w_1), (m_2, w_2), (m_3, w_3)\}$,

$\mu_2 = \{(m_1, w_2), (m_2, w_1), (m_3, w_3)\}$ and $\mu_3 = \{(m_1, w_2), (m_2, w_3), (m_3, w_1)\}$.

Proof. It is easy to check that μ_1, μ_2 and μ_3 are weakly stable. To rule out other matchings, first notice that matchings in which w_3 is matched to m_1 or m_3 is matched to w_2 are not in the weak core. That rules out the remaining matches wherein all the agents are matched. Second, if there is a matching in \mathcal{W} wherein one man-woman pair is single it can either involve w_2 and m_3 being single or w_3 and m_1 being single. Suppose w_2 and m_3 are single. Then, either we have $\{(m_1, w_1), (m_2, w_3)\}$ or $\{(m_1, w_3), (m_2, w_1)\}$ as the possible matches. In the first matching, (m_2, w_2) constitute a strong blocking pair while in the second, (m_1, w_3) is ruled out as w_3 prefers remaining single over being matched to m_1 . Similarly, it is easy to check that matchings wherein m_1 and w_3 are weakly unstable too. Lastly, it is obvious that there cannot be a weakly stable matching with two or more men unmatched. □

CLAIM: The Fishburn uncovered set in Example 4 is $\{\mu_1, \mu_3\}$ while $\mathcal{C} = \{\mu_2, \mu_3\}$.

Proof. It is easy to check that $\mu_2 \triangleright \mu_1$ and $\mu_3 \triangleright \mu_2$. However, μ_1 and μ_3 are not \triangleright comparable. Therefore, $\mathcal{C} = \{\mu_3, \mu_2\}$ as $\mu_2 \triangleright \mu_1$ but not $\mu_3 \triangleright \mu_1$.

However, if we used Fishburn covering instead of Miller covering, then μ_2 is covered by μ_3 . Recall that x F-covers y if, $z \triangleright x \implies z \triangleright y$. Since no matching \triangleright -dominates μ_3 , μ_3 F-covers μ_2 trivially. On the other hand, μ_1 is F-uncovered. Therefore, the Fishburn uncovered set here is $\{\mu_1, \mu_3\}$. □

A.1. Example illustrating the compromise core

EXAMPLE 5: Let $M = \{m_1, m_2, m_3, m_4\}$ and $W = \{w_1, w_2, w_3, w_4\}$. Consider the preferences of men given by the Hasse diagram in Figure 1.




Figure 1: Preferences represented via Hasse diagram

Moreover, w_1 and w_4 strictly prefer being matched to remaining single, but find every man incomparable.

The Hasse diagram offers a succinct way of representing preferences. For the reader unfamiliar with it, here are the preferences. For $m_1, w_1, w_4 \succ_{m_1} w_2 \succ_{m_1} w_3$ and $w_1 \otimes_{m_1} w_4$. For $m \in \{m_2, m_3, m_4\}$, we have, $w_2 \succ_m w_3 \succ_m w_4, w_1 \succ_m w_4, w_1 \otimes_m w_2$ and $w_1 \otimes_m w_3$. For $w \in \{w_2, w_3\}$, $m_2, m_3, m_4 \succ_w m_1$ and $m_i \otimes_w m_j$ if $i \neq j$ and $i, j \in \{2, 3, 4\}$. For $w \in \{w_1, w_4\}$, $m_i \otimes_w m_j$ iff $i \neq j$ and $i, j \in \{1, 2, 3, 4\}$, and $m_i \succ_w w$ for all $m_i \in M$.

It is straightforward to see that the strong core is empty, and the weak core contains all possible matchings such that $\mu(m_1) \in \{w_1, w_4\}$; and no agent is unmatched. However, there is a sense any matching $\mu \in \mathcal{W}$, such that $\mu(m_1) = w_4$ are better than those where $\mu(m_1) = w_1$. Indeed, both of these types of matchings are in the weak core. However, when $\mu(m_1) = w_1, \mu(\{m_2, m_3, m_4\}) = \{w_2, w_3, w_4\}$.²⁷ On the other hand, when $\mu(m_1) = w_4, \mu(\{m_2, m_3, m_4\}) = \{w_1, w_2, w_3\}$. Therefore, given any matching in the weak core that matches m_1 with w_1 , all the agents could decide to move to another matching in the weak core, where m_1 is matched with w_4 . This way, it is possible to make some agent strictly better off without making any agent worse off. More importantly, a similar move in the opposite direction—from m_1 being matched with w_4 to m_1 being matched with w_1 —would always make some agent strictly worse off. We now see that the compromise core would select precisely those matchings from the weak core where $\mu(m_1) = w_4$.

LEMMA 3: *In Example 5, $\mathcal{C} = \{\mu \in \mathcal{W} : \mu(m_1) = w_4\}$.*

Proof. Below, we enumerate all the matchings in \mathcal{W} , such that $\mu(m_1) = w_1$.

$$\begin{aligned}
\mu_1 &= \{(m_1, w_1), (m_2, w_2), (m_3, w_3), (m_4, w_4)\} \\
\mu_2 &= \{(m_1, w_1), (m_2, w_2), (m_3, w_4), (m_4, w_3)\} \\
\mu_3 &= \{(m_1, w_1), (m_2, w_3), (m_3, w_2), (m_4, w_4)\} \\
\mu_4 &= \{(m_1, w_1), (m_2, w_3), (m_3, w_4), (m_4, w_2)\} \\
\mu_5 &= \{(m_1, w_1), (m_2, w_4), (m_3, w_2), (m_4, w_3)\} \\
\mu_6 &= \{(m_1, w_1), (m_2, w_4), (m_3, w_3), (m_4, w_2)\}
\end{aligned}$$

²⁷Recall that $\mu(S) := \{j : j = \mu(i) \text{ for some } i \in S\}$.

Now, we enumerate all the matchings in \mathcal{W} , such that $\mu(m_1) = w_4$.

$$\begin{aligned}\mu_7 &= \{(m_1, w_4), (m_2, w_2), (m_3, w_3), (m_4, w_1)\} \\ \mu_8 &= \{(m_1, w_4), (m_2, w_2), (m_3, w_1), (m_4, w_3)\} \\ \mu_9 &= \{(m_1, w_4), (m_2, w_3), (m_3, w_2), (m_4, w_1)\} \\ \mu_{10} &= \{(m_1, w_4), (m_2, w_3), (m_3, w_1), (m_4, w_2)\} \\ \mu_{11} &= \{(m_1, w_4), (m_2, w_1), (m_3, w_2), (m_4, w_3)\} \\ \mu_{12} &= \{(m_1, w_4), (m_2, w_1), (m_3, w_3), (m_4, w_2)\}\end{aligned}$$

Figure 2 depicts the relation \supseteq in a graph with each node being a matching in \mathcal{W} .²⁸ An edge from μ to μ' means that $\mu' \supseteq \mu$.




Figure 2: Graph with nodes as matchings in \mathcal{W} .

As can be seen, matchings from μ_1 to μ_6 are covered as they do not dominate any other matching. On the other hand, μ_7 to μ_{12} are not covered. For example, $\mu_8 \supseteq \mu_7$ and $\mu_7 \supseteq \mu_9$. However, we do not have $\mu_8 \supseteq \mu_9$. Therefore,

$$\mathcal{C} = \{\mu_7, \mu_8, \mu_9, \mu_{10}, \mu_{11}, \mu_{12}\}.$$

□

²⁸To avoid cluttering the graph, I have chosen to avoid some edges—for example, an edge from μ_1 to μ_{11} . More importantly, there is no edge from μ_i to μ_j if $i \geq 7$ and $j \leq 6$. Also, there is no edge from $\{\mu_7, \mu_8, \mu_9\}$ to $\{\mu_{10}, \mu_{11}, \mu_{12}\}$.