The Uplifton

Iosif Bena¹ and Pierre Heidmann²

¹Institut de Physique Théorique, Université Paris Saclay, CEA, CNRS, F-91191 Gif-sur-Yvette, France ²Department of Physics and Astronomy, Johns Hopkins University, Baltimore, MD 21218, USA

Almost all proposals to construct de Sitter vacua with a small cosmological constant involve flux compactifications with stabilized moduli. These give AdS vacua, which are uplifted to de Sitter by adding antibranes in certain regions of the compactification manifold. However, antibranes are charged, singular and interact nontrivially with other ingredients of the compactification; this can invalidate the de Sitter construction. In this Letter, we construct a new ingredient for uplifting AdS solutions to de Sitter, which is neutral, smooth and horizonless, and therefore bypasses some of the problems of antibrane uplift.

1. INTRODUCTION

The accelerated expansion of our Universe points to the existence of a positive vacuum energy. However, String Theory appears rather reluctant to provide fourdimensional solutions with a positive vacuum energy. Even the simplest solution with positive vacuum energy - de Sitter space – is very hard to construct. There are no-go theorems preventing the direct realization of such a space via a compactification with common ingredients [1]. The most popular scenario to bypass these theorems and construct de Sitter spaces with a small cosmological constant, proposed by Kachru, Kallosh, Linde and Trivedi [2], involves a rather intricate sequence of steps: one first stabilizes the complex-structure moduli by turning on topologically-nontrivial fluxes on the compactification manifold and then one stabilizes the Kähler moduli via nonperturbative effects. This results in an Anti de Sitter solution (with a negative cosmological constant) which should be "uplifted" to a de Sitter solution (with a positive cosmological constant) by placing D3 branes with negative charge (antibranes) in a region of large warping inside the compactification manifold.

This last step has been under a lot of intense scrutiny over the past fifteen years, because the $\overline{D3}$ branes interact non-trivially with the fluxes used to stabilize the moduli [3]. This can result in tachyons, unexpected massless modes and runaways [4, 5]. Furthermore, the interaction of the four-form field sourced by the antibrane with the other fields of the compactification can give rise to new flux components, that can affect the regime of validity of Kähler moduli stabilization [6].

The purpose of this Letter is to construct a new ingredient for uplifting the cosmological constant - an uplifton - which is neutral, smooth and horizonless, thereby avoiding some of the problems of antibranes. Furthermore, unlike antibranes [7], the uplifton cannot move around the compactification manifold.

At first glance, neutral solutions which have mass but no charge do not appear to be optimal ingredients for use in flux compactifications, because they naïvely source a metric that does not preserve Lorentz invariance in the spacetime directions. The best example is perhaps a nonextremal black D3 brane, which does have more mass than charge, but whose metric breaks the Lorentz symmetry along the spacetime direction.¹ However, as we will see, the non-extremal solutions we construct preserve this Lorentz invariance through a novel mechanism which involves the shrinking of certain compact directions.

To construct these solutions, we use the formalism that has been developed over the past few years by one of the authors and Bah [8–13]: the equations governing certain supergravity solutions with D-2 commuting Killing vectors and suitable fluxes decompose into a set of Ernst equations, thereby admitting an integrable structure. This formalism has allowed to obtain a plethora of solutions, describing both bound states of bubbles and black holes, as well as smooth horizonless solutions with multiple bubbles and topologically non-trivial fluxes. Some of these solutions are non-extremal and charged, but it is also possible to construct neutral solutions with opposite fluxes wrapping different cycles [12].

At first glance, the simplest way to construct an uplifton appears to be using smooth bubbles with D3-brane and $\overline{D3}$ brane charges. However, the mechanism by which the solutions of [12] carry charges involves topologicallynontrival cycles formed by the shrinking of at least one direction inside the brane worldvolume. Hence, bubbling solutions whose bubbles have D3 and $\overline{D3}$ charges break the SO(3, 1) Lorentz invariance.

Since we are looking for upliftons that one can add to Type IIB flux compactifications, the obvious step to bypass this problem is to use the technique of [12] to construct neutral solutions with D5 and $\overline{\text{D5}}$ bubbles that preserve the SO(3, 1) invariance.

In this Letter we present the simplest of these solutions and their use in flux compactifications, leaving the details of their construction to a companion paper. Even if the upliftons are neutral, they have a nontrivial magnetic three-form field strength profile which is both positive and negative, so locally they have D5 and $\overline{\text{D5}}$ charge corresponding to branes extending along $(t, x_1, x_2, x_3, x_4, x_5)$, where x_5 must be compact. The orthogonal space is a U(1) fibration of a compact coordinate, y, over a three-dimensional base given in spherical

¹ The time-time and the parallel space-space components of this metric are different functions of the radius.



FIG. 1: Schematic description of the uplifton internal directions. The spacetime smoothly terminates at $r = \ell + 2\sigma$ where the y and x_5 circles degenerate alternatively as smooth bolts. The x_5 -bolts carry opposite D5 brane charges.

coordinates (r, θ, ϕ) . In a flux compactification, both x_4 and x_5 , together with y, r, θ, ϕ will be part of the compactification manifold. As we will show, regularity will impose certain periodicity constraints on the x_5 and ycoordinates, so, unlike $\overline{D3}$ branes, the uplifton will not be able to move inside the compactification manifold.

2. THE UPLIFTON SOLUTION

The six-dimensional spacetime transverse to the directions of the uplifton, (t, x_1, x_2, x_3) , is made from three compact circles fibered over a three-dimensional space that becomes asymptotically \mathbb{R}^3 . Using the technology of [12] we can build upliftons with an arbitrary number of bubbles, but for simplicity we will present here the simplest upliftons. They have two identical bubbles car $\mathbf{2}$

rying opposite D5 charges, and located at the opposite ends of an uncharged bubble (see Fig.1). The solutions are determined in terms of three parameters (ℓ, m, q) : ℓ and m are related to the mass, or energy, induced by the sources, while q is related to the amplitude of the D5 charges carried by the outermost bubbles.

Being a bound state of three sources, we introduce three sets of local spherical coordinates, centered around each bubble:

$$4r_{1} \equiv r_{-}^{(0)} + r_{-}^{(1)} - 4\sigma, \quad \cos\theta_{1} \equiv \frac{r_{-}^{(0)} - r_{-}^{(1)}}{4\sigma},$$

$$4r_{2} \equiv r_{-}^{(1)} + r_{+}^{(1)} - 2(\ell - 2\sigma), \quad \cos\theta_{2} \equiv \frac{r_{-}^{(1)} - r_{+}^{(1)}}{2(\ell - 2\sigma)},$$

$$4r_{3} \equiv r_{+}^{(1)} + r_{+}^{(0)} - 4\sigma, \quad \cos\theta_{3} \equiv \frac{r_{+}^{(1)} - r_{+}^{(0)}}{4\sigma}, \quad (1)$$

where we have defined 2σ and $\ell - 2\sigma$ to be the size of the outermost bubbles and the middle bubble respectively, and the distance $(r_{\pm}^{(0)}, r_{\pm}^{(1)})$, related to the \mathbb{R}^3 spherical coordinates (r, θ) such as

$$\sigma \equiv \sqrt{m^2 - q(q - 2\gamma)}, \qquad \gamma \equiv \frac{2mq}{\ell + 2m},$$

$$r_{\pm}^{(0)} \equiv 2r - (\ell + 2\sigma)(1 \pm \cos\theta), \qquad (2)$$

$$r_{\pm}^{(1)} \equiv \sqrt{((2(r - \sigma) - \ell)\cos\theta \pm (2\sigma - \ell))^2 + 4r(r - \ell - 2\sigma)\sin^2\theta}.$$

These solutions exist when the parameters satisfy [12]:

$$\ell > 2m$$
, $q < m\sqrt{\frac{\ell+2m}{\ell-2m}}$. (3)

When the second bound is saturated, the outer bubbles degenerate into two singular five-brane sources of opposite charges. Hence, in this extreme regime, the solution can be thought of point-like D5 and $\overline{\text{D5}}$ branes at the opposite ends of a bolt.

The string-frame type IIB uplifton solution is given by

$$ds_{10}^{2} = \frac{1}{\sqrt{Z}} \left[-dt^{2} + dx_{1}^{2} + dx_{2}^{2} + dx_{3}^{2} + dx_{4}^{2} + \frac{r_{1}r_{3}}{(r_{1} + 2\sigma)(r_{3} + 2\sigma)} dx_{5}^{2} \right] + \sqrt{Z} \left[f \left(\frac{dr^{2}}{1 - \frac{\ell + 2\sigma}{r}} + r^{2} d\theta^{2} \right) + r^{2} \sin^{2} \theta d\phi^{2} + \frac{r_{2}}{r_{2} + \ell - 2\sigma} dy^{2} \right]$$
(4)
$$C^{(2)} = Hd\phi \wedge dy , \qquad e^{\Phi} = \frac{1}{\sqrt{Z}} , \qquad B_{2} = C^{(0)} = C^{(4)} = 0.$$

where the definitions of r_1, r_2, r_3 are in equation (1), and we have introduced the following warp factors and gauge potentials

$$Z = \frac{(r_1 + \sigma + m)(r_3 + \sigma + m) + (q - \gamma(1 + \cos\theta_3))(q - \gamma(1 - \cos\theta_1))}{\sqrt{\left((r_1 + 2\sigma)^2 + \gamma^2 \sin^2\theta_3 \left(1 + \frac{2\sigma}{r_1}\right)\right)\left((r_3 + 2\sigma)^2 + \gamma^2 \sin^2\theta_1 \left(1 + \frac{2\sigma}{r_3}\right)\right)}},$$

$$f^{4} = \frac{(r_{1}(r_{1}+2\sigma)+\gamma^{2}\sin^{2}\theta_{3})(r_{3}(r_{3}+2\sigma)+\gamma^{2}\sin^{2}\theta_{1})}{(1+2\delta)^{2}(r_{1}+\sigma(1-\cos\theta_{1}))^{4}(r_{3}+\sigma(1+\cos\theta_{3}))^{4}} \left(1+2\delta\frac{(q-\gamma)(r_{1}-r_{3})+(\gamma m-\ell(q-\gamma))(\cos\theta_{1}+\cos\theta_{3})}{(q-\gamma)(r_{3}-r_{1})+\gamma m(\cos\theta_{1}+\cos\theta_{3})}\right)^{2}}{\chi} \frac{r_{1}^{2}r_{2}^{4}r_{3}^{2}(r_{2}+\ell-2\sigma)^{2}(r-\ell-2\sigma)^{-2}}{\left(\left(r_{2}+\frac{\ell}{2}-\sigma(1-\cos\theta_{1})\right)^{2}-\frac{\ell^{2}}{4}\right)\left(\left(r_{2}+\frac{\ell}{2}-\sigma(1+\cos\theta_{3})\right)^{2}-\frac{\ell^{2}}{4}\right)}, \qquad \delta \equiv \frac{m^{2}(\ell+2m)^{2}+\ell^{2}q^{2}}{(\ell+2m)^{2}(\ell^{2}-2m^{2})+2\ell^{2}q^{2}}, \\ H = \frac{(q-\gamma)(r_{3}-r_{1})+\gamma m(\cos\theta_{1}+\cos\theta_{3})}{(r_{1}+\sigma+m)(r_{3}+\sigma+m)+(q-\gamma(1+\cos\theta_{3}))(q-\gamma(1-\cos\theta_{1}))}\left[\frac{(r_{1}+\sigma)\cos\theta_{1}+(r_{3}+\sigma)\cos\theta_{3}}{2} - \frac{\ell}{2}\left(1+\frac{\gamma^{2}}{m}\right)\frac{(r_{3}+\sigma+m)(2m\cos\theta_{1}+\ell)-(r_{1}+\sigma+m)(2m\cos\theta_{3}-\ell)}{\ell(r_{1}-r_{3})+2(q-\gamma)^{2}(\cos\theta_{1}+\cos\theta_{3})}\right]$$

+
$$2m(q-\gamma)\frac{2(r_1-r_3)-\ell(\cos\theta_1+\cos\theta_3)}{\ell(r_1-r_3)+2(q-\gamma)^2(\cos\theta_1+\cos\theta_3)}$$

The spacetime is smooth and terminates at $r = \ell + 2\sigma$. At this locus, either r_1 or r_2 or r_3 is zero, depending on the value of θ in terms of three intervals. These intervals are determined by the critical angle, θ_c , defined as:

$$\cos\theta_c \equiv \frac{\ell - 2\sigma}{\ell + 2\sigma}.$$
 (6)

For $0 \leq \theta \leq \theta_c$ and $\pi - \theta_c \leq \theta \leq \pi$, $r_3 = 0$ and $r_1 = 0$ respectively, such that x_5 degenerates at the origin. For $\theta_c \leq \theta \leq \pi - \theta_c$, $r_2 = 0$ and the *y* coordinate degenerates. As we will see in the next section, these coordinate degeneracies correspond to smooth bolts only if x_5 and *y* are compact.² Thus, the maximal Lorentz invariance our solutions can preserve is SO(4, 1), when (x_1, x_2, x_3, x_4) are infinite. However, we can also compactify one of these directions $(x_4$ for example) to obtain a more general solution which only preserves SO(3, 1) and which can be embedded in a flux compactification.

3. THE GLUING PROPERTIES

To analyse the regularity of the uplifton, it is useful to denote the periodicities of (y, x_4, x_5) as R_y, R_{x_4} and R_{x_5} :

$$y = y + 2\pi R_y$$
, $x_4 = x_4 + 2\pi R_{x_4}$, $x_5 = x_5 + 2\pi R_{x_5}$.
(7)

As we explained above, the shrinking of the y and x_5 coordinates at the origin $(r = \ell + 2\sigma)$ depends on the three θ -intervals $0 \leq \theta_c \leq \pi - \theta_c \leq \pi$. The local coordinates adapted for the first, second and third interval respectively, are (r_3, θ_3) , (r_2, θ_2) and (r_1, θ_1) (1). They allow us to write the constant-time slices of the metrics when each $r_i \to 0$ at the origin as

$$ds_{10}^2\Big|_{dt=0} \propto \frac{dr_i^2}{r_i} + \frac{4r_i}{C_i} dX^2 + ds(\mathcal{K}_7)^2, \qquad (8)$$

where C_i are constants that depends on (ℓ, m, q) . The periodic coordinate, X stands for x_5 when i = 1, 3 and for y when i = 2, while \mathcal{K}_7 describes a smooth orthogonal space of topology $S^3 \times S^1 \times \mathbb{R}^3$ when i = 1, 3 and $S^2 \times T^2 \times \mathbb{R}^3$ when i = 2. The bolt structure is explicit in terms of the radial coordinates $\rho_i^2 \equiv 4r_i$ and the \mathbb{R}^2 has no conical singularity if $R_X^2 = C_i$. This requires:

$$R_y^2 = \frac{4(\ell^2 - 4m^2)(\ell^2 - 4\sigma^2)}{\ell^2},$$

$$R_{x_5}^2 = \frac{4(\ell + 2m)(\ell + 2\sigma)(m + \sigma)}{\ell}.$$
(9)

Moreover, the three-form field strength is regular everywhere, and its integrals on the first and third bolts are equal and opposite. They give the D5 and $\overline{\text{D5}}$ quantized charges carried by these bolts:

Ì

$$N_{\rm D5} = \frac{1}{4\pi^2 g_s l_s^2} \int_{\theta_1 \phi_y} F_3 \big|_{r_1=0} = \frac{2R_y q}{g_s l_s^2},$$

$$N_{\rm \overline{D5}} = -\frac{1}{4\pi^2 g_s l_s^2} \int_{\theta_3 \phi_y} F_3 \big|_{r_3=0} = \frac{2R_y q}{g_s l_s^2},$$
(10)

where g_s is the string coupling and l_s is the string length. One can absorb the $\sqrt{g_s} l_s$ coefficient by expressing all length scales in units of $\sqrt{g_s} l_s$. This is done by rescaling

$$(\ell, m, q, \sigma, R_X) \equiv \sqrt{g_s} \, l_s \times (\bar{\ell}, \bar{m}, \bar{q}, \bar{\sigma}, \bar{R}_X). \tag{11}$$

Our solutions have three parameters, and two regularity constraints; it is natural to choose the free parameter to be the quantized charge of the bubbles, $N = N_{D5} =$ $N_{\overline{D5}} = 2\bar{R}_y\bar{q}$. The parameters of the regular solutions are therefore completely determined by the periodicities of y and x_5 at infinity and by the D5 quantized charge:

$$\bar{m} = \frac{1}{4\bar{R}_{x_5}} \left(\frac{\bar{R}_{x_5}^3}{\sqrt{4\bar{R}_{x_5}^2 + \bar{R}_y^2}} + \frac{N^2}{\sqrt{4N^2 + \bar{R}_{x_5}^2 \bar{R}_y^2}} \right),$$

$$\bar{\ell} = 2\bar{m} + \frac{\bar{R}_y^2}{4} \left(\frac{1}{\sqrt{4\bar{R}_{x_5}^2 + \bar{R}_y^2}} + \frac{\bar{R}_{x_5}}{\sqrt{4N^2 + \bar{R}_{x_5}^2 \bar{R}_y^2}} \right),$$

$$\bar{\sigma} = \frac{1}{4\bar{R}_{x_5}} \left(\frac{\bar{R}_{x_5}^3}{\sqrt{4\bar{R}_{x_5}^2 + \bar{R}_y^2}} - \frac{N^2}{\sqrt{4N^2 + \bar{R}_{x_5}^2 \bar{R}_y^2}} \right),$$
(12)

² This is a general feature of solutions constructed using the procedure of [8–13], and the reason why one cannot construct a bound state with D3 and $\overline{\text{D3}}$ charges without having to compactify one of the internal D3 directions and break the SO(3,1) Lorentz invariance.

and the validity bound (3) translates into

$$0 \le N < \bar{R}_{x_5}^2. \tag{13}$$

The solutions have two simple limits. When N = 0, the flux of the solution is strictly zero, and the solution becomes a pure-gravity solution describing three colinear vacuum bolts. When $N = \bar{R}_{x_5}^2$ the size of the outwards bubbles becomes zero ($\sigma = 0$) and these bubbles degenerate to singular locally-supersymmetric D5 and $\overline{\text{D5}}$ branes on a vacuum bubble.

4. UPLIFTING WITH THE UPLIFTON

In order to use the uplifton for uplifting we have to embed it in flux compactifications, and compare its energy with that of other uplifting ingredients, such as fivebranes or anti-D3 branes. Our solution has three compact circles and one can consider adding it to a region of the compactification manifold where the geometry looks locally like a $U(1)^3$ fibration over \mathbb{R}^3 . The mass of this solution is then completely determined by the size of the three U(1)'s in this region and by the quantized fivebrane and anti-five-brane charges of the bubbles.

To compute the mass, we reduce the uplifton along $x_{1,2,3,4,5}, y$, and obtain a geometry with $\mathbb{R}^{3,1}$ asymptotics.³

$$ds_4^2 = -\sqrt{\frac{1 - \frac{\ell + 2\sigma}{r}}{Z}} dt^2 + \sqrt{Z \left(1 - \frac{\ell + 2\sigma}{r}\right)} ds_3^2, \quad (14)$$

where ds_3^2 is the three-dimensional base in the bracket of (4). The ADM mass per unit of spacetime volume (parameterized by x_1, x_2 and x_3) is⁴

$$\mathcal{M} = \frac{\ell + 2m}{4G_4} = \frac{\bar{R}_{x_4}\bar{R}_{x_5}\bar{R}_y(\bar{\ell} + 2\bar{m})}{4\pi^3 l_s^4}$$
(15)
$$= \frac{\bar{R}_{x_4}\bar{R}_y\left(\bar{R}_{x_5}\sqrt{4\bar{R}_{x_5}^2 + \bar{R}_y^2} + \sqrt{4N^2 + \bar{R}_{x_5}^2\bar{R}_y^2}\right)}{16\pi^3 l_s^4}.$$

This mass formula is very illustrative. First, we can see that the mass remains finite when N = 0. So the uplifton can be thought of as a topological soliton of mass

$$\mathcal{M}_{\text{topo}} = \frac{\bar{R}_{x_4} \bar{R}_y \bar{R}_{x_5} \left(\sqrt{4\bar{R}_{x_5}^2 + \bar{R}_y^2} + \bar{R}_y \right)}{16\pi^3 l_s^4} \,, \qquad (16)$$

to which one adds fluxes corresponding to D5 charges. Remembering the factors of g_s in the definition of $\bar{R}_{x_4}, \bar{R}_{x_5}$ and \bar{R}_y (11), we can see that the mass of this soliton is proportional to g_s^{-2} , exactly as one expects for a gravitational soliton. Furthermore, since in this soliton the x_4 direction is not fibered, but x_5 and y are, the dependence of the mass on \bar{R}_{x_4} , \bar{R}_{x_5} and \bar{R}_y when the other radii are kept fixed is $(\bar{R}_{x_4})^1$, $(\bar{R}_{x_5})^2$ and $(\bar{R}_y)^2$, again as one expects.

Furthermore, the second square root of (15) looks exactly like the mass of a bound state of a topological soliton with 2N objects of mass proportional to g_s^{-1} . Hence, one may naïvely conclude that the side bubbles of the uplifton are bound states of D5 branes and topological solitons. However, this is not what happens: the ADM mass of 2N D5 branes wrapping $x_{1,2,3,4,5}$ is

$$\mathcal{M}_{\rm BPS} = 2 \times \frac{q}{4G_4} = \frac{NR_{x_4}R_{x_5}}{4\pi^3 l_s^4},\tag{17}$$

while the $N \gg \bar{R}_{x_5} \bar{R}_y$ limit⁵ of the second square root in (15) gives a mass contribution proportional to $N\bar{R}_{x_4}\bar{R}_y$ instead,

$$\mathcal{M}_{\text{flux}} = \frac{N\bar{R}_{x_4}\bar{R}_y}{8\pi^3 l_s^4},\tag{18}$$

Given that in this regime of parameters the growth of the soliton mass with N is linear, one may ask whether it is possible to have an uplifton with $\mathcal{M}_{\text{flux}} > \mathcal{M}_{\text{BPS}}$, which could lower its energy by tunneling emission of a D5 and a $\overline{\text{D5}}$ brane. Using Equation (13), this does not happen, neither in the regime where the mass of the uplifton grows linearly with N, nor in any other regime of parameters.

Since our purpose is to use the uplifton to uplift the cosmological constant of a flux compactification in which supergravity can be trusted, the parameters \bar{R}_y , \bar{R}_{x_4} and \bar{R}_{x_5} have to be large (they correspond to the extradimension sizes in units of $\sqrt{g_s}l_s$). One can check that in this regime of parameters the mass of the lightest uplifton (with N = 1) is necessarily heavier than the mass of two BPS D5 branes. However, as N increases, the mass of the uplifton can become smaller than the mass of 2N D5 branes. This happens because the binding energy of the brane and antibrane regions becomes of the same order as the energy of the branes

5. DISCUSSION

We have constructed the simplest example of an uplifton: a smooth solution that has three topologicallynontrivial cycles: a neutral one in the middle and two external ones with fluxes corresponding to D5 and $\overline{\text{D5}}$ charges. The dependence of the uplifton mass on the charges and the size of the compact directions is exactly what one expects from a topologically-nontrivial solution with fluxes.

It is remarkable that our uplifton has exactly the same structure as the solution one might expect from the geometrical transition of D5 and $\overline{\text{D5}}$ branes studied in [14] (depicted in Figure 2 in that paper): The two-cycle

³ It is also possible to construct upliftons in which y is fibered over the \mathbb{R}^3 base and the asymptotics is $\mathbb{R}^{4,1}$.

⁴ The relation between the four-dimensional and tendimensional Newton constants is $G_{10} = 8\pi^6 g_s^2 l_s^8 = (2\pi \sqrt{g_s} l_s)^3 \bar{R}_{x_4} \bar{R}_{x_5} \bar{R}_y G_4.$

⁵ This limit can be achieved when $\bar{R}_y \ll \bar{R}_{x_5}$ and $N \lesssim \bar{R}_{x_5}^2$.

wrapped by the branes shrinks at the two locations of the branes, giving rise to a flux-less topologically-nontrivial three cycle between the branes. Furthermore, the threecycles with positive and negative flux that, before the geometric transition, were shrinking at the position of the five-branes, now become large. Hence, the configuration of [14] should backreact in a three-bubble solution, with a neutral bubble in the middle and two equal and oppositely-charged ones on the sides. The only difference is that in our uplifton the three-cycles have an S^1 that is trivially fibered over an S^2 , while in a more general solution one may expect a more exotic fibration.

As we mentioned in the introduction, the smoothness and neutrality of the uplifton make it a more controlled uplift ingredient than $\overline{\text{D3}}$ branes. However, in the regime of parameters where we have supergravity control $(\bar{R}_y, \bar{R}_{x_4}, \bar{R}_{x_5} > 1)$ the uplifton is heavier than $\overline{\text{D3}}$ branes. Hence, in order to use it for uplifting one has to place it in a high-warp region of the compactification manifold. It

- J. M. Maldacena and C. Nunez, Int. J. Mod. Phys. A 16, 822 (2001), hep-th/0007018.
- [2] S. Kachru, R. Kallosh, A. D. Linde, and S. P. Trivedi, Phys. Rev. D68, 046005 (2003), hep-th/0301240.
- [3] I. Bena, M. Grana, and N. Halmagyi, JHEP 1009, 087 (2010), 0912.3519.
- [4] I. Bena, E. Dudas, M. Graña, and S. Lüst, Fortsch. Phys.
 67, 1800100 (2019), 1809.06861.
- [5] I. Bena, M. Grana, S. Kuperstein, and S. Massai (2014), 1410.7776.
- [6] I. Bena, E. Dudas, M. Graña, G. Lo Monaco, and D. Toulikas (2022), 2211.14381.
- [7] O. Aharony, Y. E. Antebi, and M. Berkooz, Phys. Rev. D 72, 106009 (2005), hep-th/0508080.
- [8] I. Bah and P. Heidmann, JHEP 09, 147 (2021), 2012.13407.
- [9] I. Bah and P. Heidmann, JHEP 09, 128 (2021), 2106.05118.
- [10] I. Bah and P. Heidmann, JHEP 10, 165 (2021),

5

would be intersting to establish whether this can be done using for example the Klabanov-Strassler throat [15].

The most important question that our analysis does not answer is whether the uplifton is perturbatively stable. The Kaluza-Klein bubbles that compose the uplifton are known to be unstable in vacuum [16], so this is a nontrivial possibility. However, as shown in [17–19], when these bubbles are wrapped by electromagnetic flux this instability can disappear. Furthermore, in the absence of fluxes it is possible that our bubbles can annihilate each other as it can happen for bound states of black holes and bubbles in vacuum [20]. It would be very interesting to explore these possibilities in future projects.

Acknowledgments We would like to thank Ibou Bah, Mariana Graña and Severin Lüst for useful discussions. This work was supported in part by the ERC Grants 772408 "Stringlandscape" and 787320 "QBH Structure", and by the NSF grant PHY-2112699.

2107.13551.

- [11] P. Heidmann, JHEP 02, 162 (2022), 2112.03279.
- [12] I. Bah, P. Heidmann, and P. Weck, JHEP 08, 269 (2022), 2203.12625.
- [13] I. Bah and P. Heidmann (2022), 2210.06483.
- [14] M. Aganagic, C. Beem, J. Seo, and C. Vafa, Nucl. Phys. B 789, 382 (2008), hep-th/0610249.
- [15] I. R. Klebanov and M. J. Strassler, JHEP 0008, 052 (2000), hep-th/0007191.
- [16] E. Witten, Nucl. Phys. B 195, 481 (1982).
- [17] I. Bah, A. Dey, and P. Heidmann (2021), 2112.11474.
- [18] S. Stotyn and R. B. Mann, Phys. Lett. B 705, 269 (2011), 1105.1854.
- [19] U. Miyamoto and H. Kudoh, JHEP **12**, 048 (2006), gr-qc/0609046.
- [20] H. Elvang and G. T. Horowitz, Phys. Rev. D 67, 044015 (2003), hep-th/0210303.