## Resonance states at Casati wave numbers for the 3-disk billiard

Roland Ketzmerick<sup>1</sup> and Jan Robert Schmidt<sup>1</sup>

<sup>1</sup>Technische Universität Dresden, Institut für Theoretische Physik and Center for Dynamics, 01062 Dresden, Germany

(Dated: December 12, 2022)

Resonance states of the 3-disk scattering system are presented for the first Casati wave number  $k \approx 912$  and the second Casati wave number  $k \approx 91242$ . They show multifractal structure in phase space, similar to the pioneering work by Casati *et al.* [Physica D **131**, 311 (1999)] for an open chaotic quantum map. In position space we observe scarring along segments of rays, related to multifractality and universal fluctuations, as recently found for dielectric cavities. To the best of our knowledge this resonance state at the second Casati wave number has a much larger wave number than published resonance states for the 3-disk scattering system or any other open or closed chaotic billiard.

The multifractal structure of resonance states in scattering systems with classically chaotic dynamics was first presented by Casati, Maspero, and Shepelyansky [1, 2] for the quantum kicked rotor with absorption. It was found that resonance states are supported by the backward trapped set of the classical scattering system, which is a fractal. The multifractal structure on the backward trapped set depends systematically on the decay rate of the resonance states, which was recently analyzed for dielectric cavities [3].

This is a brief report on the occasion of the 80th birthday of Giulio Casati, born on 9.12.42, see also [4]. We present resonance states of the 3-disk scattering system [5–7], a paradigmatic model of chaotic scattering.

A resonance state at the first Casati wave number  $k \approx$ **912** is shown in Fig. 1. We consider a distance R = 2.2 between the centers of the disks, while the radius of the disks is set to one. One can see the location of the three disks and the 6-fold symmetry of the intensity of the resonance state.

At the second Casati wave number  $k \approx 91242$  the intensity of a resonance state is shown in a symmetry region near the origin, see Fig. 2. One can observe scarring along many ray segments. Often it continues beyond a reflection on the boundary. This is in agreement with recent findings for resonance modes in dielectric cavities [3]. There it was argued, that this scarring is based on multifractality of resonances in phase space and universal fluctuations. It conceptually differs from periodic-orbit scarring and becomes even more prominent in the semiclassical limit.

To the best of our knowledge this resonance state at the second Casati wave number has a much larger wave number than published resonance states for the 3-disk scattering system or any other open or closed chaotic billiard. We implemented the matrix M [5], where det(M) = 0 gives the poles of the scattering matrix, by a fast matrix-vector multiplication [8].

The multifractal structure in phase space on the boundary of a disk is shown with unprecedented resolution in Fig. 3. One observes that the resonance state is supported by the backward trapped set of the



FIG. 1. Intensity of a resonance state at the first Casati wave number k = 912.4484 - 0.3112 i.

chaotic saddle. This is well understood since the work by Casati *et al.* [1]. Additionally, one observes a multifractal structure on the backward trapped set. This structure strongly depends on the decay rate of the resonance. Here we show just one resonance state at an exemplary decay rate related to the imaginary part of k. There exist heuristic approximations for this structure in fully open quantum maps as well as in partially open quantum maps and dielectric cavities, see references in [3]. A semiclassical theory for the multifractal structure of resonance states in chaotic scattering systems, however, is still missing more than 20 years after the pioneering work of Casati *et al.* [1].

RK thanks Giulio Casati for more than three decades of stimulating scientific discussions, his support in many ways, and his valuable and long-lasting contributions to the quantum chaos community.



FIG. 2. Intensity of a resonance state at the second Casati wave number k = 91242.2841 - 0.6951 i for a symmetry region near the origin  $[0, 0.01] \times [0, 0.01]$ , showing ray-segment scarring.

- G. Casati, G. Maspero, and D. L. Shepelyansky, Quantum fractal eigenstates, Physica D 131, 311 (1999).
- [2] For color figures of Ref. [1], see https://arxiv.org/abs/ cond-mat/9710118.
- [3] R. Ketzmerick, K. Clauß, F. Fritzsch, and A. Bäcker, Chaotic resonance modes in dielectric cavities: Product of conditionally invariant measure and universal fluctuations, Phys. Rev. Lett. **129**, 193901 (2022).
- [4] Special Issue "Quantum Chaos—Dedicated to Professor Giulio Casati on the Occasion of His 80th Birthday", edited by M. Robnik, Entropy 24 (2022).



FIG. 3. Boundary Husimi function of a resonance state at the second Casati wave number k = 91242.2841 - 0.6951i for Birkhoff coordinates  $s \in [-1, 1, 1.1]$  and  $p \in [-1, 1]$ , showing a multifractal structure on fine scales. A magnification shows that the Husimi function is smooth on the scale of a Planck cell.

- [5] P. Gaspard and S. A. Rice, Exact quantization of the scattering from a classically chaotic repellor, J. Chem. Phys. 90, 2255 (1989).
- [6] A. Wirzba, Quantum mechanics and semiclassics of hyperbolic n-disk scattering systems, Phys. Rep. 309, 1 (1999).
- [7] T. Weich, S. Barkhofen, U. Kuhl, C. Poli, and H. Schomerus, Formation and interaction of resonance chains in the open three-disk system, New J. Phys. 16, 033029 (2014).
  - [8] R. Ketzmerick and J. R. Schmidt, in preparation.