Fiber sum formulas for 4-manifolds, topological modular forms and $6d \mathcal{N} = (1, 0)$ theories

John Chae

Altadena, California, USA yjchae@formerstudents.ucdavis.edu

Abstract

Using the relation between four manifolds and topological modular form (TMF) from the six dimensional approach, we exhibit fiber sum formulas for infinite families of smooth spin four manifolds associated to compactifications of free and interacting 6d (1,0) SCFTs. We find that even the free theories have nontrivial fiber sum formulas and their forms are sensitive to an individual theory and parameters of four-manifolds. Furthermore, we reinforce the conjecture of Stolz and Teichner by expanding its evidence.

1 Introduction

Conformal field theory (CFT) is a class of quantum field theory that possesses conformal symmetry and plays an important role in physics and mathematics. From physics viewpoint, CFTs describe phenomena in high energy systems and condensed matter systems. For the latter perspective, CFTs are intimately related to vertex operator algebra, its representation theory, and topology via topological quantum field theories.

Due to its multifaceted aspects, CFTs of various dimensions have been investigated extensively over the past decades. However, a class of CFTs endowed with supersymmetry, called superconformal field theory (SCFT), in six dimensions has been mysterious (see [13] for a review). Its enigmatic aspects originate from a 6-dimensional SCFT being strongly interacting, containing tensionless strings and having no known lagrangian. Consequently, the standard approaches to conventional (S)CFTs does not apply, which makes 6d SCFTs challenging to analyze. Furthermore, its existence is rather special. By the classification of superconformal algebra [26], the maximal spacetime dimension in which superconformal algebra exists is six ($D \leq 6$). The maximal amount of supersymmetry that can be endowed to a 6d CFT is $\mathcal{N} \leq 2$ by an existence of a supermultiplet containing an energy-momentum tensor; this amounts to sixteen pseudo-real supercharges. This leads to two chiral $\mathcal{N} = (1,0)$ and (2,0) SCFTs. Their superconformal symmetry algebra is $osp(6,2|\mathcal{N})$ whose (maximal) bosonic subalgebra is $so(6,2) \times sp(\mathcal{N})_R$, where R stands for R-symmetry.

Despite 6-dimensional SCFTs are nonlagrangian theories, there are two approaches to analyze them. The first is string/M/F-theory and the second one is supermultiplets and moduli spaces. In

the latter approach, one begins with a multiplet and explores corresponding moduli space of vacuum via RG flow and analyze anomaly phenomenon to extract information about the 6d SCFT. This strategy was used in [2, 16]. In case of a 6d (1,0) SCFT, there are three kinds of supermultiplets: hypermultiplet, vector multiplet and tensor multiplet. The first two are described in Section 4 and 5. The moduli space associated to hypermultiplet is called Higgs branch while tensor branch¹ for tensor multiplet. The Higgs branch breaks $sp(\mathcal{N})_R$ symmetry whereas the tensor branch preserves it. All interacting 6d (1,0) theories have a tensor branch.

Anomaly phenomenon is a core aspect of a QFT. It provides consistency conditions and serves as a tool for extracting nonperturative information of the theory. The latter part is often called 't Hooft anomaly matching, which is utilized, for example, in examining UV and IR theories connected by renormalization group flows. In 6d SCFTs, four point correlation functions of currents associated symmetries represent anomalies. The symmetries can be flavor, gauge and/or diffeomorphsim. Roughly speaking, depending on whether the currents in the correlator are same or not, an anomaly is divided into pure and mixed types. For example, there are gauge anomaly and gravitational anomaly for the pure type and gauge-gravitational anomaly for a mixed type. Some anomalies whether pure or mixed must be canceled to obtain a consistent 6d SCFT². A key object in the anomaly analysis is an anomaly polynomial of a theory. It encodes all perturbative anomaly information (see Appendix B for a review). As we will see, anomaly polynomials play a crucial role in this paper.

A unique feature of 6d SCFTs is generating a variety of lower dimensional field theories through compactifications. An aspect of the theories relevant to our paper is a conjecture by Stolz and Teichner [36, 37]. The conjecture predicts a connection between topological modular forms (TMF) [3, 15] and a space of 2-dimensional minimally suspersymmetric quantum field theories. This conjecture was motivated by search for a geometric construction of an elliptic cohomology theory. A simplified³ version proposes a classification of the 2d suspersymmetric theories via TMF, which is a generalized cohomology theory.

Conjecture 1.1. (simplified [37])

 $\pi_d TMF \simeq \pi_0 \{ 2d \ \mathcal{N} = (0,1) \ SQFTs \text{ with gravitational anomaly } 2(c_R - c_L) = d \}.$

On the RHS, using the fact that gravitational anomaly d is a deformation invariant, it labels connected components of the space of the theories. Moreover, theories having the same value of the gravitational anomaly reside in the same connected component and they are connected by supersymmetry preserving continuous deformations, which include renormalization group flows. Namely,

RHS = $\{2d \ \mathcal{N} = (0,1) \text{ SQFTs with gravitational anomaly } 2(c_R - c_L) = d\}$ /deformations

Furthermore, the conjecture implies that geometric cocycles of the cohomology theory are represented by the field theories and predicts TMF-valued invariants for the 2d supersymmetric theories.

¹A moduli space of vacuum does not exists for the vector multiplet since it does not contain a scalar field.

 $^{^2{\}rm The}$ correlation functions involving a gauge current must be canceled for any 6d SCFTs.

³TMF is evaluated at a point in this paper, $\text{TMF}^{-d}(pt) = \text{TMF}_d(pt) = \pi_d \text{TMF}$. See [37, 6] for the general version of the conjecture.

The theory of topological modular forms have been playing a key role in various physics literature in the last few years [39, 40, 21, 7, 8].

An investigation of Conjecture 1.1 in relation to an invariant of 4-manifolds was initiated in [11]. Specifically, evidence for the conjecture was provided for a range of homotopical degrees d (see Table 2 in [11]). Moreover, a connected sum formula of the TMF-valued invariant of 4-manifolds associated to compactifications of a *nonphysical* 6d (1,0) SCFT on nonsmooth and nonspin 4-manifolds was proposed⁴ (see Section 2.7 of [11]).

In this paper, we extend the investigation of [11] by analyzing compactifications of a *physical* 6d (1,0) SCFT of both free and interacting theories on smooth, spin closed 4-manifolds. This class of 4-manifolds appear abundantly as surface bundles. We consider the fiber sum operation between these manifolds, which is more natural operation than the connected sum for the class of 4-manifolds [10] (see Appendix A for a review). We conjecture their fiber sum formulas for the above mentioned theories and provide evidence for the conjectures via the explicit examples summarized in tables. Furthermore, we strengthen Conjecture 1.1 by finding appropriate 2d theories corresponding to a broader range of homotopical degrees d than the ones considered in [11].

This paper is organized as follows. In Section 2, we expand the example list for the toy model as an exercise. We next describe allowed 4-manifolds in compactifications of (phyisical) 6d (1,0) SCFTs in Section 3. We begin with a free 6d (1,0) hypermultiplet theory compactified on 4manifolds to find corresponding fiber sum formula in Section 4. And then we move onto a free (1,0) vector multiplet theory in Section 5. We next consider an interacting theory, namely, the rank one E-string theory and obtain a fiber sum formula for specific 4-manifolds in Section 6. In Section 7, we reinforce Conjecture 1.1 by providing more evidence. We finish with open questions in Section 8.

2 The extended toy model

A toy model was introduced in [11]. Since it is not associated with a 6d (1,0) SCFT, hence 2d(0,1) theories do not arise from a 6d SCFT. Furthermore, the model deals with a broader range of 4-manifolds compared to that of Section 3, namely,

$ \begin{cases} \text{Topological compact} & X \\ \text{oriented 4-manifolds} & X \end{cases} $	Т	$ \left\{ \begin{array}{l} 2d\mathcal{N} = (0,1) \text{ lattice SCFT} \\ \Gamma = H^2(X;\mathbb{Z})/TorH^2(X;\mathbb{Z}) \end{array} \right\} $	S	π TME
homeomorphism	/	SUSY deformations	/	//* T I/II

We denote $\mathcal{T} := S \circ T$ and hence $\mathcal{T}[X] \in \pi_d$ TMF, where the homotopical degree d is given by $3b_2^+ - 2b_2^-$ and π_d denotes the stable homotopy groups.

We extend the examples of 4-manifolds in [11] for the toy model. The family of the Hirzebruch surfaces F_n , $n \ge 0^5$. Its intersection form is

$$Q_{F_n} = \begin{pmatrix} n & 1\\ 1 & 0 \end{pmatrix}$$

⁴A compactification of a *physical* 6d (1,0) SCFT requires compact 4-manifolds to be smooth and spin.

 $^{{}^{5}}F_{0} = S^{2} \times S^{2}$ was done in [11]

X	$d = 3b_2^+ - 2b_2^-$	$\mathcal{T}[X] \in \pi_d \mathrm{TMF}$
$F_n = F_n (F_0 = S^2 \times S^2)$	1	η
$-E_8$	-16	E_4/Δ
\mathbb{CP}^2	3	ν
$\overline{\mathbb{CP}^2}$	-2	0
K3	-29	0
$\overline{K3}$	51	$ u\Delta^2$
$\frac{1}{2}K3, E(1)$	-15	$\eta E_4/\Delta$
$\overline{\frac{1}{2}K3}$	25	$\eta\Delta$

Table 1: This table is an expansion of the table in [11] by adding $-E_8$, $\overline{K3}$, $\frac{1}{2}K3$, and E(1). E(1) has the same degree as that of the Enriques surface $\frac{1}{2}K3$ but their intersection forms are inequivalent, hence they are not homeomorphic. η and ν are generators of $\pi_1 \text{tmf} \simeq \mathbb{Z}_2$ and $\pi_3 \text{tmf} \simeq \mathbb{Z}_{24}$, respectively (see Appendix C for details).

where n is the Euler number of the surface. Consequently, their $b_2^{\pm}(F_n) = 1$ so all F_n have the same homotopical degree as F_0 , which means that their π_d TMF coincide. Another well-known complex surface is a family of elliptic surfaces E(n), $n \in \mathbb{Z}_+$. This family includes K3 and its orientation reversal $\overline{K3}$. Using the data in Appendix A, we get d = -14n - 1 for E(n). Although K3-surface case is trivial, $\overline{K3}$ case doesn't vanish as shown in Table 1.

3 Four manifolds for compacification

Before we analyze various physical theories, let us describe a class of 4-manifolds that can be used in compactification of a 6d (1,0) SCFT [11]. In order to obtain a sensible 2d (0,1) theory from a compactification of a *physical* 6d (1,0) theory, 4-manifolds need to be smooth and spin. In this setting, the diagram in Section 2 becomes [11]

$$\frac{\left\{ \begin{array}{c} \text{Smooth spin} \\ \text{compact 4-manifolds} \end{array} X \right\}}{\text{diffeomorphism}} \longrightarrow \frac{t}{\text{SUSY deformations}} \xrightarrow{\sigma} \pi_* \text{TMF}$$

where σ is the toplogical Witten genus⁶. Moreover, $H_2(X)$ being torsion-free is also assumed. This condition can be satisfied by choosing a simply connected X ($\pi_1(X) = 1$). Furthermore, since the torsion-free part of π_* TMF is the elliptic genus and it is nonvanishing only when the degree d of the TMF is $4\mathbb{Z}$, therefore we consider X whose $d \in 4\mathbb{Z}$ in this paper.

⁶Note there is a difference between the physical and the mathematical definitions of the map (see Appendix C). The former is slightly more general.

4 Hypermultiplet theory

In this section, we consider a free $\mathcal{N} = (1,0)$ hypermultiplet theory in 6 dimensions. Although it is a free theory, a fiber sum formula for this theory turns out to be nontrivial (i.e. not a mere multiplication). This hypermultiplet consists of complex scalar fields q, \tilde{q}^{\dagger} and their superpartner spinor fields ψ on a spacetime manifold M^6 ; they are sections of certain bundles over M^6 :

$$\left\{ (q, \tilde{q}^{\dagger}) \in \Gamma \left(L \otimes R \right), \quad \psi \in \Gamma S \right\}$$

where L is a trivial complex line bundle, R is a principal $SU(2)_R$ bundle and S is a spin bundle over M^6 . q and \tilde{q}^{\dagger} together carry a fundamental representation of $SU(2)_R$ and their vacuum expectation values parametrizing the Higgs branch of the theory.

After compactification of the 6d free hypermultiplet theory on X, 2d (0, 1) theory's field content depends the signature of X [11]:

- $\sigma > 0 : \frac{\sigma}{4}$ number of (0,1) Fermi multiplets
- $\sigma < 0 : \frac{|\sigma|}{4}$ number of (0,1) chiral multiplets

An advantage of 6-dimensional theories is that (perturbative) anomaly polynomials can be computed, which in turn can be used to obtain gravitational anomaly of compactifications of the theories (see Appendix B for review). In our context, we are interested in compatifications on smooth spin 4-manifolds, thus gravitational anomaly in 2 dimensions. The 2 dimensional gravitational anomaly associated to a free single $\mathcal{N} = (1,0)$ hypermultiplet is [11]

$$2(c_R - c_L) = -\frac{\sigma(X)}{4} \in \mathbb{Z}$$

The above formula can be straightforwardly obtained from the anomaly polynomial of the theory (see Appendix B). An importance of the above formula is that the difference of the central charges is identified with the homotopical degree d of π_d TMF for any theories.

For the theory under consideration, its TMF degree⁷ d under the fiber sum is additive,

$$d(X_1 \#_f X_2) = d(X_1) + d(X_2),$$

since σ is additive. We conjecture a (generalized) fiber sum formula for a 6d (1,0) free hypermultiplet theory.

Conjecture 4.1. Let X_1, X_2 be smooth spin closed oriented 4-manifolds and X be their (generalized) fiber sum $X = X_1 \#_f X_2$, which is smooth spin closed and oriented. Moreover, let d_1, d_2 be the TMF degrees of X_1 and X_2 , respectively. For $d_1 + d_2 \notin 24\mathbb{Z}$, a 6d (1,0) free hypermultiplet SCFT compactified on $X_i, i = 1, 2$ satisfies the following (generalized) fiber sum formula for a generator of \mathbb{Z} -polynomial ring summand of π_* TMF associated to X.

$$H(X_1 \#_f X_2) \stackrel{\circ}{=} \Delta^{\left\lfloor \frac{w_1 + w_2}{2} \right\rfloor} H(X_1) * H(X_2)$$

⁷We use TMF degree and homotopical degree interchangeably (cf. footnote 3).

X	d	H(X)
E(2)	4	$\frac{2E_4^2E_6}{\Delta}$
E(4)	8	E_4
E(6)	12	$2E_6$
E(20)	40	$E_4^2\Delta$
E(22)	44	$2E_4E_6\Delta$
E(24)	48	$12\Delta^2, (j\Delta)^2$
$\overline{E(2)}$	-4	$\frac{2E_4E_6}{\Delta}$
$\overline{E(4)}$	-8	$\frac{E_4^2}{\Delta}$
$\overline{E(6)}$	-12	$\frac{2E_6}{\Delta}$

Table 2: Generators H(X) of \mathbb{Z} -polynomial ring part of π_d TMF associated to E(2r) and its orientation reversal $\overline{E(2r)}$ for the free (1,0) hypermultiplet theory. The third column can be computed using [3] (see Chapter 13) and Appendix C.

$$E_4^{p_1} * E_4^{p_2} = E_4^{p_1+p_2}, \quad if \ p_1 + p_2 \ge 3, \ use \ E_4^3 = j\Delta$$
$$E_6^{w_1} * E_6^{w_2} = E_6^b, \quad w_1 + w_2 \equiv b \mod 2, \quad (b = 0, 1)$$
$$\Delta^{m_1} * \Delta^{m_2} = \Delta^{m_1+m_2}.$$

where the circle denotes up to the j-modular function.

Remark 1 As stated in the conjecture, the (entire) RHS is understood as extracting only generators of the polynomial ring.

Remark 2 Since we consider the total degree $\notin 24\mathbb{Z}$, the *j*-function is discarded in a final expression on the RHS.

In the next subsections, we provide evidence for this conjecture.

4.1 Elliptic surfaces

We first consider a family of smooth complex elliptic surfaces E(n). They can be obtained inductively beginning from E(1) and a fiber sum operation (see Appendix A). For even values of n, they are spin manifolds. Using their topological data in Appendix A, homotopical degree d of E(n) is

$$d = -\frac{\sigma}{4} = 4r, \quad r = 1, 2, 3, \cdots$$

We list generators of integer polynomial ring part of π_d TMF in Table 2. Conjecture 4.1 was checked, for example, using the table.

X	d	H(X)
$Z_{2,3}^3 = X_{2,3} \#_f E(6)_K$ $V_3^3 = X_3 \#_f E(6)_K$	-4	$\frac{2E_4E_6}{\Delta}$
$Z_{2,3}^2 = X_{2,3} \#_f E(4)_K$ $V_3^2 = X_3 \#_f E(4)_K$	-8	$\frac{E_4^2}{\Delta}$
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	-12	$\frac{2E_6}{\Delta}$
$V_5^{16} = X_5 \#_f E(32)_K$	-16	$\frac{E_4}{\Delta}$
$Z_{2,5}^{12} = X_{2,5} \#_f E(24)_K$	-32	$\frac{E_4^2}{\Delta^2}$
$Z_{2,5}^{11} = X_{2,5} \#_f E(22)_K$	-36	$\frac{2E_6}{\Delta^2}$
$Z_{2,5}^{10} = X_{2,5} \#_f E(20)_K$ $V_5^{10} = X_5 \#_f E(20)_K$	-40	$\frac{E_4}{\Delta^2}$

Table 3: Generators H(X) of Z-polynomial ring part of π_d TMF associated to $Z_{2,n}^r$ and V_n^r for the free (1,0) hypermultiplet theory are listed. For $X_{2,3}$ and X_3 , their TMFs are π_{-16} TMF = $\langle E_4/\Delta \rangle$. For $X_{2,5}$ and X_5 , their TMFs are π_{-80} TMF = $\langle E_4^2/\Delta^4 \rangle$. $E(2r)_K$ has the same the Euler characteristic and the signature as E(2r).

The generators in Table 2 can be computed using [3] (see Chapter 13) and Appendix C. For example, E(20) has $\sigma = -160$ and hence homotopical degree $d = 40 \equiv 0 \mod 8$. This implies that π_{40} TMF contains a $\mathbb{Z}[x]$ (see Appendix C). From [3], we see that prime 2 contribution of torsion free part of π_{40} tmf₍₂₎ is $\mathbb{Z}^2_{(2)}$ whereas prime 3 contribution of π_{40} tmf₍₃₎ is $\mathbb{Z}^2_{(3)}^{8}$. They are both generated by $E_4^2\Delta$. After localization, we arrive at the polynomial ring with \mathbb{Z} coefficient for torsion free part of π_{40} TMF.

For negative values of d such as $d = -8 \equiv 0 \mod 8$, we need to use the 576 periodic property of π_d TMF. Namely,

$$\pi_{-8}$$
TMF $\simeq \pi_{568}$ TMF.

We next use the fact that $\pi_* \text{tmf}_{(2)}$ is 192 periodic and $\pi_* \text{tmf}_{(3)}$ is 72 periodic. Hence, we easily deduce that $\pi_{184} \text{tmf}_{(2)}$ for the former and $\pi_{64} \text{tmf}_{(3)}$ for the latter. Their torsion free parts are generated by $E_4^2 \Delta^7$ and $E_4^2 \Delta^2$, respectively [3]. After localization, we arrive at E_4^2 / Δ for π_{-8} TMF.

X	d	H(X)
$Z(1,1)_n$	8	E_4
$Z(1,3)_n$	12	$2E_6$
$Z(3,3)_{n}$	16	E_4^2
$Z(1,7)_n$	20	$2E_{4}E_{6}$
$Z(3, 11)_n$	32	$E_4\Delta$

Table 4: Generators H(X) of \mathbb{Z} -polynomial ring part of π_d TMF associated to $Z(k,m)_n$ for the free (1,0) hypermultiplet theory are listed.

4.2 Fiber sum manifolds

For verifications of Conjecture 4.1, we analyze families of smooth spin irreducible 4-manifolds $Z_{2,n}^r$ and V_n^r constructed in [1]. Their constructions and properties are summarized in Appendix A. The homotopical degrees of $Z_{2,n}^r$ and V_n^r coincide

$$d = -\frac{\sigma}{4} = -\frac{2n^3}{3} + \frac{2n}{3} + 4r$$

Their generators H(X) of polynomial rings of π_* TMF are listed in Table 3. Using this table and Table 2, we verified Conjecture 4.1.

We note that $X_{2,3}$ factor of $Z_{2,3}^r$ is Σ_6 -bundle over Σ_{19} whereas $X_{2,5}$ in $Z_{2,5}^r$ is Σ_{10} -bundle over Σ_{51} . X_3 in V_3^r is Σ_9 -bundle over Σ_{19} ; X_5 in V_5^r is Σ_{15} -bundle over Σ_{51} .

4.3 Nonsymplectic manifolds

We provide further evidence of Conjecture 4.1. Specifically, we consider a family $Z(k,m)_n$ of manifolds that are smooth spin nonsymplectic 4-manifolds constructed in [38] (see Appendix A for details). Their homotopical degree is

$$d = -\frac{\sigma}{4} = 2(k+m) + 4$$

for all $k, m \ge 1$ and odd. Conjecture 4.1 was checked using Tables 4 and 2, namely, $Z(k,m)_n \#_f Z(r,s)_n$ and $Z(k,m)_n \#_f E(2r)$ and $Z(k,m)_n \#_f \overline{E(2r)}$.

5 Vector multiplet theory

In this section, we describe a free $\mathcal{N} = (1,0)$ vector multiplet theory in 6 dimensions. We note that the theory is only scale invariant (nonconformal) [2, 33]. A fiber sum formula depends on a

⁸Torsion from prime 2 is $\mathbb{Z}/4\mathbb{Z}$ whereas $\mathbb{Z}/3\mathbb{Z}$ from prime 3.

value of parameter n. This multiplet consists of a 1-form A and its superpartner spinor fields λ on M^6 :

$$\left\{A\in\Omega^1(M^6),\quad \bar\lambda\in\Gamma\left(S\otimes R\right)\right\},$$

where S is a spin bundle, R is a principal $SU(2)_R$ bundle over M^6 and $\overline{\lambda}$ carries the fundamental representation of $SU(2)_R$.

After compactification of the 6d free hypermultiplet theory on X, 2d(0, 1) theory's field content depends the Betti numbers b_i of X [11]:

- b_0 number of (0,1) vector multiplets
- b_1 number of (0,1) compact chiral multiplets
- b_2^- number of (0,1) Fermi multiplets

The gravitational anomaly of a 2d (0,1) theory obtained from a compactification of the 6D free $\mathcal{N} = (1,0)$ vector multiplet theory is [11]

$$d = 2(c_R - c_L) = -2\chi_h(X) \in \mathbb{Z}$$

where χ_h is the holomorphic Euler characteristic of X. This formula is obtained from the anomaly polynomial of the theory (see Appendix B)⁹.

For this theory, the TMF degree d under the fiber sum receives a contribution from the gluing surface Σ ,

$$d(X_1 \#_f X_2) = d(X_1) + d(X_2) + \chi(\Sigma).$$

When $\Sigma = T^2$, then it is additive.

5.1 Elliptic surfaces

We begin with the elliptic surface, whose results are useful in the next subsection. The TMF degree for E(2r) is $-\chi_h = -4r$, which is same as that of $\overline{E(2r)}$ of the free hypermultiplet in the previous section. For $\overline{E(2r)}$, its degree is $-\chi_h = -20r$ so the TMF of the orientation reversed manifold is a subset of E(2r)'s. Hence, the vector multiplet is insensitive to the orientation reversal. As an example, we choose E(26) in Table 5. From the periodic property of $\pi_* \text{TMF}$, $\pi_{-52} \text{TMF} \cong \pi_{524} \text{TMF}$. Using [3],

$$\pi_{140} \operatorname{tmf}_{(2)} = \mathbb{Z}_{(2)}^6 \qquad \pi_{20} \operatorname{tmf}_{(3)} = \mathbb{Z}/3\mathbb{Z} \oplus \mathbb{Z}_{(3)}$$

The torsion free parts are generated by $2E_4E_6\Delta^5$ and E_4E_6 , respectively. After localization and using that $d(\Delta) = 24$, we get the generator in the table.

5.2 Fiber sum manifolds

We consider $Z_{2,n}^r$, n = 3, 5 and $Z_{2,7}^r$ (see Appendix A for a review) and observe the following behaviors under the fiber sum.

⁹The anomaly polynomial takes into account the nonconformality of the theory.

X	d	V(X)	X	d	V(X)
E(4)	-8	$\frac{E_4^2}{\Delta}$	$Z_{2,3}^2 = X_{2,3} \#_f E(4)_K$	-268	$\frac{2E_4E_6}{\Delta^{12}}$
E(6)	-12	$\frac{2E_6}{\Delta}$	$Z_{2,3}^3 = X_{2,3} \#_f E(6)_K$	-272	$\frac{E_4^2}{\Delta^{12}}$
E(8)	-16	$\frac{E_4}{\Delta}$	$Z_{2,3}^4 = X_{2,3} \#_f E(8)_K$	-276	$\frac{2E_6}{\Delta^{12}}$
E(20)	-40	$\frac{E_4}{\Delta^2}$	$Z_{2.5}^{10} = X_{2,5} \#_f E(20)_K$	-1220	$\frac{2E_4^2E_6}{\Lambda^{52}}$
E(24)	-48	$\frac{E_4^3}{\Delta^3}$	$Z_{2,5}^{12} = X_{2,5} \#_f E(24)_K$	-1228	$\frac{2E_4E_6}{\Delta^{52}}$
E(26)	-52	$\frac{2E_4E_6}{\Delta^3}$	$Z_{2.5}^{13} = X_{2,5} \#_f E(26)_K$	-1232	$\frac{E_4^2}{\Lambda^{52}}$
E(28)	-56	$\frac{E_4^2}{\Delta^3}$	$Z_{2,5}^{15} = X_{2,5} \#_f E(28)_K$	-1240	$\frac{E_4}{\Delta^{52}}$

Table 5: Generators V(X) of Z-polynomial ring part of π_d TMF associated to $Z_{2,n}^r$ for the free (1,0) vector multiplet theory are listed. The third column can be computed using [3] (see Chapter 13) and Appendix C.

X	d	V(X)
$Z_{2,7}^1 = X_{2,7} \#_f E(2)_K$	-3224	$\frac{E_4^2}{\Delta^{135}}$
$Z_{2,7}^2 = X_{2,7} \#_f E(4)_K$	-3228	$\frac{2E_6}{\Delta^{135}}$
$Z_{2,7}^3 = X_{2,7} \#_f E(6)_K$	-3232	$\frac{E_4}{\Delta^{135}}$
$Z_{2,7}^4 = X_{2,7} \#_f E(8)_K$	-3236	$\frac{2E_4^2E_6}{\Delta^{136}}$
$Z_{2,7}^6 = X_{2,7} \#_f E(12)_K$	-3244	$\frac{2E_4E_6}{\Delta^{136}}$
$Z_{2,7}^7 = X_{2,7} \#_f E(14)_K$	-3248	$\frac{E_4^2}{\Delta^{136}}$

Table 6: Generators V(X) of Z-polynomial ring part of π_d TMF associated to $Z_{2,7}^r$ for the free (1,0) vector multiplet theory are listed. The third column can be computed using [3] (see Chapter 13) and Appendix C.

Conjecture 5.1. Let $X_1 = X_{2,n}, X_2 = E(2r)_K$ be smooth spin closed oriented 4-manifolds and $Z_{2,n}^r$ be their (generalized) fiber sum $X_1 #_f X_2$, which is also smooth, spin, closed and oriented. Moreover, let d_1, d_2 be the TMF degrees of X_1, X_2 , respectively. For a 6d (1,0) free vector multiplet SFT compactified on X_i , i = 1, 2 a fiber sum formula for $d_1 \notin 24\mathbb{Z}$, $d(Z_{2,n}^r) \notin 24\mathbb{Z}$ and n = 3, 5, is

$$V(Z_{2,n}^{r}) \stackrel{\circ}{=} \frac{1}{\Delta^{s}} V(X_{2,n}) * V(E(2r)_{K}) \quad \text{for all } r,$$

$$E_{4}^{p_{1}} * E_{4}^{p_{2}} = E_{4}^{p_{1}+p_{2}}, \quad \text{if } p_{1}+p_{2} \ge 3, \text{ use } E_{4}^{3} = j\Delta$$

$$E_{6}^{w_{1}} * E_{6}^{w_{2}} = E_{6}^{b}, \quad w_{1}+w_{2} \equiv b \mod 2, \quad (b=0,1)$$

$$\Delta^{m_1} * \Delta^{m_2} = \Delta^{m_1 + m_2}$$

$$s = \begin{cases} n - \left\lceil \frac{p_1 + p_2}{n - 1} \right\rceil + (w_1 + w_2), \ r = even \\ n - \left\lceil \frac{p_1 + p_2}{n - 2} \right\rceil, \ r = odd \end{cases}$$

Furthermore, in case of $Z_{2,7}^r$, for $d_1(X_{2,7}) \notin 24\mathbb{Z}$ and $d(Z_{2,7}^r) \notin 24\mathbb{Z}$, we have

$$V(Z_{2,7}^{r}) \stackrel{\circ}{=} \frac{1}{\Delta^{s}} V(X_{2,7}) * V(E(2r)_{K}) \quad \text{for all } r,$$

$$E_{4}^{p_{1}} * E_{4}^{p_{2}} = E_{4}^{p_{1}+p_{2}-1}, \quad \text{if } p_{1}+p_{2}-1 \geq 3, \text{ use } E_{4}^{3} = j\Delta$$

$$E_{6}^{w_{1}} * E_{6}^{w_{2}} = E_{6}^{b}, \quad w_{1}+w_{2} \equiv b \mod 2, \quad (b=0,1)$$

$$\Delta^{m_{1}} * \Delta^{m_{2}} = \Delta^{m_{1}+m_{2}}$$

$$s = \begin{cases} 8, \ r = \ odd \\ 9, \ r = \ even \end{cases}$$

Remark It can be inferred that the degree of $Z_{2,n}^r$ is 24Z, if the RHS consists of solely Δ .

Verifications of Conjecture 5.1 were done for instance using Tables 5 and 6.

6 E-strings

We analyzed free theories in the previous sections, we proceed to examine an interacting 6D (1,0) SCFT, E-string theory compactified on a 4-manifold. We give a brief review of the theory and then move onto their compactifications and fiber sum formula.

<u>Review</u> In M-theory, E-string theory of rank 1 is a world volume theory of a brane system consisting of a M5-brane attached to a M9-brane, which carries an E_8 gauge symmetry. Specifically, the M5 is embedded as an E8-instanton. The low energy limit yields a single free hypermultiplet characterizing the center of mass motion of the M5-brane and an interacting 6d (1,0) superconformal field theory with an E_8 flavor symmetry. Furthermore, the strings arising from M2-branes suspended between the M5-brane and the M9-brane become massless (i.e. tensionless) as the M5-brane approaches the M9-brane. These strings are called E-strings. Anomaly polynomial of this theory was found in [27, 28] and recorded in Appendix B.

6.1 Elliptic surfaces

From the anomaly polynomial of the E-strings and the topological data of $E(2r)_K$ in the appendices, we find its homotopical degree d to be

$$d = -2(5\sigma + 11\chi_h) = 116r.$$

Furthermore, using the fiber sum definition of E(n) (see Appendix A), we easily deduce $E(m)#_fE(n) = E(m+n)$.

X	d	E(X)
E(2)	116	$2E_4E_6\Delta^4$
E(4)	232	$E_4^2 \Delta^9$
E(6)	348	$2E_6\Delta^{14}$
E(8)	464	$E_4\Delta^{19}$
E(10)	580	$2E_4^2 E_6 \Delta^{23}$

Table 7: Generators E(X) of Z-polynomial ring part of π_d TMF associated to E(2r) for the E-string theory are listed. The third column can be computed using [3] (see Chapter 13) and Appendix C.

Conjecture 6.1. Let E(2m) be an elliptic surface. For the E-string theory compactified on E(2m), a fiber sum formula $E(2r) #_f E(2s)$ in case of total homotopical degree $d = d(E(2r)) + d(E(2s)) \notin 24\mathbb{Z}$ is

$$\begin{split} E(E(2r)\#_{f}E(2s)) &\stackrel{\circ}{=} \Delta^{\lfloor\frac{w_{1}+w_{2}}{2}\rfloor}E(E(2r)) * E(E(2s)) \\ E_{4}^{p_{1}} * E_{4}^{p_{2}} &= E_{4}^{p_{1}+p_{2}}, \quad if \ p_{1}+p_{2} \geq 3, \ use \ E_{4}^{3} &= j\Delta \\ E_{6}^{w_{1}} * E_{6}^{w_{2}} &= E_{6}^{b}, \quad w_{1}+w_{2} \equiv b \mod 2, \quad (b=0,1) \\ \Delta^{m_{1}} * \Delta^{m_{2}} &= \Delta^{m_{1}+m_{2}} \end{split}$$

As before, the circle denotes up to the j-modular function. This result coincides with that of the free (1,0) hypermultiplet theory in Section 4. This is because the generic fiber of E(n) is T^2 whose Euler characteristic vanishes and hence d becomes additive for this manifold. It is straightforward to check the above conjecture via Table 7.

6.2 Fiber sum manifold

We move onto the fiber sum manifold $Z_{2,n}^r$. Using Appendix A and B, its TMF degree d is easily calculated to be

$$d = -\frac{4}{3} \left(97n^3 + 2n - 87r \right).$$

We propose the following fiber sum formula.

Conjecture 6.2. Let $Z_{2,n}^r$ be the fiber sum manifold and $d = d_{r,n}$ be its TMF degree as above. For values of r, n such that $d \notin 24\mathbb{Z}$ and the E-string theory compactified on $Z_{2,n}^r$, its fiber sum formula takes the following form.

$$E(Z_{2,n}^{r}) \stackrel{\circ}{=} \Delta^{\frac{\chi(\Sigma_{f+b})}{2} + s(n) + k(r)} E(X_{2,n}) * E(E(2r)_{K})$$

$$E_{4}^{p_{1}} * E_{4}^{p_{2}} = E_{4}^{p_{1}+p_{2}+c}, \quad if \ p_{1} + p_{2} + c \geq 3, \ use \ E_{4}^{3} = j\Delta$$

$$E_{6}^{w_{1}} * E_{6}^{w_{2}} = E_{6}^{b}, \ w_{1} + w_{2} \equiv b \mod 2, \quad (b = 0, 1)$$

$$\Delta^{m_{1}} * \Delta^{m_{2}} = \Delta^{m_{1}+m_{2}}$$

X	d	E(X)	X	d	E(X)
$Z^{2}_{2,3}$	-3268	$\frac{2E_4E_6}{\Delta^{137}}$	$Z^{1}_{2,7}$	-44264	$\frac{E_4^2}{\Delta^{1845}}$
$Z^{3}_{2,3}$	-3152	$\frac{E_4^2}{\Delta^{132}}$	$Z^{2}_{2,7}$	-44148	$\frac{2E_6}{\Delta^{1840}}$
$Z^4_{2,3}$	-3036	$\frac{2E_6}{\Delta^{127}}$	$Z^{3}_{2,7}$	-44032	$\frac{E_4}{\Delta^{1835}}$
$Z^{1}_{2,5}$	-16064	$\frac{E_4^2}{\Delta^{670}}$	$Z^{4}_{2,7}$	-43916	$\frac{2E_4^2E_6}{\Delta^{1831}}$
$Z^2_{2,5}$	-15948	$\frac{2E_6}{\Delta^{665}}$	$Z^{1}_{2,9}$	-94192	$\frac{E_4}{\Delta^{3925}}$
$Z^{3}_{2,5}$	-15832	$\frac{E_4}{\Delta^{660}}$	$Z^{2}_{2.9}$	-94076	$\frac{2E_4^2E_6}{\Lambda^{3921}}$

Table 8: Generators E(X) of Z-polynomial ring part of π_d TMF associated to $Z_{2,n}^r$ for the E-string theory (rank one) are listed. Four-manifolds used in the fiber summation $X_{2,n}$ have their generators as $E(X_{2,3}) = 2E_4^2 E_6/\Delta^{125}$, $E(X_{2,5}) = 2E_4 E_6/\Delta^{620}$, $E(X_{2,7}) = 2E_6/\Delta^{1747}$, $E(X_{2,9}) = 2E_6/\Delta^{3765}$.



where f and b are genus of the fiber and base surfaces, respectively.

Remark. We note that for a fixed n, the overall factor is independent of specific value of r. It turns out that the power of the overall factor cannot be expressed as a linear combinations of the r independent quantities $\chi(X_{2,n}), \sigma(X_{2,n}), \chi(\Sigma_{f+b})$ and $c_1^2(Z_{2,n}^r)$ for both even and odd r.

The conjecture was checked using Table 8.

We illustrate the derivations behind Table 8. For example, we consider $Z_{2,3}^2$. After substituting its parameter values into the above formula, we get $d = -3268 \equiv 4 \mod 8$. Applying the same method as in Section 4.1, π_{-3268} TMF $\cong \pi_{188}$ TMF. It turns out π_{188} tmf is torsion free [3]

$$\pi_{188} \operatorname{tmf}_{(2)} = \mathbb{Z}_{(2)}^8 \qquad \pi_{44} \operatorname{tmf}_{(3)} = \mathbb{Z}_{(3)}^2$$

where the periodicity of $\pi_* \text{tmf}_{(3)}$ is used. The prime 2 generator is $2E_4E_6\Delta^7$ and the prime 3 generator is $E_4E_6\Delta$. After localization, we get the generator displayed in Table 8.

7 TMF and 2d (0,1) theories

Evidence for Conjecture 1.1 was found for the TMF degrees $d = 0, 1, \dots, 15, 24$ and $d = -1, -2, \dots, -20, -24$ in [11]. We fill the gaps between 15 and 24 and between -20 and -24. Furthermore, we extend the table, which is in Appendix C (Table 9). An important lesson from [11] is that, for d > 0 many of 2d theories are realized as a sigma model whose target space is determined by generators of π_d TMF ¹⁰ whereas this is no longer true for d < 0. In this section, we describe how Table 9 is obtained for several examples.

For $d = 16 \equiv 0 \mod 8$, we see that $\pi_{16} \operatorname{tmf}_{(2)} = \mathbb{Z}_{(2)}$ and $\pi_{16} \operatorname{tmf}_{(3)} = \mathbb{Z}_{(3)}$ are generated by E_4^2 [3] (Chapter 13). Hence $\pi_{16} \operatorname{TMF} = \mathbb{Z}[x] \langle E_4^2 \rangle$. Using $\hat{A}(Y_8) = 1$ and the multiplicative property of the \hat{A} genus under cartesian product of two manifolds, we see that $\hat{A}(Y_8 \times Y_8) = 1$ and its Witten genus of $Y_8 \times Y_8$ is E_4^2 . Then the corresponding 2d(0,1) theory is a sigma model having target space ¹¹ as Y_8^2 (see Appendix C for the construction of Y_8).

For $d = 17 \equiv 1 \mod 8$, $\pi_{17} \operatorname{tmf}_{(2)} = \mathbb{Z}/2\mathbb{Z} \langle \eta E_4^2 \rangle \oplus \mathbb{Z}/2\mathbb{Z} \langle \kappa \nu \rangle$ whereas prime 3 part is trivial. A string manifold corresponding to η is S^1 and to ν is S^3 ; a Lie group G_2 is associated to κ [11]. Consequently, we arrive at the sigma models written in Table 9.

For $d = 20 \equiv 4 \mod 8$, $\pi_{20} \operatorname{tmf}_{(2)} = \mathbb{Z}/8\mathbb{Z} \langle \bar{\kappa} \rangle \oplus \mathbb{Z}_{(2)} \langle 2E_4E_6 \rangle$ whereas $\pi_{20} \operatorname{tmf}_{(3)} = \mathbb{Z}/3\mathbb{Z} \langle \beta^2 \rangle \oplus \mathbb{Z}_{(3)} \langle E_4E_6 \rangle$. Then $\pi_{20} \operatorname{TMF} = \mathbb{Z}/24\mathbb{Z} \oplus \mathbb{Z}[x]$. A Lie group Sp(2) is associated with β [11]. Hence we arrive at the target spaces in the table.

When d = -23, we need to find a theory corresponding to $8\eta/\Delta$ generator ¹². Applying the strategy in [11], we search for a 2d CFT with central charges (23/2, 0) consisting of left moving fermions moving in \mathbb{R}^r/Λ in which Λ is an odd self dual lattice. Specifically, this is may be realized by a chiral WZW theory at level $k \in \mathbb{Z}$ associated with a Lie algebra g whose root lattice $\Lambda_R = \Lambda$. The central charge of the g WZW theory is given by

$$c = \frac{k \dim g}{k + h^{\vee}}$$

where h^{\vee} is the dual Coxeter number. We find that g = so(5) and g = sp(4) both at k = -23. The fermions are moving in \mathbb{R}^2/Λ_2 . Since they are isomorphic, pick the former and denote it by $T_{(B_2;-23)}$ in Table 9 in Appendix C.

We next consider a generator ν/Δ^2 whose d = -45. As done above, we find that WZW theories $T_{(B_3;-75)}$ and $T_{(B_{22};1)}$ have $c_L = 45/2$. They are in the same connected component and hence deformation equivalent. In the former theory, fermions are moving in \mathbb{R}^3/Λ_3 whereas they are propagating in $\mathbb{R}^{22}/\Lambda_{22}$ in the latter.

¹⁰Generators for d = 0 and d = 4 have nontrivial denominators, which make their 2d theories not sigma models.

 $^{^{11}\}mathrm{Dimension}$ of a target space is set by the degree d of TMF.

¹²For d < 0, a 2d(0, 1) theory is not given by a sigma model [11]

8 Future directions

- For $Z_{g,n}^r$ manifold in which g is odd, torsion generators of π_* tmf appear in the theories considered above. Fiber sum formulas in this case would take qualitatively different form including a multiplication rule.
- Higher rank (Q > 1) E-string theory possesses an extra symmetry and hence, its anomaly polynomial generalizes. The additional symmetry is expected to have nontrivial effects on fiber sum formula.
- Generally in order to find 2d (0,1) theories corresponding to π_* TMF for * > 24, string manifolds whose Witten genus are $E_4\Delta^m$ and $2E_6\Delta^n$, $m, n \in \mathbb{Z}_+$ are needed. Constructions of the string manifolds are not known yet.

Acknowledgments. I am grateful to Sergei Gukov, Andre Henriques, Du Pei, Pavel Putrov, and Michael Hill for the numerous explanations. I would like to thank Sergei Gukov for reading a draft of this paper.

Appendix

A Four manifolds

We review the fiber sum operation between 4-manifolds and summarize the topological data of the 4-manifolds considered in this paper.

A natural generalization of the connected sum operation for (smooth) surface bundles on a surface is the fiber sum [10]. For instance, it can be used to obtain new elliptic surfaces from given elliptic surfaces. Moreover, it was used to construct a family of mutually nondiffeomorphic 4-manifolds and to analyze the symplectic geography problem [1, 29, 30]. It appeared in disproving a conjecture (the minimal conjecture) regarding a classification of smooth closed 4-manifolds [38]. The fiber sum operation between two surface bundles on surfaces X_1 and X_2 is defined as

$$X_1 \#_f X_2 = (X_1 \setminus \nu \Sigma_g) \cup_{\phi} (X_2 \setminus \nu \Sigma'_g)$$

where $\nu \Sigma_g$ and $\nu \Sigma'_g$ are tubular neighborhoods of the surfaces and ϕ is an orientation reversing diffeomorphism.

Elliptic surfaces An elliptic surface is a compact complex surface S in which there is a holomorphic map to a complex curve $C, \pi : S \to C$ such that a generic fiber is a smooth elliptic curve [35]. Elliptic surfaces have been classified up to deformation equivalence. A well-known example is S = E(n)whose generic fiber is T^2 . They are constructed inductively using the fiber sum operation starting from E(1).

$$E(n) = E(n-1) \#_f E(1).$$

Topological data of E(n) are

$$Q_{E(n)} = n(-E_8) \oplus (2n-1)H$$
 $n = \text{even}$

$$\chi(E(n)) = 12n$$
 $\sigma(E(n)) = -8n$ for all n
 $b_2(E(n)) = 12n - 2$ $b_2^+ = 2n - 1$ $b_1 = 0$ for all n

Moreover, E(n) are all simply connected.

<u>Further Surface bundles</u> Using E(n), the knot surgery and the fiber sum operation, infinite families of smooth simply-connected spin symplectic closed 4-manifolds that are mutually nondiffeomorphic were constructed in [1]. They are all irreducible (i.e. they cannot decomposed as a connected sum of two 4-manifolds in which neither of them are homotopy 4-spheres). We summarize the families.

$$Z_{g,n}^r = X_{g,n} \#_{\Sigma_{f+b}} E(2r)_K \qquad g \ge 2 \quad n = \text{odd} \ge 3$$
$$r = \min\left\{\frac{1}{12}g(g-1)(n^2-1)n^{2g-3}, 2 + gn + g(g-1)(n-1)n^{2g-3}\right\}$$

where f is a fiber genus, b is a base genus and $X_{g,n}$ is

$$\begin{array}{c} \Sigma_{gn} \to X_{g,n} \\ \downarrow \\ \Sigma_{1+q(q-1)n^{2g-2}} \end{array}$$

Its signature $\sigma(X_{g,n})$ is $(4/3)g(g-1)(n^2-1)n^{2g-3}$. $E(2r)_K$ is the Fintushel-Stern knot surgery [5] on E(2r) along a fibered knot of a certain genus g(K) in S^3 . The signature and Euler characteristic of $Z_{g,n}^r$ are

$$\sigma(Z_{g,n}^r) = \frac{4}{3}g(g-1)(n^2-1)n^{2g-3} - 16r$$
$$e(Z_{g,n}^r) = 4gn(g(g-1)n^{2g-2}+1) + 24r$$

Another surface bundle analyzed in [1] was

$$V_n^r = X_n \#_{\Sigma_{f+b}} E(2r)_K \qquad n = \text{odd} \ge 3$$
$$r = \min\left\{\frac{1}{6}(n+1)n(n-1), 2n^2 + n + 2\right\}$$

where

$$\begin{array}{c} \Sigma_{3n} \to X_n \\ \downarrow \\ \Sigma_{2n^2+1} \end{array}$$

Its signature $\sigma(X_n)$ is (8/3)(n+1)n(n-1). The signature and Euler characteristic of V_n^r are

$$\sigma(V_n^r) = \frac{8}{3}(n+1)n(n-1) - 16r$$
$$e(V_n^r) = 12n(2n^2+1) + 24r$$

Nonsymplectic manifolds Smooth simply-connected closed oriented 4-manifolds that do not carry symplectic structures $Z(k,m)_n$ were constructed in [38] $(k,m \in \mathbb{Z}_+, n \ge 0)$. They have even intersection form for k, m odd and hence carry spin structures $(w_2 = 0)$. The construction consists three steps. The first part is

$$Y = E(1)\#_f T^4 \#_f E(1)$$

= $(E(1) \setminus ndF_1) \bigcup_{g_1} \left(T^4 \setminus (ndT_{12} \bigcup ndT_{13}) \right) \bigcup_{g_2} (E(1) \setminus ndF_2)$

where F_1 and F_2 are generic fibers in E(1)'s, T_{12} and T_{13} are symplectic submanifolds, g_1 and g_2 are orientation reversing diffeomorphisms and nd denotes an open tubular neighborhood. We again apply the fiber sum operation to Y.

$$Y(k,m) = (E(k) \setminus ndT_1) \bigcup_{g_1} \left(Y \setminus (ndL_1 \bigcup ndL_2) \right) \bigcup_{g_2} (E(m) \setminus ndT_2)$$

The final step is to use the log. transform on Y, we obtain

$$Z(k,m)_n = (E(k) \setminus ndT_1) \bigcup_{g_1} \left(X_n \setminus (ndL_1 \bigcup ndL_2) \right) \bigcup_{g_2} (E(m) \setminus ndT_2)$$

where $X_n = (Y \setminus n dT_{14}) \bigcup_{\phi_n} (T^2 \times D^2)$. Topological invariants of $Z(k, m)_n$ are

$$\chi_h(Z(k,m)_n) = k + m + 4$$
 $\sigma(Z(k,m)_n) = -8(k+m) - 16$

B Anomaly polynomials

We summarize the anomaly polynomials described in [11] (see also [2, 31]). They provide 2-dimensional gravitational anomaly $2(c_L - c_R)$ expression in terms of compact 4-manifolds' topological invariants for 2d (0, 1) theories that are obtained from compactification of 6D (1,0) SCFTs on the 4-manifolds. Perturbative anomaly of 6D (1,0) SCFTs can be characterized by polynomials in characteristic classes of various bundles on 6D spacetime manifold M. Their anomaly polynomials are differential 8-form I_8 . Its formula in our context is

$$I_8 = \alpha c_2(R)^2 + \beta c_2(R)p_1(T) + \gamma p_1(T)^2 + \delta p_2(T) + I^{\text{flavor}}, \quad \alpha, \beta, \delta, \gamma \in \mathbb{R}$$

The second Chern class $c_2(R) \in H^4(R)$ is associated with the R-symmetry bundle R. Pontryagin classes of tangent bundle TM, $p_1(T) \in H^4(T)$ and $p_2(T) \in H^8(T)$ correspond to gravitational anomaly [12]. $c_2(R)p_1(T)$ represents a mixed anomaly between R-symmetry transformation and diffeomorphism of M. The last term contains a flavor symmetry anomaly.

Compactifications to 2-dimensions result in an anomaly polynomial that is a 4-form I_4 . When a background field associated with flavor symmetry group is turned off, 2d gravitational anomaly is given by

$$c_R - c_L = 18(\beta - 8\gamma - 4\delta)\sigma(X) + 12\beta\chi(X)$$

where $\sigma(X)$ is the signature of a 4-manifold X, $\chi(X)$ is the Euler characteristic of X. For the three theories analyzed in this paper, their anomaly polynomials are [2, 27, 28, 11]

- Hypermultiplet $I_8 = \frac{7p_1(T)^2 4p_2(T)}{5760}$
- Vector multiplet $I_8 = -\frac{1}{24}c_2(R)^2 \frac{1}{48}c_2(R)p_1(T) \frac{7p_1(T)^2 4p_2(T)}{5760}$
- Tensor multiplet $I_8 = \frac{1}{24}c_2(R)^2 + \frac{1}{48}c_2(R)p_1(T) + \frac{23p_1(T)^2 116p_2(T)}{5760}$
- E-strings (rank Q = 1) $I_8 = \frac{13}{24}c_2(R)^2 \frac{11}{48}c_2(R)p_1(T) + \frac{29(7p_1(T)^2 p_2(T))}{5760}$

The fourth expression excludes the contribution of the free hypermultiplet.

C Topological modular forms

We summarize properties of topological modular forms (see [3, 15, 14, 9, 20, 11] for details). There are three kinds of topological modular forms, tmf, Tmf, and TMF. The first and the third variants are relevant for this paper.

1. tmf

- Connective property $\pi_{*<0}$ tmf = 0
- Connective cover of Tmf.
- Relation to the stable homotopy groups of spheres and the ring of modular forms

$$\pi^s_* \mathbb{S} \to \pi_* \operatorname{tmf} \to \operatorname{MF}_{\frac{*}{2}}$$

The first ring homomorphism is an isomorphism for $n = 0, 1, \dots, 6$. The second map is the boundary homomorphism/the elliptic genus. (it maps to zero for * odd, thus the kernel is the torsion part).

- Relation with string manifolds $\Omega_*^{\text{string}} \to \pi_* \text{tmf}$: every tmf classes comes from a bordism class of string manifold (this surjective map is the topological Witten genus)¹³. Ω_*^{string} is a *-dimensional string (co)bordism group.
- Witten genus of a string manifold $\phi_W(Y)$ is

$$\pi_* \mathrm{MString} \to \pi_* \mathrm{tmf} \to \mathrm{MF}_{\frac{*}{2}}$$

It is factored by the elliptic genus (MString is the Thom spectrum).

2. TMF

- Localization $\pi_* \text{TMF} \simeq \pi_* \text{tmf} \left[\Delta^{-24} \right]$
- Periodicity $\pi_* \text{TMF} \simeq \pi_{*+576} \text{TMF}$ ($\Rightarrow \pi_{*<0} \text{TMF}$ can be nontrivial).
- $\pi_n \text{TMF} \supseteq \mathbb{Z}[x] \text{ for } n \equiv 0,4 \mod 8$, $\pi_n \text{TMF} \supseteq (\mathbb{Z}/2\mathbb{Z})[x] \text{ for } n \equiv 1,2 \mod 8$.

¹³More precisely, this map $\pi_*MString \to \pi_*tmf$ is a surjective homomorphism

Relations between generators

$$\pi_1(\mathbb{S}) = \mathbb{Z}/2\mathbb{Z} \langle \eta \rangle \qquad \pi_3(\mathbb{S}) = \mathbb{Z}/24\mathbb{Z} \langle \nu \rangle$$
$$\eta^3 = 12\nu \qquad 2\eta = 24\nu = 2\nu^2 = \eta^4 = \nu^4 = 0$$

 η is associated to the S^3 -Hopf fibration, $S^1 \to S^3 \to S^2$ and ν corresponds to the S^7 -Hopf fibration, $S^3 \to S^7 \to S^4$.

String manifold We briefly review string manifolds (see [3, 4] for details). A manifold equipped with a string structure is called a string manifold Y. Equivalent characterizations of the structure are

• Lifting of the map τ to the classifying space BO(n):

$$\begin{array}{c} BString(n) \\ \downarrow \\ Y \xrightarrow{\tau} BO(n) \end{array}$$

where String(n) is the 6-connected cover of O(n).

• An extension of a spin structure of Y: $\frac{p_1(TY)}{2} = 0(w_1 = w_2 = 0).$

The string condition is an orientablility condition for a generalized cohomology theory, which is TMF in our context.

<u>Construction</u> We summarize on the construction of Y_8 described in [11]¹⁴.

- 1. Plumbing and parallelizable manifold: place a twisted D^4 -bundle E^8 on D^4 at each vertex of the $-E_8$ -graph. Applying the plumbing along each edge yields a 8-dimensional smooth manifold W^8 with an exotic S^7 -boundary [23]. Since the S^7 represents an element of a subgroup $bP_8 \subset \Theta_7$, W is parallelizable [18, 24].
- 2. Boundary sum and gluing: the cobordism group of homotopy 7-spheres is $\Theta_7 = \mathbb{Z}/28\mathbb{Z}$ [18], we take a boundary sum of 28 copies of W^8 and then glue an D^8 along the S^7 to obtain $Y_8 = \natural^{28} W \cup D^8$. The signature of Y_8 is $\sigma = -8 \times 28$.
- 3. Pontryagin number: W being parallelizable implies that Y_8 is almost parallelizable, which in turn means that only the top Pontryagin class p_2 of Y_8 is nonzero [19]. Hence, Y_8 carries a string structure (i.e. $p_1(Y_8)/2 = 0$). By the Hirzebruch signature theorem $\sigma(Y_8) = \langle L(p_1, p_2), [Y_8] \rangle$, $p_2 = -1440$ (only L_2 contributes).
- 4. \hat{A} -genus and the Witten genus ϕ_W : The above information enables to find $\hat{A}_2 = \frac{1}{5760}(-4p_2 + p_1^2) = 1$, which agrees with the leading term of $\phi_W(Y_8) = E_4$.

 $^{^{14}}Y_{12}$ is constructed similarly.

d	$\pi_d \text{TMF}$	$2d \mathcal{N} = (0,1)$ theory generators
16	$\mathbb{Z}[x]$	$\sigma(Y_8 imes Y_8)$
17	$\mathbb{Z}/2\mathbb{Z} \oplus (\mathbb{Z}/2\mathbb{Z})[x]$	$\sigma(SU(2) \times G_2), \sigma(S^1 \times Y_8 \times Y_8)$
18	$\mathbb{Z}/2\mathbb{Z}[x]$	$\sigma(T^2 imes Y_8 imes Y_8)$
19	0	-
20	$\mathbb{Z}/24\mathbb{Z}\oplus\mathbb{Z}[x]$	$\sigma(Sp(2) \times Sp(2)), \sigma(Y_8 \times Y_{12})$
21	$\mathbb{Z}/2\mathbb{Z}$	$\sigma(Sp(2) \times Sp(2) \times U(1))$
22	$\mathbb{Z}/2\mathbb{Z}$	$\sigma(SU(3) imes G_2)$
23	0	-
28	$\mathbb{Z}/2\mathbb{Z}\oplus\mathbb{Z}[x]$	$\sigma(G_2 \times G_2), \sigma(Y_8 \times Y_8 \times Y_{12})$
29	0	-
30	$\mathbb{Z}/3\mathbb{Z}$	$\sigma(Sp(2)\times Sp(2)\times Sp(2))$
31	0	-
34	$\mathbb{Z}/2\mathbb{Z} \oplus (\mathbb{Z}/2\mathbb{Z})[x]$	$\sigma(Sp(2) \times Sp(2) \times G_2), \sigma(T^2 \times Y_{32})$
-21	0	-
-22	$(\mathbb{Z}/2\mathbb{Z})[x]$	$\sigma(U(1)) \otimes T_{(B_2;-23)}$
-23	$(\mathbb{Z}/2\mathbb{Z})[x]$	$T_{(B_2;-23)}$
-45	$\mathbb{Z}/24\mathbb{Z}$	$T_{(B_{22};1)}$

Table 9: This table lists π_d TMF and corresponding 2d (0, 1) supersymmetric QFTs. It fills the gaps in the table in [11] and extend it. Although construction of Y_{32} is yet unknown, it is an element of the string bordism group Ω_*^{string} by the surjective property of the map to π_* tmf (see the review above). The second column can be obtained using [3] (Chapter 13).

References

- A. Akhmedov, B. Park, Geography of simply connected spin symplectic 4-manifolds, *Math. Res. Lett.* 17 (2010), no. 3, 483–492,
- [2] C. Cordova, T.T. Dumitrescu and K. Intriligator, Anomalies, renormalization group flows and the a-theorem in six-dimensional (1,0) theories, *Journal of High Energy Physics*, volume 2016, 80 (2016), arXiv:1506.03807
- [3] C.L. Douglas, J. Francis, A.G. Henriques, M.A. Hill, Topological modular forms, American Mathematical Soc., vol. 201 (2014).
- [4] C.L. Douglas, A.G. Henriques, M.A. Hill, Homological Obstructions to String Orientations, International Mathematics Research Notices, Volume 2011, Issue 18, 2011, Pages 4074–4088, arXiv:0810.2131
- [5] R. Fintushel, R. Stern, Knots, links, and 4-manifolds, *Inventiones mathematicae* volume 134, pages363–400 (1998) arXiv:dg-ga/9612014
- [6] D. Gaiotto, T. Johnson-Freyd, Holomorphic SCFTs with small index, Canadian Journal of Mathematics, Vol 74, Issue 2, arXiv:1811.00589
- [7] D. Gaiotto, T. Johnson-Freyd, Mock modularity and a secondary elliptic genus, arXiv:1904.05788
- [8] D. Gaiotto, T. Johnson-Freyd, E. Witten, A Note On Some Minimally Supersymmetric Models In Two Dimensions, arXiv:1902.10249
- [9] P.G. Goerss, Topological modular forms, Seminaire Bourbaki (2009) 61, arXiv:0910.5130
- [10] R. Gompf,, A new construction of symplectic manifolds, Ann. of Math. 142 (1995), 527–595.
- [11] S. Gukov, D. Pei, P. Putrov and C. Vafa, 4-manifolds and topological modular forms, *Journal of High Energy Physics* volume 2021, 84 (2021), arXiv:1811.07884
- [12] L. Alvarez-Gaume, E. Witten, Gravitational anomalies, Nuclear Physics B, Volume 234, Issue 2, 2 April 1984, Pages 269-330
- [13] J. Heckman, T. Rudelius, Top down approach to 6D SCFTs, J. Phys. A: Math. Theor. 52 093001, arXiv:1805.06467
- [14] A. Henriques, Examples of stolz-teichner cocycles, in Oberwolfach, (2015), http://www.andreghenriques.com/PDF/OW-report-STcocycles.pdf
- [15] M. Hopkins, Algebraic topology and modular forms, Proceedings of the International Congress of Mathematicians, vol. I Beijing (2002), arXiv:math/0212397
- [16] K. Intriligator, 6dN = (1,0) Coulomb branch anomaly matching, JHEP, 2014, 162
- [17] J. Kim, a S. Kim, K. Lee, J. Park, C. Vafa, Elliptic genus of E-strings, JHEP, 2017, 98

- [18] M. Kervaire, J. Milnor, Groups of Homotopy Spheres 1, Annals of Mathematics, Vol. 77, No. 3, May, 1963
- [19] M. Kervaire, J. Milnor, Bernoulli numbers, homotopy groups, and a theorem of Rohlin, 1960 Proc. Internat. Congress Math. 1958.
- [20] J. Lurie, A Survey of Elliptic Cohomology, Algebraic Topology, Abel Symposia book series.
- [21] Y. Lin, D.Pei, Holomorphic CFTs and topological modular forms, arXiv:2112.10724
- [22] M. Mahowald, M. Hopkins, The structure of 24 dimensional manifolds having normal bundles which lift to BO[8], Recent progress in homotopy theory, Contemp. Math. 293, Amer. Math. Soc., Providence, RI (2002), 89-110
- [23] J. Milnor, On Manifolds Homeomorphic to the 7-Sphere, Annals of Mathematics, Vol. 64, No. 2, Sep., 1956.
- [24] J. Milnor, Differentiable manifolds which are homotopy spheres, https://www.maths.ed.ac.uk/ v1ranick/papers/homsph.pdf
- [25] S. Minwalla, Restrictions imposed by superconformal invariance On quantum field theories, Adv. Theor. Math. Phys., 2 (1998) 783-851.
- [26] W.Nahm, Supersymmetries and their representations, Nuclear Physics B Volume 135, Issue 1, 27 March 1978, Pages 149-166
- [27] K. Ohmori, H. Shimizu, Y. Tachikawa, Anomaly polynomial of E-string theories, Journal of High Energy Physics, volume 2014, 2 (2014), arXiv:1404.3887
- [28] K. Ohmori, H. Shimizu, Y. Tachikawa, K. Yonekura, Anomaly polynomial of general 6D SCFTs, *Progress of Theoretical and Experimental Physics*, Volume 2014, Issue 10, October 2014, arXiv:1408.5572
- [29] J. Park, The geography of spin symplectic 4-manifolds, Math. Z., 240,405–421 (2002)
- [30] B. Park, Z. Szabo, THE GEOGRAPHY PROBLEM FOR IRREDUCIBLE SPIN FOUR-MANIFOLDS, TRANSACTIONS OF THE AMERICAN MATHEMATICAL SOCIETY, Volume 352, Number 8, Pages 3639–3650,
- [31] S. Razamat, E. Sabag, O. Sela, G. Zafrir, Aspects of 4d supersymmetric dynamics and geometry, arXiv:2203.06880
- [32] G. Segal, What is an elliptic object?, Elliptic Cohomology: Geometry, Applications, and Higher Chromatic Analogues Cambridge University Press, 2007
- [33] S.El-Showk, Y. Nakayama, S. Rychkov, What Maxwell theory in $d \neq 4$ teaches us about scale and conformal invariance, *Nuclear Physics B* 848 (2011) 578–593.
- [34] N. Seiberg, E. Witten, Comments on string dynamics in six dimensions, Nuclear Physics B 471 (1996) 121-134.

- [35] A. Stipsicz, Z. Szabo, The smooth classification of elliptic surfaces with $b^+ > 1$, Duke Math. J. 75(1): 1-50 (July 1994).
- [36] S. Stolz and P. Teichner, What is an elliptic object?, London Mathematical Society Lecture Note Series, Cambridge University Press (2004), pp. 247–343
- [37] S. Stolz and P. Teichner, Proceedings of Symposia in Pure Math. 83, 279 (2011), Supersymmetric field theories and generalized cohomology, arXiv:1108.0189v1.
- [38] Z. Szabo, Simply-connected irreducible 4-manifolds with no symplectic structures, *Invent. math.* 132, 457±466 (1998).
- [39] Y. Tachikawa, Topological modular forms and the absence of a heterotic global anomaly, *Progress of Theoretical and Experimental Physics*, Volume 2022, Issue 4, April 2022, arXiv:2103.12211
- [40] Y. Tachikawa, M. Yamashita, Topological modular forms and the absence of all heterotic global anomalies, arXiv:2108.13542
- [41] E. Witten, Elliptic genera and quantum field theory, Commun. Math. Phys. 109(4): 525-536 (1987).