

QUANTUM BIANCHI-VII PROBLEM, MATHIEU FUNCTIONS AND ARITHMETIC

A.P. VESELOV AND Y. YE

ABSTRACT. The geodesic problem on the compact threefolds with the Riemannian metric of Bianchi-VII₀ type is studied in both classical and quantum cases. We show that the problem is integrable and describe the eigenfunctions of the corresponding Laplace-Beltrami operators explicitly in terms of the Mathieu functions with parameter depending on the lattice values of some binary quadratic forms. We use the results from number theory to discuss the level spacing statistics in relation with the Berry-Tabor conjecture and compare the situation with Bianchi-VI₀ case (*Sol*-case in Thurston's classification) and with Bianchi-IX case, corresponding to the classical Euler top.

1. INTRODUCTION

Let M^3 be a closed Riemannian 3-fold and consider the corresponding geodesic problem. It is natural to expect that the question of its integrability is connected with special geometric properties of M^3 .

One such property is local homogeneity, meaning that any two points have isometric neighbourhoods. Under certain additional assumptions Thurston classified all special geometries in dimension 3, showing that there are 8 of them: the Euclidean E^3 , spherical S^3 and hyperbolic \mathbb{H}^3 , the product type $S^2 \times \mathbb{R}$ and $\mathbb{H}^2 \times \mathbb{R}$ and three geometries related to the 3D Lie groups: *Nil*, *Sol* and $\widetilde{SL}(2, \mathbb{R})$, where the last group is the universal covering of $SL(2, \mathbb{R})$ (see [30]). The topology of the corresponding compact quotient threefolds is discussed in [28].

On the other hand, Bianchi [2] classified all the three-dimensional Lie groups, which provide the most natural class of locally homogeneous manifolds if we supply them with a left-invariant metric. The correspondence between Thurston's and Bianchi's classification was discussed in relation with the cosmological models in [15, 16] (see the next section).

The geodesic flows for the 3-folds with *Nil*, *Sol* and $\widetilde{SL}(2, \mathbb{R})$ Thurston's geometries (corresponding to Bianchi VI₀ and VIII types respectively) were studied in [9, 6, 7], which revealed the coexistence of the chaos and integrability in these systems. The quantum version of the *Sol*-case was studied in [5] in relation with quantum monodromy and arithmetic. The general Bianchi-IX case is well-known in classical mechanics as Euler top [1] and its quantum version was studied in [21] (see also [17]).

In this paper we will discuss the last remaining Bianchi-VII₀ case. The corresponding Lie group is group $E(2) = SO(2) \times \mathbb{R}^2$ of orientation preserving isometries of Euclidean plane. This case is missing in Thurston's approach since this isometry group is not maximal and can be extended to the group $E(3)$, corresponding to the 3D Euclidean geometry.

The classical system describing geodesics of left-invariant metric on $E(2)$ is completely integrable, as it was already shown by Bolsinov and Bao [4]. We will show that, in contrast to the *Sol*-case [6], the integrability remains on the compact quotients $M_n^3 = E(2)/G_n$ of $E(2)$ by a discrete crystallographic subgroup $G_n = \mathbb{Z}_n \times \mathcal{L} \subset E(2)$, where \mathbb{Z}_n is generated by an element $R \in SO(2)$, which is an order n rotational symmetry of the corresponding 2D lattice $\mathcal{L} \subset \mathbb{R}^2$.

There are 5 types of such threefolds corresponding to possible values of $n = 1, 2, 3, 4, 6$ (see e.g. [10]). They can be viewed as torus bundles over S^1 with gluing map given by an elliptic element $A \in SL(2, \mathbb{Z})$. In that sense they are similar to the *Sol*-manifolds M_A^3 with gluing map A being a hyperbolic element of $SL(2, \mathbb{Z})$ (see [5]). However, we will see that the difference between these two cases is quite substantial. In particular, in contrast to the *Sol*-manifolds with geodesic flow being non-integrable in analytic category and having positive topological entropy [5, 6], in the Bianchi-VII case we have completely integrable geodesic flow on M_n^3 with integrals polynomial in momenta and an explicit integration in terms of the elliptic functions.

We show that in the quantum case the spectral problem for the corresponding Laplace-Beltrami operator on M_n^3 can be explicitly solved in terms of the classical Mathieu functions [31]. However, the properties of the spectrum depend on the order n of the gluing map R . We show that in the symmetric case M_n^3 with $n = 3, 4, 6$ the spectrum has nice arithmetic properties similar to the *Sol*-case studied in [5] and related to the classical problem of describing the values of certain integer quadratic forms Q on two-dimensional integer lattice.

In particular, for $n = 4$ corresponding to the square lattice the parameters of the Mathieu equation and the corresponding eigenvalues depend on the values of the quadratic form $Q(x, y) = x^2 + y^2, x, y \in \mathbb{Z}$. The study of these values is a very classical part of number theory with famous contributions from Fermat, Euler, Gauss and Jacobi. A classical result due to E. Landau [22] says that the number of integers up to a number K represented by this form grows as $K/\sqrt{\log K}$. This means that the spectrum of the corresponding symmetric Bianchi threefold M_4^3 is highly degenerate and the corresponding level space statistics does not satisfy the Berry-Tabor conjecture [3]. Similar results hold for $n = 3, 6$ related to hexagonal lattice and the integer values of the quadratic form $Q(x, y) = x^2 + xy + y^2$.

For $n = 1, 2$ the situation is more delicate and depends on the geometry of the lattice and on the diophantine properties of the corresponding quadratic form Q . In particular, from the deep results of Sarnak [27] and Eskin, Margulis and Mozes [14] it follows that the pair correlation of the corresponding

values does satisfy Berry-Tabor prediction on a set of full measure in the space of quadratic forms, in particular for $Q(x, y) = x^2 + \beta y^2$ with diophantine β . Our results suggest that the same should be true for the spectrum of the corresponding Bianchi-VII threefolds.

Finally, in the last section we make a comparison with the situation in Bianchi-VI and Bianchi-IX cases.

2. GEODESIC FLOW ON BIANCHI-VII₀ THREEFOLDS.

In 1898 Bianchi [2] completed the classification of 3D real Lie algebras by splitting them into 9 types: I-IX. Since we are interested in compact quotients of the corresponding Lie groups, we must consider only the Lie algebras admitting bi-invariant volume form, which are called *unimodular*. In Bianchi's list the unimodular Lie algebras are of types

- *I: abelian* \mathbb{R}^3
- *II: nilpotent (Heisenberg)* *nil*
- *VI₀: solvable 2D Poincare* *sol* $\cong e(1, 1)$
- *VII₀: solvable 2D Euclidean* *e(2)*
- *VIII: simple* $sl(2, \mathbb{R})$
- *IX: simple* $so(3)$.

The curvature of the corresponding left-invariant metrics was studied in details by Milnor in [25].

One can compare this list with Thurston's list of 8 special geometries in 3 dimensions [30]: *Euclidean* E^3 , *spherical* S^3 , *hyperbolic* \mathbb{H}^3 , *the product type* $S^2 \times \mathbb{R}$ and $\mathbb{H}^2 \times \mathbb{R}$ and *three geometries related to 3D Lie groups* *Nil*, *Sol*, $SL(2, \mathbb{R})$.

Recall that special geometry here means local homogeneity (any two points have isometric neighbourhoods), existence of compact models and some maximality condition. Note that general Bianchi VII₀ and IX types are missing in Thurston's list since they do not satisfy the maximality condition having more symmetric versions: Euclidean E^3 and spherical S^3 respectively.

The integrability of the geodesic flow on 3D manifolds with Thurston's geometries of group types *Nil*, *Sol*, $SL(2, \mathbb{R})$ (corresponding to Bianchi types II, VI₀, VIII) was studied by Butler [9], Bolsinov and Taimanov [6] and by Bolsinov and the authors [7] respectively. The quantum *Sol*-case (or, equivalently, Bianchi-VI₀ case) was investigated by Bolsinov, Dullin and one of the authors in [5].

The general Bianchi-IX₀ problem, corresponding to the geodesic flow on the group $SO(3)$ with left-invariant metric, is well known in classical mechanics as Euler top [1]. Its quantum version was studied at the very early age of quantum mechanics by Kramers and Ittmann [21], who discovered close relations with the famous Lamè equation (see also [17]). In Thurston's list Bianchi-IX case is represented only by its most symmetric version S^3 , which can be considered as the group $SU(2)$ with bi-invariant metric.

In this paper we will study the only remaining quantum Bianchi-VII₀ case, which has many parallels with Bianchi-VI₀ case, but deserves a special investigation.

We follow here the notations of Bolsinov and Bao [4], where one can find also the relevant references about geodesic flows on 3 dimensional Lie groups.

The Lie algebras of Bianchi type VII have 3 generators X_0, X_1, X_2 , satisfying the relations

$$[X_0, X_1] = aX_1 - X_2, [X_0, X_2] = X_1 + aX_2, [X_1, X_2] = 0,$$

depending on a parameter a . These Lie algebras are solvable, but for $a \neq 0$ are not unimodular.

When $a = 0$ we have unimodular Bianchi subcase VII₀, corresponding to the Lie group $E(2)$ of isometries of the Euclidean plane. The Lie algebra $e(2)$ is the semi-direct product of \mathbb{R}^2 by \mathbb{R} with 3 generators X_0, X_1, X_2 , satisfying the relations

$$[X_0, X_1] = -X_2, [X_0, X_2] = X_1, [X_1, X_2] = 0. \quad (1)$$

The corresponding left-invariant vector fields are

$$X_0 = \partial_0, X_1 = \cos q_0 \partial_1 - \sin q_0 \partial_2, X_2 = \sin q_0 \partial_1 + \cos q_0 \partial_2,$$

where $\partial_i = \frac{\partial}{\partial q_i}$, $i = 0, 1, 2$.

The Hamiltonians of the geodesic flow corresponding to a left-invariant metric can be reduced (see [4]) to the form

$$H = \frac{1}{2}(AX_0^2 + BX_1^2 + X_2^2), \quad (2)$$

where X_0, X_1, X_2 are considered as linear functions on the dual space $e(2)^*$. In the canonical coordinates (p, q) we have

$$H = \frac{1}{2}(Ap_0^2 + p_1^2 + p_2^2 + (B-1)(p_1 \cos q_0 - p_2 \sin q_0)^2), \quad (3)$$

where $A, B > 0$ are two real parameters.

The system is clearly completely integrable in Liouville sense with the commuting integrals H and $F_1 = p_1, F_2 = p_2$.

Corresponding system can be easily integrated in quadratures (see [4]). Indeed, fixing the values of the integrals $F_1 = p_1 = c_1, F_2 = p_2 = c_2$ and assuming without loss of generality that $c_2 = 0$, we can write the equation of motion as

$$\begin{cases} \dot{q}_0 = Ap_0 \\ \dot{p}_0 = (B-1)c_1^2 \sin q_0 \cos q_0 = \frac{1}{2}(B-1)c_1^2 \sin 2q_0 \\ \dot{q}_1 = c_1 + (B-1)c_1 \cos^2 q_0 = c_1 \sin^2 q_0 + Bc_1 \cos^2 q_0 \\ \dot{q}_2 = -(B-1)c_1 \sin q_0 \cos q_0 = -\frac{1}{2}(B-1)c_1 \sin 2q_0 \end{cases} \quad (4)$$

with $\dot{p}_1 = \dot{p}_2 = 0$. Note that the first two equations are simply the equations of motion of the mathematical pendulum with

$$H_0 = \frac{1}{2}p_0^2 + C \cos 2q_0, \quad C = \frac{1}{4}(B-1)c_1^2, \quad (5)$$

which can be integrated explicitly in terms of the Jacobi elliptic functions. Qualitatively, if the energy $H_0 = h_0 > |C|$ then we have rotations, while for $|h_0| < |C|$ the orbits are periodic. For the critical levels $h_0 = \pm|C|$ we have separatrices and the stable equilibriums respectively (see Fig.1).




FIGURE 1. Phase portrait of mathematical pendulum

For known $(p_0(t), q_0(t))$ we can find $q_1(t)$ and $q_2(t)$ by integrating the last two equations, which also can be done explicitly in terms of the Jacobi elliptic functions. Note that since $B > 0$ for positive c_1 \dot{q}_1 is always positive, so q_1 monotonically increases, while $\dot{q}_2 = -c_1^{-1}\dot{p}_0$, so

$$q_2 = -c_1^{-1}p_0 + const$$

is always bounded.

This describes the corresponding geodesics on the group $E(2)$.

Consider now the compact case $M_n^3 = E(2)/G_n$, $G_n = \mathbb{Z}_n \times \mathcal{L} \subset E(2)$, where \mathbb{Z}_n is generated by matrix $R_n \in SO(2)$, defining the rotational symmetry of the corresponding 2D lattice $\mathcal{L} \subset \mathbb{R}^2$. More precisely, in types 1 and 2 with generic lattice $R_1 = I$ and $R_2 = -I$, in types 3, 4 and 6, corresponding to square and hexagonal lattices

$$R_3 = \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{pmatrix}, R_4 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, R_6 = \begin{pmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}. \quad (6)$$

respectively. For $n = 1$ we have the *Bianchi torus*

$$M_1^3 = T_{\mathcal{L}}^3 = E(2)/\mathcal{L},$$

where $\mathcal{L} \subset \mathbb{R}^2(q_1, q_2)$ is any two-dimensional lattice, with the corresponding left-invariant $E(2)$ -metrics

$$ds^2 = A^{-1}dq_0^2 + B^{-1}(dq_1^2 + dq_2^2 + (B-1)(\sin q_0 dq_1 + \cos q_0 dq_2)^2). \quad (7)$$

In other cases $M_n^3 = T_{\mathcal{L}}^3/\mathbb{Z}_n$ are simply quotients of the Bianchi torus under additional assumption of rotational symmetry R of the lattice \mathcal{L} .

Proposition 1. *The geodesic flow on Bianchi-VII₀ threefolds $M_n^3 = T_{\mathcal{L}}^3/G_n$ is integrable in Liouville sense with three polynomial in momenta Poisson commuting integrals.*

Indeed, on the quotient threefolds M_n^3 as 3 independent Poisson commuting integrals of the geodesic problem one can choose Hamiltonian (3) and $F_1 = p_1, F_2 = p_2$ for the general case $n = 1$ and G_n -invariant functions of momentum in the symmetric cases: $F_1 = p_1^2, F_2 = p_2^2$ for $n = 2$, $F_1 = p_1^2 + p_2^2, F_2 = p_1^3 - 3p_1p_2^2$ for $n = 3$, $F_1 = p_1^2 + p_2^2, F_2 = p_1^2p_2^2$ ($n = 4$), $F_1 = p_1^2 + p_2^2, F_2 = (p_1^3 - 3p_1p_2^2)^2$ ($n = 6$).

To explain the behaviour of the corresponding geodesics, it is enough to consider the case of the Bianchi torus M_1^3 . Let us assume first that the lattice $\mathcal{L} = 2\pi\mathbb{Z}^2 \subset \mathbb{R}^2(q_1, q_2)$ and that the integral $p_2 = c_2 = 0$, but $p_1 = c_1 \neq 0$. Since q_2 is bounded, the corresponding geodesics will stay near the subtorus $T^2 \subset T_{\mathcal{L}}^3$ with $q_2 = 0$. To describe qualitatively the corresponding projections (q_0, q_1) of the geodesics consider a periodic surface of revolution in cylindrical coordinates (ρ, φ, z) in \mathbb{R}^3 having the form $\rho = f(z)$ with periodic $f(z) = a + b\sin^2 z$ with positive a, b .

The geodesics on surfaces of revolutions satisfy the classical *Clairault's relation*:

$$C := \rho \cos \theta = C \tag{8}$$

is constant along the geodesics, where θ is the angle between the geodesic and parallel (see e.g. [11]). In our case we have two special geodesics, which are parallels with maximal and minimal length (which we call equator and neck respectively). We have 5 different cases:

$$C = R, r < C < R, C = r, 0 < C < r, C = 0$$

where $r = a$ and $R = a + b$ are radii of neck and equator respectively. The behaviour of the corresponding geodesics is shown on Fig 2. In the critical case $C = r$ we have closed neck geodesics and the geodesics spiralling towards the neck.




FIGURE 2. Geodesics on periodic surface of revolution

If we put here $q_0 = z \pmod{2\pi}$, $q_1 = \varphi$, then we get a qualitative description of the motion of the projected geodesics (q_0, q_1) on the torus T^2 . More precisely, the first four cases above correspond to the four cases

$$2H = p^2, p^2 < 2H < Bp^2, 2H = Bp^2, 2H > Bp^2,$$

where we assume that $p^2 := p_1^2 + p_2^2 > 0$ and $B > 1$. When $p^2 = 0$ (meaning that $p_1 = p_2 = 0$) we have the last (meridian) case with $C = 0$, corresponding to geodesics with constant q_1, q_2 and uniformly changing q_0 .

Interestingly enough, in the *Sol*-case [6] it was the analogue of the last family of simplest geodesics on M_A^3 , which led to the chaos and positive topological entropy because of the Anosov gluing map $A : T^2 \rightarrow T^2$. In our case the gluing map R_n is simply a rotation of order n , which creates no problems for integrability.

For a generic lattice \mathcal{L} the plane $q_2 = 0$ is embedded in the Bianchi torus quasi-periodically, so for generic values of integrals with $2H > B(p_1^2 + p_2^2)$ the geodesics are quasi-periodic and dense on the torus (similarly to the flat case).

Note that for $B = 1$ we do have a flat metric, corresponding to the 3D Euclidean geometry in Thurston's classification. Bianchi geometry (7) can be viewed as special deformation of this geometry with the smaller symmetry group $E(2) \subset E(3)$, which is not maximal and thus is missing in Thurston's classification.

We consider now the quantum situation, which more essentially depends on the geometry of the lattice \mathcal{L} (as we know already in the flat torus case).

3. SPECTRAL PROBLEM ON BIANCHI-VII₀ THREEFOLDS

Consider first the case of Bianchi torus $T_{\mathcal{L}}^3$ with metric (7). Let

$$v_1 = \omega_1^1 q_1 + \omega_1^2 q_2, \quad v_2 = \omega_2^1 q_1 + \omega_2^2 q_2$$

be a basis in the corresponding dual lattice $2\pi\mathcal{L}^*$:

$$2\pi\mathcal{L}^* = \{k_1(\omega_1^1 q_1 + \omega_1^2 q_2) + k_2(\omega_2^1 q_1 + \omega_2^2 q_2), \quad (k_1, k_2) \in \mathbb{Z}^2\}.$$

By definition, the elements of \mathcal{L}^* take integer values on the lattice \mathcal{L} , so

$$(k, \Omega q) := k_1(\omega_1^1 q_1 + \omega_1^2 q_2) + k_2(\omega_2^1 q_1 + \omega_2^2 q_2) \in 2\pi\mathbb{Z} \quad (9)$$

for all $(k_1, k_2) \in \mathbb{Z}^2$ and $(q_1, q_2) \in \mathcal{L}$.

The corresponding Laplace-Beltrami operator on $T_{\mathcal{L}}^3$ has the form

$$\Delta = A \frac{\partial^2}{\partial q_0^2} + \frac{\partial^2}{\partial q_1^2} + \frac{\partial^2}{\partial q_2^2} + (B - 1) \left(\cos q_0 \frac{\partial}{\partial q_1} - \sin q_0 \frac{\partial}{\partial q_2} \right)^2. \quad (10)$$

This is a self-adjoint operator in the Hilbert space $L_2(T_{\mathcal{L}}^3)$ of functions on the torus. The spectral geodesic problem is to study the spectrum of Δ in this space:

$$-\Delta\Psi = \mathcal{E}\Psi. \quad (11)$$

Because the coefficients of Δ depend only on q_0 , we separate variables and look for the eigenfunctions of Δ of the form

$$\Psi_k(q) = e^{i(k, \Omega q)} \psi(q_0), \quad k = (k_1, k_2) \in \mathbb{Z}^2. \quad (12)$$

The condition (9) guarantees that the corresponding function is well-defined in the Bianchi torus.

Substituting this into (11) and denoting $x = q_0$ we see that the $\psi(x)$ must satisfy the equation

$$-A \frac{d^2\psi}{dx^2} + [Q(k) + (B-1)((\omega_1^1 k_1 + \omega_2^1 k_2) \cos x - (\omega_1^2 k_1 + \omega_2^2 k_2) \sin x)^2] \psi(x) = \mathcal{E} \psi(x),$$

where

$$Q(k) = (\omega_1^1 k_1 + \omega_2^1 k_2)^2 + (\omega_1^2 k_1 + \omega_2^2 k_2)^2. \quad (13)$$

Define the angle $\alpha = \alpha(k)$ by the relations

$$\omega_1^1 k_1 + \omega_2^1 k_2 = \sqrt{Q(k)} \cos \alpha, \quad \omega_1^2 k_1 + \omega_2^2 k_2 = \sqrt{Q(k)} \sin \alpha,$$

then this equation will become the classical Mathieu equation (shifted by α)

$$-\frac{d^2\psi}{dx^2} + \mu \cos 2(x + \alpha) \psi(x) = \lambda \psi(x), \quad (14)$$

where

$$\mu := \frac{(B-1)Q(k)}{2A}, \quad \lambda := \frac{2\mathcal{E} - (B+1)Q(k)}{2A}. \quad (15)$$

Proposition 2. *A function Ψ of the form (12) is an eigenfunction of the Laplace-Beltrami operator (10) if and only if $\psi(x)$ satisfies the Mathieu equation (14) with parameters (15).*

Recall that the classical *Mathieu equation* [24] has the form

$$\frac{d^2y}{dx^2} + (\lambda - \mu \cos 2x)y = 0. \quad (16)$$

Its solutions are known as *Mathieu functions*, see Whittaker-Watson [31] and, for historical perspective, recent review [8]. We should mention that our notation of the parameters λ and μ are different from a and q (used by Whittaker and Watson [31]) and h and θ (used by Erdelyi [13]).

We will be interested only in 2π -periodic Mathieu functions, which are also eigenfunctions of the Mathieu operator

$$L = -\frac{d^2}{dx^2} + \mu \cos 2x \quad (17)$$

on the circle $S^1 = \mathbb{R}/2\pi i\mathbb{Z}$. For given μ there are infinite number of values of $\lambda = \lambda_l(\mu)$, $l = 0, 1, \dots$, for which Mathieu equation has periodic solutions, denoted in a properly normalised form as $ce_m(x, \mu)$, $se_m(x, \mu)$, see [8, 13].

When $\mu = 0$ the spectrum is degenerate: $\lambda = m^2$, $m = 0, 1, \dots$ with two-dimensional space of eigenfunctions generated by $\cos mx$, $\sin mx$ (except for $m = 0$ corresponding to the ground state $\varphi \equiv 1$). The eigenfunctions $ce_m(x, \mu)$, $se_m(x, \mu)$ with the eigenvalues a_m and b_m respectively are analogues of $\cos mx$ and $\sin mx$ for $\mu \neq 0$ having the following properties:

$$\begin{aligned} ce_m(-x, \mu) &= ce_m(x, \mu), & se_m(-x, \mu) &= -se_m(x, \mu), \\ ce_m(x + \pi, \mu) &= (-1)^m ce_m(x, \mu), & se_m(x + \pi, \mu) &= (-1)^m se_m(x, \mu). \end{aligned}$$




FIGURE 3. First Mathieu functions $ce_m(x, \mu)$, $se_m(x, \mu)$ for $\mu = 1$.

For $\mu < 0$ the eigenvalues $a_m = a_m(\mu)$, $b_m = b_m(\mu)$ satisfy the inequalities

$$a_0 < b_1 < a_1 < b_2 < a_2 < b_3 < a_3 \dots,$$

while for $\mu > 0$ we have

$$a_0 < a_1 < b_1 < b_2 < a_2 < a_3 < b_3 < \dots$$

(see [13]). Asymptotically for large m $a_m(\mu)$ and $b_m(\mu)$ are exponentially close to m^2 .

We denote these eigenvalues in the increasing order as $\lambda_l(\mu)$, $l \in \mathbb{Z}_{\geq 0}$ and the eigenfunctions respectively as $\varphi_l(x, \mu)$:

$$L\varphi_l(x, \mu) = \lambda_l(\mu)\varphi_l(x, \mu). \quad (18)$$

Let

$$\Psi_{k,l}(q) := e^{i(k, \Omega q)} \varphi_l(q_0, \mu) \quad (19)$$

be the corresponding eigenfunctions of the Laplace-Beltrami operator on Bianchi torus $T_{\mathcal{L}}^3$ labelled by $k = (k_1, k_2) \in \mathbb{Z}_{\times}^2 := \mathbb{Z}^2 \setminus (0, 0)$ and $n \in \mathbb{Z}_{\geq 0}$. We consider separately the eigenfunctions related to $k = (0, 0)$

$$\Psi_{0,m}(q) := e^{imq_0}, \quad m \in \mathbb{Z} \quad (20)$$

with the eigenvalues $\lambda_m = m^2$.

Theorem 1. *The eigenfunctions (19), (20) of the Laplace-Beltrami operator (10) form a complete basis in the Hilbert space $L_2(T_{\mathcal{L}}^3)$ of functions on Bianchi torus $T_{\mathcal{L}}^3$. For $B \neq 1$ the corresponding spectrum $\text{Spec} = \text{Spec}_0 \cup \text{Spec}_M$ consists of two parts:*

$$\text{Spec}_0 = \{\mathcal{E}_m = Am^2, \quad m \in \mathbb{Z}_{\geq 0}\}, \quad (21)$$

$$\text{Spec}_M = \{\mathcal{E}_{k,l} = A\lambda_l(\mu) + \frac{(B+1)Q(k)}{2}, \quad k \in \mathbb{Z}_{\times}^2, l \in \mathbb{Z}_{\geq 0}\}, \quad (22)$$

where λ_l are the eigenvalues of the Mathieu operator (17) with $\mu = \frac{(B-1)Q(k)}{2A}$, $Q(k)$ is the quadratic form of the lattice \mathcal{L} .

Proof. We need to show the completeness of the set of our eigenfunctions. For this it is enough to prove that any smooth function $\Phi : M^3 \rightarrow \mathbb{R}$ which is orthogonal to each of these eigenfunctions is zero.

Denoting again $q_0 = x$ expand the function $\Phi(x, q_1, q_2)$ as a Fourier series with respect to q_1, q_2 :

$$\Phi(x, q_1, q_2) = \sum_{s=(s_1, s_2) \in \mathbb{Z}^2} a_s(x) e^{i(s, \Omega q)}$$

with some smooth 2π -periodic coefficients $a_s(x)$, $x \in \mathbb{R}$. Orthogonality of $\Phi(x, q_1, q_2)$ to all eigenfunctions $\Psi_{k,l}(q)$ implies that for all non-zero $s \in \mathbb{Z}^2$ Fourier coefficient $a_s(x)$ is orthogonal to all Mathieu functions $\varphi_l(x, \mu)$ for all $l \in \mathbb{Z}_{\geq 0}$. Since the system of the Mathieu functions is complete in the Hilbert space of 2π -periodic functions this implies that the corresponding $a_s(x) \equiv 0$ and thus $\Phi = a_0(x)$. Now orthogonality to $\Psi_{0,l}(q)$ means that $a_0(x)$ must vanish and thus $\Phi \equiv 0$. \square

Corollary 1. *The spectrum of the Bianchi torus with metric (γ) behaves asymptotically as the values on 3D integer lattice of the quadratic form*

$$\tilde{Q}(k_0, k_1, k_2) = Ak_0^2 + \frac{(B+1)Q(k_1, k_2)}{2}, \quad (k_0, k_1, k_2) \in \mathbb{Z}^3 \quad (23)$$

with quadratic form $Q(k_1, k_2)$ given by (13).

Indeed, it is known that the Mathieu spectrum behaves asymptotically as m^2 , $m \in \mathbb{Z}$. When $B = 1$ we have the flat torus with the spectrum given by the values of the corresponding form $\tilde{Q} = Ak_0 + Q(k_1, k_2)$, so in that case the result is exact. The modification (23) of this form for general B is non-obvious and seems to be quite interesting.

Consider now the case of the quotient of Bianchi torus $M_2^3 = T_L^3/\mathbb{Z}_2$ by the involution $\tau : (q_0, q_1, q_2) \rightarrow (q_0 + \pi, -q_1, -q_2)$. Corresponding eigenfunctions can be found by the averaging:

$$\Psi_{k,l}^{(2)}(q) := e^{i(k, \Omega q)} \varphi_l(q_0, \mu) + e^{-i(k, \Omega q)} \varphi_l(q_0 + \pi, \mu) \quad (24)$$

More explicitly, for $\mu < 0$ (corresponding to $B < 1$) we have the functions

$$\begin{aligned} \Phi_{k,4j}^{(2)} &= \cos(k, \Omega q) ce_{2j}(x, \mu), \quad \Phi_{k,4j+1}^{(2)} = \sin(k, \Omega q) se_{2j+1}(x, \mu), \\ \Phi_{k,4j+2}^{(2)} &= \sin(k, \Omega q) ce_{2j+1}(x, \mu), \quad \Phi_{k,4j+3}^{(2)} = \cos(k, \Omega q) se_{2j+2}(x, \mu), \end{aligned}$$

where $k \in \mathbb{Z}_\times^2/\pm$ is considered up to a sign and $j \in \mathbb{Z}_{\geq 0}$.

The corresponding spectrum is given by the same formula (22), although the multiplicities for a generic lattice drop from 2 to 1 (since k and $-k$ are now identified). For non-generic lattices the multiplicities could be higher, see next section.

The trivial part is to be modified to $Spec_0^{(2)} = \{\mathcal{E}_m = 4Am^2, \quad m \in \mathbb{Z}_{\geq 0}\}$, with the multiplicities 2 (except ground state $\mathcal{E} = 0$ having multiplicity 1) and the eigenfunctions $e^{\pm 2imq_0}$.

The remaining cases correspond to the symmetric lattices and have nice arithmetic properties.

4. ARITHMETIC OF THE SYMMETRIC BIANCHI-VII₀ THREEFOLDS

Consider now the symmetric cases of Bianchi tori with the lattices having additional rotational symmetries of order $n > 2$. When $n = 4$ we have a square lattice, which can be assumed to be the standard square lattice $\mathcal{L} = 2\pi\mathbb{Z}^2 \subset \mathbb{R}^2(q_1, q_2)$ with quadratic form $Q(k) = k_1^2 + k_2^2$.

The same calculations as before lead to the eigenfunctions

$$\Psi_{k,l}(q) := e^{i(k_1q_1+k_2q_2)}\varphi_l(\mu, q_0) \quad (25)$$

be the corresponding eigenfunctions of the Laplace-Beltrami operator on M^3 labelled by $k = (k_1, k_2) \in \mathbb{Z}_{\times}^2 := \mathbb{Z}^2 \setminus (0, 0)$ and $l \in \mathbb{Z}_{\geq 0}$. Together with the elementary eigenfunctions $\Psi_{0,m}(q) := e^{imq_0}$, $m \in \mathbb{Z}$ with the eigenvalues $\lambda_m = Am^2$, they form a complete basis in the L_2 Hilbert space on the corresponding Bianchi torus. The nontrivial part of the spectrum (22) has the form

$$\mathcal{E}_{K,l} = A\lambda_l(\mu) + (B+1)K/2, \quad (26)$$

where $K = k_1^2 + k_2^2$, $l \in \mathbb{Z}_{\geq 0}$ and $\lambda_l(\mu)$ are the eigenvalues of Mathieu operator (17) with $\mu = \frac{(B-1)K}{2A}$. Asymptotically the spectrum behaves like the values of the quadratic form

$$\tilde{Q} = Ak_0^2 + (B+1)(k_1^2 + k_2^2)/2, \quad (k_0, k_1, k_2) \in \mathbb{Z}^3.$$

The difference with the general case is that now we have high multiplicities. Indeed, a given integer $K = k_1^2 + k_2^2$ may be represented as sum of the squares in many ways. This classical problem was studied by many mathematicians, including Fermat, Euler and Gauss. Jacobi found the following formula for the number $N(K)$ of such representations (see e.g. [12]):

$$N(K) = 4(N_1(K) - N_3(K)), \quad (27)$$

where $N_1(K)$ and $N_3(K)$ are the numbers of the divisors of K with the residues 1 and 3 modulo 4 respectively. Another classical result due to E. Landau [22] claims that the number of integers $K \leq N$ represented as the sum of two squares grows as $N/\sqrt{\log N}$.

Similar result holds for the Bianchi torus with the regular hexagonal lattice (covering M_n^3 with $n = 3$ and $n = 6$) when the non-trivial eigenvalues have the form (26) with $K = k_1^2 + k_1k_2 + k_2^2$, $(k_1, k_2) \in \mathbb{Z}^2$. In that case the number of representations $N(K)$ of a given integer K by this form is also well-known:

$$N(K) = 6(N_1(K) - N_2(K)), \quad (28)$$

where $N_1(K)$ and $N_2(K)$ are the numbers of the divisors of K with the residues 1 and 2 modulo 3 respectively (see [12]). The corresponding generalisation of Landau result was proved by Pall [26].

Theorem 2. *For generic parameters A and B in the metric the multiplicity of the eigenvalues (26) of the Laplace-Beltrami operator on the symmetric Bianchi-VII₀ threefolds M_n^3 with $n = 3, 4, 6$ are*

$$m(\mathcal{E}_{K,l}) = \frac{N(K)}{n},$$

where $N(K)$ are given by (27), (28) respectively.

Indeed, the eigenfunctions of the Laplace-Beltrami operator on the corresponding Bianchi threefolds $M_n^3 = T_{\mathcal{L}}^3/\mathbb{Z}_n$ can be found by averaging the eigenfunctions on the corresponding Bianchi torus $T_{\mathcal{L}}$ by the group generated by $\tau : (q_0, q_1, q_2) \rightarrow (q_0 + 2\pi/n, R_n(q_1, q_2))$ with the rotation R_n given by (6). The trivial part of the spectrum consists of the eigenvalues $\mathcal{E}_m = Am^2n^2$, $m \in \mathbb{Z}_{\geq 0}$ with the multiplicities 2 (for $m > 0$), corresponding to the eigenfunctions $e^{\pm 2imnq_0}$.

As a corollary of the number-theoretic results of Landau and Pall [22, 26] we have now the following result (cf. [5]).

Corollary 2. *The level spacing distribution for the spectrum of the symmetric Bianchi tori and threefolds M_n^3 , $n = 3, 4, 6$ is not Poisson and hence the Berry-Tabor conjecture does not hold in this case.*

Recall that the Berry-Tabor conjecture [3] predicts that the quantisation of completely integrable Hamiltonian systems should have Poisson distributed level spacing, which means their consecutive spacings should be independent and identically distributed. In our case the growing degeneracy means that most of level spacings are zero, which contradicts to Berry-Tabor conjecture.

5. COMPARISON WITH BIANCHI-VI AND BIANCHI-IX CASES

The classical *Sol*-case (corresponding to Bianchi-VI₀ type) was studied by Bolsinov and Taimanov in [6]. The principal examples here are the torus T^2 -fibrations M_A^3 over S^1 with hyperbolic gluing maps $A : T^2 \rightarrow T^2$, $A \in SL(2, \mathbb{Z})$. Bolsinov and Taimanov proved a surprising fact that the corresponding geodesic flow is Liouville integrable in smooth category, but not in analytic one. The obstruction to the analytic integrability [29] is purely topological and related to the exponential growth of the fundamental group $\pi_1(M_A^3)$.¹

In the quantum *Sol*-case considered in [5] the separation of variables leads to the cosh-version of Mathieu equation with parameter μ depending on the values of integer binary quadratic form Q , which is determined by the topology of the manifold (see [5]). As a result, we always have the high degeneracy of the spectrum. As we have seen in our Bianchi-VII case, the degeneracy of the eigenvalues disappears for the generic Bianchi tori M_1^3 .

¹It is interesting that these topological manifolds were considered already in 1892 by Poincaré, who was interested in the examples of manifolds with large fundamental groups.

Indeed, in that case we have the positive definite quadratic form $Q(k)$ with generic real coefficients. The study of values of quadratic forms at integer points is a classical topic in number theory, going back to Fermat.

In particular, Sarnak showed in [27] that the pair correlation of values does satisfy Berry-Tabor prediction on a set of full measure in the space of such forms. Eskin, Margulis and Mozes [14] provided stronger result by giving explicit diophantine conditions on the coefficients of the form $Q(k)$ for which this holds. In particular, they proved that if $\beta > 0$ is diophantine then the pair correlation of the values of $Q(k) = k_1^2 + \beta k_2^2$, $k_1, k_2 \in \mathbb{Z}$ agrees with the Berry-Tabor conjecture.

As we have shown, in the Bianchi-VII₀ case the spectrum asymptotically behaves as the values of the quadratic form (23) on 3D integer lattice, so it is natural to expect that the Berry-Tabor conjecture is true for the generic Bianchi tori, but this is still to be justified.

Like in the *Sol*-case [5], we can visualise the quantum monodromy for the symmetric Bianchi threefolds M_n^3 with $n = 3, 4, 6$ by plotting the values of the \mathbb{Z}_n -invariant integrals $F_1 = \sqrt{Q(p)} \cos n\varphi(p)$, $F_2 = \sqrt{Q(p)} \sin n\varphi(p)$ with $p \in \mathcal{L}^*$ and φ being its angular coordinate. In particular, for $n = 3$ we have the grid shown on Fig. 3 against the corresponding “Sol-flower” from [5] with the demonstration of the quantum monodromy in that case. In the *Sol*-case the quantum monodromy is given by the “cat-map”

$$A = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix},$$

while in our case it is simply rotation R_3 by $2\pi/3$.




FIGURE 4. Quantum monodromy in *Sol*-case and corresponding grid for Bianchi-VII threefold M_3^3 .

Finally, let us compare our case with the Bianchi-IX case. In that case we have Lie group $SO(3)$ (or, its double cover $SU(2)$) with a left-invariant metric and the geodesics describing the motion of the classical Euler top [1], which were found explicitly in terms of elliptic functions by Jacobi [20].

In quantum case the separation of variables in spherical elliptic coordinates [21, 17] leads to the classical Lamè equation

$$L = -\frac{d^2}{dx^2} + m(m+1)k^2 sn^2(x, k), \quad m \in \mathbb{Z}_{\geq 0},$$

where $sn(x, k)$ is the classical Jacobi elliptic function [31]. Note that the set of parameters $m(m+1), m \in \mathbb{Z}_{\geq 0}$ is the spectrum of the unit sphere S^2 , while in our Bianchi-VII₀ case we have the spectrum of a flat 2D torus. The corresponding Lamè operators are known to have remarkable spectral property: in their spectrum there are only finitely many (namely, exactly m) gaps [19]. The ends of the gaps can be computed using classical methods going back to Hermite and Halphen [31] (see also [17] and references therein), but the formulae become very complicated when m grows. It is interesting to note that if $m \rightarrow \infty, k \rightarrow 0$ such that $mk \rightarrow \nu$, then

$$m(m+1)k^2 sn^2(x, k) \rightarrow \nu^2 \sin^2 x,$$

and we have as a limit a (shifted) Mathieu operator from Bianchi-VII₀ case.

In the most symmetric Bianchi-IX case we have the unit sphere S^3 with the spectrum $m(m+2), m \in \mathbb{Z}_{\geq 0}$ with the multiplicity $(m+1)^2$, while in Bianchi-VII₀ case we have flat torus T^3 with spectrum given by the values of a quadratic form on the corresponding 3D lattice. In Thurston's list these two cases are represented by S^3 and E^3 respectively.

Note that in Bianchi-IX case we have also the quotients $SU(2)/G$, where G is either cyclic group \mathbb{Z}_n , or the double cover of one of the finite subgroups of $SO(3)$: dihedral D_n , tetrahedral T , octahedral O and icosahedral I (in the last case the corresponding quotient is famous Poincaré homology sphere). The spectrum of the corresponding symmetric versions can be found in [18].

6. ACKNOWLEDGEMENTS

We are very grateful to Alexey Bolsinov for many helpful discussions.

The work of the second author (Y.Ye) was supported by the XJTU Research Development Funding (grant number: RDF-21-01-031).

REFERENCES

- [1] V.I. Arnold *Mathematical Methods of Classical Mechanics*. Springer, 1984.
- [2] L. Bianchi *Sugli spazii a tre dimensioni che ammettono un gruppo continuo di movimenti*. Soc. Ital. Sci. Mem. di Mat. **11** (1898), 267.
- [3] M.V. Berry and M. Tabor *Level clustering in the regular spectrum*. Proc. Roy. Soc. London A **356** (1977), 375-394.
- [4] A. Bolsinov, J. Bao *Note about integrable systems on low-dimensional Lie groups and Lie algebras*. Regular and Chaotic Dynamics **24** (2019), 266-280.
- [5] A.V. Bolsinov, H.R. Dullin, A.P. Veselov *Spectra of Sol-manifolds: arithmetic and quantum monodromy*. Comm. Math. Phys. **264** (2006), 583-611.
- [6] A.V. Bolsinov, I.A. Taimanov *Integrable geodesic flows with positive topological entropy*. Invent. Math. **140** (2000), 639-650.
- [7] A.V. Bolsinov, A.P. Veselov, Y. Ye *Chaos and integrability in $SL(2, \mathbb{R})$ -geometry*. Russian Math. Surveys **76(4)** (2021), 557-586.

- [8] C. Brimacombe, R.M. Corless, M. Zamir *Computation and applications of Mathieu functions: A historical perspective*. SIAM Review **63:4** (2021), 653–720.
- [9] L. Butler *A new class of homogeneous manifolds with Liouville-integrable geodesic flows*. C. R. Math. Acad. Sci. Soc. R. Can. **21**(1999), no. 4, 127-131.
- [10] H.S.M. Coxeter *Introduction to Geometry* (2nd ed.), Wiley, 1989.
- [11] M.P. do Carmo *Differential Geometry of Curves and Surfaces*. Prentice-Hall, 1976.
- [12] L.E. Dickson *Introduction to the Theory of Numbers*. Univ. of Chicago Press, 1929.
- [13] A.Erdelyi (Editor) *Higher Transcendental Functions.*, Vol.1-3. McGraw-Hill, 1953.
- [14] A. Eskin, G. Margulis and S. Mozes *Quadratic forms of signature (2,2) and eigenvalue spacings on rectangular 2-tori*. Annals of Mathematics, **161** (2005), 679–725.
- [15] H.V. Fagundes *Relativistic cosmologies with closed, locally homogeneous spatial sections*. Phys. Rev. Lett. **54** (1985), 1200-1202.
- [16] Y. Fujiwara, H. Ishihara, H. Kodama *Comments on closed Bianchi models*. Class. Quantum Grav. **10** (1993), 859–867.
- [17] M.-P. Grosset, A.P. Veselov *Lame equation, quantum Euler top and elliptic Bernoulli polynomials*. Proc. Edinb. Math. Soc. **51** (2008), 635-650.
- [18] A. Ikeda *On the spectrum of homogeneous spherical space forms*. Kodai Math. J. **18** (1995), 57-67.
- [19] E.L. Ince *Further investigations into the periodic Lamé functions*. Proc. Roy. Soc. Edinburgh 60, (1940). 83–99.
- [20] C.G.J. Jacobi *Sur la rotation d'un corps*. In: Gesammelte Werke, 2.Band, 291-352, 1881.
- [21] H.A. Kramers and G.P. Ittmann *Zur Quantelung des asymmetrischen Kreisels. I, II*. Z. Physik 53, 553-565, 58, 217-231 (1929), 217-231.
- [22] E. Landau *Über die Einteilung der positive ganzen Zahlen in vier Klassen*. Archiv der Mathematik und Physik (3), **13** (1908), 305-312.
- [23] W.J. LeVeque *Fundamentals of Number Theory*. Dover Publ. inc., 1996.
- [24] E. Mathieu *Mémoire sur le mouvement vibratoire d'une membrane de forme elliptique*. J. Math. Pure Appl. **13** (1868), 137–203.
- [25] J. Milnor *Curvatures of left invariant metrics on Lie groups*. Adv. Math. **21:3** (1976), 293-329.
- [26] G. Pall *The distribution of integers represented by binary quadratic forms*. Bull. Amer. Math. Soc. **49(6)** (1943), 447-449.
- [27] P. Sarnak *Values at integers of binary quadratic forms*. CMS Conf. Proc., 21, Amer. Math. Soc., Providence, RI, (1997), 181-203.
- [28] P. Scott *The geometries of 3-manifolds*. Bull. London Math. Soc., **15** (1983), 401-487.
- [29] I.A. Taimanov *Topological obstructions to integrability of geodesic flows on non-simply-connected manifolds*. Math. USSR Izv. **30** (1988), 403-409.
- [30] W.P. Thurston *Hyperbolic geometry and 3-manifolds*. LMS Lecture Notes Series **48**, Camb. Univ. Press, 1982.
- [31] E.T. Whittaker, G.N. Watson *Course in Modern Analysis*, Camb. Univ. Press (1990).

DEPARTMENT OF MATHEMATICAL SCIENCES, LOUGHBOROUGH UNIVERSITY, LOUGHBOROUGH LE11 3TU

Email address: A.P.Veselov@lboro.ac.uk

DEPARTMENT OF FOUNDATIONAL MATHEMATICS, XI'AN JIAOTONG-LIVERPOOL UNIVERSITY, SUZHOU, 215000, CHINA

Email address: yiru.ye@xjtlu.edu.cn