

The Bel-Robinson tensor and the classical pseudotensors

Lau Loi So

Abstract

Describing the gravitational energy and momentum, the Bel-Robinson tensor is the best tensor. However, the classical pseudotensors can also manage the the same job. As Deser mentioned in 1999, a certain linear combination between Einstein pseudotensor and Landau-Lifshitz pseudoetnsor give a pure Bel-Robinson tensor. Here we used the same idea but adapted the harmonic gauge, we found that all the classical pseudotensor alone cannot give a multiple of the Bel-Robinson tensor. But under a modification, all of them contribute the same energy-momentum and stress as the Bel-Robinson tensor does.

1 Introduction

The Bel-Robinson tensor is the best tensor to describe the energy-momentum and stress in general relativity. Senovilla had written a very nice article about this tensor [1]. This tensor has many very nice mathematical properties: completely symmetric, trace free and divergence free. In 1999, Deser [2] explained how to make a certain linear combination between Einstein [3] pseudotensor and Landau-Lifshitz [4] pseudotensor to produce the pure Bel-Robinson tensor, under the Riemann normal coordinates systems. In fact, we had examined that all the classical pseudotensor alone cannot generate a multiple of the Bel-Robinson tensor [5]. Our present paper used the same idea, but using another angle, apply the harmonic gauge and using the physical detail of the metric. As expected, we found none of the classical pseudotensor can have a multiple of the pure Bel-Robinson tensor. But after adding some modification, the results indicate that each classical pseudotensor does have the pure Bel-Robinson tensor, restricted in the energy-momentum and stress. When we say the classical pseudotensors which means Einstein, Landau-Lifshitz, Bergmann-Thomson [6], Papapetrou [7] and Weinberg [8].

2 Technical background

Throughout this work, we use the same spacetime signature and notation in MTW [9], including the geometrical units $G = c = 1$, where G and c are the Newtonian constant and the speed of light. The Greek letters denote the spacetime and Latin letters refer to spatial. In vacuum, the Bel-Robinson tensor can be defined as follows

$$B_{\alpha\beta\mu\nu} := R_{\alpha\lambda\mu\sigma}R_{\beta}{}^{\lambda}{}_{\nu}{}^{\sigma} + R_{\alpha\lambda\nu\sigma}R_{\beta}{}^{\lambda}{}_{\mu}{}^{\sigma} - \frac{1}{8}g_{\alpha\beta}g_{\mu\nu}R_{\lambda\sigma\rho\tau}^2, \quad (1)$$

where $R_{\lambda\sigma\rho\tau}^2 = R_{\lambda\sigma\rho\tau}R^{\lambda\sigma\rho\tau}$. In order to extract the energy-momentum and the stress, one can use the analog of the “electric” E_{ab} and “magnetic” B_{ab} parts of the Weyl tensor [10],

$$E_{ab} := C_{0a0b}, \quad B_{ab} := *C_{0a0b}, \quad (2)$$

where $C_{\alpha\beta\mu\nu}$ is the Weyl conformal tensor and $*C_{\alpha\beta\mu\nu}$ is its dual,

$$*C_{\alpha\beta\mu\nu} = \frac{1}{2}\epsilon_{\alpha\beta\lambda\sigma}C^{\lambda\sigma}{}_{\mu\nu}. \quad (3)$$

(Here $\epsilon_{\alpha\beta\mu\nu} = \epsilon_{[\alpha\beta\mu\nu]}$ with $\epsilon_{0123} = \sqrt{-g}$ is the totally anti-symmetric Levi-Civita tensor, see [9], in particular Eq. 8.10 and Ex. 8.3.) In vacuum using the Riemann tensor

$$E_{ab} = R_{0a0b}, \quad B_{ab} = *R_{0a0b}. \quad (4)$$

Certain commonly occurring quadratic combinations of the Riemann tensor components in terms of the electric E_{ab} and magnetic B_{ab} parts in vacuum are

$$R_{0a0b}R_0{}^a{}_0{}^b = E_{ab}E^{ab}, \quad R_{0abc}R_0{}^{abc} = 2B_{ab}B^{ab}, \quad R_{abcd}R^{abcd} = 4E_{ab}E^{ab}. \quad (5)$$

In particular, the Riemann squared tensor can then be written

$$R_{\lambda\sigma\rho\tau}^2 = 8(E_{ab}E^{ab} - B_{ab}B^{ab}). \quad (6)$$

The energy-momentum and stress for the Bel-Robinson tensor are

$$B_{0000} = E_{ab}^2 + B_{ab}^2, \quad (7)$$

$$B_{000j} = 2\epsilon_{jab}E^{ac}B^b{}_c, \quad (8)$$

$$B_{00ij} = \eta_{ij}(E_{ab}^2 + B_{ab}^2) - 2(E^c{}_iE_{cj} + B^c{}_iB_{cj}), \quad (9)$$

where $E_{ab}^2 := E_{ab}E^{ab}$ and similarly for B_{ab}^2 .

Within a weak field the metric tensor can be decomposed as $g_{\alpha\beta} = \eta_{\alpha\beta} + h_{\alpha\beta}$, and its inverse is $g^{\alpha\beta} = \eta^{\alpha\beta} - h^{\alpha\beta}$. Here we only consider the lowest order (see (8) at [11]), the metric components can be written as

$$h^{00} = -E_{ab}x^ax^b, \quad h^{0j} = \frac{2}{3}\epsilon^j{}_{pq}B^p{}_l x^q x^l, \quad h^{ij} = \eta^{ij}h^{00}. \quad (10)$$

The lowest order for the harmonic gauge is

$$\Gamma^{\alpha\beta}{}_{\beta} = \partial_{\beta}\bar{h}^{\alpha\beta} = 0, \quad (11)$$

where $\bar{h}^{\alpha\beta} = h^{\alpha\beta} - \frac{1}{2}\eta^{\alpha\beta}h$. Thus, the energy-momentum and the stress for the Bel-Robinson tensor can be written as the following form

$$B^{0000} = \frac{1}{4}\vec{\nabla}^2 \left[\frac{\alpha}{2} - \frac{A}{2} + \frac{B}{2} \right], \quad (12)$$

$$B^{000j} = \frac{1}{4}\vec{\nabla}^2(C - D), \quad (13)$$

$$B^{00ij} = \frac{1}{4}\vec{\nabla}^2 \left[-X - Y + Z - U - V + \eta^{ij} \left(\frac{\alpha}{2} + \frac{A}{2} - \frac{B}{2} \right) \right], \quad (14)$$

where

$$\alpha = (\partial_c h_{00})(\partial^c h_{00}), \quad A = (\partial_c h_{0d})(\partial^c h^{0d}), \quad B = (\partial_c h_{0d})(\partial^d h^{0c}), \quad (15)$$

$$C = (\partial_c h_{00})(\partial^c h^{0j}), \quad D = (\partial_c h_{00})(\partial^j h^{0c}), \quad (16)$$

$$X = (\partial^k h_{00})(\partial^f h_{00}), \quad Y = (\partial^k h^{0c})(\partial^f h_{0c}), \quad Z = (\partial_c h^{0k})(\partial^c h^{0f}), \quad (17)$$

$$U = (\partial^k h^{0c})(\partial_c h^{0f}), \quad V = (\partial_c h^{0k})(\partial^f h^{0c}). \quad (18)$$

On the other hand, generally speaking, regarding the classical pseudotensor [5] which can be obtained from a rearrangement of the Einstein field equation: $G^{\mu\nu} = \kappa T^{\mu\nu}$, where the constant $\kappa = 8\pi G/c^4$, $G^{\mu\nu}$ and $T^{\mu\nu}$ are the Einstein and stress tensors. We define the gravitational energy-momentum density pseudotensor in terms of a suitable superpotential $U_\alpha^{[\mu\nu]}$:

$$2\kappa\sqrt{-g}t_\alpha^\mu := \partial_\nu U_\alpha^{[\mu\nu]} - 2\sqrt{-g}G_\alpha^\mu. \quad (19)$$

Alternatively, one can rewrite (19) as $\partial_\nu U_\alpha^{[\mu\nu]} = \sqrt{-g}(2G_\alpha^\mu + 2\kappa t_\alpha^\mu)$. Using the Einstein equation, the total energy-momentum density complex can be defined as

$$\mathcal{T}_\alpha^\mu := \sqrt{-g}(T_\alpha^\mu + t_\alpha^\mu) = (2\kappa)^{-1}\partial_\nu U_\alpha^{[\mu\nu]}. \quad (20)$$

This total energy-momentum density is automatically conserved as $\partial_\mu \mathcal{T}_\alpha^\mu \equiv 0$, which can be split into two parts:

$$(\partial_\nu U_\alpha^{[\mu\nu]})_1 = 2\sqrt{-g}G_\alpha^\mu, \quad (\partial_\nu U_\alpha^{[\mu\nu]})_2 = 2\sqrt{-g}\kappa t_\alpha^\mu. \quad (21)$$

The first part indicates the contribution of matter and the second piece refers to vacuum gravity [9]. For the equivalence principle small limit at a point within matter, it is known that the Einstein, Landau-Lifshitz, Bergmann-Thomson, Papapetrou and Weinberg pseudotensors all satisfy the standard result $2G_\alpha^\mu$ at the origin in Riemann normal coordinates, while t_α^μ vanishes in this limit [5].

Actually there are three kinds of superpotentials: $U_\alpha^{[\mu\nu]}$, $U^{\alpha[\mu\nu]}$ and $H^{[\alpha\beta][\mu\nu]}$. The latter also must satisfy some further symmetries, under the interchange of the pairs, and the total antisymmetry of 3 indices (i.e., the same as the symmetries of the Riemann tensor). From them the total energy-momentum complex can be obtained as follows:

$$2\kappa\mathcal{T}_\alpha^\mu = \partial_\nu U_\alpha^{[\mu\nu]}, \quad 2\kappa\mathcal{T}^{\alpha\mu} = \partial_\nu U^{\alpha[\mu\nu]}, \quad 2\kappa\mathcal{T}^{\alpha\mu} = \partial_{\beta\nu}^2 H^{[\alpha\beta][\mu\nu]}. \quad (22)$$

In vacuum, t^{00} is the gravitational energy density, t^{0j} is the momentum and t^{ij} is the stress. In short: we use the following for the representation

$$t^{\alpha\mu} = (t^{00}, t^{0j}, t^{ij}) \quad (23)$$

3 The classical pseudotensors

3.1 Einstein pseudotensor

The following is the Freud (F) superpotential [3]:

$${}_F U_\alpha^{[\mu\nu]} := -\sqrt{-g}g^{\beta\sigma}\Gamma^\tau_{\lambda\beta}\delta_{\tau\sigma\alpha}^{\lambda\mu\nu}. \quad (24)$$

The corresponding Einstein pseudotensor is

$$\begin{aligned} t_E^{\alpha\mu} &= -2\Gamma^{\beta\lambda\alpha}\Gamma^\mu_{\beta\lambda} + \Gamma^{\mu\alpha\lambda}\Gamma^\beta_{\beta\lambda} + \Gamma^{\beta\alpha}_{\beta}\Gamma^{\mu\lambda}_{\lambda} \\ &\quad + \eta^{\alpha\mu}(\Gamma^{\beta\lambda}_{\nu}\Gamma^\nu_{\beta\lambda} - \Gamma^\beta_{\beta\nu}\Gamma^{\nu\lambda}_{\lambda}) + \Gamma^{\nu\alpha\mu}\Gamma^\beta_{\beta\nu} - \Gamma^{\beta\alpha}_{\beta}\Gamma^{\lambda\mu}_{\lambda}. \end{aligned} \quad (25)$$

Apply the harmonic gauge, the energy-momentum and the stress are

$$t_E^{\alpha\mu} = - \left[\left(-\frac{\alpha}{2} - \frac{A}{2} + \frac{B}{2} \right), 0, \eta^{kf} \left(\frac{\alpha}{2} + \frac{A}{2} - \frac{B}{2} \right) - X - Y - U \right]. \quad (26)$$

According to Table 1, we add the extra piece

$$\begin{aligned} t_{\text{New}}^{\alpha\mu} &= t_E^{\alpha\mu} + 2t_1 + 2t_2 - \frac{t_3}{2} + \frac{t_5}{2} - t_9 - \frac{t_{11}}{2} \\ &= - \left[\frac{\alpha}{2} - \frac{A}{2} + \frac{B}{2}, C - D, \eta^{kf} \left(\frac{\alpha}{2} + \frac{A}{2} - \frac{B}{2} \right) - X - Y + Z - U - V \right] \end{aligned} \quad (27)$$

Then

$$-\frac{1}{4} \vec{\nabla}^2 t_{\text{New}}^{\alpha\mu} = (B^{0000}, B^{000j}, B^{00ij}). \quad (28)$$

3.2 Landau-Lifshitz pseudotensor

The Landau-Lifshitz (LL) superpotential [4]

$$U_{LL}^{\alpha[\mu\nu]} := -(-g)g^{\alpha\beta}g^{\pi\sigma}\Gamma^\tau_{\lambda\pi}\delta_{\tau\sigma\beta}^{\lambda\mu\nu}, \quad (29)$$

and the pseudotensor becomes

$$\begin{aligned} t_{LL}^{\alpha\mu} &= g^{\alpha\mu}(\Gamma^\gamma_{\gamma\nu}\Gamma^{\beta\nu}_{\beta} + \Gamma^{\nu\beta}_{\rho}\Gamma^{\rho}_{\nu\beta} - 2\Gamma^\gamma_{\gamma\nu}\Gamma^{\nu\beta}_{\beta}) + 2\Gamma^{\nu\alpha\mu}\Gamma^{\beta}_{\beta\nu} - \Gamma^{\rho\alpha}_{\rho}\Gamma^{\beta\mu}_{\beta} \\ &\quad + \Gamma^{\alpha\rho}_{\beta}\Gamma^{\mu\beta}_{\rho} - \Gamma^{\rho\alpha}_{\beta}\Gamma^{\beta\mu}_{\rho} - \Gamma^{\alpha\rho}_{\beta}\Gamma^{\beta\mu}_{\rho} - \Gamma^{\rho\alpha}_{\beta}\Gamma^{\mu\beta}_{\rho} - (\Gamma^{\alpha\mu}_{\nu} + \Gamma^{\mu\alpha}_{\nu})\Gamma^{\beta\nu}_{\beta} \\ &\quad + \Gamma^{\alpha\rho}_{\rho}\Gamma^{\beta\mu}_{\beta} + \Gamma^{\rho\alpha}_{\rho}\Gamma^{\mu\beta}_{\beta} - \Gamma^{\alpha\rho}_{\rho}\Gamma^{\mu\beta}_{\beta} + (\Gamma^{\alpha\mu}_{\rho} + \Gamma^{\mu\alpha}_{\rho})\Gamma^{\rho\beta}_{\beta}, \end{aligned} \quad (30)$$

Apply the harmonic gauge, we have the following results

$$t_{LL}^{\alpha\mu} = - \left[\frac{7\alpha}{2} + \frac{A}{2} + \frac{B}{2}, 2C - 2D, \eta^{kf} \left(\frac{\alpha}{2} + \frac{A}{2} - \frac{B}{2} \right) - X - Y + Z - U - V \right]. \quad (31)$$

This result is not good, then consider the modification

$$\begin{aligned} t_{\text{New}}^{\alpha\mu} &= t_{LL}^{\alpha\beta} + 2t_1 + 2t_2 - t_3 + \frac{3}{2}t_5 - t_{11} \\ &= - \left[\frac{\alpha}{2} - \frac{A}{2} + \frac{B}{2}, C - D, \eta^{kf} \left(\frac{\alpha}{2} + \frac{A}{2} - \frac{B}{2} \right) - X - Y + Z - U - V \right] \end{aligned} \quad (32)$$

Then we have the expected result

$$-\frac{1}{4} \vec{\nabla}^2 t_{\text{New}}^{\alpha\mu} = (B^{0000}, B^{000j}, B^{00ij}). \quad (33)$$

3.3 Bergmann-Thomson pseudotensor

The Bergmann-Thomson (BT) superpotential [6] is defined as

$$U_{BT}^{\alpha[\mu\nu]} := -\sqrt{-g}g^{\alpha\beta}g^{\pi\sigma}\Gamma^\tau_{\lambda\pi}\delta_{\tau\sigma\beta}^{\lambda\mu\nu}, \quad (34)$$

and the associated pseudotensor is

$$\begin{aligned} t_{BT}^{\alpha\mu} &= g^{\alpha\mu}\Gamma^{\beta\lambda}_{\rho}\Gamma^{\rho}_{\beta\lambda} - (g^{\alpha\mu}\Gamma^{\beta}_{\beta\rho} - \Gamma^{\alpha\mu}_{\rho} - \Gamma^{\mu\alpha}_{\rho})\Gamma^{\rho\nu}_{\nu} + \Gamma^{\alpha\nu}_{\nu}(\Gamma^{\beta\mu}_{\beta} - \Gamma^{\mu\beta}_{\beta}) \\ &\quad + (\Gamma^{\rho\alpha\mu} - \Gamma^{\alpha\mu\rho})\Gamma^{\beta}_{\beta\rho} + \Gamma^{\alpha\rho}_{\beta}(\Gamma^{\mu\beta}_{\rho} - \Gamma^{\beta\mu}_{\rho}) - \Gamma^{\rho\alpha}_{\beta}(\Gamma^{\mu\beta}_{\rho} + \Gamma^{\beta\mu}_{\rho}). \end{aligned} \quad (35)$$

Apply the harmonic gauge, the results are

$$t_{\text{New}}^{\alpha\mu} = - \left[\frac{3\alpha}{2} + \frac{A}{2} + \frac{B}{2}, C - D, \eta^{kf} \left(\frac{\alpha}{2} + \frac{A}{2} - \frac{B}{2} \right) - X - Y + Z - U - V \right]. \quad (36)$$

This result requires some modification

$$\begin{aligned} t_{\text{New}}^{\alpha\mu} &= t_{BT}^{\alpha\mu} + 2t_1 + 2t_2 - \frac{t_3}{2} + \frac{3}{2}t_5 - \frac{t_{11}}{2} \\ &= - \left[\frac{\alpha}{2} - \frac{A}{2} + \frac{B}{2}, C - D, \eta^{kf} \left(\frac{\alpha}{2} + \frac{A}{2} - \frac{B}{2} \right) - X - Y + Z - U - V \right] \end{aligned} \quad (37)$$

Then we have the expected result

$$-\frac{1}{4}\vec{\nabla}^2 t_{\text{New}}^{\alpha\mu} = (B^{0000}, B^{000j}, B^{00ij}). \quad (38)$$

3.4 Papapetrou pseudotensor

The Papapetrou superpotential [7] is defined as

$$H_P^{[\mu\nu][\alpha\beta]} := -\sqrt{-g}g^{\rho\pi}\eta^{\tau\gamma}\delta_{\pi\gamma}^{\nu\mu}\delta_{\rho\tau}^{\alpha\beta}, \quad (39)$$

equivalently one can use $U_P^{\alpha[\mu\nu]} \equiv \partial_\beta H^{[\mu\nu][\alpha\beta]}$. The corresponding pseudotensor is

$$\begin{aligned} t_P^{\alpha\mu} &= \eta^{\alpha\mu}(\Gamma^{\beta\lambda}{}_\rho\Gamma^\rho{}_{\beta\lambda} + \Gamma^{\nu\beta}{}_\nu\Gamma^\lambda{}_{\lambda\beta} - \Gamma^\beta{}_{\beta\rho}\Gamma^{\rho\lambda}{}_\lambda + h^{\beta\nu}\Gamma^\lambda{}_{\lambda\beta,\nu}) \\ &\quad + 2\Gamma^{\alpha}{}_{\beta\lambda}\Gamma^{\mu\beta\lambda} + \Gamma^{\alpha\beta}{}_\beta\Gamma^{\lambda\mu}{}_\lambda + \Gamma^{\beta\alpha}{}_\beta\Gamma^{\mu\lambda}{}_\lambda - (\Gamma^{\alpha\mu}{}_\beta + \Gamma^{\mu\alpha}{}_\beta)(2\Gamma^{\lambda\beta}{}_\lambda + \Gamma^{\beta\lambda}{}_\lambda) \\ &\quad + h^{\alpha\beta}\partial_\beta\Gamma^{\mu\lambda}{}_\lambda + h^{\beta\mu}\partial_\beta\Gamma^{\alpha\lambda}{}_\lambda - h^{\alpha\mu}\partial_\beta\Gamma^{\beta\lambda}{}_\lambda - h^{\beta\lambda}(\Gamma^{\alpha\mu}{}_{\beta,\lambda} + \Gamma^{\mu\alpha}{}_{\beta,\lambda}). \end{aligned} \quad (40)$$

Apply the harmonic gauge, we have the following results

$$t_P^{\alpha\mu} = - \left[\frac{7\alpha}{2} + \frac{A}{2} + \frac{3B}{2}, 2C - 2D, \eta^{kf} \left(\frac{\alpha}{2} + \frac{A}{2} - \frac{B}{2} \right) - X - Y + Z - U - V \right]. \quad (41)$$

Adding some extra pieces

$$\begin{aligned} t_{\text{New}}^{\alpha\mu} &= t_P^{\alpha\mu} + t_1 + t_2 - \frac{3}{2}t_3 + \frac{3}{2}t_5 - \frac{t_{11}}{2} \\ &= - \left[\frac{\alpha}{2} - \frac{A}{2} + \frac{B}{2}, C - D, \eta^{kf} \left(\frac{\alpha}{2} + \frac{A}{2} - \frac{B}{2} \right) - X - Y + Z - U - V \right] \end{aligned} \quad (42)$$

Then we have desired result

$$-\frac{1}{4}\vec{\nabla}^2 t_{\text{New}}^{\alpha\mu} = (B^{0000}, B^{000j}, B^{00ij}). \quad (43)$$

3.5 Weinberg pseudotensor

The superpotential for Weinberg(W) [8] is

$$H_W^{[\mu\nu][\alpha\beta]} := \sqrt{-\eta}\eta^{\alpha\pi}\eta^{\beta\xi}\eta^{\lambda\kappa}\delta_{\pi\xi\kappa}^{\sigma\mu\nu}g_{\lambda\sigma}. \quad (44)$$

The associated Weinberg pseudotensor becomes

$$t_W^{\alpha\mu} = (2\Gamma^{\beta\alpha\mu} - g^{\alpha\mu}\Gamma^{\beta\pi}{}_\pi)\Gamma_{\beta}{}^\nu{}_\nu + g^{\alpha\mu}\Gamma^{\rho\beta\nu}\Gamma_{\rho\beta\nu} - 2\Gamma^{\beta\nu\alpha}\Gamma_{\beta\nu}{}^\mu - 2h^{\beta\lambda}(\partial_\lambda\Gamma^{\alpha\mu}{}_\beta - \partial^\mu\Gamma^\alpha{}_{\beta\lambda}). \quad (45)$$

Apply the harmonic gauge, we have the following results

$$t_W^{\alpha\mu} = - \left[\left(\frac{3\alpha}{2} + \frac{A}{2} + \frac{B}{2} \right), D, \eta^{kf} \left(-\frac{3\alpha}{2} - \frac{3A}{2} + \frac{B}{2} \right) + X + Y - Z \right]. \quad (46)$$

Adding some modification

$$\begin{aligned} t_{\text{New}}^{\alpha\mu} &= t_W^{\alpha\mu} + 2t_1 + t_2 - \frac{t_3}{2} + \frac{5}{2}t_5 - t_6 - t_8 - t_9 - t_{10} + t_{11} \\ &= - \left[\frac{\alpha}{2} - \frac{A}{2} + \frac{B}{2}, C - D, \eta^{kf} \left(\frac{\alpha}{2} + \frac{A}{2} - \frac{B}{2} \right) - X - Y + Z - U - V \right] \end{aligned} \quad (47)$$

Then we have desired result

$$-\frac{1}{4}\vec{\nabla}^2 t_{\text{New}}^{\alpha\mu} = (B^{0000}, B^{000j}, B^{00ij}). \quad (48)$$

Table 1: Extra superpotential for U_i .

Superpotential		
$U_1 = h^{\mu\pi}\Gamma^{\alpha\nu}_{\pi} - (\mu \leftrightarrow \nu),$	$U_2 = h^{\mu\pi}\Gamma^{\nu\alpha}_{\pi} - (\mu \leftrightarrow \nu),$	$U_3 = g^{\alpha\mu}h^{\beta\nu}\Gamma^{\lambda}_{\lambda\beta} - (\mu \leftrightarrow \nu),$
$U_4 = h^{\alpha\nu}\Gamma^{\mu\lambda}_{\lambda} - (\mu \leftrightarrow \nu),$	$U_5 = h^{\alpha\mu}\Gamma^{\lambda\nu}_{\lambda} - (\mu \leftrightarrow \nu),$	$U_6 = g^{\alpha\mu}h^{\beta\lambda}\Gamma_{\beta\lambda}^{\nu} - (\mu \leftrightarrow \nu),$
$U_7 = g^{\alpha\nu}h^{\beta\mu}\Gamma_{\beta}^{\lambda\alpha} - (\mu \leftrightarrow \nu),$	$U_8 = g^{\alpha\nu}h^{\beta\lambda}\Gamma_{\beta\lambda}^{\mu} - (\mu \leftrightarrow \nu),$	$U_9 = h^{\alpha\lambda}\Gamma^{\nu\mu}_{\lambda} - (\mu \leftrightarrow \nu),$
$U_{10} = h^{\nu\pi}\Gamma_{\pi}^{\alpha\mu} - (\mu \leftrightarrow \nu),$	$U_{11} = h\Gamma^{\nu\mu\alpha} - (\mu \leftrightarrow \nu),$	$U_{12} = g^{\alpha\mu}h\Gamma^{\lambda\nu}_{\lambda} - (\mu \leftrightarrow \nu),$
$U_{13} = g^{\alpha\mu}h\Gamma^{\nu\lambda}_{\lambda} - (\mu \leftrightarrow \nu).$		

4 Conclusion

In 1999, Deser proposed a certain linear combination between Einstein and Landau Lifshitz pseudotensors to generate a multiple of the pure Bel-Robinson tensor using the Riemann normal coordinates. Here we use a similar idea but just try to obtain the energy-momentum and the stress, under a harmonic gauge condition. We found that none of the classical pseudotensor can have the expected result by itself. But after adding some modification, all of them have the desired results.

References

- [1] Senovilla J M M 2000 *Class. Quantum Grav.* **17** 2799
- [2] Deser S, Franklin J S and Seminaea D 1999 *Class. Quantum Grav.* **16** 2815
- [3] Freud Ph 1939 *Ann. Math.* **40** 417
- [4] Landau L D and Lifshitz E M 1975 *The classical theory of fields* (Oxford: Pergamon)
- [5] So L L, Nester J M and Chen H 2009 *Class. Quantum Grav.* **26** 085004
- [6] Bergmann P G and Thomson R 1953 *Phys. Rev.* **89** 400

- [7] Papapetrou A 1948 *Proc. R. Irish. Acad.* **A52** 11
- [8] Weinberg S 1972 *Gravitation and Cosmology* (New York: Wiley) p371
- [9] Misner C W, Thorne K S and Wheeler J A 1973 *Gravitation* (San Francisco, CA: Freeman)
- [10] Carmeli M “*Classical Fields General relativity and Gauge Theory*” (John Wiley & Sons 1982)
- [11] Zhang X H 1985 *Phys. Rev. D* **31** 3130