# Interaction phenomenon for variable coefficient Kadomtsev-Petviashvili equation by utilizing variable coefficient bilinear neural network method

Jian-Guo Liu<sup>1\*</sup>, Wen-Hui Zhu<sup>2\*</sup>

<sup>1</sup> College of Computer, Jiangxi University of Chinese Medicine, Jiangxi 330004, China;

<sup>2</sup> Institute of artificial intelligence, Nanchang Institute of Science and Technology, Jiangxi 330108, China;

Abstract: In this paper, a variable coefficient Bilinear neural network method is proposed to deal with the analytical solutions of variable coefficient nonlinear partial differential equations. As an example, a Kadomstev-Petviashvili equation with variable coefficients is investigated by using the variable coefficient Bilinear neural network method. By establishing "3-2-2-1" and "3-3-2-1" models respectively, rich analytical solutions of the variable coefficient Kadomstev-Petviashvili equation are obtained. By choosing some special values of the parameters, the dynamics properties are demonstrated in some three-dimensional and density graphics.

**Keywords:** variable coefficient Bilinear neural network method, variable coefficient Kadomstev-Petviashvili equation, analytical solutions, dynamic properties.

2010 Mathematics Subject Classification: 35Q51, 35G99, 33F10

### 1. Introduction

1

Partial differential equations (PDEs) have a wide range of applications, from the simulation of seismic waves on Earth to the spread of infectious diseases through human populations. Engineers, scientists, and mathematicians have resorted to PDEs to describe complex phenomena involving many independent variables [1,2]. However, the process of solving PDEs is extremely difficult, traditional numerical methods are very complex and require a lot of computation. Researchers have proved that it can obtain approximate solutions faster than traditional PDE solvers through neural network experiments [3], such as Convolutional Neural Networks [4], Recurrent Neural Network [5], Generative Adversarial Networks [6], DeepONet [7], physic informed neural network (PINN) [8-10] and so on. Recently, Zhang [11] et al proposed the Bilinear Neural Network Method (BNNM) for investigating the analytical solutions of PDEs based on the bilinear method and the Neural Network model. This method was used in many constant coefficient PDEs, such as p-generalized

<sup>&</sup>lt;sup>1</sup>\*Corresponding author: Jian-Guo Liu, Email: 20101059@jxutcm.edu.cn; Wen-Hui Zhu, Email: 415422402@qq.com.

B-type Kadomstev-Petviashvili (KP) equation [12], Caudrey-Dodd-Gibbon-Kotera-Sawada-like equation [13], (3+1)-dimensional Jimbo-Miwa equation [14], Boiti-Leon-Manna-Pempinelli equation [15] and so on. However, the BNNM has never been used in variable coefficient PDEs. Prompted by this, we modify this method and apply it to study the analytical solutions of the KP equation with variable coefficients.

In this article, under consideration is the following KP equation with variable coefficients [16]

$$\alpha u_x^2 + \alpha u u_{xx} + \beta u_{xxxx} - \gamma u_{yy} + u_{xt} = 0, \qquad (1)$$

where u = u(x, y, t).  $\alpha = \alpha(t)$ ,  $\beta = \beta(t)$  and  $\gamma = \gamma(t)$  mean the coefficients of nonlinearity, dispersion and disturbed wave velocity, respectively. Eq. (1) represents waves in ferromagnetic media, water waves of long wavelength with weakly non-linear restoring forces and frequency dispersion. Wu et al. [16-18] have studied the Bäcklund transformation, soliton, Wronskian and Gramian solutions for Eq. (1). Jia [19] et al. have investigated the lump and rogue waves of Eq. (1). The lump, breather wave and interaction solutions have been obtained in Ref. [20,21]. Multiple rogue wave solutions were derived based on the variable-coefficient symbolic computation approach [22]. Nonautonomous lump solutions were presented in Ref. [23]. Double-periodic soliton and nontraveling wave solutions have been given in Ref. [24]. Solitary and lump waves interaction were investigated by using a modified Ansätz with variable coefficients [25]. However, the bilinear neural network method has not been used in Eq. (1), which will become our main work.

Let's do the following transformation

$$\alpha = \frac{6\beta}{\Theta_0}, \quad u = 2\Theta_0 (ln\xi)_{xx}, \quad \xi = \xi(x, y, t), \tag{2}$$

Eq. (1) has the bilinear form

$$(\beta D_x^4 - \gamma D_y^2 + D_t D_x)\xi \cdot \xi = \xi(\beta \xi_{xxxx} - \gamma \xi_{yy} + \xi_{xt}) + 3\beta \xi_{xx}^2 - 4\beta \xi_x \xi_{xxx} + \gamma \xi_y^2 - \xi_t \xi_x = 0,$$
(3)

where  $\Theta_0$  is an arbitrary constant.

The paper is organized as follows. Section 2 proposes a variable coefficient Bilinear neural network method (vcBNNM); Section 3 obtains the analytical solutions of the KP equation with variable coefficients by using the vcBNNM. Meanwhile, the dynamic properties are described in some three-dimensional and contour plots. Section 4 makes a summary.

# 2. vcBNNM

In order to apply the BNNM in Ref. [11] to the variable coefficient PDEs, we will improve the BNNM as follows (named vcBNNM) (I) The tensor formula of vcBNNM is supposed as follows

$$\xi = \omega_{l_n,\xi} F_{l_n}(\xi_{l_n}), \tag{4}$$

where  $\xi = \xi(x, y, \dots, t)$ ,  $\xi_{l_n} = \xi_{l_n}(x, y, \dots, t)$ ,  $\omega_{l_n,\xi}$  is the weight coefficient of neuron  $l_n$  to  $\xi$ . F is a generalized activation function that can be defined arbitrarily and must be satisfied that  $F_{l_n}(\xi_{l_n}) \ge 0$  in the last layer.  $l_n =$  $m_{n-1} + 1, m_{n-1} + 2, \dots, n$  denotes the nth layer space of the neural network model.  $\xi_{l_i}$  is assumed as follows

$$\xi_{l_i} = \omega_{l_{i-1}, l_i} F_{l_{i-1}}(\xi_{l_{i-1}}) + \omega_{l_i}(t), \quad i = 1, 2, \cdots, n,$$
(5)

where  $l_0 = x, y, \dots, \omega_{l_0}(t), l_1 = 1, 2, \dots, m_1, l_i = m_{i-1} + 1, m_{i-1} + 2, \dots, m_i (i = 2, 3, \dots, n-1), \omega_{l_i}(t)$  represents the threshold, which is different from the constant hypothesis in Ref. [11]. This neural network tensor model is shown in Fig. 1.

(II) Substituting Eq. (4) into the Hirota bilinear form of variable coefficient PDEs and making the coefficient of  $F_{l_i}(\xi_{l_i})$  and its derivatives of all orders equal to zero, a sets of algebraic equations will be obtained. With the aid of Mathematica, the values of the parameters in Eq. (4) and Eq. (5) can be derived. Finally, substituting these values and nonlinear neural network tensor for- mula Eq. (10) into Eq. (4) and Eq. (5), the corresponding analytical solutions for the variable coefficient PDEs can be presented.



**Fig.** 1. Nonlinear neural network of Eq. (4) and Eq. (5).



Fig. 2.  $\eta_1 = \omega_{1,2} = \omega_{2,1} = \omega_{2,3} = \Theta_0 = 1$ ,  $\eta_2 = \omega_{1,1} = \omega_{1,3} = 2$ ,  $\omega_{2,2} = -2$ , y = 0, (a) three-dimensional plot, (b) density plot.

#### 3. Application

To verify the effectiveness of the vcBNNM in Section 2, we use the variable coefficient KP equation (1) as an example and select a "3-2-2-1" variable coefficient neural network model, which represents that the input layer  $l_0 = \{x, y, \omega_{l_i}(t)\}(i = 1, 2)$  has 3 neurons, the hidden layer  $l_1 = \{1, 2\}$  has 2 neurons, the hidden layer  $l_2 = \{3, 4\}$  has 2 neurons and the print layer  $\xi$  has 1 neuron. According to the idea of vcBNNM, we suppose

$$\begin{aligned} \xi_1 &= \omega_{3,1}(t) + x\omega_{1,1} + y\omega_{2,1}, \\ \xi_2 &= \omega_{3,2}(t) + x\omega_{1,2} + y\omega_{2,2}, \\ \xi_3 &= \omega_{1,3}F_1(\xi_1) + \omega_{2,3}F_2(\xi_2) + \omega_{3,3}(t), \\ \xi_4 &= \omega_{1,4}F_1(\xi_1) + \omega_{2,4}F_2(\xi_2) + \omega_{3,4}(t), \\ \xi &= \omega_{1,5}F_3(\xi_3) + \omega_{2,5}F_4(\xi_4) + \omega_{3,5}(t), \end{aligned}$$
(6)

where  $F_1(\xi_1) = \cos \xi_1$ ,  $F_2(\xi_2) = \exp \xi_2$ ,  $F_3(\xi_3) = \xi_3^2$ ,  $F_4(\xi_4) = \xi_4^2$ ,  $\omega_{i,j}(i = 1, 2; j = 1, 2, 3, 4, 5)$  is unknown constant,  $\omega_{3,j}(t)(j = 1, 2, 3, 4, 5)$  is unknown function, which is different from the constant hypothesis in Ref. [11], and the purpose is to deal with the variable coefficients in Eq. (1). This is also the biggest difference and innovation between the vcBNNM and the BNNM. Substituting Eq. (6) into Eq. (3) by using Mathematica software, we have

$$\begin{split} \omega_{2,4} &= \frac{\omega_{1,4}\omega_{2,3}}{\omega_{1,3}}, \omega_{2,5} = -\frac{\omega_{1,3}^2\omega_{1,5}}{\omega_{1,4}^2}, \omega_{3,1}(t) = \eta_2 + \\ \frac{\left(-\omega_{2,2}^2\omega_{1,1}^4 + 2\omega_{1,2}\omega_{2,1}\omega_{2,2}\omega_{1,1}^3 + \omega_{1,2}^2 \left(2\omega_{1,1}^2 + 3\omega_{1,2}^2\right)\omega_{2,1}^2\right)\int\gamma\,dt}{3\omega_{1,1}\omega_{1,2}^2 \left(\omega_{1,1}^2 + \omega_{1,2}^2\right)}, \\ \omega_{3,2}(t) &= \eta_1 + \left[\left[\omega_{1,2}^2 \left(\omega_{1,2}^2 - 6\omega_{1,1}^2\right)\omega_{2,1}^2 - 2\omega_{1,1}\omega_{1,2} \left(\omega_{1,2}^2 - 6\omega_{1,1}^2\right)\omega_{2,2}\omega_{2,1}\right.\right] \\ &+ \omega_{1,1}^2 \left(4\omega_{1,2}^2 - 3\omega_{1,1}^2\right)\omega_{2,2}^2\right]\int\gamma\,dt\right]/\left[3\omega_{1,1}^2\omega_{1,2} \left(\omega_{1,1}^2 + \omega_{1,2}^2\right)\right], \\ \omega_{3,5}(t) &= \omega_{1,5}\left[\frac{\omega_{1,3}^2\omega_{3,4}(t)^2}{\omega_{1,4}^2} - \omega_{3,3}(t)^2\right], \beta = -\frac{\left(\omega_{1,2}\omega_{2,1} - \omega_{1,1}\omega_{2,2}\right)^2\gamma}{3\omega_{1,1}^2\omega_{1,2}^2 \left(\omega_{1,1}^2 + \omega_{1,2}^2\right)}, \end{split}$$
(7)

where  $\eta_1$  and  $\eta_2$  are integration constants. Substituting Eq. (6) and Eq. (7) into Eq. (2), the corresponding analytical solution for Eq. (1) can be obtained as follows

$$u = [2\Theta_{0}[(\omega_{1,5}F_{3}(\xi_{3}) + \omega_{2,5}F_{4}(\xi_{4}) + \omega_{3,5}(t)) [\omega_{1,5}F_{3}''(\xi_{3}) [\omega_{1,1}\omega_{1,3}F_{1}'(\xi_{1}) + \omega_{1,2}\omega_{2,3}F_{2}''(\xi_{2})]^{2} + \omega_{1,5}F_{3}'(\xi_{3}) [\omega_{1,3}F_{1}''(\xi_{1}) \omega_{1,1}^{2} + \omega_{1,2}^{2}\omega_{2,3}F_{2}''(\xi_{2})] + \omega_{2,5}F_{4}'(\xi_{4}) [\omega_{1,4}F_{1}''(\xi_{1}) \omega_{1,1}^{2} + \omega_{1,2}^{2}\omega_{2,4}F_{2}''(\xi_{2})] + \omega_{2,5}[\omega_{1,1}\omega_{1,4}F_{1}'(\xi_{1}) + \omega_{1,2}\omega_{2,4}F_{2}''(\xi_{2})]^{2}F_{4}'''(\xi_{4})] - [\omega_{1,5}[\omega_{1,1}\omega_{1,3}F_{1}'(\xi_{1}) + \omega_{1,2}\omega_{2,3}F_{2}'(\xi_{2})]F_{3}'(\xi_{3}) + \omega_{2,5}[\omega_{1,1}\omega_{1,4}F_{1}'(\xi_{1}) + \omega_{1,2}\omega_{2,4}F_{2}'(\xi_{2})]F_{4}'(\xi_{4})]^{2}]] / [[\omega_{1,5}F_{3}(\xi_{3}) + \omega_{2,5}F_{4}(\xi_{4}) + \omega_{3,5}(t)]^{2}], \qquad (8)$$

where  $F_i^{(n)}(\xi_i) = \frac{d^n F_i(\xi_i)}{d\xi_i^n}$  (i = 1, 2). All parameters are arbitrary except Eq. (7). In order to discuss the influence of variable coefficients  $\gamma$ , we take  $\gamma = 1$  and  $\gamma = \cos t$  respectively to show the dynamic properties of solution (8) (see Fig. 2 and Fig. 3).



Fig. 3.  $\eta_1 = \omega_{1,2} = \omega_{2,1} = \omega_{2,3} = \Theta_0 = 1$ ,  $\eta_2 = \omega_{1,1} = \omega_{1,3} = 2$ ,  $\omega_{2,2} = -2$ , y = 0, (a) three-dimensional plot, (b) density plot.

Choosing  $F_2(\xi_2) = \cosh \xi_2$  in Eq. (6), we have

$$\begin{split} \omega_{2,4} &= \frac{\omega_{1,4}\omega_{2,3}}{\omega_{1,3}}, \omega_{2,5} = -\frac{\omega_{1,5}\omega_{2,3}^2}{\omega_{2,4}^2}, \omega_{3,1}(t) = \eta_4 + \left[ \left[ \omega_{1,1} \left[ -4\omega_{2,2}^2 \omega_{1,1}^4 \right] \right] \right] \right] \\ &+ 8\omega_{1,2}\omega_{2,1}\omega_{2,2}\omega_{1,1}^3 + \left( 3\omega_{1,1}^4 + 2\omega_{1,2}^2 \omega_{1,1}^2 + 3\omega_{1,2}^4 \right) \right] \\ &+ \left( \omega_{1,1}^4 - 6\omega_{1,2}^2 \omega_{1,1}^2 - 3\omega_{1,2}^4 \right) \right] \\ &+ 3\omega_{1,2}^4 (\omega_{2,1}\omega_{2,2}) \right] \\ &= \eta_3 + \left[ \left[ \left[ 4\omega_{1,1}^2 \omega_{2,2}^2 \omega_{1,2}^3 + \left( -3\omega_{1,1}^4 - 6\omega_{1,2}^2 \omega_{1,1}^2 + \omega_{1,2}^2 \omega_{2,2}^2 \right) \right] \right] \right] \\ &+ 3\omega_{1,2}(t) = \eta_3 \\ &+ \left[ \left[ \left[ 4\omega_{1,1}^2 \omega_{2,2}^2 \omega_{1,2}^3 + \left( -3\omega_{1,1}^4 - 6\omega_{1,2}^2 \omega_{1,1}^2 + \omega_{1,2}^2 \omega_{2,2}^2 \right) \right] \right] \right] \\ &+ 2\omega_{1,1} \left( 3\omega_{1,1}^4 + 6\omega_{1,2}^2 \omega_{1,1}^2 - \omega_{1,2}^4 \right) \right] \\ &+ 2\omega_{1,1} \left( 3\omega_{1,1}^4 + 6\omega_{1,2}^2 \omega_{1,1}^2 - \omega_{1,2}^4 \right) \\ &+ 8\omega_{1,1}\omega_{2,1}\omega_{2,2}\omega_{1,2}^3 + \left( 3\omega_{1,1}^4 + 2\omega_{1,2}^2 \omega_{2,1}^2 + 3\omega_{1,2}^4 \right) \right] \\ &+ 8\omega_{1,1}\omega_{2,1}\omega_{2,2}\omega_{1,2}^3 + \left( 3\omega_{1,1}^4 + 2\omega_{1,2}^2 \omega_{2,3}^2 \right) \right] , \\ &\omega_{3,5}(t) = \omega_{1,5} \left[ \frac{\omega_{1,3}^2 \omega_{3,4}(t)^2}{\omega_{1,4}^2} - \omega_{3,3}(t)^2 \right] , \\ &\beta = -\frac{\left( \omega_{1,2}\omega_{2,1} - \omega_{1,1}\omega_{2,2} \right)^2 \left( \omega_{1,3}^2 - \omega_{2,3}^2 \right) \gamma}{3 \left( \omega_{1,1}^2 + \omega_{1,2}^2 \right)^2 \left( \omega_{1,1}^2 - \omega_{1,3}^2 + \omega_{1,2}^2 \omega_{2,3}^2 \right)}, \end{aligned}$$
(9)

where  $\eta_3$  and  $\eta_4$  are integration constants. Substituting Eq. (6) and Eq. (9) into Eq. (2), the corresponding analytical solution of the second kind for Eq.

(1) can be given as follows

$$u = [2\Theta_{0}[(\omega_{1,5}F_{3}(\xi_{3}) + \omega_{2,5}F_{4}(\xi_{4}) + \omega_{3,5}(t)) [\omega_{1,5}F_{3}''(\xi_{3}) [\omega_{1,1}\omega_{1,3}F_{1}'(\xi_{1}) + \omega_{1,2}\omega_{2,3}F_{2}''(\xi_{2})]^{2} + \omega_{1,5}F_{3}'(\xi_{3}) [\omega_{1,3}F_{1}''(\xi_{1}) \omega_{1,1}^{2} + \omega_{1,2}^{2}\omega_{2,3}F_{2}''(\xi_{2})] + \omega_{2,5}F_{4}'(\xi_{4}) [\omega_{1,4}F_{1}''(\xi_{1}) \omega_{1,1}^{2} + \omega_{1,2}^{2}\omega_{2,4}F_{2}''(\xi_{2})] + \omega_{2,5}[\omega_{1,1}\omega_{1,4}F_{1}'(\xi_{1}) + \omega_{1,2}\omega_{2,4}F_{2}''(\xi_{2})]^{2}F_{4}''(\xi_{4})] - [\omega_{1,5}[\omega_{1,1}\omega_{1,3}F_{1}'(\xi_{1}) + \omega_{1,2}\omega_{2,3}F_{2}'(\xi_{2})]F_{3}'(\xi_{3}) + \omega_{2,5}[\omega_{1,1}\omega_{1,4}F_{1}'(\xi_{1}) + \omega_{1,2}\omega_{2,4}F_{2}'(\xi_{2})]F_{4}'(\xi_{4})]^{2}]] / [[\omega_{1,5}F_{3}(\xi_{3}) + \omega_{2,5}F_{4}(\xi_{4}) + \omega_{3,5}(t)]^{2}].$$

$$(10)$$

All parameters are arbitrary except Eq. (9). When  $\gamma = 1$  and  $\gamma = \sin t$ , the dynamic properties of solution (10) are demonstrated in Fig. 4 and Fig. 5.



Fig. 4.  $\eta_3 = \omega_{1,3} = \omega_{2,1} = \omega_{2,3} = \Theta_0 = 1$ ,  $\eta_4 = \omega_{1,1} = \omega_{1,2} = 2$ ,  $\omega_{2,2} = -2$ , y = 0, (a) three-dimensional plot, (b) density plot.



Fig. 5.  $\eta_3 = \omega_{1,3} = \omega_{2,1} = \omega_{2,3} = \Theta_0 = 1$ ,  $\eta_4 = \omega_{1,1} = \omega_{1,2} = 2$ ,  $\omega_{2,2} = -2$ , y = 0, (a) three-dimensional plot, (b) density plot.

Finally, choose a "3-3-2-1" variable coefficient neural network model, which denotes that the input layer  $l_0 = \{x, y, \omega_{l_i}(t)\}(i = 1, 2, 3)$  has 3 neurons, the

6

hidden layer  $l_1 = \{1, 2, 3\}$  has 3 neurons, the hidden layer  $l_2 = \{4, 5\}$  has 2 neurons and the print layer  $\xi$  has 1 neuron. According to the step of vcBNNM, we suppose

$$\begin{aligned} \xi_1 &= \omega_{3,1}(t) + x\omega_{1,1} + y\omega_{2,1}, \\ \xi_2 &= \omega_{3,2}(t) + x\omega_{1,2} + y\omega_{2,2}, \\ \xi_3 &= \omega_{3,3}(t) + x\omega_{1,3} + y\omega_{2,3}, \\ \xi_4 &= \omega_{1,4}F_1(\xi_1) + \omega_{2,4}F_2(\xi_2) + \omega_{3,4}F_3(\xi_3) + \omega_{4,4}(t), \\ \xi_5 &= \omega_{1,5}F_1(\xi_1) + \omega_{2,5}F_2(\xi_2) + \omega_{3,5}F_3(\xi_3) + \omega_{4,5}(t), \\ \xi &= \omega_{1,6}F_4(\xi_4) + \omega_{2,6}F_5(\xi_5) + \omega_{3,6}(t), \end{aligned}$$
(11)

where  $F_1(\xi_1) = \cos \xi_1$ ,  $F_2(\xi_2) = \cosh \xi_2$ ,  $F_3(\xi_3) = \cosh \xi_3$ ,  $F_4(\xi_4) = \xi_4^2$ ,  $F_5(\xi_5) = \xi_5^2$ ,  $\omega_{i,j}(i = 1, 2; j = 1, 2, 3, 4, 5, 6)$  and  $\omega_{3,j}(j = 4, 5)$  are unknown constants,  $\omega_{3,j}(t)(j = 1, 2, 3, 6)$  and  $\omega_{4,j}(t)(j = 4, 5)$  are unknown functions. Substituting Eq. (11) into Eq. (3) by utilizing Mathematica software, we get

$$\begin{split} \omega_{2,6} &= -\frac{\omega_{1,6}\omega_{2,4}^2}{\omega_{2,5}^2}, \\ \omega_{3,5} &= \frac{\omega_{1,5}\omega_{3,4}}{\omega_{1,4}}, \\ \omega_{2,4} &= \frac{\omega_{1,4}\omega_{2,5}}{\omega_{1,5}}, \\ \omega_{3,1}(t) &= [[\omega_{1,1}](\omega_{1,1}^2) \\ -3\omega_{1,2}^2)\omega_{1,2}^2 + 3(\omega_{1,1}^2 + \omega_{1,2}^2)\omega_{1,3}^2]\omega_{2,1}^2 + 2\omega_{1,2}[-\omega_{1,1}^4 + 3(\omega_{1,2}^2 + \omega_{1,3}^2)\omega_{1,1}^2] \\ &+ 3\omega_{1,2}^2\omega_{1,3}^2]\omega_{2,2}\omega_{2,1} + \omega_{1,1}[\omega_{1,1}^4 - 3(\omega_{1,2}^2 + \omega_{1,3}^2)\omega_{1,1}^2 - 3\omega_{1,2}^2\omega_{1,3}^2]\omega_{2,2}^2] \\ &* \int \gamma dt] / [3(\omega_{1,1}^2 + \omega_{1,2}^2)^2\omega_{1,3}^2] + \eta_6, \\ \omega_{3,2}(t) &= [[-\omega_{1,2}[\omega_{1,2}^4 - 3\omega_{1,1}^2\omega_{1,2}^2] \\ &+ 3(\omega_{1,1}^2 + \omega_{1,2}^2)\omega_{1,3}^2]\omega_{2,1}^2 + 2\omega_{1,1}[\omega_{1,2}^4 - 3\omega_{1,1}^2\omega_{1,2}^2 + 3(\omega_{1,1}^2 + \omega_{1,2}^2)\omega_{1,3}^2] \\ &+ 3(\omega_{1,1}^2 + \omega_{1,2}^2)\omega_{1,3}^2]\omega_{2,1}^2 + 2\omega_{1,1}[\omega_{1,2}^4 - 3\omega_{1,1}^2\omega_{1,2}^2 + 3(\omega_{1,1}^2 + \omega_{1,2}^2)\omega_{1,3}^2] \\ &+ \omega_{2,2}\omega_{2,1} + \omega_{1,2}[3\omega_{1,1}^4 - \omega_{1,2}^2\omega_{1,1}^2 + 3(\omega_{1,1}^2 + \omega_{1,2}^2)\omega_{1,3}^2] \int \gamma dt] \\ / [3(\omega_{1,1}^2 + \omega_{1,2}^2)^2\omega_{1,3}^2] + \eta_5, \\ \omega_{3,4}(u_{1,1}^2 + \omega_{1,2}^2)^2\omega_{1,3}^2] + \eta_5, \\ \omega_{3,4}(u_{1,1}^2 + \omega_{1,2}^2)\omega_{1,4}\sqrt{(\omega_{1,1}^2 + \omega_{1,2}^2)^2\omega_{1,3}^2] + \eta_7, \\ \omega_{3,4}(u_{1,1}^2 + \omega_{1,2}^2)\omega_{1,4}\sqrt{(\omega_{1,1}^2 + \omega_{1,3}^2)\omega_{2,2}^2, \\ \sqrt{(\omega_{1,2}^2 - \omega_{1,3}^2)(\omega_{1,1}^2 + \omega_{1,3}^2)(\omega_{1,1}^2 + \omega_{1,3}^2)(\omega_{1,2}^2, - \omega_{1,3}^2)(\omega_{2,5}^2, \\ \omega_{3,4}(u_{1,1}^2 + \omega_{1,2}^2)\omega_{1,4}\sqrt{(\omega_{1,1}^2 + \omega_{1,3}^2)(\omega_{1,1}^2 + \omega_{1,3}^2)(\omega_{2,5}^2, \\ \sqrt{(\omega_{1,2}^2 - \omega_{1,3}^2)(\omega_{1,1}^2 + \omega_{1,3}^2)(\omega_{1,5}^2, - \omega_{1,3}^2)(\omega_{2,5}^2, \\ \sqrt{(\omega_{1,2}^2 - \omega_{1,3}^2)(\omega_{1,1}^2 + \omega_{1,3}^2)(\omega_{1,5}^2, - \omega_{1,4})} \\ \omega_{3,4}(u_{1,1}^2 + \omega_{1,3}^2)^2\gamma, \\ \omega_{3,4}(u_{1,1}^2 + \omega_{1,3}^2)^2\gamma, \\ \omega_{3,6}(u_{1,1}^2 + \omega_{1,3}^2)(\omega_{1,1}^2 + \omega_{1,3}^2)(\omega_{1,5}^2, - \omega_{1,4})(u_{1,1}^2, - u_{1,3}^2)(\omega_{1,5}^2, - \omega_{1,4})(u_{1,1}^2, - u_{1,3}^2)(\omega_{1,5}^2, - \omega_{1,4})(u_{1,5}^2, - \omega_{1,4})(u_{1,5}^2, - \omega_{1,5}^2, - \omega_{1,4})(u_{1,1}^2, - u_{1,3}^2)(\omega_{1,5}^2, - \omega_{1,4})(u_{1,1}^2, - u_{1,3}^2)(\omega_{1,5}^2, - \omega_{1,4})(u_{1,5}^2, - \omega_{1,4})(u_{1,5}^2, - \omega_{1,5}^2, - \omega_{1,4})(u_{1,5}^2, - \omega_{1,5}^2, - \omega_{1,4})(u_{1,5}^2, - \omega_{1,5}^2$$

where  $\eta_5$ ,  $\eta_6$  and  $\eta_7$  are integration constants. Substituting Eq. (11) and Eq. (12) into Eq. (2), the corresponding analytical solution for Eq. (1) can be

shown as follows

$$u = [2\Theta_{0}[[\omega_{1,6}F_{4}(\xi_{4}) + \omega_{2,6}F_{5}(\xi_{5}) + \omega_{3,6}(t)][\omega_{1,6}F_{4}''(\xi_{4})[\omega_{1,1}\omega_{1,4}F_{1}'(\xi_{1}) + \omega_{1,2}\omega_{2,4}F_{2}'(\xi_{2}) + \omega_{1,3}\omega_{3,4}F_{3}'(\xi_{3})]^{2} + \omega_{1,6}F_{4}'(\xi_{4})[\omega_{1,4}F_{1}''(\xi_{1})\omega_{1,1}^{2} + \omega_{1,2}^{2}\omega_{2,4}F_{2}''(\xi_{2}) + \omega_{1,3}^{2}\omega_{3,4}F_{3}''(\xi_{3})] + \omega_{2,6}F_{5}'(\xi_{5})[\omega_{1,5}F_{1}''(\xi_{1})\omega_{1,1}^{2} + \omega_{1,2}^{2}\omega_{2,5}F_{2}''(\xi_{2}) + \omega_{1,3}^{2}\omega_{3,5}F_{3}''(\xi_{3})] + \omega_{2,6}[\omega_{1,1}\omega_{1,5}F_{1}'(\xi_{1}) + \omega_{1,2}\omega_{2,5}F_{2}'(\xi_{2}) + \omega_{1,3}\omega_{3,5}F_{3}'(\xi_{3})]^{2}F_{5}''(\xi_{5})] - [\omega_{1,6}[\omega_{1,1}\omega_{1,4}F_{1}'(\xi_{1}) + \omega_{1,2}\omega_{2,4}F_{2}'(\xi_{2}) + \omega_{1,3}\omega_{3,4}F_{3}'(\xi_{3})]F_{4}'(\xi_{4}) + \omega_{2,6}[\omega_{1,1}\omega_{1,5}F_{1}'(\xi_{1}) + \omega_{1,2}\omega_{2,5}F_{2}'(\xi_{2}) + \omega_{1,3}\omega_{3,5}F_{3}'(\xi_{3})]F_{5}'(\xi_{5})]^{2}]]/[[\omega_{1,6}F_{4}(\xi_{4}) + \omega_{2,6}F_{5}(\xi_{5}) + \omega_{3,6}(t)]^{2}].$$
(13)

All parameters are arbitrary except Eq. (12). When  $\gamma = 1$  and  $\gamma = \cos t$ , the dynamic properties of solution (13) are demonstrated in Fig. 6 and Fig. 7.



**Fig.** 6.  $\eta_5 = \omega_{1,3} = \omega_{1,4} = \omega_{1,5} = \omega_{2,1} = \omega_{2,3} = \Theta_0 = 1$ ,  $\eta_6 = \omega_{1,1} = \omega_{1,2} = 2$ ,  $\omega_{2,2} = -3$ ,  $\eta_7 = \omega_{2,5} = -1$ , (a)(d) t = -2, (b)(e) t = 0, (c)(f) t = 2.

## 4. Conclusion

The BNNM is a very direct and effective method to obtain the analytical solutions of nonlinear PDEs with constant coefficients, but it is not suitable for solving the analytical solutions of nonlinear PDEs with variable coefficients. Therefore, we propose an improved BNNM named vcBNNM to deal with the problem of analytical solutions of nonlinear PDEs with variable coefficients. To verify the validity of the method, we take the variable coefficient KP equation as an example. By establishing "3-2-2-1" and "3-3-2-1" models, we successfully obtain a rich analytical solutions of the KP equation with variable coefficients, which contains more arbitrary parameters. Meanwhile, by selecting some special values of the parameters, the dynamics properties of these derived solutions are shown in Fig. 1. Compared with the previous BNNM, we change the parameter  $\omega_{l_i}(t)$  in Eq. (5) from constant to function, and the variable t in the input layer  $l_0$  also becomes a nonlinear function. In this way, the existence of variable coefficients in the PDEs with variable coefficients can be well handled, and more analytical solutions can be obtained.



Fig. 7.  $\eta_5 = \omega_{1,3} = \omega_{1,4} = \omega_{1,5} = \omega_{2,1} = \omega_{2,3} = \Theta_0 = 1$ ,  $\eta_6 = \omega_{1,1} = \omega_{1,2} = 2$ ,  $\omega_{2,2} = -3$ ,  $\eta_7 = \omega_{2,5} = -1$ , y = 0, (a) three-dimensional plot, (b) density plot.

**Funding:** Project supported by National Natural Science Foundation of China (Grant No. 12161048), Doctoral Research Foundation of Jiangxi University of Chinese Medicine (Grant No. 2021WBZR007) and Jiangxi University of Chinese Medicine Science and Technology Innovation Team Development Program (Grant No. CXTD22015).

#### **Data Availability Statements**

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

### Declaration

**Conflict of interests** The authors declare that there is no conflict of interests regarding the publication of this article.

**Ethical standard** The authors state that this research complies with ethical standards. This research does not involve either human participants or animals.

#### References

- Sirignano J, Spiliopoulos K. DGM: a deep learning algorithm for solving partial differential equations. J. Comput. Phys. 2018;375:1339-1364.
- [2] Wang Z, Weng F, Liu J, et al. Numerical solution for high-dimensional partial differential equations based on deep learning with residual learning and data-driven learning. Int. J. Mach. Learn. Cyb. 2022;134:108340.
- [3] Li J, Chen Y. A deep learning method for solving third-order nonlinear evolution equations. Commun. Theor. Phys. 2020;72:115003.
- [4] Gao P, Zhao Z, Yang Y. Study on Numerical Solutions to Hyperbolic Partial Differential Equations Based on the Convolutional Neural Network Model. Appl. Math. Mech. 2021;42(9):932-947.
- [5] Li Y, Wang X. Almost periodic solutions in distribution of Clifford-valued stochastic recurrent neural networks with time-varying delays. Chaos Soliton. Fract. 2021;153(Part 2):111536.
- [6] Kadeethum T, ÓMalley D, Choi Y, et al. Continuous conditional generative adversarial networks for data-driven solutions of poroelasticity with heterogeneous material properties. Comput. Geosci. 2022;167:105212.
- [7] Wang S, Wang H, Perdikaris P. Learning the solution operator of parametric partial differential equations with physics-informed DeepONets. Sci. Adv. 2021;7(40):eabi8605.
- [8] Bai Y, Temuer C, Sudao B. Solving Huxley equation using an improved PINN method. Nonlinear Dyn. 2021;105(4)3439-3450.
- [9] Fang Y, Wang YY, Liu W, Dai CQ. Data-driven prediction of soliton solutions of the higher-order NLSE via the strongly-constrained PINN method. Comput. Math. Appl. 2022(127):144-153.
- [10] Peng WQ, Pu JC, Chen Y. PINN deep learning method for the Chen-Lee-Liu equation: Rogue wave on the periodic background. Commun. Nonlinear Sci. 2022;105:106067.
- [11] Zhang RF, Bilige SD. Bilinear neural network method to obtain the exact analytical solutions of nonlinear partial differential equations and its application to p-gBKP equation. Nonlinear Dyn. 2019;95:3041-3048.
- [12] Zhang RF, Bilige SD, Liu JG, Li M. Bright-dark solitons and interaction phenomenon for p-gBKP equation by using bilinear neural network method. Phys. Scr. 2021;96:025224.
- [13] Zhang RF, Li MC, Mohammed A, Zheng FC, Lan ZZ. Generalized lump solutions, classical lump solutions and rogue waves of the (2+1)-dimensional Caudrey-Dodd-Gibbon-Kotera-Sawada-like equation. Appl. Math. Comput. 2021;403:126201.
- [14] Zhang RF, Li MC, Yin HM. Rogue wave solutions and the bright and dark solitons of the (3+1)-dimensional Jimbo-Miwa equation. Nonlinear Dyn. 2021;103:1071-1079.
- [15] Qiao JM, Zhang RF, Yue RX, Rezazadeh H, Seadawy AR. Three types of periodic solutions of new (3+1)-dimensional Boiti-Leon-Manna-Pempinelli equation via bilinear neural network method. 2022;45(9):5612-5621.
- [16] Wu JP. Bilinear bäcklund transformation for a variable-coefficient Kadomtsev-Petviashvili equation. Chin. Phys. Lett., 2011;28(6):060207.
- [17] Wang YY, Zhang JF. Variable-coefficient KP equation and solitonic solution for twotemperature ions in dusty plasma. Phys. Lett. A. 2006;352(1):155-162.
- [18] Yao ZZ, Zhang CY, Zhu HW, et al. Wronskian and grammian determinant solutions for a variable-coefficient Kadomtsev-Petviashvili equation. Commun. Theor. Phys. 2008;49(5):1125-1128.
- [19] Jia XY, Tian B, Du Z, Sun Y, Liu L. Lump and rogue waves for the variable-coefficient Kadomtsev-Petviashvili equation in a fluid. Mod. Phys. Lett. B. 2018;32(10):1850086.
- [20] Liu JG, Zhu WH, Zhou L. Interaction Solutions for Kadomtsev-Petviashvili Equation with Variable Coefficients. Commun. Theor. Phys. 2019;71:793-797.

- [21] Liu JG, Zhu WH, Zhou L. Breather wave solutions for the Kadomtsev-Petviashvili equation with variable coefficients in a fluid based on the variable-coefficient threewave approach. Math. Meth. Appl. Sci. 2020;43(1):458-465.
- [22] Liu JG, Zhu WH, He Y. Variable-coefficient symbolic computation approach for finding multiple rogue wave solutions of nonlinear system with variable coefficients. Z. Angew. Math. Phys. 2021;72:154.
- [23] Wang YH. Nonautonomous lump solutions for a variable-coefficient Kadomtsev- Petviashvili equation. Appl. Math. Lett. 2021;119:107201.
- [24] Qin CR, Liu JG. Study on double-periodic soliton and non-traveling wave solutions of integrable systems with variable coefficients. Results Phys. 2022;34;105254.
- [25] Liu JG, Wazwaz AM, Zhu WH. Solitary and lump waves interaction in variablecoefficient nonlinear evolution equation by a modified Ansätz with variable coefficients. J. Appl. Anal. Comput. 2022;12(2):517-532.