

The Poisson Geometry of Plancherel Formulas for Triangular Groups

Nicholas M. Ercolani[†]

Department of Mathematics, University of Arizona, 617 N. Santa Rita Ave., Tucson, 85721-0089, AZ, USA

*Corresponding author. E-mail: ercolani@math.arizona.edu
ORCID: 0000-0003-2010-4205[†]

Abstract

In this paper we establish the existence of canonical coordinates for generic co-adjoint orbits on triangular groups. These orbits correspond to a set of full Plancherel measure on the associated dual groups. This generalizes a well-known coordinatization of co-adjoint orbits of a minimal (non-generic) type originally discovered by Flaschka. The latter had strong connections to the classical Toda lattice and its associated Poisson geometry. Our results develop connections with the Full Kostant-Toda lattice and its Poisson geometry. This leads to novel insights relating the details of Plancherel theorems for Borel Lie groups to the invariant theory for Borels and their subgroups. We also discuss some implications for the quantum integrability of the Full Kostant Toda lattice.

Keywords: orbit method, Dixmier-Pukanszky operator, Toda lattice, polarizations, invariant theory

Contents

1	Introduction	2
2	Background	5
2.1	Decomposition of a Real Semisimple Lie Groups	5
2.2	Decomposition of Unipotent Triangular subgroups	6
2.3	Schrödinger Representations of Heisenberg Groups	6
2.4	Stepwise Square Integrable Representations	7
2.4.1	Real Polarizations	7
2.4.2	Constructing the Essential Part of \widehat{N}	8
2.4.3	The Plancherel Formula for N	8
2.5	Poisson Geometry of Triangular Nilpotent Algebras	9
3	Borel Subgroups	9
3.1	Dixmier-Pukanszky Operator	10
3.2	Plancherel Theorem	11
4	Poisson Structures, Orbits and Invariant Theory	11
4.1	Poisson Reduction and Induction	11
4.2	Invariant Theory of N	14
4.3	The Dixmier-Pukanszky Operator Revisited	16
4.3.1	Some History of Two Perspectives	17
5	Polarization on Orbits and Canonical Structures	18
5.1	Pukanszky Polarization for Borels	18
5.2	Canonical Representation for the Poisson Structure of \mathfrak{b}^*	21
5.3	Plancherel's Formula Revisited	23

6	Conclusions	23
6.1	Canonical Coordinates and Standard Hamiltonian Systems	23
6.2	Quantization of Invariants and Quantum Integrability	24
A	Appendix: Enveloping Algebras	25
	References	26

1 Introduction

The purpose of this paper is to amplify and further explore the deep connections between group representations and Poisson geometry (which is the geometry underlying Hamiltonian mechanics). The perspectives we bring to this are grounded in the theory of integrable systems especially as related to generalizations of the Toda lattice. More specifically we study the bridge between the Plancherel theorem for the real (upper) triangular matrix groups, i.e., those of the form (with $b_{ii} > 0$)

$$B = \begin{bmatrix} b_{11} & * & \dots & \dots & * \\ & b_{22} & * & & \vdots \\ & & \ddots & \ddots & \vdots \\ & & & \ddots & * \\ & & & & b_{nn} \end{bmatrix}$$

and the phase space geometry of generic coadjoint orbits for those groups, which is the phase space geometry underlying the full Kostant Toda lattice [EFS93]. For groups and phase geometries in general, this bridge is referred to as the *orbit method* program and so this work provides further explicit examples for that program, illuminating some recent developments there as well as showing new results. Though we restrict to the case of triangular groups, our presentation is framed to make it natural to extend our results to general Borel subgroups of $GL(n, \mathbb{R})$ as well as Borels for other real semisimple Lie groups and so we will often refer to the triangular group as a Borel subgroup. This work also opens a door for applications to quantum Toda lattices which will be discussed in our Conclusions.

Before getting to the outline of our results we will provide some general motivation concerning Plancherel theory. (For more details on this background we refer the reader to [DM72].) Recall the original Plancherel theorem which is one of the cornerstones of classical harmonic analysis related to Fourier theory. In the case of the one-dimensional unitary group, S^1 , Plancherel states that for a function $f(\theta) \in L^2(S^1)$ the map from f to its Fourier coefficients, $\hat{f}(n) \in \ell^2(\mathbb{Z})$, is an isometry ($\|f\|_2 = \|\hat{f}\|_2$). Moreover, this isometry is explicitly given by

$$f(\theta) = \sum_{n=-\infty}^{\infty} \hat{f}(n)e^{2\pi in\theta} \tag{1.1}$$

in the sense that the series converges to $f(\theta)$ in the L^2 -norm. (In what follows we will refer to formulas like (1.1) as *Plancherel formulas*.) From the group theoretic perspective, (1.1) is seen as an expansion over all homomorphisms, $e^{2\pi in\theta}$, indexed by $n \in \mathbb{Z}$, from S^1 to \mathbb{C}^* , which are usually referred to as *characters*. Since S^1 is a commutative group, one knows that its irreducible representations are all 1-dimensional and these characters constitute all of the irreducible unitary representations of S^1 . The latter set, corresponding to the additive group \mathbb{Z} , is referred to as the dual group and denoted, in this case, by $\widehat{S^1}$. A similar story to the above holds for $L^2(\mathbb{R})$ where \mathbb{R} is regarded as a group with respect to translation, corresponding to the Fourier integral representation of $f(x) \in L^2(\mathbb{R})$.

However the story changes significantly when considering harmonic analysis on a non-commutative group G . One dimensional homomorphisms no longer suffice to give a complete decomposition of $L^2(G)$. This was realized, even going back to Frobenius, in the case of finite groups, who understood that in the non-commutative setting there were irreducible representations (homomorphisms from G to $GL(d, \mathbb{C})$) with $d > 1$, whose matrix coefficients needed to be involved in any Fourier-like decomposition of L^2 . For compact groups

the description of how this goes is based on the Peter-Weyl theorem [Kn86] which classifies all the irreducible unitary representations of G . Though no longer always one dimensional, these are still finite dimensional and constitute a discrete measure space, \widehat{G} , of equivalence classes labelled by λ . For each λ , an element of the equivalence class is a matrix representation $\rho_\lambda(g)$ acting on a finite dimensional Hilbert space V_λ . One still refers to \widehat{G} as the dual group even though it will typically no longer be a group. In this setting the Fourier coefficient of $f(g) \in L^2(G)$ at representation λ is a matrix gotten by integrating the adjoint of the representation matrix for λ against $f(g)$ with respect to Haar measure on the group:

$$\hat{f}(\lambda) = \int_G f(g) \rho_\lambda(g)^* d\mu_{\text{Haar}}(g)$$

and the Plancherel formula is

$$f(g) = \sum_{\lambda \in \widehat{G}} \dim(V_\lambda) \langle \hat{f}(\lambda), \rho_\lambda(g)^* \rangle. \quad (1.2)$$

The contraction, $\langle \cdot, \cdot \rangle$ appearing here is defined by taking the trace of the product of the two matrices appearing in the pairing. The discrete measure, $\sum_{\lambda \in \widehat{G}} \dim(V_\lambda) \delta_\lambda$, is the measure on \widehat{G} , referred to as the Plancherel measure for G and $\dim(V_\lambda)$ is the *multiplicity* with which the representation λ appears as an isotypic (meaning equivalent) copy in the Plancherel formula. We also have the statement that this is an isometry,

$$\|f\|_{L^2(G)}^2 = \sum_{\lambda \in \widehat{G}} \dim(V_\lambda) \|\hat{f}(\lambda)\|^2.$$

The entry of Poisson geometry into this story may be seen as stemming from quantum mechanics. We will again say just a few words of motivation here but refer the reader to [GuSt84] for further details. A quantum system with many symmetries typically has a Hamiltonian, given as a differential operator, that commutes with these symmetries. In cases where this leads to a maximal commuting set of observables (operators) one can expect to have a complete description of the coherent states of the system in terms of the spectra of these operators acting on their common eigenfunctions. This decomposition parallels that of the Plancherel formulae just discussed. For instance, in the case of $G = SO(3)$ these spectra are the quantum numbers of the hydrogen atom [Si05]. The semiclassical limits of these quantum systems correspond to classical Hamiltonian dynamical systems in which the commuting operators correspond to Poisson commuting functions. In a series of papers from the '60s and '70s, Kirillov and Kostant [Ki62, Ko70] introduced the idea of reversing this process thereby bypassing most of the complications of quantum mechanics in favor of classical dynamical formulations on phase spaces built directly from the Lie algebra of the associated symmetry group of the system. In a number of cases they were able to show that the orbits of the co-adjoint action of G on the dual of its Lie algebra \mathfrak{g}^* are in one-to-one correspondence with the elements of \widehat{G} . (Physically these orbits are the level sets of energy Casimirs for the system.) In these descriptions the Liouville measure on \mathfrak{g}^* is related to the Plancherel measure on \widehat{G} . So it became of interest to extend this idea to more general groups which gave rise to the orbit method program mentioned at the outset. This has been an active area of research for many years now [Ki99].

As a point of departure we make an aside here to note that for S^1 the Lie algebra is $i\mathbb{R}$. Its dual Lie algebra is comprised of the linear functionals $2\pi i\gamma\theta$. The induced action of conjugation on these functionals, which is the co-adjoint action, is trivial since this group is commutative. So the co-adjoint orbits in this case are just the singleton linear functionals. However, due to the topological non-triviality of S^1 , a condition is required to ensure that the characters, which are the exponentials of these functionals, be well-defined on S^1 . This requirement is that $\gamma = n \in \mathbb{Z}$. So, though this case is a bit degenerate, it is true that the elements of the dual group, $\widehat{S^1}$, are in one-to-one correspondence with appropriately *quantized* co-adjoint orbits. This quantization condition is the last vestige of the quantum mechanical connection in the orbit method. However, for the non-compact groups that we now turn to, no quantization conditions are in fact required.

In this paper we will be concerned with the case of non-compact, non-commutative groups, such as B . This presents new challenges including the fact that irreducible representations will now typically come in continuous series (analogous to Fourier integrals for \mathbb{R}) so that the Plancherel measure is not simply discrete and may not even be, *prima facie*, well-defined. Moreover, the irreducible representations will for the most part be infinite

dimensional (and required to be unitary). Our particular interest is in Plancherel theorems for nilpotent and solvable groups. There has been progress here too, though only comparatively recently, thanks to the pioneering work of Moore, Wolf and their collaborators [MW73, LW78]. The result of [LW78] for the Plancherel formula of B takes the form stated in Theorem 3.4. This result contains a description of the dual group \widehat{B} in terms of parameters denoted \mathfrak{a}_\diamond^* with associated Plancherel measure as well as a new ingredient for which there is no classical analogue in (1.2). This new ingredient is a *Dixmier-Pukanzsky operator*, denoted D . It is necessitated because, unlike the compact groups or even nilpotent groups, B is not unimodular (i.e., its left and right Haar measures are not the same). We will review the derivation of this result and explain the meaning of all these terms in the first two sections of this paper. However, our goal is not simply to reproduce this result. Rather it is to understand the essential ingredients involved in a purely Poisson theoretic way that has clear significance for integrable systems theory. In particular we express \mathfrak{a}_\diamond^* explicitly in terms of the Casimirs that cut out the symplectic leaves of the Lie-Poisson structure associated to B . We also explicitly define the Dixmier-Pukanzsky operator for B in terms of the invariant theory associated to the maximal unipotent subgroup of B . To our knowledge this is the first time this particular kind of link between representation theory and Poisson geometry has been made in the literature. These results then yield some novel insights into the integrable systems structure of more general Toda lattices. This leads to the extension of a signature result from the classical tri-diagonal Toda lattice. We very briefly describe that now; the details are part of the primary content of this paper.

The Toda lattice [To67] is a dynamical system on \mathbb{R}^{2n} , with coordinates $(p_1, \dots, p_n, q_1, \dots, q_n)$. The system is Hamiltonian with respect to the standard symplectic structure on \mathbb{R}^{2n} with Hamiltonian

$$H(p_1, \dots, p_n, q_1, \dots, q_n) = \frac{1}{2} \sum_{j=1}^n p_j^2 + \sum_{j=1}^{n-1} e^{q_j - q_{j+1}}. \quad (1.3)$$

Under a remarkable coordinate transformation found by Flaschka [F174], (1.3) can be transformed to a Hamiltonian system with respect to the Lie-Poisson structure on the dual to the Lie algebra of B (see Section 4 for a precise description of this Poisson structure). In natural coordinates on the dual algebra, the phase space for this system can be taken to be tridiagonal Hessenberg matrices; i.e., matrices X of the form

$$X = \begin{pmatrix} a_1 & 1 & & & \\ b_1 & a_2 & \ddots & & \\ & \ddots & \ddots & \ddots & \\ & & & b_{n-1} & a_n \end{pmatrix}. \quad (1.4)$$

In these variables the Hamiltonian for the classical Toda lattice becomes

$$H = \frac{1}{2} \text{Tr} X^2.$$

This result is the signature result alluded to in the previous paragraph. There is a family of coadjoint orbits on this phase space parametrized by values of $\text{Tr} X$ which is an invariant of the coadjoint action (equivalently a Casimir of the Toda system). Corresponding to these orbits are infinite dimensional irreducible unitary representations of B related to the action of B on classical *Whittaker vectors*. This correspondence was worked out in detail by Kostant [Ko78] as a significant illustration of the power of the orbit method in providing explicit descriptions of representations of solvable Lie groups. However, these orbits are among the smallest dimensional coadjoint orbits. In this paper we initiate a similar analysis for the *largest dimensional* orbits. These orbits correspond to a set of full measure in the dual group of B which connects this analysis to a main theme of this paper concerning the Plancherel formula. Our approach may be understood as a kind of reversal of the approach taken by Flaschka. We will seek a coordinate transformation under which the full Kostant Toda lattice and its underlying Poisson geometry is expressed in terms of canonical coordinates with respect to a standard symplectic structure. We will prove the existence of such a transformation for the maximal dimensional (generic) orbits and some implications this has for the representation theory. We briefly describe some of those implications in the next paragraph.

The outline for this paper is as follows. In Section 2 we develop the necessary background on the construction of unitary irreducible representations for unipotent triangular matrices, in terms of a hierarchy of Heisenberg algebras. In Section 3 we develop the form of the Plancherel formula for B that anchors our work. This entails

developing some structures that make the Plancherel measure well-defined and get around a central problem related to the non-unimodularity of B . In particular this leads to the introduction of a Dixmier-Pukanszky operator. These two sections present material that already exists in the literature but which we summarize in some detail here in order to make our exposition self-contained. The approach we take there follows more recent treatments given in [Wo14, Wo16]. In Section 4 we present the essential results on the Poisson structure of the dual Lie algebra for B from [EFS93]. In Proposition 4.3 the results on semi-invariants are presented differently and in more detail than in [EFS93] as suits our needs here. We make use of more recent developments in invariant theory due to Kostant that appeared after publication of our previous work. We then derive, in Theorem 4.6 and Corollary 4.7, the completely Poisson-theoretic characterization of the Dixmier-Pukanszky operator associated with the Plancherel formula for B . Section 5 contains the remainder of our main results. In Theorem 5.5 we determine a particularly natural polarization for the symplectic structure on the orbits enabling us to interpret these orbits as cotangent bundles on a B -homogeneous space, $T^*(B/H)$ where H is the subgroup associated to the polarization. The symplectic structure on the orbit then gets identified with the canonical symplectic structure of this cotangent bundle by a theorem of Pukanszky. In Theorem 5.6 the polarization is used to establish the existence of canonical coordinates, mentioned above, that interpolate between the symplectic leaf structure described in Section 4 on the one hand, and the Heisenberg hierarchy outlined in Section 2 on the other. Then in Section 5.3 we relate all these Poisson perspectives back to the Plancherel formula for B stated in Section 3 and describe their significance. We will see that the Hilbert space for the representations corresponding to the generic orbits is naturally identified with $L^2(B/H)$. Finally, in Section 6 we outline some potential directions for future investigation that our work here may lead to. One of those directions is to return to the quantum systems with symmetries that gave rise to the orbit method in the first place. It will be of interest to consider quantum Toda lattices associated to the generic orbits. The results of this paper are a first step towards assessing whether or not these systems are quantum integrable.

2 Background

2.1 Decomposition of a Real Semisimple Lie Groups

In this section we will introduce the basic Lie group and algebra structures that we will be dealing with. We consider the Lie algebra decomposition of $n \times n$ matrices

$$\mathfrak{gl} = \mathfrak{gl}(n, \mathbb{R}) = \mathfrak{n}_- \oplus \mathfrak{b}_+$$

where \mathfrak{n}_- is the lower triangular nilpotent sub-algebra (nilradical) and \mathfrak{b}_+ is the complementary maximal solvable sub-algebra, referred to as a *Borel sub-algebra*:

$$\mathfrak{n}_- = \begin{pmatrix} 0 & & & & & \\ * & 0 & & & & \\ \vdots & \ddots & \ddots & \ddots & & \\ \vdots & & \ddots & \ddots & & \\ * & \dots & \dots & * & & 0 \end{pmatrix}, \quad \mathfrak{b}_+ = \begin{pmatrix} * & * & \dots & \dots & * \\ & * & * & & \vdots \\ & & \ddots & \ddots & \vdots \\ & & & \ddots & * \\ & & & & * \end{pmatrix}.$$

We will also use \mathfrak{n} to denote the transpose of \mathfrak{n}_- and \mathfrak{b}_- to denote the transpose of \mathfrak{b}_+ . Employing the *principal nilpotent* element,

$$\varepsilon = \begin{pmatrix} 0 & 1 & & & \\ & 0 & 1 & & \\ & & \ddots & \ddots & \\ & & & \ddots & 1 \\ & & & & 0 \end{pmatrix} \tag{2.1}$$

we introduce the affine translate,

$$\varepsilon + \mathfrak{b}_- = \begin{pmatrix} * & 1 & & & \\ * & * & 1 & & \\ \vdots & \ddots & \ddots & \ddots & \\ \vdots & & \ddots & \ddots & 1 \\ * & \dots & \dots & * & * \end{pmatrix},$$

to represent the dual algebra \mathfrak{b}_+^* with respect to the G -invariant, non-degenerate inner product (Killing form), $(X, Y) = \text{Tr}XY$, on \mathfrak{gl} . The affine space $\varepsilon + \mathfrak{b}_-$ is the space of all lower Hessenberg matrices which we'll denote by \mathcal{H} . It will be useful to introduce the following algebra and group projections,

$$\begin{aligned} \pi_- : \mathfrak{gl} &\rightarrow \mathfrak{n}_-, & \Pi_- : G &\rightarrow N_- \\ \pi_+ : \mathfrak{gl} &\rightarrow \mathfrak{b}_+, & \Pi_+ : G &\rightarrow B_+ \end{aligned}$$

where $G = GL(n, \mathbb{R})$, N_- , the lower unipotent matrices, is the exponential group of the algebra \mathfrak{n}_- and B_+ , the connected component of the identity in the invertible upper triangular matrices, is the exponential group of the algebra \mathfrak{b}_+ . Π_{\pm} are defined on the open dense subset of G where there is an LU factorization. We also let N denote the upper unipotent matrices.

2.2 Decomposition of Unipotent Triangular subgroups

We introduce a further decomposition of the nilpotent algebra \mathfrak{n} which will play a fundamental role in the remainder of this paper. Let $e_{i,j}$ denote the elementary matrix whose (i, j) -entry is 1 with all other entries being 0. Then we make the following definitions for $1 \leq r \leq \lfloor \frac{n}{2} \rfloor := R$.

$$\begin{aligned} \mathfrak{m}_r &:= \text{span}\{e_{r,j}, e_{i,n-r+1} | r < i, j < n - r + 1\} \cup \mathbb{R}e_{r,n-r+1} \\ \mathfrak{n}_r &:= \bigcup_{s \leq r} \mathfrak{m}_s. \end{aligned}$$

\mathfrak{m}_r is a sub-algebra of \mathfrak{n} isomorphic to the Heisenberg algebra of dimension $2(n - 2r) + 1$, for $0 < r \leq R$ and \mathfrak{n}_r is an ideal in \mathfrak{n} . The center of \mathfrak{m}_r is one-dimensional: $\mathfrak{z}_r = \mathbb{R} \cdot e_{r,n-r}$. Note that when n is even, $\mathfrak{m}_R = \mathfrak{z}_R$. We also define the following decompositions.

$$\mathfrak{m}_r = \mathfrak{z}_r + \mathfrak{v}_r \tag{2.2}$$

$$\mathfrak{n} = \mathfrak{s} + \mathfrak{v} \text{ where} \tag{2.3}$$

$$\mathfrak{s} = \bigoplus \mathfrak{z}_r \text{ and } \mathfrak{v} = \bigoplus \mathfrak{v}_r \tag{2.4}$$

$$d_r = \frac{1}{2} \dim(\mathfrak{m}_r/\mathfrak{z}_r) (= n - 2r) \implies \frac{1}{2} \dim(\mathfrak{n}/\mathfrak{s}) = d_1 + \dots + d_R \tag{2.5}$$

At the group level one has, correspondingly,

$$N_r = M_1 M_2 \cdots M_r \text{ is normal in } N \text{ and} \tag{2.6}$$

$$N_r = N_{r-1} \rtimes M_r \tag{2.7}$$

$$N = M_1 M_2 \cdots M_R \tag{2.8}$$

$$S = Z_1 Z_2 \cdots Z_R = Z_1 \times \cdots \times Z_R \text{ where } Z_r = \text{the center of } M_r. \tag{2.9}$$

2.3 Schrödinger Representations of Heisenberg Groups

The representation theory of Heisenberg groups such as M_r , corresponding to the Heisenberg algebra \mathfrak{m}_r is well-established and enshrined in the Stone-von Neumann theorem. We consider the irreducible unitary representation $\pi_{\lambda_r}, \lambda_r \in \mathfrak{z}_r^* \equiv \mathbb{R}$, of M_r , known as a *Schrödinger representation*, acting on the Hilbert space $L^2(\mathbb{R}^{d_r})$ and defined by

$$[\pi_{\lambda_r}(\vec{q}, \vec{p}, c)f](\xi) = e^{i\lambda_r(\vec{p} \cdot \xi + \frac{1}{2}\vec{p} \cdot \vec{q} + c)} f(\xi + \vec{q})$$

where \vec{p} is the coordinate vector associated to the basis elements $e_{r,r+1}, \dots, e_{r,r+d_r}$, \vec{q} is the coordinate vector associated to the basis elements $e_{r+1,n-r}, \dots, e_{r+d_r,n-r}$ and c is the coordinate associated to $e_{r,n-r+1}$. (Note that although these elements were introduced as associated to a basis for the algebra \mathfrak{m}_r , they also serve as coordinates on M_r since this is an exponential group.)

Theorem 2.1. [Fo89] *Excluding $\lambda_r = 0$, up to unitary equivalence, the π_{λ_r} are all the infinite dimensional irreducible unitary representations of M_r .*

From this, standard results on Plancherel's theorem for the Heisenberg group apply [MW73]. The Plancherel measure on the unitary dual \widehat{M}_r is supported on the π_{λ_r} with $\lambda_r \neq 0$ and given by $2^{d_r} d_r! |\lambda_r| d\lambda_r$. For $f \in L^1(M_r)$, define $\hat{f}(\lambda_r) = \int_{M_r} f(\vec{p}, \vec{q}, c) \pi_{\lambda_r}(\vec{p}, \vec{q}, c) d\vec{p} d\vec{q} dc$. The version of Plancherel's theorem here that corresponds to the isometry of the Fourier transform mentioned in Section 1 is given by

$$\|f\|_{L^2(M_r)}^2 = 2^{d_r} d_r! \int_{\mathfrak{3}^*} \|\hat{f}(\lambda_r)\|_{HS}^2 |\lambda_r|^{d_r} d\lambda_r$$

for $f \in L^1(M_r) \cap L^2(M_r)$ and $\|\cdot\|_{HS}$ denoting the Hilbert-Schmidt norm. The result corresponding to (1.2) is

$$f(x) = 2^{d_r} d_r! \int_{\mathfrak{3}^*} \Theta_{\pi_{\lambda_r}}(R_x f) |\lambda_r|^{d_r} d\lambda_r$$

where x is the group element corresponding to (\vec{p}, \vec{q}, c) , R_x denotes right translation acting on functions $((R_x f)(g) = f(gx))$ and $\Theta_{\pi_{\lambda_r}}$ is the distribution character of π_{λ_r} defined by

$$\Theta_{\pi_{\lambda_r}}(f) = \text{Tr } \pi_{\lambda_r}(f)$$

for $f \in C_c^\infty(M_r)$.

2.4 Stepwise Square Integrable Representations

The structure summarized in (2.2) - (2.9) sets one up to carry out a recursive construction of the irreducible unitary representations (referred to as *unirreps*) of N from the Schrödinger representations discussed in the previous subsection. The full details of this construction are carried out in the foundational paper of Moore and Wolf [MW73] and subsequent work nicely reviewed in [Wo16]. Here we will provide some background and then present those results needed for our work in this paper. For ease of comparison we, for the most part, adopt the notation used in [Wo16]. The basic inductive link for the recursion is (2.7) : $N_r = N_{r-1} \rtimes M_r$. In the following subdivisions of this subsection we will first review the mechanism of real polarizations by which one lifts an irreducible representation from N_{r-1} to N_r , then state the consequence of Mackey's *little group method* by which an irreducible unitary representation of N_r is realized by the Hilbert space tensor of the lifted representation with one of the Schrödinger representations of M_r described in Section 2.3. It is noted that this a set of full measure in \widehat{N}_r . Finally we present the explicit statement of Plancherel's formula for N .

2.4.1 Real Polarizations

Consider a Lie algebra \mathfrak{g} with connected Lie group G . In the matrix group case (which will be the case for us) G acts naturally on \mathfrak{g} by conjugation and we denote this *adjoint action* by Ad_g for $g \in G$ ($Ad_g(x) \doteq gxg^{-1} \doteq g \cdot x$ for $x \in \mathfrak{g}$). This action induces a corresponding *co-adjoint* action on linear functionals $\ell \in \mathfrak{g}^*$; viz., $Ad_g^*(\ell)(x) \doteq \ell(Ad_g(x)) \doteq g \cdot \ell$. Let G_ℓ denote the isotropy subgroup of G at ℓ under the co-adjoint action.

Definition 2.2. *Fix a linear functional ℓ on \mathfrak{g} . One says that a subalgebra $\mathfrak{k} \subset \mathfrak{g}$ is a real polarization at ℓ if*

1. $\{Ad_g \mathfrak{k} | g \in G_\ell\} = \mathfrak{k}$
2. $\ell([\mathfrak{k}, \mathfrak{k}]) = 0$
3. $2 \dim \mathfrak{k} = \dim \mathfrak{g} + \dim \mathfrak{g}_\ell$, where \mathfrak{g}_ℓ is the Lie algebra of G_ℓ .

In doing calculations with co-adjoint actions for Lie algebras, \mathfrak{g} , that are subalgebras of $\mathfrak{gl}(n, \mathbb{R})$, it is usually most convenient to again use the invariant inner product, $(X, Y) = \text{Tr}XY$ to represent \mathfrak{g}^* as we did in Section 2.1. As a first example consider $\mathfrak{g} = \mathfrak{m}_r$. Here we will represent \mathfrak{m}_r^* in terms of the *opposite* algebra \mathfrak{m}_r^\dagger , the transpose within $\mathfrak{gl}(n, \mathbb{R})$. We consider the case where ℓ is given by $\ell(X) = (e_{n-r+1, r}, X)$ for $X \in \mathfrak{m}_r$. Then the representation of the co-adjoint action on ℓ is given by $\pi_{\mathfrak{m}_r^\dagger} Ad_g^{-1}(e_{n-r+1, r})$ for $g \in M_r$ where $\pi_{\mathfrak{m}_r^\dagger}$ denotes projection into $\pi_{\mathfrak{m}_r^\dagger}$. A straightforward calculation shows that $(M_r)_\ell = \mathbb{I} + \mathfrak{z}_r$. We claim that a polarization at ℓ is given by $\mathfrak{k} = \mathfrak{z}_r \oplus \mathfrak{v}_r^+$ where $\mathfrak{v}_r^+ = \text{span}\{e_{r, j} | r < j < n - r + 1\}$. From Definition 2.2, condition (1) is obvious, The bracket, $[\mathfrak{k}, \mathfrak{k}] = 0$ and so condition (2) is immediate as well. Finally (3) holds since $\dim \mathfrak{k} = d_r + 1$ while $\dim \mathfrak{m}_r + \dim(\mathfrak{m}_r)_\ell = 2d_r + 1 + \dim \mathfrak{z}_r = 2d_r + 1 + 1 = 2 \dim \mathfrak{k}$. This establishes the claim. We note that in terms of the Schrödinger representations of M_r described in Section 2.3, \mathfrak{v}_r^+ is coordinatized by \vec{p} corresponding to the classical momenta of a free particle. Moreover, the Hilbert space for π_{λ_r} is $L^2(M_r/K) = L^2(\mathbb{R}^{d_r})$ where K is the connected group corresponding to \mathfrak{k} and \mathbb{R}^{d_r} is coordinatized by \vec{q} corresponding to the classical position variables of a free particle.

The construction of unirreps in \widehat{N} will be built on a ladder of analogous polarizations. We now describe the first rung of that ladder from which steps onto the higher rungs will be clear. So take $\mathfrak{g} = \mathfrak{n}_2$ and take $r = 1$ in the construction of the previous paragraph so that $\mathfrak{n}_1 = \mathfrak{m}_1$, λ_1 be the linear functional corresponding to $e_{n, 1}$ with \mathfrak{w}_1 denoting the polarization we found for this case. Extending it by zero, λ_1 is a linear functional on \mathfrak{n}_2 still represented by $e_{n, 1}$, but now regarded as an element of \mathfrak{n}_2^\dagger . To emphasize the distinction, this extension will be denoted by λ_1' . One may again calculate its co-adjoint isotropy to find that $(N_2)_{\lambda_1'} = \mathbb{I} + \mathfrak{z}_1 \oplus \mathfrak{m}_2$. Proceeding as before one may establish that $\mathfrak{w}_1' = \mathfrak{z}_1 \oplus \mathfrak{m}_2 \oplus \mathfrak{v}_1^+$ is a polarization at λ_1' . From this one sees that $N_2/K_1' = N_1/K_1$ and consequently $L^2(N_2/K_1') = L^2(N_1/K_1)$ which is the representation space of π_{λ_1} . It follows that this representation extends to a unirep, π_{λ_1}' in \widehat{N}_2 such that $d\pi_{\lambda_1}'$ vanishes on \mathfrak{z}_2 .

2.4.2 Constructing the Essential Part of \widehat{N}

The key step for going forward is Mackey's Little Group Method which applies in the setting of inducing representations from a normal subgroup. This applies to the setting of the previous subsection to yield, for N_2 , the following result.

Proposition 2.3. [Ma58] *Elements of \widehat{N}_2 whose restrictions to N_1 are multiples of π_{λ_1} are Hilbert space tensor products $\pi_{\lambda_1}' \widehat{\otimes} \gamma$ where $\gamma \in \widehat{M}_2 = \widehat{N}_2/N_1$.*

From Section 2.1 we know that the representations π_{λ_2} , associated to $\lambda_2 \in \mathfrak{z}_2^*$ corresponding to a non-zero multiple of $e_{n-r+1, r}$, constitute a set of full measure in \widehat{M}_2 . It follows from Proposition 2.3 that the representations $\pi_{\lambda_1}' \widehat{\otimes} \pi_{\lambda_2}$, which we will denote by $\pi_{\lambda_1 + \lambda_2}$, are inequivalent unirreps in \widehat{N}_2 . It is evident that this construction can be inductively continued to arrive at a family of unirreps on \widehat{N} determined by a sequence of linear functionals $0 \neq \lambda_r \in \mathfrak{z}_r^*$, $r = 1, \dots, R$ denoted by π_λ where $\lambda = \lambda_1 + \dots + \lambda_R$. The full description of properties related to Plancherel's formulae for N are then given by the following fundamental result of Moore and Wolf.

Theorem 2.4. [MW73] *Denote $\deg(\pi_\lambda) = \deg(\pi_{\lambda_1}) \dots \deg(\pi_{\lambda_R})$ with $\deg(\pi_{\lambda_r}) = |\lambda_r|^{d_r}$. Then the coefficients $f_{z, w}(x) = \langle z, \pi_\lambda(x)w \rangle$ of the irreducible unitary representation π_λ on N are in $L^2(N/S)$ and satisfy*

$$\|f_{z, w}\|_{L^2(N/S)}^2 = \frac{\|z\|^2 \|w\|^2}{\deg(\pi_\lambda)} = \frac{\|z\|^2 \|w\|^2}{\prod |\lambda_r|^{d_r}}.$$

Due to the recursive structure of the construction leading to these representations and the previous theorem, the π_λ are referred to as the *stepwise square integrable* representations of N relative to the decompositions (2.6) - (2.9).

Proposition 2.5. [Wo16] *Plancherel measure on \widehat{N} (to be specified in Section 2.4.3) is concentrated on the stepwise square integrable representations π_λ .*

2.4.3 The Plancherel Formula for N

To now fully state the Plancherel formula for N one would like to have a more explicit description of the distribution character for π_λ as a tempered distribution. We start at the level of the Heisenberg algebra \mathfrak{m}_r on

which one may regard λ_r as a linear functional whose kernel is the non-central elements of \mathfrak{m}_r . Consider the coadjoint orbit, $Ad^*(M_r)\lambda_r$, of λ_r in \mathfrak{m}_r^* . The distribution character, originally expressed as a trace, may be re-expressed as an integral over this orbit. One uses Lebesgue measure, $d\nu_r$ on $(\mathfrak{m}_r/\mathfrak{z}_r)^*$ normalized so that the Fourier transform is an isometry from $L^2(\mathfrak{m}_r/\mathfrak{z}_r)$ to $L^2(\mathfrak{m}_r/\mathfrak{z}_r)^*$. The tangent space to the orbit is an affine translate of a hyperplane in \mathfrak{m}_r^* and so $d\nu_r$ may be translated to a measure $d\nu_{\lambda_r}$ on the orbit. The orbital integral expressing the distribution character may then be written as

$$\Theta_{\pi_{\lambda_r}}(f) = c_r^{-1} |\lambda_r|^{-d_r} \int_{Ad^*(M_r)\lambda_r} \widehat{f}_1(\xi) d\nu_{\lambda_r}(\xi)$$

where $c_r = 2^{d_r} d_r!$, $f_1(\xi) = f(\exp(\xi))$ and \widehat{f}_1 is the Fourier transform with respect to the Lebesgue measure on the orbit. This all extends to N by defining

$$\mathfrak{t}^* = \{\lambda = \lambda_1 + \dots + \lambda_R \in \mathfrak{s}^*, 0 \neq \lambda_r \in \mathfrak{z}_r^* \forall r\} \quad (2.10)$$

$$\rho(\lambda) = \lambda_1^{d_1} \lambda_2^{d_2} \dots \lambda_R^{d_R} \quad (2.11)$$

$$c = c_1 c_2 \dots c_R = 2^{d_1 + \dots + d_R} d_1! d_2! \dots d_R! \quad (2.12)$$

$$d\nu_\lambda = d\nu_{\lambda_1} \times \dots \times d\nu_{\lambda_R} \quad (2.13)$$

$$\mathcal{O}(\lambda) = Ad^*(N)\lambda = Ad^*(M_1)\lambda_1 \times \dots \times Ad^*(M_R)\lambda_R. \quad (2.14)$$

It is straightforward to check that \mathfrak{t}^* is a cross-section for the generic (maximal dimensional) co-adjoint orbits of N . One then has

Theorem 2.6. [[Wol16](#)] *The distribution character of $\pi_\lambda \in \widehat{N}$ is*

$$\Theta_{\pi_\lambda}(f) = Tr \pi_\lambda(f) = \frac{1}{c} \frac{1}{|\rho(\lambda)|} \int_{\mathcal{O}(\lambda)} \widehat{f}_1(\xi) d\nu_\lambda(\xi) \quad (2.15)$$

and N has the Plancherel formula

$$f(x) = c \int_{\mathfrak{t}^*} \Theta_{\pi_\lambda}(R_x f) |\rho(\lambda)| d\lambda. \quad (2.16)$$

2.5 Poisson Geometry of Triangular Nilpotent Algebras

From the construction in the previous subsection it is natural to consider

$$b_{\lambda_r} := (x, y) \mapsto \lambda_r([x, y])$$

which are symplectic bilinear forms, nondegenerate on $\mathfrak{m}_r/\mathfrak{z}_r$, respectively. Each defines a Poisson bracket for functions on \mathfrak{m}_r^* referred to as the Lie-Poisson bracket and defined by $\{F, G\} = b_{\lambda_r}(\nabla F, \nabla G)$. With respect to this bracket the variables \vec{q}, \vec{p} appearing in the Schrödinger representations of Section 2.3 are canonical, corresponding to position and momentum. This all extends naturally to N as

$$b_\lambda := b_{\lambda_1} \oplus \dots \oplus b_{\lambda_R}, \quad (2.17)$$

a symplectic bilinear form, nondegenerate on $\mathfrak{n}/\mathfrak{s}$ determining a Poisson bracket for functions on \mathfrak{n}^* . The co-adjoint orbits, $\mathcal{O}(\lambda)$, appearing in (2.14) and (2.15) are the symplectic leaves for this Poisson structure, parametrized by \mathfrak{t}^* . (Functions on \mathfrak{s}^* are the Casimirs of this Poisson structure.) This perfectly realizes the promise of the Kirillov-Kostant orbit method. The equivalence classes of irreducible unitary representations on a set of full measure in \widehat{N} are in 1:1 correspondence with the generic co-adjoint orbits of N (equivalently the generic symplectic leaves of the associated Poisson bracket), as expressed in the Plancherel formula of Theorem 2.6. It is also of interest to note that $\rho(\lambda)$ introduced in the previous subsection equals the Pfaffian of b_λ .

3 Borel Subgroups

The restricted triangular group, N , considered in the previous section, is a connected unipotent subgroup of the Borel subgroup, B_+ of real upper triangular matrices in $GL(n, \mathbb{R})$. We will restrict attention to the connected

component of the identity in B_+ which is comprised of elements of B_+ whose diagonal entries are all positive. For simplicity of notation we will just denote this connected component by B .

An obstacle arises in trying to extend, to B , the inductive types of constructions used in deriving the Plancherel formula for N ; namely, whereas N is unimodular, B is not. A locally compact group, Y , is unimodular if its left invariant Haar measure equals its right invariant Haar measure; i.e., it has a bi-invariant measure. In the case of non-unimodular groups the two invariant measures are multiplicatively related by a function, δ_Y on the group, referred to as the modular function. Abelian groups are clearly unimodular. A group Y is unimodular if and only if $Y/Z(Y)$ is, where $Z(Y)$ is the center of Y . Thus a straightforward induction on the length of the central series for the n -step nilpotent group N establishes that N is unimodular. However, B is not and has a non-constant modular function that will be presented shortly.

3.1 Dixmier-Pukanszky Operator

Non-unimodularity poses an issue for the formulation of a Plancherel formula. Suppose relations such as those in Theorem 2.6 were to hold for a non-unimodular group. Then for $x = 1$, the identity element, the corresponding version of (2.16) would state that $f(1) = C \int_{\widehat{G}} \Theta_{\pi_\lambda}(f) d\mu_K$, where μ_K is either the left or right invariant measure. The left hand side is manifestly invariant under conjugation whereas, in general, conjugation transforms $\pi_\lambda(f)$, and hence (2.15), by the modular function. In the case of B (and certain other parabolic subgroups) this mismatch can be compensated by the introduction of a semi-invariant operator, the Dixmier-Pukanszky operator, into the Plancherel formula. This is thanks to the work carried out by Wolf and collaborators [LW78, Wo14], the result of which we'll now summarize.

Note that $B = AN$, where A is the split part of the maximal torus represented within in the diagonal matrices. The calculation of the modular function of B is expedited by using the fact that

$$\delta_B(an) = \alpha(a)\delta_A(a)\delta_N(n)$$

where $\alpha(a)$ is a positive function on A such that for all $f \in C_c(N)$, $\int_N f(Ad_a(n))d\mu_N = \alpha(a) \int_N f(n)d\mu_N$ and $d\mu_N$ is left Haar measure on N . Applying the change of variables formula one sees that $\alpha(a)$ is simply given by the inverse determinant of the Jacobian of the linear map $Ad_a|_N$. Since $\delta_A(a) = 1 = \delta_N(n)$,

$$\begin{aligned} \delta_B(an) &= (\det Ad_a|_N)^{-1} \\ &= \prod_{i=1}^n a_i^{2i-n-1} \end{aligned}$$

for $a = \text{diag}(a_1, \dots, a_n)$. For later use we write this in a form more naturally related to the root structure of \mathfrak{gl} ,

$$\begin{aligned} \delta_B(an) &= \prod_{r=1}^R \left(\frac{a_r}{a_{n-r+1}} \right)^{d_r+1} \\ &= \prod_{r=1}^R (\exp \beta_r(\log a))^{d_r+1} \end{aligned} \tag{3.1}$$

where $\{\beta_1, \dots, \beta_R\}$ are the roots $\beta_r = e_{r,r}^* - e_{n-r+1, n-r+1}^*$.

A Dixmier-Pukanszky operator is a positive, self-adjoint invertible differential operator acting on $L^2(B)$, defined in terms of the Fourier transform and semi-invariant under the adjoint action of B . To define this for the triangular case Wolf [Wo16] notes from (2.10) and (2.11) that the Pfaffian $\rho(\lambda)$ may naturally be regarded as a polynomial on \mathfrak{s}^* , non-vanishing on \mathfrak{t}^* . By duality one may also regard β_r as a linear functional on $\lambda \in \mathfrak{s}^*$ and then define $\text{Det}_{\mathfrak{s}^*}(\lambda) := \prod_r \beta_r(\lambda)$. Then one can confirm the following by direct calculation.

Proposition 3.1. *The product $\rho \cdot \text{Det}_{\mathfrak{s}^*}$ is an $\text{Ad}(B)$ -semi-invariant polynomial on \mathfrak{s}^* of degree $\frac{1}{2}(\dim \mathfrak{n} + \dim \mathfrak{s})$ and weight equal to the modular function δ_B .*

With $V = \exp(\mathfrak{v})$ one defines the differential operator D as the Fourier transform of this polynomial acting on $B = AN = AVS$ by acting on $S = \exp(\mathfrak{s})$:

$$\mathcal{F}(Df)(\lambda, w) = \rho(\lambda)\text{Det}_{\mathfrak{s}^*}(\lambda)\mathcal{F}(f)(\lambda, w) \tag{3.2}$$

$$\mathcal{F}(f)(\lambda, w) = \int_S f(z, w)e^{-i(\lambda, z)} dz \tag{3.3}$$

where z is a Euclidean coordinate on S and w is a coordinate on AV . Finally one has

Theorem 3.2. [Wo16] *D is an invertible self-adjoint differential operator of order $\frac{1}{2}(\dim \mathfrak{n} + \dim \mathfrak{s})$ on $L^2(B)$ with dense domain of Schwarz class on B and is $\text{Ad}(B)$ semi-invariant of weight equal to the modular function δ_B . So $|D|$ is a Dixmier-Pukanszky operator on B .*

There is an alternative formulation of this D that has a natural extension to more general minimal parabolic subalgebras developed in [LW78] and [Ko12]. We will present this in Section 4 and reveal its interpretation in terms of the Casimirs for the Lie-Poisson structure on \mathfrak{b}_+ .

3.2 Plancherel Theorem

One can now formulate the Plancherel theorem for B . Any $\lambda \in \mathfrak{t}^*$ has a non-zero projection on each summand \mathfrak{z}_r^* of \mathfrak{s}^* . The action by conjugation of $a \in A$ on \mathfrak{z}_r^* amounts to multiplication by $\exp(\beta_r(\log a))$. It follows that the isotropy algebra and subgroup of the coadjoint action of A on \mathfrak{t}^* are, respectively,

$$\mathfrak{a}_\diamond = \{\xi \in \mathfrak{a} \mid \beta_r(\xi) = 0, r = 1, \dots, R\} \quad (3.4)$$

$$A_\diamond = \exp(\mathfrak{a}_\diamond). \quad (3.5)$$

Note that these isotropies are the *same* at all $\lambda \in \mathfrak{t}^*$. The dual group \widehat{A}_\diamond is comprised of unitary characters, $\exp(i\phi(\log a))$ with $\phi \in \mathfrak{a}_\diamond^*$. Irreducible, unitary representations can then be constructed, as before, by Mackey's little group method.

$$\pi_{\lambda, \phi} = \text{Ind}_{NA_\diamond}^{NA} (\pi'_\lambda \otimes \exp(i\phi)); \quad \lambda \in \mathfrak{t}^*, \phi \in \mathfrak{a}_\diamond^* \quad (3.6)$$

where π'_λ is the natural extension of the stepwise square integrable representation π_λ from N to NA_\diamond . $\pi_{\lambda, \phi}$ and $\pi_{\lambda', \phi'}$ are equivalent if and only if $\lambda' \in \text{Ad}^*(A)\lambda$ and $\phi = \phi'$.

Proposition 3.3. [Wo16] *Plancherel measure for $B = AN = NA$ is concentrated on the set of equivalence classes of the irreducible representations $\{\pi_{\lambda, \phi}\}$ defined above.*

Theorem 3.4. [Wo16] *The distribution character, $\Theta_{\pi_{\lambda, \phi}}(f) : f \mapsto \text{Tr } \pi_{\lambda, \phi}(f)$ is a tempered distribution. If f is of Schwarz class on B , then*

$$f(x) = c \int_{\mathfrak{a}_\diamond^*} \int_{\mathfrak{t}^*} \Theta_{\pi_{\lambda, \phi}}(D(R_x f) |\rho(\lambda)| d\lambda d\phi.$$

We note that this realization of Plancherel's theorem reveals a relation between the Dixmier-Pukanszky operator D constructed in (3.2) and the Plancherel measure $d\mu = |\rho(\lambda)| d\lambda d\phi$. In general neither of these is unique by themselves; however, $D \otimes \mu$ is unique. The choices made here are natural in terms of the constructions used. The nature of D as a semi-invariant also suggests a connection to the universal enveloping algebra of B which further suggests a relation to the Lie-Poisson structure on \mathfrak{b}_+^* . This encourages one to expect a correspondence between \widehat{B} as a measure space and the symplectic leaf (co-adjoint orbit) foliation for that Poisson structure with its associated Liouville measure.

4 Poisson Structures, Orbits and Invariant Theory

In this section we develop the Poisson geometry that relates to the Plancherel theorem for B that was presented in the previous section. This section culminates with results that give a Poisson semi-invariant description of the Dixmier-Pukanszky and related operators. We will also point out how this description connects the perspectives of the representation theory community with those of the integrable systems community.

4.1 Poisson Reduction and Induction

As mentioned in Section 2.1, we represent the dual space of \mathfrak{b}_+ , \mathfrak{b}_+^* , with respect to the Killing form on \mathfrak{gl} , as $\varepsilon + \mathfrak{b}_-$ which is just the space of lower Hessenberg matrices, \mathcal{H} . From a Lie theoretic perspective, this space is

the translate of \mathfrak{b}_- by the sum of the simple positive root vectors, ε . In this representation of the dual space, the Lie-Poisson bracket may be concretely expressed, for functions f, g on \mathfrak{gl} , as

$$\{\tilde{f}, \tilde{g}\}(X) = (X, [\pi_+ \nabla f(X), \pi_+ \nabla g(X)]) \quad (4.1)$$

where $\tilde{f} = f|_{\mathcal{H}}$ and ∇ denotes the gradient with respect to the Killing form. Determining the symplectic geometry underlying this structure is intimately related to the existence of commuting families of functions (Hamiltonians) with respect to the bracket and Casimirs (functions that Poisson-commute with *all* other functions). A method due to Thimm [Th82] provides a method of constructing maximal commuting families of Hamiltonians (also known as *completely integrable Hamiltonian systems*), in our case, from Casimirs of subalgebras of \mathfrak{gl} that contain \mathfrak{b}_+ , referred to as *parabolic subalgebras*. The method proceeds by considering inclusion chains of such parabolic subalgebras and the corresponding chain of projections between their duals.

$$\begin{aligned} \mathfrak{b}_+ &= \mathfrak{p}_m \subset \cdots \subset \mathfrak{p}_k \subset \cdots \subset \mathfrak{p}_0 = \mathfrak{gl} \\ \mathfrak{b}_+^* &= \mathfrak{p}_m^* \leftarrow \cdots \leftarrow \mathfrak{p}_k^* \leftarrow \cdots \leftarrow \mathfrak{p}_0^* = \mathfrak{gl}^* \end{aligned} \quad (4.2)$$

The projections in (4.2) are *Poisson maps*, meaning that the pullback of the Lie-Poisson bracket of two functions on \mathfrak{p}_{k-1}^* equals the Lie-Poisson bracket of their pullbacks on \mathfrak{p}_k^* . We let P_k denote the parabolic group (which contains B_+) corresponding to \mathfrak{p}_k .

One can show that functions on \mathfrak{p}_{k-1}^* , invariant under the coadjoint action of P_{k-1} , Poisson commute with *all* functions on \mathfrak{p}_{k-1}^* ; i.e., they are Casimirs. Since the projections in (4.2) are all Poisson maps; one can collectively pull back all these sequentially invariant functions to \mathfrak{gl}^* to form a family of involutive (commuting) functions there. Then restricting this family to \mathfrak{b}_+^* gives an involutive family of functions on \mathcal{H} . For the details on all of this we refer the reader to [EFS93]. In that paper a particular chain of parabolics is chosen as well as a class of invariant functions on each parabolic yielding Casimirs. It is then shown that the resulting involutive family on \mathcal{H} is maximal. The chain of parabolics used are those which are symmetric with respect to the anti-diagonal of the matrices in $\mathfrak{gl} : \mathfrak{p}_r = [\mathfrak{g} \setminus \mathfrak{n}_r]^\dagger, r = 0, \dots, \lfloor \frac{n-1}{2} \rfloor$, where for a subalgebra $\mathfrak{h} \subset \mathfrak{gl}$, \mathfrak{h}^\dagger denotes the subalgebra which is the transpose of \mathfrak{h} . The corresponding parabolic subgroup, P_r , is the set of elements in G whose entries below the diagonal in the first r columns and to the left of the diagonal in the last r rows are zero. The maximal involutive family on $\varepsilon + \mathfrak{b}_-$ just described is realized in terms of the following construction. The essential ideas underlying this construction were due to [DLNT86] who used it to prove the complete integrability of the full symmetric Toda lattice and were adapted, in [EFS93], to establish this for the full Kostant Toda lattice. We reproduce the main details of the latter setting here since that will be fundamental for what we do.

Definition 4.1. For $r = 0, \dots, R$, and $X \in \mathfrak{gl}$, denote by $(X - \eta\mathbb{1})_{(r)}$ the result of removing the first r rows and last r columns of $(X - \eta\mathbb{1})$. We refer to this as the *r-chop* of X and denote the coefficients in η of its determinant as follows

$$\det(X - \eta\mathbb{1})_{(r)} = E_{0,r}\eta^{n-2r} + \cdots + E_{n-2r,r}. \quad (4.3)$$

To make the connection to invariants we require an intermediate notion. We will again use the Killing form to make an identification of the dual algebra \mathfrak{p}_r^* with a translate of \mathfrak{p}_{r-} , the transpose, in \mathfrak{gl} , of \mathfrak{p}_r ; specifically $\mathfrak{p}_r^* \simeq \tau_r + \mathfrak{p}_{r-}$ where $\tau_r = e_{1,2} + \cdots + e_{r,r+1} + e_{n-r,n-r+1} + \cdots + e_{n-1,n}$. Put differently, τ_r is the sum over the root vectors of the positive simple roots whose negatives do *not* belong to \mathfrak{p}_r . Note also that $\tau_R = \varepsilon$. For $X \in \tau_r + \mathfrak{p}_{r-}$, the co-adjoint action of P_r is realized by

$$Ad_p^* X = \tau_r + \pi_{\mathfrak{p}_{r-}} p^{-1} \cdot (X - \tau_r), \quad X \in \mathfrak{p}_{r-}, \quad p \in P_r \quad (4.4)$$

where $g \cdot Y = gYg^{-1}$.

Definition 4.2. We will say that a function f on $\tau_r + \mathfrak{p}_{r-}$ is a *co-adjoint semi-invariant* for P_r if

$$f(Ad_p^* X) = \chi(p)f(X) \quad \forall p \in P_r$$

for some character χ which we refer to as the *weight* of the semi-invariant. When χ is the trivial character we say that f is a *co-adjoint invariant*. Alternatively we may refer to it as a *parabolic invariant* or *parabolic Casimir*.

Proposition 4.3. *The $E_{m,r}(X)$ are all semi-invariants of the co-adjoint action of P_r on \mathfrak{p}_r^* , with weight*

$$\chi_r(p) = \frac{p_{1,1} \cdots p_{r,r}}{p_{n-r+1,n-r+1} \cdots p_{n,n}}. \quad (4.5)$$

- i) For $r = 0$, $\{E_{m,0}\}$ are invariants under conjugation by $GL(n, \mathbb{R})$ and generate the full Poisson-commutative algebra of Casimirs on \mathfrak{gl}^* . This is equivalent to the family generated by $\{TrX^m\}$ for $X \in \mathfrak{gl}$. Upon restriction to $\mathfrak{b}^*(X \in \varepsilon + \mathfrak{b}_- = \mathcal{H})$, these are no longer all Casimirs but they do constitute an involutive family. TrX^2 is the Hamiltonian for the famous Toda Lattice system—more precisely the Full Kostant Toda System [EFS93].
- ii) Since, for each fixed r , the weight is common, one has collections of P_r -Casimirs

$$I_{m,r} = E_{m,r}/E_{0,r}, \quad r = 1, \dots, \left\lfloor \frac{n-1}{2} \right\rfloor$$

that are rational functions of the coordinates on \mathfrak{p}_{r-} . Define the subset of generic matrices in \mathcal{H} to be

$$\mathcal{H}_{gen} = \left\{ X \in \varepsilon + \mathfrak{b}_- \mid E_{0,r}(X) \neq 0 \quad r = 1, \dots, \left\lfloor \frac{n-1}{2} \right\rfloor \right\}$$

This is a Zariski open subset of \mathcal{H} . Taken all together, $\{TrX^m, I_{m,r}; r = 1, \dots, \left\lfloor \frac{n-1}{2} \right\rfloor\}$ is a maximal involutive family for the Lie-Poisson structure on \mathcal{H}_{gen} (though the elements of this family are not all B -invariants). It is also a complete family of commuting constants of motion for the Full Kostant Toda flow, demonstrating the complete integrability of that system on generic co-adjoint orbits.

- iii) The leading coefficients $\{E_{0,r}\}$ are invariants for the co-adjoint action of N represented on \mathfrak{n}_- . Note that here we allow r to run from 1 to R and R may be larger than $\left\lfloor \frac{n-1}{2} \right\rfloor$.
- iv) The subfamily generated by $\{TrX, I_{1,r}\}$ is a complete family of Casimirs for the Lie-Poisson bracket on \mathcal{H} . As such, the generic level sets of these Casimirs are the maximal dimensional symplectic leaves of the bracket and also the principal co-adjoint orbits of B .

Proof. To establish that the $E_{m,r}(X)$ are indeed semi-invariants for P_r with weight (4.5) we begin by rewriting (4.3) in terms of a contraction on the exterior space $V_{n-r} = \bigwedge^{n-r} \mathbb{R}^n$. This is a fundamental representation space for $GL(n, \mathbb{R})$ given by

$$g(v_1 \wedge \cdots \wedge v_{n-r}) = gv_1 \wedge \cdots \wedge gv_{n-r}.$$

Since $GL(n, \mathbb{R})$ is a matrix subgroup of \mathfrak{gl} this definition extends to all elements of \mathfrak{gl} . By restriction this also applies to P_r, \mathfrak{p}_r and \mathfrak{p}_{r-} . With respect to the standard basis e_1, \dots, e_n of \mathbb{C}^n one defines a Hermitian inner product on V_n by

$$\langle e_{i_1} \wedge \cdots \wedge e_{i_n}, e_{j_1} \wedge \cdots \wedge e_{j_n} \rangle = \delta_{i_1, j_1} \cdots \delta_{i_n, j_n}.$$

Set $v^{(n-r)} = e_1 \wedge \cdots \wedge e_{n-r}$ and $v_{(n-r)} = e_{r+1} \wedge \cdots \wedge e_n$. These are, respectively, the highest and lowest weight vectors, with respect to lexicographic order, of this fundamental representation [FuHa91]. It is now straightforward to see that

$$\det(X - \eta\mathbb{I})_{(r)} = \langle (X - \eta\mathbb{I}) v^{(n-r)}, v_{(n-r)} \rangle. \quad (4.6)$$

Applying the co-adjoint action (4.4) to this representation one has

$$\langle (Ad_p^* X - \eta \mathbb{I}) v^{(n-r)}, v_{(n-r)} \rangle = \langle (\tau_r + (\pi_{\mathfrak{p}_r} p^{-1} \cdot (X - \tau_r)) - \eta \mathbb{I}) v^{(n-r)}, v_{(n-r)} \rangle \quad (4.7)$$

$$= \langle ((\pi_{\mathfrak{p}_r} p^{-1} \cdot (X - \tau_r)) - \eta \mathbb{I}) v^{(n-r)}, v_{(n-r)} \rangle \quad (4.8)$$

$$= \langle ((p^{-1} \cdot (X - \tau_r)) - \eta \mathbb{I}) v^{(n-r)}, v_{(n-r)} \rangle \quad (4.9)$$

$$= \langle p^{-1} \cdot (X - \tau_r - \eta \mathbb{I}) v^{(n-r)}, v_{(n-r)} \rangle \quad (4.10)$$

$$= \langle p^{-1} (X - \tau_r - \eta \mathbb{I}) p v^{(n-r)}, v_{(n-r)} \rangle \quad (4.11)$$

$$= \langle (X - \tau_r - \eta \mathbb{I}) v^{(n-r)}, (p^{-1})^\dagger v_{(n-r)} \rangle \quad (4.12)$$

$$= p_{1,1} \cdots p_{r,r} \det C_r \langle (X - \tau_r - \eta \mathbb{I}) v^{(n-r)}, (p^{-1})^\dagger v_{(n-r)} \rangle \quad (4.13)$$

$$= \frac{p_{1,1} \cdots p_{r,r} \det C_r}{p_{n-r+1, n-r+1} \cdots p_{n,n} \det C_r} \langle (X - \tau_r - \eta \mathbb{I}) v^{(n-r)}, v_{(n-r)} \rangle \quad (4.14)$$

$$= \chi_r(p) \langle (X - \eta \mathbb{I}) v^{(n-r)}, v_{(n-r)} \rangle. \quad (4.15)$$

where C_r denotes the Levi factor of p (the $(n-r) \times (n-r)$ central core of p). In passing from (4.7) to (4.8) we simply observed that τ_r is an element of \mathfrak{n}_r . All elements of \mathfrak{n}_r annihilate e_1 and so also $v^{(n-r)}$. The difference between (4.8) to (4.9) is again an element of \mathfrak{n}_r and so these two lines are also equal. In (4.10) the conjugation is simply passed through the identity and in (4.11) that conjugation is written out explicitly. In (4.12) p^{-1} is moved from the "bra" to the "ket" by transpose. The highest and lowest weight vectors are respectively stabilized by p and p^{-1} with multipliers as shown in (4.13) - (4.14) and which may be read of from the block form of P_r . Finally, as at the outset, τ_r may be removed to get (4.15). This shows that (4.3) is a semi-invariant with weight (4.5) for all η . Hence this is also the case for all $E_{m,r}(X)$.

For $r = 0$ (4.3) is just the characteristic polynomial for $X \in \mathfrak{gl}$ whose coefficients are clearly invariant under conjugation and generate the same algebra as $\{\text{Tr} X^m\}$. The rest of statement (i) follows from the Adler-Kostant-Symes Theorem [EFS93].

The statement in (ii) about P_r Casimirs is now immediate. The involutivity follows from Thimm's construction outlined earlier and the Adler-Kostant-Symes theorem. The maximality follows from the complete integrability on generic co-adjoint orbits of B for which we refer the reader again to [EFS93].

Statement (iii) is of importance for what comes in the remainder of this section, As a consequence of the form of the weight (4.5), the $E_{m,r}(X)$ depend only on the A -factor of $B = AN$ and so they are N -invariant on $\tau_r + \mathfrak{p}_{r-}$. In general they will not be invariant when restricted to \mathfrak{n}_- . However, since the unrestricted $E_{0,r}(X)$ only depend on coordinates in \mathfrak{n}_- the N -invariance persists for those coefficients.

In a similar way one may observe that $E_{1,r}(X)$ depends only on coordinates in \mathfrak{b}_- and so it continues to be a B -semi-invariant when restricted to $\mathcal{H}(= \varepsilon + \mathfrak{b}_-)$. The independence of the $\{\text{Tr} X, I_{1,r}\}$ on \mathcal{H}_{gen} holds as in (ii) and the completeness of this family of Casimirs asserted in Statement (iv) is a consequence of the polarization to be described in Theorem 5.5. \square

A key feature of these systems is that the invariants, $I_{m,r}$, are rational functions for $r > 0$. The pole locus, where the $I_{m,r}$ are undefined, corresponds to the locus in \mathcal{H} where the co-adjoint orbits are not principal and so the corresponding symplectic leaves are not generic and one would not expect them to contribute non-negligible support to the Plancherel measure on \widehat{B} . On the other hand these singularities could potentially pose problems for an orbit-theoretic formulation of Plancherel and its applications. As a first step toward addressing such issues we now bring in the invariant theory that plays a central role in the Lie theory and, particularly, in the representation theory discussed in Section 3.

4.2 Invariant Theory of N

We consider, for the moment, the general setting of a Lie algebra \mathfrak{g} and its associated (connected) Lie group G . There is a linear isomorphism

$$\sigma : \mathbb{S}(\mathfrak{g}) \rightarrow \mathbb{D}(G)$$

between $\mathbb{S}(\mathfrak{g})$, the symmetric algebra over \mathfrak{g} , and $\mathbb{D}(G)$, the space of all left-invariant differential operators, D , on G . There is also a canonical isomorphism between $\mathbb{S}(\mathfrak{g})$ and $\mathbb{S}(\mathfrak{g}^*)$, the latter being the space of polynomials on \mathfrak{g} . Then σ essentially amounts to the correspondence between the differential operator and its symbol. We refer to Appendix A for a precise description and details. Let $\mathbb{S}(\mathfrak{g}^*)^G$ denote the co-adjoint invariant polynomials which, under the isomorphism σ , corresponds to the bi-invariant differential operators on G . One of the main points in this article is that this isomorphism lies at the heart of the relation between the Plancherel measure and the Dixmier-Pukanszky operator in our setting.

We will now invoke a theorem of Kostant which describes these spaces of invariants in the cases of interest to us. To be more self-contained we will present this result with some detail and then apply it to the semi-invariant structures presented in Proposition 4.3. Kostant begins by letting \mathfrak{n}_- represent \mathfrak{n}^* with respect to the Killing form and considers the co-adjoint action of N which is represented as

$$\begin{aligned} N \times \mathfrak{n}_- &\rightarrow \mathfrak{n}_- \\ (n, \ell) &\mapsto \pi_{\mathfrak{n}_-}(Ad_{n^{-1}}\ell). \end{aligned}$$

One then restricts to the principal (maximal dimensional) orbits which comprise a Zariski open subset, X , of \mathfrak{n}_- (so $\overline{X} = \mathfrak{n}_-$). One may further observe that the dual of \mathfrak{t}^* , which we denote by \mathfrak{t}^\times , provides a cross-section of this orbit. So $X = \cup_{\lambda \in \mathfrak{t}^\times} \mathcal{O}(\lambda)$. Moreover, all the principal orbits, $\mathcal{O}(\lambda)$, are isomorphic to N/S . All this is consistent with what was described in (2.10) - (2.14). It is also straightforward to check that \mathfrak{t}^* is an orbit under the co-adjoint action of A and that, for $a \in A$, $a \cdot \mathcal{O}(\lambda) = \mathcal{O}(a \cdot \lambda)$. The co-adjoint action of $B = AN$ preserves \mathfrak{n}_- with the subgroup N acting along its orbits while A acts transversely as described in the previous sentence.

Theorem 4.4. [Ko12] *One has an isomorphism*

$$X \rightarrow N/S \times \mathfrak{t}^\times.$$

X is an affine variety; letting $\mathcal{A}(X)$ denote its affine ring, one has the following isomorphism of B -modules,

$$\mathcal{A}(X) \simeq \mathcal{A}(N/S) \otimes \mathcal{A}(\mathfrak{t}^\times).$$

It follows from this that $\mathcal{A}(X)^N \simeq \mathcal{A}(\mathfrak{t}^\times)$.

The last observation of this theorem implies that the restricted action of the split torus A on the module of N -invariants $\mathcal{A}(X)^N$ decomposes as an action by A -weights on $\mathcal{A}(\mathfrak{t}^\times)$. We will refer to the following weight filtration.

$$\begin{array}{lll} A\text{-weight lattice} & \Lambda & \subset \mathfrak{a}^* \\ & \cup & \\ \text{root lattice} & \Lambda_{ad} & \\ & \cup & \\ \text{cascade sublattice} & \Lambda_{\mathcal{B}} & = \oplus_{\beta \in \mathcal{B}} \mathbb{Z}\beta, \end{array}$$

where \mathcal{B} denotes the collection of roots $\{\beta_1, \dots, \beta_R\}$ defined in (3.1). It is referred to as a *cascade of strongly orthogonal positive roots* because it is an example of Kostant's cascade construction [Jo77, Ko12]. As a consequence of the fact that these roots are mutually orthogonal, the direct sum decomposition in the last line above follows. Thus, $\Lambda_{\mathcal{B}}$ is a free module of rank $R (= \lfloor \frac{n}{2} \rfloor)$, in which each weight appearing occurs with multiplicity 1. If M is a B -module let $\Lambda(M) \subset \Lambda$ be the set of A -weights occurring in the weight decomposition of M . Since $\mathcal{A}(X)$ is an affine algebra one has natural inclusions

$$\mathbb{S}(\mathfrak{n}) \subset \mathcal{A}(X) \subset \mathbb{Q}(\mathfrak{n})$$

where we note that $\mathbb{S}(\mathfrak{n}) = \mathcal{A}(\mathfrak{n}_-)$ and $\mathbb{Q}(\mathfrak{n})$ is the quotient field of $\mathbb{S}(\mathfrak{n})$. This induces inclusions on the N -invariants as A -modules.

$$\mathbb{S}(\mathfrak{n})^N \subset \mathcal{A}(X)^N \subset \mathbb{Q}(\mathfrak{n})^N.$$

But it then follows from Theorem 4.4 that

$$\Lambda(\mathbb{S}(\mathfrak{n})^N) \subset \Lambda(\mathcal{A}(\mathfrak{t}^\times)) \subset \Lambda(\mathbb{Q}(\mathfrak{n})^N).$$

One sees from an explicit calculation that $\Lambda(\mathcal{A}(\mathfrak{t}^\times)) = \Lambda_{\mathcal{B}}$ and so

$$\Lambda(\mathbb{S}(\mathfrak{n})^N) \subset \Lambda_{\mathcal{B}} \subset \Lambda(\mathbb{Q}(\mathfrak{n})^N).$$

Observe further that since $\mathbb{S}(\mathfrak{n})$ is isomorphic to a polynomial ring, it is a unique factorization domain and so any element $q \in \mathbb{Q}(\mathfrak{n})$ is uniquely expressible, up to a scalar multiple, as $q = f/g$ where f and g are relatively prime elements of $\mathbb{S}(\mathfrak{n})$. By this uniqueness, if q were an N invariant, the action of N would also need to preserve f and g respectively up to scalar multiples; i.e., up to respective characters. However, since N is unipotent any such character is trivial and so $f, g \in \mathbb{S}(\mathfrak{n})^N$. It follows that $\mathbb{Q}(\mathfrak{n})^N$ is the quotient field of $\mathbb{S}(\mathfrak{n})^N$ and by the last chain of inclusions one has

Theorem 4.5. [Ko12] *Every A -weight in $\Lambda(\mathbb{S}(\mathfrak{n})^N)$ occurs with multiplicity 1 in $\mathbb{S}(\mathfrak{n})^N$. Moreover, $\Lambda(\mathbb{S}(\mathfrak{n})^N) = \Lambda_{\mathcal{B}} \cap \Lambda_{\text{dom}}$ where Λ_{dom} is the set of dominant weights; i.e., those weights ψ such that $(\psi, \phi) \geq 0$ for all positive roots ϕ . Every weight $\gamma \in \mathbb{Q}(\mathfrak{n})^N$ occurs with multiplicity 1 in $\mathbb{Q}(\mathfrak{n})^N$ and has the form $\gamma = \nu - \mu$ for $\mu, \nu \in \Lambda(\mathbb{S}(\mathfrak{n})^N)$. Additionally, the prime factors of any element in $\mathbb{S}(\mathfrak{n})^N$ are also in $\mathbb{S}(\mathfrak{n})^N$.*

4.3 The Dixmier-Pukanszky Operator Revisited

In our case the generators in $\Lambda_{\mathcal{B}} \cap \Lambda_{\text{dom}}$ are of the form $\nu + \nu^*$ where ν is the highest weight of one of the fundamental representations and ν^* is the lowest weight or, alternatively, it may be characterized as the highest weight of the contragredient representation. $\nu + \nu^*$ is then a dominant weight for the representation $V_\nu \otimes V_\nu^*$. But now one sees that the coefficients $E_{m,r}(X)$ are matrix elements of such a representation; precisely that representation for which $V_\nu = \bigwedge^{n-r} \mathbb{C}^n$. From this perspective one sees that it is most natural to view the r -chop $(X - \eta\mathbb{I})_{(r)}$ appearing in (4.1) as inducing a family of elements in $\text{End } V_\nu = V_\nu \otimes V_\nu^*$. All of the $E_{m,r}(X)$ are elements of this space corresponding to the weight $\nu + \nu^* \in \Lambda_{\mathcal{B}} \cap \Lambda_{\text{dom}}$; however, as a consequence of Proposition 4.3(iii), *only one* of these is the multiplicity 1 representative of that weight in $\mathbb{S}(\mathfrak{n}^*)^N$, namely $E_{0,r}(X)$. This establishes the following theorem.

Theorem 4.6. *The invariant theory underlying the center of the enveloping algebras for \mathfrak{n} is explicitly described in terms of the r -chops as follows. $\mathbb{S}(\mathfrak{n}^*)^N \simeq \mathbb{R}[E_{0,1}(X), \dots, E_{0,R}(X)]$ where $E_{0,r}(X)$ is explicitly defined in Proposition 4.3 with $X \in \mathfrak{n}_-$. $E_{0,r}(X)$ is homogeneous of degree r .*

We now observe, from its modular weight (3.1), that the Dixmier-Pukanszky operator, is N -invariant; moreover, by Proposition 3.1 it depends only on \mathfrak{n}^* and so in fact is a bi-invariant differential operator in $\mathbb{D}(N)$. Thus, its symbol, under σ corresponds to an element of $\mathbb{S}(\mathfrak{n}^*)^N$. It follows from the theorem that this symbol has a unique expression in terms of the $E_{0,r}(X)$.

Corollary 4.7. *The symbol of the operator, D , appearing in the Dixmier-Pukanszky operator for the Plancherel Theorem (Theorem 3.4) has a unique expression in $\mathbb{S}(\mathfrak{n}^*)^N$ given by*

$$\begin{aligned} \sigma(D) &= \prod_{r=1}^R E_{0,r}^2(X) \quad \text{for } n \text{ odd} \\ &= E_{0,R}(X) \prod_{r=1}^{R-1} E_{0,r}^2(X) \quad \text{for } n \text{ even} \end{aligned}$$

and so is homogeneous of degree $R(R+1)$ for n odd and degree R^2 for n even. In either case, with $R = \lfloor n/2 \rfloor$ one sees that this degree is equal to $\frac{1}{2}(\dim \mathfrak{n} + \dim \mathfrak{s})$, coinciding with the order of the operator D stated in Theorem 3.2.

Proof. By the Theorem one knows that the symbol has a unique representation of the form

$$\sigma(D) = \prod_{r=1}^R E_{0,r}^{\alpha_r}(X) \quad \text{for unique } \alpha_r \in \mathbb{N}.$$

From (4.5) we know that the weight of $E_{0,r}$ is $\sum_{\ell=1}^r \beta_\ell$. Hence

$$\begin{aligned} \text{weight}(\sigma(D)) &= \sum_{r=1}^R \alpha_r \left(\sum_{\ell=1}^r \beta_\ell \right) \\ &= \sum_{\ell=1}^R \left(\sum_{r=\ell}^R \alpha_r \right) \beta_\ell \end{aligned}$$

where in the second line one has just interchanged the order of summation. On the other hand, by definition this weight is determined by (3.1),

$$\text{weight}(\sigma(D)) = \sum_{\ell=1}^R (d_\ell + 1) \beta_\ell.$$

Comparing these two expressions yields the system of equations

$$\sum_{r=\ell}^R \alpha_r = d_\ell + 1$$

which can be rewritten using (2.1), with n replaced by R , in Toeplitz form as

$$\begin{aligned} (\mathbb{I} + \varepsilon + \varepsilon^2 + \cdots + \varepsilon^R)(\alpha_1, \dots, \alpha_R)^\dagger &= (d_1 + 1, \dots, d_R + 1)^\dagger \\ (\mathbb{I} - \varepsilon)^{-1}(\alpha_1, \dots, \alpha_R)^\dagger &= (d_1 + 1, \dots, d_R + 1)^\dagger \\ (\alpha_1, \dots, \alpha_R)^\dagger &= (\mathbb{I} - \varepsilon)(d_1 + 1, \dots, d_R + 1)^\dagger \\ &= (2, \dots, 2)^\dagger \text{ for odd } n \\ &= (2, \dots, 2, 1)^\dagger \text{ for even } n \end{aligned}$$

which establishes the stated expressions for $\sigma(D)$. The evaluation of the homogeneous degree then follows from the respective $E_{0,r}$ having degree r . \square

4.3.1 Some History of Two Perspectives

As may be seen from what has been discussed up to this point, the semi-invariants $E_{0,r}$ have played a significant role, separately, for developments both in representation theory of Borel subgroups and in natural integrable systems on generic co-adjoint orbits of these Lie groups. However, to our knowledge, Corollary 4.7 represents the first time a connection between these two roles has been explicitly identified. To help highlight this we briefly mention some historical antecedents in the separate disciplines. On the side of integral systems theory, the first reference we are aware of is due to Arhangel'skii [Arh79] who essentially identifies $E_{0,r}$ and $E_{1,r}$ by a direct inspection of minors as semi-invariants of the B -coadjoint action and applies the translation method of Miscenko-Fomenko to generate an involutive family. In [Tro80], Trofimov extends this approach to determine the dimension of symplectic leaves for other simple Lie algebras. Deift, Li, Nanda and Tomei [DLNT86] independently studied the Lie-Poisson structure for symmetric matrices that is associated to a different splitting of \mathfrak{gl} . They introduce the chop semi-invariants in this symmetric case. This provides a more conceptual framework for generating semi-invariants and involutive families which can be related to the Ritz values of X . The goal set in [EFS93] was to place the conceptual framework offered by the r -chops in its broadest possible setting by formulating it Lie theoretically. This was carried out for the triangular group, as described earlier in this section, (with extensions indicated for other classical Borel subalgebras) in terms of the invariant theory of more general parabolic subgroups. This, in turn, was used to construct a toric linearization of the chop-invariant commuting flows on the associated flag manifolds of the semisimple Lie algebra. Gekhtman and Shapiro [GS99] later extended this approach to establish complete integrability of the generic Toda flows for simple Lie algebras, including the exceptional ones. This work makes fundamental use of Lie-theoretic structure related to maximal cascades of strongly orthogonal roots, mentioned earlier, as developed by Joseph and Kostant.

In representation theory, the semi-invariants $E_{0,r}$ appear first in Dixmier's early formulation [Dix59] of a Plancherel theorem for N . Then in the seminal paper [LW78], Lipsman and Wolf develop the central role of Dixmier-Pukanszky operators in more general Plancherel theorems. They relate this to a construction of D in terms of semi-invariants for "good" parabolics and, in fact, see these arising from matrix coefficients with

highest weights associated to cascade roots analogous to what was described just before Theorem 4.6 relating to the chop semi-invariants. Kostant developed a systematic analysis of invariants in the enveloping algebra of \mathfrak{n} as we outlined earlier and subsequently [Ko13] used this to fill a gap in the Lipsman-Wolf construction. This did not alter anything in the conclusions of [LW78] but it did make clearer to us the connection of that construction to our Poisson theoretic calculations in Proposition 4.3 and was the initial motivation for us to prove Theorem 4.6 and Corollary 4.7.

5 Polarization on Orbits and Canonical Structures

The connection between the symplectic geometry on the generic co-adjoint orbits of B and the corresponding unirreps is further illuminated by determining canonical coordinates on these orbits which, intrinsically, means that the orbits may be represented as T^*B/H , the cotangent bundle of a homogenous space for an appropriate subgroup H . The means for doing this is to first find a real polarization (Definition 2.2) with respect to the symplectic structure on each orbit. To do this in a uniform way with respect to the principal orbit foliation requires an additional condition which was formulated by Pukanszky [Pu71].

5.1 Pukanszky Polarization for Borels

Again making our usual dual identification for \mathfrak{b}^* , choose an element $f \in \mathcal{H} = \varepsilon + \mathfrak{b}_-$ on a generic orbit and let B_f denote its isotropy subgroup under the co-adjoint action.

Definition 5.1. *We say that a subalgebra $\mathfrak{h} \subset \mathfrak{b}$ is a real polarization at f satisfying Pukanszky's condition if*

1. $\{Ad_g \mathfrak{h} | g \in B_f\} = \mathfrak{h}$
2. $(f, [\mathfrak{h}, \mathfrak{h}]) = 0$
3. $2 \dim \mathfrak{h} = \dim \mathfrak{b} + \dim \mathfrak{b}_f$
4. $f + \mathfrak{h}^\perp \subseteq \mathcal{O}(f)$, where $\mathfrak{h}^\perp = \{\ell \in \mathfrak{b}^* | \ell(h) = 0, \forall h \in \mathfrak{h}\}$

(4) is Pukanszky's condition.

With such a real polarization in place one can then derive a canonical description of the co-adjoint orbits. This is done in terms of the standard moment mapping on T^*B , equipped with its canonical symplectic structure, that is associated to B acting symplectomorphically by right translation. (We refer to [GuSt84] for the background.) In terms of the trivialization of the cotangent bundle with respect to left invariant vector fields this takes the simple form

$$\begin{aligned} J_R : T^*B = B \times \mathfrak{b}^* &\rightarrow \mathfrak{b}^* \\ (b, \ell) &\rightarrow \ell. \end{aligned} \tag{5.1}$$

Assume that B_f is connected (which is the case for us). Then define H to be the connected Lie subgroup of B whose Lie algebra is \mathfrak{h} . H is a closed subgroup of B . We now also consider the moment map associated to the left action of H which, in the trivialization, takes the form

$$\begin{aligned} J_L : B \times \mathfrak{b}^* &\rightarrow \mathfrak{h}^* \\ (b, \ell) &\rightarrow (Ad_{b^{-1}}^* \ell)|_{\mathfrak{h}}, \end{aligned} \tag{5.2}$$

which denotes the restriction of the coadjoint translate of ℓ to \mathfrak{h} . One is then in a position to carry out symplectic reduction with respect to J_L , at a generic value f , yielding a reduced symplectic manifold

$$(J_L^{-1}(f)/H, \omega_f) \tag{5.3}$$

where ω_f is the reduced symplectic form. The moment map J_R restricts naturally to the fiber (5.3) to give a reduced map J_R^f to \mathfrak{b}^* . The existence and general form of canonical structures on the generic coadjoint orbits of B will follow from the next proposition. We refer to [BGR17] for general background and details on this fundamental result.

for \mathfrak{gl} that

$$[\mathfrak{h}, \mathfrak{h}] \subset \mathfrak{v}^+ \quad (5.9)$$

confirming that \mathfrak{h} is a subalgebra of \mathfrak{b} . Next we calculate the isotropy subgroup of f .

Lemma 5.4.

$$B_f = A_\diamond.$$

Proof. Noting that $B = AN = NA$ it will suffice to consider the actions by A and N separately. It is straightforward to check that

$$\begin{aligned} Ad_a^* f &= \varepsilon + \pi_{\mathfrak{b}_-} a^{-1} \cdot (f - \varepsilon) \quad \text{for } a \in A_\diamond, \\ &= \varepsilon + a^{-1} \cdot (f - \varepsilon) \\ &= \varepsilon + f - \varepsilon = f. \end{aligned}$$

Hence, $A_\diamond \subset B_f$. On the other hand the action of an a from the complement of A_\diamond will non-trivially scale the anti-diagonal entries of f . Finally, for $n \in N$, $Ad_n^* f$ preserves the diagonal entries of $f - \varepsilon$ since conjugation just spreads that element into \mathfrak{b} . On the other hand there will be an element of N preserving the anti-diagonal entries of $f - \varepsilon$ if and only if there is an element $\xi \in \mathfrak{n}$ whose bracket with the anti-diagonal part of $f - \varepsilon$ is in \mathfrak{n} . However, by direct inspection one sees that this bracket acts on ξ (within \mathfrak{gl}) by interchanging the top R rows of ξ with its bottom R rows and (with a sign change) the last R columns of ξ with its first R columns. In this way one sees that every entry of ξ gets moved to at least one separate single entry in \mathfrak{b}_- . Hence the bracket will be 0 only if $\xi = 0$. \square

We can now check that \mathfrak{h} satisfies the conditions of Definition 5.1 and deduce some corresponding consequences from Proposition 5.2.

Theorem 5.5. \mathfrak{h} given by (5.7) is a real polarization at f specified by (5.4). Furthermore,

$$\{Ad_h^* f | h \in H\} = f + \mathfrak{h}^\perp$$

which implies that Pukanszky's condition is satisfied.

Proof. By Lemma 5.4, condition 1 of Definition 5.1 becomes the requirement that $\{Ad_g \mathfrak{h} | g \in A_\diamond\} = \mathfrak{h}$. However, since A_\diamond contains the identity this is immediate. It is straightforward to see that $(f, \mathfrak{v}^+) = 0$. By (5.9) it follows that condition 2 is also satisfied. Next, we calculate dimensions,

$$\begin{aligned} \dim \mathfrak{b}_f &= \dim \mathfrak{a}_\diamond \\ 2 \dim \mathfrak{h} &= 2 \dim \mathfrak{a}_\diamond + 2 \dim \mathfrak{s} + 2 \dim \mathfrak{v}^+ \\ &= 2 \dim \mathfrak{a}_\diamond + 2 \dim \mathfrak{s} + \dim \mathfrak{n}/\mathfrak{s} \\ &= 2 \dim \mathfrak{a}_\diamond + \dim \mathfrak{s} + \dim \mathfrak{n} \\ 2 \dim \mathfrak{h} - \dim \mathfrak{b}_f &= \dim \mathfrak{a}_\diamond + \dim \mathfrak{s} + \dim \mathfrak{n} \\ &= n - R + R + \dim \mathfrak{n} \\ &= n + \frac{n(n-1)}{2} = \frac{n(n+1)}{2} \\ \dim \mathfrak{b} &= \frac{n(n+1)}{2}, \end{aligned}$$

showing that condition 3 is satisfied. Finally, we address the Pukanszky condition. Applying duality with respect to the Killing form on \mathfrak{gl} it is straightforward to see that $\mathfrak{h}^\perp = [(\mathfrak{a}_\diamond^\perp \cap \mathfrak{a}) \oplus \mathfrak{v}^-]^\dagger$. We write

$$f = \varepsilon + \kappa + Z$$

where $\kappa = \text{diag}(\kappa_1, \kappa_2, \dots, \kappa_2, \kappa_1)$ is the diagonal component of f and Z is the component whose entries on the anti-diagonal below the diagonal are all 1 with all other entries 0: $Z = e_{n,1} + e_{n-1,2} + \dots + e_{n-R+1,R}$. Recall once more that the co-adjoint action is given by

$$Ad_h^* f = \varepsilon + \pi_{\mathfrak{b}_-} h^{-1} \cdot (f - \varepsilon).$$

We may consider the action on the components separately. For κ , since the h act as raising operators, it is straightforward to see that $\pi_{\mathfrak{b}_-} h^{-1} \cdot \kappa = \kappa$. To evaluate the form of $\pi_{\mathfrak{b}_-} h^{-1} \cdot Z$ it suffices to consider its infinitesimal form, $\pi_{\mathfrak{b}_-} [\mathfrak{h}, Z]$; since we are dealing with exponential groups here, the exponential of the range of the infinitesimal action maps diffeomorphically onto the image of the co-adjoint action. By linearity, this reduces one to evaluating the bracket on basis elements in (5.5) or (5.6) for which one may appeal to the structure relations (5.8). We know from Lemma 5.4 that $[\mathfrak{a}_\diamond, Z] = 0$ which implies that the exponential of this action is the identity on Z . Thus it suffices to just consider $[\mathfrak{s}, Z]$ and $[\mathfrak{v}^+, Z]$. In the first instance we evaluate

$$\begin{aligned} & [e_{r,n-r+1}, e_{n,1} + e_{n-1,2} + \cdots + e_{n-R+1,R}] \quad \text{for } 1 \leq r \leq R \\ = & e_{r,r} - e_{n-r+1,n-r+1} \quad 1 \leq r \leq R \end{aligned}$$

which spans $(\mathfrak{a}_\diamond^\perp \cap \mathfrak{a})$. In the second instance we consider

$$\begin{aligned} & \pi_{\mathfrak{b}_-} [e_{i,j}, e_{n,1} + e_{n-1,2} + \cdots + e_{n-R+1,R}] \quad \text{for } i < j < n - i + 1 \\ = & \pi_{\mathfrak{b}_-} (e_{i,n-j+1} - e_{n-i+1,j}) \quad i < j < n - i + 1 \\ = & -e_{n-i+1,j} \quad i < j < n - i + 1. \end{aligned}$$

which spans $(\mathfrak{v}^-)^\dagger$. Assembling all these observations we see that

$$\begin{aligned} \{Ad_h^* f | h \in H\} &= \varepsilon + \pi_{\mathfrak{b}_-} (\kappa + Z \oplus (\mathfrak{a}_\diamond^\perp \cap \mathfrak{a}) \oplus (\mathfrak{v}^-)^\dagger) \\ &= \varepsilon + \kappa + Z \oplus (\mathfrak{a}_\diamond^\perp \cap \mathfrak{a}) \oplus (\mathfrak{v}^-)^\dagger \\ &= f + \mathfrak{h}^\perp. \end{aligned}$$

This establishes the Theorem. □

5.2 Canonical Representation for the Poisson Structure of \mathfrak{b}^*

The real polarization just described now lends itself to a global coordinate description of \mathfrak{b}^* as a canonical phase space consistent with the Heisenberg algebras underlying the construction of the stepwise square integrable representations presented in Section 2.4. We can now give another presentation of the fibration of

$$\mathcal{X} = \{X \in \varepsilon + \mathfrak{b}_- \mid E_{0,r}(X) \neq 0, \forall r\}$$

as an explicit product. We define

$$\mathcal{F} = \{f \in \varepsilon + \mathfrak{b}_- \mid \text{such that } f \text{ is of the form (5.4)}\}.$$

We know from Proposition 4.3 (iv) that \mathcal{X} is foliated by maximal symplectic leaves, B/B_f and by Lemma 5.4 each of these is diffeomorphic to B/A_\diamond .

Theorem 5.6. \mathcal{F} is a cross-section of the symplectic leaf foliation and thus the map $\mathcal{X} \rightarrow \mathcal{F}$ which sends an element of \mathcal{X} to its unique representative in \mathcal{F} , yield the diffeomorphism

$$\mathcal{X} \rightarrow B/A_\diamond \times \mathcal{F}. \tag{5.10}$$

a) The reduced symplectic manifold (5.3) at f is explicitly realized by noting that

$$\begin{aligned} J_L^{-1}(f) &= \{(b, \ell) \in B \times (\varepsilon + \mathfrak{b}_-) \mid Ad_{b^{-1}} \ell \in f + \mathfrak{h}^\perp\} \\ &= \{(b, \ell) \in B \times (\varepsilon + \mathfrak{b}_-) \mid Ad_{b^{-1}} \ell \in f + [(\mathfrak{a}_\diamond^\perp \cap \mathfrak{a}) \oplus \mathfrak{v}^-]^\dagger\} \end{aligned}$$

where $\mathfrak{v}^- = \text{span}\{e_{i,j} \mid n - j + 1 < i < j\}$. Since f is a regular value, H acts freely and properly on this fiber, the quotient $J_L^{-1}(f)/H$ is a smooth manifold and ω_f is the unique symplectic form on this quotient whose pull-back to $J_L^{-1}(f)$ equals the restriction of the canonical form on T^*B to $J_L^{-1}(f)$.

b) The tangent space to $B/B_f (= B/A_\circ)$ at f splits as

$$T_f(B/B_f) = [(\mathfrak{a}_\circ^\perp \cap \mathfrak{a}) \oplus \mathfrak{v}^-] \bigoplus [\mathfrak{s} \oplus \mathfrak{v}^+].$$

Under the exponential map the first factor maps locally diffeomorphically into B/H while the second factor is diffeomorphic to H/B_f .

c) \mathcal{X} has a natural coordinatization of the form

$$\left(\begin{array}{cccccccc} a_1 + q_{0,1} & 1 & & & & & & \\ p_{1,1} & a_2 + q_{0,2} & 1 & & & & & \\ p_{2,1} & p_{1,2} & a_3 + q_{0,3} & 1 & & & & \\ p_{3,1} & p_{2,2} & p_{1,3} & a_4 & 1 & & & \\ p_{4,1} & p_{3,2} & p_{0,3} & q_{1,3} & a_3 - q_{0,3} & 1 & & \\ p_{5,1} & p_{0,2} & q_{1,2} & q_{2,2} & q_{3,2} & a_2 - q_{0,2} & 1 & \\ p_{0,1} & q_{1,1} & q_{2,1} & q_{3,1} & q_{4,1} & q_{5,1} & a_1 - q_{0,1} & \end{array} \right) \quad (5.11)$$

in which the $q_{i,j}$ may be taken as local coordinates on B/H near f and the $p_{i,j}$ may be taken as local coordinates on the cotangent fiber to B/H at f . There are, however, global constraints between these coordinates coming from the Casimirs on each orbit.

Proof. The map from \mathcal{X} to \mathcal{F} in (5.10) is explicitly determined by the relations [Ra20],

$$\kappa_1 + \cdots + \kappa_r = \frac{\text{Tr}X - I_{1,r}(X)}{2}$$

which, given the fixed values of the Casimirs, $I_{1,r}(X)$ and $\text{Tr}(X)$, on the orbit, determine the diagonal entries $\vec{\kappa}$ of f in (5.4). These Casimirs are independent on \mathcal{X} and there are $\lfloor \frac{n+1}{2} \rfloor$ of them. Note also that $\vec{\kappa}$ runs from κ_1 to $\kappa_{\lfloor \frac{n+1}{2} \rfloor}$ and that $\dim \mathfrak{a}_\circ = n - R = \lfloor \frac{n+1}{2} \rfloor$. It follows from Lemma 5.4 that \mathfrak{a}_\circ parametrizes the generic symplectic leaves. This is consistent with f providing a cross-section of the coadjoint action and also confirms the statement in Proposition 4.3 (iv) that the $\{\text{Tr}X, I_{1,r}\}$ are a maximal independent set of Casimirs for \mathcal{X} . Again by Lemma 5.4 each B/B_f is diffeomorphic to B/A_\circ , which also gives the projection onto the first factor in (5.10).

In part (a) of the theorem, for the initial equality, we first note that in (5.2),

$$T^*B = B \times \mathfrak{b}^* = B \times (\varepsilon + \mathfrak{h}),$$

so that the moment map may be re-expressed, using duality, as

$$(Ad_{b^{-1}}^* \ell)|_{\mathfrak{h}} = \pi_{\mathfrak{h}^\dagger} (Ad_{b^{-1}}(\ell - \varepsilon)) + \varepsilon,$$

where ℓ on the RHS denotes the element in $\varepsilon + \mathfrak{b}_-$ dual, with respect to the Killing form, to the linear functional ℓ on the LHS. The projection $\pi_{\mathfrak{h}^\dagger}$ is projection onto the transpose of \mathfrak{h} which is contained in \mathfrak{b}_- . From our arguments in the proof of Theorem 5.5 one sees that the kernel of $\pi_{\mathfrak{h}^\dagger}$ within \mathfrak{b}_- is precisely \mathfrak{h}^\perp . Hence the inverse image of f under the moment map J_L is given by the RHS of the first equality of (a). The second equality just uses the explicit determination of \mathfrak{h}^\perp found previously in the proof of Theorem 5.5. The remaining statements in (a) just follow from standard facts about symplectic reduction.

Part (b) summarizes the Lie algebraic calculations made in Section 5.1. The coordinatization described in part (c) follows from applying the exponential map locally to (b). \square

Remark 5.7. It should be noted that the coordinatization in (5.11) reflects the n -step Heisenberg structure of the nilradical N contained in B .

5.3 Plancherel's Formula Revisited

We recall from Theorem 3.4, in the statement of Plancherel's formula for B , that for $f(x)$ a Schwarz class function on B one has

$$f(x) = c \int_{\mathfrak{a}_\diamond^*} \int_{\mathfrak{t}^*} \Theta_{\pi_\lambda, \phi}(D(R_x f)) |P(\lambda)| d\lambda d\phi.$$

Also, from (3.6) we know that $\pi_{\lambda, \phi}$ and $\pi_{\lambda', \phi'}$ are equivalent if and only if $\lambda' \in \text{Ad}^*(A)\lambda$ and $\phi = \phi'$. Using the Killing form on \mathfrak{gl} we may identify elements $\lambda \in \mathfrak{t}^*$ with anti-diagonal elements below the diagonal, none of whose entries vanish. As noted in the proof of Lemma 5.4, conjugation by elements a in the complement of A_\diamond will non-trivially scale the anti-diagonal entries of the element representing λ . In fact given any other such λ' there is an a that will move λ to λ' . It follows that for fixed $\phi \in \mathfrak{a}_\diamond^*$ all $\pi_{\lambda, \phi}$ are equivalent and so represent just one unirep equivalence class in \widehat{B} . Their distribution characters appear in the Plancherel formula as isotypic summands in an integral direct sum, which one may regard as a representation of their multiplicity in the Plancherel measure. On the other hand the $\pi_{\lambda, \phi}$ are inequivalent for different ϕ . It follows from Proposition 3.3 that these parametrize a set of distinct equivalence classes comprising a set of full measure in \widehat{B} .

We may also view this from the Poisson perspective. From Theorem 5.6 we know that the principal co-adjoint orbits in \mathfrak{b}^* are parametrized by the diagonal parameters $\vec{\kappa} = (\kappa_1, \dots, \kappa_{n-R})$ appearing in f (5.4). However, under duality $\vec{\kappa}$ corresponds 1:1 with \mathfrak{a}_\diamond^* . So we have

Theorem 5.8. *The just described correspondence between an open dense subset of \widehat{B} and generic co-adjoint orbits in \mathfrak{b}^* explicitly verifies the validity of the orbit method that was abstractly established by Auslander and Kostant [AK71]. Moreover, in direct relation to Proposition 5.2 (2) we may identify the Hilbert space for the representation $\pi_{\lambda, \phi}$ with $L^2(B/H)$. Also the symbol of the Dixmier-Pukanszky operator, D appearing in the Plancherel formula is explicitly given by Corollary 4.7.*

6 Conclusions

In [AK71] Auslander and Kostant established, on general grounds, that for type I solvable Lie groups there is a one-to-one correspondence between equivalence classes of its unireps, the dual group, and its co-adjoint orbits. Kostant [Ko78] went on to make this explicit in the special case of the so-called Whittaker representations. These correspond to the *smallest* indecomposable co-adjoint orbits which are the tridiagonal (Jacobi) orbits in the case of triangular groups. That work opened the door for deep connections between representation theory and integrable systems theory related to the classical Toda lattices. More recently this has led to important relations with probability theory [BBO05, O13], number theory [Sh10] and combinatorics [BBF11].

The results in this paper lay the groundwork for the kind of explicit description that Kostant carried out but for generic elements of the dual group, those corresponding to the *largest* dimensional co-adjoint orbits. Specifically we have made explicit connections between all the elements of the Plancherel formulae for triangular groups, B , as developed in the works of [LW78, Wo14] on the one hand and the Poisson geometry of co-adjoint orbits on \mathfrak{b}^* as developed within the framework of the full Kostant Toda lattice. This has been done in a way that will make it natural to extend our results to the Borel subgroups of general complex semi-simple Lie groups. This now positions one to explore deeper connections with integrable systems and allied fields similar to what has been seen for the Whittaker-classical Toda case. We expect this will be pursued in future investigations, but for now we will just take some space below to point out how the work in this paper may specifically contribute to those investigations.

6.1 Canonical Coordinates and Standard Hamiltonian Systems

As with the construction of polarizations and canonical structures for generic orbits described in Section 5, one may carry out an analogous construction for the minimal orbits of tridiagonal form. We refer the reader to [BGR17] for a detailed description of this. Under the corresponding coordinate transformation the tridiagonal Toda system determined by (1.4) with Hamiltonian $Tr X^2$ takes the form (1.3) in terms of *standard* canonical variables $\{p_1, \dots, p_n; q_1, \dots, q_n\}$. By "standard" here one means that in these variables the Poisson bracket "diagonalizes" as $\{p_i, p_j\} = 0 = \{q_i, q_j\}, \{p_i, q_j\} = \delta_{i,j}$.

As a consequence of Theorem 5.5, Proposition 5.2 implies that such coordinates exist in the case of generic orbits; however, it is not immediately clear what the explicit form of the coordinate transformation is or what form the Toda Hamiltonian, or any of the other Hamiltonians that commute with it, will take. However, in [Sy80] Symes outlines a general method for constructing the transformation based on the polarization. Moreover, the stratified Heisenberg structure evident in (5.11) suggests that the canonical coordinates we seek may have a close relation to canonical Heisenberg variables. So this is a promising beginning that will be pursued in a future study.

6.2 Quantization of Invariants and Quantum Integrability

As mentioned in the introduction, the orbit method had its origins in the quantum mechanics of systems with many symmetries. Having worked out the details of this method in the case of triangular groups it is natural to go back and ask what implications this might have concerning associated quantum systems and in particular the quantum analogue of the Full Kostant Toda lattice. In the case of the tridiagonal orbits this has been done for the original Toda lattice. In this case the quantum Toda lattice takes the form

$$\widehat{H} = -\frac{\hbar^2}{2} \sum_{j=1}^n \frac{\partial^2}{\partial q_j^2} + \sum_{j=1}^{n-1} e^{q_j - q_{j+1}} \quad (6.1)$$

following the formal prescription of canonical quantization which replaces p_j by $i\hbar\partial/\partial q_j$. A number of obstacles arise in trying to place this formal procedure on a rigorous footing. The first potential issue is that the choice of quantization is in general not unique; there may be several choices of operator whose semi-classical limit leads to the same H . (See [Fo89] for further details on this issue.) However, in the case of (1.3) the evident separation of variables obviates any such ambiguity. The relation between H and (6.1) is consistent with the symbol map as described in Appendix A. The next potential obstacle concerns the extension of classical integrability, and in particular Poisson commutativity, to the quantum level. For the tridiagonal case we may consider the Poisson commuting invariants given by $\{E_{m,0}(\vec{p}, \vec{q})\}$ in Proposition 4.3 (i). These are the coefficients of $\det(X - \eta\mathbb{I})$ for X of the form (1.4) when $a_i = p_i$ and $b_i = \exp q_i - q_{i+1}$. We note that when restricted to this tridiagonal phase space all the other semi-invariants, $\{E_{m,r}\}$ for $r > 0$, that were mentioned in Proposition 4.3 vanish identically. However, the invariants for $r = 0$ suffice for the complete integrability of tridiagonal Toda. Once again, remarkably, the quantization of these invariants, denoted $\widehat{E}_{m,0}$ are uniquely determined, gotten again by simply replacing the p_j by $i\hbar\partial/\partial q_j$ [Gi97]. However, Poisson commutativity does not necessarily imply that these associated operators commute since multiplication of symbols does not pass, mutatis mutandis, over to composition of operators. The only thing one may conclude, without further examination, is that commutators vanish to order \hbar^2 [Fo89]. Nevertheless one *does* find that $[\widehat{H}, \widehat{E}_{m,0}] = 0$ for all m , meaning that these operators play the role of quantum conservation laws. They are also referred to as a *complete set of commuting observables*, and they all have common eigenstates. This is what is meant by quantum integrability. The role of the common eigenstates is what is played by the Whittaker functions alluded to at the start of these Conclusions.

In trying to extend quantization and related matters to the setting of the full Kostant Toda lattice all the above mentioned obstacles rear their heads again. Since the Hamiltonian and related invariants will be evaluated on generic orbits rather than just tridiagonal ones (and will now involve all the $\{E_{m,r}\}$) what made things work out in the tridiagonal case may no longer be available. The passage to canonical variables based on our results here and envisioned in Section 6.1 may be of decisive help, but that remains to be determined. There is one further potential obstacle in this extension to quantum full Toda, for which the results of this paper may help. As we saw, the invariants for $r > 0$ are necessarily rational functions whose associated quantized operators should be given by pseudo-differential operators in terms of an appropriate Fourier transform. To formulate this we can make use of the Plancherel formula for B (Theorem 3.4) and define, using the invariance of $I_{m,r}$,

$$\widehat{I}_{m,r}f(x) = c \int_{\mathfrak{a}_\diamond^*} \int_{\mathfrak{t}^*} \Theta_{\pi_\lambda, \phi}(D\widehat{I}_{m,r}(R_x f)|\rho(\lambda)|d\lambda d\phi$$

where $D\widehat{I}_{m,r}$ is a *rational* semi-invariant operator with the same modular weight as D and defined in the same manner as D was in (3.2) - (3.3) with symbol $\sigma(D\widehat{I}_{m,r})$. (We note that the linear bijection (A.1) extends to the function fields of the enveloping algebras involved.) The concern here is that the pseudo-differential representations of the quantizations of $I_{m,r}$ might involve singular integrals. By the results of this paper, and in particular Corollary 4.7, one knows that

$$\begin{aligned}\sigma\left(\widehat{I}_{m,r}\right) &= \frac{E_{m,r}}{E_{0,r}} \\ \sigma(D) &= E_{0,R}^p(X) \prod_{r=1}^{R-1} E_{0,r}^2(X)\end{aligned}$$

where $p = 1$ for n even and $p = 2$ for n odd. If (A.1) were an algebra homomorphism, then the symbol of the product would be the product of the symbols and in our case it is evident that this product of the symbols is a polynomial and therefore non-singular. However σ is only a linear bijection and the symbol of the product only equals the product of the symbols modulo terms of lower order. However, restricted to *invariants* it is known that a modification of σ does realize an algebra isomorphism [Du77]. Whether or not something like this could be extended to *semi-invariants* of the type we consider here is currently under consideration. That is something that could also help with the issues raised earlier in this Section.

Acknowledgements

I wish to take this opportunity to express my deep and abiding gratitude to Hermann Flaschka for introducing me, many years ago, to the fascinating connections between Lie theory and Poisson geometry, as revealed by the Toda lattice. Hermann himself viewed the Toda lattice as a “vehicle” for exploring many fascinating areas of mathematics, which often had unexpected and beautiful relations with this nonlinear system. The work presented in this paper, motivated by our earlier study [EFS93] with Stephanie Singer, had its inception in extended explorations that Hermann and I undertook starting from early 2019 up to the time of his sudden passing in early 2021. I hope he would have liked this article, and that the ideas discussed here might help continue the mathematical legacy Hermann set in motion with his early seminal work on the Toda lattice.

A Appendix: Enveloping Algebras

We recall some basic definitions and facts. Associated to any Lie algebra, $(\mathfrak{g}, [\cdot, \cdot])$ one has its *universal enveloping algebra* $\mathfrak{U}(\mathfrak{g})$ defined as the associative algebra

$$\mathfrak{U}(\mathfrak{g}) = T(\mathfrak{g})/I$$

where $T(\mathfrak{g})$ is the tensor algebra of \mathfrak{g} and I is the two-sided ideal generated by $\{X \otimes Y - Y \otimes X - [X, Y]\}$ for $X, Y \in \mathfrak{g}$. \mathfrak{g} is naturally included in $\mathfrak{U}(\mathfrak{g})$ through $T^1(\mathfrak{g})$. This has an analytical interpretation in terms of left-invariant differential operators. Given $X \in \mathfrak{g}$, one defines a first order operator acting on smooth functions, f on the associated group manifold G by

$$\tilde{X}f(g) = \frac{d}{dt}f(g \exp tX)|_{t=0}.$$

(In our case we only be considered exponential groups so that $G = \exp(\mathfrak{g})$.) For any diffeomorphism $\gamma : G \rightarrow G$, one defines an action of γ on \tilde{X} by $\tilde{X}^\gamma f(g) = Xf(g \circ \gamma) \circ \gamma^{-1}$. \tilde{X} is left-invariant in the sense that taking γ_h to be the diffeomorphism of left translation on G by a fixed element $h \in G$ one has $\tilde{X}^{\gamma_h} = \tilde{X}$ for all such h .

This extends naturally to monomial elements of $\mathfrak{U}(\mathfrak{g})$ by

$$(X_1 X_2 \dots X_k)^\sim \cdot f = \frac{\partial^k}{\partial t_1 \partial t_2 \dots \partial t_k} f(g \exp tX_1 \exp tX_2 \dots \exp tX_k)_{t_1=\dots=t_k=0}$$

and then to all of $\mathfrak{U}(\mathfrak{g})$ by linearity. In fact one has an isomorphism of associative algebras [He78],

$$\mathfrak{U}(\mathfrak{g}) \simeq \mathbb{D}(G),$$

where $\mathbb{D}(G)$ denotes the space of all left-invariant differential operators, D on G (meaning $D^{\gamma_h} = D$). Along these same lines the adjoint action of G on \mathfrak{g} extends to $\mathfrak{U}(\mathfrak{g})$,

$$Ad_g(X_1 X_2 \dots X_k) = (Ad_g X_1)(Ad_g X_2) \dots (Ad_g X_k)$$

which corresponds to an adjoint action, $Ad_{g^{-1}}D$ on left-invariant differential operators. (Since D is left-invariant this amounts to the action induced through right translation by g .) One then defines

$$ad_X D \doteq \frac{d}{dt} Ad_{\exp(tX)} D|_{t=0}$$

which coincides with ad on \mathfrak{g} . It then follows from the product rule that

$$ad_Y X_1 \dots X_k = Y X_1 X_2 \dots X_k - X_1 X_2 \dots X_k Y.$$

It follows that if an element of $\mathfrak{U}(\mathfrak{g})$ commutes with all elements of \mathfrak{g} then it is an element of the center of $\mathfrak{U}(\mathfrak{g})$ denoted $\mathfrak{Z}(\mathfrak{g})$ which in turn is isomorphic to the commutative algebra of bi-invariant differential operators.

Having made these identifications one is led to a direct connection to invariant theory. Let $\mathbb{S}(\mathfrak{g})$ be the symmetric algebra over \mathfrak{g} . Then [He84] $\exists!$ linear bijection

$$\begin{aligned} \sigma : \mathbb{S}(\mathfrak{g}) &\rightarrow \mathbb{D}(G) \\ \ni \sigma(X^m) &= \tilde{X}^m. \end{aligned} \tag{A.1}$$

This is not an algebra isomorphism; however, σ commutes with the Ad_G action on $\mathbb{S}(\mathfrak{g})$ (which preserves $\mathbb{S}(\mathfrak{g})$). It is natural to identify $\mathbb{S}(\mathfrak{g})$ with the space of polynomial functions on \mathfrak{g}^* . Thus Ad_G -invariant polynomials on \mathfrak{g} map to $\mathfrak{Z}(\mathfrak{g})$. Moreover, if the polynomials $\{P_1, \dots, P_m\}$ generate the subalgebra of invariants, $\mathbb{S}(\mathfrak{g})^G$, then $\{\sigma(P_1), \dots, \sigma(P_m)\}$ generate $\mathfrak{Z}(\mathfrak{g})$ as an algebra of bi-invariant operators.

References

- [Arh79] A.A. Arhangel'skii. *Completely Integrable Hamiltonian Systems on the Group of Triangular Matrices*. (Russian) Mat. Sb. (N.S.) **108 (150)** 134-142 (1979).
- [AK71] L. Auslander and B. Kostant. *Polarization and Unitary Representations of Solvable Lie Groups*. *Invent. Math.* **14**, 255-354 (1971).
- [BBO05] P. Biane, P. Bougerol, and N. O'Connell. *Littleman Paths and Brownian Paths*. *Duke Math. J.* **130**, 127-167, (2005).
- [BGR17] A.M. Bloch, F. Gay-Balmaz, and T.S. Ratiu. *The Geometric Nature of the Flaschka Transformation*. *Communications in Mathematical Physics* **352**, 457-517 (2017).
- [BBF11] B. Brubaker, D. Bump and S. Friedberg. *Eisenstein Series, Crystals, and Ice*. *Notices Amer. Math. Soc.* **58**, 1563-1571, (2011).
- [DLNT86] P.Deift, L.C. Li, T. Nanda, and C. Tomei. *The Toda flow on a generic orbit is integrable*. *Communications on Pure and Applied Mathematics* **39**, 183-232 (1986).
- [Dix59] J. Dixmier. *Sur les Représentations Unitaires des Groupes de Lie Nilpotents IV*. *Canadian Journal of Mathematics*, **11**, 321-344 (1959).
- [Du77] M. Duflo. *Opérateurs Différentiels Bi-invariants sur un Groupe de Lie*. *Ann. Sci. École Norm. Sup.*, **10**, 265-288 (1977).
- [DM72] H. Dym and H. P. McKean. *Fourier Series and Integrals*. Probability and Mathematical Statistics, **14**, Academic Press, New York (1972).
- [EFS93] N. M. Ercolani, H. Flaschka, and S. Singer. *The Geometry of the Full Kostant-Toda Lattice*. In *Integrable systems (Luminy, 1991)*, vol. 115 of *Progr. Math.* Birkhäuser Boston, Boston, MA. 181-225 (1993).
- [Fl74] H. Flaschka. *The Toda lattice II. Existence of Integrals*. *Physical Review B*, **9**, 1924-1925 (1974).
- [Fo89] G.B. Folland. *Harmonic Analysis in Phase Space*. , **129**, Princeton University Press (1989).
- [FuHa91] W. Fulton and J. Harris. *Representation Theory: A First Course*. *Annals of Mathematics Studies*, **122**, Springer-Verlag, New York (1991).

- [GS99] M. I. Gekhtman and M. Z. Shapiro. *Non-Commutative and Commutative Integrability of Generic Toda Flows in Simple Lie Algebras*. *Communications on Pure and Applied Mathematics*, **52**(1), 53 - 84 (1999).
- [Gi97] A. Givental. *Stationary Phase Integrals, Quantum Toda Lattices, Flag Manifolds and the Mirror Conjecture*. In *Topics in Singularity Theory*, American Mathematical Society Translations, Ser. 2, AMS, Providence, RI (1997).
- [GuSt84] V. Guillemin and S. Sternberg. *Symplectic Techniques in Physics*. Cambridge University Press, (1984).
- [He78] S. Helgason. *Differential Geometry, Lie Groups and Symmetric Spaces*. AMS, (1978).
- [He84] S. Helgason. *Groups and Geometric Analysis*. AMS, (1984).
- [Jo77] A.A. Joseph. *A Preparation Theorem for the Prime Spectrum of a Semisimple Lie Algebra*. *Journal of Algebra* **48**, 241-289 (1977).
- [Ki62] A.A. Kirillov. *Unitary Representations of nilpotent Lie Groups*. *Russian Math Surveys* **17**, 57 - 110 (1962).
- [Ki99] A.A. Kirillov. *Merits and Demerits of the Orbit Method*. *Bulletin of the AMS* **36**, 433 - 488 (1999).
- [Ko70] B. Kostant. *Quantization and Unitary Representations*. *Lecture Notes in Math.* **170**, 87 - 208 (1970).
- [Ko78] B. Kostant. *On Whittaker Vectors and Representation Theory*. *Inventiones Math.* **48**, 101-184 (1978).
- [Ko79] B. Kostant. *The Solution to a Generalized Toda Lattice and Representation Theory*. *Advances in Mathematics* **34**, 195-338 (1979).
- [Ko12] B. Kostant. *The Cascade of Orthogonal Roots and the Coadjoint Structure of the Nilradical of a Borel Subgroup of a Semisimple Lie Group*. *Moscow Math. Journal* **12**, 1-16 (2012).
- [Ko13] B. Kostant. *Center $U(\mathfrak{n})$, Cascade of Orthogonal Roots, and a Construction of Lipsman-Wolf*. In *Lie Groups: Structure, Actions and Representations, In Honor of Joseph A. Wolf on the Occasion of his 75th Birthday*, vol. 306 of *Progr. Math.* Springer Science+Business Media, New York, NY 163 - 173 (2013).
- [Kn86] A. Knapp *Representation Theory of Semisimple Groups*. Princeton University Press (1986).
- [LW78] R.L. Lipsman and J.A. Wolf. *The Plancherel Formula for Parabolic Subgroups of the Classical Groups*. *Journal D'Analyse Mathématique* **34**, 120 - 161 (1978).
- [Ma58] G. Mackey. *Unitary Representations of Group Extensions I*. *Acta Math.*, **99**, 265 - 311 (1958).
- [MW73] C.C. Moore and J.A. Wolf. *Square Integrable Representations of nilpotent Groups*. *Transactions of the AMS* **185**, 445 - 462 (1973).
- [O13] N. O'Connell. *Geometric RSK and the Toda lattice*. *Illinois Journal of Mathematics* , **57**(3), 883-918 (2013).
- [Pu71] L. Pukanszky. *On the Theory of Exponential Groups*. *Trans. Amer. Math. Soc.*, **126**, 487-507 (1967).
- [Ra20] J. Ramalheira-Tsu. *The Kostant-Toda Lattice, Combinatorial Algorithms and Ultradiscrete Dynamics*. Available from ProQuest Dissertations & Theses Global (2461615150). (2020).
- [Sh10] F. Shahidi. *Eisenstein Series and Automorphic L-functions*. American Mathematical Society Colloquium Publications, Providence, RI (2010).
- [Si05] S.F. Singer. *Linearity, Symmetry and Prediction in the Hydrogen Atom*. Undergraduate Texts in Mathematics, Springer-Verlag, New York (2005).
- [Sy80] W. W. Symes. *Hamiltonian group actions and integrable systems*. *Physica D*, **1**, 339-374 (1980).
- [Th82] A. Thimm. *Integrable geodesic flows on homogeneous spaces*. *Ergod. Th. & Dynam. Sys.* **1**, 495-517 (1982).

- [To67] M. Toda. *Vibration of a chain with a non-linear interaction*. J. Phys. Soc. Jpn., **22**(2), 431-436 (1967).
- [Tro80] V.V. Trofimov. *Finite-Dimensional Representations of Lie Algebras and Completely Integrable Systems*. (Russian) Mat. Sb. (N.S.) **111 (153)** 610-621 (1980).
- [Wo14] J.A. Wolf. *Plancherel Formula for Minimal Parabolic Subgroups*. *Journal of Lie Theory* **24**, 791 - 808 (2014).
- [Wo16] J.A. Wolf. *Stepwise Square Integrable Representations: The Concept and Some Consequences*. in: Lie Theory and its Applications in Physics, Varna, Bulgaria, June 2015, Springer Proceedings in Mathematics & Statistics, Vol. 191, Springer Nature, Singapore, 181-202 (2016).