

Impact of Spatial Inhomogeneity on Excitation Energy Transport in the Fenna-Matthews-Olson Complex

Amartya Bose*

Department of Chemical Sciences, Tata Institute of Fundamental Research, Mumbai 400005, India

Peter L. Walters†

*Department of Chemistry, University of California, Berkeley, California 94720, USA and
Miller Institute for Basic Research in Science, University of California Berkeley, Berkeley, California 94720, USA*

The dynamics of the excitation energy transfer (EET) in photosynthetic complexes is an interesting question both from the perspective of fundamental understanding and the research in artificial photosynthesis. Challenges persist in numerically simulating these systems both in parameterizing them and following their dynamics over long periods of time. Over the past decade, very accurate spectral densities have been developed to capture spatial inhomogeneities in the Fenna-Matthews-Olson (FMO) complex. We investigate the dynamics of FMO with an exact treatment of various theoretical spectral densities. Because FMO has Hamiltonian elements that connect most of the bacteriochlorophyll sites together, it becomes difficult to rigorously identify the energy transport pathways in the complex. We use the recently introduced ideas of relating coherence to population derivatives to analyze the transport process and reveal some of the pathways.

Light harvesting complexes (LHCs) play an important role in photosynthesis in a host of plants, bacteria and algae. The so-called “antenna complexes” capture solar energy, converting it into an electronic excitation, and carries it to the reaction center where charge separation leads to further chemistry. Crucially, these systems form the basis and inspiration for attempts at artificial photosynthesis. Thus, understanding the mechanisms that allow for the efficient transport of the molecular excitation from the point of creation to the reaction center is of fundamental importance. A lot of work has been done in simulating the excitation energy transport (EET) and characterizing the vibronic couplings [1–6]. Early experiments [7, 8] seemed to provide evidence of quantum beating. Theoretical studies were performed around the same time to shed light on the origins of these long-lived electronic oscillations [9–12]. It was hypothesized that this oscillatory dynamics could be the reason behind the efficiency of EET in biosystems. However, more recent experimental investigations [13, 14] have shown that the optical 2D photon echo spectra at ambient temperature does not show long-lived electronic quantum coherence.

Theoretical studies have been widely performed using Redfield [9] and Förster theory [15]. However, the applicability of these approximate perturbative techniques is not always guaranteed *a priori*. Simulations of thermal dynamics at the ambient temperature is optimally performed using reduced density matrix-based approaches like the hierarchical equations of motion [16, 17] (HEOM) or path integrals using the Feynman-Vernon influence functional [18]. While the quasi-adiabatic propagator path integral [19, 20] method (QuAPI) has been used to study the Fenna-Matthews-Olson (FMO) complex [21], almost every other contemporary and more recent studies of the exact dynamics seem to have focused on using HEOM as the method of choice [6, 11, 22–24]. EET prob-

lems have also been studied using semiclassical methods [25–29]. Advances in path integral-based methods [30, 31] have recently made them lucrative for studying EET systems [32–36]. Among these advances, there have been developments that combine ideas from tensor network and influence functional to help alleviate the cost of path integral calculations in different ways [34, 37–42].

In this work, we simulate the EET dynamics in the FMO complex with accurate descriptions of the vibrational degrees of freedom. The usual practice is to focus the numerical studies on the evolution of the population in the excited states of each of the chromophores. Here, we additionally use the recently developed coherence-based analytic techniques [43–46] to unravel the pathways that the excitation takes and the impact of the vibrational modes. Baker and Habershon [47] have explored these pathways in FMO using the Lindblad master equation. Their approach based on knocking particular chromophores out of the system and evaluating the resulting change in the transport properties. Thus, they were able to ascertain the importance of particular chromophores to the transport process.

The methods that simulate the time evolution of the reduced density matrix, typically, require the calculation of spectral densities to characterize the effect of the protein scaffolding and the localized vibrations of the chlorophyll molecules. The spectral density can be calculated as the Fourier transform of the energy gap correlation function [48, 49]. In case of an EET, this is equivalent to the autocorrelation function of excitation energy fluctuations of individual chromophores. While a lot of work has been done trying to characterize the site-dependent spectral densities for the FMO [50–53], there seem to be significant disagreements on the best way of obtaining them. Zerner’s intermediate neglect of differential orbital method (ZINDO) and time-dependent density func-




FIG. 1. Fenna-Matthews-Olson Complex in its trimeric and monomeric forms. Blue: BChl 1. Orange: BChl 2. Green: BChl 3. Red: BChl 4. Purple: BChl 5. Brown: BChl 6. Pink: BChl 7. Gray: BChl 8.

tional theory (TD-DFT) have been used to calculate the excited state energy fluctuations along classical molecular dynamics (MD) trajectories run on the ground Born-Oppenheimer (BO) surface [51, 54, 55]. Extensive analysis has been done over the past decade on how to best and most consistently capture the fine interplay of dynamics and electronic structure that goes into these spectral densities. Because of the so-called “geometry mismatch” problem stemming from the inconsistencies between the conformations obtained from these purely classical MD trajectories and the subsequent quantum mechanical calculations, mixed QM/MM dynamics simulations on the ground BO surface have also been performed [55, 56].

Under physiological conditions, FMO exists as a trimer with each monomer consisting of eight bacteriochlorophyll-a (BChl a) units. The structure of the complex is shown in Fig. 1. The Hamiltonian describing the EET process in one 8-unit monomer can be expressed by a Frenkel model,

$$\hat{H}_0 = \sum_{j=1}^8 \epsilon_j |j\rangle\langle j| + \sum_{k \neq j} h_{k,j} |j\rangle\langle k|, \quad (1)$$

where ϵ_j is electronic excitation energy of the j th BChl molecule in the absence of the protein environment, and $h_{k,j}$ represents the electronic couplings. The state where only BChl j is excited is denoted by $|j\rangle = |e_j\rangle \otimes \prod_{k \neq j} |g_k\rangle$. Here, $|g_j\rangle$ and $|e_j\rangle$ represents the local ground and excited states of the j th BChl unit. The full dissipative environment including contributions from both the local, rigid vibrations and the dynamical environment of the protein scaffolding is characterized by a harmonic

bath on each site, j :

$$\hat{H}_B^j = \sum_{\xi} \frac{p_{j\xi}^2}{2m_{j\xi}} + \frac{1}{2} m_{j\xi} \omega_{j\xi}^2 \left(x_{j\xi} - \frac{c_{j\xi} \hat{s}_j}{m_{j\xi} \omega_{j\xi}^2} \right)^2, \quad (2)$$

where $\omega_{j\xi}$ and $c_{j\xi}$ are the frequency and coupling of the ξ th mode on BChl j . The bath interacts with the j th BChl unit through the diagonal operator \hat{s}_j specified by $\hat{s}_j |g_j\rangle = 0$ and $\hat{s}_j |e_j\rangle = |e_j\rangle$. The frequencies and couplings of the bath are related to the spectral density as follows:

$$J_j(\omega) = \frac{\pi}{2} \sum_{\xi} \frac{c_{j\xi}^2}{m_{j\xi} \omega_{j\xi}} \delta(\omega_{j\xi} - \omega). \quad (3)$$

Thus, the full universe including the system and the bath is defined by the following Hamiltonian, which has a Frenkel-Holstein structure:

$$\hat{H} = \hat{H}_0 + \sum_{j=1}^8 \hat{H}_B^j. \quad (4)$$

In many cases, the system Hamiltonian, \hat{H}_0 , is specified in terms of the site energy (a.k.a the optical excitation energy), E_j . This optical excitation energy is however dependent upon the environment. Therefore, ϵ_j is obtained after shifting the optical excitation energy by the corresponding reorganization energy [10].

Under the influence of the thermal vibrational baths, the reduced density matrix corresponding to the EET system at time $t = N\Delta t$ is given by a path integral expression,

$$\tilde{\rho}(S_N^{\pm}, N\Delta t) = \sum_{S_0^{\pm}} \sum_{S_1^{\pm}} \cdots \sum_{S_{N-1}^{\pm}} \tilde{\rho}(S_0^{\pm}, 0) P_{S_0^{\pm}, S_1^{\pm}, \dots, S_N^{\pm}} \quad (5)$$





FIG. 2. Matrix product representations of the path amplitude tensor and the influence functional operators.


$$= \sum_{S_0^\pm} \sum_{S_1^\pm} \cdots \sum_{S_{N-1}^\pm} \tilde{\rho}(S_0^\pm, 0) P_{S_0^\pm, S_1^\pm \dots S_N^\pm}^{(0)} F[\{S_n^\pm\}], \quad (6)$$

where $P_{S_0^\pm, S_1^\pm \dots S_N^\pm}^{(0)}$ is the path amplitude tensor which represents the amplitude of the system for moving along the specified sequence of forward-backward states in presence of the solvent. In the notation used here, S_n^\pm represents the collective forward-backward state of the system at the n th time point. (I.e., $S_n^\pm = \{s_{1,n}^\pm, s_{2,n}^\pm \dots s_{8,n}^\pm\}$, where $s_{j,n}^\pm$ is the forward-backward state of the j th site at the n th time point.) The path amplitude tensor is a product of the “bare” path amplitude tensor, $P_{S_0^\pm, S_1^\pm \dots S_N^\pm}^{(0)}$, representing the amplitude of the isolated system for moving along the same points and the Feynman-Vernon influence functional [18], F , representing the impact of the solvent degrees of freedom on the system. The dynamics of the isolated system is Markovian. It is the presence of the influence functional, F , that induces non-Markovianness in the dynamics. While formally this expression depends on the entire history of any path going back to time step 0, in condensed phases the memory dies out and is calculated only till a finite number of time-steps, L . This L is a convergence parameter.

The dimensionality of the path amplitude tensor grows as d^{2L} where the system is d -dimensional (in this case, $d = 8$), and L is the memory length. In the time-evolving matrix product operators (TEMPO) [37] method, this exponential growth is heavily compressed by recognizing that the correlations between points separated by large time-spans becomes negligible even in the presence of a non-Markovian bath. This suggests that one can use a matrix product state (MPS) to efficiently represent the path amplitude tensor. An MPS or a tensor train form is obtained by performing a sequential truncated singular value decompositions, so that each of the indices on the original tensor can be attributed to separate low-ranked tensors. In this representation, the terminal tensors are rank-2 and the intermediate tensors are rank-3 as shown in Fig. 2 (a). The common index between any two neighboring tensors and its dimension are called the bond in-



(a) Comparison with ZINDO-based classical MD spectral density from Ref. [51]



(b) Comparison with experiment [58, 59]

FIG. 3. Comparison of spectral densities averaged across all the eight BChl molecules obtained from Maity *et al.* [55]

dex and bond dimension respectively. In such a representation, it can be shown that the influence functional can similarly be analytically written as a matrix product operator (MPO) [39], which is shown in Fig. 2 (b). This analytic form optimizes the representation accounting for the multiple baths and the symmetries in the influence functional expression [39]. While the bare path amplitude has zero long-distance correlations, and consequently a very compact MPS representation, subsequent applications of the influence functional MPO builds up these correlations, leading to an increase in the bond dimension. However, efficient algorithms exist to minimize the growth of the bond dimension of the MPS on application of an MPO based on convergence parameters [57]. This combined with an optimal representation of the influence functional MPO makes the tensor network approach especially efficient at simulating these systems. This method is used for all the simulations in this paper.

For the purposes of this exploration, we use the

spectral densities calculated via QM/MM trajectories. The two cases considered correspond to energy-gap autocorrelation functions calculated using the TD-LC-DFTB method (average reorganization energy of $\lambda = 572.976 \text{ cm}^{-1}$) and the ZINDO/S-CIS semi-empirical method (average reorganization energy of $\lambda = 839.032 \text{ cm}^{-1}$) reported by Maity *et al.* [55]. In this study, the site energies for the system Hamiltonian were calculated using the same QM/MM MD trajectories and the off-diagonal terms were calculated using TrESP [55]. Comparison of the dynamics under these recent spectral densities with that under the influence of a spectral density calculated using classical trajectories and ZINDO/S-CIS [51] is also shown. In many of the later studies, the spectral densities were resolved for each BChl unit and for each of the FMO monomers, effectively leading to 24 spectral densities. However, for simplicity we are using the BChl site-specific spectral densities averaged across the three monomers in FMO. As discussed, the optical excitation energies include influence from the solvents. Consequently, the electronic excitations, ϵ_j , in the system Hamiltonian, \hat{H}_0 , are obtained from the site energies in Ref. [55] by subtracting the site-specific reorganization energy. We used the correct system Hamiltonian corresponding to the QM/MM MD spectral densities.

The various spectral densities, averaged across all the BChls, are plotted in Fig. 3. We also show a spectral density obtained using classical MD trajectories on ZINDO. The classical MD simulation of the spectral density suffers from a significant blue shift of the high frequency vibrations. Both the QM/MM spectral densities alleviate this problem. Though the alignment of the peaks is not perfect even when using QM/MM MD simulations, it is quite close to the experimental fluorescence line narrowing spectral density [58, 59]. (Notice that the spectral density obtained using classical MD and ZINDO is far smoother than the more recent spectral densities. This is a result of a fitting procedure that was used by Olbrich *et al.* [51].)

There are a few questions that we want to focus our exploration on: (1) How important is choice of the density functional for characterizing the vibrations and the protein scaffolding vis-à-vis the EET dynamics? (2) What is the impact of the spatial inhomogeneity on the EET dynamics? (3) How does one analyze the individual pathways and routes of energy flow using exact dynamics (4) Given that BChl 3 is the sink of the EET process, how much of the excitation energy goes into BChl 3 and how quickly? The last two questions, and their connection with the features of spectral densities, is especially important in terms of efficiency of the EET process.

First, let us consider the population dynamics corresponding to an initial excitation of a single BChl unit. In Fig. 4 and Fig. 5, we show the dynamics that happen from each of the different QM/MM MD spectral densities at an ambient temperature of $T = 300 \text{ K}$ with initial




FIG. 4. Comparison of dynamics starting from $\tilde{\rho}(0) = |1\rangle\langle 1|$ for the different *ab initio* methods and with or without the spatial inhomogeneities. Blue: BChl 1. Orange: BChl 2. Green: BChl 3. Red: BChl 4. Purple: BChl 5. Brown: BChl 6. Pink: BChl 7. Gray: BChl 8.




FIG. 5. Comparison of dynamics starting from $\tilde{\rho}(0) = |6\rangle\langle 6|$ for the different *ab initio* methods and with or without the spatial inhomogeneities. Colors same as Fig. 4.

conditions $\tilde{\rho}(0) = |1\rangle\langle 1|$ and $\tilde{\rho}(0) = |6\rangle\langle 6|$ respectively. Figures 4 and 5 (a) and (c) correspond to the site specific spectral densities, whereas Figs. 4 and 5 (b) and (d) correspond to the dynamics happening in the presence of the average environments. It is interesting to note that while for both the methods, the transfer into site 3 is faster in presence of the average spectral density when $\tilde{\rho}(0) = |1\rangle\langle 1|$, this is not the case when $\tilde{\rho}(0) = |6\rangle\langle 6|$. This seems to indicate that possibly not all evolutionary modifications to these systems are geared towards an increased transport into the sink site 3. It is instructive to note that it is not possible to come up with a very simple conclusion about the effect of the spatial inhomogeneity, which seems to be initial condition dependent.

The second aspect to consider from the Figs. 4 and 5 is the effect of changing the method of calculating the energy-gap. The most prominent difference between the spectral densities corresponding to the two approaches is the higher intensity and consequently a larger reorganization energy of the ZINDO spectral density [55, 60]. This leads to a slowing down of dynamics under ZINDO. It is noteworthy that despite this slowdown of the dynamics, the transfer from site 1 to site 2 in Fig. 4 is increased from the TD-LC-DFTB functional to the ZINDO semi-empirical method. Also, note that the bath inhomogeneities bring about a smaller difference in the dynamics compared to the effect of changing from TD-LC-DFTB to ZINDO.

While it is known that resorting to classical MD trajectory-based simulations generally leads to geometry mismatch problems, the actual impact of these errors on the dynamics in this case has not been evaluated. The spectral density of the classical ZINDO bath is shown in Fig. 3 (a). The resultant spectral densities are clearly different from the QM/MM ones. However, the effect of these differences on the dynamics is far from obvious. In Fig. 6, we show the dynamics corresponding to the system coupled to site-local baths described by the classical ZINDO spectral density. The system Hamiltonian corresponding to the ZINDO QM/MM MD simulation is used with the classical spectral density as well. This has been done to ensure that the effects we see are only coming from the spectral density. In fact it is surprising, that despite the enormity of the differences at the spectral density level, the dynamics is relatively similar to the QM/MM calculation. The differences are quite subtle. In fact, it is arguable whether the differences in dynamics caused by doing a qualitatively incorrect calculation is greater than the ones seen by using a different method of estimating the energy gap.

The effects of the method behind the excited state calculation and the spatial inhomogeneity has been demonstrated on the EET. It is interesting to probe more deeply into the mechanism behind these effects. Such a probe will reveal subtler features of the dynamics than what was seen through the population dynamics. Crucial to




FIG. 6. Comparison of dynamics corresponding to ZINDO QM/MM MD spectral density versus ZINDO classical MD spectral density. Colors same as in Fig. 4.

such an exploration is an understanding of the pathway-dependent population transfer for each of the cases. Wu *et al.* [43] have built flux networks to analyze the flows along different pathways. More recently, Dani and Makri have shown that the time-derivative of the on-site populations is related to linear combinations of the off-diagonal terms of the reduced density matrix [44], and visualized the coherences in forms of maps that encode information about the dynamics of the system [45]. We extended these ideas to partition the time-dependent population change on a site in terms of transport along different state-to-state channels, which is useful for understanding the instantaneous population transfer in these systems with complex interconnects [46]. To explore the overall importance of the different pathways using this coherence-based analysis, we use a modified version of the integrated flux approach [43] and define a time-averaged state-to-state population transfer:

$$\bar{P}_{k\leftarrow j} = \frac{1}{T} \int_0^T dt P_{k\leftarrow j}(t), \quad (7)$$

where $P_{k\leftarrow j}(t) = -\frac{2}{\hbar} \langle k|\hat{H}_0|j\rangle \int_0^t dt' \text{Im} \langle k|\tilde{\rho}(t')|j\rangle$ is the population transferred directly from the j th site to the k th at time t . Here the time of integration is taken to be $T \approx 1500$ fs. The main benefit of using a time-average instead of a simple integration of the flux is that using the average one can capture the relative speeds of transfer as




FIG. 7. State-to-state population transfer in presence of the TD-LC-DFTB baths for different initial conditions. The $(k, j)^{\text{th}}$ cell in the image represents the average population transferred directly from site j to site k , $\bar{P}_{k \leftarrow j}$.

well.

The complete information about the average population transfer along each of these channels for the different initial conditions is presented in Fig. 7. This data corresponds to the TD-LC-DFTB spectral density though the main trends carry over to the ZINDO spectral density as well. Notice that for the case when $\tilde{\rho}(0) = |1\rangle\langle 1|$, the most important pathway is clearly $1 \rightarrow 2 \rightarrow 3$, with a secondary component coming from the $1 \rightarrow 6 \rightarrow 5 \rightarrow 4 \rightarrow 3$. This is known in the literature. Non-insignificant contributions also happen along $1 \rightarrow 5 \rightarrow 4 \rightarrow 3$ and $1 \rightarrow 2 \rightarrow 4 \rightarrow 3$, both short-circuiting site 6. On the other hand, if $\tilde{\rho}(0) = |6\rangle\langle 6|$, the principle pathway




FIG. 8. Time integrated transport to the 3rd site, $\bar{P}^{(3)}$, when $\tilde{\rho}(0) = |1\rangle\langle 1|$. Dark blue: $\bar{P}_{3 \leftarrow 2}$. Light blue: $\bar{P}_{3 \leftarrow 1}$. Red: $\langle \text{rest} \rangle$.




FIG. 9. Time integrated transport to the 3rd site, $\bar{P}^{(3)}$, when $\tilde{\rho}(0) = |6\rangle\langle 6|$. Dark blue: $\bar{P}_{3 \leftarrow 2}$. Dark red: $\bar{P}_{3 \leftarrow 6}$. Red: $\bar{P}_{3 \leftarrow 4}$.

is $6 \rightarrow 5 \rightarrow 4 \rightarrow 3$, with very tiny proportions of $6 \rightarrow 1 \rightarrow 2 \rightarrow 3$ and $6 \rightarrow 2 \rightarrow 3$.

Now let us concentrate on contributions of the most important pathways to the sink site 3. For our analysis, we divide the population transfer from 1 to 3 into three components: (1) direct transport from 1 to 3 ($\bar{P}_{3 \leftarrow 1}$); (2) transport via 2 to 3 ($\bar{P}_{3 \leftarrow 2}$); (3) transport from all other sites combined which encodes the possibility of transport in the $1 \rightarrow 6 \rightarrow 5 \rightarrow 4 \rightarrow 3$ (represented as $\langle \text{rest} \rangle$). The contributions of these three pathways into site 3 are shown in Fig. 8.

The classical MD spectral density completely wipes out the direct transfer from site 1 to 3. Though this direct route, $\bar{P}_{3 \leftarrow 1}$, is not exceptionally important, for the newer QM/MM-based spectral densities, it gives a non-negligible contribution to the total population of site 3. Also, note that the biggest impact of the spatial inhomogeneties is on the $1 \rightarrow 2 \rightarrow 3$ pathway. More specifically, the transfer from 2 to 3 gets substantially inhibited by the presence of different site-specific spectral densities. It is this inhibition of $\bar{P}_{3 \leftarrow 2}$ that is primarily responsible for the decrease of the overall transport into site 3.

In order to further explore the effect of the spatially inhomogeneous spectral densities, let's consider in a bit more detail the percentage contribution of the different

components to the total transport. We have already mentioned the overall increase in the transport to the 3rd site for the average spectral densities. We notice that another effect that is consistently reproduced is the fact the percentage contribution of the channel $\bar{P}_{3\leftarrow 2}$ consistently goes up from the site-dependent bath to the average bath. (The percentage contribution of $\bar{P}_{3\leftarrow 2}$ goes up from approximately 87% to 90% for TD-LC-DFTB and from approximately 81% to 88% for ZINDO.) In contrast, the proportion of population along $\langle \text{rest} \rangle$ is decreased quite significantly by the average bath (close to 7% to 3% for the TD-LC-DFTB spectral density and from around 13% to 6% for ZINDO). The direct transport along $\bar{P}_{3\leftarrow 1}$ also decreases in the presence of the average spectral density, but not by as much a margin. (While here we have been talking in terms of percentage contributions, the story is slightly different when it comes to the absolute transfers along these channels. While $\bar{P}_{3\leftarrow 2}$ and $\bar{P}_{3\leftarrow 1}$ both increase, the former much more significantly than the latter, the absolute contribution along $\langle \text{rest} \rangle$ actually decreases.) These subtle features would have been inaccessible in absence of a method to probe the pathways of EET.

A similar analysis can be done for an initial excitation on the 6th monomer, $\tilde{\rho}(0) = |6\rangle\langle 6|$. Notice that in Fig. 9, the patterns are even more convoluted than before. The most obvious thing that one can report is that the major contributor to site 3 is site 4. Consulting Fig. 7, we realize that this is must be coming from the major pathway of $6 \rightarrow 5 \rightarrow 4 \rightarrow 3$. Contributions of $6 \rightarrow 4 \rightarrow 3$ is minimal. The direct transport from site 2 to site 3 is negligible in all cases. The conclusions with any degree of universality stop there. Transport from 6 directly to 3 may be an important factor, however, its importance seems to be dependent on the particular excited state method used. Whereas the average spectral density gives a lesser transfer into site 3 for the TD-LC-DFTB spectral densities, the reverse is true for ZINDO. Additionally notice that the magnitude of differences between the TD-LC-DFTB and the ZINDO bars in Figs. 8 and 9 is larger than that between the ZINDO and the classical MD bars. Thus, the change between the spectral densities is probably significantly less important than that of changing the electronic excitation energies in the system Hamiltonian. This once again demonstrates the growing need for accurate parameters for these complex systems.

In this paper, we have evaluated the exact dynamics of the EET process in the FMO complex when the rovibrational modes of the molecules and the protein scaffold are described by accurate *ab initio* molecular dynamics computations. We have tried to shed light on the effects of the inhomogeneities in the solvent by comparing with the dynamics corresponding to the average bath. Surprisingly, the effects of changing the excited state method seem to be possibly of a larger magnitude to that of removing spatial inhomogeneities or even that

of changing from QM/MM simulations to classical MD simulations. Therefore, the noticeable differences in the spectral density must get washed out when it comes to the dynamics. Additionally, this seems to imply that estimating the correct electronic excitation energies of the BChl monomers is more important than simulating an accurate spectral density.

There is a limit to the amount of information that can be extracted from the direct population dynamics. We have used the recently introduced ideas of relating the coherence to the population transport [44–46] to explore the routes of transfer that the molecular excitation takes. Given the very complex interactions between these chromophores, it is not trivial to use an exact numerical computation to attribute the excitation transport to the different directed path-ways that exist. An analysis of the coherences is uniquely capable of answering these questions, and we have used this technique to present a static picture of the transport as it happens. In doing so, we have uncovered a couple of previously unnoticed transport pathways for the FMO. An analysis of the pathway specific contributions to the transport problem has yielded further evidence to support the greater importance of the system Hamiltonian and electronic excitation energies as opposed to the spectral density. These numerically “exact” approaches, in combination with better descriptions of the vibronic interactions, promise to provide in-depth understandings of transport in similar complex open systems.

ACKNOWLEDGMENT

We thank Prof. Ulrich Kleinekathöfer and Sayan Maity for sharing their spectral densities with us and for helpful discussions. AB acknowledges the support of Princeton University and the Computational Chemical Science Center: Chemistry in Solution and at Interfaces funded by the U.S. Department of Energy, under Award No. DESC0019394 for providing resources for the simulation of some of the dynamics.

* amartya.bose@tifr.res.in; Both authors contributed equally to this work.

† peter.l.walters2@gmail.com; Both authors contributed equally to this work.

- [1] T. Renger, Theory of excitation energy transfer: From structure to function, *Photosynthesis Research* **102**, 471 (2009).
- [2] J. Cao, R. J. Cogdell, D. F. Coker, H.-G. Duan, J. Hauer, U. Kleinekathöfer, T. L. C. Jansen, T. Mančal, R. J. D. Miller, J. P. Ogilvie, V. I. Prokhorenko, T. Renger, H.-S. Tan, R. Tempelaar, M. Thorwart, E. Thyryhaug, S. Westenhoff, and D. Zigmantas, Quantum biology revisited, *Science Advances* **6**, eaaz4888.

- [3] S. Saito, M. Higashi, and G. R. Fleming, Site-dependent fluctuations optimize electronic energy transfer in the Fenna–Matthews–Olson protein, *The Journal of Physical Chemistry B* **123**, 9762 (2019), <https://doi.org/10.1021/acs.jpcc.9b07456>.
- [4] E. Cignoni, V. Slama, L. Cupellini, and B. Mennucci, The atomistic modeling of light-harvesting complexes from the physical models to the computational protocol, *The Journal of Chemical Physics* **156**, 120901 (2022).
- [5] S. Maity and U. Kleinekathöfer, Recent progress in atomistic modeling of light-harvesting complexes: A mini review, *Photosynthesis Research* 10.1007/s11120-022-00969-w (2022).
- [6] X. Cui, Y. Yan, and J. Wei, Role of Pigment–Protein Coupling in the Energy Transport Dynamics in the Fenna–Matthews–Olson Complex, *The Journal of Physical Chemistry B* **125**, 11884 (2021).
- [7] S. Savikhin, D. R. Buck, and W. S. Struve, Oscillating anisotropies in a bacteriochlorophyll protein: Evidence for quantum beating between exciton levels, *Chemical Physics* **223**, 303 (1997).
- [8] G. S. Engel, T. R. Calhoun, E. L. Read, T.-K. Ahn, T. Mančal, Y.-C. Cheng, R. E. Blankenship, and G. R. Fleming, Evidence for wavelike energy transfer through quantum coherence in photosynthetic systems, *Nature* **446**, 782 (2007).
- [9] A. Ishizaki and G. R. Fleming, On the adequacy of the Redfield equation and related approaches to the study of quantum dynamics in electronic energy transfer, *The Journal of Chemical Physics* **130**, 234110 (2009).
- [10] A. Ishizaki and G. R. Fleming, Unified treatment of quantum coherent and incoherent hopping dynamics in electronic energy transfer: Reduced hierarchy equation approach, *The Journal of Chemical Physics* **130**, 234111 (2009).
- [11] A. Ishizaki and G. R. Fleming, Theoretical examination of quantum coherence in a photosynthetic system at physiological temperature, *Proceedings of the National Academy of Sciences* **106**, 17255 (2009).
- [12] M. Sarovar, A. Ishizaki, G. R. Fleming, and K. B. Whaley, Quantum entanglement in photosynthetic light-harvesting complexes, *Nature Physics* **6**, 462 (2010).
- [13] H.-G. Duan, V. I. Prokhorenko, R. J. Cogdell, K. Ashraf, A. L. Stevens, M. Thorwart, and R. J. D. Miller, Nature does not rely on long-lived electronic quantum coherence for photosynthetic energy transfer, *Proceedings of the National Academy of Sciences* **114**, 8493 (2017).
- [14] E. Thyryhaug, R. Tempelaar, M. J. P. Alcocer, K. Židek, D. Bína, J. Knoester, T. L. C. Jansen, and D. Zigmantas, Identification and characterization of diverse coherences in the Fenna–Matthews–Olson complex, *Nature Chemistry* **10**, 780 (2018).
- [15] T. Förster, Zwischenmolekulare Energiewanderung und Fluoreszenz, *Annalen der Physik* **437**, 55 (1948).
- [16] Y. Tanimura and R. Kubo, Time Evolution of a Quantum System in Contact with a Nearly Gaussian-Markoffian Noise Bath, *Journal of the Physical Society of Japan* **58**, 101 (1989).
- [17] Y. Tanimura, Numerically “exact” approach to open quantum dynamics: The hierarchical equations of motion (HEOM), *The Journal of Chemical Physics* **153**, 20901 (2020).
- [18] R. P. Feynman and F. L. Vernon, The theory of a general quantum system interacting with a linear dissipative system, *Annals of Physics* **24**, 118 (1963).
- [19] N. Makri and D. E. Makarov, Tensor propagator for iterative quantum time evolution of reduced density matrices. I. Theory, *The Journal of Chemical Physics* **102**, 4600 (1995).
- [20] N. Makri and D. E. Makarov, Tensor propagator for iterative quantum time evolution of reduced density matrices. II. Numerical methodology, *The Journal of Chemical Physics* **102**, 4611 (1995).
- [21] P. Nalbach and M. Thorwart, The role of discrete molecular modes in the coherent exciton dynamics in FMO, *Journal of Physics B: Atomic, Molecular and Optical Physics* **45**, 154009 (2012).
- [22] J. Strümpfer and K. Schulten, The effect of correlated bath fluctuations on exciton transfer, *The Journal of Chemical Physics* **134**, 95102 (2011).
- [23] J. Strümpfer and K. Schulten, Excited state dynamics in photosynthetic reaction center and light harvesting complex 1, *The Journal of Chemical Physics* **137**, 65101 (2012).
- [24] D. M. Wilkins and N. S. Dattani, Why Quantum Coherence Is Not Important in the Fenna–Matthews–Olsen Complex, *Journal of Chemical Theory and Computation* **11**, 3411 (2015).
- [25] G. Tao and W. H. Miller, Semiclassical Description of Electronic Excitation Population Transfer in a Model Photosynthetic System, *The Journal of Physical Chemistry Letters* **1**, 891 (2010).
- [26] P. Nalbach, A. Ishizaki, G. R. Fleming, and M. Thorwart, Iterative path-integral algorithm versus cumulant time-nonlocal master equation approach for dissipative biomolecular exciton transport, *New Journal of Physics* **13**, 63040 (2011).
- [27] M. K. Lee, P. Huo, and D. F. Coker, Semiclassical Path Integral Dynamics: Photosynthetic Energy Transfer with Realistic Environment Interactions, *Annual Review of Physical Chemistry* **67**, 639 (2016).
- [28] M. K. Lee and D. F. Coker, Modeling Electronic-Nuclear Interactions for Excitation Energy Transfer Processes in Light-Harvesting Complexes, *The Journal of Physical Chemistry Letters* **7**, 3171 (2016).
- [29] E. Mulvihill, K. M. Lenn, X. Gao, A. Schubert, B. D. Dunietz, and E. Geva, Simulating energy transfer dynamics in the Fenna–Matthews–Olson complex via the modified generalized quantum master equation, *The Journal of Chemical Physics* **154**, 204109 (2021).
- [30] N. Makri, Small matrix disentanglement of the path integral: Overcoming the exponential tensor scaling with memory length, *The Journal of Chemical Physics* **152**, 41104 (2020).
- [31] N. Makri, Small Matrix Path Integral for System-Bath Dynamics, *Journal of Chemical Theory and Computation* **16**, 4038 (2020).
- [32] S. Kundu and N. Makri, Real-Time Path Integral Simulation of Exciton-Vibration Dynamics in Light-Harvesting Bacteriochlorophyll Aggregates, *The Journal of Physical Chemistry Letters* **11**, 8783 (2020).
- [33] A. Bose and N. Makri, All-Mode Quantum–Classical Path Integral Simulation of Bacteriochlorophyll Dimer Exciton-Vibration Dynamics, *The Journal of Physical Chemistry B* **124**, 5028 (2020).
- [34] A. Bose and P. L. Walters, Tensor Network Path Integral Study of Dynamics in B850 LH2 Ring with Atomistically Derived Vibrations, *Journal of Chemical Theory*

- and Computation **18**, 4095 (2022).
- [35] S. Kundu, R. Dani, and N. Makri, B800-to-B850 relaxation of excitation energy in bacterial light harvesting: All-state, all-mode path integral simulations, *The Journal of Chemical Physics* **157**, 015101 (2022), <https://doi.org/10.1063/5.0093828>.
- [36] S. Kundu, R. Dani, and N. Makri, Tight inner ring architecture and quantum motion of nuclei enable efficient energy transfer in bacterial light harvesting, *Science Advances* **8**, eadd0023 (2022).
- [37] A. Strathearn, P. Kirton, D. Kilda, J. Keeling, and B. W. Lovett, Efficient non-Markovian quantum dynamics using time-evolving matrix product operators, *Nature Communications* **9**, 10.1038/s41467-018-05617-3 (2018).
- [38] M. R. Jørgensen and F. A. Pollock, Exploiting the Causal Tensor Network Structure of Quantum Processes to Efficiently Simulate Non-Markovian Path Integrals, *Physical Review Letters* **123**, 240602 (2019).
- [39] A. Bose and P. L. Walters, A tensor network representation of path integrals: Implementation and analysis, arXiv pre-print server arXiv:2106.12523 (2021), arXiv:2106.12523.
- [40] A. Bose, Pairwise connected tensor network representation of path integrals, *Physical Review B* **105**, 024309 (2022).
- [41] A. Bose and P. L. Walters, A multisite decomposition of the tensor network path integrals, *The Journal of Chemical Physics* **156**, 024101 (2022).
- [42] A. Bose and P. L. Walters, Effect of temperature gradient on quantum transport, *Physical Chemistry Chemical Physics* **24**, 22431 (2022).
- [43] J. Wu, F. Liu, J. Ma, R. J. Silbey, and J. Cao, Efficient energy transfer in light-harvesting systems: Quantum-classical comparison, flux network, and robustness analysis, *The Journal of Chemical Physics* **137**, 174111 (2012).
- [44] R. Dani and N. Makri, Quantum State-to-State Rates for Multistate Processes from Coherences, *The Journal of Physical Chemistry Letters* **13**, 8141 (2022).
- [45] R. Dani and N. Makri, Time-Evolving Quantum Superpositions in Open Systems and the Rich Content of Coherence Maps, *The Journal of Physical Chemistry B* **126**, 9361 (2022).
- [46] A. Bose and P. L. Walters, Impact of solvent on state-to-state population transport in multistate systems using coherences, arXiv pre-print server arXiv:2301.12712 (2023), arXiv:2301.12712.
- [47] L. A. Baker and S. Habershon, Robustness, efficiency, and optimality in the Fenna-Matthews-Olson photosynthetic pigment-protein complex, *The Journal of Chemical Physics* **143**, 105101 (2015).
- [48] N. Makri, The Linear Response Approximation and Its Lowest Order Corrections: An Influence Functional Approach, *The Journal of Physical Chemistry B* **103**, 2823 (1999).
- [49] A. Bose, Zero-cost corrections to influence functional coefficients from bath response functions, *The Journal of Chemical Physics* **157**, 054107 (2022).
- [50] A. Klinger, D. Lindorfer, F. Müh, and T. Renger, Normal mode analysis of spectral density of FMO trimers: Intra- and intermonomer energy transfer, *The Journal of Chemical Physics* **153**, 215103 (2020).
- [51] C. Olbrich, J. Strümpfer, K. Schulten, and U. Kleinekathöfer, Theory and Simulation of the Environmental Effects on FMO Electronic Transitions, *The Journal of Physical Chemistry Letters* **2**, 1771 (2011).
- [52] T. Renger, A. Klinger, F. Steinecker, M. Schmidt am Busch, J. Numata, and F. Müh, Normal Mode Analysis of the Spectral Density of the Fenna-Matthews-Olson Light-Harvesting Protein: How the Protein Dissipates the Excess Energy of Excitons, *The Journal of Physical Chemistry B* **116**, 14565 (2012).
- [53] M. L. Chaillet, F. Lengauer, J. Adolphs, F. Müh, A. S. Fokas, D. J. Cole, A. W. Chin, and T. Renger, Static Disorder in Excitation Energies of the Fenna-Matthews-Olson Protein: Structure-Based Theory Meets Experiment, *The Journal of Physical Chemistry Letters* **11**, 10306 (2020).
- [54] C. Olbrich and U. Kleinekathöfer, Time-Dependent Atomistic View on the Electronic Relaxation in Light-Harvesting System II, *The Journal of Physical Chemistry B* **114**, 12427 (2010).
- [55] S. Maity, B. M. Bold, J. D. Prajapati, M. Sokolov, T. Kubař, M. Elstner, and U. Kleinekathöfer, DFTB/MM Molecular Dynamics Simulations of the FMO Light-Harvesting Complex, *The Journal of Physical Chemistry Letters* **11**, 8660 (2020).
- [56] S. M. Blau, D. I. G. Bennett, C. Kreisbeck, G. D. Scholes, and A. Aspuru-Guzik, Local protein solvation drives direct down-conversion in phycobiliprotein PC645 via incoherent vibronic transport, *Proceedings of the National Academy of Sciences of the United States of America* **115**, E3342 (2018).
- [57] S. Paeckel, T. Köhler, A. Swoboda, S. R. Manmana, U. Schollwöck, and C. Hubig, Time-evolution methods for matrix-product states, *Annals of Physics* **411**, 167998 (2019).
- [58] M. Rätsep and A. Freiberg, Electron-phonon and vibronic couplings in the FMO bacteriochlorophyll a antenna complex studied by difference fluorescence line narrowing, *Journal of Luminescence* **127**, 251 (2007).
- [59] A. Kell, X. Feng, M. Reppert, and R. Jankowiak, On the Shape of the Phonon Spectral Density in Photosynthetic Complexes, *The Journal of Physical Chemistry B* **117**, 7317 (2013).
- [60] S. Chandrasekaran, M. Aghtar, S. Valleau, A. Aspuru-Guzik, and U. Kleinekathöfer, Influence of Force Fields and Quantum Chemistry Approach on Spectral Densities of BChl a in Solution and in FMO Proteins, *The Journal of Physical Chemistry B* **119**, 9995 (2015).