

SUFFICIENT STATISTIC AND RECOVERABILITY VIA QUANTUM FISHER INFORMATION METRICS

LI GAO, HAOJIAN LI, IMAN MARVIAN, AND CAMBYSE ROUZÉ

ABSTRACT. We prove that for a large class of quantum Fisher information, a quantum channel is sufficient for a family of quantum states, i.e., the input states can be recovered from the output, if and only if the quantum Fisher information is preserved under the quantum channel. This class, for instance, includes Winger-Yanase-Dyson skew information. On the other hand, interestingly, the SLD quantum Fisher information, as the most popular example of quantum analog of Fisher information, does not satisfy this property.

Our recoverability result is obtained by studying Riemannian monotone metrics on the quantum state space, i.e. metrics monotone decreasing under the action of quantum channels, a property often called data processing inequality. For two quantum states, the monotone metric gives the corresponding quantum χ^2 divergence. We obtain an approximate recovery result in the sense that, if the quantum χ^2 divergence is approximately preserved by a quantum channel, then two states can be approximately recovered by the Petz recovery map. We also obtain a universal recovery bound for the $\chi_{\frac{1}{2}}$ divergence.

Finally, we discuss applications in the context of quantum thermodynamics and the resource theory of asymmetry.

1. INTRODUCTION

Quantum metrology studies high-resolution measurements of physical parameters of quantum systems. In both classical and quantum metrology, the Fisher information plays an important role as a metric measuring the amount of information a system carries about a parameter θ . The concept of Fisher information goes back to mathematical statistics [Fis22]: let (Ω, μ) be a probability space and $X(\theta) : \Omega \rightarrow \mathbb{R}$ be a family of random variables depending on an unknown parameter θ . The Fisher information of X at θ is defined as

$$I_X(\theta) := \mathbb{E} \left[\left(\partial_\theta \log p_X(\theta) \right)^2 \middle| \theta \right] = \int_{\Omega} \frac{|\partial_\theta p_X(\theta, \omega)|^2}{p_X(\theta, \omega)} d\mu(\omega), \quad (1.1)$$

where $\omega \mapsto p_X(\theta, \omega)$ is the probability density function of $X(\theta)$ with respect to μ . By the famous Cramér–Rao bound [Rao45, Cra16], the Fisher information gives a fundamental limit on the precision of parameter estimation: for any unbiased estimator $\hat{\theta}$ of θ ¹, it holds that $\text{Var}(\hat{\theta}) \geq 1/I_X(\theta)$. The Cramér–Rao bound has been extended to the quantum setting [Hol11, BC94, BCM96]: for a family of quantum states ρ_θ the variance of any unbiased estimator $\hat{\theta}$ of θ satisfies

$$\text{Var}(\hat{\theta}) \geq \frac{1}{I_{\text{SLD}, \rho}(\theta)}. \quad (1.2)$$

where

$$I_{\text{SLD}, \rho}(\theta) = \text{tr}(\dot{\rho}_\theta \mathbb{J}_{\rho_\theta}(\dot{\rho}_\theta)), \quad (1.3)$$

¹Here an unbiased estimator satisfies $\mathbb{E}(\hat{\theta}|\theta) = \theta$.

and \mathbb{J}_ρ is the inverse of the symmetric multiplication map $\mathbb{J}_\rho^{-1}(A) = \frac{1}{2}(\rho A + A\rho)$. In the physics literature I_{SLD} is often called the quantum Fisher information (QFI). Following the quantum statistics literature, we call this quantity the symmetric logarithmic derivative (SLD) QFI.

A nice property of the quantum Cramér–Rao bound with SLD QFI is that, similar to its classical version, it is asymptotically achievable; that is, given state $\rho(\theta)^{\otimes n}$ there exists a measurement for which the above bound becomes tight in the limit $n \rightarrow \infty$ [BNG00, BC94]. This essentially follows from the fact that (i) both quantum and classical Fisher information are additive, and (ii) there exists a measurement on a single copy of $\rho(\theta)$, for which the classical Fisher information of the outcome is equal to $I_{\text{SLD},\rho}(\theta)$. Then, combining these facts with the asymptotic achievability of the classical Cramér–Rao bound, one can establish the achievability of the Quantum Cramér–Rao bound.² Hence SLD QFI has a distinguished role in statistics that puts it on par with classical Fisher information.

1.1. Sufficient Statistic. Another important property of classical Fisher information is in the context of sufficient statistic. For the family of random variables X with distribution $P_X(\theta)$, a statistic $t = T(X)$ is called sufficient for parameter θ if it contains all the information in X about θ , such that given $t = T(X)$, the random variable X becomes independent of θ .

It is known (see c.f. [Sch12]) that under certain regularity conditions, e.g., if $P_X(\theta)$ has full support for all θ , then

$$t = T(X) \text{ is a sufficient statistic for } \theta \iff I_{T(X)}(\theta) = I_X(\theta). \quad (1.4)$$

In the quantum setting, the notion of sufficient statistic can be defined in terms of recoverability with quantum channels: given a family of quantum states ρ_θ , a quantum channel Φ is sufficient for the family ρ_θ if and only if there exists a quantum channel \mathcal{R} such that $\mathcal{R} \circ \Phi(\rho_\theta) = \rho_\theta$ for all θ . Such \mathcal{R} is called a recovery map, which means the original family ρ_θ can be fully recovered from the channel output $\Phi(\rho_\theta)$. Jenčová and Petz [JP06] showed that such quantum sufficiency can be characterized via the relative entropy³: Φ is sufficient if and only if there exists some state σ such that

$$D(\rho_\theta \parallel \sigma) = D(\Phi(\rho_\theta) \parallel \Phi(\sigma)), \forall \theta. \quad (1.5)$$

In this situation, Petz’s work [Pet88, Pet86] showed that there is a canonical recovery map,

$$\mathcal{R}_{\sigma,\Phi}(\cdot) = \sigma^{\frac{1}{2}} \Phi^\dagger(\Phi(\sigma)^{-\frac{1}{2}} \cdot \Phi(\sigma)^{-\frac{1}{2}}) \sigma^{\frac{1}{2}}, \quad (1.6)$$

called Petz recovery map.

1.2. Failure of SLD QFI in characterizing sufficiency. A natural question is whether, similar to the classical case, the sufficiency of statistics can be determined based on SLD QFI. In other words, does $I_{\text{SLD},\rho}(\theta) = I_{\text{SLD},\Phi(\rho)}(\theta)$ imply that there exists a recovery channel \mathcal{R} such that $\mathcal{R} \circ \Phi(\rho_\theta) = \rho_\theta$? Surprisingly, it turns out that the answer is negative.

Proposition 1.1. *There exists a smooth family of full-rank qubit state ρ_θ and a quantum channel Φ ,*

$$I_{\text{SLD},\rho}(\theta) = I_{\text{SLD},\Phi(\rho)}(\theta), \forall \theta$$

²As it was noted in [BNG00], the measurement achieving this bound, in general, depends on the unknown parameter θ . [BNG00] shows how this measurement can be determined by consuming a sublinear number of copies.

³For two quantum states with density operators ρ and σ , $D(\rho \parallel \sigma) = \text{tr}(\rho \log \rho - \rho \log \sigma)$ if $\text{supp}(\rho) \subseteq \text{supp}(\sigma)$, and is infinite otherwise.

yet there does not exist a recovery channel \mathcal{R} , such that $\mathcal{R} \circ \Phi(\rho_\theta) = \rho_\theta$ for all θ .

Without the full-rank assumption, such non-recovery examples has been observed in [Mar22] (see [KS05, Pol13] for the counter-example to the full support assumption in the classical case). It was shown in [Mar22] that for any system A with density operator ρ_A and Hamiltonian H_A , there exists a purification $|\psi\rangle_{AB}$ and Hamiltonian H_B on the purifying system B , such that the SLD QFI for the family of pure states

$$|\psi(t)\rangle_{AB} = (e^{-iH_A t} \otimes e^{-iH_B t})|\psi\rangle_{AB} : t \in \mathbb{R}$$

is equal to the SLD QFI for the family of reduced density $\rho_A(t) = e^{-iH_A t} \rho_A e^{iH_A t}$, which can be obtained from the first family by discarding system B . However, despite preservation of SLD QFI under partial trace, it is impossible to recover the original state $|\psi(t)\rangle_{AB}$ from $\rho_A(t)$. Theorem 1.1 shows that this phenomenon also happens for smooth families of full-rank states, in contrast to the classical Fisher information (1.4).

1.3. Sufficiency via regular QFI metrics. The failure of SLD QFI in characterizing sufficient statistic motivates us to consider other quantum analogs of classical Fisher information that may satisfy this property. Indeed, quantum extensions of Fisher information in the context of information geometry have been intensively studied in [Pet96, PH96, LR99, PG11, Hay02, Hol11, Kos05, Pet02]. Viewing the space of all positive quantum states as a manifold, (1.3) induces a Riemannian metric, defined for any quantum state ρ as

$$\gamma_\rho(A) := \text{tr}(A \mathbb{J}_\rho(A)) , \quad (1.7)$$

for all traceless, Hermitian operators A , interpreted as tangent vectors at ρ . This metric is of special interests, as it is monotone under any quantum channel Φ ,

$$\gamma_\rho(A) \geq \gamma_{\Phi(\rho)}(\Phi(A)) . \quad (1.8)$$

Such an inequality is commonly called data processing inequality (DPI).

In the classical setting, it was proved by Čencov [Čen78] that the Fisher information metric is the unique Riemannian metric (up to scaling) satisfying the monotonicity (1.8). From this perspective, there are more than one quantum analog of the classical Fisher information whose corresponding metrics satisfy DPI. This family of metrics, called monotone metrics or quantum Fisher information (QFI) metrics, were first proposed by Čencov and Morozova [MC89], and later fully classified by Petz [Pet96]. It was observed by Lesniewski and Ruskai [LR99] that any QFI metric correspond to the Hessian of a given quantum f -divergence. In the special case where all the density operators in the family are diagonal in a fixed basis, QFI metrics all reduce to the classical Fisher information of the probability distribution defined by the eigenvalues of the density operators.

Interestingly, it turns out that SLD QFI in (1.3) is indeed the smallest QFI, which explains its special role in the Cramér–Rao bound. There is also a largest QFI, namely the Right-Logarithmic Derivative (RLD) Fisher information defined by

$$I_{\text{RLD},\rho}(\theta) = \text{tr}(\rho^{-1}(\theta)|\dot{\rho}(\theta)|^2) . \quad (1.9)$$

While SLD QFI fails to characterize sufficient statistics, our first main result shows that a large class of QFI metrics, which we call them “regular” metrics, characterize sufficiency (see Section 3 for the definition of regular QFI metrics).

Theorem 1.2. *Given a smooth family of quantum states $(\rho_\theta)_{\theta \in (a,b)}$ with full support, a quantum channel Φ is sufficient for θ if and only if*

$$I_\rho(\theta) = I_{\Phi(\rho)}(\theta) , \quad \forall \theta \in (a, b) ,$$

holds for all/any **regular** quantum Fisher information I . In particular, for any $\theta \in (a, b)$ in this family, the corresponding Petz recovery map of state ρ_θ defined in Eq. (1.6) recovers⁴ the full family, i.e.,

$$\mathcal{R}_{\rho_\theta, \Phi} \circ \Phi(\rho_\theta) = \rho_\theta, \quad \forall \theta \in (a, b).$$

The following well-known metrics are all regular QFI and therefore, according to our theorem, characterize sufficient statistic.

- a) Wigner-Yanase-Dyson (WYD) skew information: Given a density operator ρ and Hermitian operator H , the WYD skew information is defined as

$$W_H^{(\alpha)}(\rho) = -\frac{1}{2} \text{tr}([\rho^\alpha, H][\rho^{1-\alpha}, H]), \quad 0 < \alpha < 1. \quad (1.10)$$

$W_{\alpha, H}(\rho)$ is the QFI of the family $\rho_t = e^{-itH} \rho e^{itH}$, $t \in \mathbb{R}$ as the Petz-Rényi divergence $Q_\alpha(\rho, \sigma) = \text{tr}(\rho^{1-\alpha} \sigma^\alpha)$.

- b) x^α -metric: for $0 < \alpha < 1$,

$$I_{\alpha, \rho}(\theta) = \text{tr}(\dot{\rho}_\theta \rho^{-\alpha} \dot{\rho}_\theta \rho^{\alpha-1}),$$

for the special case $\alpha = \frac{1}{2}$, we call it symmetric inverse QFI.

- c) Bogoliubov-Kubo-Mori (BKM) Fisher information: BKM QFI is the negative Hessian of the relative entropy, defined as

$$I_{\text{BKM}, \rho}(\theta) := - \left. \frac{\partial^2}{\partial \theta_1 \partial \theta_2} D(\rho_{\theta_1} \| \rho_{\theta_2}) \right|_{\theta_1 = \theta_2 = \theta} \quad (1.11)$$

$$= \int_0^\infty \text{tr}(\dot{\rho}_\theta (\rho_\theta + r1)^{-1} \dot{\rho}_\theta (\rho_\theta + r1)^{-1}) dr, \quad (1.12)$$

is a regular QFI satisfying the theorem above.

In Section 1.6 we discuss implications of this result in the context of quantum thermodynamics and the resource theory of asymmetry. In conclusion, while to this date most applications of QFI in physics have been based on the special case of SLD QFI, our results clearly demonstrate operational and physical relevance of general QFI metrics, beyond this special case.

1.4. Approximate Recoverability. Over the last decade, a series of works established a stronger notion of recoverability, namely approximate recoverability [FR15, JRS⁺18, SFR16, SBT17, CV20a, GW21]. This line of research was initiated by a work of Fawzi and Renner [FR15] on approximate quantum Markov chains. The notion of approximate recoverability has found various applications in different areas of physics, including high energy physics and condensed matter theory [CHP⁺19, HPS21, HP19]. A notable result is by Junge *et al* [JRS⁺18], who proved that

$$D(\rho \| \sigma) - D(\Phi(\rho) \| \Phi(\sigma)) \geq -2 \log F(\rho, \mathcal{R}_{\sigma, \Phi}^{\text{uni}} \circ \Phi(\rho)) \geq \|\rho - \mathcal{R}_{\sigma, \Phi}^{\text{uni}} \circ \Phi(\rho)\|_1^2, \quad (1.13)$$

where $F(\rho, \sigma) := \|\sqrt{\rho} \sqrt{\sigma}\|_1$ is the fidelity between two quantum states ρ, σ and $\|\cdot\|_1$ denotes the trace norm. Here $\mathcal{R}_{\sigma, \Phi}^{\text{uni}}$ is called the universal recovery map given by the integral

$$\mathcal{R}_{\sigma, \Phi}^{\text{uni}} = \int_{\mathbb{R}} \mathcal{R}_{\sigma, \Phi}^{\frac{t}{2}} d\beta(t), \quad d\beta(t) = \frac{\pi}{2(\cosh(\pi t) + 1)} dt \quad (1.14)$$

where $\mathcal{R}_{\sigma, \Phi}^t$ is the rotated version of Petz Recovery map,

$$\mathcal{R}_{\sigma, \Phi}^t(\cdot) = \sigma^{-it} \mathcal{R}_{\sigma, \Phi}(\Phi(\sigma)^{it} \cdot \Phi(\sigma)^{-it}) \sigma^{it}. \quad (1.15)$$

⁴As we show in Theorem 5.1, this result also holds for rotated Petz maps.

Despite the progress on approximate recoverability via entropic quantities, it remains open whether it can be characterized using QFI. Our second main result address this question.

Theorem 1.3. *Suppose $\rho_\theta \geq \lambda I$ for some $\lambda > 0$ and all $\theta \in (a, b)$, then for any $s < r$*

$$\lambda^{-\frac{1}{2}} \int_s^r \sqrt{I_{\text{BKM}, \rho}(\theta) - I_{\text{BKM}, \Phi(\rho)}(\theta)} d\theta \geq D(\rho_r \| \rho_s) - D(\Phi(\rho_r) \| \Phi(\rho_s)).$$

Combining this with the existing results on approximate recoverability in terms of relative entropy, such as Eq.(1.13), one can obtain recovery bounds in terms of BKM metric.

1.5. Quantum χ^2 divergence. The proof of the above (approximate) recoverability results follows from a simpler setting, namely that of quantum χ^2 divergence. The classical χ^2 divergence of two distributions P and Q is defined by

$$\chi^2(P, Q) = \mathbb{E}_Q \left| \frac{dP}{dQ} - 1 \right|^2. \quad (1.16)$$

For two quantum states ρ, σ and a given QFI metric γ , the quantum analog of χ^2 divergence is

$$\chi^2(\rho, \sigma) = \gamma_\sigma(\rho - \sigma), \quad (1.17)$$

which can be understood as the QFI for the linear interpolation family $\rho_t = t\sigma + (1-t)\rho$. In the special case where ρ and σ commute this quantity reduces to the classical χ^2 divergence in Eq.(1.16) for the distributions defined by the eigenvalues of ρ and σ . The above definition based on QFI metrics guarantees that the χ^2 divergence associated to each QFI metric inherits the nice properties of the metric, such as monotonicity under DPI. These χ^2 divergences have found applications in characterizing the mixing time of a quantum Markov process [TKR⁺10, GR22].

It is natural to ask whether the recoverability or the stronger notion of approximate recoverability can be characterized with quantum χ^2 divergences or the corresponding monotone metric γ , as Hessians of quantum f -divergence. Again, the answer can be negative or positive depending on the choice of the metric γ . In the case of regular quantum χ^2 divergences, i.e., those corresponding to regular QFI we show that

Theorem 1.4. *Given two states ρ and σ with $\text{supp}(\rho) \subseteq \text{supp}(\sigma)$, a quantum channel Φ is sufficient for $\{\rho, \sigma\}$ if and only if*

$$\chi^2(\rho, \sigma) = \chi^2(\Phi(\rho), \Phi(\sigma)),$$

*holds for any/all **regular** quantum χ^2 -divergence. Moreover, if $\sigma > \lambda I$, there exists an explicit constant $K(\chi^2, \lambda)$ depending on χ^2 and λ such that for all $t \in \mathbb{R}$,*

$$\chi^2(\rho, \sigma) - \chi^2(\Phi(\rho), \Phi(\sigma)) \geq \frac{\pi}{\cosh(\pi t)} K(\chi^2, \lambda) \|\rho - \mathcal{R}_{\sigma, \Phi}^t(\rho)\|_1^2,$$

where $\mathcal{R}_{\sigma, \Phi}^t$ is the rotated Petz map defined in (1.15).

On the other hand, the above property does not generally hold for non-regular χ^2 -divergences, even though they still satisfy DPI inequality. Two notable examples are the SLD and RLD χ^2 -divergences (\mathbb{J}_σ is as in (1.3))

$$\chi_{\text{SLD}}^2(\rho, \sigma) = \text{tr}(\mathbb{J}_\sigma(\rho)^2 \sigma) - 1, \quad \chi_{\text{RLD}}^2(\rho, \sigma) = \text{tr}(\rho^2 \sigma^{-1}) - 1.$$

In the special case of symmetric inverse metric

$$\chi_{\frac{1}{2}}^2(\rho, \sigma) = \text{tr}((\rho - \sigma)\sigma^{-1/2}(\rho - \sigma)\sigma^{-1/2}) = \text{tr}(\rho\sigma^{-\frac{1}{2}}\rho\sigma^{-\frac{1}{2}}) - 1,$$

which is related to the Sandwiched 2-Rényi relative entropy $D_2(\rho\|\sigma) = \log \text{tr}(\rho\sigma^{-\frac{1}{2}}\rho\sigma^{-\frac{1}{2}})$. The $\chi_{\frac{1}{2}}^2$ divergence is known to enjoy some special properties (see [CL19]). Here, we achieve a universal approximate recovery bound.

Theorem 1.5. *For two states ρ and σ ,*

$$\chi_{\frac{1}{2}}^2(\rho, \sigma) - \chi_{\frac{1}{2}}^2(\Phi(\rho), \Phi(\sigma)) \geq \|\rho - \mathcal{R}_{\sigma, \Phi} \circ \Phi(\rho)\|_1^2,$$

where $\mathcal{R}_{\sigma, \Phi}$ is the Petz recovery map defined in (1.6).

The above bound improves the result of [CS22] by removing a state dependent constant $\|\sigma^{-1}\|$, which can be unbounded in infinite dimensions. It also applies to classical Fisher information as all quantum χ^2 -divergence reduces to the classical χ^2 -divergence in the commutative setting. We also note that here our recovery map $\mathcal{R}_{\sigma, \Phi}$ is the original Petz map, while it remains open whether the recovery bound (1.13) for relative entropy can be achieved with $\mathcal{R}_{\sigma, \Phi}$.

1.6. Applications: Quantum Thermodynamics and the resource theory of asymmetry. An important application of QFI is in the context of quantum thermodynamics and the closely-related resource theory of asymmetry. QFI metrics provide a useful way of quantifying the amount of coherence of a system with respect to its energy eigenbasis and, more generally, the amount of asymmetry (symmetry-breaking) of the system with respect to a given symmetry group [Mar12, MS14, Gir14, YV16, KJJ⁺18, Mar22]. Below we illustrate the application of our results for coherence, and refer to Section 6 for the more general setting of asymmetry.

For a system with density operator ρ and Hamiltonian H , consider the family of time evolution of this system, namely states $\rho(t) = e^{-iHt}\rho e^{iHt}$ for $t \in \mathbb{R}$. For any QFI metric γ , the QFI of this family with respect to the time parameter t is time-independent, that is, for any $t \in \mathbb{R}$

$$I_\rho(t) = \gamma_{\rho(t)}(\dot{\rho}(t), \dot{\rho}(t)) = \gamma_\rho(i[H, \rho], i[H, \rho]) := I_H(\rho). \quad (1.18)$$

This quantity determines the asymmetry of the system with respect to the time translation symmetry, or equivalently, the energetic coherence of the system with respect to the eigenbasis of Hamiltonian H . In particular, $I_H(\rho) = 0$ if and only if $[\rho, H] = 0$, namely, the state is diagonal in the energy-eigenbasis.

An important and useful property of this function is its monotonicity under any quantum operation \mathcal{E} (CPTP map) that respects the covariance condition

$$\mathcal{E}(e^{-iH_{\text{in}}t}(\cdot)e^{iH_{\text{in}}t}) = e^{-iH_{\text{out}}t}\mathcal{E}(\cdot)e^{iH_{\text{out}}t}, \quad \forall t \in \mathbb{R}, \quad (1.19)$$

where H_{in} and H_{out} are Hamiltonians for the input and output system, respectively. Operations satisfying this property are sometimes called time-translation invariant operations. This property implies that under channel \mathcal{E} the family of states $e^{-iH_{\text{in}}t}\rho e^{iH_{\text{in}}t}$ is mapped to $e^{-iH_{\text{out}}t}\mathcal{E}(\rho)e^{iH_{\text{out}}t}$. Then, the DPI for QFI metrics immediately implies that I_H is monotone under any such map \mathcal{E} ,

$$I_{H_{\text{out}}}(\mathcal{E}(\rho)) \leq I_{H_{\text{in}}}(\rho).$$

Any function satisfying this monotonicity is called a measure of asymmetry with respect to the symmetry group under consideration, which in this case is the time translation symmetry.⁵

⁵It is often also required that a measure of asymmetry should vanish for all states that are invariant under the action of symmetry, which in this case are states satisfying $e^{-iHt}\rho e^{iHt} = \rho$ for all $t \in \mathbb{R}$. It can be easily seen that function I_H satisfies this property.

While all QFI metrics can be used to quantify asymmetry and coherence, recent work [Mar22] has singled out SLD QFI, as the measure of asymmetry with an operational interpretation: namely, it quantifies the *coherence cost* of preparing a general mixed state from pure coherent states. RLD QFI has also shown to be useful for characterizing the distillation of energetic coherence, i.e., time-translation asymmetry [Mar20].

The present work reveals that, in addition to SLD and RLD QFI, regular QFI metrics are particularly useful for quantifying asymmetry and energetic coherence. In particular, they can determine whether the resourcefulness of the system has been degraded by noise or any process that respects the symmetry. We show that

Theorem 1.6. *Consider systems A and B with Hamiltonians H_A and H_B , respectively. Let $\rho \in \mathcal{B}(\mathcal{H}_A)$ be a full-rank density operator and let $\mathcal{E} : \mathcal{B}(\mathcal{H}_A) \rightarrow \mathcal{B}(\mathcal{H}_B)$ be time-translation invariant quantum channel. Then, there exists a time-translation invariant channel \mathcal{R} such that $\mathcal{R}(\mathcal{E}(\rho)) = \rho$, if and only if*

$$I_{H_A}(\rho) = I_{H_B}(\mathcal{E}(\rho)) , \quad (1.20)$$

where I_H is the QFI defined in Eq.(1.18) with respect to some/all regular QFI metric γ .

An important example of regular QFI metric is Wigner-Araki-Yanase skew information

$$W_H^{(\alpha)}(\rho) := \text{Tr}(\rho H^2) - \text{Tr}(\rho^{1-\alpha} H \rho^\alpha H) , \quad (1.21)$$

for $0 < \alpha < 1$, which have been previously studied as a measure of coherence and asymmetry [Mar12, MS14, Tak19]. It is worth mentioning that the above property of regular QFI metrics, that their conservation implies reversibility, was shown [ML16] to hold for the relative entropy of asymmetry (also known as the asymmetry).⁶

Theorem 1.6 follows from applying Theorem 1.2 for the family $\rho(t) = e^{-iH_A t} \rho e^{iH_A t}$ with $t \in \mathbb{R}$. By the time-translation invariance of \mathcal{E} , $\mathcal{E}(\rho(t)) = e^{-iH_B t} \mathcal{E}(\rho) e^{iH_B t} : t \in \mathbb{R}$. Furthermore, since QFI $I_\rho(t) = I_{H_A}(\rho)$ is time-independent, Eq.(1.20) implies that the QFI preserves under $I_\rho(t) = I_{H_A}(\rho) = I_{H_B}(\rho) = I_{\mathcal{E}(\rho)}(t)$ for all $t \in \mathbb{R}$. Then, our Theorem 1.2 implies that the Petz recovery of ρ recovers the states for all time t , i.e.,

$$\mathcal{R}_{\rho, \mathcal{E}}(\mathcal{E}(\rho(t))) = \rho(t) , \quad \forall t \in \mathbb{R} \quad (1.22)$$

Furthermore, for any finite $T > 0$ the time-averaged version of Petz map

$$\mathcal{R}_{\text{avg}, T}(\cdot) = \frac{1}{T} \int_{-T/2}^{T/2} dt e^{iH_A t} \mathcal{R}_{\rho, \mathcal{E}}(e^{-iH_B t}(\cdot) e^{iH_B t}) e^{-iH_A t} , \quad (1.23)$$

also satisfies (1.22) (Note that $\rho(t+s) = e^{iH_A t} \rho(s) e^{-iH_A t}$ and similar for $\mathcal{E}(\rho(t))$). In general, $\mathcal{R}_{\text{avg}, T}$ is not covariant. Nevertheless, since for finite-dimensional Hilbert space \mathcal{H}_A and \mathcal{H}_B , the set of CPTP maps is compact, there exists a limit point $\mathcal{R}_{\text{avg}, \infty} = \lim_{n \rightarrow \infty} \mathcal{R}_{\text{avg}, T_n}$. It is straightforward to show that $\mathcal{R}_{\text{avg}, \infty}$ satisfies the time translation invariant condition Eq.(1.19). This proves the Theorem 1.6. We refer to Section 6 for the more general cases of asymmetry of compact Lie groups.

Outline of the rest of the paper. We first discuss the non-recoverability of SLD and RLD QFI metric in the Section 2. We briefly review in Section 3 the definitions of general quantum monotone metrics, Fisher information and χ^2 divergences. Section 4 is devoted to the recoverability of regular χ^2 divergences (Theorem 1.4) and the universal recoverability bound for $\chi_{1/2}^2$ (Theorem 1.4). Based on that we prove the recoverability

⁶Unlike metrics is not additive in tensor-product states, the relative entropy of asymmetry grow logarithmically with the number of copies.

(Theorem 1.2) and approximate recoverability (Theorem 1.3) of regular QFI in Section 5. Section 6 discusses the application of our results in quantum coherence and asymmetry.

Acknowledgement. LG is partially supported by NSF grant DMS-2154903. IM is supported by NSF grants FET-1910571, Phy-2046195, FET-2106448. C.R. acknowledges the support of the Munich Center for Quantum Sciences and Technology, funding from the Humboldt Foundation as well as by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germanys Excellence Strategy EXC-2111 390814868. H.L. acknowledges support by the DFG cluster of excellence 2111 (Munich Center for Quantum Science and Technology).

Notations. We write \mathbb{M}_n for the set of $n \times n$ complex matrices. Given a finite dimensional Hilbert space \mathcal{H} , we denote $\mathcal{B}(\mathcal{H})$, $\mathcal{B}(\mathcal{H})_{\text{sa}}$ and $\mathcal{B}(\mathcal{H})_+$ as the set of bounded, Hermitian and positive (semi-definite) operators respectively. We write $\langle A, B \rangle = \text{tr}(A^*B)$ for the Hilbert-Schmidt inner product, where tr stands for the standard matrix trace. The Schatten norm of order $p \geq 1$ is defined as $\|A\|_p := \text{tr}(|A|^p)^{1/p}$ and $\mathcal{S}_p(\mathcal{H})$ denotes the Schatten- p space. We denote by $\mathcal{D}(\mathcal{H})$ the subset of density operators (positive semi-definite and trace 1) on \mathcal{H} , $\mathcal{D}_+(\mathcal{H})$ by the subset of invertible density operators on \mathcal{H} . We use I for the identity operator in $\mathcal{B}(\mathcal{H})$ and id for the identity map on $\mathcal{B}(\mathcal{H})$. We write A^* as the adjoint of an operator A and Φ^\dagger as the adjoint of a map Φ with respect to Hilbert-Schmidt inner product. Given two finite dimensional Hilbert spaces \mathcal{H} and \mathcal{K} , a quantum channel $\Phi : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{K})$ is a completely positive trace preserving map. In particular, $\Phi(\mathcal{D}(\mathcal{H})) \subset \mathcal{D}(\mathcal{K})$ preserves the density operators.

2. INSUFFICIENT STATISTIC PRESERVING SLD QUANTUM FISHER INFORMATION

In this section, we present the counter-example that quantum sufficiency cannot be characterized via SLD QFI. To show this it is useful to compare SLD and RLD QFI. For a family of states ρ_θ , let $\mathbb{J}_{\rho_\theta}^{-1}(\dot{\rho}_\theta) = L_\theta$. Recall that

$$I_{\text{RLD},\rho}(\theta) = \text{tr}(\rho_\theta^{-1}|\dot{\rho}_\theta|^2), \quad I_{\text{SLD},\rho}(\theta) = \text{tr}(\dot{\rho}_\theta \mathbb{J}_{\rho_\theta}(\dot{\rho}_\theta) = \text{tr}(L_\theta^2 \rho_\theta).$$

Note that I_{RLD} is infinite whenever $\text{supp}(|\dot{\rho}_\theta|) \not\subseteq \text{supp}(\rho_\theta)$, but I_{SLD} can be finite as long as $\text{supp}(\dot{\rho}_\theta)^\perp \subseteq \text{supp}(\rho_\theta)^\perp$. Moreover, RLD QFI can be rewritten as

$$I_{\text{RLD},\rho}(\theta) = \text{tr}(\rho_\theta^{-1}|\dot{\rho}_\theta|^2) = \frac{1}{4} \text{tr} \left(\rho^{-1} (L\rho + \rho L)^2 \right) \quad (2.1)$$

$$= \frac{3}{4} I_{\text{SLD},\rho}(\theta) + \frac{1}{4} \text{tr}(\rho^{-1} L \rho^2 L), \quad (2.2)$$

where to simplify the notation we have omitted the parameter θ in the second equation. We conclude that

$$I_{\text{RLD},\rho}(\theta) - I_{\text{SLD},\rho}(\theta) = \frac{1}{4} P_{L_\theta}(\rho_\theta), \quad (2.3)$$

where the quantity

$$P_L(\rho) = \text{tr}(\rho^{-1} L \rho^2 L) - \text{Tr}(\rho L^2) = -\text{tr}(\rho^{-1} [\rho, L]^2), \quad (2.4)$$

is indeed the RLD QFI for the family of states $e^{itL} \rho e^{-itL}$ with respect to parameter t , called the *purity of coherence* of ρ with respect to L [Mar20]. In particular, $P_L(\rho) \geq 0$ and is zero if and only if $[\rho, L] = 0$ commute.

Fix a parameter value θ_o and let \mathcal{L}_o be the pinching map that dephases its input with respect to the spectrum of ρ_{θ_o} . Define $\sigma_\theta = \mathcal{L}_o(\rho_\theta)$. Note that

$$\dot{\sigma}_\theta = \mathcal{L}_o(\dot{\rho}_\theta) = \frac{1}{2}\mathcal{L}_o(L_{\theta_o}\rho_{\theta_o} + \rho_{\theta_o}L_{\theta_o}) = \frac{1}{2}(L_{\theta_o}\sigma_{\theta_o} + \sigma_{\theta_o}L_{\theta_o})$$

Thus, L_o remain the SLD for σ_θ at $\theta = \theta_o$ and the SLD QFI at θ_o does not change under this map, i.e.,

$$I_{\text{SLD},\sigma}(\theta_o) = \text{tr}(L_{\theta_o}^2 \sigma_{\theta_o}) = \text{tr}(L_{\theta_o}^2 \rho_{\theta_o}) = I_{\text{SLD},\rho}(\theta_o) . \quad (2.5)$$

Furthermore, the state σ_{θ_o} commutes with L_{θ_o} and therefore $P_{L_{\theta_o}}(\sigma_{\theta_o}) = 0$, which means the gap between RLD and SLD QFI vanishes, i.e.,

$$I_{\text{RLD},\sigma}(\theta_o) = I_{\text{SLD},\sigma}(\theta_o) = I_{\text{SLD},\rho}(\theta_o) . \quad (2.6)$$

Note that unless $[L_{\theta_o}, \rho_{\theta_o}] = 0$, then $I_{\text{SLD},\rho}(\theta_o) < I_{\text{RLD},\rho}(\theta_o)$ and hence \mathcal{L}_o is not sufficient because $I_{\text{RLD},\sigma}(\theta_o) < I_{\text{RLD},\rho}(\theta_o)$. In summary, we observe

Proposition 2.1. *For a family of states ρ_θ , the difference between RLD and SLD QFI equal to $P_{L_\theta}(\rho_\theta)$, which is zero if and only if $[L_\theta, \rho_\theta] = 0$, where L_θ is the symmetric logarithm derivative of ρ_θ .*

Fix a parameter value θ_o , the dephasing map \mathcal{L}_o relative to the eigen-subspaces of L_{θ_o} always preserved SLD QFI at θ_o , but strictly decrease RLD QFI if $[L_{\theta_o}, \rho_{\theta_o}] \neq 0$, hence not recoverable.

In general, because SLD operator L_{θ_o} depends on the parameter o , so does the map \mathcal{L}_o . However, as we will see in the following example, it is possible to have a family of states ρ_θ such that the SLD operator L_{θ_o} share the same eigen-subspaces, which means the dephasing map $\mathcal{L}_\theta = \mathcal{L}$ is independent of θ . Furthermore, this family can be chosen to be full-rank. Such a family satisfies the full-rank condition in Theorem 1.2 while SLD QFI remains conserved under the map \mathcal{L} , but its conservation does not imply sufficiency of the output statistic. We emphasize that this contrasts with the classical case, where conservation of Fisher information for probability distributions with full-support implies sufficiency [Sch12, Theorem 2.8].

Qubit example: Consider the family of qubit density operators

$$\rho_\theta = \begin{pmatrix} p(\theta) & \epsilon \times r(\theta) \\ \epsilon \times r(\theta) & 1 - p(\theta) \end{pmatrix} : \quad \theta \in [a, b] ,$$

defined for $\theta \in [a, b]$, where $p : [a, b] \rightarrow (0, 1)$ is an arbitrary function with finite, non-zero derivative $\dot{p}(\theta)$. The function r is determined by p via equation

$$r(\theta) = \exp \int_a^\theta \frac{\dot{p}(s)(1 - 2p(s))}{2p(s)(1 - p(s))} ds , \quad (2.7)$$

and ϵ is chosen such that

$$0 < \epsilon^2 < \min_{\theta \in [a, b]} \frac{p(\theta)(1 - p(\theta))}{r^2(\theta)} . \quad (2.8)$$

The latter condition guarantees that ρ_θ is positive and full-rank for all $\theta \in [a, b]$.

Then, one can easily check that the Hermitian operator

$$L_\theta = \begin{pmatrix} \dot{p}(\theta)/p(\theta) & 0 \\ 0 & -\dot{p}(\theta)/(1 - p(\theta)) \end{pmatrix} ,$$

satisfies the equation

$$\dot{\rho}_\theta = \frac{1}{2} [\rho_\theta L_\theta + L_\theta \rho_\theta] , \quad (2.9)$$

and therefore is the SLD of the family ρ_θ . Note that L_θ is always diagonal. Furthermore, the assumption that $\dot{p}(\theta) \neq 0$ implies that $L(\theta)$ is non-degenerate. Let \mathcal{L} be the dephasing map in $\{|0\rangle, |1\rangle\}$ basis. Applying this map to state ρ_θ we obtain the family of states

$$\sigma_\theta = \mathcal{L}(\rho_\theta) = \begin{pmatrix} p(\theta) & 0 \\ 0 & 1 - p(\theta) \end{pmatrix} : \theta \in [a, b].$$

Under this dephasing, SLD QFI remains conserved. However, because the original family ρ_θ is not diagonal in $\{|0\rangle, |1\rangle\}$ basis, RLD QFI for the family of states σ_θ is strictly less than the RLD QFI for the family of states ρ_θ . Therefore, even though SLD QFI is preserved under \mathcal{L} , the RLD QFI decreases strictly. Hence $\sigma_\theta = \mathcal{L}(\rho_\theta)$ is not a sufficient statistic for the original family ρ_θ .

3. MONOTONE METRICS

The faithful state space $\mathcal{D} := \mathcal{D}_+(\mathcal{H})$ is a submanifold in the real Euclidean space $\mathcal{B}(\mathcal{H})$. At each point $\rho \in \mathcal{D}$, the tangent space

$$T_\rho \mathcal{D} = \{A \in \mathcal{B}(\mathcal{H}) \mid A = A^*, \text{tr}(A) = 0\}$$

is the subspace of traceless Hermitian operators. A Riemannian metric on \mathcal{D} is a smooth assignment $\rho \mapsto \gamma_\rho$ to a positive bilinear form $\gamma_\rho : T_\rho \mathcal{D} \times T_\rho \mathcal{D} \rightarrow \mathbb{R}$.

Definition 3.1. *We say a Riemannian metric γ is a monotone metric, if for any quantum channel $\Phi : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{K})$,*

$$\gamma_\rho(A, A) \geq \gamma_{\Phi(\rho)}(\Phi(A), \Phi(A)), \quad \forall A \in T_\rho \mathcal{D}.$$

We will use the short notation $\gamma_\rho^g(A) := \gamma_\rho^g(A, A)$.

The monotone metrics are classified by operator anti-monotone (i.e. decreasing) functions $g : (0, \infty) \rightarrow (0, \infty)$. Given $\rho \in \mathcal{D}$, we define

$$\mathbb{J}_\rho^g = g(L_\rho R_\rho^{-1}) R_\rho^{-1},$$

where $L_\rho(A) = \rho A$, $R_\rho(A) = A\rho$ are left and right multiplications respectively. Based on the work of Chentsov and Morozova [MC89], Petz in [Pet96] proved that every monotone metric admits the following form

$$\gamma_\rho^g(A, B) = \langle A, \mathbb{J}_\rho^g(B) \rangle = \langle A\rho^{-\frac{1}{2}}, g(L_\rho R_\rho^{-1})(B\rho^{-\frac{1}{2}}) \rangle. \quad (3.1)$$

If $\rho = \sum_i \lambda_i |\phi_i\rangle\langle\phi_i|$, the metric is explicit with matrix coefficients

$$\gamma_\rho(A, A) = \sum_{i,j} c(\lambda_i, \lambda_j) |\langle\phi_i|A|\phi_j\rangle|^2$$

where $c(x, y) = y^{-1}g(xy^{-1})$ is called Morozova-Chentsov function. If we assume that the definition of γ_ρ^g for $A, B \in \mathcal{B}(\mathcal{H})$ has the symmetric property

$$\gamma_\rho(A, B) = \gamma_\rho(B^*, A^*),$$

this corresponds to operator anti-monotone functions satisfying

$$g(x^{-1}) = xg(x), \quad g(1) = 1. \quad (3.2)$$

These \mathbb{J}_ρ^g operators are inverses to operator means between L_ρ and R_ρ , and admit the following integral form (see [LR99])

$$\mathbb{J}_\rho^g = R_\rho^{-1} \int_0^\infty \frac{1}{s + \Delta_\rho} \nu_g(s) ds \quad (3.3)$$

where $\Delta_\rho = L_\rho R_\rho^{-1}$ is the relative modular operator and $\nu_g(s) ds$ is a finite positive measure satisfying $\nu_g(s^{-1}) = s\nu_g(s)$.

We will encounter the following special cases in our discussion.

Example 3.2. (1) SLD metric: for $g(x) = \frac{2}{x+1}$ and $\nu_g = 2\delta_1$ being the point mass at 1,

$$\gamma_\rho^{\text{SLD}}(A, B) = 2 \operatorname{tr}(A^*(L_\rho + R_\rho)^{-1}B). \quad (3.4)$$

γ_ρ^{SLD} is also called Bures metric in the literature.

(2) RLD metric: for $\nu_g = \delta_0$ being point mass at $s = 0$

$$\gamma_\rho^{\text{RLD}}(A, B) = \frac{1}{2} \operatorname{tr}(A^* B \rho^{-1}) \quad (3.5)$$

This corresponds to the RLD Fisher information introduced in (1.9).

(3) BKM metric: for $g(x) = \frac{\log x}{x-1}$ and $\nu_g(s) = \frac{1}{s+1}$,

$$\gamma_\rho^{\text{BKM}}(A, B) = \int_0^\infty \operatorname{tr}(A^*(\rho + s1)^{-1}B(\rho + s1)^{-1}) ds. \quad (3.6)$$

(4) x^α -metrics: for $g(x) = \frac{1}{2}(x^{-\alpha} + x^{\alpha-1})$, $\alpha \in (0, 1)$ and $\nu_g(s) = \frac{\sin \pi s}{2\pi}(s^{-\alpha} + s^{\alpha-1})$,

$$\gamma_\rho^{(\alpha)}(A, B) = \operatorname{tr}(A^* \rho^{-\alpha} B \rho^{\alpha-1}). \quad (3.7)$$

A special case is $\alpha = \frac{1}{2}$, which we write as

$$\gamma_\rho^s(A, B) := \operatorname{tr}(A^* \rho^{-\frac{1}{2}} B \rho^{-\frac{1}{2}}). \quad (3.8)$$

(5) WYD metric: for $\alpha \in (0, 1)$, $g(x) = \frac{(1-x^\alpha)(1-x^{1-\alpha})}{\alpha(1-\alpha)(1-x^2)}$ and $\nu_g(s) = \frac{\sin(\pi s)}{\pi}(1+s)^{(\alpha-2)} + \frac{\sin(\pi s^{-1})}{s\pi}(1+s^{-1})^{(\alpha-2)}$,

$$\gamma_\rho^{\text{WYD}}(A, B) = \frac{\partial^2}{\partial s \partial t} \operatorname{tr}((\rho + sA)^\alpha (\rho + tB)^{1-\alpha})|_{s=t=0} \quad (3.9)$$

This gives the WYD skew information as in Eq. (1.10).

It is clear from functional calculus that if $g_1 \leq g_2$, then

$$\gamma_\rho^{g_1}(A, A) \leq \gamma_\rho^{g_2}(A, A), \quad \forall A \in \mathcal{B}(\mathcal{H}).$$

From this perspective, for any monotone metric γ^g ,

$$\gamma_\rho^{\text{Bures}}(A, A) \leq \gamma_\rho^g(A, A) \leq \gamma_\rho^{\text{R}}(A, A)$$

and the $\frac{1}{2}$ -metric is the smallest among all α -metrics.

Definition 3.3. We say a monotone metric γ_g or its associated operator anti-monotone function g is regular if the Lebesgue measure ds is absolutely continuous with respect to $\nu_g(s) d(s)$.

The metric γ^{BKM} , $\gamma^{(\alpha)}$ and γ^{WYD} are regular but $\gamma_\rho^{\text{Bures}}$ and γ_ρ^{RLD} are not. We will see later that this is the reason for the non-recoverability for the latter two.

We note that for a general density operator ρ , the monotone metric γ^g are well-defined and finite for A with $s(A) \leq s(\rho)$, where $s(\rho)$ is the support of ρ . For example, the RLD metric $\gamma_\rho^{\text{RLD}}(A) = +\infty$ as long as $s(A) \not\leq s(\rho)$. If in addition, $\lim_{x \rightarrow 0^+} g(x)$ exists and finite, γ^g is also finite for self-adjoint A if $(1 - s(\rho))A(1 - s(\rho)) = 0$. This is the case for SLD metric.

4. RECOVERY VIA MONOTONE METRICS

4.1. Exact Recovery via monotone metrics. Let γ^g be a monotone metric. For two quantum states $\rho, \sigma \in \mathcal{D}(\mathcal{H})$, the quantum χ^2 divergence is

$$\chi_g^2(\rho, \sigma) := \gamma_\sigma^g(\rho - \sigma). \quad (4.1)$$

It satisfies the data processing inequality: for any quantum channel Φ ,

$$\chi_g^2(\rho, \sigma) \geq \chi_g^2(\Phi(\rho), \Phi(\sigma)).$$

The main goal of this section is to prove that for a regular g ,

$$\chi_g^2(\rho, \sigma) = \chi_g^2(\Phi(\rho), \Phi(\sigma)) \iff \rho = \mathcal{R}_{\sigma, \Phi}^t \circ \Phi(\rho)$$

for any/all $t \in \mathbb{R}$, where $\mathcal{R}_{\sigma, \Phi}^t$ is the rotated Petz map of σ and channel Φ :

$$\mathcal{R}_{\sigma, \Phi}^t(A) = \sigma^{\frac{1}{2}-it} \Phi^\dagger(\Phi(\sigma)^{-\frac{1}{2}+it} A \Phi(\sigma)^{-\frac{1}{2}-it}) \sigma^{\frac{1}{2}+it}. \quad (4.2)$$

Indeed, we will show a quantitative version of the above result. Our argument is inspired from [CV20b] and [GW21].

We start with a lemma on the Stinespring dilation of the rotated Petz map.

Lemma 4.1. *Let $\rho \in \mathcal{D}_+(\mathcal{H})$ and $\Phi : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{K})$ be a quantum channel. For any $t \in \mathbb{R}$, we define the linear map*

$$V_{\rho, t} : \mathcal{B}(\mathcal{K}) \rightarrow \mathcal{B}(\mathcal{H}), \quad V_{\rho, t}(A) = \Phi^\dagger(A \Phi(\rho)^{-\frac{1}{2}-it}) \rho^{\frac{1}{2}+it},$$

Then

- i) $V_{\rho, t}^* V_{\rho, t}$ is a contraction on $\mathcal{S}_2(\mathcal{H})$, i.e. $V_{\rho, t}^* V_{\rho, t} \leq I$.
- ii) $V_{\rho, t}^* \Delta_\rho V_{\rho, t} \leq \Delta_{\Phi(\rho)}$ as positive operators on $\mathcal{S}_2(\mathcal{H})$

Proof. For any $A \in \mathcal{B}(\mathcal{H})$, we have

$$\begin{aligned} \langle A, V_{\rho, t}^* V_{\rho, t}(A) \rangle &= \langle V_{\rho, t}(A), V_{\rho, t}(A) \rangle \\ &= \langle \Phi^\dagger(A \Phi(\rho)^{-\frac{1}{2}-it}) \rho^{\frac{1}{2}+it}, \Phi^\dagger(A \Phi(\rho)^{-\frac{1}{2}-it}) \rho^{\frac{1}{2}+it} \rangle \\ &= \text{tr} \left(\rho^{\frac{1}{2}-it} \Phi^\dagger(\Phi(\rho)^{-\frac{1}{2}+it} A^*) \Phi^\dagger(A \Phi(\rho)^{-\frac{1}{2}-it}) \rho^{\frac{1}{2}+it} \right) \\ &\leq \text{tr} \left(\rho \Phi^\dagger \left(\Phi(\rho)^{-\frac{1}{2}+it} A^* A \Phi(\rho)^{-\frac{1}{2}-it} \right) \right) \\ &= \langle \Phi(\rho), \Phi(\rho)^{-\frac{1}{2}+it} A^* A \Phi(\rho)^{-\frac{1}{2}-it} \rangle \\ &= \langle A, A \rangle, \end{aligned}$$

where the above inequality follows from the operator Schwarz inequality

$$\Phi^\dagger(X^*) \Phi^\dagger(X) \leq \Phi^\dagger(X^* X), \quad \forall X \in \mathcal{B}(\mathcal{H}).$$

Similarly,

$$\begin{aligned} \langle A, V_{\rho, t}^* \Delta_\rho V_{\rho, t}(A) \rangle &= \langle V_{\rho, t}(A), \Delta_\rho V_{\rho, t}(A) \rangle \\ &= \langle \Phi^\dagger(A \Phi(\rho)^{-\frac{1}{2}-it}) \rho^{\frac{1}{2}+it}, \rho \Phi^\dagger(A \Phi(\rho)^{-\frac{1}{2}-it}) \rho^{\frac{1}{2}+it} \rho^{-1} \rangle \\ &= \text{tr} \left(\Phi^\dagger(\Phi(\rho)^{-\frac{1}{2}+it} A^*) \rho \Phi^\dagger(A \Phi(\rho)^{-\frac{1}{2}-it}) \right) \\ &= \text{tr} \left(\rho \Phi^\dagger(A \Phi(\rho)^{-\frac{1}{2}-it}) \Phi^\dagger(\Phi(\rho)^{-\frac{1}{2}+it} A^*) \right) \\ &\leq \text{tr} \left(\rho \Phi^\dagger(A \Phi(\rho)^{-1} A^*) \right) \\ &= \langle A, \Delta_{\Phi(\rho)}(A) \rangle, \end{aligned}$$

where again the inequality follows from the operator Schwarz inequality. ■

Our next lemma is a modification of [CV20b, Lemma 2.1].

Lemma 4.2. *Let $\Delta : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$, $\tilde{\Delta} : \mathcal{B}(\mathcal{K}) \rightarrow \mathcal{B}(\mathcal{K})$, $V : \mathcal{B}(\mathcal{K}) \rightarrow \mathcal{B}(\mathcal{H})$ be positive linear maps and V^*V be a contraction. If $\tilde{\Delta} \geq V^*\Delta V$ as positive operators on $\mathcal{S}_2(\mathcal{H})$, then for any $s \geq 0$ and $h \in \mathcal{B}(\mathcal{H})$,*

$$\langle h, (s + \Delta)^{-1}(h) \rangle - \langle h, V(s + \tilde{\Delta})^{-1}V^*(h) \rangle \geq \langle h_s, (s + \Delta)(h_s) \rangle,$$

where

$$h_s = (s + \Delta)^{-1}h - V(s + \tilde{\Delta})^{-1}V^*(h). \quad (4.3)$$

Proof. Let us first calculate the right hand side using the definition of h_s ,

$$\begin{aligned} \langle h_s, (s + \Delta)(h_s) \rangle &= \langle h, (s + \Delta)^{-1}(h) \rangle - 2\langle h, V(s + \tilde{\Delta})^{-1}V^*(h) \rangle \\ &\quad + \langle V(s + \tilde{\Delta})^{-1}V^*(h), (s + \Delta)V(s + \tilde{\Delta})^{-1}V^*(h) \rangle. \end{aligned}$$

Then since $V^*V \leq I$ and $V^*(\Delta + s)V \leq (\tilde{\Delta} + s)$ by the assumption, we obtain an upper bound of the last term

$$\begin{aligned} &\langle V(s + \tilde{\Delta})^{-1}V^*(h), (s + \Delta)V(s + \tilde{\Delta})^{-1}V^*(h) \rangle \\ &= \langle (s + \tilde{\Delta})^{-1}V^*(h), V^*(s + \Delta)V(s + \tilde{\Delta})^{-1}V^*(h) \rangle \\ &\leq \langle (s + \tilde{\Delta})^{-1}V^*(h), (s + \tilde{\Delta})(s + \tilde{\Delta})^{-1}V^*(h) \rangle \\ &= \langle h, V(s + \tilde{\Delta})^{-1}V^*(h) \rangle, \end{aligned}$$

which yields the desired inequality in the lemma. ■

Applying the above lemmas to $V_{\rho,t}$ and Δ_ρ , we have

$$\langle A, (s + \Delta_\rho)^{-1}(A) \rangle - \langle A, V_{\rho,t}(s + \Delta_{\Phi(\rho)})^{-1}V_{\rho,t}^*(A) \rangle \geq \langle A_{s,t}, (s + \Delta_\rho)(A_{s,t}) \rangle,$$

where

$$A_{s,t} = (s + \Delta_\rho)^{-1}A - V_{\rho,t}(s + \Delta_{\Phi(\rho)})^{-1}V_{\rho,t}^*(A). \quad (4.4)$$

The next lemma shows the above expression upper bounds the approximate recoverability.

Lemma 4.3. *Let $\rho \in \mathcal{D}_+(\mathcal{H})$ and $\Phi : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{K})$ be a quantum channel. Let $\mathcal{R}_{\rho,\Phi}^t$ be as defined in (4.2).*

$$\|A - \mathcal{R}_{\rho,\Phi}^t(\Phi(A))\|_1 \leq \left\| \left(A - \mathcal{R}_{\rho,\Phi}^t(\Phi(A)) \right) \rho^{-1/2} \right\|_2 \leq \frac{\cosh(\pi t)}{\pi} \left\| \int_0^\infty s^{-\frac{1}{2}+it} \Delta_\rho^{\frac{1}{2}} \tilde{A}_{s,t} ds \right\|_2,$$

where

$$\tilde{A}_{s,t} = (s + \Delta_\rho)^{-1}(A\rho^{-\frac{1}{2}+it}) - V_{\rho,t}(s + \Delta_{\Phi(\rho)})^{-1}V_{\rho,t}^*(A\rho^{-\frac{1}{2}+it}). \quad (4.5)$$

Proof. By the integral representation $r^{-\frac{1}{2}+it} = \frac{\cosh(\pi t)}{\pi} \int_0^\infty \frac{s^{-\frac{1}{2}+it}}{r+s} ds$ from [Kom66], we have

$$\begin{aligned} &\frac{\cosh(\pi t)}{\pi} \int_0^\infty s^{-\frac{1}{2}+it} \Delta_\rho^{\frac{1}{2}} \tilde{A}_{s,t} ds \\ &= \Delta_\rho^{\frac{1}{2}} \frac{\cosh(\pi t)}{\pi} \int_0^\infty s^{-\frac{1}{2}+it} (s + \Delta_\rho)^{-1} (A\rho^{-\frac{1}{2}+it}) ds \\ &\quad - \Delta_\rho^{\frac{1}{2}} V_{\rho,t} \frac{\cosh(\pi t)}{\pi} \int_0^\infty s^{-\frac{1}{2}+it} (s + \Delta_{\Phi(\rho)})^{-1} V_{\rho,t}^*(A\rho^{-\frac{1}{2}+it}) ds \\ &= \Delta_\rho^{it} (A\rho^{-\frac{1}{2}+it}) - \Delta_\rho^{\frac{1}{2}} V_{\rho,t} \Delta_{\Phi(\rho)}^{-\frac{1}{2}+it} V_{\rho,t}^*(A\rho^{-\frac{1}{2}+it}) \end{aligned}$$

$$\begin{aligned}
&= \rho^{it} A \rho^{-\frac{1}{2}} - \rho^{\frac{1}{2}} \Phi^\dagger (\Phi(\rho)^{-\frac{1}{2}+it} \Phi(A) \Phi(\rho)^{-\frac{1}{2}-it}) \rho^{it} \\
&= \rho^{it} (A - \mathcal{R}_{\rho, \Phi}^t(\Phi(A))) \rho^{-\frac{1}{2}}.
\end{aligned}$$

Then the inequality follows from the Hölder inequality,

$$\|\rho^{it} (A - \mathcal{R}_{\rho, \Phi}^t(\Phi(A))) \rho^{-\frac{1}{2}}\|_2 \|\rho^{\frac{1}{2}}\|_2 \geq \|A - \mathcal{R}_{\rho, \Phi}^t(\Phi(A))\|_1. \quad \blacksquare$$

We shall now estimate the above recoverability bounds via the monotone metrics. This is our main technical lemma.

Lemma 4.4. *Let γ^g be a monotone metric given by the integral representation (3.3) with measure $\nu_g(s)ds$. For any $0 < a \leq b < \infty$, suppose there exists a function $w : [0, \infty) \rightarrow \mathbb{R}_+$ such that*

$$W_{a,b} := \int_a^b \frac{w(s)}{s} ds < \infty$$

and a constant $C_g(a, b) > 0$ such that on the interval $[a, b]$

$$\frac{1}{w(s)} ds \leq C_g(a, b) \nu_g(s) ds.$$

Then we have for any $\rho \in \mathcal{D}_+(\mathcal{H})$, $A \in \mathcal{B}(\mathcal{H})$ and $t \in \mathbb{R}$

$$\|A - \mathcal{R}_{\rho, \Phi}^t(\Phi(A))\|_1 \leq \frac{\cosh(\pi t)}{\pi} \left(4\sqrt{a}h_1 + \frac{4}{\sqrt{b}}h_2 + C_g(a, b)^{\frac{1}{2}} W_{a,b}^{\frac{1}{2}} h_3 \right),$$

where

$$h_1 := \text{tr}(A^* \rho^{-1} A)^{\frac{1}{2}}, \quad h_2 := \frac{1}{2} \left(\text{tr}(\rho^{-2} A^* \rho A)^{\frac{1}{2}} + \text{tr}(\Phi(\rho)^{-2} \Phi(A)^* \Phi(\rho) \Phi(A))^{\frac{1}{2}} \right)$$

$$h_3 := \left(\gamma_\rho^g(A) - \gamma_{\Phi(\rho)}^g(\Phi(A)) \right)^{\frac{1}{2}}.$$

Proof. By Lemma 4.3, it is enough to bound $\| \int_0^\infty s^{-\frac{1}{2}+it} \Delta_\rho^{\frac{1}{2}} \tilde{A}_{s,t} ds \|_2$ from above. We split the integral into three terms

$$\begin{aligned}
\int_0^\infty s^{-\frac{1}{2}+it} \Delta_\rho^{\frac{1}{2}} \tilde{A}_{s,t} ds &= \int_0^a s^{-\frac{1}{2}+it} \Delta_\rho^{\frac{1}{2}} \tilde{A}_{s,t} ds + \int_a^b s^{-\frac{1}{2}+it} \Delta_\rho^{\frac{1}{2}} \tilde{A}_{s,t} ds + \int_b^\infty s^{-\frac{1}{2}+it} \Delta_\rho^{\frac{1}{2}} \tilde{A}_{s,t} ds \\
&= \text{I} + \text{II} + \text{III}.
\end{aligned}$$

Let $h_{t,a}(x) = \int_0^a \frac{s^{-\frac{1}{2}+it} x^{\frac{1}{2}}}{(x+s)} ds$. Then

$$\|\text{I}\|_2 \leq \|h_{t,a}(\Delta_\rho)(A \rho^{-\frac{1}{2}+it})\|_2 + \left\| \int_0^a s^{-\frac{1}{2}+it} \Delta_\rho^{\frac{1}{2}} V_{\rho,t}(s + \Delta_{\Phi(\rho)})^{-1} V_{\rho,t}^*(A \rho^{-\frac{1}{2}+it}) ds \right\|_2.$$

Next, we notice that

$$\begin{aligned}
&\left\| \int_0^a s^{-\frac{1}{2}+it} \Delta_\rho^{\frac{1}{2}} V_{\rho,t}(s + \Delta_{\Phi(\rho)})^{-1} V_{\rho,t}^*(A \rho^{-\frac{1}{2}+it}) ds \right\|_2^2 \\
&= \langle V_{\rho,t}^* \Delta_\rho V_{\rho,t} \int_0^a \frac{s^{-\frac{1}{2}+it}}{s + \Delta_{\Phi(\rho)}} V_{\rho,t}^*(A \rho^{-\frac{1}{2}+it}) ds, \int_0^a \frac{s^{-\frac{1}{2}+it}}{s + \Delta_{\Phi(\rho)}} V_{\rho,t}^*(A \rho^{-\frac{1}{2}+it}) ds \rangle \\
&\stackrel{(1)}{\leq} \langle \Delta_{\Phi(\rho)} \int_0^a \frac{s^{-\frac{1}{2}+it}}{s + \Delta_{\Phi(\rho)}} V_{\rho,t}^*(A \rho^{-\frac{1}{2}+it}) ds, \int_0^a \frac{s^{-\frac{1}{2}+it}}{s + \Delta_{\Phi(\rho)}} V_{\rho,t}^*(A \rho^{-\frac{1}{2}+it}) ds \rangle \\
&\leq \langle \int_0^a \frac{s^{-\frac{1}{2}+it} \Delta_{\Phi(\rho)}^{\frac{1}{2}}}{s + \Delta_{\Phi(\rho)}} V_{\rho,t}^*(A \rho^{-\frac{1}{2}+it}) ds, \int_0^a \frac{s^{-\frac{1}{2}+it} \Delta_{\Phi(\rho)}^{\frac{1}{2}}}{s + \Delta_{\Phi(\rho)}} V_{\rho,t}^*(A \rho^{-\frac{1}{2}+it}) ds \rangle
\end{aligned}$$

$$= \|h_{t,a}(\Delta_{\Phi(\rho)})V_{\rho,t}^*(A\rho^{-\frac{1}{2}+it})\|_2^2,$$

where the inequality in (1) above follows from Lemma 4.1. Moreover, for $x > 0$, we have

$$|h_{t,a}(x)| \leq \int_0^a \left| \frac{s^{-\frac{1}{2}+it}x^{\frac{1}{2}}}{(x+s)} \right| ds = 2 \arctan \sqrt{\frac{a}{x}} \leq 2 \sqrt{\frac{a}{x}}.$$

Together with the inequality above and another use of Lemma 4.1, we can hence bound $\|I\|_2$ from above:

$$\begin{aligned} \|I\|_2 &\leq 2\sqrt{a} \langle A\rho^{-\frac{1}{2}+it}, \Delta_{\rho}^{-1}(A\rho^{-\frac{1}{2}+it}) \rangle^{\frac{1}{2}} + 2\sqrt{a} \langle V_{\rho,t}^*(A\rho^{-\frac{1}{2}+it}), \Delta_{\Phi(\rho)}^{-1}V_{\rho,t}^*(A\rho^{-\frac{1}{2}+it}) \rangle^{\frac{1}{2}} \\ &= 2\sqrt{a} \operatorname{tr}(A^*\rho^{-1}A)^{\frac{1}{2}} + 2\sqrt{a} \operatorname{tr}(\Phi(A^*)\Phi(\rho)^{-1}\Phi(A))^{\frac{1}{2}} \\ &\leq 4\sqrt{a} \operatorname{tr}(A^*\rho^{-1}A)^{\frac{1}{2}}, \end{aligned}$$

where the last inequality follows from the Schwarz-type operator inequality (see [LR74], [Wol12, Theorem 5.3]):

$$\Phi(A^*)\Phi(\rho)^{-1}\Phi(A) \leq \Phi(A^*\rho^{-1}A).$$

(or simply the monotonicity of Right operator mean metric). Let us now consider $\|III\|_2$.

Define $\tilde{h}_{t,b}(x) = \int_b^\infty \frac{x^{\frac{1}{2}}s^{-\frac{1}{2}+it}}{(x+s)} ds$. Then for $x > 0$,

$$|\tilde{h}_{t,b}(x)| \leq \int_b^\infty \left| \frac{x^{\frac{1}{2}}s^{-\frac{1}{2}+it}}{(x+s)} \right| ds = 2 \arctan(\sqrt{\frac{x}{b}}) \leq 2\sqrt{\frac{x}{b}}$$

Similar to I, we obtain that

$$\|III\|_2 \leq \frac{2}{\sqrt{b}} \operatorname{tr}(\rho^{-2}A^*\rho A)^{\frac{1}{2}} + \frac{2}{\sqrt{b}} \operatorname{tr}(\Phi(\rho)^{-2}\Phi(A^*)\Phi(\rho)\Phi(A))^{\frac{1}{2}}.$$

Finally, we consider the second integral:

$$\begin{aligned} \|II\|_2 &\leq \int_a^b \sqrt{\frac{1}{s}} \|\Delta_{\rho}^{\frac{1}{2}}\tilde{A}_{s,t}\|_2 ds \\ &\leq \left(\int_a^b \frac{w(s)}{s} ds \right)^{\frac{1}{2}} \left(\int_a^b \frac{1}{w(s)} \|\Delta_{\rho}^{\frac{1}{2}}\tilde{A}_{s,t}\|_2^2 ds \right)^{\frac{1}{2}} \\ &= W_{a,b}^{\frac{1}{2}} \left(\int_a^b \frac{1}{w(s)} \|\Delta_{\rho}^{\frac{1}{2}}\tilde{A}_{s,t}\|_2^2 ds \right)^{\frac{1}{2}}. \end{aligned}$$

Now, since by assumption $\frac{1}{w(s)}ds \leq C_g(a,b)\nu_g(s)ds$ on the interval $[a,b]$, we have

$$\begin{aligned} \|II\|_2 &\leq W_{a,b}^{\frac{1}{2}} C_g(a,b)^{\frac{1}{2}} \left(\int_0^\infty \|\Delta_{\rho}^{\frac{1}{2}}\tilde{A}_{s,t}\|_2^2 \nu_g(s) ds \right)^{\frac{1}{2}} \\ &\stackrel{(2)}{\leq} W_{a,b}^{\frac{1}{2}} C_g(a,b)^{\frac{1}{2}} \left(\gamma_{\rho}^g(A) - \gamma_{\Phi(\rho)}^g(\Phi(A)) \right)^{\frac{1}{2}}. \end{aligned}$$

The inequality (2) above is justified from the rewriting of $\gamma_{\rho}^g(A) - \gamma_{\Phi(\rho)}^g(\Phi(A))$ as follows:

$$\begin{aligned} &\gamma_{\rho}^g(A, A) - \gamma_{\Phi(\rho)}^g(\Phi(A), \Phi(A)) \\ &= \int_0^\infty \left(\left\langle A, \frac{R_{\rho}^{-1}}{s + \Delta_{\rho}}(A) \right\rangle - \left\langle \Phi(A), \frac{R_{\Phi(\rho)}^{-1}}{s + \Delta_{\Phi(\rho)}}(\Phi(A)) \right\rangle \right) \nu_g(s) ds \\ &= \int_0^\infty \langle A\rho^{-\frac{1}{2}+it}, (s + \Delta_{\rho})^{-1}(A\rho^{-\frac{1}{2}+it}) \rangle \nu_g(s) ds \end{aligned}$$

$$- \int_0^\infty \langle A\rho^{-\frac{1}{2}+it}, V_{\rho,t}(s + \Delta_{\Phi(\rho)})^{-1}V_{\rho,t}^*(A\rho^{-\frac{1}{2}+it}) \rangle \nu_g(s) ds.$$

Then by Lemma 4.2,

$$\begin{aligned} \gamma_\rho^g(A, A) - \gamma_{\Phi(\rho)}^g(\Phi(A), \Phi(A)) &\geq \int_0^\infty \langle \tilde{A}_{s,t}, (s + \Delta_\rho)(\tilde{A}_{s,t}) \rangle \nu_g(s) ds \\ &\geq \int_0^\infty \langle \tilde{A}_{s,t}, \Delta_\rho(\tilde{A}_{s,t}) \rangle \nu_g(s) ds \\ &\geq \int_a^b \|\Delta_\rho^{\frac{1}{2}} \tilde{A}_{s,t}\|_2^2 \nu_g(s) ds \end{aligned}$$

from which (2) follows. This finishes the proof by adding up the bounds found for I, II and III. \blacksquare

Remark 4.5. The quantity $h_1 = \text{tr}(A^*\rho^{-1}A)^{1/2} = \gamma_\rho^{\text{RLD}}(\rho)$ is the RLD metric, which satisfies the monotonicity. On the other hand,

$$\text{tr}(\rho^{-2}A^*\rho A) = \langle A, \Delta_\rho R_\rho^{-1}A \rangle$$

corresponds to γ^g with $g(x) = x$, which is not a monotone metric. So we have to also include the term $\text{tr}(\Phi(\rho)^{-2}\Phi(A)^*\Phi(\rho)\Phi(A))$ in the estimate.

We are now ready to prove the recovery of monotone metric.

Theorem 4.6. *Let $\rho \in \mathcal{D}_+(\mathcal{H})$ and $\Phi : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{K})$ be a quantum channel. Then for any $A \in \mathcal{B}(\mathcal{H})$, the following are equivalent*

- i) $\gamma_\rho^g(A) = \gamma_{\Phi(\rho)}^g(\Phi(A))$ for all monotone metric γ_g .
- ii) $\gamma_\rho^g(A) = \gamma_{\Phi(\rho)}^g(\Phi(A))$ for some regular monotone metric γ_g .
- iii) $A = \mathcal{R}_{\rho,\Phi}^t \circ \Phi(A)$ for some/all $t \in \mathbb{R}$ where $\mathcal{R}_{\rho,\Phi}^t$ is the rotated Petz map.
- iv) There exists a quantum channel \mathcal{R} such that $\rho = \mathcal{R} \circ \Phi(\rho)$ and $A = \mathcal{R} \circ \Phi(A)$.

Proof. The i) \Rightarrow ii) and iii) \Rightarrow iv) are trivial. The direction iv) \Rightarrow i) follows from monotonicity. The direction ii) \Rightarrow iii) follows from Lemma 4.4. Indeed, note that for a regular g , one can choose $w(s) = s$, and $W_{a,b}, C_g(a, b)$ are finite for any $0 < a < b < \infty$. Choosing $a = \frac{1}{b}$ arbitrarily small yields the equality iii). \blacksquare

Recall that we say a quantum channel Φ is sufficient for a pair $\{\rho, \sigma\}$ of states whenever there exists a quantum channel \mathcal{R} such that $\mathcal{R} \circ \Phi(\rho) = \rho$ and $\mathcal{R} \circ \Phi(\sigma) = \sigma$.

Corollary 4.7. *Let $\rho, \sigma \in \mathcal{D}(\mathcal{H})$ be two quantum states with $\text{supp}(\sigma) \subseteq \text{supp}(\rho)$. For a quantum channel $\Phi : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{K})$, the following are equivalent*

- i) $\chi_g^2(\rho, \sigma) = \chi_g^2(\Phi(\rho), \Phi(\sigma))$ for all quantum χ^2 -divergence χ_g^2 ;
- ii) $\chi_g^2(\rho, \sigma) = \chi_g^2(\Phi(\rho), \Phi(\sigma))$ for some regular quantum χ^2 -divergence χ_g^2 ;
- iii) $\rho = \mathcal{R}_{\sigma,\Phi}^t \circ \Phi(\rho)$ for some/all $t \in \mathbb{R}$ where $\mathcal{R}_{\sigma,\Phi}^t$ is the rotated Petz map.
- iv) Φ is sufficient for $\{\rho, \sigma\}$.

Proof. This follows from Theorem 4.6 by choosing $A = \rho - \sigma$. Note that for non-faithful σ , we can always restrict the discussion on the support of σ . \blacksquare

Remark 4.8. The above recovery results does not hold for SLD and RLD metric

$$\gamma_\rho^{\text{RLD}}(A) = \text{tr}(A^*A\rho^{-1}), \quad \gamma_\rho^{\text{SLD}}(A) = \langle A, \mathbb{J}_\rho^{\text{SLD}}(A) \rangle$$

where $L = \mathbb{J}_\rho^{\text{SLD}}(A)$ is the operator L such that

$$A = \frac{1}{2}(L\rho + \rho L). \quad (4.6)$$

It was argued in Section 2 that if $[A, L] \neq 0$,

$$\gamma_{\mathcal{L}(\rho)}^{\text{RLD}}(\mathcal{L}(A)) = \gamma_{\mathcal{L}(\rho)}^{\text{SLD}}(\mathcal{L}(A)) = \gamma_{\rho}^{\text{SLD}}(A) < \gamma_{\rho}^{\text{RLD}}(A),$$

where \mathcal{L} is the pinching map for the spectrum of L . Thus γ^{SLD} is preserved under \mathcal{L} but \mathcal{L} is not sufficient for ρ and A . For RLD metric, we note that

$$\chi_{\text{RLD}}^2(\rho, \sigma) = \gamma_{\sigma}^{\text{RLD}}(\rho - \sigma) = \text{tr}((\rho - \sigma)^2 \sigma^{-1}) = \text{tr}(\rho^2 \sigma^{-1}) - 1$$

which is essentially the Petz-Rényi 2-divergence $Q_2(\rho||\sigma) = \text{tr}(\rho^2 \sigma^{-1})$. It has been observed in [HM17, Example 4.8] there exists states ρ, σ and quantum channel Φ such that

$$Q_2(\rho||\sigma) = Q_2(\Phi(\rho)||\Phi(\sigma)),$$

but Φ is not sufficient for $\{\rho, \sigma\}$.

4.2. Approximate Recovery of monotone metrics. In this section, we discuss approximate recovery bounds for specific metrics. Our starting point is a simple argument for general monotone metrics. Given a quantum channel Φ , the data processing inequality

$$\langle \Phi(A), \mathbb{J}_{\Phi(\rho)}^g(\Phi(A)) \rangle = \gamma_{\Phi(\rho)}^g(\Phi(A), \Phi(A)) \leq \gamma_{\rho}^g(A, A) = \langle A, \mathbb{J}_{\rho}^g(A) \rangle. \quad (4.7)$$

can be interpreted as

$$\Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi \leq \mathbb{J}_{\rho}^g,$$

as positive operators on the Hilbert-Schmidt space $\mathcal{S}_2(\mathcal{H})$. By the operator anti-monotonicity of $x \mapsto x^{-1}$, we have

$$(\Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi)^{-1} \geq (\mathbb{J}_{\rho}^g)^{-1}.$$

This leads to the following lemma.

Lemma 4.9. *For any density operator $\rho \in \mathcal{D}_+(\mathcal{H})$ and $A \in \mathcal{B}(\mathcal{H})$,*

$$\gamma_{\rho}^g(A) - \gamma_{\Phi(\rho)}^g(\Phi(A)) \geq \gamma_{\rho}^g(A - (\mathbb{J}_{\rho}^g)^{-1} \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A)) \geq \|A - \mathbb{J}_{\rho}^g \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A)\|_1^2 \quad (4.8)$$

Proof. We have

$$\begin{aligned} \gamma_{\rho}^g(A) - \gamma_{\Phi(\rho)}^g(\Phi(A)) &= \langle A, \mathbb{J}_{\rho}^g A \rangle - \langle \Phi(A), \mathbb{J}_{\Phi(\rho)}^g \Phi(A) \rangle \\ &= \langle A, \mathbb{J}_{\rho}^g(A - (\mathbb{J}_{\rho}^g)^{-1} \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A)) \rangle \\ &\stackrel{(1)}{\geq} \langle A - (\mathbb{J}_{\rho}^g)^{-1} \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A), \mathbb{J}_{\rho}^g(A - (\mathbb{J}_{\rho}^g)^{-1} \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A)) \rangle \\ &= \gamma_{\rho}^g\left(A - (\mathbb{J}_{\rho}^g)^{-1} \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A)\right) \\ &\stackrel{(2)}{\geq} \|A - (\mathbb{J}_{\rho}^g)^{-1} \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A)\|_1^2 \end{aligned}$$

Here the first inequality follows from

$$\begin{aligned} &\langle A - (\mathbb{J}_{\rho}^g)^{-1} \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A), \mathbb{J}_{\rho}^g(A - (\mathbb{J}_{\rho}^g)^{-1} \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A)) \rangle \\ &= \langle A, \mathbb{J}_{\rho}^g(A - (\mathbb{J}_{\rho}^g)^{-1} \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A)) \rangle - \langle (\mathbb{J}_{\rho}^g)^{-1} \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A), \mathbb{J}_{\rho}^g(A - (\mathbb{J}_{\rho}^g)^{-1} \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A)) \rangle \\ &= \langle A, \mathbb{J}_{\rho}^g(A - (\mathbb{J}_{\rho}^g)^{-1} \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A)) \rangle - \langle \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A), A \rangle + \langle \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A), (\mathbb{J}_{\rho}^g)^{-1} \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A) \rangle \end{aligned}$$

and for the last two terms we have

$$\begin{aligned} \langle \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A), (\mathbb{J}_{\rho}^g)^{-1} \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A) \rangle &\leq \langle \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A), \left(\Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi\right)^{-1} \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A) \rangle \\ &= \langle A, \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g \Phi(A) \rangle. \end{aligned}$$

The second inequality was found in [TKR⁺10], the proof of which we include for completeness: it is sufficient to consider the Bures metric

$$\gamma_{\rho}^{\text{Bures}}(X) = \langle X, \mathbb{J}_{\rho}^b(X) \rangle, \quad \mathbb{J}_{\rho}^b = 2(L_{\rho} + R_{\rho})^{-1},$$

since $\gamma_\rho^{\text{Bures}}(X) \leq \gamma_\rho^g(X)$ for any g . Then

$$\begin{aligned} \|X\|_1^2 &= \sup_U |\text{tr}(XU)|^2 = \sup_U |\langle U, X \rangle|^2 \\ &= \sup_U \langle (\mathbb{J}_\rho^b)^{-\frac{1}{2}} U, (\mathbb{J}_\rho^b)^{\frac{1}{2}} X \rangle \\ &\leq \sup_U \langle U, (\mathbb{J}_\rho^b)^{-1}(U) \rangle \cdot \langle X, \mathbb{J}_\rho^b(X) \rangle \\ &= \sup_U \frac{1}{2} \text{tr}(U^* U \rho + U^* \rho U) \cdot \gamma_\rho^{\text{Bures}}(X) = \gamma_\rho^{\text{Bures}}(X). \end{aligned}$$

That completes the proof. ■

The above estimate is nice but not necessarily gives a recovery bound. The reason is that the map

$$(\mathbb{J}_\rho^g)^{-1} \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^g$$

is not necessarily a channel. Indeed, such a map is always trace preserving because the adjoint is unital

$$\mathbb{J}_{\Phi(\rho)}^g \Phi(\mathbb{J}_\rho^g)^{-1}(1) = \mathbb{J}_{\Phi(\rho)}^g(\Phi(\rho)) = 1.$$

But the map $(\mathbb{J}_\rho^g)^{-1}$ may not be positive, as it is the case for the Bures metric

$$(\mathbb{J}_\rho^b)^{-1}(A) = \frac{1}{2}(\rho A + A \rho)$$

Nevertheless, the situation simplifies in the case of the $\frac{1}{2}$ -metric

$$\gamma_\rho^s(A, B) = \text{tr}(A^* \rho^{-\frac{1}{2}} B \rho^{-\frac{1}{2}}) = \langle A, \mathbb{J}_\rho^s B \rangle,$$

where the multiplication operator is

$$\mathbb{J}_\rho^s(A) = \rho^{-\frac{1}{2}} A \rho^{-\frac{1}{2}}.$$

The inverse operator is

$$(\mathbb{J}_\rho^s)^{-1}(A) = \rho^{\frac{1}{2}} A \rho^{\frac{1}{2}}.$$

Then it is clear that for $\frac{1}{2}$ -metric

$$(\mathbb{J}_\rho^s)^{-1} \Phi^\dagger \mathbb{J}_{\Phi(\rho)}^s = \mathcal{R}_{\rho, \Phi}$$

gives the Petz recovery map. Thus we have the following simple recovery bound for $\frac{1}{2}$ -metric

Theorem 4.10. *For any density operator $\rho \in \mathcal{D}_+(\mathcal{H})$ and $A \in \mathcal{B}(\mathcal{H})$,*

$$\gamma_\rho^s(A) - \gamma_{\Phi(\rho)}^s(\Phi(A)) \geq \gamma_\rho^s(A - \mathcal{R}_{\rho, \Phi} \circ \Phi(A)) \geq \|A - \mathcal{R}_{\rho, \Phi} \circ \Phi(A)\|_1^2 \quad (4.9)$$

In terms of $\chi_{1/2}^2$ divergence, for two quantum states ρ and σ with $\text{supp}(\sigma) \leq \text{supp}(\rho)$,

$$\chi_{1/2}^2(\sigma, \rho) - \chi_{1/2}^2(\Phi(\sigma), \Phi(\rho)) \geq \|\sigma - \mathcal{R}_{\rho, \Phi} \circ \Phi(\sigma)\|_1^2$$

A weaker inequality

$$\|\sigma - \mathcal{R}_{\rho, \Phi} \circ \Phi(\sigma)\|_1^2 \leq \|\sigma^{-1}\|_\infty \left(\chi_{\frac{1}{2}}^2(\sigma, \rho) - \chi_{\frac{1}{2}}^2(\Phi(\sigma), \Phi(\rho)) \right)$$

was obtained in [CS22]. Here we removed the singularity term $\|\sigma^{-1}\|_\infty$.

We now discuss some approximate recovery results using Lemma 4.4. Recall the Bogolyubov-Kubo-Mori metric is

$$\gamma_\rho^{\text{BKM}}(A) = \int_0^\infty \text{tr}(A^*(\rho + s)^{-1} A(\rho + s)^{-1}) ds.$$

The associated measure is $\nu(s)ds = \frac{1}{1+s}ds$.

Corollary 4.11. *For all $\rho \in \mathcal{D}_+(\mathcal{H})$ and $A \in \mathcal{B}(\mathcal{H})$, we have*

$$\gamma_\rho^{\text{BKM}}(A) - \gamma_{\Phi(\rho)}^{\text{BKM}}(\Phi(A)) \geq \sup_{0 < \epsilon < \frac{1}{2}} \left(\frac{\pi}{\cosh(\pi t)} \frac{\|A - \mathcal{R}_{\rho, \Phi}^t(\Phi(A))\|_1}{K(\rho, A, \epsilon)} \right)^{\frac{4}{1-2\epsilon}},$$

where

$$K(\rho, A, \epsilon) := 4 \operatorname{tr}(A^* \rho^{-1} A)^{\frac{1}{2}} + 2 \operatorname{tr}(\rho^{-2} A^* \rho A)^{\frac{1}{2}} + 2 \operatorname{tr}(\Phi(\rho)^{-2} \Phi(A)^* \Phi(\rho) \Phi(A))^{\frac{1}{2}} + 1 + (\epsilon e)^{-\frac{1}{2}}.$$

Proof. By choosing $w(s) = 1 + s$, Lemma 4.4 applies with the global constant $C_g = 1$. With the notations of Lemma 4.4, we hence have that, by choosing $\delta := \min\{h_3, 1\}$, $a = b^{-1} = \delta$,

$$\begin{aligned} \|A - \mathcal{R}_{\rho, \Phi}^t(\Phi(A))\|_1 &\leq \frac{\cosh(\pi t)}{\pi} \delta^{\frac{1}{2}} \left(4h_1 + 4h_2 + \left(-2 \ln \delta - \delta + \frac{1}{\delta} \right)^{\frac{1}{2}} \delta^{\frac{1}{2}} \right) \\ &\leq \frac{\cosh(\pi t)}{\pi} \delta^{\frac{1}{2}} \left(4h_1 + 4h_2 + \left(-2 \ln \delta + \frac{1}{\delta} \right)^{\frac{1}{2}} \delta^{\frac{1}{2}} \right) \\ &\stackrel{(1)}{\leq} \frac{\cosh(\pi t)}{\pi} \delta^{\frac{1}{2}} \left(4h_1 + 4h_2 + (-2 \ln \delta)^{\frac{1}{2}} \delta^{\frac{1}{2}} + 1 \right) \end{aligned}$$

where in (1) we have used the two-point inequality $(a + b)^{\frac{1}{2}} \leq a^{\frac{1}{2}} + b^{\frac{1}{2}}$ for any $a, b \geq 0$. Finally, since $2 \ln(\delta) + (\epsilon e)^{-1} \delta^{-2\epsilon} \geq 0$ for any $\epsilon > 0$ and $\delta \in (0, 1]$. Together with the fact that $\delta^{-\epsilon} \geq 1$ for $\delta \in (0, 1]$ and $\epsilon > 0$, we obtain that

$$\|A - \mathcal{R}_{\rho, \Phi}^t(\Phi(A))\|_1 \leq \frac{\cosh(\pi t)}{\pi} \delta^{\frac{1}{2}-\epsilon} \left(4h_1 + 4h_2 + (\epsilon e)^{-\frac{1}{2}} \right),$$

which completes the proof. ■

Another example is the α -metric

$$\gamma_\rho^{(\alpha)}(A) = \operatorname{tr}(A^* \rho^{\alpha-1} A \rho^{-\alpha})$$

The associated measure is $\nu(s)ds = \frac{\sin \pi \alpha}{\pi} s^\alpha ds$.

Corollary 4.12. *Let $\alpha \in (0, \frac{1}{2})$. Then, for all $\rho \in \mathcal{D}_+(\mathcal{H})$ and $A \in \mathcal{B}(\mathcal{H})$, we have*

$$\gamma_\rho^{(\alpha)}(A) - \gamma_{\Phi(\rho)}^{(\alpha)}(\Phi(A)) \geq \left(\frac{\pi}{\cosh(\pi t)} \frac{\|A - \mathcal{R}_{\rho, \Phi}^t(\Phi(A))\|_1}{K(\rho, A, \epsilon)} \right)^{\frac{2}{\alpha}},$$

where

$$\begin{aligned} K(\rho, A, \epsilon) &= 4 \operatorname{tr}(A^* \rho^{-1} A)^{\frac{1}{2}} + 2 \operatorname{tr}(\rho^{-2} A^* \rho A)^{\frac{1}{2}} + 2 \operatorname{tr}(\Phi(\rho)^{-2} \Phi(A)^* \Phi(\rho) \Phi(A))^{\frac{1}{2}} \\ &\quad + \sqrt{\frac{\pi}{\alpha \sin(\pi \alpha)}}. \end{aligned}$$

Proof. We take $w(s) = s^\alpha$ and $C_\alpha = \frac{\pi}{\sin(\pi \alpha)}$ so that $W_{a,b} = \int_a^b x^{\alpha-1} dx = \frac{1}{\alpha}(b^\alpha - a^\alpha)$. Then by Lemma 4.4, we have that, given $\delta := \min\{h_3, 1\}$,

$$\begin{aligned} \|A - \mathcal{R}_{\rho, \Phi}^t(\Phi(A))\|_1 &\leq \frac{\cosh(\pi t)}{\pi} \left(4h_1 \delta^{1/2} + 4h_2 \delta^{1/2} + \sqrt{\alpha^{-1} C_\alpha (\delta^{-\alpha} - \delta^\alpha) \delta} \right) \\ &\leq \frac{\cosh(\pi t)}{\pi} \left(4h_1 + 4h_2 + \sqrt{\frac{\pi}{\alpha \sin(\pi \alpha)}} \right) \delta^{\frac{\alpha}{2}}. \end{aligned} \quad \blacksquare$$

For $\alpha = \frac{1}{2}, t = 0$, the above bound is of course worse than our Theorem 4.10.

5. SUFFICIENCY OF QUANTUM FISHER INFORMATION

In this section, we discuss sufficiency of quantum Fisher information. Let (a, b) be an interval and $\{\rho_\theta\}_{\theta \in (a, b)}$ be a smooth one-parameter family of quantum states. Given a monotone metric γ^g in Definition 3.1, the associated quantum Fisher information of the family $\{\rho_\theta\}$ is defined as

$$I_\rho^g(\theta) = \gamma_{\rho_\theta}^g(\dot{\rho}_\theta),$$

where $\dot{\rho}_\theta = \frac{d}{d\theta}\rho_\theta$ is the derivative of ρ_θ w.r.t the parameter θ . It is inherited from γ^g that I^g satisfies the data processing inequality: for any quantum channel Φ

$$I_\rho^g(\theta) \geq I_{\Phi(\rho)}^g(\theta)$$

where the right-hand side is the Fisher information of the family $\theta \mapsto \Phi(\rho_\theta)$. Recall that we say a channel Φ is sufficient for $\{\rho_\theta\}_{\theta \in (a, b)}$ if there exists a recovery channel \mathcal{R} such that $\mathcal{R} \circ \Phi(\rho_\theta) = \rho_\theta$ for any $\theta \in (a, b)$.

Theorem 5.1. *Let $(\rho_\theta)_{\theta \in (a, b)}$ be a smooth family of full-rank quantum states in $\mathcal{D}(\mathcal{H})$. A quantum channel $\Phi : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{K})$ is sufficient for $\{\rho_\theta\}_\theta$ if and only if*

$$I_\rho^g(\theta) = I_{\Phi(\rho)}^g(\theta), \quad \forall \theta$$

for all/some regular quantum Fisher information I_ρ^g . Moreover, the recovery map \mathcal{R} can be chosen as the (rotated) Petz recovery map $\mathcal{R}_{\rho_o, \Phi}^t$ for any $o \in (a, b)$.

Proof. As Theorem 4.6 is valid on the support of ρ_θ , $I_\rho^g(\theta) = I_{\Phi(\rho)}^g(\theta)$ implies that for any $t \in \mathbb{R}$ and $\theta \in (a, b)$

$$\dot{\rho}_\theta = \mathcal{R}_{\rho_o, \Phi}^t(\Phi(\dot{\rho}_\theta)).$$

Thus for each $t \in \mathbb{R}$,

$$\rho_\theta^{-\frac{1}{2}+it} \dot{\rho}_\theta \rho_\theta^{-\frac{1}{2}-it} = \Phi^* \left(\Phi(\rho_\theta)^{-\frac{1}{2}+it} \Phi(\dot{\rho}_\theta) \Phi(\rho_\theta)^{-\frac{1}{2}-it} \right)$$

Recall that

$$\frac{d}{d\theta} \log \rho_\theta = \int_{\mathbb{R}} \rho_\theta^{-\frac{1}{2}+it} \dot{\rho}_\theta \rho_\theta^{-\frac{1}{2}-it} d\beta(t)$$

where t is integrating over the probability measure $d\beta(t) = \frac{\pi}{2(\cosh(\pi t)+1)} dt$, for example see [SBT17]. Thus we have

$$\frac{d}{d\theta} \log \rho_\theta = \Phi^* \left(\frac{d}{d\theta} \log \Phi(\rho_\theta) \right).$$

Integrating the above equality over any finite interval $[o, \theta] \subset (a, b)$, we have

$$\log \rho_\theta - \log \rho_o = \Phi^*(\log \Phi(\rho_\theta) - \log \Phi(\rho_o)).$$

We therefore obtain

$$D(\rho_\theta \| \rho_o) = D(\Phi(\rho_\theta) \| \Phi(\rho_o))$$

which implies $\rho_\theta = \mathcal{R}_{\rho_o, \Phi}^t \circ \Phi(\rho_\theta)$ for any θ and $t \in \mathbb{R}$. ■

The above result can be easily extended to the case of multivariate parameters. Let $\Theta \subset \mathbb{R}^n$ be a simple connected region and $\{\rho_\theta\}_{\theta \in \Theta}$ be a smooth family of quantum states. The Fisher information matrix is defined as

$$I_\rho^g(\theta) := [I_\rho^g(\theta)]_{ij} = [\gamma_{\rho_\theta}^g(\partial_i \rho_\theta, \partial_j \rho_\theta)]_{1 \leq i, j \leq n}.$$

$I_\rho^g(\theta)$ is a (real) $n \times n$ positive matrix. It is known that for any quantum channel Φ

$$I_\rho^g(\theta) \geq I_{\Phi(\rho)}^g(\theta)$$

as positive semi-definite matrix for every θ .

Theorem 5.2. *Let $(\rho_\theta)_{\theta \in \Theta}$ be a smooth family of full-rank quantum states in $\mathcal{D}(\mathcal{H})$. A quantum channel Φ is sufficient for $\{\rho_\theta\}$ if and only if*

$$I_\rho^g(\theta) = I_{\Phi(\rho)}^g(\theta), \quad \forall \theta,$$

for all/some regular quantum Fisher information I^g .

Proof. Take a smooth path $\eta : [0, 1] \rightarrow \Theta$ such that $\eta(0) = o$ and $\eta(1) = \theta$. Then $\rho_s := \rho_{\eta(s)}$ is a one parameter family of states and

$$\dot{\eta}(s) = D_{\dot{\eta}(s)} \rho_\theta = \sum_j \dot{\eta}_j(s) \partial_j \rho_\theta$$

where $\dot{\eta}(s)$ is the derivative of η at s . Then

$$I_\rho^g(s) = \gamma_{\rho_s}(D_{\dot{\eta}(s)} \rho_\theta, D_{\dot{\eta}(s)} \rho_\theta).$$

If

$$I_\rho(\theta) = I_{\Phi(\rho)}(\theta)$$

as positive semi-definite matrix, we have

$$I_\rho(s) = \sum_{ij} [I_\rho(\theta)]_{ij} \dot{\eta}_i(s) \dot{\eta}_j(s) = \sum_{ij} [I_{\Phi(\rho)}(\theta)]_{ij} \dot{\eta}_i(s) \dot{\eta}_j(s) = I_{\Phi(\rho)}(s)$$

where the right hand side is the QFI for the family $\Phi(\rho)_s := \Phi(\rho)_{\eta(s)}$. Then the assertion follows from the one dimensional case. \blacksquare

We now discuss a recovery bound using BKM Fisher information. Recall the BKM metric is

$$\gamma_\rho^{\text{BKM}}(X) = \langle X, \mathbb{J}_\rho^{\text{BKM}}(X) \rangle = \int_0^\infty \text{tr}(X^*(\rho + s)^{-1} X(\rho + s)^{-1}) ds.$$

The inverse map is

$$(\mathbb{J}_\rho^{\text{BKM}})^{-1}(X) = \int_0^1 \rho^t X \rho^{1-t} dt$$

It is easy to see that if $\rho \leq C\sigma$ for $C > 0$, then

$$\langle X, (\mathbb{J}_\rho^{\text{BKM}})^{-1} X \rangle \leq C \langle X, (\mathbb{J}_\sigma^{\text{BKM}})^{-1} X \rangle$$

Note that here the optimal constant C is the D_{\max} relative entropy up to a logarithm

$$D_{\max}(\rho \parallel \sigma) = \log \inf \{C > 0 \mid \rho \leq C\sigma\}$$

Theorem 5.3. *Let $\{\rho_\theta\}_{\theta \in [0,1]}$ be a smooth family of quantum states. Then for any quantum channel Φ , denoting $\Phi(\rho)_\theta := \Phi(\rho_\theta)$,*

$$D(\rho_1 \parallel \rho_0) - D(\Phi(\rho)_1 \parallel \Phi(\rho)_0) \leq \int_0^1 e^{\frac{1}{2} D_{\max}(\rho_1 \parallel \rho_\theta)} \sqrt{I_\rho^{\text{BKM}}(\theta) - I_{\Phi(\rho)}^{\text{BKM}}(\theta)} d\theta. \quad (5.1)$$

In particular, if $\rho_\theta \geq \lambda 1$ for any θ ,

$$D(\rho_1 \parallel \rho_0) - D(\Phi(\rho)_1 \parallel \Phi(\rho)_0) \leq \lambda^{-\frac{1}{2}} \int_0^1 \sqrt{I_\rho^{\text{BKM}}(\theta) - I_{\Phi(\rho)}^{\text{BKM}}(\theta)} d\theta.$$

Proof. In the following, we use the short notations $\gamma = \gamma^{\text{BKM}}$, $I = I^{\text{BKM}}$ and $\mathbb{J} = \mathbb{J}^{\text{BKM}}$. By Lemma 4.9, we have

$$\begin{aligned} I_\rho(\theta) - I_{\Phi(\rho)}(\theta) &\geq \|\mathbb{J}_{\rho_\theta}^{-\frac{1}{2}}(\mathbb{J}_{\rho_\theta}(\dot{\rho}_\theta) - \Phi^\dagger \mathbb{J}_{\Phi(\rho_\theta)} \Phi(\dot{\rho}_\theta))\|_2^2 \\ &\geq e^{-D_{\max}(\rho_1 \parallel \rho_\theta)} \|\mathbb{J}_{\rho_1}^{-\frac{1}{2}}(\mathbb{J}_{\rho_\theta}(\dot{\rho}_\theta) - \Phi^\dagger \mathbb{J}_{\Phi(\rho_\theta)}(\Phi(\dot{\rho}_\theta)_\theta))\|_2^2 \\ &= e^{-D_{\max}(\rho_1 \parallel \rho_\theta)} \|\mathbb{J}_{\rho_1}^{-\frac{1}{2}}((\log \dot{\rho}_\theta) - \Phi^\dagger(\log \dot{\Phi}(\rho)_\theta))\|_2^2 \end{aligned}$$

Then

$$\begin{aligned} & \left\| \mathbb{J}_{\rho_1}^{-\frac{1}{2}} \left(\log \rho_1 - \log \rho_0 - \Phi^\dagger(\log \Phi(\rho_1) - \log \Phi(\rho_0)) \right) \right\|_2 \\ & \leq \int_0^1 \left\| \mathbb{J}_{\rho_1}^{-\frac{1}{2}} ((\log \dot{\rho}_\theta) - \Phi^\dagger(\log \dot{\Phi}(\rho)_\theta)) \right\|_2 d\theta \\ & \leq \int_0^1 e^{\frac{1}{2}D_{\max}(\rho_1 \parallel \rho_\theta)} \sqrt{I_\rho(\theta) - I_{\Phi(\rho)}(\theta)} d\theta. \end{aligned}$$

Therefore,

$$\begin{aligned} D(\rho_1 \parallel \rho_0) - D(\Phi(\rho)_1 \parallel \Phi(\rho)_0) &= \text{tr} \left(\rho_1 (\log \rho_1 - \log \rho_0 - \Phi^\dagger(\log \Phi(\rho_1) - \log \Phi(\rho_0))) \right) \\ &\leq \left\| \mathbb{J}_{\rho_1}^{-\frac{1}{2}} \left(\log \rho_1 - \log \rho_0 - \Phi^\dagger(\log \Phi(\rho_1) - \log \Phi(\rho_0)) \right) \right\|_2^2 \\ &\leq \int_0^1 e^{\frac{1}{2}D_{\max}(\rho_1 \parallel \rho_\theta)} \sqrt{I_\rho(\theta) - I_{\Phi(\rho)}(\theta)} d\theta \quad \blacksquare \end{aligned}$$

Remark 5.4. The above estimate gives a recovery bound via

$$D(\rho \parallel \sigma) - D(\Phi(\rho) \parallel \Phi(\sigma)) \geq \frac{1}{4} \left\| \rho - \mathcal{R}_{\sigma, \Phi}^{\text{uni}} \circ \Phi(\rho) \right\|_1^2, \quad (5.2)$$

where $\mathcal{R}_{\sigma, \Phi}^{\text{uni}}$ is the universal recovery map. In the case when $\rho_1 = \rho_t$ is a small perturbation of ρ_0 and $t \rightarrow 0$, the estimate (5.2) is not of optimal asymptotic order, because

$$D(\rho_t \parallel \rho_0) - D(\Phi(\rho)_t \parallel \Phi(\rho)_0) = (I_\rho(0) - I_{\Phi(\rho)}(0))t^2 + O(t^3)$$

6. RECOVERABILITY OF ASYMMETRY

In general, the resource theory of asymmetry can be defined for all symmetry groups. Let G be a compact Lie group and \mathfrak{g} be its Lie algebra. We assume each system under consideration has a given unitary representation of G . To quantify the asymmetry of a state ρ with respect to symmetry G , we consider the QFI metric for the family of states $\rho_g = U(g)\rho U^*(g)$, where U is the unitary representation on the system. To express the metric, it suffices to consider a neighborhood of the identity element e of the Lie group, which can be smoothly parameterized by n real parameters, denoted by $\Theta = (\theta_1, \dots, \theta_n) \in \Omega \subset \mathbb{R}^n$, where $n = \dim(\mathfrak{g})$. Then for the parametrized family $\rho_\Theta = U(g(\Theta))\rho U^*(g(\Theta))$, the QFI matrix at ρ is the $n \times n$ matrix

$$I_G(\rho) := \left[\gamma_\rho(\partial_i \rho_g, \partial_j \rho_g) \Big|_{g=e} \right]_{1 \leq i, j \leq n} = \left[\gamma_\rho([L_i, \rho], [L_j, \rho]) \right]_{1 \leq i, j \leq n}, \quad (6.1)$$

where $\partial_i \rho_g = \frac{\partial \rho_g(\Theta)}{\partial \theta_i}$ is the partial derivatives and the skew-Hermitian operators

$$L_i = \frac{\partial U(g(\Theta))}{\partial \theta_i} \Big|_{g=e} : i = 1, \dots, n, \quad (6.2)$$

form a basis for the representation of the Lie algebra \mathfrak{g} induced by the unitary representation of G .

As an example, one can choose the local parametrization by the exponential map such that for $\Theta = (\theta_1, \dots, \theta_n)$, the corresponding unitary is $U(g(\Theta)) = \exp(\sum_i \theta_i L_i)$.⁷ For instance, in the case of $SO(3)$ symmetry corresponding to rotations in $3D$ space, the parametrization can be chosen such that operators L_1, L_2, L_3 become the angular momentum operators in x, y, z directions. Then, the unitary $\exp(\phi \sum_i n_i L_i)$ for a real vector $\hat{n} = (n_1, n_2, n_3)$ with the normalization $\|\hat{n}\|_2 = 1$, corresponds to the rotation by

⁷Note that in the case of compact connected Lie groups, the exponential map from the Lie algebra to the Lie group is surjective.

angle ϕ around the axis \hat{n} . In this case the QFI metric will be a 3×3 real matrix that determines the sensitivity of state ρ under rotations around x, y, z axes.

As we saw in Eq.(1.18) in the case of time translation symmetry, the value of QFI with respect to the time parameter t is constant for the entire family of time-evolved versions of state. However, this is not the case for a general group G , where QFI matrix will depend on the group element $g \in G$. Nevertheless, it turns out that for compact connected Lie groups, the QFI matrix $I_G(\rho)$ determines the QFI matrix for the entire family $\rho_g = U(g)\rho U^*(g) : g \in G$. Let $\eta = (\eta_1, \dots, \eta_n) \in \mathbb{R}^n$ be a local coordinate at point $g \in G$. Then, the QFI matrix defined relative to this coordinate system is related to the QFI matrix at the identity via the congruence transformation

$$\left[\gamma_\rho(\partial_i \rho_g, \partial_j \rho_g) \right]_{i,j} = V^T(g) \left[\gamma_\rho(\partial_i \rho_g, \partial_j \rho_g) \Big|_{g=e} \right]_{i,j} V(g) = V^T(g) I_G(\rho) V(g) , \quad (6.3)$$

where $\partial_i \rho_g = \partial \rho_{g(\Phi)} / \partial \eta_i$ and $V(g)$ is an invertible $n \times n$ real matrix defined by equation

$$U^*(g) \frac{\partial}{\partial \eta_i} U(g(\Phi)) = \sum_{r=1}^n V_{ri}(g) L_r . \quad (6.4)$$

Note that here the matrix V is only determined by the local coordinate (or equivalently the basis in Lie algebra) but independent of the representation U ⁸.

A CPTP map $\mathcal{E} : \mathcal{B}(\mathcal{H}_A) \rightarrow \mathcal{B}(\mathcal{H}_B)$ respects this symmetry, or is called covariant, if it satisfies the covariance condition

$$\mathcal{E}(U_A(g)(\cdot)U_A^*(g)) = U_B(g)\mathcal{E}(\cdot)U_B^*(g) , \forall g \in G \quad (6.5)$$

where U_A and U_B are the given representations of G on the input system \mathcal{H}_A and output systems \mathcal{H}_B , respectively. Then, for the family $\rho_g = U_A(g)\rho U_A^*(g)$, $\rho \in \mathcal{B}(\mathcal{H}_A)$,

$$\mathcal{E}(\rho_g) = \mathcal{E}(U_A(g)\rho U_A^*(g)) = U_B(g)\mathcal{E}(\rho)U_B^*(g) = \mathcal{E}(\rho)_g$$

The monotonicity of the QFI matrix under data processing then implies

$$I_G(\rho) \geq I_G(\mathcal{E}(\rho)) . \quad (6.6)$$

as matrices, for any covariant map \mathcal{E} and any QFI metric I_g . In the case of regular QFI metrics, our result in Theorem 5.1 implies that conservation of QFI metric guarantees reversibility.

Theorem 6.1. *Let G be a compact connected Lie group and let \mathcal{H}_A and \mathcal{H}_B be two systems with representations U_A and U_B of the group G . Let $\rho \in \mathcal{B}(\mathcal{H}_A)$ be a full rank density operator and $\mathcal{E} : \mathcal{B}(\mathcal{H}_A) \rightarrow \mathcal{B}(\mathcal{H}_B)$ be a covariant quantum channel. Then, this process is reversible with a covariant operation \mathcal{R} from B to A , such that $\mathcal{R}(\mathcal{E}(\rho)) = \rho$, if and only if*

$$I_G(\rho) = I_G(\mathcal{E}(\rho))$$

where I_G can be the QFI matrix in Eq.(6.1) defined by some/all regular QFI metric

The proof of this theorem is similar to the case of time-translation symmetry for the special case of a periodic representation, in which it suffices to consider the average Petz map Eq. 1.23 with the time T being the period.

⁸The exact form of matrix $V(g)$ can be obtained, e.g., via equation

$$\frac{d}{ds} \exp(X(s)) = \exp(X(s)) \frac{1 - \exp(-\text{ad}_{X(s)})}{\text{ad}_{X(s)}} \frac{d}{ds} X(s) ,$$

where $\text{ad}_X(Y) = [X, Y]$.

Proof. By the congruence transformation (6.3), we see $I_G(\rho) = I_G(\mathcal{E}(\rho))$ implies the QFI matrix

$$I_\rho(g) = I_{\mathcal{E}(\rho)}(g) ,$$

are preserved at $\rho_g = U_A(g)\rho U_A(g)^*$ for every $g \in G$. By Theorem (5.1), this implies the Petz recovery map $\mathcal{R}_{\rho,\mathcal{E}}$ satisfies

$$\mathcal{R}_{\rho,\mathcal{E}}(U_B(g)\mathcal{E}(\rho)U_B(g)^*) = U_A(g)\rho U_A(g)^* . \quad (6.7)$$

In general, this Petz map $\mathcal{R}_{\rho,\mathcal{E}}$ is not covariant. By twirling this map with the uniform (Haar) measure over the group G , we obtain the map

$$\mathcal{R}_{\text{avg}}(\cdot) := \int_G U_A(g)^* \mathcal{R}_{\rho_B,\mathcal{E}}(U_B(g)(\cdot)U_B(g)^*) U_A(g) d\mu(g) , \quad (6.8)$$

which is covariant and satisfies $\mathcal{R}_{\text{avg}}(\mathcal{E}(\rho)) = \rho$. This completes the proof. \blacksquare

REFERENCES

- [BC94] Samuel L Braunstein and Carlton M Caves. Statistical distance and the geometry of quantum states. *Physical Review Letters*, 72(22):3439, 1994.
- [BCM96] Samuel L Braunstein, Carlton M Caves, and Gerard J Milburn. Generalized uncertainty relations: theory, examples, and lorentz invariance. *annals of physics*, 247(1):135–173, 1996.
- [BNG00] OE Barndorff-Nielsen and RD Gill. Fisher information in quantum statistics. *Journal of Physics A: Mathematical and General*, 33(24):4481, 2000.
- [Čen78] NN Čencov. Algebraic foundation of mathematical statistics. *Statistics: A Journal of Theoretical and Applied Statistics*, 9(2):267–276, 1978.
- [CHP⁺19] Jordan Cotler, Patrick Hayden, Geoffrey Penington, Grant Salton, Brian Swingle, and Michael Walter. Entanglement wedge reconstruction via universal recovery channels. *Physical Review X*, 9(3):031011, 2019.
- [CL19] Yu Cao and Jianfeng Lu. Tensorization of the strong data processing inequality for quantum chi-square divergences. *Quantum*, 3:199, 2019.
- [Cra16] Harald Cramér. Mathematical methods of statistics (pms-9), volume 9. In *Mathematical Methods of Statistics (PMS-9), Volume 9*. Princeton university press, 2016.
- [CS22] Sam Cree and Jonathan Sorce. Approximate Petz recovery from the geometry of density operators. *Communications in Mathematical Physics*, pages 1–13, 2022.
- [CV20a] Eric A Carlen and Anna Vershynina. Recovery map stability for the data processing inequality. *Journal of Physics A: Mathematical and Theoretical*, 53(3):035204, 2020.
- [CV20b] Eric A Carlen and Anna Vershynina. Recovery map stability for the data processing inequality. *Journal of Physics A: Mathematical and Theoretical*, 53(3):035204, 2020.
- [Fis22] Ronald A Fisher. On the mathematical foundations of theoretical statistics. *Philosophical transactions of the Royal Society of London. Series A, containing papers of a mathematical or physical character*, 222(594-604):309–368, 1922.
- [FR15] Omar Fawzi and Renato Renner. Quantum conditional mutual information and approximate markov chains. *Communications in Mathematical Physics*, 340(2):575–611, 2015.
- [Gir14] Davide Girolami. Observable measure of quantum coherence in finite dimensional systems. *Physical review letters*, 113(17):170401, 2014.
- [GR22] Li Gao and Cambyse Rouzé. Complete entropic inequalities for quantum markov chains. *Archive for Rational Mechanics and Analysis*, pages 1–56, 2022.
- [GW21] Li Gao and Mark M Wilde. Recoverability for optimized quantum f-divergences. *Journal of Physics A: Mathematical and Theoretical*, 54(38):385302, 2021.
- [Hay02] Masahito Hayashi. Two quantum analogues of fisher information from a large deviation viewpoint of quantum estimation. *Journal of Physics A: Mathematical and General*, 35(36):7689, 2002.
- [HM17] Fumio Hiai and Milán Mosonyi. Different quantum f-divergences and the reversibility of quantum operations. *Reviews in Mathematical Physics*, 29(07):1750023, 2017.
- [Hol11] Alexander S Holevo. *Probabilistic and statistical aspects of quantum theory*, volume 1. Springer Science & Business Media, 2011.
- [HP19] Patrick Hayden and Geoffrey Penington. Learning the alpha-bits of black holes. *Journal of High Energy Physics*, 2019(12):1–55, 2019.

- [HPS21] Patrick Hayden, Onkar Parrikar, and Jonathan Sorce. The markov gap for geometric reflected entropy. *Journal of High Energy Physics*, 2021(10):1–58, 2021.
- [JP06] Anna Jenčová and Dénes Petz. Sufficiency in quantum statistical inference: A survey with examples. *Infinite Dimensional Analysis, Quantum Probability and Related Topics*, 9(03):331–351, 2006.
- [JRS⁺18] Marius Junge, Renato Renner, David Sutter, Mark M Wilde, and Andreas Winter. Universal recovery maps and approximate sufficiency of quantum relative entropy. In *Annales Henri Poincaré*, volume 19, pages 2955–2978. Springer, 2018.
- [KJJ⁺18] Hyukjoon Kwon, Hyunseok Jeong, David Jennings, Benjamin Yadin, and MS Kim. Clock–work trade-off relation for coherence in quantum thermodynamics. *Physical review letters*, 120(15):150602, 2018.
- [Kom66] Hikosaburo Komatsu. Fractional powers of operators. *Pacific Journal of Mathematics*, 19(2):285–346, 1966.
- [Kos05] Hideki Kosaki. Matrix trace inequalities related to uncertainty principle. *International Journal of Mathematics*, 16(06):629–645, 2005.
- [KS05] Abram Kagan and Lawrence A Shepp. A sufficiency paradox: an insufficient statistic preserving the fisher information. *The American Statistician*, 59(1):54–56, 2005.
- [LR74] Elliott H Lieb and Mary Beth Ruskai. Some operator inequalities of the Schwarz type. *Advances in Mathematics*, 12(2):269–273, 1974.
- [LR99] Andrew Lesniewski and Mary Beth Ruskai. Monotone riemannian metrics and relative entropy on noncommutative probability spaces. *Journal of Mathematical Physics*, 40(11):5702–5724, 1999.
- [Mar12] Iman. Marvian. *Symmetry, Asymmetry and Quantum Information*, PhD thesis. PhD thesis, University of Waterloo, <https://uwspace.uwaterloo.ca/handle/10012/7088>, 2012.
- [Mar20] Iman Marvian. Coherence distillation machines are impossible in quantum thermodynamics. *Nature communications*, 11(1):1–9, 2020.
- [Mar22] Iman Marvian. Operational interpretation of quantum fisher information in quantum thermodynamics. *Physical Review Letters*, 129(19):190502, 2022.
- [MC89] Elena Aleksandrovna Morozova and Nikolai Nikolaevich Chentsov. Markov invariant geometry on state manifolds. *Itogi Nauki i Tekhniki. Seriya " Sovremennye Problemy Matematiki. Noveishie Dostizheniya"*, 36:69–102, 1989.
- [ML16] Iman Marvian and Seth Lloyd. From clocks to cloners: Catalytic transformations under covariant operations and recoverability. *arXiv preprint arXiv:1608.07325*, 2016.
- [MS14] Iman Marvian and Robert W Spekkens. Extending noether’s theorem by quantifying the asymmetry of quantum states. *Nature communications*, 5(1):1–8, 2014.
- [Pet86] Dénes Petz. Sufficient subalgebras and the relative entropy of states of a von neumann algebra. *Communications in mathematical physics*, 105(1):123–131, 1986.
- [Pet88] Dénes Petz. Sufficiency of channels over von neumann algebras. *The Quarterly Journal of Mathematics*, 39(1):97–108, 1988.
- [Pet96] Dénes Petz. Monotone metrics on matrix spaces. *Linear algebra and its applications*, 244:81–96, 1996.
- [Pet02] Dénes Petz. Covariance and fisher information in quantum mechanics. *Journal of Physics A: Mathematical and General*, 35(4):929, 2002.
- [PG11] Dénes Petz and Catalin Ghinea. Introduction to quantum fisher information. In *Quantum probability and related topics*, pages 261–281. World Scientific, 2011.
- [PH96] Dénes Petz and Hiroshi Hasegawa. On the riemannian metric of α -entropies of density matrices. *Letters in Mathematical Physics*, 38(2):221–225, 1996.
- [Pol13] David Pollard. A note on insufficiency and the preservation of fisher information. *Probability to Statistics and Back: High-Dimensional Models and Processes-A Festschrift in Honor of Jon A. Wellner*,. *Beachwood: Institute of Mathematical Statistics*, 2013.
- [Rao45] CR Rao. Information and accuracy attainable in the estimation of statistical parameters. kots & johnson nl (eds.), *breakthroughs in statistics volume i: Foundations and basic theory*, 235–248, 1945.
- [SBT17] David Sutter, Mario Berta, and Marco Tomamichel. Multivariate trace inequalities. *Communications in Mathematical Physics*, 352(1):37–58, 2017.
- [Sch12] Mark J Schervish. *Theory of statistics*. Springer Science & Business Media, 2012.

- [SFR16] David Sutter, Omar Fawzi, and Renato Renner. Universal recovery map for approximate markov chains. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 472(2186):20150623, 2016.
- [Tak19] Ryuji Takagi. Skew informations from an operational view via resource theory of asymmetry. *Scientific reports*, 9(1):1–12, 2019.
- [TKR⁺10] Kristan Temme, Michael James Kastoryano, Mary Beth Ruskai, Michael Marc Wolf, and Frank Verstraete. The χ^2 -divergence and mixing times of quantum markov processes. *Journal of Mathematical Physics*, 51(12):122201, 2010.
- [Wol12] M. M. Wolf. Quantum channels & operations: Guided tour. *Lecture notes available at [http://www-m5.ma.tum.de/foswiki/pub M, 5](http://www-m5.ma.tum.de/foswiki/pub/M,5)*, 2012.
- [YV16] Benjamin Yadin and Vlatko Vedral. General framework for quantum macroscopicity in terms of coherence. *Physical Review A*, 93(2):022122, 2016.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HOUSTON, HOUSTON, TX 77204, USA
Email address, Li Gao: `lgao12@uh.edu`

ZENTRUM MATHEMATIK, TECHNISCHE UNIVERSITÄT MÜNCHEN, 85748 GARCHING, GERMANY
Email address, Haojian Li: `haojian.li@tum.de`

Email address, Iman Marvian: `iman.marvian@duke.edu`

ZENTRUM MATHEMATIK, TECHNISCHE UNIVERSITÄT MÜNCHEN, 85748 GARCHING, GERMANY
Email address, Cambyse Rouzé: `cambyse.rouze@tum.de`