Perspectives on fundamental cosmology from Low Earth Orbit and the Moon

Gianfranco Bertone,^{1, *} Oliver L. Buchmueller,^{2, †} and Philippa S. $Cole^{1, \ddagger}$

¹Gravitation Astroparticle Physics Amsterdam (GRAPPA),

Institute for Theoretical Physics Amsterdam and Delta Institute for Theoretical Physics,

University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands

²Imperial College London, Exhibition Rd, South Kensington, London SW7 2BX, United Kingdom

The next generation of space-based experiments will go hunting for answers to cosmology's key open questions which revolve around inflation, dark matter and dark energy. Low earth orbit and lunar missions within the European Space Agency's Human and Robotic Exploration programme can push our knowledge forward in all of these three fields. A radio interferometer on the Moon, a cold atom interferometer in low earth orbit and a gravitational wave interferometer on the Moon are highlighted as the most fruitful missions to plan and execute in the mid-term.

Keywords: dark matter — inflation — dark energy — space-based experiments

INTRODUCTION

The standard cosmological model provides a simple framework to explain a variety of observations, ranging from sub-galactic scales to the size of the observable universe. Yet many open questions remain: the model relies on an unknown mechanism for the production of perturbations in the early universe, on an unknown matter component, generically referred to as dark matter, and on an unknown mechanism that leads to an accelerated expansion of the universe, generically referred to as dark energy.

The next generation of space-based experiments are our best chance of unveiling these mysteries. A united front of low earth orbit and lunar missions, as outlined in the European Space Agency's (ESA) Human and Robotic Exploration (HRE) [1], will break unprecedented ground on all of these fronts. Alongside the Laser Interferometer Space Antenna [2], a radio interferometer on the Moon, a cold atom interferometer in low earth orbit and a gravitational wave interferometer on the Moon would provide a full-coverage approach to unravelling the key open questions in cosmology today.

In section the key knowledge gaps in cosmology are highlighted, in section specific suggestions for experiments that should be the priorities for ESA's space programme and which questions they will answer are laid out, before concluding and discussing the future outlook in section.

KEY KNOWLEDGE GAPS

Inflation

The theory of inflation is arguably the most promising model of the physics of the early universe [3]. The paradigm postulates that quantum fluctuations went on to seed the cosmological perturbations that we see imprinted on the Cosmic Microwave Background (CMB) and were the beginnings of all of the structure in the universe today. And yet, much remains to be understood about the properties of the quantum field that supposedly led to the initial period of exponential expansion of the universe. Whilst the paradigm is fully consistent with cosmological data [4–6], we still currently lack direct smoking-gun evidence supporting it, as well as a specific model for how one or more scalar fields drove the expansion.

On large scales, $k \sim 10^{-3} - 0.1 \,\mathrm{Mpc}^{-1}$, observations of the CMB temperature anisotropies by Planck [6] have confirmed to incredible precision that density perturbations were small (fluctuations of order 10^{-5}) and almost scale-invariant. The simplest single-field, slow-roll models of inflation are able to describe this spectrum of the density perturbations. However, deviations from scaleinvariance on small scales could indicate a more complicated model that exhibits a feature in the inflationary potential. Such models could have interesting observational signatures, such as ultra-compact mini-haloes [7, 8] or primordial black holes [9]. Furthermore, primordial non-Gaussianity has been constrained to be small, $f_{\rm NL,local} = -0.9 \pm 5.1$, on large scales [10]. This constraint has limited the viability of many models of inflation that predicted larger values of primordial non-Gaussianity, for example DBI inflation and EFT inflation [11, 12]. However, reaching the $f_{\rm NL,local} < 1$ threshold will provide strong evidence that observations are not consistent with multi-field models of inflation [13]. The final piece of the puzzle can be provided by the tensorto-scalar ratio, which is currently constrained to be less than 0.1 [6], a measurement of which would indicate the energy scale at which inflation happened.

Dark matter

Similarly, the existence of dark matter is supported by a wide array of independent observations, but we

^{*} g.bertone@uva.nl

[†] o.buchmueller@imperial.ac.uk

[‡] p.s.cole@uva.nl

still know very little about the fundamental nature of this elusive component of the universe. In the past four decades, a strong effort went into the search for a particular class of candidates: weakly interacting massive particles [14]. However, no experiment has yet found evidence for these particles, and attention has turned to different classes of dark matter candidates in regions of parameter space where they would have evaded strong constraints from direct detection before now, for example axion-like-particles (ALPs) [15–17] or primordial black holes (PBHs) [18].

Axion-like-particles are in particular a popular dark matter candidate [15-17]. The QCD (quantum chromodynamics) axion was first postulated in the 70s to solve the strong CP problem [19]. However, ALPs more generally, often motivated by string theories in which ultralight particles are ubiquitous, display the qualities required to explain all or part of the dark matter. Whilst searches for the standard axion with a mass of order of a few hundred keV have yielded no detections, "invisible" axions with very small masses are still viable candidates. Search strategies vary depending on the mass of the axion, which can't be theoretically predicted, but the most common approach is to probe their interactions with electromagnetic fields and constrain the axion-photon coupling [20]. Astrophysical observations are able to look for signatures of axion to photon conversion in the presence of electromagnetic fields, for example, by looking for such processes in the vicinity of the magnetosphere of neutron stars [21-23], or their production in the solar core, triggered by X-rays scattering off electrons and protons in the presence of the Sun's strong magnetic fields [24].

For masses less than 1eV, axions are a sub-set of the broader class of ultra-light dark matter models, with masses down to (theoretically) 10^{-24} eV, although Lyman-alpha forest constraints have ruled out axion masses less than 2×10^{-20} eV [25], see [26] for a review. Ultra-light dark matter models postulate a new ultralight boson, which displays wave-like properties on galactic scales, but behaves like cold dark matter on larger scales where the cold dark matter (CDM) paradigm has strong support from observations. The behaviour on galactic scales, due to the Bose-Einstein condensate which forms, can have interesting signatures that could explain small-scale problems with CDM [27, 28] and would have distinctive features for distinguishing between models such as fuzzy dark matter, self-interacting fuzzy dark matter and superfluid dark matter [29, 30].

Another promising candidate that received a lot of renewed attention after the LIGO/Virgo observations of order 10 solar mass binary black hole mergers is primordial black holes [31–33]. They are the only proposed explanation of dark matter that requires no new physics beyond the standard model, which makes them an attractive candidate. However, strong constraints have now been placed across the parameter space via microlensing, gravitational waves and CMB observations which have essentially ruled them out as making up the entirety of the dark matter budget in all but one window around an asteroid mass. See e.g. [34] for a review of current constraints. There is also the possibility of two-component dark matter models that include primordial black holes and another particle, with interesting signatures of interaction between the two [35, 36].

Dark energy

Dark energy is a generic term for the mechanism responsible for the observed accelerated expansion of the universe. One of the key questions that may bring us closer to the identification of dark energy is whether its energy density has remained constant throughout the history of the universe, as would be the case if it arises from the so-called vacuum energy, or whether it evolves with time, as appropriate for an evolving quantum field. See e.g. [37] for a review.

A diversified experimental approach involving astronomical surveys and gravitational waves searches are arguably our best hope to make progress in the search for smoking-gun evidence of inflation, identification of dark matter, and understanding of dark energy. We will highlight possibilities for future space-based experiments that can break new ground on these frontiers.

PRIORITIES FOR THE SPACE PROGRAMME

Space experiments may soon provide important clues on the nature of inflation, dark matter and dark energy. As an overview of the current context, we list in Table 1 some experiments that might in particular enable gravitational wave searches for signatures of dark matter and primordial gravitational waves with space-borne interferometers as well as indirect detection of dark matter and probing primordial fluctuations with Moon-based radio telescopes. We choose to focus on three key probes as most relevant in the framework of ESA's Human and Robotic Exploration Directorate [1] to address the knowledge gaps discussed above:

- A) a radio interferometer on the Moon (RIM)
- B) a space gravitational wave detector using cold atoms (AEDGE)
- C) a gravitational waves interferometer on the Moon (GWIM)

Radio interferometer on the Moon

The most relevant experiment for ESA's Directorate of Human and Robotic Exploration is the radio interferometer on the Moon [38–41]. The rationale for this experiment is that placing a radio telescope on the far side

Open fundamental scientific question	Focus of ESA HRE platform :	Context of related recent and	Short, middle or long term
	LEO, Moon, Mars, BLEO	future space experiments	
Origin of primordial fluctuations	Moon	LISA, AEDGE, RIM	Middle term
Nature of dark matter	LEO	LISA, AEDGE, AEDGE pathfinder, RIM, GWIM	Middle term
Phase transitions	LEO	LISA, AEDGE	Middle term
Existence of primordial black holes	Moon	LISA, AEDGE, GWIM	Middle term

TABLE I. Recommendations for addressing key questions in fundamental cosmology with the ESA HRE programme in the short, middle and long term.

of the Moon would give it access to wavelengths shorter than 30 MHz. Radiation at these frequencies is distorted or completely absorbed by the Earth's ionosphere. An interferometer on the Moon will bypass this limitation, as well as shield the instruments from the background generated by terrestrial radio sources. Furthermore, the size of the array, which determines the resolution of the detector, is less restricted than an Earth-based detector like the Hydrogen Epoch of Reionization Array (HERA) [42], the Low Frequency Array (LoFAR) [43] or the Square Kilometre Array (SKA) [44].

A radio telescope on the Moon would allow us to peer into the so-called dark ages of the universe [45], i.e. the epoch between the emission of the CMB and the reionization of the universe, triggered by the formation of the first stars. By studying the redshifted 21-cm line absorption feature, we can obtain unprecedented information on the history of reionization, and search for the signatures of dark matter annihilation or decay by looking for specific forms of the absorption feature [46, 47].

Furthermore, a radio interferometer on the Moon could in principle have baselines as long as 300km, which would enable unprecedented access to information about very small scales [48–50]. Measuring the 21cm power spectrum would provide a tracer for the underlying dark matter power spectrum. This could unlock information about dark matter sub-structures, the existence of primordial black holes, and the validity of slow-roll inflationary models. Tomographic analysis will also enable information to be gathered at a range of redshifts, providing new insight into the evolution of our Universe between the time of the CMB and today.

Finally, a lunar radio telescope would allow us also to efficiently probe the so-called primordial non-gaussianity via the 21cm bispectrum [51–53], which would provide a powerful test of the theory of inflation. 21cm observations will complement upcoming large-scale structure surveys which will help us to understand how structures are evolving, allowing for even more information to be extracted from the as yet un-probed region both in terms of redshift and access to small scales. 21cm observations from the Moon therefore add to the line of inquiry on all three fronts: inflation, dark matter and dark energy.

Arguably the largest challenge to overcome will be how to deal with extremely large foregrounds. They are expected to be 6 or 7 orders of magnitude larger than the signal being sought, and therefore systematics will need to be incredibly well understood. Extremely careful subtraction of galactic foregrounds [54, 55] will need to be performed so that any signal found in the data can be confidently interpreted [56–58].

Cold atom interferometer

The direct detection of gravitational waves (GW) with LIGO/Virgo, that led to the Nobel prize for Physics in 2017, has opened new opportunities for cosmology. The space interferometer LISA has been selected to be ESA's third large-class mission, and it is scheduled to be launched in 2034. Experience with the electromagnetic spectrum shows the importance of measurements over a range of frequencies, and we note that there is a gap between the frequencies covered by LIGO/Virgo, as well as proposed detectors Einstein Telescope and Cosmic Explorer on Earth, and LISA in space, in the deci-Hz band.

A promising candidate to explore this frequency band is the AEDGE mission concept [59], which is based on novel quantum technology utilising Cold Atom (CA) techniques. Gravitational waves alter the distance between the cold atom clouds as they pass through the interferometer.

AEDGE will be able to probe the gravitational waves due to coalescing intermediate mass black holes with masses $100 - 10^5 M_{\odot}$. This could shed light on the existence of intermediate black holes, and their potential role as seeds for the growth of supermassive black holes [60]. Additionally, black holes in the pair-instability mass gap will enter the deci-Hz range. If this gap is populated then it will motivate a deeper understanding of supernova collapse, or the need to invoke alternative mechanisms for producing such large black holes such as hierarchical mergers [61–63]. Improved constraints on order $100M_{\odot}$ primordial black holes should also be possible, which would complement constraints from the CMB on this mass range [64].

AEDGE can also probe a wide array of dark matter candidates. Scalar field dark matter, for instance, causes quantities such as the electron mass and the fine structure constant to oscillate with frequency and amplitude determined by the dark matter mass and local density. This leads to variation in the atomic transition frequencies, which imprint on the relative phase difference between cold atom clouds that atom interferometers measure. Proposed AEDGE sensitivities will enable the coupling between scalar dark matter and electrons, photons or via the Higgs-portal to be probed with up to 10 orders of magnitude improvement, with respect to current constraints from MICROSCOPE [65], on mass ranges between 10^{-18} and 10^{-12} eV. Other couplings can also be probed, for example the axion-nucleon coupling for axion-like DM lighter than 10^{-14} eV, or the coupling between a dark vector boson and the difference between baryon and lepton number [66, 67].

Furthermore, AEDGE offers a new channel for probing strong gravity regimes, where any deviations from general relativity are most likely to be noticeable. For example, precise measurements of post-Newtonian parameters would offer powerful probes of the predictions of general relativity and searches for deviations due to, for example, a graviton mass [68]. Comparing measurements in different frequency ranges will also make possible sensitive probes of Lorentz invariance [69].

An opportunity to learn about the nature of dark matter from gravitational wave signals can also be realised by observations of coalescing black holes in the deci-Hz band. Intermediate mass black holes $(10^3 - 10^5 M_{\odot})$ and primordial black holes (of any mass) may be surrounded by dense dark matter spikes [70, 71]. If these black holes form binaries with a much lighter companion, i.e. intermediate and extreme mass ratios $(q = m_2/m_1 < 10^{-2.5}),$ then the dark matter may imprint a dephasing on the gravitational waveform of the inspiralling binary. The amount of dephasing could teach us something about the nature of the dark matter surrounding the black holes, for example whether it's cold and collisionless, or whether it's an ultra-light scalar field. LISA will have sensitivity to such effects for larger mass systems [72–74], future ground-based gravitational wave observatories will have sensitivity to low-mass systems, and an experiment such as AEDGE could bridge the gap to cover the entire frequency range and promote multi-band searches for the same signal. The gravitational wave signal could also be accompanied by an electromagnetic counterpart that might provide information about an environment involving dark matter if the particle is able to convert to radiation or interact with it in some way like it does in the case of, for example, axions.

Possible non-black hole binary cosmological targets for the deci-Hz band include GWs from first-order phase transitions in the early Universe, e.g., during electroweak symmetry breaking in modifications of the Standard Model with additional interactions, or during the breaking of higher gauge symmetries [75]. The deci-Hz frequency range could also probe different parameter ranges for such transitions, and combining measurements with those by other experiments like LISA or LIGO/Virgo could help unravel different contributions, e.g., from bubble collisions, sound waves and turbulence [76, 77]. Another cosmological target for the deci-Hz band is the possible GW spectrum produced by cosmic strings [78]. In standard cosmology this spectrum would be almost scaleinvariant, but there could be modifications due to a nonstandard evolution of the early Universe. A detection or constraint on such observables could additionally provide clues as to the nature of dark matter, especially in the case of ultra-light dark matter particles which are expected to be produced alongside GWs from phase transitions in the early Universe and topological defects such as cosmic strings.

As outlined in the Cold Atom in Space Community Roadmap [79], AEDGE, its pathfinder experiments, and other cold-atom experiments in space would also be able to make sensitive measurements relevant to several other aspects of fundamental physics, including the gravitational redshift, the equivalence principle, possible longrange fifth forces [80, 81], variations in fundamental constants and popular models of dark energy [82].

With the AION experiment in the UK [83], the MAGIS experiment in the US [84], the MIGA experiment in France [85], the ZAIGA experiment in China [86], as well as the proposed European ELGAR project [87], there is already a large programme of terrestrial cold atom experiments in place. These experiments serve as terrestrial pathfinders for a large-scale mission like AEDGE, and it would be important to complement those with a dedicated technology development programme to pave the way for space-based cold atom pathfinder experiments. First, dedicated pathfinders could be hosted at the International Space Station, building the foundation of a medium-class mission. This could then lead in the longterm to a large-class mission such as AEDGE to explore the ultimate physics potential of the deci-Hz band.

A gravitational waves interferometer on the Moon

In addition to a cold atom interferometer such as AEDGE, a lunar-based gravitational wave (GW) interferometer is ideal for probing GW frequencies in the range between deci-Hz to 5 Hz, which is where both Earthand space-based detectors have less sensitivity [88, 89]. Preliminary estimates suggest that such an instrument would allow us to trace the expansion rate of the universe up to redshift $z \sim 3$, test General Relativity and the standard cosmological model up to redshift $z \sim 100$, and probe the existence of primordial black holes, as well as dark matter in neutron star cores [90, 91].

FUTURE OUTLOOK AND SUMMARY

We have outlined recommendations for how the future experiments in ESA's Human and Robotic Exploration Directorate fit into a strategy for answering the key fundamental questions in cosmology that we have highlighted in table I. These experiments will operate from either low earth orbit or the Moon, and are ambitious plans which should take place in the middle term, to give time for pathfinders and terrestrial experiments to pave the way for the development of the necessary technologies to make these moonshot missions possible. With many complementarities between experiments, European Space Agency's (ESA) Human and Robotic Exploration programme has the potential to make dramatic and perspective-changing discoveries in cosmology and astroparticle physics.

ACKNOWLEDGEMENTS

The authors thank the European Space Agency for the opportunity to contribute this review article. P.C. acknowledges funding from the Institute of Physics, University of Amsterdam.

Competing interests

The authors declare no competing financial or nonfinancial interests.

Author contributions

Gianfranco Bertone and Oliver Buchmueller wrote the ESA science community white paper related to this topic, which Philippa Cole adapted to form this perspective piece.

- [1] European Space Agency, "European Space Agency Agenda 2025," (2021).
- [2] Pau Amaro-Seoane al., et "Laser interferometer space antenna," (2017).
- [3] Alan H. Guth, "Inflationary universe: A possible solution to the horizon and flatness problems," Phys. Rev. D 23, 347–356 (1981).
- [4] A. Kogut et al., "First-year wilkinson mianisotropy probe (WMAP crowave) observations: Temperature-polarization correlation," The Astrophysical Journal Supplement Series 148, 161–173 (28) Sianfranco Bertone and Tim M. P. Tait, "A new era in
- [5] G. Hinshaw et al., "First-year wilkinson microwave anisotropy probe (WMAP) observations: Data processing methods and systematic error limits," The Astrophysical Journal Supplement Series 148, 63–95 (2003).(2021).
- [6] Planck Collaboration. "Planck 2018results." Astronomy & amp Astrophysics 641, A10 (2020).
- "Ultracompact Grigor Aslanvan et al., mini-[7]halos as probes of inflationary cosmology," Physical Review Letters 117 (2016), 10.1103/physrevlett.117.141602 version in neutron star magnetospheres:
- [8] Torsten Bringmann, Pat Scott, and Yashar Akrami, "Improved constraints on the primordial power spectrum at small scales from ultracompact minihalos," Physical Review D 85 (2012), 10.1103/physrevd.85.125027.
- [9] Juan García-Bellido and Ester Ruiz Morales, "Primordial black holes from single field models of inflation," Physics of the Dark Universe 18, 47–54 (2017).
- [10] Planck Collaboration, "Planck 2018 results. IX. Constraints on primordial non-Gaussianity," Astronomy & Astrophysics 641, A9 (2020), arXiv:1905.05697 [astro-ph.CO].
- "Primordial [11] Xingang Chen, nonmodels," gaussianities from inflation Advances in Astronomy 2010, 1–43 (2010).
- [12] Sébastien Renaux-Petel, "Primordial non-gaussianities after planck 2015 : An introductory review," Comptes Rendus Physique 16, 969–985 (2015).
- [13] Christian T. Byrnes and Ki-Young Choi, "Review of local non-gaussianity from multifield inflation," Advances in Astronomy **2010**, 1–18 (2010).
- [14] Marc Schumann, "Direct detection of WIMP status," dark matter: concepts and Journal of Physics G: Nuclear and Particle Physics 46, 103003 (2692) all-scale challenges to the λ CDM paradigm,"

- [15] Kiwoon Choi, Sang Hui Im, and Chang Sub Shin, "Recent progress in the physics of axions and axion-like particles," Annual Review of Nuclear and Particle Science 71, 225–252 (2021).
- [16] A. Ringwald, "Axions and axion-like particles," (2014), 10.48550/ARXIV.1407.0546.
- [17] Igor G. Irastorza and Javier Redondo, "New experimental approaches in the search for axion-like particles," Prog. Part. Nucl. Phys. 102, 89–159 (2018), arXiv:1801.08127 [hep-ph].
- the search for dark matter," Nature 562, 51–56 (2018).
- [19] Francesca Chadha-Day, John Ellis, and David J. E. Marsh, "Axion dark matter: What is it and why now?"
- [20] Yannis Semertzidis SungWoo Κ. and Youn. "Axion dark matter: How to see it?" (2021).
- [21]Samuel J. Witte, Dion Noordhuis, Thomas D. P. and Christoph Weniger, "Axion-photon Edwards.
- The role of the plasma in the goldreich-julian model," Physical Review D 104 (2021), 10.1103/physrevd.104.103030.
- [22]Anson Hook and Junwu Huang, "Probing axions with neutron star inspirals and other stellar processes," Journal of High Energy Physics 2018, 36 (2018), arXiv:1708.08464 [hep-ph].
- [23] Joshua W. Foster et al., "Extraterrestrial axion search with the breakthrough listen galactic center survey," (2022), 10.48550/ARXIV.2202.08274.
- [24] Shigetaka Moriyama et al., "Direct search for solar axions by using strong magnetic field and Xray detectors," Physics Letters B **434**, 147–152 (1998), arXiv:hep-ex/9805026 [hep-ex].
- [25] Keir Κ. Rogers Hiranya V. Peiris, and "Strong bound on canonical ultralight axion dark matter from the lyman-alpha forest," Physical Review Letters 126 (2021), 10.1103/physrevlett.126.071302
- [26] Elisa G. M. Ferreira, "Ultra-light dark matter," (2020).
- [27] Antonino Del Popolo and Morgan Le Delliou, "Small scale problems of the λ CDM model: A short review," Galaxies 5, 17 (2017).
- [28] James S. Bullock and Michael Boylan-Kolchin,

Annual Review of Astronomy and Astrophysics 55, 343–387 (420)1K) atie Short,

- and Jens C. [29] Xiaolong Du, Christoph Behrens, "Substructure of fuzzy Niemeyer, dark mathaloes," MNRAS 465, 941-951 (2017), ter arXiv:1608.02575 [astro-ph.CO].
- [30] Justin Khoury, "Dark superfluidity," matter SciPost Physics Lecture Notes (2022), 10.21468/scipostphyslectnbdes42.as seeds of supermassive black holes,'
- [31] Simeon Bird et al., "Did LIGO detect dark Phys. Rev. Lett. 116, 201301 (2016), matter?' arXiv:1603.00464 [astro-ph.CO].
- [32] Misao Sasaki, Teruaki Suyama, Takahiro Tanaka, and Shuichiro Yokoyama, "Primordial black hole scenario for the gravitational-wave event GW150914," Physical Review Letters 117 (2016), 10.1103/physrevlett.117.061Ddmingo,
- [33] Misao Sasaki, Teruaki Suyama, Takahiro Tanaka, "Primordial and Shuichiro Yokoyama, black holes—perspectives in gravitational wave astronomy," Classical and Quantum Gravity 35, 063001 (2018).
- [34] Anne M. Green and Bradley J. Kavanagh, "Primordial Black Holes as a dark matcandidate," J. Phys. G 48, 043001 (2021), ter arXiv:2007.10722 [astro-ph.CO].
- [35] Julian Adamek, Christian T. Byrnes, Mateja Shaun Hotchkiss, "WIMPs and Gosenca. and stellar-mass primordial black holes are incompatible,"
- [36] Gianfranco Bertone, Adam M. Coogan, Daniele Gaggero, Bradley J. Kavanagh, and Christoph "Primordial black silver Weniger, holes as bullets for new physics at the weak scale," Physical Review D 100 (2019), 10.1103/physrevd.100.123013[55] G.
- [37] Michael J. Mortonson, David H. Weinberg, and Martin White, "Dark energy: A short review," (2014).
- [38] Amin Aminaei al.. "Basic radio et interferometry for future lunar missions." in 2014 IEEE Aerospace Conference (IEEE Aerospace and Electronic Systems Society, 2014) pp. 1-19, eemcseprint-24892 ; 2014 IEEE Aerospace Conference ; Conference date: 01-03-2014 Through 08-03-2014.
- [39] Steven Furlanetto et al., "Astro 2020 Science White Paper: Fundamental Cosmology in the Dark Ages with 21cm Line Fluctuations," arXiv e-prints, arXiv:1903.06212 (2019), arXiv:1903.06212 [astro-ph.CO].
- [40] Jack О. Burns et al. "Dark cosmology: Investigating dark matter & amp; exotic physic Minthly diokiages fut heght wated shifted not did and a state of the st (2019).
- [41] Michael Duke, Wendell Mendell, and B.B. Roderts, "Lunar bases and space activities of the 21st century," Strategies for A Permanent Lunar Base, 57–68 (1985).
- [42] David R. DeBoer et al.. "Hvdrogen array epoch of reionization (HERA)," Publications of the Astronomical Society of the Pacific 129, 64500 go 201 N). Di
- [43] van Haarlem et al., M. P., "Lofar: The low-frequency array," A&A 556, A2 (2013).
- [44] Peter E. Dewdney, Peter J. Hall, Richard T. Schilizzi, and T. Joseph L. W. Lazio, "The square kilometre array," Proceedings of the IEEE 97, 1482–1496 (2009).
- [45] Leon Koopmans et al. "Peering into the dark (ages) with low-frequency space interferonerserf,"stellar-mass black holes and their gravitational-(2019).
- [46] M. Valdes, A. Ferrara, M. Mapelli, and E. Ripamonti, "Constraining dark matter through 21-cm observations," Monthly Notices of the Royal Astronomical Society 377, 245-2520(20073)k-accreting massive primordial black holes,"

- José Luis Bernal, Alvise Rac-Verde, canelli. Licia and Jens Chluba, "Enages with dark matter," lightening the dark Journal of Cosmology and Astroparticle Physics 2020, 020–020 (202
- Luis Bernal, Alvise Raccanelli, Licia Verde, [48] José and Joseph Silk, "Signatures of primordial black
- Journal of Cosmology and Astroparticle Physics 2018, 017–017 (201
- [49] Philippa S. Cole and Joseph Silk, "Small-scale primordial fluctuations in the 21 cm Dark Ages signal," Mon. Not. Roy. Astron. Soc. 501, 2627-2634 (2021), arXiv:1912.02171 [astro-ph.CO].
- [50] Olga Mena, Sergio Palomares-Ruiz, Pablo Villanueva-
- and Samuel J. Witte, "Constraining the primordial black hole abundance with 21cosmology," cm Phys. Rev. D 100, 043540 (2019), arXiv:1906.07735 [astro-ph.CO].
- [51] Annalisa Pillepich, Cristiano Porciani, and Sabino Matarrese, "The bispectrum of redshifted 21 centimeter fluctuations from the dark ages," The Astrophysical Journal 662, 1–14 (2007).
- [52] Julian B. Muñ oz, Yacine Ali-Haïmoud, and Marc Kamionkowski, "Primordial non-gaussianity from the bispectrum of 21-cm fluctuations in the dark ages," Physical Review D 92 (2015), 10.1103/physrevd.92.083508. Physical Review D 100 (2019), 10.1103/physrevd.100.023506[53] P. Daniel Meerburg et al., "Primordial non-gaussianity,"
 - (2019).[54] G. "Foregrounds Bernardi etal., for obline," servations cosmological 21 of the cmAstronomy & amp Astrophysics 500, 965–979 (2009).
 - "Foregrounds Bernardi et al., for obline," servations of the cosmological 21 cm Astronomy & amp Astrophysics 522, A67 (2010).
 - [56] J. C. Pober et al., "THE IMPORTANCE OF WIDE-FIELD FOREGROUND REMOVAL FOR 21 cm COSMOLOGY: A DEMONSTRATION WITH EARLY MWA EPOCH OF REIONIZATION OBSER-VATIONS," The Astrophysical Journal 819, 8 (2016).
 - [57] T. Lucas Makinen et al., "deep21: a deep learning method for 21 cm foreground removal," Journal of Cosmology and Astroparticle Physics 2021, 081 (2021).
 - [58] Adrian Liu, Max Tegmark, Judd Bowman, Jacqueand Matias Zaldarriaga, "An imline Hewitt,

proved method for 21-cm foreground removal,"

- [59] Yousef Abou El-Neaj et al., "AEDGE: Atomic experiment for dark matter and gravity exploration in space," EPJ Quantum Technology 7 (2020), 10.1140/epjqt/s40507-020-0080-
- "For-[60] Muhammad A. Latif and Andrea Ferrara, seeds," mation black hole of supermassive Publications of the Astronomical Society of Australia 33 (2016), 10.
- et "Binary Carlo al., black holes the pair instability mass in gap," Monthly Notices of the Royal Astronomical Society 497, 1043–1049
- [62] Ajit Kumar Mehta et al., "Observing intermediate-mass black holes and the upper stellar-mass gap with LIGO and virgo," The Astrophysical Journal 924, 39 (2022).
- [63] Davide Gerosa and Maya Fishbach, "Hierarchical merg-
- - wave signatures," Nature Astronomy 5, 749–760 (2021). [64]Vivian Poulin, Pasquale D. Serpico, Francesca Calore,
 - Sé bastien Clesse, and Kazunori Kohri, "CMB bounds
 - - Physical Review D 96 (2017), 10.1103/physrevd.96.083524.

- [65] Joel Bergé al., "Microscope first et constraints on the violation of the weak equivalence principle by а light scalardilaton," Physical Review Letters 120 (2017), 10.1103/PhysRevLett.1278]4R103/u
- David E. Kaplan, [66] Peter W. Graham, Jeremy Mardon, Surjeet Rajendran, and William A. Terrano, "Dark matter direct detection with accelerometers," Phys. Rev. D 93, 075029 (2016), arXiv:1512.06165 [hep-ph].
- [67] Tanmay Kumar Poddar, Subhendra Mohanty, and Soumya Jana, "Vector gauge boson radiation from compact binary systems in a gauged L_{μ} - L_{τ} scenario," Physical Review D 100 (2019), 10.1103/physrevd.100.123023.
- [68] Clifford M. Will. "Bounding $_{\mathrm{the}}$ mass of graviton gravitational-wave the using observations of inspiralling compact binaries," Physical Review D 57, 2061–2068 (1998).
- [69] V. Alan KosteleckAce and Matthew Mewes, "Testing local lorentz invariance with gravitational waves," Physics Letters B 757, 510–514 (2016).
- [70] Kazunari Eda, Yousuke Itoh, Sachiko Kuroyanagi, and Joseph Silk, "New probe of dark-matter properties: Gravitational waves from an intermediate-mass black hole embedded in a dark-matter minispike," Physical Review Letters 110 (2013), 10.1103/physrevlett.110.221100ferometric sensor (magis-100)," Quantum Science and
- [71] Kazunari Eda, Yousuke Itoh, Sachiko Kuroyanagi, and Joseph Silk, "Gravitational waves probe of dark matter minispikes," as a Physical Review D **91** (2015), 10.1103/physrevd.91.044045.
- [72] Bradley J. Kavanagh, David A. Nichols, Gianfranco Bertone, and Daniele Gaggero, "Detecting dark matter around black holes with gravitational waves: Effects of dark-matter dynamics on the gravitational waveform," Physical Review D 102 (2020), 10.1103/physrevd.102.083006.
- [73] Adam Coogan, Gianfranco Bertone, Daniele Gaggero, Bradley J. Kavanagh, and David A. Nichols, "Measuring the dark matter environments of black hole binaries with gravitational waves," Phys. Rev. D 105, 043009 (2022), arXiv:2108.04154 [gr-qc].
- [74] Ken K. Y. Ng, Maximiliano Carl-Johan Isi. and Salvatore "Multiband Haster. Vitale, gravitational-wave searches for ultralight bosons," Physical Review D 102 (2020), 10.1103/physrevd.102.083020[90]
- [75] Ruiyu Zhou, Ligong Bian, Huai-Ke Guo, and Yongcheng Wu, "Gravitational wave and collider searches for electroweak symmetry breaking patterns," Physical Review D **101** (2020), 10.1103/physrevd.101.035011.
- [76] Ryusuke Jinno and Masahiro Takimoto, "Gravitational waves from bubble collisions: An analytic derivation," Physical Review D 95 (2017), 10.1103/physrevd.95.024009.

- [77] Sé bastien Galtier and Sergey V. Nazarenko, "Direct evidence of a dual cascade in gravitational wave turbulence," Physical Review Letters **127** (2021), 10.1103/physrevlett.127.131101
- Zhou Ligong Bian, and "Gravitational waves from cosmic strings and first-order phase trans (2020).
- [79] Ivan Alonso et al., "Cold atoms in space: Community workshop summary and proposed road-map," (2022), 10.48550/ARXIV.2201.07789.
- [80] Baptiste Battelier al., "Exploring et the physical foundations of the universe with space tests of the equivalence principle," Experimental Astronomy 51, 1695–1736 (2021).
- [81] Benjamin Elder, Justin Khoury, Philipp Haslinger, Matt Jaffe, Holger Müller, and Paul Hamilton, "Chameleon dark energy and atom interferometry," Physical Review D 94 (2016), 10.1103/physrevd.94.044051.
- [82] D. O. Sabulsky et al., "Experiment to detect dark energy forces using atom interferometry," Physical Review Letters 123 (2019), 10.1103/physrevlett.123.061102
- [83] L. Badurina et al., "AION: An Atom Interferometer Observatory and Network," JCAP 05, 011 (2020), arXiv:1911.11755 [astro-ph.CO].
- [84] Mahiro Abe et al., "Matter-wave atomic gradiometer in-
- Technology 6, 044003 (2021).
- [85] B. Canuel et al., "Exploring gravity with the MIGA large scale atom interferometer," Sci. Rep. 8, 14064 (2018), arXiv:1703.02490 [physics.atom-ph].
- [86] Ming-Sheng Zhan et al., "ZAIGA: Zhaoshan longbaseline atom interferometer gravitation antenna," Int. J. Mod. Phys. D 29, 1940005 (2020).
- [87] B Canuel et al., "ELGAR a European Laboratory for Gravitation and Atom-interferometric Research." Class. Quantum Gravity **37**, 225017 (2020).
- [88] N. L. Lafave and T. L. Wilson, "The Role of the Moon in Gravitational Wave Astronomy," in Lunar and Planetary Science Conference, Lunar and Planetary Science Conference, Vol. 23 (1992) p. 751.
- [89] K. "Gravitational-Jani and Α. Loeb, for wave lunar observatory cosmology," Journal of Cosmology and Astroparticle Physics 2021, 044 (2021).
- Basudeb Dasgupta, Ranjan Laha, and Anupam Ray, "Low mass black holes from dark core collapse," Phys. Rev. Lett. **126**, 141105 (2021).
- [91] Basudeb Dasgupta, Aritra Gupta, and Anu-Ray, "Dark matter capture celespam in self-interactions, $_{\rm tial}$ objects: light mediators, and complementarity with direct detection," Journal of Cosmology and Astroparticle Physics 2020, 023–023 (202