

Study of chaos in rotating galaxies using extended force-gradient symplectic methods

Li-Na Zhang^{1,2,3}, Wen-Fang Liu¹, and Xin Wu^{1,2*}

1. School of Mathematics, Physics and Statistics & Center of Application and Research of Computational Physics, Shanghai University of Engineering Science, Shanghai 201620, China
2. School of Physical Science and Technology, Guangxi University, Nanning 530004, China
3. School of Physics and Electronics, Hunan Normal University, Changsha 410081, China

We take into account the dynamics of three types of models of rotating galaxies in polar coordinates in a rotating frame. Due to non-axisymmetric potential perturbations, the angular momentum varies with time, and the kinetic energy depends on the momenta and spatial coordinate. The existing explicit force-gradient symplectic integrators are not applicable to such Hamiltonian problems, but the recently extended force-gradient symplectic methods proposed in a previous work are. Numerical comparisons show that the extended force-gradient fourth-order symplectic method with symmetry is superior to the standard fourth-order symplectic method but inferior to the optimized extended force-gradient fourth-order symplectic method in accuracy. The optimized extended algorithm with symmetry is used to explore the dynamical features of regular and chaotic orbits in these rotating galaxy models. The gravity effects and the degree of chaos increase with an increase of the number of the radial terms in the series expansions of the potential. There are similar dynamical structures of regular and chaotic orbits in the three types of models for the same number of the radial terms in the series expansions, energy and initial conditions.

Keywords: symplectic integrators; chaos; galaxies; gravity

I. INTRODUCTION

Chaos in a dynamical system means that the final state of the dynamical system displays exponentially sensitive dependence on the initial state. Based on the importance of chaos, many studies have focused on the subject of chaos in the solar system [1,2] and galactic dynamics [3-7].

Regular or chaotic motions of particles in galactic dynamics may affect the fraction of mass. The authors of Refs. [3,5-7] investigated mass components of non-rotating N-body models of *elliptical galaxies* in ordered and in chaotic motion. Voglis et al. [3] found that the fraction of mass in chaotic motion is about 24% of the total mass in one non-rotating triaxial equilibrium model with smooth centers, and 32% in another non-rotating triaxial equilibrium model with smooth centers. This shows that the spatial distribution of the mass in chaotic motion is in disagreement with that in ordered motion. Muzzio et al. [7] pointed out that the fraction of mass in chaotic motion of about 53% in models of non-rotating galaxies with smooth centers. On the other hand, the fraction of mass in ordered or chaotic motion was also studied in *spiral galaxies*. Voglis et al. [8] further showed that rotation leads to increasing the fraction of mass in chaotic motion (up to the level of $\approx 65\%$) and shifting the Lyapunov numbers to larger values in N-body models of rotating galaxies. In other words, the extent of chaos is substantially enhanced by the rotation, and the fractions of mass in chaotic motion in the rotating models are larger than those in the non-rotating models. The spatial distributions of the dynamical structures along the spiral arms at the ends of the bar in the barred-spiral galaxy correspond to those of particles with masses in regular and in chaotic orbits. In fact, the mass in chaotic motion can almost completely form spiral arms emanating from the neighborhood of the Lagrangian points L_1 and L_2 at the ends of the bar in a barred-spiral galaxy. The distribution of mass must be associated with gravitational forces of particles. The gravity is the main driving mechanism for the formation and the stability of spiral arms in galaxies. The authors of Refs. [9,10] developed the Moser analytic series representing the invariant manifolds near the unstable Lagrangian equilibrium points L_1 , L_2 in a rotating barred galaxy. In this way, these series can represent the spiral arms, which are density waves and are composed of chaotic orbits. Besides the analytical theory, other methods such as the specific finite time Lyapunov characteristic number, the smaller alignment index, the surface of section and the frequency analysis were used to classify the orbits in regular and chaotic cases in barred galaxies [11-15].

The onset of chaos in two-dimensional Hamiltonian systems of rotating galaxies in the disc plane in polar coordinates [10] is due to the inclusion of non-axisymmetric potential perturbations. These perturbations cause the angular momentum with respect to the angle to vary with time and therefore the Hamiltonian systems are not integrable. This non-integrability leads to chaos under some circumstances. Reliable numerical results are always required to detect

*Electronic address: wuxin1134@sina.com

the chaotical behavior. In some cases, extremely long integration times are also required. The adopted computational schemes for the long-term integration of the Hamiltonian systems become crucial to reach better stability and higher precisions. The proper choice of the integrators should naturally be symplectic schemes, which preserve the symplectic nature of Hamiltonian dynamics [16,17]. Symplectic methods are a class of geometric integration algorithms [18] and make the local error in the total energy not grow with time. There are standard symplectic methods [19,20] that require some evaluations of the force, and force-gradient symplectic integrators [16,21-24] that require some evaluations of force gradient in addition to several evaluations of the force. Because of good long-term behavior, the standard symplectic methods have been used in the solar system [1, 25] and black hole spacetimes [26-31]. They are also suitable for the two-dimensional Hamiltonian systems of rotating galaxies in polar coordinates. However, the force-gradient symplectic integrators are not applicable to such Hamiltonian problems because the kinetic energies of the Hamiltonian systems depend on the momenta and spatial coordinates. In fact, they are only adapted to the integrations of Hamiltonian systems, where the kinetic energies are quadratic functions of momenta and the potential energies are functions of coordinates. Energy errors of the force-gradient symplectic integrators are several orders of magnitude smaller than those of the standard symplectic algorithms of the same order, as was confirmed in the literature [22-24]. Recently, our group extended these force-gradient integrators to the explicitly integrable kinetic energies, which are not only quadratic functions of momenta but also depend on coordinates [32]. When the original force-gradient operator is adjusted appropriately, the adjusted operator lacks the concept of force gradient and belongs to the momentum operator like the operator corresponding to the potential. As a result, the existing explicit force-gradient symplectic integrators [21-24] are still available in the extended Hamiltonian systems. The extended force-gradient symplectic integrators do not alter their symmetry and time reversibility compared with the existing force-gradient algorithms. The authors of Ref. [32] used a modified Hénon-Heiles system and a spring pendulum as two toy models to test the numerical performance of the extended algorithms. They showed that the fourth-order Forest-Ruth standard symplectic method does not give true dynamical properties of order and chaos to the modified Hénon-Heiles system under some circumstances, whereas the fourth-order extended force-gradient symplectic methods do. The obtained results are because the Forest-Ruth method performs poorer accuracy than the extended force-gradient algorithms. In fact, the optimized fourth-order extended force-gradient symplectic methods have energy errors that are three orders of magnitude smaller than those of the Forest-Ruth method.

Note that the extended force-gradient symplectic methods proposed in Ref. [32] were shown to have good numerical performance in the simulations of the two toy models. Now, we wonder whether the extended algorithms still exhibit excellent performance in the application of real astronomical and astrophysical problems. To evaluate the performance of the extended force-gradient symplectic methods applicable to the three types of models of rotating barred galaxies in Refs. [8,10] is one of the main aims in the present paper. On the other hand, we are interested in studying a distribution of dynamical structures regarding regular and chaotic orbits along the radial direction for a given number of the radial terms of the potential in one of the models using the techniques of Poincaré surface of section and fast Lyapunov indicators (FLIs) [33]. In this way, we desire to know how the number of the radial terms of the potential affects chaos in one of the models. We also desire to know what differences in the dynamics exist among the three types of models with the same number of the radial terms of the potential, energy and initial conditions.

To implement the three aims, we organize this paper as follows. In Section 2, we introduce three types of models of rotating barred galaxies. In Section 3, we apply the extended fourth-order force-gradient symplectic methods to Model A4 and evaluate the performance of these algorithms. In section 4, we explore the dynamical structures in these models using the techniques of Poincaré surface of section and FLIs. Finally, the main results are summarized in Section 5.

II. MODELS OF ROTATING BARRED GALAXIES

The motion of a test particle of mass $m_p = 1$ in the plane of a galaxy with a rotating bar is described in polar coordinates (r, ϕ) in the rotating frame by a two-dimensional Hamiltonian [8,10]

$$H = \frac{p_r^2}{2} + \frac{p_\phi^2}{2r^2} - \Omega p_\phi + \Phi(r, \phi). \quad (1)$$

The above notations are given here. Ω is a pattern speed of the rotating frame. p_r represents a canonical momentum vs the coordinate r , and p_ϕ is a canonical momentum of the coordinate ϕ corresponding to the angular momentum in the rest frame. $\Phi(r, \phi)$ denotes the gravitational potential of the galaxy in the rotating frame and satisfies the Poisson equation

$$\nabla^2 \Phi = 4\pi G \rho, \quad (2)$$

where G is the constant of gravity and ρ stands for the density of matter.

A complete expression of the solution of the Poisson equation is long and complicated. Harsoula et al. [10] took a simple solution of Eq. (2) as the potential for the $m = 2$ mode of the galactic bar

$$\Phi(r, \phi) = \Phi_0(r) + \Phi_1(r) \cos 2\phi + \Phi_2(r) \sin 2\phi, \quad (3)$$

where $\Phi_0(r)$, $\Phi_1(r)$ and $\Phi_2(r)$ are functions of r . If $\Phi_1(r) = \Phi_2(r) = 0$, then the potential $\Phi_0(r) = \Phi(r, \phi)$ is axisymmetric and the angular momentum p_ϕ is conserved. When $\Phi_1(r) \neq 0$ or $\Phi_2(r) \neq 0$, the potential $\Phi(r, \phi)$ is non-axisymmetric and the angular momentum p_ϕ is not conserved. Namely, the second and third terms of Eq. (3) act as an $m = 2$ mode of the non-axisymmetric potential perturbation. The three potential functions are given in Ref. [10] by

$$\Phi_0(r) = -\frac{1}{R} \left(A_{00} + \frac{1}{4}A_{20} - \frac{3}{2}A_{22} \right), \quad (4)$$

$$\Phi_1(r) = -\frac{3}{2R} \left(\frac{1}{2}A_{20} + A_{22} \right), \quad (5)$$

$$\Phi_2(r) = \frac{3}{2R}A_{21}, \quad (6)$$

where R is a numerical constant for the description of the size of the system. The size is regarded as the N-body code boundary corresponding to the solutions of the Poisson equation matched with the solutions of the Laplace equation. The other notations such as A_{00} are expanded in terms of the series of spherical Bessel functions j_0 and j_2 [34]:

$$A_{00} = \sum_{i=0}^n [B_{i00} \cdot j_0(\xi_{i0})], \quad (7)$$

$$A_{20} = \sum_{i=0}^n [B_{i20} \cdot j_2(\xi_{i2})], \quad (8)$$

$$A_{21} = \sum_{i=0}^n [C_{i21} \cdot j_2(\xi_{i2})], \quad (9)$$

$$A_{22} = \sum_{i=0}^n [B_{i22} \cdot j_2(\xi_{i2})]. \quad (10)$$

In the above series expansions of the potential, n represents the number of the radial terms; that is, each of the functions like A_{00} in the potential has $(n + 1)$ terms. In addition, $\xi_{il} = a_{il}r/R$. For $l = 0$, $a_{i0} = (i + 1/2)\pi$ is the $(i + 1)$ th root of the equation $j_{-1}(a_{i0}) = 0$. For $l = 2$, a_{i2} is the solution of the equation $\tan(a_{i2}) = a_{i2}$. The solution a_{i2} should be restricted in the range $(i - 1/2)\pi < a_{i2} < (i + 1/2)\pi$ and is solved by the Newton iteration method.

The coefficients B_{i00} , B_{i20} , C_{i21} and B_{i22} in Eqs. (7)-(10) are obtained through the N-body code on the positions of the N-body particles and Eq. (2). They are different for three types of galactic models, which are called as models A, B and C in Ref. [10]. The properties of the three models were described in the papers [8-10]. Here are some of them. Firstly, the total angular momentum is larger in model B than in model A, but smaller than in model C. This leads to decreasing the size of a rotating central bar formed by density waves and increasing the density in the region of the bar along the sequence of the models A, B and C. Secondly, the pattern speed of the bar in each of the models becomes smaller, whereas the corotation radius gets larger at the end of a Hubble time. Thirdly, the degree of chaos is enhanced by rotation increasing the fraction of mass in chaotic motion. Fourthly, mass in chaotic motion almost completely dominates the formation of spiral arms. Fifthly, invariant manifolds of all unstable periodic orbits near and beyond corotation support both the outer edge of the bar and the spiral arms. Harsoula provided the coefficients in the three models to the first author of the present paper via private communication. The coefficients with $n = 19$ are listed in Tables 1-3. The 20 coefficients B_{i00} , the 20 coefficients B_{i20} and the 40 coefficients C_{i21} , B_{i22} ($i = 0, \dots, 19$) in each model are associated to monopole terms, quadrupole terms and triaxial terms, respectively. In practice, each group of the coefficients determine the potential $\Phi(r, \phi)$. Different potentials $\Phi(r, \phi)$ correspond to different galactic models. Hereafter, the three models for $n = 19$ respectively correspond to models A19, B19 and C19. Similarly, models A9, B9 and C9 are called when $n = 9$; models A4, B4 and C4 are also called when $n = 4$.

The unit systems are those of Refs. [8,10]. The half mass radius R_h is used as a scaling unit of length, i.e. $r_{scal} = r/R_h$, where r stands for the real distance and r_{scal} is the scaling radial distance. For convenience, the scaled radial distance is still written as r in the later discussions. The half mass radius is $R_h = 0.1006$ for Model A, $R_h = 0.0926$ for Model B and $R_h = 0.1167$ for Model C. The size of the system is $R = 0.85$. Here, one unit of length is 8kpc. The unit of time is the half mass crossing time $T_{hmct} = [2R_h^3/(GM)]^{1/2} = t_{HUB}/300$, where t_{HUB} represents

TABLE I: Coefficients of the series (7)-(10) in Model A.

i	0	1	2	3	4	5	6	7	8	9
<i>Bi00</i>	250670.00	214640.00	175380.00	146580.00	126550.00	112500.00	102250.00	93269.00	84659.00	76738.00
<i>Bi20</i>	16599.00	25348.00	25460.00	25455.00	27770.00	31416.00	35390.00	38187.00	39353.00	39760.00
<i>Bi21</i>	616.35	1477.80	3093.20	4784.80	5723.80	5186.10	3564.00	1766.40	220.39	-632.86
<i>Bi22</i>	-5808.00	-9799.20	-11023.00	-10369.00	-8597.30	-6413.80	-4878.40	-4209.20	-3989.60	-3932.60
i	10	11	12	13	14	15	16	17	18	19
<i>Bi00</i>	69503.00	62789.00	56553.00	50735.00	45419.00	40595.00	36128.00	31941.00	27988.00	24271.00
<i>Bi20</i>	40303.00	41137.00	41938.00	42164.00	41642.00	40746.00	39745.00	38828.00	37960.00	36816.00
<i>Bi21</i>	-833.16	-816.34	-786.25	-741.98	-684.44	-601.88	-463.25	-324.97	-283.78	-344.23
<i>Bi22</i>	-3882.30	-3768.40	-3624.80	-3470.20	-3355.40	-3299.00	-3289.50	-3298.10	-3315.100	-3304.00

TABLE II: Coefficients of the series (7)-(10) in Model B.

i	0	1	2	3	4	5	6	7	8	9
<i>Bi00</i>	249850.00	208750.00	166240.00	136810.00	119540.00	109650.00	102040.00	94581.00	87464.00	80875.00
<i>Bi20</i>	19761.00	30778.00	32613.00	28562.00	24516.00	25984.00	30721.00	34872.00	37154.00	38234.00
<i>Bi21</i>	-674.22	-2590.80	-4227.30	-4246.90	-1974.70	118.46	87.73	-1623.40	-3752.10	-5087.90
<i>Bi22</i>	-6781.10	-10962.00	-11493.00	-9732.80	-7244.90	-5062.20	-3654.80	-2783.10	-2305.50	-2196.80
i	10	11	12	13	14	15	16	17	18	19
<i>Bi00</i>	74509.00	68185.00	61997.00	56146.00	50708.00	45630.00	40856.00	36367.00	32180.00	28326.00
<i>Bi20</i>	39040.00	40062.00	41339.00	42345.00	42784.00	42869.00	42847.00	42584.00	41768.00	40414.00
<i>Bi21</i>	-5447.70	-5355.50	-5293.10	-5374.40	-5514.50	-5533.80	-5337.50	-5081.30	-4882.80	-4719.30
<i>Bi22</i>	-2336.40	-2490.40	-2534.30	-2495.10	-2474.00	-2468.80	-2445.60	-2417.60	-2425.40	-2463.00

a Hubble time. The pattern speeds of the rotating frame in the three models A, B and C are taken as $\Omega_A = 5886.65$, $\Omega_B = 6010.36$ and $\Omega_C = 6137.14$, which correspond to $20 \sim 25$ km sec⁻¹ kpc⁻¹ in real units.

As is mentioned above, the second and third terms of Eq. (3) destroy the axial-symmetry of the system (1) such that the angular momentum p_ϕ varies with time. Thus, no additional constants of motion but the Hamiltonian H of Eq. (1) exists. This implies the nonintegrability of the system (1) with two degrees of freedom in a four-dimensional phase space. A numerical method is a good tool to solve this nonintegrable problem.

III. A CHOICE OF NUMERICAL INTEGRATOR

A prior choice to an integrator for a long-term integration of the Hamiltonian (1) is naturally a symplectic method with the conservation of the Hamiltonian flow. Following this idea, we consider the application of symplectic integration to this Hamiltonian problem.

A. Generalized force-gradient symplectic integrators

For convenience, we take $\mathbf{p} = (p_1, p_2) = (p_r, p_\phi)$ and $\mathbf{q} = (q_1, q_2) = (r, \phi)$. We rewrite the Hamiltonian (1) as

$$H(\mathbf{p}, \mathbf{q}) = T(\mathbf{p}, q_1) + \Phi(\mathbf{q}), \quad (11)$$

TABLE III: Coefficients of the series (7)-(10) in Model C.

i	0	1	2	3	4	5	6	7	8	9
<i>Bi00</i>	249390.00	205340.00	159420.00	129900.00	114510.00	106340.00	100510.00	95098.00	89366.00	83133.00
<i>Bi20</i>	13998.00	21217.00	22367.00	25113.00	29066.00	30605.00	30582.00	31466.00	33769.00	36355.00
<i>Bi21</i>	89.08	1336.80	4085.00	5662.00	3913.40	969.22	-1143.70	-2174.50	-2900.50	-3381.60
<i>Bi22</i>	-8969.80	-14504.00	-14213.00	-10170.00	-5852.70	-3207.10	-2049.70	-1654.10	-1631.90	-1714.40
i	10	11	12	13	14	15	16	17	18	19
<i>Bi00</i>	76784.00	70724.00	65078.00	59703.00	54546.00	49594.00	44931.00	40471.00	36143.00	31955.00
<i>Bi20</i>	38644.00	40349.00	41348.00	42023.00	42823.00	43574.00	43919.00	43850.00	43519.00	42874.00
<i>Bi21</i>	-3685.00	-3800.20	-3575.40	-3274.30	-3104.70	-2945.90	-2835.40	-2807.90	-2761.70	-2731.70
<i>Bi22</i>	-1706.80	-1606.50	-1571.10	-1578.70	-1567.60	-1505.90	-1508.40	-1617.50	-1786.00	-1900.20

where the kinetic energy T is a function of the momenta \mathbf{p} and coordinate r

$$T(\mathbf{p}, q_1) = \frac{p_r^2}{2} + \frac{p_\phi^2}{2r^2} - \Omega p_\phi. \quad (12)$$

The lie derivative operators of T and Φ are defined as

$$\mathcal{A} = \{, T\} = \sum_{i=1}^2 \left(T_{p_i} \frac{\partial}{\partial q_i} - T_{q_i} \frac{\partial}{\partial p_i} \right) = p_r \frac{\partial}{\partial r} + \left(\frac{p_\phi}{r^2} - \Omega \right) \frac{\partial}{\partial \phi} + \frac{p_\phi^2}{r^3} \frac{\partial}{\partial p_r}, \quad (13)$$

$$\mathcal{B} = \{, \Phi\} = -\frac{\partial \Phi}{\partial r} \frac{\partial}{\partial p_r} - \frac{\partial \Phi}{\partial \phi} \frac{\partial}{\partial p_\phi}, \quad (14)$$

where the symbols $\{, \}$ represent the Poisson brackets. Obviously, T and Φ can be exactly, analytically solvable, and \mathcal{A} , \mathcal{B} correspond to their solvers.

The solvers \mathcal{A} and \mathcal{B} can symmetrically compose a second-order symplectic leapfrog integrator

$$\text{M2} = e^{\frac{\tau}{2}\mathcal{A}} e^{\tau\mathcal{B}} e^{\frac{\tau}{2}\mathcal{A}} = e^{\tau(\mathcal{A}+\mathcal{B}) - \frac{\tau^3}{12}[\mathcal{B}, [\mathcal{A}, \mathcal{B}]] + \frac{\tau^3}{24}[\mathcal{A}, [\mathcal{B}, \mathcal{A}]]}, \quad (15)$$

where τ is a time step, and two commutators are

$$\mathcal{C} = [\mathcal{B}, [\mathcal{A}, \mathcal{B}]] = [\mathcal{B}, \mathcal{A}\mathcal{B} - \mathcal{B}\mathcal{A}] = 2\mathcal{B}\mathcal{A}\mathcal{B} - \mathcal{B}\mathcal{B}\mathcal{A} - \mathcal{A}\mathcal{B}\mathcal{B}, \quad (16)$$

$$\mathcal{D} = [\mathcal{A}, [\mathcal{B}, \mathcal{A}]] = 2\mathcal{A}\mathcal{B}\mathcal{A} - \mathcal{A}\mathcal{A}\mathcal{B} - \mathcal{B}\mathcal{A}\mathcal{A}. \quad (17)$$

In the second line of Eq. (15), the first term $\tau(\mathcal{A} + \mathcal{B})$ corresponds to the numerical solution of the Hamiltonian (1), and the second and third terms correspond to local truncation errors of the numerical solution remaining at an order $\mathcal{O}(\tau^3)$. Because of such local truncation errors, the algorithm M2 can give a second-order accuracy to the explicit numerical solution. In terms of the solvers \mathcal{A} and \mathcal{B} , a fourth-order symplectic method of Forest and Ruth [19] is established by

$$\text{M4} = e^{\alpha\tau\mathcal{A}} e^{\beta\tau\mathcal{B}} e^{(\frac{1}{2}-\alpha)\tau\mathcal{A}} e^{(1-2\beta)\tau\mathcal{B}} e^{(\frac{1}{2}-\alpha)\tau\mathcal{A}} e^{\beta\tau\mathcal{B}} e^{\alpha\tau\mathcal{A}}, \quad (18)$$

where $\beta = 1/(2 - \sqrt[3]{2})$ and $\alpha = \beta/2$.

If r^2 in the second term of Eq. (12) is absent, \mathcal{A} of Eq. (13) is a position operator which acts on only the position coordinates. $\mathcal{C} = 2\mathcal{B}\mathcal{A}\mathcal{B}$ is similar to the momentum operator \mathcal{B} , which acts on only the momenta. In fact, \mathcal{C} is an exactly, analytically solvable operator corresponding to the force gradient of the gravitational potential. However, \mathcal{D} is a momentum and position mixed operator and is not an exactly analytical solver. In view of these facts, a composition of \mathcal{B} and \mathcal{C} appears in a class of explicit symplectic integrators, called the force-gradient symplectic integration algorithms [16,21-24].

Now, r^2 in the second term of Eq. (12) remains. Although \mathcal{A} of Eq. (13) is a momentum and position mixed operator, it is still an exactly analytical solver in this case. Is \mathcal{C} an exactly analytical solver? We said yes as an answer to this question in our previous work [32]. In fact, the answer can be given through the operators acting on the momenta and positions in the second line of Eq. (16). It is easy to check that $\mathcal{A}\mathcal{B}\mathcal{B}q_i = \mathcal{A}\mathcal{B}\mathcal{B}p_i \equiv 0$, $\mathcal{B}\mathcal{A}\mathcal{B}q_i \equiv 0$, $\mathcal{B}\mathcal{B}\mathcal{A}q_i \equiv 0$, and

$$\mathcal{B}\mathcal{A}\mathcal{B}p_i = \sum_{j=1}^2 \sum_{k=1}^2 \Phi_{q_i q_j} \Phi_{q_k} T_{p_j p_k}, \quad (19)$$

$$\mathcal{B}\mathcal{B}\mathcal{A}p_i = -\sum_{j=1}^2 \sum_{k=1}^2 \Phi_{q_j} \Phi_{q_k} T_{q_i p_j p_k}, \quad (20)$$

where $\Phi_{q_j} = \partial\Phi/\partial q_j$, $\Phi_{q_k} = \partial\Phi/\partial q_k$, and $\Phi_{q_i q_j} = \frac{\partial^2\Phi}{\partial q_i \partial q_j}$, $T_{p_j p_k} = \frac{\partial^2 T}{\partial p_j \partial p_k}$, $T_{q_i p_j p_k} = \frac{\partial^3 T}{\partial q_i \partial p_j \partial p_k}$. Clearly, \mathcal{C} is also a momentum operator:

$$\mathcal{C} = 2\mathcal{B}\mathcal{A}\mathcal{B} - \mathcal{B}\mathcal{B}\mathcal{A} = \sum_{i=1}^2 \sum_{j=1}^2 \sum_{k=1}^2 (2\Phi_{q_i q_j} \Phi_{q_k} T_{p_j p_k} + \Phi_{q_j} \Phi_{q_k} T_{q_i p_j p_k}) \frac{\partial}{\partial p_i}.$$

The momentum operator \mathcal{C} of Eq. (21) is no longer the force gradient of the gravity potential, but can be regarded as an extension of the force-gradient factor in Refs. [16,21-24]. Without doubt, it is an exactly analytical solver similar to the momentum operator \mathcal{B} .

Symmetric composition methods of the three operators \mathcal{A} , \mathcal{B} and \mathcal{C} can yield various explicit symplectic algorithms. For instance, a five-stage fourth-order symplectic method including the operator \mathcal{C} was proposed by Chin [22] in the form

$$N4 = e^{\frac{\tau}{2}\left(1-\frac{1}{\sqrt{3}}\right)\mathcal{A}} e^{\frac{\tau}{2}\left(\mathcal{B}+\frac{\tau^2}{24}(2-\sqrt{3})\mathcal{C}\right)} e^{\frac{\tau}{\sqrt{3}}\mathcal{A}} e^{\frac{\tau}{2}\left(\mathcal{B}+\frac{\tau^2}{24}(2-\sqrt{3})\mathcal{C}\right)} e^{\frac{\tau}{2}\left(1-\frac{1}{\sqrt{3}}\right)\mathcal{A}}.$$

Omelyan et al. [24] also constructed a seven-stage fourth-order optimized symplectic algorithms with the inclusion of the operator \mathcal{C} :

$$N4P = e^{\theta\tau\mathcal{A}} e^{\lambda\tau[\mathcal{B}+(2\xi+\chi)\tau^2\mathcal{C}]} e^{(1-2\theta)\frac{\tau}{2}\mathcal{A}} e^{(1-2\lambda)\tau[\mathcal{B}+(2\xi+\chi)\tau^2\mathcal{C}]} e^{(1-2\theta)\frac{\tau}{2}\mathcal{A}} e^{\lambda\tau[\mathcal{B}+(2\xi+\chi)\tau^2\mathcal{C}]} e^{\theta\tau\mathcal{A}}, \quad (21)$$

where the time coefficients are

$$\begin{aligned} \theta &= 0.1159953608486416 \times 10^0, \\ \lambda &= 0.2825633404177051 \times 10^0, \\ \chi &= 0.3035236056708454 \times 10^{-2}, \\ \xi &= 0.1226088989536361 \times 10^{-2}. \end{aligned}$$

The concept of optimization is the time coefficients minimizing the norm of the leading term of fifth-order truncation errors.

The construction mechanisms of the algorithms with the extended operator \mathcal{C} was discussed in the previous work [32]. In fact, the extended force-gradient algorithms acting on the original system (1) are equivalent to the standard symplectic methods (like the method M4 without the extended operator \mathcal{C}) acting on the modified Hamiltonian systems.

B. Numerical tests

Model A4 is used as a test model to evaluate the performance of the algorithms M4, N4 and N4P. For comparison, an eighth- and ninth-order Runge-Kutta-Fehlberg integrator [RKF8(9)] with adaptive step sizes is also employed. The time step is $\tau = 0.0005 \times T_{hmct} = 0.0005 \times t_{hub}/300 = 0.0005/82.4/300 \approx 2.0 \times 10^{-8}$ (sec·Mpc/km) = 2.0×10^{-5} (sec·kpc/km) $\approx 2.0 \times 10^{-5}$ (9.5×10^8 y) = 0.19×10^5 y. In short, $\tau \approx 2.0 \times 10^{-5}$ (sec·kpc/km) is used in our codes. The authors of Ref. [8] pointed out that the time step is a good choice because the hysteresis between the bar of the potential and the bar of the real density of particles during the time is minimal and the cumulative effect of a numerical retarding torque in a Hubble time is small.

The energy of the system (1) is $E = H = -7 \times 10^6$. The initial conditions are $r = 0.164/R_h = 0.164/0.1006$, $\phi = 1.89\pi$, $p_r = 0$. The initial momentum $p_\phi > 0$ is determined by $E = H$. Fig. 1(a) plots the relative energy errors $|\Delta H/H_0| = |(E_t - E)/E|$ for these several algorithms, where E_t is the numerical energy at time t . Here, each algorithm has 2.5×10^8 steps, which correspond to time $t \approx 5056.63$ (sec·kpc/km). The three symplectic methods M4, N4 and N4P show no secular growth in the energy errors. This is an inherent advantage of these symplectic methods. Among the three symplectic integrators, the standard fourth-order symplectic method M4 without the extended operator \mathcal{C} performs the poorest accuracy, whereas the fourth-order optimized symplectic algorithm N4P with the extended force-gradient operator \mathcal{C} exhibits the best accuracy. These numerical results show that the inclusion of the extended force-gradient operator has an advantage over the exclusion of the extended force-gradient operator in the accuracy. The optimized method is also better than the corresponding non-optimized method. Unlike these symplectic methods, the non-symplectic integrator RKF8(9) makes the energy error grow with time. RKF8(9) is superior to N4P in the accuracy, but inferior to N4P in the computational efficiency, as is shown in Table 4. Taking the solutions of RKF8(9) as reference solutions, we obtain the relative position errors of the three symplectic algorithms M4, N4 and N4P in Fig. 1(b). As a result, the method M4 still has the largest position error, while the method N4P yields the smallest one. When many other values of the energy and initial conditions are also used, they do not affect the numerical performances of these integrators. In fact, the numerical performance of an integrator is independent of a choice of the parameters and initial conditions [16-32]. In spite of this, it is necessary to choose bounded orbits as testing the integrator's performance. Thus, an appropriate choice of the parameters and initial conditions is still necessary.

Because of the fourth-order optimized symplectic algorithm N4P with the extended force-gradient operator \mathcal{C} having the best performance in the stabilization of energy errors, it is used to study the dynamics of orbits.

TABLE IV: Relative energy errors, relative radial errors and CPU times for the algorithms in Fig. 1. Note that -12.2 denotes the error with an order of $10^{-12.2}$, and -2.1 means the error being an order of $10^{-2.1}$. 17'42'' in CPU times means 17 minutes and 42 seconds.

Method	RKF8(9)	M4	N4	N4P
$ \Delta H/E $	-12.2	-8.6	-9.5	-11.0
$ \Delta r/r_0 $	/	-2.1	-2.9	-3.4
Time	17'42''	4'30''	5'43''	6'57''

IV. REGULAR AND CHAOTIC DYNAMICS

The authors of Ref. [10] applied the Moser theory of invariant manifolds to the chaotic spiral arms. The invariant manifolds starting at the Lagrangian points L_1 , L_2 , or unstable periodic orbits around L_1 and L_2 are described in terms of series. The convergence of the series is an analytical means for studying chaotic orbits with initial conditions in the neighborhood of the invariant manifolds of the unstable point L_1 or L_2 . A domain of the convergence of the series around every Lagrangian point is a Moser domain. The intersection of the orbits inside the Moser domain with an apocentric section produces the spiral structure of this domain. The chaotic orbits with initial conditions near, but in the exterior of the boundary of the Moser domain of convergence become chaotic attractors. Such chaotic orbits are also identified by the techniques of Poincaré sections and fast Lyapunov indicators (FLIs) [33]. The method of Poincaré sections, which shows intersections of the particles' trajectories with the surface of section in phase space, can clearly describe the phase space structure of a conservative 4-dimensional system. One or several isolated points on the Poincaré sections correspond periodic orbits. Kolmogorov-Arnold-Moser (KAM) tori on the Poincaré sections correspond regular quasi-periodic orbits. If there are many plotted points that are distributed randomly in an area, the motion is chaotic. In a word, the distribution of the points in the Poincaré map can show whether or not the motion is chaotic. For ordered and chaotic orbits, the length of a deviation vector increases in completely different time rates. The FLI uses the completely different time rates to distinguish between the ordered and chaotic cases. The Moser theory of invariant manifolds for allowing to study chaotic orbits requires that the series describing the Hamiltonian dynamics near an unstable equilibrium point or an unstable periodic orbit be convergent. It may not work well for the non-convergent Birkhoff normal form series around stable invariant points, or stable periodic orbits. The techniques of Poincaré sections and FLIs for finding chaos do not have this restriction. Clearly, they can detect chaotic orbits in larger regions compared with the Moser theory of invariant manifolds. Here, we employ the techniques of Poincaré sections and FLIs to investigate the dynamical behavior of the three types of bar spiral galaxy models A, B and C. The considered orbits are not restricted to those around the unstable Lagrangian points L_1 and L_2 . We are mainly interested in comparing the differences in the dynamical behavior among these types of models. The effect of the number $(n + 1)$ of the radial terms in the series expansions of the potential on the dynamical behavior in each set of models is also considered.

A. Models A4, A9 and A19

The parameters and initial conditions are those of Fig. 1, but the initial values of r and p_ϕ are different. Model A4 exhibits different phase space structures in two ranges of the initial separations r , as is shown through Poincaré sections at the plane $\phi = \pi/2$ with $p_\phi > 0$ in Figs. 2 (a) and (b). The three orbits in Fig. 2(a) are KAM tori, which correspond to the regularity of the orbits. One orbit with the initial separation $r = 4.430/0.1006$ and another orbit with the initial separation $r = 4.424/0.1006$ in Fig. 2(b) seem to have no explicit difference from the phase space structures on the section, but they exhibit different regular and chaotic dynamical features, which can be identified clearly by means of the technique of FLI in Fig. 2(c). The FLI is that with two nearby orbits defined in Ref. [35]:

$$\text{FLI} = \log_{10} \frac{d(t)}{d(0)}, \quad (22)$$

where $d(0)$ and $d(t)$ are the distances between two nearby orbits at time 0 and t , respectively. The orbit with the initial separation $r = 4.430/0.1006$ corresponds to the FLI increasing in a power law with time $\log_{10} t$ and is regular. However, the orbit with the initial separation $r = 4.424/0.1006$ has the FLI increasing in an exponential law with time and should be chaotic. After 5×10^4 integration steps, the FLIs with completely different increasing laws with time are sensitive to distinguish the two cases of order and chaos.

The phase space structures for Model A9 in Fig. 3(a) are somewhat unlike those for Model A4 in Fig. 2. When the initial separation r is given in a small range of $1.45 < r < 1.7$ for Model A4, the orbits are regular KAM tori (not

plotted). However, the orbit with the initial separation $r = 0.153/0.1006$ in Model A9 has several hyperbolic points. Its chaoticity is confirmed by the FLI of Fig. 3(c). There are no chaotic orbits in a range of $10 < r < 12.3$ for Model A4 in Fig. 2(a), but there are two chaotic orbits with the initial separations $r = 1.210/0.1006$ and $r = 1.235/0.1006$ in the range of $10 < r < 12.3$ for Model A9 in Fig. 2 (b) and (c).

When Model A19 is considered in Fig. 4(a), an orbit with the initial separation $r = 0.163/0.1006$ is a regular KAM torus. Another orbit with the initial separation $r = 0.154/0.1006$ consists of many islands and therefore is still regular. However, the third orbit with the initial separation $r = 0.160/0.1006$ is chaotic because it has many discrete points which are randomly filled with an area. The degree of chaos seems to be strengthened in the small range of $1.45 < r < 1.7$ from A4 to A9 and to A19. Chaos in the range of $10 < r < 12.5$ seems to be stronger for A19 in Fig. 4(b) than for A9 in Fig. 3(b). When the initial separation r spans from 20 to 23 in Fig. 2(c), a lot of regular KAM torus orbits exist in the interior of the ordered orbit with the initial separation $r = 2.230/0.1006$. There is a small chaotic region between the two orbits of the initial separations $r = 2.230/0.1006$ and $r = 2.290/0.1006$. The regularity of the orbit with the initial separation $r = 2.230/0.1006$ and the chaoticity of the orbit with the initial separation $r = 2.290/0.1006$ can be identified clearly by means of the technique of FLIs in Fig. 4(d).

Seen from Figs. 2-4, the number and degree of chaotic orbits seem to increase as the number of the radial terms n gets large. This result is clearly shown in Fig. 5 that lists the dependence of FLI on the initial separation r in each of Models A4, A9 and A19. In other words, Fig. 5 shows that a distribution of the phase space structures of regular and chaotic orbits for each model depends on a range of the initial separation r . Each of the FLIs is obtained after 5×10^4 integration steps. 7 is a critical value of the FLI between the ordered and chaotic two cases. The FLIs not more than 7 indicate the regularity of bounded orbits, whereas those larger than 7 correspond to the chaoticity of bounded orbits. The result on an increase of n increasing the number and degree of chaotic orbits is based on n corresponding to the number of the radial terms in the series expansions of the potential. An increase of n means that of the number of the radial terms. Equivalently, the nonlinear gravitational interaction effects in the potential are enhanced.

B. Models B4, B9 and B19

Fig. 6 describes the phase space structures Model B4 in two ranges of the initial separations r . There is a weak chaotic orbit with the initial separation $r = 0.750/0.0926$ in the range of $7.2 < r < 8.2$ in Fig 6(a). The two orbits in the range of $10.0 < r < 12.2$ are regular KAM tori in Fig 6(b). However, chaos occurs for the initial separation $r = 3.746/0.0926$ in the range of $40.25 < r < 43$ in Fig. 4(c). The dynamical features of the chaotic orbit in panel (a) and the two orbits in panel (c) are also shown in Fig. 6(d). The phase space structures of Model B4 in Fig. 6 (b) and (c) are similar to those of Model A4 in Fig. 2 (a) and (b).

The phase space structures of Model B9 in Fig. 7 (a) and (b) also look like those of Model A9 in Fig. 3 (a) and (b). The initial separations $r = 0.157/0.0926$ and $r = 0.139/0.0926$ correspond to two regular orbits consisting of three loops in Fig. 7(a). Chaos is present for $r = 0.141/0.0926$ in Fig. 7(a) and $r = 1.128/0.0926$ in Fig. 7(b), as the FLIs of Fig. 7(c) show.

Model B19 seem to have stronger chaos in the ranges of $1.425 < r < 1.7$ and $10 < r < 12.25$ in Fig. 8 (a) and (b) than Model B9 in Fig. 7 (a) and (b). For the initial separation r belonging to the range of $1.425 < r < 1.7$, the orbital dynamical behavior for Model B19 in Fig. 8(a) is greatly different from that for Model A19 in Fig. 4(a). However, the phase space structures of Model B19 in Fig. 8 (b) and (c) also resemble those of Model A19 in Fig. 4 (b) and (c). The FLIs of Fig. 8(d) support the chaoticity of the two orbits for $r = 2.368/0.0926$ and $r = 2.420/0.0926$ in Fig. 8(c).

The relation between the FLI and the initial separation r in Fig. 9 clearly shows that the number and degree of chaotic orbits in Models B increase with n increasing.

C. Models C4, C9 and C19

As far as Models C are concerned, chaos seems to get stronger in the range of $3.5 < r < 5.1$ from $n = 4$ in Fig. 10(a) to $n = 9$ in Fig. 10(c) and $n = 19$ in Fig. 10(e). The result is also suitable for the range of $10 < r < 12.2$ in Fig. 10 (b), (d) and (f). The relation between the FLI and the initial separation r in Fig. (11) supports that chaos becomes stronger with n increasing.

Several points can be concluded from the above demonstrations. The three types of models A, B and C have the same expressions, but have differences in the pattern speeds Ω and the coefficients B_{i00} , B_{i20} , C_{i21} and B_{i22} . The three models have similar dynamical behaviors when they have same radial term number n , energy E and initial conditions. The dynamical structures of ordered and chaotic orbits in each model have different distributions along the radial

direction. When the radial term number n increases, the gravity effects are enhanced. This leads to strengthening the degree of chaos. It is worth noting that the main structures of the spiral arms in the models A, B and C are formed at radial distances between 3 and 5 scaled radii, as is shown in Refs. [8] and [10]. Some ranges of the scaled radial distances, such as the range of the scaled radial distances from 40 to 43, seem to be very far from the main structures of the spiral arms. These results show that chaos can occur at the radial distances close to the main structures of the spiral arms and those far from the main structures of the spiral arms. In addition, the chaotic orbits we find are not restricted in the regions near the unstable equilibrium points L_1 and L_2 . Thus, the energies selected in this paper are unlike those of [8-10].

V. CONCLUSIONS

This paper mainly focuses on three classes of models of rotating galaxies in the polar coordinates in the rotating frame. They are models A, B and C, which have the same expressions but have different pattern speeds Ω and different coefficients B_{i00} , B_{i20} , C_{i21} and B_{i22} . When the potentials with large deviations from axial symmetry are included in these models, the angular momentum p_ϕ is not a constant of motion. In this case, the kinetic energy is a function of the momenta and spatial coordinate.

The existing explicit force-gradient symplectic integrators [21-24] do not work for the present Hamiltonian problems. However, our extended force-gradient symplectic methods in the previous work [32] are still available. Numerical tests show that the fourth-order symmetric symplectic method without the extended force-gradient operator performs poorer accuracy than that with the extended force-gradient operator. In particular, the optimized extended force-gradient symplectic method is superior to the corresponding non-optimized one in accuracy.

The fourth-order optimized symplectic algorithm with the extended force-gradient operator is applied to survey the dynamical features of regular and chaotic orbits in these rotating galaxy models. It is shown through the techniques of Poincaré sections and fast Lyapunov indicators that an increase of the radial term number of the potential strengthens the gravity effects and the degree of chaos. The three types of models have similar dynamical structures for the same radial term number, energy and initial conditions.

Author Contributions: Software and Writing-original draft, L. Z.; Methodology, W. L.; Supervision, Conceptualization, Writing - Review & Editing and Funding Acquisition, X. W. All authors have read and agreed to the published version of the manuscript.

Funding: This research has been supported by the National Natural Science Foundation of China (Grant Nos. 11973020 and 11533004), and the Natural Science Foundation of Guangxi (Grant No. 2019GXNSFDA245019).

Data Availability Statement: Our paper is a theoretical work. All of the data are calculated and given in the paper.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Acknowledgments: The first author is very grateful to Dr. Harsoula for the series coefficients in Tables 1-3. The authors also thank anonymous referees for useful suggestions.

Conflicts of Interest: The authors declare no conflict of interest.

-
- [1] Sussman, G.T.; Wisdom, J. Numerical evidence that Pluto is chaotic, *Science* **1988**, 241, 433-437.
 - [2] Murray, C.D.; Dermott, S.F. Solar system dynamics, Cambridge University Press (1999).
 - [3] Voglis, N.; Kalapotharakos, C.; Stavropoulos, I. Mass components in ordered and in chaotic motion in galactic N-body models. *Mon. Not. R. Astron. Soc.* **2002**, 337, 619.
 - [4] Manos, T.; Athanassoula, E. Regular and chaotic orbits in barred galaxies - I. Applying the SALI/GALI method to explore their distribution in several models. *Mon. Not. R. Astron. Soc.* **2011**, 415, 629-642.
 - [5] Kalapotharakos, C.; Voglis, N.; Contopoulos, G. Chaos and secular evolution of triaxial N-body galactic models due to an imposed central mass. *Astron. & Astrophys.* **2004**, 428, 905.
 - [6] Kalapotharakos, C.; Voglis, N. Global Dynamics in Self-Consistent Models of Elliptical Galaxies. *Celest. Mech. Dyn. Astron.* **2005**, 92, 157.
 - [7] Muzzio, J.C.; Carpintero, D.D.; Wachlin, F.C. Spatial structure of regular and chaotic orbits in a self-consistent triaxial stellar system. *Cel. Mech. Dyn. Astr.* **2005**, 91, 173.
 - [8] Voglis, N.; Stavropoulos, I.; Kalapotharakos, C. Chaotic motion and spiral structure in self-consistent models of rotating galaxies. *Mon. Not. R. Astron. Soc.* **2006**, 372, 901.
 - [9] Tsoutsis, P.; Efthymiopoulos, C.; Voglis, N. The coalescence of invariant manifolds and the spiral structure of barred galaxies. *Mon. Not. R. Astron. Soc.* **2008**, 387, 1264.

- [10] Harsoula, M.; Efthymiopoulos, C.; Contopoulos, G. Analytical forms of chaotic spiral arms. *Mon. Not. R. Astron. Soc.* **2016**, 459, 748.
- [11] Voglis, N.; Harsoula, M.; Contopoulos, G. Orbital structure in barred galaxies. *Mon. Not. R. Astron. Soc.* **2007**, 381, 757.
- [12] Athanassoula, E.; Romero-Gómez, M.; Bosma, A.; Masdemont, J.J. Rings and spirals in barred galaxies - III. Further comparisons and links to observations. *Mon. Not. R. Astron. Soc.* **2010**, 407, 1433-1448.
- [13] Bountis, T.; Manos, T.; Antonopoulos, C. Complex statistics in Hamiltonian barred galaxy models. *Celestial Mechanics and Dynamical Astronomy* **2012**, 113, 63-80.
- [14] Machado, R.E.G.; Manos, T. Chaotic motion and the evolution of morphological components in a time-dependent model of a barred galaxy within a dark matter halo. *Mon. Not. R. Astron. Soc.* **2016**, 458, 3578-3591.
- [15] Mondal, D.; Chattopadhyay, T. Role of galactic bars in the formation of spiral arms: a study through orbital and escape dynamics-I. *Celestial Mechanics and Dynamical Astronomy* **2021**, 133, 43.
- [16] Ruth, R.D. A Canonical Integration Technique. *IEEE Trans. Nucl. Sci.* **1983**, 30, 2669.
- [17] Feng, K.; Qin, M. The symplectic methods for the computation of hamiltonian equations. *Lecture Note in Math.* **1987**, 1297, 1.
- [18] Hairer, E.; Lubich, C.; Wanner, G. *Geometric Numerical Integration: Structure-Preserving Algorithms for Ordinary Differential Equations*, 2nd ed. (Springer, Berlin, 2006).
- [19] Forest, E.; Ruth, R. Fourth-order symplectic integration. *Physica D* **1990**, 43, 105.
- [20] Yoshida, H. Construction of higher order symplectic integrators. *Phys. Lett. A* **1990**, 150, 262.
- [21] Suzuki, M. Hybrid exponential product formulas for unbounded operators with possible applications to Monte Carlo simulations. *Physics Letters A* **1995**, 201, 425-428.
- [22] Chin, S. A. Symplectic integrators from composite operator factorizations. *Phys. Lett. A* **1997**, 226, 344.
- [23] Omelyan, I.P.; Mryglod, I.M.; Folk, R. Construction of high-order force-gradient algorithms for integration of motion in classical and quantum systems. *Phys. Rev. E* **2002**, 66, 026701.
- [24] Omelyan, I.P.; Mryglod, I.M.; Folk, R. Symplectic analytically integrable decomposition algorithms: classification, derivation, and application to molecular dynamics, quantum and celestial mechanics simulations. *Comput. Phys. Commun.* **2003**, 151, 272.
- [25] Wisdom, J.; Holman, M. Symplectic maps for the n-body problem. *Astron. J.* **1991**, 102, 1528-1538.
- [26] Wang, Y.; Sun, W.; Liu, F.; Wu, X. Construction of Explicit Symplectic Integrators in General Relativity. I. Schwarzschild Black Holes. *The Astrophysical Journal* **2021**, 907, 66.
- [27] Wang, Y.; Sun, W.; Liu, F.; Wu, X. Construction of Explicit Symplectic Integrators in General Relativity. II. Reissner-Nordström Black Holes. *The Astrophysical Journal* **2021**, 909, 22.
- [28] Wang, Y.; Sun, W.; Liu, F.; Wu, X. Construction of Explicit Symplectic Integrators in General Relativity. III. Reissner-Nordström-(anti)-de Sitter Black Holes. *The Astrophysical Journal Supplement Series* **2021**, 254, 8.
- [29] Wu, X.; Wang, Y.; Sun, W.; Liu, F. Construction of Explicit Symplectic Integrators in General Relativity. IV. Kerr Black Holes. *The Astrophysical Journal* **2021**, 914, 63.
- [30] Sun, W.; Wang, Y.; Liu, F.; Wu, X. Applying explicit symplectic integrator to study chaos of charged particles around magnetized Kerr black hole. *European Physical Journal C* **2021**, 81, 785.
- [31] Zhou, N.; Zhang, H.; Liu, W.; Wu, X. A Note on the Construction of Explicit Symplectic Integrators for Schwarzschild Spacetimes. *The Astrophysical Journal* **2022**, 927, 160.
- [32] Zhang, L.; Wu, X.; Liang, E. Adjustment of Force-Gradient Operator in Symplectic Methods. *Mathematics* **2021**, 9, 2718.
- [33] Froeschlé, C.; Lega, E. On the Structure of Symplectic Mappings. The Fast Lyapunov Indicator: a Very Sensitive Tool. *Celest. Mech. Dyn. Astron.* **2000**, 78, 167.
- [34] Allen, A.J.; Palmer, P.L.; Papaloizou, J. A conservative numerical technique for collisionless dynamical systems: comparison of the radial and circular orbit instabilities. *Mon. Not. R. Astron. Soc.* **1990**, 242, 576.
- [35] Wu, X.; Huang, T.; Zhang, H. Lyapunov indices with two nearby trajectories in a curved spacetime. *Phys. Rev. D* **2006**, 74, 083001.




Fig. 1: (a) The relative energy errors $\Delta H/H_0 = (E_t - E)/E$ for the standard fourth-order symplectic method M4, the extended force-gradient fourth-order symplectic method N4, the optimized extended force-gradient fourth-order symplectic method N4P and the non-symplectic integrator RKF8(9) independently acting on Model A4. E_t is the numerical energy of the system (1) at time t , and the energy is $E = -7 \times 10^6$. The time step is $\tau = 0.0005 \times T_{hmct} \approx 2 \times 10^{-5}$ (sec-kpc/km). The initial conditions are $r = 0.164/R_h = 0.164/0.1006$, $\phi = 1.89\pi$ and $p_r = 0$. The positive initial value of p_ϕ is determined by $E = H$. The three symplectic methods do not give secular drifts to the energy errors, whereas the non-symplectic integrator RKF8(9) does. In accuracy, M4 is the poorest one among these symplectic integrators, while N4P is the best one. (b) The relative position error $\Delta r/r_0 = (r - r_0)/r_0$ between the radial separation r_0 given by RKF8(9) and the radial separation r given by one of the symplectic methods. The position error for N4 is smaller than for M4 but larger than for N4P.




Fig. 2: (a) and (b): Phase space structures of Model A4 on the Poincaré section at the plane $\phi = \pi/2$ with $p_\phi > 0$. The dynamical structures of ordered and chaotic orbits have different spatial distributions along the radial distance r . The parameters and the initial conditions (except r and p_ϕ) are those of Fig.1. The structures are considered in the range of $10 < r < 12.4$ (a) and $43.5 < r < 46.5$ (b). (c): FLIs of the two orbits in panel (b). The FLIs show the regularity of the orbit for $r = 4.430/0.1006$ and the chaoticity of the orbit for $r = 4.424/0.1006$.




Fig. 3: (a) and (b): Phase space structures of Model A9 on the Poincaré section. The structures are considered in the ranges of $1.5 < r < 1.7$ (a) and $10 < r < 12.4$ (b). The initial conditions in Fig. 3(b) are the same as those in Fig. 2(a). However, the orbits for $r = 1.210/0.1006$ and $r = 1.235/0.1006$ are chaotic, whereas the orbit for $r = 1.200/0.1006$ is regular. (c): FLIs of several orbits in panels (a) and (b). The FLIs show the regularity of the orbit for $r = 1.200/0.1006$ and the chaoticity of the three orbits for $r = 0.153/0.1006$, $r = 1.210/0.1006$ and $r = 1.235/0.1006$.




Fig. 4: (a)-(c): Phase space structures of Model A19 on the Poincaré section. (a): The structures are considered in the range of $1.45 < r < 1.7$. The orbit for $r = 0.163/0.1006$ is an ordered KAM torus, and the orbit for $r = 0.154/0.1006$ with many islands is still regular. However, the orbit for $r = 0.160/0.1006$ is chaotic. (b): The structures are considered in the range of $10 < r < 12.4$. The initial conditions in Fig. 4(b) are the same as those in Fig. 2(a). (c): The structures are considered in the range of $20 < r < 23$. There are a number of regular KAM torus orbits in the interior of the ordered orbit for $r = 2.230/0.1006$. A small chaotic region between the two orbits for $r = 2.230/0.1006$ and $r = 2.290/0.1006$ exists. (d): Fast Lyapunov indicators (FLIs) of the two orbits in panel (c). The orbit for $r = 2.230/0.1006$ having a power law increase of the FLI with time $\log_{10} t$ is ordered, but the orbit for $r = 2.290/0.1006$ having an exponential law increase of the FLI with time is chaotic. The FLIs are calculated until the number of integration steps reaches 5×10^4 .




Fig. 5: The dependence of FLI on the initial separation r in Models (a) A4, (b) A9 and (c) A19. Each of the FLIs is obtained after 5×10^4 integration steps. The FLIs not more than 7 show the regularity of bounded orbits, whereas those larger than 7 indicate the chaoticity of bounded orbits. Many values of r correspond to chaos in Model A19, but only minor values of r do in Model A4. It is clear that chaos becomes easier and its degree is enhanced as the radial term number n increases.




Fig. 6: Similar to Fig. 2 but Model B4 is considered. (a): The structures are considered in the range of $7.2 < r < 8.2$. (b): The structures are considered in the range of $12.0 < r < 12.2$. (c): The structures are considered in the range of $40 < r < 43$ and resemble those in Fig. 2(b). (d): The FLIs show the chaoticity of the two orbits in panels (a) and (c) and the regularity of the orbit in panel (c).




Fig. 7: Similar to Fig. 3 but Model B9 is considered. The phase space structures in panels (a) and (b) respectively look like those in Fig. 3 (a) and (b). The initial conditions in Fig. 7(b) are the same as those in Fig. 6(b). The orbit for $r = 1.128/0.0926$ is chaotic, but the orbit for $r = 1.120/0.0926$ is regular. (c): The FLIs show the chaoticity of the two orbits in panels (a) and (b) and the regularity of the orbit in panel (b).




Fig. 8: Similar to Fig. 4 but Model B19 is considered. The phase space structures in panel (a) are somewhat different from those in Fig. 4(a), but these structures in panels (b) and (c) respectively resemble those in Fig. 2 (b) and (c). The initial conditions in Fig. 8(b) are the same as those in Fig. 6(b). (d): The FLIs show the chaoticity of the two orbits in panel (c) and the regularity of the orbit in panel (c).




Fig. 9: Similar to Fig. 5 but Models A are replaced with Models B. The degree of chaos becomes stronger with an increase of the radial term number n .




Fig. 10: (a) and (b): Phase space structures of Model C4 on the Poincaré section. The structures are considered in the ranges of $3.5 < r < 5.1$ and $11.9 < r < 12.2$. (c) and (d): Phase space structures of Model C9 on the Poincaré section. (e) and (f): Phase space structures of Model C19 on the Poincaré section.




Fig. 11: Similar to Fig. 5 but Models A are replaced with Models C. The degree of chaos increases with the radial term number n increasing.