arXiv:2302.06574v2 [quant-ph] 23 Oct 2023

Hierarchies among Genuine Multipartite Entangling Capabilities of Quantum Gates

Samir Kumar Hazra^{1,2}, Aditi Sen(De)²

¹ Gandhi Institute of Technology and Management (Deemed to be University) Bengaluru Campus, Bengaluru-562163, India ²Harish-Chandra Research Institute, A CI of Homi Bhabha National Institute, Chhatnag Road, Jhunsi, Allahabad - 211019, India

We categorize quantum gates according to their capability to generate genuine multipartite entanglement based on the hierarchy of multipartite separable states. In particular, when a fixed unitary operator acts on the set of k-separable states, the maximal (average) genuine multipartite entanglement (GME) content produced via that particular unitary operator is determined after maximizing over the set of k-separable input states. We identify unitary operators that are beneficial for generating high GME when the input states are entangled in some bipartition, although the picture can also be reversed in which entanglement in inputs does not help. We characterize maximum entangling power of a variety of unitary operators including special classes of quantum gates, diagonal, permutation and Haar uniformly generated unitary operators by computing generalized geometric measure (GGM) as GME quantifier. We determine the unitary operators and their corresponding inputs which can create the resulting states having maximum GGM.

I. INTRODUCTION

Quantum correlations [1-3] present in shared multipartite quantum states are shown to be responsible for obtaining advantages in quantum information processing tasks over classical counterparts. Among all quantum features, multipartite entanglement emerges as the key ingredient of quantum protocols including quantum networks [4] that can transmit both classical and quantum information [5-10], quantum secret sharing involving a single sender and multiple receivers [11], and measurement-based quantum computation [12–14]. Additionally, it has also been demonstrated that the ground, thermal or the dynamical state of quantum many-body systems [15–17] possesses multipartite entanglement which can also be used to identify quantum phenomena [18-20] present in both equilibrium and nonequilibrium situations [21, 22]. More importantly, multipartite entangled states can be produced in laboratories; examples include twelve photon genuine multipartite entangled states [23, 24], trapped ions having twenty qubits [25], twenty entangled superconducting qubits [26] etc. This opens up the possibility of realizing quantum information protocols involving multipartite entangled states in laboratories.

In recent times, there have been a number of techniques developed to create multipartite entangled states in quantum many-body systems. The prominent ones involve different kinds of measurements such as projective as well as positive operator valued measurement [13, 14, 27–35], and quantum gates [36–40]. The former protocols are probabilistic in nature while the later ones are deterministic. It was demonstrated that both strategies are indeed capable of producing highly genuine multipartite entangled states.

In the context of gate-based schemes that can be a part of quantum circuits performing particular quantum algorithms [41-43], it is fascinating to characterize the unitary operators with respect to their ability to create multipartite entangled states starting from the product states. In particular, it has been shown that the average entangling power of a given two-qubit unitary operator is the mean linear entropy acquired following the action of unitary operator on a set of Haar uniformly distributed product states [44-50]. By exploiting the Cartan

decomposition from theory of Lie groups and algebras, it is possible to obtain an elementary decomposition of an arbitrary unitary operator acting on an arbitrary number of qubits [51–53]. Such connection to Cartan decomposition also sheds light on the geometric structure of the nonlocal unitary operators which are the only responsible operators to create any entanglement. In particular, Cartan decomposition applied to the Lie algebra $\mathfrak{su}(4)$ gives that the geometric structure of nonlocal gates is a 3-torus [54]. Moreover, estimates of the number of two-qubit gates in near-optimal circuits can be determined by applying Cartan decomposition recursively [55]. However, in contrast to average entangling power, the maximum entangling power of a fixed unitary operator can also reveal the optimal strength to create an entangled state from the initial product state [56] or a set of specific states having certain amount of entanglement [57]. Both directions lead to the quantification of a resource theory for operations [58].

To evaluate the strength of two-qubit quantum gates, quantum correlation measures, such as entanglement [1] and quantum discord [3] have already been used [45, 46, 59]. Apart from pure product states and unitary operators, impact of unitary operators on mixed states as well as the entangling power of non-unitary operators [60] are also examined for two-party systems [61]. In a slightly different direction, ordering between unitary operators is studied considering the simulation time required via Hamiltonian [62–64]. If one restricts only to symmetric two-qubit unitary operators which operate on the set of all symmetric separable states, the explicit form of average entangling power of these kinds of quantum gates are obtained again by utilizing Cartan decomposition of semisimple Lie algebra [65]. Beyond the two-party scenario, the entangling capacity of multipartite gates in terms of linear entropy is studied when they act on the set of fully separable states [66–68].

In this work, we deviate from the traditional analysis of determining entangling power of a given quantum gate in two ways – firstly, we change the set of input product states on which the optimization has to be performed to generate entanglement via quantum gates; secondly, the quantification of entanglement created in this process is also considered differently than how it was studied earlier in the literature. In particular, we first take various classes of separable states, referred to as "k-separable states" for different values of k, as the input states [69]. For example, in the case of three parties, we optimize over a set of 3-separable states representing fully separable states having vanishing genuine and bipartite entanglement [69, 70] as well as a set of biseparable states which possess nonvanishing bipartite entanglement in some bipartitions, called as 2-separable states. Notice that according to our definition, the set of 2-separable states does not contain the set of 3-separable states. Secondly, we determine the entangling power of a given unitary operator by the maximum amount of genuine multipartite entanglement it can create as measured by the generalized geometric measure (GGM) [10, 71] where the maximization is performed over the set of k-separable inputs. We refer to it as *genuine multipartite entangling capability* of the operator.

We identify unitary operators as well as input N-party states with vanishing GGM which together can create a maximum amount of genuine multipartite entanglement (GME). Specifically, we prove that there are unitary operators, namely diagonal and permutation unitaries, in which parameters can be tuned in such a way that the states having maximum GGM can be produced from some particular kind of k-separable states $(2 \le k \le N)$. We also classify the unitary operators based on their capability to generate GME from certain classes of input separable states. In this respect, one might expect that a global unitary operator with k-separable states as inputs generates more genuine multipartite entanglement as opposed to k + 1-separable states. However, we demonstrate that such intuition is too simplistic! Specifically, there exist unitary operators which can not take the advantage of initial entanglement present in input states (in certain bipartitions) and produce higher entanglement. Having said that, we determine unitary operators such as diagonal, permutation and Haar uniformly generated unitary operators which indeed provide the benefit of initial entanglement present in bipartition. We establish these results by considering different classes of eight- and higher-dimensional unitary operators which act on three- and higher-qubit input states respectively, and the optimization is performed over the sets of corresponding separable states.

The paper is organized as follows. In Sec. II, we introduce the notion of genuine multipartite entangling capability of unitary operators based on different classes of multipartite separable inputs and their properties. We identify classes of input states and the corresponding classes of unitary operators which can create maximally genuine multipartite entangled states in Sec. III. Sec. IV computes entangling power of randomly generated diagonal operators, Haar uniformly produced random unitary operators and special classes of operators [52] for three-qubit input states. The concluding remarks are included in Sec. V.

II. HIERARCHIES IN ENTANGLING POWER OF UNITARY OPERATORS

An *N*-party pure state, $|\psi\rangle$, is said to be *k*-separable (k = 2, ..., N) when it can be written as a product in the *k* number



FIG. 1. (Color online.) A schematic diagram on the notion of hierarchies among entangling powers of a fixed unitary operator, U, based on the inputs chosen. Each column represents different sets of separable states. From left to right, set of inputs are chosen to be Nseparable (i.e., fully separable) to k-separable. The parties entangled are marked with blue circles while the brown circles are for the rest. After the action of U on k-separable inputs, the maximum genuine multipartite entanglement of the resulting states, \mathcal{E}_{max}^k in Eq. (2), is computed which is represented by the vertical green color. Here the superscript represents the class of separable states considered for maximization. Deeper green indicates more expected GME created in the outputs.

of partitions [69]. Mathematically,

$$|\psi\rangle = \otimes_{m=1}^{k} |\psi_m\rangle. \tag{1}$$

We denote such a k-separable pure state as $|\psi^{\text{k-sep}}\rangle$. When k = N, the state is called fully separable while a pure state is called genuinely multipartite entangled when it can not be written as a product in any bipartition. The definition of k-separability for pure states can be extended to mixed states by using convex roof extension. In particular, a mixed N-party state ρ is said to be k-separable if it can be written as a convex combination of k-separable pure states, i.e., $\rho^{k-\text{sep}} = \sum_i p_i |\psi_i^{k-\text{sep}}\rangle \langle \psi_i^{k-\text{sep}}|$. Note that, in this decomposition, the states, $|\psi_i^{k-\text{sep}}\rangle$ can be k-separable in different partitions.

Definition of multipartite entangling power. A given unitary operator, U, acting on a k-separable state can, in general, produce a state having genuine multipartite entanglement. When the genuine multipartite entanglement content of the resulting state is maximized over the set of k-separable states (with a fixed value of k > 1), we call it k-maximal entangling power of U. Specifically, for a given genuine multipartite entanglement measure, \mathcal{E} , a given unitary operator, U, and a fixed value of k, it can be mathematically represented as

$$\mathcal{E}_{\max}^k(U) = \max_{\mathcal{S}^k} \mathcal{E}(U\rho^{k\text{-sep}}U^{\dagger}), \qquad (2)$$

where the maximization is performed over the set S^k containing all k-separable states. Clearly, the superscript "k" in $\mathcal{E}_{\max}^k(U)$ refers to the fact that the optimization is performed over the set of k-separable states. Depending on the set of k-separable states, a set of k-maximal entangling powers, $\{\mathcal{E}_{\max}^k\}_{k=2}^N$, for a fixed unitary operator emerges (as depicted in Fig. 1). In this context, it is important to note that in the definition of entangling power of a unitary operator, no GME measure has been used so far in the literature [45, 66, 68].

Sets for optimization. Here our main goal is to find out the hierarchies among the k-maximal entangling powers of a unitary operator U in $d = 2^N$ dimensions. Such a problem can be addressed when U acts on the set of k-separable states for $1 < k \leq N$, giving us a set of numbers $\{\mathcal{E}_{\max}^k(U) : 1 < k \leq N\}$. For a given unitary operator U, we will examine the ordering between the elements of this set. Since the set of k-separable states is naturally a subset of the set of k-1separable states, i.e. we have $\mathcal{S}^N \subseteq \ldots \subseteq \mathcal{S}^k \subseteq \mathcal{S}^{k-1} \subseteq$ $\ldots \subseteq \mathcal{S}^2$, such an ordering can be trivial as we shall obviously have $\mathcal{E}_{\max}^k(U) \leq \mathcal{E}_{\max}^{k-1}(U)$.

In order to avoid this triviality and reveal the significant information about entangling power of U, one needs to modify the sets of separable states on which the maximization is performed. In particular, for carrying out the investigation, we impose the restriction that the set of k-separable states should not contain states belonging to the sets S^{k+i} $(\forall i \ge 1)$. We denote these new sets by \mathbb{S}^k which can be given by $\mathbb{S}^k := S^k \setminus S^{k+1}$. Clearly, \mathbb{S}^k are now all disjoint. Here we keep our previous notation $\mathcal{E}^k_{\max}(U)$ for k-maximal entangling power unchanged even though the maximization is now considered over the new sets \mathbb{S}^k .

Now since the number of parties which are entangled in the states of \mathbb{S}^k is higher than the number of those in \mathbb{S}^{k+1} , one may expect higher production of entanglement from \mathbb{S}^k , i.e., $\mathcal{E}_{\max}^{k+1}(U) \leq \mathcal{E}_{\max}^k(U)$. We will illustrate that this is not always the case. It is important to stress here that all the previous studies considered the entangling power of unitary operators with their action only on the set of fully separable states [66, 68] which obviously does not allow any comparison of our kind.

Note further that with the increase in number of parties, there can be different kinds of entangled states among k-separable states themselves. For example, with N = 6, states having bipartite, genuine tripartite and genuine quadripartite entanglement can all be 3-separable states. Hence, more fine-grained entangling power of a fixed unitary operator can also be considered, depending on the different kinds of entanglement present in the input states, which is beyond the scope of this work.

Properties of multipartite entangling power. Let us discuss some of the properties of $\mathcal{E}_{\max}^k(U)$.

(1) It is nonvanishing for all Us which create GME states as outputs. In other words, $\mathcal{E}_{\max}^k(U) = 0$ if and only if all the output states have vanishing genuine multipartite entanglement. The upper bound of $\mathcal{E}_{\max}^k(U)$ is fixed by the choice of the entanglement measure, \mathcal{E} .

(2) $\mathcal{E}_{\max}^{k}(U)$ remains invariant under local unitary operations, i.e., $\mathcal{E}_{\max}^{k}(\bigotimes_{i=1}^{N}U_{i}U) = \mathcal{E}_{\max}^{k}(U)$. If the entanglement measure chosen is local unitarily equivalent, it can be shown that multipartite entangling power also remains so. Suppose before local unitary operation, the optimal state that maximizes entangling power is ρ^{opt} , i.e., $\mathcal{E}_{\max}^{k}(U) = \mathcal{E}(U\rho^{opt}U^{\dagger})$ while after local unitary operations, $\mathcal{E}_{\max}^{k}((\bigotimes_{i=1}^{N}U_{i}U) =$ $\mathcal{E}((\bigotimes_{i=1}^{N} U_{i} U \tilde{\rho}^{opt} U^{\dagger} \bigotimes_{i=1}^{N} U_{i}^{\dagger})$. Since \mathcal{E} is local unitary invariant, $\rho^{opt} = \tilde{\rho}^{opt}$ and hence the proof.

Replacing maximum by average in Eq. (2) where the averaging is performed over the set of k-separable states, we can obtain average entangling power of a unitary operator, U, as

$$\overline{\mathcal{E}^{k}(U)} = \int_{\mathbb{S}^{k}} \mathcal{E}(U\rho_{N}^{k\text{-sep}}U^{\dagger})d\Omega_{N}, \qquad (3)$$

where $d\Omega_N$ represents the Haar measure over \mathbb{S}^k . In this context also, for a fixed unitary operator U, a similar comparison between $\overline{\mathcal{E}^k}(U)$ and $\overline{\mathcal{E}^{k+1}}(U)$ can also be investigated.

In order to make circuits, our operators can be realized in terms of single- and two-qubit operations as is well known from Ref. [42]. In particular, the operators we consider here are all in $SU(2^N)$ which is a semi-simple Lie group. Any such operator can be decomposed into a product of elements of SU(2) and SU(4) following a method described in Ref. [51]. This method is based on successive application of the Cartan decomposition of the Lie group $SU(2^N)$ which also uses geometric facts about $\frac{SU(2^N)}{SU(2^{N-1})\otimes SU(2^{N-1})\otimes U(1)}$. The successive way of writing an N-dimensional operator in low dimensional operations also provides an algorithmic approach to the problem.

Quantification of GME. We use generalized geometric measure to compute genuine multipartite entanglement content of the output states [10]. For an N-party state, $|\psi\rangle$, it is defined as $G(|\psi\rangle) = 1 - \max_{|\phi\rangle \in \mathcal{D}} |\langle\phi|\psi\rangle|^2$, where \mathcal{D} denotes the set of all non-genuinely multipartite entangled states. It has been proven that the maximization involved in GGM can also be performed by using Schmidt coefficients of $|\psi\rangle$ in different bipartitions [10]. Note that $G(|\psi\rangle) = 0$ for any k-separable state $|\psi\rangle$, in particular for any state in \mathbb{S}^k (k > 1). In this work, we will be computing maximal GGM produced by the action of U on the states from the set \mathbb{S}^k , denoted by G_{\max}^k for different k values of initial separable states or alternatively G_{\max} .

III. UNITARY OPERATORS WITH MAXIMAL MULTIPARTITE ENTANGLING POWER AND THEIR INPUTS

Towards characterizing unitary operators (quantum gates) in terms of their entangling power, we identify the operators that create multipartite resulting states having *maximum* GGM. In the process, we also determine the optimal inputs which give us such outputs. Further, these investigations provide a deterministic method to produce states having maximum GGM useful in quantum information processing tasks.

Setting up the stage for three-qubits

Let us first describe how one can determine both optimal inputs and the corresponding outputs for a fixed unitary operator which is capable of producing maximum GGM for threequbits. A pure three-qubit separable state can be either fully separable,

$$|\psi^{3\text{-sep}}\rangle_{\mathbf{3}} = |\psi_1\rangle \otimes |\psi_2\rangle \otimes |\psi_3\rangle = \otimes_{i=1}^3 (\cos\theta_i |0\rangle + \sin\theta_i e^{i\xi_i} |1\rangle)$$

with $0 \le \theta_i \le \pi/2$ and $0 \le \xi_i \le 2\pi$ or a biseparable state [70] of the form

$$|\psi^{2\text{-sep}}\rangle_3 = |\psi_i\rangle \otimes |\psi_{jk}\rangle, i \neq j \neq k,$$
 (4)

where $|\psi_{jk}\rangle$ is a two-qubit entangled state. For example, considering i = 1, $|\psi^{2\text{-sep}}\rangle_3$ can be parametrized as $(\cos \theta'_1|0\rangle + \sin \theta'_1 e^{i\xi'_1}|1\rangle) \otimes \sum_{j,k=0}^1 a_{jk}|jk\rangle$ with $a_{00} = \cos \theta'_2, a_{01} = e^{i\xi'_1} \sin \theta'_2 \cos \theta'_3, a_{10} = e^{i\xi'_2} \sin \theta'_2 \sin \theta'_3 \cos \theta'_4, a_{11} = e^{i\xi'_3} \sin \theta'_2 \sin \theta'_3 \sin \theta'_4$ in terms of higher dimensional spherical polar coordinates and phases. Due to the constraint that sets of biseparable (S²) and fully separable states (S³) are disjoint, θ'_i 's have to be chosen in such a way that the bipartite states in S² have nonvanishing entanglement [1].

Let us identify the unitary operators on eight-dimensional complex Hilbert space, \mathbb{C}^8 , denoted by U(8), which act on the states chosen from \mathbb{S}^2 or \mathbb{S}^3 such that the resulting states after maximization over the inputs reach the maximal value of GGM, i.e., 0.5. In other words, our aim is to find out whether there exist any biseparable (optimal) input, $|\psi_{opt}^{2\text{-sep}}\rangle$, for a fixed element of U(8), again denoted by U(8), so that

$$G_{\max}^2(U(8)) \equiv G(U(8)|\psi_{opt}^{2\text{-sep}}\rangle_3) = 0.5,$$
(5)

and in the case of fully separable ones,

$$G_{\max}^{3}(U(8)) \equiv G(U(8))|\psi_{opt}^{3\text{-sep}}\rangle_{3}) = 0.5.$$
 (6)

We will demonstrate that, indeed, there are unitary operators and corresponding inputs for which both the above equations hold true. However, there exist eight dimensional unitary operators for which only one of the above equations can be satisfied. In what follows, we will encounter such unitary operators.

A. Multipartite entangling power of diagonal unitary operators

Let us illustrate the significance of inputs in the context of deciding multipartite entangling power of diagonal unitary operators. A diagonal unitary operator on eight-dimensional space can be written as

$$U_D^{gen} = \operatorname{diag}(e^{i\phi_1}, e^{i\phi_2}, e^{i\phi_3}, e^{i\phi_4}, e^{i\phi_5}, e^{i\phi_6}, e^{i\phi_7}, e^{i\phi_8}), (7)$$

where $\phi_j \in [0, 2\pi)$ [67] $(j = 1, \dots, 8)$.

We concentrate here on the diagonal unitary operators with $\phi_j = 0$ except ϕ_8 , thereby reducing to the case $U_D = \text{diag}(1, 1, 1, 1, 1, 1, 1, e^{i\phi})$. The entangling power of general diagonal unitary operators with arbitrary ϕ_j s will be investigated in the next section.

After application of such an operator, U_D , on general fully separable or biseparable states, the resulting states can be read as $\sum_{i,j,k=0}^{1} a_{ijk} e^{i\phi_{ijk}} |ijk\rangle$, where $e^{i\phi_{ijk}} = 1$ except $e^{i\phi_{111}} \equiv e^{i\phi}$ and a_{ijk} s are functions of θ_i s, θ'_i s, ξ_i s and ξ'_i s. To obtain GGM for this resulting three-qubit state, we need to calculate all the single qubit reduced density matrices and their eigenvalues. We first notice that during optimization, ξ_i s and ξ'_i s are not playing any role and therefore, we set them to be zero, i.e., in the optimization process, phases present in $|\psi^{2\text{-sep}}\rangle$ and $|\psi^{3\text{-sep}}\rangle$ are not involved. This implies the coefficients a_{ijk} are now real valued functions.

We first consider applying the operators of elements of \mathbb{S}^2 . After application of U_D on an arbitrary biseparable state, the output takes the form,

$$\begin{aligned} |\psi_{out}\rangle &\equiv U_D |\psi^{2\text{-sep}}\rangle \\ &= \cos\theta'_1 \cos\theta'_2 |000\rangle + \cos\theta'_1 \cos\theta'_3 \sin\theta'_2 |001\rangle \\ &+ \dots + e^{i\phi} \sin\theta'_1 \sin\theta'_2 \sin\theta'_3 \sin\theta'_4 |111\rangle, \end{aligned}$$
(8)

where only the last term has ϕ -dependence. In order to compute GGM for these states, the corresponding single-site local density matrices can be obtained which are functions of the form $G(\theta'_1, \ldots, \theta'_4, \phi)$. For a fixed ϕ , after optimizing over θ'_i s, the optimal input biseparable state turns out to be

$$|\psi_{opt}^{2\text{-sep}}\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)\frac{1}{\sqrt{2}}(|00\rangle + |11\rangle),\tag{9}$$

which leads to the resulting state, $|\psi_{out}\rangle = \frac{1}{2}(|000\rangle + |011\rangle + |100\rangle + e^{i\phi}|111\rangle)$. When $\phi = \pi$, all single-qubit reduced density matrices of $|\psi^{out}\rangle$ are maximally mixed. Therefore, the GGM of the output state achieves its maximal value, i.e.,

$$G_{\max}^2(U_D(\phi=\pi)) \equiv G(U_D(\phi=\pi)|\psi_{opt}^{2\text{-sep}}\rangle) = 0.5.$$
 (10)

In general, for arbitrary ϕ , all the local density matrices of $|\psi_{out}\rangle$ are $\mathbb{I}/2$ except the first one which is given by $\frac{1-\cos\frac{\phi}{2}}{2}|0\rangle\langle 0| + \frac{1+\cos\frac{\phi}{2}}{2}|1\rangle\langle 1|$. Therefore, in this case, we obtain

$$G_{\max}^2(U_D) \equiv G(U_D |\psi_{opt}^{2\text{-sep}}\rangle) = \sin^2 \frac{\phi}{4}, \qquad (11)$$

when $0 \le \phi \le \pi$. From the above expression, it is clear that G_{\max}^2 reaches its maximum value only for $\phi = \pi$ and trivially vanishes at $\phi = 0$.

On the other hand, if one applies U_D to the set of fully separable states, $\bigotimes_{i=1}^{3} (\cos \theta_i |0\rangle + \sin \theta_i |1\rangle)$, the output states take the form,

$$U_{\phi}|\psi^{3\text{-sep}}\rangle = \Pi_{i=1}^{3}\cos\theta_{i}|000\rangle + \cos\theta_{1}\cos\theta_{2}\sin\theta_{3}|001\rangle + \dots + e^{i\phi}\sin\theta_{1}\sin\theta_{2}\sin\theta_{3}|111\rangle,$$
(12)

where only the coefficient of $|111\rangle$ depends on ϕ . Again finding all the reduced density matrices and performing maximization over θ_i s, we find that maximum eigenvalues of all the single-party reduced density matrices exceed 1/2, i.e., $G_{\max}^3(U_{\phi})$ never reaches the maximum value of 0.5 for any values of ϕ . For example, when $\phi = \pi$, $G_{\max}^3(U_{\phi})(\phi = \pi)) = 0.34$ (see Fig. 2). This diagonal unitary operator clearly



FIG. 2. (Color online.) Enhancement in entanglement generation with initial entanglement. Biseparable inputs are better than the fully separable ones when a special diagonal unitary matrix, $U_{\phi} = \text{diag}(1, 1, 1, 1, 1, 1, e^{i\phi})$ with $\phi \in (0, 2\pi)$ [67] acts on three-party input states. The maximum GGM produced (ordinate) against ϕ (abscissa). The optimization is performed over the set of biseparable (circles) and fully separable (triangles) states to obtain G_{max} . Both the axis are dimensionless.

demonstrates how initial entanglement may assist in the generation of high GME states, although it is not true in general for an arbitrary unitary operator of U(8) as we will show below.

Remark. It was known that the diagonal unitary, U_{π} , in four-dimensions acted (at a time) on nearest neighbor sites of a suitably chosen fully separable state add up to a cluster state, having maximal GGM, useful for building one-way quantum computer [12]. However, the total operator which acts on the above *N*-party state is no more a unitary operator. Our results described above show that the action of a single diagonal unitary operator of the form U_{ϕ} in eight-dimensions on a set of fully separable three-qubit states can never produce maximum GGM. We will manifest below that the same remains true even in higher dimensions as well.

Action of higher dimensional unitaries on multiqubit inputs

Let us now examine the action of a class of diagonal unitary operators in the sixteen-dimensional complex Hilbert space, given by $U_{\phi}^{16} = \text{diag}(1, \dots, e^{i\phi})$. In a four-party system, 2-sep states of \mathbb{S}^2 can be written as

$$\begin{split} |\psi^{2\text{-sep}}\rangle_4 &= |\psi_i\rangle \otimes |\psi_{jkl}^{gen}\rangle, \text{or} \\ |\psi^{2\text{-sep}}\rangle_4 &= |\psi_{ij}\rangle \otimes |\psi_{kl}\rangle; (i, j, k, l \text{ are different}), (13) \end{split}$$

where $|\psi_{jkl}^{gen}\rangle$ represents genuine tripartite entangled states, while $|\psi_{**}\rangle$ s are the bipartite entangled states. When U_{ϕ}^{16} acts on the initial four-party states, the resulting states take the form, $|\psi^{out}\rangle = \sum_{i,j,k,l=0}^{1} a_{ijkl} e^{i\phi_{ijkl}} |ijkl\rangle$ with $e^{i\phi_{ijkl}} = 1$ except $e^{i\phi_{1111}} = e^{i\phi}$, i.e., after its action on the states, the coefficient of $|1111\rangle$ only depends on ϕ present in U_{ϕ}^{16} and $a_{ijkl}\mathbf{s}$, which depend on the input states, are typically products of cosine and sine functions involved e.g., in $|\psi^{2\text{-sep}}\rangle$. As in the case of three-qubits, the maximization process reveals that the optimal inputs are either $|\psi^{2\text{-sep}}_{opt}\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)\frac{1}{\sqrt{2}}(|00\rangle + |11\rangle$ or $|\psi^{2\text{-sep}}_{opt}\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)\frac{1}{\sqrt{2}}(|00\rangle + |11\rangle$. Therefore, the multipartite entangling power of U_{ϕ}^{16} turns out to be

$$G_{\max}^2(U_{\phi}^{16}) \equiv G(U_{\phi}^{16}|\psi_{opt}^{2\text{-sep}}\rangle) = \min[\sin^2\frac{\phi}{4}, \cos^2\frac{\phi}{4}] \ (14)$$

since all the reduced density matrices are either $\frac{\mathbb{I}}{2}$ or their eigenvalues are $\frac{1\pm\cos\frac{\phi}{2}}{2}$. It again attains maximum value, 0.5 with $\phi = \pi$. The numerical simulations performed on the set of k-separable (k > 2) inputs, i.e., over \mathbb{S}^3 and \mathbb{S}^4 , exhibit that U_{ϕ}^{16} cannot produce output states having maximal GGM from the set of 3- and 4-separable input states.

The above analysis clearly gives an indication that multipartite entangling power of the class of diagonal unitary operators in \mathbb{C}^{2^N} of the form $U_{\phi}^{2^N} = \text{diag}(1, 1, \dots, e^{i\phi})$ always achieve maximum value of 0.5 with input $|\psi^{2\text{-sep}}\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)\frac{1}{\sqrt{2}}(|0\dots0\rangle + |1\dots1\rangle$. Moreover, starting with any *N*-party input state of the form, $|\psi^{2\text{-sep}}\rangle = \frac{1}{\sqrt{2}}(|0\rangle^{\otimes k} + |1\rangle^{\otimes k})\frac{1}{\sqrt{2}}(|0\rangle^{\otimes N-k} + |1\rangle^{\otimes N-k})$ for different *k*, we obtain $G_{\max}^2(U_{\phi=\pi}^{2^N}) = 0.5$ while for arbitrary ϕ , the formula for GGM is as given in Eq. (14). In this manner, we identify both, a class of diagonal unitary operators and their optimal inputs which are capable of producing maximum GGM in a given dimension.

B. Entangling power for a set of permutation unitary operators

Let us concentrate on the set of permutation unitary operators in the eight dimensional space. In this case, unlike diagonal unitaries, we will illustrate that both fully separable and biseparable states can create outputs having maximum GGM. Let us first fix a product basis in \mathbb{C}^8 , $\mathcal{E} = \{\mathcal{E}_0 \equiv [\alpha_1 \alpha_2 \alpha_3], \mathcal{E}_1 \equiv [\alpha_1 \alpha_2 \alpha_3^{\perp}], \ldots, \mathcal{E}_8 \equiv [\alpha_1^{\perp} \alpha_2^{\perp} \alpha_3^{\perp}]\}$. We will consider the permutation operators of the form (i, j) where $i, j = 1, \ldots, 8$, called transpositions. They act by permuting the basis elements \mathcal{E}_i and \mathcal{E}_j while keeping other basis elements unchanged. In eight dimensional space, 28 such permutation operators exist for a given basis. Our aim is to identify all such permutation operators that produce maximally entangled state, i.e., G(*) = 0.5.

To obtain maximum GGM, we first notice that all the reduced density matrices of a tripartite state should be maximally mixed [72]. E.g., the three-qubit state of the form $|\psi\rangle = \frac{1}{\sqrt{2}}(|\psi_1\psi_2\psi_3\rangle + |\psi_1^{\perp}\psi_2^{\perp}\psi_3^{\perp}\rangle)$ achieves maximum GGM which is known as Greenberger-Horne-Zeilinger(GHZ) state [73].

Fully separable inputs: Among twenty eight transpositions mentioned above, we identify that twelve of them can produce maximum GGM after acting on fully separable states. Specifically, we obtain

$$G_{\max}^{3}((i,j)) \equiv G((i,j)|\psi_{opt}^{3\text{-sep}}\rangle) = 0.5.$$
 (15)

No.	Unitary	Fully Separable Inputs
1	(1, 4)	$\frac{1}{\sqrt{2}}(011\rangle + 111\rangle)$
2	(1, 6)	$\frac{1}{\sqrt{2}}(101\rangle + 111\rangle)$
3	(1,7)	$\frac{1}{\sqrt{2}}(110\rangle + 111\rangle)$
4	(2, 3)	$\frac{1}{\sqrt{2}}(001\rangle + 101\rangle)$
5	(2, 5)	$\frac{1}{\sqrt{2}}(100\rangle + 110\rangle)$
6	(2, 8)	$\frac{1}{\sqrt{2}}(110\rangle + 111\rangle)$
7	(3, 5)	$\frac{1}{\sqrt{2}}(100\rangle + 101\rangle)$
8	(3, 8)	$\frac{1}{\sqrt{2}}(101\rangle + 111\rangle)$
9	(4, 6)	$\frac{1}{\sqrt{2}}(100\rangle + 101\rangle)$
10	(4,7)	$\frac{1}{\sqrt{2}}(100\rangle + 110\rangle)$
11	(5, 8)	$\frac{1}{\sqrt{2}}(000\rangle + 100\rangle)$
12	(6,7)	$\frac{1}{\sqrt{2}}(001\rangle + 101\rangle)$

TABLE I. The permutation operators and their corresponding fully separable inputs which maximize GGM are listed in the second and third columns respectively. Note that one of the parties in the fully separable inputs is in $|+\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$ state which has (nonvanishing) maximum coherence [74].

The list of transposition unitary operators and their respective optimal fully separable inputs are given in Table I. Without loss of generality, we denote $|\psi_i\rangle$ and $|\psi_i^{\perp}\rangle$ as $|0\rangle$ and $|1\rangle$ respectively. The above table identifies the transposition operators which have maximum multipartite entangling power if the input set of optimization is restricted to fully separable states. On the other hand, if the optimization is carried out with the set of biseparable states as inputs, the maximum entangling power of the above twelve transposition operators cannot be achieved.

Remark 1. The optimal fully separable input states shown in Table I are not unique. For example, the transposition operator (1, 4) also produces maximal GGM from the state $\frac{1}{\sqrt{2}}(|000\rangle + |100\rangle)$.

Remark 2. To generate maximum GGM, one of the parties in the initial optimal state should have maximum coherence [74] with respect to the basis in which the permutation operator is written.

Biseparable inputs. Let us now change the set of input states to be the set of biseparable states. Again, we determine all the twelve transposition operators (i, j) which achieve the maximal entangling power, 0.5 (see Table II). To generate maximum GGM from the transposition operators, we find that one of the reduced two-party density matrices of the optimal biseparable input state has to be maximally entangled. We also observe that when a transposition(i, j) of Table II acts on fully separable inputs, the GGM of the resulting states are always same which turn out to be 0.33 and never reach the maximum possible value of 0.5. In other words, to have maximum GGM upon action of the transposition operators, it requires for the input states to have entanglement in some bipartition. Like the diagonal unitary operators, we notice that bipartite entanglement can sometimes provide advantages for a set of transposition operators, although it is not true, in general.

No.	Unitary	Biseparable Inputs
1	(1, 2)	$\frac{1}{\sqrt{2}}(000\rangle + 110\rangle)$
2	(1, 3)	$\frac{1}{\sqrt{2}}(000\rangle + 101\rangle)$
3	(1, 5)	$\frac{1}{\sqrt{2}}(000\rangle + 011\rangle)$
4	(2, 4)	$\frac{1}{\sqrt{2}}(001\rangle + 100\rangle)$
5	(2, 6)	$\frac{1}{\sqrt{2}}(001\rangle + 010\rangle)$
6	(3, 4)	$\frac{1}{\sqrt{2}}(010\rangle + 100\rangle)$
7	(3,7)	$\frac{1}{\sqrt{2}}(001\rangle + 010\rangle)$
8	(4, 8)	$\frac{1}{\sqrt{2}}(000\rangle + 011\rangle)$
9	(5, 6)	$\frac{1}{\sqrt{2}}(010\rangle + 100\rangle)$
10	(5,7)	$\frac{1}{\sqrt{2}}(001\rangle + 100\rangle)$
11	(6, 8)	$\frac{1}{\sqrt{2}}(000\rangle + 101\rangle)$
12	(7, 8)	$\frac{1}{\sqrt{2}}(001\rangle + 111\rangle)$

TABLE II. The second and third columns correspond to the transformation operators and their corresponding biseparable inputs respectively which lead to maximum GGM, 0.5. In this case, two-qubit reduced state is always maximally entangled, $\frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$. Note that the entangling power of the above unitaries cannot reach maximum when optimizing over the set of fully separable states.

We will now report an opposite picture, i.e., fully separable input states can give rise to a maximal GME state which biseparable states fail to attain.

C. Entangling power of special classes of unitary operators

Let us consider a specific class of unitary operators in \mathbb{C}^8 . It was shown [53] that in the case of three-qubits, any unitary operator can be decomposed into a number of two-qubit and single qubit, along with two special kinds of three-qubit unitary operators, given by

$$U_{sp}^{1} = \exp\left[i\left(\sum_{i=x,y,z} J_{i}\sigma_{i}\otimes\sigma_{i}\otimes\sigma_{z}\right)\right],$$
 (16)

and

$$U_{sp}^{2} = \exp\left[i\left(\sum_{i=x,y,z} J_{i}\sigma_{i}\otimes\sigma_{i}\otimes\sigma_{x} + J_{4}\mathbb{I}\otimes\mathbb{I}\otimes\sigma_{x}\right)\right].(17)$$

Here $\sigma_i s$ (i = x, y, z) are the Pauli matrices, \mathbb{I} is the identity operator and $J_i s$ are constants.

By examining optimizations over the set of fully separable states, we find that $|+\rangle^{\otimes 3}$ is an optimal input for the operators U_{sp}^1 with $(J_y - J_z) = n\pi/4$, $(n = \pm 1, \pm 3, \pm 5, ...)$ which leads to maximum GGM. In particular, we obtain

$$\begin{aligned} |\psi_{out}\rangle &\equiv U_{sp}^{1}|+\rangle^{\otimes 3} = \frac{1}{2\sqrt{2}} [a_{1}(|000\rangle + |110\rangle) \\ &+ \overline{a_{1}}(|001\rangle + |111\rangle) + a_{3}(|010\rangle + |100\rangle) \\ &+ \overline{a_{3}}(|011\rangle + |101\rangle)], \end{aligned}$$
(18)

$$G(|\psi_{out}\rangle) = 1 - \max[\frac{1}{2}(1 \pm \cos 2(J_y - J_z))], \quad (19)$$

which imposes the above mentioned restriction. Thus, we have identified a special subset $\tilde{U}_{sp}^1 \subset U_{sp}^1$ given by $\tilde{U}_{sp}^1 := \{U \in U_{sp}^1 | J_y - J_z = n\pi/4, n = \pm 1, \pm 3, \pm 5, \ldots\}.$

In contrast, by numerical simulation, we find that the elements of \tilde{U}_{sp}^1 do not always produce maximal GGM from the biseparbale states. We give below examples of both the two scenarios which can occur. When the operator in \tilde{U}_{sp}^1 with $(J_x = \pi/4, J_y = \pi/2, J_z = \pi/4)$ acts on $|+\rangle \otimes \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$, the GGM of the resulting state reaches maximum, 0.5. However, the operator with $(J_x = \pi/4, J_y = 11\pi/40, J_z = \pi/40)$ cannot find any biseparable state from \mathbb{S}^2 which can create maximal GGM.

On the other hand, starting from $|000\rangle$ as a fully separable and $\frac{1}{\sqrt{2}}(|0\rangle+|1\rangle)\frac{1}{\sqrt{2}}(|00\rangle+|11\rangle)$ as a biseparable input, U_{sp}^2 is capable of generating maximal GGM, when J_i s satisfy certain conditions. In particular, for the fully separable state, $J_1 - J_2 = \pi/4$ and $J_3 + J_4 = 0$ lead to maximal GME, while the conditions in case of the biseparable state are $J_2 = \pi/4$ and $J_3 - J_4 = \pi/4$.

IV. STATISTICAL ANALYSIS OF MULTIPARTITE ENTANGLING POWER OF UNITARY OPERATORS

Let us now move to consider more general scenarios. In particular, for a few classes of unitary operators, we want to assess their entangling power in terms of GGM after performing optimizations over the sets of fully separable and biseparable states. In contrast to our previous studies, we remove the constraint of achieving maximum GGM in this section. More specifically, the previous section was devoted to identify input states, both fully separable and biseparable, which are capable of creating states having maximum GGM, 0.5, after the action of different kinds of unitary operators. The study also revealed the interplay between entangling power of unitaries and the entanglement present in the inputs. The following analysis carried out on generic states and unitary operators can shed some more light on this interplay.

A. Entangling capability of diagonal quantum gates

A diagonal unitary matrix is a special kind of unitary operator having only nonvanishing diagonal elements, $U_D^{gen} =$ diag $(e^{i\phi_1}, \ldots, e^{i\phi_8})$. To find G_{\max} , we randomly choose ϕ_i s from uniform distribution over $[0, 2\pi]$. As we have shown in the preceding section (SubSec. III A), for a class of diagonal unitary operators, given by $U_D = \text{diag}(1, 1, 1, 1, 1, 1, 1, e^{i\phi})$, G_{\max}^2 obtained from the set of biseparable states is always higher than G_{\max}^3 obtained from the set of fully separable states for all values of ϕ (as depicted in Fig. 2). Let us now



FIG. 3. (Color online.) Normalized frequency distribution, $f_{G_{\text{max}}}$ (vertical axis) in Eq. (20) against G_{max} (horizontal axis) where the unitary is chosen to be diagonal, U^{diag} and $\phi \in [0, 2\pi]$ s are chosen from uniform distribution randomly. Green solid line represents $f_{G_{\text{max}}}$, when the optimization is performed over biseparable states while blue dashed line is for the set of fully-separable states. Both the axes are dimensionless.

elaborate on whether such hierarchies are maintained for randomly generated diagonal unitary operators.

We choose a set $\{\phi_i\}_{i=1}^8$ of random values of ϕ_i s to generate a fixed eight-dimensional diagonal unitary operator U_D^{gen} . For that operator, we compute the multipartite entangling power G_{\max}^i (i=2,3) by performing maximization over the inputs from \mathbb{S}^2 and \mathbb{S}^3 . We repeat this process for 5×10^3 sets of random choices of ϕ_i s, or equivalently, for 5×10^3 randomly generated diagonal unitary operators. Unlike the diagonal unitary operator U_D , here we observe that there are diagonal unitary operators U_D^{gen} , which produce higher GGM from fully separable inputs than that from the biseparable ones. However, such unitary operators are very rare. To analyze the situation more carefully, we study the normalized frequency distribution of G_{\max} , denoted as $f_{G_{\max}}$. Mathematically,

$$f_{G_{\max}} = \frac{\text{Number of states having } G_{\max}}{\text{Total number of states simulated}},$$
 (20)

which is depicted in Fig. 3. By examining the averages of the distributions, we find that $\overline{G}_{\max}^2 = 0.334$ for biseparable inputs and $\overline{G}_{\max}^3 = 0.237$ for fully separable ones. This result possibly indicates that, on an average, biseparable states are more effective inputs to create high GGM states than the fully separable inputs in case of diagonal quantum gates.

B. Specific kinds of unitary operators

We study entangling powers of two different classes of eight-dimensional unitary operators – (1) the unitary operators U_{sp}^1 in SubSec. III C and (2) a certain class of unitary operators, again involving Pauli matrices. In contrast to unitary operators described in SubSecs. IV A and IV C, we illustrate



FIG. 4. (Color online.) Contour plots in slices of multipartite entangling power, G_{\max}^3 of U_{sp}^1 , given in Eq. (16) with respect to the parameters J_x (x-axis), J_y (x-axis) and J_z (z-axis). The optimization is performed over the set of fully separable states. Both the axes are dimensionless.



FIG. 5. (Color online.) **Biseparable vs fully separable states as inputs.** When a special unitary operator, given in Eq. (21) acts on fully separable (triangles) or biseparable states (circles), the maximum GGM, G_{max} (ordinate) produced is plotted with respect to J_x (abscissa) for fixed $J_y = J_z = 0.1$. This is an example of an unitary operator for which higher GGM can be obtained from fully separable inputs compared to the biseparable ones for certain values of J_x . Both the axes are dimensionless.

some unitary operators belonging to these classes which can create higher GME states from fully separable inputs compared to biseparable ones.

Let us first determine the entangling power of $U_{sp}^1 = \exp\left[i(\sum_{i=x,y,z} J_i \sigma_i \otimes \sigma_i \otimes \sigma_z)\right]$, given in Eq. (16). In the previous section, we have found the condition on the parameters J_i s of U_{sp}^1 which leads to $G_{\max}^3 = 0.5$ for fully separable states although $G_{\max}^2 < 0.5$ for such operators, \tilde{U}_{sp}^1 . Hence, fully separable states are better inputs for this class of uni-

tary operators, \tilde{U}_{sp}^1 , than the biseparable ones. Let us find out G_{\max}^3 for arbitrary U_{sp}^1 in the (J_x, J_y, J_z) -slices. In Fig. 4, we illustrate the contours of entangling power, $G_{\max}^3(U_{sp}^1)$ by varying (J_x, J_y, J_z) . It clearly manifests that these special unitaries can typically create moderate amount of GGM from fully separable inputs, except some small regions.

Now consider the other class of special eight-dimensional unitary operators whose motivation comes from many-body physics. Those operators can be represented as

$$U_{sp}^{3} = \exp\left(-i\sum_{i=x,y,z} J_{i}\sigma_{i}\otimes\sigma_{i}\otimes\sigma_{i}\right), \qquad (21)$$

where J_i s can take any real value. By fixing the values of J_i s, a single unitary operator is obtained for which G^i_{\max} have to be determined by maximizing over the sets \mathbb{S}^i , i = 2, 3.

Case 1 (exactly one $J_i \neq 0$). Consider U_{sp}^3 for which exactly one of the J_i s is nonvanishing. In this case, for $J_i \neq 0$; i = x, y, z, we have $G_{\max}^3 = \cos^2 J_i$ when $(4n + 1)\pi/4 \leq J_i \leq (4n + 3)\pi/4$ and it is $\sin^2 J_i$ when $(4n+3)\pi/4 \leq J_i \leq (4n+5)\pi/4$, *n* is any integer. Note that it achieves maximum GGM when the nonvanishing $J_i = n\pi/4$ $(n = \pm 1, \pm 3, \ldots)$.

When we optimize on only biseparable states, it also exhibits almost a similar amount of entangling power as U_{sp}^3 , although not exactly the same. We notice that the optimal input biseparable states in this case are very close to become fully separable. Suppose we consider the biseparable state of the form $|1\rangle \otimes (\cos \theta |00\rangle + \sin \theta |11\rangle$. For a single nonvanishing J_i , we find that it gives almost the same amount of GGM as $|100\rangle$ when θ is chosen to be very small. In fact, G_{\max}^2 converges to G_{\max}^3 when $\theta \to 0$.

Case 2 (more than one $J_i \neq 0$). When more than one of the J_i s are nonvanishing, we do not have such simplified results like we had in the previous case, the picture is more involved here. To gain some knowledge about their entangling power, let us choose $J_y = J_z = 0.1$ while J_x varies. Like the operators U_{sp}^1 , these are another class of unitary operators, which can create higher GGM from fully separable inputs in comparison to the biseparable inputs (see Fig. 5). For example, when $J_x = \pi/4$, from the set of fully separable states, one obtains $G_{\max}^3 = 0.495$ while it is 0.469 when optimization is performed on \mathbb{S}^2 .

C. Randomly generated unitary operators

In this subsection, we scrutinize the situation when 10^4 Haar uniformly generated unitary matrices, U(8) [75] act on the sets of fully separable and biseparable input states. We perform optimization over inputs for each and every choice of such unitary operator. Like diagonal quantum gates, randomly generated unitary operators acted on optimal biseparable states are again capable of producing higher GME states than that from the optimal fully separable states. In this case, the normalized frequency distribution, defined in Eq. (20), depicted in Fig. 6 confirms that to generate high GGM in the



FIG. 6. (Color online.) $f_{G_{\text{max}}}$ (ordinate) against G_{max} (abscissa) where the unitary, U(8) is generated Haar uniformly. Dark (blue) and green represent $f_{G_{\text{max}}}$ obtained from the set of fully separable states biseparable states respectively. Both the axes are dimensionless.

resulting states, initial entanglement can be useful. Moreover, we observe that GGM created by Haar uniformly generated unitary operators is, on an average, higher than what can be obtained by the action of diagonal unitary operators. In this case, the mean of the distribution, $\overline{G}_{\max}^2 = 0.44$ for biseparable inputs and $\overline{G}_{\max}^3 = 0.39$ for the fully separable ones while the corresponding standard deviations are 0.018 and 0.03 respectively.

V. CONCLUSION

Entanglement produced at the different stages of quantum information processing play a significant role to achieve quantum advantages. To implement a quantum algorithm, a quantum circuit has to be designed which is composed of a set of gates, represented by unitary operators. The entangling power of a fixed quantum gate can be defined as the average or the maximum entanglement created upon action on the set of product states. In case of unitary operators acting on multipartite separable states, genuinely multipartite entangled (GME) states can, in general, be produced, thereby demonstrating their entangling capability. However, there can be different classes of separable states in a multipartite domain including fully separable states which have no entanglement in any bipartition, and states having some entanglement present in one of the bipartitions.

In literature, entangling power of a unitary operator in a multipartite system was determined by its entanglement generation capability in its reduced subsystems or in all of its bipartitions [66, 68]. Going beyond this conventional approach, we quantified multipartite entangling power of a given unitary operator via its potential to create genuine multipartite entangled outputs after the action on multipartite separable states. We proposed that the way the operators act on various classes of separable states can be utilized to introduce the concept of hierarchies among those operators. To make the hierarchies meaningful, we suggested that the sets of states on which the maximization is performed should ideally be disjoint. We also proved that the entangling power is invariant under the local unitary transformations.

In order to generate genuine multipartite entanglement, we find that the input entanglement present in some of the bipartitions of the input states is not always helpful. Specifically, we identified several classes of unitary operators and their corresponding k-separable inputs which lead to maximum genuine multipartite entangled states as outputs. We discovered that in three-qubits, there are unitary operators such as diagonal and Haar uniformly generated unitary operators that can result in the generation of higher genuine multipartite entanglement, on an average, from biseparable states than that from the fully separable ones. However, the opposite also holds on a substantial number of occasions. Our studies introduce a novel approach to classify multipartite unitary operators in terms of their capability to create genuine multipartite entanglement and the inputs which are responsible to achieve that.

ACKNOWLEDGMENTS

The authors acknowledge the support from the Interdisciplinary Cyber Physical Systems (ICPS) program of the Department of Science and Technology (DST), India, Grant No.: DST/ICPS/QuST/Theme- 1/2019/23 and the use of cluster computing facility at the Harish-Chandra Research Institute.

- R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Rev. Mod. Phys. 81, 865 (2009).
- [2] K. Modi, A. Brodutch, H. Cable, T. Paterek, and V. Vedral, Rev. Mod. Phys. 84, 1655 (2012).
- [3] A. Bera, T. Das, D. Sadhukhan, S. S. Roy, A. Sen(De), and U. Sen, Reports on Progress in Physics 81, 024001 (2017).
- [4] H. J. Kimble, Nature (London) 453, 1023 (2008), arXiv:0806.4195 [quant-ph].
- [5] C. H. Bennett and S. J. Wiesner, Phys. Rev. Lett. 69, 2881 (1992).
- [6] D. Bruß, G. M. D'Ariano, M. Lewenstein, C. Macchiavello, A. Sen(De), and U. Sen, Phys. Rev. Lett. 93, 210501 (2004).
- [7] D. Bruß, M. Lewenstein, A. Sen(De), U. Sen, G. M. D'Ariano, and C. Macchiavello, Int. J. Quant. Info. 4 (2006), 10.1142/S0219749906001888.
- [8] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, Phys. Rev. Lett. **70**, 1895 (1993).
- [9] M. Murao, D. Jonathan, M. B. Plenio, and V. Vedral, Phys. Rev. A 59, 156 (1999).
- [10] A. Sen(De) and U. Sen, Phys. Rev. A 81, 012308 (2010).

- [11] M. Hillery, V. Bužek, and A. Berthiaume, Phys. Rev. A 59, 1829 (1999).
- [12] R. Raussendorf and H. J. Briegel, Phys. Rev. Lett. 86, 5188 (2001).
- [13] H. J. Briegel and R. Raussendorf, Phys. Rev. Lett. 86, 910 (2001), arXiv:quant-ph/0004051 [quant-ph].
- [14] H. J. Briegel, D. E. Browne, W. Dür, R. Raussendorf, and M. Van den Nest, Nature 5, 19 (2009).
- [15] M. Lewenstein, A. Sanpera, V. Ahufinger, B. Damski, A. Sen(De), and U. Sen, Advances in Physics 56, 243 (2007), https://doi.org/10.1080/00018730701223200.
- [16] L. Amico, R. Fazio, A. Osterloh, and V. Vedral, Rev. Mod. Phys. 80, 517 (2008).
- [17] G. D. Chiara and A. Sanpera, Reports on Progress in Physics 81, 074002 (2018).
- [18] T.-C. Wei, D. Das, S. Mukhopadyay, S. Vishveshwara, and P. M. Goldbart, Phys. Rev. A 71, 060305 (2005).
- [19] A. Biswas, R. Prabhu, A. Sen(De), and U. Sen, Phys. Rev. A 90, 032301 (2014).
- [20] S. Haldar, S. Roy, T. Chanda, A. Sen(De), and U. Sen, Phys. Rev. B 101, 224304 (2020).
- [21] S. Sachdev, *Quantum Phase Transitions*, 2nd ed. (Cambridge University Press, 2011).
- [22] M. Heyl, Reports on Progress in Physics 81, 054001 (2018).
- [23] H.-S. Zhong, Y. Li, W. Li, L.-C. Peng, Z.-E. Su, Y. Hu, Y.-M. He, X. Ding, W. Zhang, H. Li, L. Zhang, Z. Wang, L. You, X.-L. Wang, X. Jiang, L. Li, Y.-A. Chen, N.-L. Liu, C.-Y. Lu, and J.-W. Pan, Phys. Rev. Lett. **121**, 250505 (2018).
- [24] X.-L. Wang, Y.-H. Luo, H.-L. Huang, M.-C. Chen, Z.-E. Su, C. Liu, C. Chen, W. Li, Y.-Q. Fang, X. Jiang, J. Zhang, L. Li, N.-L. Liu, C.-Y. Lu, and J.-W. Pan, Phys. Rev. Lett. **120**, 260502 (2018).
- [25] N. Friis, O. Marty, C. Maier, C. Hempel, M. Holzäpfel, P. Jurcevic, M. B. Plenio, M. Huber, C. Roos, R. Blatt, and B. Lanyon, Phys. Rev. X 8, 021012 (2018).
- [26] C. SONG, K. XU, Y.-R. ZHANG, X. ZHANG, W. LIU, Q. GUO, Z. WANG, W. REN, J. HAO, H. FENG, H. FAN, D. ZHENG, D.-W. WANG, H. WANG, and S.-Y. ZHU, Science **365** (2019).
- [27] M. Żukowski, A. Zeilinger, M. A. Horne, and A. K. Ekert, Phys. Rev. Lett. 71, 4287 (1993).
- [28] A. Sen(De), U. Sen, i. c. v. Brukner, V. Bužek, and M. Żukowski, Phys. Rev. A 72, 042310 (2005).
- [29] P. Walther, K. J. Resch, and A. Zeilinger, Phys. Rev. Lett. 94, 240501 (2005).
- [30] A. Acín, J. I. Cirac, and M. Lewenstein, Nature Physics 3, 256 (2007).
- [31] D. Cavalcanti, M. L. Almeida, V. Scarani, and A. Acín, Nature Communications 2, 184 (2011).
- [32] D. Sadhukhan, S. S. Roy, A. K. Pal, D. Rakshit, A. Sen(De), and U. Sen, Phys. Rev. A 95, 022301 (2017).
- [33] R. Banerjee, S. Ghosh, S. Mal, and A. Sen(De), Phys. Rev. Research 2, 043355 (2020).
- [34] P. Halder, S. Mal, and A. Sen(De), Phys. Rev. A **104**, 062412 (2021).
- [35] C. Mahto, V. Pathak, A. K. S., and A. Shaji, Phys. Rev. A 106, 012427 (2022).
- [36] B. Kraus and J. I. Cirac, Phys. Rev. A 63, 062309 (2001).
- [37] X. Zang, C. Yang, F. Ozaydin, W. Song, and Z. L. Cao, Scientific Reports 5 (2015), 10.1038/srep16245.
- [38] A. Sharma and A. A. Tulapurkar, Phys. Rev. A 101, 062330 (2020).
- [39] S. Daiss, S. Langenfeld, S. Welte, E. Distante, P. Thomas, L. Hartung, O. Morin, and G. Rempe, Science 371, 614 (2021),

https://www.science.org/doi/pdf/10.1126/science.abe3150.

- [40] P. Halder, R. Banerjee, S. Ghosh, A. K. Pal, and A. Sen(De), Phys. Rev. A 106, 032604 (2022).
- [41] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information: 10th Anniversary Edition* (Cambridge University Press, 2010).
- [42] A. Barenco, C. H. Bennett, R. Cleve, D. P. DiVincenzo, N. Margolus, P. Shor, T. Sleator, J. A. Smolin, and H. Weinfurter, Phys. Rev. A 52, 3457 (1995).
- [43] J. J. Vartiainen, M. Möttönen, and M. M. Salomaa, Phys. Rev. Lett. 92, 177902 (2004).
- [44] Y. Makhlin, Quantum Information Processing 1, 243 (2002).
- [45] P. Zanardi, C. Zalka, and L. Faoro, Phys. Rev. A 62, 030301 (2000).
- [46] P. Zanardi, Phys. Rev. A 63, 040304 (2001).
- [47] X. Wang and P. Zanardi, Phys. Rev. A 66, 044303 (2002).
- [48] S. Balakrishnan and R. Sankaranarayanan, Phys. Rev. A 79, 052339 (2009).
- [49] S. Balakrishnan and R. Sankaranarayanan, Phys. Rev. A 82, 034301 (2010).
- [50] L. Clarisse, S. Ghosh, S. Severini, and A. Sudbery, Phys. Rev. A 72, 012314 (2005).
- [51] N. Khaneja and S. J. Glaser, Chemical Physics 267, 11 (2001).
- [52] F. Vatan and C. Williams, Phys. Rev. A 69, 032315 (2004).
- [53] F. Vatan and C. P. Williams, "Realization of a general threequbit quantum gate," (2004).
- [54] J. Zhang, J. Vala, S. Sastry, and K. B. Whaley, Phys. Rev. A 67, 042313 (2003).
- [55] M. B. Mansky, S. L. Castillo, V. R. Puigvert, and C. Linnhoff-Popien, "Near-optimal circuit construction via cartan decomposition," (2022).
- [56] J. I. Cirac, W. Dür, B. Kraus, and M. Lewenstein, Phys. Rev. Lett. 86, 544 (2001).
- [57] M. S. Leifer, L. Henderson, and N. Linden, Phys. Rev. A 67, 012306 (2003).
- [58] M. A. Nielsen, C. M. Dawson, J. L. Dodd, A. Gilchrist, D. Mortimer, T. J. Osborne, M. J. Bremner, A. W. Harrow, and A. Hines, Phys. Rev. A 67, 052301 (2003).
- [59] F. Galve, F. Plastina, M. G. A. Paris, and R. Zambrini, Phys. Rev. Lett. **110**, 010501 (2013).
- [60] F.-Z. Kong, J.-L. Zhao, M. Yang, and Z.-L. Cao, Phys. Rev. A 92, 012127 (2015).
- [61] Z. Guan, H. He, Y.-J. Han, C.-F. Li, F. Galve, and G.-C. Guo, Phys. Rev. A 89, 012324 (2014).
- [62] W. Dür, G. Vidal, J. I. Cirac, N. Linden, and S. Popescu, Phys. Rev. Lett. 87, 137901 (2001).
- [63] G. Vidal, K. Hammerer, and J. I. Cirac, Phys. Rev. Lett. 88, 237902 (2002).
- [64] K. Hammerer, G. Vidal, and J. I. Cirac, Phys. Rev. A 66, 062321 (2002).
- [65] D. Morachis Galindo and J. A. Maytorena, Phys. Rev. A 105, 012601 (2022).
- [66] A. J. Scott, Phys. Rev. A 69, 052330 (2004).
- [67] S. A. Fenner, "A note on the entangling properties of the c-sign and related quantum gates," (2019).
- [68] T. Linowski, G. Rajchel-Mieldzioć, and K. Życzkowski, Journal of Physics A: Mathematical and Theoretical 53, 125303 (2020).
- [69] M. Blasone, F. Dell'Anno, S. De Siena, and F. Illuminati, Phys. Rev. A 77, 062304 (2008).
- [70] W. Dür, G. Vidal, and J. I. Cirac, Phys. Rev. A 62, 062314 (2000).
- [71] T.-C. Wei and P. M. Goldbart, Phys. Rev. A 68, 042307 (2003).
- [72] S. K. Hazra and A. Sen(De), in preparation (2023).

- [73] D. M. Greenberger, M. A. Horne, and A. Zeilinger, *Bell's theorem, quantum theory and conceptions of the universe* (Kluwer, Netherlands, 1989).
- [74] A. Streltsov, G. Adesso, and M. B. Plenio, Rev. Mod. Phys. 89, 041003 (2017).
- [75] K. Zyczkowski and M. Kus, Journal of Physics A: Mathematical and General 27, 4235 (1994).