

Grouped Domination Parameterized by Vertex Cover, Twin Cover, and Beyond*

Tesshu Hanaka¹[0000-0001-6943-856X], Hirotaka Ono²[0000-0003-0845-3947], Yota Otachi²[0000-0002-0087-853X], and Saeki Uda²

¹ Kyushu University, Fukuoka, Japan hanaka@inf.kyushu-u.ac.jp

² Nagoya University, Nagoya, Japan ono@nagoya-u.jp, otachi@nagoya-u.jp, uda.saeki.z4@s.mail.nagoya-u.ac.jp

Abstract. A dominating set S of graph G is called an r -grouped dominating set if S can be partitioned into S_1, S_2, \dots, S_k such that the size of each unit S_i is r and the subgraph of G induced by S_i is connected. The concept of r -grouped dominating sets generalizes several well-studied variants of dominating sets with requirements for connected component sizes, such as the ordinary dominating sets ($r = 1$), paired dominating sets ($r = 2$), and connected dominating sets (r is arbitrary and $k = 1$). In this paper, we investigate the computational complexity of r -GROUPED DOMINATING SET, which is the problem of deciding whether a given graph has an r -grouped dominating set with at most k units. For general r , r -GROUPED DOMINATING SET is hard to solve in various senses because the hardness of the connected dominating set is inherited. We thus focus on the case in which r is a constant or a parameter, but we see that r -GROUPED DOMINATING SET for every fixed $r > 0$ is still hard to solve. From the observations about the hardness, we consider the parameterized complexity concerning well-studied graph structural parameters. We first see that r -GROUPED DOMINATING SET is fixed-parameter tractable for r and treewidth, which is derived from the fact that the condition of r -grouped domination for a constant r can be represented as monadic second-order logic (MSO_2). This fixed-parameter tractability is good news, but the running time is not practical. We then design an $O^*(\min\{(2\tau(r+1))^\tau, (2\tau)^{2\tau}\})$ -time algorithm for general $r \geq 2$, where τ is the twin cover number, which is a parameter between vertex cover number and clique-width. For paired dominating set and trio dominating set, i.e., $r \in \{2, 3\}$, we can speed up the algorithm, whose running time becomes $O^*((r+1)^\tau)$. We further argue the relationship between FPT results and graph parameters, which draws the parameterized complexity landscape of r -GROUPED DOMINATING SET.

Keywords: Dominating Set · Paired Dominating Set · Parameterized Complexity · Graph Structural Parameters.

* Partially supported by JSPS KAKENHI Grant Numbers JP17H01698, JP17K19960, JP18H04091, JP20H05793, JP20H05967, JP21K11752, JP21H05852, JP21K17707, JP21K19765, and JP22H00513.

1 Introduction

1.1 Definition and motivation

Given an undirected graph $G = (V, E)$, a vertex set $S \subseteq V$ is called a *dominating set* if every vertex in V is either in S or adjacent to a vertex in S . The dominating set problem is the problem of finding a dominating set with the minimum cardinality. Since the definition of dominating set, i.e., covering all the vertices via edges, is natural, many practical and theoretical problems are modeled as dominating set problems with additional requirements; many variants of dominating set are considered and investigated. Such variants somewhat generalize or extend the ordinary dominating set based on theoretical or applicational motivations. In this paper, we focus on variants that require the dominating set to satisfy specific connectivity and size constraints. One example considering connectivity is the connected dominating set. A dominating set is called a *connected dominating set* if the subgraph induced by a dominating set is connected. Another example is the paired dominating set. A paired dominating set is a dominating set of a graph such that the subgraph induced by it admits a perfect matching.

This paper introduces the r -grouped dominating set, which generalizes the connected dominating set, the paired dominating set, and some other variants. A dominating set S is called an *r -grouped dominating set* if S can be partitioned into $\{S_1, S_2, \dots, S_k\}$ such that each S_i is a set of r vertices and $G[S_i]$ is connected. We call each S_i a *unit*. The r -grouped dominating set generalizes both the connecting dominating set and the paired dominating set in the following sense: a connecting dominating set with r vertices is equivalent to an r -grouped dominating set of one unit, and a paired dominating set with k pairs is equivalent to a 2-grouped dominating set with k units.

This paper investigates the parameterized complexity of deciding whether a given graph has an r -grouped dominating set with k units. The parameters that we focus on are so-called graph structural parameters, such as vertex cover number and twin-cover number. The results obtained in this paper are summarized in Our Contribution (Section 1.3).

1.2 Related work

An enormous number of papers study the dominating set problem, including the ones strongly related to the r -grouped dominating set.

The dominating set problem is one of the most important graph optimization problems. Due to its NP-hardness, its tractability is finely studied from several aspects, such as approximation, solvable graph classes, fast exact exponential-time solvability, and parameterized complexity. Concerning the parameterized complexity, the dominating set problem is W[2]-complete for solution size k ; it is unlikely to be fixed-parameter tractable [15]. On the other hand, since the dominating set can be expressed in MSO_1 , it is FPT when parametrized by clique-width or treewidth (see, e.g., [27]).

The connected dominating set is a well-studied variant of dominating set. This problem arises in communication and computer networks such as mobile ad hoc networks. It is also $W[2]$ -hard when parameterized by the solution size [15]. Furthermore, the connected dominating set also can be expressed in MSO_1 ; it is FPT when parametrized by clique-width and treewidth as in the ordinary dominating set problem. Furthermore, single exponential-time algorithms for connected dominating set parameterized by treewidth can be obtained by the Cut & Count technique [16] or the rank-based approach [4].

The notion of the paired dominating set is introduced in [25,26] by Haynes and Slater as a model of dominating sets with pairwise backup. It is NP-hard on split graphs, bipartite graphs [10], graphs of maximum degree 3 [8], and planar graphs of maximum degree 5 [36], whereas it can be solved in polynomial time on strongly-chordal graphs [9], distance-hereditary graphs [30], and AT-free graphs [36]. There are several graph classes (e.g., strongly orderable graphs [33]) where the paired dominating set problem is tractable, whereas the ordinary dominating set problem remains NP-hard. For other results about the paired dominating set, see a survey [17].

1.3 Our contributions

This paper provides a unified view of the parameterized complexity of dominating set problem variants with connectivity and size constraints.

As mentioned above, an r -grouped dominating set of G with 1 unit is equivalent to a connected dominating set with size r , which implies that some hardness results of r -GROUPED DOMINATING SET for general r are inherited directly from CONNECTED DOMINATING SET. From these, we mainly consider the case where r is a constant or a parameter.

Unfortunately, r -GROUPED DOMINATING SET for $r = 1, 2$ is also hard to solve again because 1-GROUPED DOMINATING SET and 2-GROUPED DOMINATING SET are respectively the ordinary dominating set problem and the paired dominating set problem. Thus, it is worth considering whether a larger but constant r enlarges, restricts, or leaves unchanged the graph classes for which similar hardness results hold. A way to classify or characterize graphs of certain classes is to focus on graph-structural parameters. By observing that the condition of r -grouped dominating set can be represented as monadic second-order logic (MSO_2), we can see that r -GROUPED DOMINATING SET is fixed-parameter tractable for r and treewidth. Recall that the condition of the connected dominating set can be represented as monadic second-order logic (MSO_1), which implies that there might exist a gap between $r = 1$ and 2, or between $k = 1$ and $k > 1$. Although this FPT result is good news, its time complexity is not practical. From these observations, we focus on less generalized graph structural parameters, vertex cover number ν or twin cover number τ as a parameter, and design single exponential fixed-parameter algorithms for r -GROUPED DOMINATING SET.

Our algorithm is based on dynamic programming on nested partitions of a vertex cover, and its running time is $O^*(\min\{(2\nu(r+1))^\nu, (2\nu)^{2\nu}\})$ for general

$r \geq 2$. For paired dominating set and trio dominating set, i.e., $r \in \{2, 3\}$, we can tailor the algorithm to run in $O^*((r+1)^\nu)$ time by observing that the nested partitions of a vertex cover degenerate in some sense.

We then turn our attention to a more general parameter, the twin cover number. We show that, given a twin cover, r -GROUPED DOMINATING SET admits an optimal solution in which twin-edges do not contribute to the connectivity of r -units. This observation implies that these edges can be removed from the graph, and thus we can focus on the resultant graph of bounded vertex cover number. Hence, we can conclude that our algorithms still work when the parameter ν in the running time is replaced with twin cover number τ .

We further argue the relationship between FPT results and graph parameters. The perspective is summarized in Figure 1, which draws the parameterized complexity landscape of r -GROUPED DOMINATING SET.




Fig. 1. The complexity of r -GROUPED DOMINATING SET with respect to structural graph parameters. An edge between two parameters indicates that there is a function in the one above that lower-bounds the one below (e.g., treewidth \leq pathwidth).

2 Preliminaries

Let $G = (V, E)$ be an undirected graph. For a vertex subset $V' \subseteq V$, the subgraph induced by V' is denoted by $G[V']$. Also, let us denote by $N(v)$ and $N[v]$ the open neighborhood and the closed neighborhood of v , respectively. The degree of a vertex v is defined by $d(v) = |N(v)|$. The maximum degree of G is denoted by Δ .

A vertex set S is a *vertex cover* of G if for every edge $\{u, v\} \in E$, at least one of u, v is in S . The *vertex cover number* ν of G is defined by the size of

a minimum vertex cover of G . A minimum vertex cover of G can be found in $O^*(1.2738^\nu)$ time [7].³

Two vertices u and v are (*true*) *twins* if $N[u] = N[v]$. An edge $\{u, v\} \in E$ is a *twin edge* if u and v are true twins. A vertex set S is a *twin cover* if for every edge $\{u, v\} \in E$, either $\{u, v\}$ is a twin edge, or at least one of u, v is in S . The size τ of a minimum twin cover of G is called the *twin cover number* of G . A minimum twin cover of G can be found in $O^*(1.2738^\tau)$ time [21].

We briefly introduce basic terminology of parameterized complexity. Given an input size n and a parameter k , a problem is *fixed-parameter tractable (FPT)* if it can be solved in $f(k)n^{O(1)}$ time where f is some computable function. Also, a problem is *slice-wise polynomial (XP)* if it can be solved in $n^{f(k)}$ time. See standard textbooks (e.g., [15]) for more details.

2.1 r -Grouped Dominating Set

An r -grouped dominating set with k units in G is a family $\mathcal{D} = \{D_1, \dots, D_k\}$ of subsets of V such that D_i 's are mutually disjoint, $|D_i| = r$, $G[D_i]$ is connected for $1 \leq i \leq k$, and $\bigcup_{D \in \mathcal{D}} D$ is a dominating set of G . For simplicity, let $\bigcup \mathcal{D}$ denote $\bigcup_{D \in \mathcal{D}} D$. We say that \mathcal{D} is a minimum r -grouped dominating set if it is an r -grouped dominating set with the minimum number of units.

r -GROUPED DOMINATING SET

Input: A graph G and positive integers r and k .

Question: Is there an r -grouped dominating set with at most k units in G ?

3 Basic Results

In this section, we prove r -GROUPED DOMINATING SET is $W[2]$ -hard but XP when parameterized by $k + r$ and it is NP-hard even on planar bipartite graphs of maximum degree 3.

We first observe that finding an r -grouped dominating set with at most 1 unit is equivalent to finding a connected dominating set of size r . Thus, the $W[2]$ -hardness of r -GROUPED DOMINATING SET parameterized by r follows the one of CONNECTED DOMINATING SET parameterized by the solution size. Also, the case $r = 1$ follows immediately from the hardness of the ordinary DOMINATING SET, which is $W[2]$ -complete on split graphs and bipartite graphs [34]. In the remaining part of this section, we discuss the hardness results only for the cases $r \geq 2$ and $k \geq 2$.

Theorem 3.1. *For every fixed $k \geq 1$, r -GROUPED DOMINATING SET is $W[2]$ -hard when parameterized by r even on split graphs.*

Proof. We give a reduction from DOMINATING SET on split graphs. Let $\langle G = (C \cup I, E), r \rangle$ be an instance of DOMINATING SET where C forms a clique and

³ The O^* notation suppresses the polynomial factors of the input size.

I forms an independent set. Without loss of generality, we suppose that $|C| \geq 2$ and $|I| \geq 2$. We create k copies $G_1 = (V_1, E_1), \dots, G_k = (V_k, E_k)$ of G where $V_i = \{v^{(i)} \mid v \in V\}$ and $E_i = \{e^{(i)} \mid e \in E\}$. Note that $V_i = C_i \cup I_i$. Finally, we make $\bigcup_i C_i$ a clique. The resulting graph G' is clearly a split graph.

We show that there is a dominating set of size at most r in G if and only if there is an r -grouped dominating set with at most k units in G' .

Suppose that there is a dominating set D of size at most r in G . Without loss of generality, we can assume that $D \subseteq C$ [3]. Then we define $D'_i = \{v^{(i)} \mid v \in D\}$ for $1 \leq i \leq k$ and $D' = \bigcup_i D'_i$. Since D is a dominating set in G , so is D'_i on G_i for each i . Thus, D' is a dominating set in G' . Because G_i is a split graph and a clique and $D'_i \subseteq C_i$, D'_i is a connected dominating set of G_i of size at most r . If $|D'_i| < r$, we arbitrarily add $r - |D'_i|$ vertices in G_i to D'_i . Then, we have a connected dominating set D'_i of G_i of size exactly r for each i , which can be regarded as a unit of size r of an r -grouped dominating set. Clearly, $\{D'_i \mid 1 \leq i \leq k\}$ is an r -grouped dominating set with k units in G' .

Conversely, suppose that there is an r -grouped dominating set \mathcal{D} with at most k units in G' . Then there is a vertex set $D_i = \bigcup \mathcal{D} \cap V_i$ of size at most r in some G_i by $|\bigcup \mathcal{D}| \leq rk$. Since $|I_i| \geq 2$, D_i contains at least one vertex in C_i . Moreover, any vertex not in V_i cannot dominate vertices in I_i . This means that D_i is a dominating set in G_i . Since G_i is a copy of G , there is a dominating set of size at most r in G . \square

By a similar reduction, we also show that r -GROUPED DOMINATING SET is $W[2]$ -hard when parameterized by k .

Theorem 3.2. *For every fixed $r \geq 1$, r -GROUPED DOMINATING SET is $W[2]$ -hard when parameterized by k even on split graphs.*

Proof. We give a reduction from DOMINATING SET on split graphs. Let $\langle G = (C \cup I, E), k \rangle$ be an instance of DOMINATING SET where C forms a clique and I forms an independent set. Without loss of generality, we suppose that $|C| \geq 2$ and $|I| \geq 2$. We create r copies $G_1 = (V_1, E_1), \dots, G_r = (V_r, E_r)$ of G where $V_i = \{v^{(i)} \mid v \in V\}$ and $E_i = \{e^{(i)} \mid e \in E\}$. Note that $V_i = C_i \cup I_i$. Then we make $\bigcup_i C_i$ a clique. The resulting graph G' is clearly a split graph.

We show that there is a dominating set of size at most k in G if and only if there is an r -grouped dominating set with at most k units in G' .

Suppose that there is a dominating set D of size at most k in G . Without loss of generality, we can assume that $D \subseteq C$. For each $v \in D$, we define $D_v = \{v^{(i)} \mid 1 \leq i \leq r\}$. Furthermore, let $\mathcal{D} = \{D_v \mid v \in D\}$. We see that \mathcal{D} is an r -grouped dominating set with at most k units in G' . Since D is a dominating set in G and G' consists of r copies of G , $\bigcup \mathcal{D}$ is clearly a dominating set in G' . Furthermore, because $\bigcup_i C_i$ is a clique, each D_v forms a clique of size r , which can be regarded as a unit. By the assumption that $|D| \leq k$, we conclude that \mathcal{D} is an r -grouped dominating set with at most k units in G' .

Conversely, suppose that there is an r -grouped dominating set \mathcal{D} with at most k units in G' . We see that there is a dominating set D of size at most k in some G_i . Indeed, $\bigcup \mathcal{D} \cap V_i$ is a dominating set D of size at most k in G_i because

$|\bigcup \mathcal{D}| \leq rk$ and there is a vertex in $\bigcup \mathcal{D} \cap C_i$ by $|I| \geq 2$. This completes the proof. \square

Furthermore, we show the $W[2]$ -hardness of r -GROUPED DOMINATING SET on bipartite graphs.

Theorem 3.3. *For every fixed $k \geq 1$, r -GROUPED DOMINATING SET is $W[2]$ -hard when parameterized by r even on bipartite graphs.*

Proof. We reduce DOMINATING SET on split graphs to r -GROUPED DOMINATING SET.

We are given an instance $\langle G = C \cup I, E \rangle, r$ of DOMINATING SET. Without loss of generality, we assume $|C| \geq 2$, $|I| \geq 2$. Moreover, we assume that if $\langle G = C \cup I, E \rangle, r$ is a yes-instance, there is a dominating set D of size at most r such that $D \subseteq C$ [3].

We first delete all the edges in the clique C , and add two edges $\{s_1, t\}, \{s_2, t\}$ and connect t to all the vertices in C . The obtained graph is bipartite. We then create k copies $G_1 = (V_1 \cup \{s_1^{(1)}, s_2^{(1)}, t^{(1)}\}, E_1), \dots, G_k = (V_k \cup \{s_1^{(k)}, s_2^{(k)}, t^{(k)}\}, E_k)$ of the graph where $V_i = \{u^{(i)} \mid u \in V\}$ for $1 \leq i \leq k$. To connect G_1, \dots, G_k , we add edges $\{s_1^{(i)}, s_1^{(i+1)}\}$ for $1 \leq i \leq k-1$. The resulting graph denoted by G' remains bipartite.

In the following, we show that there is a dominating set of size at most r in G if and only if there is an $(r+1)$ -grouped dominating set with k units in G' . Suppose that there is a dominating set $D \subseteq C$ of size at most r in G . We assume that $|D| = r$ because otherwise we only have to add $r - |D|$ vertices in G to D arbitrarily. For each graph G_i , define $D_i = \{v^{(i)} \mid v \in D\}$. Since $t^{(i)}$ is connected to $s_1^{(i)}, s_2^{(i)}$, and all the vertices in the clique part C_i of G_i and D_i is a dominating set in $G_i[V_i]$, $D_i \cup \{t^{(i)}\}$ is a connected dominating set of size $r+1$ in G_i . Therefore, $\{D_i \cup \{t^{(i)}\} \mid 1 \leq i \leq k\}$ is an $(r+1)$ -grouped dominating set with k units in G' .

Conversely, let \mathcal{D} be an $(r+1)$ -grouped dominating set with k units in G' . Since $|\bigcup \mathcal{D}| \leq (r+1)k$, some G_i satisfies $|\bigcup \mathcal{D} \cap V_i \cup \{s_1^{(i)}, s_2^{(i)}, t^{(i)}\}| \leq (r+1)$. To dominate $s_2^{(i)}$, $\bigcup \mathcal{D}$ must contain $t^{(i)}$. Note that $r \geq 2$. Thus, $|\bigcup \mathcal{D} \cap V_i| \leq r$. Furthermore, G_i is bipartite and $|I| \geq 2$, hence there is a vertex $v^{(i)}$ in $\bigcup \mathcal{D} \cap C_i$. Let $D \subseteq V$ be a set in G corresponding to $\bigcup \mathcal{D} \cap V_i$. Then D is a dominating set of size r in G . Indeed, since $\bigcup \mathcal{D} \cap V_i$ dominates the independent set part of G_i , D also dominates the independent set part of G . Moreover, since G is a split graph, vertex $v \in V$ corresponding to $v^{(i)}$ dominates all the vertices in D . This completes the proof. \square

Theorem 3.4. *For every fixed $r \geq 1$, r -GROUPED DOMINATING SET is $W[2]$ -hard when parameterized by k even on bipartite graphs.*

Proof. We reduce DOMINATING SET on split graphs to r -GROUPED DOMINATING SET.

We are given an instance $\langle G = C \cup I, E \rangle, k$ of DOMINATING SET. Without loss of generality, if $\langle G = C \cup I, E \rangle, k$ is a yes-instance, there is a dominating

set D of size at most k such that $D \subseteq C$ [3]. Then we construct a bipartite graph G' as follows. First, delete all the edges in the clique C . Then G becomes a bipartite graph. We next add k paths of length r and connect an endpoint of each path to all the vertices in C . Let $P_i = (u_1^{(i)}, \dots, u_r^{(i)})$ denote such paths for $1 \leq i \leq k$, and $u_1^{(i)}$'s are the endpoints connected to C . The resulting graph G' is bipartite.

Suppose that G has a dominating set $D = \{v_1, \dots, v_{|D|}\} \subseteq C$ of size at most k . From D , we construct an r -grouped dominating set with k units. For each $v_i \in D$, we choose a path $v_i, u_1^{(i)}, \dots, u_{r-1}^{(i)}$ of length r as one unit of the r -grouped dominating set. If $|D| < k$, we choose the remaining $k - |D|$ paths $u_1^{(i)}, \dots, u_r^{(i)}$ for $|D| + 1 \leq i \leq r$. Let \mathcal{D} be the set of such k paths. Then \mathcal{D} is an r -grouped dominating set with k units because the length of each path in \mathcal{D} is r and $\bigcup \mathcal{D}$ contains D , which dominates all the vertices in the original G and the vertices in P_i 's.

Conversely, let \mathcal{D} be an r -grouped dominating set with k units in G' . To dominate an endpoint $u_r^{(i)}$ in P_i , $\bigcup \mathcal{D}$ must contain $u_{r-1}^{(i)}$, which implies $|\bigcup \mathcal{D} \cap \bigcup_i P_i| \geq k(r-1)$. Thus, we have $|(C \cup I) \cap \bigcup \mathcal{D}| \leq k$. Since $\bigcup \mathcal{D}$ is a dominating set of G' and any vertex in P_i 's cannot dominate I , $(C \cup I) \cap \bigcup \mathcal{D}$ is a dominating set of size k in G . \square

On the other hand, we can show that the problem is XP when parameterized by $k + r$.

Theorem 3.5. *r -GROUPED DOMINATING SET can be solved in $O^*(\Delta^{O(kr^2)})$ time.*

Proof. We guess the candidates of r -grouped dominating sets with at most k units. We first pick an arbitrary vertex v and branch $d(v) + 1$ cases. One case is that v is contained in \mathcal{D} . The vertices in the unit containing v is reachable from v via at most $r - 1$ edges. Since the number of such vertices is at most Δ^{r-1} , the choice of the other $r - 1$ vertices is at most $\binom{\Delta^{r-1}}{r-1} = \Delta^{O(r^2)}$. Thus the number of candidates of units that contains v is $\Delta^{O(r^2)}$. Another case is that v is not contained in \mathcal{D} . Then at least one neighbor of v is contained in \mathcal{D} . The number of candidates of units that contain it is also $\Delta^{O(r^2)}$. Therefore, the total number of candidates of units that dominate v is $\Delta^{O(r^2)}$. After guessing one unit, we repeatedly pick a non-dominated vertex and branch as above. The repetition occurs at most k times. Thus, the total running time is $\Delta^{O(kr^2)}$. \square

Corollary 3.6. *r -GROUPED DOMINATING SET belongs to XP when parameterized by $k + r$.*

Tripathi et al. [36] showed that PAIRED DOMINATING SET (equivalently, 2-GROUPED DOMINATING SET) is NP-complete for planar graphs with maximum degree 5. We show that r -GROUPED DOMINATING SET is NP-hard even on planar bipartite graphs of maximum degree 3 for every fixed $r \geq 1$. This strengthens the result by Tripathi et al. [36].

Theorem 3.7. *For every fixed $r \geq 1$, r -GROUPED DOMINATING SET is NP-complete on planar bipartite graphs of maximum degree 3.*

Proof. We reduce RESTRICTED PLANAR 3-SAT to r -GROUPED DOMINATING SET. RESTRICTED PLANAR 3-SAT is a variant of PLANAR 3-SAT such that each variable occurs in exactly three clauses, in at most two clauses positively and in at most two clauses negatively. It is known that RESTRICTED PLANAR 3-SAT is NP-complete [31].

Let ϕ be an instance of RESTRICTED PLANAR 3-SAT, n and m be the number of variables and clauses of ϕ , respectively. The incidence graph of ϕ is a bipartite graph such that it consists of variable vertices v_{x_i} 's corresponding to variables and clause vertices c_j 's corresponding to clauses. A variable vertex v_{x_i} is connected to a clause vertex c_j if C_j has a literal of x_i . The incidence graph of ϕ is planar.

For the incidence graph of ϕ , we construct the graph $G = (V, E)$ by replacing variable vertices by variable gadgets. For each variable x_i , its variable gadget is constructed as follows. We create three vertices $v_{x_i}, v_{\bar{x}_i}, y_i$, and then add edges $\{v_{x_i}, y_i\}, \{v_{\bar{x}_i}, y_i\}$. Furthermore, we attach a path $P_i^r = y_i z_i^{(1)} z_i^{(2)} \dots z_i^{(r-1)}$ of length $r - 1$ for each y_i . Here, we define $z_i^{(0)} = y_i$. Let $V_X = \{v_{x_i}, v_{\bar{x}_i} \mid i \in \{1, \dots, n\}\}$ and $V_C = \{c_j \mid j \in \{1, \dots, m\}\}$. For each variable x_i , v_{x_i} is connected to c_j if C_j has a positive literal of x_i , and $v_{\bar{x}_i}$ is connected to c_j if C_j has a negative literal of x_i . We complete the construction of the graph $G = (V, E)$. Figure 2 shows a concrete example of $G = (V, E)$ for ϕ . Notice that G is bipartite because V_X and V_C form independent sets, respectively, and P_i^r is a path. Furthermore, G is planar because the incidence graph of ϕ and the variable gadgets are planar. Finally, since each variable occurs in exactly three clauses, in at most two clauses positively and in at most two clauses negatively, the maximum degree of G is at most 3.

We are ready to show that ϕ is a yes-instance if and only if there is an r -grouped dominating set with at most n in G .

Suppose that we are given a truth assignment of ϕ . For each variable x_i , we select path $v_{x_i} y_i z_i^{(1)} z_i^{(2)} \dots z_i^{(r-2)}$ as a unit of an r -grouped dominating set if x_i is assigned to true. Otherwise, we select path $v_{\bar{x}_i} y_i z_i^{(1)} z_i^{(2)} \dots z_i^{(r-2)}$. The number of vertices in each unit is r . The unit of x_i dominates vertex $z_i^{(r-1)}$. Since each clause has at least one truth literal for the truth assignment, each clause vertex c_j is dominated by some unit. Therefore, the set of selected paths is an r -grouped dominating set with at most n units.

Conversely, we are given an r -grouped dominating set \mathcal{D} with at most n in G . For each i , $z_i^{(r-2)}$ must be contained in $\bigcup \mathcal{D}$. If not, $z_i^{(r-1)}$ is not dominated because of $r \geq 2$. Since P_i^r is a path of length $r - 1$ and the number of units is n , the vertices of a unit are selected from $\{v_{x_i}, v_{\bar{x}_i}\} \cup \{y_i, z_i^{(1)}, z_i^{(2)}, \dots, z_i^{(r-1)}\}$ for each i and the unit forms a path of length r . If vertex $z_i^{(r-1)}$ is contained in $\bigcup \mathcal{D}$, $\bigcup \mathcal{D}$ does not contain v_{x_i} and $v_{\bar{x}_i}$. Thus, we can remove $z_i^{(r-1)}$ from $\bigcup \mathcal{D}$ and add either v_{x_i} or $v_{\bar{x}_i}$ arbitrarily to $\bigcup \mathcal{D}$. Since $v_{x_i} y_i z_i^{(1)} z_i^{(2)} \dots z_i^{(r-2)}$ is a




Fig. 2. The graph G obtained by the reduction from an instance $\phi = (x_1 \vee x_2 \vee \bar{x}_4)(\bar{x}_1 \vee x_2 \vee \bar{x}_3)(\bar{x}_2 \vee x_3 \vee x_4)(\bar{x}_1 \vee \bar{x}_3 \vee \bar{x}_4)$ of RESTRICTED PLANAR 3-SAT to 3-GROUPED DOMINATING SET.

path of length r , this replacement does not collapse the property of r -grouped dominating set. Thus, we can suppose that the path of i th unit of \mathcal{D} has either v_{x_i} or $v_{\bar{x}_i}$ as an endpoint. Since $\bigcup \mathcal{D}$ is a dominating set, each vertex in V_C has at least one vertex in $\mathcal{D} \cap V_X$ as a neighbor. This implies that the assignment corresponding to the selection of endpoints of units is a truth assignment. This completes the proof. \square

In the proof of Theorem 3.7, the size of the constructed graph for ϕ is $O(rn + m)$. Thus, we have the following corollary.

Corollary 3.8. *For every fixed $r \geq 1$, r -GROUPED DOMINATING SET cannot be solved in time $2^{o(n+m)}$ on bipartite graphs unless ETH fails.*

4 Fast Algorithms Parameterized by Vertex Cover Number and by Twin Cover Number

In this section, we present FPT algorithms for r -GROUPED DOMINATING SET parameterized by vertex cover number ν . Our algorithm is based on dynamic programming on nested partitions of a vertex cover, and its running time is $O^*((2\nu(r+1))^\nu)$ for general $r \geq 2$. For the cases of $r \in \{2, 3\}$, we can tailor the algorithm to run in $O^*((r+1)^\nu)$ time by focusing on the fact that the nested partitions of a vertex cover degenerate in some sense.

We then turn our attention to a more general parameter twin cover number. We show that, given a twin cover, r -GROUPED DOMINATING SET admits an optimal solution in which twin-edges do not contribute to the connectivity of r -units. This implies that these edges can be removed from the graph, and thus we can focus on the resultant graph of bounded vertex cover number. Hence, we can conclude that our algorithms still work when the parameter ν in the running time is replaced with twin cover number τ .

Theorem 4.1. *For graphs of twin cover number τ , r -GROUPED DOMINATING SET can be solved in $O^*((2\tau(r+1))^\tau)$ time. For the cases of $r \in \{2, 3\}$, it can be solved in $O^*((r+1)^\tau)$ time.*

With a simple observation, Theorem 4.1 implies that r -GROUPED DOMINATING SET parameterized solely by τ is fixed-parameter tractable.

Corollary 4.2. *For graphs of twin cover number τ , r -GROUPED DOMINATING SET can be solved in $O^*((2\tau)^{2\tau})$ time.*

Proof. If $r < 2\tau - 1$, then the problem can be solved in $O^*((2\tau)^{2\tau})$ time by Theorem 4.1. Assume that $r \geq 2\tau - 1$. Let C be a connected component of the input graph. If $|V(C)| < r$, then we have a trivial no-instance. Otherwise, we construct a connected dominating set D of C with size exactly r , which works as a unit dominating C . We initialize D with a non-empty twin cover of size at most τ . Note that such a set can be found in $O^*(1.2738^\tau)$ time: if C is a complete graph, then we pick an arbitrary vertex $v \in V(C)$ and set $D = \{v\}$; otherwise, just find a minimum twin cover. Since C is connected, D is a dominating set of C . If $C[D]$ is not connected, we update D with a new element v adjacent to at least two connected components of $C[D]$. Since $|D| \leq \tau$ at the beginning, we can repeat this update at most $\tau - 1$ times, and after that $C[D]$ becomes connected and $|D| \leq 2\tau - 1 \leq r$. We finally add $r - |D|$ vertices arbitrarily and obtain a desired set. \square

In the next subsection, we first present an algorithm for 2-GROUPED DOMINATING SET parameterized by vertex cover number, which gives a basic scheme of our dynamic programming based algorithms. We then see how we extend the idea to 3-GROUPED DOMINATING SET. As explained above, these algorithms are based on dynamic programming (DP), and they compute certain function values on partitions of a vertex cover. Unfortunately, it is not obvious how to extend the strategy to general r . Instead, we consider nested partitions of a vertex cover for DP tables, which makes the running time a little slower though. In the last subsection, we see how a vertex cover can be replaced with a twin cover in the same running time in terms of order.

4.1 Algorithms parameterized by vertex cover number

Algorithm for 2-Grouped Dominating Set We first present an algorithm for the simplest case $r = 2$, i.e., the paired dominating set. Let $G = (V, E)$ be a graph and J be a vertex cover of G . Then, $I = V \setminus J$ is an independent set. The basic scheme of our algorithm follows the algorithm for the dominating set problem by Liedloff [29], which focuses on a partition of a given vertex cover J . For a minimum dominating set D , the vertex cover J is partitioned into three parts: $J \cap D$; $(J \setminus D) \cap N(J \cap D)$, that is, the vertices in $J \setminus D$ that are dominated by $J \cap D$; and $J \setminus N[J \cap D]$, that is, the remaining vertices. Note that the remaining vertices in $J \setminus N[J \cap D]$ are dominated by $I \cap D$. Once $J \cap D$ is fixed, a minimum $I \cap D$ is found by solving the set cover problem that reflects




Fig. 3. Partitioning a vertex cover into three parts.

the condition that $J \setminus N[J \cap D]$ must be dominated by $I \cap D$. The algorithm computes a minimum dominating set by solving set cover problems defined by all candidates of $J \cap D$.

To adjust the algorithm to 2-GROUPED DOMINATING SET, we need to handle the condition that a dominating set contains a perfect matching.

For each subset $J_D \subseteq J$, we find a subset $I_D \subseteq I$ (if any exists) of the minimum size such that $J_D \cup I_D$ can form a 2-grouped dominating set. Let X and Y be disjoint subsets of J , and let $I = \{v_1, v_2, \dots, v_{|I|}\}$ (see Fig. 3). For $j = 0, \dots, |I|$, we define an auxiliary table $A[X, Y, j]$ as the minimum size of $I' \subseteq \{v_1, v_2, \dots, v_j\}$ that satisfies the following conditions.

1. $Y \subseteq N(I')$,
2. $I' \cup X$ has a partition $\mathcal{D}^{(2)} = \{D_1^{(2)}, D_2^{(2)}, \dots, D_p^{(2)}\}$ with $p \leq k$ such that for all $i = 1, \dots, p$, $|D_i^{(2)}| = 2$ and $G[D_i^{(2)}]$ is connected.

We set $A[X, Y, j] = \infty$ if no $I' \subseteq \{v_1, v_2, \dots, v_j\}$ satisfies the conditions. We can easily compute $A[X, Y, 0] \in \{0, \infty\}$ as $A[X, Y, 0] = 0$ if and only if $G[X]$ has a perfect matching and $Y = \emptyset$. Now the following recurrence formula computes A :

$$A[X, Y, j+1] = \min \left\{ A[X, Y, j], \min_{u \in N(v_{j+1}) \cap X} A[X \setminus \{u\}, Y \setminus N(v_{j+1}), j] + 1 \right\}.$$

The recurrence finds the best way under the condition that we can use vertices from $v_1, v_2, \dots, v_j, v_{j+1}$ in a dominating set: not using v_{j+1} , or pairing v_{j+1} with $u \in N(v_{j+1}) \cap X$. We can compute all entries of A in $O^*(3^{|J|})$ time in a DP manner as there are only $3^{|J|}$ ways for choosing disjoint subsets X and Y of J .

Now we compute the minimum number of units in a 2-grouped dominating set of G (if any exists) by looking up some appropriate table entries of A . Let \mathcal{D} be a 2-grouped dominating set of G with $J_D = J \cap \bigcup \mathcal{D}$ and $I_D = I \cap \bigcup \mathcal{D}$. Since $\bigcup \mathcal{D}$ is a dominating set with no isolated vertex in $G[\bigcup \mathcal{D}]$, J_D dominates all vertices in I . Let $J_Y = J \setminus N[J_D]$. Then the definition of A implies that $A[J_D, J_Y, |I|] = |I_D|$. Conversely, if $X \subseteq J$ dominates I , $Y = J \setminus N[X]$, and $A[X, Y, |I|] \neq \infty$, then there is a 2-grouped dominating set with $(|X| + A[X, Y, |I|])/2$ units. Therefore, the minimum number of units in a 2-grouped dominating set of G is $\min\{(|X| + A[X, J \setminus N[X], |I|])/2 \mid X \subseteq J \text{ and } I \subseteq N(X)\}$, which can be computed in $O^*(2^{|J|})$ time given the table A . Thus the total running time of the algorithm is $O^*(3^{|J|})$.

Algorithm for 3-Grouped Dominating Set Next, we consider the case $r = 3$, i.e., the trio dominating set. Let $G = (V, E)$ be a graph, J be a vertex cover of G , and $I = V \setminus J$. The basic idea is the same as the case $r = 2$ except that we partition the vertex cover into four parts in the DP, and thus the recurrence formula for A is different. In the DP, the vertex cover J is partitioned into four parts depending on the partial solution corresponding to each table entry.

For each subset $J_D \subseteq J$, we find a subset $I_D \subseteq I$ (if any exists) of the minimum size such that $J_D \cup I_D$ can form a 3-grouped dominating set. Intuitively, the set F represents partial units that will later be completed to full units. Let X , F , and Y be disjoint subsets of J , and let $I = \{v_1, v_2, \dots, v_{|I|}\}$. For $j = 0, \dots, |I|$, we define $A[X, F, Y, j]$ as the minimum size of $I' \subseteq \{v_1, v_2, \dots, v_j\}$ that satisfies the following conditions:

1. $Y \subseteq N(I')$,
2. I' can be partitioned into two parts I'_2, I'_3 satisfying the following conditions:
 - $I'_2 \cup F$ has a partition $\mathcal{D}^{(2)} = \{D_1^{(2)}, D_2^{(2)}, \dots, D_p^{(2)}\}$ with $p \leq k$ such that for all $i = 1, \dots, p$, $|D_i^{(2)}| = 2$ and $G[D_i^{(2)}]$ is connected.
 - $I'_3 \cup X$ has a partition $\mathcal{D}^{(3)} = \{D_1^{(3)}, D_2^{(3)}, \dots, D_q^{(3)}\}$ with $q \leq k$ such that for all $i = 1, \dots, q$, $|D_i^{(3)}| = 3$ and $G[D_i^{(3)}]$ is connected.

We set $A[X, F, Y, j] = \infty$ if no $I' \subseteq \{v_1, v_2, \dots, v_j\}$ satisfies the conditions. We can easily compute $A[X, F, Y, 0] \in \{0, \infty\}$ as $A[X, F, Y, 0] = 0$ if and only if $F = Y = \emptyset$ and $G[X]$ admits a partition into connected graphs of 3-vertices. The last condition can be checked in $O(2^{|J|} \cdot |J|^3)$ time for all $X \subseteq J$ by recursively considering all possible ways for removing three vertices from X ; that is, $A[X, \emptyset, \emptyset, 0] = \min_{\{x, y, z\} \in \binom{X}{3}} A[X \setminus \{x, y, z\}, \emptyset, \emptyset, 0]$ if $|X| \geq 3$. The following recurrence formula holds: $A[X, F, Y, j + 1] = \min\{f_1, f_2, f_3, f_4\}$, where

$$\begin{aligned} f_1 &= A[X, F, Y, j], \\ f_2 &= \min_{\alpha, \beta \in X, |E(G[\{\alpha, \beta, v_{j+1}\}])| \geq 2} A[X \setminus \{\alpha, \beta\}, F, Y \setminus \{N(\{\alpha, \beta, v_{j+1}\})\}, j] + 1, \\ f_3 &= \min_{\alpha \in X \cap N(v_{j+1})} A[X \setminus \{\alpha\}, F \cup \{\alpha\}, Y \setminus N(v_{j+1}), j] + 1, \\ f_4 &= \min_{\beta \in F \cap N(v_{j+1})} A[X, F \setminus \{\beta\}, Y \setminus N(\{\beta, v_{j+1}\}), j] + 1. \end{aligned}$$

The four options f_1, f_2, f_3 , and f_4 assume different ways of the role of v_{j+1} and compute the optimal value under the assumptions (see Fig. 4). Concretely, f_1 reflects the case when v_{j+1} does not belong to the solution, and f_2 reflects the case when v_{j+1} belongs to the solution together with two vertices in J in a connected way. In f_3 , it reflects that v_{j+1} forms a triple in the solution with a vertex in F and a vertex in I_j . In f_4 , it reflects that v_{j+1} currently forms a pair in J and will form a triple with a vertex in $I \setminus I_j$. We can compute all entries of A in $O^*(4^{|J|})$ time as the number of combinations of three disjoint sets X, F, Y of J is $4^{|J|}$.

Similarly to the previous case of $r = 2$, we can compute the minimum number of units in a 3-grouped dominating set as $\min\{(|X| + A[X, \emptyset, J \setminus N[X], |I|])/3 \mid$




Fig. 4. How v_{j+1} is used. (The white vertices belong to a dominating set.)

$X \subseteq J$ and $I \subseteq N(X)$. Given the table A , this can be done in $O^*(2^{|J|})$ time. Thus the total running time of the algorithm is $O^*(4^{|J|})$.

Algorithm for r -Grouped Dominating Set We now present our algorithm for general $r \geq 4$. Let $G = (V, E)$ be a graph, J be a vertex cover of G , and $I = V \setminus J$. This case still allows an algorithm based on a similar framework to the previous cases, though connected components of general r can be built up from smaller fragments of connected components; this yields an essential difference that worsens the running time. In the DP, the vertex cover J is partitioned into $r + 1$ parts depending on the partial solution corresponding to each table entry, and then some of the parts in the partition are further partitioned into smaller subsets. In other words, each table entry corresponds to a nested partition of the vertex cover.

As in the previous algorithms, for each subset $J_D \subseteq J$, we find a subset $I_D \subseteq I$ (if any exists) of the minimum size such that $J_D \cup I_D$ can form an r -grouped dominating set. Let $X, F^{(r-1)}, \dots, F^{(3)}, F^{(2)}, Y$ be disjoint subsets of J , and let $I = \{v_1, v_2, \dots, v_{|I|}\}$. For $i = 2, \dots, r - 1$, let $\mathcal{F}^{(i)}$ be a partition of $F^{(i)}$, where $\mathcal{F}^{(i)} = \{F_1^{(i)}, F_2^{(i)}, \dots, F_{|\mathcal{F}^{(i)}|}^{(i)}\}$. The number of such nested partitions $(X, \mathcal{F}^{(r-1)}, \dots, \mathcal{F}^{(2)}, Y)$ is at most $(r + 1)^{|J|} |J|^{|J|}$. For $j = 0, \dots, |I|$, we define $A[X, \mathcal{F}^{(r-1)}, \dots, \mathcal{F}^{(2)}, Y, j]$ as the minimum size of $I' \subseteq \{v_1, v_2, \dots, v_j\}$ that satisfies the following conditions:

1. $Y \subseteq N(I')$,
2. I' can be partitioned into $r - 1$ parts I'_2, I'_3, \dots, I'_r satisfying the following conditions:
 - for $i = 2, \dots, r - 1$, $I'_i \cup F^{(i)}$ has a partition $\mathcal{D}^{(i)} = \{D_1^{(i)}, D_2^{(i)}, \dots, D_{|\mathcal{F}^{(i)}|}^{(i)}\}$ such that for all $p = 1, \dots, |\mathcal{F}^{(i)}|$, $D_p^{(i)}$ includes at least one vertex of I' and is a superset of $F_p^{(i)}$, and $|D_p^{(i)}| = i$ and $G[D_p^{(i)}]$ is connected.




Fig. 5. A nested partition of a vertex cover.

- $I'_r \cup X$ has a partition $\mathcal{D}^{(r)} = \{D_1^{(r)}, D_2^{(r)}, \dots, D_q^{(r)}\}$ such that for all $i = 1, \dots, q$, $|D_i^{(r)}| = r$ and $G[D_i^{(r)}]$ is connected.

We set $A[X, \mathcal{F}^{(r-1)}, \dots, \mathcal{F}^{(2)}, Y, j] = \infty$ if no $I' \subseteq \{v_1, v_2, \dots, v_j\}$ satisfies the conditions. We can compute $A[X, \mathcal{F}^{(r-1)}, \dots, \mathcal{F}^{(2)}, Y, 0]$, which is 0 or ∞ , as it is 0 if and only if $F^{(r-1)} = \dots = F^{(2)} = Y = \emptyset$ and $G[X]$ admits a partition into connected graphs of r vertices. The last condition can be checked in $O(|J|^{|J|})$ time for all $X \subseteq J$ by checking all possible partitions of J .

Assume that all entries of A with $j \leq c$ for some c are computed. Since the degree of v_{c+1} is at most $|J|$, the number of possible ways of how v_{c+1} extends a partial solution is at most $2^{|J|}$. Thus from each table entry of A with $j = c$, we obtain at most $2^{|J|}$ candidates of the table entries with $j = c + 1$. Thus, we can compute all entries of A in $O^*(2^{|J|}(r+1)^{|J|}|J|^{|J|})$ time.

Given A , we can compute the minimum number of units in an r -grouped dominating set as $\min\{(|X| + A[X, \emptyset, \dots, \emptyset, J \setminus N[X], |I|])/r \mid X \subseteq J \text{ and } I \subseteq N(X)\}$. Again this takes only $O^*(2^{|J|})$ time. Thus the total running time of the algorithm is $O^*(2^{|J|}(r+1)^{|J|}|J|^{|J|}) = O^*((2\nu(r+1))^\nu)$.

4.2 Algorithms parameterized by twin cover number

In this subsection, we show that the algorithms presented above still work when the parameter ν in the running time is replaced with twin cover number τ . To show this, we prove the following lemma. It says that twin-edges do not contribute to the connectivity of units for some minimum r -grouped dominating sets and can be removed from the graph. As a result, the vertex cover number can be replaced with the twin cover number.

Lemma 4.3. *Let G be a graph and K be a twin cover of G . If G has an r -grouped dominating set, then there exists a minimum r -grouped dominating set such that every unit has at least one vertex in K .*

Proof. Let $G = (V, E)$ be a graph, and K be a twin cover of G . Suppose that a minimum r -grouped dominating set \mathcal{D} exists and one of its units $D =$

$\{v_1, v_2, \dots, v_r\}$ has no vertex in K . Since K is a twin cover, $N[v_1] = N[v_2] = \dots = N[v_r]$ holds. Let $K_D = K \cap N(v_1)$. Then, there is at least one vertex x in K_D such that $x \notin \bigcup \mathcal{D}$. Suppose to the contrary that there is no such x , and thus all vertices in K_D belong to $\bigcup \mathcal{D}$. This implies that no vertex is dominated only by D and that D itself is dominated by some vertices in K_D . Thus, $\mathcal{D} \setminus \{D\}$ is an r -grouped dominating set. This contradicts the minimality of \mathcal{D} . Let $D' = D \setminus \{v_1\} \cup \{x\}$, then $\mathcal{D}' = \mathcal{D} \setminus \{D\} \cup \{D'\}$ is also a minimum r -grouped dominating set of G (see Fig. 6). By repeating this process, we can obtain a minimum r -grouped dominating set such that every unit has at least one vertex in K . \square




Fig. 6. An example for exchange. White vertices belong to a dominating set.

5 Beyond Vertex Cover and Twin Cover

In this section, we further explore the parameterized complexity of r -GROUPED DOMINATING SET with respect to structural graph parameters that generalize vertex cover number and twin cover number. We show that if we do not try to optimize the running time of algorithms, then we can use known algorithmic meta-theorems that automatically give fixed-parameter algorithms parameterized by certain graph parameters.

For the sake of brevity, we define only the parameters for which we need their definitions. For example, we do not need the definition of treewidth for applying the meta-theorem described below. On the other hand, to contrast the results here with the ones in the previous sections, it is important to see the picture of the relationships between the parameters. See Fig. 1 for the hierarchy of the graph parameters we deal with.

Roughly speaking, the algorithmic meta-theorems we use here say that if a problem can be expressed in a certain logic (e.g., FO, MSO₁, or MSO₂), then the problem is fixed-parameter tractable parameterized by a certain graph parameter (e.g., twin-width, treewidth, or clique-width). Such theorems are extremely powerful and used widely for designing fixed-parameter algorithms [27]. On the other hand, the generality of the meta-theorems unfortunately comes with very

high dependency on the parameters [19]. When our target parameter is vertex cover number, the situation is slightly better, but still a double-exponential $2^{2^{\Omega(\nu)}}$ lower bound of the parameter dependency is known under ETH [28]. This implies that our “slightly superexponential” $2^{O(\tau \log \tau)}$ algorithm in Section 4 cannot be obtained by applications of known meta-theorems.

In the rest of this section, we first introduce FO, MSO₁, and MSO₂ on graphs. We then observe that the problem can be expressed in FO when r and k are part of the parameter and in MSO₂ when r is part of the parameter. These observations combined with known meta-theorems immediately imply that r -GROUPED DOMINATING SET is fixed-parameter tractable when

- parameterized by $r + k$ on nowhere dense graph classes;
- parameterized by $r + k$ + twin-width if a contraction sequence of the minimum width is given as part of the input; and
- parameterized by $r + \text{treewidth}$.

We then consider the parameter $k + \text{treewidth}$ and show that this case is intractable. More strongly, we show that r -GROUPED DOMINATING SET is W[1]-hard when the parameter is $k + \text{treedepth} + \text{feedback vertex set number}$.

We finally consider the parameter modular-width, a generalization of twin cover number, and show that r -GROUPED DOMINATING SET parameterized by modular-width is fixed-parameter tractable.

5.1 Results based on algorithmic meta-theorems

The *first-order logic* on graphs (FO) allows variables representing vertices of the graph under consideration. The atomic formulas are the equality $x = y$ of variables and the adjacency $E(x, y)$ meaning that $\{x, y\} \in E$. The FO formulas are defined recursively from atomic formulas with the usual Boolean connectives ($\neg, \wedge, \vee, \Rightarrow, \Leftrightarrow$), and quantification of variables (\forall, \exists). We also use the existential quantifier with a dot ($\dot{\exists}$) to quantify distinct objects. For example, $\dot{\exists}a, b: \phi$ means $\exists a, b: (a \neq b) \wedge \phi$. We write $G \models \phi$ if G satisfies (or *models*) ϕ . Given a graph G and an FO formula ϕ , FO MODEL CHECKING asks whether $G \models \phi$.

It is straightforward to express the property of having an r -grouped dominating set of k units with an FO formula whose length depends only on $r + k$:

$$\phi_{r,k} = \dot{\exists}v_1, v_2, \dots, v_{rk} : \text{dominating}(v_1, \dots, v_{rk}) \wedge \bigwedge_{0 \leq i \leq k-1} \text{connected}(v_{ir+1}, \dots, v_{ir+r}),$$

where $\text{dominating}(\dots)$ is a subformula expressing that the rk vertices form a dominating set and $\text{connected}(\dots)$ is the one expressing that the r vertices induce a connected subgraph (see Section 5.4 for the expressions of the subformulas). This implies that r -GROUPED DOMINATING SET parameterized by $r + k$ is fixed-parameter tractable on graph classes on which FO MODEL CHECKING parameterized by the formula length $|\phi|$ is fixed-parameter tractable. Such graph classes include nowhere dense graph classes [23] and graphs of bounded twin-width (given with so called contraction sequences) [5].

Corollary 5.1. *r -GROUPED DOMINATING SET parameterized by $r+k$ is fixed-parameter tractable on nowhere dense graph classes.*

Corollary 5.2. *r -GROUPED DOMINATING SET parameterized by $r+k+\text{twin-width}$ is fixed-parameter tractable if a contraction sequence of the minimum width is given as part of the input.*

The *monadic second-order logic* on graphs (MSO_1) is an extension of FO that additionally allows variables representing vertex sets and the inclusion predicate $X(x)$ meaning that $x \in X$. MSO_2 is a further extension of MSO_1 that also allows edge variables, edge-set variables, and an atomic formula $I(e, x)$ representing the edge-vertex incidence relation. Given a graph G and an MSO_1 (MSO_2 , resp.) formula $\phi(X)$ with a free set variable X , MSO_1 (MSO_2 , resp.) OPTIMIZATION asks to find a minimum set S such that $G \models \phi(S)$.

It is not difficult to express the property of a vertex set being the union of r -units of a r -grouped dominating set with an MSO_2 formula whose length depending only on r :⁴

$$\begin{aligned} \psi_r(X) = & \text{dominating}(X) \wedge \\ & (\exists F \subseteq E: \text{span}(F, X) \wedge (\forall C \subseteq X: \text{cc}(F, C) \Rightarrow \text{size}_r(C))), \end{aligned}$$

where $\text{dominating}(X)$ is a subformula expressing that X is a dominating set, $\text{span}(F, X)$ is the one expressing that X is the set of all endpoints of the edges in F , $\text{cc}(F, C)$ expresses that C is the vertex set of a connected component of the subgraph induced by F , and $\text{size}_r(C)$ means that C contains exactly r elements (again, see Section 5.4 for the expressions of the subformulas). Since MSO_2 OPTIMIZATION parameterized by treewidth is fixed-parameter tractable [1,6,11], we have the following result.

Corollary 5.3. *r -GROUPED DOMINATING SET parameterized by $r + \text{treewidth}$ is fixed-parameter tractable.*

5.2 Hardness parameterized by $k + \text{treewidth}$

Now the natural question regarding treewidth and r -GROUPED DOMINATING SET would be the complexity parameterized by $k + \text{treewidth}$. Unfortunately, this case is $W[1]$ -hard even if treewidth is replaced with a possibly much larger parameter $\text{pathwidth} + \text{feedback vertex set number}$ and the graphs are restricted to be planar. Furthermore, if the planarity is not required, we can replace pathwidth in the parameter with treedepth .

Theorem 5.4. *r -GROUPED DOMINATING SET parameterized by $k + \text{pathwidth} + \text{feedback vertex set number}$ is $W[1]$ -hard on planar graphs.*

⁴ Note that there is no equivalent MSO_1 formula of length depending only on r . This is because $G \models \psi_2(V)$ expresses the property of having a perfect matching, for which an MSO_1 formula does not exist (see e.g., [12]).

Proof. Given a graph $G = (V, E)$ and an integer $r \geq 2$, EQUITABLE CONNECTED PARTITION asks whether there exists a partition of V into $k = |V|/r$ sets V_1, \dots, V_k such that $G[V_i]$ is connected and $|V_i| = r$ for $1 \leq i \leq k$. It is known that EQUITABLE CONNECTED PARTITION parameterized by $k + \text{pathwidth} + \text{feedback vertex set number}$ is W[1]-hard even on planar graphs [18]. We reduce this problem to ours.

Let $\langle G = (V, E), r \rangle$ be an instance of EQUITABLE CONNECTED PARTITION. To each vertex v of G , we attach a new vertex of degree 1, which we call a *pendant* at v . This modification does not change the feedback vertex number and may increase the pathwidth by at most 1 (see e.g., [2, Lemma A.2]). Let H be the resultant graph, which is planar. To prove the lemma, it suffices to show that $\langle H, k \rangle$ is a yes-instance of r -GROUPED DOMINATING SET if and only if $\langle G, r \rangle$ is a yes-instance of EQUITABLE CONNECTED PARTITION.

To prove the if direction, assume that $\langle G, r \rangle$ is a yes-instance of EQUITABLE CONNECTED PARTITION and that V_1, \dots, V_k certificate it. Clearly, $\{V_1, \dots, V_k\}$ is an r -grouped dominating set of H .

To prove the only-if direction, assume that H has an r -grouped dominating set \mathcal{D} with at most k units. Let $v \in V$ and p be the pendant at v . Observe that $\bigcup \mathcal{D}$ contains exactly one of v and p since it needs at least one of them for dominating p and $|\bigcup \mathcal{D}| \leq rk = |V|$. Furthermore, the assumption $r \geq 2$ implies that $\bigcup \mathcal{D}$ cannot contain p as it has no neighbor other than v . This implies that $\bigcup \mathcal{D} = V$ and that \mathcal{D} contains exactly $|V|/r = k$ units. Therefore, the family \mathcal{D} is a certificate that $\langle G, r \rangle$ is a yes-instance of EQUITABLE CONNECTED PARTITION. \square

It is known that on general (not necessarily planar) graphs, EQUITABLE CONNECTED PARTITION parameterized by $k + \text{treedepth} + \text{feedback vertex set number}$ is W[1]-hard [22]. Since adding pendants to all vertices increases treedepth by at most 1 (see e.g., [32]), the same reduction shows the following hardness.

Theorem 5.5. *r -GROUPED DOMINATING SET parameterized by $k + \text{treedepth} + \text{feedback vertex set number}$ is W[1]-hard.*

5.3 Fixed-parameter tractability parameterized by modular-width

Let $G = (V, E)$ be a graph. A set $M \subseteq V$ is a *module* if for each $v \in V \setminus M$, either $M \subseteq N(v)$ or $M \cap N(v) = \emptyset$ holds. The *modular-width* of G , denoted $\text{mw}(G)$, is the minimum integer k such that either $|V| \leq k$ or there exists a partition of V into at most k modules M_1, \dots, M_k of G such that each $G[M_i]$ has modular-width at most k . It is known that the modular-width of a graph and a recursive partition certifying it can be computed in linear time [14,24,35].

Observe that if V is partitioned into modules M_1, \dots, M_k of G , then for two distinct modules M_i and M_j , we have either no or all possible edges between them. If there are all possible edges between M_i and M_j , then we say that M_i and M_j are adjacent.

Theorem 5.6. *r -GROUPED DOMINATING SET parameterized by modular-width is fixed-parameter tractable.*

Proof. Let $\langle G = (V, E), k \rangle$ be an instance of r -GROUPED DOMINATING SET. We only consider the case of $r \geq 2$ since the other case of $r = 1$ is known (see [13,20]). We may assume that G is connected since otherwise we can solve the problem on each connected component separately. We also assume that G has at least r vertices as otherwise the problem is trivial. Let M_1, \dots, M_μ be a partition of V into modules with $2 \leq \mu \leq \text{mw}(G)$. For each module M_i , there is at least one adjacent module M_j as G is connected.

We first assume that $r \geq \mu$. Let $D \subseteq V$ be an arbitrary set of size r that takes at least one vertex from each module M_i . Recall that we have either no or all possible edges between two distinct modules. Thus, the connectivity of G implies that $G[D]$ is connected and D is a dominating set of G . This implies that $\{D\}$ is an r -grouped dominating set with one unit.

Next assume that $r < \mu$. In this case, we show below that if G has an r -grouped dominating set, then G has an r -grouped dominating set with at most μ units. This implies that $r + k < 2\mu$, and thus the problem can be solved as FO MODEL CHECKING with a formula of length depending only on μ , which is fixed-parameter tractable parameterized by μ (see [13,20]).

Before proving the upper bound of k , we show that if G has an r -grouped dominating set, then there is a minimum one such that no unit is entirely contained in a module M_i . Assume that \mathcal{D} is a minimum r -grouped dominating set of G . If $D \subseteq M_i$ holds for some i and $D \in \mathcal{D}$, then there is a vertex v in a module M_j adjacent to M_i that does not belong to $\bigcup \mathcal{D}$. This is because, otherwise, $\mathcal{D} \setminus \{D\}$ is still an r -grouped dominating set. Let u be an arbitrary vertex in D and set $D' = D \setminus \{u\} \cup \{v\}$. As $r \geq 2$, D' intersects both M_i and M_j . Also we can see that $|D'| = r$, D' is connected (as u is adjacent to all vertices in M_i), and all vertices dominated by D are dominated by D' as well. Thus, $\mathcal{D} \setminus \{D\} \cup \{D'\}$ is an r -grouped dominating set. We can repeat this process until we have the claimed property.

As discussed above, it suffices to show the upper bound k for the number of units. Let \mathcal{D} be a minimum r -grouped dominating set of G such that no unit is entirely contained in a module M_i . We say that a module M_i is *private* for a unit $D \in \mathcal{D}$ if D is the only one in \mathcal{D} that intersects M_i . Suppose to the contrary that $|\mathcal{D}| > \mu$. Then, there is $D \in \mathcal{D}$ such that no module M_i is private for D . If a module M_i is adjacent to a module M_j that intersects D , then since M_j is not private for D , $\mathcal{D} \setminus \{D\}$ contains a unit intersecting M_j , which dominates M_i . If a module M_i intersects D , then since D intersects at least two modules and $G[D]$ is connected, there is a module M_j adjacent to M_i and intersecting D . Hence, as the previous case, $\mathcal{D} \setminus \{D\}$ contains a unit dominating M_i . Therefore, we can conclude that $\mathcal{D} \setminus \{D\}$ is an r -grouped dominating set. This contradicts the minimality of \mathcal{D} . \square

5.4 Auxiliary subformulas

Here we present FO or MSO₂ expressions of some of the subformulas in Section 5.1. All of them are standard and presented only to show basic ideas.

The following formulas expressing dominating sets are almost direct translation of the definition and should be easy to read.

$$\begin{aligned} \text{dominating}(X) &= \forall u \exists v: X(v) \wedge ((u = v) \vee E(u, v)). \\ \text{dominating}(v_1, \dots, v_p) &= \forall u: (u = v_1) \vee (u = v_2) \vee \dots \vee (u = v_p) \\ &\quad \vee E(u, v_1) \vee E(u, v_2) \vee \dots \vee E(u, v_p). \end{aligned}$$

The connectivity of $G[X]$ is a little bit tricky to state. We state that for each nonempty proper subset Y of X , there is an edge between Y and $X \setminus Y$. See e.g., [15] for the full expression of `connected`. The FO version of `connected` can be expressed based on the same idea but the length of the formula depends on the number r of vertices it can take (which is fine for us as r is part of the input when we use this formula). In [15], an MSO₂ formula expressing the connectivity of the graph induced by an edge set is also presented. We call it `connectedE` and use it below.

Recall that `span`(F, X) expresses that X is the set of all endpoints of the edges in F and that `cc`(F, C) expresses that C is the vertex set of a connected component of the subgraph induced by F . They can be stated as follows:

$$\begin{aligned} \text{span}(F, X) &= \forall v: X(v) \Leftrightarrow (\exists e: F(e) \wedge I(e, v)), \\ \text{cc}(F, C) &= \exists F': (F' \subseteq F) \wedge \text{span}(F', C) \wedge \text{connectedE}(F') \\ &\quad \wedge (\forall F'': (F' \subseteq F'' \subseteq F) \wedge \neg \text{connectedE}(F'')), \end{aligned}$$

where the inclusion relation $F \subseteq F'$ can be stated as $\forall e: F(e) \Rightarrow F'(e)$.

Finally, when r is part of the parameter, `sizer`(C) meaning that $|C| = r$ can be stated as follows:

$$\text{size}_r(C) = \exists v_1, \dots, v_r: \bigwedge_{1 \leq i \leq r} C(v_i) \wedge \left(\neg \exists v: C(v) \wedge \bigwedge_{1 \leq i \leq r} v \neq v_i \right).$$

References

1. Arnborg, S., Lagergren, J., Seese, D.: Easy problems for tree-decomposable graphs. *J. Algorithms* **12**(2), 308–340 (1991). [https://doi.org/10.1016/0196-6774\(91\)90006-K](https://doi.org/10.1016/0196-6774(91)90006-K)
2. Belmonte, R., Hanaka, T., Kanzaki, M., Kiyomi, M., Kobayashi, Y., Kobayashi, Y., Lampis, M., Ono, H., Otachi, Y.: Parameterized complexity of (A, ℓ) -path packing. *Algorithmica* **84**(4), 871–895 (2022). <https://doi.org/10.1007/s00453-021-00875-y>
3. Bertossi, A.A.: Dominating sets for split and bipartite graphs. *Information Processing Letters* **19**(1), 37–40 (1984). [https://doi.org/10.1016/0020-0190\(84\)90126-1](https://doi.org/10.1016/0020-0190(84)90126-1)
4. Bodlaender, H.L., Cygan, M., Kratsch, S., Nederlof, J.: Deterministic single exponential time algorithms for connectivity problems parameterized by treewidth. *Information and Computation* **243**, 86–111 (2015)
5. Bonnet, É., Kim, E.J., Thomassé, S., Watrigant, R.: Twin-width I: tractable FO model checking. *J. ACM* **69**(1), 3:1–3:46 (2022). <https://doi.org/10.1145/3486655>

6. Borie, R.B., Parker, R.G., Tovey, C.A.: Automatic generation of linear-time algorithms from predicate calculus descriptions of problems on recursively constructed graph families. *Algorithmica* **7**(5&6), 555–581 (1992). <https://doi.org/10.1007/BF01758777>
7. Chen, J., Kanj, I.A., Xia, G.: Improved upper bounds for vertex cover. *Theor. Comput. Sci.* **411**(40-42), 3736–3756 (2010). <https://doi.org/10.1016/j.tcs.2010.06.026>
8. Chen, L., Lu, C., Zeng, Z.: Hardness results and approximation algorithms for (weighted) paired-domination in graphs. *Theoretical Computer Science* **410**(47), 5063–5071 (2009). <https://doi.org/10.1016/j.tcs.2009.08.004>
9. Chen, L., Lu, C., Zeng, Z.: A linear-time algorithm for paired-domination problem in strongly chordal graphs. *Information Processing Letters* **110**(1), 20–23 (2009). <https://doi.org/10.1016/j.ipl.2009.09.014>
10. Chen, L., Lu, C., Zeng, Z.: Labelling algorithms for paired-domination problems in block and interval graphs. *Journal of Combinatorial Optimization* **19**(4), 457–470 (2010). <https://doi.org/10.1007/s10878-008-9177-6>
11. Courcelle, B.: The monadic second-order logic of graphs. I. recognizable sets of finite graphs. *Inf. Comput.* **85**(1), 12–75 (1990). [https://doi.org/10.1016/0890-5401\(90\)90043-H](https://doi.org/10.1016/0890-5401(90)90043-H)
12. Courcelle, B., Engelfriet, J.: *Graph Structure and Monadic Second-Order Logic - A Language-Theoretic Approach*. Cambridge University Press (2012), <https://www.cambridge.org/knowledge/isbn/item5758776/>
13. Courcelle, B., Makowsky, J.A., Rotics, U.: Linear time solvable optimization problems on graphs of bounded clique-width. *Theory Comput. Syst.* **33**(2), 125–150 (2000). <https://doi.org/10.1007/s002249910009>
14. Cournier, A., Habib, M.: A new linear algorithm for modular decomposition. In: *CAAP 1994. Lecture Notes in Computer Science*, vol. 787, pp. 68–84 (1994). <https://doi.org/10.1007/BFb0017474>
15. Cygan, M., Fomin, F.V., Kowalik, L., Lokshantov, D., Marx, D., Pilipczuk, M., Pilipczuk, M., Saurabh, S.: *Parameterized Algorithms*. Springer (2015). <https://doi.org/10.1007/978-3-319-21275-3>
16. Cygan, M., Nederlof, J., Pilipczuk, M., Pilipczuk, M., Van Rooij, J.M.M., Wojtaszczyk, J.O.: Solving connectivity problems parameterized by treewidth in single exponential time. *ACM Trans. Algorithms* **18**(2) (2022). <https://doi.org/10.1145/3506707>
17. Desormeaux, W.J., Haynes, T.W., Henning, M.A.: Paired domination in graphs. In: Haynes, T.W., Hedetniemi, S.T., Henning, M.A. (eds.) *Topics in Domination in Graphs*, pp. 31–77. Springer International Publishing, Cham (2020). https://doi.org/10.1007/978-3-030-51117-3_3
18. Enciso, R., Fellows, M.R., Guo, J., Kanj, I.A., Rosamond, F.A., Suchý, O.: What makes equitable connected partition easy. In: *IWPEC 2009. Lecture Notes in Computer Science*, vol. 5917, pp. 122–133 (2009). https://doi.org/10.1007/978-3-642-11269-0_10
19. Frick, M., Grohe, M.: The complexity of first-order and monadic second-order logic revisited. *Ann. Pure Appl. Log.* **130**(1-3), 3–31 (2004). <https://doi.org/10.1016/j.apal.2004.01.007>
20. Gajarský, J., Lampis, M., Ordyniak, S.: Parameterized algorithms for modular-width. In: *IPEC 2013. Lecture Notes in Computer Science*, vol. 8246, pp. 163–176 (2013). https://doi.org/10.1007/978-3-319-03898-8_15
21. Ganian, R.: Improving vertex cover as a graph parameter. *Discret. Math. Theor. Comput. Sci.* **17**(2), 77–100 (2015). <https://doi.org/10.46298/dmtcs.2136>

22. Gima, T., Otachi, Y.: Extended MSO model checking via small vertex integrity. *CoRR* **abs/2202.08445** (2022)
23. Grohe, M., Kreutzer, S., Siebertz, S.: Deciding first-order properties of nowhere dense graphs. *J. ACM* **64**(3), 17:1–17:32 (2017). <https://doi.org/10.1145/3051095>
24. Habib, M., Paul, C.: A survey of the algorithmic aspects of modular decomposition. *Comput. Sci. Rev.* **4**(1), 41–59 (2010). <https://doi.org/10.1016/j.cosrev.2010.01.001>
25. Haynes, T.W., Slater, P.J.: Paired-domination and the paired-domatic number. *Congressus Numerantium* pp. 65–72 (1995)
26. Haynes, T.W., Slater, P.J.: Paired-domination in graphs. *Networks* **32**(3), 199–206 (1998). [https://doi.org/10.1002/\(SICI\)1097-0037\(199810\)32:3;199::AID-NET4;3.0.CO;2-F](https://doi.org/10.1002/(SICI)1097-0037(199810)32:3;199::AID-NET4;3.0.CO;2-F)
27. Kreutzer, S.: Algorithmic meta-theorems. In: Esparza, J., Michaux, C., Steinhorn, C. (eds.) *Finite and Algorithmic Model Theory*, London Mathematical Society Lecture Note Series, vol. 379, pp. 177–270. Cambridge University Press (2011)
28. Lampis, M.: Algorithmic meta-theorems for restrictions of treewidth. *Algorithmica* **64**(1), 19–37 (2012). <https://doi.org/10.1007/s00453-011-9554-x>
29. Liedloff, M.: Finding a dominating set on bipartite graphs. *Information Processing Letter* **107**(5), 154–157 (2008). <https://doi.org/10.1016/j.ipl.2008.02.009>
30. Lin, C.C., Ku, K.C., Hsu, C.H.: Paired-domination problem on distance-hereditary graphs. *Algorithmica* **82**(10), 2809–2840 (2020). <https://doi.org/10.1007/s00453-020-00705-7>
31. Middendorf, M., Pfeiffer, F.: On the complexity of the disjoint paths problems. *Combinatorica* **13**(1), 97–107 (1993). <https://doi.org/10.1007/BF01202792>
32. Nešetřil, J., Ossona de Mendez, P.: *Sparsity: Graphs, Structures, and Algorithms*. Algorithms and combinatorics, Springer (2012). <https://doi.org/10.1007/978-3-642-27875-4>
33. Pradhan, D., Panda, B.: Computing a minimum paired-dominating set in strongly orderable graphs. *Discrete Applied Mathematics* **253**, 37–50 (2019). <https://doi.org/10.1016/j.dam.2018.08.022>
34. Raman, V., Saurabh, S.: Short cycles make W -hard problems hard: FPT algorithms for W -hard problems in graphs with no short cycles. *Algorithmica* **52**(2), 203–225 (2008). <https://doi.org/10.1007/s00453-007-9148-9>, <https://doi.org/10.1007/s00453-007-9148-9>
35. Tedder, M., Corneil, D.G., Habib, M., Paul, C.: Simpler linear-time modular decomposition via recursive factorizing permutations. In: *ICALP 2008*. Lecture Notes in Computer Science, vol. 5125, pp. 634–645 (2008). https://doi.org/10.1007/978-3-540-70575-8_52
36. Tripathi, V., Kloks, T., Pandey, A., Paul, K., Wang, H.: Complexity of paired domination in AT-free and planar graphs. In: *CALDAM 2022*. Lecture Notes in Computer Science, vol. 13179, pp. 65–77 (2022). https://doi.org/10.1007/978-3-030-95018-7_6