

Towards a universal representation of statistical dependence

GERY GEENENS*

School of Mathematics and Statistics,

UNSW Sydney, Australia

December 25, 2023

Abstract

Dependence is undoubtedly a central concept in statistics. Though, it proves difficult to locate in the literature a formal definition which goes beyond the self-evident ‘dependence = non-independence’. This absence has allowed the term ‘dependence’ and its declination to be used vaguely and indiscriminately for qualifying a variety of disparate notions, leading to numerous incongruities. For example, the classical Pearson’s, Spearman’s or Kendall’s correlations are widely regarded as ‘dependence measures’ of major interest, in spite of returning 0 in some cases of deterministic relationships between the variables at play – evidently not measuring dependence at all. Arguing that research on such a fundamental topic would benefit from a slightly more rigid framework, this paper suggests a general definition of the dependence between two random variables defined on the same probability space. Natural enough for aligning with intuition, that definition is still sufficiently precise for allowing unequivocal identification of a ‘universal’ representation of the dependence structure of any bivariate distribution. Links between this representation and familiar concepts are highlighted, and ultimately, the idea of a dependence measure based on that universal representation is explored and shown to satisfy Rényi’s postulates.

1 Introduction

Scientific progress relies on identifying if, how, why, and to which extent, several factors influence each other. Across all fields, studies looking for establishing an effect of an input on an output must statistically demonstrate evidence of association / correlation / relationship / causality between input and output; i.e., to establish some sort of dependence between the two. Hence statistics, the art of turning empirical evidence (data) into information, has always kept the concept of dependence at its core.

*Corresponding author: ggeenens@unsw.edu.au, School of Mathematics and Statistics, UNSW Sydney, Australia, tel +61 2 938 57032, fax +61 2 9385 7123

In spite of this, it may prove difficult to delineate ‘dependence’ precisely. It is noteworthy that, notwithstanding the frequent use of the term, it is rarely defined formally in the literature – not even in major references entirely dedicated to the topic such as Drouet Mari and Kotz (2001) or Joe (2015). Neither the ‘Cambridge Dictionary of Statistics’ (Everitt and Skrondal, 2010) nor the ‘Dictionary of Statistics’ (Upton and Cook, 2014) include an entry for dependence. The ‘Dictionary of Statistical Terms’ (Kendall and Buckland, 1971, Dodge, 2006) does, but gives the seemingly circular ‘*Dependence: quantities are dependent when they are not independent*’.

Admittedly, *independence* has an unambiguous mathematical meaning, as when the joint distribution of two variables of interest can be suitably factorised; see (3.4) below. Writing this in terms of conditional distributions allows an interpretation along the line of ‘variation in one variable does not disturb the stochastic behaviour of the other’. Thus, according to the above (pseudo-)definition, two variables are dependent as soon as there is such (loosely defined, not necessarily causal) influence of one on the other, and dependence amounts to the *existence* of the said influence. In effect such ‘*non-independence*’ interpretation of dependence makes it a binary concept – two variables do or do not satisfy the property of independence, so they are dependent or not without any room for nuance. If such a binary-in-nature view may be suitable for the *independence testing* problem, it quickly falls short when it comes to other questions. In particular, quantifying dependence appears essential in many situations, as testified by the abundance of dependence measures found in the literature – see Tjøstheim et al. (2022) for a recent review. Though, measuring a trait is incompatible with an alleged binary nature. If dependence is to be quantifiable, then clearly the non-independence definition appears inadequate – and this, without any obvious substitute.

This absence has permitted some looseness when addressing dependence-related questions, inducing confusion between disparate notions, and causing incongruities. The most obvious of those may be that the most popular ‘dependence measures’, namely Pearson’s correlation (ρ), Spearman’s rho (ρ_S) and Kendall’s tau (τ), do *not* measure dependence at all – for example, they may be 0 between two variables deterministically bound,¹ i.e., the strongest influence that one can imagine. In fact, the elusiveness of a quantifiable version of dependence has allowed for subjective interpretation to resolve what was to be measured, and how. This explains the profusion of diverse, more or less pertinent, ‘dependence measures’ (more on this in Section 6).

Arguing that further progress in an area as important as that of statistical dependence would benefit from a slightly more rigid framework, this paper attempts to provide a clearer understanding of the concept, starting from first principles (Section 2). We then propose a general definition of the dependence between two variables defined on the same probability space, which leads, implication after implication (through

¹See the textbook example $X \sim \mathcal{N}(0, 1)$, $Y = X^2 \Rightarrow \rho(X, Y) = \rho_S(X, Y) = \tau(X, Y) = 0$.

Section 3), to a unequivocal representation of the said concept (Section 4). This representation is ‘universal’ in the sense that it is valid for any random vector, regardless of its nature (discrete, continuous, mixed or hybrid) – we only require a joint distribution dominated by a reference product measure. Section 5 illustrates the distinction between dependence and other related notions by examining closely the important concept of concordance. Section 6 suggests a general measure of dependence articulated around the obtained universal representation, and Section 7 concludes.

2 Heuristic

Let (X, Y) be a bivariate random vector with unknown joint distribution π_{XY} . Suppose that an oracle gives us full knowledge of the two univariate marginal distributions π_X and π_Y . Is it enough to reconstruct π_{XY} entirely? The answer is no: this information allows us to conclude that π_{XY} belongs to the Fréchet class $\mathcal{F}(\pi_X, \pi_Y)$ – the set of all bivariate distributions with margins π_X and π_Y – but surely such class contains more than one element. Then, what is the extra piece of information, call it \mathfrak{D} , that the oracle is still concealing, and which would allow unequivocal identification of π_{XY} in $\mathcal{F}(\pi_X, \pi_Y)$?

Specifically, \mathfrak{D} stands for what is necessary and sufficient for identifying π_{XY} once π_X and π_Y are known: ‘necessary’ as it does not include any redundant information already contained in the margins, and ‘sufficient’ as nothing relevant to π_{XY} is possibly left out. This breaks down the bivariate distribution π_{XY} into three constituents showing no overlap: π_X , describing the behaviour of X alone; π_Y , describing the behaviour of Y alone; and \mathfrak{D} , describing ‘the rest’.

The marginal distributions π_X and π_Y may be given various mathematical representations, such as cumulative distribution functions, probability mass/density functions or characteristic functions, the suitability of which depending on the nature of X and Y and the assumptions made a priori about them. E.g., if π_X is supported on a finite set, then exhaustive enumeration of the relevant probabilities (‘probability mass function’) is possible, but not if π_X has uncountable support; while if π_X is assumed to be Gaussian, then the pair (mean, variance) provides an adequate description by itself. We anticipate that the third element \mathfrak{D} is no different, with potentially several mathematical representations differing according to the nature of (X, Y) – this will be formally confirmed in the following sections. For now we may just understand \mathfrak{D} informally as what remains of π_{XY} when entirely stripped from its margins, i.e., the ‘glue’ that keeps π_X and π_Y together in π_{XY} .

Clearly, if X and Y happen to be independent, \mathfrak{D} contains that information – and only that, as it is enough for identifying $\pi_{XY} = \pi_X \times \pi_Y$ in $\mathcal{F}(\pi_X, \pi_Y)$. If X and Y are not independent, then \mathfrak{D} should provide the

complete recipe as to how to interlock π_X and π_Y for giving rise to the specific π_{XY} and making it different to any other distribution of $\mathcal{F}(\pi_X, \pi_Y)$. In all cases, \mathfrak{D} describes what happens ‘between’ π_X and π_Y , which it seems fair to call the dependence structure – or just dependence – of the distribution π_{XY} . Identifying a random vector to its distribution, we may also say that \mathfrak{D} is the dependence of the vector (X, Y) . We, therefore, propose the following definition:

Definition 2.1. *The **dependence** between two random variables $X \sim \pi_X$ and $Y \sim \pi_Y$ defined on the same probability space is the information \mathfrak{D} which is necessary and sufficient to unequivocally identify their joint distribution π_{XY} in its Fréchet class $\mathcal{F}(\pi_X, \pi_Y)$.*

At this stage, this definition is purposely qualitative and very general. Yet, through a sequence of logical implications, it will translate into a precise mathematical representation (Corollary 4.1) – identifying such a representation from the intuitive Definition 2.1 is actually the main goal of this paper. As-is, it suggests the equivalence

$$\pi_{XY} \longleftrightarrow (\pi_X, \pi_Y; \mathfrak{D}), \quad (2.1)$$

where $\pi_{XY} \longrightarrow (\pi_X, \pi_Y; \mathfrak{D})$ means that any distribution π_{XY} admits one and only one decomposition in terms of its three constituents π_X , π_Y and \mathfrak{D} ; while $\pi_{XY} \longleftarrow (\pi_X, \pi_Y; \mathfrak{D})$ means that any triplet $(\pi_X, \pi_Y; \mathfrak{D})$ defines one and only one distribution π_{XY} with the corresponding margins and dependence structure. The (\longrightarrow) -part is true because any distribution in a given Fréchet class must be unequivocally distinguished from other distributions in the same class by some features not perceptible from the marginals, establishing the existence of a third element \mathfrak{D} specific to π_{XY} . In Section 3 below, we prove that the (\longleftarrow) -part holds true, too, up to some natural conditions on the cardinality and shape of the supports of the distributions at play. E.g., as anticipated above, \mathfrak{D} may be of different nature when π_X, π_Y are discrete and when they are continuous, hence a ‘continuous \mathfrak{D} ’ cannot be combined with discrete marginals (or vice-versa). But, dependence structures ‘of the right type’ may be freely interchanged for cementing any given marginals π_X, π_Y and giving rise to various distributions of $\mathcal{F}(\pi_X, \pi_Y)$.

For this to be possible, though, it is necessary that no specification of any of the three constituents alters or imposes restrictions on the others. In a sense, a distribution π_{XY} should be a point represented by a triplet $(\pi_X, \pi_Y, \mathfrak{D})$ of orthogonal coordinates in a some sort of rectangular-prismatic Cartesian space (X -margin, Y -margin, dependence). This is comparable to the concept of variation-independence between parameters (Barndorff-Nielsen, 1978, p. 26): in a parametric model, two parameters θ and θ' are variation-independent if setting the value of one does not impose any new logical constraints on the range of the other. If the decomposition $(\pi_X, \pi_Y; \mathfrak{D})$ is regarded as a parametrisation of π_{XY} , then the above discussion implies that

π_X , π_Y and \mathfrak{D} should be variation-independent. In particular, if any specific feature of π_{XY} happens not to be variation-independent of π_X and π_Y , then it cannot enter any valid description of \mathfrak{D} as such since it would automatically carry redundant information about the margins, in violation of Definition 2.1. Here three simple examples are used to illustrate the point.

Example 2.1. Bivariate Gaussian. Let (X, Y) be a bivariate Gaussian vector. Then π_{XY} is entirely described by its mean vector and its variance-covariance matrix, that is,

$$\boldsymbol{\mu} = \begin{pmatrix} \mu_X \\ \mu_Y \end{pmatrix}, \quad \boldsymbol{\Sigma} = \begin{pmatrix} \sigma_X^2 & \sigma_{XY} \\ \sigma_{XY} & \sigma_Y^2 \end{pmatrix} = \begin{pmatrix} \sigma_X^2 & \rho\sigma_X\sigma_Y \\ \rho\sigma_X\sigma_Y & \sigma_Y^2 \end{pmatrix}. \quad (2.2)$$

Once π_X and π_Y have been specified – i.e., $X \sim \mathcal{N}(\mu_X, \sigma_X^2)$ and $Y \sim \mathcal{N}(\mu_Y, \sigma_Y^2)$ – the only free parameter left in (2.2) is $\rho = \frac{\sigma_{XY}}{\sigma_X\sigma_Y}$ (Pearson’s correlation). As such, ρ alone allows unequivocal identification of π_{XY} among the bivariate Gaussian distributions of $\mathcal{F}(\mathcal{N}(\mu_X, \sigma_X^2), \mathcal{N}(\mu_Y, \sigma_Y^2))$ but does not contain any information about the marginal parameters. Therefore, ρ appropriately captures *the whole dependence structure* of (X, Y) , as per Definition 2.1 – and is not just a quantification of the strength of association. Irrespective of (μ_X, σ_X) and (μ_Y, σ_Y) , for any $\rho \in [-1, 1]$ there exists one and only one bivariate Gaussian distribution with those margins and correlation ρ ; whence (μ_X, σ_X) , (μ_Y, σ_Y) and ρ are indeed variation-independent. Obviously, any one-to-one function of ρ would allow identification of π_{XY} as well: it is the case of $z = \operatorname{atanh} \rho$ (Fisher’s z -transform), Kendall’s $\tau = \frac{2}{\pi} \operatorname{asin} \rho$, and Spearman’s $\rho_S = \frac{6}{\pi} \operatorname{asin} \frac{\rho}{2}$ (Kruskal, 1958), which therefore qualify all the same to represent the dependence here. By contrast, the covariance σ_{XY} does not, as it contains redundant marginal information: σ_X , σ_Y and σ_{XY} are not variation-independent, as $|\sigma_{XY}| \leq \sigma_X\sigma_Y$.

Naturally the parity ‘ $\mathfrak{D} \sim$ Pearson’s correlation ρ ’ follows from the rigid specification of the Gaussian model, but breaks as soon as we leave it. This explains why $\rho = 0 \iff X \perp\!\!\!\perp Y$ (independence) if (X, Y) is Gaussian, but not in general. In fact, the value of ρ does not unequivocally single out a distribution in the whole of $\mathcal{F}(\mathcal{N}(\mu_X, \sigma_X^2), \mathcal{N}(\mu_Y, \sigma_Y^2))$, as this class contains non-Gaussian elements with identical correlation but different dependence structures. E.g., there exist distributions with Gaussian marginals and null correlation, which are not the bivariate Gaussian distribution with independent components. \square

Example 2.2. Bivariate Bernoulli. Let (X, Y) be a bivariate Bernoulli vector; that is, for $p_{1\bullet}, p_{\bullet 1} \in$

$(0, 1)$, $X \sim \text{Bern}(p_{1\bullet})$, $Y \sim \text{Bern}(p_{\bullet 1})$ and the joint distribution π_{XY} is described by a (2×2) -table like

$$\begin{array}{c|cc|c}
 X \setminus Y & 0 & 1 & \\
 \hline
 0 & p_{00} & p_{01} & p_{0\bullet} \\
 1 & p_{10} & p_{11} & p_{1\bullet} \\
 \hline
 & p_{\bullet 0} & p_{\bullet 1} & 1
 \end{array}, \tag{2.3}$$

where² $p_{xy} = \mathbb{P}(X = x, Y = y) > 0$, $p_{x\bullet} = p_{x0} + p_{x1}$ and $p_{\bullet y} = p_{0y} + p_{1y}$ ($x, y \in \{0, 1\}$). As $\sum p_{xy} = 1$, there are initially 3 free parameters for π_{XY} . Once we specify the margins π_X and π_Y , that is, once we fix the two values $p_{1\bullet}$ and $p_{\bullet 1}$, only one free parameter is left – the one which would identify π_{XY} among all the distributions of $\mathcal{F}(\text{Bern}(p_{1\bullet}), \text{Bern}(p_{\bullet 1}))$. Thus, as in the Gaussian framework (Example 2.1), the dependence \mathfrak{D} is fully represented by one single number. However, that number cannot be Pearson’s correlation which is here:

$$\rho = \frac{p_{11} - p_{1\bullet}p_{\bullet 1}}{\sqrt{p_{1\bullet}(1 - p_{1\bullet})p_{\bullet 1}(1 - p_{\bullet 1})}}.$$

Indeed, $p_{11} \in (\max(0, p_{1\bullet} + p_{\bullet 1} - 1), \min(p_{1\bullet}, p_{\bullet 1}))$ – the so-called ‘Fréchet bounds’ (Fréchet, 1951, Hoeffding, 1940) in $\mathcal{F}(\text{Bern}(p_{1\bullet}), \text{Bern}(p_{\bullet 1}))$ – which in turn imposes

$$- \left(\min \left\{ \frac{p_{1\bullet}p_{\bullet 1}}{(1 - p_{1\bullet})(1 - p_{\bullet 1})}, \frac{(1 - p_{1\bullet})(1 - p_{\bullet 1})}{p_{1\bullet}p_{\bullet 1}} \right\} \right)^{1/2} < \rho < \left(\min \left\{ \frac{p_{1\bullet}(1 - p_{\bullet 1})}{(1 - p_{1\bullet})p_{\bullet 1}}, \frac{p_{\bullet 1}(1 - p_{1\bullet})}{(1 - p_{\bullet 1})p_{1\bullet}} \right\} \right)^{1/2}.$$

So, not all combinations $(p_{1\bullet}, p_{\bullet 1}, \rho) \in (0, 1) \times (0, 1) \times (-1, 1)$ produce a valid bivariate Bernoulli distribution, and ρ is not variation-independent of the marginal parameters $p_{1\bullet}$ and $p_{\bullet 1}$. The same holds for any quantity related to the χ^2 -statistic such as the contingency coefficient Φ^2 , or any one of the ‘association factors’/‘Pearson ratios’ (Good, 1956, Goodman, 1996):

$$\psi_{xy} \doteq \frac{p_{xy}}{p_{x\bullet}p_{\bullet y}} \in \left(\max \left(0, \frac{1}{p_{x\bullet}} + \frac{1}{p_{\bullet y}} - \frac{1}{p_{x\bullet}p_{\bullet y}} \right), \frac{1}{\max(p_{x\bullet}, p_{\bullet y})} \right) \quad (x, y) \in \{0, 1\} \times \{0, 1\}. \tag{2.4}$$

Likewise, the above bounds constrain Spearman’s ρ_S and Kendall’s τ , both equal to $p_{11} - p_{1\bullet}p_{\bullet 1}$ in this case (Genest and Nešlehová, 2007, Example 7), to be such that

$$\max \left(-p_{1\bullet}p_{\bullet 1}, -(1 - p_{1\bullet})(1 - p_{\bullet 1}) \right) \leq \rho_S, \tau \leq \min(p_{1\bullet}, p_{\bullet 1})(1 - \max(p_{1\bullet}, p_{\bullet 1})),$$

thus ρ_S and τ are not variation-independent to $p_{1\bullet}$ and $p_{\bullet 1}$.³ Consequently, none of these can be thought of as appropriately describing \mathfrak{D} in a distribution such as (2.3).

In fact, results of Edwards (1963) and Rudas (2018, Theorem 6.3) establish that any parameter allowing identification of the distribution (2.3) in $\mathcal{F}(\text{Bern}(p_{1\bullet}), \text{Bern}(p_{\bullet 1}))$, while being variation-independent of

²Here we consider only full tables (all positive entries) for simplicity. Presence of structural zeros is treated in full generality in the following sections.

³This remains true for any ‘tie-corrected’ version such as τ_b (Kendall, 1955, Chapter 3) or τ_c (Stuart, 1953).

$(p_{1\bullet}, p_{\bullet 1})$, must be a one-to-one function of the odds-ratio

$$\omega = \frac{p_{00}p_{11}}{p_{10}p_{01}} \in (0, \infty).$$

Indeed any triplet $(p_{1\bullet}, p_{\bullet 1}, \omega) \in (0, 1) \times (0, 1) \times (0, \infty)$ defines one and only one table (2.3) (Mosteller, 1968, equation (2*)). Any one-to-one function of ω is thus a valid representation of \mathfrak{D} as well; this includes $\log \omega$, Yule (1912)'s ‘association’ and ‘colligation’ coefficients $Q = (\omega - 1)/(\omega + 1)$ and $\Upsilon = (\sqrt{\omega} - 1)/(\sqrt{\omega} + 1)$. \square

Example 2.3. Bivariate copula model. Let the cumulative distribution function (cdf) of π_{XY} be F_{XY} , and the cdf's of π_X and π_Y be F_X and F_Y , respectively. It follows from Sklar (1959, *Théorème 2*) that there exists a *copula* C – that is, a bivariate cdf supported on $[0, 1]^2$ with standard uniform margins – such that

$$F_{XY}(x, y) = C(F_X(x), F_Y(y)), \quad \forall(x, y). \quad (2.5)$$

This establishes such C as a natural candidate for representing the dependence \mathfrak{D} in any bivariate distribution, as it appears to capture how to reconstruct F_{XY} from F_X and F_Y . If X and Y are both continuous variables, C is unique and can easily be seen to be the distribution of the vector $(F_X(X), F_Y(Y))$ (Schweizer and Wolff, 1981, Theorem 3(iii)). Therefore, by the virtue of the Probability Integral Transform result $(F_X(X), F_Y(Y)) \sim \mathcal{U}_{[0,1]}$, it is ‘marginal-distribution-free’,⁴ that is, ‘margin-free’. Thus, C provides the margin-free information which allows unequivocal identification of π_{XY} in $\mathcal{F}(\pi_X, \pi_Y)$, which is \mathfrak{D} by Definition 2.1. This is consistent with Example 2.1, as the copula of a bivariate Gaussian distribution is by definition the Gaussian copula (Joe, 2015, Section 4.3.1), whose only free parameter is the correlation ρ . By contrast, the dependence of the non-Gaussian distributions of $\mathcal{F}(\mathcal{N}(\mu_X, \sigma_X^2), \mathcal{N}(\mu_Y, \sigma_Y^2))$ is characterised by other copulas, not driven by ρ .

In non-continuous cases, though, the copula is not unique, meaning that any C satisfying (2.5) automatically contains irrelevant information not related to \mathfrak{D} . Evidently, what (2.5) may uniquely identify is only the restriction of C to⁵ $\text{Ran } F_X \times \text{Ran } F_Y$ (Sklar, 1959, *Théorème 1*), which is commonly referred to as the *subcopula* (Schweizer and Sklar, 1974, Definition 3). Defined on $\text{Ran } F_X \times \text{Ran } F_Y$, the subcopula must adjust to F_X and F_Y , and thus *cannot* be margin-free – and neither can any copula subtended by it. In particular, the copula obtained by ‘jittering’ discrete variables with uniform noise to make them artificially continuous, known as the *multilinear* or *checkerboard copula* C^{\boxtimes} , admits a density which amounts to the collection of Pearson ratios such as (2.4) (Genest et al., 2014, Definition 2.1), and these were seen not to be variation-independent to the marginal parameters.

⁴Where ‘distribution-free’ is understood in the sense of Kendall and Sundrum (1953): *free of the parent distribution*. Thus, more specifically here, ‘marginal-distribution-free’ means *free of the marginal distributions of the parent distribution* π_{XY} .

⁵ Ran denotes the range: $\text{Ran } F = \{t \in [0, 1] : \exists x \in \overline{\mathbb{R}} \text{ s.t. } F(x) = t\}$.

This may be illustrated further Within the bivariate Bernoulli framework (Example 2.2), where (2.5) reduces down to $p_{00} = C(1 - p_{1\bullet}, 1 - p_{\bullet 1})$, for given $p_{1\bullet}, p_{\bullet 1} \in (0, 1)$ and copula C . The value of C at $(1 - p_{1\bullet}, 1 - p_{\bullet 1})$ is the subcopula, and is enough to identify the other values p_{01}, p_{10} and p_{11} – hence the whole distribution – by substitution. The odds ratio, entirely capturing the dependence \mathfrak{D} here, is

$$\omega \doteq \omega(p_{1\bullet}, p_{\bullet 1}) = \frac{C(1 - p_{1\bullet}, 1 - p_{\bullet 1})(C(1 - p_{1\bullet}, 1 - p_{\bullet 1}) + p_{1\bullet} + p_{\bullet 1} - 1)}{(1 - p_{1\bullet} - C(1 - p_{1\bullet}, 1 - p_{\bullet 1}))(1 - p_{\bullet 1} - C(1 - p_{1\bullet}, 1 - p_{\bullet 1}))},$$

which clearly depends on $p_{1\bullet}$ and $p_{\bullet 1}$ in general – in fact, the only copula C guaranteeing this function $\omega(p_{1\bullet}, p_{\bullet 1})$ to be constant in $p_{1\bullet}$ and $p_{\bullet 1}$ is the *Plackett copula*, which was precisely designed in that purpose (Plackett, 1965). Thus, we may easily construct two bivariate Bernoulli distributions using the *same copula* C in (2.5), but showing very *different* dependence structures. Marshall (1996, Example 1.1) shows an extreme example of this.

All in all, when both X and Y are continuous variables, then the unique copula C of their distribution is a valid representation of their dependence. In non-continuous cases, on the other hand, neither the subcopula nor any of its copula extension may isolate the dependence structure, so none of these are valid representations of \mathfrak{D} . \square

As will be seen, the parities ‘ $\mathfrak{D} \sim$ Pearson’s correlation ρ ’, ‘ $\mathfrak{D} \sim$ odds ratio ω ’, ‘ $\mathfrak{D} \sim$ copula C ’ in the Gaussian, Bernoulli and unspecified continuous models, respectively, are just particular cases of a very general result: there exists a universal representation of the dependence \mathfrak{D} which is valid for any bivariate distribution π_{XY} , and ρ , ω and C will be shown (in Section 4.3) to be equivalent to that universal representation in the specific situations of Examples 2.1, 2.2 and 2.3.

3 The dependence \mathfrak{D} and its defining properties

3.1 Framework

We consider a general measure-theoretic framework, so as to provide in a unified way results valid across all cases of potential interest (discrete, continuous, mixed or hybrid random vectors alike). Let $(\mathfrak{X}, \mathcal{B}_{\mathfrak{X}}, \mu_{\mathfrak{X}})$ and $(\mathfrak{Y}, \mathcal{B}_{\mathfrak{Y}}, \mu_{\mathfrak{Y}})$ be two finite measure spaces, and let $\mathcal{P}_{\mathfrak{X} \times \mathfrak{Y}}$ be the set of all probability distributions on $(\mathfrak{X} \times \mathfrak{Y}, \mathcal{B}_{\mathfrak{X}} \otimes \mathcal{B}_{\mathfrak{Y}})$ which are absolutely continuous with respect to $\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}}$. For any random vector (X, Y) with distribution $\pi_{XY} \in \mathcal{P}_{\mathfrak{X} \times \mathfrak{Y}}$, denote $\varphi_{XY} \doteq d\pi_{XY} / d(\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}})$. The marginal distributions π_X on $(\mathfrak{X}, \mathcal{B}_{\mathfrak{X}})$ and π_Y on $(\mathfrak{Y}, \mathcal{B}_{\mathfrak{Y}})$ are absolutely continuous with respect to $\mu_{\mathfrak{X}}$ and $\mu_{\mathfrak{Y}}$, respectively, and we let $\varphi_X \doteq d\pi_X / d\mu_{\mathfrak{X}}$ and $\varphi_Y \doteq d\pi_Y / d\mu_{\mathfrak{Y}}$. We call these functions the joint, X -marginal and Y -marginal

densities of π_{XY} . Also, let the conditional densities be

$$\varphi_{Y|X} = \frac{d\pi_{XY}}{d(\pi_X \times \mu_{\mathfrak{Y}})} \quad \text{and} \quad \varphi_{X|Y} = \frac{d\pi_{XY}}{d(\mu_{\mathfrak{X}} \times \pi_Y)}, \quad (3.1)$$

defined on the sets $\mathcal{S}_X \times \mathfrak{Y}$ and $\mathfrak{X} \times \mathcal{S}_Y$, respectively, where $\mathcal{S}_X = \{x \in \mathfrak{X} : \pi_X(A) > 0 \forall A \in \mathcal{B}_{\mathfrak{X}} \text{ s.t. } A \ni x\} \subseteq \mathfrak{X}$ and $\mathcal{S}_Y = \{y \in \mathfrak{Y} : \pi_Y(B) > 0 \forall B \in \mathcal{B}_{\mathfrak{Y}} \text{ s.t. } B \ni y\} \subseteq \mathfrak{Y}$. Finally, denote $\mathcal{S}_{XY} = \{(x, y) \in \mathfrak{X} \times \mathfrak{Y} : \pi_{XY}(A \times B) > 0 \forall (A, B) \in \mathcal{B}_{\mathfrak{X}} \otimes \mathcal{B}_{\mathfrak{Y}} \text{ s.t. } A \ni x, B \ni y\} \subseteq \mathcal{S}_X \times \mathcal{S}_Y \subseteq \mathfrak{X} \times \mathfrak{Y}$. We will refer to the sets $\mathcal{S}_X, \mathcal{S}_Y$ and \mathcal{S}_{XY} as the (marginal and joint) supports of π_X, π_Y and π_{XY} . The above densities are unique up to relevant almost-everywhere equalities.

3.2 Conditional densities and dependence

It appears from (3.1) that

$$\pi_{XY}(\cdot) = \iint \varphi_{Y|X} \mathbb{I}_{\{\cdot\}} d(\pi_X \times \mu_{\mathfrak{Y}}) = \iint \varphi_{X|Y} \mathbb{I}_{\{\cdot\}} d(\mu_{\mathfrak{X}} \times \pi_Y) \quad (3.2)$$

where $\mathbb{I}_{\{\cdot\}}$ is the indicator function of a measurable set. Take two distinct distributions $\pi_{XY}^{(1)}$ and $\pi_{XY}^{(2)}$ in the Fréchet class $\mathcal{F}(\pi_X, \pi_Y)$. As $\pi_X^{(1)} = \pi_X^{(2)} = \pi_X$, it follows from the first equality that what differentiates $\pi_{XY}^{(1)}$ and $\pi_{XY}^{(2)}$ must be identifiable only from contrasting $\{\varphi_{Y|X}^{(1)}(\cdot|x) : x \in \mathcal{S}_X\}$ and $\{\varphi_{Y|X}^{(2)}(\cdot|x) : x \in \mathcal{S}_X\}$. It must also be identifiable only from contrasting $\{\varphi_{X|Y}^{(1)}(\cdot|y) : y \in \mathcal{S}_Y\}$ and $\{\varphi_{X|Y}^{(2)}(\cdot|y) : y \in \mathcal{S}_Y\}$, from the second equality, as $\pi_Y^{(1)} = \pi_Y^{(2)} = \pi_Y$. By Definition 2.1, what distinguishes $\pi_{XY}^{(1)}$ and $\pi_{XY}^{(2)}$ is their dependence. So:

Fact 3.1. The dependence \mathfrak{D} of a bivariate distribution π_{XY} must be entirely recoverable from either of its sets of conditional densities $\{\varphi_{Y|X}\} \doteq \{\varphi_{Y|X}(\cdot|x) : x \in \mathcal{S}_X\}$ or $\{\varphi_{X|Y}\} \doteq \{\varphi_{X|Y}(\cdot|y) : y \in \mathcal{S}_Y\}$. We will say that \mathfrak{D} is *encapsulated* in $\{\varphi_{Y|X}\}$ and in $\{\varphi_{X|Y}\}$.

The link between conditional densities and independence/non-independence is well understood. For example, Wermuth and Cox (2005) describe: ‘If proper variables are statistically independent, then the distribution of one of them is the same no matter at which fixed levels the other variable is considered’; whereas variables not independent are ‘such that there are differences in the distributions of one variable for at least some of the levels of the other’. That is, if the value taken by one of the variables changes, the stochastic behaviour of the other is disturbed – this is the ‘influence’ mentioned in Section 1. In our notation, this amounts to:⁶

$$X \perp\!\!\!\perp Y \iff \exists f_Y \in L_1(\mathfrak{Y}, \mu_{\mathfrak{Y}}) \text{ s.t. } \varphi_{Y|X}(y|x) = f_Y(y) \quad (\pi_X \times \mu_{\mathfrak{Y}})\text{-a.e.},$$

⁶For any measure space $(\mathcal{S}, \mathcal{B}, \mu)$, we define, for $p > 0$, $L_p(\mathcal{S}, \mu) \doteq \{\text{measurable } f : \mathcal{S} \rightarrow \mathbb{R} : (\int |f|^p d\mu)^{1/p} < \infty\}$. Likewise, $L_\infty(\mathcal{S}, \mu) \doteq \{\text{measurable } f : \mathcal{S} \rightarrow \mathbb{R} : \text{ess sup}|f| < \infty\}$.

i.e., $\varphi_{Y|X}(y|x)$ is essentially constant in x . It is easily shown (Dawid, 1979) that:

$$X \perp\!\!\!\perp Y \iff \exists f_Y \in L_1(\mathfrak{Y}, \mu_{\mathfrak{Y}}) \text{ s.t. } \varphi_{Y|X}(y|x) = f_Y(y) \quad (\pi_X \times \mu_{\mathfrak{Y}})\text{-a.e.} \quad (3.3)$$

$$\iff \varphi_{Y|X}(y|x) = \varphi_Y(y) \quad (\pi_X \times \mu_{\mathfrak{Y}})\text{-a.e.}$$

$$\iff \varphi_{XY}(x, y) = \varphi_X(x)\varphi_Y(y) \quad (\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}})\text{-a.e.} \quad (3.4)$$

$$\iff \varphi_{X|Y}(x|y) = \varphi_X(x) \quad (\mu_{\mathfrak{X}} \times \pi_Y)\text{-a.e.}$$

$$\iff \exists f_X \in L_1(\mathfrak{X}, \mu_{\mathfrak{X}}) \text{ s.t. } \varphi_{X|Y}(x|y) = f_X(x) \quad (\mu_{\mathfrak{X}} \times \pi_Y)\text{-a.e.} \quad (3.5)$$

From (3.3) and (3.5), it is clear that (non-)independence is determined exclusively by the conditional densities, without any reference to the marginals. Fact 3.1 actually strengthens this statement, showing that *full characterisation* of the dependence \mathfrak{D} of π_{XY} is possible from either one of $\{\varphi_{Y|X}\}$ or $\{\varphi_{X|Y}\}$, and only from it. It follows that:

Corollary 3.1. *Two distributions $\pi_{XY}^{(1)}$ and $\pi_{XY}^{(2)}$ sharing the same $X|Y$ -conditional density (i.e., $\{\varphi_{X|Y}^{(1)}\} \equiv \{\varphi_{X|Y}^{(2)}\}$) or the same $Y|X$ -conditional density (i.e., $\{\varphi_{Y|X}^{(1)}\} \equiv \{\varphi_{Y|X}^{(2)}\}$) have identical dependence structures ($\mathfrak{D}^{(1)} = \mathfrak{D}^{(2)}$).*

3.3 One-to-one marginal transformations and canonical supports

An immediate consequence of Fact 3.1 is that \mathfrak{D} is essentially invariant under one-to-one ‘marginal transformations’. Formally, let $(\mathfrak{X}', \mathcal{B}_{\mathfrak{X}'})$ be a measurable space isomorphic to $(\mathfrak{X}, \mathcal{B}_{\mathfrak{X}})$ (Dudley, 2002, Chapter 13). For a given one-to-one function $\Phi : \mathfrak{X} \rightarrow \mathfrak{X}'$, such that both Φ and Φ^{-1} are measurable (i.e., Φ is bimeasurable), define the random variable $X' \doteq \Phi(X)$, with distribution $\pi_{X'} = \pi_X \circ \Phi^{-1}$ (‘push-forward’) on the support $\mathcal{S}_{X'} = \{x' \in \mathfrak{X}' : \pi_{X'}(A') > 0 \forall A' \in \mathcal{B}_{\mathfrak{X}'} \text{ s.t. } A' \ni x'\} = \{x' \in \mathfrak{X}' : \pi_X(A) > 0 \forall A \in \mathcal{B}_{\mathfrak{X}} \text{ s.t. } A \ni \Phi^{-1}(x')\}$. The distribution $\pi_{X'Y}$ of the vector (X', Y) admits as conditional density $\{\varphi_{Y|X'}\} = \{\varphi_{Y|X'}(\cdot|x') : x' \in \mathcal{S}_{X'}\}$, with

$$\varphi_{Y|X'}(\cdot|x') = \varphi_{Y|\Phi^{-1}(X')}(\cdot|\Phi^{-1}(x')) = \varphi_{Y|X}(\cdot|\Phi^{-1}(x')).$$

From a conditioning-on- X perspective, the transformation $X' = \Phi(X)$ only changes the ‘label’ assigned to the value on which we condition, but leaves the effective conditional density of Y unaffected. Similarly to (3.2) we can write

$$\pi_{X'Y}(\cdot) = \iint \varphi_{Y|X'}(y|\Phi(x)) \mathbb{I}_{\{\cdot\}} d(\pi_X \times \mu_{\mathfrak{Y}})(x, y),$$

showing that \mathfrak{D} would be entirely recoverable from $\varphi_{Y|X'}$, too, provided we know Φ . Also, as \mathfrak{D} provides all required information for unequivocally identifying π_{XY} in $\mathcal{F}(\pi_X, \pi_Y)$, it inherently allows unequivocal identification of $\pi_{X'Y}$ in $\mathcal{F}(\pi_{X'}, \pi_Y)$ as well, as distributions of these two Fréchet classes are in one-to-one

correspondence through the given Φ .

We can replicate this with a bimeasurable bijection $\Psi : \mathfrak{Y} \rightarrow \mathfrak{Y}'$ which defines a random variable $Y' = \Psi(Y)$ living on another measurable space $(\mathfrak{Y}', \mathcal{B}_{\mathfrak{Y}'})$. Consequently, for any given mappings Φ and Ψ as above, the respective dependence structures \mathfrak{D} and \mathfrak{D}' of the vectors (X, Y) and (X', Y') are equivalent (one can be recovered from the other). Thus, any valid representation of \mathfrak{D} must be equivariant (Lehmann and Casella, 2006, Chapter 3) under such one-to-one marginal transformations: the information they contain is effectively the same, it just need be encoded differently so as to keep track of the ‘changes of label’ $x' = \Phi(x)$ and $y' = \Psi(y)$.

It follows that the dependence structures of all bivariate distributions with *marginal supports of the same cardinality* (the mappings Φ and Ψ are one-to-one) are objects of the same kind. Therefore, dependence structures of distributions with marginals supported on arbitrary ‘canonical’ sets $\mathcal{S}'_X \subseteq \mathfrak{X}'$ and $\mathcal{S}'_Y \subseteq \mathfrak{Y}'$ are enough for characterising the dependence between marginals supported on any sets $\mathcal{S}_X \subseteq \mathfrak{X}$ and $\mathcal{S}_Y \subseteq \mathfrak{Y}$ such that $|\mathcal{S}_X| = |\mathcal{S}'_X|$ and $|\mathcal{S}_Y| = |\mathcal{S}'_Y|$.⁷ E.g., the dependence between discrete variables supported on sets \mathcal{S}_X and \mathcal{S}_Y of finite cardinality R and S can be entirely described by the dependence of a distribution with margins supported on $\mathcal{S}'_X = \{0, \dots, R-1\}$ and $\mathcal{S}'_Y = \{0, \dots, S-1\}$. Likewise, dependence structures between continuous variables both supported on $\mathcal{S}'_X = \mathcal{S}'_Y = [0, 1]$ are sufficient for describing dependence in any bivariate continuous distribution. We shall call versions of π_{XY} with marginal supports adjusted to the agreed canonical sets of the right cardinality, ‘*canonical versions*’ of π_{XY} .

3.4 Joint support and regional dependence

Clearly, X and Y can only be independent if $\mathcal{S}_{XY} = \mathcal{S}_X \times \mathcal{S}_Y$ (‘rectangular support’): if not, there exist sets $A \in \mathcal{B}_{\mathfrak{X}}$ and $B \in \mathcal{B}_{\mathfrak{Y}}$ of positive π_X - and π_Y -measure such that $\varphi_{XY}(x, y) = 0$ for almost all $(x, y) \in A \times B$, violating (3.4). Thus, any non-rectangular support \mathcal{S}_{XY} automatically implies non-independence, through the fact that some values of X and Y are not allowed simultaneously – a strong form of mutual ‘influence’, indeed. Holland and Wang (1987b) refer to this as ‘*regional dependence*’.

Regional dependence is dictated by the shape of \mathcal{S}_{XY} , but not by the actual distribution π_{XY} supported on that set. As the shape of \mathcal{S}_{XY} cannot be recovered from marginal specifications only, regional dependence is part of \mathfrak{D} , as per Definition 2.1. The shape of \mathcal{S}_{XY} should be distinguished from the set \mathcal{S}_{XY} itself, though. In particular, \mathcal{S}_{XY} contains marginal information, such as the supports \mathcal{S}_X and \mathcal{S}_Y , so cannot be entirely part of \mathfrak{D} . This said, the shape of \mathcal{S}_{XY} can be inferred from the joint support \mathcal{S}'_{XY} of any canonical version π'_{XY} of π_{XY} (Section 3.3) – and this one can only reveal the uninformative \mathcal{S}'_X and \mathcal{S}'_Y .

⁷ $|\mathcal{S}|$ denotes the cardinality of the set \mathcal{S} .

Thus, to comply with Definition 2.1, the dependence \mathfrak{D} is always to be described in terms of canonical versions of distributions.

Therefore, all distributions coming into play below will be assumed to be ‘canonical’, meaning that their marginal supports have been preliminarily transformed to some universally agreed sets of the right cardinality.⁸ Thus, it will always be assumed that the joint support of a distribution does not contain any crucial information about its marginals (as $\mathcal{S}_X, \mathcal{S}_Y$ are universally known), and that marginal supports of the same cardinality are equal; that is, for any two distributions $\pi_{XY}^{(1)}$ and $\pi_{XY}^{(2)}$, $|\mathcal{S}_X^{(1)}| = |\mathcal{S}_X^{(2)}| \Rightarrow \mathcal{S}_X^{(1)} = \mathcal{S}_X^{(2)}$ and $|\mathcal{S}_Y^{(1)}| = |\mathcal{S}_Y^{(2)}| \Rightarrow \mathcal{S}_Y^{(1)} = \mathcal{S}_Y^{(2)}$. In particular, as any two distributions $\pi_{XY}^{(1)}, \pi_{XY}^{(2)}$ sharing the same dependence automatically share the same regional dependence, this assumption allows us to write:

$$\mathfrak{D}^{(1)} = \mathfrak{D}^{(2)} \Rightarrow \mathcal{S}_{XY}^{(1)} = \mathcal{S}_{XY}^{(2)}. \quad (3.6)$$

All this understood, no further explicit reference to the canonical versions/supports will be made.

3.5 Marginal replacements

Another consequence of Fact 3.1 is that \mathfrak{D} must be invariant to ‘marginal replacement’ (Holland and Wang, 1987a, Definition 4.1). Let $\pi_{XY} \in \mathcal{F}(\pi_X, \pi_Y)$ and consider another X -marginal measure π_X^* equivalent to π_X ($\pi_X \ll \pi_X^* \ll \pi_X$, denoted here $\pi_X \approx \pi_X^*$). One can use the conditional density $\{\varphi_{Y|X}\}$ of π_{XY} as a Markov kernel, and construct the distribution π_{XY}^* with density

$$\varphi_{XY}^* = \varphi_{Y|X} \frac{d\pi_X^*}{d\mu_{\mathfrak{X}}} = \varphi_{XY} \frac{d\pi_X^*}{d\pi_X}. \quad (3.7)$$

Although $\pi_X^* \neq \pi_X$ and $\pi_Y^*(\cdot) = \iint \varphi_{XY} \frac{d\pi_X^*}{d\pi_X} \mathbb{I}_{\{\mathcal{S}_X \times \cdot\}} d(\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}}) \neq \iint \varphi_{XY} \mathbb{I}_{\{\mathcal{S}_X \times \cdot\}} d(\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}}) = \pi_Y(\cdot)$ in general, the dependence structure of π_{XY} and π_{XY}^* is identical as they share the same $\{\varphi_{Y|X}\}$ (Corollary 3.1). Re-writing (3.7) as

$$\frac{d\pi_{XY}^*}{d\pi_{XY}} = \frac{d\pi_X^*}{d\pi_X} \quad (3.8)$$

shows that $\pi_{XY}^* \approx \pi_{XY}$ (as $\pi_X^* \approx \pi_X$) and thus $\pi_Y^* \approx \pi_Y$. From any other Y -marginal measure $\pi_Y^{**} \approx \pi_Y$, one can produce by the same token yet another joint distribution π_{XY}^{**} with density

$$\varphi_{XY}^{**} = \varphi_{X|Y}^* \frac{d\pi_Y^{**}}{d\mu_{\mathfrak{Y}}} = \varphi_{XY}^* \frac{d\pi_Y^{**}}{d\pi_Y^*} = \varphi_{XY} \frac{d\pi_X^*}{d\pi_X} \frac{d\pi_Y^{**}}{d\pi_Y^*}. \quad (3.9)$$

Hence

$$\frac{d\pi_{XY}^{**}}{d\pi_{XY}} = \frac{d\pi_X^*}{d\pi_X} \frac{d\pi_Y^{**}}{d\pi_Y^*}, \quad (3.10)$$

showing that $\pi_{XY}^{**} \approx \pi_{XY}$. This time $\{\varphi_{X|Y}^{**}\} = \{\varphi_{X|Y}^*\}$, and π_{XY}^{**} has the same dependence as π_{XY}^* (Corollary 3.1), and hence as π_{XY} too. We have thus created two distributions π_{XY}^* and π_{XY}^{**} with respective X - and

⁸Their exact specification is irrelevant, as they are arbitrary.

Y -marginals set to arbitrary $\pi_X^* \approx \pi_X$ and $\pi_Y^{**} \approx \pi_Y$ while maintaining the same dependence \mathfrak{D} as the initial π_{XY} . Note that π_{XY}^{**} does not share any of the conditional densities $\{\varphi_{Y|X}\}$ or $\{\varphi_{X|Y}\}$ of π_{XY} , illustrating that Corollary 3.1 provides a sufficient but not necessary condition for equal dependence structures. In (3.8) and (3.10), the Radon-Nikodym derivatives of π_{XY}^* and π_{XY}^{**} with respect to π_{XY} have a product form – Theorem 4.1 below establishes that it is not coincidental. Finally, as $\pi_{XY}^{**} \approx \pi_{XY}^* \approx \pi_{XY}$, the joint support \mathcal{S}_{XY} is preserved through the successive replacements, in agreement with (3.6).

3.6 Iterative Proportional Fitting procedure and \mathcal{I} -projections

The marginal replacement of π_X by π_X^* via (3.7) affects the Y -margin (it was noted that $\pi_Y^* \neq \pi_Y$). Likewise, the marginal replacement of π_Y^* by π_Y^{**} in (3.9) affects the X -margin (in particular: $\pi_X^{**} \neq \pi_X^*$). Thus, for a couple of targeted marginal distributions (π_X^*, π_Y^*) , it is not clear if we can obtain from π_{XY} a distribution of $\mathcal{F}(\pi_X^*, \pi_Y^*)$ via a similar dependence-preserving process. The following lemma gives a first element of answer. Let $\pi_{XY} \in \mathcal{O}_{\mathfrak{X} \times \mathfrak{Y}}$, and define

$$N_{\pi_{XY}} = \{(A, B) \in \mathcal{B}_{\mathfrak{X}} \otimes \mathcal{B}_{\mathfrak{Y}} \text{ s.t. } \pi_{XY}(A \times B) = 0\},$$

the collection of rectangular subsets of $\mathfrak{X} \times \mathfrak{Y}$ on which π_{XY} is null. Also, let $\Sigma_{\pi_{XY}} = \{\pi \in \mathcal{O}_{\mathfrak{X} \times \mathfrak{Y}} : \pi \ll \pi_{XY}\}$, that is the distributions of $\mathcal{O}_{\mathfrak{X} \times \mathfrak{Y}}$ with support contained entirely in \mathcal{S}_{XY} .

Lemma 3.1. *Let $\pi_{XY} \in \mathcal{O}_{\mathfrak{X} \times \mathfrak{Y}} \cap \mathcal{F}(\pi_X, \pi_Y)$ have support \mathcal{S}_{XY} . For any couple of marginal distributions (π_X^*, π_Y^*) such that $\pi_X^* \approx \pi_X$ and $\pi_Y^* \approx \pi_Y$, we have:*

1. $\mathcal{O}_{\mathfrak{X} \times \mathfrak{Y}} \cap \Sigma_{\pi_{XY}} \cap \mathcal{F}(\pi_X^*, \pi_Y^*) \neq \emptyset \iff \forall (A, B) \in N_{\pi_{XY}}, \pi_X^*(A) + \pi_Y^*(B) \leq 1;$

2. If $\mathcal{O}_{\mathfrak{X} \times \mathfrak{Y}} \cap \Sigma_{\pi_{XY}} \cap \mathcal{F}(\pi_X^*, \pi_Y^*) \neq \emptyset$:

- (a) $\exists \bar{\pi}_{XY}^* \in \mathcal{O}_{\mathfrak{X} \times \mathfrak{Y}} \cap \Sigma_{\pi_{XY}} \cap \mathcal{F}(\pi_X^*, \pi_Y^*)$ such that $\pi_{XY}^* \in \mathcal{O}_{\mathfrak{X} \times \mathfrak{Y}} \cap \Sigma_{\pi_{XY}} \cap \mathcal{F}(\pi_X^*, \pi_Y^*) \Rightarrow \pi_{XY}^* \ll \bar{\pi}_{XY}^*;$

- (b) $\bar{\pi}_{XY}^* \approx \pi_{XY} \iff [\forall (A, B) \in N_{\pi_{XY}}, \pi_X^*(A) + \pi_Y^*(B) = 1 \Rightarrow (A^c, B^c) \in N_{\pi_{XY}}],$ where A^c and B^c are the complements of A and B in \mathfrak{X} and \mathfrak{Y} , respectively.

Part 1 gives a necessary and sufficient condition for the existence of some distribution of $\mathcal{F}(\pi_X^*, \pi_Y^*)$ with support contained in \mathcal{S}_{XY} . If so, Part 2-2a) says that there exists one of them whose support contains the support of all the others. Part 2-2b) gives a necessary and sufficient condition for that ‘maximal support’ to be \mathcal{S}_{XY} – in other words, for the existence of a distribution of $\mathcal{F}(\pi_X^*, \pi_Y^*)$ with support \mathcal{S}_{XY} . We conclude:

Corollary 3.2. *Let $\pi_{XY} \in \mathcal{O}_{\mathfrak{X} \times \mathfrak{Y}} \cap \mathcal{F}(\pi_X, \pi_Y)$. For any couple of marginal distributions (π_X^*, π_Y^*) such that*

$\pi_X^* \approx \pi_X$ and $\pi_Y^* \approx \pi_Y$, there exists a distribution of $\pi_{XY}^* \in \mathcal{F}(\pi_X^*, \pi_Y^*)$ such that $\pi_{XY}^* \approx \pi_{XY}$ if and only if

$$\begin{aligned} \forall (A, B) \in \mathcal{N}_{\pi_{XY}}, \pi_X^*(A) + \pi_Y^*(B) \leq 1, \text{ and} \\ \text{any } (A, B) \in \mathcal{N}_{\pi_{XY}} \text{ s.t. } \pi_X^*(A) + \pi_Y^*(B) = 1 \text{ is such that } (A^c, B^c) \in \mathcal{N}_{\pi_{XY}}. \end{aligned} \quad (3.11)$$

The importance of Corollary 3.2 stems from (3.6), establishing the connection between the joint support of a distribution and its dependence. Hence (3.11) provides a necessary condition for finding a distribution π_{XY}^* in $\mathcal{F}(\pi_X^*, \pi_Y^*)$ with the same dependence as π_{XY} : marginals (π_X^*, π_Y^*) violating (3.11) cannot co-exist with the support of π_{XY} , that is, with its specific regional dependence. Condition (3.11) is trivially fulfilled for all (π_X^*, π_Y^*) if π_{XY} has rectangular support, as in that case any $(A, B) \in \mathcal{N}_{\pi_{XY}}$ is such that $\pi_X(A) = \pi_Y(B) = 0$, thus $\pi_X^*(A) = \pi_Y^*(B) = 0$ as $\pi_X^* \ll \pi_X$ and $\pi_Y^* \ll \pi_Y$. In fact, (3.11) is a rather mild condition, as breaking it requires the existence of a ‘large’ area in $\mathcal{S}_X \times \mathcal{S}_Y$ on which π_{XY} is null, imposing extensive incompatibility between values of X and Y (‘extreme’ regional dependence).

An example may be the case of a bivariate Bernoulli distribution (2.3) with the main diagonal as support, that is, $p_{10} = p_{01} = 0$. This evidently implies $p_{1\bullet} = p_{\bullet 1}$, that is, identical margins $\pi_X = \pi_Y$. Effectively this specific regional dependence structure imposes $X = Y$, therefore it cannot be found in Fréchet classes $\mathcal{F}(\text{Bern}(p_{1\bullet}), \text{Bern}(p_{\bullet 1}))$ such that $p_{1\bullet} \neq p_{\bullet 1}$, for the obvious reason.

We show below that (3.11) is actually also sufficient for finding a distribution π_{XY}^* in $\mathcal{F}(\pi_X^*, \pi_Y^*)$ with the same dependence as π_{XY} , with a few added mild technical conditions on π_{XY} .⁹ The subset of distributions of $\mathcal{P}_{\mathfrak{X} \times \mathfrak{Y}}$ satisfying those conditions, listed in Appendix A, will be denoted $\overline{\mathcal{P}}_{\mathfrak{X} \times \mathfrak{Y}}$. First, we need the following lemma, which easily ensues from Nussbaum (1993)’s Theorem 3.9, Lemma 3.11 and Corollary 3.13 (his ‘compatibility’ condition amounts to (3.11), while his Hypotheses 3.6-3.10 correspond to $\pi_{XY} \in \overline{\mathcal{P}}_{\mathfrak{X} \times \mathfrak{Y}}$).

Lemma 3.2. *Let $\pi_{XY} \in \overline{\mathcal{P}}_{\mathfrak{X} \times \mathfrak{Y}} \cap \mathcal{F}(\pi_X, \pi_Y)$ and (π_X^*, π_Y^*) be a couple of marginal distributions such that $\pi_X^* \approx \pi_X$ and $\pi_Y^* \approx \pi_Y$. Under condition (3.11), there exists a unique distribution $\pi_{XY}^* \in \mathcal{F}(\pi_X^*, \pi_Y^*)$ such that*

$$\frac{d\pi_{XY}^*}{d\pi_{XY}} = \alpha_X^* \beta_Y^* \quad (3.12)$$

for (a.e.-)positive functions $\alpha_X^* \in L_1(\mathfrak{X}, \mu_X)$ and $\beta_Y^* \in L_1(\mathfrak{Y}, \mu_Y)$ unique up to a multiplicative constant.

According to Nussbaum (1993, Theorem 4.19 and Remark 4.20), the distribution π_{XY}^* in (3.12) can be obtained as the limit of a (geometrically convergent) iterative procedure consisting in successive X - and

⁹These conditions are those given in Nussbaum (1993). In fact, on page 81, Nussbaum (1993) mentioned that these extra assumptions may well be an artefact of the argument, and the result of main interest [his Theorem 3.9] may remain true without them – this author has not attempted to prove or disprove that claim, arguing in Appendix A that these requirements are, in any case, all but constraining in the considered framework.

Y -marginal replacements *à la* (3.7)-(3.9) alternately. Specifically, if one sets

$$\forall n \geq 0, \quad \begin{cases} \frac{d\pi_{XY}^{(2n)}}{d\pi_{XY}} = \alpha_X^{(n)} \beta_Y^{(n)}, & \frac{d\pi_{XY}^{(2n+1)}}{d\pi_{XY}} = \alpha_X^{(n+1)} \beta_Y^{(n)} \\ \alpha_X^{(n+1)} = \frac{d\pi_X^*}{d\pi_X^{(2n)}} \alpha_X^{(n)}, & \beta_Y^{(n+1)} = \frac{d\pi_Y^*}{d\pi_Y^{(2n+1)}} \beta_Y^{(n)} \end{cases}, \quad (3.13)$$

with $\alpha_X^{(0)} = \beta_Y^{(0)} = 1$ and $\pi_X^{(0)} = \pi_X$, $\pi_Y^{(0)} = \pi_Y$ (and $\forall n' \geq 0$, $\pi_X^{(n')}, \pi_Y^{(n')}$ are the margins of $\pi_{XY}^{(n')}$), then $\pi_{XY}^{(n')} \rightarrow \pi_{XY}^*$ in total variation as $n' \rightarrow \infty$. Given that each step corresponds to a dependence-preserving marginal replacement and that $\pi_{XY}^{(n')} \approx \pi_{XY} \forall n'$ including in the limit ($\pi_{XY}^* \approx \pi_{XY}$), the initial dependence of π_{XY} is maintained throughout the process. Thus:

Corollary 3.3. *Let $\pi_{XY} \in \overline{\mathcal{D}}_{\mathfrak{X} \times \mathfrak{Y}} \cap \mathcal{F}(\pi_X, \pi_Y)$ and (π_X^*, π_Y^*) be marginal distributions such that $\pi_X^* \approx \pi_X$ and $\pi_Y^* \approx \pi_Y$. Under condition (3.11), there exists a unique distribution π_{XY}^* in $\mathcal{F}(\pi_X^*, \pi_Y^*)$ sharing the same dependence \mathfrak{D} as π_{XY} ; and that distribution satisfies (3.12).*

The recursive process (3.13) is known as the Iterative Proportional Fitting (IPF) procedure (Deming and Stephan, 1940, Sinkhorn, 1964, Ireland and Kullback, 1968, Kullback, 1968, Fienberg, 1970, Založnik, 2011, Idel, 2016). It has strong links to the so-called ‘ \mathcal{I} -geometry’ (Csiszár, 1975). For two distributions $\tilde{\pi}, \pi$ on $(\mathfrak{X} \times \mathfrak{Y}, \mathcal{B}_{\mathfrak{X}} \otimes \mathcal{B}_{\mathfrak{Y}})$, define

$$\mathcal{I}(\tilde{\pi} \parallel \pi) = \begin{cases} \iint \log \left(\frac{d\tilde{\pi}}{d\pi} \right) d\tilde{\pi} = \iint \log \left(\frac{d\tilde{\pi}}{d\pi} \right) \frac{d\tilde{\pi}}{d\pi} d\pi & \text{if } \tilde{\pi} \ll \pi \\ \infty & \text{if } \tilde{\pi} \not\ll \pi \end{cases},$$

the relative entropy of $\tilde{\pi}$ with respect to π – also called the \mathcal{I} -divergence, or Kullback-Leibler divergence. Properties of this quantity are well-known (Cover and Thomas, 2006). In particular, $\mathcal{I}(\tilde{\pi} \parallel \pi) \geq 0$ always, with $\mathcal{I}(\tilde{\pi} \parallel \pi) = 0 \iff \pi = \tilde{\pi}$. Hence, although it is not symmetric (and does not satisfy a triangular inequality), $\mathcal{I}(\cdot \parallel \cdot)$ may be thought of as some sort of distance between distributions, which defines a specific geometry. *Inter alia*, we may define a concept of projection as follows:

Definition 3.1. *Let \mathcal{E} be some set of distributions on $(\mathfrak{X} \times \mathfrak{Y}, \mathcal{B}_{\mathfrak{X}} \otimes \mathcal{B}_{\mathfrak{Y}})$, assumed to be convex and closed in total variation. Then, for a distribution π , any distribution $\tilde{\pi} \in \mathcal{E}$ such that $\mathcal{I}(\tilde{\pi} \parallel \pi) = \inf_{\tilde{\pi} \in \mathcal{E}} \mathcal{I}(\tilde{\pi} \parallel \pi)$ is called an \mathcal{I} -projection of π on \mathcal{E} .*

An \mathcal{I} -projection exists and is unique as soon as $\inf_{\tilde{\pi} \in \mathcal{E}} \mathcal{I}(\tilde{\pi} \parallel \pi) < \infty$, an obvious ‘reachability’ condition (Csiszár, 1975). Noting that a Fréchet class of distributions is convex and closed in total variation (Cramer, 2000, Lemma 4.1), we can define the \mathcal{I} -projection of π_{XY} on $\mathcal{F}(\pi_X^*, \pi_Y^*)$ as the distribution $\tilde{\pi}_{XY}^*$ satisfying

$$\mathcal{I}(\tilde{\pi}_{XY}^* \parallel \pi_{XY}) = \inf_{\pi \in \mathcal{F}(\pi_X^*, \pi_Y^*)} \mathcal{I}(\pi \parallel \pi_{XY}).$$

When $\inf_{\pi \in \mathcal{F}(\pi_X^*, \pi_Y^*)} \mathcal{I}(\pi \| \pi_{XY}) < \infty$ (i.e., when $\tilde{\pi}_{XY}^*$ exists and is unique), Rüschemdorf (1995) showed that the IPF procedure (3.13) converges to $\tilde{\pi}_{XY}^*$. From Csiszár (1975, Corollary 3.2) and Rüschemdorf and Thomsen (1993, Theorem 3), we can further deduce that, under condition (3.11), $\tilde{\pi}_{XY}^*$ is in fact π_{XY}^* , the distribution defined by (3.12); that is, the distribution of $\mathcal{F}(\pi_X^*, \pi_Y^*)$ with the same dependence as π_{XY} (Corollary 3.3). Hence, *\mathcal{I} -projections on Fréchet classes maintain dependence.*

Therefore, the \mathcal{I} -geometry seems particularly well suited for providing a formal geometrical framework around (2.1). For example, Csiszár (1975) proved a Pythagoras-like theorem for \mathcal{I} : for two arbitrary distributions $\pi_{XY} \in \mathcal{F}(\pi_X, \pi_Y)$ and $\check{\pi}_{XY}^* \in \mathcal{F}(\pi_X^*, \pi_Y^*)$, we have

$$\mathcal{I}(\tilde{\pi}_{XY}^* \| \pi_{XY}) = \mathcal{I}(\check{\pi}_{XY}^* \| \tilde{\pi}_{XY}^*) + \mathcal{I}(\tilde{\pi}_{XY}^* \| \pi_{XY}),$$

where $\tilde{\pi}_{XY}^*$ is the \mathcal{I} -projection of π_{XY} on $\mathcal{F}(\pi_X^*, \pi_Y^*)$. Hence $\tilde{\pi}_{XY}^*$ has the same marginals as $\check{\pi}_{XY}^*$ (but not the same dependence) while, from the previous argument, $\tilde{\pi}_{XY}^*$ has the same dependence as π_{XY} (but not the same marginals). Thus the ‘distance’ from π_{XY} to $\check{\pi}_{XY}^*$ can be broken down into two orthogonal contributions: what it takes to adjust the marginals (maintaining the dependence) plus what it takes to adjust the dependence (maintaining the marginals).

3.7 Compatibility of conditional densities

Fact 3.1 establishes that full description of \mathfrak{D} may be obtained from either $\{\varphi_{Y|X}\}$ or $\{\varphi_{X|Y}\}$. Therefore we may posit the existence of a mapping, say Δ , taking as argument conditional densities such as $\{\varphi_{Y|X}\}$ or $\{\varphi_{X|Y}\}$, and returning (a complete representation of) the dependence \mathfrak{D} which they encapsulate. Informally, we would have, for any specification of a $X|Y$ -conditional density $\{\varphi_{X|Y}\}$:

$$\Delta : \{\varphi_{X|Y}\} \rightarrow \Delta(\{\varphi_{X|Y}\}) = \text{dependence } \mathfrak{D} \text{ encapsulated in } \{\varphi_{X|Y}\},$$

and for any specification of a $Y|X$ -conditional density $\{\varphi_{Y|X}\}$:

$$\Delta : \{\varphi_{Y|X}\} \rightarrow \Delta(\{\varphi_{Y|X}\}) = \text{dependence } \mathfrak{D} \text{ encapsulated in } \{\varphi_{Y|X}\}.$$

That map should necessarily be such that

$$\Delta(\{\varphi_{X|Y}\}) = \Delta(\{\varphi_{Y|X}\}) \tag{3.14}$$

when $\{\varphi_{X|Y}\}$ and $\{\varphi_{Y|X}\}$ are the two ‘twin’ sets of conditional densities of a common bivariate distribution π_{XY} , as they encapsulate the same dependence – that of π_{XY} .

From a modelling perspective, it is sometimes more natural to specify an appropriate model for a vector (X, Y) through its two sets of conditional distributions (X given Y , and Y given X), rather than through

a joint distribution directly (Arnold et al., 2001). Yet, two separately postulated conditional models, say $\{\varphi_{X|Y}^*\}$ and $\{\varphi_{Y|X}\}$, do not necessarily interlock into a proper bivariate distribution. When there exists a bivariate distribution $\tilde{\pi}_{XY}$ with $\{\varphi_{X|Y}^*\}$ and $\{\varphi_{Y|X}\}$ as conditional densities, these two are called compatible (Arnold and Press, 1989, Arnold and Sarabia, 2022, Wang and Ip, 2008). Here we establish that two sets of conditional densities are compatible if and only if they encapsulate the same dependence.

Proposition 3.1. *For $\mathcal{S}_X \subset \mathfrak{X}$ and $\mathcal{S}_Y^* \subset \mathfrak{Y}$, let $\{\varphi_{Y|X}\} = \{\varphi_{Y|X}(\cdot|x) : x \in \mathcal{S}_X\}$ and $\{\varphi_{X|Y}^*\} = \{\varphi_{X|Y}^*(\cdot|y) : y \in \mathcal{S}_Y^*\}$ be some $Y|X$ - and a $X|Y$ -conditional density specifications. Then $\{\varphi_{Y|X}\}$ and $\{\varphi_{X|Y}^*\}$ are two compatible sets of conditional densities if and only if $\{\varphi_{Y|X}\}$ and $\{\varphi_{X|Y}^*\}$ both encapsulate the same dependence structure; i.e., $\Delta(\{\varphi_{Y|X}\}) = \Delta(\{\varphi_{X|Y}^*\})$.*

Intuitively, any arbitrary X -marginal distribution $\tilde{\pi}_X$ and $Y|X$ -conditional density $\varphi_{Y|X}$ can be freely combined as in (3.7) to form a bivariate distribution – provided that the support of $\tilde{\pi}_X$ matches \mathcal{S}_X in $\{\varphi_{Y|X}\}$. Likewise, any arbitrary Y -marginal and $X|Y$ -conditional density can be freely combined, up to the same support restriction. Thus, if we set the putative $\tilde{\pi}_{XY} = (\tilde{\pi}_X, \tilde{\pi}_Y; \tilde{\mathfrak{D}})$ as in (2.1), $\{\varphi_{Y|X}\}$ does not restrict $\tilde{\pi}_X$ while $\{\varphi_{X|Y}^*\}$ does not restrict $\tilde{\pi}_Y$, and (in)compatibility of $\{\varphi_{Y|X}\}$ and $\{\varphi_{X|Y}^*\}$ may only be dictated by their encapsulated dependence \mathfrak{D} and \mathfrak{D}^* . Clearly $\mathfrak{D} = \mathfrak{D}^*$ (which implies matching supports, as per (3.6)) is all what is needed for completing the representation $\tilde{\pi}_{XY} = (\tilde{\pi}_X, \tilde{\pi}_Y; \tilde{\mathfrak{D}})$ with $\tilde{\mathfrak{D}} = \mathfrak{D} = \mathfrak{D}^*$. On the other hand, if $\mathfrak{D} \neq \mathfrak{D}^*$, then $\{\varphi_{Y|X}\}$ and $\{\varphi_{X|Y}^*\}$ have no common ground around which articulate the desired bivariate distribution.

4 Universal representation of the dependence \mathfrak{D}

4.1 Equivalence classes of dependence structure

Combining the results of Section 3, we can now establish:

Theorem 4.1. *Let π_{XY} and π_{XY}^* be two distributions in $\overline{\mathfrak{D}}_{\mathfrak{X} \times \mathfrak{Y}}$. Then π_{XY} and π_{XY}^* admit the same dependence $\mathfrak{D} = \mathfrak{D}^*$ if and only if there exist (a.e.-)positive functions $\alpha_X^* \in L_1(\mathfrak{X}, \pi_X)$ and $\beta_Y^* \in L_1(\mathfrak{Y}, \pi_Y)$ such that*

$$\frac{d\pi_{XY}^*}{d\pi_{XY}} = \alpha_X^* \beta_Y^*. \quad (4.1)$$

The Radon-Nikodym derivative (4.1) is (a.e.-)positive, hence π_{XY} and π_{XY}^* have the same support – in agreement with (3.6). The product form $\alpha_X^* \beta_Y^*$, reminiscent of (3.4), implies that the relative difference between π_{XY} and π_{XY}^* shows up as two independent marginal adjustments, unable to affect the inner dependence structure. Evidently (4.1) defines an equivalence relationship between distributions, which in

turn defines equivalence classes of dependence:

$$\pi_{XY} \sim \pi_{XY}^* \iff \mathfrak{D} = \mathfrak{D}^* \iff \frac{d\pi_{XY}^*}{d\pi_{XY}} = \alpha_X^* \beta_Y^* \quad (\alpha_X^* \beta_Y^* > 0 \text{ a.e.}) \iff \pi_{XY}, \pi_{XY}^* \in [\mathfrak{D}], \quad (4.2)$$

where we define $[\mathfrak{D}]$ as the equivalence class containing all the distributions sharing the dependence \mathfrak{D} . Accordingly, the dependence can be thought of as a maximal invariant under the group operation (4.1). This allows the identification of a universal representation of \mathfrak{D} .

4.2 A universal representation of \mathfrak{D}

Let $\Gamma_{\mathfrak{X} \times \mathfrak{Y}}$ be the space of all finite signed measures γ on $(\mathfrak{X} \times \mathfrak{Y}, \mathcal{B}_{\mathfrak{X}} \otimes \mathcal{B}_{\mathfrak{Y}})$, and let

$$\Gamma_{\mathcal{S}_{XY}}^\circ = \{\gamma \in \Gamma_{\mathfrak{X} \times \mathfrak{Y}} : |\gamma| \ll \pi_{XY}, \gamma(A, \mathcal{S}_Y) = \gamma(\mathcal{S}_X, B) = 0 \quad \forall A \in \mathcal{B}_{\mathfrak{X}}, \forall B \in \mathcal{B}_{\mathfrak{Y}}\}, \quad (4.3)$$

the closed linear subspace of $\Gamma_{\mathfrak{X} \times \mathfrak{Y}}$ containing the measures absolutely continuous with respect to π_{XY} and with marginals both identically null – the notation $\Gamma_{\mathcal{S}_{XY}}^\circ$ makes it clear that this set is exclusively determined by the support \mathcal{S}_{XY} of π_{XY} . Equipped with the total variation norm $\|\gamma\| = |\gamma|(\mathfrak{X} \times \mathfrak{Y})$, $\Gamma_{\mathfrak{X} \times \mathfrak{Y}}$ and $\Gamma_{\mathcal{S}_{XY}}^\circ$ are Banach spaces (Folland, 2013, Section 5.1). Assume that we may find a basis for $\Gamma_{\mathcal{S}_{XY}}^\circ$, that is, a countable collection $\{\gamma_1, \gamma_2, \dots\} \in \Gamma_{\mathcal{S}_{XY}}^\circ$, such that any $\gamma \in \Gamma_{\mathcal{S}_{XY}}^\circ$ admits the representation

$$\gamma = \sum_k a_k \gamma_k$$

for a (unique) real vector (a_1, a_2, \dots) (Albiac and Kalton, 2016, Chapter 1). If so, call the number of elements in $\{\gamma_1, \gamma_2, \dots\}$ the dimension of $\Gamma_{\mathcal{S}_{XY}}^\circ$, denoted $\dim(\Gamma_{\mathcal{S}_{XY}}^\circ)$ (which may be infinite). Identifying such a basis is relatively straightforward in cases of interest – see Section 4.3. Then we have:

Theorem 4.2. *Let $\pi_{XY} \in \overline{\mathcal{P}}_{\mathfrak{X} \times \mathfrak{Y}}$ be a distribution with density $\varphi_{XY} = d\pi_{XY}/d(\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}})$ on its support \mathcal{S}_{XY} . For any basis $\{\gamma_1, \gamma_2, \dots\}$ of the set $\Gamma_{\mathcal{S}_{XY}}^\circ$ (4.3), the collection*

$$\mathbf{\Delta}_{XY} \doteq \left(\iint \log \varphi_{XY} d\gamma_1, \iint \log \varphi_{XY} d\gamma_2, \dots \right) \in \mathbb{R}^{\dim(\Gamma_{\mathcal{S}_{XY}}^\circ)} \quad (4.4)$$

is a maximal invariant under the group operation (4.1).

It directly follows:

Corollary 4.1. *Under the conditions of Theorem 4.2, the pair $(\mathcal{S}_{XY}, \mathbf{\Delta}_{XY})$ provides a complete representation of the dependence \mathfrak{D} of the vector (X, Y) .*

The representation $(\mathcal{S}_{XY}, \mathbf{\Delta}_{XY})$ is unique up to one-to-one correspondences. In particular, different bases $\{\gamma_1, \gamma_2, \dots\}$ lead to different $\mathbf{\Delta}_{XY}$'s, but all are equivalent representations of \mathfrak{D} . As $\{\gamma_k\} \in \Gamma_{\mathcal{S}_{XY}}^\circ$, we have

by definition $\iint \log \varphi_X d\gamma_k = \iint \log \varphi_Y d\gamma_k = 0 \forall k$ for any $\pi_{XY} \in \overline{\mathcal{D}}_{\mathfrak{X} \times \mathfrak{Y}}$, so that

$$\begin{aligned} \mathbf{\Delta}_{XY} &= \left(\iint \log \varphi_{XY} d\gamma_1, \iint \log \varphi_{XY} d\gamma_2, \dots \right) = \left(\iint \log \varphi_{Y|X} d\gamma_1, \iint \log \varphi_{Y|X} d\gamma_2, \dots \right) \\ &= \left(\iint \log \varphi_{X|Y} d\gamma_1, \iint \log \varphi_{X|Y} d\gamma_2, \dots \right). \end{aligned}$$

Also, $\mathcal{S}_{XY} = \{(x, y) \in \mathcal{S}_X \times \mathfrak{Y} : \varphi_{Y|X}(y|x) > 0\} = \{(x, y) \in \mathfrak{X} \times \mathcal{S}_Y : \varphi_{X|Y}(x|y) > 0\}$ (up to sets of measure 0), so the support \mathcal{S}_{XY} can be written in terms of $\{\varphi_{Y|X}\}$ or $\{\varphi_{X|Y}\}$ alone. Hence we can identify the pair $(\mathcal{S}_{XY}, \mathbf{\Delta}_{XY})$ to the mapping Δ introduced in Section 3.7:

$$(\mathcal{S}_{XY}, \mathbf{\Delta}_{XY}) = \Delta(\varphi_{Y|X}) = \Delta(\varphi_{X|Y}),$$

as prescribed by (3.14). In other words, examining only one of the sets $\{\varphi_{Y|X}\}$ or $\{\varphi_{X|Y}\}$ of a distribution π_{XY} is sufficient for identifying the dependence $\mathfrak{D} = (\mathcal{S}_{XY}, \mathbf{\Delta}_{XY})$ of π_{XY} , as per Fact 3.1.

It is also clear that, if φ_{XY} satisfies (3.4), then all elements of $\mathbf{\Delta}_{XY}$ are null. Hence, if $X \perp\!\!\!\perp Y$ then $\mathbf{\Delta}_{XY} \equiv \mathbf{0}$ and $\mathcal{S}_{XY} = \mathcal{S}_X \times \mathcal{S}_Y$ (rectangular support, Section 3.4). Conversely, if $\mathbf{\Delta}_{XY} = \mathbf{0}$ and $\mathcal{S}_{XY} = \mathcal{S}_X \times \mathcal{S}_Y$, then $X \perp\!\!\!\perp Y$. However, we may have $\mathcal{S}_{XY} \neq \mathcal{S}_X \times \mathcal{S}_Y$ and $\mathbf{\Delta}_{XY} = \mathbf{0}$: then X and Y are not independent. This is called ‘quasi-independence’ (Goodman, 1968):

Definition 4.1. *Let $(X, Y) \sim \pi_{XY}$ with density φ_{XY} on the support \mathcal{S}_{XY} . Then, X and Y are quasi-independent if there exist (a.e.-)positive functions $q_X \in L_1(\mathfrak{X}, \mu_{\mathfrak{X}})$ and $q_Y \in L_1(\mathfrak{Y}, \mu_{\mathfrak{Y}})$ such that $\varphi_{XY}(x, y) = q_X(x)q_Y(y) \mathbb{I}_{\{(x, y) \in \mathcal{S}_{XY}\}}$.*

Independence implies quasi-independence, while quasi-independence implies independence only on a rectangular support. If the support is non-rectangular, the distribution π_{XY} shows quasi-independence when the source of non-independence is purely ‘regional’, i.e. exclusively due to the shape of the support (Section 3.4). Extending the results of Caussinus (1965) in discrete settings (in particular, his ‘*Théorème IX*’ and the following ‘*Remarque*’), we conclude:

Corollary 4.2. *Let $(X, Y) \sim \pi_{XY}$. Under the conditions of Theorem 4.2:*

1. X and Y are quasi-independent $\iff \mathbf{\Delta}_{XY} = \mathbf{0}$;
2. X and Y are independent $\iff \mathbf{\Delta}_{XY} = \mathbf{0}$ and $\mathcal{S}_{XY} = \mathcal{S}_X \times \mathcal{S}_Y$.

Finally, it seems important to emphasise the following obvious consequence of Theorem 4.1:

Corollary 4.3. *Any valid representation or description of the dependence of a bivariate distribution must be maintained over any equivalence class of dependence $[\mathfrak{D}]$ defined by (4.2). In other words, any valid representation of \mathfrak{D} must be invariant under the group operation (4.1) or, yet equivalently, it must be recoverable from $(\mathcal{S}_{XY}, \mathbf{\Delta}_{XY})$ only.*

4.3 Particular cases

The representation of dependence provided by Corollary 4.1 aligns with known results in familiar situations. For less well-studied cases, it allows unequivocal identification of how to capture and/or describe the dependence. Below some of those particular cases are explored.

4.3.1 Two discrete variables

Let $\mathfrak{X} = \mathfrak{Y} = \mathbb{N}_0 = \{0, 1, 2, \dots\}$ and $\mu_{\mathfrak{X}} = \mu_{\mathfrak{Y}}$ be the counting measure on $(\mathbb{N}_0, 2^{\mathbb{N}_0})$, meaning that $\varphi_{XY}(x, y) = \mathbb{P}(X = x, Y = y) \doteq p_{xy}$, for $(x, y) \in \mathbb{N}_0 \times \mathbb{N}_0$. In this case, the set $\Gamma_{\mathfrak{X} \times \mathfrak{Y}}$ introduced in Section 4.2 is isometrically isomorphic to $\ell_1(\mathbb{N}_0 \times \mathbb{N}_0)$, which makes it easy to identify bases for $\Gamma_{\mathcal{S}_{XY}}^\circ$.

(a) X and Y have finite supports \mathcal{S}_X and \mathcal{S}_Y , $\mathcal{S}_{XY} = \mathcal{S}_X \times \mathcal{S}_Y$. Assume $\mathcal{S}_X = \{0, 1, \dots, R-1\}$ and $\mathcal{S}_Y = \{0, 1, \dots, S-1\}$, for $2 \leq R, S < \infty$, and $p_{xy} > 0 \forall (x, y) \in \{0, 1, \dots, R-1\} \times \{0, 1, \dots, S-1\} = \mathcal{S}_{XY}$. Then $\Gamma_{\mathcal{S}_{XY}}^\circ$ can be identified to the space $\mathcal{M}_{R \times S}^\circ$ of $(R \times S)$ -matrices with all rows and columns summing to 0. It is algebraic routine to show that $\dim(\Gamma_{\mathcal{S}_{XY}}^\circ) = \dim(\mathcal{M}_{R \times S}^\circ) = (R-1)(S-1)$, so the dependence in (X, Y) can be entirely described by this number of parameters – the χ^2 -test of independence in such a table articulates around $(R-1)(S-1)$ degrees of freedom, indeed. With e_{xy} the $(R \times S)$ -matrix whose all entries are zero except the (x, y) th entry which is 1, one can show that $\{E_{xy}^{(00)} \doteq (e_{xy} - e_{x0} - e_{0y} + e_{00}), x = 1, \dots, R-1, y = 1, \dots, S-1\}$ is a basis of $\mathcal{M}_{R \times S}^\circ$. Thus, the $(R-1)(S-1)$ components of Δ_{XY} in (4.4) can be

$$\log \frac{p_{00}p_{xy}}{p_{x0}p_{0y}}, \quad x = 1, \dots, R-1, y = 1, \dots, S-1, \quad (4.5)$$

in which we recognise the collection of log-odds ratios with respect to the pivot point $(0, 0)$ – any other pivot point may be chosen (Agresti, 2013, equation (2.11)). Another basis of $\mathcal{M}_{R \times S}^\circ$ is $\{E_{xy}^{(\text{loc})} \doteq (e_{x-1, y-1} - e_{x, y-1} - e_{x-1, y} + e_{xy}), x = 1, \dots, R-1, y = 1, \dots, S-1\}$, which would describe the dependence in terms of the ‘local odds ratios’ (Agresti, 2013, equation (2.10)):

$$\log \omega_{xy} \doteq \log \frac{p_{x-1, y-1}p_{xy}}{p_{x-1, y}p_{x, y-1}}, \quad x = 1, \dots, R-1, y = 1, \dots, S-1. \quad (4.6)$$

It is known that the two sets of odds-ratios are equivalent (one may be recovered from the other) – and so would be any other set built from yet a different basis of $\mathcal{M}_{R \times S}^\circ$. In agreement with Corollary 4.2, any collection of such log-odds-ratios is identically null when $X \perp\!\!\!\perp Y$. In the case $R = S = 2$, the dependence is fully captured by any one-to-one function of

$$\log \omega = \log \omega_{11} = \log \frac{p_{00}p_{11}}{p_{10}p_{01}},$$

as announced in Example 2.2.

(b) X and Y have finite supports \mathcal{S}_X and \mathcal{S}_Y , $\mathcal{S}_{XY} \neq \mathcal{S}_X \times \mathcal{S}_Y$. A non-rectangular support for a bivariate discrete distribution (‘incomplete table’) is known to cause specific issues (Bishop et al., 1975, Chapter 5). Here, the layout of the ‘forbidden’ entries is simply incorporated in $\Gamma_{\mathcal{S}_{XY}}^\circ$. As an example, consider a (3×3) -distribution π_{XY} supported on

$$\mathcal{S}_{XY} \sim \begin{pmatrix} \times & 0 & 0 \\ 0 & \times & \times \\ 0 & \times & \times \end{pmatrix} \quad (4.7)$$

(\times ’s stand for non-zero entries). It can be seen that the space $\Gamma_{\mathcal{S}_{XY}}^\circ$ is one-dimensional, with basis vector $E_{22}^{(\text{loc})} = e_{11} - e_{21} - e_{12} + e_{22}$. Hence, given the support (4.7), the dependence in such a distribution is fully characterised by (any one-to-one transformation of)

$$\log \frac{p_{11}p_{22}}{p_{21}p_{12}}. \quad (4.8)$$

The value of p_{00} does not enter the description of dependence; indeed, on support (4.7) it is entirely fixed by the marginal distributions: $p_{00} = \mathbb{P}(X = 0) = \mathbb{P}(Y = 0)$ – the regional dependence structure reflected by (4.7) imposes these two probabilities to be equal.

As another example, consider π_{XY} supported on

$$\mathcal{S}_{XY} \sim \begin{pmatrix} 0 & \times & \times \\ \times & 0 & \times \\ \times & \times & 0 \end{pmatrix}. \quad (4.9)$$

As \mathcal{S}_{XY} does not comprise any positive rectangular sub-table, a representation of dependence in terms of (log-)odds-ratios such as (4.5), (4.6) or (4.8) is not possible here. Though, Corollary 4.1 applies: the space $\Gamma_{\mathcal{S}_{XY}}^\circ$ is again unidimensional, with basis vector $e_{01} - e_{02} - e_{10} + e_{12} + e_{20} - e_{21}$. Thus, on support (4.9), the dependence is entirely captured by the single parameter

$$\log \frac{p_{01}p_{12}p_{20}}{p_{02}p_{10}p_{21}}. \quad (4.10)$$

In particular, X and Y are quasi-independent if and only if (4.10) is 0 (Corollary 4.2), that is, $p_{01}p_{12}p_{20} = p_{02}p_{10}p_{21}$, confirming Goodman (1968, equation (2.23)).

When the structural zeros are too prominent, the space $\Gamma_{\mathcal{S}_{XY}}^\circ$ may contain only the null element. Then, the dependence is entirely determined by the specific support. An obvious example is a diagonal support

such as

$$\mathcal{S}_{XY} \sim \begin{pmatrix} \times & 0 & 0 \\ 0 & \times & 0 \\ 0 & 0 & \times \end{pmatrix}. \quad (4.11)$$

Clearly $\Gamma_{\mathcal{S}_{XY}}^\circ = \{\mathbf{0}\}$, the (3×3) -null matrix. In other words, the fact that π_{XY} is supported on the main diagonal is what characterises the dependence entirely – concretely, it imposes $X = Y$ – without any need for refinement: p_{00}, p_{11} and p_{22} are fixed by the margins, which need to be identical in this case.

(c) X and Y have infinite (countable) supports \mathcal{S}_X and \mathcal{S}_Y . Assume $\mathcal{S}_X = \mathcal{S}_Y = \mathbb{N}_0$. If $\mathcal{S}_{XY} = \mathcal{S}_X \times \mathcal{S}_Y$ (rectangular support), then the dependence in (X, Y) is entirely described by an (infinite countable) collection of log-odds ratios, such as (4.5) or (4.6). If $\mathcal{S}_{XY} \neq \mathbb{N}_0 \times \mathbb{N}_0$, then the layout of structural zeros is part of the dependence. In case of scattered zeros not forming any particular pattern, the dependence may well remain captured by an (infinite countable) collection of (2×2) -odds ratios, which essentially ‘avoid’ the structural zeros. More specific patterns may give rise to other peculiar descriptions of the dependence structure, following the basis of $\Gamma_{\mathcal{S}_{XY}}^\circ$ which we can form.

4.3.2 Two (absolutely) continuous variables

Let $\mathfrak{X} = \mathfrak{Y} = [0, 1]$ and $\mu_{\mathfrak{X}} = \mu_{\mathfrak{Y}}$ be the Lebesgue measure. Denote f_{XY} the joint probability density of (X, Y) , which is essentially bounded for $\pi_{XY} \in \overline{\mathcal{D}}_{\mathfrak{X} \times \mathfrak{Y}}$ (see Appendix A). The set $\Gamma_{\mathcal{S}_{XY}}^\circ$ can now be identified with the subset of functions ψ of $L_1([0, 1]^2)$ such that $\psi(x, y) = 0$ for $(x, y) \notin \mathcal{S}_{XY}$, $\int_{\mathcal{S}_X} \psi(x, y) dx = 0 \forall y \in \mathcal{S}_Y$ and $\int_{\mathcal{S}_Y} \psi(x, y) dy = 0 \forall x \in \mathcal{S}_X$. The tensorised Haar system is a (Schauder) basis for $L_1([0, 1]^2)$ (Albiac and Kalton, 2016, Chapter 6). Appropriate amendments of it provide bases for $\Gamma_{\mathcal{S}_{XY}}^\circ$.

(a) $\mathcal{S}_{XY} = [0, 1]^2$. Call $\psi_0(x, y) = (\mathbb{I}_{\{x \in [0, 1/2)\}} - \mathbb{I}_{\{x \in [1/2, 1)\}})(\mathbb{I}_{\{y \in [0, 1/2)\}} - \mathbb{I}_{\{y \in [1/2, 1)\}})$, and define $\psi_{k, \ell}(x, y) = 2^k \psi_0(2^k(x, y) - \ell)$ for $k \in \mathbb{N}_0$ and $\ell \in \{0, \dots, 2^k - 1\}^2$. Then $\{\psi_{k, \ell}\}_{k, \ell}$ is a basis of $\Gamma_{\mathcal{S}_{XY}}^\circ$ – see that it is the usual bivariate tensorised Haar basis, amputated from the scaling function (the ‘father wavelet’) in both x - and y -directions so as to comply with $\int_0^1 \psi(x, y) dx = 0$ and $\int_0^1 \psi(x, y) dy = 0$.

Then the dependence in (X, Y) is characterised by the collection $\Delta_{XY} = \{\iint \log f_{XY} \psi_{k, \ell} dx dy\}_{k, \ell}$ (Corollary 4.1). The support of $\psi_{k, \ell}$ is the square $2^{-k}(\ell + [0, 1]^2)$, which it partitions into 4 sub-squares, taking the value 2^k on the top-right and bottom-left sub-squares and the value -2^k on the top-left and bottom-right sub-squares. Thus:

$$\begin{aligned} \iint \log f_{XY} \psi_{k, \ell} dx dy &= 2^k \iint_{2^{-k}(\ell + [0, 1/2)^2)} \log f_{XY} dx dy + 2^k \iint_{2^{-k}(\ell + [1/2, 1)^2)} \log f_{XY} dx dy \\ &\quad - 2^k \iint_{2^{-k}(\ell + [0, 1/2) \times [1/2, 1))} \log f_{XY} dx dy - 2^k \iint_{2^{-k}(\ell + [1/2, 1) \times [0, 1/2))} \log f_{XY} dx dy, \end{aligned} \quad (4.12)$$

for $k \in \mathbb{N}_0$ and $\ell \in \{0, \dots, 2^k - 1\}^2$. We may recognise in (4.12) some sort of log-odds ratio reminiscent of (4.6): here the 4 relevant ‘probabilities’ are akin to geometric means of f_{XY} (i.e., average on the log-scale) over the corresponding sub-squares. As these get from coarser to finer, the complete collection $\{\iint \log f_{XY} \psi_{k,\ell} dx dy\}_{k,\ell}$ provides an appealing ‘multi-scale’ representation of dependence.

Remark 4.1. Such ‘multi-scale’ approach, through artificially created (2×2) -tables partitioning \mathcal{S}_{XY} , resonates with recent ideas of Ma and Mao (2019), Zhang (2019) and Gorsky and Ma (2022) for independence testing. Yet, those papers consider test statistics involving regular probabilities of (X, Y) belonging to any of the above sub-squares; e.g., in our notation, odds ratios like

$$\frac{\iint_{2^{-k}(\ell+[0,1/2]^2)} f_{XY} dx dy \times \iint_{2^{-k}(\ell+[1/2,1]^2)} f_{XY} dx dy}{\iint_{2^{-k}(\ell+[0,1/2] \times [1/2,1])} f_{XY} dx dy \times \iint_{2^{-k}(\ell+[1/2,1] \times [0,1/2])} f_{XY} dx dy};$$

e.g., see Zhang (2019, Definition 3.6). As opposed to (4.12), though, these quantities are not invariant under the group transformation (4.1), hence do not describe \mathfrak{D} (Corollary 4.3). \square

It so appears that $\Delta_{XY} = \{\iint \log f_{XY} \psi_{k,\ell} dx dy\}_{k,\ell}$ consists of the coefficients representing

$$\bar{\lambda}_{XY} \doteq \log f_{XY} - \int \log f_{XY} dx - \int \log f_{XY} dy + \iint \log f_{XY} dx dy \quad (4.13)$$

in the regular bivariate tensorised Haar basis. The function $\bar{\lambda}_{XY}$, called ‘centred log-odds ratio function’ in Osius (2004, 2009), is thus in one-to-one correspondence with Δ_{XY} , and is an equally valid representation of the dependence in (X, Y) . Clearly $\bar{\lambda}_{XY}$ is akin to the association parameters in log-linear parametrisations of contingency tables (Agresti, 2013, Chapter 7). For any arbitrary choice of reference point $(x_0, y_0) \in [0, 1]^2$, we can also define the ‘odds ratio function’ (Chen, 2007, 2021, Chen et al., 2015)

$$\Omega_{XY}(x, y; x_0, y_0) \doteq \frac{f_{XY}(x, y)f_{XY}(x_0, y_0)}{f_{XY}(x, y_0)f_{XY}(x_0, y)}, \quad (4.14)$$

which is in one-to-one correspondence with $\bar{\lambda}_{XY}$:

$$\begin{aligned} \bar{\lambda}_{XY}(x, y) &= \log \Omega_{XY}(x, y; x_0, y_0) - \int \log \Omega_{XY}(x, y; x_0, y_0) dx \\ &\quad - \int \log \Omega_{XY}(x, y; x_0, y_0) dy + \iint \log \Omega_{XY}(x, y; x_0, y_0) dx dy, \\ \Omega_{XY}(x, y; x_0, y_0) &= \exp(\bar{\lambda}_{XY}(x, y) - \bar{\lambda}_{XY}(x, y_0) - \bar{\lambda}_{XY}(x_0, y) + \bar{\lambda}_{XY}(x_0, y_0)). \end{aligned}$$

Hence for any $(x_0, y_0) \in [0, 1]^2$, $\Omega_{XY}(\cdot, \cdot; x_0, y_0)$ is also a valid representation of dependence.

Furthermore, if we let $k \rightarrow \infty$ in (4.12) – i.e., if we increase the resolution indefinitely – the relevant squares become infinitesimally small and the limiting log-odds ratios turn into

$$(\log f_{XY}(x, y) + \log f_{XY}(x + dx, y + dy) - \log f_{XY}(x, y + dy) - \log f_{XY}(x + dx, y)) / (dx dy).$$

If f_{XY} is mixed-differentiable, then this leads to the ‘local dependence function’ (Holland and Wang, 1987a),

viz.

$$\gamma_{XY}(x, y) \doteq \frac{\partial^2 \log f_{XY}(x, y)}{\partial x \partial y}, \quad (x, y) \in [0, 1]^2. \quad (4.15)$$

It can be verified that

$$\gamma_{XY}(x, y) = \frac{\partial^2 \log \Omega_{XY}(x, y; x_0, y_0)}{\partial x \partial y} = \frac{\partial^2 \bar{\lambda}_{XY}(x, y)}{\partial x \partial y},$$

so γ_{XY} is in one-to-one correspondence with $\Omega_{XY}(\cdot, \cdot; x_0, y_0)$ and $\bar{\lambda}_{XY}$ and is, therefore, a valid representation of dependence as well. Indeed, for rectangular supports, Wang (1993, Lemma 3.2) proved that γ_{XY} is maximal invariant under ‘marginal replacements’, which we showed to be equivalent to (4.1). This local dependence function was investigated further in Jones (1996, 1998), Jones and Koch (2003), Molenberghs and Lesaffre (1997), and it was shown that $\gamma_{XY} \equiv 0$ if and only if $X \perp Y$ (Holland and Wang, 1987a, Lemma 4.2). Clearly, γ_{XY} is the continuous analogue of the family of local (log-)odds ratios (4.6) while Ω_{XY} corresponds to the odds ratios (4.5) with respect to a pivot point.

Consider now Sklar’s representation (2.5) of the distribution π_{XY} . As the copula C is itself the distribution of an absolutely continuous vector (U, V) , its dependence may entirely be described by its odds ratio function (4.14) as well, viz.

$$\Omega_{UV}(u, v; u_o, v_o) = \frac{c(u, v)c(u_o, v_o)}{c(u_o, v)c(u, v_o)},$$

where $(u_o, v_o) \in [0, 1]^2$ is any arbitrary pivot point, and $c(u, v) \doteq \partial^2 C(u, v) / \partial u \partial v$ is the copula density. Differentiating (2.5) yields $f_{XY}(x, y) = c(F_X(x), F_Y(y))f_X(x)f_Y(y)$, which, plugged into (4.14), immediately gives

$$\Omega_{XY}(x, y; x_o, y_o) = \Omega_{UV}(F_X(x), F_Y(y); F_X(x_o), F_Y(y_o)). \quad (4.16)$$

Thus Ω_{XY} and Ω_{UV} are identical up to the usual one-to-one marginal re-labelling $u = F_X(x)$ and $v = F_Y(y)$, meaning that F_{XY} and C share the same \mathfrak{D} in agreement with Section 3.3: the change in marginal scales does not affect the inner structure. As the marginals of a copula are fixed by definition ($C \in \mathcal{F}(\mathcal{U}_{[0,1]}, \mathcal{U}_{[0,1]})$), it follows from Definition 2.1 that there is one and only one copula with a given dependence structure/odds ratio function Ω_{UV} . In view of this one-to-one correspondence, positing the copula C in (2.5) is perfectly equivalent to positing the odds ratio function Ω_{XY} of F_{XY} as this one is Ω_{UV} after proper marginal re-labelling (4.16). Thus, the dependence in (X, Y) is unequivocally described by C as anticipated in Example 2.3.

More specifically, the dependence structure of a bivariate Gaussian vector (X, Y) (Example 2.1) is thus driven by that of $(U = \Phi(\frac{X-\mu_X}{\sigma_X}), V = \Phi(\frac{Y-\mu_Y}{\sigma_Y}))$, where Φ denotes the standard normal cumulative distribution function (Probability Integral Transforms, PIT). The distribution of (U, V) is then the Gaussian

copula with parameter ρ , whose odds ratio function (4.14) is (with $u_0 = 1/2$ and $v_0 = 1/2$):

$$\Omega_{UV}(u, v; \frac{1}{2}, \frac{1}{2}) = \frac{c_\rho(u, v)c_\rho(\frac{1}{2}, \frac{1}{2})}{c_\rho(u, \frac{1}{2})c_\rho(\frac{1}{2}, v)} = \exp\left(\frac{\rho}{1-\rho^2}\Phi^{-1}(u)\Phi^{-1}(v)\right), \quad (u, v) \in [0, 1]^2.$$

This parametric form of Ω_{UV} naturally follows from the Gaussian assumption, with its exact behaviour entirely dictated by ρ (given that $\rho/(1-\rho^2)$ is a one-to-one function of ρ). Therefore, in a Gaussian vector, Pearson's correlation ρ fully describes the dependence, as announced in Example 2.1. In particular, $X \perp\!\!\!\perp Y \iff U \perp\!\!\!\perp V \iff \log \Omega_{UV} \equiv 0 \iff \rho = 0$.

We may want to write the odds ratio function on the original support of (X, Y) . Around $x_0 = \mu_X$ and $y_0 = \mu_Y$, we obtain by substituting the bivariate Gaussian density in (4.14):

$$\Omega_{XY}(x, y; \mu_X, \mu_Y) = \exp\left(\frac{\rho}{1-\rho^2}\left(\frac{x-\mu_X}{\sigma_X}\right)\left(\frac{y-\mu_Y}{\sigma_Y}\right)\right), \quad (x, y) \in \mathbb{R}^2,$$

showing the same univocal effect of ρ . It may be surprising to see the marginal parameters (μ_X, σ_X) and (μ_Y, σ_Y) appear in this representation of dependence, though. Yet, this is just the principle of equivariance exposed in Section 3.3 coming into play. In fact, for any two bimeasurable bijections Φ, Ψ , we have (similarly to (4.16)):

$$\Omega_{\Phi(X)\Psi(Y)}(x', y'; x'_o, y'_o) = \Omega_{XY}(\Phi^{-1}(x'), \Psi^{-1}(y'); \Phi^{-1}(x'_o), \Psi^{-1}(y'_o)). \quad (4.17)$$

Any arbitrary parameters may hide into Φ and Ψ , without affecting the dependence captured by Ω_{XY} . With the above choice of PIT's, Φ and Ψ involve the marginal parameters (μ_X, σ_X) and (μ_Y, σ_Y) , but this should not be confused with some interference between these parameters and the dependence.

Now, it is shown in Jones (1996) that $\gamma_{XY}(x, y)$ may be interpreted as an (appropriately scaled) localised version of Pearson's correlation around the point (x, y) – hence the name ‘local dependence’. In general, such local dependence is not uniform over the whole support \mathcal{S}_{XY} ; e.g., some distributions show ‘tail dependence’ (Joe, 2015, Section 2.13), meaning that the variables tend to be more strongly tied in their areas of extreme values than around their median point. Yet, for a Gaussian vector, we see that

$$\gamma_{XY}(x, y) = \frac{\partial^2 \log \Omega_{XY}(x, y; \mu_X, \mu_Y)}{\partial x \partial y} = \frac{\rho}{1-\rho^2} \frac{1}{\sigma_X \sigma_Y} \quad \forall (x, y) \in \mathbb{R}^2,$$

indicating that the local dependence *is* uniform over the whole of \mathbb{R}^2 . In agreement with (2.1), we can mount arbitrary \mathbb{R} -supported marginals on this specific dependence structure for creating a multitude of distributions having this trait in common with the bivariate Gaussian.

(b) $\mathcal{S}_{XY} \subseteq [0, 1]^2$ (non-rectangular support). The support \mathcal{S}_{XY} of any $\pi_{XY} \in \overline{\mathcal{P}}_{\mathfrak{X} \times \mathfrak{Y}}$ can be expressed as a countable union of rectangles (see Appendix A). Thus, at resolution high enough, the elements of a Haar basis for $\Gamma_{\mathcal{S}_{XY}}^\circ$ may cover the whole of \mathcal{S}_{XY} arbitrarily close by a rectangular pavement. In particular, if we

push the resolution to the limit $k \rightarrow \infty$, then (assuming f_{XY} is mixed-differentiable) we recover the local dependence function (4.15) on \mathcal{S}_{XY} , irrespective of the shape of that set. In agreement with Corollary 4.1, the couple $(\mathcal{S}_{XY}, \gamma_{XY})$ provides a complete characterisation of the dependence in (X, Y) . E.g., the vector (X, Y) will show quasi-independence on its support \mathcal{S}_{XY} if and only if $\gamma_{XY} \equiv 0$ on \mathcal{S}_{XY} (Corollary 4.2).

4.3.3 One (absolutely) continuous and one discrete variable

Let $\mathfrak{X} = \mathbb{N}_0 = \{0, 1, \dots\}$, $\mu_{\mathfrak{X}}$ be the counting measure, $\mathfrak{Y} = [0, 1]$ and $\mu_{\mathfrak{Y}}$ be the Lebesgue measure. Assume for simplicity that π_{XY} is supported on $\mathcal{S}_{XY} = \mathbb{N}_0 \times [0, 1]$ (rectangular support). In such a mixed vector, the density φ_{XY} takes the form

$$\varphi_{XY}(x, y) = \mathbb{P}(X = x|Y = y)f_Y(y) = f_{Y|X}(y|x)\mathbb{P}(X = x), \quad (x, y) \in \mathbb{N}_0 \times [0, 1],$$

where the notations $\mathbb{P}(X = \cdot|Y = \cdot)$, $f_{Y|X}(\cdot|x)$, $\mathbb{P}(X = \cdot)$ and $f_Y(\cdot)$ are self-evident. Ideas from Sections 4.3.1 and 4.3.2 can be combined to identify a basis for $\Gamma_{\mathcal{S}_{XY}}^{\circ}$, of the form $\{(\delta_x - \delta_{x-1})\psi_{k\ell} : x = 1, 2, \dots, k \in \mathbb{N}_0, \ell \in \{0, \dots, 2^k - 1\}\}$, where δ_x is the unit mass at $x \in \mathbb{N}_0$, and for $k \in \mathbb{N}_0, \ell \in \{0, \dots, 2^k - 1\}$, $\psi_{k\ell}(y) = 2^k \psi_0(2^k y - \ell)$ with $\psi_0(y) = \mathbb{I}_{\{y \in [0, 1/2)\}} - \mathbb{I}_{\{y \in [1/2, 1)\}}$. The dependence in (X, Y) is then characterised (Corollary 4.1) by the collection

$$\left\{ 2^k \int_{2^{-k}(\ell+[0, 1/2))} \log \frac{\mathbb{P}(X = x|Y = y)}{\mathbb{P}(X = x-1|Y = y)} dy - 2^k \int_{2^{-k}(\ell+[1/2, 1))} \log \frac{\mathbb{P}(X = x|Y = y)}{\mathbb{P}(X = x-1|Y = y)} dy \right\}, \quad (4.18)$$

for $x = 1, 2, \dots, k \in \mathbb{N}_0, \ell \in \{0, \dots, 2^k - 1\}$. If we increase arbitrarily the resolution level of the univariate Haar system $\{\psi_{k\ell}\}$ ($k \rightarrow \infty$), the elements of (4.18) become

$$\gamma_{XY}(x, y) = \frac{\partial}{\partial y} \log \frac{\mathbb{P}(X = x|Y = y)}{\mathbb{P}(X = x-1|Y = y)}, \quad (x, y) \in \{1, 2, \dots\} \times [0, 1], \quad (4.19)$$

or equivalently

$$\gamma_{XY}(x, y) = \frac{\partial}{\partial y} \log \frac{f_{Y|X}(y|x)}{f_{Y|X}(y|x-1)}, \quad (x, y) \in \{1, 2, \dots\} \times [0, 1],$$

assuming that $\mathbb{P}(X = x|Y = \cdot)$ and $f_{Y|X}(\cdot|x)$ are differentiable as functions of y for all $x \in \mathbb{N}_0$. In such a case, either of these two functions represents unequivocally the dependence in (X, Y) .

Example 4.1. Consider the simple logistic regression framework (Agresti, 2013, Chapter 4), with $X \in \{0, 1\}$ and $\mathbb{P}(X = 1|Y = y) = \exp(\alpha + \beta y)/(1 + \exp(\alpha + \beta y))$, for $\alpha, \beta \in \mathbb{R}$. With $X \in \{0, 1\}$, (4.19) reduces down to a univariate function $\gamma_{XY}(y)$, which is here constant:

$$\gamma_{XY}(y) = \frac{\partial}{\partial y} \log \exp(\alpha + \beta y) = \beta.$$

Therefore, under this model, the dependence between X and Y is captured by one single parameter, which should be any one-to-one transformation of the slope parameter β . The constancy of γ_{XY} in this case

complements the results of Jones (1998), who showed that, in continuous situations, the local dependence function (4.15) is constant in the case of a generalised linear model (GLM) with canonical link – this includes the bivariate Gaussian case of Section 4.3.2. We note that, effectively, (4.19) can be regarded as a ‘mixed’ version of (4.15).

By contrast, in the probit model $\mathbb{P}(X = 1|Y = y) = \Phi(\alpha + \beta y)$, $\gamma_{XY}(y)$ is not constant and involves both α and β , making dependence in ‘logit’ and ‘probit’ vectors different. This could be inferred directly from Definition 2.1: given that logit and probit links may be used for defining different joint distributions for (X, Y) with the same marginals $X \sim \text{Bern}(p)$ and $Y \sim \pi_Y$, the difference between both models must be distinct dependence structures. More generally, it appears that the choice of a link function in a GLM is essentially one of dependence structure between response and predictor, with the canonical link accounting for a constant local dependence entirely describable by a single parameter. \square

4.3.4 Hybrid variables

By ‘hybrid’ we mean a distribution which is a mixture of a discrete component and an absolutely continuous component. For illustration, consider the case of a vector (X, Y) where both π_X and π_Y are made up of a probability atom at 0 and an absolutely continuous part over \mathbb{R}^+ . These marginal distributions are absolutely continuous with respect to the reference measures $\mu_X = \mu_Y = \delta_0 + \lambda$ (where δ_0 is the Dirac measure at 0 and λ is the Lebesgue measure on \mathbb{R}^+). The density φ_{XY} of such a vector takes the form

$$\varphi_{XY}(x, y) = \begin{cases} p_{00} & \text{if } x = 0, y = 0 \\ p_{10}f_{X0}(x) & \text{if } x > 0, y = 0 \\ p_{01}f_{0Y}(y) & \text{if } x = 0, y > 0 \\ p_{11}f_{XY}(x, y) & \text{if } x > 0, y > 0 \end{cases},$$

where $p_{00}, p_{10}, p_{01}, p_{11}$ are probabilities summing to 1, f_{X0} is the conditional density of X given $(X > 0, Y = 0)$, f_{0Y} is the conditional density of Y given $(X = 0, Y > 0)$ and f_{XY} is the conditional density of (X, Y) given that $(X > 0, Y > 0)$. We assume that $f_{X0}(x) > 0, f_{0Y}(y) > 0$ and $f_{XY}(x, y) > 0 \forall (x, y) \in \mathbb{R}^+ \times \mathbb{R}^+$ (rectangular support). In this case, the most natural representation of the dependence in (X, Y) seems to be the odds-ratio function with respect to the pivot point $(0, 0)$:

$$\Omega_{XY}(x, y; 0, 0) = \begin{cases} \frac{p_{00}p_{11}}{p_{10}p_{01}} \frac{f_{XY}(x, y)}{f_{X0}(x)f_{0Y}(y)} & (x, y) \in \mathbb{R}^+ \times \mathbb{R}^+ \\ 1 & (x, y) \in \{x = 0\} \cup \{y = 0\} \end{cases}.$$

We recognise the usual odds ratio $(p_{00}p_{11})/(p_{10}p_{01})$ formed from the (2×2) -table $(X = 0, X > 0) \times (Y = 0, Y > 0)$, as well as a continuous contribution for $(x, y) \in \mathbb{R}^+ \times \mathbb{R}^+$.

5 Dependence versus concordance

Most of statistical theory developed from the initial works of Galton and Pearson on the Gaussian distribution (Stigler, 2002). As a result, much of the statistical jargon in use today retains a strong Gaussian connotation. For example, the fact that the dependence reduces down to the parameter ρ in a bivariate Gaussian distribution (Example 2.1) largely explains why the correlation coefficient has been referred to as a ‘dependence measure’ universally, including outside the Gaussian model. In fact, a bivariate Gaussian vector (X, Y) is so defined that any relation which may exist between X and Y must agree with ρ . This has caused, through a similar process of conflation, most of those relations to be indiscriminately labelled as ‘dependence’. Here we elaborate on the specific case of ‘concordance’.

Two variables X and Y show concordance when large (small) values of X occur typically with large (small) values of Y ; i.e., when X and Y vary in the same direction: increasing the value of one tends to increase the value of the other. Discordance is the reverse effect: X and Y vary in opposite directions (increasing one tends to decrease the other). If X and Y are concordant, then X and $-Y$ are discordant; and discordance may be regarded as ‘negative concordance’. Note that, for introducing ‘small’ and ‘large’ values for X and Y , we need to explicitly assume in this section that \mathfrak{X} and \mathfrak{Y} are ordered sets; say, \mathfrak{X} and \mathfrak{Y} are subsets of \mathbb{R} .

Concordance is formally defined through a partial ordering of the distributions of a given Fréchet class. Specifically, for $\pi_{XY}^{(1)}, \pi_{XY}^{(2)} \in \mathcal{F}(\pi_X, \pi_Y)$ with respective cdf’s $F_{XY}^{(1)}$ and $F_{XY}^{(2)}$, we say that $\pi_{XY}^{(1)}$ is ‘less concordant’ than $\pi_{XY}^{(2)}$, denoted $\pi_{XY}^{(1)} \lesssim_c \pi_{XY}^{(2)}$, when

$$F_{XY}^{(1)}(x, y) \leq F_{XY}^{(2)}(x, y), \quad \forall (x, y) \in \mathfrak{X} \times \mathfrak{Y} \quad (5.1)$$

(Scarsini, 1984, Tchen, 1980). Any numerical description of bivariate distributions which happens to be monotonic with respect to the ordering \lesssim_c can, therefore, be thought of as a measure of concordance. This is the case of Spearman’s ρ_S and Kendall’s τ , explicitly defined as a difference between probabilities of ‘concordance’ and ‘discordance’ (Hoeffding, 1947, Kruskal, 1958), viz.

$$\begin{aligned} \rho_S &= 3(\mathbb{P}((X_1 - X_2)(Y_1 - Y_3) > 0) - \mathbb{P}((X_1 - X_2)(Y_1 - Y_3) < 0)), \\ \tau &= \mathbb{P}((X_1 - X_2)(Y_1 - Y_2) > 0) - \mathbb{P}((X_1 - X_2)(Y_1 - Y_2) < 0), \end{aligned}$$

for independent copies $(X_1, Y_1), (X_2, Y_2), (X_3, Y_3) \sim \pi_{XY}$. Indeed ρ_S and τ are such that:

$$\pi_{XY}^{(1)} \lesssim_c \pi_{XY}^{(2)} \quad \Rightarrow \quad \begin{cases} \rho_S^{(1)} \leq \rho_S^{(2)} \\ \tau^{(1)} \leq \tau^{(2)} \end{cases}$$

(Tchen, 1980, Corollary 3.2). As in Section 3.3, introduce $(\mathfrak{X}', \mathcal{B}_{\mathfrak{X}'})$ and $(\mathfrak{Y}', \mathcal{B}_{\mathfrak{Y}'})$ two measurable spaces isomorphic to $(\mathfrak{X}, \mathcal{B}_{\mathfrak{X}})$ and $(\mathfrak{Y}, \mathcal{B}_{\mathfrak{Y}})$, and define two bimeasurable bijections $\Phi : \mathfrak{X} \rightarrow \mathfrak{X}'$ and $\Psi : \mathfrak{Y} \rightarrow \mathfrak{Y}'$. Then, for $k = 1, 2$, let $\pi_{\Phi(X)\Psi(Y)}^{(k)}(A' \times B') = \pi_{XY}^{(k)}(\Phi^{-1}(A') \times \Psi^{-1}(B'))$, $\forall (A', B') \in \mathcal{B}_{\mathfrak{X}'} \otimes \mathcal{B}_{\mathfrak{Y}'}$. If Φ and Ψ are order-preserving (that is, if they are increasing), then $F_{XY}(x, y) = F_{\Phi(X)\Psi(Y)}(\Phi(x), \Psi(y)) \forall (x, y) \in \mathfrak{X} \times \mathfrak{Y}$, and it is clear from (5.1) that

$$\pi_{\Phi(X)\Psi(Y)}^{(1)} \underset{c}{\succsim} \pi_{\Phi(X)\Psi(Y)}^{(2)} \iff \pi_{XY}^{(1)} \underset{c}{\succsim} \pi_{XY}^{(2)}.$$

In other words, concordance is a trait invariant to increasing transformations of the marginals. Evidently ρ_S and τ are invariant under such transformations, too.

Now, the Gaussianity of (X, Y) parameterised as in (2.2) imposes the linear relationship

$$Y = \mu_Y + \rho \frac{\sigma_Y}{\sigma_X} (X - \mu_X) + \epsilon, \quad \text{with } \epsilon \sim \mathcal{N}(0, \sigma_X^2(1 - \rho^2)).$$

Clearly, a positive (resp. negative) slope for this straight line implies concordance (resp. discordance) between X and Y . Hence the value of ρ (i.e., the dependence) directly dictates the concordance/discordance in (X, Y) – both through its sign and its absolute value, as a reduced variability for ϵ strengthens the sense of concordance/discordance between X and Y . So, it is not surprising that ρ_S and τ are one-to-one functions of ρ in this model, and thus valid representations of \mathfrak{D} (Example 2.1). That the two concepts coincide in the Gaussian case may explain why concordance is commonly understood as ‘some form’ of dependence. For example, the term ‘monotone dependence’ (‘positive’/‘negative’, for concordance/discordance) is often used, while Spearman’s ρ_S and Kendall’s τ are routinely referred to as ‘dependence measures’, in spite of specifically accounting for concordance. Yet, in general, dependence (understood through Definition 2.1) and concordance may well be unrelated.

Concordance/discordance explicitly implies an ‘influence’ of one variable on the other (‘if X increases, Y tends to increase/decrease’), hence non-independence (Section 3.2). Accordingly, $X \perp Y$ implies X and Y not in concordance/discordance. But X and Y not in concordance/discordance does not imply their independence (Scarsini, 1984, Theorem 4). Hence the same concordance status may exist with different dependence structures (e.g., ‘not concordant/discordant’ may be observed in vectors showing independence as well as in vectors showing non-independence). Reversely, the same dependence can exist with different levels of concordance, as illustrated by the following example.

Example 5.1. Let $X \sim \mathcal{U}_{[0,1]}$ and $Y = ZX$, where Z is a Rademacher random variable¹⁰ independent of X . The distribution π_{XY} is supported on the two diagonals of the unit square:¹¹ call the 4 arms of

¹⁰ $\frac{Z+1}{2} \sim \text{Bern}(1/2)$.

¹¹ The argument is clearer when examining this singular distribution, although it may not perfectly fit in the general

this cross ‘top right’, ‘bottom right’, ‘bottom left’ and ‘top left’. Due to the perfect symmetry, (X, Y) cannot show any concordance or discordance, and indeed, $\tau = 0$ in this case. Now, define π_{XY}^* through (4.1) with $\alpha_X^*(x) = 2(1 - a)\mathbb{I}_{\{x \in [0, 1/2]\}} + 2a\mathbb{I}_{\{x \in (1/2, 1]\}}$ and $\beta_Y^*(y) = 2(1 - b)\mathbb{I}_{\{y \in [0, 1/2]\}} + 2b\mathbb{I}_{\{y \in (1/2, 1]\}}$, for $a, b \in (0, 1)$. According to Theorem 4.1, this maintains the dependence $(\mathfrak{D} = \mathfrak{D}^*)$ regardless of the exact values of a and b – in particular, the support is the same. However, it can be verified that Kendall’s tau is $\tau^* = (2a - 1)(2b - 1)$ for π_{XY}^* . If both a and b are close to 1, π_{XY}^* is mostly concentrated on the top right arm of the cross, and thus shows strong concordance ($\tau^* \simeq 1$). By contrast, if a is close to 1 but b is close to 0, then π_{XY}^* is mostly concentrated on the bottom right arm, and thus shows strong discordance ($\tau^* \simeq -1$). In fact, varying the values of $a, b \in (0, 1)$, we can obtain any value of $\tau^* \in (-1, 1)$ while keeping the initial dependence \mathfrak{D} . \square

Thus, beyond the triviality ‘independence’ \Rightarrow ‘no concordance/discordance’, there is in general no link between concordance and dependence. This can be appreciated pragmatically: for defining \mathfrak{D} we could remain oblivious of the ordering in \mathfrak{X} and \mathfrak{Y} – in fact, we allowed such an ordering to be entirely reshuffled in Section 3.3 – hence concordance, exclusively based on such ordering, must be of different nature. For example, the dependence in a discrete distribution is characterised by a collection of log-odds ratios such as (4.5), which each captures how likely the specific values $X = x$ and $Y = y$ are to occur together. Clearly, these log-odd ratios can be listed in any order for covering all combinations of X - and Y -values in \mathcal{S}_{XY} . In fact, we could describe the dependence between two categorical *non-ordinal* variables by a similar collection of odds-ratios. The same observation may be made in the continuous case as well, from the odds ratio function Ω_{XY} (4.14): the assumed order on \mathfrak{X} and \mathfrak{Y} may be arbitrarily altered through one-to-one bimeasurable Lebesgue-measure preserving functions $\Phi : \mathfrak{X} \rightarrow \mathfrak{X}$ and $\Psi : \mathfrak{Y} \rightarrow \mathfrak{Y}$ while keeping the same information about the dependence *à la* (4.17). Notably, this agrees with Rényi (1959)’s views, which require valid dependence measures to be invariant under one-to-one (but not necessarily increasing) transformations of the margins (his axiom F).

Now, some specific dependence structures do dictate the concordance. For example, a vector $(X, Y) \sim \pi_{XY}$ is said to show ‘Positive Likelihood Ratio Dependence’ (PLRD) (Lehmann, 1966, Section 8) when, for all $x_1, x_2 \in \mathfrak{X}$ and $y_1, y_2 \in \mathfrak{Y}$ such that $x_1 < x_2$ and $y_1 < y_2$,

$$\varphi_{XY}(x_1, y_1)\varphi_{XY}(x_2, y_2) \geq \varphi_{XY}(x_2, y_1)\varphi_{XY}(x_1, y_2) \quad (5.2)$$

(with strict inequality for at least one pair $(x_1, y_1), (x_2, y_2)$). Evidently (5.2) remains true for the density φ_{XY}^* of any distribution π_{XY}^* satisfying (4.1), hence PLRD is truly a dependence property: in a given framework ($\pi_{XY} \notin \mathcal{P}_{\mathfrak{X} \times \mathfrak{Y}}$). An absolutely continuous version may easily be constructed, though.

equivalence class $[\mathfrak{D}]$, either all distributions show PLRD, or none does. It is clear that (5.2) induces a sense of concordance, and indeed, PLRD implies what has been called ‘Positive Quadrant Dependence’ (PQD) (Joe, 1997, Theorem 2.3), which itself implies $\rho_S > 0$ and $\tau > 0$ (Lehmann, 1966, Corollary 1). Note that PQD, defined as when $F_{XY}(x, y) \geq F_X(x)F_Y(y) \forall (x, y) \in \mathfrak{X} \times \mathfrak{Y}$, is not necessarily preserved by (4.1), hence is *not* a dependence property: within the same equivalence class $[\mathfrak{D}]$, some distributions may show PQD while others don’t, in violation of Corollary 4.3. The fact that all bivariate Gaussian distributions with $\rho \neq 0$ show either PLRD ($\rho > 0$) or its negative counterpart ($\rho < 0$) (Lehmann, 1966, Example 10) explains why dependence and concordance merge in that case. The same may be said about the bivariate Bernoulli model (Example 2.2), from Lehmann (1966, Example 11).

6 Measuring the dependence

6.1 General comments

It appears clearly from Corollary 4.1 that dependence is a polymorphic concept that cannot be entirely characterised by one single parameter in general (parametrically restricted environments such as in Examples 2.1, 2.2 and 4.1 are exceptions). This said, it often remains essential to express succinctly, by a suitable index, how strongly two variables ‘influence’ each other. This is the main motivation behind the numerous dependence measures mentioned in Section 1. Yet, most of those measures do not actually deal with dependence as delineated by Definition 2.1. For example, Section 5 described how concordance measures such as Spearman’s ρ_S or Kendall’s τ may hardly be thought of as measuring dependence, despite being non-null only in case of non-independence. In particular, $\rho_S = 0$ or $\tau = 0$ does not mean ‘no dependence’.

Failure to identify independence is not the only impediment, though. Consider the Mutual Information

$$\text{MI}(X, Y) = \mathcal{I}(\pi_{XY} \| \pi_X \times \pi_Y) = \iint \log \frac{d\pi_{XY}}{d(\pi_X \times \pi_Y)} d\pi_{XY}, \quad (6.1)$$

a popular ‘dependence measure’ owing to the fact that $\text{MI}(X, Y) = 0 \iff X \perp\!\!\!\perp Y$ (Cover and Thomas, 2006, p.28). Yet, it appears directly from (6.1) that $\text{MI}(X, Y)$ is not invariant under the group operation (4.1); meaning that it may vary from a distribution to another within the same equivalence class of dependence $[\mathfrak{D}]$. Hence the Mutual Information violates Corollary 4.3 and cannot be thought of as quantifying dependence as such. In fact, as its name suggests, $\text{MI}(X, Y)$ measures the *information* shared between X and Y , a specific notion defined by Shannon (1948) and interpreted in terms of the reduction of the uncertainty surrounding Y once we observe X (or vice-versa). While such shared information may only be null when $X \perp\!\!\!\perp Y$, this concept should not be amalgamated with dependence. For example,

Geenens and Lafaye de Micheaux (2022, p. 643) exhibit a vector (X, Y) in which X and Y are ‘nearly-independent’ while their Mutual Information is arbitrarily large, illustrating the discrepancy between the value of $\text{MI}(X, Y)$ and the underlying dependence structure.

Similar observations may be made about many other alleged ‘dependence measures’. It can reasonably be admitted that, if X and Y are independent, then ‘nothing’ may exist between them. Any index quantifying the magnitude of a trait which cannot be present between two independent variables is automatically null for $X \perp\!\!\!\perp Y$; and consequently, a non-null value for that index reveals non-independence. Yet, this does not mean that the said index relates to Definition 2.1. If it does not, calling it a ‘dependence measure’ is misleading – such an index should rather be called a ‘*non-independence indicator*’.¹²

In fact, Corollary 4.3 imposes that any valid description of the dependence in the vector (X, Y) – which includes any attempt to quantify its strength – must be based on $(\mathcal{S}_{XY}, \mathbf{\Delta}_{XY})$, and on this only. Both components \mathcal{S}_{XY} and $\mathbf{\Delta}_{XY}$ may characterise a deviation from the state of independence, irrespective of the other: if \mathcal{S}_{XY} is not rectangular, then X and Y cannot be independent even when $\mathbf{\Delta}_{XY} = \mathbf{0}$ (Section 3.4), and $\mathbf{\Delta}_{XY} \neq \mathbf{0}$ precludes independence, even on a rectangular support (Corollary 4.2). Hence a proper quantification of the overall dependence must involve two contributions: one for the effect of the shape of the support \mathcal{S}_{XY} (‘regional dependence’), the other one assessing through $\mathbf{\Delta}_{XY}$ the deviation from ‘quasi-independence’ on that support. When \mathcal{S}_{XY} is rectangular, the first contribution should be 0 and the second would then measure deviation from independence as such.

Below we explore this idea for the case of a bivariate discrete distribution on a finite support – a more thorough treatment is left for future research. So we assume here that π_{XY} is the distribution of a vector supported on $\mathcal{S}_{XY} \subseteq \mathcal{S}_X \times \mathcal{S}_Y = \{0, 1, \dots, R-1\} \times \{0, 1, \dots, S-1\}$ ($2 \leq R, S < \infty$). The distribution π_{XY} may thus be identified to a non-negative matrix of $\mathcal{M}_{R \times S}$ with entries p_{xy} summing to 1.

6.2 Regional dependence

If $\mathcal{S}_{XY} = \mathcal{S}_X \times \mathcal{S}_Y$ (rectangular support), the dependence in π_{XY} is entirely characterised by $(R-1)(S-1)$ parameters (the components of $\mathbf{\Delta}_{XY}$); see Section 4.3.1(a). By contrast, if the support is not rectangular (Section 4.3.1(b)), then the dimension of $\mathbf{\Delta}_{XY}$ is $\dim(\Gamma_{\mathcal{S}_{XY}}^\circ) < (R-1)(S-1)$, where $\Gamma_{\mathcal{S}_{XY}}^\circ$ is the subspace of $\mathcal{M}_{R \times S}$ consisting of matrices with rows and columns summing to 0 and support contained in \mathcal{S}_{XY} . It is not so much that the dependence can be described by less parameters, it is rather that the support restrictions consume some of the initial $(R-1)(S-1)$ degrees of freedom. For example, once a support like (4.7) or (4.9) is identified for a (3×3) -distribution, only one parameter suffices for completing the

¹²Note that it would be perfectly valid to use such indicator as the test statistic for independence testing.

full description of dependence. Thus, compared to the $(R - 1)(S - 1) = 4$ necessary parameters on a rectangular support of that size, the supports (4.7) and (4.9) accounts for 3 dependence parameters. If the support is (4.11), all 4 initial dependence parameters are consumed by the support constraints.

Distributions with rectangular support do not show any regional dependence, while for distributions such that Δ_{XY} is empty (i.e., $\Gamma_{\mathcal{S}_{XY}}^\circ = \{\mathbf{0}\}$), dependence is exclusively regional in nature. It seems, therefore, natural to quantify the amount of regional dependence in π_{XY} by the fraction of dependence parameters consumed by the support constraints. Thus, we define

$$R(X, Y) = \frac{(R - 1)(S - 1) - \dim(\Gamma_{\mathcal{S}_{XY}}^\circ)}{(R - 1)(S - 1)}$$

as a measure of regional dependence. Evidently this index solely reflects (through $\Gamma_{\mathcal{S}_{XY}}^\circ$) the shape of \mathcal{S}_{XY} inside $\mathcal{S}_X \times \mathcal{S}_Y$, but does not involve the distribution π_{XY} as such – in agreement with our description of regional dependence (Section 3.4). Hence $R(X, Y)$ is automatically invariant under the group transformation (4.1), which preserves supports. Also, it is clear that $R(X, Y) \in [0, 1]$, with $R(X, Y) = 0 \iff \mathcal{S}_{XY} = \mathcal{S}_X \times \mathcal{S}_Y$ and $R(X, Y) = 1 \iff \Gamma_{\mathcal{S}_{XY}}^\circ = \{\mathbf{0}\}$.

Remark 6.1. Holland and Wang (1987b) proposed a measure of regional dependence which, in the notation of this paper, would amount to $1 - (\pi_X \times \pi_Y)(\mathcal{S}_{XY})$. However, this measure explicitly depends on the marginal distributions π_X and π_Y , and fails the basic test grounded in Corollary 4.3. \square

6.3 Deviation from quasi-independence on \mathcal{S}_{XY}

Define $\Lambda_{XY} \in \mathcal{M}_{R \times S}$ as the matrix with component $(x, y) \in \{0, 1, \dots, R - 1\} \times \{0, 1, \dots, S - 1\}$ equal to¹³

$$[\Lambda_{XY}]_{xy} = \begin{cases} \log p_{xy} & \text{if } (x, y) \in \mathcal{S}_{XY} \\ 0 & \text{if } (x, y) \notin \mathcal{S}_{XY} \end{cases}.$$

Denote $d_o \doteq \dim(\Gamma_{\mathcal{S}_{XY}}^\circ)$, which we assume positive here (the dependence is not exclusively regional), and consider an arbitrary orthonormal basis $\{E_k : k = 1, \dots, d_o\}$ of $\Gamma_{\mathcal{S}_{XY}}^\circ$. Then we can write $\Delta_{XY} = (\delta_{XY;1}, \dots, \delta_{XY;d_o})$ with $\delta_{XY;k} \doteq \Lambda_{XY} \cdot E_k$ ($k = 1, \dots, d_o$) and $A \cdot B = \sum_{x,y} A_{xy} B_{xy}$ the Frobenius inner product between two matrices $A, B \in \mathcal{M}_{R \times S}$. Clearly, Δ_{XY} is the vector of coordinates of the orthogonal projection, say Λ_{XY}° , of Λ_{XY} onto $\Gamma_{\mathcal{S}_{XY}}^\circ$; that is $\Lambda_{XY}^\circ = \sum_{k=1}^{d_o} \delta_{XY;k} E_k$. If X and Y are quasi-independent on \mathcal{S}_{XY} , then $\Delta_{XY} \equiv \mathbf{0}$ (Corollary 4.2), meaning that the projection of Λ_{XY}° is the null matrix. The Frobenius norm

$$\|\Lambda_{XY}^\circ\|_F = \sqrt{\sum_{k=1}^{d_o} \delta_{XY;k}^2} = \|\Delta_{XY}\|_2 \quad (6.2)$$

¹³Here we arbitrarily set $[\Lambda_{XY}]_{xy} = 0$ for $(x, y) \notin \mathcal{S}_{XY}$, but this is inconsequential.

can therefore be thought of as quantifying how remote from quasi-independence is the dependence structure of π_{XY} on \mathcal{S}_{XY} . The norm $\|\Delta_{XY}\|_2$ is invariant to a change of (orthonormal) basis. Note that the bases $\{E_{xy}^{(00)}\}$ and $\{E_{xy}^{(\text{loc})}\}$ leading to the usual sets of odds ratios (Section 4.3.1) are not orthonormal, but orthonormal versions may easily be obtained via Gram-Schmidt. Other ‘interpretable’ bases may be constructed as in Egozcue et al. (2015) and Fačevićová et al. (2018).

Example 6.1. Consider a bivariate Bernoulli distribution with full support ($p_{xy} > 0 \forall(x, y)$) as in Example 2.2. Here $d_o = 1$ with (normalised) basis element $\frac{1}{2}\{e_{00} + e_{11} - e_{10} - e_{01}\}$, in the notation of Section 4.3.1. Hence $\Delta_{XY} = \log \frac{\sqrt{p_{00}p_{11}}}{\sqrt{p_{10}p_{01}}} = \frac{1}{2} \log \omega$, and $\|\Delta_{XY}\|_2 = \frac{1}{2} |\log \omega|$. For ease of interpretation, it may be desirable to transform monotonically this value to $[0, 1]$. E.g., we could take $\tanh(\|\Delta_{XY}\|_2/2) = |\Upsilon| \in [0, 1]$, where Υ is Yule’s colligation coefficient (Yule, 1912) (see Geenens (2020, Section 5.5) for reasons why this choice may be natural). The dependence is properly quantified by an absolute value, but characterising it entirely requires the signed number Υ : the distribution cannot be fully identified from $p_{\bullet 1}, p_{1 \bullet}$ and $|\Upsilon|$ only. \square

When $d_o > 1$, $\|\Delta_{XY}\|_2$ provides a single-number quantification of the magnitude of the multiple entries which characterise the part of the dependence of π_{XY} which is not regional. We may want to consider an increasing transformation $T: \mathbb{R}^+ \rightarrow [0, 1]$, with $T(0) = 0$, and propose

$$Q(X, Y) = T(\|\Delta_{XY}\|_2) \in [0, 1] \tag{6.3}$$

as a measure of deviation from quasi-independence on \mathcal{S}_{XY} . As a function of Δ_{XY} only, $Q(X, Y)$ is invariant under the group transformation (4.1), and clearly, $Q(X, Y) = 0 \iff X$ and Y are quasi-independent. The transformation T is arbitrary and only aims at calibration.

6.4 An overall measure of dependence

Combining the previous two contributions, one may suggest the following overall measure of dependence:

$$D(X, Y) = R(X, Y) + (1 - R(X, Y))Q(X, Y). \tag{6.4}$$

This is the weighted average of $Q(X, Y)$ and 1 with weights $1 - R(X, Y)$ and $R(X, Y)$, respectively. In a sense, this amounts to setting the ‘missing’ entries of Δ_{XY} (those dependence parameters consumed by the support constraints) to their maximal value.

This may be justified by referring to Example 6.1. As all probabilities p_{xy} are positive, the support is rectangular so that $R(X, Y) = 0$ and $D(X, Y) = Q(X, Y) = |\Upsilon|$ (using the above suggested transformation $T(\cdot) = \text{atanh}(\cdot/2)$). If one of the probabilities p_{xy} approaches 0 (while staying positive), then $\log \omega \rightarrow \pm\infty$ and $|\Upsilon| \rightarrow 1$, so that $D(X, Y) \rightarrow 1$. Now, suppose that one of the probabilities p_{xy} is effectively 0: then

$\Gamma_{XY}^\circ = \{\mathbf{0}\}$, and $R(X, Y) = 1$. The induced dependence is entirely regional, and consequently $D(X, Y) = 1$, too. The proposed measure (6.4) is, therefore, continuous in the probabilities p_{xy} , including at 0.

Further properties of $D(X, Y)$ easily follow from those of $R(X, Y)$ and $Q(X, Y)$. In particular, $D(X, Y)$ is invariant under the group transformation (4.1), as both $R(X, Y)$ and $Q(X, Y)$ are, and hence properly describes dependence: $D(X, Y)$ is constant over any class $[\mathfrak{D}]$. Also, $D(X, Y)$ fulfils the relevant requirements of Rényi (1959), qualifying valid dependence measures:

- (A) $D(X, Y)$ is defined for any pair of random variables X and Y , neither of them being constant with probability 1;
- (B) $D(X, Y) = D(Y, X)$;
- (C) $0 \leq D(X, Y) \leq 1$;
- (D) $D(X, Y) = 0 \iff X \perp\!\!\!\perp Y$;
- (E) $D(X, Y) = 1$ if there is a strict dependence between X and Y , i.e., either $X = f(Y)$ or $Y = g(X)$, where f and g are Borel-measurable functions;
- (F) If the Borel-measurable functions f and g map the real axis in a one-to-one way onto itself, $D(f(X), g(Y)) = D(X, Y)$;

Properties (A),¹⁴ (B) and (C) are obvious. Property (D) follows as $D(X, Y) = 0 \iff R(X, Y) = 0$ and $Q(X, Y) = 0 \iff \mathcal{S}_{XY} = \mathcal{S}_X \times \mathcal{S}_Y$ and $\Delta_{XY} = 0 \iff X \perp\!\!\!\perp Y$, by Corollary 4.2. What Rényi (1959) called ‘strict dependence’ in (E) here reduces to the situation where there is exactly one positive probability p_{xy} for each $x \in \mathcal{S}_X$ if $R \geq S$ or exactly one positive probability p_{xy} for each $y \in \mathcal{S}_Y$ if $S \geq R$ (if $R = S$ then there is a one-to-one relationship between X and Y). In that case $\Gamma_{XY}^\circ = \{\mathbf{0}\}$, hence $R(X, Y) = 1$ and $D(X, Y) = 1$. Property (F) was mentioned in Section 5, and in terms of $D(X, Y)$ is reflected by the fact that $\dim(\Gamma_{\mathcal{S}_{XY}}^\circ)$ and hence $R(X, Y)$ are invariant under permutations of the row and/or columns of π_{XY} , and $Q(X, Y)$ is invariant to changes of the (orthonormal) basis $\{\gamma_1, \gamma_2, \dots\}$ (which also covers permutations of the row and/or columns of π_{XY}). Note that Rényi (1959)’s last requirement is:

- (G) If the joint distribution of X and Y is normal, then $D(X, Y) = |\rho|$, where ρ is Pearson’s correlation between X and Y .

Evidently this does not apply to the above discrete setting. This said, a continuous analogue of $Q(X, Y) = D(X, Y)$ (as the support of the bivariate Gaussian is rectangular, we would have $R(X, Y) = 0$) may easily be devised. Like in Section 4.3.2(a), consider the representation of a bivariate Gaussian distribution with

¹⁴Admittedly we only consider discrete variables, here.

correlation $\rho \in (-1, 1)$ as the Gaussian copula density c_ρ on $[0, 1]^2$. With the ‘amputated’ Haar basis considered in Section 4.3.2 (which happens to be orthonormal in $L_2([0, 1]^2)$), the entries of $\mathbf{\Delta}_{UV}$ are $\delta_{UV;k\ell} = \iint \log c_\rho \psi_{k\ell} du dv$, for $k \in \mathbb{N}_0$ and $\ell \in \{0, \dots, 2^k - 1\}^2$. Hence the continuous analogue to (6.2) would be

$$\|\mathbf{\Delta}_{UV}\|_2^2 = \iint_{[0,1]^2} \bar{\lambda}_\rho^2 du dv,$$

where, similarly to (4.13), we have

$$\bar{\lambda}_\rho \doteq \log c_\rho - \int \log c_\rho du - \int \log c_\rho dv + \iint \log c_\rho du dv,$$

the centred log-odds ratio function of the copula density c_ρ . As $\log c_\rho \in L_2([0, 1]^2)$ for all $\rho \in (-1, 1)$, $\|\mathbf{\Delta}_{UV}\|_2 < \infty$, and can be verified to be an increasing function of $|\rho|$, indeed. With the appropriate transformation T in (6.3), we may make it happen that $D(U, V) = Q(U, V) = |\rho|$, fulfilling Rényi (1959)’s postulate (G). This is similar to some ideas recently explored in Genest et al. (2022).

7 Conclusion

Close to 50 years ago, Mosteller and Tukey (1977, p.262) already regretted the ‘troublesome, if not dangerous, confusion’ surrounding the usage of the word ‘dependence’ in statistics. The comment remains very topical, as ‘dependence’ and its declension continue to be used confusedly for referring to both the state of ‘non-independence’ of two variables *and* the quantifiable strength of the mutual influence between them necessarily induced by that state. If the former unambiguously describes the negative of the precisely defined ‘independence’ concept, a proper definition of the latter seems to be lacking. As coherently attacking many important problems requires the dependence between variables to be plainly understood as a specific structure of which we should be able to provide a precise and nuanced description, the lack of any clear framework around this leads to undesirable looseness and subjectivity when addressing essential questions. In response, this paper proposes a general definition for the dependence between two random variables defined on the same probability space.

If one is willing to accept that definition (Definition 2.1), then a cascade of implications establishes that the dependence \mathfrak{D} in a bivariate distribution π_{XY} admits a general representation (Corollary 4.1), ‘universal’ in the sense that it is valid irrespective of the nature of the variables X and Y at play. This representation consists of two elements. The first one accounts for ‘regional dependence’, the contribution to \mathfrak{D} due to a non-rectangular joint support for π_{XY} , reflecting the incompatibility between some values of X and some values of Y . The second one expresses in which way specific values of X are more or less likely to occur with specific values of Y on that support, and can be generally described as an ‘odds-ratio-like’ object (or

equivalent). The universal representation allows unequivocal identification of the proper way of capturing the dependence in particular situations of interest. Known results are recovered in familiar cases, but novel perspectives arise for less well-studied ones.

The proposed framework opens up numerous avenues for future research. For example, here we have considered exclusively bivariate distributions, that is, we focused on the dependence between two scalar variables. Though, Definition 2.1 can be extended naturally to the very general situation of a collection of random objects for which we may define a joint distribution. This would allow the identification of dependence between an arbitrary number of random vectors, for example. Also, Section 6 explored the idea of a dependence measure based on the obtained representation – thus being consistent with the given definition of dependence – in the simplest case of a bivariate discrete distribution with finite support. An important question is how to extend this to make it more ‘universal’.

Acknowledgments

The author thanks Pierre Lafaye de Micheaux (UNSW Sydney, Australia) and Ivan Kojadinovic (Université de Pau et des Pays de l’Adour, France) for useful comments and discussions.

References

- Agresti, A., 2013. *Categorical data analysis*, 3rd Edition. John Wiley & Sons.
- Albiac, F., Kalton, N., 2016. *Topics in Banach Space Theory*. Graduate Texts in Mathematics. Springer International Publishing.
- Arnold, B. C., Castillo, E., Sarabia, J. M., 2001. Conditionally specified distributions: an introduction. *Statistical Science* 16 (3), 249–274.
- Arnold, B. C., Press, S. J., 1989. Compatible conditional distributions. *Journal of the American Statistical Association* 84 (405), 152–156.
- Arnold, B. C., Sarabia, J. M., 2022. Conditional specification of statistical models: Classical models, new developments and challenges. *Journal of Multivariate Analysis* 188.
- Barndorff-Nielsen, O., 1978. *Information and Exponential Families*. Wiley, New York.
- Bishop, Y. M., Fienberg, S. E., Holland, P. W., 1975. *Discrete multivariate analysis: theory and practice*. Springer Science & Business Media.
- Caussinus, H., 1965. Contribution à l’analyse statistique des tableaux de corrélation. In: *Annales de la Faculté des sciences de Toulouse: Mathématiques*. Vol. 29. pp. 77–183.
- Chen, H. Y., 2007. A semiparametric odds ratio model for measuring association. *Biometrics* 63 (2), 413–421.
- Chen, H. Y., 2021. *Semiparametric Odds Ratio Model and Its Applications*. Chapman and Hall/CRC.

- Chen, H. Y., Rader, D. E., Li, M., 2015. Likelihood inferences on semiparametric odds ratio model. *Journal of the American Statistical Association* 110 (511), 1125–1135.
- Cover, T., Thomas, J., 2006. Elements of Information Theory (2nd Edition). Wiley, New York.
- Cramer, E., 2000. Probability measures with given marginals and conditionals: I-projections and conditional Iterative Proportional Fitting. *Statistics & Risk Modeling* 18 (3), 311–330.
- Craven, B., 1978. Mathematical Programming and Control Theory. Chapman and Hall Mathematics Series. Springer Netherlands.
- Csiszár, I., 1975. I-divergence geometry of probability distributions and minimization problems. *The Annals of Probability* (3), 146–158.
- Dawid, A. P., 1979. Conditional independence in statistical theory. *Journal of the Royal Statistical Society: Series B (Methodological)* 41 (1), 1–15.
- Deming, W. E., Stephan, F. F., 1940. On a least squares adjustment of a sampled frequency table when the expected marginal totals are known. *The Annals of Mathematical Statistics* 11 (4), 427–444.
- Dodge, Y., 2006. The Oxford Dictionary of Statistical Terms. Oxford University Press.
- Drouot Mari, D., Kotz, S., 2001. Correlation and Dependence. Imperial College Press.
- Dudley, R., 2002. Real Analysis and Probability. Cambridge Studies in Advanced Mathematics. Cambridge University Press.
- Edwards, A. W., 1963. The measure of association in a 2×2 table. *Journal of the Royal Statistical Society: Series A (General)* 126 (1), 109–114.
- Egozcue, J. J., Pawlowsky-Glahn, V., Templ, M., Hron, K., 2015. Independence in contingency tables using simplicial geometry. *Communications in Statistics - Theory and Methods* 44 (18), 3978–3996.
- Everitt, B., Skrondal, A., 2010. The Cambridge Dictionary of Statistics. Cambridge University Press.
- Fačevicová, K., Hron, K., Todorov, V., Templ, M., 2018. General approach to coordinate representation of compositional tables. *Scandinavian Journal of Statistics* 45 (4), 879–899.
- Fienberg, S. E., 1970. An iterative procedure for estimation in contingency tables. *The Annals of Mathematical Statistics* 41 (3), 907–917.
- Folland, G., 2013. Real Analysis: Modern Techniques and Their Applications. Pure and Applied Mathematics: A Wiley Series of Texts, Monographs and Tracts. Wiley.
- Fréchet, M., 1951. Sur les tableaux de corrélation dont les marges sont données. *Annals of the University of Lyon, Section A* 14.
- Geenens, G., 2020. Copula modeling for discrete random vectors. *Dependence Modeling* 8 (1), 417–440.
- Geenens, G., Lafaye de Micheaux, P., 2022. The Hellinger correlation. *Journal of the American Statistical Association* (117), 639–653.
- Genest, C., Hron, K., Nešlehová, J. G., 2022. Orthogonal decomposition of multivariate densities in Bayes spaces and its connection with copulas. *arXiv preprint arXiv:2206.13898*.
- Genest, C., Nešlehová, J., 2007. A primer on copulas for count data. *ASTIN Bulletin* 37, 475–515.
- Genest, C., Nešlehová, J., Rémillard, B., 2014. On the empirical multilinear copula process for count data. *Bernoulli* (20), 1344–1371.
- Good, I., 1956. On the estimation of small frequencies in contingency tables. *Journal of the Royal Statistical Society: Series B (Methodological)* 18 (1), 113–124.
- Goodman, L. A., 1968. The analysis of cross-classified data: Independence, quasi-independence, and interactions in contingency

- tables with or without missing entries. *Journal of the American Statistical Association* 63 (324), 1091–1131.
- Goodman, L. A., 1996. A single general method for the analysis of cross-classified data: Reconciliation and synthesis of some methods of Pearson, Yule, and Fisher, and also some methods of correspondence analysis and association analysis. *Journal of the American Statistical Association* 91 (433), 408–428.
- Gorsky, S., Ma, L., 2022. Multi-scale Fisher’s independence test for multivariate dependence. *Biometrika* 109 (3), 569–587.
- Hoeffding, W., 1940. Masstabinvariante korrelationstheorie. *Schriften des Mathematischen Instituts und des Instituts für Angewandte Mathematik der Universität Berlin* 5 (3), 181–233.
- Hoeffding, W., 1947. On the distribution of the rank correlation coefficient τ when the variates are not independent. *Biometrika* 34 (3-4), 183–196.
- Holland, P. W., Wang, Y. J., 1987a. Dependence function for continuous bivariate densities. *Communications in Statistics - Theory and Methods* 16 (3), 863–876.
- Holland, P. W., Wang, Y. J., 1987b. Regional dependence for continuous bivariate densities. *Communications in Statistics - Theory and Methods* 16 (1), 193–206.
- Idel, M., 2016. A review of matrix scaling and Sinkhorn’s normal form for matrices and positive maps. *arXiv e-prints arXiv:1609.06349*.
- Ireland, C. T., Kullback, S., 1968. Contingency tables with given marginals. *Biometrika* 55 (1), 179–188.
- Joe, H., 1997. *Multivariate Models and Multivariate Dependence Concepts*. Chapman & Hall/CRC. Taylor & Francis.
- Joe, H., 2015. *Dependence Modeling with Copulas*. Chapman and Hall/CRC.
- Jones, M. C., 1996. The local dependence function. *Biometrika* 83 (4), 899–904.
- Jones, M. C., 1998. Constant local dependence. *Journal of Multivariate Analysis* 64 (2), 148–155.
- Jones, M. C., Koch, I., 2003. Dependence maps: Local dependence in practice. *Statistics and Computing* 13 (3), 241–255.
- Kendall, M., 1955. *Rank Correlation Methods. Theory & Applications of Rank Order-Statistics*. C. Griffin.
- Kendall, M. G., Buckland, W. R., 1971. *A dictionary of statistical terms*, 3rd Edition (revised).
- Kendall, M. G., Sundrum, R., 1953. Distribution-free methods and order properties. *Revue de l’Institut International de Statistique* 21, 124–134.
- Kruskal, W. H., 1958. Ordinal measures of association. *Journal of the American Statistical Association* 53 (284), 814–861.
- Kullback, S., 1968. Probability densities with given marginals. *The Annals of Mathematical Statistics* 39 (4), 1236–1243.
- Lehmann, E., Casella, G., 2006. *Theory of Point Estimation*. Springer Texts in Statistics. Springer New York.
- Lehmann, E. L., 1966. Some concepts of dependence. *The Annals of Mathematical Statistics* 37 (5), 1137–1153.
- Ma, L., Mao, J., 2019. Fisher exact scanning for dependency. *Journal of the American Statistical Association* 114 (525), 245–258.
- Marshall, A., 1996. Copulas, marginals and joint distributions. In: Rüschendorf, L., Schweizer, B., Taylor, M. (Eds.), *Distributions with Fixed Marginals and Related Topics*. Institute of Mathematical Statistics, pp. 213–222.
- Molenberghs, G., Lesaffre, E., 1997. Non-linear integral equations to approximate bivariate densities with given marginals and dependence function. *Statistica Sinica*, 713–738.
- Mosteller, F., 1968. Association and estimation in contingency tables. *Journal of the American Statistical Association* 63 (321), 1–28.
- Mosteller, F., Tukey, J., 1977. *Data Analysis and Regression: A Second Course in Statistics*. Addison-Wesley Publishing Company.
- Nussbaum, R. D., 1993. Entropy minimization, Hilbert’s projective metric, and scaling integral kernels. *Journal of Functional*

- Analysis* 115 (1), 45–99.
- Osius, G., 2004. The association between two random elements: A complete characterization and odds ratio models. *Metrika* 60 (3), 261–277.
- Osius, G., 2009. Asymptotic inference for semiparametric association models. *The Annals of Statistics* 37 (1), 459–489.
- Plackett, R. L., 1965. A class of bivariate distributions. *Journal of the American Statistical Association* 60 (310), 516–522.
- Rényi, A., 1959. On measures of dependence. *Acta mathematica hungarica* 10 (3-4), 441–451.
- Rudas, T., 2018. Lectures on Categorical Data Analysis. Springer.
- Rudin, W., 1991. Functional Analysis. International series in pure and applied mathematics. McGraw-Hill.
- Rüschendorf, L., 1995. Convergence of the iterative proportional fitting procedure. *The Annals of Statistics* 23 (4), 1160–1174.
- Rüschendorf, L., Thomsen, W., 1993. Note on the Schrödinger equation and I-projections. *Statistics and Probability Letters* 17 (5), 369–375.
- Scarsini, M., 1984. On measures of concordance. *Stochastica* 8 (3), 201–218.
- Schweizer, B., Sklar, A., 1974. Operations on distribution functions not derivable from operations on random variables. *Studia Mathematica* 52 (1), 43–52.
- Schweizer, B., Wolff, E. F., 1981. On nonparametric measures of dependence for random variables. *The Annals of Statistics* 9 (4), 879–885.
- Shannon, C. E., 1948. A mathematical theory of communication. *The Bell system technical journal* 27 (3), 379–423.
- Sinkhorn, R., 1964. A relationship between arbitrary positive matrices and doubly stochastic matrices. *The Annals of Mathematical Statistics* 35 (2), 876–879.
- Sklar, M., 1959. Fonctions de répartition à n dimensions et leurs marges. *Publ. inst. statist. univ. Paris* 8, 229–231.
- Stigler, S., 2002. *Statistics on the Table: The History of Statistical Concepts and Methods*. Harvard University Press.
- Stuart, A., 1953. The estimation and comparison of strengths of association in contingency tables. *Biometrika* 40 (1/2), 105–110.
- Tchen, A. H., 1980. Inequalities for distributions with given marginals. *The Annals of Probability* 8 (4), 814–827.
- Tjøstheim, D., Otneim, H., Støve, B., 2022. Statistical dependence: Beyond Pearson’s ρ . *Statistical Science* 37 (1), 90–109.
- Upton, G., Cook, I., 2014. *A Dictionary of Statistics (3e Edition)*. Oxford Paperback Reference. OUP Oxford.
- Wang, Y. J., 1993. Construction of continuous bivariate density functions. *Statistica Sinica* 3 (1), 173–187.
- Wang, Y. J., Ip, E. H., 2008. Conditionally specified continuous distributions. *Biometrika* 95 (3), 735–746.
- Webster, R., 1994. *Convexity*. Oxford science publications. Oxford University Press.
- Wermuth, N., Cox, D., 2005. Statistical dependence and independence. *Encyclopedia of Biostatistics* 7, 4260–4264.
- Yule, G. U., 1912. On the methods of measuring association between two attributes. *Journal of the Royal Statistical Society* 75 (6), 579–652.
- Založnik, M., 2011. Iterative proportional fitting: Theoretical synthesis and practical limitations. Ph.D. thesis, University of Liverpool.
- Zhang, K., 2019. BET on independence. *Journal of the American Statistical Association* 114 (528), 1620–1637.

A Definition of the set $\overline{\mathcal{P}}_{\mathfrak{X} \times \mathfrak{Y}}$

Let $\mathcal{P}_{\mathfrak{X} \times \mathfrak{Y}}$ be the set of all probability distributions π_{XY} on $\mathfrak{X} \times \mathfrak{Y}$ such that $\pi_{XY} \ll \mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}}$. Let $\varphi_{XY} = d\pi_{XY}/d(\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}})$ be the corresponding density, \mathcal{S}_{XY} , \mathcal{S}_X and \mathcal{S}_Y be the joint and marginal supports, as described in Section 3.1. The subset $\overline{\mathcal{P}}_{\mathfrak{X} \times \mathfrak{Y}}$ consists of the distributions $\pi_{XY} \in \mathcal{P}_{\mathfrak{X} \times \mathfrak{Y}}$ such that:

- (1) $\varphi_{XY} \in L_{\infty}(\mathfrak{X} \times \mathfrak{Y}, \mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}})$;
- (2) there exist rectangles $\{(E_i, F_i) \in \mathcal{B}_{\mathfrak{X}} \otimes \mathcal{B}_{\mathfrak{Y}}\}_{i=1}^m$, $\{(G_j, H_j) \in \mathcal{B}_{\mathfrak{X}} \otimes \mathcal{B}_{\mathfrak{Y}}\}_{j=1}^n$ of positive $(\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}})$ -measure, such that $\mathcal{S}_X = \bigcup_{i=1}^m E_i$ and $\mathcal{S}_Y = \bigcup_{j=1}^n G_j$ (up to sets of measure 0); and a constant $c > 0$ such that $\varphi_{XY}(x, y) \geq c$ $(\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}})$ -a.e. over $\bigcup_{i=1}^m (E_i \times F_i)$ and $\bigcup_{j=1}^n (G_j \times H_j)$;
- (3) there exists a sequence of measurable sets $K_j \in \mathcal{B}_{\mathfrak{X}} \otimes \mathcal{B}_{\mathfrak{Y}}$ ($j = 1, 2, \dots$) and a sequence of scalars $\varepsilon_j > 0$ such that $\lim_{j \rightarrow \infty} \varepsilon_j = 0$, with
 - (a) $\varphi_{XY} > \varepsilon_j$ $(\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}})$ -a.e. over K_j ;
 - (b) K_j is a countable union of measurable rectangles;
 - (c) $\mathcal{S}_{XY} = \bigcup_{j=1}^{\infty} K_j$.

These basic requirements are reasonably mild. For example, they are automatically fulfilled if \mathcal{S}_X and \mathcal{S}_Y are compact and φ_{XY} is continuous (Nussbaum, 1993, Theorem 2.34) – this is the case of any discrete distribution with finite support, among others. In our context, the above constraints may even be thought of as being much milder, in the following sense: we are actually allowed flexibility for choosing the working density φ_{XY} . Indeed we are free to pick any convenient ‘canonical’ marginal supports and any one-to-one mappings Φ and Ψ for moving to those supports (Section 3.3), producing any hand-picked ‘nice’ marginal distributions π_X and π_Y which may be convenient. In addition, the measures $\mu_{\mathfrak{X}}$ and $\mu_{\mathfrak{Y}}$ (Section 3.1) are left free, too; so that the density itself may be taken with respect to any convenient reference. In fact, Corollary 2.31 and Remark 3.14 in Nussbaum (1993) guarantee that, if Lemma 3.2 is valid for a distribution π_{XY} , then it would also hold true for any distribution π_{XY}^* such that $\pi_{XY}^* \approx \pi_{XY}$ and $\varphi_{XY}^* \in L_{\infty}(\mathfrak{X} \times \mathfrak{Y}, \mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}})$. So, all what is required is to find a joint measure equivalent to (some marginally-transformed version of) π_{XY} with a density with respect to ‘some’ product measure satisfying the requirements. This flexibility is justified by the fact that the above choices do not disturb the dependence of a given distribution.

B Proofs of the main results

Proof of Lemma 3.1

Any distribution $\pi_{\mathfrak{X}\times\mathfrak{Y}}^*$ in $\mathcal{P}_{\mathfrak{X}\times\mathfrak{Y}} \cap \Sigma_{\pi_{\mathfrak{X}\mathfrak{Y}}} \cap \mathcal{F}(\pi_{\mathfrak{X}}^*, \pi_{\mathfrak{Y}}^*)$ would have a density $\varphi_{\mathfrak{X}\mathfrak{Y}}^* = d\pi_{\mathfrak{X}\mathfrak{Y}}^*/d(\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}})$ such that

$$\begin{cases} \int \varphi_{\mathfrak{X}\mathfrak{Y}}^* d\mu_{\mathfrak{Y}} = \varphi_{\mathfrak{X}}^* & (\pi_{\mathfrak{X}}\text{-a.e.}) \\ \int \varphi_{\mathfrak{X}\mathfrak{Y}}^* d\mu_{\mathfrak{X}} = \varphi_{\mathfrak{Y}}^* & (\pi_{\mathfrak{Y}}\text{-a.e.}) \\ \iint \varphi_{\mathfrak{X}\mathfrak{Y}}^* \mathbb{I}_{\{\mathcal{S}_{\mathfrak{X}\mathfrak{Y}}^c\}} d(\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}}) = 0 \end{cases}, \quad (\text{B.1})$$

where $\varphi_{\mathfrak{X}}^* = d\pi_{\mathfrak{X}}^*/d\mu_{\mathfrak{X}}$, $\varphi_{\mathfrak{Y}}^* = d\pi_{\mathfrak{Y}}^*/d\mu_{\mathfrak{Y}}$ and $\mathcal{S}_{\mathfrak{X}\mathfrak{Y}}^c$ is the complement in $\mathfrak{X} \times \mathfrak{Y}$ of the support $\mathcal{S}_{\mathfrak{X}\mathfrak{Y}}$ of $\pi_{\mathfrak{X}\mathfrak{Y}}$. Define the closed convex cone $C = \{\varphi \in L_1(\mathfrak{X} \times \mathfrak{Y}, \mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}}) : \varphi \geq 0 \text{ a.e.}\} \subset L_1(\mathfrak{X} \times \mathfrak{Y}, \mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}})$, and let \mathcal{A} be the linear operator from $\Xi \doteq L_1(\mathfrak{X} \times \mathfrak{Y}, \mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}})$ to $\Upsilon \doteq (L_1(\mathfrak{X}, \mu_{\mathfrak{X}}), L_1(\mathfrak{X}, \mu_{\mathfrak{X}}), L_1(\mathfrak{Y}, \mu_{\mathfrak{Y}}), L_1(\mathfrak{Y}, \mu_{\mathfrak{Y}}), \mathbb{R})$ defined as

$$\mathcal{A}\varphi = \left(\int \varphi d\mu_{\mathfrak{Y}}, -\int \varphi d\mu_{\mathfrak{Y}}, \int \varphi d\mu_{\mathfrak{X}}, -\int \varphi d\mu_{\mathfrak{X}}, \iint \varphi \mathbb{I}_{\{\mathcal{S}_{\mathfrak{X}\mathfrak{Y}}^c\}} d(\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}}) \right).$$

The existence of a solution to (B.1) means that $\mathbf{b} \doteq (\varphi_{\mathfrak{X}}^*, -\varphi_{\mathfrak{X}}^*, \varphi_{\mathfrak{Y}}^*, -\varphi_{\mathfrak{Y}}^*, 0) \in \Upsilon$ belongs to the (closed convex) cone $\mathcal{A}(C) \doteq \{v \in \Upsilon : \exists \varphi \in C \text{ s.t. } v = \mathcal{A}\varphi\}$. Following Farkas lemma (see, e.g., Craven (1978, Theorem 2.2.6)), we have the equivalence

$$\mathbf{b} \in \mathcal{A}(C) \iff (u \in \Upsilon', \mathcal{A}^T u \in C' \implies u(\mathbf{b}) \geq 0), \quad (\text{B.2})$$

where Υ' is the dual of Υ , \mathcal{A}^T is the adjoint operator of \mathcal{A} and C' is the dual cone of C , i.e., $C' = \{\xi \in \Xi' : \xi(\varphi) \geq 0 \forall \varphi \in C\}$, with Ξ' the dual of Ξ .

Now, $\mathcal{A}^T u \in C' \iff (\mathcal{A}^T u)(\varphi) \geq 0$ for all $\varphi \in C \iff u(\mathcal{A}\varphi) \geq 0$ for all $\varphi \in C$, where

$$\begin{aligned} u(\mathcal{A}\varphi) &= \iint a^{(+)} \varphi d\mu_{\mathfrak{Y}} d\mu_{\mathfrak{X}} - \iint a^{(-)} \varphi d\mu_{\mathfrak{Y}} d\mu_{\mathfrak{X}} \\ &\quad + \iint b^{(+)} \varphi d(\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}}) - \iint b^{(-)} \varphi d(\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}}) \\ &\quad + k \iint_{\mathcal{S}_{\mathfrak{X}\mathfrak{Y}}^c} \varphi d(\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}}) \end{aligned}$$

for some $a^{(+)}, a^{(-)} \in L_{\infty}(\mathfrak{X}, \mu_{\mathfrak{X}})$, $b^{(+)}, b^{(-)} \in L_{\infty}(\mathfrak{Y}, \mu_{\mathfrak{Y}})$ and $k \geq 0$. Setting $a = a^{(+)} - a^{(-)}$, $b = b^{(+)} - b^{(-)}$, we have

$$u(\mathcal{A}\varphi) = \iint (a + b + k \mathbb{I}_{\{\mathcal{S}_{\mathfrak{X}\mathfrak{Y}}^c\}}) \varphi d(\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}}).$$

Likewise,

$$u(\mathbf{b}) = \int a \varphi_{\mathfrak{X}}^* d\mu_{\mathfrak{X}} + \int b \varphi_{\mathfrak{Y}}^* d\mu_{\mathfrak{Y}}.$$

(a) $\mathcal{P}_{\mathfrak{X}\times\mathfrak{Y}} \cap \Sigma_{\pi_{\mathfrak{X}\mathfrak{Y}}} \cap \mathcal{F}(\pi_{\mathfrak{X}}^*, \pi_{\mathfrak{Y}}^*) = \emptyset$ means that there is no solution to (B.1), that is, $\mathbf{b} \notin \mathcal{A}(C)$, which by

(B.2) is equivalent to the existence of some $u \in \Upsilon'$ such that $\mathcal{A}^T u \in C'$ and $u(\mathbf{b}) < 0$. From above,

$$\begin{aligned} \mathcal{A}^T u \in C' &\iff \iint (a + b + k \mathbb{I}_{\{\mathcal{S}_{XY}^c\}}) \varphi \, d(\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}}) \geq 0 \quad \forall \varphi \in C \\ &\iff a(x) + b(y) + k \mathbb{I}_{\{(x,y) \notin \mathcal{S}_{XY}\}} \geq 0 \quad (\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}})\text{-a.e.} \end{aligned} \quad (\text{B.3})$$

while

$$u(\mathbf{b}) < 0 \iff \int a \varphi_X^* \, d\mu_{\mathfrak{X}} + \int b \varphi_Y^* \, d\mu_{\mathfrak{Y}} < 0. \quad (\text{B.4})$$

Let X^* (resp. Y^*) be a random variable defined on \mathfrak{X} (resp. \mathfrak{Y}) with distribution π_X^* (resp. π_Y^*), and define $U = a(X^*)$ and $V = -b(Y^*)$, for a, b two functions satisfying (B.3) and (B.4). We have:

$$\int_{-\infty}^{\infty} (\mathbb{P}(U > t) - \mathbb{P}(V > t)) \, dt = \mathbb{E}(U) - \mathbb{E}(V) = \int a \, d\pi_X^* + \int b \, d\pi_Y^* < 0$$

(from (B.4)). Thus $\mathfrak{T} \doteq \{t \in \mathbb{R} : \mathbb{P}(U > t) - \mathbb{P}(V > t) < 0\} \neq \emptyset$. For any $t \in \mathfrak{T}$, define $A_t \doteq \{x \in \mathfrak{X} : a(x) \leq t\}$ and $B_t \doteq \{y \in \mathfrak{Y} : b(y) < -t\}$ so that for any $(x, y) \in A_t \times B_t$, $a(x) + b(y) < 0$. From (B.3), such (x, y) cannot be in \mathcal{S}_{XY} . Thus for any $t \in \mathfrak{T}$, $(A_t, B_t) \in \mathcal{N}_{\pi_{XY}}$, while

$$\pi_X^*(A_t) + \pi_Y^*(B_t) = \int_{A_t} \varphi_X^* \, d\mu_{\mathfrak{X}} + \int_{B_t} \varphi_Y^* \, d\mu_{\mathfrak{Y}} = \mathbb{P}(U \leq t) + \mathbb{P}(V > t) = 1 + \mathbb{P}(V > t) - \mathbb{P}(U > t) > 1.$$

(b) (i) The claim is trivial, by the convexity of the sets $\Sigma_{\pi_{XY}}$ and $\mathcal{F}(\pi_X^*, \pi_Y^*)$.

(ii) (\Leftarrow) Let $(A, B) \in \mathcal{N}_{\pi_{XY}}$ such that $\pi_X^*(A) + \pi_Y^*(B) = 1$ and $(A^c, B^c) \notin \mathcal{N}_{\pi_{XY}}$. For any $\pi_{XY}^* \in \mathcal{D}_{\mathfrak{X} \times \mathfrak{Y}} \cap \Sigma_{\pi_{XY}} \cap \mathcal{F}(\pi_X^*, \pi_Y^*)$ (including $\bar{\pi}_{XY}^*$ from (b)(i)), $\pi_{XY}^*(A \times B^c) = \pi_X^*(A)$ and $\pi_{XY}^*(A^c \times B^c) = \pi_Y^*(B^c) - \pi_{XY}^*(A \times B^c) = \pi_X^*(A) - \pi_{XY}^*(A \times B^c) = 0$. Hence, as $\pi_{XY}(A^c \times B^c) \neq 0$ by assumption, $\pi_{XY} \not\ll \pi_{XY}^*$ and they are not measure-theoretically equivalent.

(ii) (\Rightarrow) By definition we have $\bar{\pi}_{XY}^* \ll \pi_{XY}$, but assume that $\pi_{XY} \not\ll \bar{\pi}_{XY}^*$. Then there exists a measurable set $S_0 \subset \mathcal{S}_{XY}$ such that $\bar{\pi}_{XY}^*(S_0) = 0$ while $\pi_{XY}(S_0) > 0$. By (i), all distributions $\pi_{XY}^* \in \mathcal{D}_{\mathfrak{X} \times \mathfrak{Y}} \cap \Sigma_{\pi_{XY}} \cap \mathcal{F}(\pi_X^*, \pi_Y^*)$ are such that $\pi_{XY}^* \ll \bar{\pi}_{XY}^*$, so $\pi_{XY}^* \in \mathcal{D}_{\mathfrak{X} \times \mathfrak{Y}} \cap \Sigma_{\pi_{XY}} \cap \mathcal{F}(\pi_X^*, \pi_Y^*) \Rightarrow \pi_{XY}^*(S_0) = 0$. Now, $\pi_{XY}^* \in \mathcal{D}_{\mathfrak{X} \times \mathfrak{Y}} \cap \Sigma_{\pi_{XY}} \cap \mathcal{F}(\pi_X^*, \pi_Y^*)$ means that its density φ_{XY}^* satisfies (B.1), of which $\iint \varphi_{XY}^* \mathbb{I}_{\{S_0\}} \, d(\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}}) = 0$ should therefore be a consequence. Thus the system (B.1) augmented by the strict inequality

$$\iint \varphi_{XY}^* \mathbb{I}_{\{S_0\}} \, d(\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}}) > 0 \quad (\text{B.5})$$

must be inconsistent. As above, we can resort to Farkas lemma to show that such inconsistency is equivalent to the existence of some functions $a \in L_{\infty}(\mathfrak{X}, \mu_{\mathfrak{X}})$, $b \in L_{\infty}(\mathfrak{Y}, \mu_{\mathfrak{Y}})$, $k \geq 0$ and $\ell > 0$ such that

$$a + b + k \mathbb{I}_{\{\mathcal{S}_{XY}^c\}} - \ell \mathbb{I}_{\{S_0\}} = 0 \quad (\mu_{\mathfrak{X}} \times \mu_{\mathfrak{Y}})\text{-a.e.} \quad (\text{B.6})$$

and

$$\int a \, d\pi_X^* + \int b \, d\pi_Y^* \leq 0. \quad (\text{B.7})$$

(Note that the inequality is not strict here, as opposed to (B.4), as a consequence of the strict inequality in (B.5). Also, $\ell > 0$; *à la* Webster (1994, Theorem 4.2.4).)

Let X^* (resp. Y^*) be a random variable defined on \mathfrak{X} (resp. \mathfrak{Y}) with distribution π_X^* (resp. π_Y^*), and define $U = a(X^*)$ and $V = -b(Y^*)$, for a, b two functions satisfying (B.6) and (B.7). We have:

$$\int_{-\infty}^{\infty} (\mathbb{P}(U > t) - \mathbb{P}(V > t)) \, dt = \mathbb{E}(U) - \mathbb{E}(V) = \int a \, d\pi_X^* + \int b \, d\pi_Y^* \leq 0.$$

Denote u_0 the lower bound of the support of U , and v_0 the upper bound of the support of V (these bounds are finite as a, b are essentially bounded). As $\mathbb{P}(U > t) - \mathbb{P}(V > t) \geq 0$ for $t < u_0$ or $t \geq v_0$, there necessarily exists $t \in [u_0, v_0)$ such that $\mathbb{P}(U > t) - \mathbb{P}(V > t) \leq 0$. For such t , define $A_t \doteq \{x \in \mathfrak{X} : a(x) \leq t\}$ and $B_t \doteq \{y \in \mathfrak{Y} : b(y) < -t\}$ so that for any $(x, y) \in A_t \times B_t$, $a(x) + b(y) < 0$ (a.e.), implying that

$$k \mathbb{I}_{\{S_{XY}^c\}} - \ell \mathbb{I}_{\{S_0\}} > 0$$

from (B.6). As $S_0 \subset S_{XY}$, it follows that (x, y) cannot belong to S_{XY} when $(x, y) \in A_t \times B_t$; that is, $\pi_{XY}(A_t \times B_t) = 0$, and $(A_t, B_t) \in N_{\pi_{XY}}$. In addition,

$$\pi_X^*(A_t) = \mathbb{P}(U \leq t) = \mathbb{P}(U \in [u_0, t]) > 0 \quad \text{and} \quad \pi_Y^*(B_t) = \mathbb{P}(V > t) = \mathbb{P}(V \in (t, v_0]) > 0$$

and

$$\pi_X^*(A_t) + \pi_Y^*(B_t) = \mathbb{P}(U \leq t) + \mathbb{P}(V > t) = 1 - \mathbb{P}(U > t) + \mathbb{P}(V > t) \geq 1.$$

It must be that $\pi_X^*(A_t) + \pi_Y^*(B_t) = 1$, as by assumption $\pi_X^*(A) + \pi_Y^*(B) \leq 1 \, \forall (A, B) \in N_{\pi_{XY}}$.

As $(A_t \times B_t) \subset S_{XY}^c$ and $S_0 \subset S_{XY}$, $(A_t \times B_t) \subset S_0^c$, that is, $S_0 \subset (A_t \times B_t)^c = (A_t \times B_t^c) \cup (A_t^c \times B_t) \cup (A_t^c \times B_t^c)$. Assume that $(A_t^c, B_t^c) \in N_{\pi_{XY}}$. Then $S_{XY} \subseteq (A_t \times B_t^c) \cup (A_t^c \times B_t)$. As $\pi_X^*(A_t) + \pi_Y^*(B_t) = 1$, we know that $\pi_X^*(A_t) = \pi_Y^*(B_t^c)$, and $\pi_{XY}^*(A_t \times B_t^c) = \pi_X^*(A_t)$. Then there cannot be $\tilde{A} \subset A_t$ and $\tilde{B} \subset B_t^c$ such that $(\tilde{A}, \tilde{B}) \in N_{\pi_{XY}}$ and $\pi_X^*(\tilde{A}) + \pi_Y^*(\tilde{B}) > \pi_X^*(A_t)$, otherwise $(\tilde{A}, \tilde{B} \cup B_t) \in N_{\pi_{XY}}$ and $\pi_X^*(\tilde{A}) + \pi_Y^*(\tilde{B}) + \pi_Y^*(B_t) > \pi_X^*(A_t) + \pi_Y^*(B_t) = 1$, which is not allowed by assumption. So the restriction of π_{XY} , π_X^* and π_Y^* may be handled in the same way as the above argument. If for all $\tilde{A} \subset A_t$ and $\tilde{B} \subset B_t^c$ such that $(\tilde{A}, \tilde{B}) \in N_{\pi_{XY}}$, $\pi_X^*(\tilde{A}) + \pi_Y^*(\tilde{B}) < \pi_X^*(A_t)$, then $\bar{\pi}_{XY}^* \approx \pi_{XY}$ on $A_t \times B_t^c$ and thus $S_0 \not\subset A_t \times B_t^c$ – we come to the same conclusion recursively if there exists $\tilde{A} \subset A_t$ and $\tilde{B} \subset B_t^c$ such that $(\tilde{A}, \tilde{B}) \in N_{\pi_{XY}}$, $\pi_X^*(\tilde{A}) + \pi_Y^*(\tilde{B}) = \pi_X^*(A_t)$. Similarly, one shows that $S_0 \not\subset A_t^c \times B_t$. But if $(A_t^c, B_t^c) \in N_{\pi_{XY}}$, $\pi_{XY}(A_t^c \times B_t^c) = 0$ and S_0 cannot be in $A_t^c \times B_t^c$, leading to a contradiction. Hence $(A_t^c, B_t^c) \notin N_{\pi_{XY}}$.

Thus $\pi_{XY} \not\ll \bar{\pi}_{XY}^*$ implies the existence of a set $(A, B) \in N_{\pi_{XY}}$ such that $\pi_X^*(A) + \pi_Y^*(B) = 1$, and

$(A^c, B^c) \notin N_{\pi_{XY}}$.

Proof of Proposition 3.1

(\Rightarrow) Assume that $\{\varphi_{Y|X}\}$ and $\{\varphi_{X|Y}^*\}$ are compatible. By definition, there exists a bivariate distribution $\tilde{\pi}_{XY}$ admitting $\{\varphi_{Y|X}\}$ and $\{\varphi_{X|Y}^*\}$ as conditional densities. Thus $\{\varphi_{Y|X}\}$ and $\{\varphi_{X|Y}^*\}$ must encapsulate the same dependence structure, which is the dependence structure of $\tilde{\pi}_{XY}$. Hence $\Delta(\{\varphi_{Y|X}\}) = \Delta(\{\varphi_{X|Y}^*\})$.

(\Leftarrow) Assume that $\Delta(\{\varphi_{Y|X}\}) = \Delta(\{\varphi_{X|Y}^*\})$. By (3.6), this implies that any distribution admitting $\{\varphi_{Y|X}\}$ as $Y|X$ -conditional density shares the same joint support (and hence marginal supports) as any other distribution admitting $\{\varphi_{X|Y}^*\}$ as $X|Y$ -conditional density. In particular, take a distribution π_X on \mathcal{S}_X such that $d\pi_X/d\mu_{\mathfrak{X}} \in L_\infty(\mathfrak{X}, \mu_{\mathfrak{X}})$, and a distribution π_Y^* on \mathcal{S}_Y such that $d\pi_Y^*/d\mu_{\mathfrak{Y}} \in L_\infty(\mathfrak{Y}, \mu_{\mathfrak{Y}})$, and form

$$\pi_{XY}(\cdot) \doteq \iint \varphi_{Y|X} \mathbb{I}_{\{\cdot\}} d(\pi_X \times \mu_{\mathfrak{Y}}) \quad \text{and} \quad \pi_{XY}^*(\cdot) \doteq \iint \varphi_{X|Y}^* \mathbb{I}_{\{\cdot\}} d\pi_Y^* d\mu_{\mathfrak{X}}.$$

As these two distributions have conditional densities $\{\varphi_{Y|X}\}$ and $\{\varphi_{X|Y}^*\}$, respectively, they have the same support. See from Corollary 3.2 that condition (3.11) is automatically fulfilled for π_{XY} and the marginals π_X^*, π_Y^* of π_{XY}^* , given that $N_{\pi_{XY}} = N_{\pi_{XY}^*}$ and there exists a distribution in $\mathcal{F}(\pi_X^*, \pi_Y^*)$ with the suitable support. Then, according to Corollary 3.3, there is a unique distribution in $\mathcal{F}(\pi_X^*, \pi_Y^*)$ with the same dependence as π_{XY} – this must be π_{XY}^* , by assumption – and it is such that

$$\frac{d\pi_{XY}^*}{d\pi_{XY}} = \alpha_X^* \beta_Y^*$$

for some positive functions $\alpha_X^* \in L_1(\mathfrak{X}, \mu_{\mathfrak{X}})$ and $\beta_Y^* \in L_1(\mathfrak{Y}, \mu_{\mathfrak{Y}})$. This implies that

$$\frac{\varphi_{X|Y}^*}{\varphi_{Y|X}} = \frac{d\pi_{XY}^*}{d(\mu_{\mathfrak{X}} \times \pi_Y^*)} \frac{d(\pi_X \times \mu_{\mathfrak{Y}})}{d\pi_{XY}} = \left(\alpha_X^* \frac{d\pi_X}{d\mu_{\mathfrak{X}}} \right) \left(\beta_Y^* \frac{d\mu_{\mathfrak{Y}}}{d\pi_Y^*} \right) \doteq \tilde{\alpha}_X^* \tilde{\beta}_Y^*.$$

As $d\pi_X/d\mu_{\mathfrak{X}} \in L_\infty(\mathfrak{X}, \mu_{\mathfrak{X}})$, $\tilde{\alpha}_X^* \in L_1(\mathfrak{X}, \mu_{\mathfrak{X}})$. Arnold and Press (1989, Theorem 4.1) establishes this as a necessary and sufficient condition for $\{\varphi_{X|Y}^*\}$ and $\{\varphi_{Y|X}\}$ to be compatible. \square

Proof of Theorem 4.1

Let $\{\varphi_{Y|X}\}$ be the $Y|X$ -conditional density of π_{XY} , and $\{\varphi_{X|Y}^*\}$ be the $X|Y$ -conditional density of π_{XY}^* . As per Proposition 3.1, $\mathfrak{D} = \mathfrak{D}^*$ if and only if $\{\varphi_{Y|X}\}$ and $\{\varphi_{X|Y}^*\}$ are compatible. From Arnold and Press (1989, Theorem 4.1), $\{\varphi_{Y|X}\}$ and $\{\varphi_{X|Y}^*\}$ are compatible if and only if (i) $\{(x, y) \in \mathcal{S}_X \times \mathfrak{Y} : \varphi_{Y|X}(y|x) > 0\} = \{(x, y) \in \mathfrak{X} \times \mathcal{S}_Y : \varphi_{X|Y}^*(x|y) > 0\}$, which is equivalent to $\mathcal{S}_{XY} = \mathcal{S}_{XY}^*$ and therefore follows from (3.6); and (ii) there exist (a.e.-)positive functions $\phi_X \in L_1(\mathfrak{X}, \mu_{\mathfrak{X}})$ and $\phi_Y \in L_1(\mathfrak{Y}, \mu_{\mathfrak{Y}})$, such that

$$\frac{\varphi_{X|Y}^*}{\varphi_{Y|X}} = \frac{\phi_X}{\phi_Y} \quad \text{a.e. on } \mathcal{S}_{XY}.$$

As

$$\frac{\varphi_{\mathfrak{X}|\mathfrak{Y}}^*}{\varphi_{\mathfrak{Y}|\mathfrak{X}}} = \frac{d\pi_{\mathfrak{X}\mathfrak{Y}}^*}{d(\mu_{\mathfrak{X}} \times \pi_{\mathfrak{Y}}^*)} \frac{d(\pi_{\mathfrak{X}} \times \mu_{\mathfrak{Y}})}{d\pi_{\mathfrak{X}\mathfrak{Y}}}$$

directly from (3.1), we have

$$\frac{d\pi_{\mathfrak{X}\mathfrak{Y}}^*}{d\pi_{\mathfrak{X}\mathfrak{Y}}} = \frac{\phi_{\mathfrak{X}} d\mu_{\mathfrak{X}}}{d\pi_{\mathfrak{X}}} \frac{d\pi_{\mathfrak{Y}}^*}{\phi_{\mathfrak{Y}} d\mu_{\mathfrak{Y}}} \doteq \alpha_{\mathfrak{X}}^* \beta_{\mathfrak{Y}}^*. \quad (\text{B.8})$$

See that $\alpha_{\mathfrak{X}}^* = \phi_{\mathfrak{X}} d\mu_{\mathfrak{X}}/d\pi_{\mathfrak{X}} \in L_1(\mathfrak{X}, \pi_{\mathfrak{X}})$, as $\int \alpha_{\mathfrak{X}}^* d\pi_{\mathfrak{X}} = \int \phi_{\mathfrak{X}} d\mu_{\mathfrak{X}} < \infty$. Likewise, $1/\beta_{\mathfrak{Y}}^* = \phi_{\mathfrak{Y}} d\mu_{\mathfrak{Y}}/d\pi_{\mathfrak{Y}}^* \in L_1(\mathfrak{Y}, \pi_{\mathfrak{Y}}^*)$. From (B.8) we have directly that $d\pi_{\mathfrak{X}\mathfrak{Y}}/d\pi_{\mathfrak{X}\mathfrak{Y}}^* = 1/\alpha_{\mathfrak{X}}^* \times 1/\beta_{\mathfrak{Y}}^*$, while, swapping $\pi_{\mathfrak{X}\mathfrak{Y}}^*$ and $\pi_{\mathfrak{X}\mathfrak{Y}}$ from the beginning of the argument, we would have obtained instead of (B.8):

$$\frac{d\pi_{\mathfrak{X}\mathfrak{Y}}}{d\pi_{\mathfrak{X}\mathfrak{Y}}^*} = \frac{\phi_{\mathfrak{X}}^* d\mu_{\mathfrak{X}}}{d\pi_{\mathfrak{X}}^*} \frac{d\pi_{\mathfrak{Y}}}{\phi_{\mathfrak{Y}}^* d\mu_{\mathfrak{Y}}}$$

for some functions $\phi_{\mathfrak{X}}^* \in L_1(\mathfrak{X}, \mu_{\mathfrak{X}})$ and $\phi_{\mathfrak{Y}}^* \in L_1(\mathfrak{Y}, \mu_{\mathfrak{Y}})$. Identifying both expressions, we see that $\beta_{\mathfrak{Y}}^*$ can be written $\phi_{\mathfrak{Y}}^* d\mu_{\mathfrak{Y}}/d\pi_{\mathfrak{Y}}$, showing that $\beta_{\mathfrak{Y}}^* \in L_1(\mathfrak{Y}, \pi_{\mathfrak{Y}})$. The claim follows. \square

Proof of Theorem 4.2

We show that, for two distributions $\pi_{\mathfrak{X}\mathfrak{Y}}, \pi_{\mathfrak{X}\mathfrak{Y}}^* \in \overline{\mathcal{D}}_{\mathfrak{X} \times \mathfrak{Y}}$,

$$\pi_{\mathfrak{X}\mathfrak{Y}} \sim \pi_{\mathfrak{X}\mathfrak{Y}}^* \iff \iint \log \varphi_{\mathfrak{X}\mathfrak{Y}} d\gamma_k = \iint \log \varphi_{\mathfrak{X}\mathfrak{Y}}^* d\gamma_k \quad \forall k \in \{1, 2, \dots\},$$

where $\pi_{\mathfrak{X}\mathfrak{Y}} \sim \pi_{\mathfrak{X}\mathfrak{Y}}^*$ is defined as per (4.2) and $\{\gamma_1, \gamma_2, \dots\}$ is any basis of $\Gamma_{\mathfrak{X}\mathfrak{Y}}^\circ$ defined in (4.3).

(\Rightarrow) Assume $\pi_{\mathfrak{X}\mathfrak{Y}} \sim \pi_{\mathfrak{X}\mathfrak{Y}}^*$. This means that $\varphi_{\mathfrak{X}\mathfrak{Y}}^* = \varphi_{\mathfrak{X}\mathfrak{Y}} \alpha_{\mathfrak{X}}^* \beta_{\mathfrak{Y}}^*$ for two (a.e.-)positive functions $\alpha_{\mathfrak{X}}^* \in L_1(\mathfrak{X}, \pi_{\mathfrak{X}})$ and $\beta_{\mathfrak{Y}}^* \in L_1(\mathfrak{Y}, \pi_{\mathfrak{Y}})$. Note that, by symmetry (see Proof of Theorem 4.1), $\alpha_{\mathfrak{X}}^*$ and $\beta_{\mathfrak{Y}}^*$ are also essentially bounded. Then, $\forall k \in \{1, 2, \dots\}$,

$$\iint \log \varphi_{\mathfrak{X}\mathfrak{Y}}^* d\gamma_k = \iint \log \varphi_{\mathfrak{X}\mathfrak{Y}} d\gamma_k + \iint \log \alpha_{\mathfrak{X}}^* d\gamma_k + \iint \log \beta_{\mathfrak{Y}}^* d\gamma_k = \iint \log \varphi_{\mathfrak{X}\mathfrak{Y}} d\gamma_k$$

by definition, as any $\gamma_k \in \Gamma_{\mathfrak{X}\mathfrak{Y}}^\circ$ has null marginals, making $\iint \log \alpha_{\mathfrak{X}}^* d\gamma_k = \iint \log \beta_{\mathfrak{Y}}^* d\gamma_k = 0$.

(\Leftarrow) Assume that $\iint \log \varphi_{\mathfrak{X}\mathfrak{Y}} d\gamma_k = \iint \log \varphi_{\mathfrak{X}\mathfrak{Y}}^* d\gamma_k$, $\forall k \in \{1, 2, \dots\}$. As $\{\gamma_1, \gamma_2, \dots\}$ is a basis of $\Gamma_{\mathfrak{X}\mathfrak{Y}}^\circ$, it follows that

$$\iint \log \frac{\varphi_{\mathfrak{X}\mathfrak{Y}}^*}{\varphi_{\mathfrak{X}\mathfrak{Y}}} d\gamma = 0 \quad \forall \gamma \in \Gamma_{\mathfrak{X}\mathfrak{Y}}^\circ.$$

Hence $\log \frac{\varphi_{\mathfrak{X}\mathfrak{Y}}^*}{\varphi_{\mathfrak{X}\mathfrak{Y}}}$ must belong to the annihilator (Rudin, 1991, Section 4.6) of $\Gamma_{\mathfrak{X}\mathfrak{Y}}^\circ$, viz.

$$\Psi_{\mathfrak{X}\mathfrak{Y}}^\circ \doteq \{\psi \in L_\infty(\mathfrak{X} \times \mathfrak{Y}) : \iint \psi d\gamma = 0, \forall \gamma \in \Gamma_{\mathfrak{X}\mathfrak{Y}}^\circ\},$$

which can be verified to be $\Psi_{\mathfrak{X}\mathfrak{Y}}^\circ = L_\infty(\mathfrak{X}, \mu_{\mathfrak{X}}) \oplus L_\infty(\mathfrak{Y}, \mu_{\mathfrak{Y}})$. Hence there exist $\psi_{\mathfrak{X}} \in L_\infty(\mathfrak{X}, \mu_{\mathfrak{X}})$ and $\psi_{\mathfrak{Y}} \in L_\infty(\mathfrak{Y}, \mu_{\mathfrak{Y}})$ such that $\log \frac{\varphi_{\mathfrak{X}\mathfrak{Y}}^*}{\varphi_{\mathfrak{X}\mathfrak{Y}}} = \psi_{\mathfrak{X}} + \psi_{\mathfrak{Y}}$, and finally $\frac{d\pi_{\mathfrak{X}\mathfrak{Y}}^*}{d\pi_{\mathfrak{X}\mathfrak{Y}}} = \frac{\varphi_{\mathfrak{X}\mathfrak{Y}}^*}{\varphi_{\mathfrak{X}\mathfrak{Y}}} = \exp(\psi_{\mathfrak{X}} + \psi_{\mathfrak{Y}}) \doteq \alpha_{\mathfrak{X}}^* \beta_{\mathfrak{Y}}^*$, where $\alpha_{\mathfrak{X}}^* \beta_{\mathfrak{Y}}^* > 0$. As $\alpha_{\mathfrak{X}}^*$ and $\beta_{\mathfrak{Y}}^*$ are essentially bounded, they are $\pi_{\mathfrak{X}}$ - and $\pi_{\mathfrak{Y}}$ -integrable. \square