

**The cosmological constant, dark matter, supersymmetry,
and other unsolved problems from a fresh perspective**

Roland E. Allen

Department of Physics and Astronomy

Texas A&M University, College Station, Texas 77843, USA

allen@tamu.edu, orcid 0000-0001-5951-4717

arXiv:2302.10241v4 [physics.gen-ph] 12 Jan 2024

Abstract

Quantum theory, general relativity, the standard model of particle physics, and the Λ CDM model of cosmology have all been spectacularly successful within their respective regimes of applicability, but each of these descriptions also has clear limitations. The most popular attempts to extend these theories and unify them have, after decades of initial optimism, turned out so far to be failures: There is still no widely accepted solution to the cosmological constant problem; dark matter has not yet been definitively detected; supersymmetry has not yet been observed; and string theory has yielded no testable predictions. These long-term failures (after intense efforts by many of the most mathematically talented theorists) suggest that a completely new perspective is needed. Here we propose a fundamental theory which might be regarded as a cousin to string theory, since it also is based on higher dimensions (with an internal space), a form of supersymmetry, important topological structures, and the implication of a multiverse. Our universe is topologically stable because it is the product of two vortex-like (or instanton-like) field configurations – one in 4-dimensional external spacetime, with the big bang at its origin, and the other in a $(D-4)$ -dimensional naturally compactified internal space, which automatically yields an $SO(D-4)$ unified gauge theory. The fundamental theory cannot even be formulated without a primitive form of supersymmetry (susy) among initial primitive fields. A stable vacuum with Lorentz invariance requires a breaking of this initial susy, as the primitive (and unphysical) bosonic fields are transformed into physical fields. There is then a new interpretation of all scalar boson sectors – including but extending the Higgs and sfermion sectors. This last feature predicts a novel dark matter WIMP with no (nongravitational) interactions except second-order gauge couplings to W and Z bosons. Calculations and estimates of the relevant cross-sections for this particle demonstrate that (1) it may be detectable within the next 5-15 years in Xe-based direct-detection experiments, (2) it may be observable within about 12-20 years at the high-luminosity LHC, and (3) it may already have been detected in the gamma rays observed by Fermi-LAT and antiprotons observed by AMS-02. The reinterpretation of scalar boson fields also implies a new phenomenology for sfermions, with reduced cross-sections. There is then a unified picture which may explain why dark matter WIMPs and electroweak-scale sparticles have not yet been detected. Many other central theoretical issues are also discussed in the present paper. But since it is already unusually long and complicated, nonessential topics are omitted here, and quantitative predictions for specific detection cross-sections are left for papers with student collaborators, as cited in the text.

I. INTRODUCTION

The great achievements of modern physics and astronomy have actually *increased* the number of perceived mysteries as we strive for a fundamental understanding of Nature. Some of these mysteries were already appreciated about a century ago, by Einstein and others, and some have arisen from recent discoveries. Many excellent reviews have been given of the current situation [1], but it may be worthwhile to begin with a brief summary.

Most recently, the particle discovered by the ATLAS and CMS collaborations at the LHC is now known to be a Higgs boson [1–3]. A naive conclusion is that the standard model of particle physics is now complete. But the more profound interpretation is that the discovery of a scalar boson immediately points to physics beyond the standard model, since otherwise radiative corrections should push the mass of this particle up to an absurdly large value. The most natural candidate for such new physics is supersymmetry (susy) [4–7], for which there is already indirect experimental evidence: The coupling constants of the 3 nongravitational forces are found to converge to a common value, as they are run up to high energy in a grand unified theory, only if the calculation includes susy. So, instead of acting as an endpoint for physics, and a mere capstone of the standard model, the observation of a Higgs boson may open the door to a plethora of new particles and effects. But susy has not yet been observed, despite widespread initial optimism, and even at the theoretical level an accepted mechanism for susy breaking has not been found.

Another major advance has been the discovery and exploration of neutrino masses [8], which appear to open the door to a more fundamental understanding of forces and matter via grand unification [5, 6, 9–12]. There are two possibilities for a neutrino mass, either of which is inconsistent with the requirements of the standard model: For a Dirac mass an extra field has to be added for each generation of fermions, and for a Majorana mass lepton number conservation has to be violated, whereas either or both types of mass are natural with grand unification, and in addition a seesaw mechanism can explain the small observed values of neutrino masses. Still, it is not yet demonstrated whether neutrinos have Majorana masses or Dirac masses or both, and the unification of coupling constants at high energy appears to require susy, which again has an uncertain status currently.

There are many other gaps in fundamental understanding. For example, the discovery and exploration of cosmic acceleration [13, 14] has suggested the need for truly revolutionary new

physics. The cause of this acceleration has increasingly been found to resemble a cosmological constant Λ , and has therefore been a strong reminder of the original cosmological constant problem [15]: Because of the various contributions to the vacuum energy, conventional general relativity predicts that Λ should be vastly larger than permitted by observation, unless there is a somewhat magical cancellation of positive and negative contributions. This problem is so difficult that a number of outstanding theorists, including Steven Weinberg [16, 17], have felt that there is no convincing solution other than an anthropic argument.

A parallel astronomical mystery is the origin of dark matter [18]. Many dark matter candidates for which there was initial optimism have now been ruled out by the remarkable achievements of the experimentalists, in obtaining ever-increasing sensitivity and sophistication.

Yet another and even older theoretical problem is the difficulty of reconciling general relativity with quantum mechanics: A new fundamental theory must somehow regularize quantum gravity, in addition to reducing the value of Λ by many orders of magnitude.

The next level of theoretical understanding is not likely to be a “theory of everything”, since “everything” surely transcends our current observational capabilities and imagination. But the following are a few among the many problems that a fundamental theory might hope to explain: the absence of an enormous cosmological constant, the origin of dark matter, the role and nature of supersymmetry, the origin of Lorentz invariance, the origin of gravitational and gauge interactions, the gravitational metric and its signature (which distinguishes time from space and characterizes spacetime as 4-dimensional), the fundamental action for fermionic and bosonic fields, the regularization of quantum gravity, the origin of quantum fields, and the origin of spacetime coordinates.

The present theory addresses all of the issues listed above and leads to a substantial number of predictions. The most quantitative are those for the dark matter candidate, as discussed below and in our papers on this one aspect of the present theory [19–22].

II. OVERVIEW

DeWitt has provided an elegant survey of contemporary fundamental physics [23], which is based on path-integral quantization over classical trajectories in the combined space of coordinates and fields: “A classical dynamical system is described globally by a *trajectory*

or *history*. A history is a section of a fibre bundle E having the manifold M of spacetime as its base space. The typical fibre is known as configuration space and will be denoted by C Denote by Φ the set, or space, of all possible field histories, both those that do and those that do not satisfy the dynamical equations The nature and dynamical properties of a classical dynamical system are completely determined by specifying an *action functional* S for it.”

This is the basic picture used in most versions of fundamental physics that are investigated by sizable communities of physicists. Notice that coordinates and fields have essentially the same status. This is consistent with the way they are defined in the present theory, starting near the beginning of the next section. In standard field theory, the spacetime coordinates x^μ correspond to M and the fields to C . In conventional string (or p -brane) theory, M corresponds to a 2-dimensional worldsheet (or $(p + 1)$ -dimensional worldvolume) and C to bosonic and fermionic coordinates. Note also that coordinates (in nonrelativistic quantum mechanics) and fields (in quantum field theory) are treated essentially the same way in both canonical and path-integral quantization.

The arguments below involve many unfamiliar definitions and redefinitions of fields, with physically accessible fields emerging from more primitive “hidden” degrees of freedom. But this kind of treatment is already necessary and familiar in standard physics – for example, when the fundamental gauge fields are redefined after Higgs condensation, and when many kinds of elementary excitations are defined in condensed matter physics. More broadly, the deeper hidden origin of observed phenomena is a common theme in science.

The physical fields that have emerged by the end of this paper are interpreted as those chosen by Nature to yield a stable vacuum $|0\rangle$ satisfying $a|0\rangle = 0$, where a is a typical destruction operator for one of these fields. For example, the excitations $a^\dagger|0\rangle$ must have positive energy.

The principal ideas and results of this paper are as follows.

(1) In standard physics, all the events of the world are simply a progression through the states of fields on a spacetime manifold. In the present picture, they are a progression through the states of a single fundamental system. This second unified picture leads back to (and beyond) the first via the arguments below.

Time will be defined as a parameter describing a trajectory through the space of states. There are, of course, precedents for defining time internally, within a stationary state, using

some internal parameter. For example, in minisuperspace models based on the Wheeler-DeWitt equation [24, 25], this parameter is the cosmic scale factor, and time is defined by the expansion of the universe. In the present picture, *all* coordinates and fields are defined internally, with the progression of events in Nature regarded as progression through a space of states. In the most primitive space, each point is defined by a microstate $|m\rangle$. In the emergent space corresponding to DeWitt's Φ (see above), each point is defined by a macrostate, as described below.

For a concrete characterization of both microstates and macrostates, we adopt the picture that Nature is composed of discrete distinguishable constituents that are called “dits” because each can exist in any of d states labeled by $i = 1, 2, \dots, d$. The number of dits in the i th state determines the size of the field or coordinate labeled by i .

The fact that fields and coordinates have the same basic status in standard physics, as noted above, is then explained by the fact that they have the same fundamental origin.

This appears to be the simplest imaginable picture that can still lead back to standard physics. The fundamental objects are dits (a generalization of Wheeler's “it from bit” [26]), and these are the “atoms” of the fundamental system. Coordinates and fields are then emergent quantities, associated with the possible paths through the full state vector of the fundamental system. It is somewhat miraculous that such a simple picture can lead back to standard physics and its natural generalizations.

(2) The action is defined to be essentially the negative of the entropy for an observable macrostate (which corresponds to a particular set of values of coordinates and fields). Action conventionally has the units of \hbar , and entropy the units of the Boltzmann constant k , but here we use natural units, with $\hbar = k = c = 1$.

(3) Much later in the paper, the resulting path integrals with Euclidean form are transformed to equivalent Lorentzian path integrals, with the action left unchanged.

The standard formulation of quantum field theory has then been regained. One can subsequently transform from path-integral quantization to canonical quantization in the usual way (since the action has a standard form).

Since spacetime points are discrete (although very closely spaced) in the present description, path integrals are well-defined. All aspects of the theory are in fact clearly mathematically consistent, since both coordinates and fields are initially defined on a lattice specified by integers, and the continuum is only an approximation. In contrast, both the mean-

ing of path integrals and overall mathematical consistency are issues in field theory and in extensions such as string theory.

(4) The assumption that all states of the fundamental system are realized leads inevitably to a multiverse picture. There are still many who reject the possibility of a multiverse, but one should recall that most people at the time of Galileo would have rejected the possibility of hundreds of billions of galaxies, or a single galaxy, or even a heliocentric Solar System, and that the history of physics shows a steady progression toward more expansive views of Nature.

Among the vast number of states of the fundamental system (in the full path integral), there are some trajectories through these states which can legitimately be assigned to universes, in the sense that the states or configurations can be coherently connected with high probability. This will be the case if there is a path through the space of these configurations along which a local minimum is stably maintained in the action.

In the present picture our own universe is stabilized *topologically* – specifically, it has an extremely stable geometry determined by two topological defects in a primordial condensate, one in 4-dimensional external spacetime and the other in a $(D-4)$ -dimensional internal space. Each of these is “vortex-like” (or “instanton-like”), in the sense that they involve circulation of the primordial condensate around a central point. The stability of our universe is then analogous to the stability of a vortex in a 2-dimensional fluid.

This stability has an additional feature, in that there is a natural mechanism for compactification of the internal space, as mentioned at the end of Appendix D. (The term “compactification” as used here means that the internal space is restricted to a finite and very small volume.) If the relevant functions in the internal space are exponentially decaying solutions to the fundamental higher-order differential equation, they will be confined to an extremely small region (with a length scale $\sim a_0$, the fundamental length defined below). There is then a natural explanation of the difference between external spacetime and the internal space: The condensate and basis functions in internal space decay exponentially (with distance from the internal topological singularity), but the external condensate begins in one of the nondecaying solutions, with normal and unlimited expansion of the universe afterward.

Despite early attempts by Candelas and Weinberg [27], and others, a convincing mechanism for compactification in Kaluza-Klein, supergravity, and string theories is an unsolved

problem, and the qualitatively different nature of the internal space is ordinarily just postulated.

There is an additional nuance, in that a universe can be made stable through an effect which is exhibited in the behavior of the quartic self-coupling of the recently discovered Higgs boson: Within the standard model, the unrenormalized value of this coupling appears to be very nearly equal to zero [28–31], but at low energies it is made appreciably finite by radiative corrections, so that a stable Higgs condensate forms. This suggests that more generally there will be configurations, within the complete path integral of all possibilities, where radiative corrections will similarly yield a nonzero quartic self-coupling for a primordial condensate which allows it to form. In order to search for such a possible solution (i.e. a persistent local minimum in the action), we adopt the artifice of adding an imaginary random potential to the action, proportional to a parameter b which is ultimately taken to go to zero: $b \rightarrow 0+$.

Known physics is regained in a very simple picture based on a primordial condensate with an $SU(2) \times U(1)$ order parameter in external spacetime and an $SO(D-4) \times U(1)$ (or more precisely $Spin(D-4) \times U(1)$) order parameter in the internal space. Each of these factors in the overall order parameter has a vortex-like topological singularity at the origin. The external topological defect is interpreted as the big bang, and the internal topological defect gives rise to an $SO(D-4)$ grand-unified gauge group, with $D-4 \geq 10$.

The gravitational vierbein and the gauge fields of other forces are interpreted as “superfluid velocities”, with arbitrary curvatures permitted by a background of “rapidly fluctuating” topological defects that are analogous to vortices and vortex rings (or extended and closed flux tubes) – in roughly the same way that, as shown by Feynman and Onsager, vortices permit rotation of a superfluid.

(5) In the detailed treatment, the fermionic fields and scalar bosonic fields are found to automatically couple in the correct way to both the gravitational field and the gauge fields of the other forces.

(6) At the same time, local Lorentz invariance automatically emerges (rather than being postulated), and external spacetime is automatically $(3+1)$ -dimensional.

(7) The present theory unavoidably predicts $SO(D-4)$ grand unification. This is consistent with the fact that many regard $SO(10)$ as the most appealing gauge group for unifying standard model forces. Family replication can result from $D-4 > 10$, via a horizontal group.

Gauge symmetry results from rotational symmetry in the internal space, and this ex-

plains why forces are described by a basic gauge symmetry which is so similar to rotational symmetry.

(8) The present theory also unavoidably predicts a radically modified version of supersymmetry: There is initially a primordial mathematical susy involving standard fermion fields but unphysical boson fields, which must be transformed to obtain a stable vacuum with Lorentz invariance. The final action resembles a standard supersymmetric action, automatically including auxiliary fields F , but there is not invariance under the mathematical transformations of conventional susy. Instead there is only a *physical* susy at high energy [32] – with the full range of superpartners – which still protects the Higgs boson mass-squared from a quadratic divergence due to radiative corrections, and also can lead to unification of coupling constants at high energy.

Breaking of susy is an unsolved problem in conventional models, but here the primordial (mathematical) susy is automatically broken because the initial bosonic and fermionic fields must be treated quite differently: To satisfy Lorentz invariance, the initial primordial bosonic fields must be transformed into scalar boson fields of two kinds – complex Higgs-like fields ϕ and real fields φ with novel properties.

There will still be a lightest supersymmetric particle [4–7] which (as a subdominant component) can stably coexist with the dark matter candidate of the present theory, as can axion-like particles [33–37], because these various species are stable for different reasons [19–22].

(9) The usual cosmological constant (regarded as one of the deepest problems in standard physics) automatically vanishes for two independent reasons:

(i) For fermion fields and scalar boson fields, there is initially no factor of $e = \sqrt{-g}$ in the integrals giving their action. As shown below, this implies that their conventional contribution to a cosmological constant vanishes. The standard familiar form of the action can be regained,

$$S_{matter} = \int d^4x e \bar{\mathcal{L}}_{matter} , \quad (2.1)$$

but the stress-energy tensor for a fermion or scalar boson field must then be calculated from

$$T_{\varphi}^{\mu\nu} = 2 \varphi^{\dagger} \frac{\delta A_{\varphi}}{\delta g_{\mu\nu}} \varphi \quad (2.2)$$

and not the conventional

$$T_{\varphi}^{\mu\nu} = 2 e^{-1} \frac{\delta (e\varphi^{\dagger} A_{\varphi} \varphi)}{\delta g_{\mu\nu}} \quad (2.3)$$

where the notation is defined below.

(ii) When the gauge-field action is quantized, the operators must be normal-ordered, in accordance with the interpretation of the origin of this action in the present theory: It arises from the response of vacuum fields to the curvature of the external gauge fields, and it must therefore vanish when there are no external fields. The vacuum stress-energy tensor for the gauge fields then also vanishes.

Standard physics is regained in each case:

(i) Classical matter (which follows the on-shell classical equations of motion) acts as a source for Einstein gravity in the same way as in standard physics, and all matter and fields move in the same way.

(ii) The results are consistent with experiment and observation, even though there is no vacuum zero-point energy for the gauge fields.

Many people (including some who are otherwise expert in this area) will naively object that the Casimir effect, as verified experimentally, demonstrates that the electromagnetic field does have a zero-point energy in the vacuum.

This belief is common but incorrect [38]. The experimentally-observed Casimir effect demonstrates only that the static electromagnetic field energy is *changed* by the modification of boundary conditions [39, 40]. In the simplest model, two metal plates are inserted and the force between them calculated. There are two ways to do the calculation: The first is indeed to assume zero-point vibrations of the electromagnetic field, whose energy is modified when the boundary conditions are modified. The second approach is instead to consider the processes involving virtual photons which mediate the interaction of the plates, with no reference to zero-point vibrations and no need for a vacuum energy. The first method is more popular because it is easier (and has been assigned as a homework problem by the present author). But the two methods give the same basic answer, as they should. (The second method regards the force as mediated by virtual photons, and the first obtains the force from the derivative of the energy with respect to a displacement.) The second method is more difficult, but is consistent with the way other virtual processes are calculated in e.g. quantum electrodynamics. Of course, the second method also implies a change in the static

electromagnetic field energy (interpreted as a van der Waals interaction), but this *change* does not imply an initially nonzero vacuum zero-point energy.

In summary, the observed Casimir effect is perfectly consistent with the present theory, in which there is no vacuum zero-point energy due to the electromagnetic field or other gauge fields.

(10) Although the usual cosmological constant vanishes, there will still be a weaker response of the vacuum to the imposition of external fields. Within the present picture, the Maxwell-Yang-Mills action for gauge fields and the Einstein-Hilbert action for the gravitational field must arise from the modification of vacuum states (and the terms in their action) when they adjust to the curvature of these fields, just as the free energy of a metal associated with Landau diamagnetism arises from the modification of the electronic ground state when it qualitatively changes to accommodate the gauge curvature of a magnetic field.

Quantum gravity is regularized by an energy and momentum cutoff a_0^{-1} , where a_0 is an extremely small minimum length defined below.

(11) Topological singularities, including those associated with black holes and the big bang, are admissible in the present theory, for the same reason as in a standard superfluid: The density of the primordial condensate underlying standard physics goes to zero at each point where there is a gravitational or gauge-field singularity. (These singularities are present in the continuum approximation, on a length scale far larger than a_0 .)

(12) The present theory predicts new particles, including a new dark matter WIMP which is consistent with experiment and observation because it has no couplings other than its second-order gauge couplings to W and Z bosons. It should be barely observable in the high-luminosity large hadron collider (LHC) within roughly the next 12-20 years, and in direct-detection experiments within roughly the next 5-15 years, and it may already have been detected via the gamma rays observed by Fermi-LAT and antiprotons observed by AMS-02 [19–22]. This particle is unique among viable dark matter candidates in that both its couplings and its mass are precisely determined, making clean experimental tests possible in the near future.

Let us now proceed to the detailed arguments behind the above claims.

III. STATISTICAL ORIGIN OF THE INITIAL ACTION

For a theory to be viable, it must be mathematically (and philosophically) consistent, its premises must lead to testable predictions, and these predictions must be consistent with experiment and observation. The theory presented here satisfies these requirements, but it starts with an extremely unfamiliar point of view: There are initially no laws, and instead all possibilities are realized with equal weight. The observed laws of nature are emergent phenomena, which result from statistical counting and the geography (i.e. specific features) of our particular universe in D dimensions. In other words, standard physics (plus natural extensions such as grand unification and a modified supersymmetry) emerges as an effective field theory at relatively low energies.

Our starting point is a single fundamental system which consists of identical (but distinguishable) irreducible objects, which we will call “dits”. Each dit can exist in any of d states, with the number of dits in the i th state represented by n_i . An unobservable microstate of the fundamental system is specified by the number of dits and the state of each dit. An observable macrostate is specified by only the occupancies n_i of the states.

As discussed below, D of the states are used to define D spacetime coordinates x^M , and N_F of the states are used to define fields ϕ_k .

Let us begin with the coordinates:

$$x^M = \Delta n_M a_0 \quad , \quad \Delta n_M = n_M - \bar{n}_M \quad , \quad M = 0, 1, \dots, D - 1 \quad (3.1)$$

so that \bar{n}_M specifies a shift in the origin of coordinates. It is convenient to include a (very small) fundamental length a_0 in this definition, so that we can later express the coordinates in conventional units.

In parallel with the treatment of fields in Appendix A, we eventually take the limit $\bar{n}_M \rightarrow \infty$, with Δn_M finite, and there will then be no lower bound to negative coordinates. I.e., Δn_M can have any integer value. (A central feature of the present theory is that both coordinates and physical fields are defined by relatively small perturbations $\Delta n_i = n_i - \bar{n}_i$, analogous to waves on a deep ocean.)

Now define a set of initial fields ϕ_k by

$$\phi_k^2(x) = \rho_k(x) \quad , \quad k = 1, 2, \dots, N_F \quad (3.2)$$

where

$$\rho_k(x) = n_k(x)/a_0^D \quad (3.3)$$

and x represents all the coordinates. (To avoid awkward notation, we write n_k for $n_{i=D+k}$.)

The most primitive fields ϕ_k are then real, and defined only up to a phase factor ± 1 .

In Appendices A and B we obtain the entropy S for a given configuration of the ϕ_k at all points in spacetime, which then leads to the action for a given path (i.e. specific classical field configuration) in the quantum path integral:

$$S_E = \int d^D x \left\{ \frac{1}{2m_0} \left[\frac{\partial \Psi^\dagger}{\partial x^M} \frac{\partial \Psi}{\partial x^M} + \frac{a_0^2}{16} \frac{\partial^2 \Psi^\dagger}{\partial (x^M)^2} \frac{\partial^2 \Psi}{\partial (x^M)^2} \right] - \mu_0 \Psi^\dagger \Psi + \frac{1}{2} b (\Psi^\dagger \Psi)^2 \right\} \quad (3.4)$$

with a sum over the repeated index M in the first two terms. Here Ψ contains primordial bosonic and fermionic fields Ψ_b and Ψ_f

$$\Psi = \begin{pmatrix} \Psi_b \\ \Psi_f \end{pmatrix} \quad (3.5)$$

which, after the transformations below, will become the more familiar spin 0 and spin 1/2 fields of standard physics. For any functional F of these fields we have

$$\langle F \rangle = \frac{\int \mathcal{D} \Psi^\dagger \mathcal{D} \Psi F e^{-S_E}}{\int \mathcal{D} \Psi^\dagger \mathcal{D} \Psi e^{-S_E}}. \quad (3.6)$$

Ordinarily we can let $a_0 \rightarrow 0$ in (3.4), so that

$$S_E = \int d^D x \left[\frac{1}{2m_0} \partial_M \Psi^\dagger \partial_M \Psi - \mu_0 \Psi^\dagger \Psi + \frac{1}{2} b (\Psi^\dagger \Psi)^2 \right]. \quad (3.7)$$

However, the higher-derivative term in (3.4) is relevant in the internal space defined below, and a finite a_0 also automatically provides an ultimate ultraviolet cutoff.

IV. ORIGIN OF FERMION ACTION AND (3+1) DIMENSIONAL SPACETIME

The present theory is based on (1) statistical counting (which ultimately produces the results of (3.4)-(3.7) via the arguments of Appendices A and B) and (2) the geography (or specific features) of our universe, to which we now turn.

The most central assumption is that

$$\Psi_b = \Psi'_0 + \Psi'_b \quad (4.1)$$

where Ψ'_0 contains the order parameter Ψ_0 for a primordial bosonic condensate which is present from the beginning of the universe, and Ψ'_b contains all the other bosonic fields. I.e., in Ψ'_0 only one set of components is nonzero and equal to Ψ_0 , and this set of components is zero in Ψ'_b . The treatment of Appendix C implies that

$$\Psi'_0{}^\dagger \Psi'_b = 0 \quad (4.2)$$

everywhere. (The fields in other representations do not overlap the representation containing Ψ_0 . Fields in the same representation are orthogonal according to (C11) and the comments above (C2) and (C9).)

As mentioned at the end of Section IX, in the simplest description this primordial condensate is essentially an extended version of the singlet scalar boson field in a 16 representation of $SO(10)$. In conventional treatments of $SO(10)$ grand unification, the condensation of this scalar field provides the first stage of the symmetry breakings that ultimately result in the $SU(3) \times SU(2) \times U(1)$ symmetry of the standard model [5]. Here a more general initial symmetry breaking occurs at very high energy, with the resulting vacuum expectation value of the field Ψ_0 (i.e. the order parameter) having multiple components and therefore able to rotate as a function of the coordinates x^M in D dimensions. At the earliest stage of cosmology, as described in detail below, Ψ_0 breaks down to a product of a 2-component field Ψ_{ext} , which is a function of 4 external coordinates x^μ , and an N_g -component order parameter Ψ_{int} , which is primarily a function of $D - 4$ internal coordinates x^m . The initial gauge symmetry (resulting from the rotational symmetry of the internal space, as described in Section V) includes $SO(10)$ – but this symmetry is itself broken to $SU(5) \times U(1)$ by the condensation of Ψ_{ext} . At lower energies (and lower cosmological temperatures) there are further symmetry breakings that are not considered in the present paper.

The action can now be written as

$$S_E = S_{cond} + S_b + S_f + S_{int} \quad (4.3)$$

$$S_{cond} = \int d^D x \left[\frac{1}{2m_0} \partial_M \Psi_0^\dagger \partial_M \Psi_0 - \mu_0 \Psi_0^\dagger \Psi_0 + \frac{1}{2} b (\Psi_0^\dagger \Psi_0)^2 \right] \quad (4.4)$$

$$S_b = \int d^D x \left[\frac{1}{2m_0} \partial_M \Psi_b'^\dagger \partial_M \Psi_b' + (V_0 - \mu_0) \Psi_b'^\dagger \Psi_b' + \frac{1}{2} b (\Psi_b'^\dagger \Psi_b')^2 \right] \quad (4.5)$$

$$S_f = \int d^D x \left[\frac{1}{2m_0} \partial_M \Psi_f^\dagger \partial_M \Psi_f + (V_0 - \mu_0) \Psi_f^\dagger \Psi_f + \frac{1}{2} b (\Psi_f^\dagger \Psi_f)^2 \right] \quad (4.6)$$

$$S_{int} = \int d^D x b (\Psi_f^\dagger \Psi_f) (\Psi_b'^\dagger \Psi_b') \quad (4.7)$$

$$V_0 = b \Psi_0^\dagger \Psi_0. \quad (4.8)$$

In the remainder of this paper, (4.7) and the last terms in (4.5) and (4.6) will be neglected; we are then considering the theory prior to formation of further condensates beyond the primordial Ψ_0 .

For a static condensate we could write $\Psi_0 = n_0^{1/2} \eta_0$, where η_0 is constant, $\eta_0^\dagger \eta_0 = 1$, and $n_0 = \Psi_0^\dagger \Psi_0$ is the condensate density. This picture is too simplistic, however, since the order parameter can exhibit rotations that are analogous to the rotations in the complex plane of the order parameter $\psi_s = e^{i\theta_s} n_s^{1/2}$ for an ordinary superfluid:

$$\Psi_0(x) = U_0(x) n_0(x)^{1/2} \eta_0 \quad , \quad U_0^\dagger U_0 = 1 . \quad (4.9)$$

After an integration by parts in (4.4) (with boundary terms usually neglected in the present paper), extremalization of the action gives the classical equation of motion for the order parameter:

$$-\frac{1}{2m_0} \partial_M \partial_M \Psi_0 + (V_0 - \mu_0) \Psi_0 = 0 . \quad (4.10)$$

Now an important nuance, which requires some references to the gauge potentials A_μ^i , the metric tensor $g_{\mu\nu}$, and the covariant derivative ∇_μ defined below: Because the primordial condensate density is extremely large, (4.10) is assumed to always hold. (More precisely, at normal energies this constraint is imposed in the configurations that are included in the path integral over Ψ_0 .) However, consistent with this constraint, its “phase” and “superfluid velocities” are allowed to vary within the path integral. This means that A_μ^i and $g_{\mu\nu}$ are quantized. I.e., the path integral over the original field Ψ_0 is replaced by path integrals over A_μ^i and $g_{\mu\nu}$ (with the usual gauge-fixing). In the present treatment, we additionally assume, as usual, the minimal case of a torsion-free universe, so that

$$\nabla_\mu e_\nu^\alpha = 0 \quad (4.11)$$

always holds. I.e., the path integral over $g_{\mu\nu}$ is restricted to torsion-free spacetime geometries.

In specifying the geography of our universe, it will be assumed that Ψ_0 can be written as the product of a 2-component external order parameter Ψ_{ext} , which is a function of 4 external coordinates x^μ , and an internal order parameter Ψ_{int} , which is primarily a function of $D - 4$ internal coordinates x^m , but which also varies with x^μ :

$$\Psi_0 = \Psi_{ext}(x^\mu) \Psi_{int}(x^m, x^\mu) \quad (4.12)$$

$$\Psi_{ext}(x^\mu) = U_{ext}(x^\mu) n_{ext}(x^\mu)^{1/2} \eta_{ext} \quad , \quad \mu = 0, 1, 2, 3 \quad (4.13)$$

$$\Psi_{int}(x^m, x^\mu) = U_{int}(x^m, x^\mu) n_{int}(x^m, x^\mu)^{1/2} \eta_{int} \quad , \quad m = 4, \dots, D - 1 \quad (4.14)$$

where again η_{ext} and η_{int} are constant, and $\eta_{ext}^\dagger \eta_{ext} = \eta_{int}^\dagger \eta_{int} = 1$. Here, according to a standard notation, x^μ actually represents the set of x^μ , and x^m the set of x^m .

Let us define external and internal ‘‘superfluid velocities’’ by

$$m_0 v_\mu = -i U_{ext}^{-1} \partial_\mu U_{ext} \quad , \quad m_0 v_m = -i U_{int}^{-1} \partial_m U_{int} . \quad (4.15)$$

The fact that $U_{ext}^\dagger U_{ext} = 1$ implies that $(\partial_\mu U_{ext}^\dagger) U_{ext} = -U_{ext}^\dagger (\partial_\mu U_{ext})$ with $U_{ext}^\dagger = U_{ext}^{-1}$, or $m_0 v_\mu = i (\partial_\mu U_{ext}^\dagger) U_{ext}$, so that

$$v_\mu^\dagger = v_\mu . \quad (4.16)$$

For simplicity, let us first consider the case

$$\partial_\mu U_{int} = 0 \quad (4.17)$$

for which there are separate external and internal equations of motion:

$$\left(-\frac{1}{2m_0} \partial_\mu \partial_\mu - \mu_{ext} \right) \Psi_{ext} = 0 \quad , \quad \left(-\frac{1}{2m_0} \partial_m \partial_m - \mu_{int} + V_0 \right) \Psi_{int} = 0 \quad (4.18)$$

with

$$\mu_{int} = \mu_0 - \mu_{ext} . \quad (4.19)$$

The quantities μ_{int} and V_0 have a relatively slow parametric dependence on x^μ .

When (4.13) and (4.15) are used in (4.18), we obtain

$$\eta_{ext}^\dagger n_{ext}^{1/2} \left[\left(\frac{1}{2} m_0 v_\mu v_\mu - \frac{1}{2m_0} \partial_\mu \partial_\mu - \mu_{ext} \right) - i \left(\frac{1}{2} \partial_\mu v_\mu + v_\mu \partial_\mu \right) \right] n_{ext}^{1/2} \eta_{ext} = 0 \quad (4.20)$$

and its Hermitian conjugate

$$\eta_{ext}^\dagger n_{ext}^{1/2} \left[\left(\frac{1}{2} m_0 v_\mu v_\mu - \frac{1}{2m_0} \partial_\mu \partial_\mu - \mu_{ext} \right) + i \left(\frac{1}{2} \partial_\mu v_\mu + v_\mu \partial_\mu \right) \right] n_{ext}^{1/2} \eta_{ext} = 0 . \quad (4.21)$$

Subtraction gives the equation of continuity

$$\partial_\mu j_\mu^{ext} = 0 \quad , \quad j_\mu^{ext} = n_{ext} \eta_{ext}^\dagger v_\mu \eta_{ext} \quad (4.22)$$

and addition gives the Bernoulli equation

$$\frac{1}{2} m_0 \bar{v}_{ext}^2 + P_{ext} = \mu_{ext} \quad (4.23)$$

where

$$\bar{v}_{ext}^2 = \eta_{ext}^\dagger v_\mu v_\mu \eta_{ext} \quad , \quad P_{ext} = -\frac{1}{2m_0} n_{ext}^{-1/2} \partial_\mu \partial_\mu n_{ext}^{1/2} . \quad (4.24)$$

Since the order parameter Ψ_{ext} in external spacetime has 2 components, its “superfluid velocity” v_μ can be written in terms of the identity matrix σ^0 and Pauli matrices σ^a :

$$v_\mu = v_{\alpha\mu}\sigma^\alpha \quad , \quad \alpha = 0, 1, 2, 3 . \quad (4.25)$$

Let us now transform to a coordinate system in which

$$v_{0k} = v_0^k = v_{a0} = v_a^0 = 0 \quad , \quad k = 1, 2, 3 \quad \text{and} \quad a = 1, 2, 3 \quad (4.26)$$

(with the volume element held constant) so that (4.23) becomes

$$\frac{1}{2}m_0 v_\alpha^\mu v_{\alpha\mu} + P_{ext} = \mu_{ext} . \quad (4.27)$$

To avoid notational complexity we will still use x^μ to label the new coordinates. At this point a physically meaningful metric tensor has not yet been introduced, and the notation in (4.27) merely indicates that $v_\alpha^\mu v_{\alpha\mu}$ is to be kept constant under coordinate transformations. Later in the development, where Lorentz invariance holds (at energies far below m_0), the usual conventions are used for raising and lowering indices.

The transformation to (4.26) is trivial in, e.g., a cosmological model in which the big bang is at the origin of the new coordinates, with the $U(1)$ phase of Ψ_0 varying only with respect to the radial coordinate x^0 , and the “ $SU(2)$ phase” involving the Pauli matrices varying within successive 3-spheres with coordinates x^k , so that v_{ak} has a vortex-like (or instanton-like) configuration.

More generally, the time coordinate x^0 is distinguished from the spatial coordinates x^k in (4.26) because it is the direction of $U(1)$ rather than $SU(2)$ rotations of the order parameter.

As $v_\alpha^\mu v_{\alpha\mu}$ varies, μ_{ext} varies in response, with μ_{int} determined by (4.19).

Now expand Ψ'_b in terms of a complete set of basis functions $\tilde{\psi}_{int}^r$ in the internal space:

$$\Psi'_b(x^\mu, x^m) = \tilde{\psi}_b^r(x^\mu) \tilde{\psi}_{int}^r(x^m) \quad (4.28)$$

with

$$\left(-\frac{1}{2m_0} \partial_m \partial_m - \mu_{int} + V_0 \right) \tilde{\psi}_{int}^r(x^m) = \varepsilon_r \tilde{\psi}_{int}^r(x^m) \quad (4.29)$$

$$\int d^{D-4}x \tilde{\psi}_{int}^{r\dagger}(x^m) \tilde{\psi}_{int}^{r'}(x^m) = \delta_{rr'} \quad (4.30)$$

and with the usual summation over repeated indices in (4.28). For reasons that will become fully apparent below, but which are already suggested by the form of the order parameter,

each $\tilde{\psi}_b^r(x^\mu)$ has two components. As usual, only the zero ($\varepsilon_r = 0$) modes will be kept. (To simplify the presentation, the higher-derivative terms are not explicitly shown in the present section; they will be restored in the next section.) When (4.28)-(4.30) are then used in (4.5) (with the last term neglected), the result is

$$S_b = \int d^4x \tilde{\psi}_b^\dagger \left(-\frac{1}{2m_0} \partial^\mu \partial_\mu - \mu_{ext} \right) \tilde{\psi}_b \quad (4.31)$$

where $\tilde{\psi}_b$ is the vector with components $\tilde{\psi}_b^r$.

Let $\tilde{\psi}_b$ be written in the form

$$\tilde{\psi}_b(x^\mu) = U_{ext}(x^\mu) \psi_b(x^\mu) \quad (4.32)$$

or equivalently

$$\tilde{\psi}_b^r(x^\mu) = U_{ext}(x^\mu) \psi_b^r(x^\mu) . \quad (4.33)$$

Here ψ_b has a simple interpretation: It is the field seen by an observer in the frame of reference that is moving with the condensate. In the present theory, a (very high density) condensate Ψ_0 is present from the beginning of the universe, and the other bosonic and fermionic fields are subsequently born into it. It is therefore natural to define the fields ψ_b^r in the condensate's frame.

Equation (4.32) is, in fact, exactly analogous to rewriting the wavefunction of a particle in an ordinary superfluid moving with velocity v_s : $\tilde{\psi}_{par}(x) = \exp(imv_s x) \psi_{par}(x)$. Here ψ_{par} is the wavefunction in the superfluid's frame of reference.

The frame of the rotating order parameter may be interpreted as the frame of an observer composed of the physical fields defined below (with (4.32) or(4.33) very roughly analogous to shifting to the frame of reference of a human observer as the Solar System revolves around the galaxy, or the Earth revolves within the Solar System, or a point on the Earth follows its daily rotation, although these are coordinate rather than field rotations).

When (4.32) is substituted into (4.31), the result is

$$S_b = \int d^4x \psi_b^\dagger \left[\left(\frac{1}{2} m_0 v^\mu v_\mu - \frac{1}{2m_0} \partial^\mu \partial_\mu - \mu_{ext} \right) - i \left(\frac{1}{2} \partial_\mu v^\mu + v^\mu \partial_\mu \right) \right] \psi_b . \quad (4.34)$$

We assume that, in ordinary circumstances, the extremely dense primordial condensate can be treated as an ‘‘incompressible fluid’’ with the condensate density n_{ext} constant, so that $P_{ext} = 0$ (according to (4.24)), and with

$$\partial_\mu v^\mu = 0 \quad (4.35)$$

(as is consistent with (4.22)). Then (4.26), (4.27), and (4.35) lead to the simplification

$$S_b = - \int d^4x \psi_b^\dagger \left(\frac{1}{2m_0} \partial^\mu \partial_\mu + i v_\alpha^\mu \sigma^\alpha \partial_\mu \right) \psi_b . \quad (4.36)$$

In most of the remainder of the paper it will be assumed that the first term in parentheses is negligible compared to the second for states ψ with energies ~ 1 TeV or less (as would be the case if we had, e.g., $m_0 = a_0^{-1} \gtrsim 10^{15}$ TeV and $v_\alpha^\mu \sim 1$ for $\mu = \alpha$), so that (4.36) reduces to just

$$S_b = \int d^4x \psi_b^\dagger i e_\alpha^\mu \sigma^\alpha \partial_\mu \psi_b \quad (4.37)$$

$$e_\alpha^\mu = -v_\alpha^\mu . \quad (4.38)$$

With this choice all fields are initially right-handed. With the choice $e_\alpha^0 = -v_\alpha^0$, $e_\alpha^k = v_\alpha^k$ all fields would be initially left-handed, as they are for fermions in conventional $SU(5)$ and $SO(10)$ grand-unified theories [9, 10]. It is trivial to change from one convention to the other, of course.

A central feature of the present theory is that the *primitive* bosonic fields ψ_b are 2-component spinors but the final *physical* fields derived from these are the 1-component scalar boson fields defined below – the Higgs and higgson fields of an extended Higgs sector.

To permit local Lorentz transformations as well as general coordinate transformations, let us rewrite (4.37) as

$$S_b = \int d^4x \bar{\mathcal{L}}_b \quad , \quad \bar{\mathcal{L}}_b = \psi_b^\dagger i e_\alpha^\mu \sigma^\alpha \nabla_\mu^R \psi_b \quad (4.39)$$

where ∇_μ^R gives the standard (curved spacetime) covariant derivative for a right-handed Weyl field. The full covariant derivative for a Dirac field, with right-handed and left-handed Weyl components, is given by [44, 45]

$$\nabla_\mu^D = \partial_\mu + i \omega_\mu^{\alpha\beta} \Sigma_{\alpha\beta} . \quad (4.40)$$

(The notation and conventions in this context usually follow those most common in the gravitational and string theory communities, as in Refs. [45–48], rather than the particle physics and field theory communities, as in Ref. [44]. Mainly, the metric tensor convention is $(-+++)$ throughout this paper. However, the Dirac gamma matrices are defined as in most field theory textbooks [49, 50].) The origin and meaning of curvature involving the $\omega_\mu^{\alpha\beta}$ will be considered below. For a vector field, the usual covariant derivative provides invariance

under a coordinate transformation. The vierbein introduced above has both coordinate and tangent-space indices, so [44, 45]

$$\nabla_\mu e_\nu^\alpha = \partial_\mu e_\nu^\alpha - \Gamma_{\mu\nu}^\rho e_\rho^\alpha + \omega_\mu^\alpha{}_\beta e_\nu^\beta . \quad (4.41)$$

In the following we will interpret ∇_μ as an operator which yields the appropriate potentials and generators for each representation – i. e., when acting on each specific field.

The above arguments also hold for fermions, with (in the initial notation)

$$S_f = \int d^D x \left(-\frac{1}{2m_0} \Psi_f^\dagger \partial^M \partial_M \Psi_f - \mu_0 \Psi_f^\dagger \Psi_f + V_0 \Psi_f^\dagger \Psi_f \right) \quad (4.42)$$

$$\Psi_f(x^\mu, x^m) = \tilde{\psi}_f^r(x^\mu) \tilde{\psi}_{int}^r(x^m) \quad (4.43)$$

leading to the result

$$S_f = \int d^4 x \bar{\mathcal{L}}_f \quad , \quad \bar{\mathcal{L}}_f = \psi_f^\dagger i e_\alpha^\mu \sigma^{\alpha} \nabla_\mu \psi_f . \quad (4.44)$$

Eqs. (4.39) and (4.44) are still restricted to the coordinate system of (4.26), with a volume element equal to that of the original coordinate system. Each of these actions has the form

$$S_\varphi = \int d^4 x \bar{\mathcal{L}}_\varphi \quad , \quad \bar{\mathcal{L}}_\varphi = \varphi^\dagger A_\varphi \varphi \quad (4.45)$$

and this form will still hold after mass and interaction terms are added. To permit transformations to a general coordinate system, let

$$\bar{\varphi} = e^{-1/2} \varphi \quad (4.46)$$

$$e = |\det e_\mu^\alpha| = (-\det g_{\mu\nu})^{1/2} \quad (4.47)$$

where the metric tensor $g_{\mu\nu}$ is defined as usual by

$$g^{\mu\nu} = \eta^{\alpha\beta} e_\alpha^\mu e_\beta^\nu \quad , \quad \eta^{\alpha\beta} = \text{diag}(-1, 1, 1, 1) \quad , \quad g_{\mu\nu} g^{\nu\rho} = \delta_\mu^\rho \quad , \quad e_\mu^\alpha e_\beta^\mu = \delta_\beta^\alpha . \quad (4.48)$$

Each operator A_φ considered here or in the following contains a term with one or two factors of ∇_μ . But even this term commutes with $e^{-1/2}$, because (4.11) implies that the covariant derivative of a function of only the vierbein (or metric tensor) is zero:

$$\nabla_\mu e^{-1/2} \varphi = [\nabla_\mu e^{-1/2}] \varphi + e^{-1/2} \nabla_\mu \varphi = e^{-1/2} \nabla_\mu \varphi . \quad (4.49)$$

Then (4.45) can be replaced by

$$S_{\bar{\varphi}} = \int d^4x e \bar{\mathcal{L}}_{\bar{\varphi}} \quad , \quad \bar{\mathcal{L}}_{\bar{\varphi}} = \bar{\varphi}^\dagger A_\varphi \bar{\varphi} . \quad (4.50)$$

According to (4.45), the contribution of φ to the gravitational energy-momentum tensor in the present theory is

$$T_\varphi^{\mu\nu} = 2e^{-1} \frac{\delta \bar{\mathcal{L}}_\varphi}{\delta g_{\mu\nu}} = 2e^{-1} \varphi^\dagger \frac{\delta A_\varphi}{\delta g_{\mu\nu}} \varphi = 2\bar{\varphi}^\dagger \frac{\delta A_\varphi}{\delta g_{\mu\nu}} \bar{\varphi} \quad (4.51)$$

since each term in $\delta A_\varphi / \delta g_{\mu\nu}$, involving functions and covariant derivatives, also commutes with $e^{-1/2}$.

In (4.51) and the remainder of this paper, a covariant derivative ∇_μ is treated as an operator which is independent of $g_{\mu\nu}$, with the specific representation of $\nabla_\mu \varphi$ determined by the character of φ and its geometrical setting. In other words, $\delta / \delta g_{\mu\nu}$ operates on the functions in A_φ but not on ∇_μ .

If $\bar{\varphi}$ is renamed φ for simplicity of notation, (4.45) is changed to

$$S_\varphi = \int d^4x e \bar{\mathcal{L}}_\varphi \quad , \quad \bar{\mathcal{L}}_\varphi = \varphi^\dagger A_\varphi \varphi \quad (4.52)$$

and in particular

$$S_b = \int d^4x e \bar{\mathcal{L}}_b \quad , \quad \bar{\mathcal{L}}_b = \psi_b^\dagger i e_\alpha^\mu \sigma^\alpha \nabla_\mu \psi_b \quad (4.53)$$

$$S_f = \int d^4x e \bar{\mathcal{L}}_f \quad , \quad \bar{\mathcal{L}}_f = \psi_f^\dagger i e_\alpha^\mu \sigma^\alpha \nabla_\mu \psi_f . \quad (4.54)$$

The present theory thus yields the basic form of the standard Lagrangian (4.54) for Weyl fermions. The path integral still has a Euclidean form and the action for bosons is also not yet in standard form, but we will return to these points below.

Notice that, for the fields of (4.53) and (4.54), the stress-energy tensor must now be calculated from

$$T_\varphi^{\mu\nu} = 2\varphi^\dagger \frac{\delta A_\varphi}{\delta g_{\mu\nu}} \varphi \quad (4.55)$$

and not from $T_\varphi^{\mu\nu} = 2e^{-1} \delta(e \bar{\mathcal{L}}_\varphi) / \delta g_{\mu\nu}$. It will be seen below that the two formulas give the same result for particles satisfying their classical equations of motion, but not in general, and not for the zero-point (vacuum) action of fermions and scalar bosons (if used in the conventional way). For fermions or scalar bosons with a bilinear action (4.55) is a natural definition of the stress-energy, or energy-momentum, tensor in a quantum description.

In the present picture spacetime is consistently 4-dimensional with one time coordinate because there are 3 Pauli matrices and one 2×2 unit matrix, and because an instanton-like topological defect can be defined in 4 dimensions with a preferred radial coordinate. An instanton-like defect can also be naturally defined in a $(D - 4)$ -dimensional internal space if, e.g., $D - 4 = 10$. In the present theory, our universe is a product of these two spaces.

V. GAUGE FIELDS

In most of this section and the following two sections, we will temporarily ignore the spin connection and write ∂_μ instead of ∇_μ , to avoid irrelevant complications in notation. Let us now relax assumption (4.17) and allow U_{int} to vary with the external coordinates x^μ . The more general version of (4.10) is satisfied if (4.18) is generalized to

$$\left(-\frac{1}{2m_0}\partial^\mu\partial_\mu - \mu_{ext}\right)\Psi_{ext}(x^\mu)\Psi_{int}(x^m, x^\mu) = 0 \quad (5.1)$$

with Ψ_{int} required to satisfy the internal equation of motion (at each x^μ)

$$\left[\sum_m \frac{1}{2m_0} \left(-\frac{\partial^2}{\partial(x^m)^2} + \frac{a_0^2}{16} \frac{\partial^4}{\partial(x^m)^4}\right) + V_0(x^m) - \mu_{int}\right]\Psi_{int}(x^m, x^\mu) = 0. \quad (5.2)$$

The higher-derivative term of (B14) has been retained and two integrations by parts have been performed. (In order to simplify the notation, we do not explicitly show the weak parametric dependence of μ_{int} , V_0 , and n_{int} on x^μ .) This is a nonlinear equation because (at each x^μ) $V_0(x^m)$ is mainly determined by $n_{int} = \Psi_{int}^\dagger \Psi_{int}$.

The internal basis functions satisfy the more general version of (4.29) with $\varepsilon_r = 0$:

$$\left[\sum_m \frac{1}{2m_0} \left(-\frac{\partial^2}{\partial(x^m)^2} + \frac{a_0^2}{16} \frac{\partial^4}{\partial(x^m)^4}\right) + V_0(x^m) - \mu_{int}\right]\tilde{\psi}_{int}^r(x^m, x^\mu) = 0. \quad (5.3)$$

This is a linear equation because $V_0(x^m)$ is now regarded as a known function.

The full path integral involving (B14) contains all configurations of the fields, including those with nontrivial topologies. In the present theory, the geography of our universe includes a topological defect in the $(D - 4)$ -dimensional internal space which is analogous to a vortex. (See Appendix C.) The standard features of four-dimensional physics arise from the presence of this internal topological defect. For example, it compels the initial gauge symmetry to be $SO(D - 4)$. The $SU(3) \times SU(2) \times U(1)$ structure of the standard model is obtained if $D - 4 = 10$, but $D > 10$ may permit an initially higher gauge group which breaks down to

$SO(10)$ (which itself breaks down to the groups of the standard model) plus a horizontal group that explains the 3 generations of fermions.

The behavior of the condensate and basis functions in the internal space is discussed in Appendices C and D. In (C15), the parameters $\bar{\phi}_i$ specify a rotation of $\Psi_{int}(x^m, x^\mu)$ as the external coordinates x^μ are varied, and according to (C16) the \bar{J}_i satisfy the $SO(D-4)$ algebra

$$\bar{J}_i \bar{J}_j - \bar{J}_j \bar{J}_i = i c_{ij}^k \bar{J}_k . \quad (5.4)$$

For simplicity of notation, let

$$\langle r | Q | r' \rangle = \int d^{D-4} x \tilde{\psi}_{int}^{r\dagger} Q \tilde{\psi}_{int}^{r'} \quad \text{with} \quad \langle r | r' \rangle = \delta_{rr'} \quad (5.5)$$

for any operator Q , and in particular let

$$t_i^{rr'} = \langle r | \bar{J}_i | r' \rangle \quad (5.6)$$

with the matrices $t_i^{rr'}$ (which are constant according to (C17)) inheriting the $SO(D-4)$ algebra:

$$(t_i t_j - t_j t_i)^{rr'} = \sum_{r''} \langle r | \bar{J}_i | r'' \rangle \langle r'' | \bar{J}_j | r' \rangle - \sum_{r''} \langle r | \bar{J}_j | r'' \rangle \langle r'' | \bar{J}_i | r' \rangle \quad (5.7)$$

$$= \langle r | \bar{J}_i \bar{J}_j | r' \rangle - \langle r | \bar{J}_j \bar{J}_i | r' \rangle \quad (5.8)$$

$$= i c_{ij}^k t_k^{rr'} . \quad (5.9)$$

The t_i are the generators in the N_g -dimensional reducible representation determined by the physically significant solutions to (5.3), which spans all the irreducible (physical) gauge representations.

When $x^\mu \rightarrow x^\mu + \delta x^\mu$, Ψ_{int} and $\tilde{\psi}_{int}^r$ rotate together, and (C15) implies that

$$\partial_\mu \tilde{\psi}_{int}^r(x^m, x^\mu) = \frac{\partial \bar{\phi}_i}{\partial x^\mu} \frac{\partial}{\partial \bar{\phi}_i} \tilde{\psi}_{int}^r(x^m, x^\mu) \quad (5.10)$$

$$= -i A_\mu^i \bar{J}_i \tilde{\psi}_{int}^r(x^m, x^\mu) \quad (5.11)$$

where

$$A_\mu^i = \frac{\partial \bar{\phi}_i}{\partial x^\mu} . \quad (5.12)$$

The A_μ^i will be interpreted below as gauge potentials. In other words, the gauge potentials are simply the rates at which the internal order parameter $\Psi_{int}(x^m, x^\mu)$ is rotating as a function of the external coordinates x^μ .

Let us return to the fermionic action (4.42). If (4.43) is written in the more general form

$$\Psi_f(x^\mu, x^m) = \tilde{\psi}_f^r(x^\mu) \tilde{\psi}_{int}^r(x^m, x^\mu) = U_{ext}(x^\mu) \psi_f^r(x^\mu) \tilde{\psi}_{int}^r(x^m, x^\mu) \quad (5.13)$$

we now have, from (4.15) and (5.11),

$$\partial_\mu \Psi_f = U_{ext}(x^\mu) \left(\partial'_\mu + im_0 v_{\alpha\mu} \sigma^\alpha - i A_\mu^i \bar{J}_i \right) \psi_f^r \tilde{\psi}_{int}^r \quad (5.14)$$

where the prime indicates that ∂'_μ does not operate on $\tilde{\psi}_{int}^r$.

Straightforward repetition of the treatment in the preceding section, with inclusion of the additional term involving

$$\int d^{D-4}x \tilde{\psi}_{int}^{r\dagger} \left(-i A_\mu^i \bar{J}_i \right) \tilde{\psi}_{int}^{r'} = -i A_\mu^i \langle r | \bar{J}_i | r' \rangle = -i A_\mu^i t_i^{rr'} \quad (5.15)$$

and with the ψ_f and ψ_b again redefined to include a factor of $e^{-1/2}$, gives

$$S_f = \int d^4x e \bar{\mathcal{L}}_f \quad , \quad \bar{\mathcal{L}}_f = \psi_f^\dagger i e_\alpha^\mu \sigma^\alpha \tilde{D}_\mu \psi_f \quad (5.16)$$

$$S_b = \int d^4x e \bar{\mathcal{L}}_b \quad , \quad \bar{\mathcal{L}}_b = \psi_b^\dagger i e_\alpha^\mu \sigma^\alpha \tilde{D}_\mu \psi_b . \quad (5.17)$$

where

$$\tilde{D}_\mu = \nabla_\mu - i A_\mu^i t_i . \quad (5.18)$$

This is the generalization of (4.53) and (4.54) when the internal order parameter is permitted to vary as a function of the external coordinates x^μ .

For simplicity, we adopt a locally inertial coordinate system in the following two sections, where $e \rightarrow 1$ and the covariant derivative is just

$$D_\mu = \partial_\mu - i A_\mu^i t_i . \quad (5.19)$$

VI. TRANSFORMATION TO LORENTZIAN PATH INTEGRAL: FERMIONS

All of the foregoing is within a Euclidean picture, but we will now show that, in the case of fermions, there is a relatively trivial transformation to the more familiar Lorentzian description. A key point is that the low-energy operator $i e_\alpha^\mu \sigma^\alpha D_\mu$ in S_f is automatically

in the correct Lorentzian form, even though the initial *path integral* is in Euclidean form. It is this fact which permits the following transformation to a Lorentzian path integral. Within the present theory, neither the fields nor the operators (nor the meaning of the time coordinate) need to be modified in performing this transformation.

The operator within S_f can be diagonalized to give

$$S_f = \sum_s \bar{\psi}_f^*(s) a(s) \bar{\psi}_f(s) \quad (6.1)$$

where

$$\psi_f(x) = \sum_s U(x, s) \bar{\psi}_f(s) \quad , \quad \bar{\psi}_f(s) = \int d^4x U^\dagger(x, s) \psi_f(x) \quad (6.2)$$

with

$$ie_\alpha^\mu \sigma^\alpha D_\mu U(x, s) = a(s) U(x, s) \quad (6.3)$$

$$\int d^4x U^\dagger(x, s) U(x, s') = \delta_{ss'} \quad , \quad \sum_s U(x, s) U^\dagger(x', s) = \delta(x - x') \quad (6.4)$$

Here, and in the following, x represents a point in external spacetime, and $U(x, s)$ is a multicomponent eigenfunction. There is an implicit inner product in

$$U^\dagger(x, s) \psi_f(x) = \sum_r U_r^\dagger(x, s) \psi_f^r(x) \quad (6.5)$$

with the $2N_g$ components of $\psi_f(x)$ labeled by $r = 1, \dots, N_g$ (spanning all components of all irreducible gauge representations) and $a = 1, 2$ (labeling the components of Weyl spinors), and with s and (x, r, a) each having N values. Also, the delta function in (6.4) implicitly multiplies the $2N_g \times 2N_g$ identity matrix.

Evaluation of the present Euclidean path integral (a Gaussian integral with Grassmann variables) is then trivial for fermions; as usual,

$$Z_f = \int \mathcal{D}\psi_f^\dagger(x) \mathcal{D}\psi_f(x) e^{-S_f} \quad (6.6)$$

$$= \prod_{x, r, a} \int d\psi_f^{ra*}(x) \int d\psi_f^{ra}(x) e^{-S_f} \quad (6.7)$$

$$= \prod_s z_f(s) \quad (6.8)$$

with

$$z_f(s) = \int d\bar{\psi}_f^*(s) \int d\bar{\psi}_f(s) e^{-\bar{\psi}_f^*(s) a(s) \bar{\psi}_f(s)} \quad (6.9)$$

$$= a(s) \quad (6.10)$$

since the transformation is unitary [49]. Now let

$$Z_f^L = \int \mathcal{D}\bar{\psi}_f^\dagger(s) \mathcal{D}\bar{\psi}_f(s) e^{iS_f} \quad (6.11)$$

$$= \prod_s z_f^L(s) \quad (6.12)$$

where

$$z_f^L(s) = \int d\bar{\psi}_f^*(s) \int d\bar{\psi}_f(s) e^{i\bar{\psi}_f^*(s)a(s)\bar{\psi}_f(s)} \quad (6.13)$$

$$= -ia(s) \quad (6.14)$$

so that

$$Z_f^L = c_f Z_f \quad , \quad c_f = \prod_s (-i) . \quad (6.15)$$

This result holds for the path integral over an arbitrary time interval, with the fields, operator, and meaning of time left unchanged.

The transition amplitude from an initial state to a final state is equal to the path integral between these states, so transition probabilities are the same with the Lorentzian and Euclidean forms of the path integral. This result is consistent with the fact that the classical equations of motion are also the same, since they follow from extremalization of the same action. Furthermore, using the method on pp. 290-291 or 302-303 of Ref. [49], it is easy to show that the magnitude $|G(x, x')|$ of the 2-point function is again the same, so particles propagate the same way in both descriptions. This result is also obtained in Appendix E with a different method.

It may seem strange that the Lorentzian and Euclidean forms of the path integral yield the same physical results, but perusal of the standard arguments in e.g. field theory textbooks shows that the physically significant features of the results derive from the Lorentzian form of the *action* rather than the path integral.

When the inverse transformation from $\bar{\psi}_f$ to ψ_f is performed, we obtain

$$Z_f^L = \int \mathcal{D}\psi_f^\dagger(x) \mathcal{D}\psi_f(x) e^{iS_f} \quad (6.16)$$

with S_f having its form (5.16) in the coordinate representation.

One may perform calculations in either the path-integral formulation or the equivalent canonical formulation, which can now be obtained in the standard way: Let us use the notation \int_a^b to indicate that the fields in a path integral are specified to begin in a state $|a\rangle$

at time t_a and end in state $|b\rangle$ at time t_b , and also to indicate that a path integral showing these limits has its conventional definition (so that it may differ by a normalization constant from Z_f^L as defined above). Then the Hamiltonian H_f is defined by

$$\langle b| U_f(t_b, t_a) |a\rangle = \int_a^b \mathcal{D}\psi_f^\dagger(x) \mathcal{D}\psi_f(x) e^{iS_f} \quad (6.17)$$

$$i\frac{d}{dt}U_f(t, t_a) = H_f(t)U_f(t, t_a) \quad , \quad U_f(t_a, t_a) = 1 . \quad (6.18)$$

I.e., the time evolution operator $U_f(t_b, t_a)$ is defined to have the same effect as the path integral over intermediate states, and it is then straightforward to reverse the usual logic which leads from canonical quantization to path-integral quantization [49, 51]. The Schrödinger equation

$$i\frac{d}{dt}|\psi_f(t)\rangle = H_f(t)|\psi_f(t)\rangle \quad (6.19)$$

follows from (6.18) and

$$|\psi_f(t)\rangle = U_f(t, t_a)|\psi_f(t_a)\rangle . \quad (6.20)$$

The Hamiltonian H_f can be written in terms of field operators, and then creation and destruction operators, in a standard way (and essentially just as for the harmonic oscillator).

VII. TRANSFORMATION TO LORENTZIAN PATH INTEGRAL: BOSONS

For bosons we can again perform the transformation (6.2) to obtain

$$S_b = \sum_s \bar{\psi}_b^*(s) a(s) \bar{\psi}_b(s) . \quad (7.1)$$

In Appendix F it is shown that this action can be put into a form that involves boson fields Φ_r which are intermediate in transforming the original unphysical fields ψ_b^r into the physical scalar boson fields ϕ_R , φ_S , and F_R defined below. The resulting Lagrangian is

$$\bar{\mathcal{L}}_\Phi = \frac{1}{2}\Phi^\dagger(x) i\bar{\sigma}^\mu D_\mu i\sigma^\nu D_\nu \Phi(x) + \frac{1}{2}\Phi^\dagger(x) i\sigma^\mu D_\mu i\bar{\sigma}^\nu D_\nu \Phi(x) . \quad (7.2)$$

To simplify the mathematics below, it is convenient to temporarily write

$$\Phi_b = \begin{pmatrix} \Phi \\ \bar{\Phi} \end{pmatrix} \quad (7.3)$$

and to use the same Weyl representation as is used for Dirac fermions, with

$$\gamma^\mu = \begin{pmatrix} 0 & \sigma^\mu \\ \bar{\sigma}^\mu & 0 \end{pmatrix} \quad (7.4)$$

so that (7.2) can be written as

$$\bar{\mathcal{L}}_\Phi = -\frac{1}{2}\Phi_b^\dagger(x)\gamma^\mu D_\mu\gamma^\nu D_\nu\Phi_b(x) \quad (7.5)$$

$$= -\frac{1}{2}\Phi_b^\dagger(x)\not{D}^2\Phi_b(x) . \quad (7.6)$$

A standard result [49, 50] which can be extended to the nonabelian case (see (16.107) of Ref. [49]) gives

$$-\not{D}^2 = D^\mu D_\mu - S^{\mu\nu}F_{\mu\nu} \quad (7.7)$$

with the present convention for the metric tensor. (See pp. 173-174 of [50] for this result and those immediately below.) Here the field strength tensor $F_{\mu\nu}$ spans all the irreducible (physical) gauge representations, and the second term can be rewritten with “magnetic” and “electric” fields B_k and E_k defined by

$$F_{kk'} = -\varepsilon_{kk'k''}B_{k''} \quad , \quad F_{0k} = E_k \quad (7.8)$$

since [50]

$$-S^{\mu\nu}F_{\mu\nu} = \begin{pmatrix} (\vec{B} + i\vec{E}) \cdot \vec{\sigma} & 0 \\ 0 & (\vec{B} - i\vec{E}) \cdot \vec{\sigma} \end{pmatrix} \quad (7.9)$$

where $a \cdot b = a_k b^k$. (Recall that $\mu = 0, 1, 2, 3$ and $k = 1, 2, 3$.) We then obtain

$$\bar{\mathcal{L}}_\Phi = \Phi^\dagger(x) D^\mu D_\mu \Phi(x) + \Phi^\dagger(x) \vec{B} \cdot \vec{\sigma} \Phi(x) . \quad (7.10)$$

The second term above is invariant under a rotation, but not under a boost, so it still breaks Lorentz invariance for the intermediate bosonic fields in Φ . This issue is considered in the next section, where the truly physical scalar boson fields are defined.

All of the above is in a locally inertial frame of reference. Now let us return to a general coordinate system and initially assume no nongravitational gauge fields, so that $D_\mu \rightarrow \nabla_\mu$ and

$$\bar{\mathcal{L}}_\Phi = -\frac{1}{2}\Phi_b^\dagger(x)\underline{\gamma}^\mu\nabla_\mu\underline{\gamma}^\nu\nabla_\nu\Phi_b(x) = -\frac{1}{2}\Phi_b^\dagger(x)\not{\nabla}^2\Phi_b(x) \quad (7.11)$$

where

$$\not{\Psi} \equiv \underline{\gamma}^\mu \nabla_\mu \quad , \quad \underline{\gamma}^\mu = \begin{pmatrix} 0 & \underline{\sigma}^\mu \\ \overline{\underline{\sigma}}^\mu & 0 \end{pmatrix} \quad (7.12)$$

with $\underline{\sigma}^\mu = e_\alpha^\mu \sigma^\alpha$ and $\overline{\underline{\sigma}}^\alpha = e_\alpha^\mu \overline{\sigma}^\alpha$. But

$$-\not{\Psi}^2 = g^{\mu\nu} \nabla_\mu \nabla_\nu - \frac{1}{4} R \quad (7.13)$$

follows from e.g. (5.293) of Ref. [44] with our $(-+++)$ convention for the metric tensor, where R is the curvature scalar in 4-dimensional spacetime, so

$$\overline{\mathcal{L}}_\Phi = \Phi^\dagger(x) \left(g^{\mu\nu} \nabla_\mu \nabla_\nu - \frac{1}{4} R \right) \Phi(x) . \quad (7.14)$$

(See also (15.5.5) of Ref. [45], with the present convention for the metric tensor, but the opposite convention of their p. 274 for the Dirac gamma matrices.)

It is well known that a scalar field can have a Lagrangian which includes a term $-\xi R$ (as in (2.35) of Ref. [44]), with ξ undetermined in standard physics, but the present picture yields $\xi = 1/4$.

VIII. SCALAR BOSONS

Let Φ_r be one of the intermediate 2-component spin 1/2 bosonic fields in Φ . We can construct physical fields, which satisfy Lorentz invariance, by eliminating the anomalous second term of (7.10) in either of two ways.

Higgs fields can be constructed by combining 2-component fields Φ_r and $\Phi_{r'}$ with the same gauge quantum numbers but opposite spins (and equal amplitudes):

$$\Phi_R = \begin{pmatrix} \Phi_r \\ \Phi_{r'} \end{pmatrix} \quad (8.1)$$

so that

$$\begin{aligned} \Phi_R^\dagger(x) \overrightarrow{\sigma} \Phi_R(x) &= \begin{pmatrix} \Phi_r^\dagger & \Phi_{r'}^\dagger \end{pmatrix} \begin{pmatrix} \overrightarrow{\sigma} & 0 \\ 0 & \overrightarrow{\sigma} \end{pmatrix} \begin{pmatrix} \Phi_r \\ \Phi_{r'} \end{pmatrix} \\ &= \Phi_r^\dagger \overrightarrow{\sigma} \Phi_r + \Phi_{r'}^\dagger \overrightarrow{\sigma} \Phi_{r'} \end{aligned} \quad (8.2)$$

$$= 0 . \quad (8.3)$$

In this case we write for each gauge component

$$\Phi_R(x) = \phi_R(x) \xi_R \quad \text{with} \quad \xi_R^\dagger \xi_R = 1 \quad (8.4)$$

where ξ_R has 4 constant components and $\phi_R(x)$ is a 1-component complex amplitude which is roughly analogous to the Higgs/amplitude modes predicted and observed in superconductors (and possibly other systems) [52–54]. If this can be done for a full gauge multiplet $\tilde{\Phi}_R$ containing Φ_R , yielding a multiplet $\tilde{\phi}_R$ of 1-component scalar boson fields, then (7.10) implies that

$$\tilde{S}_R = \int d^4x e \tilde{\phi}_R^\dagger(x) D^\mu D_\mu \tilde{\phi}_R(x) \quad (8.5)$$

is the action for this multiplet of Higgs fields (before masses and other interaction terms are added). More generally, we can regard the generators in D_μ as operators (rather than matrices in a specific representation) and write

$$S_R = \int d^4x e \bar{\mathcal{L}}_R \quad , \quad \bar{\mathcal{L}}_R = \phi_R^\dagger(x) D^\mu D_\mu \phi_R(x) \quad . \quad (8.6)$$

If a Higgs field has a vacuum expectation value V_R , we can also write

$$\phi_R(x) = V_R + H_R \quad (8.7)$$

where the excitation H_R corresponds to a Higgs boson. (Here and in our more recent papers [21, 22], in deference to the notation of Ref. [1], the notation of our previous papers [19, 20] has been reversed, so that $H \leftrightarrow$ Higgs boson and $h \leftrightarrow$ higgson.)

Higgson fields can be constructed by combining a 2-component field Φ_s and a charge conjugate field Φ_s^c (with opposite gauge quantum numbers but here taken to have the same spin):

$$\Phi_S = \begin{pmatrix} \Phi_s \\ \Phi_s^c \end{pmatrix} \quad (8.8)$$

The subscripts s and S are used in this context to avoid confusion.

Φ_s and Φ_s^c have opposite expectation values for the generators t^j (which are again treated as operators rather than matrices):

$$\Phi_s^c t^j \Phi_s^c = -\Phi_s^\dagger t^j \Phi_s \quad (8.9)$$

Since

$$B_{k''} = -\varepsilon_{k''kk'} F_{kk'} \quad , \quad F_{kk'} = F_{kk'}^j t^j \quad (8.10)$$

we have

$$\Phi_S^\dagger(x) \vec{B} \Phi_S(x) = \begin{pmatrix} \Phi_S^\dagger & \Phi_S^{c\dagger} \end{pmatrix} \begin{pmatrix} \vec{B} & 0 \\ 0 & \vec{B} \end{pmatrix} \begin{pmatrix} \Phi_S \\ \Phi_S^c \end{pmatrix} \quad (8.11)$$

$$= \Phi_S^\dagger \vec{B} \Phi_S + \Phi_S^{c\dagger} \vec{B} \Phi_S^c \quad (8.12)$$

$$= 0 . \quad (8.13)$$

As in Ref. [19] the complex 4-component higgson field Φ_S can be separated into real and imaginary parts for each of its two polarizations, and each of the four resulting 4-component real fields will be called $\overline{\Phi}_S$ (in a slight change of notation from that of Ref. [19], with $H \rightarrow \overline{\Phi}_S$). Before mass is acquired, the Lagrangian for such a field in the electroweak sector contains the terms [19]

$$\overline{\mathcal{L}}_S^0 = \overline{\Phi}_S^\dagger \partial^\mu \partial_\mu \overline{\Phi}_S \quad , \quad \overline{\mathcal{L}}_S^Z = -\frac{g_Z^2}{4} \overline{\Phi}_S^\dagger Z^\mu Z_\mu \overline{\Phi}_S \quad , \quad \overline{\mathcal{L}}_S^W = -\frac{g^2}{2} \overline{\Phi}_S^\dagger W^{\mu+} W_\mu^- \overline{\Phi}_S \quad (8.14)$$

where $g_Z = g/\cos\theta_W$. Since $\overline{\Phi}_S$ has spin 0 and is real [19], we can write

$$\overline{\Phi}_S(x) = \varphi_S(x) \zeta_S \quad \text{with} \quad \zeta_S^\dagger \zeta_S = 1 \quad (8.15)$$

where ζ_S has 4 constant components which incorporate the quantum numbers of $\overline{\Phi}_S$, and $\varphi_S(x)$ is again an amplitude. Each Φ_S , with four degrees of freedom, has first become four independent 4-component real fields $\overline{\Phi}_S$, and then four 1-component real fields φ_S , which are the physically relevant higgson fields. To allow for the possibility of a vacuum expectation value, we can write

$$\varphi_S(x) = v_S + h_S \quad (8.16)$$

where the excitation h_S corresponds to a higgson. (But it is assumed that $v_S = 0$ for the lowest-mass higgson field, which yields the dark matter candidate below.)

Each higgson field φ_S can be treated (and quantized) like a standard real scalar field, but with no net quantum numbers and no interactions other than those in the reduced version of (8.14):

$$\overline{\mathcal{L}}_S^Z = -\frac{g_Z^2}{4} \varphi_S Z^\mu Z_\mu \varphi_S \quad , \quad \overline{\mathcal{L}}_S^W = -\frac{g^2}{2} \varphi_S W^{\mu+} W_\mu^- \varphi_S . \quad (8.17)$$

Note that (i) the second term in (7.14) is obtained for all scalar boson fields and (ii) gravitational and gauge cross terms in the full \mathcal{D}^2 of (9.1) below ultimately give zero in all physical fields, because the factor related to $\Sigma_{\alpha\beta}$ (containing Pauli matrices) produces cancellation in Higgs fields and the factor involving A_μ produces cancellation in higgson fields. For higgson fields we then have, in a general coordinate system,

$$\overline{\mathcal{L}}_S^h = \varphi_S \left(g^{\mu\nu} \nabla_\mu \nabla_\nu - \frac{1}{4} R \right) \varphi_S + \overline{\mathcal{L}}_S^Z + \overline{\mathcal{L}}_S^W . \quad (8.18)$$

Also note that some of the Higgs fields that are normally expected in a grand-unified theory may not exist, because their degrees of freedom may instead have been absorbed in higgson fields when the mass eigenstates are formed. This fact may be relevant to the well-known doublet-triplet problem [10].

The present reinterpretation of scalar boson sectors can be extended to supersymmetric partners [32]. A detailed discussion is beyond the scope of the present paper, but the main results of Ref. [32], and of quantitative calculations of detection cross-sections by student collaborators [68], will be mentioned in the Conclusion.

IX. FUNDAMENTAL ACTION FOR FERMIONS AND SCALAR BOSONS

In the following we change to a simplified notation with $\tilde{D}_\mu \rightarrow D_\mu$:

$$D_\mu = \nabla_\mu - iA_\mu \quad (9.1)$$

where ∇_μ and A_μ are regarded as operators which yield the appropriate generators and potentials for each representation. When the various components in the preceding sections are assembled, the total action for fermion fields ψ_f^r , Higgs fields ϕ_R , auxiliary fields F_R , and higgson fields φ_S is

$$S_{matter} = \int d^4x e \overline{\mathcal{L}}_{matter} \quad (9.2)$$

$$\begin{aligned} \overline{\mathcal{L}}_{matter} = \sum_r \psi_f^{r\dagger}(x) i e_\alpha^\mu \sigma^\alpha D_\mu \psi_f^r(x) + \sum_R \phi_R^\dagger(x) \left(g^{\mu\nu} D_\mu D_\nu - \frac{1}{4} R \right) \phi_R(x) \\ + \sum_R F_R^\dagger(x) F_R(x) + \sum_S \overline{\mathcal{L}}_S^h \end{aligned} \quad (9.3)$$

or

$$\overline{\mathcal{L}}_{matter} = \psi_f^\dagger i e_\alpha^\mu \sigma^\alpha D_\mu \psi_f + \phi^\dagger \left(g^{\mu\nu} D_\mu D_\nu - \frac{1}{4} R \right) \phi + F^\dagger F + \overline{\mathcal{L}}^h \quad (9.4)$$

after transformation to a general coordinate system, and before masses and further interactions result from symmetry breakings, radiative corrections, and other effects.

The spin 1/2 fermion fields in ψ_f span the various physical representations of the most fundamental gauge group, which must be $SO(D-4)$ in the present theory. (More precisely, the group for standard model fermions is $Spin(D-4)$, but $SO(D-4)$ is conventional terminology.) F_R is the amplitude of a component of \mathcal{F} , defined in the same way as ϕ_R . The last term in (9.3) or (9.4) contains the action for all higgsion fields, including those which lie outside the electroweak sector. All of (9.3) has the basic form for a Lorentz-invariant, gauge-invariant, and prima facie even potentially supersymmetric action which is also invariant under general coordinate transformations. (The gauge invariance of the last term is discussed in Ref. [32]: If the gauge fields are transformed, it can be seen that the scalar boson fields change appropriately to keep the action invariant, and this is what is meant by gauge invariance.) However, the present theory implies a radically modified version of susy, which has the necessary *physical* consequences of conventional susy – protection of the Higgs mass and unification of coupling constants – as well as Lorentz and gauge invariance, but is *mathematically* quite different; see Ref. [32] for a detailed discussion.

Gravitational and gauge curvatures must ultimately originate from a background (in the path integral) of “rapidly fluctuating” 4-dimensional topological defects (analogous to vortices and vortex rings, or extended and closed flux tubes) associated with the gauge potentials of (5.12) and the vierbein of (4.38). Here “rapidly fluctuating” means that the A_μ and $g_{\mu\nu}$ (or e^μ_α) are actually averages over many topological configurations of the field Ψ_0 . As mentioned below (4.10), the path integral over all these configurations is replaced by a path integral over the A_μ and $g_{\mu\nu}$ with an effective action. The gauge and gravitational fields then vary over all possibilities, and this is how these force fields are quantized in the present theory. A detailed discussion of the topological defects is beyond the scope of this paper, but all that is required here is the fact that topological defects permit the curvature associated with the the gauge potentials and vierbein to be nonzero. I.e., the gauge curvature (in a locally inertial frame, and with the coupling constant no longer absorbed into A_μ)

$$F_{\mu\nu}^i = \partial_\mu A_\nu^i - \partial_\nu A_\mu^i + g f_{jk}^i A_\mu^j A_\nu^k \quad (9.5)$$

and the gravitational spin connection and curvature (see p. 274 of Ref. [45])

$$\omega_{\mu}^{\alpha\beta} = \frac{1}{2}e^{\nu\alpha}(\partial_{\mu}e_{\nu}^{\beta} - \partial_{\nu}e_{\mu}^{\beta}) - \frac{1}{2}e^{\nu\beta}(\partial_{\mu}e_{\nu}^{\alpha} - \partial_{\nu}e_{\mu}^{\alpha}) - \frac{1}{2}e^{\rho\alpha}e^{\sigma\beta}(\partial_{\rho}e_{\sigma\gamma} - \partial_{\sigma}e_{\rho\gamma})e_{\mu}^{\gamma} \quad (9.6)$$

$$R_{\mu\nu}{}^{\alpha}{}_{\beta} = \partial_{\mu}\omega_{\nu}{}^{\alpha}{}_{\beta} - \partial_{\nu}\omega_{\mu}{}^{\alpha}{}_{\beta} + [\omega_{\mu}, \omega_{\nu}]_{\beta}{}^{\alpha} \quad , \quad R_{\mu\nu}{}^{\alpha}{}_{\beta} = e_{\sigma}^{\alpha}e_{\beta}^{\tau}R_{\mu\nu}{}^{\sigma}{}_{\tau} \quad (9.7)$$

originate from vortex-like configurations in the same way that the vorticity of a superfluid

$$\omega_{k\ell}^s = \partial_k v_{\ell}^s - \partial_{\ell} v_k^s \quad \text{or} \quad \omega_z^s = \partial_x v_y^s - \partial_y v_x^s \quad (9.8)$$

originates from ordinary vortices, as pointed out by Feynman and Onsager. The simplest case is a magnetic field with

$$B_z = \partial_x A_y - \partial_y A_x \quad (9.9)$$

but all the force fields above have the same basic form. In each case, the curvature is nonzero in a region penetrated by flux lines that are interpreted as vortex lines.

From a more fundamental point of view, it is not necessary to consider specific topological defects, since, as mentioned below (4.10), the path integral automatically includes all possible configurations of the condensate order parameter Ψ_0 consistent with this equation of motion, and thus all physically independent (and physically acceptable) values of the gauge potentials and metric tensor. These configurations of Ψ_0 span all topologies, including those with vortex-like and monopole-like topological defects. The path integral over Ψ_0 can then be replaced by a path integral over the gauge potentials A_{μ}^i and metric tensor $g_{\mu\nu}$, with a global transformation of integration variables

$$d\Psi_0^I(x) = \sum_{x'} \sum_{\mu} \sum_i \frac{\partial \Psi_0^I(x)}{\partial A_{\mu}^i(x')} dA_{\mu}^i(x') + \sum_{x'} \sum_{\mu} \sum_{\nu} \frac{\partial \Psi_0^I(x)}{\partial g_{\mu\nu}(x')} dg_{\mu\nu}(x') \quad (9.10)$$

where I labels a component of the full order parameter. Here, in this one context, x represents all of the D coordinates x^M , but x' is limited to the 4 external spacetime coordinates x'^{μ} . When (9.10) is substituted for each component of Ψ_0 , one obtains the path integral over the gauge and gravitational fields. (If the values of x' were continuous, $\sum_{x'}$ would be replaced by $\int d^4x'$ and the partial derivatives by functional derivatives. But in the present description a path integral is over discrete spacetime points.) A detailed treatment is beyond the scope of the present paper, but it is only required that such a transformation is possible.

In Appendix C it is pointed out that spinorial fields $\tilde{\psi}_{sp}^r$ in the internal space are appropriate basis functions for ordinary fermion and sfermion fields, and for the primordial

condensate Ψ_0 . If we ignore further quantum numbers arising from a larger internal space for $D - 4 > 10$, the rotational symmetry of a 10-dimensional internal space implies $SO(10)$ gauge symmetry in external spacetime, according to the arguments of Section V. Since the vacuum should initially have no quantum numbers corresponding to those of the standard model, Ψ_0 must then be the singlet field in a spinorial representation of $SO(10)$ [5] – but with the 2-component description of (4.13).

X. COSMOLOGICAL CONSTANT, GRAVITATIONAL ACTION, AND DARK MATTER

A. Cosmological constant and gravitational action

In conventional physics, the contribution of fermion and scalar boson fields to the vacuum energy corresponds to a Lagrangian $e\bar{\mathcal{L}}_{vac}$, with $\bar{\mathcal{L}}_{vac}$ constant. The resulting gravitational stress-energy tensor is

$$T_{vac}^{\mu\nu} = 2e^{-1}\delta(e\bar{\mathcal{L}}_{vac})/\delta g_{\mu\nu} = g^{\mu\nu}\bar{\mathcal{L}}_{vac} \quad , \quad \bar{\mathcal{L}}_{vac} = -(8\pi\ell_P^2)^{-1}\Lambda \quad (10.1)$$

since

$$\delta e = \frac{1}{2} e g^{\mu\nu} \delta g_{\mu\nu} \quad (10.2)$$

where ℓ_P is the Planck length, and this produces a term $\Lambda g_{\mu\nu}$ in the Einstein field equations with a cosmological constant Λ .

In the present description, on the other hand, the left-hand side of (4.51) implies that

$$T_{vac}^{\mu\nu} = 2e^{-1}\delta\bar{\mathcal{L}}_{vac}/\delta g_{\mu\nu} = 0 \quad (10.3)$$

so there is no direct contribution to a cosmological constant from these fields.

The predictions of the present theory are identical to those of standard general relativity for the motion of all particles and waves in gravitational fields, and for gauge bosons acting as a source of gravity. We will now show that they are also the same for matter acting as a classical gravitational source. The usual energy-momentum tensor is

$$T_{\varphi}^{\mu\nu} = 2e^{-1}\frac{\delta(e\bar{\mathcal{L}}_{\varphi})}{\delta g_{\mu\nu}} = 2\bar{\mathcal{L}}_{\varphi}e^{-1}\frac{\delta e}{\delta g_{\mu\nu}} + 2\frac{\delta\bar{\mathcal{L}}_{\varphi}}{\delta g_{\mu\nu}} = 2\frac{\delta\bar{\mathcal{L}}_{\varphi}}{\delta g_{\mu\nu}} = 2\varphi^{\dagger}\frac{\delta A_{\varphi}}{\delta g_{\mu\nu}}\varphi \quad (10.4)$$

since the bilinear forms of (9.3) – or similar bilinear forms including mass and interaction terms, or for composite objects like protons or planets – imply that $\overline{\mathcal{L}}_\varphi = 0$ if the classical equations of motion are satisfied. (Classical in the present context means that particles or composite bodies remain on the mass shell, and the energy-momentum tensor here corresponds to the energy and momentum of quantum fields, particles, or composite objects satisfying their quantum equations of motion, which yield classical trajectories according to Ehrenfest’s theorem.) Since the result of (10.4) is identical to that of (4.55), the role of classical matter in acting as a source of gravity is unchanged.

The predictions of classical general relativity are thus unchanged in the present picture, except that the cosmological constant of (10.3) is zero if the vacuum Lagrangian density $\overline{\mathcal{L}}_{vac}$ is taken to be fixed.

On the other hand, $\overline{\mathcal{L}}_{vac}$ is not really fixed, since the fields in the vacuum will be modified when they are forced to accommodate the curvature associated with the gauge potentials of (5.12) and the vierbein of (4.38) (or metric tensor of (4.48)). Within the present description, the Maxwell-Yang-Mills action

$$S_g = \int d^4x \mathcal{L}_g \quad , \quad \mathcal{L}_g = -\frac{1}{4} g_0^{-2} e g^{\mu\rho} g^{\nu\sigma} F_{\mu\nu}^i F_{\rho\sigma}^i \quad (10.5)$$

and the Einstein-Hilbert action

$$S_G = \int d^4x \mathcal{L}_G \quad , \quad \mathcal{L}_G = (16\pi\ell_P^2)^{-1} e R \quad (10.6)$$

must result from this modification of the vacuum states (and the terms in their action).

This form is analogous to the shift in the free energy

$$\Omega(B) - \Omega(0) = \frac{e^2 A}{24\pi m c^2} B^2 \quad , \quad (10.7)$$

when electrons in a metal (with area A) respond to the gauge curvature of an applied magnetic field \vec{B} , exhibiting Landau diamagnetism.

It appears that it is nontrivial to obtain (10.5) and (10.6) from a proper calculation, and that a detailed knowledge of the vacuum states is required. However, (i) these states will surely be even qualitatively modified when they are perturbed by the curvature of the vierbein and gauge fields (as are the states of electrons in a metal), (ii) (10.5) and (10.6) are the simplest forms consistent with the symmetries of the vacuum (including invariance under coordinate, Lorentz, and gauge transformations), and (iii) within the present picture (10.5) and (10.6) can originate only from the response of the vacuum to external fields.

With this interpretation, \mathcal{L}_g must necessarily vanish when these fields vanish – i.e., in the vacuum itself:

$$\langle \mathcal{L}_g \rangle_{vac} = 0. \quad (10.8)$$

This means that when (10.5) is quantized, the field operators must be normal-ordered. It follows that there is no cosmological constant resulting from the gauge fields. On the other hand, virtual processes will still be affected by a change in their boundary conditions; a detailed treatment of this aspect, and of the observed Casimir effect [38–40], would be inappropriately long here, but see the discussion of this point in Section II.

B. Dark matter

There are a vast number of hypothetical dark matter candidates, most of which do not have well-defined masses or couplings, and many of which have already been ruled out by experiment – or at least found to be subdominant species in a multicomponent scenario. For example, the simplest susy models which have “natural” values for the parameters, and which are also compatible with limits from the LHC, are found to be in disagreement with both the abundance of dark matter and the limits from direct-detection experiments [55–59] – if the lightest supersymmetric particle (LSP) is assumed to be the dominant constituent. But, as mentioned in Section II, the present picture requires (unconventional and broken) susy at some energy scale, and the LSP (as a subdominant component) can stably coexist with the present dark matter candidate – the lightest higgson, h^0 .

Fortunately the enthusiasm of experimentalists searching for new physics has not been notably diminished by the past lack of success. For example, LZ and XENONnT have begun taking data with larger and more sophisticated detectors than those of past experiments. They should ultimately be able to detect a dark matter WIMP with a mass of $\sim 50 \text{ GeV}/c^2$ if the Xe collision cross-section is larger than $\sim 1.4 \times 10^{-48} \text{ cm}^2$ [61–66], within roughly the next 5 years. The projected future detector DARWIN [67] will have an even greater sensitivity.

For the WIMP predicted by the present theory – the lowest-mass higgson of Section VIII – this cross-section is estimated to be $\sim 10^{-48} \text{ cm}^2$ [21], with a mass of about $70 \text{ GeV}/c^2$, so it may be barely within reach of both LZ and XENONnT.

More generally, this dark matter candidate is consistent with all current experiments, and

observable in the near or foreseeable future through a wide variety of direct, indirect, and collider detection experiments. To review the conclusions of Refs. [19–22]: This particle is unique in that it has precisely defined couplings and a well-defined mass, providing specific cross-sections and other experimental signatures as targets for clean experimental tests. It has not yet been detected because it has no interactions other than second-order gauge couplings, to W and Z bosons. However, these weak couplings are still sufficient to enable observation by experiments which are currently taking data and planned for the foreseeable future.

This lightest higgson h^0 is stable because of the form of the interaction in (8.17): It can radiate gauge bosons, annihilate into gauge bosons, scatter via exchange of gauge bosons, and be created in pairs, but not decay, since a single initial h^0 implies a final state containing h^0 and two gauge bosons.

The cross-section for collider detection at LHC energies is small, ~ 10 femtobarn [68], but observation may ultimately be achievable at the high-luminosity LHC – if it can reach $\sim 3000 \text{ fb}^{-1}$ of integrated luminosity – and should certainly be within reach of the even more powerful colliders now being planned.

There is already a strong case that the present dark matter candidate has been observed via indirect detection: Several analyses of gamma rays from the Galactic center, observed by Fermi-LAT [73–78], and of antiprotons, observed by AMS-02 [79–83], have shown consistency with the interpretation that these result from annihilation of dark matter particles having approximately the same mass and annihilation cross-section as the present candidate.

Finally, there is consistency with the observations of Planck, which have ruled out various possible candidates with larger masses and cross-sections [84].

XI. CONCLUSION

Starting with the simplest imaginable picture, and interpreting our universe as the product of two spaces with topological singularities, we obtain the following results: 4-dimensional spacetime with one time coordinate; spin 1/2 fermion and spin zero boson fields defined on this spacetime; path-integral quantization of these fields; gauge fields and a fundamental gauge theory which is necessarily $SO(D-4)$; correct couplings of matter fields to the gauge fields; a gravitational vierbein; correct couplings of matter fields to gravity;

Lorentz invariance; broken supersymmetry; and elimination of the usual enormous cosmological constant.

The theory is ultimately founded on counting arguments with positive integers, and the continuum is only an approximation to a discrete set of (closely spaced) points. The mathematical foundations are therefore consistent and well-defined, including the meaning of path integrals. Singularities are not a problem in the present theory, because the density of the primordial condensate underlying standard physics goes to zero at each point where there is a singularity.

However, the above are primarily theoretical issues, and the ultimate objective of any theory should be testable experimental predictions. The predictions already obtained for the present theory, obtained here and elsewhere [19–22, 32, 68], can now be summarized:

(1) A dark matter WIMP with well-defined mass and couplings, which is currently consistent with all experiments and observations. It should be within reach of direct-detection experiments in the near future and collider-detection experiments on a longer time scale, and it may have already been observed via indirect detection. There are no free parameters and the predictions for cross-sections are quantitative.

(2) The full range of superpartners in a modified version of susy, possibly observable in the near future through a modified phenomenology [32]. Quantitative predictions of cross-sections for LHC and e^+e^- production of sparticles have been obtained via MadGraph by Tallman et al. and others [68], and will be published separately.

(3) A further enlargement of the complete particle spectrum, because the full $16 + \overline{16}$ fundamental gauge representation is required for both fermion and sfermion fields [32], rather than just the conventional 16. (Since all fields have the same initial chirality in the present theory, the 16 and $\overline{16}$ are not equivalent.) The arguments of Ref. [32] require that many of these new particles have masses that are not far above the TeV scale.

It should be emphasized, in conclusion and in anticipating future work, that many remaining fundamental problems are not considered here. Examples are the origin of Yukawa couplings, the origin of the Higgs boson mass, the nature of dark energy, an inflation-like mechanism in the early universe, a new mechanism for CP violation to explain the prevalence of matter over antimatter in the current universe, $3-5\sigma$ results of various experiments and observations which suggest that the standard models of both particle physics and cosmology need to be extended [1, 85], and the values of the fundamental gauge and gravitational

coupling constants, g_0 and G , which in the present theory would require a detailed understanding of the very complex vacuum fields after multiple symmetry breakings at various energy scales. These problems are all challenging, but the present picture does not appear to be inconsistent with potential solutions.

Appendix A: Primordial fields and action

We begin with the initial picture of Section III, in which coordinates and fields are defined by (3.1), (3.2), and (3.3).

Let $S(x)$ be the entropy at a fixed point x , as defined by $S(x) = \log W(x)$. Here $W(x)$ is the total number of microstates for fixed occupation numbers n_i : $W(x) = N(x)!/\prod_i n_i(x)!$, with

$$N(x) = \sum_i n_i(x) \quad , \quad i = 1, 2, \dots, d. \quad (\text{A1})$$

The total number of available microstates for all points x is $W = \prod_x W(x)$, so the total entropy is

$$S = \sum_x S(x) \quad , \quad S(x) = \log \Gamma(N(x) + 1) - \sum_i \log \Gamma(n_i(x) + 1). \quad (\text{A2})$$

We will see below that $n_k(x)$ can be approximately treated as a continuous variable when it is extremely large, with

$$\frac{\partial S}{\partial n_k(x)} = \psi(N(x) + 1) - \psi(n_k(x) + 1) \quad (\text{A3})$$

$$\frac{\partial^2 S}{\partial n_{k'}(x) \partial n_k(x)} = \psi^{(1)}(N(x) + 1) - \psi^{(1)}(n_k(x) + 1) \delta_{k'k}. \quad (\text{A4})$$

The functions $\psi(z) = d \log \Gamma(z) / dz$ and $\psi^{(1)}(z) = d^2 \log \Gamma(z) / dz^2$ have the asymptotic expansions

$$\psi(z) = \log z - \frac{1}{2z} - \sum_{l=1}^{\infty} \frac{B_{2l}}{2l z^{2l}} \quad , \quad \psi^{(1)}(z) = \frac{1}{z} + \frac{1}{2z^2} + \sum_{l=1}^{\infty} \frac{B_{2l}}{z^{2l+1}} \quad (\text{A5})$$

as $z \rightarrow \infty$. It will be assumed that each $n_k(x)$ has some characteristic value $\bar{n}_k(x)$ which is vastly larger than nearby values:

$$n_k(x) = \bar{n}_k(x) + \Delta n_k(x) \quad , \quad \bar{n}_k(x) \gg \gg |\Delta n_k(x)| \quad (\text{A6})$$

where “ $\gg \gg$ ” means “is vastly greater than”, as in $10^{1000} \gg \gg 1$. (This assumption is consistent with the fact that the initial action below has no lower bound as $n_k(x) \rightarrow \infty$ before an extra

stochastic term is added.) Then it is an extremely good approximation to use the asymptotic formulas above and write

$$S = S_0 + \sum_{x,k} a_k(x) \Delta n_k(x) - \sum_{x,k} a'_k(x) [\Delta n_k(x)]^2 + \sum_{x,k,k' \neq k} a'_{kk'}(x) \Delta n_k(x) \Delta n_{k'}(x) \quad (\text{A7})$$

$$a_k(x) = \log \bar{N}(x) - \log \bar{n}_k(x) \quad (\text{A8})$$

$$a'_k(x) = (2\bar{n}_k(x))^{-1} - (2\bar{N}(x))^{-1} \quad , \quad a'_{kk'}(x) = (2\bar{N}(x))^{-1} \quad (\text{A9})$$

where $\bar{N}(x)$ is the value of $N(x)$ when $n_k(x) = \bar{n}_k(x)$ for all k , and the higher-order terms have been separately neglected in $a_k(x)$ and $a'_k(x)$. (The above results then also follow immediately from Stirling's approximation for factorials.) For simplicity, we will also neglect the terms involving $(2\bar{N}(x))^{-1}$. (If these small terms are retained, the conclusions below still hold with some trivial redefinitions, but the notation and algebra become much more tedious.) Since there is initially no distinction between the fields labeled by k , it is consistent to assume that they all have the same $\bar{n}_k(x) = \bar{n}(x)$, and that $\bar{n}(x)$ is independent of x : $\bar{n}(x) = \bar{n}$ and $\bar{N}(x) = \bar{N}$, so that

$$a_k(x) = a = \log(\bar{N}/\bar{n}) \quad (\text{A10})$$

$$a'_k(x) = a' = (2\bar{n})^{-1} \quad . \quad (\text{A11})$$

The above assumptions are actually needed only to simplify the presentation, and they have no effect on the final results below with an appropriate limiting procedure as \bar{n}_k and $\bar{n}_M \rightarrow \infty$.

It is not conventional or convenient to deal with $\Delta n_k(x)$ and $[\Delta n_k(x)]^2$, so let us instead write S in terms of the fields ϕ_k and their derivatives $\partial\phi_k/\partial x^M$ via the following procedure: First, we can switch to a new set of points \bar{x} , defined to be the corners of the D -dimensional hypercubes centered on the original points x . It is easy to see that

$$S = S_0 + \sum_{\bar{x},k} a \langle \Delta n_k(x) \rangle - \sum_{\bar{x},k} a' \langle [\Delta n_k(x)]^2 \rangle \quad (\text{A12})$$

where $\langle \dots \rangle$ in the present context indicates an average over the 2^D boxes labeled by x which have the common corner \bar{x} . Second, at each point x we can write $\Delta n_k = \Delta\rho_k a_0^D = (\langle \Delta\rho_k \rangle + \delta\rho_k) a_0^D$, with $\langle \delta\rho_k \rangle = 0$:

$$S = S_0 + \sum_{\bar{x},k} a \langle (\langle \Delta\rho_k \rangle + \delta\rho_k) a_0^D \rangle - \sum_{\bar{x},k} a' \langle ((\langle \Delta\rho_k \rangle + \delta\rho_k)^2) (a_0^D)^2 \rangle \quad (\text{A13})$$

$$= S_0 + \sum_{\bar{x},k} a \langle \Delta\rho_k \rangle a_0^D - \sum_{\bar{x},k} a' [\langle (\Delta\rho_k)^2 \rangle + \langle (\delta\rho_k)^2 \rangle] (a_0^D)^2 \quad . \quad (\text{A14})$$

Each of the points x surrounding \bar{x} is displaced by $\delta x^M = \pm a_0/2$ along each of the x^M axes, so

$$\langle (\delta \rho_k)^2 \rangle = \langle (\delta \phi_k^2)^2 \rangle \quad (\text{A15})$$

$$= \left\langle \sum_M \left(\frac{\partial \phi_k^2}{\partial x^M} \delta x^M + \frac{1}{2} \frac{\partial^2 \phi_k^2}{\partial (x^M)^2} (\delta x^M)^2 \right)^2 \right\rangle \quad (\text{A16})$$

$$= \left\langle \sum_M \left(2\phi_k \frac{\partial \phi_k}{\partial x^M} \delta x^M + \left(\frac{\partial \phi_k}{\partial x^M} \right)^2 (\delta x^M)^2 + \phi_k \frac{\partial^2 \phi_k}{\partial (x^M)^2} (\delta x^M)^2 \right)^2 \right\rangle \quad (\text{A17})$$

to lowest order, where it is now assumed that at normal energies the fields are slowly varying over the extremely small distance a_0 . This assumption is justified by the prior assumption that \bar{n} is extremely large: $\phi_k^2(x) = \rho_k(x) = n_k(x)/a_0^D$ implies that $2\delta\phi_k/\phi_k \approx \delta n_k/n_k$ and $\phi_k = n_k^{1/2} a_0^{-D/2}$, so that $\delta\phi_k \sim \delta n_k n_k^{-1/2} a_0^{-D/2}$. The minimum change in ϕ_k is given by $\delta n_k = 1$:

$$\delta\phi_k^{\min} \sim n_k^{-1/2} a_0^{-D/2} \quad (\text{A18})$$

which means that $\delta\phi_k^{\min}$ is extremely small if n_k is extremely large.

In other words, the fields ϕ_k have effectively continuous values as $\bar{n} \rightarrow \infty$.

For extremely large \bar{n} it is an extremely good approximation to neglect the middle term in (A17), and to replace ϕ_k^2 by

$$\bar{\phi}^2 = \bar{\rho} = \bar{n}/a_0^D \quad (\text{A19})$$

giving

$$a' \langle (\delta \rho_k)^2 \rangle = \frac{1}{2a_0^D} \sum_M \left[\left(\frac{\partial \phi_k}{\partial x^M} \right)^2 a_0^2 + \left(\frac{\partial^2 \phi_k}{\partial (x^M)^2} \right)^2 \frac{a_0^4}{16} \right]. \quad (\text{A20})$$

It is similarly an extremely good approximation to neglect the term in (A14) involving $a' (a_0^D)^2 \langle \Delta \rho_k \rangle^2 = \langle \Delta n_k \rangle^2 / 2\bar{n}$ in comparison to that involving $\langle \Delta \rho_k \rangle a_0^D = \langle \Delta n_k \rangle$, so that

$$S = S_0 + \sum_{\bar{x}, k} a_0^D \frac{\mu_0}{m_0} \left(\phi_k^2 - \bar{\phi}^2 \right) - \sum_{\bar{x}, k} \sum_M a_0^D \frac{1}{2m_0^2} \left[\left(\frac{\partial \phi_k}{\partial x^M} \right)^2 + \frac{a_0^2}{16} \left(\frac{\partial^2 \phi_k}{\partial (x^M)^2} \right)^2 \right] \quad (\text{A21})$$

where

$$m_0 = a_0^{-1} \quad , \quad \mu_0 = m_0 \log(\bar{N}/\bar{n}) \quad (\text{A22})$$

The philosophy behind the above treatment is simple: We essentially wish to replace $\langle f^2 \rangle$ by $(\partial f / \partial x)^2$, and this can be accomplished because

$$\langle f^2 \rangle - \langle f \rangle^2 = \langle (\delta f)^2 \rangle \approx \langle (\partial f / \partial x)^2 (\delta x)^2 \rangle = (\partial f / \partial x)^2 (a_0/2)^2 \quad (\text{A23})$$

The form of (A21) also has a simple interpretation: The entropy S increases with the number of dits, but decreases when the dits are not uniformly distributed.

In the continuum limit,

$$\sum_{\bar{x}} a_0^D \rightarrow \int d^D x \quad (\text{A24})$$

(A21) becomes

$$S = S_0 + \int d^D x \sum_k \left\{ \frac{\mu_0}{m_0} (\phi_k^2 - \bar{\phi}^2) - \frac{1}{2m_0^2} \sum_M \left[\left(\frac{\partial \phi_k}{\partial x^M} \right)^2 + \frac{a_0^2}{16} \left(\frac{\partial^2 \phi_k}{\partial (x^M)^2} \right)^2 \right] \right\}. \quad (\text{A25})$$

A physical configuration of all the fields $\phi_k(x)$ corresponds to a specification of all the density variations $\Delta\rho_k(x)$. In the present picture, the probability of such a configuration is proportional to $W = e^S$. In a path integral with Euclidean form, the probability is proportional to $e^{-\bar{S}_b}$, where \bar{S}_b is the action for these bosonic fields. We conclude that

$$\bar{S}_b = -S + \text{constant} \quad (\text{A26})$$

and we will choose the constant to equal S_0 .

In the following it will be convenient to write the action in terms of $\tilde{\phi}_k = m_0^{-1/2} \phi_k$. For simplicity, we assume that the number of relevant $\tilde{\phi}_k$ is even, so that we can group these real fields in pairs to form N_f complex fields $\tilde{\Psi}_{b,k}$. It is also convenient to subtract out the enormous contribution of $\bar{\phi}$ by defining

$$\Psi_b = \tilde{\Psi}_b - \bar{\Psi}_b \quad (\text{A27})$$

where $\tilde{\Psi}_b$ is the vector with components $\tilde{\Psi}_{b,k}$ and $\bar{\Psi}_b$ is similarly defined with $\phi_k \rightarrow \bar{\phi}$. Then the action can be written

$$\bar{S}_b = \int d^D x \left\{ \frac{1}{2m_0} \left[\frac{\partial \Psi_b^\dagger}{\partial x^M} \frac{\partial \Psi_b}{\partial x^M} + \frac{a_0^2}{16} \frac{\partial^2 \Psi_b^\dagger}{\partial (x^M)^2} \frac{\partial^2 \Psi_b}{\partial (x^M)^2} \right] - \mu_0 (\tilde{\Psi}_b^\dagger \tilde{\Psi}_b - \bar{\Psi}_b^\dagger \bar{\Psi}_b) \right\} \quad (\text{A28})$$

since $\bar{\Psi}_b$ is constant, with summation now implied over repeated indices like M .

As described above, in Section II, we now add an extra imaginary term $i\tilde{V} \Psi_b^\dagger \Psi_b$ in the integral giving the action. Here \tilde{V} is a potential which has a Gaussian distribution, with

$$\langle \tilde{V} \rangle = 0 \quad , \quad \langle \tilde{V}(x) \tilde{V}(x') \rangle = b \delta(x - x') \quad (\text{A29})$$

where b is a constant, with

$$b \rightarrow 0+ \quad (\text{A30})$$

at the end of the calculations.

Then the complete action has the form

$$\begin{aligned} \tilde{S}_B [\Psi_b^\dagger, \Psi_b] = \int d^D x \left\{ \frac{1}{2m_0} \left[\frac{\partial \Psi_b^\dagger}{\partial x^M} \frac{\partial \Psi_b}{\partial x^M} + \frac{a_0^2}{16} \frac{\partial^2 \Psi_b^\dagger}{\partial (x^M)^2} \frac{\partial^2 \Psi_b}{\partial (x^M)^2} \right] \right. \\ \left. - \mu_0 \left(\tilde{\Psi}_b^\dagger \tilde{\Psi}_b - \overline{\Psi}_b^\dagger \overline{\Psi}_b \right) + i\tilde{V} \Psi_b^\dagger \Psi_b \right\}. \end{aligned} \quad (\text{A31})$$

In the following we will assume that the only fields which make an appreciable contribution in (A31) are those for which $\int d^D x \overline{\Psi}_b^\dagger \Psi_b = \overline{\Psi}_b^\dagger \int d^D x \Psi_b = 0$. This assumption is justified by the fact that $\overline{\Psi}_b$ is constant with respect to all the coordinates and, in the present picture, fields Ψ_b corresponding to physical gauge representations have nonzero angular momenta in the internal space of Appendices C and D. Then (A31) simplifies to

$$\tilde{S}_B [\Psi_b^\dagger, \Psi_b] = \int d^D x \left\{ \frac{1}{2m_0} \left[\frac{\partial \Psi_b^\dagger}{\partial x^M} \frac{\partial \Psi_b}{\partial x^M} + \frac{a_0^2}{16} \frac{\partial^2 \Psi_b^\dagger}{\partial (x^M)^2} \frac{\partial^2 \Psi_b}{\partial (x^M)^2} \right] - \mu_0 \Psi_b^\dagger \Psi_b + i\tilde{V} \Psi_b^\dagger \Psi_b \right\}. \quad (\text{A32})$$

Appendix B: Primordial supersymmetry

If F is any functional of the fundamental fields Ψ_b , its average value is given by

$$\langle F \rangle = \left\langle \frac{\int \mathcal{D} \Psi_b^\dagger \mathcal{D} \Psi_b F [\Psi_b^\dagger, \Psi_b] e^{-\tilde{S}_B [\Psi_b^\dagger, \Psi_b]}}{\int \mathcal{D} \underline{\Psi}_b^\dagger \mathcal{D} \underline{\Psi}_b e^{-\tilde{S}_B [\underline{\Psi}_b^\dagger, \underline{\Psi}_b]}} \right\rangle \quad (\text{B1})$$

where $\langle \dots \rangle$ now represents an average over the perturbing potential $i\tilde{V}$ and $\int \mathcal{D} \Psi_b^\dagger \mathcal{D} \Psi_b$ is to be interpreted as $\prod_{x,k} \int_{-\infty}^{\infty} d \text{Re} \Psi_{b,k}(x) \int_{-\infty}^{\infty} d \text{Im} \Psi_{b,k}(x)$. The transition from the original summation over $n_k(x)$ to this Euclidean path integral has the form (with $\Delta n = 1$ here)

$$\sum_{n=0}^{\infty} f(n) \Delta n \rightarrow \int_0^{\infty} f dn \rightarrow \int_0^{\infty} f d(a_0^D \phi^2) \rightarrow 2\bar{\phi} a_0^D \int_0^{\infty} f d\phi \rightarrow 2\bar{\phi} a_0^D m_0^{1/2} \int_{-\infty}^{\infty} f d\phi' \quad (\text{B2})$$

where $\phi' = \tilde{\phi} - m_0^{-1/2} \bar{\phi}$, since $d(\phi^2) \approx 2\bar{\phi} d\phi$ is an extremely good approximation for physically relevant fields, and since ϕ' effectively ranges from $-\infty$ to $+\infty$. Each ϕ' then becomes a $\text{Re} \Psi_{b,k}(x)$ or $\text{Im} \Psi_{b,k}(x)$, and the constant factors cancel in the numerator and denominator of (B1).

The presence of the denominator makes it difficult to perform the average of (B1), but there is a trick for removing the bosonic degrees of freedom $\underline{\Psi}_b$ in the denominator and

replacing them with fermionic degrees of freedom Ψ_f in the numerator [41–43]: After integration by parts (with boundary terms usually assumed either to vanish or to be irrelevant in this paper), (A32) can be written in the form $\tilde{S}_B = \int d^D x \Psi_b^\dagger A \Psi_b$. Then, since

$$\int \mathcal{D} \underline{\Psi}_b^\dagger \mathcal{D} \underline{\Psi}_b e^{-\tilde{S}_B[\underline{\Psi}_b^\dagger, \underline{\Psi}_b]} = C (\det \mathcal{A})^{-1} \quad (\text{B3})$$

$$\int \mathcal{D} \Psi_f^\dagger \mathcal{D} \Psi_f e^{-\tilde{S}_B[\Psi_f^\dagger, \Psi_f]} = \det \mathcal{A} \quad (\text{B4})$$

where the matrix \mathcal{A} corresponds to the operator A and C is a constant, it follows that

$$\langle F \rangle = \frac{1}{C} \left\langle \int \mathcal{D} \Psi_b^\dagger \mathcal{D} \Psi_b \mathcal{D} \Psi_f^\dagger \mathcal{D} \Psi_f F e^{-\tilde{S}_B[\Psi_b^\dagger, \Psi_b]} e^{-\tilde{S}_B[\Psi_f^\dagger, \Psi_f]} \right\rangle \quad (\text{B5})$$

$$= \frac{1}{C} \left\langle \int \mathcal{D} \Psi^\dagger \mathcal{D} \Psi F e^{-\tilde{S}_{bf}[\Psi^\dagger, \Psi]} \right\rangle \quad (\text{B6})$$

where Ψ_b and Ψ_f have been combined into

$$\Psi = \begin{pmatrix} \Psi_b \\ \Psi_f \end{pmatrix} \quad (\text{B7})$$

and

$$\tilde{S}_{bf}[\Psi^\dagger, \Psi] = \int d^D x \left\{ \frac{1}{2m_0} \left[\frac{\partial \Psi^\dagger}{\partial x^M} \frac{\partial \Psi}{\partial x^M} + \frac{a_0^2}{16} \frac{\partial^2 \Psi^\dagger}{\partial (x^M)^2} \frac{\partial^2 \Psi}{\partial (x^M)^2} \right] - \mu_0 \Psi^\dagger \Psi + i\tilde{V} \Psi^\dagger \Psi \right\}. \quad (\text{B8})$$

In (B7), Ψ_f consists of Grassmann variables $\Psi_{f,k}$, just as Ψ_b consists of ordinary variables $\Psi_{b,k}$, and $\int \mathcal{D} \Psi^\dagger \mathcal{D} \Psi$ is to be interpreted as

$$\prod_{x,k} \int_{-\infty}^{\infty} d\text{Re} \Psi_{b,k}(x) \int_{-\infty}^{\infty} d\text{Im} \Psi_{b,k} \int d\Psi_{f,k}^*(x) \int d\Psi_{f,k}(x). \quad (\text{B9})$$

Recall that Ψ_b and Ψ_f each have N_f components.

In this early stage of the theory, the bosonic fields in the denominator – which are of critical importance for proper normalization – have effectively been transformed into fermionic fields in the numerator, where they perform the same function. This is the origin of supersymmetry in the present picture.

For a Gaussian random variable v whose mean is zero, the result

$$\langle e^{-iv} \rangle = e^{-\frac{1}{2}\langle v^2 \rangle} \quad (\text{B10})$$

implies that

$$\left\langle e^{-\int d^D x i\tilde{V} \Psi^\dagger \Psi} \right\rangle = e^{-\frac{1}{2} \int d^D x d^D x' \Psi^\dagger(x) \Psi(x) \langle \tilde{V}(x) \tilde{V}(x') \rangle \Psi^\dagger(x') \Psi(x')} \quad (\text{B11})$$

$$= e^{-\frac{1}{2} b \int d^D x [\Psi^\dagger(x) \Psi(x)]^2}. \quad (\text{B12})$$

It follows that

$$\langle F \rangle = \frac{1}{C} \int \mathcal{D}\Psi^\dagger \mathcal{D}\Psi F e^{-S_E} \quad (\text{B13})$$

with

$$S_E = \int d^D x \left\{ \frac{1}{2m_0} \left[\frac{\partial\Psi^\dagger}{\partial x^M} \frac{\partial\Psi}{\partial x^M} + \frac{a_0^2}{16} \frac{\partial^2\Psi^\dagger}{\partial (x^M)^2} \frac{\partial^2\Psi}{\partial (x^M)^2} \right] - \mu_0 \Psi^\dagger \Psi + \frac{1}{2} b (\Psi^\dagger \Psi)^2 \right\}. \quad (\text{B14})$$

A special case (with $F = 1$) is

$$Z = \frac{1}{C} \int \mathcal{D}\Psi^\dagger \mathcal{D}\Psi e^{-S_E} \quad (\text{B15})$$

but according to (B1) $Z = 1$, so $C = \int \mathcal{D}\Psi^\dagger \mathcal{D}\Psi e^{-S_E}$ and

$$\langle F \rangle = \frac{\int \mathcal{D}\Psi^\dagger \mathcal{D}\Psi F e^{-S_E}}{\int \mathcal{D}\Psi^\dagger \mathcal{D}\Psi e^{-S_E}}. \quad (\text{B16})$$

Again, notice that the fermionic fields Ψ_f are effectively a transformed version of the bosonic fields $\underline{\Psi}_b$. The coupling between the fields Ψ_b and Ψ_f (or $\underline{\Psi}_b$) is due to the random perturbing potential $i\tilde{V}$. In the replacement of (B1) by (B16), F essentially serves as a test functional. The meaning of this replacement is that the action (B14), with both bosonic and fermionic fields, must be used instead of the original action (A32), with only bosonic fields, in treating all physical quantities and processes, if the average over random fluctuations in (B1) is to disappear from the theory.

Notice that the two steps above serve two independent purposes: The transformation of $\underline{\Psi}_b$ in the denominator to Ψ_f in the numerator provides a more convenient formulation because all fields now have equal status in the numerator, and can be treated in the same way. The introduction of an infinitesimal perturbing potential is then preparation for the formation of a condensate at finite energy (in the very early universe), as discussed in Section II. Of course, it is the conjunction of these two steps that makes the following developments possible.

Appendix C: The internal space

The internal space of Section V is $(D-4)$ -dimensional, with $SO(D-4)$ and $Spin(D-4)$ rotation groups and their vector, spinor, adjoint, etc. representations. For a 10-dimensional space (which may be a subset of the full internal space) these are the 10, 16, 45, $16 + \overline{16}$, 16×16 , etc. representations (in a simplified notation).

It may be helpful to begin with an analogy, in which external spacetime is replaced by the z -axis. The internal space is replaced by an xy -plane, with internal states described by 2-dimensional vector fields (rather than the higher-dimension vector and spinor fields considered below). One of these states is occupied by the condensate, and is represented by a vector \mathbf{v}_1 which points radially outward from the origin at all points in the xy -plane when $z = 0$. The other state is an additional basis function, represented by a vector \mathbf{v}_2 which is everywhere perpendicular to \mathbf{v}_1 . But \mathbf{v}_1 is allowed to rotate as a function of z , so it has both radial and tangential components after a displacement along the z -axis. Then \mathbf{v}_2 is forced to rotate with \mathbf{v}_1 – i.e., the condensate – in order to preserve orthogonality.

Now let us turn to the actual internal space, first considering a set of $(D-4)$ -dimensional vector fields $\tilde{\psi}_{vec}^r$. Let $\tilde{\psi}_{vec}^0$ represent the state occupied by a bosonic condensate. In the simplest picture, and at some fixed x_0^μ , only the r th component of the field $\tilde{\psi}_{vec}^r$ is nonzero along some radial direction in the internal space, making the fields trivially orthogonal in that direction. Then, with x^μ still fixed, $\tilde{\psi}_{vec}^r(x^m)$ in all other radial directions is obtained from the original $\tilde{\psi}_{vec}^r(x_0^m)$ by rotating it to x^m . In other words, the field at each point in the internal space is identical to the field that would be obtained at that point if the original field $\tilde{\psi}_{vec}^r(x_0^m)$ were subjected to a rotation about the origin. This produces an isotropic configuration for the condensate and each basis function. As in (4.14) we can write

$$\tilde{\psi}_{vec}^r(x^m) = U_{vec}(x^m, x_0^m) \tilde{\psi}_{vec}^r(x_0^m) . \quad (C1)$$

Just as in the analogy, a field that is radial at x_0^m will also be radial at all other points x^m . However, a general $\tilde{\psi}_{vec}^r(x_0^m)$ permits a general vortex-like configuration of the condensate.

Also as in the analogy, the state $\tilde{\psi}_{vec}^0$ of the condensate is allowed to rotate as a function of x^μ (because such a rotation does not alter the internal action). Since the other basis functions $\tilde{\psi}_{vec}^r$ are required to remain orthogonal to $\tilde{\psi}_{vec}^0$ and each other, they are required to rotate with the condensate. Then (C1) becomes more generally

$$\tilde{\psi}_{vec}^r(x^m, x^\mu) = U_{vec}(x^m, x_0^m; x^\mu, x_0^\mu) \tilde{\psi}_{vec}^r(x_0^m, x_0^\mu) \quad (C2)$$

with

$$\tilde{\psi}_{vec}^{r\dagger}(x^m, x^\mu) \tilde{\psi}_{vec}^{r'}(x^m, x^\mu) = \tilde{\psi}_{vec}^{r\dagger}(x_0^m, x_0^\mu) \tilde{\psi}_{vec}^{r'}(x_0^m, x_0^\mu) = \delta_{rr'} \quad (C3)$$

since

$$U_{vec}^\dagger(x^m, x_0^m; x^\mu, x_0^\mu) U_{vec}(x^m, x_0^m; x^\mu, x_0^\mu) = 1 . \quad (C4)$$

In general (with x^μ fixed), let $\tilde{\psi}(\mathbf{x})$ represent a multicomponent basis function with angular momentum j at a point \mathbf{x} in the $(D-4)$ -dimensional internal space. After a rotation about the origin specified by the $(D-4) \times (D-4)$ matrix \mathbf{R} , it is transformed to

$$\tilde{\psi}'(\mathbf{x}) = \mathcal{R}(\mathbf{R}) \tilde{\psi}(\mathbf{R}^{-1}\mathbf{x}) \quad (\text{C5})$$

where $\mathcal{R}(\mathbf{R})$ belongs to the appropriate representation of the group $SO(D-4)$ or $Spin(D-4)$. However, we require that the field be isotropic, so that it is left unchanged after a rotation:

$$\tilde{\psi}'(\mathbf{x}) = \tilde{\psi}(\mathbf{x}) . \quad (\text{C6})$$

Then we can define $\tilde{\psi}(\mathbf{x})$ at each value of the radial coordinate r by starting with a $\tilde{\psi}(\mathbf{x}_0)$ and requiring that

$$\tilde{\psi}(\mathbf{x}) = \mathcal{R}(\mathbf{R}) \tilde{\psi}(\mathbf{x}_0) \quad , \quad \mathbf{x} = \mathbf{R} \mathbf{x}_0 . \quad (\text{C7})$$

With this definition, each component $\tilde{\psi}_p(\mathbf{x})$ is a single-valued function of the coordinates only if j is an integer. If $j = 1/2$, e.g., $\tilde{\psi}_p(\mathbf{x})$ acquires a minus sign after a rotation of 2π , but it is single-valued on the $Spin(D-4)$ group manifold.

Multivalued functions are well-known in other similar contexts, such as the behavior of the phase of an ordinary superfluid order parameter $\psi_s = e^{i\theta_s} n_s^{1/2}$ around a vortex, which becomes discontinuous if it is required to be a single-valued function of the coordinates [86]. In the same way, $z^{1/2}$ exhibits a discontinuity across a branch cut if it is required to be a single-valued function and z is restricted to a single complex plane. I.e., $z^{1/2} = |z|^{1/2} e^{i\phi/2}$ gives $+|z|^{1/2}$ for $\phi = 0$ and $-|z|^{1/2}$ for $\phi = 2\pi$. But when defined on a pair of Riemann sheets, $z^{1/2}$ is a continuous function, and the same is true of $\tilde{\psi}(\mathbf{x})$ or each component $\tilde{\psi}_p(\mathbf{x})$ as we have defined it above, on the group manifold. The key idea in either case is to extend the manifold over which the function is defined, so that there are no artificial discontinuities. A similar principle holds in condensed matter physics, where a spinor can be a multivalued function of position.

The above discussion, however, is only relevant to mathematical consistency. Although the mathematical quantity $\tilde{\psi}_p(\mathbf{x})$ can be multivalued, the more fundamental and physically significant quantity $\tilde{\psi}_p^*(\mathbf{x}) \tilde{\psi}_p(\mathbf{x})$ is a single-valued function of position. The same is true regarding the $SU(2) \times U(1)$ order parameter and fields in external spacetime.

In the 10-dimensional internal space, spinorial fields $\tilde{\psi}_{sp}^r$ are appropriate basis functions for ordinary fermions and sfermions, and for the primordial bosonic condensate of Section IV. Again, let $\tilde{\psi}_{sp}^r(x_0^m)$ represent a field along some radial direction in the internal space at some fixed x_0^μ . Then the field configuration for every point x^m is obtained by taking $\tilde{\psi}_{sp}^r(x^m)$ to be identical to the field that would be obtained at that point if $\tilde{\psi}_{sp}^r(x_0^m)$ were subjected to a rotation, with

$$\tilde{\psi}_{sp}^r(x^m) = U_{sp}(x^m, x_0^m) \tilde{\psi}_{sp}^r(x_0^m) \quad (\text{C8})$$

as in (C7). Again, the state $\tilde{\psi}_{sp}^0$ of the condensate is allowed to rotate as a function of x^μ , and since the other basis functions $\tilde{\psi}_{sp}^r$ must remain orthogonal to $\tilde{\psi}_{sp}^0$ they are required to rotate with the condensate. The general version of (C8) is then

$$\tilde{\psi}_{sp}^r(x^m, x^\mu) = U_{sp}(x^m, x_0^m; x^\mu, x_0^\mu) \tilde{\psi}_{sp}^r(x_0^m, x_0^\mu) . \quad (\text{C9})$$

The same reasoning applies to each representation, and thus to the combined set of all physically relevant basis functions (or internal fields) $\tilde{\psi}_{int}^r(x^m, x^\mu)$:

$$\tilde{\psi}_{int}^r(x'^m, x'^\mu) = U_{int}(x'^m, x^m; x'^\mu, x^\mu) \tilde{\psi}_{int}^r(x^m, x^\mu) \quad (\text{C10})$$

with

$$\tilde{\psi}_{int}^{r\dagger}(x'^m, x'^\mu) \tilde{\psi}_{int}^{r'}(x'^m, x'^\mu) = \tilde{\psi}_{int}^{r\dagger}(x^m, x^\mu) \tilde{\psi}_{int}^{r'}(x^m, x^\mu) = \delta_{rr'} . \quad (\text{C11})$$

So that the internal action will be unaffected as $x^\mu \rightarrow x'^\mu$, we require that the order parameter experience a uniform rotation, described by a matrix $\overline{\mathcal{R}}_{int}$ which is independent of x^m . Then U_{int} has the form

$$U_{int}(x'^m, x^m; x'^\mu, x^\mu) = \overline{\mathcal{R}}_{int}(x'^\mu, x^\mu) \mathcal{R}_{int}(x'^m, x^m) . \quad (\text{C12})$$

(Notice that (C12) is to be distinguished from a rotation about the origin, which is given by (C5), and which according to (C6) would leave $\tilde{\psi}(x^m)$ unchanged rather than rotated at each point x^m .) It follows that

$$\tilde{\psi}_{int}^r(x^m, x^\mu) = \overline{\mathcal{R}}_{int}(x^\mu, x_0^\mu) \tilde{\psi}_{int}^r(x^m, x_0^\mu) . \quad (\text{C13})$$

At each fixed x^m , the order parameter has been rotated as $x_0^\mu \rightarrow x^\mu$. We define the parameters $\delta\overline{\phi}_i$ by

$$\overline{\mathcal{R}}_{int}(x^\mu + \delta x^\mu, x_0^\mu) = \overline{\mathcal{R}}_{int}(x^\mu, x_0^\mu) (1 - i \delta\overline{\phi}_i J_i) \quad (\text{C14})$$

or

$$\delta\tilde{\psi}_{int}^r(x^m) = -i\delta\bar{\phi}_i\bar{J}_i\tilde{\psi}_{int}^r(x^m) \quad \text{as } x^\mu \rightarrow x^\mu + \delta x^\mu \quad (\text{C15})$$

$$\bar{J}_i = \bar{\mathcal{R}}_{int}(x^\mu, x_0^\mu) J_i \bar{\mathcal{R}}_{int}^{-1}(x^\mu, x_0^\mu) \quad (\text{C16})$$

where the matrices J_i are the generators in the reducible representation of $Spin(D-4)$ corresponding to $\tilde{\psi}_{int}^r$. The matrix elements of \bar{J}_i are independent of x^μ :

$$\int d^{D-4}x \tilde{\psi}_{int}^{r\dagger}(x^m, x^\mu) \bar{J}_i \tilde{\psi}_{int}^{r'}(x^m, x^\mu) = \int d^{D-4}x \tilde{\psi}_{int}^{r\dagger}(x^m, x_0^\mu) J_i \tilde{\psi}_{int}^{r'}(x^m, x_0^\mu). \quad (\text{C17})$$

The primordial condensate is in a specific representation, but the basis functions in other representations are chosen to rotate with it according to (C13) and (C15).

It may be helpful to illustrate the above ideas by returning to the 2-dimensional analogy. Equation (C7) becomes

$$\mathbf{v}(\mathbf{x}) = \mathcal{R}_{vec} \mathbf{v}(\mathbf{x}_0), \quad \mathcal{R}_{vec} = \begin{pmatrix} \cos\phi & -\sin\phi \\ \sin\phi & \cos\phi \end{pmatrix}, \quad \mathbf{v}(\mathbf{x}_0) = \begin{pmatrix} R(r) \\ 0 \end{pmatrix} \text{ or } \begin{pmatrix} 0 \\ R(r) \end{pmatrix} \quad (\text{C18})$$

for the vector representation and

$$s(\mathbf{x}) = \mathcal{R}_{sp} s(\mathbf{x}_0), \quad \mathcal{R}_{sp} = e^{-i\sigma_3\phi/2}, \quad s(\mathbf{x}_0) = \begin{pmatrix} R(r) \\ 0 \end{pmatrix} \text{ or } \begin{pmatrix} 0 \\ R(r) \end{pmatrix} \quad (\text{C19})$$

for the spinor representation. The matrices corresponding to the J_i are

$$J_{vec} = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \text{and} \quad J_{sp} = \frac{\sigma_3}{2} = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (\text{C20})$$

Notice that ϕ_i is an angular coordinate in the internal space, whereas $\bar{\phi}_i$ is a parameter specifying the rotation of $\tilde{\psi}_{int}^r$ at fixed x^m as x^μ is varied.

Appendix D: Solutions in the internal space

Our goal in this appendix is merely to show that there are solutions with the form required in Appendix C, so we will look first for solutions with the higher-derivative terms in (5.2) and (5.3) neglected, and with Ψ_{int} sufficiently small that $V_0(x^m)$ can also be neglected. Then (5.2) and (5.3) become

$$\left(-\frac{1}{2m_0}\partial_m\partial_m - \mu_{int}\right)\Psi_{int}(x^m, x^\mu) = 0 \quad (\text{D1})$$

$$\left(-\frac{1}{2m_0}\partial_m\partial_m - \mu_{int}\right)\tilde{\psi}_{int}^r(x^m, x^\mu) = 0. \quad (\text{D2})$$

For simplicity of notation, let $\tilde{\psi}_{int}^r(x^m, x^\mu)$ again be represented by $\tilde{\psi}(\mathbf{x})$, with components $\tilde{\psi}_p(\mathbf{x})$. Each component varies with position in the way specified by (C7) (together with the radial dependence of $\tilde{\psi}(\mathbf{x}_0)$). It therefore has an orbital angular momentum given by the usual orbital angular momentum operators \hat{J}_i in \bar{d} dimensions [87–92], which essentially measure how rapidly $\tilde{\psi}_p(\mathbf{x})$ varies as a function of the angles ϕ_i .

The Laplacian $\partial_m \partial_m$ can be rewritten in terms of radial derivatives and the usual \hat{J}^2 , giving [87–89]

$$\left(-\frac{1}{r^{2K}} \frac{\partial}{\partial r} \left(r^{2K} \frac{\partial}{\partial r} \right) + \frac{\hat{J}^2}{r^2} - 1 \right) \tilde{\psi}_p(\mathbf{x}) = 0 \quad , \quad K = \frac{\bar{d} - 1}{2} \quad (\text{D3})$$

after rescaling of the radial coordinate r , where

$$\bar{d} = D - 4 . \quad (\text{D4})$$

In addition, it is shown in Narumi and Nakau [87], Gallup [88], and Louck [89] that

$$\hat{J}^2 \tilde{\psi}_p(\mathbf{x}) = j(j + \bar{d} - 2) \tilde{\psi}_p(\mathbf{x}) \quad (\text{D5})$$

where j is the orbital angular momentum quantum number, as defined on p. 677 of Gallup, but with this definition extended to half-integer values of m_α and j . Normally, of course, only integer values of these orbital quantum numbers are permitted. However, the functions $\tilde{\psi}_p(\mathbf{x})$ as defined in C can have $j = 1/2$ etc. (in which case they are multivalued functions of the coordinates but single-valued functions on the group manifold, as discussed below (C7)). Also, the demonstration of (D5) of Gallup [88] can be extended in the present context to half-integer j , because it employs raising and lowering operators. (At each \mathbf{x} , $\tilde{\psi}_p$ is a linear combination of states with different values of m_α , but (D5) still holds.) For each $\tilde{\psi}_p(\mathbf{x})$ the radial wavefunction then satisfies

$$\left[-\frac{1}{r^{2K}} \frac{d}{dr} \left(r^{2K} \frac{d}{dr} \right) + \frac{j(j + \bar{d} - 2)}{r^2} - 1 \right] R(r) = 0 . \quad (\text{D6})$$

It may be helpful once again to consider the 2-dimensional analogy of C, where the orbital angular momentum operator is

$$\hat{J} = -i\partial/\partial\phi . \quad (\text{D7})$$

For the vector representation, (C18) implies

$$\partial_m \partial_m \mathbf{v}(\mathbf{x}) = \left[\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2}{\partial \phi^2} \right] \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix} \mathbf{v}(\mathbf{x}_0) \quad (\text{D8})$$

$$= \left[\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial}{\partial r} \right) - \frac{1}{r^2} \right] \mathbf{v}(\mathbf{x}) \quad (\text{D9})$$

in agreement with (D6) for $j = 1$. For the spinor representation, (C19) gives

$$\partial_m \partial_m s(\mathbf{x}) = \left[\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2}{\partial \phi^2} \right] e^{-i\sigma_3 \phi/2} s(\mathbf{x}_0) \quad (\text{D10})$$

$$= \left[\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial}{\partial r} \right) - \frac{1/4}{r^2} \right] s(\mathbf{x}) \quad (\text{D11})$$

in agreement with (D6) for $j = 1/2$.

Equation (D6) can be further reduced to [90, 92]

$$\left[-\frac{d^2}{dr^2} + \frac{k(k-1)}{r^2} - 1 \right] \chi(r) = 0 \quad , \quad k = j + K = j + \frac{\bar{d}-1}{2} \quad (\text{D12})$$

where $\chi(r) \equiv r^K R(r)$. It is then easy to show that

$$\chi(r) \propto r^k \quad \text{as } r \rightarrow 0 \quad , \quad \chi(r) \propto \sin(r + \delta) \quad \text{as } r \rightarrow \infty \quad (\text{D13})$$

where δ is a phase.

The higher derivatives in the full internal wave equation (5.3) permit exponentially decaying solutions which are then normalizable and have finite action. (For completeness, it may be worth noting that the resulting differential equation would be even higher than fourth-order if higher than second-order terms were kept in the expansion of (A16).) Suppose that the above equation at large r is modified to

$$\left[\alpha^2 \frac{d^4}{dr^4} - \frac{d^2}{dr^2} - 1 \right] \chi(r) = 0. \quad (\text{D14})$$

The solutions are

$$\chi(r) \propto e^{iqr} \quad , \quad q^2 = -\frac{1}{2\alpha^2} \pm \frac{\sqrt{1+4\alpha^2}}{2\alpha^2}. \quad (\text{D15})$$

There is then an exponentially decaying solution with the form $q = i/\bar{\alpha}$ and

$$\chi(r) \propto e^{-r/\bar{\alpha}} \quad (\text{D16})$$

so both the order parameter and the basis functions fall to zero as $r \rightarrow \infty$.

In general, differential equations with higher derivatives will asymptotically have exponentially decaying, exponentially growing, and oscillating solutions. This suggests that the following picture defines our universe, or any with both a very large external spacetime and a very small internal space: The solution for the internal condensate decays exponentially, while the external condensate is born in a nondecaying solution. Here the internal space is confined on an extremely small length scale of roughly a_0 .

Appendix E: Euclidean and Lorentzian Propagators

For Weyl fermions, the Euclidean 2-point function is

$$\begin{aligned} G_f(x_1, x_2) &= \langle \psi_f(x_1) \psi_f^\dagger(x_2) \rangle = \frac{\int \mathcal{D}\psi_f^\dagger \mathcal{D}\psi_f \psi_f(x_1) \psi_f^\dagger(x_2) e^{-S_f}}{\int \mathcal{D}\psi_f^\dagger \mathcal{D}\psi_f e^{-S_f}} \\ &= \frac{\prod_s \int d\bar{\psi}_f^*(s) \int d\bar{\psi}_f(s) e^{-\bar{\psi}_f^*(s)a(s)\bar{\psi}_f(s)} \sum_{s_1, s_2} \bar{\psi}_f^*(s_1) \bar{\psi}_f^*(s_2) U(x_1, s_1) U^\dagger(x_2, s_2)}{\prod_s \int d\bar{\psi}_f^*(s) \int d\bar{\psi}_f(s) e^{-\bar{\psi}_f^*(s)a(s)\bar{\psi}_f(s)}} \end{aligned} \quad (\text{E1})$$

where (6.1) and (6.2) have been used. In a term with $s_2 \neq s_1$, the numerator contains the factor

$$\int d\bar{\psi}_f^*(s_1) \int d\bar{\psi}_f(s_1) e^{-\bar{\psi}_f^*(s_1)a(s_1)\bar{\psi}_f(s_1)} \bar{\psi}_f^*(s_1) \bar{\psi}_f(s_1) = 0 \quad (\text{E2})$$

according to the rules for Berezin integration. But a term with $s_2 = s_1$ contributes

$$\begin{aligned} &\frac{\int d\bar{\psi}_f^*(s_1) \int d\bar{\psi}_f(s_1) e^{-\bar{\psi}_f^*(s_1)a(s_1)\bar{\psi}_f(s_1)} \bar{\psi}_f^*(s_1) \bar{\psi}_f^*(s_1)}{\int d\bar{\psi}_f^*(s_1) \int d\bar{\psi}_f(s_1) e^{-\bar{\psi}_f^*(s_1)a(s_1)\bar{\psi}_f(s_1)}} U(x_1, s_1) U^\dagger(x_2, s_1) \\ &= a(s_1)^{-1} U(x_1, s_1) U^\dagger(x_2, s_1) \end{aligned} \quad (\text{E3})$$

so

$$G_f(x_1, x_2) = \sum_s \bar{G}_f(s) U(x_1, s) U^\dagger(x_2, s) \quad , \quad \bar{G}_f(s) = a(s)^{-1} . \quad (\text{E4})$$

If the $U(x, s)$ used to represent $\psi_f(x)$ are a complete set, the propagator $G_f(x, x')$ is a true Green's function:

$$L_f(x) U(x, s) = a(s) U(x, s) \quad , \quad \psi_f(x) = \sum_s U(x, s) \bar{\psi}_f(s) \quad (\text{E5})$$

and $\sum_s U(x, s) U^\dagger(x', s) = \delta(x - x')$ imply that

$$L_f(x) G_f(x, x') = \delta(x - x') \quad (\text{E6})$$

as usual.

The treatment for scalar bosons is similar:

$$G_b(x_1, x_2) = \langle \phi_b(x_1) \phi_b^\dagger(x_2) \rangle = \frac{\int \mathcal{D}\phi_b^\dagger \mathcal{D}\phi_b \phi_b(x_1) \phi_b^\dagger(x_2) e^{-S_f}}{\int \mathcal{D}\phi_b^\dagger \mathcal{D}\phi_b e^{-S_f}} \quad (\text{E7})$$

$$= \frac{\prod_s \int_{-\infty}^{\infty} d\text{Re} \bar{\phi}_b(s) \int_{-\infty}^{\infty} d\text{Im} \bar{\phi}_b(s) e^{-\tilde{a}(s)[(\text{Re} \bar{\phi}_b(s))^2 + (\text{Im} \bar{\phi}_b(s))^2]} \sum_{s_1, s_2} \bar{\phi}_b(s_1) \bar{\phi}_b^*(s_2)}{\prod_s \int_{-\infty}^{\infty} d\text{Re} \bar{\phi}_b(s) \int_{-\infty}^{\infty} d\text{Im} \bar{\phi}_b(s) e^{-\tilde{a}(s)[(\text{Re} \bar{\phi}_b(s))^2 + (\text{Im} \bar{\phi}_b(s))^2]} \times U_b(x_1, s_1) U_b^\dagger(x_2, s_2)} \quad (\text{E8})$$

where

$$L_b(x) U_b(x, s) = \tilde{a}(s) U_b(x, s) \quad , \quad \phi_b(x) = \sum_s U_b(x, s) \bar{\phi}_b(s) . \quad (\text{E9})$$

In a term with $s_2 \neq s_1$, the numerator contains the factor

$$\int_{-\infty}^{\infty} d\text{Re} \bar{\phi}_b(s_1) \int_{-\infty}^{\infty} d\text{Im} \bar{\phi}_b(s_1) e^{-\tilde{a}(s_1)[(\text{Re} \bar{\phi}_b(s_1))^2 + (\text{Im} \bar{\phi}_b(s_1))^2]} [\text{Re} \bar{\phi}_b(s_1) + i \text{Im} \bar{\phi}_b(s_1)] = 0 \quad (\text{E10})$$

since the integrand is odd. But a term with $s_2 = s_1$ contains the factor

$$\frac{\int_{-\infty}^{\infty} d\text{Re} \bar{\phi}_b(s_1) e^{-\tilde{a}(s_1)(\text{Re} \bar{\phi}_b(s_1))^2} (\text{Re} \bar{\phi}_b(s_1))^2}{\int_{-\infty}^{\infty} d\text{Re} \bar{\phi}_b(s_1) e^{-\tilde{a}(s_1)(\text{Re} \bar{\phi}_b(s_1))^2}} + \frac{\int_{-\infty}^{\infty} d\text{Im} \bar{\phi}_b(s_1) e^{-\tilde{a}(s_1)(\text{Im} \bar{\phi}_b(s_1))^2} (\text{Im} \bar{\phi}_b(s_1))^2}{\int_{-\infty}^{\infty} d\text{Im} \bar{\phi}_b(s_1) e^{-\tilde{a}(s_1)(\text{Im} \bar{\phi}_b(s_1))^2}} = \tilde{a}(s_1)^{-1} \quad (\text{E11})$$

so

$$G_b(x_1, x_2) = \sum_s \bar{G}_b(s) U_b(x_1, s) U_b^\dagger(x_2, s) \quad , \quad \bar{G}_b(s) = \tilde{a}(s)^{-1} . \quad (\text{E12})$$

As usual, $a(s)$ and $\tilde{a}(s)$ contain a $+i\epsilon$ which is associated with a convergence factor in the path integral (and which gives a well-defined inverse).

The above are the propagators in the Euclidean formulation. The Lorentzian propagators are obtained through the same procedure with $a(s) \rightarrow -ia(s)$ and $\tilde{a}(s) \rightarrow -i\tilde{a}(s)$:

$$\bar{G}_f^L(s) = ia(s)^{-1} \quad , \quad \bar{G}_b^L(s) = i\tilde{a}(s)^{-1} . \quad (\text{E13})$$

The propagators in the Euclidean and Lorentzian formulations thus differ by only a factor of i . More generally, in the present picture, the action, fields, operators, classical equations of motion, quantum transition probabilities, propagation of particles, and meaning of time are the same in both formulations.

For a single noninteracting bosonic field with a mass m_b , the basis functions are

$$U_b(x, p) = \mathcal{V}^{-1/2} e^{ip \cdot x} = \mathcal{V}^{-1/2} e^{-i\omega t} e^{i\vec{p} \cdot \vec{x}} \quad (\text{E14})$$

so with $s \rightarrow p$ we have $\tilde{a}(p) = \omega^2 - |\vec{p}|^2 - m_b^2 + i\epsilon$ and

$$\overline{G}_b(p) = \frac{1}{\omega^2 - |\vec{p}|^2 - m_b^2 + i\epsilon} \quad , \quad \overline{G}_b^L(p) = \frac{i}{\omega^2 - |\vec{p}|^2 - m_b^2 + i\epsilon} . \quad (\text{E15})$$

Appendix F: Transformation to intermediate bosonic fields

In this appendix, still working in a locally inertial frame, we perform the most complicated and difficult of the steps required in the transformation from initial unphysical bosonic fields ψ_b^r to physical scalar boson fields ϕ_R , φ_S , and F_R .

First, if the gauge potentials A_μ^i were zero, (6.3) would become

$$i\sigma^\mu \partial_\mu U^0(x, s) = a_0(s) U^0(x, s) . \quad (\text{F1})$$

Then

$$U^0(x, s) = \mathcal{V}^{-1/2} u(s) e^{ip_s \cdot x} \quad , \quad p_s \cdot x = \eta_{\mu\nu} p_s^\mu x^\nu \quad , \quad \eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1) \quad (\text{F2})$$

(with \mathcal{V} a four-dimensional normalization volume) gives

$$-\eta_{\mu\nu} \sigma^\mu p_s^\nu U^0(x, s) = a_0(s) U^0(x, s) \quad (\text{F3})$$

where σ^μ implicitly multiplies the identity matrix for the multicomponent function $U^0(x, s)$.

A given 2-component spinor $u_r(s)$ has two eigenstates of $p_s^k \sigma^k$:

$$p_s^k \sigma^k u_r^+(s) = |\vec{p}_s| u_r^+(s) \quad , \quad p_s^k \sigma^k u_r^-(s) = -|\vec{p}_s| u_r^-(s) \quad (\text{F4})$$

where \vec{p}_s is the 3-momentum, with magnitude $|\vec{p}_s|$. The multicomponent eigenstates of $i\sigma^\mu \partial_\mu$ and their eigenvalues $a_0(s) = p_s^0 \mp |\vec{p}_s|$ thus come in pairs, corresponding to opposite helicities.

For nonzero A_μ^i , the eigenvalues $a(s)$ will also come in pairs, with one growing out of $a_0(s)$ and the other out of its partner $a_0(s')$ as the A_μ^i are turned on. To see this, first write (6.3) as

$$(i\partial_0 + A_0^i t_i) U(x, s) + \sigma^k (i\partial_k + A_k^i t_i) U(x, s) = a(s) U(x, s) \quad (\text{F5})$$

or

$$(i\partial_0\delta_{rr'} + A_0^i t_i^{rr'}) U_{r'}(x, s) - P_{rr'} U_{r'}(x, s) - a(s) \delta_{rr'} U_{r'}(x, s) = 0 \quad (\text{F6})$$

$$P_{rr'} \equiv -\sigma^k (i\partial_k \delta_{rr'} + A_k^i t_i^{rr'}) \quad (\text{F7})$$

with the usual implied summations over repeated indices. At fixed r, r' (and x, s), apply a matrix s which will diagonalize the 2×2 matrix $P_{rr'}$, bringing it into the form $p_{rr'}\sigma^3 + \bar{p}_{rr'}\sigma^0$, where $p_{rr'}$ and $\bar{p}_{rr'}$ are 1-component operators, while at the same time rotating the 2-component spinor $U_{r'}$:

$$sP_{rr'}s^{-1} = P'_{rr'} = p_{rr'}\sigma^3 + \bar{p}_{rr'}\sigma^0, \quad U'_{r'} = sU_{r'} \quad (\text{F8})$$

$$\sigma^0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (\text{F9})$$

But $P_{rr'}$ is traceless, and the trace is invariant under a similarity transformation, so $\bar{p}_{rr'} = 0$. Then the second term in (F6) (for fixed r and r') becomes $s^{-1}p_{rr'}\sigma^3 U'_{r'}(x, s)$. The two independent choices

$$U'_{r'}(x, s) \propto \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \sigma^3 U'_{r'}(x, s) = +U'_{r'}(x, s) \quad (\text{F10})$$

$$U'_{r'}(x, s) \propto \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad \sigma^3 U'_{r'}(x, s) = -U'_{r'}(x, s) \quad (\text{F11})$$

give $\pm s^{-1}p_{rr'}U'_{r'}(x, s)$. Now use $s^{-1}U'_{r'} = U_{r'}$ to obtain for (F6)

$$(i\partial_0\delta_{rr'} + A_0^i t_i^{rr'}) U_{r'}(x, s) \mp p_{rr'} U_{r'}(x, s) - a(s) \delta_{rr'} U_{r'}(x, s) = 0 \quad (\text{F12})$$

so (F5) reduces to two sets of equations with different eigenvalues $a(s)$ and $a(s')$:

$$a(s) = a_1(s) + a_2(s), \quad a(s') = a_1(s) - a_2(s) \quad (\text{F13})$$

where these equations define $a_1(s)$ and $a_2(s)$. Notice that letting $\sigma^k \rightarrow -\sigma^k$ in (F5) reverses the signs in (F12), and results in $a(s) \rightarrow a(s')$:

$$(i\partial_0 + A_0^i t_i) U(x, s) - \sigma^k (i\partial_k + A_k^i t_i) U(x, s) = a(s') U(x, s). \quad (\text{F14})$$

The action for a single eigenvalue $a(s)$ and its partner $a(s')$ is

$$\tilde{\mathcal{S}}_b(s) = \bar{\psi}_b^*(s) a(s) \bar{\psi}_b(s) + \bar{\psi}_b^*(s') a(s') \bar{\psi}_b(s') \quad (\text{F15})$$

$$= \bar{\psi}_b^*(s) (a_1(s) + a_2(s)) \bar{\psi}_b(s) + \bar{\psi}_b^*(s') (a_1(s) - a_2(s)) \bar{\psi}_b(s'). \quad (\text{F16})$$

In the following we will circumvent singularities by implicitly following the standard prescription $a_1 \rightarrow a_1 + i\epsilon$, $\epsilon \rightarrow 0+$, which reduces to $\omega \rightarrow \omega + i\epsilon$ when there are no gauge fields.

For $a_1(s) \geq 0$, let us choose $a_2(s) \geq 0$ and define

$$\bar{\psi}_b(s') = a(s)^{1/2} \bar{\phi}_b(s') = (a_1(s) + a_2(s))^{1/2} \bar{\phi}_b(s') \quad (\text{F17})$$

$$\bar{\psi}_b(s) = a(s)^{-1/2} \bar{F}_b(s) = (a_1(s) + a_2(s))^{-1/2} \bar{F}_b(s) \quad (\text{F18})$$

so that

$$\tilde{s}_b(s) = \bar{\phi}_b^*(s') \tilde{a}(s) \bar{\phi}_b(s') + \bar{F}_b^*(s) \bar{F}_b(s) \quad , \quad a_1(s) \geq 0 \quad (\text{F19})$$

where

$$\tilde{a}(s) = a(s) a(s') = a_1(s)^2 - a_2(s)^2 \quad . \quad (\text{F20})$$

For $a_1(s) < 0$, let us choose $a_2(s) \leq 0$ and write

$$\bar{\psi}_b(s') = (-a(s))^{1/2} \bar{\phi}_b(s') = (-a_1(s) - a_2(s))^{1/2} \bar{\phi}_b(s') \quad (\text{F21})$$

$$\bar{\psi}_b(s) = (-a(s))^{-1/2} \bar{F}_b(s) = (-a_1(s) - a_2(s))^{-1/2} \bar{F}_b(s) \quad (\text{F22})$$

so that

$$\tilde{s}_b(s) = - \left[\bar{\phi}_b^*(s') \tilde{a}(s) \bar{\phi}_b(s') + \bar{F}_b^*(s) \bar{F}_b(s) \right] \quad , \quad a_1(s) < 0 \quad . \quad (\text{F23})$$

Then we have

$$S_b = \sum'_s \tilde{s}_b(s) \quad (\text{F24})$$

$$= \sum'_{a_1(s) \geq 0} \left[\bar{\phi}_b^*(s') \tilde{a}(s) \bar{\phi}_b(s') + \bar{F}_b^*(s) \bar{F}_b(s) \right] - \sum'_{a_1(s) < 0} \left[\bar{\phi}_b^*(s') \tilde{a}(s) \bar{\phi}_b(s') + \bar{F}_b^*(s) \bar{F}_b(s) \right] \quad (\text{F25})$$

where a prime on a summation or product over s means that only one member of an s, s' pair (as defined in (F12) and (F13)) is included. Let us separate the non-negative contribution S_+ from the anomalous negative contribution S_- :

$$S_b = S_+ + S_- \quad (\text{F26})$$

$$S_+ = \sum'_{s \geq 0} \bar{\phi}_b^*(s') |\tilde{a}(s)| \bar{\phi}_b(s') + \sum'_{a_1(s) \geq 0} \bar{F}_b^*(s) \bar{F}_b(s) \quad (\text{F27})$$

$$S_- = - \left[\sum'_{s < 0} \bar{\phi}_b^*(s') |\tilde{a}(s)| \bar{\phi}_b(s') + \sum'_{a_1(s) < 0} \bar{F}_b^*(s) \bar{F}_b(s) \right] \quad (\text{F28})$$

where

$$s < 0 \iff \tilde{a}(s) = a_1(s)^2 - a_2(s)^2 < 0 \text{ if } a_1(s) \geq 0 \quad (\text{F29})$$

$$\iff \tilde{a}(s) = a_1(s)^2 - a_2(s)^2 > 0 \text{ if } a_1(s) < 0 \quad (\text{F30})$$

with $s \geq 0$ otherwise.

Recall that if the gauge potentials A_μ^i were zero, we would have $a_1 = \omega$ and $a_2 = \mp |\vec{p}|$, where ω is the frequency and \vec{p} the 3-momentum.

The negative-action modes of (F28) are discussed in Appendix G, where it is found that their *excitations* can be treated in the same way as the positive-action modes. (Here and below, “positive-action modes” means those which do not have negative action in the present context, before mass and interaction terms are acquired. After the transformation to a Lorentzian path integral below, some fields may come to have negative action due to symmetry breaking and condensation, but this is permissible within a Lorentzian description.) It follows that, in the treatment below, no degrees of freedom are lost in the bosonic *excitations*. On-shell modes already have non-negative action, so both the excitation and condensation of scalar boson fields remain exactly the same as in standard physics.

The path integral for positive-action modes in S_+ is

$$Z_+ = \int \mathcal{D}\psi_b^\dagger(x) \mathcal{D}\psi_b(x) e^{-S_+} \quad (\text{F31})$$

$$\begin{aligned} &= \prod_{x,ra} \int_{-\infty}^{\infty} d(\text{Re } \psi_{b,ra}(x)) \int_{-\infty}^{\infty} d(\text{Im } \psi_{b,ra}(x)) e^{-S_+} \\ &= \prod_s \int_{-\infty}^{\infty} d(\text{Re } \bar{\psi}_b(s)) \int_{-\infty}^{\infty} d(\text{Im } \bar{\psi}_b(s)) e^{-S_+} . \end{aligned} \quad (\text{F32})$$

Each of the transformations above from $\bar{\psi}_b$ to $\bar{\phi}_b$ and \bar{F}_b has the form

$$\bar{\psi}_b(s') = A(s)^{1/2} \bar{\phi}_b(s') \quad , \quad \bar{\psi}_b(s) = A(s)^{-1/2} \bar{F}_b(s) \quad (\text{F33})$$

so that $d\bar{\psi}_b(s') = A(s)^{1/2} d\bar{\phi}_b(s')$, $d\bar{\psi}_b(s) = A(s)^{-1/2} d\bar{F}_b(s)$, and the Jacobian is $\prod'_s A(s)^{1/2} A(s)^{-1/2} = 1$. These transformations then lead to

$$Z_{sb} = \prod'_{s \geq 0} z_\phi(s) \cdot \prod'_{s < 0} z_\phi(s) \cdot \prod'_{a_1(s) \geq 0} z_F(s) = \prod'_s z_\phi(s) \cdot \prod'_{a_1(s) \geq 0} z_F(s) \quad (\text{F34})$$

where the excitations of negative-action modes have been added and

$$z_\phi(s) = \int_{-\infty}^{\infty} d(\text{Re } \bar{\phi}_b(s')) \int_{-\infty}^{\infty} d(\text{Im } \bar{\phi}_b(s')) e^{-|\tilde{a}(s)|[(\text{Re } \bar{\phi}_b(s'))^2 + (\text{Im } \bar{\phi}_b(s'))^2]} \quad (\text{F35})$$

$$= \frac{\pi}{|\tilde{a}(s)|} \quad (\text{F36})$$

$$z_F(s) = \int_{-\infty}^{\infty} d(\text{Re } \bar{F}_b(s)) \int_{-\infty}^{\infty} d(\text{Im } \bar{F}_b(s)) e^{-[(\text{Re } \bar{F}_b(s))^2 + (\text{Im } \bar{F}_b(s))^2]} \quad (\text{F37})$$

$$= \pi. \quad (\text{F38})$$

Now let

$$S_{sb} = \sum'_s \bar{\phi}_b^*(s') \tilde{a}(s) \bar{\phi}_b(s') + \sum'_{a_1(s) > 0} \bar{F}_b^*(s) \bar{F}_b(s) \quad (\text{F39})$$

$$Z_{sb}^L = \int \mathcal{D} \bar{\phi}_b^\dagger(s') \mathcal{D} \bar{\phi}_b(s') \mathcal{D} \bar{F}_b^\dagger(s) \mathcal{D} \bar{F}_b(s) e^{iS_{sb}} \quad (\text{F40})$$

$$= \prod'_s z_\phi^L(s) \cdot \prod'_{a_1(s) \geq 0} z_F^L(s) \quad (\text{F41})$$

where

$$z_\phi^L(s) = \int_{-\infty}^{\infty} d(\text{Re } \bar{\phi}_b(s')) \int_{-\infty}^{\infty} d(\text{Im } \bar{\phi}_b(s')) e^{i\tilde{a}(s)[(\text{Re } \bar{\phi}_b(s'))^2 + (\text{Im } \bar{\phi}_b(s'))^2]} \quad (\text{F42})$$

$$= i \frac{\pi}{\tilde{a}(s)} \quad (\text{F43})$$

$$z_F^L(s) = \int_{-\infty}^{\infty} d(\text{Re } \bar{F}_b(s)) \int_{-\infty}^{\infty} d(\text{Im } \bar{F}_b(s)) e^{i[(\text{Re } \bar{F}_b(s))^2 + (\text{Im } \bar{F}_b(s))^2]} \quad (\text{F44})$$

$$= i\pi \quad (\text{F45})$$

since $\int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy \exp(ia(x^2 + y^2)) = i\pi/a$. (Nuances of Lorentzian path integrals are discussed in, e.g., Peskin and Schroeder [49], p. 286.) We have then obtained

$$Z_{sb}^L = c_b Z_{sb} \quad (\text{F46})$$

where c_b is a product of factors of i and -1 .

To return to the coordinate representation, let us define the intermediate fields

$$\Phi(x) = \sum'_s U(x, s') \bar{\phi}_b(s') \quad (\text{F47})$$

and

$$\mathcal{F}(x) = \sum'_s U(x, s) \bar{F}_b(s). \quad (\text{F48})$$

Recall that these fields include only positive-action modes and *excitations* of negative-action modes.

As a reminder of the notation, recall that, according to (F5) and (F14),

$$i\sigma^\mu D_\mu U(x, s') = a(s') U(x, s') \quad , \quad i\bar{\sigma}^\mu D_\mu U(x, s') = a(s) U(x, s') \quad (\text{F49})$$

with $\bar{\sigma}^0 = \sigma^0$, $\bar{\sigma}^k = -\sigma^k$, $a(s) = a_1(s) + a_2(s)$, $a(s') = a_1(s) - a_2(s)$, $\tilde{a}(s) = a(s)a(s') = a_1(s)^2 - a_2(s)^2$, and $s > 0$ or < 0 defined by (F29)-(F30) and the line following. Again, in the absence of gauge potentials we have $a_1 = \omega$ and $a_2 = \mp |\vec{p}|$.

We could return to the original coordinate system, with the action in (F39) becoming

$$S_{sb} = S_\Phi + S_{\mathcal{F}} \quad (\text{F50})$$

where

$$S_\Phi = \int d^4x e \bar{\mathcal{L}}_\Phi \quad , \quad S_{\mathcal{F}} = \int d^4x e \bar{\mathcal{L}}_{\mathcal{F}} \quad , \quad \bar{\mathcal{L}}_{\mathcal{F}} = \mathcal{F}^\dagger(x) \mathcal{F}(x) \quad (\text{F51})$$

and $\bar{\mathcal{L}}_\Phi$ is obtained via $\sigma^\mu \rightarrow e^\mu_\alpha \sigma^\alpha$, as in (6.3). However, in the following it is more convenient to remain in the locally inertial frame used above, where

$$\bar{\mathcal{L}}_\Phi = \frac{1}{2} \Phi^\dagger(x) i\bar{\sigma}^\mu D_\mu i\sigma^\nu D_\nu \Phi(x) + \frac{1}{2} \Phi^\dagger(x) i\sigma^\mu D_\mu i\bar{\sigma}^\nu D_\nu \Phi(x) \quad . \quad (\text{F52})$$

Appendix G: Negative-action modes

According to (F28), there is a negative contribution

$$S_-^\phi = - \sum'_{s < 0} \bar{\phi}_b^*(s') |\tilde{a}(s)| \bar{\phi}_b(s') \quad (\text{G1})$$

to the action for what will become scalar boson fields. In the present context we will consider the version of this expression which corresponds to the original fields and coordinates of (4.45), rather than the redefined fields of (4.46) in a locally inertial frame. The results are essentially the same in either version.

It will be assumed that the action for each independent degree of freedom $x_s = \text{Re } \bar{\phi}_b(s')$ or $\text{Im } \bar{\phi}_b(s')$ receives a quartic contribution $\frac{1}{2} b_s x_s^4$ from radiative corrections, stabilizing the vacuum, so that the total action for x_s is

$$-|\tilde{a}(s)| x_s^2 + \frac{1}{2} b_s x_s^4 \quad . \quad (\text{G2})$$

The results below are obtained even in the limit $b_s \rightarrow 0$.

Minimization gives the value

$$\bar{x}_s^2 = |\tilde{a}(s)|/b_s . \quad (\text{G3})$$

An excitation out of the vacuum can be represented by $x_s = \bar{x}_s + \delta x_s$, and the lowest-order additional action is

$$\delta S_-^\phi = \left[\frac{\partial S_-^\phi}{\partial x_s} \right]_{\bar{x}_s} \delta x_s + \frac{1}{2} \left[\frac{\partial^2 S_-^\phi}{\partial x_s^2} \right]_{\bar{x}_s} (\delta x_s)^2 \quad (\text{G4})$$

$$= 0 + [-|\tilde{a}(s)| + 3b_s \bar{x}_s^2] (\delta x)^2 \quad (\text{G5})$$

$$= 2|\tilde{a}(s)| (\delta x)^2 . \quad (\text{G6})$$

Notice that the quartic coefficient b_s is not present in this result (so we could even take $b_s \rightarrow 0$). For an excitation $\delta \bar{\phi}_b(s') = \text{Re } \delta \bar{\phi}_b(s') + i \text{Im } \delta \bar{\phi}_b(s')$, therefore, the action is

$$\delta S_-^\phi = 2|\tilde{a}(s)| \left[(\text{Re } \delta \bar{\phi}_b(s'))^2 + (\text{Im } \delta \bar{\phi}_b(s'))^2 \right] \quad (\text{G7})$$

$$= 2|\tilde{a}(s)| \delta \bar{\phi}_b^*(s') \delta \bar{\phi}_b(s') \quad (\text{G8})$$

$$= |\tilde{a}(s)| \bar{\phi}_b'^*(s') \bar{\phi}_b'(s') \quad (\text{G9})$$

where

$$\bar{\phi}_b'(s') = \sqrt{2} \delta \bar{\phi}_b(s') . \quad (\text{G10})$$

This result also holds if we revert to the fields and coordinates in the specific context of (F28). When $\bar{\phi}_b'(s')$ is renamed $\bar{\phi}_b(s')$, as in (F27), the excitations of negative-action modes have the same form for the action as those of positive-action modes. Of course, there still remains a negative-action “condensate density” buried in the vacuum (and there is also a trivial change in the measure within the path integral).

With $|\tilde{a}(s)| = b_s \bar{x}_s^2$, the contribution of \bar{x}_s to the action is properly written as

$$-\frac{1}{2} b_s \bar{x}_s^4 . \quad (\text{G11})$$

In classical general relativity the matter fields, and thus \bar{x}_s , are held constant while their stress-energy tensor is evaluated according to the left-most equation in (4.51). (This also follows from the prescription that the matter fields and metric tensor vary independently in the path integral. The classical extremalization with respect to matter fields and metric tensor essentially involves partial derivatives with respect to independent variables.) Then

(G11) implies that the classical stress-energy tensor of the negative-action modes is zero, just as for the other vacuum fields.

-
- [1] E.g., R.L. Workman et al. (Particle Data Group), “The Review of Particle Physics (2022)”, Prog. Theor. Exp. Phys. 2022, 083C01 (2022), <https://pdg.lbl.gov/>.
 - [2] ATLAS Collaboration, “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC”, Phys. Lett. B716, 1 (2012), arXiv:1207.7214 [hep-ex].
 - [3] CMS Collaboration, “Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC”, Phys. Lett. B716, 30 (2012), arXiv:1207.7235 [hep-ex].
 - [4] H. Baer and X. Tata, *Weak Scale Supersymmetry: From Superfields to Scattering Events* (Cambridge University Press, 2006).
 - [5] Pran Nath, *Supersymmetry, Supergravity, and Unification* (Cambridge University Press, 2016).
 - [6] R. N. Mohapatra, *Unification and Supersymmetry: The Frontiers of Quark-Lepton Physics* (Springer, 2010).
 - [7] *Perspectives on Supersymmetry II*, edited by G. L. Kane (World Scientific, 2010).
 - [8] Joseph A. Formaggio, André Luiz C. de Gouvêa, and R.G. Hamish Robertson, “Direct measurements of neutrino mass”, Physics Reports 914, 1 (2021), and references therein.
 - [9] T.-P. Cheng and L.-F. Li, *Gauge Theory of Elementary Particle Physics* (Clarendon, Oxford, 1984).
 - [10] C. Kounnas, A. Masiero, D. V. Nanopoulos, and K. A. Olive, *Grand Unification with and without Supersymmetry and Cosmological Implications* (World Scientific, 1984).
 - [11] G. G. Ross, *Grand Unified Theories* (Benjamin, 1985), and references therein.
 - [12] V. D. Barger and R. J. N. Phillips, *Collider Physics Updated Edition* (Addison-Wesley, 1997).
 - [13] A. G. Riess et al., “Observational evidence from supernovae for an accelerating universe and a cosmological constant”, Astron. J. 116, 1009 (1998).
 - [14] S. Perlmutter et al., “Measurement of Ω and Λ from 42 high-redshift supernovae”, Astrophys. J. 517, 565 (1999).
 - [15] S. Weinberg, “The cosmological constant problem”, Rev. Mod. Phys. 61, 1 (1989).

- [16] S. Weinberg, “Anthropic Bound on the Cosmological Constant”, *Phys. Rev. Lett.* 59, 2607 (1987).
- [17] Steven Weinberg, “The Cosmological Constant Problems (Talk given at Dark Matter 2000, February, 2000)”, arXiv:astro-ph/0005265.
- [18] Gianfranco Bertone and Dan Hooper, “History of dark matter”, *Rev. Mod. Phys.* 90, 45002 (2018), arXiv:1605.04909 [astro-ph.CO].
- [19] Reagan Thornberry, Maxwell Throm, John Killough, Dylan Blend, Michael Erickson, Brian Sun, Brett Bays, Gabriel Frohaug, and Roland E. Allen, “Experimental signatures of a new dark matter WIMP”, *EPL [European Physics Letters]* 134, 49001 (2021), arXiv:2104.11715 [hep-ph], and references therein.
- [20] Caden LaFontaine, Bailey Tallman, Spencer Ellis, Trevor Croteau, Brandon Torres, Sabrina Hernandez, Diego Cristancho Guerrero, Jessica Jaksik, Drue Lubanski, and Roland E. Allen, “A Dark Matter WIMP That Can Be Detected and Definitively Identified with Currently Planned Experiments”, *Universe* 7, 270 (2021), arXiv:2107.14390 [hep-ph], and references therein.
- [21] Bailey Tallman, Alexandra Boone, Caden LaFontaine, Trevor Croteau, Quinn Ballard, Sabrina Hernandez, Spencer Ellis, Adhithya Vijayakumar, Fiona Lopez, Samuel Apata, Jehu Martinez, and Roland Allen, “Indirect detection, direct detection, and collider detection cross-sections for a 70 GeV dark matter WIMP”, *Proceedings of Science*, <https://pos.sissa.it/414/988/pdf> [proceedings of the 41st International Conference on High Energy Physics, ICHEP 2022], arXiv:2210.05380 [hep-ph].
- [22] Bailey Tallman, Alexandra Boone, Adhithya Vijayakumar, Fiona Lopez, Samuel Apata, Jehu Martinez, and Roland Allen, “Potential for definitive discovery of a 70 GeV dark matter WIMP with only second-order gauge couplings”, *Letters in High Energy Physics LHEP-342* (2023), doi.org/10.31526/lhep.2023, arXiv:2210.15019 [hep-ph].
- [23] B. S. DeWitt, *The Global Approach to Quantum Field Theory* (Oxford University Press, 2014).
- [24] B. S. DeWitt, “Quantum Theory of Gravity. I. The Canonical Theory”, *Phys. Rev.* 160, 1113 (1967).
- [25] J. B. Hartle and S. W. Hawking, “Wave function of the Universe”, *Phys. Rev. D.* 28, 2960 (1983), and references therein.
- [26] J. A. Wheeler, “Information, physics, quantum: the search for links”, *Proceedings of the*

- 3rd International Symposium on Foundations of Quantum Mechanics in the Light of New Technology: Central Research Laboratory, Hitachi, Tokyo, Japan, August 28-31, 1989, pp. 354-368, <https://philpapers.org/archive/WHEIPQ.pdf>.
- [27] Philip Candelas and Steven Weinberg, “Calculation of gauge couplings and compact circumferences from self-consistent dimensional reduction”, Nucl. Phys. B 237, 397 (1984).
- [28] J. Elias-Miro, J. R. Espinosa, G. F. Giudice, G. Isidori, A. Riotto, and A. Strumia, “Higgs mass implications on the stability of the electroweak vacuum”, Phys. Lett. B 709, 222 (2012).
- [29] S. Alekhin, A. Djouadi, and S. Moch, “The top quark and Higgs boson masses and the stability of the electroweak vacuum”, Phys. Lett. B 716, 214 (2012), arXiv:1207.0980 [hep-ph], and references therein.
- [30] G. Degrassi, S. Di Vita, J. Elias-Miró, J. R. Espinosa, G. F. Giudice, G. Isidori, and A. Strumia, “Higgs mass and vacuum stability in the Standard Model at NNLO”, JHEP 1208, 098 (2012), arXiv:1205.6497 [hep-ph].
- [31] D. Buttazzo, G. Degrassi, P. P. Giardino, G. F. Giudice, F. Sala, A. Salvio, and A. Strumia, “Investigating the near-criticality of the Higgs boson”, JHEP 12, 089 (2013), arXiv:1307.3536 [hep-ph].
- [32] Roland E. Allen, “An alternative form of supersymmetry with reduced cross-sections and modified experimental signatures”, arXiv:2307.04255.
- [33] R. D. Peccei and Helen R. Quinn, “ CP Conservation in the Presence of Pseudoparticles”, Phys. Rev. Lett. 38, 1440 (1977).
- [34] Steven Weinberg, “A New Light Boson?”, Phys. Rev. Lett. 40, 223 (1978).
- [35] F. Wilczek, “Problem of Strong P and T Invariance in the Presence of Instantons”, Phys. Rev. Lett. 40, 279 (1978).
- [36] P. Sikivie, “Experimental Tests of the “Invisible” Axion”, Phys. Rev. Lett. 51, 1415 (1983), and references therein.
- [37] P. Sikivie, N. Sullivan, and D. B. Tanner, “Proposal for Axion Dark Matter Detection Using an LC Circuit”, Phys. Rev. Lett. 112,131301 (2014), arXiv:1310.8545 [hep-ph], and references therein.
- [38] R. L. Jaffe, “Casimir effect and the quantum vacuum”, Phys. Rev. D 72, 021301 (2005), arXiv:hep-th/0503158.
- [39] M. Bordag, G. L. Klimchitskaya, U. Mohideen, and V. M. Mostepanenko, *Advances in the*

- Casimir Effect* (Oxford University Press, 2015). See, e.g., the paragraph on p. 6 concerning the real-world Casimir effect.
- [40] W. M. R. Simpson and U. Leonhardt, *Forces of the Quantum Vacuum: An Introduction to Casimir Physics* (World Scientific, 2015). See, e.g., p. 53.
- [41] G. Parisi, *Field Theory, Disorder and Simulations* (World Scientific, 1992).
- [42] K. Efetov, *Supersymmetry in Disorder and Chaos* (Cambridge University Press, 1999).
- [43] K. Huang, *Quantum Field Theory From Operators to Path Integrals* (Wiley, 1998), p. 292.
- [44] Leonard Parker and David Toms, *Quantum Field Theory in Curved Spacetime: Quantized Fields and Gravity* (Cambridge University Press, 2009).
- [45] Michael B. Green, John H. Schwarz, and Edward Witten, *Superstring Theory: Volume 2, Loop Amplitudes, Anomalies and Phenomenology* (Cambridge University Press, 1988).
- [46] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (W. H. Freeman, 1973).
- [47] Steven Weinberg, *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity* (Wiley, 1972).
- [48] Robert M. Wald, *General Relativity* (University of Chicago Press, 1984).
- [49] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory* (Perseus, 1995).
- [50] M. D. Schwartz, *Quantum Field Theory and the Standard Model* (Cambridge University Press, 2014).
- [51] S. Weinberg, *The Quantum Theory of Fields*, Vol. I (Cambridge University Press, 1995).
- [52] P. B. Littlewood and C. M. Varma, “Amplitude collective modes in superconductors and their coupling to charge-density waves”, *Phys. Rev. B* 26, 4883 (1982).
- [53] David Pekker and C.M. Varma, “Amplitude/Higgs Modes in Condensed Matter Physics”, *Annu. Rev. Condens. Matter Phys.* 6, 269 (2015), arXiv:1406.2968.
- [54] Ryo Shimano and Naoto Tsuji, “Higgs Mode in Superconductors”, *Annu. Rev. Condens. Matter Phys.* 11, 103 (2020), arXiv:1906.09401.
- [55] Howard Baer, Vernon Barger, and Hasan Serce, “SUSY under siege from direct and indirect WIMP detection experiments”, *Phys. Rev. D* 94, 115019 (2016), arXiv:1609.06735 [hep-ph].
- [56] Howard Baer, Vernon Barger, Dibyashree Sengupta, and Xerxes Tata, “Is natural higgsino-only dark matter excluded?”, *Eur. Phys. J. C* 78, 838 (2018), arXiv:1803.11210 [hep-ph].
- [57] Leszek Roszkowski, Enrico Maria Sessolo, and Sebastian Trojanowski, “WIMP dark matter candidates and searches - current status and future prospects”, *Rept. Prog. Phys.* 81, 066201

- (2018), arXiv:1707.06277 [hep-ph].
- [58] Howard Baer, Dibyashree Sengupta, Shadman Salam, Kuver Sinha, and Vernon Barger, “Mid-review: Status of weak scale supersymmetry after LHC Run 2 and ton-scale noble liquid WIMP searches”, *Eur. Phys. J. Spec. Top.* 229, 3085 (2020), arXiv:2002.03013 [hep-ph].
- [59] Xerxes Tata, “Natural Supersymmetry: Status and Prospects”, *Eur. Phys. J. Spec. Top.* 229, 3061 (2020), arXiv:2002.04429 [hep-ph].
- [60] G. Belanger, F. Boudjema, A. Pukhov, and A. Semenov, “micrOMEGAs: a tool for dark matter studies”, *Nuovo Cimento* 33, 111 (2010), arXiv:1005.4133 [hep-ph], and references therein; <http://lapth.cnrs.fr/micromegas/>.
- [61] D. S. Akerib et al. [LUX-ZEPLIN Collaboration], “Projected WIMP sensitivity of the LUX-ZEPLIN dark matter experiment”, *Phys. Rev. D* 101, 052002 (2020), arXiv:1802.06039 [astro-ph.IM].
- [62] Kelly Stifter [on behalf of the LZ collaboration], “First dark matter search results from the LUX-ZEPLIN (LZ) experiment”, talk at International Conference on Neutrinos and Dark Matter (NuDM-2022), September 25-28, 2022, Sharm El-Sheikh, Egypt.
- [63] J. Aalbers et al. [LUX-ZEPLIN (LZ) Collaboration], “First Dark Matter Search Results from the LUX-ZEPLIN (LZ) Experiment”, arXiv:2207.03764 [hep-ex].
- [64] E. Aprile et al. [XENON collaboration], “Projected WIMP Sensitivity of the XENONnT Dark Matter Experiment”, *JCAP* 11, 031 (2020), arXiv:2007.08796 [physics.ins-det].
- [65] E. Aprile et al. [XENON collaboration], “An approximate likelihood for nuclear recoil searches with XENON1T data”, arXiv:2210.07231 [hep-ex].
- [66] See also Yue Meng et al. [PandaX collaboration], “Dark Matter Search Results from the PandaX-4T Commissioning Run”, *Phys. Rev. Lett.* 127, 261802 (2021), arXiv:2107.13438 [hep-ex].
- [67] J. Aalbers et al. (DARWIN collaboration), “DARWIN: towards the ultimate dark matter detector”, *JCAP* 1611, 017 (2016), arXiv:1606.07001 [astro-ph.IM] and “A Next-Generation Liquid Xenon Observatory for Dark Matter and Neutrino Physics”, *J. Phys. G: Nucl. Part. Phys.* 50, 013001 (2023), arXiv:2203.02309 [physics.ins-det].
- [68] Bailey Tallman et al.; Kane Rylander et al.; and Jehu Martinez et al., private communications based on MadGraph calculations, and to be published.
- [69] J. Billard, E. Figueroa-Feliciano, and L. Strigari, “Implication of neutrino backgrounds on the

- reach of next generation dark matter direct detection experiments”, *Phys. Rev. D* 89, 023524 (2014), arXiv:1307.5458 [hep-ph].
- [70] Alexandre Alvarez, Francesca Calore, Anna Genina, Justin Read, Pasquale Dario Serpico, and Bryan Zaldivar, “Dark matter constraints from dwarf galaxies with data-driven J-factors”, *JCAP* 09, 004 (2020), arXiv:2002.01229 [astro-ph.HE].
- [71] G. Steigman, B. Dasgupta, and J. F. Beacom, “Precise relic WIMP abundance and its impact on searches for dark matter annihilation”, *Phys. Rev. D* 86 023506 (2012), arXiv:1204.3622 [hep-ph].
- [72] Planck Collaboration, “Planck 2018 results. VI. Cosmological parameters”, arXiv:1807.06209 [astro-ph.CO].
- [73] Rebecca K. Leane, “Indirect Detection of Dark Matter in the Galaxy”, arXiv:2006.00513 [hep-ph].
- [74] Lisa Goodenough and Dan Hooper, “Possible Evidence For Dark Matter Annihilation In The Inner Milky Way From The Fermi Gamma Ray Space Telescope”, arXiv:0910.2998 [hep-ph].
- [75] Vincenzo Vitale and Aldo Morselli (for the Fermi/LAT Collaboration), “Indirect Search for Dark Matter from the center of the Milky Way with the Fermi-Large Area Telescope”, arXiv:0912.3828 [astro-ph.HE].
- [76] Christopher Karwin, Simona Murgia, Tim M. P. Tait, Troy A. Porter, and Philip Tanedo, “Dark matter interpretation of the Fermi-LAT observation toward the Galactic Center”, *Phys. Rev. D* 95, 103005 (2017), arXiv:1612.05687 [hep-ph], and references therein.
- [77] Rebecca K. Leane, Tracy R. Slatyer, John F. Beacom, and Kenny C. Y. Ng, “GeV-scale thermal WIMPs: Not even slightly ruled out”, *Phys. Rev. D* 98, 023016 (2018), arXiv:1805.10305 [hep-ph].
- [78] Rebecca K. Leane and Tracy R. Slatyer, “Revival of the Dark Matter Hypothesis for the Galactic Center Gamma-Ray Excess”, *Phys. Rev. Lett.* 123, 241101 (2019), arXiv:1904.08430 [astro-ph.HE], and references therein.
- [79] Alessandro Cuoco, Jan Heisig, Michael Korsmeier, and Michael Krämer, “Probing dark matter annihilation in the Galaxy with antiprotons and gamma rays”, *JCAP* 10, 053 (2017), arXiv:1704.08258 [astro-ph.HE].
- [80] Alessandro Cuoco, Michael Krämer, and Michael Korsmeier, “Novel dark matter constraints from antiprotons in the light of AMS-02”, *Phys. Rev. Lett.* 118, 191102 (2017),

- arXiv:1610.03071 [astro-ph.HE].
- [81] Ming-Yang Cui, Qiang Yuan, Yue-Lin Sming Tsai, and Yi-Zhong Fan, “Possible dark matter annihilation signal in the AMS-02 antiproton data”, *Phys. Rev. Lett.* 118, 191101 (2017), arXiv:1610.03840 [astro-ph.HE].
- [82] Ilias Cholis, Tim Linden, and Dan Hooper, “A Robust Excess in the Cosmic-Ray Antiproton Spectrum: Implications for Annihilating Dark Matter”, *Phys. Rev. D* 99, 103026 (2019), arXiv:1903.02549 [astro-ph.HE].
- [83] Alessandro Cuoco, Jan Heisig, Lukas Klamt, Michael Korsmeier, and Michael Krämer, “Scrutinizing the evidence for dark matter in cosmic-ray antiprotons”, *Phys. Rev. D* 99, 103014 (2019), arXiv:1903.01472 [astro-ph.HE].
- [84] See also Dan Hooper, Ilias Cholis, Tim Linden, and Ke Fang, “HAWC Observations Strongly Favor Pulsar Interpretations of the Cosmic-Ray Positron Excess”, *Phys. Rev. D* 96, 103013 (2017), arXiv:1702.08436 [astro-ph.HE].
- [85] L. Perivolaropoulos and F. Skara, “Challenges for Λ CDM: An update”, doi:10.1016/j.newar.2022.101659, arXiv:2105.05208 [astro-ph.CO].
- [86] H. Kleinert, *Multivalued Fields: in Condensed Matter, Electromagnetism, and Gravitation* (World Scientific, 2008).
- [87] H. Narumi and T. Nakau, “On the Quantization of Angular Momentum in Multi-Dimensional Space”, *Bull. Inst. Chem. Research Kyoto Univ.* 26, 26 (1951); 27, 29 (1951).
- [88] G. A. Gallup, “Angular Momentum in n-Dimensional Spaces”, *J. Mol. Spec.* 3, 673 (1959).
- [89] J. D. Louck, “Generalized Orbital Angular Momentum and the n-Fold Degenerate Quantum-Mechanical Oscillator, Part II, The n-Fold Degenerate Oscillator”, *J. Mol. Spec.* 4, 298 (1960).
- [90] D. R. Herrick, “Degeneracies in energy levels of quantum systems of variable dimensionality”, *J. Math. Phys.* 16, 281 (1975).
- [91] S.-H. Dong, *Wave Equations in Higher Dimensions* (Springer, 2011), pp. 41-45.
- [92] R. Friedberg, T. D. Lee, and W. Q. Zhao, “Convergent iterative solutions for a Sombrero-shaped potential in any space dimension and arbitrary angular momentum”, *Annals of Physics* 321, 1981 (2006).