Entangled probability distributions

Vladimir N. Chernega,¹ Olga V. Man'ko,^{2,3} Vladimir I. Man'ko^{2,4}

¹Russian University of Transport (MIIT), Moscow, Russia

²Lebedev Physical Institute, Russian Academy of Sciences Leninskii Prospect 53, Moscow 119991, Russia

³Bauman Moscow State Technical University The 2nd Baumanskaya Street 5, Moscow 105005, Russia

⁴Moscow Institute of Physics and Technology (State University) Institutskii per. 9, Dolgoprudny, Moscow Region 141700, Russia

*Corresponding author e-mail: mankoov@lebedev.ru

Abstract

Concept of entangled probability distribution of several random variables is introduced. These probability distributions describe multimode quantum states in probability representation of quantum mechanics. Example of entangled probability distribution is considered.

Keywords: probability distribution, entanglement, tomograms.

1 Introduction.

The foundations of conventional probability theory were considered long ago and are described, e.g. in books [1, 2, 3]. The probability theory is necessary to be used to describe random process [3] both classical and quantum. Also to construct the theory of quantum phenomena like quantum mechanics, quantum statistics, quantum field theory one needs to apply all the aspects of the conventional probability theory. The aim of our work is to study the possibility to use the problem related to back action of the results obtained in quantum mechanics to clarify some aspects which are new in conventional probability theory. The point is that in quantum mechanics new interesting phenomenon called entanglement of quantum system states was found [4, 5]. It

example, the review of the entanglement phenomenon in information theory was discussed in [6]. The phenomenon of entanglement is related to the properties of quantum systems like Bell inequalities [7]. For example, quantum properties reflected from Bell inequalities were discussed in [8, 9, 10, 11, 12]. On the other hand recently it was clarified that all quantum states of physical systems can be described by conventional probability distributions using the introduced probability representation of quantum mechanics [13, 14, 15, 16, 17]. The spin tomography was introduced in [18, 19] and discussed in [20, 21, 22, 23]. In [9] in the frame of probability representation of quantum mechanics the spin tomography was applied to Bell's inequality problem, and was shown that the satisfaction or violation of Bell's inequalities can be understood as properties of tomographic functions for joint probability distributions for two spins.

In the probability representation entanglement phenomenon is also associated with the specific properties of the quantum states which may be considered as properties determined by the specific properties of used probability distributions which we call entangled probability distributions. In our work we introduce autonomous notion of the entangled probability distribution and consider their properties on few examples induced by the existed entanglement phenomena in quantum mechanics like these phenomena in oscillator system states.

In quantum mechanics very important phenomenon is the superposition principle. The superposition principle of wave functions [24] can be discussed using density matrices describing quantum states introduced by [25, 26]. The nonlinear addition of density matrices describing quantum states corresponds to the linear combination of the wave function [27, 28, 29]. We will use the map of tomograms and density matrices [30, 31, 32, 33] to study properties corresponding to entanglement phenomenon. The tomograms and the entanglement phenomenon in the two mode squeezed states and two-mode even and odd coherent states was considered in [34, 35]. The application of the symplectic tomography scheme to the Stimulated Raman Scattering (SRS) and to the Stimulated Brillouin Scattering was done in [36, 37, 38, 39] and the entanglement in these processes were discussed. The classical propagator of SRS was introduced in [40]. The tomographic entropy, information, entanglement phenomenon and quantum distances between quantum states of photon-phonon mode of SRS in the probability

The quantum entanglement in the Raman Scattering was also investigated in [42, 43]. It is necessary to add that optical tomograms of the time-evolved states generated by the evolution of different kinds of initial wave packets in a Kerr medium and optical tomograms of maximally entangled states generated at the output modes of a beam splitter was theoretically studied in [44], [45]. The entanglement criterion in probability representation was suggested in [46, 47]. In [48] a review of the Radon transform and the instability of the tomographic reconstruction process were discussed. In [49] it was shown that in classical mechanics can be introduced the Hermitian operators and the concepts of classical mechanics can be formulated using the inverse Wigner–Weyl transform of the classical probability distributions. In [50] the attempts to study the problems of complex biosystems composed of many subsystems and in [51] the attempts to model social process using the language of quantum mechanics, thermodynamics, quantum information and field theory were done. In [52] a review of classical probability representations of quantum states and observables is done and in [53] quantum nonlocality is discussed. New fundamental aspects of quantum mechanics based on groupoid approach are investigated in [54]. Development of the tomographic probability distribution method and different aspects of its applications are presented in [55]. The classical and quantum aspects of tomography were discussed in [56] and classical-quantum dynamics were considered in [57]. The probability representation of quantum states is used recently to study properties of cosmology [58, 59, 60]. The review of the harmonic analyses of the density matrix properties including the method of symplectic tomography is given in [61, 62, 63]. The methods of star-product, tomography and probability representation of quantum mechanics were applied to different problems of quantum phenomena in [64, 65, 66, 67]. The geometrical interpretation of spin state in probability representation with the help of Malevich's squares is done in [68, 69, 70, 71, 72, 73, 74].

We give a review of the tomographic probability representation of quantum mechanics. In [75] the formalism of quantum states and quantum observables using the formalism of standard probability distributions and classical-like random variables is presented and the evolution equation and energy spectra in the form of equations for probability distributions determining the coherent and number states of photons are Applications of quantum tomographic approach to different kinds of experiments as well as theoretical researches were discussed in [76, 77, 78, 79, 80]. The tomographic causal analysis of two-qubit states was done and theoretically was studied operations with a four-level superconducting circuit as a two-qubit system in [81]. The tomograms of the state of a quantum circuit and of the Josephson junction were considered in [82]. The probability representation of the nonlinear coherent states of an ion in the Paul trap was introduced in [83], the Schrödinger cat states of a trapped ion in the Paul trap were discussed in [84], the tomograms of an ion in the Paul and Penning traps were considered in [85]. The tomograms of the states of the damped oscillator with Kronig–Penney excitation were considered in [86]. The tomograms of the quark states were introduces in [87].

The consideration of classical mechanics within the framework of the tomographic representation was done in [16], [88], [89], [90], [91]. The relation between the quantum state description and the classical state description is elucidated in [92]. The quantum-classical limits for quantum tomograms are studied and compared with the corresponding classical tomograms in [93]. The inverted oscillator was discussed recently in [94, 95] in connection with relation to cosmological problems. The inverted oscillator in the probability representation of quantum mechanics was considered in [96]. Quantum evolution equation for open system was considered in [97, 98, 99]. Schrödinger and von Neumann equations, as well as equations for the evolution of open systems, are written in the form of linear classicallike equations for the probability distributions determining the quantum system states in [100]. The evolution of tomograms for different quantum systems, both finite and infinite dimensional was considered in [101]. A method based on tomographic representation to simulate the quantum dynamics was applied to the wave packet tunneling of one and two interacting particles in [102].

The paper is organized as follows. In Sec.2 we provide definition of entangled probability distributions and describe some their properties. In Sec.3 we present the examples of entangled probability distributions of two random continious variables. The conclusions and prospectives are given in Sec.4.

2 Probability distributions

Let us consider the set of continious random variables $\vec{x} = (x_1, x_2, \dots, x_N)$ and parameter $\vec{a} = (a_1, a_2, \dots, a_N)$ which can take some values and we consider a function of these variables $0 \le w(\vec{x}|\vec{a}) \le 1$. This function satisfies the condition

$$\int w(\vec{x}|\vec{a}) \prod_{k=1}^{N} dx_k = 1$$

We consider this nonnegative normalized function as a conditional probability distribution of N random variables. Then we can introduce nonnegative functions $w_k(x_k|a_k)$ using the definition

$$w_k(x_k|a_k) = \int w(\vec{x}|\vec{a}) \prod_{k' \neq k=1}^N dx'_k.$$
 (1)

We call the probability distribution $w(\vec{x}|\vec{a})$ separable one if it equals to convex sum of the products of functions (1)

$$w(\vec{x}|\vec{a}) = \sum_{s} \mathcal{P}_s \prod_{s=1}^{N} w_s(x_s|a_s), \quad \mathcal{P}_s \ge 0, \quad \sum_{s} \mathcal{P}_s = 1.$$
⁽²⁾

If the function $w(\vec{x}|\vec{a})$ can not be represented in the form of convex sum of the functions (1) we call the probability distribution $w(\vec{x}|\vec{a})$ entangled probability distribution, i.e.

$$w(\vec{x}|\vec{a}) \neq \sum_{s} \mathcal{P}_{s} \prod_{s=1}^{N} w_{s}(x_{s}|a_{s}).$$
(3)

In the case of two variables x_1, x_2 and parameters a_1, a_2 we have for entangled probability distribution $w(x_1, x_2|a_1, a_2)$ the definition

$$w(x_1, x_2|a_1, a_2) \neq \sum_s \mathcal{P}_s w_s(x_1|a_1) w_s(x_2|a_2).$$
(4)

3 Examples of the entangled probability distributions

Let us remind the value of the propagator for the inverted oscillator evolution

$$G(x, y, t) = \frac{1}{\sqrt{2\pi \sinh t}} \exp\left[\frac{i}{2}\left(\coth t(y^2 + x^2) - \frac{2xy}{\sinh t}\right)\right].$$
(5)

We will use this formula later. In order to consider examples we will study properties

form

$$\psi(x,y) = \frac{1}{\sqrt{2}} \left(\psi_0(x)\psi_1(y) + \psi_1(x)\psi_0(y) \right) = \frac{x+y}{\sqrt{\pi}} \exp\left(-\frac{x^2}{2} - \frac{y^2}{2}\right).$$
(6)

The function (6) is the superposition of the wave functions of one-mode oscillators, where first function is ground state of the first oscillator

$$\psi_0(x) = \frac{e^{-\frac{x^2}{2}}}{\pi^{1/4}} \tag{7}$$

and the second function is the first excited state of the second oscillator

$$\psi_1(y) = \frac{\sqrt{2}y}{\pi^{1/4}} e^{-\frac{y^2}{2}}.$$
(8)

The normalization constant is such that we have

$$\int_{-\infty}^{\infty} \psi(x,y)\psi^*(x,y)dxdy = 1.$$
(9)

We calculated this normalization condition using the known formula

$$\int_{-\infty}^{\infty} \exp\left(-ax^2 + bx\right) dx = \sqrt{\frac{\pi}{a}} \exp\left(\frac{b^2}{4a}\right).$$
(10)

Considering the derivative of both sides of the above equality with respect to the parameter b we obtain the integral relation

$$\int_{-\infty}^{\infty} x \exp\left(-ax^2 + bx\right) dx = \sqrt{\frac{\pi}{a}} \frac{b}{2a} \exp\left(\frac{b^2}{4a}\right).$$
(11)

To calculate another integral we calculate derivative of the both sides of equality (10) with respect to the parameter -a then we get the value of the integral

$$\int_{-\infty}^{\infty} x^2 \exp\left(-ax^2 + bx\right) dx = \frac{1}{2a} \sqrt{\frac{\pi}{a}} \exp\left(\frac{b^2}{4a}\right) \left(1 + \frac{b^2}{2a}\right). \tag{12}$$

Using the values of calculating integrals we get the normalization condition

$$\frac{1}{\pi} \int_{-\infty}^{\infty} (x+y)^2 \exp\left(-x^2 - y^2\right) dx dy = 1.$$
(13)

To consider the map of the wave function $\psi(x, y)$ onto the probability distribution let us study the integral

$$A = \int_{-\infty}^{\infty} \psi(x, y) \exp\left[i\left(\frac{\mu_1 x^2}{2\nu_1} - \frac{Xx}{\nu_1} + \frac{\mu_2 y^2}{2\nu_2} - \frac{Yy}{\nu_2}\right)\right] dxdy.$$
 (14)

The integrals (10)-(12) give the possibility to obtain the relation

$$A = \frac{\sqrt{\pi}}{\sqrt{\left(1 - \frac{i\mu_1}{\nu_1}\right)\left(1 - \frac{i\mu_2}{\nu_2}\right)}} \left(-\frac{2iX}{\nu_1\left(1 - \frac{i\mu_1}{\nu_1}\right)} - \frac{2iY}{\nu_2\left(1 - \frac{i\mu_2}{\nu_2}\right)}\right)$$
$$\times \exp\left[-\frac{X^2}{2\nu_1^2\left(1 - \frac{i\mu_1}{\nu_1}\right)} - \frac{Y^2}{2\nu_2^2\left(1 - \frac{i\mu_2}{\nu_2}\right)}\right]. \tag{15}$$

The probability distribution related to these formulae reads (see, for example, [100])

$$w = \frac{AA^*}{4\pi^2 |\nu_1| |\nu_2|}.$$
(16)

In explicit form the probability distribution $w(X, Y | \mu_1, \nu_1, \mu_2, \nu_2)$ is given by the relation

$$w(X,Y|\mu_1,\nu_1,\mu_2,\nu_2) = \frac{(\nu_2^2 + \mu_2^2) X^2 + 2 (\nu_1 \nu_2 + \mu_1 \mu_2) XY + (\nu_1^2 + \mu_1^2) Y^2}{\pi (\nu_1^2 + \mu_1^2)^{3/2} (\nu_2^2 + \mu_2^2)^{3/2}} \times \exp\left[-\frac{X^2}{\mu_1^2 + \nu_1^2} - \frac{Y^2}{\mu_2^2 + \nu_2^2}\right].$$
(17)

For particular case $\nu_1 = \nu_2 = 1$, $\mu_1 = \mu_2 = 0$, one gets

$$w(X, Y|\mu_1 = 0, \nu_1 = 1, \mu_2 = 0, \nu_2 = 1) = \frac{1}{\pi} (X + Y)^2 \exp\left(-X^2 - Y^2\right).$$
 (18)

One can check, that the function $w(X, Y, |\mu_1, \nu_1, \mu_2, \nu_2)$ (18) satisfy the condition

$$\int \int w(X, Y, |\mu_1, \nu_1, \mu_2, \nu_2) dX dY = 1.$$
(19)

As we know, this probability distribution function determines the quantum states which are entangled states. Due to this we call this probability distribution entangled probability distribution. In quantum mechanics the wave functions of two-mode oscillators which are obtained by means of superposition of two different wave functions are entangled pure states. In connection with this the tomographic probability distribution is described by the probability distribution function (17) and it cannot be represented in the form of equation (2). On the other hand the integral

$$w(X|\mu_1,\nu_1) = \int w(X,Y|\mu_1,\nu_1,\mu_2,\nu_2)dY = \frac{\exp\left(-\frac{X^2}{\mu_1^2+\nu_1^2}\right)}{\sqrt{\pi\left(\mu_1^2+\nu_1^2\right)}} \left[\frac{1}{2} + \frac{X^2}{\mu_1^2+\nu_1^2}\right].$$
 (20)

One can check, that

This function $w(X|\mu_1,\nu_1)$ is marginal conditional probability distribution of position X which is the position of the first oscillator and the conditions are labeled by the real parameters μ_1 , ν_1 . Also, if we repeat analogous calculations for the second oscillator we will get

$$w(Y|\mu_2,\nu_2) = \int w(X,Y|\mu_1,\nu_1,\mu_2,\nu_2) dX = \frac{\exp\left(-\frac{Y^2}{\mu_2^2+\nu_2^2}\right)}{\sqrt{\pi\left(\mu_2^2+\nu_2^2\right)}} \left[\frac{1}{2} + \frac{Y^2}{\mu_2^2+\nu_2^2}\right].$$
 (22)

One can check, that

$$\int w(Y|\mu_2, \nu_2) dY = 1.$$
(23)

This function $w(Y|\mu_2, \nu_2)$ is marginal conditional probability distribution of position Y which is the position of the second oscillator and the conditions are labeled by the real parameters μ_2 , ν_2 .

Let us calculate other probability distributions given by two density operators $\hat{\bar{\rho}}_0,$ $\hat{\bar{\rho}}_1$

$$\hat{\rho}(0,1) = \frac{1}{2} \left(\hat{\bar{\rho}}_0 \hat{\tilde{\rho}}_1 + \hat{\bar{\rho}}_1 \hat{\tilde{\rho}}_0 \right),$$
(24)

here $\hat{\rho}_0(x, x') = \psi_0(x)\psi_0^*(x')$ and $\hat{\rho}_1(y, y') = \psi_1(y)\psi_1^*(y')$. This density operator $\hat{\rho}(0, 1)$ describes the separable states of the two-mode oscillator. Other two separable states are described by operators $\hat{\rho}_0\hat{\rho}_1$ and $\hat{\rho}_1\hat{\rho}_0$. The state with probability distribution (17) is the entangled state. The three states which are separable states are described by the probability distributions

$$w_{\hat{\rho}(0,1)}(X,Y|\mu_1,\nu_1,\mu_2,\nu_2) = \frac{\exp\left(-\frac{X^2}{\mu_1^2+\nu_1^2} - \frac{Y^2}{\mu_2^2+\nu_2^2}\right)}{\pi\left(\mu_1^2+\nu_1^2\right)^{1/2}\left(\mu_2^2+\nu_2^2\right)^{1/2}} \left(\frac{Y^2}{\mu_2^2+\nu_2^2} + \frac{X^2}{\mu_1^2+\nu_1^2}\right), \quad (25)$$

$$w_{\hat{\bar{\rho}}_{0}\hat{\bar{\rho}}_{1}}(X,Y|\mu_{1},\nu_{1},\mu_{2},\nu_{2}) = \frac{2Y^{2}}{\pi \left(\mu_{1}^{2}+\nu_{1}^{2}\right)^{1/2} \left(\mu_{2}^{2}+\nu_{2}^{2}\right)^{3/2}} \exp\left(-\frac{X^{2}}{\mu_{1}^{2}+\nu_{1}^{2}}-\frac{Y^{2}}{\mu_{2}^{2}+\nu_{2}^{2}}\right),$$

$$w_{\hat{\bar{\rho}}_{1}\hat{\bar{\rho}}_{0}}(X,Y|\mu_{1},\nu_{1},\mu_{2},\nu_{2}) = \frac{2X^{2}}{\pi \left(\mu_{2}^{2}+\nu_{2}^{2}\right)^{1/2} \left(\mu_{1}^{2}+\nu_{1}^{2}\right)^{3/2}} \exp\left(-\frac{X^{2}}{\mu_{1}^{2}+\nu_{1}^{2}}-\frac{Y^{2}}{\mu_{2}^{2}+\nu_{2}^{2}}\right).$$

$$(26)$$

$$(26)$$

$$(27)$$

Comparing (17) with the (25), (26), 27) we see that the entangled state probability distribution (17) is essentially different from the separable state probability distributions

Let us calculate the marginal distributions for the probability distributions given by the formulae (25), (26), (27). We get

$$w_{\hat{\rho}(0,1)}(X|\mu_1,\nu_1) = \int w_{\hat{\rho}(0,1)}(X,Y|\mu_1,\nu_1,\mu_2,\nu_2)dY = \frac{1}{2\sqrt{\pi (\nu_1^2 + \mu_1^2)}} \exp\left(-\frac{X^2}{\mu_1^2 + \nu_1^2}\right) \left(1 + \frac{2X^2}{\mu_1^2 + \nu_1^2}\right),$$
(28)

$$w_{\hat{\rho}(0,1)}(Y|\mu_2,\nu_2) = \int w_{\hat{\rho}(0,1)}(X,Y|\mu_1,\nu_1,\mu_2,\nu_2)dX = \frac{1}{2\sqrt{\pi(\nu_2^2 + \mu_2^2)}} \exp\left(-\frac{Y^2}{\mu_2^2 + \nu_2^2}\right) \left(1 + \frac{2Y^2}{\mu_2^2 + \nu_2^2}\right),$$
(29)

$$w_{\hat{\rho}_{0}\hat{\rho}_{1}}(X|\mu_{1},\nu_{1}) = \int w_{\hat{\rho}_{0}\hat{\rho}_{1}}(X,Y|\mu_{1},\nu_{1},\mu_{2},\nu_{2})dY = \frac{1}{\sqrt{\pi (\nu_{1}^{2} + \mu_{1}^{2})}} \exp\left(-\frac{X^{2}}{\mu_{1}^{2} + \nu_{1}^{2}}\right),$$
(30)

$$w_{\hat{\bar{\rho}}_{0}\hat{\bar{\rho}}_{1}}(Y|\mu_{2},\nu_{2}) = \int w_{\hat{\bar{\rho}}_{0}\hat{\bar{\rho}}_{1}}(X,Y|\mu_{1},\nu_{1},\mu_{2},\nu_{2})dX = \frac{2Y^{2}}{\sqrt{\pi}\left(\nu_{2}^{2}+\mu_{2}^{2}\right)^{3/2}}\exp\left(-\frac{Y^{2}}{\mu_{2}^{2}+\nu_{2}^{2}}\right),$$
(31)

$$w_{\hat{\rho}_{1}\hat{\rho}_{0}}(Y|\mu_{2},\nu_{2}) = \int w_{\hat{\rho}_{1}\hat{\rho}_{0}}(X,Y|\mu_{1},\nu_{1},\mu_{2},\nu_{2})dX = \frac{1}{\sqrt{\pi(\nu_{2}^{2}+\mu_{2}^{2})}}\exp\left(-\frac{Y^{2}}{\mu_{2}^{2}+\nu_{2}^{2}}\right),$$
(32)

$$w_{\hat{\bar{\rho}}_{1}\hat{\bar{\rho}}_{0}}(X|\mu_{1},\nu_{1}) = \int w_{\hat{\bar{\rho}}_{1}\hat{\bar{\rho}}_{0}}(X,Y|\mu_{1},\nu_{1},\mu_{2},\nu_{2})dY = \frac{2X^{2}}{\sqrt{\pi}\left(\nu_{1}^{2}+\mu_{1}^{2}\right)^{3/2}}\exp\left(-\frac{X^{2}}{\mu_{1}^{2}+\nu_{1}^{2}}\right),$$
(33)

Comparing (20) and (22) with (28)-(33) we also see the difference of marginal distributions in entangled state (17) and in separable states (25), (26), (27). There exists a central mass tomography of two-mode oscillator [103, 104, 105]. These oscillator states are described by the probability distributions of one random variable (the center of mass position). We will study entangled and separable states of the oscillator in future

4 Conclusion

We conclude mentioning the main ideas to be used on the bases of the probability representation of quantum mechanics. In fact this approach provides the possibility to formulate new aspects of probability theory and group representation theory which play important role in quantum mechanics foundations. As we know the JordanSchwinger map [106, 107] provides the possibility to map the properties of matrices like the $N \times N$ matrices A, B, C with commutation relation [A, B] = C onto three operators \hat{A} , \hat{B} , \hat{C} acting in the Hilbert space of N-dimensional oscillator states, namely, $A \leftrightarrow \hat{A} = \sum_{i,k=1}^{N} A_{ik} \hat{a}_{i}^{\dagger} \hat{a}_{k}, \ B \leftrightarrow \hat{B} = \sum_{i,k=1}^{N} B_{ik} \hat{a}_{i}^{\dagger} \hat{a}_{k}, \ C \leftrightarrow \hat{C} = \sum_{i,k=1}^{N} C_{ik} \hat{a}_{i}^{\dagger} \hat{a}_{k}, \text{ where } \hat{A} \leftrightarrow \hat{A} = \sum_{i,k=1}^{N} C_{ik} \hat{a}_{i}^{\dagger} \hat{a}_{k}, \ \hat{A} \leftrightarrow \hat{A} = \sum_{i,k=1}^{N} C_{ik} \hat{a}_{i}^{\dagger} \hat{a}_{k}, \ \hat{A} \leftrightarrow \hat{A} = \sum_{i,k=1}^{N} C_{ik} \hat{a}_{i}^{\dagger} \hat{a}_{k}, \ \hat{A} \leftrightarrow \hat{A} = \sum_{i,k=1}^{N} C_{ik} \hat{a}_{i}^{\dagger} \hat{a}_{k}, \ \hat{A} \leftrightarrow \hat{A} = \sum_{i,k=1}^{N} C_{ik} \hat{a}_{i}^{\dagger} \hat{a}_{k}, \ \hat{A} \leftrightarrow \hat{A} = \sum_{i,k=1}^{N} C_{ik} \hat{a}_{i}^{\dagger} \hat{a}_{k}, \ \hat{A} \leftrightarrow \hat{A} = \sum_{i,k=1}^{N} C_{ik} \hat{a}_{i}^{\dagger} \hat{a}_{k}, \ \hat{A} \leftrightarrow \hat{A} = \sum_{i,k=1}^{N} C_{ik} \hat{a}_{i}^{\dagger} \hat{a}_{k}, \ \hat{A} \leftrightarrow \hat{A} = \sum_{i,k=1}^{N} C_{ik} \hat{a}_{i}^{\dagger} \hat{a}_{k}, \ \hat{A} \leftrightarrow \hat{A} \to \hat{A}$ $[\hat{a}_i, \hat{a}_k^{\dagger}] = \delta_{ik}$ and $\hat{a}_i, \hat{a}_k^{\dagger}$ are the annihilation and creation boson operators of the Nmode oscillator. The commutator $[\hat{A}, \hat{B}] = \hat{C}$ of the oscillator operators repeats the commutator properties of the matrices A, B, C. For example, the spin Pauli matrices commutation relations are reproduced by the two-dimensional oscillator bosonic commutation relations and the states of s-spin are mapped onto the excited energy states of the energy levels of the oscillator. We get the results that spin states can be invertibly mapped onto energy levels states of the two-mode oscillator. From this observation follows an idea that all the spin states can be mapped on the probability distributions describing the states of the two-mode oscillator (or the Hermitte polynomials [108, 109, 110]). This provides the possibility to map all the spin states (irreducible representations of rotation SU(2) group) onto transforms of the probability distributions which is new aspect of the group representation theory. If to apply this idea to N-dimensional oscillator properties we extend the result to other group representation theory (SU(N)) group and for other Lie groups). The s-spin states can be described as the multi-mode states associated with probability distributions of multidimensional Hermite polynomials. The approach will be considered in future publications.

References

 Kolmogorov, A. N., "Foundation of the Theory of Probability", Chelsea: New York, NY, USA, 1956. ISBN-10:0821826484.
 ISBN 12:0720221826485.

- Holevo, A. S., Probabilistic and Statistical Aspects of Quantum Theory, North-Holland Publishing Company: Amsterdam, The Netherlands, 1982
 ISBN: 9780444863331/0444863338.
- [3] Khrennikov, A., Probability and Randomness. Quantum versus Classical; World Scientific: Singapore, 2016. ISBN-10:1783267968, ISBN-13:978-1783267965.
 Int. J. Quantum Information, 14, p.1640009 (2016) https://doi.org/10.1142/S0219749916400098
- [4] Schrödinger, E., "Discussion of probability relations between separated systems", Mathematical Proceedings of the Cambridge Philosophical Society, **31 (4)**, pp. 555563 (1935) doi:10.1017/S0305004100013554. S2CID 121278681.
- Schrödinger, E., "Probability relations between separated systems", Mathematical Proceedings of the Cambridge Philosophical Society 32 (3), pp. 446452 (1936) doi:10.1017/S0305004100019137. S2CID 122822435.
- [6] Nielsen, M. A., and Chuang, I. L., "Quantum Computation and Quantum Information, 10th Anniversary ed.", University Press: Cambridge, UK, 2010; p. 704
 ISBN 978-1-107-00217-3.
- Bell, J. S., "On the Einstein–Podolsky–Rosen Paradox" Physics, 1 (3), pp. 195200 (1964)
 doi:10.1103/PhysicsPhysiqueFizika.1.195.
- [8] Khrennikov, A., "Against Identification of Contextuality with Violation of the Bell Inequalities: Lessons from Theory of Randomness", J. Russ. Laser Res., 43, pp. 4859 (2022). https://doi.org/10.1007/s10946-022-10022-9 arXiv:2112.03910 [quant-ph] (2021)
- [9] Andreev, V. A., Man'ko, V. I., Man'ko, O. V., and Sckhukin, E. V., "Tomography of Spin States, the Entanglement Criterion, and Bell's Inequalities", *Theor. Math. Phys.*, **146**, N 1, pp. 140151 (2006)

- [10] Andreev, V. A., "Generalized Bell inequality and a method for its verification", Theor. Math. Phys., 152, pp. 12861298 (2007) https://doi.org/10.1007/s11232-007-0113-1
- [11] Andreev, V. A., "The correlation Bell inequalities", Theor. Math. Phys., 158, pp. 196209 (2009)
 https://doi.org/10.1007/s11232-009-0016-4
- [12] Chernega, V. N., Man'ko O. V., and Man'ko V. I., "Entangled Qubit States and Linear Entropy in the Probability Representation of Quantum Mechanics", *Entropy*, 24(4), p. 527 (2022) https://doi.org/10.3390/e24040527
- Mancini, S., Man'ko, V. I., and Tombesi, P., "Symplectic Tomography as Classical Approach to Quantum Systems", *Phys. Lett. A*, **213**, pp. 1-6 (1996) https://doi.org/10.1016/0375-9601(96)00107-7
- [14] Mancini, S., Man'ko, V.I., and Tombesi, P., "Classical-like description of quantum dynamics by means of symplectic tomography", Found. Phys., 27, pp. 801824 (1997)
 https://doi.org/10.1007/BF02550342
- [15] Mancini, S., Man'ko, V. I., and Tombesi, P., "Wigner function and probability distribution for shifted and squeezed quadratures", J. Opt. B: Quantum Semiclass. Opt., 7, p. 615 (1995)
 DOI 10.1088/1355-5111/7/4/016
- [16] Man'ko, O. V., and Man'ko, V. I., "Quantum states in probability representation and tomography, J. Rus. Laser Res., 18, pp. 407444 (1997) https://doi.org/10.1007/BF02559668
- [17] Caponigro, M., Mancini, S., and Man'ko, V. I., "A probabilistic approach to quantum mechanics based on "tomograms", Fortschritte der Physik, 54, p. 602 (2006) https://doi.org/10.1002/prop.200610304

- [18] Dodonov, V. V., and Manko, V. I., "Positive Distribution Description for Spin States", Phys. Lett. A, 229, pp. 335339 (1997)
 https://doi.org/10.1016/S0375-9601(97)00199-0
- [19] Manko, V. I., and Manko, O. V., "Spin State Tomography", J. Exp. Theor. Phys.,
 85, pp. 430434 (1997)
 https://doi.org/10.1134/1.558326
- Manko, O. V., "Tomography of Spin States and Classical Formulation of Quantum Mechanics", in: Gruber, B., Ramek, M. (eds), Symmetries in Science X, 1998, Springer, Boston, MA, 1998, p. 207 doi:10.1007/978-1-4899-1537-5-13
- [21] Terra-Cunha, M. O., Manko, V.I., and Scully, M. O., "Quasiprobability and Probability Distributions for Spin-1/2 States", Found. Phys. Lett., 14, pp. 103117 (2001) https://doi.org/10.1023/A:1012373419313
- [22] Amiet, J. P., and Weigert, S., "Coherent states and the reconstruction of pure spin states", J. Opt. B: Quantum Semiclass. Opt., 1, pp. L5 - L8 (1999) DOI 10.1088/1464-4266/1/5/101
- [23] D'Ariano, G. M., Maccone, L., and Paini, M., "Spin tomography", J. Opt. B: Quantum Semiclass. Opt., 5, p. 77 (2003)
 DOI 10.1088/1464-4266/5/1/311
- [24] Schrödinger, E., "Quantisierung als Eigenwertproblem (Erste Mitteilung)", Ann.
 Phys., 384, pp. 361376 (1926)
 https://doi.org/10.1002/andp.19263840404
- [25] Landau, L., "Das Dämpfungsproblem in der Wellenmechanik", Z. Phys., 45, pp.
 430441 (1927)
 DOI 10.1007/BF01343064
- [26] Neumann, J. von, "Wahrscheinlichkeitstheoretischer Aufbau der Quantenmechanik", Gtt. Nach., 1, pp. 245272 (1927)

- [27] Manko, V. I., Marmo, G., Sudarshan, E. C. G., and Zaccaria, F., "Interference and entanglement: an intrinsic approach", *J. Phys. A: Math. Gen.*, **35**, N 33, p. 7137 (2002)
 DOI 10.1088/0305-4470/35/33/311
- [28] Sudarshan, E. C. G., "Search for Purity and Entanglement", J. Russ. Laser Res.,
 24, pp. 195203 (2003)
 https://doi.org/10.1023/A:1024090024283
- [29] Man'ko, V. I., Marmo, G., Sudarshan, E. C. G., and Zaccaria, F., "Entanglement Structure of the Adjoint Representation of the Unitary Group and Tomography of Quantum States", J. Russ. Laser Res., 24, pp. 507543 (2003) https://doi.org/10.1023/B:JORR.0000004166.55179.aa
- [30] Man'ko, O. V., Man'ko, V. I., and Marmo, G., "Alternative commutation relations, star-produst and tomography", J. Phys. A, 35, 699-719 (2002)
 DOI 10.1088/0305-4470/35/3/315
- [31] Man'ko, O. V., Man'ko, V. I., and Marmo, G., "Star-product of generalized Weyl symbols on SU(2) group, deformations and tomographic probability distributions", *Phys. Scripta*, **62**, 446-452 (2000)
 DOI: 10.1238/Physica.Regular.062a00446
- [32] Man'ko, O. V., Man'ko, V. I.,and Marmo, G., "Tomographic map within the framework of star-product quantization", in: *Proceedings of Second International Sympo*sium on Quantum Theory and Symmetries, Krakow, Poland, 2001, (E. Kapuschik, A. Morzela, eds), World Scientific, 126-133 (2001) https://doi.org/10.1142/9789812777850-0011
- [33] Man'ko, O. V., Man'ko, V. I., Marmo, G.,and Vitale, P., "Star product, duality and double Lie algebras", Phys. Lett. A, 360, 522-532 (2007) https://doi.org/10.1016/j.physleta.2006.08.057
- [34] Kuznetsov, S. V., Kyusev, A. V., and Man'ko O. V., "Tomographic and statistical properties of superposition states for two-mode systems", *Proc. SPIE*, **5402**, *In*-

https://doi.org/10.1117/12.562263 arXiv:1903.10784 [quant-ph]

[35] Kuznetsov S. V., Kyusev A.V., Man'ko O.V., and Chernega N. V., "Tomographic and statistical properties of superposition states for two-mode systems", Bulletin of the Russian Academy of Science, Physics, 58, N 9, p.p. 1239-1243 (2004), Allerton Press

https://www.elibrary.ru/item.asp?id=17641248 (in Russian)

- [36] Kuznetsov, S. V., Man'ko, O. V., and Tcherniega, N. V., "Photon distribution function, tomograms and entanglement in Stimulated Raman Scattering", J. Opt. B: Quantum Semiclass. Opt, 5, p. 5503 (2003)
 DOI 10.1088/1464-4266/5/4/357
- [37] Kuznetsov, S. V., and Man'ko, O. V. "Entanglement and photon number probability distribution function for Stimulated Raman Scattering", Proc. SPIE, 5402, pp. 302-313 (2004)
 doi: 10.1117/12.562224
 Kuznetsov, S. V., Kyusev, A. V., and Man'ko, O.V., arXiv:1903.10784 [quant-ph]
- [38] Man'ko, O.V., and Tcherniega, N. V., "Tomographic description of Stimulated Brillouin Scattering", J. Russ. Laser Res., 22, N 3, pp. 201-218 (2001) https://doi.org/10.1023/A:1011304404336
- [39] Kuznetsov, S. V., Kusev, A. V., Man'ko, O. V., and Tcherniega, N. V., "Entanglement in the process of Stimulated Brillouin Scattering", Acta Physica Hungarica B: Series Quantum Electronics, 20/1, pp. 11-14 (2004) https://doi.org/10.1556/APH.20.2004.1-2.4
- [40] Man'ko, O. V., "Photon distribution function for stimulated Raman scattering", Proc. SPIE, 4089, Raman Scattering, (Moscow, 18 February 2000) (2000) https://doi.org/10.1117/12.378116
- [41] Man'ko, O. V., and Tcherniega N. V., "Photon-number tomography and symplectic approach to stimulated Raman scattering". *Proc. SPIE*, **6256W**, *ICONO*

and Atom Optics; Engineering of Quantum Information, pp. 62560W-1-62560W-12 (2006) https://doi.org/10.1117/12.682581

- [42] Giri, S. K., Sen, B., Pathak, A., and Jana, P. C., "Higher-order two-mode and multimode entanglement in Raman processes", *Phys. Rev. A*, 93, p. 012340 (2016) https://doi.org/10.1103/PhysRevA.93.012340
- [43] Pathak, A., Krepelka, J., and Perina, J., "Nonclassicality in Raman scattering: Quantum entanglement, squeezing of vacuum fluctuations, sub-shot noise and joint photonphonon number and integrated-intensity distributions", *Phys. Lett. A*, 377, N 38, pp. 2692-2701 (2013) https://doi.org/10.1016/j.physleta.2013.07.046
- [44] Rohith, M., and Sudheesh, C., "Visualizing revivals and fractional revivals in a Kerr medium using an optical tomogram", *Phys. Rev. A*, **92**, p. 053828 (2015) https://doi.org/10.1103/PhysRevA.92.053828
- [45] Rohith, M., and Sudheesh, C., "Signatures of entanglement in an optical tomogram," JOSA B, 33, pp. 126-133 (2016) https://doi.org/10.1364/JOSAB.33.000126
- [46] Man'ko, O. V., Man'ko, V. I., Marmo, G., Shaji, A., Sudarshan, E. C. G., and Zaccaria, F., "Partial positive scaling transform and a separability criterion", *Phys. Lett. A*, **339**, pp. 194-206 (2005) https://doi.org/10.1016/j.physleta.2005.02.070
- [47] Man'ko, O. V., "Heating map in classical and quantum mechanics", J. Russ. Laser Res., 26, pp. 109115 (2005)
 https://doi.org/10.1007/s10946-005-0010-8
- [48] Facchi, P., Ligabó, M., and Solimini, S., "Tomography: mathematical aspects and applications," *Phys. Scr.*, **90**, N 7, p. 074007 (2015)
 DOI 10.1088/0031-8949/90/7/074007

- [49] Claeyes., P. W., and Polkovnikov, A., "Quantum eigenstates from classical Gibbs distributions" SciPost Phys., 10, p. 014 (42pp.) (2021)
 doi: 10.21468/SciPostPhys.10.1.014
- [50] Khrennikov, A., Physica A: Statistical Mechanics and its Applications, 393, 207 (2014)
- [51] Khrennikov., A., *Social Laser*, Jenny Stanford Publ. 280 pp.(2020)
 ISBN 978-981-4800-83-9
- [52] Khrennikov, A., and Alodjants, A., "Classical (Local and Contextual) Probability Model for BohmBell Type Experiments: No-Signaling as Independence of Random Variables", *Entropy*, **21(2)**, p. 157 (2019) https://doi.org/10.3390/e21020157
- [53] Khrennikov., A., "Quantum postulate vs. quantum nonlocality: on the role of the Planck constant in Bells argument", Found. Phys., 51, p. 16 (12pp) (2021) https://doi.org/10.1007/s10701-021-00430-3
- [54] Ciaglia, F. M., Di Cosmo, F., Ibort, A., and Marmo, G., "Evolution of Classical and Quantum States in the Groupoid Picture of Quantum Mechanics", *Entropy*, 22(11), p. 1292 (2020) https://doi.org/10.3390/e22111292
- [55] Przhiyalkovskiy, Y. V., "Continuous Measurements in Probability Representation of Quantum Mechanics", Proc. Steklov Inst. Math., 313, pp.193202 (2021) https://doi.org/10.1134/S0081543821020188 arXiv:2101.07568v1 [quant-ph]
- [56] Facchi, P., and Ligabó, M., "Classical and quantum aspects of tomography", in:
 M. Asorey, J. Clemente-Gallardo, E. Martnez, J. F. Cariena (Ed.) Proceedings of XVIII International Fall Workshop on Geometry and Physics, Benasque, Spain, 2009, AIP Conference Proceedings, 1260 p. 3 (2010) https://doi.org/10.1063/1.3479322

- [57] Elze, H.-T., Gambarotta, G., and Vallone, F., "General Linear DynamicsQuantum, Classical or Hybrid", J. Phys. Conf. Ser., 306, p. 012010. (2011)
 DOI 10.1088/1742-6596/306/1/012010
- [58] Stornaiolo, C., "Emergent classical universes from initial quantum states in a tomographical description", Int. J. Geom. Meth. in Modern Phys., .17(11), p. 2050167 (2020)
 DOI: 10.1142/S0219887820501674
- [59] Stornaiolo, C., "Tomographic Represention of Quantum and Classical Cosmology",
 in: C. Moreno Gonzlez, J. Madriz Aguilar, L. Reyes Barrera (Ed.), Accelerated Cosmic Expansion. Proceedings of the Fourth International Meeting on Gravitation and Cosmology, Astrophysics and Space Science Proceedings, 38 pp. 1155-1157 (2014) Springer doi: 10.1142/9789814623995-0107
- [60] BerraMontiel, J.; and Molgado, A. "Tomography in loop quantum cosmology", Eur. Phys. J. Plus, 137, p. 283 (2022)
 DOI: 10.1140/epjp/s13360-022-02504-1
- [61] de Gosson, M. A., "Quantum Harmonic Analysis of the Density Matrix", Quanta,
 7, p. 74 (2018) https://doi.org/10.12743/quanta.v7i1.74
- [62] Gosson, M. A., and de Gosson, C., "Symplectic Polar Duality, Quantum Blobs, and Generalized Gaussians" Symmetry, 14(9), p. 1890 (2022) https://doi.org/10.3390/sym14091890
- [63] Gosson, M. A., "Symplectic Radon Transform and the Metaplectic Representation", Entropy, 24(6), p. 761 (2022) https://doi.org/10.3390/e24060761
- [64] Foukzon, J., Potapov, A. A., Menkova, E., and Podosenov, S. A. "A New Quantum-Mechanical Formalism Based on the Probability Representation of Quantum States," (2016) viXra:1612.0298.

- [65] Chernega, V. N., Belolipetskiy, S. N., Man'ko, O. V., and Man'ko, V. I., "Probability representation of quantum mechanics and star product quantization", J. of Physics: Conf. Ser., 1348, Irreversible Processes in Nature and Technics 2931 January 2019, Moscow, Russian Federation, p. 012101 (2019) DOI 10.1088/1742-6596/1348/1/012101
- [66] Bazrafkan, M. R., and Nahvifard, E., "Stationary perturbation theory in the probability representation of quantum mechanics", J. Russ. Laser Res., 30, pp. 392403 (2009)
 https://doi.org/10.1007/s10946-009-9079-9
- [67] Filinov, V. S., Schubert, G., Levashov, P., Bonitz, M., Fehske, H., Fortov, V. E., and Filinov, A. V., "Center-of-mass tomographic approach to quantum dynamics", *Phys. Lett. A*, **372**, p. 5064 (2008) https://doi.org/10.1016/j.physleta.2008.05.053
- [68] Chernega, V. N., Man'ko, O. V., and Man'ko, V. I., "Triangle Geometry of the Qubit State in the Probability Representation Expressed in Terms of the Triada of Malevichs Squares", J. Rus. Laser Res., 38, N 2, pp. 141149 (2017) https://doi.org/10.1007/s10946-017-9628-6
- [69] Chernega, V. N., Man'ko, O. V., and Man'ko, V. I., "Probability Representation of Quantum Observables and Quantum States", J. Rus. Laser Res., 38, N 4, pp. 324-333 (2017)
 https://doi.org/10.1007/s10946-017-9648-2
- [70] Chernega, V. N., Man'ko, O. V., and Man'ko, V. I., "Triangle Geometry for Qutrit States in the Probability Representation", J. Rus. Laser Res., 38, N 5, pp. 416425 (2017) https://doi.org/10.1007/s10946-017-9662-4
- [71] Chernega, V. N., Man'ko, O. V., and Man'ko, V. I., "God Plays Coins or Superposition Principle for Classical Probabilities in Quantum Suprematism Representation of Qubit States", J. Rus. Laser Res., 39, N 2, pp. 128139 (2017)

- [72] Chernega, V. N., Man'ko, O. V., and Man'ko, V. I., "Quantum suprematism picture of Triada of Malevich's squares for spin states and the parametric oscillator evolution in the probability representation of quantum mechanics", J. Phys.: Conf. Ser., 1071, p. 012008 (2018)
 DOI 10.1088/1742-6596/1071/1/012008
- [73] Chernega, V. N., Man'ko, O. V., and Man'ko, V. I., "Correlations in a system of classical-like coins simulating spin-1/2 states in the probability representation of quantum mechanics", *Eur. Phys. J. D*, 73, p. 10 (2019). https://doi.org/10.1140/epjd/e2018-90487-9
- [74] Chernega, V. N., Man'ko, O. V., and Man'ko, V. I., "Probability Representation of Quantum States as a Renaissance of Hidden Variables God Plays Coins", J. Rus. Laser Res., 40, N 2, pp. 107120 (2019) https://doi.org/10.1007/s10946-019-09778-4
- [75] Chernega, V. N., Man'ko, O. V., and Man'ko, V. I., "Probability Representation of Photon States and Tomography", J. Rus. Laser Res., 40, N 6, pp. 503514 (2019) https://doi.org/10.1007/s10946-019-09831-2
- [76] Miroshnichenko, G. P., "CQED Quantum Tomography of a Microwave Range", arXiv:1510.03155 [quant-ph] (2015) https://doi.org/10.48550/arXiv.1510.03155
- [77] Koczor, B., Zeier, R., and Glaser, S. J., "Continuous Phase-Space Representations for Finite-Dimensional Quantum States and their Tomography", *Phys. Rev. A*, 101, p. 022318 (2020) https://doi.org/10.1103/PhysRevA.101.022318 arXiv:1711.07994v3 [quant-ph] (2018)
- [78] Toninelli. E., Ndagano, B., Valles, A., and Forbes, A., "Concepts in quantum state tomography and classical implementation with intense light: a tutorial" Advances in Optics and Photonics, 11(1), p. 67–134 (2019) https://doi.org/10.1364/AOP.11.000067

- [79] Almarashi, A. M., Abd-Elmougod, G. A., and Raqab, M. Z., "Quantum Extropy and Statistical Properties of the Radiation Field for Photonic Binomial and Even Binomial Distributions", J. Russ. Laser Res., 41, pp. 334-343 (2020) https://doi.org/10.1007/s10946-020-09883-9
- [80] Leon, R. C. C., Yang, C. H., Hwang, J. C. C., Lemyre, J. C., Tanttu, T., Huang, W., Huang, J. H., Hudson, F. E., Itoh, K. M., Laucht, A., Pioro-Ladriére, M., Saraiva, A., and Dzurak, A. S., "Bell-state tomography in a silicon many-electron artificial molecule", *Nat. Commun.*, **12**, p. 3228 (2021) https://doi.org/10.1038/s41467-021-23437-w arXiv:2008.03968v1 (2020) https://doi.org/10.48550/arXiv.2008.03968
- [81] Kiktenko, E. O., Fedorov, A. K., Man'ko, O. V. and Man'ko, V. I., "Multilevel superconducting circuits as two-qubit systems: Operations, state preparation, and entropic inequalities", *Phys. Rev. A*, **91**, p. 042312 (2015)
 DOI:https://doi.org/10.1103/PhysRevA.91.042312
- [82] Man'ko, O.V., "Quantum tomography of current and voltage states in nanoelectriccircuits", AIP Conference Proc., 1424, p. 221-233 (2012) https://doi.org/10.1063/1.3688974
- [83] Man'ko, O. V., "Symplectic tomography of nonlinear coherent states of a trapped ion", Phys. Lett. A, 228, pp. 29-35 (1997) https://doi.org/10.1016/S0375-9601(97)00091-1
- [84] Man'ko, O. V., "Symplectic tomography of Schödinger cat states of a trapped ion", in Proc. of "New developments pon Fundamental Problems in quantum physics, Oviedo, Spain, 1996, Foundamental Theories of Physics, Kluwer Publ. M. Ferrero, A. van der Merwe Eds.p.225-229 (1997) DOI: 10.1007/978-94-011-5886-2-29
- [85] Man'ko, O. V., "Optical tomography and measuring quantum states of an ion in a Paul and in a Penning trap", Proc. SPIE, 3736, ICONO '98: Quantum Optics, Interference Phenomena in Atomic Systems, and High-Precision Measurements,

(25 February 1999) (1999) https://doi.org/10.1117/12.340153

- [86] Chernega, V. N., and Man'ko, O. V., "A Damped Oscillator with a δ-Kicked Frequency in the Probability Representation of Quantum Mechanics", J. Russ. Laser Res., 39, N 4, pp. 411-419 (2018) https://doi.org/10.1007/s10946-018-9735-z
- [87] Klimov, A. B., Man'ko, O. V., Man'ko, V. I., Smirnov, Yu. F., and Tolstoy, V. N., "Tomographic representation of spin and quark states", J. Phys. A: Math. Gen., 35, p. 6101 (2002) DOI 10.1088/0305-4470/35/29/312
- [88] Man'ko, O. V., and Man'ko, V. I., "Probability-Representation Entropy for Spin-State Tomogram", J. Russ. Laser Res., 25, pp. 115-122 (2004) https://doi.org/10.1023/B:JORR.0000018981.06864.12
- [89] Man'ko, O. V., and Man'ko, V. I., "Wave Function in Classical Statistical Mechanics", J. Russ. Laser Res., 22, pp. 149-167 (2001) https://doi.org/10.1023/A:1011360006073
- [90] Man'ko, O. V., and Man'ko, V. I., "Classical Mechanics Is not the ħ → 0 Limit of Quantum Mechanics", J. Russ. Laser Res., 25, p. 477-492 (2004) https://doi.org/10.1023/B:JORR.0000043735.34372.8f
- [91] Man'ko, O. V., Man'ko, V. I., Pilyavets, O.V., "Probability representation of classical states", J. Rus. Laser Res., v. 26, N 6, p. 429-444 (2005) https://doi.org/10.1007/s10946-005-0045-x
- [92] Asorey, M., Ibort, A., Marmo G., and Ventriglia, F., "Quantum Tomography twenty years later", *Phys. Scr.*, **90**, p. 074031 (2015)
 DOI: 10.1088/0031-8949/90/7/074031
- [93] Man'ko, V. I., Marmo, G., Simoni, A., Stern, A., and Ventriglia, F., "An Introduction to the tomographic picture of quantum mechanics", *Phys. Lett. A*, 343, p. 251 (2005)

- [94] Ullinger, F., Zimmermann, M., and Schleich, W. P., "The logarithmic phase singularity in the inverted harmonic oscillator", AVS Quantum Sci., 4(2), p. 024402 (2022)
 DOI: 10.1116/5.0074429
- [95] Subramanyan, V., Hegde, S. S., Vishveshwara, S., and Bradlyn, B., "Physics of the Inverted Harmonic Oscillator: From the lowest Landau level to event horizons", Annals Phys., 435, p. 168470 (2021) DOI: 10.1016/j.aop.2021.168470
- [96] Man'ko, V. I., and Man'ko, O.V., "Inverted Oscillator Quantum States in the Probability Representation", Entropy, 25(2), p.217 (2023) https://doi.org/10.3390/e25020217
- [97] Nosal, I. A., and Teretenkov, A. E., "Exact Dynamics of Moments and Correlation Functions for GKSL Fermionic Equations of Poisson Type", Math. Notes, 108, p. 911-925 (2020) https://doi.org/10.1134/S0001434620110358
- [98] Teretenkov, A. E., "Irreversible quantum evolution with quadratic generator: Review", Infin. Dimens. Anal. Quantum Probab. Relat. Top., 22, N 4, p. 1930001, 27 pp. (2019)
 https://doi.org/10.1142/S0219025719300019
- [99] Mancini, S., Man'ko, O. V., Man'ko, V. I., and Tombesi, P., "The Pauli equation for probability distributions", J. Phys. A: Math. Gen., 34, p. 3461. (2001)
 DOI 10.1088/0305-4470/34/16/314
- [100] Man'ko, O. V., and Man'ko, V. I., "Probability Representation of Quantum States", Entropy, 23(5), p. 549 (2021) https://doi.org/10.3390/e23050549
- [101] Thapliyal, K., Banerjee, S., and Pathak, A., "Tomograms for open quantum systems: In(finite) dimensional optical and spin systems", Annals of Physics, 366, pp. 148-167 (2016)
 https://doi.org/10.1016/j.aop.2016.01.010

- [102] Arkhipov, A. S., and Lozovik, Yu. E., "New method of quantum dynamics simulation based on the quantum tomography", Phys. Lett. A, **319**, pp. 217-224 (2003) doi:10.1016/j.physleta.2003.09.084
- [103] Arkhipov, A. S., Lozovik, Yu. E., and Man'ko, V. I., "Center of mass tomography for reconstructing quantum states of multipartite systems", *Phys. Lett.A*, **328**, N
 6, pp. 419-431 (2004)
 doi: 10.1016/j.physleta.2004.06.039
- [104] Manko, O.V., and Manko, V.I., "Fidelity in the center-of-mass tomography", J.
 Russ. Laser Res., **31**, pp. 608613 (2010)
 https://doi.org/10.1007/s10946-010-9181-z
- [105] Dudinets, I.V., and Manko, V.I., "Center-of-Mass Tomography and Wigner Function for Multimode Photon States", Int. J. Theor. Phys., 57, pp. 16311644 (2018) https://doi.org/10.1007/s10773-018-3690-x
- [106] Jordan, P., "Zusammenhang der symmetrischen und linearen Gruppen und das Mehrkorperproblem", Z. Phys., 94, pp. 531-535 (1935)
 doi: 10.1007/BF01330618
- [107] Schwinger, J.: "On angular momentum", in Quantum Theory of Angular Momentum, eds. L.C. Biedenharn and H. Van Dam, Academic, 1965.
- [108] Man'ko, M. A., and Man'ko, V. I., "Hermite Polynomial Representation of Qubit States in Quantum Suprematism Picture", In: Marmo, G., Martn de Diego, D., Muoz Lecanda, M. (eds) *Classical and Quantum Physics*, Springer Proceedings in Physics, vol 229. Springer, Cham. (2019) https://doi.org/10.1007/978-3-030-24748-5-16
- [109] Lemeshevskiy, D. B., Manko, V. I. "Hermite polynomial representation of the spin states", J. Russ. Laser Res.. 34, N 2, pp. 175-184 (2013) https://doi.org/10.1007/s10946-013-9340-0
- [110] Man'ko, M. A., "Unitary transforms of probability distributions describing quantum states of two-level systems", J. Russ. Laser Res., 43, pp. 645652 (2022)