MULTIPLICITY OF NONTRIVIAL ZEROS OF PRIMITIVE L-FUNCTIONS VIA HIGHER-LEVEL CORRELATIONS

FELIPE GONÇALVES, DAVID DE LAAT, AND NANDO LEIJENHORST

Abstract. We give universal bounds on the fraction of nontrivial zeros having given multiplicity for L-functions attached to a cuspidal automorphic representation of GL_m/\mathbb{Q} . For this, we apply the higher-level correlation asymptotic of Hejhal and Rudnick & Sarnak in conjunction with semidefinite programming bounds.

CONTENTS

1. INTRODUCTION

Let $m \geq 1$ and let π be an irreducible cuspidal automorphic representation of GL_m/\mathbb{Q} . Let $L(s,\pi)$ be its attached L-function (see [\[15\]](#page-14-0) for theoretical background). Such functions generalize the classical Dirichlet L-functions (the case $m = 1$). Let $\rho_j = 1/2 + i\gamma_j$, with $j \in \mathbb{Z}$, be an enumeration of the nontrivial zeros of $L(s, \pi)$ repeated according to multiplicity. The Generalized Riemann Hypothesis (GRH) for $L(s, \pi)$ states that all its nontrivial zeros are aligned in the line Re $s = 1/2$; that is, $\gamma_i \in \mathbb{R}$ for all j. Assuming GRH, we can enumerate the ordinates of the zeros in a nondecreasing fashion

$$
\ldots \leq \gamma_{-2} \leq \gamma_{-1} < 0 \leq \gamma_1 \leq \gamma_2 \leq \ldots
$$

and we have

$$
N(T) := \sum_{\substack{j \ge 1 \\ \gamma_j \le T}} 1 \sim \frac{m}{2\pi} T \log T.
$$

Hejhal [\[14\]](#page-14-1) and later on Rudnick $&$ Sarnak [\[20,](#page-14-2) [21\]](#page-14-3) investigated the *n*-level correlation distribution of the zeros of $L(s, \pi)$ guided by the relationship with Gaussian ensemble models in random matrix theory, pointed out by the breakthrough work

Date: March 3, 2023.

²⁰²⁰ Mathematics Subject Classification. 11M26, 90C22.

of Montgomery [\[17\]](#page-14-4). This area has a long history and we recommend [\[21\]](#page-14-3), and the references therein, for more information.

The main result of Rudnick & Sarnak reads as follows: Assuming GRH for $L(s, \pi)$ and that either $m = 1, 2, 3$ or condition (1.7) in [\[21\]](#page-14-3) is satisfied, we have

$$
\frac{1}{N(T)} \sum_{\substack{j_1,\ldots,j_n \geq 1 \text{ distinct} \\ \gamma_{j_1},\ldots,\gamma_{j_n} \leq T}} f(\frac{L}{2\pi}\gamma_{j_1},\ldots,\frac{L}{2\pi}\gamma_{j_n})
$$
\n
$$
\to \int_{\mathbb{R}^n} f(x)W_n(x)\delta\left(\frac{x_1 + \ldots + x_n}{n}\right)dx_1\cdots dx_n
$$
\n(1)

as $T \to \infty$, where $L = m \log T$,

$$
W_n(x) = \det \left[\frac{\sin(\pi(x_i - x_j))}{\pi(x_i - x_j)} \right]_{i,j=1,...,n}
$$

is Dyson's limiting n-level correlation density for the Gaussian Unitary Ensemble (GUE) model, and f is any *admissible* test function satisfying

- $f \in C^{\infty}_{\sim}(\mathbb{R}^n)$, f is symmetric, and $f(x+t(1,\ldots,1)) = f(x)$ for $t \in \mathbb{R}$;
- $\overline{\text{supp}}(f) \subseteq \{x \in \mathbb{R}^n : |x_1| + \ldots + |x_n| < 2/m\}$ (in the distributional sense);
- $f(x) \to 0$ rapidly as $|x| \to \infty$ in the hyperplane $\sum_j x_j = 0$.

Let m_ρ denote the multiplicity of ρ , and define

$$
N^*(T) := \sum_{\substack{j \ge 1 \\ \gamma_j \le T}} m_{\rho_j}.
$$

Montgomery's pair correlation approach has been used to obtain bounds of the form

$$
N^*(T) \le (c + o(1))N(T),
$$
\n(2)

where for $m = 1$ the current smallest known value for c assuming the Riemann hypothesis is 1.3208 [\[4,](#page-13-1) [5,](#page-13-2) [17,](#page-14-4) [18\]](#page-14-5). Let $Z_n(T)$ count the number of nontrivial zeros of $L(s, \pi)$ with multiplicity at most $n - 1$ up to height T:

$$
Z_n(T) = \sum_{\substack{j \ge 1 \\ \gamma_j \le T, \, m_{\rho_j} \le n-1}} 1.
$$

Then $Z_2(T) \geq 2N(T) - N^*(T)$, so [\(2\)](#page-1-0) can for instance be used to give a lower bound on the fraction of simple zeros, although the current best known bound is obtained using different techniques [\[2,](#page-13-3) [4,](#page-13-1) [5,](#page-13-2) [11\]](#page-13-4).

The goal of this paper is to use the *n*-level correlations by Hejhal and Rudnick $\&$ Sarnak to obtain bounds analogous to [\(2\)](#page-1-0), and use this to obtain universal bounds on the fraction of nontrivial zeros of $L(s, \pi)$ having multiplicity at most $n - 1$.

Theorem 1. Let $L(s, \pi)$ be an L-function attached to an irreducible cuspidal automorphic representation π of GL_m/\mathbb{Q} . Assuming GRH for $L(s,\pi)$ we have

$$
\liminf_{T \to \infty} \frac{Z_n(T)}{N(T)} \ge \begin{cases} 0.9614 & \text{if } n = 3 \text{ and } m = 1, \\ 0.2997 & \text{if } n = 3 \text{ and } m = 2, \\ 0.9787 & \text{if } n = 4 \text{ and } m = 1, \end{cases}
$$

where the results for $(n, m) = (3, 2), (4, 1)$ hold under the additional assumption that certain series of rational functions are summed correctly using Maple; see Section [6.](#page-10-0)

As far as we know, the bounds in Theorem [1](#page-1-1) are all new. The case $m = 2$ is of special interest, since for this there are no previous effective bounds in the literature. For $(n, m) = (2, 2)$ we show in Section [3](#page-5-0) our method does not produce a positive lower bound, but as shown above for $(n, m) = (3, 2)$ it does.

Montgomery's approach to finding a good value for c in (2) leads to an optimization problem that is similar to the one-dimensional version of the Cohn-Elkies bound [\[7\]](#page-13-5) for the sphere packing problem. Interestingly, however, the higher-order correlation approach in this paper is very different from the higher-order correlation approach for the sphere packing problem recently introduced in [\[6\]](#page-13-6).

In Section [2](#page-2-0) we set up the optimization problem to compute the above bounds, and we show how the optimal solution connects to reproducing kernels, inspired by the approach from [\[3\]](#page-13-7). In Section [3](#page-5-0) we give a simpler analytical derivation of the known optimal solution for the case $n = 2$.

For larger values of n we use a computational approach to obtain rigorous bounds, whereas in Section [4](#page-7-0) we use the symmetry and the rank-1 structure to find the bounds by solving linear systems. We then give two approaches for finding good solutions to the optimization problem. In Section [5](#page-9-0) we use a parametrization using polynomials to find a good value for the case $(n, m) = (3, 1)$ and explain why this approach becomes problematic for $n > 3$. In Section [6](#page-10-0) we give an alternative parametrization based on lattice shifts and use this to compute bounds for (n, m) = $(3, 2), (4, 1).$

2. Derivation of the optimization problem

Let

$$
M_n(T) = \sum_{\substack{j_1,\ldots,j_n \geq 1 \text{ distinct} \\ \gamma_{j_1},\ldots,\gamma_{j_n} \leq T}} 1 = \sum_{\substack{j \geq 1 \\ \gamma_j \leq T}} (m_{\rho_j} - 1) \cdots (m_{\rho_j} - n + 1).
$$

Then $M_1(T) = N(T)$ and $M_2(T) = N^*(T) - N(T)$. We apply [\(1\)](#page-1-2) to obtain

$$
\limsup_{T \to \infty} \frac{M_n(T)}{N(T)} \le c_{n,m}
$$

where

$$
c_{n,m} := \inf \frac{1}{f(1,\ldots,1)} \int_{\mathbb{R}^n} f(x)W_n(x)\delta\left(\frac{x_1 + \ldots + x_n}{n}\right) dx
$$

and where the infimum is taken over nonnegative admissible test functions f . We have

$$
\int_{\mathbb{R}^n} f(x)W_n(x)\delta\left(\frac{x_1 + \dots + x_n}{n}\right) dx
$$
\n
$$
= \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}} f(x + (x_n, \dots, x_n, 0))W_n(x + (x_n, \dots, x_n, 0))
$$
\n
$$
\delta\left(\frac{x_1 + \dots + x_{n-1}}{n} + x_n\right) dx_n dx_1 \cdots dx_{n-1}
$$
\n
$$
= \int_{\mathbb{R}^{n-1}} f(x_1, \dots, x_{n-1}, 0)W_n(x_1, \dots, x_{n-1}, 0) dx_1 \cdots dx_{n-1}.
$$

With $g(x_1, \ldots, x_{n-1}) = f(x_1, \ldots, x_{n-1}, 0)$, we see that $\overline{\text{supp}}(\widehat{g})$ is now contained in the interior of $\frac{2}{m}H_{n-1}$, where

$$
H_{n-1} = \{x \in \mathbb{R}^{n-1} : |x_1| + \ldots + |x_{n-1}| + |x_1 + \ldots + x_{n-1}| \leq 1\}.
$$

Note that $H_{n-1} \subseteq [-1/2, 1/2]^{n-1}$. By an approximation argument $c_{n,m}$ is given by the infimum of

$$
\nu_n(g) := \int_{\mathbb{R}^{n-1}} g(x) W_n(x, 0) \, dx
$$

over all $g \in L^1(\mathbb{R}^{n-1})$ satisfying

(i)
$$
\overline{\text{supp}}(\widehat{g}) \subseteq \frac{2}{m} H_{n-1};
$$

(ii) g is nonnegative;

(iii) $g(0) = 1$.

Let PW(Ω) be the Paley-Wiener space of functions $g \in L^2(\mathbb{R}^{n-1})$ whose Fourier transform is supported in Ω . Given a finite-dimensional subspace F of PW($\frac{1}{m}H_{n-1}$) and a basis ${g_i}$ of F, we can optimize over functions of the form

$$
g(x) = \sum_{i,i'} X_{i,i'} g_i(x) g_{i'}(x),
$$
\n(3)

where X is a positive semidefinite matrix. These are integrable functions satisfying conditions (i) and (ii). Since condition (iii) and $\nu_n(g)$ are both linear in the entries of X, this reduces the optimization problem to a semidefinite program.

We have that

$$
\sum_{\substack{j\geq 1\\m_{\gamma_j}\leq T,\,\rho_j\geq n}}1\leq \frac{M_n(T)}{(n-1)!},
$$

and therefore

$$
\frac{Z_n(T)}{N(T)} \ge 1 - \frac{M_n(T)}{N(T)(n-1)!}.
$$

Thus,

$$
\liminf_{T \to \infty} \frac{Z_n(T)}{N(T)} \ge 1 - \frac{c_{n,m}}{(n-1)!},
$$

which shows we can use the above optimization problem to obtain a lower bound on the fraction of zeros having multiplicity at most $n - 1$.

We end this section with a possible formal solution to the above optimization problem. Note that the following result is true for H_{n-1} replaced by any compact set with a nonempty interior with essentially the same proof.

Theorem 2. For $n, m > 0$ there exists a kernel $K: \mathbb{C}^{n-1} \times \mathbb{C}^{n-1} \to \mathbb{C}$ such that

$$
g(w) = \int_{\mathbb{R}^{n-1}} g(v) \overline{K(w, v)} \, d\nu_n(v) \tag{4}
$$

for all $g \in PW(\frac{1}{m}H_{n-1})$ and $w \in \mathbb{C}^{n-1}$. We have

$$
c_{n,m} \le \frac{1}{K(0,0)},
$$

and equality is attained if every nonnegative $g \in L^1(\mathbb{R}^n)$ with $\text{supp}\,\widehat{g} \subseteq \frac{2}{m}H_{n-1}$ is a sum of squares in the sense that there are $g_{j,N} \in PW(\frac{1}{m}H_{n-1})$ such that

$$
g(x) = \lim_{N \to \infty} \sum_{j=1}^{N} |g_{j,N}(x)|^2,
$$
\n(5)

with almost everywhere convergence.

Proof. We have that $0 \leq W_n(x) \leq 1$ for all x, and $W_n(x) = 0$ if and only if $x_i = x_j$ for some $i \neq j$. Using induction it can be shown that for every $\epsilon > 0$ there exists $c(\epsilon) > 0$ such that $W_n(x) \geq c(\epsilon)$ whenever $|x_i - x_j| \geq \epsilon$ for all $i \neq j$. Define

 $S_{\epsilon} = \{x \in \mathbb{R}^{n-1} : |x_i| > \epsilon \text{ and } |x_i - x_j| > \epsilon \text{ for all distinct } i, j = 1, ..., n-1\}.$

Then $\nu_n(|g|^2) \le ||g||_2^2$ and

$$
\nu_n(|g|^2) \ge c(\epsilon) \int_{S_{\epsilon}} |g(x)|^2 dx.
$$

Observe that the set S_{ϵ} is relatively dense, that is, there is a cube Q (e.g., Q = $[-3\epsilon, 3\epsilon]^{n-1}$ and $\gamma > 0$ such that

$$
\inf_{x \in \mathbb{R}^{b-1}} \text{Vol}(S_{\epsilon} \cap (Q+x)) \ge \gamma \text{Vol}(Q).
$$

We could then apply the Logvinenko-Sereda Theorem [\[13,](#page-13-8) p. 112], which says there exists $b(\epsilon) > 0$ such that

$$
\int_{S_{\epsilon}} |g(x)|^2 dx \ge b(\epsilon) \|g\|_2^2,
$$

proving the norms $\nu_n(|g|^2)$ and $||g||_2^2$ are equivalent. For completeness, we give a direct proof of the above inequality in our particular case.

We apply the Hardy-Littlewood-Sobolev theorem of fractional integration (see, e.g., [\[23,](#page-14-6) Section V.1.2]), which implies that there exists a constant $C > 0$ such that

$$
||h(\cdot)||\cdot|^{-1/4}||_2\leq C||\widehat{h}||_{4/3}
$$

for all $h \in L^2(\mathbb{R})$. If in addition $\overline{\text{supp}}(h) \subseteq [-a, a]$, then using Hölder's inequality we obtain

$$
||h(\cdot)|| \cdot |^{-1/4}||_2 \le C ||\widehat{h}||_{4/3} \le (2a)^{1/4} C ||\widehat{h}||_2 = (2a)^{1/4} C ||h||_2.
$$

Then, with $h(x_i) = \tilde{g}(x + x_i e_i)$ and $a = 1/2$, we get

$$
\int_{\{x:|x_i-x_j|<\epsilon\}} |g(x)|^2\,dx \le \epsilon^{1/2} \int_{\{x:|x_i-x_j|<\epsilon\}} \left|\frac{g(x)}{|x_i-x_j|^{1/4}}\right|^2\,dx \le C\epsilon^{1/2} \|g\|_2^2.
$$

A similar procedure shows that $\int_{\{x:|x_i|<\epsilon\}} |g(x)|^2 dx \leq C\epsilon^{1/2} \|g\|_2^2$. Hence,

$$
\int_{S_{\epsilon}} |g(x)|^2 dx \ge ||g||_2^2 - \sum_{i=1}^{n-1} \int_{\{x: |x_i| < \epsilon\}} |g(x)|^2 dx - \sum_{1 \le i < j \le n} \int_{\{x: |x_i - x_j| < \epsilon\}} |g(x)|^2 dx
$$
\n
$$
= \left(1 - {n \choose 2} C \epsilon^{1/2}\right) ||g||_2^2.
$$

By choosing $\epsilon > 0$ small enough the constant $1 - {n \choose 2} C \epsilon^{1/2}$ is positive, which shows ν_n defines a norm equivalent to the L^2 norm. Hence PW($\frac{1}{m}H_{n-1}$) is a Hilbert space with the inner product given by ν_n .

Using Fourier inversion, Cauchy's inequality, and Plancherel's theorem it can be shown that the evaluation functions $f \mapsto f(x)$ on $PW(\frac{1}{m}H_{n-1})$ are continuous in the L^2 topology, so by the Riesz representation theorem there exists a kernel $K: \mathbb{C}^{n-1} \times \mathbb{C}^{n-1} \to \mathbb{C}$ with $K(x, \cdot) \in L^2(\mathbb{C}^{n-1})$ satisfying [\(4\)](#page-3-0).

To finish, observe that if g satisfies conditions (i), (ii), and (iii) and condition (5) , then by the monotone convergence theorem, limit [\(5\)](#page-3-1) holds also in the $L^1(\mathbb{R}^{n-1})$ sense, and since $PW(\frac{1}{m}H_{n-1})\cap L^1(\mathbb{R}^{n-1})$ is a reproducing kernel space, it is simple to show that we actually have uniform convergence in compact sets. Then,

$$
1 = g(0) = \lim_{N \to \infty} \sum_{j=1}^{N} |g_{j,N}(0)|^2 = \lim_{N \to \infty} \sum_{j=1}^{N} |\nu_n(g_{j,N}(\cdot)K(0,\cdot))|^2
$$

$$
\leq \lim_{N \to \infty} \sum_{j=1}^{N} \nu_n(|g_{j,N}|^2)K(0,0) = \nu_n(g)K(0,0).
$$

This shows $c_{n,m} \geq 1/K(0,0)$. Since $g(x) = K(0,x)^2/K(0,0)^2$ satisfies conditions (i), (ii), and (iii) and $\nu_n(g) = 1/K(0,0)$, we have $c_{n,m} \le 1/K(0,0)$, which completes the proof. the proof. \square

It is worth mentioning that condition [\(5\)](#page-3-1) holds true, and so equality $c_{n,m}$ $K(0,0)^{-1}$ is true, in the one-dimensional case (Hadamard factorization), which implies this condition is true in higher dimensions if H_{n-1} is replaced by a cube. Condition [\(5\)](#page-3-1) also holds true for a ball (see [\[9\]](#page-13-9)) and radial g, which is enough to show $c_{n,m} = K(0,0)^{-1}$. It is an open problem to prove condition [\(5\)](#page-3-1) for H_{n-1} replaced by a generic convex body. J. Vaaler, via personal communication, proposed an interesting way of obtaining [\(5\)](#page-3-1). He conjectures the following: For any centrally symmetric convex body K there is a constant $c(K) \in (0,1)$ such that for any nonnegative $g \in PW(K) \cap L^1$ there is $h \in PW(K)$ such that

$$
g \ge |h|^2
$$
 and $||h||_2^2 \ge c(K)||g||_1$.

If this conjecture is true, iterating this procedure ones finds functions h_1, h_2, \ldots such that $g \ge |h_1|^2 + ... + |h_N|^2$ and

 $||h_N||_2^2 \ge c(K)(||g||_1 - ||h_1||_2^2 - \ldots - ||h_{N-1}||_2^2).$

Then it is easy to see that we must have $g = \sum_{n \geq 1} |h_n|^2$ pointwise.

3. ANALYTIC SOLUTION FOR $n = 2$

In this section, we compute the kernel K of Theorem [2](#page-3-2) for $n = 2$ and all $m \ge 1$. We note that this kernel was already computed in [\[3\]](#page-13-7) but only for $m = 1$, and it relied heavily on the theory of de Branges spaces and Fourier transform techniques. In what follows we present a direct and simple approach that avoids any such things. Moreover, if one wants to compute only $K(0,0)$, then the argument is really short.

We want to compute the reproducing kernel $K(w, z)$ such that

$$
f(w) = \int_{\mathbb{R}} f(x) \overline{K(w, x)} (1 - s(x)^{2}) dx
$$

for every $w \in \mathbb{C}$ and $f \in PW([-\frac{1}{2m}, \frac{1}{2m}])$, where $s(x) = \frac{\sin(\pi x)}{\pi x}$. Letting $k(w, z) =$ $mK(mw, mz)$ we obtain

$$
f(w) = \int_{\mathbb{R}} f(x)\overline{k(w,x)}(1 - s(mx)^2) dx =: \langle f|k(w,\cdot)\rangle
$$

for every $w \in \mathbb{C}$ and $f \in PW([-\frac{1}{2},\frac{1}{2}])$. In what follows we use the trigonometric identity $2\sin^2(\theta) = 1 - \cos(2\theta)$. Observe that if $g \in PW([-1, 1])$ is even then

$$
\int_{\mathbb{R}} g(x)s(mx)^2 dx = \int_{\mathbb{R}} \frac{g(x) - g(0)}{2(\pi mx)^2} (1 - \cos(2\pi mx)) dx + \frac{1}{m} g(0)
$$

$$
= \int_{\mathbb{R}} \frac{g(x) - g(0)}{2(\pi mx)^2} dx + \frac{1}{m} g(0).
$$

Since $1 - s(x)^2$ is even, a simple computation shows $k(-w, z) = k(w, -z)$, and so $k(0, x)$ is even. Note also that any reproducing kernel satisfies $k(w, z) = \overline{k(z, w)}$ $k(\overline{z}, \overline{w})$. For simplicity we let

$$
k_{\pm}(w, z) = \frac{1}{2}(k(w, z) \pm k(-w, z))
$$
 and $s_{\pm}(w, z) = \frac{1}{2}(s(w - z) \pm s(w + z)).$

For $x, y \in \mathbb{R}$ we obtain

$$
s_{+}(x, y) = \langle s_{+}(x, \cdot) | k_{+}(y, \cdot) \rangle
$$

\n
$$
= k_{+}(x, y) - \frac{1}{m} s(x)k(0, y) - \int_{\mathbb{R}} \frac{s_{+}(x, t)k_{+}(t, y) - s(x)k(0, y)}{2(\pi mt)^{2}} dt
$$

\n
$$
= k_{+}(x, y) - \frac{1}{m} s_{+}(x, 0)k(0, y) - \int_{\mathbb{R}} s_{+}(x, t) \frac{k_{+}(t, y) - k(0, y)}{2(\pi mt)^{2}} dt
$$

\n
$$
- k(0, y) \int_{\mathbb{R}} \frac{s_{+}(x, t) - s(x)}{2(\pi mt)^{2}} dt
$$

\n
$$
= k_{+}(x, y) - \frac{1}{m} s(x)k(0, y) - \frac{k_{+}(x, y) - k(0, y)}{2(\pi mx)^{2}}
$$

\n
$$
- k(0, y) \frac{1 - \pi x \sin(\pi x) - \cos(\pi x)}{2(\pi mx)^{2}}
$$

\n
$$
= k_{+}(x, y) \left(1 - \frac{1}{2(\pi mx)^{2}}\right) + k(0, y) \left(-\frac{1}{m} s(x) + \frac{\pi x \sin(\pi x) + \cos(\pi x)}{2(\pi mx)^{2}}\right),
$$

where we used the identity

$$
\int_{\mathbb{R}} \frac{s_+(x,t) - s(x)}{2(\pi mt)^2} dt = \frac{1 - \pi x \sin(\pi x) - \cos(\pi x)}{2(\pi mx)^2}
$$

which can be proven via the residue theorem. We now evaluate at $x = u_m$ with $u_m = (\pi m \sqrt{2})^{-1}$ to get

$$
k(0, y) = \frac{s_{+}(u_{m}, y)}{-\sqrt{2}\sin(\pi u_{m}) + \pi u_{m}\sin(\pi u_{m}) + \cos(\pi u_{m})}
$$

which gives

$$
K(0,0) = \frac{k(0,0)}{m} = \frac{1}{\frac{1}{\sqrt{2}}\cot(\frac{1}{\sqrt{2}m}) - \frac{2m-1}{2m}}.
$$

We note that from this point on one can easily extract $k_{+}(x, y)$ by back substitution, if we let $a(x) = -s(x)/m + \frac{\pi x \sin(\pi x) + \cos(\pi x)}{2(\pi m x)^2}$ $\frac{2(\pi x)+\cos(\pi x)}{2(\pi mx)^2}$ then

$$
k_{+}(x,y) = \frac{s_{+}(x,y) - s_{+}(u_{m},y)a(x)/a(u_{m})}{1 - \frac{1}{2(\pi m x)^{2}}}.
$$

For $k_-(x, y)$, one can do a similar trick to extract $\partial_t k(y, t)|_{t=0}$. We have

$$
s_{-}(x,y) = k_{-}(x,y) - \int_{\mathbb{R}} \frac{s_{-}(x,t)k_{-}(y,t)}{2(\pi mt)^{2}} dt
$$

= $k_{-}(x,y) - \frac{1}{x} \int_{\mathbb{R}} \frac{(s_{+}(x,t) - \cos(\pi x)s(t))k_{-}(y,t)/t}{2(\pi m)^{2}} dt$
= $k_{-}(x,y) \left(1 - \frac{1}{2(\pi mx)^{2}}\right) + \frac{\cos(\pi x)}{2(\pi m)^{2}x} \partial_{t}k(y,t)|_{t=0},$

where we used that $s_-(x,t)/t = s_+(x,t) - \cos(\pi x)s(t)$, hence evaluation at $x = u_m$ gives

$$
\partial_t k(y, t)|_{t=0} = \frac{s_-(u_m, y)}{u_m \cos(\pi u_m)}.
$$

Letting $b(x) = \frac{\cos(\pi x)}{2(\pi m)^2 x}$ we get

$$
k_{-}(x,y) = \frac{s_{-}(x,y) - s_{-}(u_m, y)b(x)/b(u_m)}{1 - \frac{1}{2(\pi m x)^2}}
$$

and

$$
K(x, y) = \frac{1}{m}(k_{+}(x/m, y/m) + k_{-}(x/m, y/m)).
$$

4. Symmetry and rank-1 structure

In this section we show how we can use symmetry and a rank-1 structure to solve this problem efficiently.

Let Γ_n be the transpose of the symmetry group of H_n ; that is, the group containing the matrices A^{T} where A is a matrix such that $AH_n = H_n$. If g is a function satisfying conditions (i), (ii), and (iii), then, for $\gamma \in \Gamma_n$, the function $L(\gamma)g$ defined by $L(\gamma)g(x) = g(\gamma^{-1}x)$ also satisfies these constraints and $\nu_n(g) = \nu_n(L(\gamma)g)$. It then follows by an averaging argument that we may restrict to Γ_n -invariant functions.

Now assume F is Γ_n invariant. Consider a complete set Γ_n of irreducible, unitary representations of Γ_n . Let $\{g_{\pi,i,j}\}\)$ be a symmetry adapted system of the space F. By this we mean that $\{g_{\pi,i,j}\}\$ is a basis of F such that $H_{\pi,i} := \text{span}\{g_{\pi,i,j} : j = 1\}$ $1, \ldots, d_{\pi}$ is a Γ_n -irreducible representation with $H_{\pi,i}$ equivalent to $H_{\pi',i'}$ if and only if $\pi = \pi'$, and for each π , i and i' there exists a Γ_n -equivariant isomorphism $T_{\pi,i,i'}: H_{\pi,i} \to H_{\pi,i'}$ such that $g_{\pi,i',j} = T g_{\pi,i,j}$ for all j. Then, assuming the representations of Γ_n are of real type, any function g that is a sum of squares of functions from F can be written as

$$
g(x) = \sum_{\pi} \sum_{i,i'} X_{i,i'}^{(\pi)} \sum_j g_{\pi,i,j}(x) g_{\pi,i',j}(x),
$$

for positive semidefinite matrices $X^{(\pi)}$. This follows from Schur's lemma as explained in [\[10\]](#page-13-10) (see [\[8,](#page-13-11) Proposition 4.1] for a detailed proof).

To generate the symmetry-adapted system ${g_{\pi,i,j}}$ for F we use an implementation of the projection algorithm described in [\[22\]](#page-14-7). For this, we first define the operators

$$
p_{j,j'}^{(\pi)} = \frac{d_{\pi}}{|\Gamma|} \sum_{\gamma \in \Gamma} \pi(\gamma^{-1})_{j,j'} L(\gamma), \tag{6}
$$

where d_{π} is the dimension of π and $L(\gamma)$ is the operator $L(\gamma)g(x) = g(\gamma^{-1}x)$. Then we choose bases $\{g_{\pi,i,1}\}_i$ of $\text{Im}(p_{1,1}^{(\pi)})$ and set $g_{\pi,i,j} = p_{j,1}^{(\pi)} g_{\pi,i,1}$. **Lemma 3.** If π is not the trivial representation, then $g_{\pi,i,j}(0) = 0$.

Proof. Let $p_{j,j'}^{(\pi)}$ be the operator as defined in [\(6\)](#page-7-1) and $g \in F$. Then,

$$
(p_{j,1}^{(\pi)} p_{1,1}^{(\pi)} g)(x) = \frac{d_{\pi}^2}{|\Gamma|^2} \sum_{\beta,\gamma \in \Gamma} \pi(\beta^{-1})_{j,1} \pi(\gamma^{-1})_{1,1} g(\beta^{-1} \gamma^{-1} x),
$$

and thus

$$
(p_{j,1}^{(\pi)} p_{1,1}^{(\pi)} g)(0) = \frac{d_{\pi}^2}{|\Gamma|^2} \left(\sum_{\beta \in \Gamma} \pi(\beta^{-1})_{j,1} \right) \left(\sum_{\gamma \in \Gamma} \pi(\gamma^{-1})_{1,1} \right) g(0).
$$

If π is not the trivial representation, then $\sum_{\gamma \in \Gamma} \pi(\gamma^{-1})_{j,j'} = 0$ by orthogonality of matrix coefficients. By linearity, the result then follows. \square

This shows optimizing over functions of the form [\(3\)](#page-3-3) is the same as optimizing over functions of the form

$$
g(x) = \sum_{i,i'} X_{i,i'} g_{1,i,1}(x) g_{1,i',1}(x),
$$

where X is positive semidefinite and where we denote the trivial representation by 1. That is, instead of using a Γ-invariant space of functions, we can use the much smaller space of Γ-invariant functions $g_i(x) = g_{1,i,1}(x)$.

We now obtain the semidefinite program

minimize
$$
\langle A, X \rangle
$$

subject to $\langle bb^{\mathsf{T}}, X \rangle = 1$ (7)
 $X \succeq 0$,

where $A_{i,i'} = \nu_n(g_i g_{i'})$ and $b_i = g_i(0)$. Here we use the notation $\langle A, B \rangle = \text{tr}(A^{\mathsf{T}}B)$.

A semidefinite program with m equality constraints has an optimal solution of rank r with $r(r + 1)/2 \leq m$ [\[19\]](#page-14-8). Since the above semidefinite program has only one equality constraint, we know that there exists an optimal solution of rank 1. In fact, the following lemma shows we can find this rank 1 solution by solving a linear system. We can view this lemma as a finite-dimensional version of Theorem [2.](#page-3-2)

Lemma 4. Let A be a positive semidefinite matrix and let x be a solution to the system $Ax = b$. Then $xx^T/(x^Tb)^2$ is an optimal solution to the semidefinite program

$$
minimize \quad \langle A, X \rangle
$$

subject to
$$
\langle bb^{\mathsf{T}}, X \rangle = 1
$$

 $X \succeq 0$.

Proof. Let X be feasible to the semidefinite program and consider the spectral decomposition $X = \sum_i v_i v_i^{\mathsf{T}}$. Using Cauchy-Schwarz we have

$$
1 = b^{\mathsf{T}} X b = \sum_{i} (b^{\mathsf{T}} v_i)^2 = \sum_{i} (x^{\mathsf{T}} A v_i)^2 \le \sum_{i} (x^{\mathsf{T}} A x)(v_i^{\mathsf{T}} A v_i) = x^{\mathsf{T}} b \langle A, X \rangle,
$$

$$
\langle A, X \rangle \ge \frac{1}{x^{\mathsf{T}} b} = \left\langle A, \frac{xx^{\mathsf{T}}}{(x^{\mathsf{T}} b)^2} \right\rangle.
$$

so

5. Parametrization using polynomials

Here we consider the case $(n, m) = (3, 1)$. Let $\chi_{H_2}(x)$ be the indicator function of the region H_2 and define the functions $g_\alpha(x)$ by

$$
\widehat{g}_{\alpha}(x) = x_1^{\alpha_1} x_2^{\alpha_2} \chi_{H_2}(x).
$$

Let $F = \text{span}\lbrace g_{\alpha} : \alpha_1 + \alpha_2 \leq d \rbrace$. As before we let $\lbrace g_i \rbrace$ be a basis of the Γ-invariant functions in F and we use the parametrization

$$
g(x) = \sum_{i,i'} X_{i,i'} g_i(x) g_{i'}(x).
$$

To set up the linear system for solving [\(7\)](#page-8-0) we need to compute the linear functionals $g(0)$ and $\nu_3(g)$ explicitly in the entries of the matrix X. To compute $g(0)$ we first note that with $C = [-1/2, 0] \times [0, 1/2]$ we have

$$
H_2 = C \cup r^2 C \cup r^4 C.
$$

Since g_i is of the form $\hat{g}_i(x) = p_i(x) \chi_{H_2}(x)$, with p_i a Γ -invariant polynomial, we have

$$
g_i(0) = \int_{H_2} p_i(x) dx = \int_C (p_i(x) + p_i(r^2 x) + p_i(r^4 x)) dx = 3 \int_C p_i(x) dx
$$

= $3 \int_0^{1/2} \int_{-1/2}^0 p_i(x_1, x_2) dx_1 dx_2$,

so

$$
g(0) = \sum_{i,i'} X_{i,i'} g_i(0) g_{i'}(0)
$$

is an explicitly computable linear functional in the entries of the matrices X .

As computed in Hejhal [\[14\]](#page-14-1), we have

$$
\nu_3(g) = \int_{\mathbb{R}^2} g(x_1, x_2) W_3(x_1, x_2, 0) dx_1 dx_2
$$

= $2 + \hat{g}(0) + 6 \int_0^1 \hat{g}(x, 0)(x - 1) dx - 12 \int_0^1 \int_{-x_2}^0 \hat{g}(x_1, x_2) x_2 dx_1 dx_2.$

The term $\hat{g}(0)$ can be computed in the same way $g_i(0)$ is computed above. To compute the second term we use the notation

$$
\mathcal{J}_u = \begin{cases} \int_{-\frac{1}{2}+u}^{0} \int_{-\frac{1}{2}-u}^{\frac{1}{2}-u} + \int_{0}^{u} \int_{-\frac{1}{2}}^{\frac{1}{2}-u} + \int_{u}^{\frac{1}{2}} \int_{-\frac{1}{2}}^{\frac{1}{2}-x_2} & 0 \le u \le \frac{1}{2}, \\ \int_{u-\frac{1}{2}}^{\frac{1}{2}} \int_{-\frac{1}{2}}^{\frac{1}{2}-u} & \frac{1}{2} \le u \le 1. \end{cases}
$$

By symmetry, we have $\int_0^1 \hat{g}(x,0)(x-1) dx = \int_0^1 \hat{g}(-x,x)(x-1) dx$, which gives the integral

$$
\int_0^1 \hat{g}(u, -u)(u-1) du = \sum_{i,i} X_{i,i'} \int_0^1 \mathcal{J}_u p_i(x_1, x_2) p_{i'}(x_1 + u, x_2 - u) dx_1 dx_2 du.
$$

TABLE 1. Upper bounds on the best possible value for c_3 obtained using a parametrization using polynomials of degree d.

d.	CЗ
10	0.077516654
20	0.077222625
30	0.077206761
40	0.077200000
50	0.077198398
60	0.077197284

To compute the third term we use the notation

$$
\mathcal{I}_{u} = \begin{cases}\n\int_{u_{2}-\frac{1}{2}}^{\frac{1}{2}} \int_{-\frac{1}{2}}^{u_{1}+\frac{1}{2}} \left(u_{1}, u_{2} \right) \in R_{1}, \\
\int_{u_{2}-\frac{1}{2}}^{\frac{1}{2}} \int_{-\frac{1}{2}-x_{2}+u_{1}+u_{2}}^{\frac{1}{2}-x_{2}} \left(u_{1}, u_{2} \right) \in R_{2}, \\
\int_{u_{1}+u_{2}}^{\frac{1}{2}} \int_{-\frac{1}{2}}^{-x_{2}} + \int_{-u_{1}}^{u_{1}+u_{2}} \int_{u_{1}+u_{2}-x_{2}-\frac{1}{2}}^{1-x_{2}} + \int_{u_{2}-\frac{1}{2}}^{u_{1}} \int_{u_{1}+u_{2}-x_{2}-\frac{1}{2}}^{u_{1}+\frac{1}{2}} \left(u_{1}, u_{2} \right) \in R_{3}, \\
\int_{-u_{1}}^{\frac{1}{2}} \int_{-\frac{1}{2}}^{-x_{2}} + \int_{u_{1}+u_{2}}^{-u_{1}} \int_{-\frac{1}{2}}^{u_{1}+\frac{1}{2}} + \int_{u_{2}-\frac{1}{2}}^{u_{1}+u_{2}} \int_{u_{1}+u_{2}-x_{2}-\frac{1}{2}}^{u_{1}+\frac{1}{2}} \left(u_{1}, u_{2} \right) \in R_{4}.\n\end{cases}
$$

where we have $R_1 = \{-1 \le -u_2 \le u_1 \le -\frac{1}{2}\}, R_2 = \{-\frac{1}{2} \le \frac{1}{2} - u_2 \le u_1 \le 0\},$ $R_3 = \{0 \le -2u_1 \le u_2 \le \frac{1}{2} - u_1 \le 1\}$ and $R_4 = \{0 \le -u_1 \le u_2 \le -2u_1 \le 1\}$. We then have

$$
\int_0^1 \int_{-u_2}^0 \hat{g}(u_1, u_2) u_2 du_1 du_2 = \sum_{i,i'} X_{i,i'} \int_0^1 \int_{-u_2}^0 \mathcal{I}_u p_i(x) p_{i'}(x-u) u_2 dx_1 dx_2 du_1 du_2.
$$

In Table [1](#page-10-1) we show the bounds obtained with this parametrization for degree d. This proves the $(n, m) = (3, 1)$ case of Theorem [1.](#page-1-1) As ancillary files to the arXiv version of this paper we give the GAP source code to compute the polynomials p_i in exact arithmetic and a Julia file to compute the above bound rigorously by solving a linear system in ball arithmetic.

6. Parametrization using shifts

Now we consider the functions g_{λ} obtained by shifting the Fourier transform of the indicator function of $\frac{1}{m}H_{n-1}$ by $\lambda \in m\mathbb{Z}^{n-1}$:

$$
g_{\lambda}(x) = \widehat{\chi_{\frac{1}{m}H_{n-1}}(x-\lambda)} = \frac{1}{m^{n-1}} \widehat{\chi_{H_{n-1}}}\left(\frac{1}{m}(x-\lambda)\right).
$$

Then $g_{\lambda}(x)$ is the Fourier transform of $\chi_{\frac{1}{m}H_{n-1}}(\cdot)e^{-2\pi i\langle\lambda,\cdot\rangle}$, so $g_{\lambda}(\gamma x) = g_{\gamma^{-1}\lambda}(x)$ for all γ in the group Γ_n as defined in Section [4.](#page-7-0)

We observe that the set of functions $G = \{g_{\lambda}\}_{{\lambda \in {\Lambda}}}$ with ${\Lambda} = m\mathbb{Z}^{n-1}$ is an orthogonal basis of PW($\frac{1}{m}H_{n-1}$) with respect to the Lebesgue L^2 -norm for $n=2$, but it is not for $n \geq 3$. For $n = 3$, a classical result of Venkov [\[24\]](#page-14-9) shows that G is an orthogonal basis if we take Λ as the dual lattice of $\{(u/2+v, u/2-v/2) : u, v \in \frac{1}{m}\mathbb{Z}\}.$ For $n \geq 4$, a recent result [\[12,](#page-13-12) [16\]](#page-14-10) implies that no set of translations Λ exists such that G is an orthogonal basis, otherwise H_{n-1} would need to be centrally symmetric, have only centrally symmetric faces and tile the space by translations, but none of these properties hold. In any case, in practice, using $\Lambda = m\mathbb{Z}^{n-1}$

works reasonably well since G is an orthogonal basis of $PW(\frac{1}{m}[-1/2, 1/2]^{n-1})$ which contains $PW(\frac{1}{m}H_{n-1}).$

Let F be a finite-dimensional, Γ_n -invariant subspace of span $\{g_\lambda : \lambda \in m\mathbb{Z}^{n-1}\},$ and again use the parametrization

$$
g(x) = \sum_{i,i'} X_{i,i'} g_i(x) g_{i'}(x),
$$

where ${g_i}$ is a basis for the invariants in F.

It remains to compute the matrix $A_{i,i'} = \nu_n(g_i g_{i'})$ and the vector $b_i = g_i(0)$, so that by Lemma [4](#page-8-1) the optimal function is given by

$$
g(x) = \frac{(\sum_i c_i g_i(x))^2}{(c^{\mathsf{T}} b)^2}
$$

with c a solution to $Ac = b$.

In [A](#page-12-0)ppendix A we calculate the Fourier transform of $\chi_{H_n}(x)$ for general n. This function has removable singularities whenever $x_i = 0$ for some i or $x_i = x_j$ for some $i \neq j$. The function g_i is a linear combination of shifts of the Fourier transform and thus can also have removable singularities. To compute $g_i(0)$ we consider each term individually, check whether the denominator evaluates to zero and if it does we use repeated applications of L'Hôpital's rule.

We want to compute $\int_{\mathbb{R}^{n-1}} f(x) dx$, where $f(x) = g(x)W_n(x, 0)$. To simplify the computations we change variables to $m^{-1}x$. Since the Fourier transform of $g_i(mx)$ is supported in $[-1/2, 1/2]^{n-1}$ and the Fourier transform of $W_n(mx, 0)$ is supported in $[-m, m]^{n-1}$, the Fourier transform of f is supported in $[-(m + 1), (m + 1)]^{n-1}$. By Poisson summation, we then have

$$
\int_{\mathbb{R}^{n-1}} f(x) dx = \widehat{f}(0) = \sum_{k \in (m+1)\mathbb{Z}^{n-1}} \widehat{f}(k) = \frac{1}{(m+1)^{n-1}} \sum_{k \in \frac{1}{m+1}\mathbb{Z}^{n-1}} f(k).
$$

To avoid the removable singularities in the summation we again use Poisson summation, by which

$$
\sum_{k \in \frac{1}{m+1} \mathbb{Z}^{n-1}} f(k) = (m+1)^{n-1} \widehat{f}(0)
$$
\n
$$
= (m+1)^{n-1} \sum_{k \in (m+1) \mathbb{Z}^{n-1}} \widehat{f}(k) e^{2\pi i \langle k, s \rangle} = \sum_{k \in \frac{1}{m+1} \mathbb{Z}^{n-1}} f(k+s)
$$

for any $s \in \mathbb{R}^{n-1}$. In our computations we use

$$
s = \left(\frac{1}{m(m+1)n}, \frac{2}{m(m+1)n}, \dots, \frac{n-1}{m(m+1)n}\right)
$$

.

Because of the sines and cosines in the formula of $\chi_{H_{n-1}}$, the series $\sum_{k} f(k+s)$ can be split into $(2(m+1))^{n-1}$ series of rational functions. We can then further split this into a sum of series of rational functions with constant numerators. For (n, m) = (4, 1) this results in 71568 series of rational functions with constant numerators.

Computing these series exactly is the computational bottleneck in our approach. Maple is the only system among the computer algebra systems we tried in which we could successfully compute the sums of all the series, but, unfortunately, Maple 2022.2 contains a bug where it acts nondeterministically and sometimes gives a completely wrong answer. We, therefore, check that the sum computed with Maple deviates at most 0.1% from the sum we obtain by truncating the series to the box $[-C/(m+1), C/(m+1)]^{n-1}$, where for $n = 3$ we use $C = 400$ (or $C = 1300$ for the cases where this is not sufficient) and for $n = 4$ we use $C = 25$. We give text files containing the series in Maple syntax and for each series an interval containing the correct sum given by decimal expressions of a midpoint and a radius. We also give a Julia file with the code to obtain the rigorous bounds using the data in these text files.

For $(n, m) = (3, 1)$ and $(n, m) = (3, 2)$ $(n, m) = (3, 2)$ $(n, m) = (3, 2)$, Table 2 gives the results for the subspaces $F = \text{span}{g_{\lambda} : \lambda \in S_d}$, where

$$
S_d = \Gamma_n \{ m\lambda \in \mathbb{Z}^{n-1} : ||\lambda||_1 \le d \}.
$$

This proves the case $(3, 2)$ of Theorem [1](#page-1-1) (the case $(3, 1)$ was already done in Sec-tion [5\)](#page-9-0). For $(n, m) = (4, 1)$, we only compute the series exactly for $\lambda = 0$ (partly because of the aforementioned Maple bug), which gives $c_{4,1} \leq 447/3500 \approx 0.1277$ as used in Theorem [1.](#page-1-1) Numerically we estimate that by using more values of λ this can be improved to $c_{4,1} \leq 0.026$. This would improve the lower bound in Theorem [1](#page-1-1) from 0.9787 to 0.9956.

TABLE 2. Upper bounds on the best possible value for $c_{n,m}$ obtained through a parametrization using shifts S_d^2 .

d.	$c_{3,1}$	$c_{3,2}$
Ω	0.14444445	1.48888889
1	0.077710979	1.401604735
$\mathcal{D}_{\mathcal{A}}$	0.077580416	1.401343568
3	0.077261926	1.400616000
4	0.077247720	1.400581457
5	0.077213324	1.400506625

APPENDIX A. THE FOURIER TRANSFORM OF χ _{H_{n−1}}

To compute the Fourier transform of the indicator function of H_{n-1} we use the Fourier transform of the indicator function of the simplex $S_n = \{x \in \mathbb{R}^n : x_j \geq 0\}$ 0, $x_1 + ... + x_n \le 1$ as computed in [\[1,](#page-13-13) Eq. 18]:

$$
\widehat{\chi}_{S_n}(y) = \frac{(-1)^{n+1}}{(2\pi i)^n} \sum_{j=1}^n \frac{1 - e^{-2\pi i y_j}}{y_j \prod_{i \neq j} (y_j - y_i)}.
$$

The Fourier transform of the indicator function of the cross-polytope $C_n = \{x \in$ $\mathbb{R}^n : |x_1| + \ldots + |x_n| \leq 1$ is given by

$$
\widehat{\chi}_{C_n}(y) = \sum_{\sigma \in \{\pm 1\}^n} \widehat{\chi}_{S_n}(\sigma_1 y_1, \ldots, \sigma_n y_n).
$$

If we consider each term in $\widehat{\chi}_{S_n}(y)$ individually then it is simple to deduce that

$$
\sum_{\sigma \in \{\pm 1\}^n} \frac{1 - e^{-2\pi i \sigma_j y_j}}{\sigma_j y_j \prod_{i \neq j} (\sigma_j y_j - \sigma_i y_i)} = \sum_{\sigma \in \{\pm 1\}} (2\sigma y_j)^{n-1} \frac{1 - e^{-2\pi i \sigma y_j}}{\sigma y_j \prod_{i \neq j} (y_j^2 - y_i^2)},
$$

while the latter depends on the parity of n . This implies that

$$
\widehat{\chi}_{C_n}(y) = \begin{cases}\n\pi^{-n}(-1)^{(n-1)/2} \sum_{j=1}^n \frac{y_j^{n-2} \sin(2\pi y_j)}{\prod_{i \neq j} (y_j^2 - y_i^2)} & \text{if } n \text{ is odd, and} \\
\pi^{-n}(-1)^{n/2-1} \sum_{j=1}^n \frac{2y_j^{n-2} \sin(\pi y_j)^2}{\prod_{i \neq j} (y_j^2 - y_i^2)} & \text{if } n \text{ is even.} \n\end{cases}
$$

Finally, notice that H_{n-1} can be identified with $C_n \cap \{x : x_1 + \ldots + x_n = 0\}$ and thus one can rigorously justify that

$$
\widehat{X}_{H_{n-1}}(y_1,\ldots,y_{n-1}) = \lim_{T \to \infty} \int_{-T}^T \widehat{X}_{C_n}(y_1+t,\ldots,y_{n-1}+t,t) dt
$$

The function under the integral sign above only has simple poles at $t = -\frac{y_j + y_k}{2}$ for $k \neq j$ and $t = -y_j/2$ (for generic $y \in \mathbb{R}^{n-1}$). A routine application of the residue theorem and a grouping of terms shows that $\widehat{\chi}_{H_{n-1}}(y)$ is equal to

$$
\frac{(-1)^{(n-1)/2}}{2^{n-2}\pi^{n-1}}\bigg[-\sum_{1\leq j < k\leq n-1} \frac{(y_j - y_k)^{n-3}\cos(\pi(y_j - y_k))}{y_j y_k \prod_{i\neq j,k} (y_j - y_i)(y_k - y_i)} + \sum_{j=1}^{n-1} \frac{y_j^{n-3}\cos(\pi y_j)}{\prod_{i\neq j} (y_j - y_i)y_i}\bigg]
$$

if n is odd, and equal to

$$
\frac{(-1)^{n/2-1}}{2^{n-2}\pi^{n-1}}\bigg[\sum_{1\leq j < k\leq n-1} \frac{(y_j - y_k)^{n-3}\sin(\pi(y_j - y_k))}{y_j y_k \prod_{i\neq j,k} (y_j - y_i)(y_k - y_i)} + \sum_{j=1}^{n-1} \frac{y_j^{n-3}\sin(\pi y_j)}{\prod_{i\neq j} (y_j - y_i)y_i}\bigg]
$$

if n is even.

REFERENCES

- [1] B. Borda, Lattice points in algebraic cross-polytopes and simplices, Discrete & Computational Geometry 60 (2018), 145–169.
- [2] H. M. Bui and D. R. Heath-Brown, On simple zeros of the Riemann zeta-function, Bull. Lond. Math. Soc. 45 (2013), no. 5, 953–961. MR 3104987
- [3] E. Carneiro, V. Chandee, F. Littmann, and M. B. Milinovich, Hilbert spaces and the pair correlation of zeros of the Riemann zeta-function, J. Reine Angew. Math. **725** (2017), 143– 182. MR 3630120
- [4] A. Y. Cheer and D. A. Goldston, *Simple zeros of the Riemann zeta-function*, Proc. Amer. Math. Soc. 118 (1993), no. 2, 365–372.
- [5] A. Chirre, F. Gonçalves, and D. de Laat, Pair correlation estimates for the zeros of the zeta function via semidefinite programming, Adv. Math. 361 (2020), 106926, 22.
- [6] H. Cohn, D. de Laat, and A. Salmon, Three-point bounds for sphere packing, preprint, 2022. arXiv:2206.15373
- [7] H. Cohn and N. Elkies, New upper bounds on sphere packings. I, Ann. of Math. (2) 157 (2003), no. 2, 689–714. MR 1973059
- [8] D. de Laat and N. Leijenhorst, Solving clustered low-rank semidefinite programs arising from polynomial optimization, preprint, 2022. arXiv:2202.12077
- [9] A. V. Efimov, An analog of rudin's theorem for continuous radial positive definite functions of several variables, Proceedings of the Steklov Institute of Mathematics (Supplementary issues) 284 (2014), no. 1, 79–86.
- [10] K. Gatermann and P. A. Parrilo, Symmetry groups, semidefinite programs, and sums of squares, J. Pure Appl. Algebra 192 (2004), no. 1-3, 95–128. MR 2067190
- [11] D. A. Goldston, S. M. Gonek, A. E. Ozlük, and C. Snyder, On the pair correlation of zeros of the Riemann zeta-function, Proc. London Math. Soc. (3) 80 (2000), no. 1, 31–49. MR 1719184
- [12] R. Greenfeld and N. Lev, Fuglede's spectral set conjecture for convex polytopes, Analysis and PDE (2017), no. 10, 1497–1538.
- [13] V. P. Havin and B. Joricke, he uncertainty principle in harmonic analy- sis, Springer-Verlag, Berlin Heidelberg (1994).
- [14] D. A. Hejhal, On the triple correlation of zeros of the zeta function, International Mathematics Research Notices 1994 (1994), no. 7, 293–302.
- [15] H. Iwaniec and E. Kowalski, Analytic number theory, American Mathematical Society Colloquium Publications, AMS, Providence, RI 53 (2004).
- [16] N. Lev and M. Matolcsi, The fuglede conjecture for convex domains is true in all dimensions, Acta Mathematica (2022), no. 228, 385–420.
- [17] H. L. Montgomery, The pair correlation of zeros of the zeta function, Analytic number theory (Proc. Sympos. Pure Math., Vol. XXIV, St. Louis Univ., St. Louis, Mo., 1972), 1973, pp. 181– 193. MR 0337821
- [18] H. L. Montgomery, Distribution of the zeros of the Riemann zeta function, Proceedings of the International Congress of Mathematicians (Vancouver, B. C., 1974), Vol. 1, 1975, pp. 379– 381. MR 0419378
- [19] G. Pataki, On the rank of extreme matrices in semidefinite programs and the multiplicity of optimal eigenvalues, Math. Oper. Res. 23 (1998), 339–358.
- [20] Z. Rudnick and P. Sarnak, The n-level correlations of zeros of the zeta function, C. R. Acad. Sci. Paris Sér. I Math. 319 (1994), no. 10, 1027-1032. MR 1305671
- [21] , Zeros of principal L-functions and random matrix theory, Duke Math. J. 81 (1996), no. 2, 269–322, A celebration of John F. Nash, Jr. MR 1395406
- [22] J.-P. Serre, Linear representations of finite groups, Graduate Texts in Mathematics, Vol. 42, Springer-Verlag, New York-Heidelberg, 1977, Translated from the second French edition by Leonard L. Scott. MR 0450380
- [23] E. M. Stein, Singular integrals and differentiability properties of functions, Princeton Mathematical Series, No. 30, Princeton University Press, Princeton, N.J., 1970. MR 0290095
- [24] B. Venkov, On a class of euclidean polyhedra (russian), Vestnik Leningrad. Univ. Ser. Mat. Fiz. Him. (1954), no. 9, 11–31.

IMPA, Rio de Janeiro, Brazil Email address: goncalves@impa.br

Delft Institute of Applied Mathematics, Delft University of Technology, Delft, The Netherlands

Email address: d.delaat@tudelft.nl

Delft Institute of Applied Mathematics, Delft University of Technology, Delft, The Netherlands

Email address: n.m.leijenhorst@tudelft.nl