Energy stable and L^2 norm convergent BDF3 scheme for the Swift–Hohenberg equation

Xuan Zhao^{*} Ran Yang[†] Zhongqin Xue [‡] Hong Sun[§]

Abstract

A fully discrete implicit scheme is proposed for the Swift-Hohenberg model, combining the third-order backward differentiation formula (BDF3) for the time discretization and the second-order finite difference scheme for the space discretization. Applying the Brouwer fixed-point theorem and the positive definiteness of the convolution coefficients of BDF3, the presented numerical algorithm is proved to be uniquely solvable and unconditionally energy stable, further, the numerical solution is shown to be bounded in the maximum norm. The proposed scheme is rigorously proved to be convergent in L^2 norm by the discrete orthogonal convolution (DOC) kernel, which transfer the four-level-solution form into the three-levelgradient form for the approximation of the temporal derivative. Consequently, the error estimate for the numerical solution is established by utilization of the discrete Gronwall inequality. Numerical examples in 2D and 3D cases are provided to support the theoretical results.

Keywords: Swift-Hohenberg equation; BDF3 scheme; Energy stability; Convergence;

1 Introduction

In this work, a fully discrete implicit scheme is proposed for the following Swift-Hohenberg equation with the periodic boundary conditions

$$u_t = -(1+\Delta)^2 u - f(u), (1.1)$$

where $f(u) = u^3 - gu^2 - \epsilon u$, ϵ and g are positive contants with physical significance. A closed two-dimensional domain $\Omega \subset \mathbb{R}^2$ is considered for the theoretical part.

The Swift-Hohenberg equation is a nonlinear fourth-order partial differential equation, which describes the effects of thermal fluctuations on the Rayleigh-Bénard instability [1]. As one of the paradigms of nonlinear dynamical system, the Swift-Hohenberg equation has been widely applied in physics [2], materials science [3], dynamics of ecological systems [4], and many other nonlinear fields [5–7].

^{*}Corresponding author. School of Mathematics, Southeast University, Nanjing 210096, P. R. China (xu-anzhao11@seu.edu.cn).

[†]School of Mathematics, Southeast University, Nanjing 210096, P. R. China (220191527@seu.edu.cn).

[‡]School of Mathematics, Southeast University, Nanjing 210096, P. R. China. (zqxue@seu.edu.cn).

[§]School of Mathematics, Southeast University, Nanjing 210096; Department of Mathematics and Physics, Nanjing Institute of Technology, Nanjing 211167, P. R. China (sunhongzhal@126.com).

The energy of the equation(1.1) is defined by the Lyapunov energy functional

$$E(u) := \int_{\Omega} \left(\frac{1}{2} u (1+\Delta)^2 u + \frac{1}{4} u^4 - \frac{g}{3} u^3 - \frac{\epsilon}{2} u^2 \right) \mathrm{d}\mathbf{x}, \tag{1.2}$$

and is nonincreasing in time

$$\frac{d}{dt}E(u) = -\int_{\Omega} \left(\frac{du}{dt}\right)^2 \,\mathrm{d}\mathbf{x} \le 0.$$
(1.3)

Bifurcation analysis [8] and pattern selection [9] of the equilibrium solutions for the Swift-Hohenberg equation are theoretically analyzed. Shi and Han [10] obtained a reversible homoclinic solution approaching to a periodic solution of the Swift-Hohenberg equation, and the existence of nontrivial periodic solutions was proved by Lai and Zhang [11]. In [12], the effects of rapid oscillations of the forcing term on the long-time behaviour of the solutions are studied. We also refer the reader to the recent article [13] on nonstationary case and the references therein.

Recently, the efficient numerical methods for the Swift-Hohenberg equation are considered since the analytically solutions cannot be generally obtained. A fully discrete discontinuous Galerkin scheme is proposed in [14], where the time step as well as the mesh size are irrespective. The scheme is proved uniquely solvable and unconditionally energy stable. A second-order energy-stable time-integration method that suppresses numerical instabilities is presented by Sarmiento et al. [15] and a detailed proof of the unconditional energy stability is provided. Liu [16] considered two linear, second-order and unconditionally energy stable schemes by linear invariant energy quadratization and modified scalar auxiliary variable approaches. Dehghan et al. [17] combined the proper orthogonal decomposition approach and the local discontinuous Galerkin technique, and discussed the energy stability. Sun et al. [21] proposed an adaptive BDF2 scheme for the Swift-Hohenberg equation, and proved the energy stability and the convergence. Besides, readers can refer to the references [22–30] for some related numerical methods. However, in most existing works, the convergence of the numerical algorithms for Swift-Hohenberg equation is rarely considered.

As is known that the Swift-Hohenberg equation takes a long-time to reach the steady state, the key challenge in design the numerical methods is the improvement for the computational efficiency of the numerical schemes. High order time-stepping strategies and adaptive timestepping algorithms are always suitable choice for developing the efficient numerical schemes. In the literature, we name the following methods in simplicity for the discretization of Swift-Hohenberg equation in temporal direction

1) Crank-Nicolson type of scheme (second-order) [18]

$$\frac{u^{k+1} - u^k}{\tau} = D\left(-\frac{\partial^4}{\partial x^4} - \frac{\partial^4}{\partial y^4} - 2k^2\frac{\partial^2}{\partial x^2} - 2k^2\frac{\partial^2}{\partial y^2} - 2\frac{\partial^4}{\partial x^2\partial y^2} - k^4\right)\frac{u^{k+1} + u^k}{2}$$
$$-\frac{U(u^{n+1}) - U(u^n)}{u^{n+1} - u^n}$$

where U(u) stands for the potential of the nonlinear force upon the system.

2) Convex splitting method (first-/second-order) [19]

- The first order convex splitting

$$\frac{u^{k+1} - u^k}{\tau} = -(u^{k+1})^3 + g(u^{k+1})^2 - (A + (1+\Delta)^2)u^{k+1} + (A+\epsilon)u^k,$$

- The second order convex splitting

$$\frac{u^{k+1} - u^k}{\tau} = -\left(\chi(u^{k+1}, u^k) + (A + (1 + \Delta)^2)\left(\frac{u^{k+1} + u^k}{2}\right) - (A + \epsilon)\frac{3u^k - u^{n-1}}{2}\right),$$

where $\chi(u, v) = \frac{u^2 + v^2}{2}\frac{u + v}{2} - g\frac{u^2 + uv + v^2}{3}.$

3) Semi-implicit Euler scheme (first-order) [20]

$$\frac{u^{k+1} - u^k}{\tau} + A(u^{k+1} - u^k) = 2\Delta u^k - \Delta^2 u^{k+1} - (1 - \epsilon)u^{k+1} - (u^k)^3,$$

where $A(u^{k+1} - u^k)$ is an extra artificial term to preserve the energy stability.

4) Multiple SAV scheme based on the Crank-Nicolson method (second-order) [16]

$$\begin{split} \frac{u^{k+1} - u^k}{\tau} &= M \Delta \mu^{k+\frac{1}{2}}, \\ \mu^{k+\frac{1}{2}} &= (1+\Delta)^2 \left(\frac{u^{k+1} + u^k}{2}\right) + \frac{r^{k+1} + r^k}{2\sqrt{\tilde{E}(\tilde{u}^{k+1/2})}} U(\tilde{u}^{k+1/2}) \\ &- \frac{m^{k+1} + m^k}{2\sqrt{E_0(\tilde{u}^{k+1/2})}} V(\tilde{u}^{k+1/2}), \\ \frac{r^{k+1} - r^k}{\tau} &= \frac{1}{2\sqrt{\tilde{E}(\tilde{u}^{k+1/2})}} \int_{\Omega} U(\tilde{u}^{k+1/2}) \frac{u^{k+1} - u^k}{\tau} \, \mathrm{d}\mathbf{x}, \\ \frac{m^{k+1} - m^k}{\tau} &= \frac{1}{2\sqrt{E_0(\tilde{u}^{k+1/2})}} \int_{\Omega} V(\tilde{u}^{k+1/2}) \frac{u^{k+1} - u^k}{\tau} \, \mathrm{d}\mathbf{x}, \end{split}$$

where $E_0(u) = \int_{\Omega} \frac{\mathbf{g}}{6} u^4 + \frac{\mathbf{g}}{6} u^2 + \frac{\mathbf{e}}{6} u^4 > 0$, $\tilde{E}(u) = \int_{\Omega} \frac{1}{4} u^4 - \frac{\mathbf{g}}{3} u^3 + \frac{\mathbf{g}}{6} u^4 + \frac{\mathbf{g}}{6} u^2 > 0$, $U(u) = \frac{\delta \tilde{E}}{\delta u}$, $V(u) = \frac{\delta E_0}{\delta u}$, $r(t) = \sqrt{\tilde{E}(u)}$ and $m(t) = \sqrt{E_0(u)}$ are two scalar auxiliary variables, $\tilde{u}^{k+1/2}$ is any explicit $O(\tau^2)$ approximation for $u(t^{k+1/2})$.

5) Invariant Energy Quadratization(IEQ) method (first-/second-order) [14]

$$\begin{pmatrix} u_h^{k+1} - u_h^k \\ \tau \end{pmatrix} = -A(\phi, q_h^{k+1}) - (H(u_h^k)U^{k+1}, \phi),$$
$$(q_h^k, \psi) = A(u_h^k, \psi), \qquad \text{for } \forall \phi, \psi \text{ in the space of piecewise polynomials,}$$

where $H(u) = \frac{\Phi'(u)}{\sqrt{\Phi(u) + B}}$ for the constant B > 0, $\Phi = \frac{1}{4}u^4 - \frac{g}{3}u^3 - \frac{\epsilon}{2}u^2$, $U = \sqrt{\Phi(u) + B}$,

besides, $A(\cdot, \cdot)$ is a biliner operator corresponding to the operator $-(\Delta + \frac{a}{2})$. Once the q_h^{k+1} , U^{k+1} and $H(u_h^k)$ are replaced by $(q_h^{k+1} + q_h^k)/2$, $(U^{k+1} + U_h^k)/2$ and $H(u_h^{k,*})$ respectively, with $u_h^{k,*} = \frac{3}{2}u_h^k - \frac{1}{2}u_h^{k-1}$, a second order discretization in time is obtained.

6) Adaptive BDF2 scheme (second-order) [21]

$$\frac{1+2r_k}{\tau_k(1+r_k)}(u^k-u^{k-1}) - \frac{r_k^2}{\tau_k(1+r_k)}(u^{k-1}-u^{k-2}) = -(1+\Delta)^2 u^k - f(u^k)$$

where $\tau_k = t_k - t_{k-1}$ is the time step and $r_k = \tau_k - \tau_{k-1}$. The adaptive time step allows rather larger time step than the uniform time discretizations.

As is seen that most of the existing approaches proposed in the literatures keeps first-/secondorder accurate in time direction. Whereas, we consider a third order time-integration approach, namely the BDF3 formula [31], to solve the Swift-Hohenberg equation for the time discretization with central difference approximation in space. This third-order accurate method has very satisfactory stability properties [32], and was found to be the most efficient in terms of the computational cost for a given accuracy level compared to the lower-order schemes in [33]. However, it is known that the L^2 -norm stability and convergence for the BDF3 formula are difficult to obtain as the scheme is not A-stable. Relying on the equivalence between A-stability and G-stability [34], some useful tools for the numerical analysis of the BDF3 scheme were proposed [35, 36]. Furthermore, Liao et al. [37] achieved a novel yet straightforward discrete energy analysis for the BDF3 schemes using the discrete orthogonal convolution (DOC) kernel technique, and the L^2 -norm stability and convergence of the BDF3 scheme has already been rigorously proved for the linear reaction-diffusion equation.

We focus on developing a fully discrete implicit scheme for the Swift-Hohenberg equation. By virtue of the DOC kernel, the optimal L^2 error estimate is carried out for the proposed scheme, and the scheme is rigorously proved to be third order in time. To the best of our knowledge, this is the first time that the L^2 norm convergence of BDF3 method is obtained for the Swift-Hohenberg equation. Besides, by applying the Brouwer fixed-point theorem and other analytical methods, the unique solvability and the energy stability of the proposed scheme are also provided.

The remainder of the paper is organized as follows. In Section 2, with the introduction of some notations, the construction of the BDF3 difference scheme for the Swift-Hohenberg equation (1.1) is shown in detail. Section 3 presents the results that the proposed scheme is uniquely solvable and maintains the properties of global energy stability. In Section 4, using the DOC kernel proposed in [38], the L^2 -norm convergence of the BDF3 scheme is rigorously proved. Numerical tests of 2D and 3D are performed in Section 5 to verify the accuracy, the stability and the efficiency of the proposed scheme. Some concluding remarks are given in Section 6.

2 Construction of the BDF3 difference scheme

In this section, we set up a fully discrete scheme for the Swift-Hohenberg equation described as follows

$$u_t = -(1 + \Delta)^2 u - f(u), \qquad (x, y) \in \Omega, \ 0 < t \le T, \qquad (2.1)$$

$$u(0, y, t) = u(1, y, t), \ u(x, 0, t) = u(x, 1, t), \qquad (x, y) \in \Omega, \ 0 < t \le T, \qquad (2.2)$$

$$u(x, y, 0) = \varphi_0(x, y), \qquad (x, y) \in \overline{\Omega}, \qquad (2.3)$$

where $\Omega = (0, 1)^2$, $\overline{\Omega} = [0, 1]^2$, the nonlinear term $f(u) = u^3 - gu^2 - \epsilon u$ with two positive physical parameters g and ϵ , and φ_0 is a known smooth function.

Now we consider the difference schemes of (2.1)-(2.3). For the discretization of time direction, the BDF3 formula is applied. Take a positive integer N, and denote $\tau = \frac{T}{N}$, $t_k = \tau k$ for $0 \le k \le N$, then the well-known BDF3 formula can be described as

$$D_3 v^k := \frac{11}{6\tau} \nabla_\tau v^k - \frac{7}{6\tau} \nabla_\tau v^{k-1} + \frac{1}{3\tau} \nabla_\tau v^{k-2}, \qquad \text{for } k \ge 3,$$
(2.4)

where $\nabla_{\tau} v^k := v^k - v^{k-1}$ for any sequence $v^k = v(t_k)$. Referring to [38], the formula (2.4) can be rewritten as a discrete convolution summation,

$$D_3 v^k := \sum_{j=1}^k b_{k-j} \nabla_\tau v^j, \quad k \ge 3,$$
(2.5)

where the convolution kernels b_{k-j} are defined by

$$b_0 := \frac{11}{6\tau}, \quad b_1 := -\frac{7}{6\tau} \quad \text{and} \quad b_2 := \frac{1}{3\tau}, \quad \text{togeter with} \quad b_j := 0, \quad \text{for} \quad 3 \le j \le k - 1.$$
 (2.6)

Since the BDF3 formula requires three previous levels of unknowns, extra schemes should be used to compute u^1 and u^2 . Taylor expansion leads to

$$\frac{1}{2}[u_t(t_1) + u_t(t_0)] = \frac{1}{\tau}[u(t_1) - u(t_0)] + O(\tau^2),$$

thus, a second-order scheme can be obtained

$$D_3 u^1 = \frac{2}{\tau} \nabla_\tau u^1, \tag{2.7}$$

where $u^0 := u(t_0) + \frac{\tau}{2} u_t(t_0)$. Then, the BDF2 formula is used to calculate u^2 as

$$D_3 u^2 = -\frac{1}{2\tau} \nabla_\tau u^1 + \frac{3}{2\tau} \nabla_\tau u^2.$$
 (2.8)

For the spatial direction discretization, let M be a positive integer, the spatial lengths $h_x = h_y = h := L/M$, the grid points $x_i = ih$, $y_j = jh$ and $x_h = (x_i, y_j)$. The discrete

spatial grid $\Omega_h := \{ \mathbf{x}_h = (x_i, y_j) \mid 1 \le i, j \le M - 1 \}$ and $\overline{\Omega}_h := \{ \mathbf{x}_h = (x_i, y_j) \mid 0 \le i, j \le M \}.$ Consider the periodic function space

 $\mathcal{V}_h := \{ v_h = v(\mathbf{x}_h) \mid \mathbf{x}_h \in \overline{\Omega}_h \text{ and } v_h \text{ is periodic in each direction} \}.$

Given a grid function $v \in \mathcal{V}_h$, introduce the following notations $\delta_x v_{i+\frac{1}{2},j} = (v_{i+1,j} - v_{ij})/h$, and $\delta_x^2 v_{ij} = (\delta_x v_{i+\frac{1}{2},j} - \delta_x v_{i-\frac{1}{2},j})/h$. The discrete notations $\delta_y v_{i,j+\frac{1}{2}}$ and $\delta_y^2 v_{ij}$ can be defined similarly. Also, the discrete Laplacian operator $\Delta_h v_{ij} = \delta_x^2 v_{ij} + \delta_y^2 v_{ij}$ and the discrete gradient vector $\nabla_h v_{ij} = (\delta_x v_{ij}, \ \delta_y v_{ij})^T$.

Considering Eq. (2.1) at the grid point (x_h, t_n) , we have

$$u_t(\mathbf{x}_h, t_n) + (1 + \Delta)^2 u(\mathbf{x}_h, t_n) + f(u(\mathbf{x}_h, t_n)) = 0.$$

It follows from (2.5)-(2.8) for the time discretization and the difference formula for the space that the fully discrete scheme is obtained in the following

$$D_{3}u_{h}^{n} + (1 + \Delta_{h})^{2}u_{h}^{n} + f(u_{h}^{n}) = 0 \quad \text{for } \mathbf{x}_{h} \in \Omega_{h}, \ 1 \le n \le N,$$
(2.9)

subjected to the periodic boundary conditions and the following initial condition

$$u_h^0 = \varphi_0(\mathbf{x}_h) + \frac{\tau}{2} \varphi_1(\mathbf{x}_h) \quad \text{for } \mathbf{x}_h \in \bar{\Omega}_h,$$
(2.10)

where $\varphi_1 := f(\varphi_0) + (1 + \Delta_h)^2 \varphi_0$ for the smooth data $\varphi_0 \in H^4(\Omega)$. For any grid functions $v, w \in \mathcal{V}_h$, we define the inner product $\langle v, w \rangle := h^2 \sum_{\mathbf{x}_h \in \Omega_h} v_h w_h$, the

associated L^2 norm $||v||_{l^2} := \sqrt{\langle v, v \rangle}$ and the discrete L^q norm

$$\|v\|_q := \sqrt[q]{h^2 \sum_{\mathbf{x}_h \in \Omega_h} |v_h|^q} \quad \text{for } v \in \mathcal{V}_h$$

To simplify the notation, we write $\|v\| := \|v\|_{l^2}$ and the discrete L^{∞} norm $\|v\|_{\infty} := \max_{\mathbf{x}_h \in \Omega_h} |v_h|$ in the following analysis. The discrete Green's formula with periodic boundary conditions yields $\langle -\Delta_h v, w \rangle = \langle \nabla_h v, \nabla_h w \rangle$ and $\langle \Delta_h^2 v, w \rangle = \langle \Delta_h v, \Delta_h w \rangle$.

Solvability and the energy dissipation law 3

In this section, we demonstrate the unique solvability of the BDF3 implicit scheme (2.9) based on the Brouwer fixed-point theorem, and then, using the positive definiteness of the convolution coefficients, the energy stability is proved in detail, which further deduces the boundedness of the numerical solution in L^{∞} norm.

3.1Unique solvability

For simplicity, we introduce notations that $b_0^{(1)} = \frac{2}{\tau}, b_0^{(2)} = \frac{3}{2\tau}$ and $b_0^{(n)} = b_0$, for any fixed index n, define the map $\Pi_n: \mathcal{V}_h \to \mathcal{V}_h$ as follows

$$\Pi_n(w_h) := b_0^{(n)} w_h - g_h^{n-1} + (1 + \Delta_h)^2 w_h + f(w_h), \quad \mathbf{x}_h \in \bar{\Omega}_h, \quad \text{for} \quad n \ge 1,$$
(3.1)

where $g_h^{n-1} = b_0 u_h^{n-1} + b_1 \nabla_\tau u_h^{n-1} - b_2 \nabla_\tau u_h^{n-2}$ for $n \ge 3$, $g_h^1 = \frac{3}{2\tau} u_h^1 + \frac{1}{2\tau} \nabla_\tau u_h^1$, and $g_h^0 = \frac{2}{\tau} \varphi_0 + \varphi_1$. It is easily seen that the equation $\Pi_n(u^n) = 0$ is equivalent to the proposed nonlinear implicit

scheme (2.9). Thus, the solvability of BDF3 scheme (2.9) can be verified via the equation $\Pi_n(w_h) = 0$, $n \ge 1$ in the following theorem.

Theorem 3.1 Suppose the time step τ holds the condition that $\tau < \frac{3}{2(g^2+\epsilon)}$, the difference scheme (2.9) is uniquely solvable.

Proof Firstly, we prove the existence of the solution.

Suppose u^{n-1} , u^{n-2} have been determined, taking the inner product of (3.1) with w, it yields

$$\langle \Pi_n(w), w \rangle = b_0^{(n)} \langle w, w \rangle + \left\langle (1 + \Delta_h)^2 w, w \right\rangle + \left\langle f(w), w \right\rangle - \left\langle g^{n-1}, w \right\rangle, \text{ for } n \ge 1.$$

For the second term, using discrete Green's formula, we obtain

$$\langle (1+\Delta_h)^2 w, w \rangle = \langle (1+\Delta_h)w, (1+\Delta_h)w \rangle \ge 0.$$

Then, it follows that

$$\begin{split} \langle \Pi_n(w), w \rangle &\geq b_0^{(n)} \|w\|^2 + \|w\|_4^4 - g\left\langle w^2, w \right\rangle - \epsilon \|w\|^2 - \|g^{n-1}\| \cdot \|w\| \\ &\geq \left(b_0^{(n)} - \epsilon\right) \|w\|^2 + \|w\|_4^4 - \left(\frac{g^2}{4}\|w\|^2 + \|w\|_4^4\right) - \|g^{n-1}\| \cdot \|w\| \\ &= \left(b_0^{(n)} - \epsilon - \frac{g^2}{4}\right) \|w\|^2 - \|g^{n-1}\| \cdot \|w\|. \end{split}$$

Evidently, it follows from the condition $\tau < \frac{3}{2(g^2 + \epsilon)}$ that

 $\langle \Pi_n(w), w \rangle \ge 0,$

with $||w|| = \frac{1}{b_0^{(n)} - \epsilon - \frac{g^2}{4}} ||g^{n-1}||$. By means of the well-known Brouwer fixed-point theorem, there exists a w^* such that

$$\Pi_n(w^*) = 0,$$

which implies the BDF3 scheme (2.9) is solvable.

Then, we are prepared to prove the uniqueness of the solution. Suppose w_h and v_h are the solutions of the difference scheme (2.9). Denote $\zeta_h = w_h - v_h$, we start from the following equation with respect to ζ_h ,

$$b_0^{(n)}\zeta_h + (1+\Delta_h)^2\zeta_h + f(w_h) - f(v_h) = 0.$$
(3.2)

Taking the inner product of (3.2) with ζ , we have

$$b_0^{(n)} \|\zeta\|^2 + \|(1+\Delta_h)\zeta\|^2 + \langle f(w) - f(v), \zeta \rangle = 0.$$
(3.3)

For the nonlinear term in (3.3), the further estimation reads

 $\langle f(w) - f(v), \zeta \rangle = \langle w^3 - v^3, \zeta \rangle - g \langle w^2 - v^2, \zeta \rangle - \epsilon \|\zeta\|^2$

$$= \langle w^2 + wv + v^2 - gw - gv, \zeta^2 \rangle - \epsilon \|\zeta\|^2$$

$$\geq \frac{1}{2} \langle (w - g)^2 + (v - g)^2, \zeta^2 \rangle - g^2 \|\zeta\|^2 - \epsilon \|\zeta\|^2$$

Substituting the above inequality into (3.3), it yields

$$(b_0^{(n)} - g^2 - \epsilon) \|\zeta\|^2 \leq 0.$$

From $b_0^{(n)} > \mathbf{g}^2 + \epsilon$, it follows that

 $\|\zeta\|=0,$

which implies that the scheme (2.9) has a unique solution.

3.2 Energy dissipation law

In order to show the energy stability of the proposed scheme (2.9), the following lemmas are needed.

Lemma 3.1 [21] For any grid function $v \in \mathcal{V}_h$, it has

$$\|v\|_{\infty} \le C_{\Omega} \big(\|v\| + \|(1 + \Delta_h)v\| \big), \tag{3.4}$$

where \tilde{C}_{Ω} is a constant depending on the size of space domain Ω but independent of the grid size.

Lemma 3.2 For any real sequence $\{w_k\}_{k=3}^n$ with n-3 entries, it holds that

$$6\tau w_n \sum_{j=3}^n b_{n-j} w_j = \frac{9}{2} (w_n^2 + \frac{2}{9} w_{n-1}^2) - \frac{9}{2} \left(w_{n-1}^2 + \frac{2}{9} w_{n-2}^2 \right) + 2w_n^2 + (w_n + w_{n-2})^2 \qquad (3.5)$$
$$+ \frac{7}{2} (w_n - w_{n-1})^2, \qquad for \ n \ge 3.$$

Thus, the following discrete convolution kernels b_{n-k} are positive definite,

$$2\sum_{k=3}^{n} w_k \sum_{j=3}^{k} b_{k-j} w_j \ge 0.$$
(3.6)

Proof With the definition of b_{n-k} in (2.6), we begin with a decomposition

$$6\tau w_n \sum_{j=3}^n b_{n-j}^{(n)} w_j = w_n (11w_n - 7w_{n-1} + 2w_{n-2}),$$

$$= \frac{9}{2} (w_n^2 + \frac{2}{9}w_{n-1}^2) - \frac{9}{2} \left(w_{n-1}^2 + \frac{2}{9}w_{n-2}^2 \right) + 2w_n^2 + (w_n + w_{n-2})^2 + \frac{7}{2} (w_n - w_{n-1})^2.$$

Making use of (3.5), we are able to prove the positive definiteness as follows

$$6\tau \sum_{k=3}^{n} w_k \sum_{j=3}^{k} b_{k-j} w_j = 6\tau \left(\sum_{k=5}^{n} w_k \sum_{j=3}^{k} b_{k-j} w_j + w_4 (b_1 w_3 + b_0 w_4) + b_0 w_3^2 \right)$$

$$\geq \frac{9}{2} (w_n^2 + \frac{2}{9} w_{n-1}^2) - \frac{9}{2} (w_4^2 + \frac{2}{9} w_3^2) - 7w_3 w_4 + 11 w_4^2 + 11 w_3^2$$

$$= \frac{9}{2} (w_n^2 + \frac{2}{9} w_{n-1}^2) - 7w_3 w_4 + \frac{13}{2} w_4^2 + 10 w_3^2$$

$$\geq \frac{9}{2} (w_n^2 + \frac{2}{9} w_{n-1}^2) - \frac{7}{2} (w_3^2 + w_4^2) + \frac{13}{2} w_4^2 + 10 w_3^2 \geq 0.$$

Lemma 3.3 [21] For any $a, b \in \mathbb{R}$, the following inequality holds

$$(a^3 - ga^2)(a - b) \ge \frac{1}{4}(a^4 - b^4) - \frac{g}{3}(a^3 - b^3) - \frac{2g^2}{3}(a - b)^2.$$

Now, we are in the position to present the energy stability of the BDF3 formula (2.9). Let $E[u^n]$ be the discrete version of energy functional (1.2)

$$E[u^n] := \frac{1}{2} \left\| (1 + \Delta_h) u^n \right\|^2 + \frac{1}{4} \|u^n\|_4^4 - \frac{g}{3} \left\langle (u^n)^2, u^n \right\rangle - \frac{\epsilon}{2} \|u^n\|^2.$$

Denote the modified discrete energy

$$\mathcal{E}[u^{0}] = E[u^{0}], \ \mathcal{E}[u^{1}] = E[u^{1}] + \frac{3}{4\tau} \|\nabla_{\tau} u^{1}\|^{2},$$

$$\mathcal{E}[u^{n}] = E[u^{n}] + \frac{3}{4\tau} \|\nabla_{\tau} u^{n}\|^{2} + \frac{1}{6\tau} \|\nabla_{\tau} u^{n-1}\|^{2}, \text{ for } n \ge 2,$$

Then, the discrete energy dissipation law with respect to the above modified discrete energies is shown in the following theorem.

Theorem 3.2 Suppose the time step τ satisfies

$$\tau \le \frac{1}{3(\frac{2g^2}{3} + \frac{\epsilon}{2})},\tag{3.7}$$

the solution of the adaptive BDF3 scheme (2.9) satisfies

$$E[u^n] \le \mathcal{E}[u^n] \le \mathcal{E}[u^{n-1}] \le \dots \le \mathcal{E}[u^0] = E[u^0], \quad 1 \le n \le N.$$

Proof Taking the inner product of (2.9) with $\nabla_{\tau} u^n$, it yields

$$\langle D_3 u^n, \nabla_\tau u^n \rangle + \langle (1 + \Delta_h)^2 u^n, \nabla_\tau u^n \rangle + \langle f(u^n), \nabla_\tau u^n \rangle = 0, \quad n \ge 3.$$
(3.8)

According to Lemma (3.2), the first term on the left-hand side can be rewritten as

$$6\tau \langle D_3 u^n, \nabla_\tau u^n \rangle \ge \frac{9}{2} (\|\nabla_\tau u^n\|^2 + \frac{2}{9} \|\nabla_\tau u^{n-1}\|^2) - \frac{9}{2} \left(\|\nabla_\tau u^{n-1}\|^2 + \frac{2}{9} \|\nabla_\tau u^{n-2}\|^2 \right) + 2 \|\nabla_\tau u^n\|^2.$$

For the second term, a direct application of the identity $2a(a-b) = a^2 - b^2 + (a-b)^2$ and the summation by parts gives

$$\langle (1+\Delta_h)^2 u^n, \nabla_\tau u^n \rangle = \langle (1+\Delta_h) u^n, \nabla_\tau (1+\Delta_h) u^n \rangle$$

= $\frac{1}{2} \Big(\| (1+\Delta_h) u^n \|^2 - \| (1+\Delta_h) u^{n-1} \|^2 + \| \nabla_\tau ((1+\Delta_h) u^n) \|^2 \Big).$

For the nonlinear term in (3.8), with the help of the equality $2a(a-b) = a^2 - b^2 + (a-b)^2$ and Lemma 3.3, it yields

$$\begin{split} \langle f(u^{n}), \nabla_{\tau} u^{n} \rangle &= \langle (u^{n})^{3} - g(u^{n})^{2}, \nabla_{\tau} u^{n} \rangle - \epsilon \langle u^{n}, \nabla_{\tau} u^{n} \rangle \\ &\geq \frac{1}{4} (\|u^{n}\|_{4}^{4} - \|u^{n-1}\|_{4}^{4}) - \frac{g}{3} \Big(\langle (u^{n})^{2}, u^{n} \rangle - \langle (u^{n-1})^{2}, u^{n-1} \rangle \Big) \\ &- \frac{\epsilon}{2} (\|u^{n}\|^{2} - \|u^{n-1}\|^{2}) - \Big(\frac{2g^{2}}{3} + \frac{\epsilon}{2} \Big) \|\nabla_{\tau} u^{n}\|^{2}. \end{split}$$

As a consequence, a substitution of the above three inqualities into (3.8), results in

$$\left(\frac{1}{3\tau} - \frac{2g^2}{3} - \frac{\epsilon}{2}\right) \|\nabla_{\tau} u^n\|^2 + \mathcal{E}[u^n] - \mathcal{E}[u^{n-1}] \le 0.$$

Under the condition (3.7), it implies that $\mathcal{E}[u^n] \leq \mathcal{E}[u^{n-1}]$ with $3 \leq n \leq N$.

For n = 2, applying Cauchy-Schwarz inequality and the condition (3.7), we obtain

$$\begin{split} \left\langle D_{3}u^{2}, \nabla_{\tau}u^{2}\right\rangle &= \frac{3}{2\tau} \|\nabla_{\tau}u^{2}\|^{2} - \frac{1}{2\tau} \left\langle \nabla_{\tau}u^{1}, \nabla_{\tau}u^{2} \right\rangle \\ &\geq \frac{3}{2\tau} \|\nabla_{\tau}u^{2}\|^{2} - \frac{1}{2\tau} \left(\frac{7}{6} \|\nabla_{\tau}u^{1}\|^{2} + \frac{3}{14} \|\nabla_{\tau}u^{2}\|^{2}\right) \\ &= \frac{3}{4\tau} \left(\|\nabla_{\tau}u^{2}\|^{2} + \frac{2}{9} \|\nabla_{\tau}u^{1}\|^{2} \right) - \frac{3}{4\tau} \|\nabla_{\tau}u^{1}\|^{2} + \frac{9}{14\tau} \|\nabla_{\tau}u^{2}\|^{2} \end{split}$$

Obviously, it yields that $\mathcal{E}[u^2] \leq \mathcal{E}[u^1]$. For n = 1, using the condition (3.7), it has

$$\langle D_3 u^1, \nabla_\tau u^1 \rangle = \frac{3}{4\tau} \| \nabla_\tau u^1 \|^2 + \frac{5}{4\tau} \| \nabla_\tau u^1 \|^2 = \frac{2}{\tau} \| \nabla_\tau u^1 \|^2,$$

which can easily deduce $\mathcal{E}[u^1] \leq \mathcal{E}[u^0]$. It completes the proof.

It follows from Theorem 3.2 that the solution of BDF3 scheme (2.9) is bounded. We further state the result in the following lemma.

Lemma 3.4 Assume the time step τ satisfies the condition (3.7). The solution of BDF3 scheme (2.9) is stable in the L^{∞} norm with $||u^n||_{\infty} \leq c_0$ for $n \geq 1$, where c_0 is independent of the time-step sizes τ .

Proof Utilizing Theorem 3.2 and Young inequality, it follows from $E[u^n] \leq E[u^0]$ that

$$4E[u^0] \ge 2\|(1+\Delta_h)u^n\|^2 + \|u^n\|_4^4 - \frac{4g}{3}\left\langle (u^n)^2, u^n \right\rangle - 2\epsilon \|u^n\|^2$$

$$\geq 2 \| (1 + \Delta_h) u^n \|^2 + \| u^n \|_4^4 - \frac{2}{9} \| u^n \|_4^4 - 2g^2 \| u^n \|^2 - 2\epsilon \| u^n \|^2 \\ \geq 2 \| (1 + \Delta_h) u^n \|^2 + \frac{7}{9} \| u^n \|_4^4 - 2(g^2 + \epsilon) \| u^n \|^2, \quad n \geq 1.$$
(3.9)

For any real value $a \in R$, noticing $\left[a^2 - \frac{9}{7}(1 + \epsilon + g^2)\right]^2 \ge 0$, it yields

$$||u^{n}||_{4}^{4} \ge \frac{18}{7}(1+\epsilon+g^{2})||u^{n}||^{2} - \frac{81}{49}(1+\epsilon+g^{2})^{2}|\Omega_{h}|.$$

Substituting the above inequality into (3.9), it follows that

$$4E[u^{0}] \geq 2\|(1+\Delta_{h})u^{n}\|^{2} + 2\|u^{n}\|^{2} - \frac{9}{7}(1+\epsilon+g^{2})^{2}|\Omega_{h}|$$
$$\geq (\|(1+\Delta_{h})u^{n}\| + \|u^{n}\|)^{2} - \frac{9}{7}(1+\epsilon+g^{2})^{2}|\Omega_{h}|.$$

By Lemma 3.1, we have

$$\|u^{n}\|_{\infty} \leq \tilde{C}_{\Omega}(\|(1+\Delta_{h})u^{n}\|+\|u^{n}\|)$$

$$\leq \tilde{C}_{\Omega}\sqrt{4E[u^{0}]+\frac{9}{7}(1+\epsilon+g^{2})^{2}|\Omega_{h}|}$$

$$:= c_{0}.$$

4 L^2 norm error estimate

In this section, we investigate the error estimates for the difference scheme(2.9) and prove the L^2 norm covergence of the scheme.

4.1 the DOC kernels

To establish the L^2 norm estimate, we introduce the DOC kernels ϑ_{n-k} with

$$\vartheta_0 := \frac{1}{b_0} \quad \text{and} \quad \vartheta_{n-k} := -\frac{1}{b_0} \sum_{j=k+1}^n \vartheta_{n-j} b_{j-k}, \quad \text{for } 3 \le k \le n-1,$$
(4.1)

where $b_0 = \frac{11}{6\tau}$, $b_1 = -\frac{7}{6\tau}$, $b_2 = \frac{1}{3\tau}$, and $b_j = 0$, for $3 \le j \le n-3$. In general, assume the summation $\sum_{k=i}^{j}$, i > j to be zero. Obviously, rewritting the (4.1), a discrete orthogonal identity can be obtained

$$\sum_{j=k}^{n} \vartheta_{n-j} b_{j-k} \equiv \delta_{nk}, \ 3 \le k \le n.$$
(4.2)

The following lemma presents the properties of the DOC kernels ϑ_{n-k} , which plays a key role in the convergence analysis of the scheme (2.9).

Lemma 4.1 (I) For any sequences $\{w_k\}_{k=3}^n$, we have

$$\sum_{k=3}^{n} w_k \sum_{j=3}^{k} \vartheta_{k-j} w_j \ge 0, \qquad \text{for } 3 \le n.$$

$$(4.3)$$

(II) The DOC kernels ϑ_{n-k} in (4.1) have an explicit formula

$$\vartheta_{n-k} = \frac{6\tau}{11} \left[\frac{39 + 7\sqrt{39}i}{78} \left(\frac{7 - \sqrt{39}i}{22} \right)^{n-k} + \frac{39 - 7\sqrt{39}i}{78} \left(\frac{7 + \sqrt{39}i}{22} \right)^{n-k} \right], \quad (4.4)$$

and the following bounds

$$|\vartheta_{n-k}| \le (\frac{2}{11})^{\frac{n-k}{2}}\tau, \quad and \quad \sum_{k=3}^{n} |\vartheta_{n-k}| \le \frac{22}{9}\tau, \quad for \ 3 \le k \le n.$$
 (4.5)

Proof (1) Based on any given real sequence $\{w_k\}_{k=3}^n$, the sequence $\{v_k\}_{k=3}^n$ can be defined using the discrete BDF3 kernels b_{n-k}

$$v_k = -\frac{1}{b_0} \sum_{j=3}^{k-1} b_{k-j} w_j + \frac{w_k}{b_0}, \quad \text{for} \quad k \ge 3,$$

which is equivalent to

$$w_k = \sum_{j=3}^k b_{k-j} v_j, \quad \text{for} \quad k \ge 3.$$
 (4.6)

Multiplying the DOC kernels ϑ_{n-k} on both sides of the formula(4.6) and summing k form 3 to n, it yields

$$\sum_{k=3}^{n} \vartheta_{n-k} w_k = \sum_{k=3}^{n} \vartheta_{n-k} \sum_{j=3}^{k} b_{k-j} v_j = \sum_{j=3}^{n} v_j \sum_{k=j}^{n} \vartheta_{n-k} b_{k-j} = v_n, \quad \text{for} \quad n \ge 3.$$
(4.7)

In turn, a combination of (4.6) and (4.7) leads to the proof of (4.3)

$$\sum_{k=3}^{n} w_k \sum_{j=3}^{k} \vartheta_{k-j} w_j = \sum_{k=3}^{n} v_k \sum_{j=3}^{k} b_{k-j} v_j \ge 0, \quad \text{for} \quad n \ge 3,$$

in which Lemma 3.2 is applied.(2) Identity (4.2) indicates that

when n = k, $\vartheta_0 b_0 = 1$, for $k \ge 3$, (4.8)

when
$$n = k + 1$$
, $\vartheta_1 b_0 + \vartheta_0 b_1 = 0$, for $k \ge 3$, (4.9)

when
$$n \ge k+2$$
, $\vartheta_{n-k}b_0 + \vartheta_{n-k-1}b_1 + \vartheta_{n-k-2}b_2 = 0$, for $k \ge 3$. (4.10)

Substituting BDF3 kernels b_{n-k} into (4.8)-(4.10), the recursive formula becomes available

$$11\vartheta_{n-k} - 7\vartheta_{n-k-1} + 2\vartheta_{n-k-2} = 0, \quad \text{for } 3 \le k \le n-2, \tag{4.11}$$

where $\vartheta_0 = \frac{6\tau}{11}$ and $\vartheta_1 = \frac{42\tau}{121}$. The explicit formula of $\{\vartheta_{n-k}\}_{k=3}^n$ is obtained by the characteristic equation method

$$\vartheta_{n-k} = \frac{\vartheta_1 - p\vartheta_0}{q - p}q^{n-k} + \frac{\vartheta_1 - q\vartheta_0}{p - q}p^{n-k}, \qquad \text{for} \quad 3 \le k \le n, \tag{4.12}$$

where p and q are two distinct roots of the characteristic equation $11\lambda^2 - 7\lambda + 2 = 0$. Furthermore, taking absolute values for ϑ_{n-k} , it leads to

$$\begin{aligned} |\vartheta_{n-k}| &= \left| \frac{6\tau}{11} \left[\frac{39 + 7\sqrt{39}i}{78} \left(\frac{7 - \sqrt{39}i}{22} \right)^{n-k} + \frac{39 - 7\sqrt{39}i}{78} \left(\frac{7 + \sqrt{39}i}{22} \right)^{n-k} \right] \right| \\ &\leq \frac{12\tau}{11} \sqrt{(\frac{39}{78})^2 + (\frac{7\sqrt{39}}{78})^2} \left(\sqrt{(\frac{7}{22})^2 + (\frac{\sqrt{39}}{22})^2} \right)^{n-k} \\ &= \tau \sqrt{\frac{96}{143}} \left(\frac{2}{11} \right)^{\frac{n-k}{2}} \\ &\leq \left(\frac{2}{11} \right)^{\frac{n-k}{2}} \tau, \qquad \text{for} \quad 3 \le k \le n. \end{aligned}$$

Summing k from 3 to n separately for the both sides of the above inequality, it yields

$$\sum_{k=3}^{n} |\vartheta_{n-k}| \le \tau \sum_{k=3}^{n} \left(\frac{2}{11}\right)^{\frac{n-k}{2}} \le \tau \sum_{k=3}^{\infty} \left(\frac{2}{11}\right)^{\frac{n-k}{2}} \le \frac{\tau}{1-\sqrt{\frac{2}{11}}} \le \frac{22}{9}\tau.$$

It completes the proof.

4.2**Convergence** analysis

For the regularity of the exact solution U^n , suppose there exists a constant $c_1 > 0$ such that $\{\|U^n\|_{\infty}, \|\partial_t U^n\|_{\infty}\} \leq c_1$, where c_1 is a constant independent of the spatial step h and the time step τ .

Lemma 4.2 [38] Let $\lambda \ge 0$, the sequences $\{p_k\}_{k=1}^N$ and $\{V_k\}_{k=1}^N$ be nonnegative. If the sequence satisfies

$$V_n \le \lambda \sum_{j=1}^{n-1} \tau_j V_j + \sum_{j=1}^n p_j, \text{ for } 1 \le n \le N,$$

then it holds that

$$V_n \le \exp(\lambda t_{n-1}) \sum_{j=1}^n p_j, \text{ for } 1 \le n \le N$$

Now we are ready to show the convergence analysis of the BDF3 difference scheme (2.9). Denote the error $e_h^n = U_h^n - u_h^n$, $x_h \in \overline{\Omega}_h$, $0 \le n \le N$. The error equations is shown as follows

$$D_3 e_h^n + (1 + \Delta_h)^2 e_h^n + \left(f(U_h^n) - f(u_h^n) \right) = \xi_h^n + \eta_h^n, \quad \mathbf{x}_h \in \bar{\Omega}_h, \ 1 \le n \le N,$$
(4.13)

with $e_h^0 = 0$, $\mathbf{x}_h \in \overline{\Omega}_h$, where ξ_h^n and η_h^n denote the local consistency error in time and space, respectively.

Theorem 4.1 Suppose the problem (1.1) has a unique smooth solution and $u_h^n \in \mathcal{V}_h$ is the solution of the difference scheme (2.9). If the time step τ satisifies the condition that

$$\tau \le \frac{9}{88\rho},\tag{4.14}$$

then, the numerical scheme is convergent in L^2 norm,

 $||e^k|| \le C (\tau^3 + h^2),$

where $\rho = c_1^2 + c_1c_0 + c_0^2 + g(c_0 + c_1) + \epsilon$, C is a positive constant independent of the time steps τ and the space length h.

Proof For n = 1, taking the discrete inner product with (4.13) by e^1 leads to

$$\langle D_3 e^1, e^1 \rangle + \langle (1 + \Delta_h)^2 e^1, e^1 \rangle + \langle (f(U^1) - f(u^1), e^1) \rangle = \langle \xi^1 + \eta^1, e^1 \rangle,$$
 (4.15)

where $\xi_h^1 = O(\tau^2), \ \eta_h^1 = O(h^2).$

For the nonlinear term in (4.15), denote

$$f_h^1 = (U_h^1)^2 + U_h^1 u_h^1 + (u_h^1)^2 - g(U_h^1 + u_h^1) - \epsilon,$$

then it yields that $f(U_h^1) - f(u_h^1) = f_h^1 e_h^1$. It follows from Lemma 3.4 and $||U^n||_{\infty} \leq c_1$ that

 $\|f^1\|_{\infty} < \rho,$

where $\rho = c_1^2 + c_1 c_0 + c_0^2 + g(c_0 + c_1) + \epsilon$. Noticing $\langle (1 + \Delta_h)^2 e^1, e^1 \rangle = ||(1 + \Delta)e^1||^2 \ge 0$ with $e^0 = 0$, and using Cauchy-Schwarz inequality, (4.15) can be rewritten as

$$\frac{2}{\tau} \|e^1\|^2 - \rho \|e^1\|^2 \le (\|\xi^1\| + \|\eta^1\|) \|e^1\|,$$

that is

$$(2 - \rho\tau) \|e^1\| \le \tau(\|\xi^1\| + \|\eta^1\|).$$

In view of the condition (4.14), it yields

$$||e^1|| \le C_1(\tau^3 + h^2), \tag{4.16}$$

where C_1 is a positive constant independent of τ and h.

While for n = 2, taking the inner product of (4.13) with e^2 , we obtain

$$\langle D_3 e^2, e^2 \rangle + \langle (1+\Delta)^2 e^2, e^2 \rangle + \langle (f(U^1) - f(u^2), e^2) \rangle = \langle \xi^2 + \eta^2, e^2 \rangle,$$
 (4.17)

where $\xi_h^2 = O(\tau^2)$, $\eta_h^2 = O(h^2)$. An application of Cauchy-Schwarz inequality for the first term on the left hand side directly leads to the following inquality

$$\langle D_3 e^2, e^2 \rangle = \frac{3}{2\tau} \langle e^2, e^2 \rangle - \frac{2}{\tau} \langle e^1, e^2 \rangle \ge \frac{3}{2\tau} ||e^2||^2 - \frac{2}{\tau} ||e^1|| ||e^2||.$$

Similar to the case of n = 1, equation (4.17) can be rewritten as

$$\left(\frac{3}{2} - \rho\tau\right) \|e^2\| \le 2\|e^1\| + \tau(\|\xi^1\| + \|\eta^1\|).$$

According to the condition (4.14), the following convergence estimate is obtained

$$||e^2|| \le C_2(\tau^3 + h^2), \tag{4.18}$$

where C_2 is a positive constant independent of τ and h.

For $n \geq 3$, replacing n by l in (4.13) and multiplying both sides of (4.13) by the DOC kernels ϑ_{k-l} , and summing l from 3 to k, for $\mathbf{x}_h \in \overline{\Omega}_h$, we have the following identity for $k \geq 3$

$$\sum_{l=3}^{k} \vartheta_{k-l} D_3 e_h^l + \sum_{l=3}^{k} \vartheta_{k-l} (1+\Delta_h)^2 e_h^l - \sum_{l=3}^{k} \vartheta_{k-l} \left(f(U_h^l) - f(u_h^l) \right) = P_h^k + Q_h^k, \tag{4.19}$$

where $P_h^k = \sum_{l=3}^k \vartheta_{k-l} \xi_h^l$ and $Q_h^k = \sum_{l=3}^k \vartheta_{k-l} \eta_h^l$. It follows from Lemma 4.1 (II) and $\xi_h^l = O(\tau^3), \eta_h^l = O(h^2)$ that

$$\sum_{k=3}^{n} \|P^{k}\| \le \frac{22}{9} t_{n} \tau^{3} \le C_{3} \tau^{3}, \text{ for } 3 \le n \le N,$$
(4.20)

$$\sum_{k=3}^{n} \|Q^{k}\| \le \frac{22}{9} t_{n} \tau^{3} \le C_{4} h^{2}, \text{ for } 3 \le n \le N,$$
(4.21)

where C_3, C_4 are positive constants independent of τ and h.

Recalling the orthogonal identity (4.2), it yields

$$\begin{split} \sum_{l=3}^{k} \vartheta_{k-l} D_{3} e_{h}^{l} &= \sum_{l=3}^{k} \vartheta_{k-l} \sum_{j=1}^{l} b_{l-j} \nabla_{\tau} e_{h}^{j} = \sum_{l=3}^{k} \vartheta_{k-l} \left(\sum_{j=3}^{l} b_{l-j} \nabla_{\tau} e_{h}^{j} + b_{l-1} \nabla_{\tau} e_{h}^{1} + b_{l-2} \nabla_{\tau} e_{h}^{2} \right) \\ &= \sum_{j=3}^{k} \nabla_{\tau} e_{h}^{j} \sum_{l=j}^{k} \vartheta_{k-l} b_{l-j} + \sum_{l=3}^{k} \vartheta_{k-l} b_{l-1} \nabla_{\tau} e_{h}^{1} + \sum_{l=3}^{k} \vartheta_{k-l} b_{l-2} \nabla_{\tau} e_{h}^{2} \\ &= \nabla_{\tau} e_{h}^{k} + \sum_{l=3}^{k} \vartheta_{k-l} b_{l-1} \nabla_{\tau} e_{h}^{1} + \sum_{l=3}^{k} \vartheta_{k-l} b_{l-2} \nabla_{\tau} e_{h}^{2}, \quad \text{ for } 3 \le k \le n, \end{split}$$

where the orthogonal identy (4.2) is applied.

Taking the inner product of (4.19) with e^k and summing up k from 3 to n, it yields

$$\sum_{k=3}^{n} \langle \nabla_{\tau} e^{k}, e^{k} \rangle + \sum_{k=3}^{n} \sum_{l=3}^{k} \vartheta_{k-l}^{(k)} \langle (1+\Delta_{h})^{2} e^{l}, e^{k} \rangle + \sum_{k=3}^{n} \sum_{l=3}^{k} \vartheta_{k-l}^{(k)} \langle f(U^{l}) - f(u^{l}), e^{k} \rangle$$
$$= -\sum_{k=3}^{n} \sum_{l=3}^{k} \vartheta_{k-l} b_{l-1} \langle \nabla_{\tau} e^{1}, e^{k} \rangle - \sum_{k=3}^{n} \sum_{l=3}^{k} \vartheta_{k-l} b_{l-2} \langle \nabla_{\tau} e^{2}, e^{k} \rangle + \sum_{k=3}^{n} \langle P^{k} + Q^{k}, e^{k} \rangle.$$
(4.22)

In the following, each term in (4.22) will be estimated rigorously. For the first term on the left hand side, the following identity holds

$$\sum_{k=3}^{n} \left\langle \nabla_{\tau} e^{k}, e^{k} \right\rangle = \frac{1}{2} (\|e^{n}\|^{2} - \|e^{2}\|^{2} + \sum_{k=3}^{n} \|\nabla_{\tau} e^{k}\|^{2}).$$
(4.23)

For the second one on the left hand side, with the application of the positive definiteness of the discrete kernels ϑ_{n-k} in (4.3), we have

$$\sum_{k=3}^{n} \sum_{l=3}^{k} \vartheta_{k-l} \langle (1+\Delta_h)^2 e^l, e^k \rangle = \sum_{k=3}^{n} \sum_{l=3}^{k} \vartheta_{k-l} \langle (1+\Delta_h) e^l, (1+\Delta_h) e^k \rangle \ge 0.$$
(4.24)

For the third nonlinear term on the left-hand side, we begin with a estimate

$$f(U_h^l) - f(u_h^l) = f_h^l e_h^l,$$

where we denote

$$f_h^l = (U_h^l)^2 + U_h^l u_h^l + (u_h^l)^2 - g(U_h^l + u_h^l) - \epsilon.$$

It follows from Lemma 3.4 that

 $\|f^l\|_{\infty} \le \rho,$

where $\rho = c_1^2 + c_1c_0 + c_0^2 + g(c_0 + c_1) + \epsilon$. Consequently, it yields

$$\sum_{k=3}^{n} \sum_{l=3}^{k} \vartheta_{k-l} \langle f(U^{l}) - f(u^{l}), e^{k} \rangle = \sum_{k=3}^{n} \sum_{l=3}^{k} \vartheta_{k-l} \langle f^{l} e^{l}, e^{k} \rangle \le \rho \sum_{k=3}^{n} \|e^{k}\| \sum_{l=3}^{k} |\vartheta_{k-l}| \|e^{l}\|.$$
(4.25)

Noticing $b_j = 0$ when $j \ge 3$, the first term on the right hand side of (4.22) gives

$$-\sum_{k=3}^{n}\sum_{l=3}^{k}\vartheta_{k-l}b_{l-1}\langle\nabla_{\tau}e^{1},e^{k}\rangle = -\sum_{k=3}^{n}\vartheta_{k-3}b_{2}\langle\nabla_{\tau}e^{1},e^{k}\rangle \le \frac{1}{3\tau}\sum_{k=3}^{n}|\vartheta_{k-3}|\|\nabla_{\tau}e^{1}\|\|e^{k}\|, \quad (4.26)$$

where Cauchy-Schwarz inequality is used. Similarly, the second term on the right hand side of (4.22) can be estimated as

$$-\sum_{k=3}^{n}\sum_{l=3}^{k}\vartheta_{k-l}b_{l-2}\langle\nabla_{\tau}e^{2},e^{k}\rangle = -\sum_{k=3}^{n}\vartheta_{k-3}b_{1}\langle\nabla_{\tau}e^{2},e^{k}\rangle - \sum_{k=4}^{n}\vartheta_{k-4}b_{2}\langle\nabla_{\tau}e^{2},e^{k}\rangle$$

$$\leq \frac{7}{6\tau} \sum_{k=3}^{n} |\vartheta_{k-3}| \|\nabla_{\tau} e^2\| \|e^k\| + \frac{1}{3\tau} \sum_{k=4}^{n} |\vartheta_{k-4}| \|\nabla_{\tau} e^2\| \|e^k\|.$$
(4.27)

Substituting (4.23)-(4.27) into (4.22), one arrives at

$$\|e^{n}\|^{2} - \|e^{2}\|^{2} \leq 2\rho \sum_{k=3}^{n} \|e^{k}\| \sum_{l=3}^{k} |\vartheta_{k-l}| \|e^{l}\| + \frac{2}{3\tau} \sum_{k=3}^{n} |\vartheta_{k-3}| \|\nabla_{\tau}e^{1}\| \|e^{k}\| + \frac{7}{3\tau} \sum_{k=3}^{n} |\vartheta_{k-3}| \|\nabla_{\tau}e^{2}\| \|e^{k}\| + \frac{2}{3\tau} \sum_{k=4}^{n} |\vartheta_{k-4}| \|\nabla_{\tau}e^{2}\| \|e^{k}\| + 2\sum_{k=3}^{n} \|e^{k}\| (\|P^{k}\| + \|Q^{k}\|), \quad (4.28)$$

where the last item is obtained by Cauchy-Schwarz inequality.

There exists an integer n_0 such that $||e^{n_0}|| = \max_{0 \le k \le n} ||e^k||$. Taking $n = n_0$ in (4.28), we have

$$\begin{aligned} \|e^{n_0}\| &\leq \|e^2\| + 2\rho \sum_{k=3}^{n_0} \|e^k\| \sum_{l=3}^k |\vartheta_{k-l}| + \frac{2}{3\tau} \sum_{k=3}^{n_0} |\vartheta_{k-3}| \|\nabla_{\tau} e^1\| \\ &+ \frac{7}{3\tau} \sum_{k=3}^{n_0} |\vartheta_{k-3}| \|\nabla_{\tau} e^2\| + \frac{2}{3\tau} \sum_{k=4}^{n_0} |\vartheta_{k-4}| \|\nabla_{\tau} e^2\| + 2 \sum_{k=3}^{n_0} (\|P^k\| + \|Q^k\|), \end{aligned}$$

When $\tau \leq \frac{9}{88\rho}$, it follows from Lemma 4.1 (II) that

$$\|e^{n}\| \leq \|e^{n_{0}}\| \leq 2\|e^{2}\| + \frac{88}{9}\rho\tau \sum_{k=3}^{n-1} \|e^{k}\| + \frac{88}{27}\|\nabla_{\tau}e^{1}\| + \frac{44}{3}\|\nabla_{\tau}e^{2}\| + 4\sum_{k=3}^{n}(\|P^{k}\| + \|Q^{k}\|), \quad 1 \leq n \leq N.$$

$$(4.29)$$

Applying Lemma 4.2, and combining the estimates (4.16) and (4.18) leads to the following overall estimation

$$\begin{aligned} \|e^{n}\| &\leq 2\exp(\frac{88}{9}\rho t_{n-1}) \Big(\|e^{2}\| + \frac{44}{27} \|\nabla_{\tau}e^{1}\| + \frac{22}{3} \|\nabla_{\tau}e^{2}\| + 2\sum_{k=3}^{n} (\|P^{k}\| + \|Q^{k}\|) \Big) \\ &\leq 2\exp(\frac{88}{9}\rho t_{n-1}) \Big(\|e^{2}\| + \frac{44}{27} \|e^{1}\| + \frac{22}{3}\sqrt{2}(\|e^{2}\| + \|e^{1}\|) + 2C_{3}\tau^{3} + 2C_{4}h^{2} \Big) \\ &\leq C(\tau^{3} + h^{2}), \qquad 3 \leq n \leq N. \end{aligned}$$

where C is a positive constant independent of the time step τ and the space length h.

5 Numerical results

In this section, numerical examples in 2D and 3D are used to verify the convergence order in time direction, global energy stability and evolution properties of the numerical solution. The L^2 norm error between the exact solution U_h^N and the numerical solution u_h^N is denoted by $e(N) := ||U^N - u^N||$, and the order of convergence in temporal direction is estimated by order $= \log(e(N)/e(2N))$. **Example 5.1** (Temporal Accuracy Test) Consider the Swift-Hohenberg equation $u_t + (1+\Delta)^2 u + f(u) = g(x, y, t), (x, y) \in \Omega, \ 0 < t \leq T$, in the two-dimensional domain $\Omega = (0, 2\pi)^2$ with T = 10, subjected to the periodic boundary conditions and the initial data

$$u(x, y, 0) = \sin(2x)\sin(2y), \quad (x, y) \in \Omega,$$

such that it has an exact solution

$$u(x, y, t) = \cos(t)\sin(2x)\sin(2y).$$

This example is also used in [21] for the numerical test. The spatial grid points are fixed to be M=10000, and the temporal steps are varied by N=10,20,30,40. Model parameters g and ϵ are specified in different cases that $(1)\epsilon = 0.25$, g = 1, $(2)\epsilon = 0.15$, g = 1 and $(3)\epsilon = 0.25$, g = 0to test the convergence of the method under different conditions. The experimental results are displayed in Table 1, which shows the order of convergence in time direction matches well with the theoretical analysis in Section 4.

Table 1: L^2 errors and convergence orders of BDF3 scheme (2.9) with different ϵ and g.

	$\epsilon = 0.25, \ g = 1$		$\epsilon = 0.15, g = 1$		$\epsilon = 0.25, g = 0$	
Ν	e(N)	Order	e(N)	Order	e(N)	Order
10	3.167 e-01	*	3.154e-01	*	3.127e-01	*
20	5.699e-02	2.47	5.647 e-02	2.48	5.315e-02	2.56
40	7.321e-03	2.96	7.246e-03	2.96	6.625 e- 03	3.00
80	9.311e-04	2.98	9.309e-04	2.96	8.443e-04	2.97

Example 5.2 (Evolution of energy and solution in 2D) We consider the Swift-Hohenberg equation (1.1) on $\Omega = (0, 100)^2$ with different parameters ϵ and g, subjected to the periodic boundary conditions and an initial data [?]

$$u(x, y, 0) = 0.1 + 0.02 \cos\left(\frac{\pi x}{100}\right) \sin\left(\frac{\pi y}{100}\right) + 0.05 \sin\left(\frac{\pi x}{20}\right) \cos\left(\frac{\pi y}{20}\right), \qquad (x, y) \in \Omega.$$

In this example, we will show the evolution of both the energy and the numerical solution based on spatial meshes 128×128 with $\tau = 0.1$. The energy evolution in time with T = 50for different parameters ϵ and g are shown in Figure 1, from which we see that scheme (2.9) is always energy dissipating for any parameters as tested. What's more, the values of ϵ and g appear to manipulate the decay rate of the energy.

The parameters ϵ and g have great impact on the dissipation property according to Figure 1. For fixed $\tau = 0.1$ and $\epsilon = 0.25$, Figure 2 and Figure 3 show the numerical solutions before stabilization at different time t = 0.1, 16, 32, 64 with parameters g = 0 and g = 1 respectively. We find that the evolved patterns of the numerical solution are effected by the parameter g, which reveals cylindrical patterns when g = 1 and curvilinear patterns when g = 0 at the saturation time.



Figure 1: Evolution of energy with different parameters ϵ and g.



Example 5.3 (Evolution of energy and solution in 3D) In this example, in order to further test the evolutionary properties of the BDF3 scheme (2.9), we consider the Swift-Hohenberg equation (1.1) on three-dimensional space $\Omega = (0, 48)^3$ with $\epsilon = 0.1$, g = 0, subjected to the periodic boundary conditions and random initial condition

$$u(x, y, z, 0) = 0.01 * rand(x, y, z), \quad (x, y, z) \in \Omega,$$

where rand(x,y,z) is a random number between -0.01 and 0.01.

For fixed h = 1, $\tau = 0.01$, $\epsilon = 0.1$, g = 0, the numerical solutions are reported in Figure 4, while the snapshots from t = 0.01 to t = 5000 reveal vividly the formation and evolution of the curvilinear patterns. The pattern evolution looks slow in the beginning, however, we observe that at a certain point, before t = 100 in this case, the states of circular aggregation break up giving way to the curvilinear patterns. A stable curvilinear pattern is taking its shape after

 $t \geq 5000$, and the steady state is approached. The energy evolution in Figure 5 clearly confirms this.



Figure 4: Evolution of u(x, y, t) at t = 0.1, 10, 100, 500, 2000, 5000 with g = 0



Figure 5: Evolution of energy using the BDF3 scheme (2.9).

6 Concluding remarks

In this paper, we presented and analyzed an energy stable, uniquely solvable, and convergent numerical scheme for the Swift-Hohenberg equation. The energy stability and unique solvability

of the fully discrete scheme were derived using the Brouwer fixed-point theorem and some analytical techniques. The L^2 norm convergence was then proved utilizing the DOC kernel. Numerical results showed the convergence order and dissipative properties of the proposed BDF3 scheme in 2D and 3D simulations. The analysis framework developed in this work can be further extended to some other problems with gradient flow structure.

Acknowledgement

We would like to acknowledge support by the National Natural Science Foundation of China (No. 11701081,11861060), the Fundamental Research Funds for the Central Universities, the Jiangsu Provincial Key Laboratory of Networked Collective Intelligence (No. BM2017002), Key Project of Natural Science Foundation of China (No. 61833005) and ZhiShan Youth Scholar Program of SEU, China Postdoctoral Science Foundation (No. 2019M651634), High-level Scientific Research foundation for the introduction of talent of Nanjing Institute of Technology (No. YKL201856).

References

- J. Swift, P. C. Hohenberg, Hydrodynamic fluctuations at the convective instability. Phys. Rev. A, 15 (1977) 319-328.
- [2] S. Li, F. Denner, N. Morgan et al., Transient structures in rupturing thin films: Marangoniinduced symmetry-breaking pattern formation in viscous fluids, Sci. Adv., 6 (2020) eabb0597.
- [3] S. M. Wise, C. Wang and J. S. Lowengrub, An Energy-Stable and Convergent Finite-Difference Scheme for the Phase Field Crystal Equation, SIAM J. Numer. Anal., 47(2009) 2269-2288.
- [4] M. Tlidi, E. Berríos-Caro, D. Pinto-Ramo et al., Interaction between vegetation patches and gaps: A self-organized response to water scarcity, Physica D, 414 (2020) 132708.
- [5] P. Stefanovic, M. Haataja and N. Provatas, Phase field crystal study of deformation and plasticity in nanocrystalline materials, Phys. Rev. E, 80 (2009) 046107.
- [6] R. R. Rosa, J. Pontes, C. I. Christov, F. M. Ramos, C. Rodrigues Neto, E. L. Rempel and D. Walgraef, Gradient pattern analysis of Swift-Hohenberg dynamics: phase disorder characterization, Physica A, 283 (2000) 156-159.
- [7] A. Hutt and A. Longtin and L. Schimansky-Geier, Additive noise-induced Turing transitions in spatial systems with application to neural fields and the Swift-Hohenberg equation, Physica D, 237 (2008) 755-773.
- [8] Y. Choi, T. Ha, J. Han and D. S. Lee, Bifurcation and final patterns of a modified Swift-Hohenberg equation, Discret. Contin. Dyn. Syst.-Ser. B, 22 (2017) 2543.
- [9] L. A. Peletier and V. Rottschäfer, Pattern selection of solutions of the Swift-Hohenberg equation, Physica D, 194 (2004) 95-126.

- [10] Y. X. Shi and M. A. Han, Existence of generalized homoclinic solutions for a modified Swift-Hohenberg equation, Discret. Contin. Dyn. Syst.-Ser. S, 103 (2020) 3189.
- [11] B. S. Lai and L. L. Zhang, Existences of periodic solutions to the generalized Swift-Hohenberg equation on symmetry-breaking model, Appl. Math. Lett., 13 (2020) 106206.
- [12] P. Gao, Averaging principles for the Swift-Hohenberg equation, Commun. Pure Appl. Anal., 19 (2020) 293.
- [13] D. V. Kostin, Initial-boundary value problems for Fuss-Winkler-Zimmermann and Swift-Hohenberg nonlinear equations of 4th order, Matematicki Vesnik, 70 (2018) 26-39.
- [14] H. L. Liu and P. M. Yin, Unconditionally energy stable DG schemes for the Swift-Hohenberg equation, J. Sci. Comput., 81 (2019) 789-819.
- [15] A. F. Sarmiento, L. F. R. Espath, P. Vignal et al., An energy-stable generalized- α method for the Swift-Hohenberg equation, J. Comput. Appl. Math., 344 (2018) 836-851.
- [16] Z. G. Liu, Novel energy stable schemes for Swift-Hohenberg model with quadratic-cubic nonlinearity based on the H^{-1} -gradient flow approach, Numer. Algorithms, 3 (2020)
- [17] M. Dehghan, M. Abbaszadeh, A. Khodadadian et al., proper orthogonal decompositionreduced order method (POD-ROM) for solving generalized Swift-Hohenberg equation, Int. J. Numer. Methods Heat Fluid Flow, 29(2019) 2642-2665.
- [18] C. I. Christov and J. Pontes, Numerical scheme for Swift-Hohenberg equation with strict implementation of Lyapunov functional, Math. Comput. Modelling, 35 (2002) 87-99.
- [19] H. G. Lee, An energy stable method for the Swift-Hohenberg equation with quadratic-cubic nonlinearity, Comput. Meth. Appl. Mech. Eng., 343 (2019) 40-51.
- [20] Z. R. Zhang and Y. Z. Ma, On a Large Time-Stepping Method for the Swift-Hohenberg Equation, Adv. Appl. Math. Mech., 8 (2018) 992-1003.
- [21] H. Sun, X. Zhao, H. Y. Cao et al., Stability and convergence analysis of adaptive BDF2 scheme for the Swift-Hohenberg equation, Communications in Nonlinear Science and Numerical Simulation, 111 (2022) 106412.
- [22] J. Su, W. W. Fang, Q. Yu and Y. B. Li, Numerical simulation of Swift-Hohenberg equation by the fourth-order compact scheme, Comput. and Appl. Math., 38 (2019) 54.
- [23] H. G. Lee, A semi-analytical Fourier spectral method for the Swift-Hohenberg equation, Comput. Math. Appl.,74 (2017) 1885-1896.
- [24] M. Dehghan and M. Abbaszadeh, The meshless local collocation method for solving multidimensional Cahn-Hilliard, Swift-Hohenberg and phase field crystal equations, Eng. Anal. Bound. Elem., 78 (2017) 49-64.
- [25] H. Gomez and X. Nogueira, A new space-time discretization for the Swift-Hohenbergequation that strictly respects the Lyapunov functional, Commun. Nonlinear Sci. Numer. Simul., 17 (2012) 4930-4946.

- [26] H. Q. Wang and L. Yanti, An Efficient Numerical Method for the Quintic Complex Swift-Hohenberg Equation, Numer. Math. Theor. Meth. Appl., 4 (2011) 237-254,
- [27] X. P. Zhao, B. Liu, P. Zhang et al., Fourier spectral method for the modified Swift-Hohenberg equation, Adv. Difference Equ., 2013 (2013) 156.
- [28] H. G. Lee, A new conservative Swift-Hohenberg equation and its mass conservative method, J. Comput. Appl. Math., 375 (2020) 112815.
- [29] J. Y. Wang and S. Y. Zhai, A Fast and Efficient Numerical Algorithm for the Nonlocal Conservative Swift-Hohenberg Equation, Math. Probl. Eng., 2020 (2020) 7012483.
- [30] J. Zhou and X. M. Dai, An energy-stable pseudospectral scheme for Swift-Hohenberg equation with its Lyapunov functional, Therm. Sci., 23 (2019) 975-982
- [31] E. Hairer and K. Wanner, Solving ordinary differential equations: Stiff and differentialalgebraic problems, Heidelberg: Springer Verlag, 1996.
- [32] P. Kim, D. Kim, X. F. Piao and S. Bak, A completely explicit scheme of Cauchy problem in BSLM for solving the Navier-Stokes equations, J. Comput. Phys., 401 (2020) 109028.
- [33] Q. F. Zhang and C. J. Zhang, Block preconditioning strategies for nonlinear viscous wave equations, Appl. Math. Model., 37 (2013) 5801-5813.
- [34] G. Dahlquist, G-stability is equivalent to A-stability, BIT, 18 (1978) 384-401.
- [35] J. Liu, Simple and Efficient ALE Methods with Provable Temporal Accuracy up to Fifth Order for the Stokes Equations on Time Varying Domains, SIAM J. Numer. Anal., 51 (2013) 743-772.
- [36] O. Nevanlinna and F. Odeh, Multiplier techniques for linear multistep methods, Numer. Funct. Anal. Optim., 3 (1981) 377-423.
- [37] H. L. Liao, T. Tang and T. Zhou, A new discrete energy technique for multi-step backward difference formulas, CSIAM Transactions on Applied Mathematics, 3(2) (2022) 318-334.
- [38] H. L. Liao and Z. Zhang, Analysis of adaptive BDF2 scheme for diffusion equations, Math. Comp., 329 (2021) 1207-1226.