Estimation of Long-Range Dependent Models with Missing Data: to Impute or not to Impute?

Guilherme Pumi^{a,*}, Gladys Choque Ulloa^a and Taiane Schaedler Prass^a

Abstract

Among the most important models for long-range dependent time series is the class of ARFIMA(p, d, q) (Autoregressive Fractionally Integrated Moving Average) models. Estimating the long-range dependence parameter d in ARFIMA models is a well-studied problem, but the literature regarding the estimation of d in the presence of missing data is very sparse. There are two basic approaches to dealing with the problem: missing data can be imputed using some plausible method, and then the estimation can proceed as if no data were missing; or we can use a specially tailored methodology to estimate d in the presence of missing data. In this work, we review some of the methods available for both approaches. One of the methods is an adaptation of a certain copula-based method for estimation of long-range dependent univariate time series, for which we provide a rigorous asymptotic theory in the presence of missing data. We also present an extensive Monte Carlo simulation study to compare these methods, considering among 35 different setups to estimate d, under tenths of different scenarios contemplating percentages of missing data ranging from as few as 10% up to 70% and several levels of dependence.

Keywords: Long-range dependence; time series analysis; missing data; semiparametric estimation; copulas.

MSC: 62M10, 62D10, 62F10, 62E20, 60G15.

Statements and Declarations: The authors declare that they have NO affiliations with or involvement in any organization or entity with any financial interests in the subject matter or materials discussed in this manuscript.

1 Introduction

Long-range dependent processes have a long history and through time the subject has evolved into an essential component of time series analysis. We refer the reader to the book by Palma (2007), the compilations by Doukhan et al. (2003) and Robinson (2003), and, for an account of the history and early days' developments, the book by Beran (1994). In this work we are interested in the class of ARFIMA(p, d, q) models introduced by Honsking (1981) and Granger and Joyeux (1980), which is one of the most applied and studied classes of long-range dependent models in the literature. There are numerous estimation procedures for the long-range dependence parameter d, including methods based on state-space representation, spectral density, approximations to the likelihood, wavelets, detrended fluctuation analysis (DFA), and copulas (Hurst, 1951; Geweke and Porter-Hudak, 1983; Peng et al., 1994; Robinson, 1995a,b; Abry and Veitch, 1998; Chan and Palma, 1998; Palma, 2007; Faÿ et al., 2009; Pumi et al., 2023, and references therein). Monte Carlo simulation studies comparing different estimators have also been conducted (Taqqu et al., 1995; Kokoszka and Bhansali, 2001; Reisen et al., 2001; Faÿ et al., 2009; Rea et al., 2013).

^{*}Corresponding author. This Version: January 3, 2025

^aPrograma de Pós-Graduação em Estatística - Universidade Federal do Rio Grande do Sul.

E-mails: guilherme.pumi@ufrgs.br (G. Pumi), gladyschoqueulloa7@gmail.com (G.U. Choque) and ta-ianeprass@ufrgs.br (T.S. Prass)

ORCIDs: 0000-0002-6256-3170 (G. Pumi); 0000-0003-3136-909X (T.S. Prass).

Estimating d in the presence of missing data, on the other hand, is a much less studied problem with sparse literature. There are two approaches to handling missing data in time series. The first one, imputation, is the most widely used. The basic idea is to replace missing data with plausible values, and then proceed with the analysis as if no data were missing. Of course, there are many ways to do this. One of the simplest is to replace the missing values with the sample mean or median calculated for the non-missing cases, especially when in the context of stationary time series. For nonstationary time series, a missing value can be replaced with the average of previous values up to that point, or by calculating the mean locally using a sliding window approach. When there are just a few missing values corresponding to a small percentage of the total sample size, any reasonable imputation method applied to a stationary time series tends to yield good results in the sense that estimated quantities should be close to those obtained if no data were missing, especially for point estimation. However, as the number of missing values increases, the quality of imputation-based estimation becomes poor. Imputation has the advantage of being quick and simple to implement, which adds to its appeal.

Despite their strengths, almost all conventional deterministic imputation methods suffer from three problems in the context of time series. First, variances tend to be underestimated, leading to biases in other parameters (like correlations) that depend on variances. Mean/median substitution, for example, replaces the presumably different missing values with a single value, reducing variance. It also has an effect on stationary distributions, artificially creating a point of mass in an otherwise continuous distribution. The second problem is that imputing an exogenous value into a time series changes the dependence structure in ways difficult to quantify or even understand. This is especially problematic considering that estimation in models of practical interest takes into account the time series' dependence structure. The third problem is equally serious. Standard error calculations presume that all data is accurate. The inherent uncertainty and sampling variability in the imputed values are not taken into account. As a result, standard errors and p-values may be distorted, leading to incorrect confidence intervals and hypothesis testing, as well as potentially misspecified models.

The second method for estimating d in the presence of missing data is to use an estimator that has been modified or designed specifically to handle missing data. An estimator in this group obviously has none of the three problems that imputation methods have since any reasonable estimator in this group should account for the absence of some observations and bypass their implications. The problem is that there are few options in the literature for estimators in that category, and the available options are usually more complex to implement and slower to run when compared to an imputation alternative. In this work we consider 3 different methods in this category. One of the novelty of the paper is to propose an adaptation of the copula-based method for estimation of long-range dependent univariate time series introduced in Pumi et al. (2023) in the context of fully observed time series, to the case were data is missing. For this particular estimator, we develop a rigorous asymptotic theory guaranteeing its consistency and deriving its limiting distribution under mild conditions on the number of missing data.

It is intuitive to expect that, if the percentage of missing data is too high, the methodologies discussed above should break down in the sense that the estimated values are either so biased that they are practically useless, or cannot be computed due to numerical instability. In this direction, an important question is how high the percentage of missing data must be so that we start loosing trust on (or stop being able to compute) a given methodology? Does the strength of the dependence affect this percentage at all? Must we always use an estimator adapted or specially made for missing data, or can we use the much simpler and faster approach of imputation? Is this answer influenced by the percentage of missing and/or strength of the dependence? In this work, we present a Monte Carlo simulation study to shed some light on these questions. We consider three different estimators specially tailored for missing data in different configurations, five of the most commonly used semiparametric estimators in the literature, paired with three different approaches to imputation, under 28 different scenarios of dependence strength and percentage of missing values. We also introduce a random imputation method especially tailored to closely mimic the original variance of the time series without introducing any potential outliers and while taking into account the time series' local dependence structure.

The paper is organized as follows. In the next Section, we introduce the estimators and the imputation methods applied in the simulation and discuss some of their properties. In Section 3 we present an extensive Monte Carlo simulation study to answer the questions posed in the introduction. Section 4 discusses our findings and presents our conclusions. The asymptotic theory of the proposed adaptation of Pumi et al. (2023)'s estimator in the context of missing data is developed in the Appendix. This paper also accompanies supplementary material containing additional simulation results.

2 Framework

We say that $\{Y_t\}_{t\in\mathbb{Z}}$ is an ARFIMA(p, d, q) if it satisfies the difference equation

$$\phi(L)Y_t = \theta(L)(1-L)^{-d}\varepsilon_t, \qquad (2.1)$$

where $\{\varepsilon_t\}$ is a zero mean white noise with $\sigma_{\varepsilon}^2 := \operatorname{Var}(\varepsilon_t) < \infty$, L is the backward shift operator, $\phi(L) := 1 - \phi_1 L - \cdots - \phi_p L^p$ and $\theta(L) := 1 + \theta_1 L + \cdots + \theta_q L^q$ are the AR and MA operators, respectively; $(1-L)^{-d}$ is a fractional differencing operator defined by the binomial expansion

$$(1-L)^{-d} = \sum_{j=0}^{\infty} \eta_j L^j, \quad \text{with} \quad \eta_j := \frac{\Gamma(j+d)}{\Gamma(j+1)\Gamma(d)},$$

for $-1 < d < \frac{1}{2}$. It is usual to assume that ϕ and θ have no common roots and all roots of ϕ lie outside the unitary circle $\{z \in \mathbb{C} : |z| = 1\}$. Under these conditions, there exists a unique stationary solution for (2.1) with autocovariance satisfying

$$\gamma(h) \sim \kappa_d |h|^{2d-1}, \quad \text{with} \quad \kappa_d := \sigma_{\varepsilon}^2 \left| \frac{\theta(1)}{\phi(1)} \right|^2 \frac{\Gamma(1-2d)}{\Gamma(1-d)\Gamma(d)}, \quad \text{as } h \to \infty.$$
 (2.2)

For 0 < d < 1/2, the covariance in (2.2) decays at a hyperbolic rate so that the autocorrelation is not absolutely summable.

In this work, our main interest is estimating the long-range parameter d under the presence of missing values. We assume that missing values occur completely at random, in the sense that the epoch at which a missing value occurs is independent of the time series' past and future values. We also assume, without loss of generality, that the time series' first and last values are never missing. We consider two approaches for dealing with missing values. The first is by employing a method specifically designed to estimate d without the need for missing data completion. Estimators in this class will be called *intrinsic estimators*. The second approach involves using an estimator that requires the time series to have no missing data. We proceed by imputing the missing values and then applying the estimator as if no values were missing. In this work, we shall consider five of the most commonly applied estimators for d, besides the intrinsic estimators, which can also be used when the time series is complete, and three methods for data imputation. In total, 35 different estimation procedures will be compared. In the following sections, we will briefly review each procedure.

2.1 Intrinsic methods

To the best of our knowledge, the earliest intrinsic estimator for d in the presence of missing values is Chan and Palma (1998), which introduces a state-space representation of an ARFIMA model and a modification of the Kalman filter to approximate the likelihood function in the presence of missing data. Although the methodology performed reasonably well in the authors' simulations, it suffers from being very slow, especially when compared to other alternatives, and as a result, it was not included in our simulation study.

In this work, we consider three different intrinsic estimators for d. The first method is an adaptation to the context of missing values of the estimator proposed by Pumi et al. (2023), which is the only copula-based estimator for the long-range parameter d for univariate time series in the literature. In Pumi et al. (2023), the authors derive a novel relationship between the decay of covariance $\gamma(h)$ in a time series $\{X_t\}_{t\in\mathbb{Z}}$ and the behavior of the copulas of pairs (X_0, X_h) as h increases. The relationship is used to propose an estimation procedure for any quantity of interest identifiable through the covariance decay. One special case is the parameter d in view of (2.2). This makes Pumi et al. (2023)'s the only copula-based estimator for the long-range parameter d for univariate time series in the literature. The estimator is defined in the context of complete time series, but, being copula-based, the procedure can be adapted to handle missing data. Besides presenting this adaptation, we take the idea further and derive a rigorous asymptotic theory for the proposed estimator.

Two semiparametric (first and second generation) wavelet-based estimators proposed by Knight et al. (2017) and Craigmile and Mondal (2020) are also considered. They rely on a relationship between the undecimated wavelet variance and the parameter d. The former estimates the wavelet variance using wavelet lifting while the latter uses a specially designed estimator. In the next two sections, we present details regarding these intrinsic estimators.

2.1.1 A copula-based method

In this section, we review the copula-based estimator of d, introduced by Pumi et al. (2023), in the particular case in which it is applied here. In the Appendix A we present the general definition of the estimator, which is also needed for the asymptotic distribution there developed. For a comprehensive introduction to copulas, as well as more details about the theory, we refer the reader to the monographes by Joe (1997) and Nelsen (2013). Let $\{X_t\}_{t\in\mathbb{Z}}$ be a stationary long-range dependent time series of interest and let $\{C_\theta\}_{\theta\in\Theta}$ be a parametric family of copulas such that there exists a point $a \in \Theta$ for which C_a is the independent copula. Suppose that the copula associated to (X_{n_0}, X_{n_0+h}) , for $n_0 \in \mathbb{Z}$ and all h > 0 is C_{θ_h} . In Pumi et al. (2023), the authors uncover a relationship between the behavior of θ_h and the decay of $\gamma(h) = \text{Cov}(X_t, X_{t+h})$ as h increases, and based on this relationship, propose a copula-based estimator for the long-range parameter d. To the best of our knowledge, this is the first (and, as of today, the only one available) copula-based estimator of d in the context of univariate time series in the literature.

More specifically, let x_1, \dots, x_n be a sample from a long-range dependent time series with parameter d and let \hat{F}_n be the empirical distribution calculated from the sample, \hat{F}'_n be an

estimator of the density of X_t (such as a kernel density estimator) and let

$$\hat{K} = \iint_{(0,1)^2} \frac{1}{\hat{F}'_n(\hat{F}_n^{(-1)}(u))\hat{F}'_n(\hat{F}_n^{(-1)}(v))} \lim_{\theta \to a} \frac{\partial C_\theta(u,v)}{\partial \theta} du dv.$$

Let $y_k := F_n^{-1}(x_k), k \in \{1, \dots, n\}$. For $s \in \{1, \dots, n-1\}$, we form a new bivariate time series $\{u_k^{(s)}\}_{k=1}^{n-s}$ by setting $u_i^{(s)} := (y_i, y_{i+s}), i = 1, \dots, n-s$. Observe that $\{u_k^{(s)}\}_{k=1}^{n-s}$ can be regarded as a (correlated) sample from C_{θ_s} , by Sklar's theorem. From these pseudo observations, θ_s can be estimated by a reasonable method, such as the inversion of Kendall's τ or Spearman's ρ , or by maximum pseudo-likelihood. We choose two positive integers 0 < s < m < n and define the estimator of d as

$$\hat{d} := \underset{|d|<0.5}{\operatorname{argmin}} \bigg\{ \sum_{h=s}^{m} \bigg[\hat{K}\hat{\theta}_h - \frac{\Gamma(1-d)}{\Gamma(d)} h^{2d-1} \bigg]^2 \bigg\}.$$
(2.3)

For time series presenting missing data, the estimator is defined by applying the copula estimator only to complete pseudo-observations, which are used to obtain $\{\hat{\theta}_s, \dots, \hat{\theta}_m\}$. Once this sequence is obtained, the estimation procedure remains the same. The asymptotic theory related to the proposed procedure for time series with missing data is presented in the Appendix A, where we show that the limiting distribution and convergence rate are the same as in the complete data case, as long as the information regarding lagged pairs (X_t, X_{t+h}) in the sample increases as n increases.

To apply the estimator (2.3) a few choices need to be made. First, we must choose the parametric family of copulas to apply: a common problem in copula-based methodologies. However, simulation results presented in Pumi et al. (2023) show that the procedure is robust against copula misspecification. It is also necessary to choose estimators for the density, the distribution function, the quantile function, and the estimator for the copula parameter. The authors show that as long as consistent estimators are chosen, under mild smoothness conditions, the methodology yields good results. The authors provide a simulation comparing three types of copula estimators, the methods based on the inversion of Kendall's τ and Spearman's ρ and the pseudo maximum likelihood. The results show little difference between the three. They also provide a comparison between using the correct marginal distribution against using the empirical distribution for \hat{F}_n , concluding that the difference is barely noticeable. As for the values of s and m, the simulation results presented in Pumi et al. (2023) suggest that s = 1 and m = 24 should yield generally good estimates. In the presence of missing data, the estimators of F, F', and F^{-1} are calculated considering only observed values.

2.1.2 Knight et al. (2017)'s LoMPE estimator

A first generation wavelet-based estimator for the long memory parameter d is presented in Knight et al. (2017). The method is based on a multiscale lifting transformation known as LOCAAT (lifting one coefficient at a time) proposed by Jansen et al. (2001). The LoMPE's idea is to apply the LOCAAT method to obtain a collection of lifting coefficients, which, after suitable normalization, are used to estimate the wavelet's coefficient variance. To estimate d, a regression approach similar to (2.4) is applied. Bootstrapped lifting trajectories can be used to improve estimation. The authors present a simulation considering small missing data percentages, from 5% to 20% using 50 bootstrapped trajectories. The method's main problem is its computational cost, which is high compared to other methods considered here. More details can be found in Jansen et al. (2001) and Knight et al. (2017).

2.1.3 Craigmile and Mondal (2020)'s wavelet method

Let $\{X_t\}_{t\in\mathbb{Z}}$ be a stationary Gaussian long-range dependent time series of interest. The idea behind Craigmile and Mondal (2020)'s estimator is based on undecimated wavelet analysis of X_t using the Daubechies' class of wavelets, characterized by the filter width L > 2. A fully detailed explanation of Craigmile and Mondal (2020)'s estimator would be lengthly. However, for completeness, a brief description will be provided here. Let $\{h_{j,l}\}_{j,l}$ for $j \in \{0, 1, \dots\}$ and $l \in \{0, \dots, L-1\}$ be a Daubechies' wavelet filter of even width L. The undecimated wavelet representation of $\{X_t\}_{t\in\mathbb{Z}}$ has coefficients given by

$$W_{j,t} = \sum_{l=0}^{L-1} h_{j,l} X_{t-l}.$$

Since $\{X_t\}_{t\in\mathbb{Z}}$ is a Gaussian process, for each j, $\{W_{j,t}\}_{t\in\mathbb{Z}}$ is a zero mean stationary Gaussian process. Let $v_j^2 := \operatorname{Var}(W_{j,t})$. Craigmile and Mondal (2020) show that, in this case, for $d \in (0, 1/2)$,

$$\log(v_i^2) \approx C + (2d - 1)j\log(2), \tag{2.4}$$

for large j and some constant C. Given an estimator of v_j^2 , \hat{v}_j^2 for $j \in \{j_0, \dots, j_0 + m\}$ for positive j_0 and m, d can be estimated by regressing $\log(\hat{v}_{j_0}^2), \dots, \log(\hat{v}_{j_0+m}^2)$ in j_0, \dots, j_{0+m} plus an intercept. The main contribution in Craigmile and Mondal (2020) is to propose an unbiased, consistent, and asymptotically normally distributed estimator for v_j^2 in longrange dependent processes containing missing data. However, to estimate d from \hat{v}_j^2 , using a regression approach will depend on an unknown dispersion matrix. The authors propose an approximation for this matrix, leading to an estimator of d called the full estimator (called C.full here). A second estimator is also inspired by the work of Abry and Veitch (1998) for complete long-range dependent processes, using only the diagonal elements of the full dispersion matrix to estimate d. The authors called it the diagonal estimator (called C.abry here). We refer the reader to Craigmile and Mondal (2020) for more details.

2.2 Traditional methods

As mentioned in the introduction, there are several estimators to estimate d when the time series is complete. These estimators typically cannot handle missing data naturally. However, they can still be used after the imputation of the missing data. In this Section, we review five well-known semiparametric estimators for d.

2.2.1 Rescaled Range Method (R/S)

Consider a sample $\{y_1, \dots, y_n\}$ from a stationary long-memory process and let $x_t := \sum_{j=1}^t y_j$ for $t \in \{1, \dots, n\}$ be the partial sums of the y_j 's and let $s_n^2 := \frac{1}{n-1} \sum_{t=1}^n (y_t - \overline{y})^2$ be the sample variance, where $\overline{y} = x_n/n$. The rescaled range statistic (R/S), introduced by Hurst (1951), is defined by

$$R_n := \frac{1}{s_n} \left[\max_{1 \le t \le n} \left\{ x_t - \frac{t}{n} x_n \right\} - \min_{1 \le t \le n} \left\{ x_t - \frac{t}{n} x_n \right\} \right].$$

Denoting $Q_n := n^{-\frac{1}{2}-d}R_n$, it can be shown that $\log(R_n) = \mathbb{E}(Q_n) + (d+\frac{1}{2})\log(n) + [\log(Q_n) - \mathbb{E}(Q_n)]$. So that we can obtain an estimator of the long-memory parameter d by least squares. For instance, if $R_{t,k}$ is the R/S statistic based on a sample of size $k, \{y_t, \dots, y_{t+k-1}\}$, for

 $1 \leq t \leq n-k+1$, then an estimator of d can be obtained by regressing $\log(R_{t,k})$ on $\log(k)$ plus an intercept, for $1 \leq t \leq n-k+1$. Some asymptotic results of the R/S statistics are presented in Mandelbrot (1975).

2.2.2 Geweke and Porter-Hudak (1983)'s estimator (GPH)

Let $\{Y_t\}_{t\in\mathbb{Z}}$ be a stationary long-range dependent process with spectral density satisfying

$$f(\lambda) = f_0(\lambda) [2\sin(\lambda/2)]^{-2d}, \qquad (2.5)$$

for some continuous function f_0 . Taking logarithms on both sides of (2.5) evaluated at the Fourier sequences $\lambda_j := 2\pi j/n$, we obtain

$$\log\left(f(\lambda_j)\right) = \log\left(f_0(0)\right) - 2d\log\left(2\sin(\lambda_j/2)\right) + \log\left(\frac{f_0(\lambda_j)}{f_0(0)}\right).$$
(2.6)

From (2.6), Geweke and Porter-Hudak (1983) suggested a regression approach to estimate d. We start by estimating the spectral density using the periodogram, defined by

$$I(\lambda) := \left| \frac{1}{\sqrt{2\pi n}} \sum_{t=1}^{n} e^{it\lambda} Y_t \right|^2.$$
(2.7)

Upon writing

$$\log (I(\lambda_j)) = \log \left(\frac{I(\lambda_j)}{f(\lambda_j)}\right) + \log (f(\lambda_j)),$$

and combining it with (2.6), we obtain

$$\log\left(I(\lambda_j)\right) = \log\left(f_0(0)\right) - 2d\log\left(2\sin(\lambda_j/2)\right) + \log\left(\frac{I(\lambda_j)[2\sin(\lambda/2)]^{2d}}{f_0(0)}\right).$$
(2.8)

From (2.8), we can estimate d considering the first m ordinates of the periodogram by regressing log $(I(\lambda_1)), \dots, \log(I(\lambda_m))$ on $2\log(2\sin(\lambda_1/2)), \dots, 2\log(2\sin(\lambda_m/2))$ plus an intercept. The asymptotic properties of the GPH have been studied by Robinson (1995b) and Hurvich et al. (1998), among others. Under mild conditions, the GPH's estimator is consistent and asymptotically normally distributed with a convergence rate of $n^{4/5}$, independently of d.

2.2.3 Local Whittle estimator (LW)

Let $\{Y_t\}_{t\in\mathbb{Z}}$ be a stationary long-range dependent process with parameter d and spectral density f satisfying

$$f(\lambda) \sim G\lambda^{-2d},$$
 (2.9)

as $\lambda \to 0^+$ with G > 0. For an ARFIMA(p, d, q) process, $G = \left[\frac{\sigma_{\varepsilon}\theta(1)}{2\pi\phi(1)}\right]^2$. The semiparametric estimator, known as the local Whittle estimator, is defined by

$$\hat{d} = \underset{|d|<1/2}{\operatorname{argmin}} \{ R(d) \},$$
 (2.10)

where

$$R(d) := \log\left(\widehat{G}(d)\right) - 2d\frac{1}{m}\sum_{j=1}^{m}\log(\lambda_j), \quad \text{for} \quad \widehat{G}(d) := \frac{1}{m}\sum_{j=1}^{m}\lambda_j^{2d}I(\lambda_j),$$
(2.11)

I is the periodogram defined in (2.7) and 0 < m < n/2 is an integer. Estimator (2.10) was introduced and studied in Robinson (1995a), which proved its consistency and asymptotic normality under mild assumptions, with a \sqrt{m} convergence rate.

2.2.4 Exact Local Whittle estimator (ELW)

The semiparametric estimator introduced by Shimotsu and Phillips (2005), called the exact local Whittle estimator (ELW) is very similar in nature to the local Whittle. The main difference is that the estimator is derived from algebraic manipulation of the Whittle likelihood without relying on approximations of the periodogram, being exact in this sense. The ELW is also applicable to the non-stationary region d > 0.5, but for simplicity, we will consider the range |d| < 0.5. Let

$$I_{\Delta^d}(\lambda) := \left| \frac{1}{\sqrt{2\pi n}} \sum_{t=1}^n e^{it\lambda} (1-L)^d (Y_t) \right|^2.$$

The estimator is defined in the same fashion as the local Whittle, but using $I_{\Delta d}$ instead of I in (2.11), that is

$$\hat{d} = \underset{|d|<1/2}{\operatorname{argmin}} \{ R(d) \},$$

where

$$R(d) := \log\left(\widehat{G}(d)\right) - 2d\frac{1}{m}\sum_{j=1}^{m}\log(\lambda_j), \quad \text{for} \quad \widehat{G}(d) := \frac{1}{m}\sum_{j=1}^{m}\lambda_j^{2d}I_{\Delta^d}(\lambda_j)$$

The asymptotic theory of the ELW is very similar to the local Whittle's but the conditions required on m are slightly stronger. Computationally, according to the authors, the ELW is about 10 times slower than the local Whittle, but it is still a very fast method in comparison to others (see Section 3.5). A discussion of the advantages and disadvantages of the ELW over the GPH and local Whittle as well as more details can be found in Shimotsu and Phillips (2005).

2.2.5 DFA-based estimator

The so-called *detrended fluctuation analysis* (DFA) method was introduced by Peng et al. (1994) and is based on the behavior of the detrended variance in long-range dependent time series. Before proceeding with the definition of the estimator, we introduce some notation, following the more general work of Prass and Pumi (2021).

Let $\{Y_t\}_{t\in\mathbb{Z}}$ be a stationary long-range dependent process with parameter d and let Y_1, \dots, Y_n be a sample from it. Let $R_t := \sum_{j=1}^t Y_j$ for $t \in \{1, \dots, n\}$ denote the integrated signals. Divide the integrated signals into $k = \lfloor n/(m+1) \rfloor$ non-overlapping boxes each containing m + 1 values, which we denote by $\mathbf{R}_i := (R_{(m+1)(i-1)+1}, \dots, R_{i(m+1)})'$, for $i \in \{1, \dots, k\}$. Next, in each box we fit a polynomial of degree $\nu + 1$ via least squares, considering

$$D_{m+1}^{\top} := \begin{pmatrix} 1 & 1 & \dots & 1 \\ 1 & 2 & \dots & m+1 \\ \vdots & \vdots & \ddots & \vdots \\ 1^{\nu+1} & 2^{\nu+1} & \dots & (m+1)^{\nu+1} \end{pmatrix}, \qquad P_{m+1} := D_{m+1} (D_{m+1}^{\top} D_{m+1})^{-1} D_{m+1}^{\top},$$

Define

$$\boldsymbol{\mathcal{E}}_i := Q_{m+1} \boldsymbol{R}_i = (\mathcal{E}_i, \cdots, \mathcal{E}_{m+i})^\top,$$

the vector of residuals at the *i*-th box. Let $f_{DFA}^2(m,i) := \frac{1}{m+1} \mathcal{E}'_i \mathcal{E}_i$, i.e. the sample variance of the residual in the *i*-th fit. The detrended variance F_{DFA}^2 is defined by

$$F_{DFA}^2(m) := \frac{1}{k} \sum_{i=1}^k f_{DFA}^2(m, i).$$

As described by Peng et al. (1994), for a long-range dependent process,

$$\sqrt{F_{DFA}^2(m)} \sim cm^{d+1/2}$$

for large *m*. Consequently, taking logarithms and denoting $L_m := \log \left(\sqrt{F_{DFA}^2(m)} \right)$, we obtain

$$L_m \sim \log(c) + \left(d + \frac{1}{2}\right)\log(m), \qquad (2.12)$$

from which we can estimate d by regressing L_{s+1}, \dots, L_{s+l} in $\log(s+1), \dots, \log(s+l)$, for s, l > 0, plus an intercept. In Bardet and Kammoun (2008), the consistency of the DFA-based estimator for d in the context of fractional Brownian motion is demonstrated under mild conditions, with a convergence rate of $\sqrt{n/m}$ and some control over how m increases. The authors also claim that the DFA-based estimator satisfies a CLT, but provide no proof of it.

2.3 Imputation methods

In this section, we review some of the imputation methods applied in the Monte Carlo simulation and introduce a new one. For a review of some of the R packages available for time series imputation and their performance in the estimation of ARMA models, see Moritz et al. (2015) and Moritz and Bartz-Beielstein (2017). An interesting alternative time series imputation method based on genetic (evolutionary) algorithms is presented in García et al. (2010). The proposed algorithm treats every missing point as a parameter to be estimated in order to minimize an objective depending on the mean, variance, and covariance structure of the time series. Its goal is to obtain a set of values to replace the missing data such that the observed time series' mean, variance and covariance structure are preserved. The method, however, is complex and extremely slow in comparison to other approaches, so it was not considered here.

2.3.1 Mean Substitution

The mean substitution is among the simplest imputation methods available. It is based on substituting missing data with the mean calculated over the observed time series values. In other words, if y_1, \dots, y_n is a sample from a time series for which there are m missing values and denoting by $M := \{i : y_i \text{ is missing}\}$ the set containing the time epochs for which a value is missing (observed), we substitute a missing value y_i by

$$y_i = \frac{1}{\operatorname{card}(M^{\complement})} \sum_{k \in M^{\complement}} y_k, \quad i \in M,$$

where for a set A, card(A) denotes the number of elements (cardinality) in A. As mentioned in the introduction, the mean substitution's strength lies in its simplicity, but since a single value replaces all missing data, it induces some problems: points of masses are produced which affects the time series' variance, dependence structure, and standard errors calculation (for estimators). It also drastically affects estimators based on distributions, such as the estimator in subsection 2.1.1. Alternatively, it could be used the median instead of the mean to impute missing values. Since in this work, we are considering only symmetric time series, the median and mean should perform the same, so we choose the latter since it is faster to calculate.

2.3.2 Linear interpolation

Interpolation using some simple model is quite a common practice in the literature. This can be achieved by a simple linear interpolation in the vicinity of the missing data. If y_t is missing, we apply a simple linear interpolation between the two nearest observed points. Let y_{t_1} and y_{t_2} be the two closest observed points in time satisfying $t_1 < t < t_2$. We impute y_t as

$$y_t = y_{t_1} + \left(\frac{y_{t_2} - y_{t_1}}{t_2 - t_1}\right)(t - t_1).$$

Linear interpolation is also a very simple method that, contrary to the mean substitution, imputes different values for each missing data, when the underlying distribution is absolutely continuous. However, it still underestimates the variance and affects the time series dependence structure. It also imputes wide gaps as a straight line, potentially affecting standard error calculation (for estimators).

2.3.3 Random Substitution

Random substitution is an imputation method based on drawing from a predetermined distribution to substitute a given missing value. The most common is to replace a missing value with a random value drawn from a uniform distribution, typically between the minimum and maximum observed values. This simple method has the advantage that no matter the size of a gap, the imputed values will never be equal. However, since values near the minimum and maximum observed values occur with the same probability as any other interval of the same length, this imputation method tends to inflate the variance of the time series, altering its underlying distribution and affecting its dependence structure.

In what follows, we propose a random substitution method that inherits information regarding the dependence structure on the immediate vicinity of the missing value being imputed. The proposed method is a hybrid of the last observation carried forward method, which consists in substituting each missing value with the most recent observed value, and the random substitution. Let $tN(\mu, \sigma^2, a, b)$ denote the truncated normal distribution, truncated in the interval (a, b), with mean $\mu \in \mathbb{R}$ and variance $\sigma^2 > 0$. If $Z \sim tN(\mu, \sigma^2, a, b)$, Z has density

$$f(x;\mu,\sigma^2,a,b) = \frac{\phi\left(\frac{x-\mu}{\sigma}\right)}{\sigma\left[\Phi\left(\frac{b-\mu}{\sigma}\right) - \Phi\left(\frac{a-\mu}{\sigma}\right)\right]}I(a < x < b),$$

where ϕ and Φ denote the density and distribution of a standard normal distribution, respectively. Let y_t , t > 1 be a missing value to be imputed. We propose imputing y_t by drawing $y_t \sim tN(y_{t-1}, \sigma^2, y_{(1)}, y_{(n)})$, where $y_{(1)} = \min\{y_i : i \in M^{\complement}\}$ and $y_{(n)} = \max\{y_i : i \in M^{\complement}\}$, $M := \{i : y_i \text{ is missing}\}$. The variance parameter σ^2 can be tuned in order to match the variance in the observed time series (see Section 3.2).

3 Simulation

In this section, we present a Monte Carlo simulation study to compare the different approaches to estimate the long-range dependent parameter d in the context of ARFIMA processes. We consider a total of 35 different combinations of estimation and imputation methods. In the simulation, we examine three primary contexts for estimating long-range dependence in time series with missing data. The first involves investigating the impact of imputing the data and subsequently estimating the long-range parameter using methods that require complete data to operate. The second context involves exploring how estimators that can handle missing data perform under different scenarios and various percentages of missing data. These first two contexts are the most natural to consider.

To motivate the third context, recall that methods that can handle time series with missing data can also tackle complete data. Since it takes great effort to develop such estimators, papers introducing these approaches often overlook the performance of an estimator designed for gappy time series when utilized in conjunction with imputation. This idea may go against the rationale of developing a method that can inherently handle missing data, but holds relevance from a practical point of view. Typically, preference is given to the method that produces the most accurate estimate, regardless of its derivation. Hence, the third context we investigate involves assessing the performance of estimators designed to handle time series with missing data when the time series is imputed first before estimation.

3.1 Data generation process (DGP)

In the Monte Carlo simulation study, we consider Gaussian ARFIMA processes generated using R package arfima (Veenstra, 2012), which, according to the package's documentation, generates "a sample from a multivariate normal distribution that has a covariance structure defined by the autocovariances generated for given parameters". The innovation variance was taken as 1. We generate samples of length 2,000 and discarded the first 1,000 observations as burn-in.

To generate the time series containing missing values, we first generate the complete time series, as described above, and then sample the appropriate number of time indexes for which the data is to be missing from a discrete uniform distribution in the set $\{2, \dots, 999\}$. By design, the first and last values of the time series are never missing. Finally, the observations related to the sampled time indexes are set to NA. This procedure is repeated for each replication, resulting in a distinct missing data pattern for each generated time series. However, because we used the same random seed in each session, the pattern of missing data for scenarios with identical n and the percentage of missing data is the same. The missing data pattern for scenario d = 0.1 with 10%, for instance, is the same as the missing data pattern for scenario d = 0.4 with 10% missing data.

3.2 Implementation details

In this Section, we shall review some computational details about each estimator's implementation and the imputation method used in the simulations.

Intrinsic methods

The R code for the intrinsic Craigmile and Mondal's wavelet method presented in Section 2.1.3 can be found in Craigmile's Github: github.com/petercraigmile/GappyLRD. We have made a few changes to the code to suit our purposes better. We apply Daubechi's D4 wavelet considering levels from 1 to 7. A set of R orthonormal Slepian tapers is used to estimate a specific dispersion matrix in the presence of missing data. The argument R in the code was kept at its default value of 7. More information can be found in Craigmile and Mondal (2020).

The copula-based estimator presented in Section 2.1.1 is implemented in the R package

PPMiss (Prass and Pumi, 2023). We consider two configurations for the estimator. In both cases, the copula parameter is estimated using the inversion of Spearman's ρ , and, for the estimation window, we apply s = 1 and m = 24. For the first one, called PP.G, we consider the Gaussian copula as the underlying copula family, while for the second one, called PP.F, we consider Frank's family. This choice was made because since we are simulating Gaussian ARFIMA processes, the Gaussian family is correctly specified. In practice, however, we can never be sure about the copula choice. By considering the misspecified Frank's family, we examine a more realistic scenario.

As for the intrinsic method LoMPE presented in Section 2.1.2, the method is implemented in R package liftLRD (Knight and Nunes, 2018). We consider the implemented bias-corrected estimator in the simulation. The long-range dependence parameter is estimated using a weighted least squares approach, in which the slope of the log-linear relationship between the artificial scales and the log of the integrals is used to re-weight the estimates. The slope in the energy-scale relationship is calculated using all wavelet levels. The LoMPE method is slow compared to the other methods considered, so we only calculate the minimum integral lifting trajectory to estimate d, as bootstrapping makes the method too slow for simulation purposes.

Traditional methods

The R/S method of section 2.2.1 was implemented by the authors. The estimator is obtained through ordinary least squares using the R function 1m. Geweke and Porter-Hudak's estimator presented in Section 2.2.2 is implemented in R package LongMemoryTS (Leschinski, 2019). The first $m = \lfloor 1 + \sqrt{n} \rfloor$ Fourier frequencies were considered in the estimation. The Local Whittle and Exact local Whittle estimators, presented in Section 2.2.3 and 2.2.4, respectively, are also implemented in the package LongMemoryTS. In both cases, we considered $m = \lfloor 1 + \sqrt{n} \rfloor$. In the ELW, the initial value, Y_0 is considered known (or estimated). To fulfill this requirement, we consider the time series $\tilde{Y}_{t-1} := Y_t - Y_1$ for $t \in \{2, \dots, n\}$ so that $\tilde{Y}_0 = 0$ is used. See Remark 2 in Shimotsu and Phillips (2005) for more details and the package LongMemoryTS's documentation.

The DFA method presented in section 2.2.5 was implemented by the authors. Function F_{DFA}^2 is calculated using the R package DCCA (Prass and Pumi, 2020), considering nonoverlapping windows. Regression (2.12) is estimated via ordinary least squares using R function 1m and considering $h \in \{50, \dots, 100\}$, (i.e., l = 50 and s = 50).

Imputation methods

The mean substitution method was implemented by the authors. The Linear interpolation method is implemented in R package zoo (Zeileis and Grothendieck, 2005) through function na.approx. The proposed random interpolation method was implemented by the authors. Hyperparameters a and b were taken as the minimum and maximum observed values, respectively. The variance hyperparameter σ is user-chosen. The goal in defining σ is to use a value that closely approximates the sample standard deviation calculated over the observed values (S) for a variety of values of d and missing data proportions.

We conducted a Monte Carlo Simulation to determine the best value for σ in the context of Gaussian ARFIMA(0, d, 0) processes. We generate time series of length n = 1,000 for $d \in \{0.1, 0.2, 0.3, 0.4\}$. A proportion of 30%, 50%, and 70% of missing data is induced in each time series. In each case, we use the proposed method for imputation considering $\sigma = S/\varsigma$, for $\varsigma \in \{4, 6, 8, 10\}$. The standard deviation of the imputed time series is then computed. We repeat the experiment 1,000 times for each scenario.

The simulation results are summarized in Table 1, which shows the average standard deviation of the generated (complete) time series, the standard deviation calculated from the observed values after introducing missing data (S), and the standard deviation calculated from the data after imputation for each scenario. In parentheses, we present the standard deviation of the estimates alongside the estimated standard deviation. The average standard deviation obtained from the generated time series and S are very similar in all scenarios, a consequence of the model's stationarity. The results show that $\varsigma = 10$ produces the closest results to S while introducing no additional variability. As a result, in the simulation we set $\varsigma = 10$.

Table 1: Simulation results presenting the average standard deviation calculated from the complete generated time series (complete), the observed time series after introducing missing data (S), and the imputed data using the proposed methodology. Also presented are the standard deviation of the estimates (in parentheses).

| d | Missing | C | | imputed t | ime series | | complete |
|-----|---------|---------|-----------------|-----------------|-----------------|------------------|----------|
| | Missing | 3 | $\varsigma = 4$ | $\varsigma = 6$ | $\varsigma = 8$ | $\varsigma = 10$ | complete |
| | 0.2 | 1.007 | 1.017 | 1.011 | 1.008 | 1.007 | |
| | 0.5 | (0.027) | (0.031) | (0.030) | (0.030) | (0.030) | |
| 0.1 | 0.5 | 1.007 | 1.032 | 1.017 | 1.012 | 1.009 | 1.007 |
| 0.1 | 0.5 | (0.032) | (0.041) | (0.040) | (0.040) | (0.040) | (0.023) |
| | 0.7 | 1.009 | 1.059 | 1.028 | 1.018 | 1.012 | |
| | 0.7 | (0.040) | (0.056) | (0.054) | (0.053) | (0.053) | |
| | 0.2 | 1.039 | 1.050 | 1.043 | 1.041 | 1.040 | |
| | 0.5 | (0.030) | (0.034) | (0.033) | (0.033) | (0.033) | |
| 0.2 | 0.5 | 1.039 | 1.065 | 1.050 | 1.045 | 1.042 | 1.040 |
| 0.2 | 0.5 | (0.035) | (0.043) | (0.043) | (0.042) | (0.042) | (0.026) |
| | 0.7 | 1.040 | 1.096 | 1.062 | 1.050 | 1.045 | |
| | 0.7 | (0.044) | (0.061) | (0.058) | (0.057) | (0.056) | |
| | 0.2 | 1.112 | 1.124 | 1.117 | 1.114 | 1.113 | |
| | 0.5 | (0.042) | (0.045) | (0.045) | (0.044) | (0.044) | |
| 0.3 | 0.5 | 1.112 | 1.140 | 1.125 | 1.119 | 1.116 | 1.113 |
| 0.5 | 0.5 | (0.046) | (0.053) | (0.052) | (0.052) | (0.051) | (0.040) |
| | 0.7 | 1.111 | 1.170 | 1.137 | 1.125 | 1.120 | |
| | 0.1 | (0.054) | (0.069) | (0.066) | (0.065) | (0.065) | |
| | 0.2 | 1.256 | 1.270 | 1.261 | 1.258 | 1.257 | |
| | 0.5 | (0.076) | (0.079) | (0.078) | (0.077) | (0.077) | |
| 0.4 | 0.5 | 1.256 | 1.289 | 1.271 | 1.264 | 1.261 | 1.256 |
| 0.4 | 0.5 | (0.078) | (0.084) | (0.082) | (0.082) | (0.082) | (0.075) |
| | 0.7 | 1.255 | 1.320 | 1.282 | 1.270 | 1.262 | |
| | 0.7 | (0.082) | (0.094) | (0.090) | (0.091) | (0.090) | |

3.3 ARFIMA(0, d, 0) scenario

In this section we discuss the estimation of the long-range dependence parameter d in the context of Gaussian ARFIMA(0, d, 0) processes for $d \in \{0.1, 0.2, 0.3, 0.4\}$ and missing data proportions $\{0.1, 0.2, \dots, 0.7\}$. The intrinsic estimators presented in section 2.1 are used to estimate parameter d for each generated time series with missing values. We then proceed with the estimation of d for each estimator considering the originally generated time series,

with no missing data, henceforth referred to as the "original" time series. Next, the time series with missing data are imputed using the three methods discussed in section 2.3. We estimate d for each imputation method and percentage of missing values using the intrinsic and the five estimators discussed in Section 2.2. In total 35 different estimation methods were considered. We repeat the experiment 1,000 times. Due to space limitations, we only present the results for $d \in \{0.1, 0.4\}$. The other cases are presented in the supplementary material that accompanies the paper.

Case d = 0.1

Table 2 shows the simulation results for d = 0.1. The estimated value of d for each estimator is presented in blocks organized by the type of time series considered, namely, intrinsic (methods applied to the time series with missing data), mean, linear, and random (methods applied to the time series imputed using the mean, linear and the proposed random methods, respectively). We present the estimation for each percentage of missing values along with the estimated values for the original time series, presented in column "0". These values are repeated throughout the blocks for convenience. The best estimate in each block is highlighted in bold.

There is a lot to discuss from Table 2. First, we look at the best results. Interestingly, when there is no missing data, the intrinsic methods produce the best results. The copulabased methods presented the best results for complete data among the intrinsic estimators, followed closely by the LoMPE. When missing data is taken into account, for all percentages up to 50% the intrinsic estimators produce the best results overall by a wide margin. The behavior is somewhat wild in extreme cases (60% and 70%).

The effects of using the mean imputation method are very noticeable for all estimators. Among the imputation methods, the mean performs the worst for all percentages of missing, except for 70% for which it presented the best overall result. To begin, all methods underestimate d, with the exception of the R/S, which does the opposite. For all estimators, with the exception of the R/S, using the mean imputation method degrades the estimated values as the percentage of missing data increases. Most estimated values present a relative bias of over 50% in this case. The methods that suffer the most are the frequency domain ones, namely GPH, LW, and ELW. The R/S, however, behaves exactly the opposite and produces the best results when the mean imputation method is applied, even when the percentage of missing data increases.

When the best results from the linear and random imputation methods are compared, we find that they perform similarly, with the linear method having a slight advantage. The estimated values of intrinsic methods and the R/S for both imputation methods greatly overestimate the parameter and always increase with the percentage of missing data, rendering them effectively useless when more than 20% of the data is missing. With 10% of missing data, the copula-based ones (PP.F and PP.G) present the best overall performance. With 20% of missing data or more, the DFA is the one that performs the best for both imputation methods. The frequency domain estimators (GPH, LW, and ELW) consistently underestimate d and perform poorly, with relative bias exceeding 50%.

Boxplots of the results are presented in Figure 1. The columns represent the percentage of missing data (10%, 40%, and 70%). The first row presents the intrinsic methods applied to the data with missing values. From the second row on, we present the boxplot of the estimated values using each imputation method. As expected, the variability of the intrinsic estimators in the case of missing data increases as the percentage of missing data increases.

| | | | | d = 0.1 | | | | | |
|-----------|-----------|-------|-------|---------|-------|--------|--------|--------|--------|
| | | | | | Mis | sing | | | |
| Type | Estimator | 0 | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 0.6 | 0.7 |
| | Full | 0.076 | 0.077 | 0.086 | 0.088 | 0.109 | 0.113 | 0.116 | 0.152 |
| | Abry | 0.081 | 0.083 | 0.086 | 0.089 | 0.096 | 0.103 | 0.116 | 0.142 |
| Intrinsic | PP.G | 0.092 | 0.091 | 0.090 | 0.089 | 0.089 | 0.088 | 0.082 | 0.074 |
| | PP.F | 0.086 | 0.086 | 0.085 | 0.084 | 0.084 | 0.083 | 0.079 | 0.071 |
| | LoMPE | 0.084 | 0.079 | 0.076 | 0.071 | 0.070 | 0.063 | 0.060 | 0.057 |
| | Full | 0.076 | 0.068 | 0.060 | 0.052 | 0.045 | 0.035 | 0.027 | 0.019 |
| | Abry | 0.081 | 0.073 | 0.064 | 0.056 | 0.049 | 0.039 | 0.031 | 0.022 |
| | PP.G | 0.092 | 0.074 | 0.058 | 0.044 | 0.067 | 0.050 | 0.010 | 0.006 |
| | PP.F | 0.086 | 0.069 | 0.052 | 0.037 | 0.055 | 0.049 | 0.008 | 0.004 |
| Moon | LoMPE | 0.084 | 0.077 | 0.070 | 0.062 | 0.053 | 0.046 | 0.036 | 0.029 |
| Mean | DFA | 0.073 | 0.066 | 0.059 | 0.052 | 0.043 | 0.030 | 0.023 | 0.016 |
| | GPH | 0.034 | 0.020 | 0.013 | 0.001 | -0.013 | -0.033 | -0.054 | -0.076 |
| | LW | 0.048 | 0.045 | 0.041 | 0.038 | 0.034 | 0.030 | 0.028 | 0.022 |
| | ELW | 0.046 | 0.044 | 0.039 | 0.041 | 0.039 | 0.042 | 0.059 | 0.073 |
| | R/S | 0.145 | 0.140 | 0.136 | 0.131 | 0.125 | 0.118 | 0.112 | 0.101 |
| | Full | 0.076 | 0.133 | 0.197 | 0.269 | 0.350 | 0.441 | 0.546 | 0.673 |
| | Abry | 0.081 | 0.131 | 0.188 | 0.254 | 0.331 | 0.418 | 0.521 | 0.650 |
| | PP.G | 0.092 | 0.120 | 0.143 | 0.163 | 0.182 | 0.201 | 0.221 | 0.247 |
| | PP.F | 0.086 | 0.114 | 0.139 | 0.161 | 0.183 | 0.206 | 0.233 | 0.266 |
| т: | LoMPE | 0.084 | 0.135 | 0.192 | 0.256 | 0.327 | 0.411 | 0.511 | 0.631 |
| Linear | DFA | 0.073 | 0.073 | 0.073 | 0.081 | 0.081 | 0.087 | 0.104 | 0.142 |
| | GPH | 0.034 | 0.028 | 0.027 | 0.024 | 0.023 | 0.025 | 0.026 | 0.046 |
| | LW | 0.048 | 0.044 | 0.041 | 0.037 | 0.034 | 0.033 | 0.032 | 0.035 |
| | ELW | 0.046 | 0.042 | 0.039 | 0.038 | 0.034 | 0.033 | 0.036 | 0.043 |
| | R/S | 0.145 | 0.151 | 0.159 | 0.168 | 0.178 | 0.193 | 0.212 | 0.237 |
| | Full | 0.076 | 0.128 | 0.184 | 0.244 | 0.311 | 0.380 | 0.454 | 0.538 |
| | Abry | 0.081 | 0.126 | 0.177 | 0.233 | 0.296 | 0.365 | 0.442 | 0.531 |
| | PP.G | 0.092 | 0.118 | 0.141 | 0.162 | 0.184 | 0.207 | 0.231 | 0.262 |
| | PP.F | 0.086 | 0.112 | 0.137 | 0.159 | 0.183 | 0.209 | 0.237 | 0.272 |
| | LoMPE | 0.084 | 0.130 | 0.181 | 0.235 | 0.292 | 0.358 | 0.431 | 0.511 |
| Kandom | DFA | 0.073 | 0.070 | 0.067 | 0.072 | 0.075 | 0.080 | 0.096 | 0.135 |
| | GPH | 0.034 | 0.025 | 0.024 | 0.019 | 0.021 | 0.023 | 0.027 | 0.039 |
| | LW | 0.048 | 0.043 | 0.038 | 0.035 | 0.031 | 0.030 | 0.028 | 0.032 |
| | ELW | 0.046 | 0.041 | 0.037 | 0.035 | 0.034 | 0.031 | 0.035 | 0.043 |
| | R/S | 0.145 | 0.148 | 0.151 | 0.156 | 0.165 | 0.179 | 0.197 | 0.224 |

Table 2: Simulation results for the ARFIMA(0, 0.1, 0) scenario.

The bias is also affected but to a lesser degree.



Figure 1: Box plot of the fitted ARFIMA (0, 0.1, 0) model.

When data imputation is in place, the variability of the estimators does not seem to be significantly impacted by the percentage of missing data. The bias, however, is very much so. The DFA, ELW, and GPH are the methods presenting the highest overall variability. It is also noteworthy that, for 10% of missing, the imputation method applied makes little difference in the boxplot within each method, but at 40% and 70% the imputation method plays an important role.

Case d = 0.4

Table 3 shows the simulation results for d = 0.4. When comparing the five intrinsic estimators, the copula-based ones consistently outperform the others, presenting a very good performance even when the percentage of missing data is 70%. When the original time series is taken into account, the GPH estimator is the best performer, followed by the copula-based ones. When there is no missing data, Craigmile and Mondal's (Full and Abry), LoMPE, and LW estimators all perform poorly.

The effects of using the mean imputation method are even more severe than in the case of d = 0.1 with the mean imputation method performing the worst of all imputation methods for all percentages of missing data. As the percentage of missing data increases, we observe a degradation of the estimated values with the mean imputation for all estimators, which uniformly underestimate d. The GPH was the best performer overall for mean imputation. Regardless of the method, when missing data reaches 30% or above, the estimated values are uniformly poor, to the point that they are useless for practical purposes.

The linear and random imputation methods perform about the same, just as in the case of d = 0.1. For percentages of missing data of 30% or above, the copula-based estimators are the best performers by a good margin, while the other intrinsic methods, the DFA, LW, ELW, and R/S increasingly degrade as the percentage of missing data increases beyond 30%, yielding useless estimations. The GPH also degrades as the percentage of missing data increases

beyond 30% but to a lower degree.

Boxplots of the results are presented in Figure 2. From the boxplots, we observe that the variability of the spectral density-based estimators (GPH, EL, and ELW) are considerably greater than in the case d = 0.1 for all imputation methods. Comparing both cases, d = 0.1 and d = 0.4, we observe somewhat similar behavior for all estimators when the percentage of missing values is 10%. For missing data percentage of 70%, all imputation methods severely affect the wavelet-based estimators (Abry, Full, and LoMPE), causing such a bias that the estimated values are useless. In the case of linear imputation, for instance, all estimated values for these estimators are higher than 0.6, so the boxplots does not even appear in the plotting region of Figure 2.

The copula-based estimators perform well and consistently in the case of linear and random imputation as the percentage of missing values increases. In the case of mean imputation, however, the estimated values are useless for percentages of missing data above 10%. The DFA, GPH, ELW, and LW methods present a somewhat comparable overall performance, with a slight advantage for the GPH in most cases. The R/S performs stably for all percentages of missing values, with considerable bias, especially for the mean imputation case.

| | | | | d = 0.4 | Ł | | | | |
|-----------------------------------|------------|-------|-------|---------|-------|-------|-------|-------|-------|
| | Estimation | | | | Mis | sing | | | |
| Type | Estimator | 0 | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 0.6 | 0.7 |
| | Full | 0.332 | 0.341 | 0.347 | 0.346 | 0.353 | 0.358 | 0.366 | 0.365 |
| Type Intrinsic Mean Linear Random | Abry | 0.344 | 0.350 | 0.352 | 0.355 | 0.358 | 0.360 | 0.362 | 0.366 |
| Intrinsic | PP.G | 0.382 | 0.381 | 0.381 | 0.381 | 0.380 | 0.380 | 0.378 | 0.376 |
| | PP.F | 0.379 | 0.379 | 0.378 | 0.379 | 0.378 | 0.378 | 0.376 | 0.375 |
| | LoMPE | 0.333 | 0.326 | 0.322 | 0.314 | 0.335 | 0.325 | 0.316 | 0.336 |
| | Full | 0.332 | 0.294 | 0.260 | 0.227 | 0.196 | 0.165 | 0.133 | 0.102 |
| | Abry | 0.344 | 0.307 | 0.274 | 0.241 | 0.209 | 0.178 | 0.144 | 0.112 |
| | PP.G | 0.382 | 0.352 | 0.316 | 0.276 | 0.249 | 0.096 | 0.113 | 0.056 |
| | PP.F | 0.379 | 0.346 | 0.305 | 0.260 | 0.225 | 0.083 | 0.090 | 0.041 |
| Moon | LoMPE | 0.333 | 0.297 | 0.265 | 0.235 | 0.205 | 0.176 | 0.147 | 0.116 |
| Mean | DFA | 0.357 | 0.344 | 0.334 | 0.318 | 0.295 | 0.276 | 0.248 | 0.211 |
| | GPH | 0.389 | 0.374 | 0.361 | 0.342 | 0.321 | 0.292 | 0.262 | 0.215 |
| | LW | 0.314 | 0.307 | 0.300 | 0.292 | 0.278 | 0.262 | 0.243 | 0.215 |
| | ELW | 0.372 | 0.355 | 0.340 | 0.323 | 0.297 | 0.266 | 0.235 | 0.198 |
| | R/S | 0.342 | 0.332 | 0.321 | 0.309 | 0.291 | 0.273 | 0.250 | 0.221 |
| | Full | 0.332 | 0.375 | 0.424 | 0.481 | 0.544 | 0.617 | 0.702 | 0.807 |
| | Abry | 0.344 | 0.381 | 0.423 | 0.473 | 0.531 | 0.599 | 0.681 | 0.786 |
| | PP.G | 0.382 | 0.383 | 0.384 | 0.385 | 0.388 | 0.391 | 0.393 | 0.400 |
| | PP.F | 0.379 | 0.381 | 0.384 | 0.386 | 0.391 | 0.397 | 0.403 | 0.416 |
| Linoar | LoMPE | 0.333 | 0.371 | 0.415 | 0.464 | 0.519 | 0.586 | 0.667 | 0.766 |
| Linear | DFA | 0.357 | 0.352 | 0.351 | 0.345 | 0.340 | 0.335 | 0.336 | 0.344 |
| | GPH | 0.389 | 0.384 | 0.381 | 0.374 | 0.373 | 0.359 | 0.355 | 0.352 |
| | LW | 0.314 | 0.310 | 0.307 | 0.300 | 0.296 | 0.286 | 0.276 | 0.267 |
| | ELW | 0.372 | 0.367 | 0.363 | 0.354 | 0.351 | 0.341 | 0.329 | 0.322 |
| | R/S | 0.342 | 0.339 | 0.338 | 0.336 | 0.333 | 0.331 | 0.332 | 0.336 |
| | Full | 0.332 | 0.363 | 0.397 | 0.436 | 0.476 | 0.522 | 0.571 | 0.628 |
| | Abry | 0.344 | 0.369 | 0.398 | 0.432 | 0.470 | 0.513 | 0.565 | 0.626 |
| | PP.G | 0.382 | 0.382 | 0.383 | 0.385 | 0.387 | 0.391 | 0.394 | 0.401 |
| | PP.F | 0.379 | 0.380 | 0.382 | 0.384 | 0.387 | 0.393 | 0.398 | 0.409 |
| Bandom | LoMPE | 0.333 | 0.360 | 0.390 | 0.423 | 0.459 | 0.501 | 0.550 | 0.605 |
| random | DFA | 0.357 | 0.349 | 0.344 | 0.335 | 0.324 | 0.317 | 0.307 | 0.313 |
| | GPH | 0.389 | 0.383 | 0.376 | 0.369 | 0.364 | 0.346 | 0.335 | 0.330 |
| | LW | 0.314 | 0.308 | 0.303 | 0.293 | 0.285 | 0.272 | 0.256 | 0.241 |
| | ELW | 0.372 | 0.365 | 0.358 | 0.347 | 0.338 | 0.325 | 0.310 | 0.298 |
| | R/S | 0.342 | 0.335 | 0.328 | 0.323 | 0.316 | 0.312 | 0.309 | 0.315 |

Table 3: Simulation results for the ARFIMA(0, 0.4, 0) scenario.

3.4 ARFIMA(1, d, 1) scenario

In this section we consider the estimation of the long-range dependence parameter d in the context of Gaussian ARFIMA(1, d, 1) processes for $d \in \{0.1, 0.2, 0.3, 0.4\}$, $\phi = 0.5$, $\theta = 0.6$ and missing data proportions $\{0.1, 0.2, \dots, 0.7\}$. We follow the same steps as in Section 3.3. Tables and plots presenting the results can be found in the supplementary material that accompanies the paper. The results for the (1, d, 1) case are very similar to the (0, d, 0)'s and the same remarks presented in Section 3.3 apply here case by case. The similarity between the results is expected since the estimators applied in the simulation are all semiparametric, focusing only on the long-range dependence structure in the time series, regardless of short-range nuisances.



Figure 2: Box plot of the fitted ARFIMA(0, 0.4, 0) model.

3.5 Time benchmarking

In this section, we compare the computational speed of each estimator considered in the simulations. Besides which estimator is the fastest to compute, there are a few other questions regarding computational speed that are of interest. For instance, does doubling the length of the time series double the time required to estimate d? Is the computational time required to estimate d affected by the strength of the dependence? Is the percentage of missing data a factor in estimation times? What about the imputation method applied? In this section, we study these questions through a series of Monte Carlo simulations.

3.5.1 Setup

We perform a series of routines with each estimator involving several different subtasks, measuring the time spent performing them for each estimator. The routine is divided into two main tasks. The first main task is intended only for the intrinsic estimators and consists of estimating d in Gaussian ARFIMA(0, d, 0) generated with 20% and 70% of missing data, considering $d \in \{0.1, 0.4\}$ data and sample size $n \in \{1000, 2000\}$. The time spent executing each subtask 1,000 times for each estimator was recorded. The time series were all generated and prepared beforehand, so the recorded times reflect the time spent actually performing the task, and nothing else.

The second main task involves all estimators. The estimators are used to estimate d in the original time series. Next, the estimators are applied to estimate d from the time series with 20% and 70% of induced time series after imputation using each of the three methods presented in section 2.3 is performed. The exercise is performed 1,000, and the time spent performing it is recorded for each subtask and each estimator. Again, time series were all generated and prepared beforehand. All simulations were run serially in a computer using R version 4.1.3 with the following configurations: Intel Core is 8600k CPU (3.6GHz) processor

(factory configurations), and 16GB RAM running on Windows 10 Pro. The simulations were performed.

3.5.2 Results

The complete results are shown in Table 4 (time series with missing data) and Table 5 (original and imputed time series). For the Pumi et al. (2023)'s copula-based estimator, there is a significant difference in speed between the two variants considered. When considering the imputed (or original) time series, using Frank's family (PP.F) to perform the subtasks takes approximately 4.1 times longer than using the Gaussian's (PP.G) for n = 1,000, and approximately 3.2 when n = 2,000. Considering time series with missing data, these ratios increase to 4.3 and 3.4, respectively. This distinction is justified by the fact that the relationship between Spearman's ρ and Frank's copula parameter is dependent on special functions (c.f. Nelsen, 2013, p. 171) so estimating the parameter via inversion of Spearman's ρ requires a somewhat lengthy numerical inversion. On the other hand, Spearman's ρ has a closed formula for the Gaussian family, given by the classical relationship $\rho = \frac{6}{\pi} \arcsin(r)$, where r denotes the (Pearson's) correlation.

Table 4: Time spent to complete the simulation task for the intrinsic estimators. Presented is the total time spent, in seconds, performing the respective subtask.

| Intrincia | | n = 1 | 1,000 | | n = 2,000 | | | | | |
|-----------|-------|-------|-------|-------|-----------|--------|---------|--------|--|--|
| Estimator | d = | 0.1 | d = | 0.4 | d = | 0.1 | d = 0.4 | | | |
| Estimator | 20% | 70% | 20% | 70% | 20% | 70% | 20% | 70% | | |
| PP.G | 94.2 | 88.4 | 93.4 | 87.7 | 130.8 | 120.7 | 129.5 | 120.0 | | |
| PP.F | 392.8 | 386.3 | 394.7 | 390.3 | 427.5 | 415.2 | 431.8 | 423.5 | | |
| Abry | 392.7 | 392.4 | 388.2 | 388.2 | 1367.6 | 1342.7 | 1334.9 | 1335.2 | | |
| Full | 392.8 | 392.4 | 388.2 | 388.3 | 1353.8 | 1341.8 | 1335.1 | 1335.0 | | |
| LoMPE | 244.8 | 68.8 | 238.1 | 68.1 | 745.6 | 162.4 | 713.2 | 161.3 | | |

Does doubling the time series' length double the time spent in estimating d?

It depends on the estimator and the setup. Considering only intrinsic estimators (Table 4), increasing the sample size from n = 1,000 to 2,000 took, on average, only 8.5% longer for PP.F to complete the tasks and 37.7% longer for PP.G. For the wavelet-based estimators, the scenario is very different. LoMPE and Craigmile and Mondal (2020)'s estimator took, on average, 2.7 and 3.4 times the amount of time to complete the task, respectively.

From Table 5, doubling the sample size from n = 1,000 to n = 2,000 requires less than twice the time to complete the task, on average, for the estimators GPH (1.81), LW (1.34) and DFA (1.65). The R/S and ELW took, on average, 4.3 and 9.3 times the amount of time to complete the task, respectively. Applying the intrinsic estimators to the original and imputed time series (Table 5) and doubling the sample size from n = 1,000 to n = 2,000 on average, produce an overall increase in the time spent to complete the tasks. For the PP.F and PP.G, this increase was small: 10% and 40.5% longer to complete the tasks, respectively. LoMPE and Craigmile and Mondal (2020)'s estimator took about 3.6 and 4.5 times the time spent to complete the tasks, on average, respectively.

| n | d | % | input. | PP.G | PP.F | Abry | Full | LoMPE | GPH | LW | ELW | DFA | R/S |
|-----|-----|-----|--------|-------|-------|-------|-------|--------|------|------|------|------|------|
| | | or | iginal | 97.0 | 396.3 | 37.1 | 37.1 | 745.6 | 0.09 | 0.14 | 2.63 | 3.36 | 8.97 |
| | | | mean | 97.7 | 397.3 | 36.9 | 37.1 | 745.9 | 0.11 | 0.16 | 2.81 | 3.26 | 8.94 |
| | | 0.2 | linear | 97.5 | 396.2 | 37.2 | 37.0 | 745.2 | 0.10 | 0.16 | 2.64 | 3.21 | 8.97 |
| | 0.1 | | rand | 96.3 | 394.4 | 36.9 | 36.9 | 744.4 | 0.11 | 0.15 | 2.65 | 3.22 | 9.07 |
| | | | mean | 98.6 | 395.6 | 36.9 | 36.9 | 747.7 | 0.11 | 0.16 | 2.73 | 3.22 | 8.94 |
| | | 0.7 | linear | 96.4 | 400.3 | 36.9 | 36.9 | 746.8 | 0.10 | 0.15 | 2.57 | 3.22 | 9.04 |
| 000 | | | rand | 95.5 | 398.0 | 36.9 | 36.9 | 782.8 | 0.11 | 0.16 | 2.58 | 3.36 | 9.28 |
| 1,(| | or | iginal | 95.9 | 397.3 | 36.6 | 36.6 | 728.3 | 0.11 | 0.11 | 2.19 | 3.19 | 9.03 |
| | | | mean | 96.6 | 395.4 | 36.5 | 36.5 | 733.1 | 0.11 | 0.11 | 2.39 | 3.20 | 8.88 |
| | | 0.2 | linear | 95.5 | 403.3 | 36.6 | 37.0 | 735.8 | 0.11 | 0.10 | 2.17 | 3.12 | 8.95 |
| | 0.4 | | rand | 95.4 | 398.0 | 36.5 | 36.5 | 732.4 | 0.10 | 0.11 | 2.20 | 3.18 | 9.00 |
| | | | mean | 97.7 | 392.8 | 36.5 | 36.4 | 732.2 | 0.11 | 0.11 | 2.65 | 3.19 | 8.87 |
| | | 0.7 | linear | 95.0 | 408.6 | 36.5 | 36.5 | 733.5 | 0.11 | 0.11 | 2.33 | 3.18 | 8.87 |
| | | | rand | 94.6 | 402.0 | 36.5 | 36.5 | 735.7 | 0.11 | 0.11 | 2.23 | 3.18 | 8.87 |
| | | or | iginal | 142.3 | 448.5 | 171.1 | 171.1 | 2784.4 | 0.21 | 0.18 | 25.5 | 5.37 | 39.4 |
| | | | mean | 141.0 | 442.5 | 170.6 | 168.7 | 2808.4 | 0.20 | 0.18 | 25.6 | 5.38 | 38.7 |
| | | 0.2 | linear | 136.5 | 431.9 | 168.8 | 166.3 | 2633.2 | 0.19 | 0.18 | 25.1 | 5.28 | 38.4 |
| | 0.1 | | rand | 134.4 | 430.8 | 166.6 | 167.8 | 2636.9 | 0.19 | 0.21 | 25.0 | 5.30 | 38.4 |
| | | | mean | 135.4 | 429.6 | 166.7 | 167.6 | 2645.8 | 0.18 | 0.22 | 25.9 | 5.28 | 38.4 |
| | | 0.7 | linear | 134.2 | 434.9 | 166.9 | 166.5 | 2661.0 | 0.21 | 0.22 | 25.1 | 5.65 | 39.7 |
| 000 | | | rand | 138.7 | 453.9 | 170.5 | 169.9 | 2690.8 | 0.19 | 0.26 | 24.4 | 5.37 | 38.4 |
| 5,0 | | or | iginal | 133.6 | 435.4 | 165.7 | 165.5 | 2638.1 | 0.17 | 0.14 | 20.4 | 5.27 | 38.4 |
| | | | mean | 134.3 | 433.3 | 165.8 | 165.4 | 2652.5 | 0.20 | 0.14 | 20.9 | 5.26 | 38.5 |
| | | 0.2 | linear | 133.5 | 438.4 | 165.8 | 165.4 | 2650.5 | 0.20 | 0.15 | 20.6 | 5.27 | 38.5 |
| | 0.4 | | rand | 133.3 | 435.9 | 165.6 | 165.6 | 2641.4 | 0.19 | 0.15 | 20.4 | 5.25 | 38.4 |
| | | | mean | 134.8 | 429.0 | 165.7 | 165.8 | 2635.3 | 0.19 | 0.14 | 23.2 | 5.29 | 38.4 |
| | | 0.7 | linear | 132.7 | 447.3 | 165.6 | 166.0 | 2638.0 | 0.18 | 0.14 | 20.8 | 5.25 | 38.4 |
| | | | rand | 132.0 | 441.4 | 165.3 | 165.7 | 2637.2 | 0.19 | 0.16 | 20.7 | 5.38 | 38.3 |

Table 5: Time spent (in seconds) to complete the simulation task considering all estimators and imputed/original time series. Presented is the total time spent (in seconds) performing the subtasks.

Does the dependence strength affect computational times?

It depends on the estimator. Completing the full task in the case d = 0.1 takes on average longer compared to d = 0.4 for all classical estimators. More precisely, comparing d = 0.1versus d = 0.4, it takes about 42% longer for the LW, 18% for the ELW, 2% for the DFA, 1% for the R/S, and 0.25% for the GPH to complete the task. On the other hand, the time spent to complete the task using the intrinsic estimators is not significantly affected by the dependence strength - the difference in completing the full task when d = 0.1 takes no more than 2.3% in absolute value compared to d = 0.4.

Does the percentage of missing data affect computational times?

It depends on the estimator and whether the time series contains missing data or not. When missing values are considered, performing the tasks is not significantly affected by the percentage of missing values for estimators PP.G, PP.F Full, and Abry. For these estimators, when 20% of the data is missing, performing the task takes no more than 2% longer when compared to 70%. However, for LoMPE, performing the task when 20% of the data is missing takes about 4 times the time spent when 70% is missing.

When the time series is the original or imputed, the Whittle estimators LW and ELW are slightly affected by the percentage of missing value, taking about 6.6% and 2% longer to complete the task when 70% of the data is missing compared to 20%. The other estimators are affected for no more than 1.2%.

Does the imputation method applied to affect computational times?

The time spent completing the task after imputation does not depend on the percentage of missing values prior to imputation nor on the imputation method applied.

Which estimator is the fastest to compute?

It depends on the metrics and scenario. The intrinsic estimators are capable of handling missing data but they are more involved to calculate than the classic estimators. Hence it is expected that the classical estimators can be computed faster.

Table 6 presents the total time spent by each estimator to complete the whole routine, along with the minimum and maximum amount of time spent in a single subtask. Both Craigmile and Mondal (2020)'s variants, Full and Abry, spent about the same amount of time performing the full task. This is expected since the only difference in the estimators is the way a certain matrix is used to calculate the estimator (whole matrix or the diagonal entries only), at the end of the estimation algorithm. If we only consider the total amount of time

Table 6: Time spent to complete the simulation task considering all estimators and imputed/original time series. Presented is the average time spent (in seconds) over the percentage of missing and imputation methods to perform the subtasks.

| motric | | Original/imputed time series | | | | | | | | | | | |
|--------|--------|------------------------------|--------|-----------|------------|----------|------|-------|-------|-------|--|--|--|
| metric | PP.G | PP.F | Abry | Full | LoMPE | GPH | LW | ELW | DFA | R/S | | | |
| total | 3246.3 | 11708.2 | 2855.4 | 2852.4 | 47743.3 | 4.18 | 4.31 | 358.3 | 119.7 | 665.8 | | | |
| max | 142.3 | 453.9 | 171.1 | 171.1 | 2808.4 | 0.21 | 0.26 | 25.9 | 5.65 | 39.7 | | | |
| min | 94.6 | 392.8 | 36.5 | 36.4 | 728.3 | 0.09 | 0.10 | 2.17 | 3.12 | 8.87 | | | |
| metric | | | r | Гime seri | es with mi | ssing da | ata | | | | | | |
| total | 864.8 | 3262.1 | 6941.9 | 6927.5 | 2402.3 | | | | | | | | |
| max | 130.8 | 431.8 | 1367.6 | 1353.8 | 745.6 | | | | | | | | |
| min | 87.7 | 386.3 | 388.2 | 388.2 | 68.1 | | | | | | | | |

spent performing the task, when missing data is considered, the fastest estimator is the PP.G followed by LoMPE, which takes about 2.8 times the time PP.G takes to perform the whole

task. In this scenario, Craigmile and Mondal (2020)'s ones were the slowest, taking about 8 times the time spent by the PP.G to complete the task. Among the classical estimators, the fastest is the GPH followed closely by the LW. The third fastest is the DFA, but taking about 27.8 times the time to complete the task of the second one. The fastest among the intrinsic when the original/imputed time series is considered are the Craigmile and Mondal (2020)'s ones, followed closely by the PP.G with LoMPE being the slowest. The fastest intrinsic estimator (Full) took an astonishing 682 times GPH's total time to complete the task.

However, looking only at totals may not be ideal. For instance, in Table 4, we observe that when n = 1,000 and the percentage of missing is 20%, the PP.G is the fastest intrinsic estimator, while LoMPE is the fastest when 70% of the data is missing. When n = 2,000, the PP.G is uniformly faster though. Looking at the results presented in Table 5, we observe that the GPH and LW are the fastest estimators by far. For n = 1,000, the GPH is as fast or faster than LW in all but one subtask, while for n = 2,000, the LW is faster in 10 out of 14 subtasks.

A curiosity is that in Shimotsu and Phillips (2005), the authors claim (page 1891), based on their simulation experience, that the ELW is about ten times more expensive to compute than the LW. In the presented simulations, we found this number to be about 83 times more expensive, according to Table 6. Looking at Table 5, we found that the ELW is never less expensive than 16.1 times the LW, with a top value of 166 times. On average, the ELW is about 76.8 times more expensive to compute than the LW. This discrepancy could be due to the efficiency of the implementation applied in the original paper and the one used here.

3.5.3 Convergence

The classical semiparametric estimators applied are known to be computationally stable. The copula-based estimator is computationally stable as well, as is the LoMPE. Only Craigmile and Mondal (2020)'s presented computational issues. For most scenarios, the estimator failed in about 6% of the trials. The problem is accentuated when d = 0.1 and the percentage of missing data is higher than 40%. For instance, when d = 0.1 and the percentage of missing is 70%, a third of all attempts to estimate d with Craigmile and Mondal (2020)'s estimators fail. The failures can be traced back to the particular wavelet-based construction employed in the estimator. In sparse scenarios, a lack of data across the wavelet scales may occur, thereby hindering the estimator's capacity to generate meaningful estimates.

4 Discussion

In this work, we presented an extensive Monte Carlo simulation study regarding the estimation of parameter d in long-range dependent time series in the presence of missing data. We considered estimators especially tailored to deal with missing data, and estimators that can only be used after the imputation of the missing values, which was done considering three different methods. A variety of scenarios were considered, including percentages of missing data ranging from 10% to 70%, different sample sizes, and values of d.

Our findings show that in the context of long-range dependent time series, mean imputation should be avoided, in favor of the linear or random methods. When the dependence strength is low applying a intrinsic estimator (other than LoMPE) usually yields the best results, with a small advantage for the copula-based estimators, especially the PP.G given its numerical stability, comparatively low computational cost, bias, and variance. In this context, if a classical estimator is to be used, the DFA paired with the random or linear imputation is the only acceptable choice as the others present an extremely poor performance.

Under strong long-range dependence, the copula-based estimators applied to the gappy time series present the overall best results, with a small advantage for the Gaussian variant, PP.G. If imputation is to be used, the copula-based estimators paired with the random or linear imputation yield the most consistent results, with a small advantage to the PP.F.

Our findings show that increasing the sample size has different effects on different estimators, for LoMPE, Full, Abry, R/S and ELW, doubling the sample size from n = 1,000 to n = 2,000 requires more than twice the time to complete the task, on average, while for the others, it requires less. We found that for most estimators, the value of d has a negligible effect on the time required to perform the estimation (exceptions: LW and ELW). The overall fastest estimators are by far the GPH and LW. Among the intrinsic estimator, the PP.G was the estimator that perform the fastest for n = 2,000. All estimators are very stable with exception of Craigmile and Mondal (2020)'s, which, due to its particular construction, is prune to fail in very high missing scenarios.

References

- Abry, P. and Veitch, D. (1998). Wavelet analysis of long-range-dependent traffic. *IEEE Transactions on Information Theory*, 44(1):2–15.
- Bardet, J.-M. and Kammoun, I. (2008). Asymptotic properties of the detrended fluctuation analysis of long range dependence processes. *IEEE Transactions on Information Theory*, 54(5):2041–2052.
- Beran, J. (1994). Statistics for Long Memory Processes. Chapman and Hall.
- Chan, N. H. and Palma, W. (1998). State space modeling of long-memory processes. *The* Annals of Statistics, 26(2):719 – 740.
- Craigmile, P. F. and Mondal, D. (2020). Estimation of long-range dependence in gappy Gaussian time series. *Statistics and Computing*, 30(1):167–185.
- Doukhan, P., Oppenheim, G., and Taqqu, M. S. (2003). Theory and Applications of Long-Range Dependence. Birkhäuser Boston, MA.
- Faÿ, G., Moulines, E., Roueff, F., and Taqqu, M. S. (2009). Estimators of long-memory: Fourier versus wavelets. *Journal of Econometrics*, 151(2):159–177.
- García, J. C. F., Kalenatic, D., and Bello, C. A. L. (2010). An evolutionary approach for imputing missing data in time series. *Journal of Circuits, Systems and Computers*, 19(01):107–121.
- Geweke, J. and Porter-Hudak, S. (1983). The estimation and application of long memory time series models. *Journal of time series analysis*, 4(4):221–238.
- Granger, C. W. J. and Joyeux, R. (1980). An introduction to long memory time series and fractional differencing. *Journal of Time Series Analysis*, 1:15–30.
- Honsking, J. R. M. (1981). Fractional differencing. Biometrika, 1(68):165–176.
- Hurst, H. E. (1951). Long-term storage capacity of reservoirs. Transactions of the American Society of Civil Engineers, 116(1):770–799.

- Hurvich, C. M., Deo, R., and Brodsky, J. (1998). The mean squared error of Geweke and Porter-Hudak's estimator of the memory parameter of a long-memory time series. *Journal* of *Time Series Analysis*, 19(1):19–46.
- Jansen, M., Nason, G. P., and Silverman, B. W. (2001). Scattered data smoothing by empirical bayesian shrinkage of second-generation wavelet coefficients. In *Wavelets: Applications in* Signal and Image Processing IX, volume 4478, pages 87–97. SPIE.
- Joe, H. (1997). Multivariate models and dependence concepts. Chapman & Hall.
- Knight, M. and Nunes, M. (2018). *liftLRD: wavelet lifting estimators of the Hurst exponent* for regularly and irregularly sampled time series. R package version 1.0-8.
- Knight, M. I., Nason, G. P., and Nunes, M. A. (2017). A wavelet lifting approach to longmemory estimation. *Statistics and Computing*, 27(6):1453–1471.
- Kokoszka, P. S. and Bhansali, R. J. (2001). Estimation of the long memory parameter: A review of recent developments and an extension. In Basawa, I. V., Heyde, C. C., and Taylor, R. L., editors, *Proceedings of the Symposium on Inference for Stochastic Processes*, IMS Lecture Notes, pages 125–150, Athens, Greece.
- Leschinski, C. (2019). LongMemoryTS: long memory time series. R package version 0.1.0.
- Mandelbrot, B. B. (1975). Limit theorem on the self-normalized range for weakly and strongly dependent process. Zeitschrift f
 ür Wahrscheinlichkeitstheorie und Verwandte Gebiete, 31:271–285.
- Moritz, S. and Bartz-Beielstein, T. (2017). impute TS: time series missing value imputation in R. *The R Journal*, 9(1):207–218.
- Moritz, S., Sardá, A., Bartz-Beielstein, T., Zaefferer, M., and Stork, J. (2015). Comparison of different methods for univariate time series imputation in R.
- Nelsen, R. (2013). An Introduction to Copulas. Lecture Notes in Statistics. Springer New York, 2nd. edition.
- Palma, W. (2007). Long-memory time series: theory and methods. John Wiley & Sons.
- Peng, C. K., Buldyrev, S. V., Havlin, S., Simons, M., Stanley, H. E., and Goldberger, A. L. (1994). Mosaic organization of DNA nucleotides. *Phys. Rev. E*, 49:1685–1689.
- Prass, T. S. and Pumi, G. (2020). DCCA: Detrended Fluctuation and Detrended Cross-Correlation Analysis. R package version 0.1.1.
- Prass, T. S. and Pumi, G. (2021). On the behavior of the DFA and DCCA in trend-stationary processes. *Journal of Multivariate Analysis*, 182:104703.
- Prass, T. S. and Pumi, G. (2023). PPMiss: Copula-Based Estimator for Long-Range Dependent Processes under Missing Data. R package version 0.1.1.
- Pumi, G., Prass, T. S., and Lopes, S. R. C. (2023). A novel copula-based approach for parametric estimation of univariate time series through its covariance decay. *Statistical Papers*, forthcoming.
- Rea, W., Oxley, L., Reale, M., and Brown, J. (2013). Not all estimators are born equal: The empirical properties of some estimators of long memory. *Mathematics and Computers in Simulation*, 93:29–42.

- Reisen, V., Abraham, B., and Lopes, S. R. C. (2001). Estimation of parameters in ARFIMA processes: a simulation study. *Communications in Statistics - Simulation and Computation*, 30(4):787–803.
- Robinson, P. M. (1995a). Gaussian semiparametric estimation of long range dependence. *The* Annals of statistics, pages 1630–1661.
- Robinson, P. M. (1995b). Log-periodogram regression of time series with long range dependence. *The annals of Statistics*, pages 1048–1072.
- Robinson, P. M. (2003). Time Series with Long Memory. Oxford University.
- Shimotsu, K. and Phillips, P. C. B. (2005). Exact local Whittle estimation of fractional integration. The Annals of Statistics, 33(4):1890 – 1933.
- Taqqu, M. S., Teverovsky, V., and Willinger, W. (1995). Estimators for long-range dependence: An empirical study. *Fractals*, 03(04):785–798.
- Veenstra, J. Q. (2012). Persistence and anti-persistence: theory and software. PhD thesis, Western University.
- Zeileis, A. and Grothendieck, G. (2005). zoo: S3 infrastructure for regular and irregular time series. Journal of Statistical Software, 14(6):1–27.

A Appendix A: Asymptotic theory of Pumi et al. (2023)'s estimator in the presence of missing data

In this section we develop the asymptotic theory of Pumi et al. (2023)'s estimator in the presence of missing data. Before we proceed with the asymptotic results, we need to introduce some notation whilst providing adequate mathematical framework and then defining the estimator Pumi et al. (2023). We develop the theory in its general form, as presented Pumi et al. (2023), but assuming that the underlying process is strongly stationary, which entails some simplifications.

Mathematical Framework

We start by specifying the underlying copulas. We choose a family of parametric copulas $\{C_{\theta}\}_{\theta\in\Theta}$, with $\Theta\subseteq\mathbb{R}$ with non-empty interior, and such that $\lim_{\theta\to a} C_{\theta} = \Pi$ (the independence copula) for some $a\in\Theta'$. The limit is to be understood as the adequate lateral limit if $a\notin \operatorname{int}(\Theta)$. Also assume that there exist a set $D\subseteq\Theta$ with non-empty interior such that $a\in D'$ and C_{θ} is twice continuously differentiable in $\theta\in D$. Let $\{\theta_n\}_{n\in\mathbb{N}^*}$ be a sequence in D such that $\lim_{n\to\infty} \theta_n = a$.

Next we specify the stochastic process we are interested in. Let $\{X_n\}_{n\in\mathbb{N}}$ be a process for which X_n is identically distributed with common absolutely continuous distribution Fsuch that $\mathbb{E}(X_1^2) < \infty$. Assume that the copula related to (X_0, X_k) is C_{θ_k} , where $\{\theta_k\}_{k\in\mathbb{N}^*}$ is a sequence in D satisfying $\lim_{n\to\infty} \theta_n = a$. Assume further that $\operatorname{Cov}(X_0, X_n) \sim R(n, \eta)$, where $R(n, \eta)$ is a given continuous function such that $R(n, \eta) \to 0$, as n goes to infinity, and $\eta \in S \subseteq \mathbb{R}^p$ is some (identifiable) parameter of interest. Also assume that $\theta_n - a \sim L(n, \eta)$, where $L(n, \eta)$ is a given continuous function satisfying $L(n, \eta) \to 0$, as n goes to infinity. Finally, assume that $0 < K_1 < \infty$ and $K_2 < \infty$, where

$$K_1 = \iint_{I^2} \frac{1}{F'(F^{(-1)}(u))F'(F^{(-1)}(v))} \lim_{\theta \to a} \frac{\partial C_\theta(u,v)}{\partial \theta_i} \, du dv, \tag{A.13}$$

and

$$K_{2} = \iint_{I^{2}} \frac{1}{F'(F^{(-1)}(u))F'(F^{(-1)}(v))} \lim_{\theta \to a} \frac{\partial^{2}C_{\theta}(u,v)}{\partial\theta^{2}} \, du dv.$$
(A.14)

Definition of the estimator

Let x_1, \dots, x_n be a realization of a strongly stationary $\{X_n\}_{n \in \mathbb{N}}$, as specified above, with no missing data. In order to estimate η , Pumi et al. (2023) suggests the following multi-step procedure.

- 1. Given $\{C_{\theta}\}_{\theta\in\Theta}$ as discussed above, chose a method to perform parameter estimation in the family.
- 2. Chose estimators \hat{F}_n , \hat{F}_n^{-1} and \hat{F}'_n of the underlying unknown distribution F, quantile function F^{-1} and density function F', respectively. Estimate \hat{K}_1 and \hat{K}_2 by plugging in these estimators into (A.13) and (A.14), respectively. We must have $0 < \hat{K}_1 < \infty$ and $\hat{K}_2 < \infty$.
- 3. Set $y_i := \hat{F}_n(x_i)$, for $i = 1, \dots, n$.
- 4. Let s and m be two integers satisfying 1 < s < m < n. For each $\ell \in \{s, \dots, m\}$, form a sequence $\{\boldsymbol{u}_k^{(\ell)}\}_{k=1}^{n-\ell}$ by setting $\boldsymbol{u}_i^{(\ell)} := (y_i, y_{i+\ell}) \in [0, 1]^2, i = 1, \dots, n - \ell$. From these pseudo-observations, estimate of the copula parameter θ_ℓ , denoted by $\hat{\theta}_\ell(n)$.
- 5. Let $\mathscr{D}: \mathbb{R}^{m-s+1} \times \mathbb{R}^{m-s+1} \to [0,\infty)$, be a given function measuring the distance between two vectors in \mathbb{R}^{m-s+1} . Let $\widehat{L}_{s,m}(n) := \widehat{K}_1(\widehat{\theta}_s(n) - a, \cdots, \widehat{\theta}_m(n) - a)'$ and $R_{s,m}(\eta) := (R(s, \eta), \cdots, R(m, \eta))'$. The estimator $\widehat{\eta}_{s,m}(n)$ of η is then defined as

$$\hat{\boldsymbol{\eta}}_{s,m}(n) := \operatorname*{argmin}_{\boldsymbol{\eta} \in S} \left\{ \mathscr{D} \left(\widehat{\boldsymbol{L}}_{s,m}(n), \boldsymbol{R}_{s,m}(\boldsymbol{\eta}) \right) \right\}.$$
(A.15)

In practice, the estimation procedure in 1 and the estimators in 2 can be any reasonable ones. For instance, 1 can be attained using inference for the margins, maximum likelihood, inversion of Kendall's tau or Spearman's rho, etc - see Joe (1997) and Nelsen (2013). Further conditions are necessary when data is missing or when large sample results are of interest (see below). The metric \mathscr{D} in 5 has little effect in parameter estimation in practice, but its form is crucial for large sample inference.

In 3, s and m delimit the number of lags used for inference regarding the copula parameter. This provides great flexibility when tailoring the methodology to a specific model. For instance, under short range dependence, it may be that only a few lags may be enough to provide good estimates of the parameters. It also helps in the context of smaller samples, since the pseudo observations in 4 contain $n - \ell$ pairs, for $s \leq \ell \leq m$, which can be very small for m close to n. In this context one can choose a smaller value of m to counter this effect. We refer the reader to Section 3 of Pumi et al. (2023) for more details regarding the definition of the estimator.

Large Sample Theory

To show the consistency and to study the asymptotic distribution of the method, we need the estimators in 2 to be able to handle missing data. These conditions are presented in assumption \mathbf{M} below, which permeates the discussion that follows.

M. Suppose that $\{X_n\}_{n \in \mathbb{N}}$ is strongly stationary and let x_1, \dots, x_n be a sample from this process with missing values. For all $s \leq h \leq m$, let

 $w_{n,h} :=$ number of pairs (x_t, x_{t+h}) with no missing entries, $1 \le t \le n-h$,

and assume that

$$w_{n,h} \longrightarrow \infty, \quad \text{as} \quad n \to \infty.$$
 (A.16)

Assume further that the estimators presented in 2 can be calculated in the context of missing data.

Now we are in conditions to state the conditions necessary to show consistency, which are essentially the same as in Pumi et al. (2023).

- **A1.** \hat{F}_n , \hat{F}'_n and \hat{F}_n^{-1} are consistent estimators of F, F' and F^{-1} in the sense that $\hat{F}_n(x) \xrightarrow{P} F(x)$, $\hat{F}'_n(x) \xrightarrow{P} F'(x)$, for all $x \in \mathbb{R}$, and $\hat{F}_n^{-1}(u) \xrightarrow{P} F^{-1}(u)$, for all $u \in I$, and such that $\hat{K}_1 \xrightarrow{P} K_1$, as n tends to infinity.
- **A2**. The estimator of the copula parameter at lag k, $\hat{\theta}_k(n)$, satisfies $\hat{\theta}_k(n) \xrightarrow{P} \theta_k^0$, as $n \to \infty$, for all $s \le k \le m$, where θ_k^0 denotes the true copula parameter at lag k.
- A3. The space $(\mathbb{R}^{m-s+1}, \mathscr{D})$ is a metric space and \mathscr{D} is equivalent to the usual metric in \mathbb{R}^{m-s+1} .

Theorem A.1. Let η_0 denote the true parameter. Under assumptions M, A1-A3, $\hat{\eta}_{s,m}(n) \xrightarrow{P} \eta_0$, as n tends to infinity.

Proof: Observe that conditions **M** and **A1** imply that $\hat{K}_1 \xrightarrow{P} K_1$, as $n \to \infty$, as the number of non-missing observations goes to infinity as $n \to \infty$ under (A.16). Additionally, under **M**, (A.16) implies that for any $s \leq \ell \leq m$, the amount of pairs $\boldsymbol{u}_i^{(\ell)} := (y_i, y_{i+\ell}) \in [0, 1]^2$, for $i = 1, \dots, n - \ell$, that contain non-missing entries goes to infinity as $n \to \infty$. Hence, **A2** implies that $\hat{\theta}_{\ell}(n) - a \xrightarrow{P} \theta_{\ell}^0 - a \in \mathbb{R}$, as n tends to infinity, so that $\hat{\boldsymbol{L}}_{s,m}(n) \xrightarrow{P} \boldsymbol{R}_{s,m}(\boldsymbol{\eta}_0)$. The rest of the proof now carries on as in the proof of theorem 4.1 in Pumi et al. (2023).

To provide a central limit theorem for the Pumi et al. (2023)'s estimator, we need to strengthen the assumptions. Again η_0 denotes the true parameter to be estimated.

- **B1**. There exist a positive integer k_0 such that, as a function of $\boldsymbol{\eta}$, $L(k, \cdot) : \mathbb{R}^p \to \mathbb{R}$ is twice differentiable in a neighborhood $\Omega_0 \subseteq \mathbb{R}^p$ of $\boldsymbol{\eta}_0$ and $\boldsymbol{a}_k \boldsymbol{a}'_k$ is positive definite, where, $\boldsymbol{a}_k = \frac{\partial L(k, \boldsymbol{\eta})}{\partial \boldsymbol{\eta}}$, for all $k > k_0$ and $\boldsymbol{\eta} \in \Omega_0$.
- **B2**. There exists a positive integer k_1 , a neighborhood $\Omega_1 \subseteq \mathbb{R}^p$ of η_0 and a sequence $b_n \to \infty$ such that the copula parameter estimator at lag k, $\hat{\theta}_k(n)$, satisfies,

$$b_n(\hat{\theta}_k(n) - a - L(k, \boldsymbol{\eta})) \stackrel{d}{\longrightarrow} Z_k, \quad \forall \ k \ge k_1, \ \boldsymbol{\eta} \in \Omega_1,$$

with $\mathbb{E}(Z_k^2) < \infty$. Furthermore, we assume that the random variables $\{\hat{\theta}_k(n)\}_{k,n}$ and $\{Z_k\}_k$ are defined in the same probability space for all $k \geq k_1$ and n.

B3. \mathscr{D} is the Euclidean distance, that is, $\mathscr{D}(\boldsymbol{u}, \boldsymbol{v}) = \sum_{i=1}^{p} (v_i - u_i)^2, \, \boldsymbol{u}, \boldsymbol{v} \in \mathbb{R}^p.$

For a complete discussion regarding these assumptions, see section 4 of Pumi et al. (2023). It is interesting to notice that the rate of convergence in assumption **B2** defines the convergence rate for the estimator. Contrarily to Theorem A.1, the limiting distribution depends heavily on the metric \mathscr{D} applied, so that Pumi et al. (2023) considered the Euclidean distance, assumption **B3** here.

Theorem A.2. Suppose that assumptions M, B1 – B3 hold. Then,

$$b_n(\widehat{\boldsymbol{\eta}}_{s,m}(n) - \boldsymbol{\eta}_0) \xrightarrow{d} \sum_{k=0}^{m-s} \boldsymbol{\tau}_{s+k} Z_{s+k}, \qquad (A.17)$$

as n tends to infinity, for all s and m, where $\boldsymbol{\tau}_{s+k} = \left[\sum_{j=0}^{m-s} \boldsymbol{a}_{s+j} \boldsymbol{a}_{s+j}'\right]^{-1} \boldsymbol{a}_{s+k}$.

Proof: Without loss of generality, assume that a = 0. Let $s > s_0 = \max\{k_0, k_1\}$ and $\Omega = \Omega_0 \cap \Omega_1$. Denote by $\hat{\theta}_{\ell}(n)$ the estimated value of the copula parameter θ_{ℓ} calculated considering non-missing pairs $\boldsymbol{u}_i^{(\ell)} := (y_i, y_{i+\ell}) \in [0, 1]^2$, for $i = 1, \dots, n - \ell$ and $s \leq \ell \leq m$. Let $\hat{\boldsymbol{\eta}}$ the estimated value of $\boldsymbol{\eta}$ obtained from these estimates. With **B3** in mind, define

$$S_{s,m}(\boldsymbol{\eta};n) = \sum_{k=0}^{m-s} \left(\widehat{\theta}_{s+k}(n) - L(s+k,\boldsymbol{\eta})\right)^2,$$

and observe that **M** implies that the number of non-missing observations goes to infinity as $n \to \infty$ in view of (A.16), so that

$$\mathbf{0} = \frac{\partial S_{s,m}(\boldsymbol{\eta};n)}{\partial \boldsymbol{\eta}}\Big|_{\widehat{\boldsymbol{\eta}}} = \frac{\partial S_{s,m}(\boldsymbol{\eta};n)}{\partial \boldsymbol{\eta}}\Big|_{\boldsymbol{\eta}_0} + \left(\frac{\partial^2 S_{s,m}(\boldsymbol{\eta};n)}{\partial \boldsymbol{\eta} \partial \boldsymbol{\eta}'}\Big|_{\overline{\boldsymbol{\eta}}}\right)(\widehat{\boldsymbol{\eta}} - \boldsymbol{\eta}_0),$$

almost surely, as *n* tends to infinity, for some $\overline{\eta} \in \Omega$ such that $\|\overline{\eta} - \eta_0\| \leq \|\widehat{\eta} - \eta_0\|$. The remaining of the proof is analogous to the proof of theorem 4.2 in Pumi et al. (2023).

Estimation of Long-Range Dependent Models with Missing Data: to Impute or not to Impute?

Supplementary Material

Guilherme Pumi^{a,*}, Gladys Choque Ulloa^a and Taiane Schaedler Prass^a

This supplementary material is intended to present some complementary results to the ones presented in the parent paper. Henceforth, We shall apply the same notation, nomenclature, and abbreviations as in the main paper without further mention or reference. To the interested reader we refer the main paper for details.

The results presented here cover two main scenarios. The first one is the case of simulated ARFIMA(0, d, 0), for which we present the cases the case $d \in \{0.1, 0.4\}$ in the main paper and $d \in \{0.2, 0.3\}$ here. The results presented here are very similar to the ones presented in the paper. The discussion presented in the paper for the case d = 0.1 apply viz-a-viz to the case d = 0.2 here. The results for d = 0.3 presented here are very similar to those for d = 0.4 discussed in the main paper.

The second main scenario is the case of simulated ARFIMA(1, d, 1), which is presented here. Since the estimators applied in the paper are semiparametric, the AR and MA contribution to the estimation of d should, in principle, be negligible. It is indeed the case and the results of every scenario simulated for the ARFIMA(1, d, 1) are similar to the respective ARFIMA(0, d, 0)counterpart. Hence, the discussion presented in the paper for the ARFIMA(0, d, 0) apply viza-viz to the (1, d, 1) case presented here.

^{*}Corresponding author. This Version: December 22, 2023

^aPrograma de Pós-Graduação em Estatística - Universidade Federal do Rio Grande do Sul.

E-mails: guilherme.pumi@ufrgs.br (G. Pumi), gladyschoqueulloa7@gmail.com (G. U. Choque) and taianeprass@ufrgs.br (T. S. Prass)

1 ARFIMA(0, d, 0) scenario



Figure 1: Box plot of the adjusted model ARFIMA (0, d, 0) for d = 0.2.



Figure 2: Box plot of the adjusted model ARFIMA (0, d, 0) for d = 0.3.

| | | | | d = 0.2 | 2 | | | | |
|-----------|------------|-------|-------|---------|-------|-------|-------|-------|-------|
| | E-time ton | | | | Mis | sing | | | |
| Type | Estimator | 0 | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 0.6 | 0.7 |
| | Full | 0.160 | 0.164 | 0.164 | 0.170 | 0.181 | 0.190 | 0.197 | 0.225 |
| Intrincia | Abry | 0.167 | 0.171 | 0.171 | 0.175 | 0.178 | 0.182 | 0.185 | 0.203 |
| Intrinsic | PP.G | 0.188 | 0.188 | 0.187 | 0.186 | 0.187 | 0.183 | 0.181 | 0.177 |
| | PP.F | 0.181 | 0.181 | 0.180 | 0.179 | 0.180 | 0.177 | 0.175 | 0.173 |
| | LoMPE | 0.164 | 0.157 | 0.151 | 0.144 | 0.145 | 0.136 | 0.127 | 0.127 |
| | Full | 0.160 | 0.146 | 0.130 | 0.114 | 0.098 | 0.083 | 0.064 | 0.048 |
| | Abry | 0.167 | 0.152 | 0.136 | 0.121 | 0.105 | 0.089 | 0.070 | 0.053 |
| | PP.G | 0.188 | 0.161 | 0.131 | 0.102 | 0.114 | 0.092 | 0.030 | 0.015 |
| | PP.F | 0.181 | 0.152 | 0.120 | 0.090 | 0.095 | 0.066 | 0.023 | 0.011 |
| Mean | LoMPE | 0.164 | 0.151 | 0.136 | 0.121 | 0.106 | 0.091 | 0.074 | 0.060 |
| | DFA | 0.164 | 0.156 | 0.143 | 0.132 | 0.121 | 0.104 | 0.087 | 0.066 |
| | GPH | 0.158 | 0.148 | 0.128 | 0.111 | 0.094 | 0.069 | 0.042 | 0.011 |
| | LW | 0.123 | 0.118 | 0.108 | 0.102 | 0.096 | 0.082 | 0.070 | 0.058 |
| | ELW | 0.130 | 0.120 | 0.106 | 0.094 | 0.085 | 0.072 | 0.071 | 0.081 |
| | R/S | 0.216 | 0.207 | 0.200 | 0.189 | 0.181 | 0.168 | 0.154 | 0.136 |
| | Full | 0.160 | 0.212 | 0.268 | 0.333 | 0.407 | 0.494 | 0.587 | 0.707 |
| | Abry | 0.167 | 0.212 | 0.262 | 0.321 | 0.389 | 0.471 | 0.563 | 0.685 |
| | PP.G | 0.188 | 0.200 | 0.210 | 0.221 | 0.233 | 0.243 | 0.258 | 0.277 |
| | PP.F | 0.181 | 0.194 | 0.206 | 0.219 | 0.234 | 0.249 | 0.269 | 0.295 |
| Linear | LoMPE | 0.164 | 0.210 | 0.260 | 0.318 | 0.383 | 0.461 | 0.551 | 0.669 |
| | DFA | 0.164 | 0.160 | 0.156 | 0.155 | 0.156 | 0.152 | 0.162 | 0.184 |
| | GPH | 0.158 | 0.155 | 0.143 | 0.135 | 0.132 | 0.118 | 0.124 | 0.116 |
| | LW | 0.123 | 0.118 | 0.109 | 0.104 | 0.096 | 0.084 | 0.080 | 0.077 |
| | ELW | 0.130 | 0.127 | 0.120 | 0.112 | 0.109 | 0.096 | 0.096 | 0.101 |
| | R/S | 0.216 | 0.217 | 0.219 | 0.222 | 0.228 | 0.234 | 0.246 | 0.263 |
| | Full | 0.160 | 0.204 | 0.250 | 0.302 | 0.358 | 0.420 | 0.485 | 0.563 |
| | Abry | 0.167 | 0.205 | 0.245 | 0.293 | 0.346 | 0.407 | 0.474 | 0.557 |
| | PP.G | 0.188 | 0.198 | 0.209 | 0.220 | 0.234 | 0.247 | 0.265 | 0.286 |
| | PP.F | 0.181 | 0.192 | 0.204 | 0.217 | 0.232 | 0.249 | 0.270 | 0.295 |
| Random | LoMPE | 0.164 | 0.203 | 0.243 | 0.290 | 0.340 | 0.397 | 0.460 | 0.536 |
| | DFA | 0.164 | 0.156 | 0.148 | 0.143 | 0.143 | 0.139 | 0.144 | 0.163 |
| | GPH | 0.158 | 0.153 | 0.136 | 0.130 | 0.123 | 0.110 | 0.111 | 0.107 |
| | LW | 0.123 | 0.116 | 0.103 | 0.098 | 0.087 | 0.074 | 0.070 | 0.065 |
| | ELW | 0.130 | 0.124 | 0.115 | 0.109 | 0.103 | 0.091 | 0.090 | 0.090 |
| | R/S | 0.216 | 0.211 | 0.210 | 0.208 | 0.212 | 0.216 | 0.226 | 0.244 |

Table 1: Simulation results for the ARFIMA $\left(0,0.2,0\right)$ scenario.

| | | | | d = 0.3 | 3 | | | | |
|-----------|-----------|-------|-------|---------|-------|-------|-------|-------|-------|
| Trues | Estimator | | | | Mis | sing | | | |
| Type | Estimator | 0 | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 0.6 | 0.7 |
| | Full | 0.247 | 0.255 | 0.264 | 0.277 | 0.241 | 0.272 | 0.269 | 0.296 |
| | Abry | 0.256 | 0.261 | 0.263 | 0.264 | 0.266 | 0.270 | 0.273 | 0.283 |
| Intrinsic | PP.G | 0.284 | 0.283 | 0.284 | 0.282 | 0.282 | 0.280 | 0.278 | 0.275 |
| | PP.F | 0.278 | 0.277 | 0.278 | 0.277 | 0.277 | 0.275 | 0.274 | 0.271 |
| | LoMPE | 0.248 | 0.240 | 0.237 | 0.227 | 0.237 | 0.226 | 0.215 | 0.223 |
| | Full | 0.247 | 0.223 | 0.200 | 0.175 | 0.151 | 0.128 | 0.103 | 0.079 |
| | Abry | 0.256 | 0.233 | 0.210 | 0.185 | 0.161 | 0.138 | 0.112 | 0.086 |
| | PP.G | 0.284 | 0.255 | 0.220 | 0.182 | 0.167 | 0.123 | 0.058 | 0.033 |
| | PP.F | 0.278 | 0.245 | 0.207 | 0.166 | 0.144 | 0.102 | 0.046 | 0.023 |
| Mean | LoMPE | 0.248 | 0.226 | 0.204 | 0.181 | 0.159 | 0.138 | 0.115 | 0.090 |
| | DFA | 0.263 | 0.251 | 0.243 | 0.225 | 0.211 | 0.190 | 0.161 | 0.129 |
| | GPH | 0.273 | 0.257 | 0.247 | 0.227 | 0.203 | 0.179 | 0.149 | 0.108 |
| | LW | 0.216 | 0.208 | 0.203 | 0.190 | 0.179 | 0.167 | 0.147 | 0.125 |
| | ELW | 0.252 | 0.235 | 0.219 | 0.199 | 0.176 | 0.159 | 0.138 | 0.121 |
| | R/S | 0.280 | 0.271 | 0.261 | 0.250 | 0.236 | 0.220 | 0.200 | 0.175 |
| | Full | 0.247 | 0.293 | 0.346 | 0.406 | 0.473 | 0.551 | 0.643 | 0.755 |
| | Abry | 0.256 | 0.296 | 0.342 | 0.395 | 0.457 | 0.531 | 0.621 | 0.733 |
| | PP.G | 0.284 | 0.288 | 0.292 | 0.297 | 0.302 | 0.308 | 0.316 | 0.328 |
| | PP.F | 0.278 | 0.283 | 0.289 | 0.295 | 0.303 | 0.312 | 0.326 | 0.345 |
| Linear | LoMPE | 0.248 | 0.289 | 0.337 | 0.388 | 0.450 | 0.522 | 0.607 | 0.712 |
| | DFA | 0.263 | 0.258 | 0.256 | 0.250 | 0.246 | 0.241 | 0.242 | 0.254 |
| | GPH | 0.273 | 0.266 | 0.262 | 0.256 | 0.248 | 0.238 | 0.234 | 0.225 |
| | LW | 0.216 | 0.210 | 0.206 | 0.197 | 0.189 | 0.178 | 0.167 | 0.157 |
| | ELW | 0.252 | 0.246 | 0.242 | 0.236 | 0.224 | 0.215 | 0.207 | 0.198 |
| | R/S | 0.280 | 0.279 | 0.279 | 0.279 | 0.280 | 0.281 | 0.286 | 0.295 |
| | Full | 0.247 | 0.284 | 0.324 | 0.367 | 0.415 | 0.468 | 0.527 | 0.593 |
| | Abry | 0.256 | 0.287 | 0.322 | 0.360 | 0.405 | 0.458 | 0.518 | 0.589 |
| | PP.G | 0.284 | 0.287 | 0.291 | 0.296 | 0.301 | 0.309 | 0.319 | 0.333 |
| | PP.F | 0.278 | 0.282 | 0.287 | 0.293 | 0.301 | 0.310 | 0.323 | 0.342 |
| Random | LoMPE | 0.248 | 0.280 | 0.317 | 0.354 | 0.398 | 0.448 | 0.503 | 0.567 |
| | DFA | 0.263 | 0.254 | 0.249 | 0.239 | 0.231 | 0.223 | 0.221 | 0.227 |
| | GPH | 0.273 | 0.263 | 0.256 | 0.250 | 0.238 | 0.223 | 0.217 | 0.209 |
| | LW | 0.216 | 0.207 | 0.200 | 0.188 | 0.176 | 0.164 | 0.151 | 0.136 |
| | ELW | 0.252 | 0.245 | 0.236 | 0.228 | 0.212 | 0.203 | 0.190 | 0.180 |
| | R/S | 0.280 | 0.274 | 0.269 | 0.265 | 0.262 | 0.260 | 0.263 | 0.276 |

Table 2: Simulated results for the $\operatorname{ARFIMA}(0,0.3,0)$ scenario

2 ARFIMA(1, d, 1) scenario



Figure 3: Box plot of the adjusted model ARFIMA (1, d, 1) for d = 0.1.



Figure 4: Box plot of the adjusted model ARFIMA (1, d, 1) for d = 0.2.



Figure 5: Box plot of the adjusted model ARFIMA (1, d, 1) for d = 0.3.



Figure 6: Box plot of the adjusted model ARFIMA (1, d, 1) for d = 0.4.

| | | | | d = 0.1 | L | | | | |
|-----------|--------------|-------|-------|---------|-------|--------|--------|--------|--------|
| | Dation of an | | | | Mis | sing | | | |
| Type | Estimator | 0 | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 0.6 | 0.7 |
| | Full | 0.076 | 0.077 | 0.086 | 0.088 | 0.109 | 0.113 | 0.116 | 0.152 |
| | Abry | 0.081 | 0.083 | 0.086 | 0.089 | 0.096 | 0.103 | 0.116 | 0.142 |
| Intrinsic | PP.G | 0.092 | 0.091 | 0.091 | 0.090 | 0.088 | 0.086 | 0.084 | 0.073 |
| | PP.F | 0.086 | 0.086 | 0.086 | 0.085 | 0.083 | 0.081 | 0.080 | 0.071 |
| | LoMPE | 0.085 | 0.080 | 0.076 | 0.071 | 0.070 | 0.066 | 0.062 | 0.057 |
| | Full | 0.076 | 0.068 | 0.060 | 0.052 | 0.045 | 0.035 | 0.027 | 0.019 |
| | Abry | 0.081 | 0.073 | 0.064 | 0.056 | 0.049 | 0.039 | 0.031 | 0.022 |
| | PP.G | 0.092 | 0.075 | 0.057 | 0.043 | 0.062 | 0.054 | 0.012 | 0.007 |
| | PP.F | 0.086 | 0.069 | 0.051 | 0.037 | 0.053 | 0.055 | 0.009 | 0.005 |
| Moon | LoMPE | 0.085 | 0.078 | 0.070 | 0.063 | 0.055 | 0.047 | 0.041 | 0.029 |
| Mean | DFA | 0.072 | 0.066 | 0.057 | 0.053 | 0.047 | 0.037 | 0.024 | 0.008 |
| | GPH | 0.034 | 0.025 | 0.012 | 0.002 | -0.010 | -0.030 | -0.047 | -0.086 |
| | LW | 0.047 | 0.045 | 0.041 | 0.039 | 0.038 | 0.032 | 0.030 | 0.020 |
| | ELW | 0.044 | 0.041 | 0.037 | 0.036 | 0.039 | 0.044 | 0.050 | 0.065 |
| | R/S | 0.143 | 0.138 | 0.134 | 0.130 | 0.125 | 0.117 | 0.109 | 0.098 |
| | Full | 0.076 | 0.133 | 0.197 | 0.269 | 0.350 | 0.441 | 0.546 | 0.673 |
| | Abry | 0.081 | 0.131 | 0.188 | 0.254 | 0.331 | 0.418 | 0.521 | 0.650 |
| | PP.G | 0.092 | 0.120 | 0.143 | 0.164 | 0.182 | 0.200 | 0.221 | 0.247 |
| | PP.F | 0.086 | 0.115 | 0.139 | 0.161 | 0.183 | 0.205 | 0.233 | 0.266 |
| Lincor | LoMPE | 0.085 | 0.135 | 0.192 | 0.256 | 0.329 | 0.412 | 0.512 | 0.632 |
| Linear | DFA | 0.072 | 0.072 | 0.072 | 0.078 | 0.083 | 0.094 | 0.108 | 0.134 |
| | GPH | 0.034 | 0.032 | 0.029 | 0.026 | 0.029 | 0.028 | 0.031 | 0.036 |
| | LW | 0.047 | 0.046 | 0.040 | 0.038 | 0.039 | 0.033 | 0.032 | 0.031 |
| | ELW | 0.044 | 0.041 | 0.037 | 0.036 | 0.039 | 0.033 | 0.034 | 0.036 |
| | R/S | 0.143 | 0.149 | 0.157 | 0.167 | 0.178 | 0.192 | 0.209 | 0.234 |
| | Full | 0.076 | 0.128 | 0.184 | 0.244 | 0.311 | 0.380 | 0.454 | 0.538 |
| | Abry | 0.081 | 0.126 | 0.177 | 0.233 | 0.296 | 0.365 | 0.442 | 0.531 |
| | PP.G | 0.092 | 0.118 | 0.141 | 0.163 | 0.184 | 0.206 | 0.231 | 0.261 |
| | PP.F | 0.086 | 0.112 | 0.136 | 0.160 | 0.183 | 0.208 | 0.236 | 0.271 |
| Dandom | LoMPE | 0.085 | 0.131 | 0.180 | 0.235 | 0.293 | 0.359 | 0.431 | 0.512 |
| nandom | DFA | 0.072 | 0.068 | 0.067 | 0.070 | 0.072 | 0.083 | 0.101 | 0.130 |
| | GPH | 0.034 | 0.030 | 0.023 | 0.021 | 0.025 | 0.021 | 0.029 | 0.041 |
| | LW | 0.047 | 0.044 | 0.038 | 0.035 | 0.033 | 0.029 | 0.028 | 0.031 |
| | ELW | 0.044 | 0.040 | 0.036 | 0.035 | 0.035 | 0.029 | 0.031 | 0.037 |
| | R/S | 0.143 | 0.145 | 0.150 | 0.156 | 0.164 | 0.177 | 0.193 | 0.222 |

Table 3: Simulation results for the $\operatorname{ARFIMA}(1, 0.1, 1)$ scenario.

| | | | | d = 0.2 | 2 | | | | |
|-----------|------------|-------|-------|---------|-------|-------|-------|-------|-------|
| | E-time ton | | | | Mis | sing | | | |
| Type | Estimator | 0 | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 0.6 | 0.7 |
| | Full | 0.160 | 0.164 | 0.164 | 0.170 | 0.181 | 0.190 | 0.197 | 0.225 |
| Intrincia | Abry | 0.167 | 0.171 | 0.171 | 0.175 | 0.178 | 0.182 | 0.185 | 0.203 |
| Intrinsic | PP.G | 0.188 | 0.187 | 0.186 | 0.185 | 0.185 | 0.182 | 0.179 | 0.177 |
| | PP.F | 0.180 | 0.180 | 0.180 | 0.179 | 0.178 | 0.176 | 0.174 | 0.172 |
| | LoMPE | 0.164 | 0.156 | 0.151 | 0.142 | 0.146 | 0.138 | 0.130 | 0.125 |
| | Full | 0.160 | 0.146 | 0.130 | 0.114 | 0.098 | 0.083 | 0.064 | 0.048 |
| | Abry | 0.167 | 0.152 | 0.136 | 0.121 | 0.105 | 0.089 | 0.070 | 0.053 |
| | PP.G | 0.188 | 0.161 | 0.131 | 0.105 | 0.108 | 0.068 | 0.027 | 0.018 |
| | PP.F | 0.180 | 0.152 | 0.120 | 0.092 | 0.091 | 0.072 | 0.021 | 0.013 |
| Mean | LoMPE | 0.164 | 0.149 | 0.136 | 0.121 | 0.106 | 0.091 | 0.075 | 0.057 |
| | DFA | 0.172 | 0.164 | 0.152 | 0.138 | 0.125 | 0.109 | 0.089 | 0.069 |
| | GPH | 0.154 | 0.141 | 0.126 | 0.106 | 0.087 | 0.068 | 0.043 | 0.003 |
| | LW | 0.124 | 0.118 | 0.110 | 0.102 | 0.094 | 0.082 | 0.072 | 0.057 |
| | ELW | 0.135 | 0.126 | 0.112 | 0.097 | 0.088 | 0.079 | 0.076 | 0.086 |
| | R/S | 0.214 | 0.207 | 0.198 | 0.189 | 0.178 | 0.167 | 0.152 | 0.135 |
| | Full | 0.160 | 0.212 | 0.268 | 0.333 | 0.407 | 0.494 | 0.587 | 0.707 |
| | Abry | 0.167 | 0.212 | 0.262 | 0.321 | 0.389 | 0.471 | 0.563 | 0.685 |
| | PP.G | 0.188 | 0.199 | 0.210 | 0.220 | 0.231 | 0.243 | 0.257 | 0.277 |
| | PP.F | 0.180 | 0.193 | 0.206 | 0.218 | 0.232 | 0.248 | 0.268 | 0.295 |
| Linear | LoMPE | 0.164 | 0.208 | 0.259 | 0.317 | 0.385 | 0.463 | 0.554 | 0.666 |
| | DFA | 0.172 | 0.170 | 0.164 | 0.159 | 0.160 | 0.161 | 0.165 | 0.189 |
| | GPH | 0.154 | 0.148 | 0.142 | 0.133 | 0.127 | 0.120 | 0.115 | 0.117 |
| | LW | 0.124 | 0.119 | 0.110 | 0.103 | 0.096 | 0.087 | 0.080 | 0.076 |
| | ELW | 0.135 | 0.131 | 0.124 | 0.118 | 0.113 | 0.102 | 0.098 | 0.101 |
| | R/S | 0.214 | 0.216 | 0.218 | 0.220 | 0.225 | 0.232 | 0.242 | 0.261 |
| | Full | 0.160 | 0.204 | 0.250 | 0.302 | 0.358 | 0.420 | 0.485 | 0.563 |
| | Abry | 0.167 | 0.205 | 0.245 | 0.293 | 0.346 | 0.407 | 0.474 | 0.557 |
| | PP.G | 0.188 | 0.198 | 0.209 | 0.219 | 0.232 | 0.246 | 0.264 | 0.287 |
| | PP.F | 0.180 | 0.191 | 0.204 | 0.216 | 0.231 | 0.248 | 0.269 | 0.297 |
| Random | Knight | 0.164 | 0.202 | 0.243 | 0.290 | 0.341 | 0.399 | 0.463 | 0.535 |
| | DFA | 0.172 | 0.166 | 0.157 | 0.149 | 0.145 | 0.145 | 0.150 | 0.172 |
| | GPH | 0.154 | 0.146 | 0.137 | 0.130 | 0.115 | 0.114 | 0.103 | 0.110 |
| | LW | 0.124 | 0.116 | 0.105 | 0.097 | 0.086 | 0.076 | 0.069 | 0.066 |
| | ELW | 0.135 | 0.129 | 0.121 | 0.113 | 0.105 | 0.095 | 0.089 | 0.092 |
| | R/S | 0.214 | 0.211 | 0.208 | 0.207 | 0.208 | 0.212 | 0.225 | 0.244 |

Table 4: Simulation results for the $\operatorname{ARFIMA}(1,0.2,1)$ scenario.

| | | | | d = 0.3 | 3 | | | | |
|-----------|-----------|-------|-------|---------|-------|-------|-------|-------|-------|
| Trme | Estimator | | | | Mis | sing | | | |
| Type | Estimator | 0 | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 0.6 | 0.7 |
| | Full | 0.247 | 0.255 | 0.264 | 0.277 | 0.241 | 0.272 | 0.269 | 0.296 |
| T . · · | Abry | 0.256 | 0.261 | 0.263 | 0.264 | 0.266 | 0.270 | 0.273 | 0.283 |
| Intrinsic | PP.G | 0.283 | 0.282 | 0.282 | 0.281 | 0.282 | 0.280 | 0.277 | 0.276 |
| | PP.F | 0.277 | 0.277 | 0.276 | 0.276 | 0.276 | 0.275 | 0.273 | 0.273 |
| | LoMPE | 0.247 | 0.239 | 0.234 | 0.225 | 0.235 | 0.224 | 0.210 | 0.218 |
| | Full | 0.247 | 0.223 | 0.200 | 0.175 | 0.151 | 0.128 | 0.103 | 0.079 |
| | Abry | 0.256 | 0.233 | 0.210 | 0.185 | 0.161 | 0.138 | 0.112 | 0.086 |
| | PP.G | 0.283 | 0.253 | 0.217 | 0.180 | 0.171 | 0.094 | 0.058 | 0.029 |
| | PP.F | 0.277 | 0.244 | 0.205 | 0.164 | 0.149 | 0.082 | 0.046 | 0.021 |
| Mean | LoMPE | 0.247 | 0.225 | 0.202 | 0.181 | 0.159 | 0.136 | 0.113 | 0.088 |
| | DFA | 0.268 | 0.257 | 0.243 | 0.229 | 0.212 | 0.196 | 0.167 | 0.138 |
| | GPH | 0.273 | 0.259 | 0.246 | 0.225 | 0.208 | 0.176 | 0.150 | 0.112 |
| | LW | 0.215 | 0.208 | 0.199 | 0.189 | 0.178 | 0.164 | 0.146 | 0.126 |
| | ELW | 0.250 | 0.235 | 0.218 | 0.192 | 0.177 | 0.150 | 0.135 | 0.120 |
| | R/S | 0.280 | 0.271 | 0.261 | 0.249 | 0.237 | 0.221 | 0.201 | 0.177 |
| | Full | 0.247 | 0.293 | 0.346 | 0.406 | 0.473 | 0.551 | 0.643 | 0.755 |
| | Abry | 0.256 | 0.296 | 0.342 | 0.395 | 0.457 | 0.531 | 0.621 | 0.733 |
| | PP.G | 0.283 | 0.287 | 0.292 | 0.295 | 0.301 | 0.307 | 0.315 | 0.328 |
| | PP.F | 0.277 | 0.282 | 0.288 | 0.293 | 0.303 | 0.312 | 0.325 | 0.345 |
| Linear | LoMPE | 0.247 | 0.288 | 0.333 | 0.386 | 0.448 | 0.520 | 0.603 | 0.711 |
| | DFA | 0.268 | 0.262 | 0.258 | 0.252 | 0.247 | 0.247 | 0.247 | 0.258 |
| | GPH | 0.273 | 0.269 | 0.265 | 0.252 | 0.250 | 0.235 | 0.227 | 0.233 |
| | LW | 0.215 | 0.210 | 0.204 | 0.193 | 0.186 | 0.173 | 0.165 | 0.160 |
| | ELW | 0.250 | 0.245 | 0.240 | 0.227 | 0.222 | 0.208 | 0.204 | 0.201 |
| | R/S | 0.280 | 0.279 | 0.279 | 0.278 | 0.280 | 0.282 | 0.286 | 0.297 |
| | Full | 0.247 | 0.284 | 0.324 | 0.367 | 0.415 | 0.468 | 0.527 | 0.593 |
| | Abry | 0.256 | 0.287 | 0.322 | 0.360 | 0.405 | 0.458 | 0.518 | 0.589 |
| | PP.G | 0.283 | 0.286 | 0.291 | 0.294 | 0.301 | 0.308 | 0.318 | 0.334 |
| | PP.F | 0.277 | 0.281 | 0.286 | 0.291 | 0.300 | 0.309 | 0.323 | 0.343 |
| Random | LoMPE | 0.247 | 0.279 | 0.313 | 0.352 | 0.398 | 0.446 | 0.500 | 0.567 |
| | DFA | 0.268 | 0.259 | 0.252 | 0.238 | 0.231 | 0.227 | 0.227 | 0.233 |
| | GPH | 0.273 | 0.266 | 0.263 | 0.245 | 0.239 | 0.221 | 0.215 | 0.216 |
| | LW | 0.215 | 0.207 | 0.200 | 0.186 | 0.175 | 0.158 | 0.149 | 0.139 |
| | ELW | 0.250 | 0.242 | 0.237 | 0.219 | 0.210 | 0.195 | 0.191 | 0.182 |
| | R/S | 0.280 | 0.274 | 0.269 | 0.263 | 0.261 | 0.261 | 0.264 | 0.277 |

Table 5: Simulation results for the $\operatorname{ARFIMA}(1, 0.3, 1)$ scenario.

| | | | | d = 0.4 | 1 | | | | |
|------------|-------------|-------|-------|---------|-------|-------|-------|-------|-------|
| | E-time to a | | | | Mis | sing | | | |
| Type | Estimator | 0 | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 0.6 | 0.7 |
| | Full | 0.332 | 0.341 | 0.347 | 0.346 | 0.353 | 0.358 | 0.366 | 0.365 |
| | Abry | 0.344 | 0.350 | 0.352 | 0.355 | 0.358 | 0.360 | 0.362 | 0.366 |
| Intrinsic | PP.G | 0.384 | 0.383 | 0.383 | 0.383 | 0.382 | 0.380 | 0.380 | 0.378 |
| | PP.F | 0.381 | 0.380 | 0.380 | 0.381 | 0.380 | 0.378 | 0.379 | 0.378 |
| | LoMPE | 0.332 | 0.325 | 0.322 | 0.314 | 0.336 | 0.325 | 0.312 | 0.334 |
| | Full | 0.332 | 0.294 | 0.260 | 0.227 | 0.196 | 0.165 | 0.133 | 0.102 |
| | Abry | 0.344 | 0.307 | 0.274 | 0.241 | 0.209 | 0.178 | 0.144 | 0.112 |
| | PP.G | 0.384 | 0.354 | 0.317 | 0.277 | 0.253 | 0.102 | 0.114 | 0.062 |
| | PP.F | 0.381 | 0.348 | 0.307 | 0.261 | 0.229 | 0.086 | 0.092 | 0.046 |
| Mean | LoMPE | 0.332 | 0.297 | 0.265 | 0.234 | 0.205 | 0.175 | 0.145 | 0.117 |
| Mean | DFA | 0.367 | 0.355 | 0.344 | 0.326 | 0.305 | 0.274 | 0.251 | 0.213 |
| | GPH | 0.393 | 0.379 | 0.363 | 0.346 | 0.323 | 0.292 | 0.267 | 0.220 |
| | LW | 0.317 | 0.311 | 0.303 | 0.295 | 0.280 | 0.263 | 0.246 | 0.218 |
| | ELW | 0.380 | 0.368 | 0.353 | 0.332 | 0.303 | 0.276 | 0.234 | 0.204 |
| | R/S | 0.341 | 0.332 | 0.321 | 0.308 | 0.291 | 0.271 | 0.249 | 0.220 |
| | Full | 0.332 | 0.375 | 0.424 | 0.481 | 0.544 | 0.617 | 0.702 | 0.807 |
| | Abry | 0.344 | 0.381 | 0.423 | 0.473 | 0.531 | 0.599 | 0.681 | 0.786 |
| | PP.G | 0.384 | 0.385 | 0.386 | 0.388 | 0.389 | 0.391 | 0.396 | 0.402 |
| | PP.F | 0.381 | 0.383 | 0.386 | 0.389 | 0.392 | 0.397 | 0.406 | 0.417 |
| Linear | LoMPE | 0.332 | 0.371 | 0.414 | 0.462 | 0.520 | 0.587 | 0.664 | 0.765 |
| Lincar | DFA | 0.367 | 0.364 | 0.362 | 0.355 | 0.350 | 0.342 | 0.345 | 0.347 |
| | GPH | 0.393 | 0.391 | 0.385 | 0.382 | 0.375 | 0.363 | 0.359 | 0.353 |
| | LW | 0.317 | 0.314 | 0.311 | 0.305 | 0.298 | 0.286 | 0.278 | 0.269 |
| | ELW | 0.380 | 0.376 | 0.372 | 0.364 | 0.356 | 0.345 | 0.336 | 0.329 |
| | R/S | 0.341 | 0.339 | 0.336 | 0.335 | 0.332 | 0.330 | 0.332 | 0.335 |
| | Full | 0.332 | 0.363 | 0.397 | 0.436 | 0.476 | 0.522 | 0.571 | 0.628 |
| | Abry | 0.344 | 0.369 | 0.398 | 0.432 | 0.470 | 0.513 | 0.565 | 0.626 |
| | PP.G | 0.384 | 0.384 | 0.386 | 0.387 | 0.388 | 0.390 | 0.397 | 0.404 |
| | PP.F | 0.381 | 0.382 | 0.384 | 0.386 | 0.389 | 0.393 | 0.401 | 0.412 |
| Bandom | LoMPE | 0.332 | 0.359 | 0.389 | 0.422 | 0.460 | 0.502 | 0.547 | 0.605 |
| Italiuolli | DFA | 0.367 | 0.361 | 0.355 | 0.344 | 0.334 | 0.323 | 0.321 | 0.315 |
| | GPH | 0.393 | 0.388 | 0.382 | 0.374 | 0.363 | 0.354 | 0.343 | 0.331 |
| | LW | 0.317 | 0.312 | 0.307 | 0.298 | 0.287 | 0.273 | 0.260 | 0.244 |
| | ELW | 0.380 | 0.373 | 0.369 | 0.358 | 0.345 | 0.329 | 0.318 | 0.305 |
| | R/S | 0.341 | 0.334 | 0.328 | 0.321 | 0.316 | 0.310 | 0.311 | 0.314 |

Table 6: Simulation results for the ARFIMA(1, 0.4, 1) scenario.