

An alternate proof of Payan's theorem on cubelike graphs

Jonathan Cervantes^a and Mike Krebs^b

^aUniversity of California, Riverside, Dept. of Mathematics, Skye Hall, 900 University Ave.,
Riverside, CA 92521, jcerv092@ucr.edu

^bCalifornia State University, Los Angeles, Dept. of Mathematics, 5151 State University
Drive, Los Angeles, CA 91711, mkrebs@calstatela.edu

November 15, 2023

Keywords— graph, chromatic number, abelian group, Cayley graph, cube-like graph, Payan's theorem

Abstract

A cubelike graph is a Cayley graph on the product $\mathbb{Z}_2 \times \cdots \times \mathbb{Z}_2$ of the integers modulo 2 with itself finitely many times. In 1992, Payan proved that no cubelike graph can have chromatic number 3. The authors of the present paper previously developed a general matrix method for studying chromatic numbers of Cayley graphs on abelian groups. In this note, we apply this method of Heuberger matrices to give an alternate proof of Payan's theorem.

1 Introduction

Given a finite set A , we take $\mathcal{P}(A)$ to be the power set of A . We have that $\mathcal{P}(A)$ is an abelian group under the operation of symmetric difference, that is, $X \triangle Y = (X \setminus Y) \cup (Y \setminus X)$. A *cubelike graph* is a Cayley graph whose underlying group is $\mathcal{P}(A)$. Equivalently, writing $A = \{x_1, \dots, x_n\}$ and identifying the set $X \subset A$ with the n -tuple whose i th component is 1 if $x_i \in X$ and is 0 otherwise, a cubelike graph can be regarded as a Cayley graph whose underlying group is an n -fold product $\mathbb{Z}_2^n = \mathbb{Z}_2 \times \cdots \times \mathbb{Z}_2$, where \mathbb{Z}_2 is the group of the integers modulo 2 under addition.

Chromatic numbers of cubelike graphs have been studied by many authors. One notable result is due to Payan [4], who proved that the chromatic number of a nonbipartite cubelike graph is always at least 4. That is, the chromatic number of a cubelike graph cannot equal 3. Publications with other results on chromatic numbers of cube-like graphs include [3] and [2, Section 9.7].

Payan's proof is rather clever. It is, however, somewhat *ad hoc*. The purpose of the present note is to furnish an

alternate proof of Payan’s theorem, one that may lend itself naturally to generalizations.

Indeed, in [1], the authors put forward a general method for approaching the problem of finding the chromatic number of a Cayley graph on an abelian group. We show that Payan’s theorem falls out quite naturally as a byproduct of this “method of Heuberger matrices.”

The other key ingredient in our proof is a special case of Payan’s theorem due to Sokolová [5], who computed that even-dimensional cubes-with-diagonals (defined below) have chromatic number 4. The key idea of our proof of Payan’s theorem is that if a cubelike graph is nonbipartite, then there is a graph homomorphism to it from an even-dimensional cube-with-diagonals. The Heuberger matrices make transparent the existence of this homomorphism.

This note depends heavily on [1], which we will refer to frequently. The reader should assume that all notation, terminology, and theorems used but not explained here are explained there.

2 Payan’s theorem

In this section we prove the following theorem.

Theorem 2.1 ([4]). *A cube-like graph cannot have chromatic number 3.*

Throughout this section we take cube-like graph to be a Cayley graph on \mathbb{Z}_2^n .

A special case of Theorem 2.1 had previously been proven by Sokolová in [5]. We will derive Payan’s theorem from Sokolová’s theorem, and for that reason we begin by discussing the latter.

For a positive integer n , the n -dimensional cube-with-diagonals graph Q_n^d is defined by

$$Q_n^d = \text{Cay}(\mathbb{Z}_2^n, \{e_1, \dots, e_n, w_n\}),$$

where e_j is the n -tuple in \mathbb{Z}_2^n with 1 in the j th entry and 0 everywhere else, and w_n is the n -tuple in \mathbb{Z}_2^n with 1 in every entry. We can visualize Q_n^d as a hypercube with edges (called “diagonals,” hence the name and the superscript ‘ d ’) added to join each pair of antipodal vertices. Sokolová proved that for n even, Q_n^d has chromatic number 4. We present here a condensed version of the proof in [5] of this result.

Theorem 2.2 ([5]). *If n is even, then $\chi(Q_n^d) = 4$.*

Proof. First observe that $(x_1, \dots, x_n) \mapsto (x_1, x_2 + \dots + x_n)$ defines a group homomorphism from \mathbb{Z}_2^n to \mathbb{Z}_2^2 mapping $\{e_1, \dots, e_n, w_n\}$ to $\{(1, 0), (0, 1), (1, 1)\}$. So this defines a graph homomorphism from Q_n^d to $Q_2^d \cong K_4$, the complete graph on 4 vertices. Hence $\chi(Q_n^d) \leq 4$.

Next we show that Q_n^d is not properly 3-colorable. We do so by induction. For the base case ($n = 2$), we saw previously that $Q_2^d \cong K_4$, which is not properly 3-colorable. Now assume that Q_n^d is not properly 3-colorable, and we will show that Q_{n+2}^d is not properly 3-colorable. Suppose to the contrary that $c: \mathbb{Z}_2^{n+2} \rightarrow \mathbb{Z}_3$ is a proper 3-coloring. For two

tuples $v = (v_1, \dots, v_j)$ and $u = (u_1, \dots, u_k)$, we define $v * u = (v_1, \dots, v_j, u_1, \dots, u_k)$. Define $c': \mathbb{Z}_2^n \rightarrow \mathbb{Z}_3$ by $c'(v) = k$ if $\{c(v * (0, 0)), c((v + w_n) * (1, 0))\}$ equals either $\{k\}$ or $\{k, k + 1\}$. A straightforward case-by-case analysis shows that c' is a proper 3-coloring of Q_n^d , which is a contradiction. \square

Remark 2.3. We briefly digress to remark that Sokolová's theorem can be restated as follows. In any (not necessarily proper) 3-coloring of the vertices of an even-dimensional hypercube, there must exist two antipodal vertices, both of which are assigned the same color. Stated this way, it brings to mind various topological theorems such as the hairy ball theorem and the Borsuk–Ulam theorem. We wonder whether there might be some connection between Sokolová's combinatorial result and one or more of these facts from topology, perhaps along the lines of the connection between Sperner's lemma and the Brouwer fixed point theorem. \square

Using Heuberger matrices, we will now see how Sokolová's theorem implies Payan's theorem. The key idea is to show that every nonbipartite cube-like graph contains a homomorphic image of an even-dimensional cube-with-diagonals graph.

Proof of Theorem 2.1. Let $X = \text{Cay}(\mathbb{Z}_2^n, S)$ be a nonbipartite cube-like graph. Because $2x = 0$ for all $x \in S$, there is a Heuberger matrix M_X associated to X whose last m columns are $2e_1, \dots, 2e_m$, where $m = |S|$. That is, M_X has the form $(A \mid 2I_m)$ for some integer matrix A . Here I_m is the $m \times m$ identity matrix. Using column operations as in [1, Lemma 2.6], we have that

$$(A \mid 2I_m)_X^{\text{SACG}} \cong (A' \mid 2I_m)_X^{\text{SACG}},$$

for some matrix A' whose entries are all in $\{0, 1\}$. Because X is nonbipartite, by [1, Lemma 2.11], some column y of A' contains an odd number z of nonzero entries. Hence by [1, Lemma 2.10, parts (4) and (5)], we have homomorphisms

$$(w_z^t \mid 2I_z)_Y^{\text{SACG}} \xrightarrow[\tau_1]{\circledast} (y \ 2e_{i_1} \cdots 2e_{i_z})^{\text{SACG}} \xrightarrow[\tau_2]{\circledast} (A' \mid 2I_m)_X^{\text{SACG}}$$

where i_1, \dots, i_z are the indices of the nonzero entries of y , and w_z^t is a column vector of length z with a 1 in every entry. For τ_1 , we insert zero rows as appropriate; for τ_2 we append the requisite columns. So $\chi(Y) \leq \chi(X)$ by [1, Lemma 2.9]. If $z = 1$, then X has loops and is not properly colorable. So assume $z \geq 3$. Observe that $Y \cong Q_{z-1}^d$. An application of Theorem 2.2 then completes the proof. \square

References

- [1] J. Cervantes and M. Krebs, *Chromatic numbers of Cayley graphs of abelian groups: A matrix method*, www.arxiv.org.
- [2] Tommy R. Jensen and Bjarne Toft, *Graph coloring problems*, Wiley-Interscience Series in Discrete Mathematics and Optimization, John Wiley & Sons, Inc., New York, 1995, A Wiley-Interscience Publication.

- [3] Janne I. Kokkala and Patric R. J. Östergård, *The chromatic number of the square of the 8-cube*, Math. Comp. **87** (2018), no. 313, 2551–2561.
- [4] Charles Payan, *On the chromatic number of cube-like graphs*, Discrete Math. **103** (1992), no. 3, 271–277.
- [5] Marie Sokolová, *The chromatic number of extended odd graphs is four*, Časopis Pěst. Mat. **112** (1987), no. 3, 308–311.