

Trinomial Planar Functions on Cubic and Quartic Extensions of Finite Fields

Ruikai Chen^{1,2} Sihem Mesnager^{1,2,3}

¹Department of Mathematics, University of Paris VIII, F-93526 Saint-Denis

²Laboratory Analysis, Geometry and Applications, LAGA, University Sorbonne Paris Nord, CNRS, UMR 7539, F-93430, Villetaneuse, France

³Telecom Paris, Polytechnic institute of Paris, 91120 Palaiseau, France

Emails: chen.rk@outlook.com smesnager@univ-paris8.fr

Abstract

Planar functions, introduced by Dembowski and Ostrom, are functions from a finite field to itself that give rise to finite projective planes. They exist, however, only for finite fields of odd characteristics. They have attracted much attention in the last decade thanks to their interest in theory and those deep and various applications in many fields. This paper focuses on planar trinomials over cubic and quartic extensions of finite fields. Our achievements are obtained using connections with quadratic forms and classical algebraic tools over finite fields. Furthermore, given the generality of our approach, the methodology presented could be employed to drive more planar functions on some finite extension fields.

Keywords. Planar function, Finite field, Extension Field, Algebraic curve, Polynomial.

Mathematics Subject Classification: 11R32, 12E10, 11G20, 11R11, 11R16, 11T06, 12E10, 51E15.

1 Introduction

Consider an extension $\mathbb{F}_{q^n}/\mathbb{F}_q$ of finite fields of odd characteristic. The function induced by a polynomial f over \mathbb{F}_{q^n} is called bent, if for every $b \in \mathbb{F}_{q^n}$,

$$\left| \sum_{t \in \mathbb{F}_{q^n}} \psi(f(t) - bt) \right|^2 = q^n,$$

where ψ is the canonical additive character of \mathbb{F}_{q^n} . In addition, it is called a planar function if $f(x+c) - f(x)$ is a permutation on \mathbb{F}_{q^n} for every $c \in \mathbb{F}_{q^n}^*$. As a well-known result, $f(x)$

is planar if and only if $cf(x)$ is bent for every $c \in \mathbb{F}_{q^n}^*$. Constructions of planar functions can be found in [11, 13, 19, 32].

Historically, planar functions with odd characteristics have been introduced by Dembowski and Ostrom ([14]), aiming to design finite projective planes with odd characteristics. These functions are also connected with important mathematical objects, such as permutation polynomials over finite fields. The notion of planar functions coincides with the one of perfect nonlinear (PN) functions from \mathbb{F}_q to itself (see, e.g., [7, Chapter 6, pp. 296], [26] and [31]). Planar functions have deep applications in different areas of mathematics for communication (coding theory, cryptography, and related topics such as combinatorics).

In particular, it has been shown that planar functions are helpful in DES-like cryptosystems ([25]). In addition, it has been demonstrated in several articles (see, e.g., [8, 16, 30]) that planar functions allow constructing error-correcting codes, which are then employed to design secret sharing schemes. Planar functions are also applied to the constructions of authentication codes ([15]), constant composition codes ([18]) and signal sets ([17]). Besides, planar functions induce many combinatorial objects such as skew Hadamard difference sets, Paley type partial difference sets ([29]), relative difference sets ([20]) and symplectic spreads ([1]). Also, it has been shown in two main papers [33] and [27] that a natural analogue notion of planar functions over fields of characteristic 2 could be introduced and studied, which is sometimes called pseudo-planar or modified planar (see [7, Chapter 6, pp. 296]). Such planar functions have similar properties concerning relative difference sets and finite geometries as their counterparts in odd characteristics. Finally, by employing various methods, much attention has been addressed to the problem of the existence and non-existence of planar mappings. For example, the classification of planar monomials over fields of prime square order has been obtained by Coulter ([10]). Very recent results on this framework are given by Bergman et al. ([6]) devoted to classifying planar monomials over fields of order a prime cubed, and by Beierle and Felke toward the classification of planar monomials over finite fields of small order ([4, 5]).

Note that most of known planar functions on \mathbb{F}_{q^n} are of the form

$$f(x) = \sum_{0 \leq j \leq i < n} a_{ij} x^{q^i + q^j},$$

also known as Dembowski-Ostrom polynomials. Some work has been done on this class of polynomials. For example, those planar binomial functions on \mathbb{F}_{q^3} are classified in [23]. Also, the sufficient and necessary conditions for the polynomial $x^{q^2+1} + x^{q+1} + ux^2$ ($u \in \mathbb{F}_{q^3}$) to be planar are given in [22] and [12], as well as the polynomial $x^{q^2+1} + ax^{q+1} + bx^2$ considered in [3], however, with coefficients restricted in \mathbb{F}_q .

In this paper, we will extend these results. In Section 3, we provide characterizations of two classes of planar trinomials on \mathbb{F}_{q^3} . The polynomials from [22], [12] and [3] are special cases of those in Theorem 4. In Section 4, we construct planar trinomials on \mathbb{F}_{q^4} . The method used there can be applied to more planar functions on extensions of finite fields.

2 An Overview of the Approach

Consider the polynomial ring $\mathbb{F}_{q^n}[x]$ with an indeterminate x . When discussing functions on \mathbb{F}_{q^n} , we identify them with polynomials in $\mathbb{F}_{q^n}[x]$ modulo $x^{q^n} - x$. We focus on planar functions on \mathbb{F}_{q^n} are of the form

$$f(x) = \sum_{0 \leq j \leq i < n} a_{ij} x^{q^i + q^j}.$$

In particular, the monomial x^{q^k+1} is planar if $\frac{n}{\gcd(k,n)}$ is odd. For the general case, we can investigate $cf(x)$ for every $c \in \mathbb{F}_{q^n}^*$. In fact, $\text{Tr}(cf(x))$ is a quadratic form on \mathbb{F}_{q^n} as a vector space over \mathbb{F}_q , where Tr denotes the trace function from \mathbb{F}_{q^n} to \mathbb{F}_q ; moreover,

$$\text{Tr}(cf(x)) = \text{Tr} \left(\sum_{0 \leq j \leq i < n} (ca_{ij})^{q^{n-j}} x^{q^i - j + 1} \right).$$

It has been proved (e.g., in [21, Proposition 2]) that $cf(x)$ is a bent function on \mathbb{F}_{q^n} if and only if the corresponding quadratic form is nondegenerate, if and only if

$$\left| \sum_{t \in \mathbb{F}_{q^n}} \psi(cf(t)) \right|^2 = q^n.$$

To determine whether f is a planar function, we calculate the determinant of the quadratic form $\text{Tr}(cf(x))$ for every $c \in \mathbb{F}_{q^n}^*$. Specifically, it can be characterized given the coefficients as follows.

Lemma 1. *Let $f(x) = ax^{q+1} + bx^2$ for some $a, b \in \mathbb{F}_{q^n}^*$. Then*

$$\left| \sum_{t \in \mathbb{F}_{q^n}} \psi(f(t)) \right|^2 = q^n$$

if and only if

- $\text{Tr}(a)^2 - 4\text{N}(b) \neq 0$ in the case $n = 2$, and
- $4\text{N}(b) + \text{N}(a) - \text{Tr}(a^2b^{q^2}) \neq 0$ in the case $n = 3$,

where Tr and N are the trace and the norm function from \mathbb{F}_{q^n} to \mathbb{F}_q respectively.

Proof. This follows from [21, Proposition 2] or [9, Section 3], where the corresponding matrix is

$$\begin{pmatrix} 2b & a^q + a \\ a^q + a & 2b^q \end{pmatrix}$$

in the case $n = 2$, and

$$\begin{pmatrix} 2b & a & a^{q^2} \\ a & 2b^q & a^q \\ a^{q^2} & a^q & 2b^{q^2} \end{pmatrix}$$

in the case $n = 3$. □

There is an equivalence relation preserving the properties of planar functions. Two Dembowski-Ostrom polynomials f_0 and f_1 over \mathbb{F}_{q^n} are called equivalent if there exist linearized permutation polynomial L_0 and L_1 over \mathbb{F}_{q^n} such that $f_0(x) = L_0(f_1(L_1(x)))$. Note that all those planar functions on \mathbb{F}_{q^3} are equivalent to either x^2 or x^{q+1} according to [12, Lemma 2.1].

The following result will be useful for the remaining part.

Lemma 2 ([24, Theorem 5.3]). *For any $r \in \mathbb{F}_q$ and $s \in \mathbb{F}_q^*$, there exists an element $\alpha \in \mathbb{F}_{q^3}^*$ with $\text{Tr}(\alpha) = r$ and $\text{N}(\alpha) = s$. In other words, given $a \in \mathbb{F}_{q^3}^*$, the function $\text{Tr}(ax^{q-1})$ takes all values in \mathbb{F}_q .*

3 Planar Trinomials on Cubic Extensions

Let Tr and N denote the trace and the norm function from \mathbb{F}_{q^3} to \mathbb{F}_q respectively. We characterize two classes of trinomial planar functions in what follows.

Proposition 3. *For $A, B \in \mathbb{F}_{q^3}^*$ and $r \in \mathbb{F}_q$, the function $\text{Tr}(Ax^{q-1} + Bx^{1-q}) + r$ has no zero in $\mathbb{F}_{q^3}^*$ if and only if $r = \frac{\text{N}(A)+\text{N}(B)}{AB} \neq 0$.*

Proof. The equation $\text{Tr}(Ax^{q-1} + Bx^{1-q}) + r = 0$ is equivalent to $Ax^{q-1} + B^{q^2}x^{q^2-1} + u = 0$ and $\text{Tr}(u) = r$ for some $u \in \mathbb{F}_{q^3}$. Here $Ax^{q-1} + B^{q^2}x^{q^2-1} + u$ has a zero in $\mathbb{F}_{q^3}^*$ if and only if $B^{q^2}x^{q^2} + Ax^q + ux$ does not permute \mathbb{F}_{q^3} ; that is,

$$\text{N}(u) + \text{N}(A) + \text{N}(B) - \text{Tr}(ABu^{q^2}) = \begin{vmatrix} u & A & B^{q^2} \\ B & u^q & A^q \\ A^{q^2} & B^q & u^{q^2} \end{vmatrix} = 0.$$

Therefore, the function $\text{Tr}(Ax^{q-1} + Bx^{1-q}) + r$ has a zero in $\mathbb{F}_{q^3}^*$ if and only if

$$\text{Tr}(u) = r \quad \text{and} \quad \text{N}(u) + \text{N}(A) + \text{N}(B) - \text{Tr}(ABu^{q^2}) = 0 \tag{1}$$

for some $u \in \mathbb{F}_{q^3}$.

Suppose $AB \in \mathbb{F}_q$. If $r = \frac{\text{N}(A)+\text{N}(B)}{AB} \neq 0$, then $\text{Tr}(u) = r$ implies that $u \neq 0$ and

$$\begin{aligned} & \text{N}(u) + \text{N}(A) + \text{N}(B) - \text{Tr}(ABu^{q^2}) \\ &= \text{N}(u) + \text{N}(A) + \text{N}(B) - ABr \\ &= \text{N}(u) \neq 0. \end{aligned}$$

If $r = \frac{N(A)+N(B)}{AB} = 0$, then $u = 0$ implies (1). If $r \neq \frac{N(A)+N(B)}{AB}$, then by Lemma 2, there exists $u \in \mathbb{F}_{q^3}$ with $\text{Tr}(u) = r$ and

$$N(u) = -N(A) - N(B) + AB r = -N(A) - N(B) + \text{Tr}(ABu^2),$$

which means (1) holds.

Suppose $AB \notin \mathbb{F}_q$ and let $\beta_0, \beta_1, \beta_2$ be the dual basis of $1, AB, (AB)^2$ in $\mathbb{F}_{q^3}/\mathbb{F}_q$. For arbitrary $r \in \mathbb{F}_q$, if $u^{q^2} = r\beta_0 + r_1\beta_1 + r_2\beta_2$ for some $r_1, r_2 \in \mathbb{F}_q$, then $\text{Tr}(u) = r$ and

$$N(u) + N(A) + N(B) - \text{Tr}(ABu^{q^2}) = N(r\beta_0 + r_1\beta_1 + r_2\beta_2) + N(A) + N(B) - r_1.$$

It remains to show that the right side is zero for some $r_1, r_2 \in \mathbb{F}_q$. Let y be an element in some extension of $\mathbb{F}_q(x)$ with

$$N'(r\beta_0 + x\beta_1 + y\beta_2) + N(A) + N(B) - x = 0, \quad (2)$$

where

$$\begin{aligned} & N'(r\beta_0 + x\beta_1 + y\beta_2) \\ &= (r\beta_0 + x\beta_1 + y\beta_2)(r\beta_0^q + x\beta_1^q + y\beta_2^q)(r\beta_0^{q^2} + x\beta_1^{q^2} + y\beta_2^{q^2}). \end{aligned}$$

Assume $y \in \overline{\mathbb{F}_q}(x)$ and let v_∞ be the valuation at the infinite place of $\overline{\mathbb{F}_q}(x)$. If $v_\infty(y) > -1$, then $v_\infty(N'(r\beta_0 + x\beta_1 + y\beta_2)) = -3$. If $v_\infty(y) < -1$, then $v_\infty(N'(r\beta_0 + y\beta_1 + x\beta_2)) = 3v_\infty(y)$. Both cases contradict the fact $v_\infty(x - N(A) - N(B)) = -1$. Thus we have $v_\infty(y) = -1$. Note that y is integral over $\overline{\mathbb{F}_q}[x]$, which is an integrally closed domain. Then $y = \lambda x + \mu$ for some $\lambda, \mu \in \overline{\mathbb{F}_q}$, and

$$\begin{aligned} & N'(r\beta_0 + x\beta_1 + y\beta_2) \\ &= ((\beta_1 + \lambda\beta_2)x + r\beta_0 + \mu\beta_2)((\beta_1^q + \lambda\beta_2^q)x + r\beta_0^q + \mu\beta_2^q)((\beta_1^{q^2} + \lambda\beta_2^{q^2})x + r\beta_0^{q^2} + \mu\beta_2^{q^2}). \end{aligned}$$

Since $v_\infty(N'(r\beta_0 + x\beta_1 + y\beta_2)) = -1$, without loss of generality we have

$$\beta_1 + \lambda\beta_2 = \beta_1^q + \lambda\beta_2^q = 0.$$

Then $\lambda \in \mathbb{F}_q$, but β_1 and β_2 are linearly independent over \mathbb{F}_q , a contradiction. This shows that $\mathbb{F}_q(x, y)/\mathbb{F}_q(x)$ is an extension of function fields of degree 3 with constant field \mathbb{F}_q .

Consider the corresponding plane projective curve defined by (2), which is absolutely irreducible, for a cubic polynomial reducible over a field must have a root in it (see also [28, Corollary 3.6.8]). As a consequence of [2], the number N of rational points over \mathbb{F}_q of the curve satisfies

$$|N - q - 1| \leq (3 - 1)(3 - 2)\sqrt{q} = 2\sqrt{q},$$

which means

$$N \geq q + 1 - 2\sqrt{q} > 0,$$

as $q \geq 3$. Clearly, there is no point at infinity on this curve, so there exists at least one element in \mathbb{F}_{q^3} with desired properties. \square

Theorem 4. For $a, b \in \mathbb{F}_{q^3}^*$, the function $x^{q^2+1} + ax^{q+1} + bx^2$ is planar on \mathbb{F}_{q^3} if and only if $a = b^{q+1}$ with $N(b) \neq 1$, or $N(a) - 2ab^{q^2} + 1 = 0$ with $N(a)^2 \neq 1$.

Proof. For $c \in \mathbb{F}_{q^3}^*$, note that

$$\mathrm{Tr}(c(x^{q^2+1} + ax^{q+1} + bx^2)) = \mathrm{Tr}((c^q + ac)x^{q+1} + bcx^2),$$

and

$$\begin{aligned} & 4N(bc) + N(c^q + ac) - \mathrm{Tr}((c^q + ac)^2(bc)^{q^2}) \\ &= 4N(bc) + N(c) + N(ac) + \mathrm{Tr}(ac^{q^2+2}) + \mathrm{Tr}(a^{q+1}c^{q+2}) \\ &\quad - \mathrm{Tr}(b^{q^2}c^{q^2+2q} + 2ab^{q^2}c^{q^2+q+1} + a^2b^{q^2}c^{q^2+2}) \\ &= N(c)(4N(b) + 1 + N(a) - 2\mathrm{Tr}(ab^{q^2})) \\ &\quad + N(c)\mathrm{Tr}((a^{q^2+q} - b^{q^2})c^{q-1} + (a - a^2b^{q^2})c^{1-q}). \end{aligned}$$

By Lemma 1, the function is planar if and only if

$$\mathrm{Tr}(Ac^{q-1} + Bc^{1-q}) + 4N(b) + 1 + N(a) - 2\mathrm{Tr}(ab^{q^2}) \neq 0$$

for all $c \in \mathbb{F}_{q^3}^*$, where $A = a^{q^2+q} - b^{q^2}$ and $B = a - a^2b^{q^2}$. This happens only if $AB \neq 0$, for otherwise either $AB = 0 \neq A + B$, or $A = B = 0 \neq 4N(b) + 1 + N(a) - 2\mathrm{Tr}(ab^{q^2})$. The former is excluded in view of Lemma 2, and the latter implies $4N(b) + 1 + N(a) - 2\mathrm{Tr}(ab^{q^2}) = 0$ by a simple investigation. Then that is equivalent to

$$AB \neq 0 \quad \text{and} \quad \frac{N(A) + N(B)}{AB} = 4N(b) + 1 + N(a) - 2\mathrm{Tr}(ab^{q^2}) \quad (3)$$

by Proposition 3.

We claim that if $\delta = ab^{q^2} \in \mathbb{F}_q$, then

$$\begin{aligned} & N(A) + N(B) - AB(4N(b) + 1 + N(a) - 2\mathrm{Tr}(ab^{q^2})) \\ &= N(a)^{-1}\delta(\delta^2 - N(a))(N(a) - 2\delta + 1)^2. \end{aligned}$$

In this case $A = a^{-1}(N(a) - \delta)$ and $B = a(1 - \delta)$. Then

$$N(A) + N(B) = N(a)^{-1}(N(a) - \delta)^3 + N(a)(1 - \delta)^3,$$

and

$$\begin{aligned} & AB(4N(b) + 1 + N(a) - 2\mathrm{Tr}(ab^{q^2})) \\ &= (N(a) - \delta)(1 - \delta)(4N(a)^{-1}\delta^3 + 1 + N(a) - 6\delta) \\ &= N(a)^{-1}(N(a) - \delta)(1 - \delta)(4\delta^3 + N(a) + N(a)^2 - 6N(a)\delta). \end{aligned}$$

It is routine to expand

$$(N(a) - \delta)^3 + N(a)^2(1 - \delta)^3 - (N(a) - \delta)(1 - \delta)(4\delta^3 + N(a) + N(a)^2 - 6N(a)\delta)$$

and

$$\delta(\delta^2 - N(a))(N(a) - 2\delta + 1)^2,$$

which are actually equal.

Assume that (3) holds. Then $(N(a) - \delta)(1 - \delta) = AB \in \mathbb{F}_q$, which means $\delta \in \mathbb{F}_{q^2} \cap \mathbb{F}_{q^3} = \mathbb{F}_q$. By the above claim we have $\delta^2 = N(a)$ or $N(a) - 2\delta + 1 = 0$. If $\delta^2 = N(a)$, then

$$a = \frac{N(a)}{a^{q^2+q}} = \frac{(ab^{q^2})^{q^2+q}}{a^{q^2+q}} = b^{q+1},$$

and $N(b) = ab^{q^2} \neq 1$ as $AB \neq 0$. If $N(a) - 2\delta + 1 = 0$, then $N(a) - 1 = 2(\delta - 1) \neq 0$ and $N(a) + 1 = 2\delta \neq 0$. For the converse, if $a = b^{q+1}$ and $N(b) \neq 1$, then $\delta = N(b) \in \mathbb{F}_q^*$, and $\delta^2 = N(b)^2 = N(a)$ with $AB = (N(b)^2 - N(b))(1 - N(b)) \neq 0$. If $N(a) - 2ab^{q^2} + 1 = 0$ and $N(a)^2 \neq 1$, then $2\delta = N(a) + 1 \in \mathbb{F}_q^*$ and $4AB = (N(a) - 1)(1 - N(a)) \neq 0$. Then the proof is complete. \square

Using the same argument, we characterize another class of trinomials.

Theorem 5. *For $a, b \in \mathbb{F}_{q^3}^*$, the function $x^{q+1} + ax^{2q} + bx^2$ is planar on \mathbb{F}_{q^3} if and only if $4ab = 1$ and $N(2a) \neq -1$, or $4ab \neq 1$ and $N(a) + N(b) + ab = 0$.*

Proof. For $c \in \mathbb{F}_{q^3}^*$, note that

$$\text{Tr}(c(x^{q+1} + ax^{2q} + bx^2)) = \text{Tr}(cx^{q+1} + (a^{q^2}c^{q^2} + bc)x^2),$$

and

$$\begin{aligned} & 4N(a^{q^2}c^{q^2} + bc) + N(c) - \text{Tr}(c^2(a^{q^2}c^{q^2} + bc)^{q^2}) \\ &= 4(N(ac) + N(bc) + \text{Tr}((ac)^{q^2+q}(bc)^q) + \text{Tr}(ac(bc)^{q^2+1})) \\ & \quad + N(c) - \text{Tr}(a^{q^2}c^{q^2+2q} + b^{q^2}c^{q^2+2}) \\ &= 4N(c)(N(a) + N(b) + \text{Tr}(a^{q^2+q}b^qc^{q-1}) + \text{Tr}(ab^{q^2+1}c^{1-q})) \\ & \quad + N(c)(1 - \text{Tr}(a^{q^2}c^{q-1} + b^{q^2}c^{1-q})). \end{aligned}$$

Clearly the function is planar if and only if

$$\text{Tr}(Ac^{q-1} + Bc^{1-q}) + 4N(a) + 4N(b) + 1 \neq 0$$

for all $c \in \mathbb{F}_{q^3}^*$, where

$$A = 4a^{q^2+q}b^q - a^{q^2} = a^{q^2}(4(ab)^q - 1)$$

and

$$B = 4ab^{q^2+1} - b^{q^2} = b^{q^2}(4ab - 1).$$

If $4ab = 1$, then

$$4a(x^{q+1} + ax^{2q} + bx^2) = 4ax^{q+1} + 4a^2x^{2q} + x^2 = (2ax^q + x)^2.$$

This is planar on \mathbb{F}_{q^3} if and only if $2ax^q + x$ is a permutation polynomial of \mathbb{F}_{q^3} , i.e., $N(2a) \neq -1$ (this is equivalent to $N(a) + N(b) + ab \neq 0$).

Suppose now $4ab \neq 1$, which means $AB \neq 0$. Then by Proposition 3, the function $x^{q+1} + ax^{2q} + bx^2$ is planar if and only if

$$\frac{N(A) + N(B)}{AB} = 4N(a) + 4N(b) + 1.$$

If $ab \in \mathbb{F}_q$, then

$$N(A) + N(B) = N(a^{q^2}(4(ab)^q - 1)) + N(b^{q^2}(4ab - 1)) = (N(a) + N(b))(4ab - 1)^3,$$

and

$$AB = (ab)^{q^2}(4(ab)^q - 1)(4ab - 1) = ab(4ab - 1)^2.$$

Consequently,

$$\begin{aligned} & N(A) + N(B) - AB(4N(a) + 4N(b) + 1) \\ &= (N(a) + N(b))(4ab - 1)^3 - ab(4ab - 1)^2(4N(a) + 4N(b) + 1) \\ &= -(4ab - 1)^2(N(a) + N(b) + ab). \end{aligned}$$

If $ab \notin \mathbb{F}_q$, then

$$AB = (ab)^{q^2} \frac{N(4ab - 1)}{4(ab)^{q^2} - 1} = \frac{N(4ab - 1)}{4 - (ab)^{-q^2}} \notin \mathbb{F}_q.$$

The results follow immediately. □

4 Planar Trinomials on Quartic Extensions

In this section, let Tr and N denote the trace and the norm function from \mathbb{F}_{q^4} to \mathbb{F}_{q^2} respectively. A class of planar trinomials on \mathbb{F}_{q^4} can be constructed in a similar way.

Theorem 6. *For $a, b \in \mathbb{F}_{q^4}^*$, the function $x^{q^3+q} + ax^{q^2+1} + bx^2$ is planar on \mathbb{F}_{q^4} if*

- (1) $\text{Tr}(a) \neq 0$, $N(b) = N(a) - \text{Tr}(a)^{1-q}$, $1 - 4\text{Tr}(a)^{-1-q}$ is a nonzero square in \mathbb{F}_q , and $N(b)\text{Tr}(a)$ is a nonzero square in \mathbb{F}_{q^2} , or

(2) $\text{Tr}(a) = 0$, $\text{N}(b) = \text{N}(a) - \theta^{1-q}$ for some $\theta \in \mathbb{F}_{q^2}^*$ such that $\theta^{\frac{q^2-1}{2}} = (-1)^{\frac{q-1}{2}}$, and $\text{N}(b)\theta$ is a nonzero square in \mathbb{F}_{q^2} .

Proof. For the canonical additive character ψ of \mathbb{F}_{q^4} , we have

$$\psi(cx^{q^3+q} + acx^{q^2+1} + bcx^2) = \psi((c^q + ac)x^{q^2+1} + bcx^2).$$

By Lemma 1, the function $x^{q^3+q} + ax^{q^2+1} + bx^2$ is planar on \mathbb{F}_{q^4} if and only if

$$\text{Tr}(c^q + ac)^2 - 4\text{N}(bc) \neq 0$$

for every $c \in \mathbb{F}_{q^4}^*$. Suppose first that $a \in \mathbb{F}_{q^2}^*$, and let $d = \text{Tr}(c)^2 - 4\text{N}(c)$ and $t = \text{Tr}(c)$. Then

$$\begin{aligned} \text{Tr}(c^q + ac)^2 - 4\text{N}(bc) &= \text{Tr}(c)^{2q} + 2a\text{Tr}(c)^{q+1} + a^2\text{Tr}(c)^2 - 4\text{N}(bc) \\ &= t^{2q} + 2at^{q+1} + a^2t^2 + \text{N}(b)(d - t^2), \end{aligned}$$

and $\text{Tr}(c^q + ac)^2 - 4\text{N}(bc) = 0$ if and only if

$$d = -\text{N}(b)^{-1}(t^{2q} + 2at^{q+1} + (a^2 - \text{N}(b))t^2).$$

Note that $d \neq 0$ when $t = 0$, $d = 0$ when $c \in \mathbb{F}_{q^2}$ and d is a non-square in \mathbb{F}_{q^2} when $c \in \mathbb{F}_{q^4} \setminus \mathbb{F}_{q^2}$, so it suffices to show that

$$\text{N}(b)(t^{2q} + 2at^{q+1} + (a^2 - \text{N}(b))t^2)$$

is a nonzero square in \mathbb{F}_{q^2} for every $t \in \mathbb{F}_{q^2}^*$.

Assume the condition (1); that is, $\text{N}(b) = a^2 - a^{1-q}$, $1 - a^{-1-q} = \delta^2$ for some $\delta \in \mathbb{F}_q^*$, and $\text{N}(b)a$ is a nonzero square in \mathbb{F}_{q^2} . Let $\alpha = \beta^{-q}a^{-1}$ for some $\beta \in \mathbb{F}_{q^2}$ such that $\beta^{q+1} = \delta + 1$, where $\beta \neq 0$, for otherwise $a^{-1-q} = 1 - \delta^2 = 0$. It follows that

$$\begin{aligned} \alpha^{q+1} + \beta^{q+1} &= \frac{1}{(\delta + 1)a^{q+1}} + \delta + 1 \\ &= \frac{1 + (\delta + 1)^2 a^{q+1}}{(\delta + 1)a^{q+1}} \\ &= \frac{1 + (1 - a^{-1-q} + 2\delta + 1)a^{q+1}}{(\delta + 1)a^{q+1}} \\ &= 2, \end{aligned}$$

and then

$$\begin{aligned} &\text{N}(b)(t^{2q} + 2at^{q+1} + (a^2 - \text{N}(b))t^2) \\ &= \text{N}(b)a(a^{-1}t^{2q} + 2t^{q+1} + a^{-q}t^2) \\ &= \text{N}(b)a(\alpha\beta^qt^{2q} + (\alpha^{q+1} + \beta^{q+1})t^{q+1} + \alpha^q\beta t^2) \\ &= \text{N}(b)a(\alpha t^q + \beta t)^{q+1}. \end{aligned}$$

This is indeed a nonzero square in \mathbb{F}_{q^2} for all $t \in \mathbb{F}_{q^2}^*$, since $\alpha^{q+1} \neq \beta^{q+1}$ (otherwise $\delta + 1 = \beta^{q+1} = 1$), which means $\alpha t^q + \beta t \neq 0$.

Suppose now that $a \notin \mathbb{F}_{q^2}$. Noting that

$$\mathrm{Tr}\left(\frac{1}{a - a^{q^2}}\right) = \mathrm{Tr}\left(a \frac{a^{q^2}}{a - a^{q^2}}\right) = 0,$$

and

$$\mathrm{Tr}\left(a \frac{1}{a - a^{q^2}}\right) = \mathrm{Tr}\left(-\frac{a^{q^2}}{a - a^{q^2}}\right) = 1,$$

we have

$$c = \frac{1}{a - a^{q^2}} \mathrm{Tr}(ac) - \frac{a^{q^2}}{a - a^{q^2}} \mathrm{Tr}(c)$$

for $c \in \mathbb{F}_{q^4}^*$. Thus,

$$N(c) = N(a - a^{q^2})^{-1} (\mathrm{Tr}(ac)^2 - \mathrm{Tr}(a) \mathrm{Tr}(ac) \mathrm{Tr}(c) + N(a) \mathrm{Tr}(c)^2).$$

Let $t_0 = \mathrm{Tr}(c)$, $t_1 = \mathrm{Tr}(ac)$ and $\gamma = 4N(b)N(a - a^{q^2})^{-1}$, so that

$$\begin{aligned} & \mathrm{Tr}(c^q + ac)^2 - 4N(bc) \\ &= \mathrm{Tr}(c)^{2q} + 2\mathrm{Tr}(c)^q \mathrm{Tr}(ac) + \mathrm{Tr}(ac)^2 - 4N(bc) \\ &= t_0^{2q} + 2t_0^q t_1 + t_1^2 - \gamma(t_1^2 - \mathrm{Tr}(a)t_0 t_1 + N(a)t_0^2). \end{aligned}$$

Since $a \notin \mathbb{F}_{q^2}$, the map $c \mapsto (t_0, t_1)$ is bijective from \mathbb{F}_{q^4} to $\mathbb{F}_{q^2} \times \mathbb{F}_{q^2}$, and thus $\mathrm{Tr}(c^q + ac)^2 - 4N(bc) \neq 0$ for all $c \in \mathbb{F}_{q^4}^*$ if and only if

$$(t_0^{2q} + 2t_0^q t_1 + t_1^2) - \gamma(t_1^2 - \mathrm{Tr}(a)t_0 t_1 + N(a)t_0^2) \neq 0$$

for all $t_0, t_1 \in \mathbb{F}_{q^2}$ with t_0 or t_1 nonzero. If $t_0 = 0$, then $\mathrm{Tr}(c^q + ac)^2 - 4N(bc) = (1 - \gamma)t_1^2$, and if $\gamma = 1$, then $\mathrm{Tr}(c^q + ac)^2 - 4N(bc) = 0$ for all $t_1 \in \mathbb{F}_{q^2}$. Therefore, assume $\gamma \neq 1$ and then

$$\mathrm{Tr}(c^q + ac)^2 - 4N(bc) = (1 - \gamma)t_1^2 + (2t_0^q + \gamma \mathrm{Tr}(a)t_0)t_1 + t_0^{2q} - \gamma N(a)t_0^2$$

can be viewed as a quadratic polynomial in t_1 over \mathbb{F}_{q^2} , whose discriminant is

$$\begin{aligned} D &= (2t_0^q + \gamma \mathrm{Tr}(a)t_0)^2 - 4(1 - \gamma)(t_0^{2q} - \gamma N(a)t_0^2) \\ &= 4\gamma t_0^{2q} + 4\gamma \mathrm{Tr}(a)t_0^{q+1} + (\gamma^2 \mathrm{Tr}(a)^2 + 4(1 - \gamma)\gamma N(a))t_0^2 \\ &= 4\gamma t_0^{2q} + 4\gamma \mathrm{Tr}(a)t_0^{q+1} + (\gamma^2(\mathrm{Tr}(a)^2 - 4N(a)) + 4\gamma N(a))t_0^2 \\ &= 4\gamma t_0^{2q} + 4\gamma \mathrm{Tr}(a)t_0^{q+1} + 4\gamma(N(a) - N(b))t_0^2. \end{aligned}$$

Altogether, it suffices to show that $\gamma \neq 1$ and D is a non-square in \mathbb{F}_{q^2} for all $t_0 \in \mathbb{F}_{q^2}^*$.

Assuming the condition (1), we have $\text{Tr}(a) \neq 0$, $1 - 4\text{Tr}(a)^{-1-q} = \delta^2$ for some $\delta \in \mathbb{F}_q^*$, and $\gamma \neq 1$ since

$$4\text{N}(b) - \text{N}(a - a^{q^2}) = 4\text{N}(a) - 4\text{Tr}(a)^{1-q} + (a - a^{q^2})^2 = \text{Tr}(a)^2(1 - 4\text{Tr}(a)^{-1-q}) \neq 0.$$

Let $\alpha = \beta^{-q} \text{Tr}(a)^{-1}$ for some $\beta \in \mathbb{F}_{q^2}$ such that $\beta^{q+1} = \frac{\delta+1}{2}$ ($\beta \neq 0$ as easily seen). Then

$$\begin{aligned} \alpha^{q+1} + \beta^{q+1} &= \frac{2}{(\delta+1)\text{Tr}(a)^{q+1}} + \frac{\delta+1}{2} \\ &= \frac{4 + (\delta+1)^2 \text{Tr}(a)^{q+1}}{2(\delta+1)\text{Tr}(a)^{q+1}} \\ &= \frac{4 + (1 - 4\text{Tr}(a)^{-1-q} + 2\delta + 1)\text{Tr}(a)^{q+1}}{2(\delta+1)\text{Tr}(a)^{q+1}} \\ &= 1. \end{aligned}$$

Hence,

$$\begin{aligned} &\text{Tr}(a)^{-1}t_0^{2q} + t_0^{q+1} + \text{Tr}(a)^{-1}(\text{N}(a) - \text{N}(b))t_0^2 \\ &= \text{Tr}(a)^{-1}t_0^{2q} + t_0^{q+1} + \text{Tr}(a)^{-q}t_0^2 \\ &= \alpha\beta^q t_0^{2q} + (\alpha^{q+1} + \beta^{q+1})t_0^{q+1} + \alpha^q \beta t_0^2 \\ &= (\alpha t_0^q + \beta t_0)^{q+1}, \end{aligned}$$

and then

$$D = 4\gamma \text{Tr}(a)(\alpha t_0^q + \beta t_0)^{q+1}$$

is a non-square in \mathbb{F}_{q^2} for all $t_0 \in \mathbb{F}_{q^2}^*$, for

$$\gamma \text{Tr}(a) = -4(a - a^{q^2})^{-2} \text{N}(b) \text{Tr}(a)$$

is a non-square in \mathbb{F}_{q^2} and $\alpha t_0^q + \beta t_0 \neq 0$ as $\alpha^{q+1} \neq \beta^{q+1}$.

Assume the condition (2). Then $\text{Tr}(a) = 0$ and $\gamma \neq 1$ as

$$4\text{N}(b) - \text{N}(a - a^{q^2}) = 4\text{N}(b) - 4\text{N}(a) = -4\theta^{1-q} \neq 0.$$

Moreover,

$$D = 4\gamma t_0^{2q} + 4\gamma(\text{N}(a) - \text{N}(b))t_0^2 = 4\gamma(t_0^{2q} + \theta^{1-q}t_0^2) = 4\gamma\theta^{-q}(\theta^q t_0^{2q} + \theta t_0^2).$$

Here $\theta^q t_0^{2q} + \theta t_0^2 \neq 0$ for $t_0 \in \mathbb{F}_{q^2}^*$, for otherwise $\theta t_0^2 \notin \mathbb{F}_q$ and $-(\theta t_0^2)^{q+1} = (\theta t_0^2)^2$, which implies

$$(-1)^{\frac{q-1}{2}} \theta^{\frac{q^2-1}{2}} = (\theta t_0^2)^{q-1} \neq 1,$$

contradicting the assumption. We conclude that D is a non-square in \mathbb{F}_{q^2} for all $t_0 \in \mathbb{F}_{q^2}^*$, since

$$\gamma\theta^{-q} = -4(a - a^{q^2})^{-2} N(b)\theta^{-q}$$

is a non-square in \mathbb{F}_{q^2} and $\theta^q t_0^{2q} + \theta t_0^2 \in \mathbb{F}_q^*$. \square

We show that this class of planar functions are not equivalent to monomials in general. For simplicity, let q be a prime, so that we only need to consider the equivalence to x^2 . Assume that

$$L_0(x^{q^3+q} + ax^{q^2+1} + bx^2) = L_1(x)^2$$

for some linearized permutation polynomials L_0 and L_1 over \mathbb{F}_{q^4} , and without loss of generality, let the coefficient of x in $L_1(x)$ be nonzero. Comparing the degree, one easily finds that $L_1(x) = \alpha x^{q^2} + \beta x$ for some $\alpha, \beta \in \mathbb{F}_{q^4}$. Then

$$L_0(x^{q^3+q} + ax^{q^2+1} + bx^2) = \alpha^2 x^{2q^2} + 2\alpha\beta x^{q^2+1} + \beta^2 x^2.$$

It follows that

$$L_0(bx^2) = \alpha^2 x^{2q^2} + \beta^2 x^2,$$

and thus,

$$L_0(x) = \alpha^2 b^{-q^2} x^{q^2} + \beta^2 b^{-1} x.$$

On the other hand,

$$L_0(1)x^{q^3+q} + L_0(a)x^{q^2+1} = L_0(x^{q^3+q} + ax^{q^2+1}) = 2\alpha\beta x^{q^2+1},$$

which means $L_0(1) = 0$, a contradiction.

Acknowledgement

The work of the first author was supported by the China Scholarship Council. The funding corresponds to the scholarship for the PhD thesis of the first author in Paris, France. The French Agence Nationale de la Recherche partially supported the second author's work through ANR BARRACUDA (ANR-21-CE39-0009).

References

- [1] K. Abdukhalikov. Symplectic spreads, planar functions and mutually unbiased bases, *J. Algebraic Comb.* (41): 1055-1077, 2015.
- [2] Y. Aubry and M. Perret. A Weil theorem for singular curves. *Contemporary Mathematics*, 1–8, 1996.

- [3] D. Bartoli and M. Bonini. Planar polynomials arising from linearized polynomials. *Journal of Algebra and Its Applications*, (21) 01, 2250002, 2022.
- [4] C. Beierle and P. Felke. Classification of planar monomials over finite fields of small order. Available in: arXiv:2202.11032.
- [5] C. Beierle and P. Felke. Exhaustive search for planar monomials. Software (Version v1). Available via <https://doi.org/10.5281/zenodo.6226425>, 2022.
- [6] E. Bergman, R. S. Coulter, and I. Villa. Classifying planar monomials over fields of order a prime cubed. *Finite Fields Their Appl.*, 78: 101959, 2022.
- [7] C. Carlet. Boolean Functions for Cryptography and Coding Theory. *Cambridge University Press*, 2021.
- [8] C. Carlet, C. Ding, and J. Yuan. Linear codes from perfect nonlinear mappings and their secret sharing schemes. *IEEE Trans. Inf. Theory* 51: 2089–2102, 2005.
- [9] R. Chen and S. Mesnager. Evaluation of weil sums for some polynomials and associated quadratic forms. *Cryptography and Communications*, 15(3):661–673, 2023.
- [10] R. S. Coulter. The classification of planar monomials over fields of prime square order. *Proc. Am. Math. Soc.*, 134 (11): 3373–3378, 2006.
- [11] R. S. Coulter and M. Henderson. Commutative presemifields and semifields. *Advances in Mathematics*, 217(1): 282–304, 2008.
- [12] R. S. Coulter and M. Henderson. On a conjecture on planar polynomials of the form $x(\text{Tr}_n(x) - ux)$. *Finite Fields and Their Applications*, 21:30–34, 2013.
- [13] R. S. Coulter and R. W. Matthews. Planar functions and planes of Lenz-Barlotti class II. *Des. Codes Cryptogr.* 10(2): 167-184, 1997.
- [14] P. Dembowski and T.G. Ostrom. Planes of order n with collineation groups of order n^2 . *Math. Z.* (103): 239-258, 1968.
- [15] C. Ding and H. Niederreiter, Systematic authentication codes from highly nonlinear functions. *IEEE Trans. Inf. Theory* 50 (10): 2421–2428, 2004.
- [16] C. Ding, Q. Xiang, J. Yuan, and P. Yuan, Explicit classes of permutation polynomials of \mathbb{F}_3^{3m} , *Sci. China Ser. A, Math.* 53 (4): 639-647, 2009.
- [17] C. Ding and J. Yin. Signal sets from functions with optimum nonlinearity. *IEEE Trans. Commun.* 55 (5): 936–940, 2007.
- [18] C. Ding and J. Yuan, A family of optimal constant-composition codes, *IEEE Trans. Inf. Theory* 51 (10): 3668–3671, 2005.

- [19] C. Ding and J. Yuan. A family of skew Hadamard difference sets. *Journal of Combinatorial Theory, Series A*, 113(7): 1526–1535, 2006.
- [20] M.J. Ganley, E. Spence, Relative difference sets and quasiregular collineation groups. *J. Comb. Theory, Ser. A* 19, 134–153, 1975.
- [21] T. Helleseht and A. Kholosha. Monomial and quadratic bent functions over the finite fields of odd characteristic. *IEEE Transactions on Information Theory*, 52(5): 2018–2032, 2006.
- [22] G. Kyureghyan and F. Özbudak. Planarity of products of two linearized polynomials. *Finite Fields and Their Applications*, 18(6):1076–1088, 2012.
- [23] M. Kyuregyan, F. Özbudak, and A. Pott. Some planar maps and related function fields. *Arithmetic, Geometry, Cryptography and Coding Theory*, 574:87–114, 2012.
- [24] M. Moisiu. Kloosterman sums, elliptic curves, and irreducible polynomials with prescribed trace and norm. *Acta Arithmetica*, 132: 329-350, 2008.
- [25] K. Nyberg and L.R. Knudsen. Provable security against differential cryptanalysis, in: *Advances in Cryptology-CRYPTO’ 92*, Santa Barbara, CA, 1992, in: *Lecture Notes in Computer Science*, vol. 740, Springer, Berlin: 566-574, 1993.
- [26] A. Pott. Almost perfect and planar functions. *Des. Codes Cryptogr.* 78 (1): 141–195, 2016.
- [27] K.-U. Schmidt and Y. Zhou. Planar functions over fields of characteristic two. *J. Algebraic Combin.* 40, 2: 503–526, 2014.
- [28] H. Stichtenoth. *Algebraic function fields and codes*. Springer Science & Business Media, 2009.
- [29] G. Weng, W. Qiu, Z. Wang and Q. Xiang. Pseudo-Paley graphs and skew Hadamard difference sets from presemifields. *Des. Codes Cryptogr.* 44 (1–3): 49–62, 2007.
- [30] J. Yuan, C. Carlet, and C. Ding. The weight distribution of a class of linear codes from perfect nonlinear functions. *IEEE Trans. Inf. Theory* 52 (2): 712–717, 2006.
- [31] M. E. Zieve. Planar functions and perfect nonlinear monomials over finite fields. *Des. Codes Cryptogr.*,75(1): 71–80, 2015.
- [32] Z. Zha, G. M. Kyureghyan, and X. Wang. Perfect nonlinear binomials and their semifields. *Finite Fields and Their Applications*, 15(2): 125–133, 2009.
- [33] Y. Zhou. $(2^n, 2^n, 2^n, 1)$ -relative difference sets and their representations. *J. Combin. Des.* 21: 563–584, 2013.