

ON THE DIOPHANTINE SYSTEM INVOLVING PAIRS OF TRIANGLES WITH THE SAME AREA AND THE SAME PERIMETER

YANGCHENG LI AND YONG ZHANG

ABSTRACT. Many authors studied the problem that rational triangle pairs (triangle-parallelogram pairs) with the same area and the same perimeter. They investigated this problem by solving the rational solutions of the corresponding Diophantine equations. In this paper, we give a unified description of this problem by using the affine transformation in a rectangular coordinate system. According to the fact that two triangles with the same area and the same perimeter determine an affine transformation, this problem can be reduced to solving a specific Diophantine system. Moreover, we will give some rational solutions to this Diophantine system.

1. INTRODUCTION

A rational (Heron) triangle is a triangle with rational (integral) sides and rational (integral) area. Many authors studied two or more triangles with the same area and the same perimeter. Prielipp [20] proved that there are no two such distinct right triangles, Bradley [2] (also see [23]) showed that there are no three such isosceles triangles, Hirakawa and Matsumura [11] obtained a unique pair of right triangle-isosceles triangle, Choudhry [6] (also see [1, 3, 13, 19]) investigated two or more distinct Heron (rational) triangles, Juyal and Moody [12] studied Heron triangle and right triangle pairs. More results can be found in [18, 22, 7].

For triangle and quadrilateral pairs, Guy [10] proved that there is no non-degenerate such integral right triangle-rectangle pair. Several authors studied other cases, such as Zhang [24] (integral right triangle-parallelogram and Heron triangle-parallelogram pairs), Lalsín and Ma [14] (θ -triangle- ω -parallelogram pairs). For more results, we can refer to [4, 5, 8, 25, 26, 27, 28].

Suppose the areas of two polygons are A_1, A_2 , and the perimeters are P_1, P_2 , respectively. We say that the areas and perimeters of these two polygons are in fixed proportions (α, β) respectively, if they satisfy the following relationship

$$A_2 = \alpha A_1, \quad P_2 = \beta P_1,$$

where α and β are positive rational numbers. When $\alpha = \beta = 1$, it corresponds to two polygons with the same area and the same perimeter.

2010 *Mathematics Subject Classification.* Primary 11D72, 51M25; Secondary 51N10, 11G05.

Key words and phrases. rational triangle, parallelogram, affine transformation, elliptic curve.

This research was supported by the National Natural Science Foundation of China (Grant No. 11501052), the Natural Science Foundation of Education Department of Hunan Province (Project No. 21C0211), the Natural Science Foundation of Hunan Province (Project No. 2021JJ30699 and No. 2022JJ40464), and Hunan Provincial Key Laboratory of Mathematical Modeling and Analysis in Engineering (Changsha University of Science and Technology).

Li and Zhang [16, 15, 17] obtained infinitely many pairs of rational triangles (pairs of cyclic quadrilaterals, pairs of triangles and some special quadrilaterals) with areas and perimeters in fixed proportions (α, β) respectively.

On an affine plane, the affine transformation is a one-to-one, onto function which preserve parallelism. In this paper, the affine transformation is confined to a rectangular coordinate system. Let f be an affine transformation and P be a point with coordinates (x, y) . Suppose that the point P becomes $f(P)$ with coordinates (x', y') . We have the following transformation formula:

$$(1.1) \quad \begin{cases} x' = a_{11}x + a_{12}y + b_1, \\ y' = a_{21}x + a_{22}y + b_2. \end{cases}$$

Let T be a triangle and $f(T)$ be the image of T . Let A and A' be the areas of T and $f(T)$ respectively, then we have $A' = |\det(M_f)|A$, where M_f is the transformation matrix of f , i.e.,

$$M_f = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}.$$

The structure of this paper is as follows. In Section 2, we give the definition of the weakly metric equivalence of two rational triangles and the set $VL_2(\mathbb{Q})$ to describe the affine transformation between two weakly metric equivalent rational triangles. In Section 3, the set $VL_2(\mathbb{Q})$ is used to uniformly describe the existed conclusions that two rational triangles are weakly metric equivalent, and to get more such pairs. In Section 4, the affine transformation is also used for describing the problem that triangle and parallelogram pairs having the same area and the same perimeter. In Section 5, we consider the rational triangle pairs or rational triangle-parallelogram pairs with areas and perimeters in fixed proportions (α, β) , respectively, where α and β are positive rational numbers.


2. EQUIVALENCE RELATION AND THE SET $VL_2(\mathbb{Q})$

The purpose of this paper is to study the problem of two rational triangles with the same area and the same perimeter. It is useful to define the following equivalence relation.

Definition 2.1. Two rational triangles are weakly metric equivalent if they have the same area and the same perimeter. In particular, two rational triangles are metric equivalent if they are congruent.

Suppose that the rational triangles T_1 and T_2 are weakly metric equivalent, we write it as $T_1 \sim T_2$. If the rational triangles T_1 and T_2 are metric equivalent, we write it as $T_1 = T_2$.

For a rational triangle T , we wish to find an affine transformation f such that $f(T)$ is weakly equivalent to T , i.e., $f(T) \sim T$. Let V_1, V_2 and V_3 be the three vertices of the rational triangle T . In a rectangular coordinate system, we put the rational triangle T in the first quadrant. The vertex V_1 is at the origin of the coordinate system, the vertex V_2 is on the positive x axis and the vertex V_3 is above the x axis (see Figure 1). Let $f(V_1), f(V_2)$ and $f(V_3)$ be the three vertices of the rational triangle $f(T)$. We can make $f(V_1)$ coincide with V_1 , that is still at the origin. $f(V_2)$ is on the positive x axis and $f(V_3)$ is above the x axis.

FIGURE 1. Rational triangles T and $f(T)$.

In other words, the rational triangles to be discussed are confined to the following set

$$X(T) = \{T \mid V_1 = (0, 0), V_2 = (r, 0), V_3 = (s, t), r \in \mathbb{Q}^*, s \in \mathbb{Q}_{\geq 0}, t \in \mathbb{Q}^*\},$$

where $\mathbb{Q}^* = \{x \in \mathbb{Q} \mid x > 0\}$ and $\mathbb{Q}_{\geq 0} = \{x \in \mathbb{Q} \mid x \geq 0\}$. It is not difficult to find that to make $f(T) \in X(T)$, we only need $b_1 = b_2 = a_{21} = 0$ in the transformation formula (1.1). In addition, to make the areas of T and $f(T)$ be the same, as long as the determinant $\det(M_f)$ of M_f is equal to ± 1 . We just take $\det(M_f) = 1$. Therefore, M_f can be assumed to be in the following form

$$M_f = \begin{bmatrix} \frac{1}{a} & b \\ 0 & a \end{bmatrix},$$

where $a \in \mathbb{Q}^*, b \in \mathbb{Q}$. Thus, the transformation formula (1.1) becomes

$$\begin{cases} x' = \frac{1}{a}x + by, \\ y' = ay. \end{cases}$$

Let

$$UL_2(\mathbb{Q}) = \left\{ \begin{bmatrix} \frac{1}{a} & b \\ 0 & a \end{bmatrix} \mid a \in \mathbb{Q}^*, b \in \mathbb{Q} \right\}.$$

It is easy to check that the set $UL_2(\mathbb{Q})$ forms a group under the multiplication of matrices. Let

$$U = \{f \mid M_f \in UL_2(\mathbb{Q})\}.$$

Obviously, an operation of U on $X(T)$ is given by the map

$$U \times X(T) \rightarrow X(T), \quad (f, T) \mapsto f(T).$$

Suppose the area of $T \in X(T)$ is A_T , then the orbit of T in $X(T)$ is the set

$$O_T = \{T' \in X(T) \mid \text{the area of } T' \text{ is } A_T\}.$$

But the group $UL_2(\mathbb{Q})$ is not the set that we need, because we cannot guarantee that T and $f(T)$ have the same perimeter.

The following definition is derived from the weakly metric equivalence of two rational triangles.

Definition 2.2. An affine transformation f is called weak metric affine transformation about rational triangle T , if there is a rational triangle T such that $f(T)$ is weakly metric equivalent to T . The corresponding transformation matrix M_f is called weak metric transformation matrix.

Let D be the set of all weak metric affine transformations. Correspondingly, let $DL_2(\mathbb{Q})$ be the set of all weak metric transformation matrices. We take the intersection of the sets $UL_2(\mathbb{Q})$ and $DL_2(\mathbb{Q})$, i.e.,

$$VL_2(\mathbb{Q}) = \{M_f \mid M_f \in UL_2(\mathbb{Q}) \cap DL_2(\mathbb{Q})\},$$

which is the set we will discuss next. Correspondingly, let

$$V = \{f \mid M_f \in VL_2(\mathbb{Q})\}.$$

By Definition 2.1 and the set V , we have

Proposition 2.3. *If $T_1 \sim T_2$, then there is a $f \in V$ such that $f(T_1) = T_2$.*

The set $VL_2(\mathbb{Q})$ is not empty, because $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \in VL_2(\mathbb{Q})$. Obviously, for any rational triangle T , $f_I(T) = T$. Hence, the set $VL_2(\mathbb{Q})$ is a monoid.

For any $f \in V$, there is a rational triangle T such that $f(T) \sim T$. Obviously, $f^{-1}(f(T)) = T$ is weakly metric equivalent to $f(T)$. Therefore, for any $M_f \in VL_2(\mathbb{Q})$, $M_f^{-1} \in VL_2(\mathbb{Q})$.

However, the set $VL_2(\mathbb{Q})$ may not form a group, because it is not clear whether the composition of weak metric affine transformation is closed. Taking any $f, g \in V$ such that $f(T_1) \sim T_1$ and $g(T_2) \sim T_2$. The question is whether we can find a rational triangle T such that $g(f(T)) \sim T$ or $f(g(T)) \sim T$. If there is a positive answer to this question, the set $VL_2(\mathbb{Q})$ can form a group.

Taking any $f \in V$, there is a rational triangle T such that $f(T) \sim T$. Assuming that rational triangle T' is similar to T , then $f(T') \sim T'$.

For the set $VL_2(\mathbb{Q})$ or V , we have the following two basic questions:

(1) Given a rational triangle T , is there a $f \in V$ such that $f(T) \sim T$?

(2) Given any $f \in V$ such that $f(T) \sim T$, is there a rational triangle $T' \in X(T)$ that is not similar to T such that $f(T') \sim T'$?

To solve question (1), we give the following two lemmas:

Lemma 2.4. [6] *Given any scalene rational triangle T_1 , whose sides do not, in any order, satisfy the Diophantine equation*

$$\begin{aligned} x^4 - x^3y - 2x^3z + 3x^2yz - xy^3 + 3xy^2z \\ - 6xyz^2 + 3xz^3 + y^4 - 2y^3z + 3yz^3 - 2z^4 = 0, \end{aligned}$$

then there exists a second rational triangle T_2 such that $T_1 \sim T_2$.

Lemma 2.5. [6] *Given an arbitrary positive integer $k \geq 2$, there exist k scalene rational triangles are weakly metric equivalent.*

Lemma 2.4 gives a positive answer to question (1), and Lemma 2.5 shows that there can be an infinite number of such $f \in V$. By Lemma 2.5, we give the concept of weak similarity of weakly metric affine transformation.

Definition 2.6. Let $f, g \in V$, if there is a rational triangle T and $h \in V$ such that $f(T) = h(g(T))$, then f and g are weakly similar, we write it as $f \sim g$.

Obviously, weak similarity is an equivalence relation. All weakly metric affine transformations f about the same rational triangle T are weakly similar and vice versa.

For question (2), it is easy to see that $f(T) \sim T$ for any rational triangle T when $f = f_I$. For a general $f \in V$, there should be $f(T) \sim T$ only for some specific rational triangles T . Therefore, we have the following conjecture:

Conjecture 2.1. *If $f \in V, f \neq f_I$, there are only a finite number of dissimilar rational triangles $T \in X(T)$ such that $f(T) \sim T$.*

It is not clear whether the set V is closed under the composition of weakly metric affine transformation. However, in some special cases, this is possible. By lemma 2.5, we can get a sequence of rational triangles $\{T_n\}, n \geq 1$, in which any two rational triangles are weakly metric equivalent. By Proposition 2.3, we can use $\{T_n\}$ to construct a sequence $\{f_n\}$ such that $f_i(T_i) = T_{i+1}, i \geq 1$. We have the following proposition:

Proposition 2.7. *If $f_1, \dots, f_k \in \{f_n\}, k \geq 1$, then $f_k \cdots f_1 \in V$. Generally, $f_k \cdots f_i \in V, k \geq i \geq 1$.*

We have a certain grasp of the properties of the set $VL_2(\mathbb{Q})$, but the conditions satisfied by a and b are not specific. Next, we show that the values of a and b are determined by a Diophantine system.

For a rational triangle $T \in X(T)$, the coordinates of the three vertices of T are

$$V_1 = (0, 0), V_2 = (r, 0), V_3 = (s, t),$$

where $r \in \mathbb{Q}^*, s \in \mathbb{Q}_{\geq 0}, t \in \mathbb{Q}^*$. Thus, the sides of the rational triangle T are

$$|V_1V_2| = r, |V_1V_3| = \sqrt{s^2 + t^2}, |V_2V_3| = \sqrt{(s-r)^2 + t^2}.$$

Since the sides of rational triangle T are rationals, we thus have

$$s^2 + t^2 = \square, (s-r)^2 + t^2 = \square, r \in \mathbb{Q}^*, s \in \mathbb{Q}_{\geq 0}, t \in \mathbb{Q}^*.$$

By the transformation f , the coordinates of the three vertices of $f(T)$ are

$$f(V_1) = (0, 0), f(V_2) = \left(\frac{r}{a}, 0\right), f(V_3) = \left(\frac{s}{a} + bt, at\right).$$

Then, the sides of the rational triangle $f(T)$ are

$$\begin{aligned} |f(V_1)f(V_2)| &= \frac{r}{a}, |f(V_1)f(V_3)| = \sqrt{\left(\frac{s}{a} + bt\right)^2 + (at)^2}, \\ |f(V_2)f(V_3)| &= \sqrt{\left(\frac{s-r}{a} + bt\right)^2 + (at)^2}. \end{aligned}$$

To ensure that the sides of $f(T)$ are rational, we have

$$\left(\frac{s}{a} + bt\right)^2 + (at)^2 = \square, \left(\frac{s-r}{a} + bt\right)^2 + (at)^2 = \square.$$

Since T and $f(T)$ are weakly metric equivalent, they have the same perimeter, i.e.,

$$r + \sqrt{s^2 + t^2} + \sqrt{(s-r)^2 + t^2} = \frac{r}{a} + \sqrt{\left(\frac{s}{a} + bt\right)^2 + (at)^2} + \sqrt{\left(\frac{s-r}{a} + bt\right)^2 + (at)^2}.$$

Therefore, for any rational triangle $T \in X(T)$, its corresponding weakly metric affine transformation f is completely characterized by the following Diophantine system

$$(2.1) \quad \begin{cases} s^2 + t^2 = w_1^2, \\ (s-r)^2 + t^2 = w_2^2, \\ \left(\frac{s}{a} + bt\right)^2 + (at)^2 = w_3^2, \\ \left(\frac{s-r}{a} + bt\right)^2 + (at)^2 = w_4^2, \\ r + w_1 + w_2 = \frac{r}{a} + w_3 + w_4, \end{cases}$$

where $b \in \mathbb{Q}$, $s \in \mathbb{Q}_{\geq 0}$, $a, r, t, w_1, w_2, w_3, w_4 \in \mathbb{Q}^*$.

Assume that r, s and t satisfy the first two equations, then $(a, b) = (1, 0)$ is a solution of Eq. (2.1). The corresponding affine transformation is f_I . Moreover, $(a, b) = (1, \frac{r-2s}{t})$ is also a solution of Eq. (2.1).

Suppose (a, b, r, s, t) is any solution of Eq. (2.1) and k is any positive rational number, then (a, b, kr, ks, kt) is also a solution of Eq. (2.1), which means that Eq. (2.1) is homogeneous with respect to r, s and t .

Let S be the solution set of Eq. (2.1). In particular, let $S_{(r,s,t)}$ be the solution set with respect to (a, b) . Thus, the set $VL_2(\mathbb{Q})$ can be expressed as

$$VL_2(\mathbb{Q}) = \{M \mid M \in UL_2(\mathbb{Q}), (a, b) \in S_{(r,s,t)}\},$$

which gives an accurate description of the set $VL_2(\mathbb{Q})$.

Now we discuss whether the composition of weakly metric affine transformations in the set V is closed according to the solution set S . The question can be described as: Suppose $(a_1, b_1, r_1, s_1, t_1) \in S$ and $(a_2, b_2, r_2, s_2, t_2) \in S$, if $(a, b) = (a_1 a_2, \frac{b_2}{a_1} + a_2 b_1)$ or $(a, b) = (a_1 a_2, \frac{b_1}{a_2} + a_1 b_2)$, does Eq. (2.1) have a rational solution? If there is a positive answer to this question, the set $VL_2(\mathbb{Q})$ can obviously form a group. Thus, we have the following question.

Question 2.8. Does the set $VL_2(\mathbb{Q})$ form a group?

3. CONCRETE WEAKLY METRIC AFFINE TRANSFORMATION

In this section, we will discuss two problems. One is using the set $VL_2(\mathbb{Q})$ to uniformly describe the existed conclusions that two rational triangles with the same area and the same perimeter. The other is applying the set $VL_2(\mathbb{Q})$ to find more weakly metric equivalent rational triangles.

3.1. Existed results. We first give a unified description of the existed results, focusing on the following four cases.

Case 3.1.1. T and $f(T)$ are right triangles.

Without losing generality, we take $r = 2km, s = 0, t = k(m^2 - 1), k > 0, m > 1$. Then the coordinates of the three vertices of T are

$$V_1 = (0, 0), V_2 = (2km, 0), V_3 = (0, k(m^2 - 1)).$$

The coordinates of the three vertices of $f(T)$ are

$$f(V_1) = (0, 0), f(V_2) = \left(\frac{2km}{a}, 0\right), f(V_3) = (bk(m^2 - 1), ak(m^2 - 1)).$$

Since $f(T)$ is a right triangle, we can take $b = 0$, then $w_3 = ak(m^2 - 1)$. Eq. (2.1) now becomes

$$(3.1) \quad \begin{cases} \frac{4k^2m^2}{a^2} + a^2k^2(m^2 - 1)^2 = w_4^2, \\ 2km(m + 1) = \frac{2km}{a} + ak(m^2 - 1) + w_4. \end{cases}$$

Eliminating w_4 in Eq. (3.1), we have

$$(a - 1)((m^2 - 1)a - 2m) = 0.$$

Then $a = 1$ or $a = \frac{2m}{m^2 - 1}$. Obviously, when $a = 1$ or $a = \frac{2m}{m^2 - 1}$, $f(T)$ and T are congruent. Therefore, there are no two distinct right triangles are weakly metric equivalent, which is obtained by Prielipp [20].

Case 3.1.2. T and $f(T)$ are isosceles triangles.

Case 3.1.2.1. Take $r = 4km, s = 2km, t = k(m^2 - 1), k > 0, m > 1$, then the coordinates of the three vertices of T are

$$V_1 = (0, 0), V_2 = (4km, 0), V_3 = (2km, k(m^2 - 1)).$$

The coordinates of the three vertices of $f(T)$ are

$$f(V_1) = (0, 0), f(V_2) = \left(\frac{4km}{a}, 0\right), f(V_3) = \left(\frac{2km}{a} + bk(m^2 - 1), ak(m^2 - 1)\right).$$

Since $f(T)$ is an isosceles triangle, we can take $b = 0$, then $w_3 = w_4$. Eq. (2.1) now becomes

$$(3.2) \quad \begin{cases} \frac{4k^2m^2}{a^2} + a^2k^2(m^2 - 1)^2 = w_4^2, \\ 2k(m + 1)^2 = \frac{4km}{a} + 2w_4. \end{cases}$$

Eliminating w_4 in Eq. (3.2), we have

$$(a - 1)((m - 1)^2a^2 + (m - 1)^2a - 4m) = 0.$$

Then

$$a = 1 \text{ or } a = \frac{1 - m \pm \sqrt{m^2 + 14m + 1}}{2(m - 1)}.$$

When $a = 1$, $f = f_I$, this is a trivial case. Since $a \in \mathbb{Q}^*$, then a has at most one nontrivial positive rational solution, which leads to the fact that there are no three isosceles triangles are weakly metric equivalent at the same time.

Case 3.1.2.2. Take $r = 2k(m^2 - 1), s = k(m^2 - 1), t = 2km, k > 0, m > 1$, then the coordinates of the three vertices of T are

$$V_1 = (0, 0), V_2 = (2k(m^2 - 1), 0), V_3 = (k(m^2 - 1), 2km).$$

The coordinates of the three vertices of $f(T)$ are

$$f(V_1) = (0, 0), f(V_2) = \left(\frac{2k(m^2 - 1)}{a}, 0\right), f(V_3) = \left(\frac{k(m^2 - 1)}{a} + 2bkm, 2akm\right).$$

Since $f(T)$ is an isosceles triangle, we can take $b = 0$, then $w_3 = w_4$. Eq. (2.1) now becomes

$$(3.3) \quad \begin{cases} \frac{k^2(m^2 - 1)^2}{a^2} + 4a^2k^2m^2 = w_4^2, \\ 4km^2 = \frac{2k(m^2 - 1)}{a} + 2w_4. \end{cases}$$

Eliminating w_4 in Eq. (3.3), we have

$$(a-1)(a^2+a-(m^2-1))=0.$$

Then

$$a=1 \text{ or } a=\frac{-1\pm\sqrt{4m^2-3}}{2}.$$

Hence, a has at most one nontrivial positive rational solution, then there are no three isosceles triangles are weakly metric equivalent at the same time, this case was studied by Bradley [2] (also see [23]).

Case 3.1.3. T is a right triangle and $f(T)$ is an isosceles triangle.

Take $r=2km, s=0, t=k(m^2-1), k>0, m>1$. Then the coordinates of the three vertices of T are

$$V_1=(0,0), V_2=(2km,0), V_3=(0,k(m^2-1)).$$

The coordinates of the three vertices of $f(T)$ are

$$f(V_1)=(0,0), f(V_2)=\left(\frac{2km}{a},0\right), f(V_3)=(bk(m^2-1),ak(m^2-1)).$$

Since $f(T)$ is an isosceles triangle, we can take $bk(m^2-1)=\frac{km}{a}$, then $w_3=w_4$. Eq. (2.1) now becomes

$$(3.4) \quad \begin{cases} \frac{k^2m^2}{a^2}+a^2k^2(m^2-1)^2=w_4^2, \\ 2km(m+1)=\frac{2km}{a}+2w_4. \end{cases}$$

Eliminating w_4 in Eq. (3.4), we have

$$(3.5) \quad (m+1)(m-1)^2a^3-m^2(m+1)a+2m^2=0.$$

The discriminant of Eq. (3.5) with respect to a is

$$4(m^2-1)^2m^4(m^4+2m^3-26m^2+54m-27)>0.$$

Eq. (3.5) is equivalent to

$$(m+1)a((m-1)^2a^2-m^2)+2m^2=0,$$

then $a<\frac{m}{m-1}$.

It is easy to verify that $(a,m)=\left(\frac{8}{3},\frac{16}{11}\right)$ ($k=121$) is the solution of Eq. (3.5), which corresponds to the unique pair of right triangle and isosceles triangle obtained by Hirakawa and Matsumura [11], i.e., $(135,352,377), (132,366,366)$. Unfortunately, we cannot prove that this solution is the only nontrivial positive rational solution of Eq. (3.5).

Case 3.1.4. T and $f(T)$ are rational triangles.

In this case, we only give the affine transformation corresponding to a parameter solution obtained in [6]. The sides of T and $f(T)$ are given respectively by

$$\begin{aligned} a_1 &= (R^2+1)(S^2+S+1), \\ b_1 &= S(S+1)(R^2+S^2+S+1), \\ c_1 &= R^2+(S^2+S+1)^2, \end{aligned}$$

and

$$\begin{aligned} a_2 &= (R^2 + S^2)(S^2 + S + 1), \\ b_2 &= (S + 1)(R^2 + S^2 + S + 1), \\ c_2 &= R^2 S^2 + (S^2 + S + 1)^2, \end{aligned}$$

where R, S are arbitrary positive rational numbers. Take

$$\begin{aligned} r &= R^2 + (S^2 + S + 1)^2, \\ s &= \frac{S(S + 1)(R^2 + S^2 + S + 1)(S^2 + R + S + 1)(S^2 - R + S + 1)}{S^4 + 2S^3 + R^2 + 3S^2 + 2S + 1}, \\ t &= \frac{(2S^2 + 2S + 2)(R^2 + S^2 + S + 1)RS(S + 1)}{S^4 + 2S^3 + R^2 + 3S^2 + 2S + 1}. \end{aligned}$$

Then

$$\begin{aligned} a &= \frac{S^4 + 2S^3 + R^2 + 3S^2 + 2S + 1}{R^2 S^2 + S^4 + 2S^3 + 3S^2 + 2S + 1}, \\ b &= \frac{(S - 1)b(R, S)}{2SR(S^4 + 2S^3 + R^2 + 3S^2 + 2S + 1)(R^2 S^2 + S^4 + 2S^3 + 3S^2 + 2S + 1)}, \end{aligned}$$

where

$$\begin{aligned} b(R, S) &= R^6 S^2 - (S^2 + S + 1)(S^4 + S^3 - S^2 + S + 1)R^4 \\ &\quad - (2S^2 + 3S + 2)(S^2 + S + 1)^3 R^2 - (S^2 + S + 1)^5. \end{aligned}$$

3.2. Other solutions of Eq. (2.1). We can get more solutions by directly solving Eq. (2.1). We are concerned with the following two cases.

Case 3.2.1. $s = 0$, T is a right triangle.

Take $r = 2km$, $s = 0$, $t = k(m^2 - 1)$, $k > 0$, $m > 1$, then

$$w_1 = k(m^2 - 1), \quad w_2 = k(m^2 + 1).$$

From the third equation of Eq. (2.1), we have

$$k^2(m^2 - 1)^2(a^2 + b^2) = w_3^2,$$

then

$$a = 2cp, \quad b = c(p^2 - 1), \quad w_3 = ck(m^2 - 1)(p^2 + 1), \quad c > 0, \quad p > 1.$$

Eq. (2.1) now becomes

$$(3.6) \quad \begin{cases} \left(ck(p^2 - 1)(m^2 - 1) - \frac{km}{cp} \right)^2 + 4c^2 p^2 k^2 (m^2 - 1)^2 = w_4^2, \\ 2km(m + 1) = \frac{km}{cp} + ck(m^2 - 1)(p^2 + 1) + w_4. \end{cases}$$

Eliminating w_4 in Eq. (3.6), we have

$$(3.7) \quad p(p^2 + 1)(m^2 - 1)c^2 - p(m^2 + (p + 1)m - p)c + m = 0,$$

then

$$c = \frac{p((m - 1)p + m(m + 1)) \pm \sqrt{\Delta(c)}}{2p(p^2 + 1)(m^2 - 1)},$$

where

$$\Delta(c) = p^2 m^4 - 2p(p^2 - p + 2)m^3 + p^2(p^2 + 1)m^2 - 2p(p^3 - p^2 - 2)m + p^4.$$

To get the rational solutions of Eq. (3.7) with respect to c , it needs $\Delta(c)$ be a perfect square, say w^2 . Using a method described by Fermat (see [9, p. 639]), it's easy to get the following solutions

$$m = \frac{p^2 + p + 1}{p^2}, \quad m = \frac{p^2 - p + 1}{p(p-2)}, \quad m = \frac{p^3 - p^2 - 1}{p^2(p+1)}$$

such that $\Delta(c)$ is a perfect square. Take $m = \frac{p^2+p+1}{p^2}$, by Eq. (3.7), we get

$$c = \frac{p}{p^2 + 1}, \quad c = \frac{p^2 + p + 1}{2p^3 + 3p^2 + 2p + 1}.$$

Take $c = \frac{p}{p^2+1}$, then

$$\begin{aligned} s &= 0, \quad r = \frac{2k(p^2 + p + 1)}{p^2}, \quad t = \frac{k(p+1)(2p^2 + p + 1)}{p^4}, \\ a &= \frac{2p^2}{p^2 + 1}, \quad b = \frac{p(p^2 - 1)}{p^2 + 1}, \\ w_1 &= \frac{k(p+1)(2p^2 + p + 1)}{p^4}, \quad w_2 = \frac{k(p^2 + 1)(2p^2 + 2p + 1)}{p^4}, \\ w_3 &= \frac{k(p+1)(2p^2 + p + 1)}{p^3}, \quad w_4 = \frac{(p^4 + 2p^3 + 4p^2 + 2p + 1)k}{p^4}, \end{aligned}$$

where $k > 0, p > 1$, this is a parametric solution of Eq. (2.1).

Let us consider the rational points on the quartic curve

$$\mathcal{C} : w^2 = \Delta(c).$$

By the map φ :

$$\begin{aligned} X &= \frac{3p(p(p^2 + 1)m^2 - 6(p^3 - p^2 - 2)m + 6p^3 + 6pw)}{m^2}, \\ Y &= -\frac{Y(m, w)}{m^3}, \end{aligned}$$

where

$$\begin{aligned} Y(m, w) &= 54p(p^2((p^2 - p + 2)m^3 - p(p^2 + 1)m^2 + 3(p^3 - p^2 - 2)m - 2p^3) \\ &\quad + ((p^3 - p^2 - 2)m - 2p^3)w). \end{aligned}$$

We can transform \mathcal{C} into the elliptic curve

$$\begin{aligned} \mathcal{E}_p : Y^2 &= X^3 - 27p^2(p^6 - 12p^5 + 38p^4 - 36p^3 + 49p^2 - 24p + 48)X \\ &\quad + 54p^4(p^3 - 6p^2 + p - 12)(p^5 - 12p^4 + 38p^3 - 36p^2 + p - 24). \end{aligned}$$

The discriminant of \mathcal{E}_p is

$$\Delta(p) = 544195584p^6(p^4 - 10p^3 + 17p^2 + 8p + 16)(p-1)^2(p^2 + 1)^2.$$

Hence, $\Delta(p) \neq 0$, so \mathcal{E}_p is non-singular.

It is easy to check that the elliptic curve \mathcal{E}_p contains two rational points

$$Q = (3p(p^3 - 6p^2 + p - 12), 0), \quad P = (3p^2(p^2 - 6p + 1), 108p^2(p + 1)).$$

By the inverse map φ^{-1} , we have

$$m = \frac{6p(Yp - 3(p^3 - p^2 - 2)X + 9p^2(p^5 - 7p^4 + 7p^3 - 15p^2 - 2))}{(X - 3p^2(p^2 - 6p + 1))(X - 3p^2(p^2 + 6p + 1))}.$$

When $p = 2$, we get

$$\mathcal{E}_2 : Y^2 = X^3 - 21168X + 494208.$$

By the package of Magma, the rank of \mathcal{E}_2 is 1. From the theorem of Silverman (see p. 457 of [21]) about the rank of elliptic curve: If an elliptic curve E defined over $\mathbb{Q}(t)$ for some values of t has positive rank, for all but finitely many $t \in \mathbb{Q}$ it has positive rank. Hence, the elliptic curve \mathcal{E}_p , for all but finitely many $p > 1$, has positive rank. Therefore, we can get more parametric solutions by the group law, such as

$$[2]P = \left(\frac{3(p^6 - 4p^5 + 38p^4 - 16p^3 + 25p^2 - 12p + 12)}{(p+1)^2}, \frac{108(2p^2+1)(2p^5-4p^4+5p^3-2p^2+3p-2)}{(p+1)^3} \right).$$

From the rational point $[2]P$, we obtain

$$m = \frac{p(p+1)}{p^2-p+1},$$

then

$$c = \frac{p+1}{2p^2+1}, \quad c = \frac{p^2-p+1}{2p^3-p^2+2p-1}.$$

Take $c = \frac{p+1}{2p^2+1}$, then

$$\begin{aligned} s &= 0, \quad r = \frac{2kp(p+1)}{p^2-p+1}, \quad t = \frac{k(2p-1)(2p^2+1)}{(p^2-p+1)^2}, \\ a &= \frac{2(p+1)p}{2p^2+1}, \quad b = \frac{(p+1)^2(p-1)}{2p^2+1}, \\ w_1 &= \frac{k(2p-1)(2p^2+1)}{(p^2-p+1)^2}, \quad w_2 = \frac{k(2p^4+4p^2-2p+1)}{(p^2-p+1)^2}, \\ w_3 &= \frac{(p^2+1)(2p-1)k(p+1)}{(p^2-p+1)^2}, \quad w_4 = \frac{kp(5p^2-2p+2)}{(p^2-p+1)^2}, \end{aligned}$$

where $k > 0, p > 1$.

Case 3.2.2. $s \neq 0$, T is a rational triangle.

Take $s = 2km, t = k(m^2 - 1), k > 0, m > 1$, then

$$w_1 = k(m^2 + 1).$$

From the second equation of Eq. (2.1), we have

$$r = \frac{2k((m^2+1)c-2m)}{c^2-1}, \quad w_2 = \frac{k((m^2+1)c^2-4mc+m^2+1)}{c^2-1},$$

where c is a rational number.

From the third equation of Eq. (2.1), we have

$$\left(\frac{2km}{a} + bk(m^2 - 1) \right)^2 + a^2k^2(m^2 - 1)^2 = w_3^2.$$

Let

$$\frac{2km}{a} + bk(m^2 - 1) = d(p^2 - 1), \quad ak(m^2 - 1) = 2dp,$$

then

$$a = \frac{2dp}{k(m^2 - 1)}, \quad b = \frac{p(p^2 - 1)d^2 - k^2m(m^2 - 1)}{dkp(m^2 - 1)}, \quad w_3 = d(p^2 + 1).$$

Eq. (2.1) now becomes

$$(3.8) \quad \begin{cases} \left(\frac{d^2p(p^2 - 1)(c^2 - 1) - (m^2 - 1)((m^2 + 1)c - 2m)k^2}{(c^2 - 1)dp} \right)^2 + 4d^2p^2 = w_4^2, \\ \frac{2k((m^2 + 1)c - 2m)}{c - 1} = \frac{(m^2 - 1)((m^2 + 1)c - 2m)k^2}{(c^2 - 1)dp} + d(p^2 + 1) + w_4 \end{cases}$$

Eliminating w_4 in Eq. (3.8), we have

$$(3.9) \quad \begin{aligned} & dp((p^2 + 1)d - k(m^2 + 1))c^2 \\ & - k(m - 1)(p((m + 1)p + m - 1)d - k(m + 1)(m^2 + 1))c \\ & - p(p^2 + 1)d^2 + kp((m^2 - 1)p + 2m)d - 2k^2m(m^2 - 1) = 0, \end{aligned}$$

then

$$c = \frac{(m^4 - 1)k^2 - p((m^2 - 1)p + (m - 1)^2)k \pm \sqrt{\Delta(c)}}{2p(k(m^2 + 1) - (p^2 + 1))},$$

where

$$\begin{aligned} \Delta(c) &= (m^4 - 1)^2k^4 - 2dp(m^4 - 1)((m^2 - 1)p + (m + 1)^2)k^3 \\ &+ d^2p((m^2 - 1)^2p^3 + 2(m^2 - 1)(3m^2 + 2m + 3)p^2 \\ &+ (m + 1)^4p + 8m(m^2 - 1))k^2 \\ &- 4d^3p^2(p^2 + 1)((m^2 - 1)p + (m + 1)^2)k + 4d^4p^2(p^2 + 1)^2. \end{aligned}$$

To get the rational solutions of Eq. (3.9) with respect to c , it needs $\Delta(c)$ be a perfect square, say w^2 . Since the discriminant $\Delta(c)$ is homogeneous with respect to k and d , we can let $d = 1$. Using a method described by Fermat (see [9, p. 639]), it's easy to get the following solution

$$k = \frac{2(m^2 + 1)p^2 + (m + 1)^2}{(m^2 + 1)((m^2 - 1)p + (m + 1)^2)}$$

such that $\Delta(c)$ is a perfect square. By Eq. (3.9), we get

$$c = \frac{(m + 1)(m^2 + 1)p^2 + (m - 1)(m^2 + 1)p + 2m(m + 1)}{(m^2 + 1)((m + 1)p^2 - (m - 1)p + m + 1)}.$$

By some calculations, we obtain

$$\begin{aligned} r &= \frac{2((m+1)p^2 - (m-1)p + m+1)(m^2+1)p}{(m+1)(2(m^2+1)p - m^2+1)}, \\ s &= \frac{2(2(m^2+1)p^2 + (m+1)^2)m}{(m^2+1)((m^2-1)p + (m+1)^2)}, \\ t &= \frac{(m-1)(2(m^2+1)p^2 + (m+1)^2)}{((m-1)p + m+1)(m^2+1)}, \\ a &= \frac{2((m-1)p + m+1)(m^2+1)p}{(m-1)(2(m^2+1)p^2 + (m+1)^2)}, \\ b &= \frac{b(m,p)}{(m^4-1)(2(m^2+1)p^2 + (m+1)^2)((m^2-1)p + (m+1)^2)p}, \end{aligned}$$

where

$$\begin{aligned} b(m,p) &= (m^4-1)^2p^5 + 2(m^2-1)(m^2+1)^3p^4 + 4(m+1)^2m(m^2+1)^2p^3 \\ &\quad - 2(m+1)^4(m^4-1)p^2 - (m+1)^4(m^2+1)^2p - (m+1)^5m(m-1), \end{aligned}$$

and $m > 1, p > 1$, this is a parametric solution of Eq. (2.1).

Let us consider the rational points on the quartic curve

$$\mathcal{C} : w^2 = \Delta(c).$$

By the map φ :

$$\begin{aligned} X &= \frac{3pX(k,w)}{k^2}, \\ Y &= -\frac{108p^2(p^2+1)Y(k,w)}{k^3}, \end{aligned}$$

where

$$\begin{aligned} X(k,w) &= ((m^2-1)^2p^3 + 2(m^2-1)(3m^2+2m+3)p^2 + (m+1)^4p \\ &\quad + 8m(m^2-1))k^2 - 12p(p^2+1)((m^2-1)p + (m+1)^2)k \\ &\quad + 24p(p^2+1)^2 - 12(p^2+1)w, \\ Y(k,w) &= (m^4-1)((m^2-1)p + (m+1)^2)k^3 - ((m^2-1)^2p^3 \\ &\quad + 2(m^2-1)(3m^2+2m+3)p^2 + (m+1)^4p + 8m(m^2-1))k^2 \\ &\quad + 6p(p^2+1)((m^2-1)p + (m+1)^2)k - 8p(p^2+1)^2 \\ &\quad - (((m^2-1)p + (m+1)^2)k - 4(p^2+1))w. \end{aligned}$$

We can transform \mathcal{C} into the elliptic curve

$$\mathcal{E}_{(m,p)} : Y^2 = X^3 - 27p^2A_4(m,p)X + 54p^3A_6(m,p),$$

where

$$\begin{aligned}
A_4(m, p) &= (m^2 - 1)^4 p^6 - 4(3m^2 - 2m + 3)(m^2 - 1)^3 p^5 \\
&\quad + 2(19m^4 - 20m^3 + 50m^2 - 20m + 19)(m^2 - 1)^2 p^4 \\
&\quad - 4(m + 1)^3 (m - 1)(3m^2 - 2m + 3)^2 p^3 \\
&\quad + (m + 1)^2 (49m^6 - 90m^5 + 223m^4 - 300m^3 + 223m^2 - 90m + 49)p^2 \\
&\quad - 8(m + 1)^5 (m - 1)(3m^2 - 2m + 3)p + 16(m^2 + 3)(3m^2 + 1)(m^2 - 1)^2, \\
A_6(m, p) &= ((m^2 - 1)^2 p^3 - 2(m^2 - 1)(3m^2 - 2m + 3)p^2 + (m + 1)^4 p \\
&\quad - 4(m^2 - 1)(3m^2 - 2m + 3)((m^2 - 1)^4 p^6 \\
&\quad - 4(3m^2 - 2m + 3)(m^2 - 1)^3 p^5 \\
&\quad + 2(19m^4 - 20m^3 + 50m^2 - 20m + 19)(m^2 - 1)^2 p^4 \\
&\quad - 4(m + 1)^3 (m - 1)(3m^2 - 2m + 3)^2 p^3 \\
&\quad + (m + 1)^2 (m^6 + 102m^5 - 113m^4 + 84m^3 - 113m^2 + 102m + 1)p^2 \\
&\quad - 8(m + 1)^5 (m - 1)(3m^2 - 2m + 3)p + 32m(3m^2 + 2m + 3)(m^2 - 1)^2).
\end{aligned}$$

By some calculations, we can check that when $m \neq \frac{p+1}{p-1}$, the discriminant $\Delta(m, p)$ of $\mathcal{E}_{(m,p)}$ is not 0, then $\mathcal{E}_{(m,p)}$ is non-singular.

It is easy to check that the elliptic curve $\mathcal{E}_{(m,p)}$ contains two rational points

$$\begin{aligned}
Q &= (X_1(m, p), 0), \\
P &= (X_2(m, p), 216p^2(p^2 + 1)(m^4 - 1)((m^2 - 1)p + (m + 1)^2)),
\end{aligned}$$

where

$$\begin{aligned}
X_1(m, p) &= 3p((m^2 - 1)^2 p^3 - 2(m^2 - 1)(3m^2 - 2m + 3)p^2 \\
&\quad + (m + 1)^4 p - 4(m^2 - 1)(3m^2 - 2m + 3)), \\
X_2(m, p) &= 3p((m^2 - 1)^2 p^3 + 2(m^2 - 1)(9m^2 + 2m + 9)p^2 \\
&\quad + (m + 1)^4 p + 4(m^2 - 1)(3m^2 + 2m + 3)).
\end{aligned}$$

By the inverse map φ^{-1} , we have

$$k = -\frac{12p(p^2 + 1)k(X, Y)}{(X - X_1(m, p))(X - X_2(m, p))},$$

where

$$\begin{aligned}
k(X, Y) &= Y + 3p((m^2 - 1)p + (m + 1)^2)(X - 3p((m^2 - 1)^2 p^3 \\
&\quad - 2(m^2 - 1)(3m^2 - 2m + 3)p^2 + (m + 1)^4 p \\
&\quad - 4(m^2 - 1)(3m^2 - 2m + 3))).
\end{aligned}$$

Similar to Case 3.2.1, we can get more parametric solutions by the group law. From the rational point $[2]P$, we get

$$k = \frac{2(p^2 + 1)p((m - 1)p + m + 1)}{(m - 1)(2(m^2 + 1)p^2 + (m + 1)^2)},$$

then

$$c = \frac{2(m^2 - 1)p^2 + 4pm + m^2 - 1}{2(m^2 + 1)p - m^2 + 1}.$$

By some calculations, we obtain

$$\begin{aligned} r &= \frac{(p^2 + 1)(2(m^2 + 1)p - m^2 + 1)}{(m^2 - 1)p^2 - (m - 1)^2p + m^2 - 1}, \\ s &= \frac{4(p^2 + 1)p((m - 1)p + m + 1)m}{(m - 1)(2(m^2 + 1)p^2 + (m + 1)^2)}, \\ t &= \frac{2(p^2 + 1)p((m^2 - 1)p + (m + 1)^2)}{2(m^2 + 1)p^2 + (m + 1)^2}, \\ a &= \frac{2(m^2 + 1)p^2 + (m + 1)^2}{(p^2 + 1)((m^2 - 1)p + (m + 1)^2)}, \\ b &= -\frac{b(m, p)}{2(m^2 - 1)p((m^2 - 1)p + (m + 1)^2)(p^2 + 1)(2(m^2 + 1)p^2 + (m + 1)^2)}, \end{aligned}$$

where

$$\begin{aligned} b(m, p) &= 4m(m^2 - 1)^2p^7 - 4(m^2 - 1)(m^4 - 2m^3 - 2m^2 - 2m + 1)p^6 \\ &\quad + 4(m + 1)^2m(3m^2 - 2m + 3)p^5 + 8m(m^2 - 1)(m^2 + 4m + 1)p^4 \\ &\quad + 4(m + 1)^2m(3m^2 + 2m + 3)p^3 + 3(m - 1)(m + 1)^5p^2 \\ &\quad + 4(m + 1)^4mp + (m - 1)(m + 1)^5, \end{aligned}$$

and $m > 1, p > 1$, this is a parametric solution of Eq. (2.1).

4. TRIANGLE AND PARALLELOGRAM PAIRS

In this section, we consider the problem that triangle and parallelogram pairs have the same area and the same perimeter.

4.1. The corresponding Diophantine system. A triangle cannot be transformed into a parallelogram by affine transformation, but the parallelogram is composed of two congruent triangles (see Figure 2), which allows us to study this problem by the affine transformation.




FIGURE 2. Rational triangle T and parallelogram L .

Let V_1, V_2 and V_3 be the three vertices of rational triangle T , and $T \in X(T)$. Let $f(V_1), f(V_2), f(V_3)$ and $f(V_4)$ be the four vertices of parallelogram L . Let $f(T)$ be a triangle with $f(V_1), f(V_2)$ and $f(V_3)$ as vertices. Let A and A' be the areas of rational triangle T and $f(T)$, respectively. Considering the affine transformation f

from T to $f(T)$ such that $A = 2A'$, then the area of parallelogram L is equal to the area of rational triangle T . The transformation matrix M_f can be taken as

$$M_f = \begin{bmatrix} \frac{1}{2a} & b \\ 0 & a \end{bmatrix},$$

where $a \in \mathbb{Q}^*$, $b \in \mathbb{Q}$. In order to ensure that rational triangle T and parallelogram L have the same perimeter, we have the following Diophantine system.

$$(4.1) \quad \begin{cases} s^2 + t^2 = w_1^2, \\ (s-r)^2 + t^2 = w_2^2, \\ \left(\frac{s}{2a} + bt\right)^2 + (at)^2 = w_3^2, \\ r + w_1 + w_2 = 2\left(\frac{r}{2a} + w_3\right), \end{cases}$$

where $b \in \mathbb{Q}$, $s \in \mathbb{Q}_{\geq 0}$, $a, r, t, w_1, w_2, w_3 \in \mathbb{Q}^*$.

4.2. Existed results. We use Eq. (4.1) to discuss the existed results, focusing on the following two cases.

Case 4.2.1. T is a right triangle and L is a rectangle.

Take $r = 2km$, $s = 0$, $t = k(m^2 - 1)$, $k > 0$, $m > 1$, then

$$w_1 = k(m^2 - 1), \quad w_2 = k(m^2 + 1).$$

Since L is a rectangle, we can take $b = 0$. Eq. (4.1) now becomes

$$(4.2) \quad 2km(m+1) = 2\left(\frac{km}{a} + ak(m^2 - 1)\right).$$

Solving Eq. (4.2), we get

$$a = \frac{m^2 + m \pm \sqrt{\Delta(a)}}{2(m^2 - 1)},$$

where

$$\Delta(a) = m^4 - 2m^3 + m^2 + 4m.$$

To get the rational solutions of Eq. (4.2), we need to make the discriminant $\Delta(a)$ be a perfect square, say w^2 . By the map

$$\begin{aligned} X &= \frac{3(m^2 + 10m - 12w + 13)}{(m-1)^2}, \\ Y &= -\frac{108(m^3 - 2m^2 + 7m + 2 - (m+3)w)}{(m-1)^3}, \end{aligned}$$

We get an elliptic curve

$$\mathcal{E} : Y^2 = X^3 - 675X + 13662.$$

The discriminant of \mathcal{E} is -3809369088 , so \mathcal{E} is nonsingular.

The inverse map is

$$\begin{aligned} m &= \frac{X^2 + 30X - 99 - 12Y}{(X+33)(X-39)}, \\ w &= -\frac{2(X^3 - 27X^2 + 2727X - 31293 - 6(X+33)Y)}{(X+33)(X-39)^2}. \end{aligned}$$

By the package of Magma, the rank of \mathcal{E} is 0, and \mathcal{E} only has degenerate solutions

$$(X, Y) = (-33, 0), (3, \pm 108), (39, \pm 216).$$

Therefore, there is no right triangle and rectangle pair with the same area and the same perimeter, this case was obtained by Guy [10].

Case 4.2.2. T is an isosceles triangle and L is a rhombus.

Case 4.2.2.1. Take $r = 4km, s = 2km, t = k(m^2 - 1), k > 0, m > 1$, then

$$w_1 = w_2 = k(m^2 + 1).$$

Since L is a rhombus, then Eq. (4.1) becomes

$$(4.3) \quad \begin{cases} \left(\frac{2km}{a}\right)^2 = \left(\frac{km}{a} + bk(m^2 - 1)\right)^2 + a^2k^2(m^2 - 1)^2, \\ 2k(m+1)^2 = \frac{8km}{a}. \end{cases}$$

Eliminating a in Eq. (4.3), we have

$$(4.4) \quad \begin{aligned} &16(m-1)^2(m+1)^4b^2 + 8(m-1)(m+1)^5b \\ &- 3m^6 - 18m^5 + 211m^4 - 572m^3 + 211m^2 - 18m - 3 = 0. \end{aligned}$$

The discriminant of Eq. (4.4) with respect to b is

$$\Delta(b) = (m^3 + 11m^2 - 5m + 1)(m^3 - 5m^2 + 11m + 1).$$

Obviously, we need to make $\Delta(b)$ be a perfect square, say w^2 . We consider the curve

$$\mathcal{C}_1 : w^2 = (m^3 + 11m^2 - 5m + 1)(m^3 - 5m^2 + 11m + 1).$$

By the package of Magma, the Mordell-Weil rank of the Jacobian variety of \mathcal{C}_1 over \mathbb{Q} is at most 1, and Magma's Chabauty routines determine the only finite rational points in \mathcal{C}_1 are

$$(m, w) = (0, \pm 1), (1, \pm 8).$$

Obviously, all points are degenerate points.

Case 4.2.2.2. Take $r = 2k(m^2 - 1), s = k(m^2 - 1), t = 2km, k > 0, m > 1$, then

$$w_1 = w_2 = k(m^2 + 1).$$

Since L is a rhombus, then Eq. (4.1) becomes

$$(4.5) \quad \begin{cases} \left(\frac{k(m^2 - 1)}{a}\right)^2 = \left(\frac{k(m^2 - 1)}{2a} + 2bkm\right)^2 + (2akm)^2, \\ 4km^2 = \frac{4k(m^2 - 1)}{a}. \end{cases}$$

Eliminating a in Eq. (4.5), we have

$$(4.6) \quad 16b^2m^4 + 8bm^5 - 3m^6 + 16m^4 - 32m^2 + 16 = 0.$$

The discriminant of Eq. (4.6) with respect to b is

$$\Delta(b) = (m^3 - 2m^2 + 2)(m^3 + 2m^2 - 2).$$

We consider the curve

$$\mathcal{C}_2 : w^2 = (m^3 - 2m^2 + 2)(m^3 + 2m^2 - 2).$$

The Mordell-Weil rank of the Jacobian variety of \mathcal{C}_2 over \mathbb{Q} is at most 1, and Magma's Chabauty routines determine the only finite rational points in \mathcal{C}_2 are

$$(m, w) = (\pm 1, \pm 1).$$

Therefore, there is no isosceles triangle and rhombus pair with the same area and the same perimeter, this case was obtained by Zhang and Peng [25].

4.3. Other solutions of Eq. (4.1). We can get more solutions by directly solving Eq. (4.1). We are concerned with the following two cases.

Case 4.3.1. $s = 0$, T is a right triangle.

Take $r = 2km$, $s = 0$, $t = k(m^2 - 1)$, $k > 0$, $m > 1$, then

$$w_1 = k(m^2 - 1), \quad w_2 = k(m^2 + 1).$$

From the third equation of Eq. (4.1), we have

$$k^2(m^2 - 1)^2(a^2 + b^2) = w_3^2,$$

then

$$a = 2cp, \quad b = c(p^2 - 1), \quad w_3 = ck(m^2 - 1)(p^2 + 1), \quad c > 0, \quad p > 1.$$

Eq. (4.1) now becomes

$$(4.7) \quad 2p(p^2 + 1)(m^2 - 1)c^2 - 2m(m + 1)pc + m = 0,$$

then

$$c = \frac{m(m + 1)p \pm \sqrt{\Delta(c)}}{2p(p^2 + 1)(m^2 - 1)},$$

where

$$\Delta(c) = p^2m^4 - 2p(p^2 - p + 1)m^3 + p^2m^2 + 2p(p^2 + 1)m.$$

To get the rational solutions of Eq. (4.7) with respect to c , it needs $\Delta(c)$ be a perfect square, say w^2 . Using a method described by Fermat (see [9, p. 639]), it's easy to get the following solutions

$$m = -\frac{(p - 1)^4}{4p(p^2 - 3p + 1)}$$

such that $\Delta(c)$ is a perfect square. By Eq. (4.7), we get

$$c = -\frac{(p - 1)^2}{(p^2 + 1)(p^2 - 4p + 1)}.$$

By some calculations, we obtain

$$s = 0, \quad r = -\frac{k(p - 1)^4}{2p(p^2 - 3p + 1)}, \quad t = \frac{k(p^2 + 2p - 1)(p^2 - 2p - 1)(p^2 - 4p + 1)^2}{16p^2(p^2 - 3p + 1)^2},$$

$$a = -\frac{2(p - 1)^2p}{(p^2 + 1)(p^2 - 4p + 1)}, \quad b = -\frac{(p - 1)^3(p + 1)}{(p^2 + 1)(p^2 - 4p + 1)},$$

where $k > 0$, $1 + \sqrt{2} < p < \frac{3 + \sqrt{5}}{2}$, this is a parametric solution of Eq. (4.1).

Let us consider the rational points on the quartic curve

$$\mathcal{C} : w^2 = \Delta(c).$$

By the map φ :

$$X = -\frac{3p(6(m^2 - 1)p^2 - p(13m^2 + 10m + 1) + 6(m^2 - 1) + 12w)}{(m - 1)^2},$$

$$Y = \frac{108pY(m, w)}{(m - 1)^3},$$

where

$$Y(m, w) = p((m - 1)(m^2 + 4m + 1)p^2 - m(m + 1)(3m + 1)p + (m - 1)(m^2 + 4m + 1)) - ((m - 1)p^2 - (3m + 1)p + m - 1)w.$$

We can transform \mathcal{C} into the elliptic curve

$$\mathcal{E}_p : Y^2 = X^3 - 27p^2(12p^4 - 12p^3 + 25p^2 - 12p + 12)X + 54p^4(12p^2 - p + 12)(6p^2 - p + 6).$$

The discriminant of \mathcal{E}_p is

$$\Delta(p) = 34012224p^6(4p^4 - 12p^3 + 9p^2 - 12p + 4)(p^2 + 1)^4.$$

Hence, $\Delta(p) \neq 0$, so \mathcal{E}_p is non-singular.

It is easy to check that the elliptic curve \mathcal{E}_p contains two rational points

$$Q = (-3p(6p^2 - p + 6), 0), P = (-3p(6p^2 - 25p + 6), -216p^2(p^2 - 3p + 1)).$$

By the inverse map φ^{-1} , we have

$$m = \frac{(X + 3p(6p^2 - p + 6))(X - 3p(6p^2 - 11p + 6)) - 12pY}{(X + 3p(6p^2 - p + 6))(X + 3p(6p^2 - 25p + 6))}.$$

When $p = 2$, we get

$$\mathcal{E}_2 : Y^2 = X^3 - 19872X + 1403136.$$

The rank of \mathcal{E}_2 is 1. From the theorem of Silverman (see p. 457 of [21]), the elliptic curve \mathcal{E}_p , for all but finitely many $p > 1$, has positive rank. Therefore, we can get more parametric solutions by the group law. From the rational point $[2]P$, we obtain

$$m = \frac{(p^2 - 4p + 1)^2}{(p^2 - 2p - 1)(p^2 + 2p - 1)},$$

then

$$c = -\frac{p^2 - 4p + 1}{4(p - 1)^2p}.$$

By some calculations, we have

$$s = 0, r = \frac{2k(p^2 - 4p + 1)^2}{(p^2 + 2p - 1)(p^2 - 2p - 1)}, t = -\frac{16k(p - 1)^4p(p^2 - 3p + 1)}{(p^2 + 2p - 1)^2(p^2 - 2p - 1)^2},$$

$$a = -\frac{p^2 - 4p + 1}{2(p - 1)^2}, b = -\frac{(p + 1)(p^2 - 4p + 1)}{4p(p - 1)},$$

where $k > 0, 1 + \sqrt{2} < p < \frac{3 + \sqrt{5}}{2}$.

Case 4.3.2. $s \neq 0$, T is a rational triangle.

Take $s = 2km, t = k(m^2 - 1), k > 0, m > 1$, then

$$w_1 = k(m^2 + 1).$$

From the second equation of Eq. (4.1), we have

$$r = \frac{2k((m^2 + 1)c - 2m)}{c^2 - 1}, \quad w_2 = \frac{k((m^2 + 1)c^2 - 4mc + m^2 + 1)}{c^2 - 1},$$

where c is a rational number.

From the third equation of Eq. (4.1), we have

$$\left(\frac{km}{a} + bk(m^2 - 1)\right)^2 + a^2k^2(m^2 - 1)^2 = w_3^2.$$

Let

$$\frac{km}{a} + bk(m^2 - 1) = d(p^2 - 1), \quad ak(m^2 - 1) = 2dp,$$

then

$$a = \frac{2dp}{k(m^2 - 1)}, \quad b = \frac{2p(p^2 - 1)d^2 - k^2m(m^2 - 1)}{2dkp(m^2 - 1)}, \quad w_3 = d(p^2 + 1).$$

Eq. (4.1) now becomes

$$(4.8) \quad \begin{aligned} & 2dp((p^2 + 1)d - k(m^2 + 1))c^2 \\ & - k(m - 1)(2p(m - 1)d - k(m + 1)(m^2 + 1))c \\ & - 2p(p^2 + 1)d^2 + 4dkmp - 2k^2m(m^2 - 1) = 0, \end{aligned}$$

then

$$c = \frac{k(2(m - 1)^2pd - (m^4 - 1)k) \pm \sqrt{\Delta(c)}}{4dp((p^2 + 1)d - k(m^2 + 1))},$$

where

$$\begin{aligned} \Delta(c) &= (m^4 - 1)^2k^4 - 4dp(m^4 - 1)(m + 1)^2k^3 \\ &+ 4d^2p(4m(m^2 - 1)p^2 + (m + 1)^4p + 4m(m^2 - 1))k^2 \\ &- 16d^3p^2(p^2 + 1)(m + 1)^2k + 16d^4p^2(p^2 + 1)^2. \end{aligned}$$

To get the rational solutions of Eq. (4.8) with respect to c , it needs $\Delta(c)$ be a perfect square, say w^2 . Since the discriminant $\Delta(c)$ is homogeneous with respect to k and d , we can let $d = 1$. Using a method described by Fermat (see [9, p. 639]), it's easy to get the following solution

$$k = \frac{p^2 + 1}{m^2 + 1}$$

such that $\Delta(c)$ is a perfect square.

Let us consider the rational points on the quartic curve

$$\mathcal{C} : w^2 = \Delta(c).$$

By the map φ :

$$\begin{aligned} X &= \frac{12pX(k, w)}{k^2}, \\ Y &= \frac{432p^2(p^2 + 1)Y(k, w)}{k^3}, \end{aligned}$$

where

$$\begin{aligned} X(k, w) &= (4m(m^2 - 1)p^2 + (m + 1)^4p + 4m(m^2 - 1))k^2 \\ &\quad - 12p(p^2 + 1)(m + 1)^2k + 24p(p^2 + 1)^2 - 6(p^2 + 1)w, \\ Y(k, w) &= (m^4 - 1)(m + 1)^2k^3 - 2(4m(m^2 - 1)p^2 + (m + 1)^4p + 4m(m^2 - 1))k^2 \\ &\quad + 12p(p^2 + 1)(m + 1)^2k - 16p(p^2 + 1)^2 - ((m + 1)^2k - 4(p^2 + 1))w. \end{aligned}$$

We can transform \mathcal{C} into the elliptic curve

$$\mathcal{E}_{(m,p)} : Y^2 = X^3 - 432p^2A_4(m,p)X - 3456p^3A_6(m,p),$$

where

$$\begin{aligned} A_4(m,p) &= 4(m^2 + 3)(3m^2 + 1)(m^2 - 1)^2p^4 - 4(m - 1)(3m^2 - 2m + 3)(m + 1)^5p^3 \\ &\quad + (25m^6 - 42m^5 + 119m^4 - 140m^3 + 119m^2 - 42m + 25)(m + 1)^2p^2 \\ &\quad - 4(m - 1)(3m^2 - 2m + 3)(m + 1)^5p + 4(m^2 + 3)(3m^2 + 1)(m^2 - 1)^2, \\ A_6(m,p) &= (2(m^2 - 1)(3m^2 - 2m + 3)p^2 - (m + 1)^4p \\ &\quad + 2(m^2 - 1)(3m^2 - 2m + 3))(8m(3m^2 + 2m + 3)(m^2 - 1)^2p^4 \\ &\quad - 4(m - 1)(3m^2 - 2m + 3)(m + 1)^5p^3 \\ &\quad + (m^6 + 54m^5 - 49m^4 + 52m^3 - 49m^2 + 54m + 1)(m + 1)^2p^2 \\ &\quad - 4(m - 1)(3m^2 - 2m + 3)(m + 1)^5p + 8m(3m^2 + 2m + 3)(m^2 - 1)^2). \end{aligned}$$

By some calculations, we can check that the discriminant $\Delta(m, p)$ of $\mathcal{E}_{(m,p)}$ is not 0, then $\mathcal{E}_{(m,p)}$ is non-singular.

It is easy to check that the elliptic curve $\mathcal{E}_{(m,p)}$ contains two rational points

$$\begin{aligned} Q &= (X_1(m, p), 0), \\ P &= (X_2(m, p), 864p^2(p^2 + 1)(m^4 - 1)(m + 1)^2), \end{aligned}$$

where

$$\begin{aligned} X_1(m, p) &= -12p(2(m^2 - 1)(3m^2 - 2m + 3)p^2 \\ &\quad - (m + 1)^4p + 2(m^2 - 1)(3m^2 - 2m + 3)), \\ X_2(m, p) &= 12p(2(m^2 - 1)(3m^2 + 2m + 3)p^2 \\ &\quad + (m + 1)^4p + 2(m^2 - 1)(3m^2 + 2m + 3)). \end{aligned}$$

By the inverse map φ^{-1} , we have

$$k = -\frac{24p(p^2 + 1)k(X, Y)}{(X - X_1(m, p))(X - X_2(m, p))},$$

where

$$\begin{aligned} k(X, Y) &= Y + 6p(m + 1)^2(X + 12p(2(m^2 - 1)(3m^2 - 2m + 3)p^2 \\ &\quad - (m + 1)^4p + 2(m^2 - 1)(3m^2 - 2m + 3))). \end{aligned}$$

By the group law, we get

$$\begin{aligned} [2]P &= (12(3(m^2 - 1)^2p^4 - 2(m^2 - 1)(3m^2 - 2m + 3)p^3 \\ &\quad + (m + 1)^2(7m^2 - 10m + 7)p^2 - 2(m^2 - 1)(3m^2 - 2m + 3)p + 3(m^2 - 1)^2), \\ &\quad - 216(p^2 + 1)^2(m^2 - 1)^2((m^2 - 1)p^2 - (3m^2 - 2m + 3)p + m^2 - 1)). \end{aligned}$$

From the rational point $[2]P$, we have

$$k = \frac{4p}{m^2 - 1},$$

then

$$c = -\frac{(m^2 - 1)p^2 + 8mp + m^2 - 1}{(m^2 - 1)p^2 - 4(m^2 + 1)p + m^2 - 1}.$$

By some calculations, we obtain

$$r = -\frac{(p^2 + 1)((m^2 - 1)p^2 - 4(m^2 + 1)p + m^2 - 1)}{(m^2 - 1)p^2 - 2(m - 1)^2p + m^2 - 1}, \quad s = \frac{8mp}{m^2 - 1}, \quad t = 4p,$$

$$a = \frac{1}{2}, \quad b = \frac{(m^2 - 1)p^2 - 8mp - m^2 + 1}{4p(m^2 - 1)},$$

where

$$1 < p < 2 + \sqrt{3}, \quad m > 1 \text{ or } p > 2 + \sqrt{3}, \quad 1 < m < \frac{\sqrt{(p^2 - 4p + 1)(p^2 + 4p + 1)}}{p^2 - 4p + 1},$$

this is a parametric solution of Eq. (4.1).

5. AREAS AND PERIMETERS ARE PROPORTIONAL

In this section, we consider the problem that rational triangle pairs or rational triangle-parallelogram pairs with areas and perimeters in fixed proportions (α, β) respectively, where α and β are positive rational numbers. The existed results can be found in [16, 15, 17]. We only give the Diophantine systems corresponding to Eq. (2.1) and Eq. (4.1), respectively.

Case 5.1. Rational triangle pairs.

The transformation matrix M_f can be taken as

$$M_f = \begin{bmatrix} \frac{\alpha}{a} & b \\ 0 & a \end{bmatrix},$$

where $a \in \mathbb{Q}^*$, $b \in \mathbb{Q}$. The corresponding Diophantine system is

$$(5.1) \quad \begin{cases} s^2 + t^2 = w_1^2, \\ (s - r)^2 + t^2 = w_2^2, \\ \left(\frac{\alpha s}{a} + bt\right)^2 + (at)^2 = w_3^2, \\ \left(\frac{\alpha(s - r)}{a} + bt\right)^2 + (at)^2 = w_4^2, \\ \frac{\alpha r}{a} + w_3 + w_4 = \beta(r + w_1 + w_2), \end{cases}$$

where $b \in \mathbb{Q}$, $s \in \mathbb{Q}_{\geq 0}$, $\alpha, \beta, a, r, t, w_1, w_2, w_3, w_4 \in \mathbb{Q}^*$.

Case 5.2. Rational triangle and parallelogram pairs.

The transformation matrix M_f can be taken as

$$M_f = \begin{bmatrix} \frac{\alpha}{2a} & b \\ 0 & a \end{bmatrix},$$

where $a \in \mathbb{Q}^*$, $b \in \mathbb{Q}$. The corresponding Diophantine system is

$$(5.2) \quad \begin{cases} s^2 + t^2 = w_1^2, \\ (s - r)^2 + t^2 = w_2^2, \\ \left(\frac{\alpha s}{2a} + bt\right)^2 + (at)^2 = w_3^2, \\ 2\left(\frac{\alpha r}{2a} + w_3\right) = \beta(r + w_1 + w_2), \end{cases}$$

where $b \in \mathbb{Q}$, $s \in \mathbb{Q}_{\geq 0}$, $\alpha, \beta, a, r, t, w_1, w_2, w_3 \in \mathbb{Q}^*$.

The solutions of Eq. (5.1) and Eq. (5.2) can be discussed similarly to Eq. (2.1) and Eq. (4.1), respectively.

REFERENCES

- [1] M. Aassila, *Some results on Heron triangles*, Elem. Math. **56** (2001), 143–146.
- [2] M. E. Bradley, *Isosceles triangles with the same perimeter and area*, Math. Teacher **73**(4) (1980), 264–266.
- [3] A. Bremner, *On Heron triangles*, Ann. Math. Inform. **33** (2006), 15–21.
- [4] A. Bremner and R. K. Guy, *Triangle-rectangle pairs with a common area and a common perimeter*, Int. J. Number Theory **2** (2006), 217–223.
- [5] S. Chern, *Integral right triangle and rhombus pairs with a common area and a common perimeter*, Forum Geom. **16** (2016), 25–27.
- [6] A. Choudhry, *Rational triangles with the same perimeter and the same area*, Hardy-Ramanujan J. **30** (2007), 19–30.
- [7] A. Choudhry and A. S. Zargar, *Pairs of equiperimeter and equiareal triangles whose sides are perfect squares*, Funct. Approx. Comment. Math. **66**(2) (2022), 245–251.
- [8] P. Das, A. Juyal and D. Moody, *Integral isosceles triangle-parallelgram and Heron triangle-rhombus pairs with a common area and common perimeter*, J. Number Theory **180** (2017), 208–218.
- [9] L. E. Dickson, *History of the Theory of Numbers, Vol. II: Diophantine Analysis*, Dover Publications, New York, 2005.
- [10] R. K. Guy, *My favorite elliptic curve: a tale of two types of triangles*, Amer. Math. Monthly **102** (1995), 771–781.
- [11] Y. Hirakawa and H. Matsumura, *A unique pair of triangles*, J. Number Theory **194** (2019), 297–302.
- [12] A. Juyal and D. Moody, *Pairs of Heron and right triangles with a common area and a common perimeter*, Publ. Math. Debrecen **100**(3-4) (2022), 449–460.
- [13] A.-V. Kramer and F. Luca, *Some remarks on Heron triangles*, Acta Acad. Paed. Agriensis, Sectio Math. **26** (2000), 1–10.
- [14] M. Lalsín and X. Ma, *θ -triangle and ω -parallelogram pairs with common area and common perimeter*, J. Number Theory. **202** (2019), 1–26.
- [15] Y. C. Li and Y. Zhang, *θ -triangle and ω -parallelogram pairs with areas and perimeters in certain proportions*, Rocky Mountain J. Math. **50**(3) (2020), 1059–1071.
- [16] Y. C. Li and Y. Zhang, *Rational triangles pairs and cyclic quadrilaterals pairs with areas and perimeters in certain proportions*, Funct. Approx. Comment. Math. **65**(1) (2021), 47–59.
- [17] Y. C. Li and Y. Zhang, *On an elliptic curve involving pairs of triangles and special quadrilaterals*, Int. J. Number Theory **18**(7) (2022), 1517–1533.
- [18] D. R. Lichtenberg, *More about triangles with the same area and the same perimeter*, Math. Teacher **67**(7) (1974), 659–660.
- [19] R. van Luijk, *An elliptic K3 surface associated to Heron triangles*, J. Number Theory **123** (2004), 92–119.
- [20] R. W. Prielipp, *Are triangles that have the same area and perimeter congruent?*, Math. Teacher **67** (1974), 157–159.
- [21] J. H. Silverman, *The Arithmetic of Elliptic Curves*, Second edition, Graduate Texts in Mathematics, 106, Springer, New York, 2009.

- [22] A. Wares, *Mutually non-congruent triangles with the same perimeter and the same area*, Inter. J. Math. Educ. **33(5)** (2002), 788–791.
- [23] P. Yiu, *Isosceles triangles equal in perimeter and area*, Missouri J. Math. Sci. **10** (1998), 106–111.
- [24] Y. Zhang, *Right triangle and parallelogram pairs with a common area and a common perimeter*, J. Number Theory **164** (2016), 179–190.
- [25] Y. Zhang and J. Y. Peng, *Heron triangle and rhombus pairs with a common area and a common perimeter*, Forum Geom. **17** (2017), 419–423.
- [26] Y. Zhang, J. Y. Peng and J. M. Wang, *Integral triangles and trapezoids pairs with a common area and a common perimeter*, Forum Geom. **18** (2018), 371–380.
- [27] Y. Zhang and A. S. Zargar, *Integral triangles and cyclic quadrilaterals pairs with a common area and a common perimeter*, to appear in Forum Geom (2019).
- [28] Y. Zhang and A. S. Zargar, *Integral triangles and perpendicular quadrilateral pairs with a common area and a common perimeter*, Funct. Approx. Comment. Math. **63(2)** (2020), 165–180.

SCHOOL OF MATHEMATICS AND STATISTICS, CHANGSHA UNIVERSITY OF SCIENCE AND TECHNOLOGY; HUNAN PROVINCIAL KEY LABORATORY OF MATHEMATICAL MODELING AND ANALYSIS IN ENGINEERING, CHANGSHA 410114, PEOPLE'S REPUBLIC OF CHINA

Email address: liyangchengmlx@163.com

SCHOOL OF MATHEMATICS AND STATISTICS, CHANGSHA UNIVERSITY OF SCIENCE AND TECHNOLOGY; HUNAN PROVINCIAL KEY LABORATORY OF MATHEMATICAL MODELING AND ANALYSIS IN ENGINEERING, CHANGSHA 410114, PEOPLE'S REPUBLIC OF CHINA

Email address: zhangyongzju@163.com